

# A bounded coarse structure for families of pseudometrics

Jesús P. Moreno-Damas

## Abstract

We define the bounded coarse structure attached to a family of pseudometrics and give some counterexamples to conjectures that arise naturally.

## 1 Introduction

We define the bounded coarse structure attached to a family of pseudometrics of a space, generalizing the one attached to a single metric. We do it in the same way as it is done in [1] for the  $C_0$  coarse structure (defined originally by Wright for a single metric in [2, 3]).

Moreover, we give some counterexamples of conjectures which arise naturally.

They are some of the conclusions of a coarse geometry's seminar with Jerzy Dydak, Manuel Alonso Morón, and Jesús P. Moreno-Damas in the Dept. of Geometría y Topología of Universidad Complutense de Madrid.

## 2 Preliminaries

A pseudometric on a set is a map  $d : X \times X \rightarrow [0, \infty)$  such that  $d(x, x) = 0$  for every  $x$ ,  $d(x, y) = d(y, x)$  for every  $x, y$  and  $d(x, z) \leq d(x, y) + d(y, z)$  for every  $x, y, z$ . If moreover  $d(x, y) = 0$  implies  $x = y$  for every  $x, y$ , then  $d$  is a metric.

## 3 The bounded coarse structure attached to a family of pseudometrics

If  $\{\mathcal{E}_i\}_{i \in I}$  is a family of coarse structures over  $X$ , then  $\bigcap_{i \in I} \mathcal{E}_i$  is a coarse structure.

If particular, if  $D$  is a family of pseudometrics of  $X$  and  $\mathcal{E}_b(d_i)$  is the bounded coarse structure of each one, then

$$\bigcap_{d \in D} \mathcal{E}_b(d) = \{E \subset X \times X : \forall d \in D \exists R_d > 0 \text{ such that } d(x, y) \leq R_d \forall (x, y) \in E\} \quad (1)$$

is a coarse structure.

**Definition 1.** Let  $D$  a family of pseudometrics of  $X$ . The bounded coarse structure attached to  $D$ , denoted by  $\mathcal{E}_b(D)$  is the one defined in (1).

**Remark 2.** Let  $X$  be a set, let  $D$  be a family of pseudometrics of  $X$ . Consider the metric on  $X$ :

$$\rho(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

and let  $D' = \{d + \rho : d \in D\}$ .

Then,  $D'$  is a family of metrics of  $X$  such that  $\mathcal{E}_b(D) = \mathcal{E}_b(D')$ .

The first natural question is if the bounded coarse structure attached to a family of pseudometrics is more general than the one attached to one metric. Example 5 will show that, in general, it doesn't happen.

**Definition 3.** Let  $(X, \mathcal{E})$  be a coarse space and let  $\{E_\lambda\} \subset \mathcal{E}$ . We say that  $\{E_\lambda\}$  generes strongly  $\mathcal{E}$  if for every  $E \in \mathcal{E}$ , there exists  $\lambda \in \Lambda$  such that  $E \subset E_\lambda$ .

**Example 4.** Let  $(X, d)$  be a pseudometric space and let, for every  $n \in \mathbb{N}$ :

$$E_n = \{(x, y) : d(x, y) \leq n\}$$

Then,  $\{E_n\}_{n=1}^\infty$  generes strongly  $(X, d)$ .

**Example 5.** For each  $n \in \mathbb{N}$ , consider in  $\mathbb{R}^\mathbb{N}$  the pseudometric  $d_n(x, y) = |x_n - y_n|$ . Let  $D = \{d_n : n \in \mathbb{N}\}$ . From the definition, we get that  $E \in \mathcal{E}_b(D)$  if and only if there are  $\{R_n\}_{n \in \mathbb{N}}$  such that  $|x_n - y_n| \leq R_n$  for every  $(x, y) \in E$  and every  $n \in \mathbb{N}$ .

There is not a pseudometric  $d$  on  $\mathbb{R}^\mathbb{N}$  such that  $\mathcal{E}_b(D) = \mathcal{E}_b(d)$ . Suppose, on the contrary, there is one. Then, by Example 4, there exists  $\{E_n\}_{n=1}^\infty \subset \mathcal{E}_b(D)$  which generes strongly  $\mathcal{E}_b(D)$ .

For each  $n$ ,  $E_n \in \mathcal{E}_b(D)$ . Let  $R_n$  be such that  $|x_i - y_i| < R_n$  for every  $(x, y) \in E_n$ . Consider the set  $E = \{(x, y) \in \mathbb{R}^\mathbb{N} : |x_n - y_n| \leq 2R_n \forall n \in \mathbb{N}\}$ . Let  $x = (0)_{n \in \mathbb{N}}$  and  $y = (2R_n)_{n \in \mathbb{N}}$ . Clearly,  $(x, y) \in E$ , but, for every  $n \in \mathbb{N}$ ,  $(x, y) \notin E_n$ . Then,  $E \not\subset E_n$  for every  $n$  and  $\{E_n\}_{n \in \mathbb{N}}$  for every  $n \in \mathbb{N}$ .

But in Example 5, in general the controlled sets are not proper, as they use to be in the usual examples of bounded coarse structures attached to metric spaces, like the one attached to a finite generated group, for example  $\mathbb{Z}$  of  $\mathbb{Z}^2$ . What happen if the coarse space  $\mathcal{E}_b(D)$  is proper? We put the question in the following terms:

**Conjecture 6.** *Suppose that  $D$  is a family of pseudometrics of  $X$  such that the topology induced by  $D$  is locally compact and the coarse structure  $\mathcal{E}_b(D)$  induced by  $D$  is proper. Then,  $(X, \mathcal{E}_b(D))$  is coarsely metrizable, i. e, there exists  $d$  such that  $\mathcal{E}_b(D) = \mathcal{E}_b(d)$ .*

The following counterexample will show that this conjecture is not true:

**Example 7.** Consider on  $\mathbb{N} \times \mathbb{N}$  the maps:

$$f_0 : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}, f_0(a, b) = 2^a$$

$$\forall n \in \mathbb{N}, f_n : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}, f_n(a, b) = \begin{cases} b & \text{if } a = n \\ 0 & \text{if } a \neq n \end{cases}$$

and, for each  $n \in \mathbb{N} \cup \{0\}$ , let  $d_n$  be the metric such that  $d_n(x, y) = |f_n(x) - f_n(y)|$ .

Let  $\mathcal{E} = \mathcal{E}_b(\{d_n\}_{n=0}^\infty)$ . Let us see that  $\mathcal{E}$  is a proper coarse structure such that there is no pseudometric  $d$  of  $X$  with  $\mathcal{E} = \mathcal{E}_b(d)$ .

Let  $M \in \mathbb{N}$  and let  $\{R_n\}_{n=1}^\infty \subset \mathbb{R}^+$ . Let  $E_{M, \{R_n\}_{n=1}^\infty}$  be the set of those  $((a, b), (a', b')) \in (\mathbb{N} \times \mathbb{N}) \times (\mathbb{N} \times \mathbb{N})$  satisfying any of this properties:

- $a, b, a', b' \leq M$
- $a = a'$  and  $|b - b'| \leq R_a$

Let us see first that

$$\{E_{M, \{R_n\}_{n=1}^\infty} : M \in \mathbb{N}, R_n \in \mathbb{R}^+ \forall n \in \mathbb{N}\}$$

generates strongly  $\mathcal{E}$ .

Fix  $M \in \mathbb{N}$  and  $\{R_n\}_{n=1}^\infty \subset \mathbb{R}^+$ . Let  $R'_0 = 2^{M+1}$  and  $R'_n = R_n + 2M$  for every  $n \in \mathbb{N}$ . Let  $((a, b), (a', b')) \in E_{M, \{R_n\}_{n=1}^\infty}$ . Suppose that  $a = a'$  and  $|b - b'| \leq R_n$ . Clearly,

$$d_a((a, b), (a', b')) = |b - b'| \leq R_a \leq R'_a$$

$$d_0((a, b), (a', b')) = |2^a - 2^a| = 0 < R'_0$$

$$d_n((a, b), (a', b')) = |0 - 0| = 0 < R_n \forall n \in \mathbb{N} \setminus \{a\}$$

Suppose now that  $a, a', b, b' \leq M$ . Then,

$$d_0((a, b), (a', b')) = |2^a - 2^{a'}| \leq 2^a + 2^{a'} \leq 2^M + 2^M = R_0$$

$$d_n((a, b), (a', b')) = |0 - 0| = 0 \forall n \in \mathbb{N} \setminus \{a, a'\}$$

If  $a = a'$ , then:

$$d_a((a, b), (a', b')) = |b - b'| \leq b + b' = 2M \leq R'_a$$

And, if  $a \neq a'$ , then:

$$d_a((a, b), (a', b')) = |b - 0| = b \leq M \leq R'_a$$

$$d_{a'}((a, b), (a', b')) = |0 - b'| = b' \leq M \leq R'_{a'}$$

Then,  $E_{M, \{R_n\}_{n=1}^\infty} \in \mathcal{E}$

Suppose now that  $E \in \mathcal{E}$ . Let  $\{R_n\}_{n=0}^\infty \subset \mathbb{R}^+$  such that  $d_n((a, b), (a', b')) \leq R_n$  for every  $((a, b), (a', b')) \in E$  and every  $n \in \mathbb{N} \cup \{0\}$ . Let  $M_0 \in \mathbb{N}$  be such that  $M_0 \geq R_0$  and let  $M \in \mathbb{N}$  be such that  $M \geq M_0$  and  $M \geq R_n$  for every  $n \leq M_0$ . Let  $((a, b), (a', b')) \in E$ .

If  $a = a'$ , then  $|b - b'| = |f_a(a, b) - f_a(a', b')| \leq R_a$  and  $((a, b), (a', b')) \in E_{M, \{R_n\}_{n=1}^\infty}$ .

Suppose now that  $a \neq a'$ . If  $a > a'$ , then  $d_0((a, b), (a', b')) = |2^a - 2^{a'}| \geq |2^a - 2^{a-1}| = 2^{a-1} \geq a > a'$ . By symmetry, if  $a < a'$ , then  $d_0((a, b), (a', b')) \geq a' > a$ . Hence:

$$a, a' \leq d_0((a, b), (a', b')) \leq R_0 \leq M_0 \leq M$$

Since  $a, a' \leq M_0$ , we have:

$$M \geq R_a \geq d_a((a, b), (a', b')) = |f_a(a, b) - f_a(a', b')| = |b - 0| = b$$

$$M \geq R_{a'} \geq d_{a'}((a, b), (a', b')) = |f_{a'}(a, b) - f_{a'}(a', b')| = |0 - b'| = b'$$

Then,  $a, a', b, b' \leq M$  and  $((a, b), (a', b')) \in E_{M, \{R_n\}_{n=1}^\infty}$ . Thus,  $E \subset E_{M, \{R_n\}_{n=1}^\infty}$ .

Let us see that  $\mathcal{E}$  is a proper coarse structure. Obviously, the diagonal is an open neighborhood of the diagonal. Let  $E \in \mathcal{E}$  be symmetric. To see that  $E$  is proper, it is enough to see that  $E_{(a', b')}$  is finite for every  $(a', b') \in \mathbb{N} \times \mathbb{N}$ .

Let  $M \in \mathbb{N}$  and  $\{R_n\}_{n=0}^\infty$  such that  $E \subset E_{M, \{R_n\}_{n=1}^\infty}$ . Let  $(a, b) \in E_{(a', b')}$ . Then,  $((a, b), (a', b')) \in E_{M, \{R_n\}_{n=1}^\infty}$ . If  $a, a', b, b' \leq M$ , then  $(a, b) \in \{(x, y) : x, y \leq M\}$ . If  $a = a'$  and  $|b - b'| \leq R_{a'}$ , then  $(a, b) \in \{(x, y) : x = a', b' - R_{a'} \leq x \leq b' - R_{a'}\}$ . Then,

$$E_{(a', b')} \subset \{(x, y) : x, y \leq M\} \cup \{(x, y) : x = a', b' - R_{a'} \leq x \leq b' - R_{a'}\}$$

which is finite. Then,  $E$  is proper and  $\mathcal{E}$  is proper.

Suppose that  $d$  is a pseudometric on  $X$  such that  $\mathcal{E} = \mathcal{E}_b(d)$ . Then, by Example 4, there is  $\{E_m\}_{m \in \mathbb{N}} \subset \mathcal{E}$  generating strongly  $\mathcal{E}$ .

For every  $m$ , let  $M^m \in \mathbb{N}$  and  $\{R_n^m\}_{n=1}^\infty$  be such that  $E_m \subset E_{M^m, \{R_n^m\}_{n=1}^\infty}$ .

For every  $n \in \mathbb{N}$ , let  $R_n = R_n^m + 1$ . Let  $E = E_{1, \{R_n\}_{n=1}^\infty}$ .

For every  $m \in \mathbb{N}$ , consider the point  $x_m = ((m, R_m^m + M_m + 2), (m, M_m + 1))$ .

We have that  $m = m$  and  $|R_m^m + M_m + 2 - (M_m + 1)| = R_m^m + 1 = R_m$ , hence  $x_m \in E$ .

In the other hand,  $m, R_m^m + M_m + 2, m, M_m + 1 \not\leq M_m$  and  $m = m$  but  $|R_m^m + M_m + 2 - (M_m + 1)| = R_m^m + 1 \not\leq R_m^m$ , hence  $x_m \notin E_m$ .

Then,  $E \not\subset E_m$  for every  $m$  and hence, there is no  $d$  such that  $\mathcal{E} = \mathcal{E}_b(d)$ .

## References

- [1] Moreno-Damas, Jesús P. *A  $C_0$  coarse structure for families of pseudometrics and the Higson-Roe functor* arXiv:1410.2756v1 [math.GN].
- [2] Nick Wright,  *$C_0$  Coarse Geometry*. PhD Thesis.
- [3] Wright, Nick  *$C_0$  coarse geometry and scalar curvature*. J. Funct. Anal. 197 (2003), no. 2, 469-488.