

THE ASYMPTOTIC BEHAVIOR OF THE REIDEMEISTER TORSION FOR SEIFERT MANIFOLDS AND $\mathrm{PSL}_2(\mathbb{R})$ -REPRESENTATIONS OF FUCHSIAN GROUPS

YOSHIKAZU YAMAGUCHI

ABSTRACT. We show that a $\mathrm{PSL}_2(\mathbb{R})$ -representation of a Fuchsian group induces the asymptotics of the Reidemeister torsion for the Seifert manifold corresponding to the euler class of the $\mathrm{PSL}_2(\mathbb{R})$ -representation. We also show that the limit of leading coefficient of the Reidemeister torsion is determined by the euler class of a $\mathrm{PSL}_2(\mathbb{R})$ -representation of a Fuchsian group. In particular, the leading coefficient of the Reidemeister torsion for the unit tangent bundle over a two-orbifold converges to $-\chi \log 2$ where χ is the Euler characteristic of the two-orbifold. We also give a relation between \mathbb{Z}_2 -extensions for $\mathrm{PSL}_2(\mathbb{R})$ -representations of a Fuchsian group and the asymptotics of the Reidemeister torsion.

1. INTRODUCTION

The previous works [Yam13, Yam] of the author have investigated the asymptotic behavior of the Reidemeister torsion for a Seifert manifold with an sequence of $\mathrm{SL}_{2N}(\mathbb{C})$ -representation of the fundamental group. Here the sequence of $\mathrm{SL}_{2N}(\mathbb{C})$ -representations starts with an $\mathrm{SL}_2(\mathbb{C})$ -representation and the remaining representations are given by the composition with the irreducible $2N$ -dimensional representations of $\mathrm{SL}_2(\mathbb{C})$. A sequence of $\mathrm{SL}_{2N}(\mathbb{C})$ -representations defines a sequence of the Reidemeister torsion of a Seifert manifold. We can consider the asymptotic behavior of the sequence given by the Reidemeister torsions. Under the constraint that $\mathrm{SL}_{2N}(\mathbb{C})$ -representations send a regular fiber to $-I_{2N}$, the observation in [Yam] revealed the growth order and the convergence of the leading coefficient of the Reidemeister torsion for a Seifert manifold.

In this paper, we observe when we have a natural situation for a Seifert manifold and $\mathrm{SL}_2(\mathbb{R})$ -representations satisfying our constraint. We will deal with closed Seifert manifolds whose base orbifolds have the negative Euler characteristics. Namely, they admit the $\mathbb{H}^2 \times \mathbb{R}$ or $\widetilde{\mathrm{SL}}_2(\mathbb{R})$ -geometry. We can regard the fundamental group of the base orbifold as a cocompact Fuchsian group Γ which can be embedded in $\mathrm{PSL}_2(\mathbb{R})$. The fundamental group of a Seifert manifold is a central extension of the Fuchsian group Γ by \mathbb{Z} . The universal cover $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ is also a central extension of $\mathrm{PSL}_2(\mathbb{R})$ by \mathbb{Z} . If we choose a homomorphism $\bar{\rho}$ from Γ to $\mathrm{PSL}_2(\mathbb{R})$ for a Seifert manifold M , then we have a lift $\tilde{\rho}$ from $\pi_1(M)$ to $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$. This is due to the work [JN85] by M. Jankins and W. Neumann (we review this in Section 2.2). Since $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ is also the universal cover of $\mathrm{SL}_2(\mathbb{R})$, we have the $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M)$ given by the composition of $\tilde{\rho}$ with the projection from $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ onto $\mathrm{SL}_2(\mathbb{R})$. The $\mathrm{SL}_2(\mathbb{R})$ -representation ρ induces the $\mathrm{SL}_{2N}(\mathbb{R})$ -representations which define the acyclic chain complexes of the Seifert manifold M . We can observe the

2010 *Mathematics Subject Classification.* Primary: 57M27, 57M05, Secondary: 57M50.

Key words and phrases. Seifert fibered spaces; geometric structure; Reidemeister torsion; asymptotic behaviors.

asymptotics of the Reidemeister torsion starting with a $\mathrm{PSL}_2(\mathbb{R})$ -representation of a Fuchsian group.

We refer the case of a Seifert homology sphere and $\mathrm{SU}(2)$ -representations to the previous work [Yam], in which the maximum and minimum of the limits have been observed in detail. The limit of the leading coefficient is determined by the images of exceptional fibers by a given $\mathrm{SU}(2)$ -representation.

First, we will show that the limit of leading coefficient in the asymptotic behavior of the Reidemeister torsions is determined by a representation of a Fuchsian group. Here we start with a $\mathrm{PSL}_2(\mathbb{R})$ -representation of the Fuchsian group. Let $\bar{\rho}$ be a $\mathrm{PSL}_2(\mathbb{R})$ -representation of a cocompact Fuchsian group $\Gamma = \Gamma(g; \alpha_1, \dots, \alpha_n)$ (for the notation, see Section 2.2). If we have the following diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} & \rightarrow & \pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})) & \rightarrow & \Gamma & \rightarrow & 1 \\ & & \parallel & & \downarrow \bar{\rho} & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\mathrm{PSL}}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) & \rightarrow & 1, \end{array}$$

then we have the $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$ which induces a sequence of the Reidemeister torsion $\mathrm{Tor}(M; \rho_{2N})$ (for the details on this sequence, we refer to Theorem 2.1 and Lemma 3.3). The sequence of $\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|$ has the growth order of $2N$ and the following leading coefficient.

Theorem (Theorem 3.5). *For the induced $\mathrm{SL}_{2N}(\mathbb{R})$ -representations ρ_{2N} , the limit of the leading coefficient of $\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|$ is expressed as*

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|}{2N} = -\left(2 - 2g - \sum_{j=1}^n \frac{\lambda_j - 1}{\lambda_j}\right) \log 2.$$

Here λ_j is the order of $\bar{\rho}(q_j)$ where q_j is the homotopy class of a loop around the j -th branched point on the two-orbifold.

Remark. The existence of a lift $\tilde{\rho}$ is determined by the euler class $e(\bar{\rho})$ in $H^2(\Gamma; \mathbb{Z})$. The explicit value of λ_j is given in Theorem 3.5. The limit of the leading coefficient is also determined by the euler class $e(\bar{\rho})$.

The unit tangent bundle over a two-orbifold \mathbb{H}^2/Γ is also a Seifert manifold with the index $((1, 2g - 2), (\alpha_1, \alpha_1 - 1), \dots, (\alpha_n, \alpha_n - 1))$. We can think of the unit tangent bundle $T^1(\mathbb{H}^2/\Gamma)$ as $\mathrm{PSL}_2(\mathbb{R})/\Gamma$. For a Seifert manifold $\mathrm{PSL}_2(\mathbb{R})/\Gamma$, we can take a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\tilde{\rho}$ of $\pi_1(\mathrm{PSL}_2(\mathbb{R})/\Gamma)$ as a lift of an embedding from Γ into $\mathrm{PSL}_2(\mathbb{R})$.

Corollary (Corollary 3.7). *Suppose that an $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(\mathrm{PSL}_2(\mathbb{R})/\Gamma)$ is the composition of $\tilde{\rho}$ with the projection from $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ onto $\mathrm{SL}_2(\mathbb{R})$. Then we have the following limit of the leading coefficient of $\log |\mathrm{Tor}(\mathrm{PSL}_2(\mathbb{R})/\Gamma; \rho_{2N})|$:*

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(\mathrm{PSL}_2(\mathbb{R})/\Gamma; \rho_{2N})|}{2N} &= -\left(2 - 2g - \sum_{j=1}^n \frac{\alpha_j - 1}{\alpha_j}\right) \log 2 \\ &= -\chi \log 2 \end{aligned}$$

where χ is the Euler characteristic of the base orbifold \mathbb{H}^2/Γ .

We also investigate which $\mathrm{PSL}_2(\mathbb{R})$ -representation $\bar{\rho}$ of Γ induces $\mathrm{SL}_2(\mathbb{R})$ -representation of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$. We show a sufficient condition for $\bar{\rho}$ to be lifted to an $\mathrm{SL}_2(\mathbb{R})$ -representation for a given Seifert manifold $M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})$ in the following diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} & \rightarrow & \pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})) & \rightarrow & \Gamma & \rightarrow & 1 \\ & & \downarrow & & \downarrow \rho & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \mathrm{SL}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) & \rightarrow & 1. \end{array}$$

Our sufficient condition is given in terms of the euler class of $\bar{\rho}$.

Theorem (Theorem 4.3). *A $\mathrm{PSL}_2(\mathbb{R})$ -representation $\bar{\rho}$ of Γ induces an $\mathrm{SL}_2(\mathbb{R})$ -one of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$ such that $\rho(h) = -I$ if the euler class $e(\bar{\rho})$ satisfies the criteria of Theorem 2.2 and gives the equivalent class $[bx_0 + \beta_1 x_1 + \dots + \beta_n x_n]$ in $\mathrm{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$.*

Moreover the limit of the leading coefficient for $\mathrm{Tor}((; M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})$ is expressed in Theorem 4.6. We will compute explicit examples for $\mathrm{SL}_2(\mathbb{R})$ -representations of Brieskorn manifolds by using Theorem 4.3

Organization. We review the previous result on the asymptotic behavior of the Reidemeister torsion in Section 2.1. Section 2.2 gives a brief review the work on $\mathrm{PSL}_2(\mathbb{R})$ -representations of a cocompact Fuchsian group by Jankins and Neumann [JN85]. In Section 3, We observe the asymptotic behavior of the Reidemeister torsion for a Seifert manifold and the $\mathrm{SL}_2(\mathbb{R})$ -representation induced by a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -one. We deal with $\mathrm{SL}_2(\mathbb{R})$ -representations for a Seifert manifold, which are given by $\mathrm{PSL}_2(\mathbb{R})$ -representations of a Fuchsian group with different euler classes in Section 4. The last Section 5 shows explicit examples of $\mathrm{SL}_2(\mathbb{R})$ -representations induced by $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -ones for Brieskorn manifolds.

2. PRELIMINARIES

For a Seifert index $(g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$, we follow the convention of Jankins and Neumann [JN85]. This notation differs from that of [Yam] in sign of b and β_1, \dots, β_n .

2.1. Asymptotic behavior of the Reidemeister torsion. Let M be a Seifert manifold of the index $(g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$. The fundamental group of M is expressed as

$$\pi_1(M) = \langle a_1, b_1, \dots, a_g, b_g, q_1, \dots, q_n, h \mid h \text{ is central, } q_j^{\alpha_j} = h^{\beta_j}, q_1 \cdots q_n \prod_{i=1}^g [a_i, b_i] = h^{-b} \rangle.$$

We use the symbol ρ for an $\mathrm{SL}_2(\mathbb{C})$ -representation of $\pi_1(M)$. We denote by ρ_{2N} the composition of ρ with the irreducible $2N$ -dimensional representation of $\mathrm{SL}_2(\mathbb{C})$ and by $\mathrm{Tor}(M; \rho_{2N})$ the Reidemeister torsion of M and ρ_{2N} . The asymptotic behavior of the Reidemeister torsion is expressed as follows.

Theorem 2.1 (Theorem 4.5 in [Yam]). *If $\pi_1(M)$ has an $\mathrm{SL}_2(\mathbb{C})$ -representation ρ sending the homotopy class h of a regular fiber to $-I$, the asymptotic behavior of the sequence given by the Reidemeister torsion $\mathrm{Tor}(M; \rho_{2N})$ is expressed as*

$$(1) \quad \begin{aligned} \lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M; \rho_{2N})|}{(2N)^2} &= 0, \\ \lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M; \rho_{2N})|}{2N} &= -\left(2 - 2g - \sum_{j=1}^n \frac{\lambda_j - 1}{\lambda_j}\right) \log 2. \end{aligned}$$

The right hand side of (1) is determined by the orders of the $\mathrm{SL}_2(\mathbb{C})$ -matrices for the exceptional fibers. When we denote by ℓ_j the homotopy class of j -th exceptional fiber, each λ_j is half the order of $\rho(\ell_j)$.

2.2. $\mathrm{PSL}_2(\mathbb{R})$ -representations of Fuchsian groups and the euler classes. We use the symbol $\Gamma = \Gamma(g; \alpha_1, \dots, \alpha_n)$ for a cocompact Fuchsian group of genus g with branch indices $\alpha_1, \dots, \alpha_n$. The Fuchsian group Γ has the following presentation:

$$(2) \quad \Gamma = \langle a_1, b_1, \dots, a_g, b_g, q_1, \dots, q_n \mid q_j^{\alpha_j} = 1, q_1 \cdots q_n \prod_{i=1}^g [a_i, b_i] = 1 \rangle.$$

In [JN85], Jankins and Neumann determined the set of components in $\mathrm{Hom}(\Gamma, \mathrm{PSL}_2(\mathbb{R}))$ by an euler class:

$$e : \mathrm{Hom}(\Gamma, \mathrm{PSL}_2(\mathbb{R})) \rightarrow H^2(\Gamma; \mathbb{Z}) = ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle$$

The Euler class e is defined as follows. Taking the pull-back central extension for $f \in \mathrm{Hom}(\Gamma, \mathrm{PSL}_2(\mathbb{R}))$ from

$$\begin{array}{ccccccc} & & & & \Gamma & & \\ & & & & \downarrow f & & \\ 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\mathrm{PSL}}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) \rightarrow 1, \end{array}$$

we have the following commutative diagram:

$$(3) \quad \begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\Gamma} & \rightarrow & \Gamma \rightarrow 1 \\ & & \parallel & & \downarrow \tilde{f} & & \downarrow f \\ 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\mathrm{PSL}}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) \rightarrow 1. \end{array}$$

The central extension $\widetilde{\Gamma}$ has a presentation:

$$\langle a_1, b_1, \dots, a_g, b_g, q_1, \dots, q_n, h \mid h \text{ is central, } q_j^{\alpha_j} = h^{\beta_j}, q_1 \cdots q_n \prod_{i=1}^g [a_i, b_i] = h^{-b} \rangle$$

with $0 \leq \beta_i < \alpha_i$. By the presentation of $\widetilde{\Gamma}$, we define $e(f)$ to be $bx_0 + \beta_1 x_1 + \cdots + \beta_n x_n$.

Theorem 2.2 (Theorem 1 in [JN85]). *Suppose that $x \in H^2(\Gamma; \mathbb{Z})$ satisfies $x = bx_0 + \beta_1 x_1 + \cdots + \beta_n x_n$ ($0 \leq \beta_i < \alpha_i$). There exists some f such that $e(f) = x$ if and only if the following holds:*

- (i) *If $g > 0$ then $2 - 2g - n \leq b \leq 2g - 2$;*
- (ii) *If $g = 0$ then either*
 - (a) *$2 - n \leq b \leq -2$ or;*
 - (b) *$b = -1$ and $\sum_{j=1}^n (\beta_j / \alpha_j) \leq 1$ or;*
 - (c) *$b = 1 - n$ and $\sum_{j=1}^n (\beta_j / \alpha_j) \geq n - 1$.*

Remark 2.3. The above definition of the euler class e is the alternative one used in the proof of [JN85, Theorem 1]. The euler class e is defined as $e(f) = f^*(c)$ where $f^* : H^2(\mathrm{BPSL}_2(\mathbb{R}); \mathbb{Z}) \rightarrow H^2(\mathrm{B}\Gamma; \mathbb{Z}) = H^2(\Gamma; \mathbb{Z})$ is the induced homomorphism by f and c is a generator of $H^2(\mathrm{BPSL}_2(\mathbb{R}); \mathbb{Z}) \simeq \mathbb{Z}$.

Jankins and Neumann also proved the following theorem on the components of the $\mathrm{PSL}_2(\mathbb{R})$ -representation space of a Fuchsian group.

Theorem 2.4 (Theorem 2 in [JN85]). *Let Γ be a cocompact Fuchsian group. The fibers of e are the components of $\mathrm{Hom}(\Gamma, \mathrm{PSL}_2(\mathbb{R}))$.*

3. ASYMPTOTICS OF THE REIDEMEISTER TORSION VIA $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -REPRESENTATIONS

In the observation of the asymptotic behavior for the Reidemeister torsion, we require that an $\mathrm{SL}_2(\mathbb{C})$ -representation ρ sends the central element h to $-I$. Since $\pi_1(M)$ is a central extension of Γ , this is equivalent to that we have the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} & \rightarrow & \pi_1(M) & \rightarrow & \Gamma & \rightarrow & 1 \\ & & \downarrow & & \downarrow \rho & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \{\pm I\} & \rightarrow & \mathrm{SL}_2(\mathbb{C}) & \rightarrow & \mathrm{PSL}_2(\mathbb{C}) & \rightarrow & 1. \end{array}$$

The limit (1) in Theorem 2.1 is determined by the induced $\mathrm{PSL}_2(\mathbb{C})$ -representation $\bar{\rho}$ of Γ .

Proposition 3.1. *The integer λ_j in the equation (1) coincides with the order of $\bar{\rho}(q_j)$.*

Proof. The order of $\rho(\ell_j)$ is equal to $2\lambda_j$. This means that λ_j is the minimum of natural numbers such that $\rho(\ell_j)^\lambda = -I$. On the other hand, the order of $\bar{\rho}(\ell_j)$ in $\mathrm{PSL}_2(\mathbb{C})$ is the minimum of natural numbers such that $\rho(\ell_j) = \pm I$. Hence the order of $\bar{\rho}(\ell_j)$ is equal to λ_j . \square

We can rewrite the statement of Theorem 2.1 in terms of $\bar{\rho}$.

Corollary 3.2. *If a Seifert manifold M has an $\mathrm{SL}_2(\mathbb{C})$ -representation ρ such that $\rho(h) = -I$, then we have*

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M; \rho_{2N})|}{2N} = -\left(2 - 2g - \sum_{j=1}^n \frac{\lambda_j - 1}{\lambda_j}\right) \log 2.$$

where λ_j is the order of $\bar{\rho}(q_j)$ in $\mathrm{PSL}_2(\mathbb{C})$.

It is natural to try to start with a $\mathrm{PSL}_2(\mathbb{C})$ -representation of Γ . Here and subsequently, we focus our attention on $\mathrm{PSL}_2(\mathbb{R})$ -representations of a Fuchsian group and the induced $\mathrm{SL}_2(\mathbb{R})$ -representation of $\pi_1(M)$.

Given a Fuchsian group $\Gamma = \Gamma(g; \alpha_1, \dots, \alpha_n)$ and a $\mathrm{PSL}_2(\mathbb{R})$ -representation $\bar{\rho}$, we have a Seifert manifold M and a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\bar{\rho}$ of $\pi_1(M)$, induced from the diagram:

$$(4) \quad \begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} = \langle h \rangle & \rightarrow & \pi_1(M) & \rightarrow & \Gamma & \rightarrow & 1 \\ & & \parallel & & \downarrow \bar{\rho} & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\mathrm{PSL}}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) & \rightarrow & 1. \end{array}$$

The Seifert index $(g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$ of M is given by the euler class $e(f)$.

We also have the $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M)$ defined by the composition with the projection $\widetilde{\mathrm{PSL}}_2(\mathbb{R}) \rightarrow \mathrm{SL}_2(\mathbb{R})$. Here we identify the universal cover $\widetilde{\mathrm{SL}}_2(\mathbb{R})$ with $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$.

Lemma 3.3. *Suppose that $(b; \beta_1, \dots, \beta_n)$ satisfies the criteria of Theorem 2.2. Then there exist $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representations $\bar{\rho}$ of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$ such that $\bar{\rho}(h) = \mathrm{sh}(1)$.*

Remark 3.4. Here for any $\gamma \in \mathbb{R}$ we write $\mathrm{sh}(\gamma)$ for the shift by γ , that is the self-homeomorphism of \mathbb{R} , $r \mapsto r + \gamma$. We can consider $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ as a subgroup of the group of homeomorphisms $f : \mathbb{R} \rightarrow \mathbb{R}$ which are lifts of homeomorphisms of the circle. The shift $\mathrm{sh}(\gamma)$ is an element in $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$, which projects to $\begin{pmatrix} \cos(2\pi\gamma) & -\sin(2\pi\gamma) \\ \sin(2\pi\gamma) & \cos(2\pi\gamma) \end{pmatrix}$ in $\mathrm{PSL}_2(\mathbb{R})$

by the projection. The center of $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ is $\{\mathrm{sh}(k) \mid k \in \mathbb{Z}\} \simeq \mathbb{Z}$.

Proof. This is a consequence of Theorems 2.2 and 2.4. \square

Theorem 3.5. *Suppose that $(b; \beta_1, \dots, \beta_n)$ satisfies the criterion of Theorem 2.2. If $\bar{\rho}$ is a $\mathrm{PSL}_2(\mathbb{R})$ -representation of Γ in $e^{-1}(bx_0 + \beta_1 x_1 + \dots + \beta_n x_n)$ and ρ is the induced $\mathrm{SL}_2(\mathbb{R})$ -representation of $M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})$ by $\bar{\rho}$, then λ_j in the asymptotic behavior (1) is expressed as*

$$\lambda_j = \frac{\alpha_j}{(\alpha_j, \beta_j)}$$

where (α_j, β_j) is the greatest common divisor of α_j and β_j .

If all (α_j, β_j) are equal to 1, then we have

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|}{2N} = -(2 - 2g - \sum_{j=1}^n \frac{\alpha_j - 1}{\alpha_j}) \log 2 = -\chi \log 2$$

where χ is the Euler characteristic of the base orbifold.

Remark 3.6. If M is the unit tangent bundle over a two-orbifold \mathbb{H}^2/Γ whose fundamental group is embedded as a Fuchsian group Γ in $\mathrm{PSL}_2(\mathbb{R})$, i.e., $M = \mathrm{T}^1\mathbb{H}^2/\Gamma (= \mathrm{PSL}_2(\mathbb{R})/\Gamma)$, then M is the Seifert manifold with the index of $(g; (2g - 2), (\alpha_1, \alpha_1 - 1), \dots, (\alpha_n, \alpha_n - 1))$ and there exists a lift $\tilde{\rho}$ of an embedding $\bar{\rho}$ of Γ into $\mathrm{PSL}_2(\mathbb{R})$ as follows:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z} & \rightarrow & \pi_1(M) & \rightarrow & \Gamma & \rightarrow & 1 \\ & & \parallel & & \downarrow \tilde{\rho} & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \mathbb{Z} & \rightarrow & \widetilde{\mathrm{PSL}}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) & \rightarrow & 1. \end{array}$$

We can regard this $\tilde{\rho}$ as an embedding of $\pi_1(M)$ into $\mathrm{Isom}^+(\widetilde{\mathrm{PSL}}_2(\mathbb{R}))$ since $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ is a subgroup of $\mathrm{Isom}^+(\widetilde{\mathrm{PSL}}_2(\mathbb{R}))$. For more details, we refer to [Sco83].

Corollary 3.7. *Suppose that M is the quotient of $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ by a Fuchsian group Γ and $\tilde{\rho}$ is an embedding of $\pi_1(M)$ into $\widetilde{\mathrm{PSL}}_2(\mathbb{R}) (\subset \mathrm{Isom}^+(\widetilde{\mathrm{PSL}}_2(\mathbb{R})))$. For the $\mathrm{SL}_2(\mathbb{R})$ -representation ρ induced by $\tilde{\rho}$, the asymptotic behavior of the Reidemeister torsion is expressed as*

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|}{2N} = -\chi \log 2$$

Proof. It follows from Euclidean algorithm that $(\alpha_i, \alpha_i - 1) = 1$. Together with Theorem 3.5, we obtain the limit in our claim. \square

Remark 3.8. The Seifert manifold $\mathrm{PSL}_2(\mathbb{R})/\Gamma$ is also regarded as $\widetilde{\mathrm{PSL}}_2(\mathbb{R})/p^{-1}(\Gamma)$ where p is the projection from $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ onto $\mathrm{PSL}_2(\mathbb{R})$.

The next lemma was shown in the proof of [JN85, Theorem 1].

Lemma 3.9. *Let $\bar{\rho}$ be a $\mathrm{PSL}_2(\mathbb{R})$ -representation of Γ in $e^{-1}(bx_0 + \beta_1 x_1 + \dots + \beta_n x_n)$. The induced $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\tilde{\rho}$ of $M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})$ satisfies that every $\tilde{\rho}(q_j)$ is conjugate to $\mathrm{sh}(\beta_j/\alpha_j)$ in $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ for $j = 1, \dots, n$.*

We enclose this section with the proof of Theorem 3.5.

Proof of Theorem 3.5. Our $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$ is given by the $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\tilde{\rho}$ as follows:

$$\rho : \pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})) \xrightarrow{\tilde{\rho}} \widetilde{\mathrm{PSL}}_2(\mathbb{R}) \rightarrow \mathrm{SL}_2(\mathbb{R}).$$

By Lemmas 3.3 and 3.9, we can see that

$$\rho(q_j) \stackrel{\text{conj.}}{\sim} \begin{pmatrix} \cos \pi \beta_j / \alpha_j & -\sin \pi \beta_j / \alpha_j \\ \sin \pi \beta_j / \alpha_j & \cos \pi \beta_j / \alpha_j \end{pmatrix}.$$

Hence the order of $\bar{\rho}(g_j)$ in $\mathrm{PSL}_2(\mathbb{R})$ is equal to $\alpha_j/(\alpha_j, \beta_j)$. Together with Proposition 3.1, we can obtain that $\lambda_j = \alpha_j/(\alpha_j, \beta_j)$. \square

4. \mathbb{Z}_2 -EXTENSIONS OF A FUCHSIAN GROUP AND $\mathrm{SL}_2(\mathbb{R})$ -REPRESENTATIONS

Let Γ be a Fuchsian group of genus g with branch indices $\alpha_1, \dots, \alpha_n$ and M be a Seifert manifold with the index $(g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$. Theorem 2.2 shows that when the integers $(b, \beta_1, \dots, \beta_n)$ corresponds to some euler class $e(\bar{\rho})$ in $H^2(\Gamma; \mathbb{Z})$, we have a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\tilde{\rho}$ of $\pi_1(M)$ induced by $\bar{\rho}$. Then $\tilde{\rho}$ also induces an $\mathrm{SL}_2(\mathbb{R})$ -representation ρ such that $\rho(h) = -I$.

We can also find $\mathrm{SL}_2(\mathbb{R})$ -representations of $\pi_1(M)$ induced by other $(b', \beta'_1, \dots, \beta'_n)$ in $H^2(\Gamma; \mathbb{Z})$. We classify $(b', \beta'_1, \dots, \beta'_n)$ which gives an $\mathrm{SL}_2(\mathbb{R})$ -representation of $\pi_1(M)$.

4.1. \mathbb{Z}_2 -extension and $\mathrm{SL}_2(\mathbb{R})$ -representation. Every irreducible $\mathrm{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M)$ factors through $\pi_1(M)/\langle h^2 \rangle$ since ρ sends the central element h into the center of $\mathrm{SL}_2(\mathbb{R})$. When we regard Γ as $\pi_1(M)/\langle h \rangle$, we have the following central extension of Γ :

$$0 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow \pi_1(M)/\langle h^2 \rangle \rightarrow \Gamma \rightarrow 1.$$

Taking the pull-back central extension from

$$\begin{array}{ccccccc} & & & & \Gamma & & \\ & & & & \downarrow \bar{\rho} & & \\ 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \mathrm{SL}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) \rightarrow 1, \end{array}$$

we have the following diagram:

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \hat{\Gamma} & \rightarrow & \Gamma \rightarrow 1 \\ & & \parallel & & \downarrow \rho & & \downarrow \bar{\rho} \\ 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \mathrm{SL}_2(\mathbb{R}) & \rightarrow & \mathrm{PSL}_2(\mathbb{R}) \rightarrow 1. \end{array}$$

Here $\hat{\Gamma}$ is isomorphic to $\pi_1(M)/\langle h^2 \rangle$ for some M .

Lemma 4.1. *The group of central extensions of Γ by $\mathbb{Z}/2\mathbb{Z}$ is expressed as*

$$\mathrm{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z}) \simeq \mathrm{Ext}(\Gamma; \mathbb{Z}) / 2 \mathrm{Ext}(\Gamma; \mathbb{Z}).$$

Remark 4.2. Lemma 4.1 is a consequence of the remark following the proof of [NJ81, Theorem 10.4].

Proof. In proving the surjectivity of the following homomorphism:

$$H^2(\Gamma; \mathbb{Z}) = \mathrm{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z}) \rightarrow ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle,$$

we define a function

$$\nu : ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle \rightarrow S(\mathbb{Z}) = \{\text{subgroups of } \mathbb{Z}\}$$

by $[s_0, \dots, s_n] \mapsto \ker(\mathbb{Z} \rightarrow \pi(s_0, \dots, s_n))$. Here $\pi(s_0, \dots, s_n)$ is a central extension of Γ given by $\pi_1(M(g; (1, s_0), (\alpha_1, s_1), \dots, (\alpha_n, s_n)))$. The function ν gives a central extension of Γ by

$$1 \rightarrow \mathbb{Z}/\nu([s_0, \dots, s_n]) \rightarrow \pi(s_0, \dots, s_n) \rightarrow \Gamma \rightarrow 1$$

for each $[s_0, \dots, s_n]$. It is shown in [NJ81, the proof of Theorem 10.4] that

$$(5) \quad \mathrm{Ext}(\Gamma; \mathbb{Z}) \simeq \nu^{-1}(\{0\}), \quad \nu^{-1}(\{0\}) = ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle.$$

In the case of $\mathrm{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$, we also define a function

$$\nu' : ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle \rightarrow S(\mathbb{Z}) = \{\text{subgroups of } \mathbb{Z}\}$$

by $[s_0, \dots, s_n] \mapsto \ker(\mathbb{Z} \rightarrow \pi(s_0, \dots, s_n)/\langle h^2 \rangle)$. One can see that

$$\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z}) \simeq v^{-1}(2\mathbb{Z}) / 2ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle$$

since the submodule $2ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle$ corresponds to the trivial extension. We can see that each element of $v^{-1}(\{0\})$ is contained in $v^{-1}(2\mathbb{Z})$ by the following diagram:

$$\begin{array}{ccccccc} 1 & \rightarrow & \mathbb{Z} & \rightarrow & \pi(s_0, \dots, s_n) & \rightarrow & \Gamma \rightarrow 1 \\ & & \downarrow & & \downarrow & \nearrow & \\ & & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \pi(s_0, \dots, s_n)/\langle h^2 \rangle. & & \end{array}$$

Hence it follows from

$$v^{-1}(\{0\}) \subset v^{-1}(2\mathbb{Z}) \subset ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle$$

and (5) that $v^{-1}(2\mathbb{Z}) = ab\langle x_0, \dots, x_n \mid \alpha_i x_i = x_0, i = 1, \dots, n \rangle \simeq \text{Ext}(\Gamma; \mathbb{Z})$. Therefore we have the isomorphism between $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$ and $\text{Ext}(\Gamma; \mathbb{Z})/2\text{Ext}(\Gamma; \mathbb{Z})$. \square

Theorem 4.3. *Suppose that both of $(b, \beta_1, \dots, \beta_n)$ and $(b', \beta'_1, \dots, \beta'_n)$ give the same class in $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$. If $(b', \beta'_1, \dots, \beta'_n)$ gives an euler class as in Theorem 2.2, then the central extension $[(b', \beta'_1, \dots, \beta'_n)]$ in $\text{Ext}(\Gamma; \mathbb{Z})$ induces an $\text{SL}_2(\mathbb{R})$ -representation ρ' of $\pi_1(M)$ such that $\rho'(h) = -I$.*

Proof. It follows from the assumption that we have an isomorphism φ from $\pi_1(M)/\langle h^2 \rangle$ to $\pi_1(M')/\langle h^2 \rangle$ as

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \pi_1(M)/\langle h^2 \rangle & \rightarrow & \Gamma \rightarrow 1 \\ & & \parallel & & \downarrow \varphi & & \parallel \\ 0 & \rightarrow & \mathbb{Z}/2\mathbb{Z} & \rightarrow & \pi_1(M')/\langle h^2 \rangle & \rightarrow & \Gamma \rightarrow 1 \end{array}$$

where M and M' are the Seifert manifolds with the indices $(g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$ and $(g; (1, b'), (\alpha_1, \beta'_1), \dots, (\alpha_n, \beta'_n))$. Since $b'x_0 + \beta'_1x_1 + \dots + \beta'_nx_n$ is an euler class in $H^2(\Gamma; \mathbb{Z}) = \text{Ext}(\Gamma; \mathbb{Z})$, there exists a $\widetilde{\text{PSL}}_2(\mathbb{R})$ -representation $\bar{\rho}'$ of $\pi_1(M')$. This $\widetilde{\text{PSL}}_2(\mathbb{R})$ -representation $\bar{\rho}'$ gives an $\text{SL}_2(\mathbb{R})$ -representation ρ' such that $\rho'(h) = -I$. Taking the pull-back of the homomorphism from $\pi_1(M')/\langle h^2 \rangle$ to $\text{SL}_2(\mathbb{R})$ by φ , we obtain a homomorphism from $\pi_1(M)/\langle h^2 \rangle$ to $\text{SL}_2(\mathbb{R})$. The composition with the projection from $\pi_1(M)$ gives an $\text{SL}_2(\mathbb{R})$ -representation of $\pi_1(M)$ sending h to $-I$. \square

Remark 4.4. Although $(b, \beta_1, \dots, \beta_n)$ is contained in the equivalent class of $(b', \beta'_1, \dots, \beta'_n)$ in $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$, the induced representation ρ by $(b, \beta_1, \dots, \beta_n)$ is not necessarily conjugate to ρ' induced by $(b', \beta'_1, \dots, \beta'_n)$. We can see an example in Section 5.

Proposition 4.5. *Suppose that $\bar{\rho}$ and $\bar{\rho}'$ are $\text{PSL}_2(\mathbb{R})$ -representations of Γ satisfying that $[e(\bar{\rho})] = [e(\bar{\rho}')] = [bx_0 + \beta_1x_1 + \dots + \beta_nx_n]$ in $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$ but $e(\bar{\rho}) \neq e(\bar{\rho}')$ in $\text{Ext}(\Gamma; \mathbb{Z})$.*

Then the $\text{SL}_2(\mathbb{R})$ -representation ρ of $\pi_1(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}))$ is not conjugate to ρ' .

Proof. If ρ were conjugate to ρ' , then $\bar{\rho}$ would be also conjugate to $\bar{\rho}'$. Since the euler class is invariant under conjugation (we refer to the proof of [JN85, Theorem 1]), the euler class $e(\bar{\rho})$ must coincide with $e(\bar{\rho}')$. This is a contradiction to the assumption that $e(\bar{\rho}) \neq e(\bar{\rho}')$. \square

Theorem 4.6. *Suppose that M denotes a Seifert manifold $M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n})$ and integers $(b', \beta'_1, \dots, \beta'_n)$ gives an euler class satisfying the criteria of Jankins and Neumann in Theorem 2.2. If $(b, \beta_1, \dots, \beta_n)$ and $(b', \beta'_1, \dots, \beta'_n)$ give the same class in $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$, then*

there exists an $\mathrm{SL}_2(\mathbb{R})$ -representation ρ' of $\pi_1(M)$ such that the asymptotic behavior of $\mathrm{Tor}(M; \rho'_{2N})$ is expressed as

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M; \rho'_{2N})|}{2N} = -\left(2 - 2g - \sum_{j=1}^n \frac{\lambda'_j - 1}{\lambda'_j}\right) \log 2$$

where $\lambda'_j = \alpha_j / (\alpha_j, \beta'_j)$.

5. EXAMPLES

5.1. $\mathrm{SU}(1, 1)$ -representations of a Brieskorn manifold. It is known that $\mathrm{SL}_2(\mathbb{R})$ is conjugate to $\mathrm{SU}(1, 1) = \left\{ \begin{pmatrix} \xi & \eta \\ \bar{\eta} & \bar{\xi} \end{pmatrix} \mid |\xi|^2 - |\eta|^2 = 1 \right\}$ by $\begin{pmatrix} 1 & -\sqrt{-1} \\ 1 & \sqrt{-1} \end{pmatrix}$. We can consider $\mathrm{SU}(1, 1)$ -representations instead of $\mathrm{SL}_2(\mathbb{R})$.

Suppose that M is a Seifert homology sphere with three exceptional fibers. The genus of the base orbifold must be zero. The Seifert index is given by

$$(0; (1, b), (\alpha_1, \beta_1), (\alpha_2, \beta_2), (\alpha_3, \beta_3)).$$

We can express $\pi_1(M)$ as

$$\pi_1(M) = \langle q_1, q_2, q_3, h \mid h: \text{central}, q_1^{\alpha_j} = h^{\beta_j} (j = 1, 2, 3), q_1 q_2 q_3 = h^{-b} \rangle.$$

We write $\begin{pmatrix} \xi_j & \eta_j \\ \bar{\eta}_j & \bar{\xi}_j \end{pmatrix}$ for $\rho(q_j)$ and $a_j + b_j \sqrt{-1}$ for each ξ_j .

We will compute irreducible $\mathrm{SU}(1, 1)$ -representations of $\pi_1(M)$ such that $\rho(h) = -I$ up to conjugation.

Definition 5.1. For an irreducible $\mathrm{SU}(1, 1)$ -representation of $\pi_1(M)$, we have the triple (k_1, k_2, k_3) of natural numbers such that

- $\mathrm{tr} \rho(q_j) = 2 \cos(k_j \pi / \alpha_j)$;
- $0 < k_j < \alpha_j$;
- $k_j \equiv \beta_j \pmod{2}$.

Lemma 5.2. Let ρ be an irreducible $\mathrm{SU}(1, 1)$ -representation of $\pi_1(M)$. The conjugacy class of ρ is determined by the pair of the triple (k_1, k_2, k_3) and the sign of b_1 .

Proof. By the relations of $\pi_1(M)$, the representation of ρ is determined by ξ_1, η_1, ξ_2 and η_2 . We can assume that $\rho(q_1)$ is diagonal i.e., $\eta_1 = 0$, and η_2 is a positive real number. The second assumption is realized by the conjugation of a diagonal matrix. Then a_1 equals to $\cos(k_1 \pi / \alpha_1)$ and $b_1 = \pm \sin(k_1 \pi / \alpha_1)$. Since $\mathrm{tr} \rho(q_2)$ equals to $2a_2$, we also have that $a_2 = \cos(k_2 \pi / \alpha_2)$. By the relation that $q_1 q_2 = h^{-b} q_3^{-1}$, we have that

$$\mathrm{tr} \rho(q_3) = \mathrm{tr} \rho(q_3)^{-1} = (-1)^b 2(a_1 a_2 - b_1 b_2).$$

Together with $\mathrm{tr} \rho(q_3) = 2 \cos(k_3 \pi / \alpha_3)$, we can see that b_2 is determined by (k_1, k_2, k_3) and the sign of b_2 . Last η_2 is given by the equality that $a_2^2 + b_2^2 - \eta_2^2 = 1$. \square

Remark 5.3. The diagonal matrix $\begin{pmatrix} e^{\sqrt{-1}\theta} & 0 \\ 0 & e^{-\sqrt{-1}\theta} \end{pmatrix}$ is not conjugate to $\begin{pmatrix} e^{-\sqrt{-1}\theta} & 0 \\ 0 & e^{\sqrt{-1}\theta} \end{pmatrix}$ in $\mathrm{SU}(1, 1)$.

Proposition 5.4. Suppose that an $\mathrm{SU}(1, 1)$ -representation ρ of $\pi_1(M)$ is irreducible and satisfies that $\rho(h) = -I$.

If b is even, then the corresponding triple (k_1, k_2, k_3) satisfies the either inequality:

$$(6) \quad 0 < \frac{k_3}{\alpha_3} \leq \left| \frac{k_1}{\alpha_1} - \frac{k_2}{\alpha_2} \right| \quad \text{or} \quad 1 - \left| \frac{k_1}{\alpha_1} + \frac{k_2}{\alpha_2} - 1 \right| \leq \frac{k_3}{\alpha_3} < 1.$$

If b is odd, then the corresponding triple (k_1, k_2, k_3) satisfies the either inequality:

$$(7) \quad 0 < \frac{k_3}{\alpha_3} \leq \left| \frac{k_1}{\alpha_1} + \frac{k_2}{\alpha_2} - 1 \right| \quad \text{or} \quad 1 - \left| \frac{k_1}{\alpha_1} - \frac{k_2}{\alpha_2} \right| \leq \frac{k_3}{\alpha_3} < 1.$$

Proof. We prove the case that b is even. In this case, we have the equality that $\rho(q_3) = \rho(q_2)^{-1}\rho(q_1)^{-1}$. We can assume $\rho(q_1)$ is diagonal, i.e.,

$$\rho(q_1) = \begin{pmatrix} \xi_1 & 0 \\ 0 & \bar{\xi}_1 \end{pmatrix}$$

without loss of generality. Hence the trace $\text{tr } \rho(q_3) = 2 \cos(k_3\pi/\alpha_3)$ equals to $\xi_1\xi_2 + \bar{\xi}_1\bar{\xi}_2 = 2(a_1a_2 - b_1b_2)$. We have that

$$(8) \quad a_1a_2 - b_1b_2 = \cos\left(\frac{k_3\pi}{\alpha_3}\right).$$

It follows from the assumption that $a_1 = \cos(k_1\pi/\alpha_1)$, $b_1 = \pm \sin(k_1\pi/\alpha_1)$. We also have the inequality that $b_2^2 \geq \sin^2(k_2\pi/\alpha_2)$ from $|\xi_2|^2 = 1 + |\eta_2|^2 \geq 1$, $|\xi_2|^2 = a_2^2 + b_2^2$ and $a_2 = \cos(k_2\pi/\alpha_2)$. Together with Eq. (8), we obtain the following constrains:

$$\begin{aligned} \cos(k_1\pi/\alpha_1) \cos(k_2\pi/\alpha_2) \mp \sin(k_1\pi/\alpha_1)b_2 &\geq \cos(k_3\pi/\alpha_3), \\ b_2^2 &\geq \sin^2(k_2\pi/\alpha_2). \end{aligned}$$

We can derive our inequality (6) from the above constrains. Similarly we can also derive the inequality (7) in the case that b is odd. \square

Remark 5.5. One can find the similar inequality for $\text{SU}(2)$ -representations in [FS90], [Sav99, §14.5].

5.2. Brieskorn manifold of type $(2, 3, 7)$. Let M be the Seifert manifold of the index $(g; (1, b), (\alpha_1, \beta_1), (\alpha_2, \beta_2), (\alpha_3, \beta_3)) = (0; (1, -1), (2, 1), (3, 1), (7, 1))$. Then the fundamental group $\pi_1(M)$ is expressed as

$$\pi_1(M) = \langle q_1, q_2, q_3, h \mid h: \text{central}, q_1^2 = h, q_3^3 = h, q_3^7 = h, q_1q_2q_3 = h^{-(-1)} \rangle.$$

Lemma 5.6. *There are two conjugacy classes of irreducible $\text{SU}(1, 1)$ -representations ρ of $\pi_1(M)$ such that $\rho(h) = -I$.*

Proof. We have the triple (k_1, k_2, k_3) of natural numbers for the $\text{SU}(1, 1)$ -representation ρ , as in Definition 5.1. This triple satisfies the constrain (7). Therefore we have only one triple $(k_1, k_2, k_3) = (1, 1, 1)$. Since we have two possibility of the sign of b_1 , we can conclude that there are the two conjugacy classes of irreducible $\text{SU}(1, 1)$ -representations of $\pi_1(M)$. \square

We see the correspondence between the conjugacy classes and the euler classes of $\text{PSL}_2(\mathbb{R})$ -representations of Γ . From Theorem 4.3 it is enough to find equivalent classes $[(b', \beta'_1, \dots, \beta'_3)]$ in $\text{Ext}(\Gamma; \mathbb{Z}/2\mathbb{Z})$, which satisfy the criteria of Theorem 2.2. Since the genus g in the Seifert index is equal to 0 and the number n of exceptional fibers is equal to 3, we consider the cases (iib) and (iic) in Theorem 2.2.

- (iib) We suppose that $b' = -1$. If we find $(\beta'_1, \beta'_2, \beta'_3)$ such that $\beta'_1/2 + \beta'_2/3 + \beta'_3/7 \leq 1$, then we have only solution $(1, 1, 1)$. Since these integers $(b', \beta'_1, \beta'_2, \beta'_3)$ coincides with $(b, \beta_1, \beta_2, \beta_3) = (-1, 1, 1, 1)$ in the Seifert index of M , this solution gives an $\widetilde{\text{PSL}}_2(\mathbb{R})$ -representation of $\pi_1(M)$.

(iic) We suppose that $b' = -2$. If we find $(\beta'_1, \beta'_2, \beta'_3)$ such that $\beta'_1/2 + \beta'_2/3 + \beta'_3/7 \geq n-1 = 2$, then we have only solution $(1, 2, 6)$. This solution induces an $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation of $\pi_1(M')$ where M' is the Seifert manifold with the index of

$$(g; (1, b), (\alpha_1, \beta_1), (\alpha_2, \beta_2), (\alpha_n, \beta_n)) = (0; (1, -2), (2, 1), (3, 2), (7, 6)).$$

Remark 5.7. The Seifert manifold M' in (iic) is obtained by the homeomorphism by reversing the orientation of a fiber in M . This is due to that the Seifert index of M' is obtained from M as follows:

$$\begin{aligned} (0; (1, -1), (2, 1), (3, 1), (7, 1)) &\xrightarrow{\text{ori. rev.}} (0; (1, 1), (2, -1), (3, -1), (7, -1)) \\ &= (0; (1, 1-3), (2, -1+2), (3, -1+3), (7, -1+7)) \\ &= (0; (1, -2), (2, 1), (3, 2), (7, 6)). \end{aligned}$$

We refer to [NJ81, Theorem 1.5] for the above operations of Seifert index.

Lemma 5.8. *The isomorphism from $\pi_1(M)$ to $\pi_1(M')$ induced by the homeomorphism in Remark 5.7. is given by the following correspondence:*

$$h \mapsto h^{-1}, \quad q_j \mapsto q'_j h^{-1}.$$

Proof. The homeomorphism by reversing the orientation of a fiber induces the correspondence sending h to h^{-1} . The change of Seifert index from $(0; (1, 1), (2, -1), (3, -1), (7, -1))$ to $(0; (1, -2), (2, 1), (3, 2), (7, 6))$ corresponds to changing the generators q_j to $q'_j h^{-1}$. If we write q'_j for the new generators, then we have the presentation of $\pi_1(M')$. \square

Proposition 5.9. *Let ρ and ρ' be irreducible $\mathrm{SU}(1, 1)$ -representations of $\pi_1(M)$ induced by $(b, \beta_1, \beta_2, \beta_3) = (-1, 1, 1, 1)$ and $(b', \beta'_1, \beta'_2, \beta'_3) = (-2, 1, 1, 1)$. The conjugacy class of ρ is different from that of ρ' .*

Proof. The euler class of $e(\bar{\rho}')$ is equal to $-e(\bar{\rho})$. It follows from Proposition 4.5 that ρ is not conjugate to ρ' . \square

Corollary 5.10. *The $\mathrm{SU}(1, 1)$ -representations ρ and ρ' give all representatives in the conjugacy classes of irreducible $\mathrm{SU}(1, 1)$ -representations sending h to $-I$.*

Therefore we have the isomorphism φ from $\pi_1(M)$ to $\pi_1(M')$ induced by the orientation reversing homeomorphism. The composition of φ gives an $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation of $\pi_1(M)$. These two $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representations give the representatives of different conjugacy classes in the set of $\mathrm{SL}_2(\mathbb{R})$ -representations of $\pi_1(M)$.

This observation can be extend to a general Seifert manifold.

Theorem 5.11. *Suppose that M is a Seifert manifold and M' is the orientation reversed manifold $-M$ along fibers. If there exists a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\bar{\rho}$ of $\pi_1(M)$, then we also have a $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation $\bar{\rho}'$ of $\pi_1(M')$ such that*

- $\bar{\rho}'$ induces an $\mathrm{SL}_2(\mathbb{R})$ -representation ρ' of $\pi_1(M)$ which arises the following asymptotic behavior of the Reidemeister torsion:

$$\lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho'_{2N})|}{2N} = \lim_{N \rightarrow \infty} \frac{\log |\mathrm{Tor}(M(\frac{1}{b}, \frac{\alpha_1}{\beta_1}, \dots, \frac{\alpha_n}{\beta_n}); \rho_{2N})|}{2N}$$

where ρ is the $\mathrm{SL}_2(\mathbb{R})$ -representation induced by $\bar{\rho}$;

- $e(\bar{\rho}') = -e(\bar{\rho})$ where $\bar{\rho}$ and $\bar{\rho}'$ are the induced $\widetilde{\mathrm{PSL}}_2(\mathbb{R})$ -representation of Γ .

Proof. Let φ denote the orientation reversing homeomorphism from M to M' . Then we can choose the composition $\rho \circ \varphi^{-1}$ as ρ' . By the construction, we can see the equality on the asymptotic behaviors of the Reidemeister torsions. We also have the equality on the euler classes of $\mathrm{PSL}_2(\mathbb{R})$ -representations of Γ . \square

ACKNOWLEDGMENT

The author wishes to express his thanks to Professor Makoto Sakuma for drawing the author's attention to $\mathrm{PSL}_2(\mathbb{R})$ -representations of a Fuchsian group and the induced asymptotic behavior of the Reidemeister torsion. This research was supported by by JSPS KAKENHI Grant Number 26800030.

REFERENCES

- [FS90] S. Fintushel and R. Stern, *Instanton homology of Seifert fibered homology three spheres*, Proc. London Math. Soc. **61** (1990), 109–137.
- [JN85] M. Jankins and W. Neumann, *Homomorphisms of fuchsian groups to $\mathrm{PSL}(2, \mathbb{R})$* , Comment. Math. Helv. **60** (1985), 480–495.
- [NJ81] W. Neumann and M. Jankins, *Seifert manifolds*, Lecture notes, Brandeis Univ., 1981.
- [Sav99] N. Saveliev, *Lectures on the topology of 3-manifolds*, de Gruyter Textbook, Walter de Gruyter & Co., Berlin, 1999.
- [Sco83] P. Scott, *The geometries of 3-manifolds*, Bull. London Math. Soc. **15** (1983), 401–487.
- [Yam] Y. Yamaguchi, *A surgery formula for the asymptotics of the higher dimensional reidemeister torsion and seifert fibered spaces*, arXiv:1210.8040.
- [Yam13] ———, *Higher even dimensional Reidemeister torsion for torus knot exteriors*, Math. Proc. Cambridge Philos. Soc. **155** (2013), 297–305.

DEPARTMENT OF MATHEMATICS, AKITA UNIVERSITY, 1-1 TEGATA-GAKUENMACHI, AKITA, 010-8502, JAPAN
E-mail address: shouji@math.akita-u.ac.jp