

CLASSIFICATION OF PSEUDO H -TYPE ALGEBRAS

CHRISTIAN AUTENRIED, KENRO FURUTANI, IRINA MARKINA

1. INTRODUCTION

A. Kaplan introduced the Lie groups of Heisenberg type or shortly H -type groups in 1980 [5] and studied in details, for instance in [6, 7]. The H -type Lie algebras of the H -type Lie groups, constructed in [5], used the presence of an inner product on the Lie algebra. Later this approach was extended by exploiting an arbitrary indefinite non-degenerate scalar product in [1, 3] and the introduced Lie algebras received the name pseudo H -type algebras. This construction is closely related to the existence of a special scalar product on the representation space of Clifford algebras. Namely the Clifford algebras $Cl_{r,0}$ generated by a positive definite scalar product space $(\mathbb{R}^{r,0}, \langle \cdot, \cdot \rangle_{r,0})$, which lead to the H -type Lie algebras $\mathfrak{n}_{r,0}$ introduced by A. Kaplan and the Clifford algebras $Cl_{r,s}$ generated by indefinite non-degenerate scalar product spaces $(\mathbb{R}^{r,s}, \langle \cdot, \cdot \rangle_{r,s})$ creating pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$.

In the present work we study the isomorphism properties of the Lie algebras $\mathfrak{n}_{r,s}$, that were constructed as pseudo H -type Lie algebras. Thus, we neglect the presence of the scalar product $\langle \cdot, \cdot \rangle_{r,s}$ on the Lie algebra $\mathfrak{n}_{r,s}$ and study isomorphisms of Lie algebras as themselves. The isomorphism of Lie algebras defines the isomorphism of the corresponding Lie groups. In the present paper we mostly concentrate on the minimal admissible modules. We showed that the Lie algebras $\mathfrak{n}_{r,s}$ can not be isomorphic to $\mathfrak{n}_{u,t}$ unless $r = t$ and $s = u$ or $r = u$ and $s = t$. The question of existence of an isomorphism between $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$ is much more complicated. We proved that if $s = 0$, then the Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ are isomorphic if the dimensions of the centers coincide. If $r, s \neq 0$, then we present examples of both cases: the isomorphic pairs and non isomorphic pairs, having equal dimensions. Some of the Lie algebras, that we call of block type, allow to use the Bott periodicity of underlying Clifford algebras and obtain more isomorphic pairs, see Theorems 5.9 and 5.10.

We stress an interesting feature, that there are no direct relations between the isomorphisms of the underlying Clifford algebras and the isomorphisms of generated pseudo H -type Lie algebras. In some cases the isomorphic Clifford algebras lead to isomorphic Lie algebras, in other cases not. For instance, in spite of the isomorphism of the Clifford algebras $Cl_{8,0}$, $Cl_{0,8}$, and $Cl_{4,4}$, the corresponding Lie algebras $\mathfrak{n}_{8,0}$, $\mathfrak{n}_{0,8}$ are isomorphic, but not isomorphic to the Lie algebra $\mathfrak{n}_{4,4}$.

The structure of the article is the following. After this Introduction we give all necessary definitions and notations in Section 2. Furthermore, the relation between Clifford algebras and pseudo H -type Lie algebras is presented. In Section 3, we discuss a necessary condition for isomorphisms of pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$, which shows

Key words and phrases. Sub-Riemannian geometry, Clifford Algebras, H -type algebras, Classification, Lie algebras.

The authors are partially supported by the NFR-FRINAT grants #204726/V30 and #213440/BG..

that the only possible algebra which is isomorphic to $\mathfrak{n}_{r,s}$ is the pseudo H -type Lie algebra $\mathfrak{n}_{s,r}$. Section 4 is devoted to the complete classification of pseudo H -type Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$. Sections 5 and 6 study two different situations revealing that the Lie algebras $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$ can be both isomorphic and non-isomorphic. In Section 7, we exhibit the strongly bracket generating property of the pseudo H -type Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,s}$ and the non-existence of this property for pseudo H -type algebras $\mathfrak{n}_{r,s}$ with $r, s \neq 0$. Furthermore, we introduce an equivalent definition for pseudo H -type Lie algebras, that was introduced in [3], and explain the equivalence of these definitions in details in subsection 7.1. In Section 8, we briefly discuss the isomorphism of pseudo H -type Lie algebras related to non-equivalent irreducible Clifford modules. Finally Appendix 9, gives the commutator tables of $\mathfrak{n}_{8,0}$, $\mathfrak{n}_{0,8}$, $\mathfrak{n}_{4,4}$ and a table of permutations for a basis of $\mathfrak{n}_{8,0}$.

2. PRELIMINARIES

2.1. Clifford algebras and their representations. Throughout this paper we assume all scalar products to be non-degenerate besides otherwise stated. We denote by \mathfrak{n} a nilpotent 2-step Lie algebra endowed with a scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{n}}$ of signature (r, s) , $r, s \in \mathbb{N}$, $r + s = n$: that means that there exists a basis $\{U_1, \dots, U_n\}$ of \mathfrak{n} which satisfies

$$\langle U_i, U_j \rangle_{\mathfrak{n}} = \epsilon_i(r, s) \delta_{ij}, \quad \text{where} \quad \epsilon_i(r, s) = \begin{cases} 1, & \text{for } i = 1, \dots, r, \\ -1, & \text{for } i = r + 1, \dots, r + s. \end{cases}$$

The definition of the index varies in the literature, thus we follow to one which is given above. Let \mathfrak{z} be the center of the 2-step nilpotent Lie algebra \mathfrak{n} and $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$ the restriction of the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{n}}$ to \mathfrak{z} . We assume that $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$ is non-degenerate. Then the orthogonal complement $\mathfrak{v} := \mathfrak{z}^{\perp}$ is also a non-degenerate scalar product space, where we use the symbol $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ to denote the restriction of $\langle \cdot, \cdot \rangle_{\mathfrak{n}}$ on \mathfrak{v} . Thus $\mathfrak{n} = \mathfrak{z} \oplus_{\perp} \mathfrak{v}$ is an orthogonal decomposition with respect to the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{z}} + \langle \cdot, \cdot \rangle_{\mathfrak{v}}$. Since \mathfrak{z} is the center of \mathfrak{n} , the commutator is a skew-symmetric bi-linear map $[\cdot, \cdot]: \mathfrak{v} \times \mathfrak{v} \rightarrow \mathfrak{z}$.

Definition 2.1. *Let $\mathfrak{n} = (\mathfrak{z} \oplus_{\perp} \mathfrak{v}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}})$ be a Lie algebra described above. We define the map $J: \mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$ by*

$$\langle J_Z v, w \rangle_{\mathfrak{v}} = \langle Z, [v, w] \rangle_{\mathfrak{z}}, \quad \text{for all } v, w \in \mathfrak{v}. \quad (1)$$

Definition 2.2. [1] *We call a 2-step nilpotent Lie algebra $\mathfrak{n} = (\mathfrak{z} \oplus_{\perp} \mathfrak{v}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}})$ with $J: \mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$ from Definition 2.1 a pseudo H -type Lie algebra if*

$$\langle J_Z v, J_Z v \rangle_{\mathfrak{v}} = \langle Z, Z \rangle_{\mathfrak{z}} \langle v, v \rangle_{\mathfrak{v}} \quad \text{for all } Z \in \mathfrak{z} \text{ and } v \in \mathfrak{v}. \quad (2)$$

We write $\mathfrak{n}_{r,s}$ to emphasize that $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$ has signature (r, s) .

We call the attention of the reader that the sign in equation (2) differs from the original work [1] due to the different agreement on the signature. It follows directly from Definition 2.1 that J_Z is skew-adjoint with respect to the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$:

$$\langle J_Z v, w \rangle_{\mathfrak{v}} = -\langle v, J_Z w \rangle_{\mathfrak{v}} \quad \text{for all } Z \in \mathfrak{z}, \quad v, w \in \mathfrak{v}. \quad (3)$$

Using polarization in (2) we obtain

$$\langle J_Z v, J_{Z'} v \rangle_{\mathfrak{v}} = \langle Z, Z' \rangle_{\mathfrak{z}} \langle v, v \rangle_{\mathfrak{v}}, \quad \text{and} \quad \langle J_Z v, J_Z v' \rangle_{\mathfrak{v}} = \langle Z, Z \rangle_{\mathfrak{z}} \langle v, v' \rangle_{\mathfrak{v}}. \quad (4)$$

Applying the skew-adjoint property (3) one also obtains

$$J_Z \circ J_Z := J_Z J_Z = J_Z^2 = -\langle Z, Z \rangle_{\mathfrak{z}} \text{Id}_{\mathfrak{v}}, \quad \text{or} \quad J_{Z'} J_Z + J_Z J_{Z'} = -2\langle Z, Z' \rangle_{\mathfrak{z}} \text{Id}_{\mathfrak{v}}, \quad (5)$$

for all $Z, Z' \in \mathfrak{z}$. Equality (5) implies that $J: \mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$ defines a representation of the Clifford algebra $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$. We recall the definition of a Clifford algebra and its representations.

Definition 2.3. *A Clifford algebra $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ generated by a scalar product space $(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ is the unital associative algebra generated by \mathfrak{z} subject to the relations:*

$$Z \otimes Z = -\langle Z, Z \rangle_{\mathfrak{z}} \mathbb{I}_{\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})},$$

for all $Z \in \mathfrak{z}$ and the unit $\mathbb{I}_{\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})}$ of the Clifford algebra.

Thus, \mathfrak{z} can be considered as a subset of $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ and

$$Z \otimes W + W \otimes Z = -2\langle Z, W \rangle_{\mathfrak{z}} \mathbb{I}_{\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})} \quad \text{for all } Z, W \in \mathfrak{z}.$$

Proposition 2.4. [4, 9] *Let $J: \mathfrak{z} \rightarrow \mathcal{A}$ be a linear map into an associative algebra \mathcal{A} with an identity element $\text{Id}_{\mathcal{A}}$ and product " $\cdot_{\mathcal{A}}$ ", such that*

$$J(Z) \cdot_{\mathcal{A}} J(Z) = -\langle Z, Z \rangle_{\mathfrak{z}} \text{Id}_{\mathcal{A}} \quad \text{for all } Z \in \mathfrak{z}.$$

Then J extends uniquely to an algebra homomorphism $\tilde{J}: \text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}}) \rightarrow \mathcal{A}$. Moreover, $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ is the unique associative algebra with this property.

Definition 2.5. *A representation of a Clifford algebra $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ is an algebra homomorphism $J: \text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}}) \rightarrow \text{End}(\mathfrak{v})$ into the algebra of linear transformations of a finite dimensional vector space \mathfrak{v} . The space \mathfrak{v} is called a $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ -module.*

Since J is an algebra homomorphism we obtain that

$$J_Z^2 := J_Z \circ J_Z = J_{Z \otimes Z} = J_{-\langle Z, Z \rangle_{\mathfrak{z}} \mathbb{I}_{\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})}} = -\langle Z, Z \rangle_{\mathfrak{z}} \text{Id}_{\mathfrak{v}}.$$

The scalar product space $(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ with the signature (r, s) is isomorphic to $\mathbb{R}^{r,s} = (\mathbb{R}^{r+s}, \langle \cdot, \cdot \rangle_{r,s})$, $r + s = n$, where the scalar product $\langle \cdot, \cdot \rangle_{r,s}$ is defined by $\langle Z, W \rangle_{r,s} = \sum_{i=1}^r Z_i W_i - \sum_{j=r+1}^{r+s} Z_j W_j$ for all $Z, W \in \mathbb{R}^{r+s}$. It allows us to use the isomorphism of the Clifford algebras $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ and $\text{Cl}(\mathbb{R}^{r,s}, \langle \cdot, \cdot \rangle_{r,s})$. The Clifford algebra $\text{Cl}(\mathbb{R}^{r,s}, \langle \cdot, \cdot \rangle_{r,s})$ is denoted by $\text{Cl}_{r,s}$, and we denote by (Z_1, \dots, Z_{r+s}) the orthonormal basis of $\mathbb{R}^{r,s}$ with $\langle Z_i, Z_j \rangle_{r,s} = \epsilon_i(r, s) \delta_{ij}$.

2.2. Admissible Clifford modules and pseudo H -type Lie algebras. In this section we explain when a representation space \mathfrak{v} of a Clifford algebra can be endowed with a scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ such that the representation map J satisfies (3). We call a positive definite scalar product we an *inner product*, and in any case we work with *only* non-degenerate scalar products.

Proposition 2.6. [4] *Let $J: \text{Cl}_{r,0} \rightarrow \text{End}(\mathfrak{v})$ be a representation. Then there exists an inner product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ on \mathfrak{v} , such that the following holds:*

$$\langle J_Z w, J_Z v \rangle_{\mathfrak{v}} = \langle w, v \rangle_{\mathfrak{v}} \quad \text{for all } w, v \in \mathfrak{v}, Z \in \mathbb{R}^{r,0} \quad \text{with } \langle Z, Z \rangle_{r,0} = 1. \quad (6)$$

Corollary 2.7. *Any representation $J: \text{Cl}_{r,0} \rightarrow \text{End}(\mathfrak{v})$ satisfies property (3) with respect to an inner product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$.*

Proof. Corollary 2.7 follows by replacing w by $J_Z w$ in (6) and applying $J_Z^2 = -\text{Id}_{\mathfrak{v}}$. \square

Thus the Clifford algebras $\text{Cl}_{r,0}$ possess always an inner product on \mathfrak{v} such that J_Z is skew-adjoint for all $Z \in \mathbb{R}^{r,0}$. A. Kaplan used inner products on \mathfrak{v} for the construction of H -type algebras, which are based on $\text{Cl}_{r,0}$ -modules, see [5, 7]. For $\text{Cl}_{r,s}$ -modules with $s \geq 1$ equation (6) is only true for orthonormal bases and in general not true for an arbitrary element of $\mathbb{R}^{r,s}$, see [4].

Definition 2.8. [1] *A pair $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$, where \mathfrak{v} is a $\text{Cl}_{r,s}$ -module is said to be an admissible $\text{Cl}_{r,s}$ -module if the representation operators $J_Z: \mathfrak{v} \rightarrow \mathfrak{v}$ are skew-adjoint with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$, i.e. satisfies (3) for all $Z \in \mathbb{R}^{r,s}$.*

The following proposition guarantees the existence of admissible $\text{Cl}_{r,s}$ -modules.

Proposition 2.9. [1] *For any given $\text{Cl}_{r,s}$ -module \mathfrak{v} the vector space \mathfrak{v} itself (or $\mathfrak{v} \oplus \mathfrak{v}$) can be equipped with a scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ (or $\langle \cdot, \cdot \rangle_{\mathfrak{v} \oplus \mathfrak{v}}$), such that*

$$\langle J_Z w, v \rangle_{\mathfrak{v}} = -\langle w, J_Z v \rangle_{\mathfrak{v}}, \quad \text{or} \quad \langle J'_Z w, v \rangle_{\mathfrak{v} \oplus \mathfrak{v}} = -\langle w, J'_Z v \rangle_{\mathfrak{v} \oplus \mathfrak{v}}$$

for all $Z \in \mathbb{R}^{r,s}$ and all $w, v \in \mathfrak{v}$ (or $w, v \in \mathfrak{v} \oplus \mathfrak{v}$, where the operator $J': \text{Cl}_{r,s} \rightarrow \text{End}(\mathfrak{v} \oplus \mathfrak{v})$ should be redefined correspondingly).

The relation between Lie algebras of Definition 2.2 and admissible Clifford modules is summarized in the following proposition.

Proposition 2.10. [1] *Let \mathfrak{v} be a $\text{Cl}_{r,s}$ -module. Then $\mathfrak{n} = \mathfrak{v} \oplus \mathbb{R}^{r,s}$ can be supplied with the structure of the pseudo H -type algebra if and only if there exists a scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ making the $\text{Cl}_{r,s}$ -module \mathfrak{v} into an admissible Clifford module $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$. The bracket $[\cdot, \cdot]: \mathfrak{v} \times \mathfrak{v} \rightarrow \mathbb{R}^{r,s}$ on \mathfrak{n} is given by (2.1) and the scalar product is $\langle \cdot, \cdot \rangle_{\mathfrak{n}} := \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{r,s}$. The decomposition $\mathfrak{n} = \mathfrak{v} \oplus \mathbb{R}^{r,s}$ is orthogonal and $\mathbb{R}^{r,s}$ is the center of \mathfrak{n} .*

Proposition 2.11. [1] *Let \mathfrak{n} be a pseudo H -type algebra. Then the corresponding admissible $\text{Cl}_{r,s}$ -module $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$ is a neutral scalar product space for $s \geq 1$, i.e. the signature of $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ is (l, l) , with $l \in \mathbb{N}$.*

2.3. Existence of the integral structure on pseudo H -type Lie algebras. In this subsection we state some necessary facts about the latest research on pseudo H -type Lie algebras based on the work [2]. The principal result of [2] states that pseudo H -type Lie algebras admit a special choice of basis giving integer structure constants.

Theorem 2.12. [2] *Let $\mathfrak{n} = (\mathfrak{v} \oplus_{\perp} \mathfrak{z}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ be a pseudo H -type Lie algebra. Then for any orthonormal basis $\{Z_1, \dots, Z_n\}$ for \mathfrak{z} there is an orthonormal basis $\{v_1, \dots, v_m\}$ for \mathfrak{v} such that $[v_{\alpha}, v_{\beta}] = \sum_{k=1}^n A_{\alpha\beta}^k Z_k$, where $A_{\alpha\beta}^k = 0, \pm 1$.*

Corollary 2.13. [2] *There exists an orthonormal basis $\mathcal{B} = \{v_1, \dots, v_m, Z_1, \dots, Z_n\}$ for any pseudo H -type Lie algebra such that $[v_{\alpha}, v_{\beta}] = \pm Z_{k_{\alpha,\beta}}$ or $[v_{\alpha}, v_{\beta}] = 0$. In particular, for every Z_k and v_{α} in \mathcal{B} there exists exactly one $\beta \in \{1, \dots, m\}$ such that $[v_{\alpha}, v_{\beta}] = \pm Z_k$.*

Definition 2.14. *We call an orthonormal basis $\{v_1, \dots, v_m, Z_1, \dots, Z_n\}$ of a pseudo H -type Lie algebra with the form of Corollary 2.13 an integral basis. The corresponding Clifford module $\mathfrak{v} = \text{span}\{v_1, \dots, v_m\}$ from Corollary 2.13 is called an integral module and if it is of minimal possible dimension we call it minimal integral module.*

Let $\{v_1, \dots, v_m, Z_1, \dots, Z_n\}$ be an orthonormal basis of \mathfrak{n} . Denote by $\epsilon_\alpha^{\mathfrak{v}}$ and $\epsilon_k^{\mathfrak{z}}$ the indices corresponding to scalar product spaces $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$ and $(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$. The structure constants and the coefficients of the representation operator $J: \mathfrak{z} \rightarrow \text{End}(\mathfrak{v})$ are

$$[v_\alpha, v_\beta] = \sum_{k=1}^n A_{\alpha\beta}^k Z_k \quad \text{and} \quad J_{Z_k} v_\alpha = \sum_{\beta=1}^m B_{\alpha\beta}^k v_\beta. \quad (7)$$

Then we obtain the relation

$$\epsilon_\beta^{\mathfrak{v}} B_{\alpha\beta}^k = \epsilon_k^{\mathfrak{z}} A_{\alpha\beta}^k \quad (8)$$

from $\langle J_{Z_k} v_\alpha, v_\beta \rangle_{\mathfrak{v}} = \langle Z_k, [v_\alpha, v_\beta] \rangle_{\mathfrak{z}}$ by [2]. Equality (8) allows to relate the structure constants $A_{\alpha\beta}^k$ of pseudo H -type Lie algebras and coefficients $B_{\alpha\beta}^k$ of the representation operator. From now on, we denote the pseudo H -type Lie algebra induced by $\text{Cl}_{r,s}$ by $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$, where $\mathfrak{v}_{r,s}$ is the minimal admissible integral module $\mathfrak{v}_{r,s}$ of $\text{Cl}_{r,s}$ and $\mathfrak{z}_{r,s} = \mathbb{R}^{r,s}$ the generator space of the Clifford algebra $\text{Cl}_{r,s}$ and the center of the Lie algebra $\mathfrak{n}_{r,s}$.

3. NECESSARY CONDITION FOR ISOMORPHISMS OF PSEUDO H -TYPE LIE ALGEBRAS

In the present section we identify the admissible module $\mathfrak{v}_{r,s}$, $s \neq 0$ with $\mathbb{R}^{l,l}$ equipped with the neutral scalar product $\langle x, y \rangle_{l,l} = \sum_{i=1}^l x_i y_i - \sum_{j=l+1}^{2l} x_j y_j$ for $x, y \in \mathbb{R}^{l,l}$. In the case $\mathfrak{v}_{r,0}$ we use the identification of $\mathfrak{v}_{r,0}$ with \mathbb{R}^{2l} endowed with the inner product $\langle x, y \rangle_{2l} = \sum_{i=1}^{2l} x_i y_i$ for $x, y \in \mathbb{R}^{2l}$. Thus a pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$ is isometric to $\mathbb{R}^{l,l} \oplus \mathbb{R}^{r,s}$. Let $A \in GL(\mathbb{R}^{2l})$. We denote by A^τ the adjoint map with respect to the neutral scalar product $\langle \cdot, \cdot \rangle_{l,l}$

$$\langle Aw, v \rangle_{l,l} = \langle w, A^\tau v \rangle_{l,l}.$$

The same symbol we use to write the adjoint map $\langle Aw, v \rangle_{2l,0} = \langle w, A^\tau v \rangle_{l,l}$ with respect to scalar products $\langle \cdot, \cdot \rangle_{2l,0}$ and $\langle \cdot, \cdot \rangle_{l,l}$.

The adjoint map C^τ for the map $C: \mathbb{R}^{t,u} \rightarrow \mathbb{R}^{r,s}$ with $t+u = r+s$ with respect to corresponding scalar products is given by

$$\langle C(Z), \zeta \rangle_{r,s} = \langle Z, C^\tau(\zeta) \rangle_{t,u}.$$

Assume that two pseudo H -type Lie algebras $\mathfrak{n}_{t,u}$ and $\mathfrak{n}_{r,s}$ are isomorphic and $f: \mathfrak{n}_{t,u} \rightarrow \mathfrak{n}_{r,s}$ is an isomorphism. Then $t+u = r+s$ and since the center of $\mathfrak{n}_{t,u}$ is mapped to the center of $\mathfrak{n}_{r,s}$, the matrix of the map f takes the form

$$M_f = \begin{pmatrix} A & 0 \\ B & C \end{pmatrix}, \quad A \in GL(\mathbb{R}^{2l}), \quad C \in GL(\mathbb{R}^{r+s}) \quad (9)$$

and B is a $((r+s) \times 2l)$ -matrix. Checking the commutation relations, we get $[Aw, Av] = C([w, v])$ for all $w, v \in \mathbb{R}^{2l}$. With this notation we prove our first classification result.

Lemma 3.1. *Let $f: \mathfrak{n}_{t,u} \rightarrow \mathfrak{n}_{r,s}$ be a Lie algebra isomorphism represented by (9). Then the matrices A and C in (9) satisfy*

$$A^\tau \circ J_Z \circ A = J_{C^\tau(Z)}, \quad \text{for all } Z \in \mathbb{R}^{r,s}. \quad (10)$$

Proof. Formula (10) follows from the calculations

$$\begin{aligned} \langle A^\tau \circ J_Z \circ Aw, v \rangle_{l,l} &= \langle J_Z Aw, Av \rangle_{l,l} = \langle Z, [Aw, Av] \rangle_{r,s} \\ &= \langle Z, C([w, v]) \rangle_{r,s} = \langle C^\tau(Z), [w, v] \rangle_{t,u} = \langle J_{C^\tau(Z)} w, v \rangle_{l,l} \end{aligned}$$

for all $w, v \in \mathbb{R}^{2l}$, $Z \in \mathbb{R}^{r,s}$ with $s \neq 0$. If $s = 0$ we use the same arguments applied for scalar products $\langle \cdot, \cdot \rangle_{2l,0}$ and $\langle \cdot, \cdot \rangle_{l,l}$. \square

Theorem 3.2. *A pseudo H -type algebra $\mathfrak{n}_{r,0}$ can only be isomorphic to $\mathfrak{n}_{0,r}$.*

Proof. Formula (10) implies that the action $J_{C^\tau(Z)}$ is singular if and only if J_Z is singular for $Z \in \mathbb{R}^{r,s}$ and this happens only if Z is a null vector in $\mathbb{R}^{r,s}$. Since the space $\mathbb{R}^{r,0}$ has no null vectors, the isomorphic Lie algebra can only have the center isomorphic to $\mathbb{R}^{0,r}$. \square

Theorem 3.3. *A pseudo H -type algebra $\mathfrak{n}_{r,s}$ can only be isomorphic to $\mathfrak{n}_{s,r}$ for $r, s \neq 0$.*

Proof. Let $Z_+, Z_- \in \mathbb{R}^{t,u}$ with $\langle Z_+, Z_+ \rangle_{t,u} > 0$ and $\langle Z_-, Z_- \rangle_{t,u} < 0$. We consider the line segment $\gamma(t) = (1-t)Z_+ + tZ_-$ for $t \in [0, 1]$. Then there exists $t_0 \in (0, 1)$ such that $\langle \gamma(t_0), \gamma(t_0) \rangle_{t,u} = 0$ by the continuity of the scalar product and as $\langle \gamma(0), \gamma(0) \rangle_{t,u} > 0$, $\langle \gamma(1), \gamma(1) \rangle_{t,u} < 0$.

Assume that $\mathfrak{B}_{\mathbb{R}^{t,u}} = \{Z_1, \dots, Z_{t+u}\}$ is an orthonormal basis in $\mathbb{R}^{t,u}$ such that $\langle Z_i, Z_j \rangle_{t,u} = \epsilon_i(t, u) \delta_{ij}$ and we denote

$$\varphi_{ij}(t) = \langle (1-t)Z_i + tZ_j, (1-t)Z_i + tZ_j \rangle_{t,u} = (1-t)^2 \langle Z_i, Z_i \rangle_{t,u} + t^2 \langle Z_j, Z_j \rangle_{t,u},$$

for $i \neq j$. Then

$$\varphi_{ij}(t) > 0 \text{ if } i, j = 1, \dots, t, \quad \text{and} \quad \varphi_{ij}(t) < 0 \text{ if } i, j = t+1, \dots, t+u \quad (11)$$

for all $t \in [0, 1]$.

Let $f: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{t,u}$ be the isomorphism represented by (9). Consider the image $\{C^\tau(Z_1), \dots, C^\tau(Z_{t+u})\} \subset \mathbb{R}^{r,s}$ under the map C^τ of the basis $\mathfrak{B}_{\mathbb{R}^{t,u}}$. First we note that $\langle C^\tau(Z_i), C^\tau(Z_i) \rangle_{r,s} \neq 0$ by Lemma 3.1.

We claim that for basis vectors Z_i , $i = 1, \dots, t$, one gets $\langle C^\tau(Z_i), C^\tau(Z_i) \rangle_{r,s} > 0$ or $\langle C^\tau(Z_i), C^\tau(Z_i) \rangle_{r,s} < 0$ for all indices $i = 1, \dots, t$, simultaneously. Indeed, assume that there are $C^\tau(Z_i)$ and $C^\tau(Z_j)$ for $i, j = 1, \dots, t$ such that products $\langle C^\tau(Z_i), C^\tau(Z_i) \rangle_{r,s}$ and $\langle C^\tau(Z_j), C^\tau(Z_j) \rangle_{r,s}$ have opposite sign. Then there exists $t_0 \in (0, 1)$ such that

$$\langle (1-t_0)C^\tau(Z_i) + t_0C^\tau(Z_j), (1-t_0)C^\tau(Z_i) + t_0C^\tau(Z_j) \rangle_{r,s} = 0,$$

which implies that $J_{(1-t_0)Z_i + t_0Z_j}$ is singular by Lemma 3.1, which contradicts (11). The same arguments are valid for the basis vectors Z_i with $i = t+1, \dots, t+u$. Thus we conclude that the scalar product $\langle \cdot, \cdot \rangle_{r,s}$ restricted to subspaces $\text{span}\{C^\tau(Z_1), \dots, C^\tau(Z_t)\}$ and $\text{span}\{C^\tau(Z_{t+1}), \dots, C^\tau(Z_{t+u})\}$ is sign definite. As $\{C^\tau(Z_1), \dots, C^\tau(Z_{t+u})\}$ is a basis of $\mathbb{R}^{r,s}$ it follows that

$$r = t \quad \text{and} \quad s = u, \quad \text{or} \quad r = u \quad \text{and} \quad s = t.$$

This implies, that the only possible isomorphic pseudo H -type algebra for $\mathfrak{n}_{r,s}$ is $\mathfrak{n}_{s,r}$. \square

Theorem 3.4. *If $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$, $r \neq s$, are isomorphic, then there exists a Lie algebra isomorphism $\varphi: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{s,r}$ given by the matrix $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$ with $CC^\tau = -\text{Id}_{\mathbb{R}^{r,s}}$. Moreover $C^\tau C = -\text{Id}_{\mathbb{R}^{r,s}}$ and C, C^τ are anti-isometries.*

Proof. Theorem 3.3 implies that for any $Z \in \mathbb{R}^{s,r}$: $\langle Z, Z \rangle_{s,r} = -\lambda \langle C^\tau(Z), C^\tau(Z) \rangle_{r,s}$ for some $\lambda > 0$. To determine λ we pick up an arbitrary $Z \in \mathbb{R}^{s,r}$ and calculate

$$(\det(A^\tau J_Z A))^2 = (\det(A^\tau A))^2 (\langle Z, Z \rangle_{s,r})^{2l}.$$

On the other hand

$$(\det(A^\tau J_Z A))^2 = (\det(J_{C^\tau(Z)}))^2 = (\langle C^\tau(Z), C^\tau(Z) \rangle_{r,s})^{2l},$$

which is equivalent to $|\det(A^\tau A)|^{1/l} \langle Z, Z \rangle_{s,r} = -\langle C^\tau(Z), C^\tau(Z) \rangle_{r,s} = -\langle Z, CC^\tau(Z) \rangle_{s,r}$. It follows that $CC^\tau = -|\det(A^\tau A)|^{1/l} \text{Id}_{\mathbb{R}^{s,r}}$.

If $\varphi: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{s,r}$ is a Lie algebra isomorphism, then $\tilde{\varphi} = \begin{pmatrix} \mu A & 0 \\ B & \mu^2 C \end{pmatrix}$ for $\mu \neq 0$ is also a Lie algebra isomorphism as

$$\begin{aligned} \tilde{\varphi}([w, v]_{r,s}) &= (\mu^2 C)([w, v]_{r,s}) = \mu^2(C([w, v]_{r,s})) = \mu^2([Aw, Av]_{s,r}) \\ &= [\mu Aw, \mu Av]_{s,r} = [\tilde{\varphi}(w), \tilde{\varphi}(v)]_{s,r} \end{aligned}$$

for all $w, v \in \mathbb{R}^{l,l}$. Hence, without loss of generality, we can assume that $|\det(A^\tau A)| = 1$, which implies that $CC^\tau = -\text{Id}_{\mathbb{R}^{s,r}}$.

To show that C and C^τ are anti-isometries we choose an arbitrary $Z \in \mathbb{R}^{s,r}$ and obtain

$$\langle C^\tau(Z), C^\tau(Z) \rangle_{r,s} = \langle CC^\tau(Z), Z \rangle_{s,r} = -\langle Z, Z \rangle_{s,r}.$$

As C^τ is an isomorphism for any $Y \in \mathbb{R}^{r,s}$ there exists a unique $Z_Y \in \mathbb{R}^{s,r}$ such that $C^\tau(Z_Y) = Y$. It follows that for any $Y \in \mathbb{R}^{r,s}$ we have the equality $\langle Y, Y \rangle_{r,s} = \langle C^\tau(Z_Y), C^\tau(Z_Y) \rangle_{r,s} = -\langle Z_Y, Z_Y \rangle_{s,r}$. Thus

$$\begin{aligned} \langle C^\tau C(Y), Y \rangle_{r,s} &= \langle C(Y), C(Y) \rangle_{s,r} = \langle CC^\tau(Z_Y), CC^\tau(Z_Y) \rangle_{s,r} = \langle Z_Y, Z_Y \rangle_{s,r} \\ &= -\langle Y, Y \rangle_{r,s}. \end{aligned}$$

Hence $C^\tau C = -\text{Id}_{\mathbb{R}^{r,s}}$ and C is an anti-isometry. \square

Theorem 3.5. *For any $\mathfrak{n}_{r,s}$, $r \neq s$ there exists a Lie algebra automorphism $f: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{r,s}$ given by the matrix $\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}$ with $CC^\tau = \text{Id}_{\mathbb{R}^{r,s}}$. Moreover $C^\tau C = \text{Id}_{\mathbb{R}^{r,s}}$ and C, C^τ are isometries.*

Proof. The proof follows analogously to the proof of Theorem 3.4 for $r \neq s$. For $r = s$ one of the possible automorphisms is the identity map. \square

Remark 3.6. *If $r = s$, then there can be cases when there exist two automorphisms: one with $CC^\tau = \text{Id}_{\mathbb{R}^{r,r}}$ and one with $CC^\tau = -\text{Id}_{\mathbb{R}^{r,r}}$, see for instance Theorem 5.6. But there exist also cases where there are only automorphisms with $CC^\tau = \text{Id}_{\mathbb{R}^{r,r}}$, see for instance Theorem 6.6.*

4. CLASSIFICATION OF $\mathfrak{n}_{r,0}$ AND $\mathfrak{n}_{0,r}$

Note that since the isomorphisms have to preserve the dimensions of Lie algebras and their centers, we only need to check algebras $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{t,u}$ with $r + s = t + u$.

4.1. Classification of $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ with $r = 1, 2, 4, 8$.

Definition 4.1. Let $\{v_1, \dots, v_m, Z_1, \dots, Z_n\}$ be an integral basis of a pseudo H -type Lie algebra \mathfrak{n} and $\{\tilde{v}_1, \dots, \tilde{v}_m, \tilde{Z}_1, \dots, \tilde{Z}_n\}$ an integral basis of a pseudo H -type Lie algebra $\tilde{\mathfrak{n}}$. If a Lie algebra isomorphism $f: \mathfrak{n} \rightarrow \tilde{\mathfrak{n}}$ satisfies $f(v_i) = \tilde{v}_i$ and $f(Z_l) = \tilde{Z}_l$, then f is called an integral isomorphism and we say that \mathfrak{n} is integral isomorphic to $\tilde{\mathfrak{n}}$.

Notation 4.2. For a given orthonormal basis $\{Z_1, \dots, Z_{r+s}\}$ of the center $\mathfrak{z}_{r,s}$ of the pseudo H -type algebra $\mathfrak{n}_{r,s}$ we simplify the notation of the operator $J_{Z_i}: \mathfrak{v}_{r,s} \rightarrow \mathfrak{v}_{r,s}$ to $J_{Z_i} := J_i$ for all $Z_i \in \{Z_1, \dots, Z_{r+s}\}$.

Theorem 4.3. The Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ with $r = 1, 2, 4, 8$ are integral isomorphic.

Proof. For all the cases we denote by $\{Z_1, \dots, Z_{r+s}\}$ an orthonormal basis of the center $\mathfrak{z}_{r,s}$ of the pseudo H -type algebra $\mathfrak{n}_{r,s}$ with $\langle Z_k, Z_k \rangle_{\mathfrak{z}_{r,s}} = \epsilon_k(r, s)$. Furthermore, all admissible modules are assumed to be minimal and all structural constants are obtained by the use of relation (8). For more details how to obtain integral bases for general pseudo H -type algebras see [2].

Isomorphism on $\mathfrak{n}_{1,0}$ and $\mathfrak{n}_{0,1}$. Let $(\mathfrak{v}_{1,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{1,0}})$ be an admissible module and $w \in \mathfrak{v}_{1,0}$ be such that $\langle w, w \rangle_{\mathfrak{v}_{1,0}} = 1$. Then the basis $w_1 = w$, $w_2 = J_1 w$ is integral and $\langle w_i, w_i \rangle_{\mathfrak{v}_{1,0}} = 1$, $i = 1, 2$.

Let $(\mathfrak{v}_{0,1}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,1}})$ be an admissible module and $\tilde{w} \in \mathfrak{v}_{0,1}$ such that $\langle \tilde{w}, \tilde{w} \rangle_{\mathfrak{v}_{0,1}} = 1$. Then the basis $\tilde{w}_1 = \tilde{w}$, $\tilde{w}_2 = \tilde{J}_1 \tilde{w}$, is integral and $\langle \tilde{w}_i, \tilde{w}_i \rangle_{\mathfrak{v}_{0,1}} = \epsilon_i(1, 1)$, $i = 1, 2$.

Calculating the commutators with respect to both integral bases, presented in Table 1, we conclude that they coincide. It follows that $\mathfrak{n}_{1,0}$ is integral isomorphic to $\mathfrak{n}_{0,1}$ under the isomorphism $\varphi_{1,0}: \mathfrak{n}_{1,0} \rightarrow \mathfrak{n}_{0,1}$ defined by $w_1 \mapsto \tilde{w}_1$, $w_2 \mapsto \tilde{w}_2$, $Z_1 \mapsto \tilde{Z}_1$.

TABLE 1. Commutation relations for $\mathfrak{n}_{1,0}$ and $\mathfrak{n}_{0,1}$

[row, col.]	w_1	w_2
w_1	0	Z_1
w_2	$-Z_1$	0

Isomorphism of $\mathfrak{n}_{2,0}$ and $\mathfrak{n}_{0,2}$. In the admissible module $(\mathfrak{v}_{2,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{2,0}})$ we pick up $w \in \mathfrak{v}_{2,0}$ such that $\langle w, w \rangle_{\mathfrak{v}_{2,0}} = 1$. Then the basis $w_1 = w$, $w_2 = J_2 J_1 w$, $w_3 = J_1 w$, and $w_4 = J_2 w$ is integral and $\langle w_i, w_i \rangle_{\mathfrak{v}_{2,0}} = 1$, $i = 1, \dots, 4$.

In the admissible module $(\mathfrak{v}_{0,2}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,2}})$ we choose $\tilde{w} \in \mathfrak{v}_{0,2}$ with $\langle \tilde{w}, \tilde{w} \rangle_{\mathfrak{v}_{0,2}} = 1$ and construct the integral basis $\tilde{w}_1 = \tilde{w}$, $\tilde{w}_2 = J_1 J_2 \tilde{w}$, $\tilde{w}_3 = J_1 \tilde{w}$, and $\tilde{w}_4 = J_2 \tilde{w}$ with $\langle \tilde{w}_i, \tilde{w}_i \rangle_{\mathfrak{v}_{0,2}} = \epsilon_i(2, 2)$.

The commutation relations with respect to both bases are equal, see Table 2, and this leads to the integral isomorphism. $\varphi_{2,0}: \mathfrak{n}_{2,0} \rightarrow \mathfrak{n}_{0,2}$ defined by

$$w_1 \mapsto \tilde{w}_1, \quad w_2 \mapsto \tilde{w}_2, \quad w_3 \mapsto \tilde{w}_3, \quad w_4 \mapsto \tilde{w}_4, \quad Z_1 \mapsto \tilde{Z}_1, \quad Z_2 \mapsto \tilde{Z}_2.$$

Isomorphism of $\mathfrak{n}_{4,0}$ and $\mathfrak{n}_{0,4}$. Let $(\mathfrak{v}_{4,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{4,0}})$ be an admissible module with $w \in \mathfrak{v}_{4,0}$ such that $J_1 J_2 J_3 J_4 w = w$ and $\langle w, w \rangle_{\mathfrak{v}_{4,0}} = 1$. Then the basis

$$\begin{aligned} w_1 &= w, & w_2 &= J_1 J_2 w, & w_3 &= J_1 J_3 w, & w_4 &= J_1 J_4 w, \\ w_5 &= J_1 w, & w_6 &= J_2 w, & w_7 &= J_3 w, & w_8 &= J_4 w, \end{aligned}$$

with $\langle w_i, w_i \rangle_{\mathfrak{v}_{4,0}} = \epsilon_i(8, 0) = 1$ is integral, with commutation relations in Table 3.

TABLE 2. Commutation relations on $\mathfrak{n}_{2,0}$ and $\mathfrak{n}_{0,2}$

[row, col.]	w_1	w_2	w_3	w_4
w_1	0	0	Z_1	Z_2
w_2	0	0	$-Z_2$	Z_1
w_3	$-Z_1$	Z_2	0	0
w_4	$-Z_2$	$-Z_1$	0	0

 TABLE 3. Commutation relations on $\mathfrak{n}_{4,0}$

[row, col.]	w_1	w_2	w_3	w_4	w_5	w_6	w_7	w_8
w_1	0	0	0	0	Z_1	Z_2	Z_3	Z_4
w_2	0	0	0	0	Z_2	$-Z_1$	$-Z_4$	Z_3
w_3	0	0	0	0	Z_3	Z_4	$-Z_1$	$-Z_2$
w_4	0	0	0	0	Z_4	$-Z_3$	Z_2	$-Z_1$
w_5	$-Z_1$	$-Z_2$	$-Z_3$	$-Z_4$	0	0	0	0
w_6	$-Z_2$	Z_1	$-Z_4$	Z_3	0	0	0	0
w_7	$-Z_3$	Z_4	Z_1	$-Z_2$	0	0	0	0
w_8	$-Z_4$	$-Z_3$	Z_2	Z_1	0	0	0	0

Let $(\mathfrak{v}_{0,4}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,4}})$ be an admissible module and $w \in \mathfrak{v}_{0,4}$ be such that $J_1 J_2 J_3 J_4 w = w$ and $\langle \tilde{w}, \tilde{w} \rangle_{\mathfrak{v}_{0,4}} = 1$. Then the basis

$$\begin{aligned} \tilde{w}_1 &= \tilde{w}, & \tilde{w}_2 &= J_1 J_2 \tilde{w}, & \tilde{w}_3 &= J_1 J_3 \tilde{w}, & \tilde{w}_4 &= J_1 J_4 \tilde{w}, \\ \tilde{w}_5 &= J_1 \tilde{w}, & \tilde{w}_6 &= J_2 \tilde{w}, & \tilde{w}_7 &= J_3 \tilde{w}, & \tilde{w}_8 &= J_4 \tilde{w}, \end{aligned}$$

with $\langle \tilde{w}_i, \tilde{w}_i \rangle_{\mathfrak{v}_{0,4}} = \epsilon_i(4, 4)$ is integral with commutation relations listed in Table 4.

 TABLE 4. Commutation relations on $\mathfrak{n}_{0,4}$

[row, col.]	\tilde{w}_1	\tilde{w}_2	\tilde{w}_3	\tilde{w}_4	\tilde{w}_5	\tilde{w}_6	\tilde{w}_7	\tilde{w}_8
\tilde{w}_1	0	0	0	0	\tilde{Z}_1	\tilde{Z}_2	\tilde{Z}_3	\tilde{Z}_4
\tilde{w}_2	0	0	0	0	$-\tilde{Z}_2$	\tilde{Z}_1	\tilde{Z}_4	$-\tilde{Z}_3$
\tilde{w}_3	0	0	0	0	$-\tilde{Z}_3$	$-\tilde{Z}_4$	\tilde{Z}_1	\tilde{Z}_2
\tilde{w}_4	0	0	0	0	$-\tilde{Z}_4$	\tilde{Z}_3	$-\tilde{Z}_2$	\tilde{Z}_1
\tilde{w}_5	$-\tilde{Z}_1$	\tilde{Z}_2	\tilde{Z}_3	\tilde{Z}_4	0	0	0	0
\tilde{w}_6	$-\tilde{Z}_2$	$-\tilde{Z}_1$	\tilde{Z}_4	$-\tilde{Z}_3$	0	0	0	0
\tilde{w}_7	$-\tilde{Z}_3$	$-\tilde{Z}_4$	$-\tilde{Z}_1$	\tilde{Z}_2	0	0	0	0
\tilde{w}_8	$-\tilde{Z}_4$	\tilde{Z}_3	$-\tilde{Z}_2$	$-\tilde{Z}_1$	0	0	0	0

We see from Tables 3 and 4 that the linear map $\varphi_{4,0}: \mathfrak{n}_{4,0} \rightarrow \mathfrak{n}_{0,4}$ defined by

$$\begin{aligned} w_i &\mapsto \tilde{w}_i & \text{if } & i = 1, 5, 6, 7, 8, \\ w_i &\mapsto -\tilde{w}_i & \text{if } & i = 2, 3, 4, \\ Z_k &\mapsto \tilde{Z}_k & \text{if } & k = 1, 2, 3, 4, \end{aligned}$$

is an integral isomorphism.

Isomorphism of $\mathfrak{n}_{8,0}$ and $\mathfrak{n}_{0,8}$. Let $\{Z_1, \dots, Z_8\}$ be an orthonormal basis for $\mathbb{R}^{8,0}$. Take a minimal admissible module $(\mathfrak{v}_{8,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{8,0}})$ and choose $w \in \mathfrak{v}_{8,0}$ with $\langle w, w \rangle_{\mathfrak{v}_{8,0}} = 1$ such that

$$J_1 J_2 J_3 J_4 w = J_1 J_2 J_5 J_6 w = J_2 J_3 J_5 J_7 w = J_1 J_2 J_7 J_8 w = w.$$

A method of work [2] shows that the basis

$$\begin{aligned}
u_1 &:= w, & u_2 &:= J_1 J_2 w, & u_3 &:= J_1 J_3 w, & u_4 &:= J_1 J_4 w, \\
u_5 &:= J_1 J_5 w, & u_6 &:= J_1 J_6 w, & u_7 &:= J_1 J_7 w, & u_8 &:= J_1 J_8 w, \\
u_9 &:= J_1 w, & u_{10} &:= J_2 w, & u_{11} &:= J_3 w, & u_{12} &:= J_4 w, \\
u_{13} &:= J_5 w, & u_{14} &:= J_6 w, & u_{15} &:= J_7 w, & u_{16} &:= J_8 w,
\end{aligned} \tag{12}$$

is orthonormal and satisfies $\langle u_i, u_i \rangle_{\mathfrak{v}_{8,0}} = \epsilon_i(16, 0) = 1$. Thus the minimal admissible module $(\mathfrak{v}_{8,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{8,0}})$ receives the integral basis (12).

Let $\{\tilde{Z}_1, \dots, \tilde{Z}_8\}$ be an orthonormal basis for $\mathbb{R}^{0,8}$. Given a minimal admissible module $(\mathfrak{v}_{0,8}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,8}})$, we choose a vector $w^1 \in \mathfrak{v}_{0,8}$ with $\langle w^1, w^1 \rangle_{\mathfrak{v}_{0,8}} = 1$ such that

$$\tilde{J}_1 \tilde{J}_2 \tilde{J}_3 \tilde{J}_4 w^1 = \tilde{J}_1 \tilde{J}_2 \tilde{J}_5 \tilde{J}_6 w^1 = \tilde{J}_2 \tilde{J}_3 \tilde{J}_5 \tilde{J}_7 w^1 = \tilde{J}_1 \tilde{J}_2 \tilde{J}_7 \tilde{J}_8 w^1 = w^1.$$

Then the orthonormal basis

$$\begin{aligned}
v_1 &:= w^1, & v_2 &:= \tilde{J}_1 \tilde{J}_2 w^1, & v_3 &:= \tilde{J}_1 \tilde{J}_3 w^1, & v_4 &:= \tilde{J}_1 \tilde{J}_4 w^1, \\
v_5 &:= \tilde{J}_1 \tilde{J}_5 w^1, & v_6 &:= \tilde{J}_1 \tilde{J}_6 w^1, & v_7 &:= \tilde{J}_1 \tilde{J}_7 w^1, & v_8 &:= \tilde{J}_1 \tilde{J}_8 w^1, \\
v_9 &:= \tilde{J}_1 w^1, & v_{10} &:= \tilde{J}_2 w^1, & v_{11} &:= \tilde{J}_3 w^1, & v_{12} &:= \tilde{J}_4 w^1, \\
v_{13} &:= \tilde{J}_5 w^1, & v_{14} &:= \tilde{J}_6 w^1, & v_{15} &:= \tilde{J}_7 w^1, & v_{16} &:= \tilde{J}_8 w^1,
\end{aligned} \tag{13}$$

with $\langle v_i, v_i \rangle_{\mathfrak{v}_{0,8}} = \epsilon_i(8, 8)$ is integral, see [2]. Tables 10 and 11 in the Appendix show the non-vanishing commutation relations on the pseudo H -type Lie algebras $\mathfrak{n}_{8,0}$ and $\mathfrak{n}_{0,8}$. It allows us to construct the Lie algebra integral isomorphism

$$\varphi_{8,0}: \begin{cases} u_i \mapsto v_i & \text{if } i = 1, 9, 10, \dots, 16, \\ u_i \mapsto -v_i & \text{if } i = 2, \dots, 8, \\ Z_k \mapsto \tilde{Z}_k & \text{if } k = 1, \dots, 8. \end{cases} \tag{14}$$

□

4.2. Structure constants for $\mathfrak{n}_{r+8,s}$ and $\mathfrak{n}_{r,s+8}$. This subsection is purely technical and auxiliary for the upcoming classification. A result of [2] gives an integral basis which satisfies Theorem 2.12 for all admissible Clifford modules $\mathfrak{v}_{r,s}$. Furthermore, the authors proved that it is possible to obtain any minimal admissible integral module $\mathfrak{v}_{t,u}$ by taking the tensor product of minimal admissible integral $\mathfrak{v}_{r,s}$ -modules $0 \leq r, s \leq 8$ by the minimal admissible integral modules $\mathfrak{v}_{8,0}$, $\mathfrak{v}_{0,8}$ or $\mathfrak{v}_{4,4}$.

Proposition 4.4. [2] *Consider two minimal admissible integral modules $(\mathfrak{v}_{r,s}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}})$ and $(\mathfrak{v}_{0,8}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,8}})$, where the representations $J_{\tilde{Z}_j}: \text{Cl}_{0,8} \rightarrow \text{End}(\mathfrak{v}_{0,8})$ permute the integral basis of $\mathfrak{v}_{0,8}$ up to sign for all orthonormal generators $\tilde{Z}_j \in \mathbb{R}^{0,8}$. Then the scalar product space given by the tensor product $(\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{0,8}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}} \cdot \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,8}})$ is a minimal admissible integral module $(\mathfrak{v}_{r,s+8}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s+8}})$.*

Remark 4.5. *In Proposition 4.4 one can change the minimal admissible integral module $(\mathfrak{v}_{0,8}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,8}})$ to the minimal admissible integral module $(\mathfrak{v}_{8,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{8,0}})$ or $(\mathfrak{v}_{4,4}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{4,4}})$ and, taking the tensor product, to obtain the modules*

$$\begin{aligned}
(\mathfrak{v}_{r+8,s}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r+8,s}}) &= (\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{8,0}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}} \cdot \langle \cdot, \cdot \rangle_{\mathfrak{v}_{8,0}}) \\
(\mathfrak{v}_{r+4,s+4}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r+4,s+4}}) &= (\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{4,4}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}} \cdot \langle \cdot, \cdot \rangle_{\mathfrak{v}_{4,4}})
\end{aligned}$$

respectively, which are minimal admissible integral. Details can be found in [2].

A pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$ is called *extended* if its minimal admissible integral module $\mathfrak{v}_{r,s}$ was constructed as in Proposition 4.4 or in Remark 4.5. The tensor products can be taken several times and with different spaces. Before we show how the structure constants for the Lie algebras $\mathfrak{n}_{r+8,s}$, $\mathfrak{n}_{r+4,s+4}$ and $\mathfrak{n}_{r,s+8}$ depend on the structure constants of $\mathfrak{n}_{r,s}$, $\mathfrak{n}_{8,0}$, $\mathfrak{n}_{0,8}$ and $\mathfrak{n}_{4,4}$, we state the notation that will be used in the forthcoming sections. We write \mathfrak{B}_V for an integral basis of the space V . Thus

$$\mathfrak{B}_{\mathfrak{z}_{r,s}} := \{Z_1^{r,s}, \dots, Z_{r+s}^{r,s}\}, \quad \langle Z_k^{r,s}, Z_m^{r,s} \rangle_{\mathfrak{z}_{r,s}} = \epsilon_k(r,s)\delta_{km},$$

$$\mathfrak{B}_{\mathfrak{v}_{r,s}} := \{w_1, \dots, w_{2l}\}, \quad \langle w_i, w_j \rangle_{\mathfrak{v}_{r,s}} = \begin{cases} \epsilon_i(l,l)\delta_{ij} & \text{for } s \neq 0, \\ \delta_{ij} & \text{for } s = 0, \end{cases}$$

and define $\mathfrak{B}_{\mathfrak{n}_{r,s}} := \mathfrak{B}_{\mathfrak{v}_{r,s}} \cup \mathfrak{B}_{\mathfrak{z}_{r,s}}$.

We fix the letters u , v , and y for the following bases given by (12), and a modification of (14) and (20)

$$\begin{aligned} \mathfrak{B}_{\mathfrak{v}_{8,0}} &= \{u_1, \dots, u_{16}\}, & \mathfrak{B}_{\mathfrak{v}_{0,8}} &= \{v_1, \dots, v_{16}\}, & \mathfrak{B}_{\mathfrak{v}_{4,4}} &= \{y_1, \dots, y_{16}\}, \\ \mathfrak{B}_{\mathfrak{z}_{8,0}} &= \{Z_1^{8,0}, \dots, Z_8^{8,0}\}, & \mathfrak{B}_{\mathfrak{z}_{0,8}} &= \{Z_1^{0,8}, \dots, Z_8^{0,8}\}, & \mathfrak{B}_{\mathfrak{z}_{4,4}} &= \{Z_1^{4,4}, \dots, Z_8^{4,4}\} \end{aligned}$$

with

$$\begin{aligned} \langle u_i, u_j \rangle_{\mathfrak{v}_{8,0}} &= \delta_{ij}, & \langle v_i, v_j \rangle_{\mathfrak{v}_{0,8}} &= \epsilon_i(8,8)\delta_{ij}, & \langle y_i, y_j \rangle_{\mathfrak{v}_{4,4}} &= \epsilon_i(8,8)\delta_{ij}, \\ \langle Z_k^{8,0}, Z_m^{8,0} \rangle_{\mathfrak{z}_{8,0}} &= \delta_{km}, & \langle Z_k^{0,8}, Z_m^{0,8} \rangle_{\mathfrak{z}_{0,8}} &= -\delta_{km}, & \langle Z_k^{4,4}, Z_m^{4,4} \rangle_{\mathfrak{z}_{4,4}} &= \epsilon_k(4,4)\delta_{km}, \end{aligned}$$

such that $\mathfrak{B}_{\mathfrak{n}_{8,0}}$ and $\mathfrak{B}_{\mathfrak{n}_{0,8}}$ have the same structural constants by Theorem 4.3, i.e.

$$\begin{aligned} \varphi_{8,0}(u_i) &= v_i, & \text{for all } i = 1, \dots, 16, \\ \varphi_{8,0}(Z_k^{8,0}) &= Z_k^{0,8}, & \text{for all } k = 1, \dots, 8. \end{aligned}$$

If $\mathfrak{v}_{t,u}$ is obtained by taking the tensor product of $\mathfrak{v}_{r,s}$ by one of the $\mathfrak{v}_{8,0}$, $\mathfrak{v}_{0,8}$ or $\mathfrak{v}_{4,4}$, we write for the basis

$$\mathfrak{B}_{\mathfrak{n}_{t,u}} = \{w_i \otimes \alpha_j, Z_m^{t,u} \mid i = 1, \dots, 2l, j = 1, \dots, 16, m = 1, \dots, r+s+8\},$$

where $w_i \in \mathfrak{B}_{\mathfrak{v}_{r,s}}$, the vectors α_j are from the corresponding integral bases $\mathfrak{B}_{\mathfrak{v}_{8,0}}$, $\mathfrak{B}_{\mathfrak{v}_{0,8}}$ or $\mathfrak{B}_{\mathfrak{v}_{4,4}}$, and $Z_m^{t,u} \in \mathfrak{B}_{\mathfrak{z}_{r+p,s+q}} = \mathfrak{B}_{\mathfrak{z}_{t,u}}$, with (p,q) equal to one of the pairs $(8,0)$, $(0,8)$, $(4,4)$. For definiteness we preserve the order of the elements in $\mathfrak{B}_{\mathfrak{z}_{t,u}}$ and we write first those which have positive squares of the scalar product and then those with negative squares of the scalar product.

Lemma 4.6. *Let $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$ be a pseudo H -type algebra and A_{ij}^m the structure constants with respect to the integral basis $\mathfrak{B}_{\mathfrak{n}_{r,s}}$. Let $\mathfrak{n}_{8,0} = \mathfrak{v}_{8,0} \oplus \mathfrak{z}_{8,0}$ has the integral basis $\mathfrak{B}_{\mathfrak{n}_{8,0}}$ with the corresponding structure constants \bar{A}_{ij}^m . Then the Lie algebra $\mathfrak{n}_{r+8,s} = (\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{8,0}) \oplus \mathfrak{z}_{r+8,s}$ has the following structure constants $\tilde{A}_{ij,pq}^m$ with respect to the integral basis $\mathfrak{B}_{\mathfrak{n}_{r+8,s}}$.*

If $s = 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} -A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 1, \dots, 8, \\ A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 9, \dots, 16, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+8 & \text{and } i = p = 1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (15)$$

If $s > 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} -A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 1, \dots, 8, \\ -A_{ip}^{m-8} & \text{if } m = r+8+1, \dots, r+8+s & \text{and } j = q = 1, \dots, 8, \\ A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 9, \dots, 16, \\ A_{ip}^{m-8} & \text{if } m = r+8+1, \dots, r+8+s & \text{and } j = q = 9, \dots, 16, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+8 & \text{and } i = p = 1, \dots, l, \\ -\bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+8 & \text{and } i = p = l+1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (16)$$

Proof. We recall that the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}_{r+8,s}}$ of $\mathfrak{v}_{r+8,s}$ is given by the product $\langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}} \cdot \langle \cdot, \cdot \rangle_{\mathfrak{v}_{8,0}}$. To shorten the notation we write

$$J_{Z_m}^{r,s} = J_{Z_m} : \mathfrak{v}_{r,s} \rightarrow \mathfrak{v}_{r,s}, \quad J_{Z_m}^{8,0} = \bar{J}_{Z_m} : \mathfrak{v}_{8,0} \rightarrow \mathfrak{v}_{8,0}, \quad J_{Z_m}^{r+8,s} = \tilde{J}_{Z_m} : \mathfrak{v}_{r+8,s} \rightarrow \mathfrak{v}_{r+8,s}$$

and the operator $E := \bar{J}_{Z_1} \cdots \bar{J}_{Z_8} : \mathfrak{v}_{8,0} \rightarrow \mathfrak{v}_{8,0}$ with the properties

$$E^2 = \text{Id}_{\mathfrak{v}_{8,0}}, \quad E\bar{J}_{Z_j} = -\bar{J}_{Z_j}E, \quad j = 1, \dots, 8, \quad \langle Eu, u^* \rangle_{\mathfrak{v}_{8,0}} = \langle u, Eu^* \rangle_{\mathfrak{v}_{8,0}}, \quad u, u^* \in \mathfrak{v}_{8,0}.$$

It leads to the following equalities:

$$\begin{aligned} \langle \tilde{J}_{Z_m} \tilde{w}_{i,j}, \tilde{w}_{p,q} \rangle_{\mathfrak{v}_{r+8,s}} &= \langle J_{Z_m} w_i, w_p \rangle_{\mathfrak{v}_{r,s}} \langle Eu_j, u_q \rangle_{\mathfrak{v}_{8,0}}, & m = 1, \dots, r, \\ \langle \tilde{J}_{Z_m} \tilde{w}_{i,j}, \tilde{w}_{p,q} \rangle_{\mathfrak{v}_{r+8,s}} &= \langle w_i, w_p \rangle_{\mathfrak{v}_{r,s}} \langle \bar{J}_{Z_{m-r}} u_j, u_q \rangle_{\mathfrak{v}_{8,0}}, & m = r+1, \dots, r+8, \\ \langle \tilde{J}_{Z_m} \tilde{w}_{i,j}, \tilde{w}_{p,q} \rangle_{\mathfrak{v}_{r+8,s}} &= \langle J_{Z_{m-8}} w_i, w_p \rangle_{\mathfrak{v}_{r,s}} \langle Eu_j, u_q \rangle_{\mathfrak{v}_{8,0}}, & m = r+9, \dots, r+s+8, \end{aligned} \quad (17)$$

with the integral basis $\{\tilde{w}_{i,j} = w_i \otimes u_j, \tilde{Z}_m | i = 1, \dots, 2l, j = 1, \dots, 16, m = 1, \dots, r+s+8\}$ by [2]. Similar equations for the cases $\mathfrak{n}_{0,8}$ and $\mathfrak{n}_{4,4}$ can be found in [2].

Let $m = 1, \dots, r$ and note that the mapping E acts on the integral basis $\mathfrak{B}_{\mathfrak{v}_{8,0}}$ by $Eu_j = -u_j \epsilon_j(8, 8)$, which follows from the permutation Table 13 in the Appendix. That leads to $\langle Eu_j, u_q \rangle_{\mathfrak{v}_{8,0}} = -\epsilon_j(8, 8) \delta_{jq}$. Then the first equation in (17) gives

$$\tilde{B}_{ij,pq}^m \epsilon_p^{\mathfrak{v}_{r+8,s}} = -B_{i,p}^m \epsilon_p^{\mathfrak{v}_{r,s}} \epsilon_j(8, 8) \delta_{jq}$$

by (7). Making use of formula (8) we obtain the following equations for structure constants

$$\tilde{A}_{ij,pq}^m (\epsilon_{pq}^{\mathfrak{v}_{r+8,s}})^2 \epsilon_m^{\mathfrak{z}_{r+8,s}} = -A_{i,p}^m (\epsilon_p^{\mathfrak{v}_{r,s}})^2 \epsilon_m^{\mathfrak{z}_{r,s}} \epsilon_j(8, 8) \delta_{jq}$$

that yields to the first two lines in formula (15) and corresponding lines in (16). Arguing in a similar way for the rest of the formulas in (17) we obtain

$$\begin{aligned} \epsilon_m^{\mathfrak{z}_{r+8,s}} \tilde{A}_{ij,pq}^m &= -A_{i,p}^m \epsilon_m^{\mathfrak{z}_{r,s}} \epsilon_j(8, 8) \delta_{jq}, & m = 1, \dots, r, \\ \epsilon_m^{\mathfrak{z}_{r+8,s}} \tilde{A}_{ij,pq}^m &= \bar{A}_{jq}^{m-r} \epsilon_{m-r}^{\mathfrak{z}_{8,0}} \epsilon_i^{\mathfrak{v}_{r,s}} \delta_{ip}, & m = r+1, \dots, r+8, \\ \epsilon_m^{\mathfrak{z}_{r+8,s}} \tilde{A}_{ij,pq}^m &= -A_{ip}^{m-8} \epsilon_{m-8}^{\mathfrak{z}_{r,s}} \epsilon_j(8, 8) \delta_{jq}, & m = r+9, \dots, r+s+8. \end{aligned}$$

This implies (15) and (16). \square

Lemma 4.7. *Let $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$ be a pseudo H -type algebra with the structure constants A_{ij}^m written with respect to $\mathfrak{B}_{\mathfrak{n}_{r,s}}$ and $\mathfrak{n}_{0,8} = \mathfrak{v}_{0,8} \oplus \mathfrak{z}_{0,8}$ with the integral basis $\mathfrak{B}_{\mathfrak{n}_{0,8}}$, and the corresponding structure constants \bar{A}_{ij}^m . Then the Lie algebra $\mathfrak{n}_{r,s+8} = (\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{0,8}) \oplus \mathfrak{z}_{r,s+8}$ has the following structure constants $\tilde{A}_{ij,pq}^m$ with respect to the integral basis $\mathfrak{B}_{\mathfrak{n}_{r,s+8}}$.*

If $s = 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} -A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 1, \dots, 16, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+8 & \text{and } i = p = 1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (18)$$

If $s > 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} -A_{ip}^m & \text{if } m = 1, \dots, r+s & \text{and } j = q = 1, \dots, 16, \\ \bar{A}_{jq}^{m-r-s} & \text{if } m = r+s+1, \dots, r+s+8 & \text{and } i = p = 1, \dots, l, \\ -\bar{A}_{jq}^{m-r-s} & \text{if } m = r+s+1, \dots, r+s+8 & \text{and } i = p = l+1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (19)$$

The proof of Lemma 4.7 is analogous to the proof of Lemma 4.6.

Before we present the structure constants for the Lie algebra $\mathfrak{n}_{r+4,s+4}$ we write the integral basis for the pseudo H -type Lie algebra $\mathfrak{n}_{4,4}$.

$$\begin{aligned} y_1 &= w, & y_2 &= J_1 w, & y_3 &= J_2 w, & y_4 &= J_3 w, \\ y_5 &= J_4 w, & y_6 &= J_1 J_2 w, & y_7 &= J_1 J_3 w, & y_8 &= J_1 J_4 w, \\ y_9 &= J_5 w, & y_{10} &= J_6 w, & y_{11} &= J_7 w, & y_{12} &= J_8 w, \\ y_{13} &= J_1 J_5 w, & y_{14} &= J_1 J_6 w, & y_{15} &= J_1 J_7 w, & y_{16} &= J_1 J_8 w, \end{aligned} \quad (20)$$

for $J_1 J_2 J_3 J_4 w = J_1 J_2 J_5 J_6 w = J_2 J_3 J_5 J_7 w = J_1 J_2 J_7 J_8 w = w$ with

$$\langle w_i, w_i \rangle_{\mathfrak{v}_{4,4}} = \epsilon_i(8, 8), \quad \langle Z_k, Z_k \rangle_{\mathfrak{z}_{4,4}} = \epsilon_k(4, 4).$$

Lemma 4.8. *Let $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$ has the structure constants A_{ij}^m with respect to $\mathfrak{B}_{\mathfrak{n}_{r,s}}$ and $\mathfrak{n}_{4,4} = \mathfrak{v}_{4,4} \oplus \mathfrak{z}_{4,4}$ has the structure constants \bar{A}_{ij}^m with respect to the integral basis $\mathfrak{B}_{\mathfrak{n}_{4,4}}$. Then the Lie algebra $\mathfrak{n}_{r+4,s+4} = (\mathfrak{v}_{r,s} \otimes \mathfrak{v}_{4,4}) \oplus \mathfrak{z}_{r+4,s+4}$ has the following structure constants $\tilde{A}_{ij,pq}^m$ with respect to the integral basis $\mathfrak{B}_{\mathfrak{n}_{r+4,s+4}}$.*

If $s = 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 2, \dots, 5, 13, \dots, 16, \\ -A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 1, 6, \dots, 12, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+8 & \text{and } i = p = 1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (21)$$

If $s > 0$, then

$$\tilde{A}_{ij,pq}^m = \begin{cases} A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 2, \dots, 5, 13, \dots, 16, \\ -A_{ip}^m & \text{if } m = 1, \dots, r & \text{and } j = q = 1, 6, \dots, 12, \\ A_{ip}^{m-8} & \text{if } m = r+9, \dots, r+s+8 & \text{and } j = q = 2, \dots, 5, 13, \dots, 16, \\ -A_{ip}^{m-8} & \text{if } m = r+5, \dots, r+s+4 & \text{and } j = q = 1, 6, \dots, 12, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+4 & \text{and } i = p = 1, \dots, l, \\ \bar{A}_{jq}^{m-r} & \text{if } m = r+5, \dots, r+8 & \text{and } i = p = 1, \dots, l, \\ -\bar{A}_{jq}^{m-r} & \text{if } m = r+1, \dots, r+4 & \text{and } i = p = l+1, \dots, 2l, \\ -\bar{A}_{jq}^{m-r} & \text{if } m = r+5, \dots, r+8 & \text{and } i = p = l+1, \dots, 2l, \\ 0 & \text{otherwise.} \end{cases} \quad (22)$$

The proof of Lemma 4.8 is analogous to the proof of Lemma 4.6.

4.3. Classification of $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ for $r > 8$. We observe an interesting property of some of the H -type Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ that will be used in the proof of Theorem 4.11.

Definition 4.9. *We say that an orthonormal basis $\{w_1, \dots, w_{2l}, Z_1, \dots, Z_n\}$ of a pseudo H -type Lie algebra is of block-type if $[w_i, w_j] = 0$ for both indices $i, j = 1, \dots, l$ and $i, j = l + 1, \dots, 2l$. We call a pseudo H -type Lie algebra of block-type if it has a block-type basis.*

Lemma 4.10. *The pseudo H -type algebras $\mathfrak{n}_{r,0}$ with $r \in \{0, 1, 2, 4\} \pmod{8}$ and $\mathfrak{n}_{0,s}$ with $s \in \mathbb{N}$ are of block-type.*

Proof. We prove by induction that $\mathfrak{n}_{r,0}$ with $r \in \{0, 1, 2, 4\} \pmod{8}$ is of block-type. The base of induction follows from Tables 1, 2, 3, 10 of non-vanishing commutators on $\mathfrak{n}_{1,0}$, $\mathfrak{n}_{2,0}$, $\mathfrak{n}_{4,0}$ and $\mathfrak{n}_{8,0}$. For the induction step we assume that $\mathfrak{n}_{r,0}$ with $r \in \{0, 1, 2, 4\} \pmod{8}$ has a block-type basis $\{w_1, \dots, w_{2l}, Z_1^{r,0}, \dots, Z_r^{r,0}\}$ with $[w_i, w_j] = 0$ when both indices $i, j = 1, \dots, l$ and $i, j = l + 1, \dots, 2l$. By extension we construct the Lie algebra $\mathfrak{n}_{r+8,0}$ of dimension $32l + r + 8$ with the basis $\{w_1 \otimes u_1, \dots, w_{2l} \otimes u_{16}, Z_1^{r+8,0}, \dots, Z_{r+8}^{r+8,0}\}$. Equations (17) imply that $[w_i \otimes u_j, w_p \otimes u_q] = 0$ for the following cases:

- $i = p$ and both $j, q = 1, \dots, 8$ and $j, q = 9, \dots, 16$,
- $j = q$ and both $i, p = 1, \dots, l$ and $i, p = l + 1, \dots, 2l$,
- $i \neq p$ and $j \neq q$ or $i = p$ and $j = q$.

We define a decomposition of the basis vectors of $\mathfrak{v}_{r+8,0}$ by

$$\begin{aligned} \mathbf{A}_{r,0}^1 &:= \{w_i \otimes u_j \mid i = 1, \dots, l, \quad j = 1, \dots, 8\}, \\ \mathbf{A}_{r,0}^2 &:= \{w_i \otimes u_j \mid i = l + 1, \dots, 2l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{r,0}^1 &:= \{w_i \otimes u_j \mid i = 1, \dots, l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{r,0}^2 &:= \{w_i \otimes u_j \mid i = l + 1, \dots, 2l, \quad j = 1, \dots, 8\}, \\ \mathbf{A}_{r,0} &:= \mathbf{A}_{r,0}^1 \cup \mathbf{A}_{r,0}^2, \quad \mathbf{B}_{r,0} := \mathbf{B}_{r,0}^1 \cup \mathbf{B}_{r,0}^2. \end{aligned} \tag{23}$$

It follows that for any $\tilde{w}, \tilde{v} \in A_{r,0}$ and for any $\tilde{x}, \tilde{y} \in B_{r,0}$ we obtain that $[\tilde{w}, \tilde{v}] = 0 = [\tilde{x}, \tilde{y}]$. As the cardinality of the basis of $\mathfrak{v}_{r+8,s}$ is $32l$ and the cardinality of each of the sets $A_{r,0}$ and $B_{r,0}$ is $16l$ we proved that $\mathfrak{n}_{r,0}$ with $r \in \{0, 1, 2, 4\} \pmod{8}$ is of block-type.

Now we consider H -type Lie algebras $\mathfrak{n}_{0,s}$. The space $\mathfrak{v}_{0,s}$ is neutral with an integral basis $\{w_1, \dots, w_{2l}\}$ satisfying $\langle w_i, w_j \rangle_{l,l} = \epsilon_i(l, l) \delta_{ij}$. Let $\mathfrak{B}_{30,s} = \{Z_1^{0,s}, \dots, Z_s^{0,s}\}$. The map $J_{Z_k^{0,s}}$ is an anti-isometry and permutes the basis $\{w_1, \dots, w_{2l}\}$ for all $k = 1, \dots, s$. It follows that $B_{ij}^k = 0$ for $i, j = 1, \dots, l$ or $i, j = l + 1, \dots, 2l$. Then by $\epsilon_\beta^v B_{\alpha\beta}^k = \epsilon_k^3 A_{\alpha\beta}^k$ we obtain that $A_{ij}^k = 0$ when both indices $i, j = 1, \dots, l$ or $i, j = l + 1, \dots, 2l$. Hence $\mathfrak{n}_{0,s}$ is a block-type Lie algebra for all $s \in \mathbb{N}$. \square

Theorem 4.11. *The Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ are integral isomorphic if and only if $r \in \{0, 1, 2, 4\} \pmod{8}$.*

Proof. The H -type Lie algebras $\mathfrak{n}_{r,0}$ are integral isomorphic to $\mathfrak{n}_{0,r}$ for $r = 1, 2, 4, 8$ by Theorem 4.3. The Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ are non-isomorphic for $r = 3, 5, 6, 7$ by Theorem 4.12 due to the different dimensions which are preserved under the extension process. Thus it remains to show that the Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ are integral isomorphic if $r \in \{0, 1, 2, 4\} \pmod{8}$.

By induction we assume that $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$ are integral isomorphic for $r \in \{0, 1, 2, 4\}$ mod 8 with the integral block-type bases

$$\mathfrak{B}_{\mathfrak{n}_{r,0}} = \{w_1, \dots, w_{2l}, Z_1^{r,0}, \dots, Z_r^{r,0}\}, \quad \mathfrak{B}_{\mathfrak{n}_{0,r}} = \{x_1, \dots, x_{2l}, Z_1^{0,r}, \dots, Z_r^{0,r}\},$$

with equal structure constants A_{ij}^m , i.e. $\varphi_{r,0}(w_i) = x_i$ for all $i = 1, \dots, 2l$ and $\varphi_{r,0}(Z_k^{r,0}) = Z_k^{0,r}$ for all $k = 1, \dots, r$. Furthermore, we recall that in the integral block-type bases $\mathfrak{B}_{\mathfrak{n}_{8,0}}$ and $\mathfrak{B}_{\mathfrak{n}_{0,8}}$ the structure constants denoted by \bar{A}_{ij}^m are equal for both Lie algebras, i.e. $\varphi_{8,0}(u_i) = v_i$ for all $i = 1, \dots, 16$ and $\varphi_{8,0}(Z_k^{8,0}) = Z_k^{0,8}$ for all $k = 1, \dots, 8$.

We exploit Proposition 4.4 and Remark 4.5 and obtain the integral bases

$$\mathfrak{B}_{\mathfrak{n}_{r+8,0}} = \{w_i \otimes u_j, Z_m^{r+8,0}\}, \quad \mathfrak{B}_{\mathfrak{n}_{0,r+8}} = \{x_i \otimes v_j, Z_m^{0,r+8}\}.$$

We define the bijective linear map $\varphi_{r+8,0}: \mathfrak{n}_{r+8,0} \rightarrow \mathfrak{n}_{0,r+8}$ by

$$\begin{aligned} w_i \otimes u_j &\mapsto x_i \otimes v_j && \text{if } i \in \{1, \dots, l\}, && j \in \{1, \dots, 16\}, \\ w_i \otimes u_j &\mapsto x_i \otimes v_j && \text{if } i \in \{l+1, \dots, 2l\}, && j \in \{1, \dots, 8\}, \\ w_i \otimes u_j &\mapsto -x_i \otimes v_j && \text{if } i \in \{l+1, \dots, 2l\}, && j \in \{9, \dots, 16\}, \\ Z_m^{r+8,0} &\mapsto Z_m^{0,r+8} && \text{if } m \in \{1, \dots, r+8\}. \end{aligned}$$

It remains to prove that $\varphi_{r+8,0}$ is a Lie algebra isomorphism, i.e. $\varphi_{r+8,0}([w_i \otimes u_j, w_p \otimes u_q]) = [\varphi_{r+8,0}(w_i \otimes u_j), \varphi_{r+8,0}(w_p \otimes u_q)]$. We know that the structural constants $\tilde{A}_{ij,pq}^m$ of $[w_i \otimes u_j, w_p \otimes u_q]$ are given by formula (15) and that the structural constants $C_{ij,pq}^m$ for $[x_i \otimes v_j, x_p \otimes v_q]$ are given by formula (19), where we have to put the index r instead of $r+s$. It follows that if $i \neq p$ and $j \neq q$ or $i = p$ and $j = q$ the commutators vanish:

$$\varphi_{r+8,0}([w_i \otimes u_j, w_p \otimes u_q]) = \varphi_{r+8,0}(0) = 0 = \pm[x_i \otimes v_j, x_p \otimes v_q].$$

Let us consider the case $i = p$ and $j \neq q$.

- if $i = p = 1, \dots, l$, then:

$$\varphi_{r+8,0}([w_i \otimes u_j, w_i \otimes u_q]) = \bar{A}_{jq}^{m-r} \varphi_{r+8,0}(Z_m^{r+8,0}) = \bar{A}_{jq}^{m-r} Z_m^{0,r+8},$$

$$[\varphi_{r+8,0}(w_i \otimes u_j), \varphi_{r+8,0}(w_i \otimes u_q)] = [x_i \otimes v_j, x_i \otimes v_q] = \bar{A}_{jq}^{m-r} Z_m^{0,r+8}$$

with $m = r+1, \dots, r+8$ by formulas (15) and (19).

- if $i = p = l+1, \dots, 2l$, then we use Lemma 4.10.

$$\varphi_{r+8,0}([w_i \otimes u_j, w_i \otimes u_q]) = \bar{A}_{jq}^{m-r} \varphi_{r+8,0}(Z_m^{r+8,0}) = \bar{A}_{jq}^{m-r} Z_m^{0,r+8},$$

$$\begin{aligned} &[\varphi_{r+8,0}(w_i \otimes u_j), \varphi_{r+8,0}(w_i \otimes u_q)] \\ &= \begin{cases} [x_i \otimes v_j, x_i \otimes v_q] & \text{if } j, q = 1, \dots, 8 \text{ or } j, q = 9, \dots, 16, \\ -[x_i \otimes v_j, x_i \otimes v_q] & \text{otherwise,} \end{cases} \\ &= \begin{cases} -\bar{A}_{jq}^{m-r} Z_m^{0,r+8} & \text{if } j, q = 1, \dots, 8 \text{ or } j, q = 9, \dots, 16, \\ \bar{A}_{jq}^{m-r} Z_m^{0,r+8} & \text{otherwise,} \end{cases} \end{aligned}$$

with $m = r+1, \dots, r+8$ by formulas (15), (19), and the definition of $\varphi_{r+8,0}$. We observe that $\bar{A}_{jq}^{m-r} = 0$ when for both indices j, q simultaneously either $j, q = 1, \dots, 8$ or $j, q = 9, \dots, 16$, since the Lie algebras $\mathfrak{n}_{8,0}$ and $\mathfrak{n}_{0,8}$ are of block type, see Table 10. We see that the map $\varphi_{r+8,0}$ satisfies the Lie algebra isomorphism properties in this case.

We turn to consider the case $i \neq p$ and $j = q$

- if $j = q = 1, \dots, 8$, then

$$\begin{aligned}\varphi_{r+8,0}([w_i \otimes u_j, w_p \otimes u_j]) &= -A_{ip}^m \varphi_{r+8,0}(Z_m^{r+8,0}) = -A_{ip}^m Z_m^{0,r+8}, \\ [\varphi_{r+8,0}(w_i \otimes u_j), \varphi_{r+8,0}(w_p \otimes u_j)] &= [x_i \otimes v_j, x_p \otimes v_j] = -A_{ip}^m Z_m^{0,r+8},\end{aligned}$$

with $m = 1, \dots, r$ by formulas (15) and (19).

- if $j = q = 9, \dots, 16$, then we use the block form of Lie algebras $\mathfrak{n}_{r,0}$ and $\mathfrak{n}_{0,r}$. We calculate as above

$$\varphi_{r+8,0}([w_i \otimes u_j, w_p \otimes u_j]) = A_{ip}^m \varphi_{r+8,0}(Z_m^{r+8,0}) = A_{ip}^m Z_m^{0,r+8}.$$

On the other side

$$\begin{aligned}& [\varphi_{r+8,0}(w_i \otimes u_j), \varphi_{r+8,0}(w_p \otimes u_j)] \\ &= \begin{cases} [x_i \otimes v_j, x_p \otimes v_q] & \text{if } i, p = 1, \dots, l \text{ or } i, p = l+1, \dots, 2l, \\ -[x_i \otimes v_j, x_p \otimes v_q] & \text{otherwise,} \end{cases} \\ &= \begin{cases} -A_{ip}^m Z_m^{0,r+8} = 0 & \text{if } i, p = 1, \dots, l \text{ or } i, p = l+1, \dots, 2l, \\ A_{ip}^m Z_m^{0,r+8} & \text{otherwise,} \end{cases}\end{aligned}$$

with $m = 1, \dots, r$ by formulas (15) and (19). This finishes the proof of the theorem. \square

Theorem 4.12. *The pseudo H-type Lie algebra $\mathfrak{n}_{r+8t,0}$ is not isomorphic to $\mathfrak{n}_{0,r+8t}$ for $r = 3, 5, 6, 7$ and a non-negative integer t .*

Proof. We prove the theorem by counting the dimensions of the Lie algebras. The dimensions of the minimal admissible modules are given by

$$\begin{aligned}\dim(\mathfrak{v}_{3+8t,0}) &= 4 \cdot 16^t \neq 8 \cdot 16^t = \dim(\mathfrak{v}_{0,3+8t}), \\ \dim(\mathfrak{v}_{r+8t,0}) &= 8 \cdot 16^t \neq 16 \cdot 16^t = \dim(\mathfrak{v}_{0,r+8t}), \quad \text{for } r = 5, 6, 7.\end{aligned}$$

\square

5. ISOMORPHISM OF LIE ALGEBRAS $\mathfrak{n}_{r,s}$ WITH $r, s \neq 0$

In this section we show, making use of the ideas developed in the previous section, that the Bott-periodicity is inherited in isomorphism properties of Lie algebras of block type.

5.1. Decompositions of bases $r, s \neq 0$. We recall the notations of the bases $\mathfrak{B}_{\mathfrak{n}_{r,s}}$ and state the result that extends the notion of the block type algebras.

Lemma 5.1. *Let us assume that the integral basis $\mathfrak{B}_{\mathfrak{v}_{r,s}}$, $r, s \neq 0$, for the pseudo H-type Lie algebra $\mathfrak{n}_{r,s}$ with $\dim(\mathfrak{v}_{r,s}) = 2l$ satisfies the following decomposition*

$$\begin{aligned}\mathfrak{B}_{\mathfrak{v}_{r,s}} &= \mathbf{A}_{r,s} \cup \mathbf{B}_{r,s}, \quad \text{card}(\mathbf{A}_{r,s}) = \text{card}(\mathbf{B}_{r,s}) = l, \\ [\mathbf{A}_{r,s}, \mathbf{A}_{r,s}] &= [\mathbf{B}_{r,s}, \mathbf{B}_{r,s}] = 0, \\ \mathbf{A}_{r,s} &= \mathbf{A}_{r,s}^+ \cup \mathbf{A}_{r,s}^-, \quad \mathbf{B}_{r,s} = \mathbf{B}_{r,s}^+ \cup \mathbf{B}_{r,s}^-, \quad \text{card}(\mathbf{A}_{r,s}^\pm) = \text{card}(\mathbf{B}_{r,s}^\pm) = \frac{l}{2},\end{aligned}\tag{24}$$

$$\text{where } \langle w_i, w_i \rangle_{\mathfrak{v}_{r,s}} = \begin{cases} 1 & \text{if } w_i \in \mathbf{A}_{r,s}^+ \cup \mathbf{B}_{r,s}^+, \\ -1 & \text{if } w_i \in \mathbf{A}_{r,s}^- \cup \mathbf{B}_{r,s}^-. \end{cases}$$

Then the extended pseudo H-type Lie algebras $\mathfrak{n}_{r+8,s}$, $\mathfrak{n}_{r,s+8}$, and $\mathfrak{n}_{r+4,s+4}$ admit a decomposition of type (24).

Proof. Define the following sets

$$\begin{aligned} \mathbf{A}_{8,0} &= \{u_1, \dots, u_8\}, & \mathbf{B}_{8,0} &= \{u_9, \dots, u_{16}\}, & \text{for } u_i &\in \mathfrak{B}_{\mathbf{v}_{8,0}} \\ \mathbf{A}_{0,8} &= \{v_1, \dots, v_8\}, & \mathbf{B}_{0,8} &= \{v_9, \dots, v_{16}\}, & \text{for } v_i &\in \mathfrak{B}_{\mathbf{v}_{0,8}} \end{aligned} \quad (25)$$

$$\begin{aligned} \mathbf{A}_{4,4}^+ &= \{y_1, y_6, y_7, y_8\}, & \mathbf{A}_{4,4}^- &= \{y_{13}, y_{14}, y_{15}, y_{16}\}, \\ \mathbf{B}_{4,4}^+ &= \{y_2, y_3, y_4, y_5\}, & \mathbf{B}_{4,4}^- &= \{y_9, y_{10}, y_{11}, y_{12}\}. \end{aligned} \quad (26)$$

for $y_i \in \mathfrak{B}_{\mathbf{v}_{4,4}}$ given by (20).

Now, making use of $\mathbf{A}_{r,s}$, $\mathbf{B}_{r,s}$ and (25), (26), we define the decompositions for the bases of the extended algebras.

$$\begin{aligned} \mathbf{A}_{r+8,s}^+ &= \mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{8,0} \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{B}_{8,0}, & \mathbf{A}_{r+8,s}^- &= \mathbf{A}_{r,s}^- \otimes \mathbf{A}_{8,0} \cup \mathbf{B}_{r,s}^- \otimes \mathbf{B}_{8,0}, \\ \mathbf{B}_{r+8,s}^+ &= \mathbf{A}_{r,s}^+ \otimes \mathbf{B}_{8,0} \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{A}_{8,0}, & \mathbf{B}_{r+8,s}^- &= \mathbf{A}_{r,s}^- \otimes \mathbf{B}_{8,0} \cup \mathbf{B}_{r,s}^- \otimes \mathbf{A}_{8,0}, \\ \mathbf{A}_{r,s+8}^+ &= \mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{0,8} \cup \mathbf{B}_{r,s}^- \otimes \mathbf{B}_{0,8}, & \mathbf{A}_{r,s+8}^- &= \mathbf{A}_{r,s}^- \otimes \mathbf{A}_{0,8} \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{B}_{0,8}, \\ \mathbf{B}_{r,s+8}^+ &= \mathbf{A}_{r,s}^+ \otimes \mathbf{B}_{0,8} \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{A}_{0,8}, & \mathbf{B}_{r,s+8}^- &= \mathbf{A}_{r,s}^- \otimes \mathbf{B}_{0,8} \cup \mathbf{B}_{r,s}^- \otimes \mathbf{A}_{0,8}, \end{aligned}$$

$$\begin{aligned} \mathbf{A}_{r+4,s+4}^+ &= \mathbf{B}_{r,s}^+ \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{B}_{r,s}^- \otimes \mathbf{B}_{4,4}^- \cup \mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{A}_{r,s}^- \otimes \mathbf{A}_{4,4}^-, \\ \mathbf{A}_{r+4,s+4}^- &= \mathbf{B}_{r,s}^- \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{B}_{4,4}^- \cup \mathbf{A}_{r,s}^- \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{4,4}^-, \\ \mathbf{B}_{r+4,s+4}^+ &= \mathbf{A}_{r,s}^+ \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{A}_{r,s}^- \otimes \mathbf{B}_{4,4}^- \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{B}_{r,s}^- \otimes \mathbf{A}_{4,4}^-, \\ \mathbf{B}_{r+4,s+4}^- &= \mathbf{A}_{r,s}^- \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{A}_{r,s}^+ \otimes \mathbf{B}_{4,4}^- \cup \mathbf{B}_{r,s}^- \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{B}_{r,s}^+ \otimes \mathbf{A}_{4,4}^-. \end{aligned}$$

All the necessary properties follows directly from the definition of the basis for the extended Lie algebras and Lemmas 4.6, 4.7, and 4.8. We illustrate only the proof of the following property $[\mathbf{A}_{r+8,s}, \mathbf{A}_{r+8,s}] = 0$, considering several cases. To show that $[\mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{8,0}, \mathbf{A}_{r,s}^+ \otimes \mathbf{A}_{8,0}] = 0$ we choose $w_i, w_j \in \mathbf{A}_{r,s}^+$ and $u_p, u_q \in \mathbf{A}_{8,0}$. Then

$$[w_i \otimes u_p, w_j \otimes u_q] = \begin{cases} 0 & \text{if } i \neq j, p \neq q, \text{ or } i = j, p = q, \\ [u_p, u_q] = 0 & \text{if } i = j, p \neq q, \text{ since } [u_p, u_q] \in [\mathbf{A}_{8,0}, \mathbf{A}_{8,0}] = 0, \\ -[w_i, w_j] = 0 & \text{if } i \neq j, p = q, \text{ since } [w_i, w_j] \in [\mathbf{A}_{r,s}^+, \mathbf{A}_{r,s}^+] = 0. \end{cases}$$

Analogously we show

$$[\mathbf{A}_{r,s}^\pm \otimes \mathbf{A}_{8,0}, \mathbf{A}_{r,s}^\pm \otimes \mathbf{A}_{8,0}] = [\mathbf{B}_{r,s}^\pm \otimes \mathbf{B}_{8,0}, \mathbf{B}_{r,s}^\pm \otimes \mathbf{B}_{8,0}] = 0$$

for any combinations of $+$ and $-$. Any term of the type $[\mathbf{A}_{r,s}^\pm \otimes \mathbf{A}_{8,0}, \mathbf{B}_{r,s}^\pm \otimes \mathbf{B}_{8,0}]$ vanishes since $\mathbf{A}_{r,s}^\pm \cap \mathbf{B}_{r,s}^\pm = \emptyset$ and $\mathbf{A}_{8,0} \cap \mathbf{B}_{8,0} = \emptyset$ and as if both $i \neq j, p \neq q$ we obtain that $[w_i \otimes u_p, w_j \otimes u_q] = 0$ for any $w_i \in \mathbf{A}_{r,s}^\pm, w_j \in \mathbf{B}_{r,s}^\pm, u_p \in \mathbf{A}_{8,0}, u_q \in \mathbf{B}_{8,0}$. \square

In the following lemma we present a list of pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$ satisfying (24), which can be used as a base for the successive extensions.

Lemma 5.2. *The pseudo H -type Lie algebras $\mathfrak{n}_{r,8}, \mathfrak{n}_{8,r}, \mathfrak{n}_{r+4,4}, \mathfrak{n}_{4,r+4}$ for $r \in \{1, 2, 4, 8\} \pmod{8}$ and $\mathfrak{n}_{11}, \mathfrak{n}_{2,2}, \mathfrak{n}_{4,4}$ admit decomposition (24) of their bases.*

Proof. DECOMPOSITIONS FOR $\mathfrak{n}_{r,8}$, $\mathfrak{n}_{8,r}$, $r \in \{1, 2, 4, 8\} \pmod 8$.

$$\begin{aligned} \mathbf{A}_{r,8}^+ &= \{w_i \otimes v_j \mid i = 1, \dots, l, \quad j = 1, \dots, 8\}, \\ \mathbf{A}_{r,8}^- &= \{w_i \otimes v_j \mid i = l+1, \dots, 2l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{r,8}^- &= \{w_i \otimes v_j \mid i = 1, \dots, l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{r,8}^+ &= \{w_i \otimes v_j \mid i = l+1, \dots, 2l, \quad j = 1, \dots, 8\}, \end{aligned}$$

for $w_i \in \mathfrak{B}_{v_{r,0}}$, $v_j \in \mathfrak{B}_{v_{0,8}}$.

$$\begin{aligned} \mathbf{A}_{8,r}^+ &= \{w_i \otimes u_j \mid i = 1, \dots, l, \quad j = 1, \dots, 8\}, \\ \mathbf{A}_{8,r}^- &= \{w_i \otimes u_j \mid i = l+1, \dots, 2l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{8,r}^+ &= \{w_i \otimes u_j \mid i = 1, \dots, l, \quad j = 9, \dots, 16\}, \\ \mathbf{B}_{8,r}^- &= \{w_i \otimes u_j \mid i = l+1, \dots, 2l, \quad j = 1, \dots, 8\}, \end{aligned}$$

for $w_i \in \mathfrak{B}_{v_{0,r}}$, $u_j \in \mathfrak{B}_{v_{8,0}}$. For the proof we use Tables (1)-(4), Tables (10), (11) and the block structure of the corresponding algebras.

DECOMPOSITIONS FOR $\mathfrak{n}_{r+4,4}$, $\mathfrak{n}_{4,r+4}$ FOR $r \in \{1, 2, 4, 8\} \pmod 8$. Recall decompositions (23) and (26) and define

$$\begin{aligned} \mathbf{A}_{r+4,4}^+ &= \mathbf{B}_{r,0} \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{A}_{r,0} \otimes \mathbf{A}_{4,4}^+, & \mathbf{A}_{r+4,4}^- &= \mathbf{B}_{r,0} \otimes \mathbf{B}_{4,4}^- \cup \mathbf{A}_{r,0} \otimes \mathbf{A}_{4,4}^-, \\ \mathbf{B}_{r+4,4}^+ &= \mathbf{B}_{r,0} \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{A}_{r,0} \otimes \mathbf{B}_{4,4}^+, & \mathbf{B}_{r+4,4}^- &= \mathbf{B}_{r,0} \otimes \mathbf{A}_{4,4}^- \cup \mathbf{A}_{r,0} \otimes \mathbf{B}_{4,4}^-, \\ \mathbf{A}_{4,r+4}^+ &= \mathbf{B}_{0,r} \otimes \mathbf{B}_{4,4}^+ \cup \mathbf{A}_{0,r} \otimes \mathbf{A}_{4,4}^+, & \mathbf{A}_{4,r+4}^- &= \mathbf{B}_{0,r} \otimes \mathbf{B}_{4,4}^- \cup \mathbf{A}_{0,r} \otimes \mathbf{A}_{4,4}^-, \\ \mathbf{B}_{4,r+4}^+ &= \mathbf{B}_{0,r} \otimes \mathbf{A}_{4,4}^+ \cup \mathbf{A}_{0,r} \otimes \mathbf{B}_{4,4}^+, & \mathbf{B}_{4,r+4}^- &= \mathbf{B}_{0,r} \otimes \mathbf{A}_{4,4}^- \cup \mathbf{A}_{0,r} \otimes \mathbf{B}_{4,4}^-. \end{aligned}$$

For the proof we use Tables (1)-(4), Tables (10), (11), (12), the block structure of the corresponding algebras, and Lemma 4.8.

DECOMPOSITIONS FOR $\mathfrak{n}_{1,1}$, $\mathfrak{n}_{2,2}$, AND $\mathfrak{n}_{4,4}$.

We define an orthonormal basis of $\mathfrak{n}_{1,1}$ by

$$\mathfrak{B}_{v_{1,1}} = \{w_1 = w, w_2 = J_1 w, w_3 = J_2 w, w_4 = J_2 J_1 w\}, \quad \mathfrak{B}_{z_{1,1}} = \{Z_1, Z_2\}, \quad (27)$$

with $\langle w_i, w_i \rangle_{v_{1,1}} = \epsilon_i(2, 2)$, $\langle Z_k, Z_k \rangle_{z_{1,1}} = \epsilon_k(1, 1)$. The commutators are in Table 5.

TABLE 5. Commutation relations on $\mathfrak{n}_{1,1}$

[row, col.]	w_1	w_4	w_2	w_3
w_1	0	0	Z_1	Z_2
w_4	0	0	$-Z_2$	$-Z_1$
w_2	$-Z_1$	Z_2	0	0
w_3	$-Z_2$	Z_1	0	0

The sets $\mathbf{A}_{1,1}$ and $\mathbf{B}_{1,1}$ are given by

$$\mathbf{A}_{1,1} = \mathbf{A}_{1,1}^+ \cup \mathbf{A}_{1,1}^- = \{w_1\} \cup \{w_4\}, \quad \mathbf{B}_{1,1} = \mathbf{B}_{1,1}^+ \cup \mathbf{B}_{1,1}^- = \{w_2\} \cup \{w_3\}.$$

We define an orthonormal basis of $\mathfrak{B}_{z_{2,2}} = \{Z_1, Z_2, Z_3, Z_4\}$ and

$$\mathfrak{B}_{v_{2,2}} = \left\{ \begin{array}{cccc} w_1 = w, & w_2 = J_1 w, & w_3 = J_2 w, & w_4 = J_1 J_2 w, \\ w_5 = J_3 w, & w_6 = J_4 w, & w_7 = J_1 J_3 w, & w_8 = J_1 J_4 w \end{array} \right\}, \quad (28)$$

for $J_1 J_2 J_3 J_4 w = w$ with $\langle w_i, w_i \rangle_{\mathfrak{v}_{2,2}} = \epsilon_i(4, 4)$, $\langle Z_k, Z_k \rangle_{\mathfrak{z}_{2,2}} = \epsilon_k(2, 2)$. The sets $\mathbf{A}_{2,2}$ and $\mathbf{B}_{2,2}$ are given by

$$\mathbf{A}_{2,2} = \mathbf{A}_{2,2}^+ \cup \mathbf{A}_{2,2}^- = \{w_1, w_4\} \cup \{w_7, w_8\}, \quad \mathbf{B}_{2,2} = \mathbf{B}_{2,2}^+ \cup \mathbf{B}_{2,2}^- = \{w_2, w_3\} \cup \{w_5, w_6\}$$

according to Table 6.

TABLE 6. Commutation relations on $\mathfrak{n}_{2,2}$

[row, col.]	w_1	w_4	w_7	w_8	w_2	w_3	w_5	w_6
w_1	0	0	0	0	Z_1	Z_2	Z_3	Z_4
w_4	0	0	0	0	Z_2	$-Z_1$	Z_4	$-Z_3$
w_7	0	0	0	0	Z_3	$-Z_4$	Z_1	$-Z_2$
w_8	0	0	0	0	Z_4	Z_3	Z_2	Z_1
w_2	$-Z_1$	$-Z_2$	$-Z_3$	$-Z_4$	0	0	0	0
w_3	$-Z_2$	Z_1	Z_4	$-Z_3$	0	0	0	0
w_5	$-Z_3$	$-Z_4$	$-Z_1$	$-Z_2$	0	0	0	0
w_6	$-Z_4$	Z_3	Z_2	$-Z_1$	0	0	0	0

The integral basis $\mathfrak{B}_{\mathfrak{n}_{4,4}}$ of $\mathfrak{n}_{4,4}$ is given in (20) and the decomposition is given in (26) according to Table 12. □

5.2. Inductive construction of isomorphisms of Lie algebras $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$. In this subsection we prove that if two pseudo H -type algebras possess decomposition (24) and they are isomorphic under a map satisfying some special conditions, then the extensions of them are also isomorphic and the corresponding isomorphism map satisfies the same properties. It allows us to perform an induction proof. Before we state the base of induction we formulate the properties we require from the isomorphism.

Remark 5.3. PROPERTIES OF THE ISOMORPHISM $\varphi_{r,s}: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{s,r}$. *Let the integral bases $\mathfrak{B}_{\mathfrak{n}_{r,s}}$, $\mathfrak{B}_{\mathfrak{n}_{s,r}}$ of the pseudo H -type algebras $\mathfrak{n}_{r,s}$, $\mathfrak{n}_{s,r}$ admit decomposition (24). Assume there exists a Lie algebra isomorphism $\varphi_{r,s}: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{s,r}$ such that*

$$\varphi_{r,s}(\mathbf{A}_{r,s}^\pm) = \mathbf{A}_{s,r}^\pm \quad \text{and} \quad \varphi_{r,s}(\mathbf{B}_{r,s}^\pm) = \mathbf{B}_{s,r}^\mp.$$

Furthermore, the restriction $\varphi_{r,s}|_{\mathfrak{z}_{r,s}}$ is an anti-isometry and is a permutation of the set $\{Z_1, \dots, Z_{r+s}\}$, i.e. $\varphi_{r,s}(Z_k) = Z_{\pi_{r,s}(k)}$ with the permutation $\pi_{r,s}: \{1, \dots, r+s\} \rightarrow \{1, \dots, r+s\}$ such that $\pi_{r,s}(\{1, \dots, r\}) = \{s+1, \dots, s+r\}$ and $\pi_{r,s}(\{r+1, \dots, r+s\}) = \{1, \dots, s\}$.

Theorem 5.4. *The Lie algebras $\mathfrak{n}_{r,8}$ and $\mathfrak{n}_{8,r}$ are integral isomorphic if and only if $r \in \{0, 1, 2, 4\} \pmod{8}$ and the Lie algebra isomorphism $\varphi_{r,8}: \mathfrak{n}_{r,8} \rightarrow \mathfrak{n}_{8,r}$ satisfies Remark 5.3 with $s = 8$.*

Proof. The H -type Lie algebras $\mathfrak{n}_{r,0}$ are integral isomorphic to $\mathfrak{n}_{0,r}$ for $r \in \{1, 2, 4, 8\} \pmod{8}$ by Theorem 4.11. Recall that we used the following integral block-type bases

$$\mathfrak{B}_{\mathfrak{n}_{r,0}} = \{w_1, \dots, w_{2l}, Z_1^{r,0}, \dots, Z_r^{r,0}\}, \quad \mathfrak{B}_{\mathfrak{n}_{0,r}} = \{x_1, \dots, x_{2l}, Z_1^{0,r}, \dots, Z_r^{0,r}\},$$

with $\langle w_i, w_i \rangle_{\mathfrak{v}_{r,0}} = 1$ for $i = 1, \dots, 2l$, $\langle x_i, x_i \rangle_{\mathfrak{v}_{0,r}} = \epsilon_i(l, l)$, $\langle Z_k^{r,0}, Z_k^{r,0} \rangle_{\mathfrak{z}_{r,0}} = 1$, and $\langle Z_k^{0,r}, Z_k^{0,r} \rangle_{\mathfrak{z}_{r,0}} = -1$ for all $k = 1, \dots, r$, where $\varphi_{r,0}(w_i) = x_i$ for all $i = 1, \dots, 2l$ and

$\varphi_{r,0}(Z_k^{r,0}) = Z_k^{0,r}$ for all $k = 1, \dots, r$. The equal structure constants are denoted by A_{ij}^m . We write $\varphi_{8,0}(u_i) = v_i$, for $u_i \in \mathfrak{B}_{\mathfrak{v}_{8,0}}$, $v_i \in \mathfrak{B}_{\mathfrak{v}_{0,8}}$, $i = 1, \dots, 16$ and $\varphi_{8,0}(Z_k^{8,0}) = Z_k^{0,8}$, $k = 1, \dots, 8$. The equal structure constants are denoted by \bar{A}_{ij}^m for both Lie algebras.

We exploit Proposition 4.4 and Remark 4.5 and obtain the integral bases

$$\begin{aligned} \{w_1 \otimes v_1, \dots, w_{2l} \otimes v_{16}, Z_1^{r,8}, \dots, Z_{r+8}^{r,8}\} & \quad \text{for } \mathfrak{n}_{r,8}, \\ \{x_1 \otimes u_1, \dots, x_{2l} \otimes u_{16}, Z_1^{8,r}, \dots, Z_{r+8}^{8,r}\} & \quad \text{for } \mathfrak{n}_{8,r}. \end{aligned}$$

Define the bijective linear map $\varphi_{r,8}: \mathfrak{n}_{r,8} \rightarrow \mathfrak{n}_{8,r}$ by

$$\begin{aligned} w_i \otimes v_j & \mapsto x_i \otimes u_j & \text{if } i \in \{1, \dots, l\}, & j \in \{1, \dots, 16\}, \\ w_i \otimes v_j & \mapsto x_i \otimes u_j & \text{if } i \in \{l+1, \dots, 2l\}, & j \in \{1, \dots, 8\}, \\ w_i \otimes v_j & \mapsto -x_i \otimes u_j & \text{if } i \in \{l+1, \dots, 2l\}, & j \in \{9, \dots, 16\}, \\ Z_m^{r,8} & \mapsto Z_{m+8}^{8,r} & \text{if } m \in \{1, \dots, r\}, \\ Z_m^{r,8} & \mapsto Z_{m-r}^{8,r} & \text{if } m \in \{r+1, \dots, r+8\}. \end{aligned}$$

It remains to prove that $\varphi_{r,8}$ is a Lie algebra isomorphism, i.e. $\varphi_{r,8}([w_i \otimes v_j, w_p \otimes v_q]) = [\varphi_{r,8}(w_i \otimes v_j), \varphi_{r,8}(w_p \otimes v_q)]$. The structure constants $\bar{A}_{ij,pq}^m$ of the commutators $[w_i \otimes v_j, w_p \otimes v_q]$ of $\mathfrak{n}_{r,8}$ are given by formula (18), and the structure constants $C_{ij,pq}^m$ for $[x_i \otimes u_j, x_p \otimes u_q]$ of the Lie algebra $\mathfrak{n}_{8,r}$ are given by formula (16). It follows that if $i \neq p$ and $j \neq q$ or $i = p$ and $j = q$ the commutators vanish:

$$\varphi_{r,8}([w_i \otimes v_j, w_p \otimes v_q]) = \varphi_{r,8}(0) = 0 = \pm[x_i \otimes u_j, x_p \otimes u_q].$$

It is left to consider the following two remaining cases.

CASE $i = p$ AND $j \neq q$. If, additionally, both indices simultaneously satisfy either $j, q = 1, \dots, 8$ or $j, q = 9, \dots, 16$, then

$$\varphi_{r,8}([w_i \otimes v_j, w_i \otimes v_q]) = \varphi_{r,8}(0) = 0 = \pm[x_i \otimes u_j, x_i \otimes u_q],$$

because of the block form of the Lie algebras $\mathfrak{n}_{8,0}$, $\mathfrak{n}_{0,8}$. Thus, we can assume without limit of generality that $j = 1, \dots, 8$ and $q = 9, \dots, 16$.

- if $i = p = 1, \dots, l$, $j = 1, \dots, 8$ and $q = 9, \dots, 16$, then:

$$\varphi_{r,8}([w_i \otimes v_j, w_i \otimes v_q]) = \bar{A}_{jq}^{m-r} \varphi_{r,8}(Z_m^{r,8}) = \bar{A}_{jq}^{m-r} Z_{m-r}^{8,r},$$

$$[\varphi_{r,8}(w_i \otimes v_j), \varphi_{r,8}(w_i \otimes v_q)] = [x_i \otimes u_j, x_i \otimes u_q] = \bar{A}_{jq}^{m-r} Z_{m-r}^{8,r}$$

with $m = r+1, \dots, r+8$ by formulas (18) and (16).

- if $i = p = l+1, \dots, 2l$, $j = 1, \dots, 8$ and $q = 9, \dots, 16$, then

$$\varphi_{r,8}([w_i \otimes v_j, w_i \otimes v_q]) = \bar{A}_{jq}^{m-r} \varphi_{r,8}(Z_m^{r,8}) = \bar{A}_{jq}^{m-r} Z_{m-r}^{8,r},$$

$$[\varphi_{r,8}(w_i \otimes v_j), \varphi_{r,8}(w_i \otimes v_q)] = [x_i \otimes u_j, -x_i \otimes u_q] = -(-\bar{A}_{jq}^{m-r} Z_{m-r}^{8,r})$$

with $m = r+1, \dots, r+8$ by formulas (18) and (16). We see that the map $\varphi_{r,8}$ satisfies the Lie algebra isomorphism properties in this case.

CASE $i \neq p$ AND $j = q$. If in addition $i, p = 1, \dots, l$ or $i, p = l+1, \dots, 2l$, then the block form of the Lie algebras $\mathfrak{n}_{r,0}$, $\mathfrak{n}_{0,r}$ implies

$$\varphi_{r,8}([w_i \otimes v_j, w_p \otimes v_j]) = \varphi_{r,8}(0) = 0 = \pm[x_i \otimes u_j, x_p \otimes u_j],$$

such that we can assume that $i = 1, \dots, l$ and $p = l+1, \dots, 2l$.

- if $j = q = 1, \dots, 8$, $i = 1, \dots, l$ and $p = l + 1, \dots, 2l$, then

$$\begin{aligned}\varphi_{r,8}([w_i \otimes v_j, w_p \otimes v_j]) &= -A_{ip}^m \varphi_{r,8}(Z_m^{r,8}) = -A_{ip}^m Z_{m+8}^{8,r}, \\ [\varphi_{r,8}(w_i \otimes v_j), \varphi_{r,8}(w_p \otimes v_j)] &= [x_i \otimes u_j, x_p \otimes u_j] = -A_{ip}^m Z_{m+8}^{8,r}.\end{aligned}$$

- if $j = q = 9, \dots, 16$, $i = 1, \dots, l$ and $p = l + 1, \dots, 2l$ then

$$\begin{aligned}\varphi_{r,8}([w_i \otimes v_j, w_p \otimes v_j]) &= -A_{ip}^m \varphi_{r,8}(Z_m^{r,8}) = -A_{ip}^m Z_{m+8}^{8,r}, \\ [\varphi_{r,8}(w_i \otimes v_j), \varphi_{r,8}(w_p \otimes v_j)] &= [x_i \otimes u_j, -x_p \otimes u_j] = -A_{ip}^m Z_{m+8}^{8,r},\end{aligned}$$

with $m = 1, \dots, r$ by formulas (18) and (16). This shows that $\varphi_{r,8}$ is a Lie algebra isomorphism. The map $\varphi_{r,8}$ satisfies Remark 5.3 by its definition. \square

Theorem 5.5. *Assume that Lie algebras $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$, $r, s \neq 0$, satisfy Remark 5.3. Then there exists a Lie algebra isomorphism $\varphi_{r+8,s}: \mathfrak{n}_{r+8,s} \rightarrow \mathfrak{n}_{s,r+8}$ and two integral bases $\mathfrak{B}_{r+8,s}$ and $\mathfrak{B}_{r,s+8}$ satisfying Remark 5.3.*

Proof. Let $\varphi_{r,s}: \mathfrak{n}_{r,s} \rightarrow \mathfrak{n}_{s,r}$ be the assumed Lie algebra isomorphism. By extension we construct the Lie algebra $\mathfrak{n}_{r+8,s}$ of dimension $32l+r+s+8$ with the basis $\mathfrak{B}_{r+8,s} = \{x_1 \otimes u_1, \dots, x_{2l} \otimes u_{16}, Z_1^{r+8,s}, \dots, Z_{r+s+8}^{r+8,s}\}$. The assumptions imply that $[x_i \otimes u_j, x_p \otimes u_q] = 0$ for the following cases:

- $x_i = x_p$ and both $u_j, u_q \in \mathbf{A}_{8,0}$ or $u_j, u_q \in \mathbf{B}_{8,0}$,
- $u_j = u_q$ and both $x_i, x_p \in \mathbf{A}_{r,s}$ or $x_i, x_p \in \mathbf{B}_{r,s}$,
- $x_i \neq x_p$ and $u_j \neq u_q$ or $x_i = x_p$ and $u_j = u_q$,

by formula (16), where $\mathbf{A}_{8,0}, \mathbf{B}_{8,0}$ are defined in (25). Then we define the bijective linear map $\varphi_{r+8,s}: \mathfrak{n}_{r+8,s} \rightarrow \mathfrak{n}_{s,r+8}$ by

$$\begin{aligned}x_i \otimes u_\alpha &\mapsto -\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_\alpha) && \text{if } x_i \in \mathbf{B}_{r,s}, \quad \text{and } u_\alpha \in \mathbf{B}_{8,0}, \\ x_i \otimes u_\alpha &\mapsto \varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_\alpha) && \text{if otherwise,} \\ Z_m^{r+8,s} &\mapsto Z_{\pi_{r,s}(m)}^{s,r+8} && \text{if } m \in \{1, \dots, r\}, \\ Z_m^{r+8,s} &\mapsto Z_{\pi_{8,0}(m-r)+r+s}^{s,r+8} && \text{if } m \in \{r+1, \dots, r+8\}, \\ Z_m^{r+8,s} &\mapsto Z_{\pi_{r,s}(m-8)}^{s,r+8} && \text{if } m \in \{r+9, \dots, r+s+8\},\end{aligned}$$

where $\varphi_{8,0}: \mathfrak{n}_{8,0} \rightarrow \mathfrak{n}_{0,8}$ is the Lie algebra isomorphism given by (14). We see that the restriction of $\varphi_{r+8,s}$ to $\mathfrak{z}_{r+8,s}$ is an anti-isometry, such that it remains to prove that $\varphi_{r+8,s}$ is a Lie algebra homomorphism.

Before we continue, we draw the attention of the reader to the following. By Lemma 4.6 we know that $[x_i \otimes u_j, x_i \otimes u_q] = \pm[u_j, u_q]_{r+8,s} \in \text{span}\{Z_k^{r+8,s} | k = r+1, \dots, r+8\}$. Since the index k belongs to the set $\{r+1, \dots, r+8\}$, the structure constants $[u_j, u_q]$ in $\mathfrak{n}_{r+8,s}$ coincide with the structure constants $[u_j, u_q]$ in $\mathfrak{n}_{8,0}$. Analogously we write $[x_i \otimes u_j, x_p \otimes u_j] = \pm[x_i, x_p]_{r+8,s} \in \text{span}\{Z_k | k = 1, \dots, r, r+9, \dots, r+s+8\}$ and observe that $[x_i, x_p]_{r+8,s} = [x_i, x_p]_{r,s}$. Thus

$$\varphi_{8,0}([u_j, u_q]_{r+8,s}) = [\varphi_{8,0}(u_j), \varphi_{8,0}(u_q)]_{r+8,s}, \quad \varphi_{r,s}([x_i, x_p]_{r+8,s}) = [\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+8,s}$$

as $\varphi_{8,0}$ and $\varphi_{r,s}$ are Lie algebra isomorphisms. Now, we consider the following cases by using formulas (16) and (19).

- If $x_i = x_p \in \mathbf{B}_{r,s}^+$, $u_j \in \mathbf{A}_{8,0}$ and $u_q \in \mathbf{B}_{8,0}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_i \otimes u_q]) = \varphi_{r+8,s}([u_j, u_q]_{r+8,s}) = \varphi_{8,0}([u_j, u_q]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_i \otimes u_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), -\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_q)] \\ &= [\varphi_{8,0}(u_j), \varphi_{8,0}(u_q)]_{r+8,s} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{B}_{s,r}^-$.

- If $x_i = x_p \in \mathbf{B}_{r,s}^-$, $u_j \in \mathbf{A}_{8,0}$ and $u_q \in \mathbf{B}_{8,0}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_i \otimes u_q]) = \varphi_{r+8,s}(-[u_j, u_q]_{r+8,s}) = -\varphi_{8,0}([u_j, u_q]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_i \otimes u_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), -\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_q)] \\ &= -[\varphi_{8,0}(u_j), \varphi_{8,0}(u_q)]_{r+8,s} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{B}_{s,r}^+$.

- If $x_i = x_p \in \mathbf{A}_{r,s}^+$, $u_j \in \mathbf{A}_{8,0}$ and $u_q \in \mathbf{B}_{8,0}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_i \otimes u_q]) = \varphi_{r+8,s}([u_j, u_q]_{r+8,s}) = \varphi_{8,0}([u_j, u_q]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_i \otimes u_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), \varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_q)] \\ &= [\varphi_{8,0}(u_j), \varphi_{8,0}(u_q)]_{r+8,s} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{A}_{s,r}^+$.

- If $x_i = x_p \in \mathbf{A}_{r,s}^-$, $u_j \in \mathbf{A}_{8,0}$ and $u_q \in \mathbf{B}_{8,0}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_i \otimes u_q]) = \varphi_{r+8,s}(-[u_j, u_q]_{r+8,s}) = -\varphi_{8,0}([u_j, u_q]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_i \otimes u_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), \varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_q)] \\ &= -[\varphi_{8,0}(u_j), \varphi_{8,0}(u_q)]_{r+8,s} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{A}_{s,r}^-$.

- If $u_j = u_q \in \mathbf{B}_{8,0}$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_p \otimes u_j]) = \varphi_{r+8,s}([x_i, x_p]_{r+8,s}) = \varphi_{r,s}([x_i, x_p]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_p \otimes u_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), -\varphi_{r,s}(x_p) \otimes \varphi_{8,0}(u_j)] \\ &= [\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+8,s} \end{aligned}$$

as $\varphi_{8,0}(u_j) \in \pm \mathbf{B}_{0,8}$.

- If $u_j = u_q \in \mathbf{A}_{8,0}$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\varphi_{r+8,s}([x_i \otimes u_j, x_p \otimes u_j]) = \varphi_{r+8,s}(-[x_i, x_p]_{r+8,s}) = -\varphi_{r,s}([x_i, x_p]_{r+8,s}),$$

$$\begin{aligned} [\varphi_{r+8,s}(x_i \otimes u_j), \varphi_{r+8,s}(x_p \otimes u_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{8,0}(u_j), \varphi_{r,s}(x_p) \otimes \varphi_{8,0}(u_j)] \\ &= -[\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+8,s} \end{aligned}$$

as $\varphi_{8,0}(u_j) \in \pm \mathbf{A}_{0,8}$.

Hence $\varphi_{r+8,s}$ is a Lie algebra isomorphism satisfying Remark 5.3. \square

Now we turn to consider the extension obtained by making use of the tensor product with $\mathfrak{v}_{4,4}$.

Theorem 5.6. *For any $\mathfrak{n}_{r,r}$ with $r = 1, 2, 4$ there exists an automorphism $\varphi_{r,r}: \mathfrak{n}_{r,r} \rightarrow \mathfrak{n}_{r,r}$ and an integral basis $\mathfrak{B}_{r,r}$ satisfying Remark 5.3.*

Proof. In this proof we explicitly state the automorphisms.

The basis of $\mathfrak{n}_{1,1}$ is given in (27) and the commutations in Table 5. The automorphism $\varphi_{1,1}: \mathfrak{n}_{1,1} \rightarrow \mathfrak{n}_{1,1}$ with anti-isometry on the center is given by

$$\begin{aligned} Z_1^{1,1} &\mapsto Z_2^{1,1}, & Z_2^{1,1} &\mapsto Z_1^{1,1}, \\ w_1 &\mapsto w_1, & w_2 &\mapsto w_3, & w_3 &\mapsto w_2, & w_4 &\mapsto w_4. \end{aligned} \quad (29)$$

We defined an orthonormal basis of $\mathfrak{n}_{2,2}$ in (28) with commutators in Table 6. The automorphism $\varphi_{2,2}: \mathfrak{n}_{2,2} \rightarrow \mathfrak{n}_{2,2}$ with anti-isometry on the center is given by

$$\begin{aligned} Z_1^{2,2} &\mapsto Z_3^{2,2}, & Z_2^{2,2} &\mapsto Z_4^{2,2}, & Z_3^{2,2} &\mapsto Z_1^{2,2}, & Z_4^{2,2} &\mapsto Z_2^{2,2}, \\ w_1 &\mapsto w_1, & w_2 &\mapsto w_5, & w_3 &\mapsto w_6, & w_4 &\mapsto w_4, \\ w_5 &\mapsto w_2, & w_6 &\mapsto w_3, & w_7 &\mapsto w_7, & w_8 &\mapsto w_8. \end{aligned} \quad (30)$$

Recalling the basis (20) and Table 12 we define the automorphism $\varphi_{4,4}: \mathfrak{n}_{4,4} \rightarrow \mathfrak{n}_{4,4}$ with anti-isometry on the center by

$$\begin{aligned} Z_1^{4,4} &\mapsto Z_5^{4,4}, & Z_2^{4,4} &\mapsto Z_6^{4,4}, & Z_3^{4,4} &\mapsto Z_8^{4,4}, & Z_4^{4,4} &\mapsto Z_7^{4,4}, \\ Z_5^{4,4} &\mapsto Z_1^{4,4}, & Z_6^{4,4} &\mapsto Z_2^{4,4}, & Z_7^{4,4} &\mapsto Z_4^{4,4}, & Z_8^{4,4} &\mapsto Z_3^{4,4}, \\ y_1 &\mapsto y_1, & y_2 &\mapsto y_9, & y_3 &\mapsto y_{10}, & y_4 &\mapsto y_{12}, \\ y_5 &\mapsto y_{11}, & y_6 &\mapsto y_6, & y_7 &\mapsto y_7, & y_8 &\mapsto -y_8, \\ y_9 &\mapsto y_2, & y_{10} &\mapsto y_3, & y_{11} &\mapsto y_5, & y_{12} &\mapsto y_4, \\ y_{13} &\mapsto y_{13}, & y_{14} &\mapsto y_{14}, & y_{15} &\mapsto -y_{15}, & y_{16} &\mapsto y_{16}. \end{aligned} \quad (31)$$

□

Theorem 5.7. *The Lie algebras $\mathfrak{n}_{r+4,4}$ and $\mathfrak{n}_{4,r+4}$ are integral isomorphic if and only if $r \in \{0, 1, 2, 4\} \pmod{8}$. In the case $r \in \{0, 1, 2, 4\} \pmod{8}$ there exists a Lie algebra isomorphism $\varphi_{r+4,4}: \mathfrak{n}_{r+4,4} \rightarrow \mathfrak{n}_{4,r+4}$ and two integral bases $\mathfrak{B}_{r+4,4}$ and $\mathfrak{B}_{4,r+4}$ satisfying Remark 5.3.*

Proof. By extension we construct the Lie algebra $\mathfrak{n}_{r+4,4}$ of dimension $32l+r+8$ with the integral basis $\{x_1 \otimes y_1, \dots, x_{2l} \otimes y_{16}, Z_1^{r+4,4}, \dots, Z_{r+8}^{r+4,4}\}$, where $\{x_1, \dots, x_{2l}\} = \mathfrak{B}_{\mathfrak{v}_{r,0}}$. Lemma 4.8 implies that $[x_i \otimes y_j, x_p \otimes y_q] = 0$ for the following cases:

- $x_i = x_p$ and both $y_j, y_q \in \mathbf{A}_{4,4}$ or $y_j, y_q \in \mathbf{B}_{4,4}$,
- $y_j = y_q$ and both $x_i, x_p \in \mathbf{A}_{r,0}$ or $x_i, x_p \in \mathbf{B}_{r,0}$,
- $x_i \neq x_p$ and $y_j \neq y_q$ or $x_i = x_p$ and $y_j = y_q$,

where $\mathbf{A}_{r,0}, \mathbf{B}_{r,0}$ are defined in (23) and $\mathbf{A}_{4,4}, \mathbf{B}_{4,4}$ are defined in (26). We define the bijective linear map $\varphi_{r+4,4}: \mathfrak{n}_{r+4,4} \rightarrow \mathfrak{n}_{4,r+4}$ by

$$\begin{aligned} x_i \otimes y_\alpha &\mapsto -\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_\alpha) && \text{if } x_i \in \mathbf{B}_{r,0}, \text{ and } y_\alpha \in \mathbf{B}_{4,4}, \\ x_i \otimes y_\alpha &\mapsto \varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_\alpha) && \text{if otherwise,} \\ Z_m^{r+4,4} &\mapsto Z_{\pi_{r,0}(m)+8}^{4,r+4} && \text{if } m \in \{1, \dots, r\}, \\ Z_m^{r+4,4} &\mapsto Z_{\pi_{4,4}(m-r)}^{4,r+4} && \text{if } m \in \{r+1, \dots, r+8\}. \end{aligned}$$

We see that the restriction of $\varphi_{r+4,4}$ to $\mathfrak{z}_{r+4,4}$ is an anti-isometry, such that it remains to prove that $\varphi_{r+4,4}$ is a Lie algebra homomorphism.

As in Theorem 5.5 we make the following observation. By Lemma 4.8 we know that $[x_i \otimes y_j, x_i \otimes y_q] = \pm[y_j, y_q]_{r+4,4} \in \text{span}\{Z_k | k = r+1, \dots, r+8\}$ and therefore $[y_j, y_q]_{r+4,4} = [y_j, y_q]_{4,4}$. Analogously, because of $[x_i \otimes y_j, x_p \otimes y_j] = \pm[x_i, x_p]_{r+4,4} \in \text{span}\{Z_k | k = 1, \dots, r\}$ we obtain $[x_i, x_p]_{r+4,4} = [x_i, x_p]_{r,0}$. Thus $\varphi_{4,4}([y_j, y_q]_{r+4,4}) = [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,4}$ and $\varphi_{r,0}([x_i, x_p]_{r+4,4}) = [\varphi_{r,0}(x_i), \varphi_{r,0}(x_p)]_{r+4,4}$, respectively, as

$\varphi_{4,4}$ and $\varphi_{r,0}$ are Lie algebra isomorphisms. We turn to consider several cases, where we use formulas (21) and (22).

- If $x_i = x_p \in \mathbf{B}_{r,0}$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,4}([y_j, y_q]_{r+4,4}) = \varphi_{4,4}([y_j, y_q]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_i \otimes y_q)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,4} \end{aligned}$$

as $\varphi_{r,0}(x_i) \in \pm \mathbf{B}_{0,r}$.

- If $x_i = x_p \in \mathbf{A}_{r,0}$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,4}([y_j, y_q]_{r+4,s+4}) = \varphi_{4,4}([y_j, y_q]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_i \otimes y_q)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,4} \end{aligned}$$

as $\varphi_{r,0}(x_i) \in \pm \mathbf{A}_{0,r}$.

- If $y_j = y_q \in \mathbf{B}_{4,4}^+$, $x_i \in \mathbf{A}_{r,0}$ and $x_p \in \mathbf{B}_{r,0}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,4}([x_i, x_p]_{r+4,4}) = \varphi_{r,0}([x_i, x_p]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_p \otimes y_j)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,0}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= [\varphi_{r,0}(x_i), \varphi_{r,0}(x_p)]_{r+4,4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{B}_{4,4}^-$.

- If $y_j = y_q \in \mathbf{B}_{4,4}^-$, $x_i \in \mathbf{A}_{r,0}$ and $x_p \in \mathbf{B}_{r,0}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,4}(-[x_i, x_p]_{r+4,4}) = -\varphi_{r,0}([x_i, x_p]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_p \otimes y_j)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,0}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= -[\varphi_{r,0}(x_i), \varphi_{r,0}(x_p)]_{r+4,4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{B}_{4,4}^+$.

- If $y_j = y_q \in \mathbf{A}_{4,4}^+$, $x_i \in \mathbf{A}_{r,0}$ and $x_p \in \mathbf{B}_{r,0}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,4}(-[x_i, x_p]_{r+4,4}) = -\varphi_{r,0}([x_i, x_p]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_p \otimes y_j)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,0}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= -[\varphi_{r,0}(x_i), \varphi_{r,0}(x_p)]_{r+4,4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{A}_{4,4}^+$.

- If $y_j = y_q \in \mathbf{A}_{4,4}^-$, $x_i \in \mathbf{A}_{r,0}$ and $x_p \in \mathbf{B}_{r,0}$, then:

$$\begin{aligned} \varphi_{r+4,4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,4}([x_i, x_p]_{r+4,4}) = \varphi_{r,0}([x_i, x_p]_{r+4,4}), \\ [\varphi_{r+4,4}(x_i \otimes y_j), \varphi_{r+4,4}(x_p \otimes y_j)] &= [\varphi_{r,0}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,0}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= [\varphi_{r,0}(x_i), \varphi_{r,0}(x_p)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{A}_{4,4}^-$. Hence $\varphi_{r+4,4}$ is a Lie algebra isomorphism which is an anti-isometry on its center $\mathfrak{z}_{r+4,4}$ satisfying Remark 5.3. \square

The generalization of these results is based on the technical Lemma 4.8.

Theorem 5.8. *Assume that the pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$ and $\mathfrak{n}_{s,r}$, $r, s \neq 0$ satisfy Remark 5.3. Then there exists a Lie algebra isomorphism $\varphi_{r+4,s+4}: \mathfrak{n}_{r+4,s+4} \rightarrow \mathfrak{n}_{s+4,r+4}$ also satisfying Remark 5.3.*

Proof. By extension we construct the Lie algebra $\mathfrak{n}_{r+4,s+4}$ of dimension $32l+r+s+8$ with the basis $\{x_1 \otimes y_1, \dots, x_{2l} \otimes y_{16}, Z_1^{r+4,s+4}, \dots, Z_{r+s+8}^{r+4,s+4}\}$, where $\{x_1, \dots, x_{2l}\} = \mathfrak{B}_{\mathfrak{v}_{r,s}}$. The assumptions imply that $[x_i \otimes y_j, x_p \otimes y_q] = 0$ for the following cases:

- $x_i = x_p$ and both $y_j, y_q \in \mathbf{A}_{4,4}$ or $y_j, y_q \in \mathbf{B}_{4,4}$,
- $y_j = y_q$ and both $x_i, x_p \in \mathbf{A}_{r,s}$ or $x_i, x_p \in \mathbf{B}_{r,s}$,
- $x_i \neq x_p$ and $y_j \neq y_q$ or $x_i = x_p$ and $y_j = y_q$,

by formula (22). We define the bijective linear map $\varphi_{r+4,s+4}: \mathfrak{n}_{r+4,s+4} \rightarrow \mathfrak{n}_{s+4,r+4}$ by

$$\begin{aligned} x_i \otimes y_\alpha &\mapsto -\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_\alpha) && \text{if } x_i \in \mathbf{B}_{r,s}, \text{ and } y_\alpha \in \mathbf{B}_{4,4}, \\ x_i \otimes y_\alpha &\mapsto \varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_\alpha) && \text{if otherwise,} \\ Z_m^{r+4,s+4} &\mapsto Z_{\pi_{r,s}(m)+8}^{r+4,s+4} && \text{if } m \in \{1, \dots, r\}, \\ Z_m^{r+4,s+4} &\mapsto Z_{\pi_{4,4}(m-r)+s}^{r+4,s+4} && \text{if } m \in \{r+1, \dots, r+8\}, \\ Z_m^{r+4,s+4} &\mapsto Z_{\pi_{r,s}(m-8)}^{r+4,s+4} && \text{if } m \in \{r+9, \dots, r+s+8\}. \end{aligned}$$

We see that the restriction of $\varphi_{r+4,s+4}$ to $\mathfrak{z}_{r+4,s+4}$ is an anti-isometry. Let us show that $\varphi_{r+4,s+4}$ is a lie algebra homomorphism.

Observe that Lemma 4.8 implies that $[x_i \otimes y_j, x_i \otimes y_q] = \pm[y_j, y_q]_{r+4,s+4} \in \text{span}\{Z_k | k = r+1, \dots, r+8\}$ and $[x_i \otimes y_j, x_p \otimes y_j] = \pm[x_i, x_p]_{r+4,s+4} \in \text{span}\{Z_k | k = 1, \dots, r, r+9, \dots, r+s+8\}$. Thus $[y_j, y_q]_{r+4,s+4} = [y_j, y_q]_{4,4}$ and $[x_i, x_p]_{r+4,s+4} = [x_i, x_p]_{r,s}$. Therefore,

$$\begin{aligned} \varphi_{4,4}([y_j, y_q]_{r+4,s+4}) &= [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,s+4}, \\ \varphi_{r,s}([x_i, x_p]_{r+4,s+4}) &= [\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+4,s+4}, \end{aligned}$$

respectively, as $\varphi_{4,4}$ and $\varphi_{r,s}$ are Lie algebra isomorphisms. The remaining cases follow from formula (22).

- If $x_i = x_p \in \mathbf{B}_{r,s}^+$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,s+4}([y_j, y_q]_{r+4,s+4}) = \varphi_{4,4}([y_j, y_q]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_i \otimes y_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{B}_{s,r}^-$.

- If $x_i = x_p \in \mathbf{B}_{r,s}^-$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,s+4}(-[y_j, y_q]_{r+4,s+4}) = -\varphi_{4,4}([y_j, y_q]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_i \otimes y_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= -[\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{B}_{s,r}^+$.

- If $x_i = x_p \in \mathbf{A}_{r,s}^+$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,s+4}([y_j, y_q]_{r+4,s+4}) = \varphi_{4,4}([y_j, y_q]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_i \otimes y_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= [\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{A}_{s,r}^+$.

- If $x_i = x_p \in \mathbf{A}_{r,s}^-$, $y_j \in \mathbf{A}_{4,4}$ and $y_q \in \mathbf{B}_{4,4}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_i \otimes y_q]) &= \varphi_{r+4,s+4}(-[y_j, y_q]_{r+4,s+4}) = -\varphi_{4,4}([y_j, y_q]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_i \otimes y_q)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_q)] \\ &= -[\varphi_{4,4}(y_j), \varphi_{4,4}(y_q)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{r,s}(x_i) \in \pm \mathbf{A}_{s,r}^-$.

- If $y_j = y_q \in \mathbf{B}_{4,4}^+$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,s+4}([x_i, x_p]_{r+4,s+4}) = \varphi_{r,s}([x_i, x_p]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_p \otimes y_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,s}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= [\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{B}_{4,4}^-$.

- If $y_j = y_q \in \mathbf{B}_{4,4}^-$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,s+4}(-[x_i, x_p]_{r+4,s+4}) = -\varphi_{r,s}([x_i, x_p]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_p \otimes y_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), -\varphi_{r,s}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= -[\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{B}_{4,4}^+$.

- If $y_j = y_q \in \mathbf{A}_{4,4}^+$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,s+4}([x_i, x_p]_{r+4,s+4}) = \varphi_{r,s}([x_i, x_p]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_p \otimes y_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,s}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= [\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{A}_{4,4}^+$.

- If $y_j = y_q \in \mathbf{A}_{4,4}^-$, $x_i \in \mathbf{A}_{r,s}$ and $x_p \in \mathbf{B}_{r,s}$, then:

$$\begin{aligned} \varphi_{r+4,s+4}([x_i \otimes y_j, x_p \otimes y_j]) &= \varphi_{r+4,s+4}(-[x_i, x_p]_{r+4,s+4}) = -\varphi_{r,s}([x_i, x_p]_{r+4,s+4}), \\ [\varphi_{r+4,s+4}(x_i \otimes y_j), \varphi_{r+4,s+4}(x_p \otimes y_j)] &= [\varphi_{r,s}(x_i) \otimes \varphi_{4,4}(y_j), \varphi_{r,s}(x_p) \otimes \varphi_{4,4}(y_j)] \\ &= -[\varphi_{r,s}(x_i), \varphi_{r,s}(x_p)]_{r+4,s+4} \end{aligned}$$

as $\varphi_{4,4}(y_j) \in \pm \mathbf{A}_{4,4}^-$.

Hence $\varphi_{r+4,s+4}$ is a Lie algebra isomorphism which satisfies Remark 5.3. \square

5.3. Main results of Section 5.

Theorem 5.9. *The H -type Lie algebras $\mathfrak{n}_{r+4t_1+8t_2,8t_3+4t_1}$ and $\mathfrak{n}_{8t_3+4t_1,r+4t_1+8t_2}$ are integral isomorphic for $r \in \{0, 1, 2, 4\} \pmod{8}$ and $t_1, t_2, t_3 \in \mathbb{N} \cup \{0\}$.*

Proof. We prove by induction. The beginning of the induction is stated in Theorem 5.4 and Theorem 5.7. Then the induction step is given by Theorem 5.5 and Theorem 5.8. \square

Theorem 5.10. *The H -type Lie algebras $\mathfrak{n}_{r+8t_1+4t_2,r+8t_3+4t_2}$ and $\mathfrak{n}_{r+8t_3+4t_2,r+8t_1+4t_2}$ are integral isomorphic for $r \in \{0, 1, 2\} \pmod{4}$ and $t_1, t_2, t_3 \in \mathbb{N} \cup \{0\}$.*

Proof. We prove by induction. The beginning of the induction is stated in Theorem 5.6. Then the induction step is given by Theorem 5.5 and Theorem 5.8. \square

6. SOME NON-ISOMORPHIC LIE ALGEBRAS $\mathfrak{n}_{r,s}$ AND $\mathfrak{n}_{s,r}$

The behavior of the Lie algebras of block type, or those that admit the decomposition (24) are very special and as we saw in the previous section it is preserved under extension. The situation is much less predictable if the Lie algebra is not of block type and different situations can occur. In the present section we show one example of non isomorphic algebras: $\mathfrak{n}_{2,3}$ and $\mathfrak{n}_{3,2}$. We show also that, in a contrary to algebras $\mathfrak{n}_{r,r}$, $r \in \{1, 2, 4\} \pmod{4}$ the pseudo H -type Lie algebra $\mathfrak{n}_{3,3}$ does not admit an automorphism such that the restriction to the center is an anti-isometry. We also observe that our method does not allow to show that, for instance, $\mathfrak{n}_{7,7}$, with $\mathfrak{v}_{7,7} = \mathfrak{v}_{3,3} \otimes \mathfrak{v}_{4,4}$, admits or does not admit an automorphism such that the restriction to the center is an anti-isometry.

6.1. Non-isomorphism of $\mathfrak{n}_{3,2}$ and $\mathfrak{n}_{2,3}$. First we introduce the integral basis of $\mathfrak{n}_{3,2}$ and $\mathfrak{n}_{2,3}$ which is essential for the proof of Theorem 6.4.

We define an orthonormal basis of $\mathfrak{n}_{3,2}$ by $\mathfrak{B}_{\mathfrak{z}_{3,2}} = \{Z_0, Z_1, Z_2, Z_3, Z_4\}$ and

$$\mathfrak{B}_{\mathfrak{v}_{3,2}} = \left\{ \begin{array}{llll} w_1 = w, & w_2 = J_1 w, & w_3 = J_2 w, & w_4 = J_1 J_2 w, \\ w_5 = J_3 w, & w_6 = J_4 w, & w_7 = J_1 J_3 w, & w_8 = J_1 J_4 w, \end{array} \right\}$$

for $J_1 J_2 J_3 J_4 w = J_0 J_1 J_2 w = w$ with $\langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} = \epsilon_i(4, 4)$, $\langle Z_k, Z_k \rangle_{\mathfrak{z}_{3,2}} = \epsilon_{k+1}(3, 2)$.

TABLE 7. Commutation relations on $\mathfrak{n}_{3,2}$

[row, col.]	w_1	w_4	w_7	w_8	w_2	w_3	w_5	w_6
w_1	0	$-Z_0$	0	0	Z_1	Z_2	Z_3	Z_4
w_4	Z_0	0	0	0	Z_2	$-Z_1$	Z_4	$-Z_3$
w_7	0	0	0	$-Z_0$	Z_3	$-Z_4$	Z_1	$-Z_2$
w_8	0	0	Z_0	0	Z_4	Z_3	Z_2	Z_1
w_2	$-Z_1$	$-Z_2$	$-Z_3$	$-Z_4$	0	$-Z_0$	0	0
w_3	$-Z_2$	Z_1	Z_4	$-Z_3$	Z_0	0	0	0
w_5	$-Z_3$	$-Z_4$	$-Z_1$	$-Z_2$	0	0	0	Z_0
w_6	$-Z_4$	Z_3	Z_2	$-Z_1$	0	0	$-Z_0$	0

We define an orthonormal basis of $\mathfrak{n}_{2,3}$ by $\mathfrak{B}_{\mathfrak{z}_{2,3}} = \{\bar{Z}_1, \bar{Z}_2, \bar{Z}_3, \bar{Z}_4, \bar{Z}_5\}$ and

$$\mathfrak{B}_{\mathfrak{v}_{2,3}} = \left\{ \begin{array}{llll} \bar{w}_1 = \bar{w}, & \bar{w}_2 = J_1 \bar{w}, & \bar{w}_3 = J_2 \bar{w}, & \bar{w}_4 = J_1 J_2 \bar{w}, \\ \bar{w}_5 = J_3 \bar{w}, & \bar{w}_6 = J_4 \bar{w}, & \bar{w}_7 = J_1 J_3 \bar{w}, & \bar{w}_8 = J_1 J_4 \bar{w}, \end{array} \right\}$$

for $J_1 J_2 J_3 J_4 \bar{w} = J_1 J_4 J_5 \bar{w} = \bar{w}$ with $\langle \bar{w}_i, \bar{w}_i \rangle_{\mathfrak{v}_{2,3}} = \epsilon_i(4, 4)$, $\langle \bar{Z}_k, \bar{Z}_k \rangle_{\mathfrak{z}_{2,3}} = \epsilon_k(2, 3)$.

Proposition 6.1. *The following is true.*

- The linear map $\text{ad}_X: \mathfrak{v}_{3,2} \rightarrow \mathfrak{z}_{3,2}$ is surjective if and only if $\langle X, X \rangle_{\mathfrak{v}_{3,2}} \neq 0$.
- The linear map $\text{ad}_X: \mathfrak{v}_{2,3} \rightarrow \mathfrak{z}_{2,3}$ is surjective if and only if $\langle X, X \rangle_{\mathfrak{v}_{2,3}} \neq 0$.

TABLE 8. Commutation relations on $\mathfrak{n}_{2,3}$

[row, col.]	\bar{w}_1	\bar{w}_4	\bar{w}_7	\bar{w}_8	\bar{w}_2	\bar{w}_3	\bar{w}_5	\bar{w}_6
\bar{w}_1	0	0	0	Z_5	Z_1	Z_2	Z_3	Z_4
\bar{w}_4	0	0	$-Z_5$	0	Z_2	$-Z_1$	Z_4	$-Z_3$
\bar{w}_7	0	Z_5	0	0	Z_3	$-Z_4$	Z_1	$-Z_2$
\bar{w}_8	$-Z_5$	0	0	0	Z_4	Z_3	Z_2	Z_1
\bar{w}_2	$-Z_1$	$-Z_2$	$-Z_3$	$-Z_4$	0	0	0	Z_5
\bar{w}_3	$-Z_2$	Z_1	Z_4	$-Z_3$	0	0	Z_5	0
\bar{w}_5	$-Z_3$	$-Z_4$	$-Z_1$	$-Z_2$	0	$-Z_5$	0	0
\bar{w}_6	$-Z_4$	Z_3	Z_2	$-Z_1$	$-Z_5$	0	0	0

Proof. First we note that ad_X is surjective for all $X \in \mathfrak{v}_{r,s}$ with $\langle X, X \rangle_{\mathfrak{v}_{r,s}} \neq 0$ by Definition 7.2, such that it suffices to prove that for $\langle X, X \rangle_{\mathfrak{v}_{3,2}} = 0$, $\langle X, X \rangle_{\mathfrak{v}_{2,3}} = 0$, respectively, the map ad_X is not surjective.

We write $X = \sum_{i=1}^8 \lambda_i w_i$ for $X \in \mathfrak{v}_{3,2}$ and define the representation matrix M_X of ad_X with respect to the orthonormal basis $\mathfrak{B}_{\mathfrak{n}_{3,2}}$ by

$$M_X = [V_1^X \quad V_4^X \quad V_7^X \quad V_8^X \quad V_2^X \quad V_3^X \quad V_5^X \quad V_6^X],$$

where V_i^X is the vector representation $\begin{pmatrix} \mu_{0i}^X \\ \vdots \\ \mu_{4i}^X \end{pmatrix}$ of $[X, w_i] = \sum_{k=0}^4 \mu_{ki}^X Z_k$. The matrix

M_X for $\mathfrak{n}_{3,2}$ is given by

$$\begin{pmatrix} \lambda_4 & -\lambda_1 & \lambda_8 & -\lambda_7 & \lambda_3 & -\lambda_2 & -\lambda_6 & \lambda_5 \\ -\lambda_2 & \lambda_3 & -\lambda_5 & -\lambda_6 & \lambda_1 & -\lambda_4 & \lambda_7 & \lambda_8 \\ -\lambda_3 & -\lambda_2 & \lambda_6 & -\lambda_5 & \lambda_4 & \lambda_1 & \lambda_8 & -\lambda_7 \\ -\lambda_5 & \lambda_6 & -\lambda_2 & -\lambda_3 & \lambda_7 & \lambda_8 & \lambda_1 & -\lambda_4 \\ -\lambda_6 & -\lambda_5 & \lambda_3 & -\lambda_2 & \lambda_8 & -\lambda_7 & \lambda_4 & \lambda_1 \end{pmatrix}.$$

Note that M_X is surjective if and only if $\det(M_X M_X^T) \neq 0$ as $\text{rank}(M_X M_X^T) = \text{rank}(M_X)$. The determinant of $M_X M_X^T$ is

$$\begin{aligned} & (\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 - \lambda_7^2 - \lambda_8^2)^2 (\lambda_1^2 + \lambda_4^2 + \lambda_7^2 + \lambda_8^2 + \lambda_2^2 + \lambda_3^2 + \lambda_5^2 + \lambda_6^2) \\ & \times \left[\lambda_1^4 + \lambda_4^4 + \lambda_7^4 + 2\lambda_7^2 \lambda_8^2 + \lambda_8^4 + 2\lambda_7^2 \lambda_2^2 + 2\lambda_8^2 \lambda_2^2 + \lambda_2^4 + 2\lambda_7^2 \lambda_3^2 + 2\lambda_8^2 \lambda_3^2 + 2\lambda_2^2 \lambda_3^2 \right. \\ & + \lambda_3^4 + 2\lambda_7^2 \lambda_5^2 + 2\lambda_8^2 \lambda_5^2 - 2\lambda_2^2 \lambda_5^2 - 2\lambda_3^2 \lambda_5^2 + \lambda_5^4 + 2(\lambda_7^2 + \lambda_8^2 - \lambda_2^2 - \lambda_3^2 + \lambda_5^2) \lambda_6^2 \\ & + \lambda_6^4 - 8\lambda_1(\lambda_7 \lambda_2 \lambda_5 + \lambda_8 \lambda_3 \lambda_5 + \lambda_8 \lambda_2 \lambda_6 - \lambda_7 \lambda_3 \lambda_6) + 8\lambda_4(-\lambda_8 \lambda_2 \lambda_5 + \lambda_7 \lambda_3 \lambda_5 \\ & + \lambda_7 \lambda_2 \lambda_6 + \lambda_8 \lambda_3 \lambda_6) + 2\lambda_4^2(-\lambda_7^2 - \lambda_8^2 + \lambda_2^2 + \lambda_3^2 + \lambda_5^2 + \lambda_6^2) \\ & \left. + 2\lambda_1^2(\lambda_4^2 - \lambda_7^2 - \lambda_8^2 + \lambda_2^2 + \lambda_3^2 + \lambda_5^2 + \lambda_6^2) \right]. \end{aligned}$$

It follows that for all $X \in \mathfrak{v}_{3,2}$ with $\langle X, X \rangle_{\mathfrak{v}_{3,2}} = \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 - \lambda_7^2 - \lambda_8^2 = 0$ the determinant $\det(M_X M_X^T)$ vanishes. This finishes the proof for $\mathfrak{n}_{3,2}$.

For $\mathfrak{n}_{2,3}$ the matrix M_X is given by

$$\begin{pmatrix} -\lambda_2 & \lambda_3 & -\lambda_5 & -\lambda_6 & \lambda_1 & -\lambda_4 & \lambda_7 & \lambda_8 \\ -\lambda_3 & -\lambda_2 & \lambda_6 & -\lambda_5 & \lambda_4 & \lambda_1 & \lambda_8 & -\lambda_7 \\ -\lambda_5 & \lambda_6 & -\lambda_2 & -\lambda_3 & \lambda_7 & \lambda_8 & \lambda_1 & -\lambda_4 \\ -\lambda_6 & -\lambda_5 & \lambda_3 & -\lambda_2 & \lambda_8 & -\lambda_7 & \lambda_4 & \lambda_1 \\ -\lambda_8 & \lambda_7 & -\lambda_4 & \lambda_1 & -\lambda_6 & -\lambda_5 & \lambda_3 & \lambda_2 \end{pmatrix},$$

and the determinant of $M_X M_X^T$ is given by

$$\begin{aligned} & (\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 - \lambda_7^2 - \lambda_8^2)^2 (\lambda_1^2 + \lambda_4^2 + \lambda_7^2 + \lambda_8^2 + \lambda_2^2 + \lambda_3^2 + \lambda_5^2 + \lambda_6^2) \\ & \times \left[\lambda_1^4 + \lambda_2^4 + \lambda_3^4 + 2\lambda_3^2\lambda_4^2 + \lambda_4^4 - 2\lambda_3^2\lambda_5^2 + 2\lambda_4^2\lambda_5^2 + \lambda_5^4 - 2\lambda_3^2\lambda_6^2 + 2\lambda_4^2\lambda_6^2 + 2\lambda_5^2\lambda_6^2 + \lambda_6^4 \right. \\ & - 8\lambda_3\lambda_4\lambda_5\lambda_7 + 2\lambda_3^2\lambda_7^2 - 2\lambda_4^2\lambda_7^2 + 2\lambda_5^2\lambda_7^2 + 2\lambda_6^2\lambda_7^2 + \lambda_7^4 + 8\lambda_3\lambda_4\lambda_6\lambda_8 + 2\lambda_3^2\lambda_8^2 - 2\lambda_4^2\lambda_8^2 \\ & + 2\lambda_5^2\lambda_8^2 + 2\lambda_6^2\lambda_8^2 + 2\lambda_7^2\lambda_8^2 + \lambda_8^4 - 8\lambda_2\lambda_4(\lambda_6\lambda_7 + \lambda_5\lambda_8) \\ & + 2\lambda_1(\lambda_2^2 + \lambda_3^2 + \lambda_4^2 + \lambda_5^2\lambda_6^2 - \lambda_7^2 - \lambda_8^2) + 2\lambda_2^2(\lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 + \lambda_7^2 + \lambda_8^2) \\ & \left. + 8\lambda_1(\lambda_3(\lambda_6\lambda_7 + \lambda_6\lambda_8) + \lambda_2(-\lambda_5\lambda_7 + \lambda_6\lambda_8)) \right]. \end{aligned}$$

Thus if $X \in \mathfrak{v}_{2,3}$ and $\langle X, X \rangle_{\mathfrak{v}_{2,3}} = \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 - \lambda_7^2 - \lambda_8^2 = 0$, then $\det(M_X M_X^T) = 0$. This finishes the proof for $\mathfrak{n}_{2,3}$. \square

We stress that the results of Proposition 6.1 and Proposition 6.5 represent quite exceptional cases. Definition 7.2 does not imply that ad_X is not surjective for any $X \in \mathfrak{v}_{r,s}$ with $\langle X, X \rangle_{\mathfrak{v}_{r,s}} = 0$. In the following we state a couple of lemmas illustrating this possibility.

Lemma 6.2. *For any of the pseudo H -type Lie algebras $\mathfrak{n}_{11,2}$, $\mathfrak{n}_{7,6}$, $\mathfrak{n}_{6,7}$ and $\mathfrak{n}_{2,11}$ there exists X in the corresponding space $\mathfrak{v}_{r,s}$ such that $\langle X, X \rangle_{\mathfrak{v}_{r,s}} = 0$ but, nevertheless, the map ad_X is surjective.*

Proof. Recall that the pseudo H -type Lie algebras $\mathfrak{n}_{11,2}$, $\mathfrak{n}_{7,6}$, $\mathfrak{n}_{6,7}$ and $\mathfrak{n}_{2,11}$ are obtained from $\mathfrak{n}_{3,2}$ and $\mathfrak{n}_{2,3}$ by extensions. We define $X = w_1 \otimes u_1 + w_7 \otimes u_2 \in \mathfrak{n}_{11,2}$ where $w_1, w_7 \in \mathfrak{B}_{\mathfrak{v}_{3,2}}$ and $u_1, u_2 \in \mathfrak{B}_{\mathfrak{v}_{8,0}}$ and note that $\langle X, X \rangle_{\mathfrak{v}_{11,2}} = 0$. Then

$$\begin{aligned} [X, w_i \otimes u_1] &= [w_1 \otimes u_1, w_i \otimes u_1] + [w_7 \otimes u_2, w_i \otimes u_1] \\ &= \begin{cases} -[w_1, w_i]_{\mathfrak{n}_{11,2}} & \text{for } i = 1, \dots, 6, 8, \\ -[w_1, w_i]_{\mathfrak{n}_{11,2}} - [u_2, u_1]_{\mathfrak{n}_{11,2}} & \text{for } i = 7, \end{cases} \\ &= -[w_1, w_i]_{\mathfrak{n}_{11,2}}, \quad \text{for } i = 1, \dots, 8. \end{aligned}$$

Hence $\text{span}\{Z_1, Z_2, Z_3, Z_{12}, Z_{13}\} \subset \text{Image}(\text{ad}_X)$. Furthermore,

$$\begin{aligned} [X, w_1 \otimes u_j] &= [w_1 \otimes u_1, w_1 \otimes u_j] + [w_7 \otimes u_2, w_1 \otimes u_j] \\ &= \begin{cases} [u_1, u_j]_{\mathfrak{n}_{11,2}} & \text{for } j = 1, 3, \dots, 16, \\ [u_1, u_j]_{\mathfrak{n}_{11,2}} - [w_7, w_1]_{\mathfrak{n}_{11,2}} & \text{for } j = 2, \end{cases} \\ &= [u_1, u_j]_{\mathfrak{n}_{11,2}}, \quad \text{for } j = 1, \dots, 16. \end{aligned}$$

Hence $\text{span}\{Z_4, \dots, Z_{11}\} \subset \text{Image}(\text{ad}_X)$, i.e. the map ad_X is surjective.

The proof for $\mathfrak{n}_{7,6}$ and $\mathfrak{n}_{6,7}$ is obtained analogously by replacing u_1 and u_2 by $y_1, y_6 \in \mathfrak{B}_{4,4}$. For the proof for $\mathfrak{n}_{2,11}$ we replace $u_1, u_2 \in \mathfrak{B}_{8,0}$ by $v_1, v_2 \in \mathfrak{B}_{0,8}$, respectively. \square

Lemma 6.3. *For any pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$ with $r, s \neq 0$, satisfying (24), there exists at least one $X \in \mathfrak{v}_{r,s}$ with $\langle X, X \rangle_{\mathfrak{v}_{r,s}} = 0$ such that the map ad_X is surjective.*

Proof. We choose the basis vectors $w_i \in \mathbf{A}_{r,s}^+$ and $w_j \in \mathbf{B}_{r,s}^-$ and define $X = w_i + w_j$ such that $\langle X, X \rangle_{\mathfrak{v}_{r,s}} = 0$. We note that the map $\text{ad}_{w_i} : \mathfrak{V}_{w_i} \rightarrow \mathfrak{z}_{r,s}$ and $\text{ad}_{w_j} : \mathfrak{V}_{w_j} \rightarrow \mathfrak{z}_{r,s}$ are surjective, where we denote by \mathfrak{V}_{w_i} the orthogonal complement to the kernel of ad_{w_i} . Therefore $\mathfrak{V}_{w_i} \subset \text{span}\{\mathbf{B}_{r,s}\}$ and $\mathfrak{V}_{w_j} \subset \text{span}\{\mathbf{A}_{r,s}\}$ as $[w_i, \mathbf{A}_{r,s}] = 0$ and $[w_j, \mathbf{B}_{r,s}] = 0$. It follows that

$$\begin{aligned} \text{span}\{[X, \mathbf{A}_{r,s}]\} &= \text{span}\{[w_i, \mathbf{A}_{r,s}] + [w_j, \mathbf{A}_{r,s}]\} = \text{span}\{[w_j, \mathbf{A}_{r,s}]\} \supset [w_j, \mathfrak{V}_{w_j}] = \mathfrak{z}_{r,s}, \\ \text{span}\{[X, \mathbf{B}_{r,s}]\} &= \text{span}\{[w_i, \mathbf{B}_{r,s}] + [w_j, \mathbf{B}_{r,s}]\} = \text{span}\{[w_i, \mathbf{B}_{r,s}]\} \supset [w_i, \mathfrak{V}_{w_i}] = \mathfrak{z}_{r,s}. \end{aligned}$$

Hence the map ad_X is surjective. \square

Theorem 6.4. *The H -type Lie algebras $\mathfrak{n}_{3,2}$ and $\mathfrak{n}_{2,3}$ are not isomorphic.*

Proof. We assume that there exists an isomorphism $\varphi_{3,2} : \mathfrak{n}_{3,2} \rightarrow \mathfrak{n}_{2,3}$ where the restriction $\varphi_{3,2}|_{\mathfrak{z}_{3,2}} : \mathfrak{z}_{3,2} \rightarrow \mathfrak{z}_{2,3}$ is an anti-isometry. The adjoint operator $\text{ad}_{w_i} : \mathfrak{V}_{w_i} \rightarrow \mathfrak{z}_{3,2}$ is an isometry or anti-isometry by Definition 7.2 for any $w_i \in \mathfrak{B}_{\mathfrak{v}_{3,2}}$, i.e.

$$\langle \text{ad}_{w_i}(X), \text{ad}_{w_i}(X) \rangle_{\mathfrak{z}_{3,2}} = \langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} \langle X, X \rangle_{\mathfrak{v}_{3,2}}$$

for all $X \in \mathfrak{V}_{w_i}$, where \mathfrak{V}_{w_i} is the orthogonal complement to the kernel of ad_{w_i} . As the map $\varphi_{3,2}|_{\mathfrak{z}_{3,2}}$ is an anti-isometry, it follows that the composition $\varphi_{3,2} \circ \text{ad}_{w_i} : \mathfrak{V}_{w_i} \rightarrow \mathfrak{z}_{2,3}$ is an anti-isometry for $\langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} = 1$ and is an isometry for $\langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} = -1$, hence

$$\begin{aligned} -\langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} \langle w_j, w_j \rangle_{\mathfrak{v}_{3,2}} &= \langle \varphi_{3,2} \circ \text{ad}_{w_i}(w_j), \varphi_{3,2} \circ \text{ad}_{w_i}(w_j) \rangle_{\mathfrak{z}_{2,3}} \\ &= \langle [\varphi_{3,2}(w_i), \varphi_{3,2}(w_j)], [\varphi_{3,2}(w_i), \varphi_{3,2}(w_j)] \rangle_{\mathfrak{z}_{2,3}}. \end{aligned}$$

As the map $\varphi_{3,2} \circ \text{ad}_{w_i}$ is surjective and

$$\varphi_{3,2} \circ \text{ad}_{w_i}(w_j) = [\varphi_{3,2}(w_i), \varphi_{3,2}(w_j)] = \text{ad}_{\varphi_{3,2}(w_i)}(\varphi_{3,2}(w_j))$$

it follows by Proposition 6.1 that $\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}} \neq 0$ for all $i = 1, \dots, 8$.

We recall that from Definition 7.2 it follows that for all $X \in \mathfrak{v}_{r,s}$ with $\langle X, X \rangle_{\mathfrak{v}_{r,s}} \neq 0$ and $Y \in \mathfrak{V}_X$:

$$\langle \text{ad}_X(Y), \text{ad}_X(Y) \rangle_{\mathfrak{z}_{r,s}} = \langle X, X \rangle_{\mathfrak{v}_{r,s}} \langle Y, Y \rangle_{\mathfrak{v}_{r,s}},$$

hence

$$\begin{aligned} -\langle w_i, w_i \rangle_{\mathfrak{v}_{3,2}} \langle w_j, w_j \rangle_{\mathfrak{v}_{3,2}} & \tag{32} \\ &= \langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}} \langle \varphi_{3,2}(w_j), \varphi_{3,2}(w_j) \rangle_{\mathfrak{v}_{2,3}}. \end{aligned}$$

We obtain the following relations for w_1 and w_4 :

$$\begin{aligned} \text{sign}(\langle \varphi_{3,2}(w_1), \varphi_{3,2}(w_1) \rangle_{\mathfrak{v}_{2,3}}) &= -\text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}), & \text{for } i = 2, 3, 4, \\ \text{sign}(\langle \varphi_{3,2}(w_1), \varphi_{3,2}(w_1) \rangle_{\mathfrak{v}_{2,3}}) &= \text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}), & \text{for } i = 5, 6, \\ \text{sign}(\langle \varphi_{3,2}(w_4), \varphi_{3,2}(w_4) \rangle_{\mathfrak{v}_{2,3}}) &= -\text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}), & \text{for } i = 1, 2, 3, \\ \text{sign}(\langle \varphi_{3,2}(w_4), \varphi_{3,2}(w_4) \rangle_{\mathfrak{v}_{2,3}}) &= \text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}), & \text{for } i = 5, 6. \end{aligned}$$

It implies that for $i = 2, 3$

$$\begin{aligned} -\text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}) &= \text{sign}(\langle \varphi_{3,2}(w_1), \varphi_{3,2}(w_1) \rangle_{\mathfrak{v}_{2,3}}) \\ &= -\text{sign}(\langle \varphi_{3,2}(w_4), \varphi_{3,2}(w_4) \rangle_{\mathfrak{v}_{2,3}}) \\ &= \text{sign}(\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}}). \end{aligned}$$

Hence $\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}} = 0$ for $i = 2, 3$. This contradicts (32):

$\langle \varphi_{3,2}(w_i), \varphi_{3,2}(w_i) \rangle_{\mathfrak{v}_{2,3}} \neq 0$ for $i = 1, \dots, 8$, as the map $\text{ad}_{\varphi_{3,2}(w_i)}$ is surjective.

Hence $\mathfrak{n}_{3,2}$ is not isomorphic to $\mathfrak{n}_{2,3}$. \square

6.2. Special features of the Lie algebra $\mathfrak{n}_{3,3}$. We introduce an integral basis of $\mathfrak{n}_{3,3}$ by $\mathfrak{B}_{\mathfrak{z}_{3,3}} = \{Z_1, Z_2, Z_3, Z_4, Z_5, Z_6\}$ and

$$\begin{aligned} \mathfrak{B}_{\mathfrak{v}_{3,3}} &= \left\{ \begin{array}{cccc} w_1 = w, & w_2 = J_1 w, & w_3 = J_2 w, & w_4 = J_3 w, \\ w_5 = J_1 J_6 w, & w_6 = J_6 w, & w_7 = J_4 w, & w_8 = J_5 w, \end{array} \right\}, \\ &\text{for } J_2 J_3 J_4 J_5 w = J_1 J_2 J_5 J_6 w = J_1 J_2 J_3 w = w, \\ &\text{and } \langle w_i, w_i \rangle_{\mathfrak{v}_{3,3}} = \epsilon_i(4, 4), \quad \langle Z_k, Z_k \rangle_{\mathfrak{z}_{3,3}} = \epsilon_k(3, 3). \end{aligned}$$

TABLE 9. Commutation relations on $\mathfrak{n}_{3,3}$

[row, col.]	w_1	w_2	w_5	w_6	w_3	w_4	w_7	w_8
w_1	0	Z_1	0	Z_6	Z_2	Z_3	Z_4	Z_5
w_2	$-Z_1$	0	$-Z_6$	0	$-Z_3$	Z_2	$-Z_5$	Z_4
w_5	0	Z_6	0	Z_1	Z_5	Z_4	Z_3	Z_2
w_6	$-Z_6$	0	$-Z_1$	0	Z_4	$-Z_5$	Z_2	$-Z_3$
w_3	$-Z_2$	Z_3	$-Z_5$	$-Z_4$	0	$-Z_1$	Z_6	0
w_4	$-Z_3$	$-Z_2$	$-Z_4$	Z_5	Z_1	0	0	$-Z_6$
w_7	$-Z_4$	Z_5	$-Z_3$	$-Z_2$	$-Z_6$	0	0	Z_1
w_8	$-Z_5$	$-Z_4$	$-Z_2$	Z_3	0	Z_6	$-Z_1$	0

Proposition 6.5. *The linear map $\text{ad}_X: \mathfrak{v}_{3,3} \rightarrow \mathfrak{z}_{3,3}$ is surjective if and only if $\langle X, X \rangle_{\mathfrak{v}_{3,3}} \neq 0$ for $X \in \mathfrak{v}_{3,3}$.*

Proof. We use similar arguments as in the proof of Proposition 6.1. The matrix M_X that we calculate by using Table 9 is given by

$$\begin{pmatrix} -\lambda_2 & \lambda_1 & -\lambda_6 & \lambda_5 & \lambda_4 & -\lambda_3 & -\lambda_8 & \lambda_7 \\ -\lambda_3 & -\lambda_4 & -\lambda_8 & -\lambda_7 & \lambda_1 & \lambda_2 & \lambda_6 & \lambda_5 \\ -\lambda_4 & \lambda_3 & -\lambda_7 & \lambda_8 & -\lambda_2 & \lambda_1 & \lambda_5 & -\lambda_6 \\ -\lambda_7 & -\lambda_8 & -\lambda_4 & -\lambda_3 & \lambda_6 & \lambda_5 & \lambda_1 & \lambda_2 \\ -\lambda_8 & \lambda_7 & -\lambda_3 & \lambda_4 & \lambda_5 & -\lambda_6 & -\lambda_2 & \lambda_1 \\ -\lambda_6 & \lambda_5 & -\lambda_2 & \lambda_1 & -\lambda_7 & \lambda_8 & \lambda_3 & -\lambda_4 \end{pmatrix}.$$

The determinant of $M_X M_X^T$ has the form

$$\begin{aligned} &(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 - \lambda_5^2 - \lambda_6^2 - \lambda_7^2 - \lambda_8^2)^4 \\ &\times ((\lambda_1 - \lambda_5)^2 + (\lambda_2 - \lambda_6)^2 + (\lambda_4 - \lambda_7)^2 + (\lambda_3 - \lambda_8)^2) \\ &\times ((\lambda_1 + \lambda_5)^2 + (\lambda_2 + \lambda_6)^2 + (\lambda_4 + \lambda_7)^2 + (\lambda_3 + \lambda_8)^2). \end{aligned}$$

Thus the map ad_X is surjective if and only if $\langle X, X \rangle_{\mathfrak{v}_{3,3}} \neq 0$. \square

Theorem 6.6. *There does not exist an automorphism $\varphi_{3,3}$ of $\mathfrak{n}_{3,3}$ such that the restriction to the center $\varphi|_{\mathfrak{z}_{3,3}}$ is an anti-isometry.*

Proof. By repeating the arguments of the proof of Theorem 6.4, we obtain equation (32). This implies the relations

$$\begin{aligned} \text{sign}(\langle \varphi_{3,3}(w_1), \varphi_{3,3}(w_1) \rangle_{\mathfrak{v}_{3,3}}) &= -\text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}), & \text{for } i = 2, 3, 4, \\ \text{sign}(\langle \varphi_{3,3}(w_1), \varphi_{3,3}(w_1) \rangle_{\mathfrak{v}_{3,3}}) &= \text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}), & \text{for } i = 6, 7, 8, \\ \text{sign}(\langle \varphi_{3,3}(w_2), \varphi_{3,3}(w_2) \rangle_{\mathfrak{v}_{3,3}}) &= -\text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}), & \text{for } i = 1, 3, 4, \\ \text{sign}(\langle \varphi_{3,3}(w_2), \varphi_{3,3}(w_2) \rangle_{\mathfrak{v}_{3,3}}) &= \text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}), & \text{for } i = 5, 7, 8. \end{aligned}$$

Thus, for $i = 3, 4$

$$\begin{aligned} -\text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}) &= \text{sign}(\langle \varphi_{3,3}(w_1), \varphi_{3,3}(w_1) \rangle_{\mathfrak{v}_{3,3}}) \\ &= -\text{sign}(\langle \varphi_{3,3}(w_2), \varphi_{3,3}(w_2) \rangle_{\mathfrak{v}_{3,3}}) \\ &= \text{sign}(\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}}). \end{aligned}$$

Hence $\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}} = 0$ for $i = 3, 4$. This contradicts to the fact that $\langle \varphi_{3,3}(w_i), \varphi_{3,3}(w_i) \rangle_{\mathfrak{v}_{3,3}} \neq 0$ for $i = 1, \dots, 8$, as the map $\text{ad}_{\varphi_{3,3}(w_i)}$ is surjective.

Hence there does not exist an automorphism $\varphi_{3,3}$ of $\mathfrak{n}_{3,3}$ such that $\varphi|_{\mathfrak{z}_{3,3}}$ is an anti-isometry. \square

Proposition 6.7. *For the pseudo H -type Lie algebras $\mathfrak{n}_{11,3}$, $\mathfrak{n}_{7,7}$ and $\mathfrak{n}_{3,11}$ there exists X in respective $\mathfrak{v}_{r,s}$ such that ad_X is surjective.*

Proof. We replace in the proof of Lemma 6.2 $w_1, w_7 \in \mathfrak{B}_{\mathfrak{v}_{2,3}}$ by $w_1, w_5 \in \mathfrak{B}_{\mathfrak{v}_{3,3}}$. \square

7. STRONGLY BRACKET GENERATING PROPERTY

In this section we study the bracket generating property of the pseudo H -type Lie algebras. For that purpose we state an equivalent definition of the pseudo H -type algebras $\mathfrak{n}_{r,s}$, which is related to the definition of the strongly bracket generating property.

Definition 7.1. *Let $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$ be a pseudo H -type Lie algebra. We call a vector space $\mathfrak{v}_{r,s}$ strongly bracket generating if for any non-zero $v \in \mathfrak{v}_{r,s}$ the linear map $\text{ad}_v = [v, \cdot]: \mathfrak{v}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ is surjective, i.e. $\text{span}\{\mathfrak{v}_{r,s}, [v, \mathfrak{v}_{r,s}]\} = \mathfrak{n}_{r,s}$ for all $v \in \mathfrak{v}_{r,s} \setminus \{0\}$. We say in this case that the pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$ has strongly bracket generating property.*

Let $N_{r,s}$ be the Lie group, corresponding to the pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$ and let \mathcal{H} be the left translation of the vector space $\mathfrak{v}_{r,s}$. If $\mathfrak{v}_{r,s}$ is strongly bracket generating, then the left invariant distribution \mathcal{H} is strongly bracket generating in a sense that $\text{span}\{\mathcal{H}, [X, \mathcal{H}]\} = TN_{r,s}$ for any smooth non-zero section X of the distribution \mathcal{H} .

Even the strongly bracket generating property seems to be just of interest from a geometrical point of view, it actually has a close relation to an equivalent definition of pseudo H -type algebras, which can be seen in the following subsection.

7.1. An equivalent definition of pseudo H -type Lie algebras. In this subsection we define general H -type algebras [3] and prove that they are equivalent to pseudo H -type algebras.

Let $\mathfrak{n} = (\mathfrak{v} \oplus_{\perp} \mathfrak{z}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ be an arbitrary 2-step nilpotent Lie algebra with center \mathfrak{z} and a non-degenerate scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{n}}$. We write $\text{ad}_v: \mathfrak{v} \rightarrow \mathfrak{z}$

for the linear map given by $\text{ad}_v w = [v, w]$. We assume that the restriction of the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ onto the subspace $\ker(\text{ad}_v) \subset \mathfrak{v}$ is non-degenerate and denote its orthogonal complement with respect to $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ by \mathfrak{V}_v , which is also non-degenerate. Thus the restricted map $\text{ad}_v: \mathfrak{V}_v \rightarrow \mathfrak{z}$ is injective.

Definition 7.2. [3] *A two-step nilpotent Lie algebra $\mathfrak{n} = (\mathfrak{v} \oplus_{\perp} \mathfrak{z}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ is of general H -type if $\text{ad}_v: \mathfrak{V}_v \rightarrow \mathfrak{z}$ is a surjective isometry for all $v \in \mathfrak{v}$ with $\langle v, v \rangle_{\mathfrak{v}} = 1$ and a surjective anti-isometry for all $v \in \mathfrak{v}$ with $\langle v, v \rangle_{\mathfrak{v}} = -1$.*

Let $(U, \langle \cdot, \cdot \rangle_U)$, $(V, \langle \cdot, \cdot \rangle_V)$ be two vector spaces with corresponding non-degenerate quadratic forms, written as bi-linear symmetric forms, or scalar products.

Definition 7.3. *A bilinear map $\mu: U \times V \rightarrow V$ is called a composition of the scalar products $\langle \cdot, \cdot \rangle_U$ of U and $\langle \cdot, \cdot \rangle_V$ of V if the equality*

$$\langle \mu(u, v), \mu(u, v) \rangle_V = \langle u, u \rangle_U \langle v, v \rangle_V \quad (33)$$

holds for any $u \in U$ and $v \in V$.

We assume that there is $u_0 \in U$ such that $\langle u_0, u_0 \rangle_U = 1$ and $\mu(u_0, v) = v$. This can always be done by normalization procedure of quadratic forms, see [8]. Let us denote by \mathcal{Z} the orthogonal complement to the non-degenerate space $\text{span}\{u_0\}$ and by J the restriction of μ to \mathcal{Z} , thus $J: \mathcal{Z} \times V \rightarrow V$. The map J is skew-adjoint in the sense that $\langle J(Z, v), v' \rangle_V = -\langle v, J(Z, v') \rangle_V$ for any $Z \in \mathcal{Z}$ and $v, v' \in V$. Therefore, the map J can be used to define a Lie algebra structure on $\mathfrak{n} = \mathcal{Z} \oplus V$ by $\langle J(Z, v), v' \rangle_V = \langle Z, [v, v'] \rangle_{\mathcal{Z}}$. The obtained Lie algebra is a general H -type algebra, see [3, Theorem 1]. Now, rephrasing Definition 2.2 of a pseudo H -type algebra we can say that a two-step nilpotent Lie algebra is a pseudo H -type algebra if the map J defined by (2.1) is the restriction on the center $\mathcal{Z} = \mathfrak{z}$ of a composition of corresponding quadratic forms for vector spaces $V = \mathfrak{v}$, $U = \text{span}\{u_0\} \oplus_{\perp} \mathcal{Z}$.

Theorem 7.4. *Definitions 2.2 and 7.2 are equivalent.*

Proof. Let us prove that Definition 7.2 implies Definition 2.2. It was shown in [3, Theorem 1] that any general H -type algebra $\mathfrak{n} = (\mathfrak{v} \oplus_{\perp} \mathfrak{z}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ defines a composition of the quadratic form $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ and another quadratic form whose restriction on \mathfrak{z} coincides with $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$. Particularly, it implies (2) and therefore a general H -type algebra is a pseudo H -type algebra.

Now we assume that we are given a pseudo H -type algebra $\mathfrak{n} = (\mathfrak{v} \oplus_{\perp} \mathfrak{z}, [\cdot, \cdot], \langle \cdot, \cdot \rangle_{\mathfrak{n}} = \langle \cdot, \cdot \rangle_{\mathfrak{v}} + \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ with center \mathfrak{z} . Let us fix $v \in \mathfrak{v}$ with $\langle v, v \rangle_{\mathfrak{v}} = \pm 1$. We need to show that $\text{ad}_v: \mathfrak{v} \rightarrow \mathfrak{z}$ is a surjective (anti-)isometry. The following equation is true

$$\langle Z, \text{ad}_v(J_{Z'}v) \rangle_{\mathfrak{z}} = \langle Z, [v, J_{Z'}v] \rangle_{\mathfrak{z}} = \langle J_{Z'}v, J_{Z'}v \rangle_{\mathfrak{v}} = \langle Z, Z' \rangle_{\mathfrak{z}} \langle v, v \rangle_{\mathfrak{v}} = \pm \langle Z, Z' \rangle_{\mathfrak{z}},$$

for all $Z, Z' \in \mathfrak{z}$ by formula (4). We use both notations $J(Z, v)$ and $J_{Z'}v$. This implies $\text{ad}_v(J_{Z'}v) = \pm Z'$ for all $Z' \in \mathfrak{z}$ by the non-degenerate property of the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$. Since $\langle J_{Z'}v, w \rangle_{\mathfrak{v}} = \langle Z', [v, w] \rangle_{\mathfrak{z}} = \langle Z', 0 \rangle_{\mathfrak{z}} = 0$, for all $w \in \ker(\text{ad}_v)$, it follows that $J_{Z'}v \in \mathfrak{V}_v = (\ker(\text{ad}_v))^{\perp}$. We showed that ad_v is surjective.

To prove that ad_v is an isometry for $\langle v, v \rangle_{\mathfrak{v}} = 1$ and an anti-isometry for $\langle v, v \rangle_{\mathfrak{v}} = -1$, we exhibit that the maps $\text{ad}_v: \mathfrak{V}_v \rightarrow \mathfrak{z}$ and $J_{(\cdot)}v: \mathfrak{z} \rightarrow \mathfrak{V}_v$ are inverse and then equality (2) implies the isometry and anti-isometry properties. Let us assume that

$\langle v, v \rangle_{\mathfrak{v}} = 1$. We proved that $\text{ad}_v: \mathfrak{V}_v \rightarrow \mathfrak{z}$ is bijective, thus the image of $J_{(\cdot)}v$ belongs to \mathfrak{V}_v , and $\text{ad}_v(J_{(\cdot)}v) = \text{Id}_{\mathfrak{z}}$, where $\text{Id}_{\mathfrak{z}}$ is the identity map on \mathfrak{z} .

We claim that the map $J_{(\cdot)}v: \mathfrak{z} \rightarrow \mathfrak{V}_v$ is bijective. Indeed if we assume that $J_{(\cdot)}v$ is not surjective, then there is $w \in \mathfrak{V}_v$ that is not in the image of $J_{(\cdot)}v$. Let $\text{ad}_v(w) = Z \in \mathfrak{z}$, then $\text{ad}_v(J_Z v) = Z$ which implies $w = J_Z v$ by injectivity of ad_v and leads to contradiction.

If we now assume that $J_{(\cdot)}v$ is not injective, then we find $Z', Z'' \in \mathfrak{z}$, $Z' \neq Z''$, such that $J_{Z'}(v) = J_{Z''}(v)$. But in this case $Z' = \text{ad}_X(J_{Z'}v) = \text{ad}_v(J_{Z''}v) = Z''$ by bijectivity of ad_v and we again get a contradiction. The proof for $\langle v, v \rangle_{\mathfrak{v}} = -1$ is analogous and we conclude that ad_v and $J_{(\cdot)}v$ are inverse maps to each other. Equality (2) becomes

$$\langle J_Z v, J_Z v \rangle_{\mathfrak{v}} = \langle Z, Z \rangle_{\mathfrak{z}} \text{ for } \langle v, v \rangle_{\mathfrak{v}} = 1, \text{ and } \langle J_Z v, J_Z v \rangle_{\mathfrak{v}} = -\langle Z, Z \rangle_{\mathfrak{z}} \text{ for } \langle v, v \rangle_{\mathfrak{v}} = -1,$$

that shows the (anti-)isometry property of the map $J_{(\cdot)}v: \mathfrak{z} \rightarrow \mathfrak{V}_v$ and its inverse $\text{ad}_v: \mathfrak{V}_v \rightarrow \mathfrak{z}$. \square

7.2. Bracket generating property of pseudo H -type algebras.

Theorem 7.5. *The pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$ with $r = 0$ or $s = 0$ have the strongly bracket generating property.*

Proof. Let $s = 0$. This implies that $\langle v, v \rangle_{\mathfrak{v}_{r,0}} > 0$ for all $v \in \mathfrak{v}_{r,0}$ with $v \neq 0$. Definition 7.2 yields that ad_v is surjective, i.e. $\mathfrak{v}_{r,0}$ is strongly bracket generating.

Let now $r = 0$. Recall that $\mathfrak{v}_{0,s}$ is a neutral space, i.e. $\langle \cdot, \cdot \rangle_{\mathfrak{v}_{0,s}}$ has index (l, l) and we can identify $\mathfrak{v}_{0,s}$ with $\mathbb{R}^{l,l}$. This implies that there exists elements $v \in \mathfrak{v}_{0,s}$, $v \neq 0$, with $\langle v, v \rangle_{l,l} = 0$. According to Definition 7.2 we need only to show that $\text{ad}_v: \mathfrak{v}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ is surjective for vectors with $\langle v, v \rangle_{l,l} = 0$, since for all other vectors the adjoint map is surjective.

We define the orthonormal basis $\{w_1, \dots, w_{2l}\}$ of $\mathfrak{v}_{0,s}$ with $\langle w_i, w_i \rangle_{\mathfrak{v}_{0,s}} = \epsilon_i(l, l)$ and fix an arbitrary $v \in \mathfrak{v}_{0,s}$ with $\langle v, v \rangle_{l,l} = 0$ and $v = \sum_{i=1}^{2l} \lambda_i w_i$. We split v in the form $v = v^+ + v^-$, with $v^+ = \sum_{i=1}^l \lambda_i w_i$, $v^- = \sum_{i=l+1}^{2l} \lambda_i w_i$ and $\langle v^+, v^+ \rangle_{l,l} = -\langle v^-, v^- \rangle_{l,l} > 0$ and $\langle v^+, v^- \rangle_{l,l} = 0$. We note that $[w_i, w_j] = 0$ if $i, j = 1, \dots, l$ or $i, j = l+1, \dots, 2l$ as

$$\langle [w_i, w_j], [w_i, w_j] \rangle_{\mathfrak{z}_{0,s}} \geq 0, \quad \text{for } i, j = 1, \dots, l, \quad \text{or} \quad i, j = l+1, \dots, 2l.$$

Hence $\mathfrak{z}_{0,s} = \text{ad}_{w_i}(\mathfrak{v}_{0,s}) = \text{ad}_{w_i}(\text{span}\{w_1, \dots, w_l\})$ for $i = l+1, \dots, 2l$. It follows that

$$[v, \text{span}\{w_1, \dots, w_l\}] = [v^-, \text{span}\{w_1, \dots, w_l\}] = \mathfrak{z}_{0,s}.$$

Hence ad_v is surjective, i.e. the pseudo H -type algebras $\mathfrak{n}_{0,s} = \mathfrak{v}_{0,s} \oplus \mathfrak{z}_{0,s}$, where $s > 0$, have strongly bracket generating property. \square

Theorem 7.6. *The pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$ with $r, s \neq 0$ do not have the strongly bracket generating property.*

Proof. We assume that $\mathfrak{n}_{r,s} = \mathfrak{v}_{r,s} \oplus \mathfrak{z}_{r,s}$ with $r, s \neq 0$ has the strongly bracket generating property, i.e. for all $v \in \mathfrak{v}_{r,s}$: $[v, \mathfrak{v}_{r,s}] = \mathfrak{z}_{r,s}$ and we show that it contradicts to the presence of nullvectors in the scalar product space $(\mathfrak{z}_{r,s}, \langle \cdot, \cdot \rangle_{r,s})$. The non-degenerate property of the indefinite scalar-product $\langle \cdot, \cdot \rangle_{r,s}$ implies that for all $v \in \mathfrak{v}_{r,s}$ and for all $Z \in \mathfrak{z}_{r,s}$ there exists $v_Z \in \mathfrak{v}_{r,s}$ such that

$$\langle [v, v_Z], Z \rangle_{r,s} \neq 0 \quad \stackrel{(1)}{\iff} \quad \langle J_Z(v), v_Z \rangle_{\mathfrak{v}_{r,s}} \neq 0.$$

It follows that $J_Z(v) \neq 0$ for all $v \in \mathfrak{v}_{r,s}$ and for all $Z \in \mathfrak{z}_{r,s}$, i.e. $\ker\{J_Z\} = \{0\}$ for all $Z \in \mathfrak{z}_{r,s}$. But there exist elements $Z_0 \in \mathfrak{z}_{r,s}$ such that $\langle Z_0, Z_0 \rangle_{r,s} = 0$ as $r, s \neq 0$. This implies that $J_{Z_0}^2 = 0$ which is equivalent to $\ker\{J_{Z_0}\} \neq \{0\}$. This is a contradiction, hence the pseudo H -type Lie algebras $\mathfrak{n}_{r,s}$ with $r, s \neq 0$ do not have the strongly bracket generating property. \square

8. NON-ISOMORPHISM PROPERTIES FOR PSEUDO H -TYPE GROUPS IN GENERAL POSITION

In this section we discuss the possible extension of our results to pseudo H -type Lie algebras, constructed from non-minimal admissible Clifford modules. Here we need to distinguish two essentially different situations. The first one when the irreducible module is unique and other one when there are two non-equivalent irreducible modules. We introduce new notations.

1. Let the Clifford algebras $\text{Cl}_{r,s}$ admit only one (up to equivalence) irreducible module and we write $\mathfrak{v}_{r,s}$ for the minimal admissible module, that could be a direct sum of two irreducible modules. This situation occurs when $r - s \not\equiv 3 \pmod{4}$. Any non-minimal admissible $\text{Cl}_{r,s}$ -module \mathfrak{v} is isomorphic (and isometric) to the direct sum of minimal admissible modules $\mathfrak{v}_{r,s}$, see [2, 9]:

$$\mathfrak{v} = \mathfrak{v}_{r,s}(\mu) \cong \bigoplus^{\mu} \mathfrak{v}_{r,s}.$$

Here and further on we use the notation $\mathfrak{v}_{r,s}(\mu)$ for the μ -fold direct sum of minimal admissible modules $\mathfrak{v}_{r,s}$. Thus the argument μ shows how many equivalent (in the sense of the representation theory) minimal admissible modules contains the sum. The lower index, as previously, indicates the index of the metric of the generating space for the Clifford algebra.

2. If $r - s \equiv 3 \pmod{4}$, then the Clifford algebra $\text{Cl}_{r,s}$ admits two non-equivalent Clifford modules. We write $\mathfrak{v}_{r,s}^1$ and $\mathfrak{v}_{r,s}^2$ for the minimal admissible modules. Recall, that in this case each of the admissible modules $\mathfrak{v}_{r,s}^l$, $l = 1, 2$ is either irreducible, or the direct sum of two equivalent irreducible modules, where the representation map is changed appropriately [1]. We emphasize that a minimal admissible module $\mathfrak{v}_{r,s}^l$, $l = 1, 2$, can not be a direct sum of two non-equivalent irreducible modules. In this case a non-minimal admissible $\text{Cl}_{r,s}$ -module \mathfrak{v} is isomorphic to

$$\mathfrak{v} = \mathfrak{v}_{r,s}(\mu, \nu) \cong (\bigoplus^{\mu} \mathfrak{v}_{r,s}^1) \bigoplus (\bigoplus^{\nu} \mathfrak{v}_{r,s}^2)$$

for some positive integers μ, ν which show the number of equivalent and non-equivalent minimal admissible modules contained in the admissible module $\mathfrak{v}_{r,s}(\mu, \nu)$. To unify the notation we always write $\mathfrak{v}_{r,s}(\mu, \nu)$, where $\nu = 0$ if $r - s \not\equiv 3 \pmod{4}$ and ν can be different from zero in the case $r - s \equiv 3 \pmod{4}$. According to this new notation we also write $\mathfrak{n}_{r,s}(\mu, \nu)$ for a pseudo H -type Lie algebra in the case if it is isomorphic to the direct sum $\mathfrak{v}_{r,s}(\mu, \nu) \oplus \mathfrak{z}_{r,s}$.

Results of [2] imply also that a non-minimal admissible module $(\mathfrak{v}_{r,s}(\mu, \nu), \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}(\mu, \nu)})$ of the Clifford algebra $\text{Cl}_{r,s}$ is given as an orthogonal sum of n -dimensional minimal admissible modules $\mathfrak{v}_{r,s} = (\mathfrak{v}_{r,s}, \langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}})$, where each scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}}$ is the restriction of $\langle \cdot, \cdot \rangle_{\mathfrak{v}_{r,s}(\mu, \nu)}$ on the corresponding copy of the vector space $\mathfrak{v}_{r,s}$. To describe the Lie bracket on $\mathfrak{n}_{r,s}(\mu, \nu)$ we proceed as follows. Let $\{Z_1, \dots, Z_m\}$ be an orthonormal basis of $\mathfrak{z}_{r,s}$. We denote a basis of j -term in the sum $\bigoplus^{\mu} \mathfrak{v}_{r,s}^l$, $l = 1, 2$ by $\{v_{1j}^l, \dots, v_{n_j}^l\}$ with

structure constants $(A_{ip}^k)_j^l$. For the sum $\mathfrak{n}_{r,s}(\mu, \nu) = ((\oplus_{j=1}^{\mu}(\mathfrak{v}_{r,s}^1)_j) \oplus (\oplus_{q=1}^{\nu}(\mathfrak{v}_{r,s}^2)_q)) \oplus \mathfrak{z}_{r,s}$ we choose the basis

$$\{v_{ij}^1, v_{pq}^2, Z_k \mid i, p = 1, \dots, n, k = 1, \dots, r+s, j = 1, \dots, \mu, q = 1, \dots, \nu\}. \quad (34)$$

The Lie bracket on $\mathfrak{n}_{r,s}(\mu, \nu)$ with respect to this basis is given by

$$[w_{ij}^{l_1}, w_{pq}^{l_2}] = \delta_{l_1 l_2} \delta_{jq} \sum_{k=1}^{r+s} (A_{ip}^k)_j^{l_t} Z_k, \quad t = 1, 2. \quad (35)$$

The bilinear maps $J_j^l: \mathfrak{z}_{r,s} \times (\mathfrak{v}_{r,s}^l)_j \rightarrow (\mathfrak{v}_{r,s}^l)_j$, $l = 1, 2$ are defined by a representation of $\text{Cl}_{r,s}$ over $(\mathfrak{v}_{r,s}^l)_j$ and are extended to $\tilde{J}: \mathfrak{z}_{r,s} \times \mathfrak{v}_{r,s}(\mu, \nu) \rightarrow \mathfrak{v}_{r,s}(\mu, \nu)$ by

$$\tilde{J} := \begin{pmatrix} J_1^1 & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & J_\nu^2 \end{pmatrix}.$$

Then the operator $\tilde{J}: \mathfrak{z}_{r,s} \rightarrow \text{End}(\mathfrak{v}_{r,s}(\mu, \nu))$ satisfies $\langle \tilde{J}_Z v, w \rangle_{\mathfrak{v}_{r,s}(\mu, \nu)} = \langle Z, [v, w] \rangle_{\mathfrak{z}_{r,s}}$, for all $Z \in \mathfrak{z}_{r,s}$, $v, w \in \mathfrak{v}_{r,s}(\mu, \nu)$ and can be extended to the representation of $\text{Cl}_{r,s}$ over $\mathfrak{v}_{r,s}(\mu, \nu)$. We can assume, by a change of coordinates, without loss of generality, that $J_1^1 = \dots = J_\mu^1$ and $J_1^2 = \dots = J_\nu^2$. If a Clifford algebra $\text{Cl}_{r,s}$ admits only one irreducible representation, then the notation simplifies due to the absence of upper indices $l = 1, 2$.

We start from general observations where the first one follows easily from the dimension argument.

Proposition 8.1. *Pseudo H -type Lie algebras $\mathfrak{n}_{r,s}(\mu_1, 0)$ and $\mathfrak{n}_{r,s}(\mu_2, 0)$ and respectively $\mathfrak{n}_{r,s}(0, \mu_1)$ and $\mathfrak{n}_{r,s}(0, \mu_2)$ for $r - s \not\equiv 3 \pmod{4}$ are isomorphic if and only if $\mu_1 = \mu_2$.*

Theorem 8.2. *Two H -type Lie algebras $\mathfrak{n}_{r,s}(\mu_1, \nu_1)$ and $\mathfrak{n}_{r,s}(\mu_2, \nu_2)$ for $r - s \equiv 3 \pmod{4}$ are isomorphic if and only if $\mu_1 = \mu_2$ and $\nu_1 = \nu_2$ or $\mu_1 = \nu_2$ and $\nu_1 = \mu_2$.*

Proof. In the first step we show that $\mathfrak{n}_{r,s}(\mu, 0)$ and $\mathfrak{n}_{r,s}(0, \mu)$ are isomorphic for $r - s \equiv 3 \pmod{4}$. Let $J^1: \mathfrak{v}_{r,s}^1 \oplus \mathfrak{z}_{r,s} \rightarrow \mathfrak{v}_{r,s}^1$ and $J^2: \mathfrak{v}_{r,s}^2 \oplus \mathfrak{z}_{r,s} \rightarrow \mathfrak{v}_{r,s}^2$ be two non-equivalent representations over two minimal admissible modules. Let $\mathfrak{n}_{r,s}(1, 0) = (\mathfrak{v}_{r,s}^1 \oplus \mathfrak{z}_{r,s}, [\cdot, \cdot]^1)$ and $\mathfrak{n}_{r,s}(0, 1) = (\mathfrak{v}_{r,s}^2 \oplus \mathfrak{z}_{r,s}, [\cdot, \cdot]^2)$ be the pseudo H -type Lie algebras, where we used the maps J^1 and J^2 to define the corresponding brackets by (2.1). We can assume that the vector spaces $\mathfrak{v}_{r,s}^1$ and $\mathfrak{v}_{r,s}^2$ are isomorphic under an isomorphism $A: \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^2$. We define a map $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ by

$$J^1(v, C(Z)) = A^\tau \circ J^2(A(v), Z), \quad \text{for any } v \in \mathfrak{v}_{r,s}^1, Z \in \mathfrak{z}_{r,s}, \quad (36)$$

where $\langle Av, u \rangle_{\mathfrak{v}_{r,s}^2} = \langle v, A^\tau u \rangle_{\mathfrak{v}_{r,s}^1}$. We claim that the map $F = A \oplus C^\tau: \mathfrak{v}_{r,s}^1 \oplus \mathfrak{z}_{r,s} \rightarrow \mathfrak{v}_{r,s}^2 \oplus \mathfrak{z}_{r,s}$ is a Lie algebra isomorphism $F: \mathfrak{n}_{r,s}(1, 0) \rightarrow \mathfrak{n}_{r,s}(0, 1)$, where C^τ is the adjoint map to C with respect to the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{z}_{r,s}}$. Indeed, the chain of equalities

$$\begin{aligned} \langle Z, C^\tau([v, w]^1) \rangle_{\mathfrak{z}_{r,s}} &= \langle C(Z), [v, w]^1 \rangle_{\mathfrak{z}_{r,s}} = \langle J_{C(Z)}^1 v, w \rangle_{\mathfrak{v}_{r,s}^1} = \langle A^\tau \circ J_Z^2(Av), w \rangle_{\mathfrak{v}_{r,s}^1} \\ &= \langle J_Z^2(Av), Aw \rangle_{\mathfrak{v}_{r,s}^2} = \langle Z, [Av, Aw]^2 \rangle_{\mathfrak{z}_{r,s}} \end{aligned}$$

for any $v, w \in \mathfrak{v}_{r,s}^1$, $Z \in \mathfrak{z}_{r,s}$ shows that $F([v, w]^1) = C^\tau([v, w]^1) = [Av, Aw]^2 = [Fv, Fw]^2$.

To show that the Lie algebras $\mathfrak{n}_{r,s}(\mu, \nu)$ and $\mathfrak{n}_{r,s}(\nu, \mu)$ are isomorphic, we choose the map $A: \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^2$ to be not only the isomorphism of vector spaces, but also an isometry between the admissible modules. It, particularly, implies that $A^\tau = A^{-1}$. The corresponding map $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ will also be an isometry by Theorem 3.5. We fix an orthonormal basis Z_1, \dots, Z_{r+s} of $\mathfrak{z}_{r,s}$, then the set $C(Z_1), \dots, C(Z_{r+s})$ also forms an orthonormal basis. We construct an integral basis $v_{11}^2, \dots, v_{n1}^2$ of $\mathfrak{v}_{r,s}^2$ by using the map $J^2: \mathfrak{z}_{r,s} \oplus \mathfrak{v}_{r,s}^2 \rightarrow \mathfrak{v}_{r,s}^2$ and the orthonormal basis Z_1, \dots, Z_{r+s} as it was done in [2]. Then, by making use of the same method, we obtain the integral basis $v_{11}^1, \dots, v_{n1}^1$ constructed from the orthonormal basis $C(Z_1), \dots, C(Z_{r+s})$ and the map $J^1: \mathfrak{z}_{r,s} \oplus \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^1$. By the choice of the map A we get $\prod_{k=1}^l J_{C(Z_{i_k})}^1 = A^{-1} \circ \left(\prod_{k=1}^l J_{Z_{i_k}}^2 \right) \circ A$ for any choice of orthonormal generators Z_{i_1}, \dots, Z_{i_l} in $\mathfrak{z}_{r,s}$. It guarantees that there is a vector $v \in \mathfrak{v}_{r,s}^1$ such that $\langle v, v \rangle_{\mathfrak{v}_{r,s}^1} = \langle Av, Av \rangle_{\mathfrak{v}_{r,s}^2}$ and $\prod_{k=1}^l J_{C(Z_{i_k})}^1 v = v$ implies $\prod_{k=1}^l J_{Z_{i_k}}^2 (Av) = Av$. The method of the construction of the integral basis in [2] implies that $v_{ij}^2 = Av_{ij}^1$. Hence the structural constants with respect to the basis $\{v_{11}^2, \dots, v_{n1}^2, Z_1, \dots, Z_{r+s}\}$ are identical to the structural constants with respect to the basis $\{v_{11}^1, \dots, v_{n1}^1, C(Z_1), \dots, C(Z_{r+s})\}$. More precise, if we write $(A_{ip}^k)_1^1 = (A_{ip}^k)_1^2 = A_{ip}^k$ in the notation (35), then

$$[v_{i1}^1, v_{p1}^1] = \sum_{k=1}^{r+s} A_{ip}^k C(Z_k) \quad \text{and} \quad [v_{i1}^2, v_{p1}^2] = \sum_{k=1}^{r+s} A_{ip}^k Z_k.$$

We can find the exact form of the map $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$. Let $\mathcal{Z} = \{Z_1, \dots, Z_{r+s}\}$ be an orthonormal basis for $\mathfrak{z}_{r,s}$. Then the volume elements have different actions on their modules, namely $\omega^1(\mathcal{Z}) = \prod_{k=1}^{r+s} J_{Z_k}^1 = \text{Id}$ on $\mathfrak{v}_{r,s}^1$ and $\omega^2(\mathcal{Z}) = \prod_{k=1}^{r+s} J_{Z_k}^2 = -\text{Id}$ on $\mathfrak{v}_{r,s}^2$, see [9]. Let $A: \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^2$ be an isometry and $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ be the mapping induced by (36). Then

$$\omega^1(C(\mathcal{Z}))v = \prod_{k=1}^{r+s} J_{C(Z_k)}^1 v = A^{-1} \circ \prod_{k=1}^{r+s} J_{Z_k}^2 \circ Av = A^{-1} \omega^2(\mathcal{Z}) Av = A^{-1}(-Av) = -v$$

for $v \in \mathfrak{v}_{r,s}^1$. Since for $r - s = 3 \pmod{4}$ we have $r + s = 2(s + 2k + 1) + 1$, $k \in \mathbb{Z}$, we conclude that $r + s$ is an odd number. Then from

$$\prod_{k=1}^{r+s} J_{Z_k}^1 = \omega^1(\mathcal{Z}) = \text{Id}_{\mathfrak{v}_{r,s}^1} = -\omega^1(C(\mathcal{Z})) = -\prod_{k=1}^{r+s} J_{C(Z_k)}^1 = \prod_{k=1}^{r+s} J_{-C(Z_k)}^1$$

we can assume that the map $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$ maps the basis $\mathcal{Z} = \{Z_1, \dots, Z_{r+s}\}$ to the basis $-\mathcal{Z} = \{-Z_1, \dots, -Z_{r+s}\}$. We write

$$\{v_{ij}^1, v_{iq}^2, Z_k | i = 1, \dots, n, \quad j = 1, \dots, \mu, \quad q = 1, \dots, \nu, \quad k = 1, \dots, r + s\}$$

for an integral basis of $\mathfrak{n}_{r,s}(\mu, \nu)$ where v_{ij}^1 is the i -th coordinate in the j -s copy of the module $\mathfrak{v}_{r,s}^1$ and v_{iq}^2 is the i -th coordinate in the q -s copy of the n -dimensional admissible module $\mathfrak{v}_{r,s}^2$. Analogously,

$$\{v_{iq}^2, v_{ij}^1, Z_k | i = 1, \dots, n, \quad q = 1, \dots, \mu, \quad j = 1, \dots, \nu, \quad k = 1, \dots, r + s\}$$

is an integral basis of $\mathfrak{n}_{r,s}(\nu, \mu)$. Recall that in both Lie algebras $\mathfrak{n}_{r,s}(\mu, \nu)$ and $\mathfrak{n}_{r,s}(\nu, \mu)$ the following relations hold: $[v_{ij}^1, v_{pq}^1] = [v_{ij}^1, v_{pr}^2] = [v_{ij}^2, v_{pq}^2] = 0$ for $j \neq q$ and for any

i, p, r . Moreover

$$[v_{ij}^1, v_{pj}^1] = [v_{i1}^1, v_{p1}^1] = -\sum_{k=1}^{r+s} A_{ip}^k Z_k, \quad [v_{ij}^2, v_{pj}^2] = [v_{i1}^2, v_{p1}^2] = \sum_{k=1}^{r+s} A_{ip}^k Z_k \quad (37)$$

for the above chosen $C: \mathfrak{z}_{r,s} \rightarrow \mathfrak{z}_{r,s}$. The bijective linear map $f: \mathfrak{n}_{r,s}(\mu, \nu) \rightarrow \mathfrak{n}_{r,s}(\nu, \mu)$ defined by

$$\begin{aligned} v_{ij}^1 &\mapsto v_{ij}^2 = A(v_{ij}^1), & \text{for } j = 1, \dots, \mu, \\ v_{ip}^2 &\mapsto v_{ip}^1 = A^{-1}(v_{ip}^2), & \text{for } p = 1, \dots, \nu, \\ Z_k &\mapsto -Z_k, & \text{for } k = 1, \dots, r+s, \end{aligned}$$

and $i, p = 1, \dots, n$ induces a Lie algebra homomorphism. Indeed, by (37)

$$\begin{aligned} f([v_{ij}^1, v_{pj}^1]) &= f([v_{i1}^1, v_{p1}^1]) = f\left(-\sum_{k=1}^{r+s} A_{ip}^k Z_k\right) = -\sum_{k=1}^{r+s} A_{ip}^k f(Z_k) \\ &= \sum_{k=1}^{r+s} A_{ip}^k Z_k = [v_{ij}^2, v_{pj}^2] = [A(v_{ij}^1), A(v_{pj}^1)] = [f(v_{ij}^1), f(v_{pj}^1)]. \end{aligned}$$

and, analogously,

$$\begin{aligned} f([v_{ij}^2, v_{pj}^2]) &= f([v_{i1}^2, v_{p1}^2]) = f\left(\sum_{k=1}^{r+s} A_{ip}^k Z_k\right) = \sum_{k=1}^{r+s} A_{ip}^k f(Z_k) \\ &= -\sum_{k=1}^{r+s} A_{ip}^k Z_k = [v_{ij}^1, v_{pj}^1] = [A^{-1}(v_{ij}^2), A^{-1}(v_{pj}^2)] = [f(v_{ij}^2), f(v_{pj}^2)]. \end{aligned}$$

To show the reverse statement we assume that Lie algebras $\mathfrak{n}_{r,s}(\mu_1, \nu_1)$ and $\mathfrak{n}_{r,s}(\mu_2, \nu_2)$ are isomorphic for some $\mu_1 > \mu_2$ and $\nu_1 > \nu_2$. Then there are bijective maps $A_{12}: \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^2$ and $A_{11}: \mathfrak{v}_{r,s}^1 \rightarrow \mathfrak{v}_{r,s}^1$ of minimal dimensional modules where the map A_{12} induces C by (36) and C induces A_{11} by (10). Then we obtain $J_{C(Z)}^1 = A_{12}^\top \circ J_Z^2 \circ A_{12} = A_{11}^\top \circ J_Z^1 \circ A_{11}$, that contradicts to the assumption that modules $\mathfrak{v}_{r,s}^1$ and $\mathfrak{v}_{r,s}^2$ are non equivalent. \square

8.1. Open problems on classification of H -type Lie algebras $\mathfrak{n}_{r,s}(\mu, \nu)$. The problem of the isomorphism of the pseudo H -type Lie algebras $\mathfrak{n}_{r,s}(\mu, \nu)$ with different signatures (r, s) turns out to be not so trivial. The increasing of dimension of admissible modules allows more freedom for action of the representation maps and some isomorphic Lie algebras can appear. For instance, it is possible to show, in the above notation, that the Lie algebras $\mathfrak{n}_{2,1}$ and $\mathfrak{n}_{1,2}(1, 1)$ are isomorphic, but the Lie algebras $\mathfrak{n}_{2,1}$ and $\mathfrak{n}_{1,2}(2, 0)$ are not isomorphic. Thus we leave the full description of classification of pseudo H -type Lie algebras for forthcoming paper.

8.2. Bracket generating properties.

Theorem 8.3. *The pseudo H -type algebras $\mathfrak{n}_{r,s}(\mu, \nu)$ possesses the strongly bracket generating property if only if $r = 0$ or $s = 0$.*

Proof. First we prove that $\mathfrak{n}_{r,0}(\mu, \nu)$ is strongly bracket generating, i.e. $[w, \mathfrak{v}_{r,0}(\mu, \nu)] = \mathfrak{z}_{r,0}$ for all $w \in \mathfrak{n}_{r,0}(\mu, \nu)$, $w \neq 0$. We recall that $\mathfrak{v}_{r,0}(\mu, \nu) = \bigoplus_{j=1}^{\mu} (\mathfrak{v}_{r,0}^1)_j \oplus \bigoplus_{j=1}^{\nu} (\mathfrak{v}_{r,0}^2)_j$.

TABLE 11. Commutation relations on $\mathfrak{n}_{0,8}$

$[r, c]$	v_1	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_{10}	v_{11}	v_{12}	v_{13}	v_{14}	v_{15}	v_{16}
v_1	0	0	0	0	0	0	0	0	Z_1	Z_2	Z_3	Z_4	Z_5	Z_6	Z_7	Z_8
v_2	0	0	0	0	0	0	0	0	$-Z_2$	Z_1	Z_4	$-Z_3$	Z_6	$-Z_5$	Z_8	$-Z_7$
v_3	0	0	0	0	0	0	0	0	$-Z_3$	$-Z_4$	Z_1	Z_2	$-Z_8$	$-Z_7$	Z_6	Z_5
v_4	0	0	0	0	0	0	0	0	$-Z_4$	Z_3	$-Z_2$	Z_1	$-Z_7$	Z_8	Z_5	$-Z_6$
v_5	0	0	0	0	0	0	0	0	$-Z_5$	$-Z_6$	Z_8	Z_7	Z_1	Z_2	$-Z_4$	$-Z_3$
v_6	0	0	0	0	0	0	0	0	$-Z_6$	Z_5	Z_7	$-Z_8$	$-Z_2$	Z_1	$-Z_3$	Z_4
v_7	0	0	0	0	0	0	0	0	$-Z_7$	$-Z_8$	$-Z_6$	$-Z_5$	Z_4	Z_3	Z_1	Z_2
v_8	0	0	0	0	0	0	0	0	$-Z_8$	Z_7	$-Z_5$	Z_6	Z_3	$-Z_4$	$-Z_2$	Z_1
v_9	$-Z_1$	Z_2	Z_3	Z_4	Z_5	Z_6	Z_7	Z_8	0	0	0	0	0	0	0	0
v_{10}	$-Z_2$	$-Z_1$	Z_4	$-Z_3$	Z_6	$-Z_5$	Z_8	$-Z_7$	0	0	0	0	0	0	0	0
v_{11}	$-Z_3$	$-Z_4$	$-Z_1$	Z_2	$-Z_8$	$-Z_7$	Z_6	Z_5	0	0	0	0	0	0	0	0
v_{12}	$-Z_4$	Z_3	$-Z_2$	$-Z_1$	$-Z_7$	Z_8	Z_5	$-Z_6$	0	0	0	0	0	0	0	0
v_{13}	$-Z_5$	$-Z_6$	Z_8	Z_7	$-Z_1$	Z_2	$-Z_4$	$-Z_3$	0	0	0	0	0	0	0	0
v_{14}	$-Z_6$	Z_5	Z_7	$-Z_8$	$-Z_2$	$-Z_1$	$-Z_3$	Z_4	0	0	0	0	0	0	0	0
v_{15}	$-Z_7$	$-Z_8$	$-Z_6$	$-Z_5$	Z_4	Z_3	$-Z_1$	Z_2	0	0	0	0	0	0	0	0
v_{16}	$-Z_8$	Z_7	$-Z_5$	Z_6	Z_3	$-Z_4$	$-Z_2$	$-Z_1$	0	0	0	0	0	0	0	0

TABLE 12. Commutation relations on $\mathfrak{n}_{4,4}$

$[r, c]$	y_1	y_6	y_7	y_8	y_{13}	y_{14}	y_{15}	y_{16}	y_2	y_3	y_4	y_5	y_9	y_{10}	y_{11}	y_{12}
y_1	0	0	0	0	0	0	0	0	Z_1	Z_2	Z_3	Z_4	Z_5	Z_6	Z_7	Z_8
y_6	0	0	0	0	0	0	0	0	Z_2	$-Z_1$	$-Z_4$	Z_3	Z_6	$-Z_5$	Z_8	$-Z_7$
y_7	0	0	0	0	0	0	0	0	Z_3	Z_4	$-Z_1$	$-Z_2$	Z_8	Z_7	$-Z_6$	$-Z_5$
y_8	0	0	0	0	0	0	0	0	Z_4	$-Z_3$	Z_2	$-Z_1$	$-Z_7$	Z_8	Z_5	$-Z_6$
y_{13}	0	0	0	0	0	0	0	0	Z_5	$-Z_6$	$-Z_8$	Z_7	Z_1	$-Z_2$	Z_4	$-Z_3$
y_{14}	0	0	0	0	0	0	0	0	Z_6	Z_5	$-Z_7$	$-Z_8$	Z_2	Z_1	$-Z_3$	$-Z_4$
y_{15}	0	0	0	0	0	0	0	0	Z_7	$-Z_8$	Z_6	$-Z_5$	$-Z_4$	Z_3	Z_1	$-Z_2$
y_{16}	0	0	0	0	0	0	0	0	Z_8	Z_7	Z_5	Z_6	Z_3	Z_4	Z_2	Z_1
y_2	$-Z_1$	$-Z_2$	$-Z_3$	$-Z_4$	$-Z_5$	$-Z_6$	$-Z_7$	$-Z_8$	0	0	0	0	0	0	0	0
y_3	$-Z_2$	Z_1	$-Z_4$	Z_3	Z_6	$-Z_5$	Z_8	$-Z_7$	0	0	0	0	0	0	0	0
y_4	$-Z_3$	Z_4	Z_1	$-Z_2$	Z_8	Z_7	$-Z_6$	$-Z_5$	0	0	0	0	0	0	0	0
y_5	$-Z_4$	$-Z_3$	Z_2	Z_1	$-Z_7$	Z_8	Z_5	$-Z_6$	0	0	0	0	0	0	0	0
y_9	$-Z_5$	$-Z_6$	$-Z_8$	Z_7	$-Z_1$	$-Z_2$	Z_4	$-Z_3$	0	0	0	0	0	0	0	0
y_{10}	$-Z_6$	Z_5	$-Z_7$	$-Z_8$	Z_2	$-Z_1$	$-Z_3$	$-Z_4$	0	0	0	0	0	0	0	0
y_{11}	$-Z_7$	$-Z_8$	Z_6	$-Z_5$	$-Z_4$	Z_3	$-Z_1$	$-Z_2$	0	0	0	0	0	0	0	0
y_{12}	$-Z_8$	Z_7	Z_5	Z_6	Z_3	Z_4	Z_2	$-Z_1$	0	0	0	0	0	0	0	0

REFERENCES

- [1] P. Ciatti, *Scalar products on Clifford modules and pseudo-H-type Lie algebras*. Ann. Mat. Pura Appl. **178** (2000), no. 4, p. 1–32.
- [2] K. Furutani, I. Markina, *Existence of the lattice on general H-type groups*. J. Lie Theory, **24**, (2014), 979–1011.
- [3] M. Godoy Molina, A. Korolko, I. Markina, *Sub-semi-Riemannian geometry of general H-type groups*. Bull. Sci. Math. **137** (2013), no. 6, 805–833.
- [4] D. Husemoller, *Fibre bundles*. Second edition. Graduate Texts in Mathematics, No. 20. Springer-Verlag, New York-Heidelberg, 1975. 327 pp.
- [5] A. Kaplan, *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*. Trans. Amer. Math. Soc. **258** (1980), no. 1, p. 147–153.
- [6] A. Kaplan, *Riemannian nilmanifolds attached to Clifford modules*. Geom. Dedicata **11** (1981), no. 2, p. 127–136.
- [7] A. Kaplan, *On the geometry of groups of Heisenberg type*. Bull. London Math. Soc. **15** (1983), no. 1, p. 35–42.

TABLE 13. Permutations of the basis of $\mathfrak{n}_{8,0}$ by J_i

$J_i u_j$	J_1	J_2	J_3	J_4	J_5	J_6	J_7	J_8
u_1	u_9	u_{10}	u_{11}	u_{12}	u_{13}	u_{14}	u_{15}	u_{16}
u_2	$-u_{10}$	u_9	u_{12}	$-u_{11}$	u_{14}	$-u_{13}$	u_{16}	$-u_{15}$
u_3	$-u_{11}$	$-u_{12}$	u_9	u_{10}	$-u_{16}$	$-u_{15}$	u_{14}	u_{13}
u_4	$-u_{12}$	u_{11}	$-u_{10}$	u_9	$-u_{15}$	u_{16}	u_{13}	$-u_{14}$
u_5	$-u_{13}$	$-u_{14}$	u_{16}	u_{15}	u_9	u_{10}	$-u_{12}$	$-u_{11}$
u_6	$-u_{14}$	u_{13}	u_{15}	$-u_{16}$	$-u_{10}$	u_9	$-u_{11}$	u_{12}
u_7	$-u_{15}$	$-u_{16}$	$-u_{14}$	$-u_{13}$	u_{12}	u_{11}	u_9	u_{10}
u_8	$-u_{16}$	u_{15}	$-u_{13}$	u_{14}	u_{11}	$-u_{12}$	$-u_{10}$	u_9
u_9	$-u_1$	$-u_2$	$-u_3$	$-u_4$	$-u_5$	$-u_6$	$-u_7$	$-u_8$
u_{10}	u_2	$-u_1$	u_4	$-u_3$	u_6	$-u_5$	u_8	$-u_7$
u_{11}	u_3	$-u_4$	$-u_1$	u_2	$-u_8$	$-u_7$	u_6	u_5
u_{12}	u_4	u_3	$-u_2$	$-u_1$	$-u_7$	u_8	u_5	$-u_6$
u_{13}	u_5	$-u_6$	u_8	u_7	$-u_1$	u_2	$-u_4$	$-u_3$
u_{14}	u_6	u_5	u_7	$-u_8$	$-u_2$	$-u_1$	$-u_3$	u_4
u_{15}	u_7	$-u_8$	$-u_6$	$-u_5$	u_4	u_3	$-u_1$	u_2
u_{16}	u_8	u_7	$-u_5$	u_6	u_3	$-u_4$	$-u_2$	$-u_1$

- [8] T. Y. Lam, *The algebraic theory of quadratic forms*. Mathematics Lecture Note Series. W. A. Benjamin, Inc., Reading, Mass., 1973, pp. 344.
- [9] H. B. Lawson, M.-L. Michelsohn, *Spin geometry*. Princeton Mathematical Series, 38. Princeton University Press, Princeton, NJ, 1989. 427 pp.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, NORWAY.
E-mail address: christian.autenried@math.uib.no

K. FURUTANI: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, SCIENCE UNIVERSITY OF TOKYO, 2641 YAMAZAKI, NODA, CHIBA (278-8510), JAPAN
E-mail address: furutani_kenro@ma.noda.tus.ac.jp

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, NORWAY.
E-mail address: irina.markina@math.uib.no