Dynamic Programming Principle for Stochastic Recursive Optimal Control Problem under G-framework

Mingshang Hu * Shaolin Ji[†]

Abstract. In this paper, we study a stochastic recursive optimal control problem in which the cost functional is described by the solution of a backward stochastic differential equation driven by G-Brownian motion. Under standard assumptions, we establish the dynamic programming principle and the related fully nonlinear HJB equation in the framework of G-expectation. Finally, we show that the value function is the viscosity solution of the obtained HJB equation.

Key words. *G*-expectation, backward stochastic differential equations, stochastic recursive optimal control, robust control, dynamic programming principle

AMS subject classifications. 93E20, 60H10, 35K15

1 Introduction

It is well known that Duffie and Epstein [7] introduced a stochastic differential recursive utility which corresponds to the solution of a particular backward stochastic differential equation (BSDE). Thus the BSDE point of view gives a simple formulation of recursive utilities (see [8]). Since then, the classical stochastic optimal control problem is generalized to a so called "stochastic recursive optimal control problem" in which the cost functional is defined by the solution of BSDE. The stochastic maximum principle and dynamic programming principle for this problem were first established in Peng [18] and [23] respectively.

 $^{^{*}}$ Qilu Institute of Finance, Shandong University, Jinan, Shandong 250100, PR China. humingshang@sdu.edu.cn. Research supported by NSF (No. 11201262, 11101242 and 11301068) and Shandong Province (No. BS2013SF020)

[†]Qilu Institute of Finance, Shandong University, Jinan, Shandong 250100, PR China. jsl@sdu.edu.cn (Corresponding author). Research supported by NSF (No. 11171187, 11222110 and 11221061), Shandong Province (No.JQ201202), Programme of Introducing Talents of Discipline to Universities of China (No.B12023) and Program for New Century Excellent Talents in University of China. Hu and Ji's research was partially supported by NSF (No. 10921101) and by the 111 Project (No. B12023)

Recently Hu et. al studied a new kind of BSDE which is driven by G-Brownian motion in [13] and [12]:

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds + \int_t^T g(s, Y_s, Z_s) d\langle B \rangle_s$$

$$- \int_t^T Z_s dB_s - (K_T - K_t).$$
(1)

They proved that there exists a unique triple of processes (Y, Z, K) which solves (1) under the standard Lipschitz conditions. This new kind of BSDE is based on the *G*-expectation theory which is introduced by Peng (see [19], [22] and the references therein). This *G*-expectation framework (*G*-framework for short) does not require the probability space and is convenient to study financial problems involving volatility uncertainty. Let us mention that there are other recent advances in this direction. Denis, Martini [5] and Denis, Hu, Peng [6] developed quasi-sure stochastic analysis. Soner et al. [26] have obtained a existence and uniqueness theorem for a new type of fully nonlinear BSDE, called 2BSDE.

An important property of the solution Y of (1) is that it can be represented as the "supremum of expectations" over a set of nondominated probability measures. For example, the solution Y of (1) at time 0 can be written as

$$Y_{0} = \hat{\mathbb{E}}[\xi + \int_{0}^{T} f(s, Y_{s}, Z_{s})ds + \int_{0}^{T} g(s, Y_{s}, Z_{s})d\langle B \rangle_{s}]$$
(2)
$$= \sup_{P \in \mathcal{P}} E_{P}[\xi + \int_{0}^{T} f(s, Y_{s}, Z_{s})ds + \int_{0}^{T} g(s, Y_{s}, Z_{s})d\langle B \rangle_{s}]$$

where \mathcal{P} is a family of weakly compact nondominant probability measures. Then, (1) can be used to define recursive utility under volatility uncertainty. It is worth to point out that the recursive utility under mean uncertainty was developed in Chen and Epstein [3]. Epstein and Ji [9, 10] introduced a particular recursive utility under both mean and volatility uncertainty.

Motivated by the recursive utility optimization under volatility uncertainty, we explore a stochastic recursive optimal control problem in which the cost functional is defined by the solution of the above new type of BSDE. In more details, the state equation is governed by the following controlled SDE driven by G-Brownian motion

$$dX_{s}^{t,x,u} = b(s, X_{s}^{t,x,u}, u_{s})ds + h_{ij}(s, X_{s}^{t,x,u}, u_{s})d\langle B^{i}, B^{j}\rangle_{s} + \sigma(s, X_{s}^{t,x,u}, u_{s})dB_{s},$$

$$X_{*}^{t,x,u} = x.$$

The cost functional is introduced by the solution $Y_t^{t,x,u}$ of the following BSDE driven by *G*-Brownian motion at time *t*:

$$\begin{array}{ll} -dY_{s}^{t,x,u} = & f(s,X_{s}^{t,x,u},Y_{s}^{t,x,u},Z_{s}^{t,x,u},u_{s})ds + g_{ij}(s,X_{s}^{t,x,u},Y_{s}^{t,x,u},Z_{s}^{t,x,u},u_{s})d\langle B^{i},B^{j}\rangle_{s} \\ & -Z_{s}^{t,x,u}dB_{s} - dK_{s}^{t,x,u}, \\ Y_{T}^{t,x,u} = & \Phi(X_{T}^{t,x,u}), \quad s \in [t,T]. \end{array}$$

We define the value function of our stochastic recursive optimal control problem as follows:

$$V(t,x) = \operatorname{ess\,inf}_{u(\cdot) \in \mathcal{U}[t,T]} Y_t^{t,x,u},$$

where the control set is in the *G*-framework. In view of (2), we essentially have to solve a "inf sup problem". Such problem is known as the robust optimal control problem, i.e., we consider the worst scenario by maximizing over a set of probability measures and then we minimize the cost functional. For recent development of robust utility maximization under volatility uncertainty, we refer the interested readers to [27], [17] and [4]. Tevzadze, Toronjadze, Uzunashvili [27] studied robust exponential and power utilities. Matoussi, Possamai, Zhou [17] related robust utility maximization problem to a particular 2BSDE with quadratic growth. In [4], Denis and Kervarec established a duality theory for this problem in nondominated models.

The objective of our paper is to establish the dynamic programming principle (DPP) for this stochastic recursive optimal control problem and investigate the value function in *G*-framework.

It is well known that DPP and related HJB equations is a powerful approach to solving optimal control problems (see [11], [28] and [23]). For the classical stochastic recursive optimal control problem, Peng [23] obtained the Hamilton– Jacobi–Bellman equation and proved that the value function is its viscosity solution. In [24], Peng generalized his results and originally introduced the notion of stochastic backward semigroups which allows him to prove DPP in a very straightforward way. This backward semigroup approach is also introduced in the theory of stochastic differential games by Buckdahn and Li in [1]. Note that Buckdahn et al. [2] obtained an existence result of the stochastic recursive optimal control problem.

In this paper, we adopt the backward semigroup approach to build the DPP in our context. At first, we need to define the essential infimum of a family of random variables in the "quasi-surely" sense (q.s. for short). Compared with classical case in [24], this kind of essential infimum may not exist in our case (the q.s. case). We define the essential infimum and prove its existence in this paper. Under a family of non-dominated probability measures, it is far from being trivial to prove that the value function V is wellposed and deterministic. Due to a new result in [16], we construct the approximation of an element of the admissible control set which is the key step to prove that $\underset{u(\cdot) \in \mathcal{U}[t,T]}{\operatorname{essinf}} Y_t^{t,x,u}$ is

a deterministic function. At last, we adopt an "implied partition" approach to prove DPP (see Lemma 22) which is completely new in the literature.

We states that V is deterministic continuous viscosity solution of the following fully nonlinear HJB equation

$$\partial_t V(t,x) + \inf_{u \in U} H(t,x,V,\partial_x V,\partial_{xx}^2 V,u) = 0,$$

$$V(T,x) = \Phi(x), \qquad x \in \mathbb{R}^n,$$

where

$$\begin{array}{ll} H(t,x,v,p,A,u) = & G(F(t,x,v,p,A,u)) + \langle p,b(t,x,u) \rangle + f(t,x,v,\sigma^{T}(t,x,u)p,u), \\ F_{ij}(t,x,v,p,A,u) = & (\sigma^{T}(t,x,u)A\sigma(t,x,u))_{ij} + 2\langle p,h_{ij}(t,x,u) \rangle \\ & + 2g_{ij}(t,x,v,\sigma^{T}(t,x,u)p,u), \end{array}$$

 $(t, x, v, p, A, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{S}_n \times U$. The main difficulty to prove this statement lies in the appearance of two decreasing *G*-martingale terms. Applying a property of decreasing *G*-martingale proved in Lemma 30, we overcome this difficulty (see Lemma 29) and obtain the result.

In conclusion, since there is no reference probability measure under the Gframework, our results generalize the results in Peng [23] and [24] which was only considered in the Wiener space (corresponding to G is linear in our paper). Compared with our earlier article [15], the problem in [15] is essentially a "sup sup problem" which is easier to deal with. And the techniques developped in this paper can also used to solve the problem in [15]. Note that G has the representation (4) which leads to that the above HJB equation can also be understood as a kind of Bellman-Issac equation. Then, it is meaningful to show the difference between our paper and some related references (see [1] and [25]) in game theory. Needless to say, the game problem is more complicated than the robust control problem since it needs to study the value of game. Buckdahn, Li [1] employed strategies and Pham, Zhang [25] formulated their game problem in a weak framework. In construct, we use controls and our formulation is a "strong" framework under the G-framework. Different from [25], as revealed in [16], our admissible control set has quasi-continuous property and in particular, it does not change with time. It is worth mentioning that, in our context, the coefficients of the state equation include the state variable X.

The paper is organized as follows. In section 2, we present some fundamental results on G-expectation theory. We formulate our stochastic recursive optimal control problem in section 3. We prove the properties of the value function in section 4 and establish the dynamic programming principle in section 5. In section 6, we first derive the fully nonlinear HJB equation and prove that the value function is the viscosity solution of the obtained HJB equation.

2 Preliminaries

We review some basic notions and results of G-expectation and the related spaces of random variables. The readers may refer to [19], [20], [21], [22] for more details.

Let $\Omega_T = C_0([0,T]; \mathbb{R}^d)$, the space of R^d -valued continuous functions on [0,T] with $\omega_0 = 0$, and $B_t(\omega) = \omega_t$ be the canonical process. Set

$$L_{ip}(\Omega_T) := \{ \varphi(B_{t_1}, ..., B_{t_n}) : n \ge 1, t_1, ..., t_n \in [0, T], \varphi \in C_{b.Lip}(\mathbb{R}^{d \times n}) \},\$$

where $C_{b.Lip}(\mathbb{R}^{d \times n})$ denotes the set of bounded Lipschitz functions on $\mathbb{R}^{d \times n}$.

We denote the *G*-expectation space by $(\Omega_T, L_{ip}(\Omega_T), \hat{\mathbb{E}})$. The function $G : \mathbb{S}_d \to \mathbb{R}$ is defined by

$$G(A) := \frac{1}{2} \hat{\mathbb{E}}[\langle AB_1, B_1 \rangle].$$
(3)

where \mathbb{S}_d denotes the collection of $d \times d$ symmetric matrices. Note that there exists a bounded and closed subset $\Gamma \subset \mathbb{R}^{d \times d}$ such that

$$G(A) = \frac{1}{2} \sup_{Q \in \Gamma} tr[AQQ^T].$$
(4)

In this paper, we only consider non-degenerate *G*-normal distribution, i.e., there exists some $\underline{\sigma}^2 > 0$ such that $G(A) - G(B) \geq \frac{1}{2}\underline{\sigma}^2 \operatorname{tr}[A - B]$ for any $A \geq B$. We denote by $L^p_G(\Omega_T)$ the completion of $L_{ip}(\Omega_T)$ under the norm $\|X\|_{p,G} = \frac{1}{2} \frac{1}$

We denote by $L_G^p(\Omega_T)$ the completion of $L_{ip}(\Omega_T)$ under the norm $||X||_{p,G} = (\hat{\mathbb{E}}[|X|^p])^{1/p}$ for $p \ge 1$. For each $t \ge 0$, the conditional *G*-expectation $\hat{\mathbb{E}}_t[\cdot]$ can be extended continuously to $L_G^1(\Omega_T)$ under the norm $||\cdot||_{1,G}$.

Definition 1 Let $M_G^0(0,T)$ be the collection of processes in the following form: for a given partition $\{t_0, \dots, t_N\} = \pi_T$ of [0,T],

$$\eta_t(\omega) = \sum_{j=0}^{N-1} \xi_j(\omega) I_{[t_j, t_{j+1})}(t),$$

where $\xi_i \in L_{ip}(\Omega_{t_i}), i = 0, 1, 2, \dots, N - 1.$

We denote by $M_G^p(0,T)$ the completion of $M_G^0(0,T)$ under the norm $\|\eta\|_{M_G^p} = \{\hat{\mathbb{E}}[\int_0^T |\eta_s|^p ds]\}^{1/p}$ for $p \ge 1$.

Theorem 2 ([6, 14]) There exists a family of weakly compact probability measures \mathcal{P} on $(\Omega, \mathcal{B}(\Omega))$ such that

$$\hat{\mathbb{E}}[\xi] = \sup_{P \in \mathcal{P}} E_P[\xi] \quad for \quad all \ \xi \in L^1_G(\Omega).$$

 \mathcal{P} is called a set that represents $\mathbb{\hat{E}}$.

For this \mathcal{P} , we define capacity

$$c(A) := \sup_{P \in \mathcal{P}} P(A), \ A \in \mathcal{B}(\Omega_T)$$

A set $A \subset \Omega_T$ is polar if c(A) = 0. A property holds "quasi-surely" (q.s. for short) if it holds outside a polar set. In the following, we do not distinguish two random variables X and Y if X = Y q.s.. We set

$$\mathbb{L}^{p}(\Omega_{T}) := \{ X \in \mathcal{B}(\Omega_{T}) : \sup_{P \in \mathcal{P}} E_{P}[|X|^{p}] < \infty \} \text{ for } p \ge 1.$$

It is important to note that $L^p_G(\Omega_T) \subset \mathbb{L}^p(\Omega_T)$. We extend *G*-expectation \mathbb{E} to $\mathbb{L}^p(\Omega_T)$ and still denote it by $\mathbb{\hat{E}}$, for each $X \in \mathbb{L}^1(\Omega_T)$, we set

$$\mathbb{\hat{E}}[X] = \sup_{P \in \mathcal{P}} E_P[X].$$

For $p \ge 1$, $\mathbb{L}^p(\Omega_T)$ is a Banach space under the norm $(\hat{\mathbb{E}}[|\cdot|^p])^{1/p}$.

Furthermore, we extend the definition of conditional G-expectation. For each fixed $t \in [0,T]$, let $(A_i)_{i=1}^n$ be a partition of $\mathcal{B}(\Omega_t)$, and set

$$\xi = \sum_{i=1}^{n} \eta_i I_{A_i},$$

where $\eta_i \in L^1_G(\Omega_T)$, $i = 1, \dots, n$. We define the corresponding generalized conditional G-expectation, still denoted by $\hat{\mathbb{E}}_s[\cdot]$, by setting

$$\hat{\mathbb{E}}_s\left[\sum_{i=1}^n \eta_i I_{A_i}\right] := \sum_{i=1}^n \hat{\mathbb{E}}_s[\eta_i] I_{A_i} \quad \text{for } s \in [t, T].$$

Then, many properties of the conditional G-expectation still hold (refer to Proposition 2.5 in [12]).

Problem 3

3.1State equations

We first give the definition of admissible controls.

Definition 3 For each $t \in [0, T]$, u is said to be an admissible control on [t, T], if it satisfies the following conditions:

(i) $u: [t,T] \times \Omega \to U$ where U is a given compact set of \mathbb{R}^m ; (ii) $u \in M^{\frac{1}{2}}_{G}(t,T;\mathbb{R}^{m}).$

The set of admissible controls on [t, T] is denoted by $\mathcal{U}[t, T]$. In the rest of this paper, we use Einstein summation convention. Let $t \in [0,T], \xi \in \bigcup_{\varepsilon > 0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$ and $u \in \mathcal{U}[t,T]$. Consider the following

forward and backward SDEs driven by G-Brownian motion:

$$dX_{s}^{t,\xi,u} = b(s, X_{s}^{t,\xi,u}, u_{s})ds + h_{ij}(s, X_{s}^{t,\xi,u}, u_{s})d\langle B^{i}, B^{j}\rangle_{s} + \sigma(s, X_{s}^{t,\xi,u}, u_{s})dB_{s},$$
(5)

 $X_t^{t,\xi,u} = \xi,$

and

$$\begin{aligned} -dY_{s}^{t,\xi,u} &= f(s, X_{s}^{t,\xi,u}, Y_{s}^{t,\xi,u}, Z_{s}^{t,\xi,u}, u_{s})ds + g_{ij}(s, X_{s}^{t,\xi,u}, Y_{s}^{t,\xi,u}, Z_{s}^{t,\xi,u}, u_{s})d\langle B^{i}, B^{j}\rangle_{s} \\ &\quad (6) \\ &\quad -Z_{s}^{t,\xi,u}dB_{s} - dK_{s}^{t,\xi,u}, \\ Y_{T}^{t,\xi,u} &= \Phi(X_{T}^{t,\xi,u}), \quad s \in [t,T]. \\ \text{Set} \\ Set \\ S_{G}^{0}(0,T) &:= \{h(t, B_{t_{1}\wedge t}, \cdots, B_{t_{n}\wedge t}) : t_{1}, \dots, t_{n} \in [0,T], h \in C_{b,Lip}(\mathbb{R}^{n+1})\}. \end{aligned}$$

For $p \ge 1$ and $\eta \in S^0_G(0,T)$, let $\|\eta\|_{S^p_G} = \{\hat{\mathbb{E}}[\sup_{t \in [0,T]} |\eta_t|^p]\}^{\frac{1}{p}}$. Denote by $S^p_G(0,T)$ the completion of $S_G^0(0,T)$ under the norm $\|\cdot\|_{S_G^p}$.

For given t, u and ξ , $(X^{t,\xi,u})$ and $(Y^{t,\xi,u}, Z^{t,\xi,u}, K^{t,\xi,u})$ are called solutions of the above forward and backward SDEs respectively if

- $\begin{array}{l} \text{(i)} \ (X^{t,\xi,u}) \in M^2_G(t,T;\mathbb{R}^n);\\ \text{(ii)} \ (Y^{t,\xi,u},Z^{t,\xi,u}) \in S^2_G(0,T) \times M^2_G(0,T;\mathbb{R}^d); \end{array}$
- (iii) $K^{t,\xi,u}$ is a decreasing *G*-martingale with $K^{t,\xi,u}_t = 0, K^{t,\xi,u}_T \in L^2_G(\Omega_T)$;
- (iv) (5) and (6) are satisfied respectively.

We assume that $b, h_{ij} : [0,T] \times \mathbb{R}^n \times U \to \mathbb{R}^n, \sigma : [0,T] \times \mathbb{R}^n \times U \to \mathbb{R}^{n \times d},$ $\Phi : \mathbb{R}^n \to \mathbb{R}, f, g_{ij} : [0,T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times U \to \mathbb{R}$ are deterministic functions and satisfy the following conditions:

(A1) There exists a constant C > 0 such that $\forall (s, x, y, z, u), (s, x', y', z', v)$ $\in [0,T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^d \times U,$

$$\begin{aligned} |b(s, x, u) - b(s, x', v)| + |h_{ij}(s, x, u) - h_{ij}(s, x', v)| + |\sigma(s, x, u) - \sigma(s, x', v)| \\ &\leq C(|x - x'| + |u - v|), \end{aligned}$$

$$|\Phi(x) - \Phi(x')| \le C|x - x'|,$$

$$|f(s, x, y, z, u) - f(s, x', y', z', v)| + |g_{ij}(s, x, y, z, u) - g_{ij}(s, x', y', z', v)| \le C(|x - x'| + |y - y'| + |z - z'| + |u - v|);$$

(A2) $b, h_{ij}, \sigma, f, g_{ij}$ are continuous in s.

Then, we have the following theorems.

Theorem 4 ([22]) Let Assumptions (A1) and (A2) hold. Then there exists a unique adapted solution X for equation (5).

Theorem 5 ([22])Let $\xi, \xi' \in L^p_G(\Omega_t; \mathbb{R}^n)$ with $p \ge 2$ and $u, v \in \mathcal{U}[t, T]$. Then we have, for each $\delta \in [0, T - t]$,

$$\begin{split} \hat{\mathbb{E}}_t[|X_{t+\delta}^{t,\xi,u} - X_{t+\delta}^{t,\xi',v}|^2] &\leq \bar{C}(|\xi - \xi'|^2 + \hat{\mathbb{E}}_t[\int_t^{t+\delta} |u_s - v_s|^2 ds]), \\ \hat{\mathbb{E}}_t[|X_{t+\delta}^{t,\xi,u}|^p] &\leq \bar{C}(1 + |\xi|^p), \\ \hat{\mathbb{E}}_t[\sup_{s \in [t,t+\delta]} |X_s^{t,\xi,u} - \xi|^p] &\leq \bar{C}(1 + |\xi|^p)\delta^{p/2}, \end{split}$$

where the constant \overline{C} depends on C, G, p, n, U and T.

Theorem 6 ([12]) Let Assumptions (A1) and (A2) hold. Then there exists a unique adapted solution (Y, Z, K) for equation (6).

Theorem 7 ([12])Let $\xi, \xi' \in \bigcup_{\varepsilon > 0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$ and $u, v \in \mathcal{U}[t, T]$. Then there exist two positive constants \overline{C}_1 and \overline{C}_2 depending on C, G and T such that

$$|Y_t^{t,\xi,u} - Y_t^{t,\xi',v}|^2 \le \bar{C}_1 \hat{\mathbb{E}}_t [|\Phi(X_T^{t,\xi,u}) - \Phi(X_T^{t,\xi',v})|^2 + (\int_t^T \hat{F}_s ds)^2] \\\le \bar{C}_2 \hat{\mathbb{E}}_t [|\Phi(X_T^{t,\xi,u}) - \Phi(X_T^{t,\xi',v})|^2 + \int_t^T |\hat{F}_s|^2 ds],$$

where

$$\begin{split} \hat{F}_s &= |f(s, X_s^{t,\xi,u}, Y_s^{t,\xi,u}, Z_s^{t,\xi,u}, u_s) - f(s, X_s^{t,\xi',v}, Y_s^{t,\xi,u}, Z_s^{t,\xi,u}, v_s)| \\ &+ \sum_{i,j=1}^d |g_{ij}(s, X_s^{t,\xi,u}, Y_s^{t,\xi,u}, Z_s^{t,\xi,u}, u_s) - g_{ij}(s, X_s^{t,\xi',v}, Y_s^{t,\xi,u}, Z_s^{t,\xi,u}, v_s)|. \end{split}$$

Theorem 8 ([16])Let b, h_{ij} , σ be independent of u and satisfy (A1) and (A2). Assume further that there exist constants L > 0, $0 < \lambda < \Lambda$ such that $|b| \leq L$, $|h_{ij}| \leq L$, $\lambda \leq |\sigma_i| \leq \Lambda$ for $i \leq n$, where σ_i is the *i*-th row of σ . Then for each x, $a, a' \in \mathbb{R}^n$ with $a \leq a', s \geq t$, we have $I_{\{X_s^{t,x} \in [a,a')\}} \in L^2_G(\Omega_s)$. In particular, for each $c, c' \in \mathbb{R}^{d \times k}$, $c \leq c'$ and $t \leq s_1 \leq \cdots \leq s_k$, we have $I_{\{(B_{s_1}-B_t,\ldots,B_{s_k}-B_t)\in [c,c')\}} \in L^2_G(\Omega_{s_k})$.

Remark 9 If there exists a $t_0 < T$ such that b, h_{ij} , σ are continuous in s just on $[t_0, T]$, then the above theorem still holds by the proof in [16].

3.2 Stochastic optimal control problem

The state equation of our stochastic optimal control problem is governed by the above forward SDE (5) and the objective functional is introduced by the solution of the BSDE (6) at time t. Let ξ equals a constant $x \in \mathbb{R}^n$. When u changes, $Y_t^{t,x,u}$ (the solution $Y^{t,x,u}$ at time t) also changes. In order to study the value function of our stochastic optimal control problem, we need to define the essential infimum of $\{Y_t^{t,x,u} \mid u \in \mathcal{U}[t,T]\}$.

Definition 10 The essential infimum of $\{Y_t^{t,x,u} \mid u \in \mathcal{U}[t,T]\}$, denoted by essinf $Y_t^{t,x,u}$, is a random variable $\zeta \in L^2_G(\Omega_t)$ satisfying:

(i) $\forall u \in \mathcal{U}[t,T], \zeta \leq Y_t^{t,x,u} \quad q.s.;$

(ii) if η is a random variable satisfying $\eta \leq Y_t^{t,x,u}$ q.s. for any $u \in \mathcal{U}[t,T]$, then $\zeta \geq \eta$ q.s..

Similarly, we can define the essential infimum of $\{Y_t^{t,\xi,u} \mid u \in \mathcal{U}[t,T]\}$, where $\xi \in \bigcup_{\varepsilon > 0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$.

The following example shows that the essential infimum may be not exist.

Example 11 Let d = 1 and $(B_t)_{t\geq 0}$ be a 1-dimensional G-Brownian motion with $G(a) = \frac{1}{2}(a^+ - \frac{1}{3}a^-)$. We first show that $I_{\{\langle B \rangle_1 = \frac{1}{2}\}}$, $I_{\{\langle B \rangle_1 \geq \frac{1}{2}\}}$ and $I_{\{\langle B \rangle_1 > \frac{1}{2}\}} \notin L^2_G(\Omega_1)$.

It is easy to verify that $h_k(\langle B \rangle_1) \downarrow I_{\{\langle B \rangle_1 = \frac{1}{2}\}}$, where

$$h_k(x) = k(x - \frac{1}{2} + \frac{1}{k})I_{\left[\frac{1}{2} - \frac{1}{k}, \frac{1}{2}\right]}(x) + \left[1 - k(x - \frac{1}{2})\right]I_{\left(\frac{1}{2}, \frac{1}{2} + \frac{1}{k}\right]}(x).$$

If $I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \in L^2_G(\Omega_1)$, then

$$h_k(\langle B \rangle_1) - I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \in L^2_G(\Omega_1) \text{ and } h_k(\langle B \rangle_1) - I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \downarrow 0.$$

By Corollary 33 in [6], we have $\hat{\mathbb{E}}[h_k(\langle B \rangle_1) - I_{\{\langle B \rangle_1 = \frac{1}{2}\}}] \downarrow 0$. On the other hand,

$$\hat{\mathbb{E}}[h_k(\langle B \rangle_1) - I_{\{\langle B \rangle_1 = \frac{1}{2}\}}]$$

$$\geq \lim_{j \to \infty} \hat{\mathbb{E}}[h_k(\langle B \rangle_1) - h_j(\langle B \rangle_1)]$$

$$= \lim_{j \to \infty} \sup\{h_k(x) - h_j(x) : x \in [0, 1]\}$$

$$= 1.$$

Thus $I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \notin L^2_G(\Omega_1)$. Similarly, we can prove that $I_{\{\langle B \rangle_1 \geq \frac{1}{2}\}}$ and $I_{\{\langle B \rangle_1 \leq \frac{1}{2}\}} \notin$ $L^2_G(\Omega_1)$, which implies that

$$I_{\{\langle B \rangle_1 > \frac{1}{2}\}} = 1 - I_{\{\langle B \rangle_1 \le \frac{1}{2}\}} \notin L^2_G(\Omega_1).$$

Set $\mathcal{H}_1 = \{h_k(\langle B \rangle_1) : k \ge 1\}$ and $\mathcal{H}_2 = \{g_k(\langle B \rangle_1) : k \ge 1\}$, where

$$g_k(x) = k(x - \frac{1}{2} + \frac{1}{k})I_{\left[\frac{1}{2} - \frac{1}{k}, \frac{1}{2}\right]}(x) + I_{\left(\frac{1}{2}, \infty\right)}(x)$$

We assert that either $\underset{\xi \in \mathcal{H}_1}{\operatorname{essinf}} \xi$ or $\underset{\xi \in \mathcal{H}_2}{\operatorname{essinf}} \xi$ does not exist. Otherwise, $\underset{\xi \in \mathcal{H}_1}{\operatorname{essinf}} \xi$ and $\underset{\xi \in \mathcal{H}_2}{\operatorname{ess\,inf} \xi \text{ belong to } L^2_G(\Omega_1). }$ By the definition we get

$$\begin{aligned} & \underset{\xi \in \mathcal{H}_1}{\operatorname{ess\,inf}} \xi \leq I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \ q.s.; \\ & \underset{\xi \in \mathcal{H}_2}{\operatorname{ess\,inf}} \xi \leq I_{\{\langle B \rangle_1 \geq \frac{1}{2}\}} \ q.s.; \\ & (\underset{\xi \in \mathcal{H}_1}{\operatorname{ess\,inf}} \xi - \underset{\xi \in \mathcal{H}_2}{\operatorname{ess\,inf}} \xi)^+ \in L^2_G(\Omega_1); \\ & (\underset{\xi \in \mathcal{H}_1}{\operatorname{ess\,inf}} \xi - \underset{\xi \in \mathcal{H}_2}{\operatorname{ess\,inf}} \xi)^- \in L^2_G(\Omega_1), \end{aligned}$$

which implies that

$$I_{\{\langle B \rangle_1 = \frac{1}{2}\}} \underset{\xi \in \mathcal{H}_2}{\operatorname{ess\,inf}} \xi = \underset{\xi \in \mathcal{H}_1}{\operatorname{ess\,inf}} \xi.$$

Note that $\tilde{h}_k(\langle B \rangle_1) \leq \underset{\xi \in \mathcal{H}_2}{\operatorname{essinf}} \xi \text{ for } k \geq 1, \text{ where }$

$$\tilde{h}_k(x) = k(x - \frac{1}{2})I_{[\frac{1}{2}, \frac{1}{2} + \frac{1}{k})}(x) + I_{[\frac{1}{2} + \frac{1}{k}, \infty)}(x).$$

It yields that $I_{\{\langle B \rangle_1 > \frac{1}{2}\}} \leq \underset{\xi \in \mathcal{H}_2}{\operatorname{essinf}} \xi \ q.s.$ Then $\underset{\xi \in \mathcal{H}_2}{\operatorname{essinf}} \xi = \underset{\xi \in \mathcal{H}_1}{\operatorname{essinf}} \xi + I_{\{\langle B \rangle_1 > \frac{1}{2}\}}$ which implies $I_{\{\langle B \rangle_1 > \frac{1}{2}\}} \in L^{2}_{G}(\Omega_1)$.

But this contradicts to $I_{\{\langle B \rangle_1 > \frac{1}{2}\}} \notin L^2_G(\Omega_1)$.

Our stochastic optimal control problem is: for given $x \in \mathbb{R}^n$, to find $u(\cdot) \in \mathcal{U}[t,T]$ so as to minimize the objective function $Y_t^{t,x,u}$.

For $x \in \mathbb{R}^n$, we define the value function

$$V(t,x) := \operatorname{ess\,inf}_{u \in \mathcal{U}[t,T]} Y_t^{t,x,u} \text{ for } x \in \mathbb{R}^n.$$

$$\tag{7}$$

In the following we will prove that $V(\cdot, \cdot)$ exists and is deterministic and for each $\xi \in \bigcup_{\varepsilon>0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$, $V(t,\xi) = \underset{u\in\mathcal{U}[t,T]}{ess} inf Y_t^{t,\xi,u}$. Furthermore, we will obtain the dynamic programming principle and the related fully nonlinear HJB equation.

4 Properties of the value function

We first give some notations:

$$\begin{split} L_{ip}(\Omega_s^t) &:= \{\varphi(B_{t_1} - B_t, ..., B_{t_n} - B_t) : n \ge 1, t_1, ..., t_n \in [t, s], \varphi \in C_{b.Lip}(\mathbb{R}^{d \times n})\};\\ L_G^2(\Omega_s^t) &:= \{\text{the completion of } L_{ip}(\Omega_s^t) \text{ under the norm } \| \cdot \|_{2,G}\};\\ M_G^{0,t}(t,T) &:= \{\eta_s = \sum_{i=0}^{N-1} \xi_i I_{[t_i,t_{i+1})}(s) : t = t_0 < \dots < t_N = T, \xi_i \in L_{ip}(\Omega_{t_i}^t)\};\\ M_G^{2,t}(t,T) &:= \{\text{the completion of } M_G^{0,t}(t,T) \text{ under the norm } \| \cdot \|_{M_G^2}\};\\ \mathcal{U}[t,T] &:= \{u : u \in M_G^2(t,T;\mathbb{R}^m) \text{ with values in } U\};\\ \mathcal{U}^t[t,T] &:= \{u : u \in M_G^{2,t}(t,T;\mathbb{R}^m) \text{ with values in } U\};\\ \mathbb{U}[t,T] &:= \{u = \sum_{i=1}^n I_{A_i} u^i : n \in \mathbb{N}, u^i \in \mathcal{U}^t[t,T], I_{A_i} \in L_G^2(\Omega_t), \Omega = \bigcup_{i=1}^n A_i\};\\ \mathbb{U}^t[t,T] &:= \{u = \sum_{i=0}^{N-1} (\sum_{j=1}^l a_j^i I_{A_j^j}) I_{[t_i,t_{i+1})}(s) : l_i \in \mathbb{N}, a_j^i \in U, I_{A_j^i} \in L_G^2(\Omega_{t_i}^t), \Omega = \bigcup_{j=1}^l A_j^i\}. \end{split}$$

Remark 12 For $t = t_0 < \cdots < t_N = T$, $\xi_i \in L^2_G(\Omega^t_{t_i})$, it is easy to check that $\sum_{i=0}^{N-1} \xi_i I_{[t_i,t_{i+1})}(s) \in M^{2,t}_G(t,T)$. From this we can deduce that $\mathbb{U}^t[t,T] \subset \mathcal{U}^t[t,T] \subset \mathcal{U}[t,T] \subset \mathcal{U}[t,T]$.

In order to prove

$$V(t,x) = \inf_{u \in \mathcal{U}^t[t,T]} Y_t^{t,x,u} = \inf_{u \in \mathbb{U}^t[t,T]} Y_t^{t,x,u},$$

we need the following lemmas.

Lemma 13 Let $u \in \mathcal{U}[t,T]$ be given. Then there exists a sequence $(u^k)_{k\geq 1}$ in $\mathbb{U}[t,T]$ such that

$$\lim_{k \to \infty} \hat{\mathbb{E}}[\int_t^T |u_s - u_s^k|^2 ds] = 0.$$

Proof. For each $\varepsilon > 0$, we only need to prove that there exists a process $v \in \mathbb{U}[t,T]$ such that $\hat{\mathbb{E}}[\int_t^T |u_s - v_s|^2 ds] \leq \varepsilon$. Since $u \in M_G^2(t,T;\mathbb{R}^m)$, there exists a sequence processes $v^k \in M_G^0(t,T;\mathbb{R}^m)$ such that $\hat{\mathbb{E}}[\int_t^T |u_s - v_s^k|^2 ds] \to 0$. Set $U^{\varepsilon} := \{a \in \mathbb{R}^m : d(a,U) \leq \frac{\sqrt{\varepsilon}}{4}\}$, then

$$\hat{\mathbb{E}}[\int_t^T |u_s - v_s^k|^2 ds] \ge \hat{\mathbb{E}}[\int_t^T |u_s - v_s^k|^2 I_{\{v_s^k \notin U^\varepsilon\}} ds] \ge \frac{\varepsilon}{16} \hat{\mathbb{E}}[\int_t^T I_{\{v_s^k \notin U^\varepsilon\}} ds],$$

which implies that $\hat{\mathbb{E}}[\int_t^T I_{\{v_s^k \notin U^{\varepsilon}\}} ds] \to 0$. Thus there exists a $k_0 \ge 1$ such that

$$\hat{\mathbb{E}}[\int_t^T |u_s - v_s^{k_0}|^2 ds] \leq \frac{\varepsilon}{4}, \ \hat{\mathbb{E}}[\int_t^T I_{\{v_s^{k_0} \notin U^\varepsilon\}} ds] \leq \frac{\varepsilon}{16M^2},$$

where $M = \sup\{|a| : a \in U\}$. Set $\bar{v} = v^{k_0}$, we can write \bar{v} as

$$\bar{v}_s = \varphi_0(\xi_0) I_{[t_0, t_1)}(s) + \sum_{i=1}^{N-1} \varphi_i(\xi_0, \xi_i) I_{[t_i, t_{i+1})}(s),$$

where $t = t_0 < t_1 < \cdots < t_N = T$, $\xi_0 = (B_{s_1^0}, \dots, B_{s_{k_0}^0})$ for $s_j^0 \in [0, t]$, $\xi_i = (B_{s_1^i} - B_t, \dots, B_{s_{k_i}^i} - B_t)$ for $s_j^i \in [t, t_i]$, $i \ge 1$, $\varphi_i \in C_{b.Lip}(\mathbb{R}^{n_i}; \mathbb{R}^m)$ with $n_0 = dk_0, n_i = d(k_0 + k_i)$ for $i \ge 1$. Obviously, we can find two constants $\overline{M} > 0$ and L > 0 such that for $i \le N - 1$,

$$|\varphi_i| \leq \overline{M}, \ |\varphi_i(x^i) - \varphi_i(\overline{x}^i)| \leq L|x^i - \overline{x}^i| \text{ for } x^i, \overline{x}^i \in \mathbb{R}^{n_i}.$$

For each $k \geq 1$, we can find finite nonempty cubes $A_j^{i,k} \subset \mathbb{R}^{dk_i}$, $i \geq 0$, $j = 1, \ldots, l_i^k - 1$, such that $[-ke^i, ke^i) = \bigcup_{j \leq l_i^k - 1} A_j^{i,k}$ with $e^i = [1, \ldots, 1]^T \in \mathbb{R}^{dk_i}$ and $\rho(A_j^{i,k}) := \sup\{|x^i - \bar{x}^i| : x^i, \bar{x}^i \in A_j^{i,k}\} \leq \frac{1}{k}$. Set $A_{l_i^k}^{i,k} = \mathbb{R}^{dk_i} \setminus [-ke^i, ke^i)$ and

$$\begin{split} \bar{v}_{s}^{k} =& (\sum_{j_{0} \leq l_{0}^{k}} \varphi_{0}(x_{j_{0}}^{0,k}) I_{\{\xi_{0} \in A_{j_{0}}^{0,k}\}}) I_{[t_{0},t_{1})}(s) \\ &+ \sum_{i=1}^{N-1} (\sum_{j_{0} \leq l_{0}^{k}, j_{i} \leq l_{i}^{k}} \varphi_{i}(x_{j_{0}}^{0,k}, x_{j_{i}}^{i,k}) I_{\{\xi_{0} \in A_{j_{0}}^{0,k}\}} I_{\{\xi_{i} \in A_{j_{i}}^{i,k}\}}) I_{[t_{i},t_{i+1})}(s) \\ &= \sum_{j_{0} \leq l_{0}^{k}} I_{\{\xi_{0} \in A_{j_{0}}^{0,k}\}} \left(\varphi_{0}(x_{j_{0}}^{0,k}) I_{[t_{0},t_{1})}(s) + \sum_{i=1}^{N-1} (\sum_{j_{i} \leq l_{i}^{k}} \varphi_{i}(x_{j_{0}}^{0,k}, x_{j_{i}}^{i,k}) I_{\{\xi_{i} \in A_{j_{i}}^{i,k}\}}) I_{[t_{i},t_{i+1})}(s) \right) \end{split}$$

where $x_j^{i,k}$ is one point belonging to $A_j^{i,k}$ for $i \ge 0$ and $j \le l_i^k$. By Theorem 8 we can get $I_{\{\xi_0 \in A_{j_0}^{0,k}\}} \in L^2_G(\Omega_t)$ and $I_{\{\xi_i \in A_{j_i}^{i,k}\}} \in L^2_G(\Omega_{t_i}^t)$ for $i \ge 1$. Then we have

$$\hat{\mathbb{E}}\left[\int_{t}^{T} |\bar{v}_{s} - \bar{v}_{s}^{k}|^{2} ds\right] \leq \sum_{i=0}^{N-1} \hat{\mathbb{E}}\left[|\bar{v}_{t_{i}} - \bar{v}_{t_{i}}^{k}|^{2}\right](t_{i+1} - t_{i})$$

$$\leq \sum_{i=0}^{N-1} \hat{\mathbb{E}}\left[\frac{2L^{2}}{k^{2}} + \frac{4\bar{M}^{2}}{k^{2}}(|\xi_{0}|^{2} + |\xi_{i}|^{2})\right](t_{i+1} - t_{i})$$

$$\to 0 \text{ as } k \to \infty.$$

We set

$$\tilde{v}_{s}^{k} = \sum_{j_{0} \leq l_{0}^{k}} I_{\{\xi_{0} \in A_{j_{0}}^{0,k}\}} \left(\tilde{\varphi}_{0}(x_{j_{0}}^{0,k}) I_{[t_{0},t_{1})}(s) + \sum_{i=1}^{N-1} (\sum_{j_{i} \leq l_{i}^{k}} \tilde{\varphi}_{i}(x_{j_{0}}^{0,k}, x_{j_{i}}^{i,k}) I_{\{\xi_{i} \in A_{j_{i}}^{i,k}\}}) I_{[t_{i},t_{i+1})}(s) \right),$$

where $\tilde{\varphi}_i(x_{j_0}^{0,k}, x_{j_i}^{i,k})$ is one point in U such that $|\varphi_i(x_{j_0}^{0,k}, x_{j_i}^{i,k}) - \tilde{\varphi}_i(x_{j_0}^{0,k}, x_{j_i}^{i,k})| = d(\varphi_i(x_{j_0}^{0,k}, x_{j_i}^{i,k}), U)$. By Remark 12, it is easy to verify that $\tilde{v}^k \in \mathbb{U}[t, T]$ and

$$\tilde{\varphi}_0(x_{j_0}^{0,k})I_{[t_0,t_1)}(s) + \sum_{i=1}^{N-1} (\sum_{j_i \le l_i^k} \tilde{\varphi}_i(x_{j_0}^{0,k}, x_{j_i}^{i,k})I_{\{\xi_i \in A_{j_i}^{i,k}\}})I_{[t_i,t_{i+1})}(s) \in \mathbb{U}^t[t,T].$$

Note that

$$\begin{aligned} |u_s - \tilde{v}_s^k|^2 &= |u_s - \tilde{v}_s^k|^2 I_{\{\bar{v}_s \notin U^\varepsilon\}} + |u_s - \tilde{v}_s^k|^2 I_{\{\bar{v}_s \in U^\varepsilon\}} \\ &\leq 4M^2 I_{\{\bar{v}_s \notin U^\varepsilon\}} + 2|u_s - \bar{v}_s|^2 + 2|\bar{v}_s - \tilde{v}_s^k|^2 I_{\{\bar{v}_s \in U^\varepsilon\}} \\ &\leq 4M^2 I_{\{\bar{v}_s \notin U^\varepsilon\}} + 2|u_s - \bar{v}_s|^2 + 2(\frac{\sqrt{\varepsilon}}{4} + \frac{2\sqrt{2}L}{k})^2, \end{aligned}$$

then we get

$$\limsup_{k \to \infty} \hat{\mathbb{E}}\left[\int_t^T |u_s - \tilde{v}_s^k|^2 ds\right] \le \frac{7}{8}\varepsilon.$$

Thus there exists a $k_1 \geq 1$ such that $\hat{\mathbb{E}}[\int_t^T |u_s - \tilde{v}_s^{k_1}|^2 ds] \leq \varepsilon$. The proof is complete by taking $v = \tilde{v}^{k_1} \in \mathbb{U}[t,T]$.

Lemma 14 Let $u \in \mathcal{U}^t[t,T]$ be given. Then there exists a sequence $(u^k)_{k\geq 1}$ in $\mathbb{U}^t[t,T]$ such that

$$\lim_{k \to \infty} \hat{\mathbb{E}}\left[\int_t^T |u_s - u_s^k|^2 ds\right] = 0.$$

Proof. The proof is the same as Lemma 13, we omit it.

Lemma 15 Let $\xi \in \bigcup_{\varepsilon>0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$, $u \in \mathcal{U}[t,T]$ and $v_s = \sum_{i=1}^N I_{A_i} v_s^i \in \mathbb{U}[t,T]$. Then there exists a constant L_1 depending on T, G and C such that

$$\hat{\mathbb{E}}[|Y_t^{t,\xi,u} - \sum_{i=1}^N I_{A_i} Y_t^{t,\xi,v^i}|^2] \le L_1 \hat{\mathbb{E}}[\int_t^T |u_s - v_s|^2 ds].$$

Proof. Consider the following equations:

$$\begin{split} dX_s^{t,\xi,v^i} &= b(s, X_s^{t,\xi,v^i}, v_s^i) ds + h_{ij}(s, X_s^{t,\xi,v^i}, v_s^i) d\langle B^i, B^j \rangle_s + \sigma(s, X_s^{t,\xi,v^i}, v_s^i) dB_s, \\ X_t^{t,\xi,v^i} &= \xi, \ s \in [t,T], \ i = 1, ..., N. \end{split}$$

$$\begin{split} -dY_{s}^{t,\xi,v^{i}} &= f(s, X_{s}^{t,\xi,v^{i}}, Y_{s}^{t,\xi,v^{i}}, Z_{s}^{t,\xi,v^{i}}, v_{s}^{i})ds + g_{ij}(s, X_{s}^{t,\xi,v^{i}}, Y_{s}^{t,\xi,v^{i}}, Z_{s}^{t,\xi,v^{i}}, v_{s}^{i})d\langle B^{i}, B^{j}\rangle_{s} \\ &- Z_{s}^{t,\xi,v^{i}}dB_{s} - dK_{s}^{t,\xi,v^{i}}, \\ Y_{T}^{t,\xi,v^{i}} &= \Phi(X_{T}^{t,\xi,v^{i}}), \quad s \in [t,T], \, i = 1, ..., N. \end{split}$$

For $s \in [t, T]$, we set

$$\bar{X}_{s}^{t,\xi,v} = \sum_{i=1}^{N} I_{A_{i}} X_{s}^{t,\xi,v^{i}}, \bar{Y}_{s}^{t,\xi,v} = \sum_{i=1}^{N} I_{A_{i}} Y_{s}^{t,\xi,v^{i}}, \bar{Z}_{s}^{t,\xi,v} = \sum_{i=1}^{N} I_{A_{i}} Z_{s}^{t,\xi,v^{i}}, \bar{K}_{s}^{t,\xi,v} = \sum_{i=1}^{N} I_{A_{i}} K_{s}^{t,\xi,v^{i}}.$$

Multiplying I_{A_i} on both sides of the above equations and summing up, we have

$$\begin{split} d\bar{X}_{s}^{t,\xi,v} &= b(s, \bar{X}_{s}^{t,\xi,v}, v_{s}) ds + h_{ij}(s, \bar{X}_{s}^{t,\xi,v}, v_{s}) d\langle B^{i}, B^{j} \rangle_{s} + \sigma(s, \bar{X}_{s}^{t,\xi,v}, v_{s}) dB_{s}, \\ \bar{X}_{t}^{t,\xi,v} &= \xi, \ s \in [t,T], \end{split}$$

$$\begin{split} -d\bar{Y}_{s}^{t,\xi,v} &= f(s, \bar{X}_{s}^{t,\xi,v}, \bar{Y}_{s}^{t,\xi,v}, \bar{Z}_{s}^{t,\xi,v}, v_{s})ds + g_{ij}(s, \bar{X}_{s}^{t,\xi,v}, \bar{Y}_{s}^{t,\xi,v}, \bar{Z}_{s}^{t,\xi,v}, v_{s})d\langle B^{i}, B^{j}\rangle_{s} \\ &- Z_{s}^{t,\xi,v^{i}}dB_{s} - d\bar{K}_{s}^{t,\xi,v}, \\ \bar{Y}_{T}^{t,\xi,v} &= \Phi(\bar{X}_{T}^{t,\xi,v}), \quad s \in [t,T]. \end{split}$$

By Theorem 7, we can obtain that there exists a constant $C_1 > 0$ depending on T, G and C such that

$$\begin{aligned} &|Y_t^{t,\xi,u} - \sum_{i=1}^N I_{A_i} Y_t^{t,\xi,v^i}|^2 = |Y_t^{t,\xi,u} - \bar{Y}_t^{t,\xi,v}|^2 \\ &\leq C_1 \hat{\mathbb{E}}_t [| \Phi(X_T^{t,\xi,u}) - \Phi(\bar{X}_T^{t,\xi,v}) |^2 + \int_t^T (|X_s^{t,\xi,u} - \bar{X}_s^{t,\xi,v}|^2 + |u_s - v_s|^2) ds] \\ &\leq C_1 \hat{\mathbb{E}}_t [C \mid X_T^{t,\xi,u} - \bar{X}_T^{t,\xi,v} \mid^2 + \int_t^T (|X_s^{t,\xi,u} - \bar{X}_s^{t,\xi,v}|^2 + |u_s - v_s|^2) ds], \end{aligned}$$

$$(8)$$

where C is the Lipschitz constant of Φ . By Theorem 5, there exists a constant $C_2 > 0$ depending on T, n, G and C such that

$$\hat{\mathbb{E}}[|X_s^{t,\xi,u} - \bar{X}_s^{t,\xi,v}|^2] \le C_2 \hat{\mathbb{E}}[\int_t^T |u_s - v_s|^2 ds].$$

Taking G-expectation on both sides of (8), we obtain the result. \blacksquare

Remark 16 By the definition of generalized conditional G-expectation and Proposition 2.5 in [12], the above analysis still holds for the case that $\{A_i\}_{i=1}^N$ is a $\mathcal{B}(\Omega_t)$ -partition of Ω .

Theorem 17 The value function V(t, x) exists and

$$V(t,x) = \inf_{u \in \mathcal{U}^t[t,T]} Y_t^{t,x,u} = \inf_{u \in \mathbb{U}^t[t,T]} Y_t^{t,x,u}.$$

Proof. For each $v \in \mathcal{U}^t[t,T]$, it is easy to check that $Y_t^{t,x,v}$ is a constant. In the following, we prove that $\underset{u \in \mathcal{U}[t,T]}{essinf} Y_t^{t,x,u} = \inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v}$, q.s.. Since $\mathcal{U}^t[t,T] \subset \mathcal{U}[t,T]$, we only need to show that $Y_t^{t,x,u} \geq \inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v}$ q.s. for each $u \in \mathcal{U}[t,T]$. For each fixed $u \in \mathcal{U}[t,T]$, by Lemma 13, there exists a sequence $u^k = \sum_{i=1}^{N_k} I_{A_i^k} v^{i,k} \in \mathbb{U}[t,T]$, k = 1, 2, ..., such that

$$\widehat{\mathbb{E}}[\int_t^T |u_s - \sum_{i=1}^{N_k} I_{A_i^k} v^{i,k}|^2 ds] \to 0.$$

By Lemma 15,

$$\hat{\mathbb{E}}[|Y_t^{t,x,u} - \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x,v^{i,k}}|^2]$$

$$\leq L_1 \hat{\mathbb{E}}[\int_t^T |u_s - \sum_{i=1}^{N_k} I_{A_i^k} v^{i,k}|^2 ds].$$

It yields that $\sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x,v^{i,k}}$ converges to $Y_t^{t,x,u}$ in \mathbb{L}^2_G . Then there exists a subsequence (for simplicity, we still denote it by $\{\sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x,v^{i,k}}\}$) which converges to $Y_t^{t,x,u}$ q.s.. Note that

$$\sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x,v^{i,k}} \ge \inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v} \text{ q.s.},$$

then we have $Y_t^{t,x,u} \geq \inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v}$ q.s.. Thus

$$V(t,x) = \operatorname{ess\,inf}_{u \in \mathcal{U}[t,T]} Y_t^{t,x,u} = \operatorname{inf}_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v}.$$

Similarly, by Lemmas 14 and 15, we can get $\inf_{u \in \mathcal{U}^t[t,T]} Y_t^{t,x,u} = \inf_{u \in \mathbb{U}^t[t,T]} Y_t^{t,x,u}$. The proof is complete.

Lemma 18 There exists a constant $L_2 > 0$ depending on T, G and C such that

$$|V(t,x) - V(t,y)| \le L_2 |x-y|$$
 for any $x, y \in \mathbb{R}^n$.

Proof. By Theorems 5 and 7, for any $x, y \in \mathbb{R}^n$ and $u \in \mathcal{U}^t[t, T]$,

$$| Y_t^{t,x,u} - Y_t^{t,y,u} |^2$$

 $\leq C_1 \hat{\mathbb{E}}[| \Phi(X_T^{t,x,u}) - \Phi(X_T^{t,y,u}) |^2 + \int_t^T | X_s^{t,x,u} - X_s^{t,y,u} |^2 ds]$
 $\leq C_2 | x - y |^2 .$

It is easy to verify that $|\inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,x,v} - \inf_{v \in \mathcal{U}^t[t,T]} Y_t^{t,y,v}| \leq \sup_{v \in \mathcal{U}^t[t,T]} |Y_t^{t,x,v} - Y_t^{t,y,v}|$. Thus by the above estimate and Theorem 17, we obtain the result.

Lemma 19 There exists a constant $L_3 > 0$ depending on T, G and C such that

$$|V(t,x)| \leq L_3(1+|x|)$$
 for any $x \in \mathbb{R}^n$.

Proof. The proof is similar to Lemma 18, we omit it.

Theorem 20 For any $\xi \in \bigcup_{\varepsilon > 0} L^{2+\varepsilon}_G(\Omega_t; \mathbb{R}^n)$, we have

$$V(t,\xi) = \operatorname{ess\,inf}_{u \in \mathcal{U}[t,T]} Y_t^{t,\xi,u}.$$

Proof. First, we prove that $\forall u \in \mathcal{U}[t,T]$, $V(t,\xi) \leq Y_t^{t,\xi,u}$ q.s.. For a fixed $\xi \in \bigcup_{\varepsilon > 0} L_G^{2+\varepsilon}(\Omega_t; \mathbb{R}^n)$, we can find a sequence $\xi^k = \sum_{i=1}^{N_k} x_i^k I_{A_i^k}$, $k = 1, 2, \dots$, where $x_i^k \in \mathbb{R}^n$ and $\{A_i^k\}_{i=1}^{N_k}$ is a $\mathcal{B}(\Omega_t)$ -partition of Ω , such that

$$\lim_{k \to \infty} \hat{\mathbb{E}}[|\xi - \xi^k|^2] = 0.$$

Here $I_{A_i^k}$ may not in $L^2_G(\Omega_t)$. By Lemma 18, we have

$$\hat{\mathbb{E}}[|V(t,\xi) - V(t,\xi^k)|^2] \le L_2^2 \hat{\mathbb{E}}[|\xi - \xi^k|^2] \to 0$$

By similar analysis as in Lemma 15 and the definition of generalized conditional G-expectation,

$$\begin{aligned} &|Y_t^{t,\xi,u} - \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x_i^k,u}|^2 \\ &\leq C_1 \hat{\mathbb{E}}_t[| \Phi(X_T^{t,\xi,u}) - \Phi(\sum_{i=1}^{N_k} I_{A_i^k} X_T^{t,x_i^k,u}) |^2 + \int_t^T | X_s^{t,\xi,u} - \sum_{i=1}^{N_k} I_{A_i^k} X_s^{t,x_i^k,u} |^2 ds] \\ &\leq C_2 \hat{\mathbb{E}}_t[| \xi - \xi^k |^2]. \end{aligned}$$

Then

$$\lim_{k \to \infty} \hat{\mathbb{E}}[|Y_t^{t,\xi,u} - \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x_i^k,u}|^2] = 0.$$

Note that

$$V(t,\xi^k) = \sum_{i=1}^{N_k} I_{A_i^k} V(t,x_i^k) \le \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x_i^k,u} \text{ q.s.}$$

Thus

$$V(t,\xi) \le Y_t^{t,\xi,u}$$
 q.s..

Second, we prove that for a given $\eta \in L^2_G(\Omega_t)$, if $\forall u \in \mathcal{U}[t,T], \eta \leq Y_t^{t,\xi,u}$ q.s., then $\eta \leq V(t,\xi)$ q.s..

By the above analysis, we know that

$$|Y_t^{t,\xi,u} - \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t,x_i^k,u}|^2 \le C_2 \hat{\mathbb{E}}_t[|\xi - \xi^k|^2] \text{ q.s.}.$$

Then, for any $u \in \mathcal{U}[t,T]$,

$$\eta \leq \sum_{i=1}^{N_k} I_{A_i^k} Y_t^{t, x_i^k, u} + \sqrt{C_2 \hat{\mathbb{E}}_t [|\xi - \xi^k|^2]}$$
q.s..

For each fixed N_k , it yields that

$$\begin{split} \eta &\leq \sum_{i=1}^{N_k} I_{A_i^k} V(t, x_i^k) + \sqrt{C_2 \hat{\mathbb{E}}_t[|\xi - \xi^k|^2]} \\ &= V(t, \xi^k) + \sqrt{C_2 \hat{\mathbb{E}}_t[|\xi - \xi^k|^2]} \;\; \text{q.s..} \end{split}$$

Note that

$$\lim_{k \to \infty} \hat{\mathbb{E}}[|V(t,\xi) - V(t,\xi^k)|^2] = 0$$

and

$$\lim_{k \to \infty} \hat{\mathbb{E}}[|\xi - \xi^k|^2] = 0.$$

Then there exists a subsequence (ξ^{k_i}) of (ξ^k) such that as $k_i \longrightarrow \infty$,

$$V(t,\xi^{k_i}) \longrightarrow V(t,\xi), \ \hat{\mathbb{E}}_t[|\xi - \xi^{k_i}|^2] \longrightarrow 0, \ \text{q.s.}.$$

Thus $\eta \leq V(t,\xi)$ q.s.. This completes the proof.

5 Dynamic programming principle

For given initial data (t, x), a positive real number $\delta \leq T - t$ and $\eta \in \bigcup_{\varepsilon > 0} L_G^{2+\varepsilon}(\Omega_{t+\delta})$, we define

$$\mathbb{G}_{t,t+\delta}^{t,x,u}[\eta] := \tilde{Y}_t^{t,x,u}$$

where $(X_s^{t,x,u}, \tilde{Y}_s^{t,x,u}, \tilde{Z}_s^{t,x,u})_{t \le s \le t+\delta}$ is the solution of the following forward and backward equations:

$$\begin{array}{lll} dX_s^{t,x,u} = & b(s, X_s^{t,x,u}, u_s)ds + h_{ij}(s, X_s^{t,x,u}, u_s)d\langle B^i, B^j\rangle_s + \sigma(s, X_s^{t,x,u}, u_s)dB_s, \\ X_t^{t,x,u} = & x, \end{array}$$

and

$$\begin{aligned} -d\tilde{Y}_{s}^{t,x,u} &= f(s, X_{s}^{t,x,u}, \tilde{Y}_{s}^{t,x,u}, \tilde{Z}_{s}^{t,x,u}, u_{s})ds + g_{ij}(s, X_{s}^{t,x,u}, \tilde{Y}_{s}^{t,x,u}, \tilde{Z}_{s}^{t,x,u}, u_{s})d\langle B^{i}, B^{j} \rangle_{s} \\ &- \tilde{Z}_{s}^{t,x,u}dB_{s} - d\tilde{K}_{s}^{t,x,u}, \\ \tilde{Y}_{t+\delta}^{t,x,u} &= \eta, \quad s \in [t, t+\delta]. \end{aligned}$$

$$(9)$$

(9) Note that $\mathbb{G}_{t,t+\delta}^{t,x,u}[\cdot]$ is a (backward) semigroup which was first introduced by Peng in [24].

Our main result in this section is the following dynamic programming principle.

Theorem 21 Let Assumptions (A1) and (A2) hold. Then for any $t \leq s \leq T$, $x \in \mathbb{R}^n$, we have

$$V(t,x) = \underset{u(\cdot) \in \mathcal{U}[t,s]}{\operatorname{essinf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})]$$

$$= \underset{u(\cdot) \in \mathcal{U}^t[t,s]}{\operatorname{inf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})].$$
(10)

In order to prove Theorem 21, we need the following lemmas.

Lemma 22 Let Assumptions (A1) and (A2) hold. Assume further that there exist constants L > 0, $\Lambda > 0$ such that $|b| \leq L$, $|h_{ij}| \leq L$ and $|\sigma_i| \leq \Lambda$ for $i \leq n$ and $(t, x, u) \in [0, T] \times \mathbb{R}^n \times U$, where σ_i is the *i*-th row of σ . Then for any $t < s \leq T$, $x \in \mathbb{R}^n$, we have

$$V(t,x) \leq \inf_{u(\cdot) \in \mathbb{U}^t[t,s]} \mathbb{G}^{t,x,u}_{t,s}[V(s,X^{t,x,u}_s)].$$

Proof. For each $\varepsilon > 0$, there exists a $u(\cdot) \in \mathbb{U}^t[t,s]$ such that

$$\mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})] - \varepsilon \leq \inf_{v(\cdot) \in \mathbb{U}^t[t,s]} \mathbb{G}_{t,s}^{t,x,v}[V(s,X_s^{t,x,v})].$$

We can write $u(\cdot)$ as

$$u_r = a_1^0 I_{[t_0, t_1)}(r) + \sum_{i=1}^{N-1} (\sum_{j=1}^{l_i} a_j^i I_{A_j^i}) I_{[t_i, t_{i+1})}(r),$$

where $t = t_0 < t_1 < \cdots < t_N = s$, $l_i \in \mathbb{N}$, $a_j^i \in U$, $I_{A_j^i} \in L^2_G(\Omega_{t_i}^t)$ and $\{A_j^i\}_{j=1}^{l_i}$ is a partition of Ω . Consider the following SDE: for any $v(\cdot) \in \mathbb{U}^t[t, s]$,

$$\begin{split} d\tilde{X}_{r}^{t,x,v} &= b(r, \tilde{X}_{r}^{t,x,v} - \tilde{X}_{r}e, v_{r})dr + h_{ij}(r, \tilde{X}_{r}^{t,x,v} - \tilde{X}_{r}e, v_{r})d\langle B^{i}, B^{j}\rangle_{r} \\ &+ \sigma(r, \tilde{X}_{r}^{t,x,v} - \tilde{X}_{r}e, v_{r})dB_{r} + (\Lambda + 1)edB_{r}^{1}, \\ d\tilde{X}_{r} &= (\Lambda + 1)dB_{r}^{1}, \\ \tilde{X}_{t}^{t,x,v} &= x, \ \tilde{X}_{t} = 0, \ r \in [t,s], \end{split}$$

where $e = [1, ..., 1]^T \in \mathbb{R}^n$. It is easy to verify that

$$\tilde{X}_{r}^{t,x,v} = X_{r}^{t,x,v} + (\Lambda + 1)(B_{r}^{1} - B_{t}^{1})e, \ \tilde{X}_{r} = (\Lambda + 1)(B_{r}^{1} - B_{t}^{1}), \ r \in [t,s]$$

is the solution of the above SDE. We set

$$I_{i+1} = \{J_{i+1} = (j_0, j_1, \dots, j_i) : 1 \le j_k \le l_k, 0 \le k \le i\},\$$

where $l_0 = 1, i \leq N-1$. For each given $J_{i+1} = (j_0, j_1, \dots, j_i) \in I_{i+1}$, we denote

$$\tilde{X}_r^{t,x,J_{i+1}} := \tilde{X}_r^{t,x,u^{J_{i+1}}}, \ r \in [t, t_{i+1}]$$

where $u_r^{J_{i+1}} = \sum_{k=0}^i a_{j_k}^k I_{[t_k, t_{k+1})}(r)$ is deterministic. We claim that

$$\tilde{X}_{t_{i+1}}^{t,x,u} = \sum_{J_{i+1} \in I_{i+1}} (\prod_{k=1}^{i} I_{A_{j_k}^k}) \tilde{X}_{t_{i+1}}^{t,x,J_{i+1}} \text{ for } 0 \le i \le N-1,$$
(11)

where $\Pi_{k=1}^{0} I_{A_{j_k}^k} = 1$. It is easy to check that the equality (11) holds for i = 0. Suppose that the equality (11) holds for $i_0 \ge 0$, then by the similar analysis as in the proof of Lemma 15, we can get

$$\tilde{X}_{t_{i_{0}+2}}^{t,x,u} = \sum_{j=1}^{t_{i_{0}+1}} I_{A_{j}^{i_{0}+1}} \tilde{X}_{t_{i_{0}+2}}^{t_{i_{0}+1},\xi,a_{j}^{i_{0}+1}}, \ \tilde{X}_{t_{i_{0}+2}}^{t_{i_{0}+1},\xi,a_{j}^{i_{0}+1}} = \sum_{J_{i_{0}+1} \in I_{i_{0}+1}} (\prod_{k=1}^{i_{0}} I_{A_{j_{k}}^{k}}) \tilde{X}_{t_{i_{0}+2}}^{t_{i_{0}+1},\xi^{J_{i_{0}+1}},a_{j}^{i_{0}+1}},$$

where $\xi = \tilde{X}_{t_{i_0+1}}^{t,x,u}, \xi^{J_{i_0+1}} = \tilde{X}_{t_{i_0+1}}^{t,x,J_{i_0+1}}$. It is easy to verify that $\tilde{X}_{t_{i_0+2}}^{t_{i_0+1},\xi^{J_{i_0+1}},a_j^{i_0+1}} = \tilde{X}_{t_{i_0+2}}^{t,x,(J_{i_0+1},a_j^{i_0+1})}$. Thus the equality (11) holds for $i_0 + 1$. From this we can deduce that

$$\tilde{X}_{s}^{t,x,u} = \sum_{J_{N} \in I_{N}} (\prod_{k=1}^{N-1} I_{A_{j_{k}}^{k}}) \tilde{X}_{s}^{t,x,J_{N}}.$$

It is easy to check that

$$\sqrt{|\sigma_{i1} + \Lambda + 1|^2 + |\sigma_{i2}|^2 + \dots + |\sigma_{id}|^2} \ge 1,$$

where σ_{ij} is the *i*-th row and *j*-th column of σ . Then by Theorem 8, we have $I_{\{\tilde{X}_s^{t,x,J_N} \in [a,a')\}} \in L^2_G(\Omega^t_s)$ for $a, a' \in \mathbb{R}^n$ with $a \leq a'$. Thus

$$I_{\{\tilde{X}^{t,x,u}_{s}\in[a,a')\}} = \sum_{J_{N}\in I_{N}} (\prod_{k=1}^{N-1} I_{A^{k}_{j_{k}}}) I_{\{\tilde{X}^{t,x,J_{N}}_{s}\in[a,a')\}} \in L^{2}_{G}(\Omega^{t}_{s}).$$

For each integer $k \geq 1$, we can choose finite nonempty cubes $D_j^k \subset \mathbb{R}^n$, $E_j^k \subset \mathbb{R}$ and $x_j^k \in D_j^k$, $q_j^k \in E_j^k$ for $j = 1, \ldots, l^k$, such that $[-ke, ke) = \bigcup_{j \leq l^k} D_j^k$, $[-k,k) = \bigcup_{j \leq l^k} E_j^k$, $\rho(D_j^k) = \sup\{|x - \bar{x}| : x, \bar{x} \in D_j^k\} \leq \frac{1}{k}$ and $\rho(E_j^k) = \sup\{|q - \bar{q}| : q, \bar{q} \in E_j^k\} \leq \frac{1}{\sqrt{n}(\Lambda + 1)k}$. Set

$$\xi^{k,u} = \sum_{j_1=1}^{l^k} \sum_{j_2=1}^{l^k} (x_{j_1}^k - (\Lambda+1)q_{j_2}^k e) I_{\{\tilde{X}_s^{t,x,u} \in D_{j_1}^k\}} I_{\{B_s^1 - B_t^1 \in E_{j_2}^k\}}.$$

Note that $\tilde{X}_s^{t,x,u} = X_s^{t,x,u} + (\Lambda + 1)(B_s^1 - B_t^1)e$, then we get

$$|X_s^{t,x,u} - \xi^{k,u}| \le \frac{2}{k} + |X_s^{t,x,u}| \frac{|X_s^{t,x,u}| + \sqrt{n}(\Lambda + 1)|B_s^1 - B_t^1|}{k}.$$

By Theorem 5, there exists a constant $C_1 > 0$ depending on T, n, U, G and C such that

$$\hat{\mathbb{E}}[|X_s^{t,x,u}|^4] \le C_1(1+|x|^4).$$

Thus we obtain

$$\hat{\mathbb{E}}[|X_s^{t,x,u} - \xi^{k,u}|^2] \le \hat{\mathbb{E}}[\frac{8(1 + |X_s^{t,x,u}|^4) + n^2(\Lambda + 1)^4|B_s^1 - B_t^1|^4}{k^2}] \le \frac{C_2(1 + |x|^4)}{k^2}.$$

Set $\tilde{x}_{j_1j_2}^k = x_{j_1}^k - (\Lambda + 1)q_{j_2}^k e$ for $j_1, j_2 \leq l^k$, by Theorem 17, $V(s, \tilde{x}_{j_1j_2}^k) = \inf_{u \in \mathbb{U}^s[s,T]} Y_s^{s, \tilde{x}_{j_1j_2}^k, u}$ and $V(s, 0) = \inf_{u \in \mathbb{U}^s[s,T]} Y_s^{s,0,u}$, then we can choose $\bar{u}^{j_1j_2,k}$, $\bar{u}^{0,k} \in \mathbb{U}^s[s,T]$ such that

$$V(s, \tilde{x}_{j_1 j_2}^k) \le Y_s^{s, \tilde{x}_{j_1 j_2}^k, \bar{u}^{j_1 j_2, k}} \le V(s, \tilde{x}_{j_1 j_2}^k) + \varepsilon, \ V(s, 0) \le Y_s^{s, 0, \bar{u}^{0, k}} \le V(s, 0) + \varepsilon.$$

Set

$$\bar{u}^{k} = \sum_{j_{1}=1}^{l^{k}} \sum_{j_{2}=1}^{l^{k}} I_{\{\tilde{X}_{s}^{t,x,u} \in D_{j_{1}}^{k}\}} I_{\{B_{s}^{1} - B_{t}^{1} \in E_{j_{2}}^{k}\}} \bar{u}^{j_{1}j_{2},k} + I_{\{\tilde{X}_{s}^{t,x,u} \in [-ke,ke)^{c}\} \cup \{B_{s}^{1} - B_{t}^{1} \in [-k,k)^{c}\}} \bar{u}^{0,k},$$

by the similar analysis as in the proof of Lemma 15, we can get

$$V(s,\xi^{k,u}) \le Y_s^{s,\xi^{k,u},\bar{u}^k} \le V(s,\xi^{k,u}) + \varepsilon.$$

By Theorem 7, there exists a constant $C_3 > 0$ depending on T, G and C such that Ŕ

$$\hat{\mathbb{E}}[|Y_s^{s,X_s^{t,x,u},\bar{u}^k} - Y_s^{s,\xi^{k,u},\bar{u}^k}|^2] \le C_3 \hat{\mathbb{E}}[|X_s^{t,x,u} - \xi^{k,u}|^2].$$

Set $\tilde{u}(r) = u(r)I_{[t,s]}(r) + \bar{u}^k(r)I_{(s,T]}(r)$, it is easy to check that $\tilde{u} \in \mathbb{U}^t[t,T]$ and

$$V(t,x) \le Y_t^{t,x,\tilde{u}} = \mathbb{G}_{t,s}^{t,x,u} [Y_s^{s,X_s^{t,x,u},\bar{u}^k}].$$

By Theorem 7, we get

$$\begin{split} |\mathbb{G}_{t,s}^{t,x,u}[Y_s^{s,X_s^{t,x,u},\bar{u}^k}] - \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})]|^2 \\ &\leq C_3 \hat{\mathbb{E}}[|Y_s^{s,X_s^{t,x,u},\bar{u}^k} - V(s,X_s^{t,x,u})|^2] \\ &\leq 2C_3 (\hat{\mathbb{E}}[|Y_s^{s,X_s^{t,x,u},\bar{u}^k} - Y_s^{s,\xi^{k,u},\bar{u}^k}|^2] + \hat{\mathbb{E}}[|Y_s^{s,\xi^{k,u},\bar{u}^k} - V(s,X_s^{t,x,u})|^2]) \\ &\leq 2C_3 (C_3 \hat{\mathbb{E}}[|X_s^{t,x,u} - \xi^{k,u}|^2] + \hat{\mathbb{E}}[(|V(s,\xi^{k,u}) - V(s,X_s^{t,x,u})| + \varepsilon)^2]) \\ &\leq \frac{C_4 (1 + |x|^4)}{k^2} + 4C_3 \varepsilon^2. \end{split}$$

Thus

$$V(t,x) - \frac{\sqrt{C_4(1+|x|^4)}}{k} - (2\sqrt{C_3}+1)\varepsilon \le \inf_{v(\cdot)\in\mathbb{U}^t[t,s]} \mathbb{G}_{t,s}^{t,x,v}[V(s,X_s^{t,x,v})].$$

Letting $k \to \infty$ first and then $\varepsilon \downarrow 0$, we obtain $V(t, x) \leq \inf_{v(\cdot) \in \mathbb{U}^t[t,s]} \mathbb{G}_{t,s}^{t,x,v}[V(s, X_s^{t,x,v})]$. The proof is complete. \blacksquare

Remark 23 In the above proof, $\xi^{k,u}$ is called an "implied partition" of $X_s^{t,x,u}$. **Lemma 24** Let Assumptions (A1) and (A2) hold. Then for any $t < s \leq T$, $x \in \mathbb{R}^n$, we have

$$V(t,x) \leq \inf_{u(\cdot) \in \mathbb{U}^t[t,s]} \mathbb{G}^{t,x,u}_{t,s}[V(s,X^{t,x,u}_s)].$$

Proof. For each fixed N > 0, we set $b^{i_1,N} = (b^{i_1} \wedge N) \vee (-N)$, $h^{i_1,N}_{ij} = (h^{i_1}_{ij} \wedge N) \vee (-N)$, $\sigma^N_{i_1i_2} = (\sigma_{i_1i_2} \wedge N) \vee (-N)$ for $i_1 \leq n$, $i_2 \leq d$ and $b^N = (b^{1,N}, \dots, b^{n,N})^T$, $h^N_{ij} = (h^{1,N}_{ij}, \dots, h^{n,N}_{ij})^T$, $\sigma^N = (\sigma^N_{i_1i_2})_{i_1i_2}$. Consider the following FBSDEs: $dX^{t,x,u,N}_s = b^N(s, X^{t,x,u,N}_s, u_s)ds + h^N_{ij}(s, X^{t,x,u,N}_s, u_s)d\langle B^i, B^j \rangle_s + \sigma^N(s, X^{t,x,u,N}_s, u_s)dB_s, X^{t,x,u,N}_t = x,$

and

$$\begin{aligned} -dY_{s}^{t,x,u,N} &= f(s, X_{s}^{t,x,u,N}, Y_{s}^{t,x,u,N}, Z_{s}^{t,x,u,N}, u_{s})ds \\ &+ g_{ij}(s, X_{s}^{t,x,u,N}, Y_{s}^{t,x,u,N}, Z_{s}^{t,x,u,N}, u_{s})d\langle B^{i}, B^{j}\rangle_{s} - Z_{s}^{t,x,u,N}dB_{s} - dK_{s}^{t,x,u,N}, \\ Y_{T}^{t,x,u,N} &= \Phi(X_{T}^{t,x,u,N}), \quad s \in [t,T]. \end{aligned}$$

We define

$$V^{N}(t,x) = \inf_{u(\cdot) \in \mathbb{U}^{t}[t,T]} Y_{t}^{t,x,u,N}.$$

By Lemma 22, we get for any $t < s \leq T, x \in \mathbb{R}^n$,

$$V^{N}(t,x) \leq \inf_{u(\cdot) \in \mathbb{U}^{t}[t,s]} \mathbb{G}^{t,x,u,N}_{t,s}[V^{N}(s,X^{t,x,u,N}_{s})],$$
(12)

where $\mathbb{G}_{t,s}^{t,x,u,N}[\cdot]$ is defined as in $\mathbb{G}_{t,s}^{t,x,u}[\cdot]$. By Theorem 5, there exists a constant $C_1 > 0$ depending on T, n, U, G and C such that for any $u(\cdot) \in \mathbb{U}^t[t,T], r \in [t,T]$,

$$\hat{\mathbb{E}}[|X_r^{t,x,u}|^4] \le C_1(1+|x|^4),$$

$$\begin{split} \hat{\mathbb{E}}[|X_{r}^{t,x,u,N} - X_{r}^{t,x,u}|^{2}] &\leq C_{1}\hat{\mathbb{E}}[\int_{t}^{T}(|b - b^{N}|^{2} + |h_{ij} - h_{ij}^{N}|^{2} + |\sigma - \sigma^{N}|^{2})(s, X_{s}^{t,x,u}, u)ds] \\ &\leq C_{1}\hat{\mathbb{E}}[\int_{t}^{T}\frac{1}{N^{2}}(|b|^{4} + |h_{ij}|^{4} + |\sigma|^{4})(s, X_{s}^{t,x,u}, u)ds] \\ &\leq \frac{C_{2}(1 + |x|^{4})}{N^{2}}, \end{split}$$

where C_2 depending on T, n, U, G and C. By Theorem 7, there exists a constant $C_3 > 0$ depending on T, G and C such that for any $u(\cdot) \in \mathbb{U}^t[t, T]$,

$$\begin{aligned} |Y_t^{t,x,u,N} - Y_t^{t,x,u}|^2 &\leq C_3 \hat{\mathbb{E}}[|\Phi(X_T^{t,x,u,N}) - \Phi(X_T^{t,x,u})|^2 + \int_t^T |X_r^{t,x,u,N} - X_r^{t,x,u}|^2 dr] \\ &\leq \frac{C_4(1+|x|^4)}{N^2}, \end{aligned}$$

where C_4 depending on T, n, U, G and C. Thus we get

$$|V^{N}(t,x) - V(t,x)| \le \sup_{u(\cdot) \in \mathbb{U}^{t}[t,T]} |Y_{t}^{t,x,u,N} - Y_{t}^{t,x,u}| \le \frac{\sqrt{C_{4}(1+|x|^{4})}}{N}.$$

It is easy to verify that Lemma 18 still holds for V^N . Then we can get

$$\begin{aligned} &|V^{N}(s, X^{t,x,u,N}_{s}) - V(s, X^{t,x,u}_{s})|^{2} \\ &\leq 2(|V^{N}(s, X^{t,x,u,N}_{s}) - V^{N}(s, X^{t,x,u}_{s})|^{2} + |V^{N}(s, X^{t,x,u}_{s}) - V(s, X^{t,x,u}_{s})|^{2}) \\ &\leq 2(L_{2}^{2}|X^{t,x,u,N}_{s} - X^{t,x,u}_{s}|^{2} + \frac{C_{4}(1 + |X^{t,x,u}_{s}|^{4})}{N^{2}}). \end{aligned}$$

Similar to the proof of Lemma 22, we can obtain for any $u(\cdot) \in \mathbb{U}^t[t, s]$,

$$|\mathbb{G}_{t,s}^{t,x,u,N}[V^{N}(s,X_{s}^{t,x,u,N})] - \mathbb{G}_{t,s}^{t,x,u}[V(s,X_{s}^{t,x,u})]|^{2} \\ \leq C_{3}\hat{\mathbb{E}}[|V^{N}(s,X_{s}^{t,x,u,N}) - V(s,X_{s}^{t,x,u})|^{2} + \int_{t}^{s} |X_{r}^{t,x,u,N} - X_{r}^{t,x,u}|^{2}dr].$$

Thus

$$|\inf_{u(\cdot)\in\mathbb{U}^t[t,s]}\mathbb{G}^{t,x,u,N}_{t,s}[V^N(s,X^{t,x,u,N}_s)] - \inf_{u(\cdot)\in\mathbb{U}^t[t,s]}\mathbb{G}^{t,x,u}_{t,s}[V(s,X^{t,x,u}_s)]| \le \frac{\sqrt{C_5(1+|x|^4)}}{N},$$

where C_5 depending on T, n, U, G and C. Taking $N \to \infty$ in inequality (12), we obtain the result. The proof is complete.

Now we give the proof of Theorem 21:

Proof. (1) By Lemma 18, we have for any $u(\cdot) \in \mathcal{U}[t, T]$,

$$Y_s^{s, X_s^{t, x, u}, u} \ge V(s, X_s^{t, x, u})$$
q.s.,

where $Y_s^{s,X_s^{t,x,u},u} = Y_s^{t,x,u}$ is the solution of equation (9) at time s. Then, by the comparison theorem of G-BSDE, we obtain

$$Y_t^{t,x,u} \geq \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})] \text{ q.s.},$$

which leads to

$$V(t,x) \ge \underset{u(\cdot) \in \mathcal{U}[t,s]}{\operatorname{ess\,inf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})]$$

- (2) Now we prove the converse inequality.
- By Theorem 17, we get

$$\begin{split} \underset{u(\cdot)\in\mathcal{U}[t,s]}{\operatorname{ess\,inf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})] &= \underset{u(\cdot)\in\mathcal{U}^t[t,s]}{\operatorname{inf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})] \\ &= \underset{u(\cdot)\in\mathbb{U}^t[t,s]}{\operatorname{inf}} \mathbb{G}_{t,s}^{t,x,u}[V(s,X_s^{t,x,u})]. \end{split}$$

By Lemma 24, we obtain

$$V(t,x) \leq \inf_{u(\cdot) \in \mathbb{U}^t[t,s]} \mathbb{G}^{t,x,u}_{t,s}[V(s,X^{t,x,u}_s)].$$

This completes the proof. \blacksquare

The following lemma shows the continuity of V in t.

Lemma 25 The value function V is $\frac{1}{2}$ Hölder continuous in t.

Proof. Set $(t, x) \in \mathbb{R}^n \times [0, T]$ and $\delta > 0$. By dynamic programming principle, we have

$$V(t,x) = \inf_{u(\cdot) \in \mathcal{U}^t[t,t+\delta]} \mathbb{G}_{t,t+\delta}^{t,x,u}[V(t+\delta,X_{t+\delta}^{t,x,u})].$$

Set $\overline{f} = \overline{g}_{ij} = 0$, it is easy to verify that $(V(t + \delta, x), 0, 0)_{s \in [t,t+\delta]}$ is the solution of *G*-BSDE (9) with terminal condition $\overline{Y}_{t+\delta} = V(t+\delta, x)$. Thus by Proposition 5.1 in [12], there exists a constant $C_1 > 0$ depending on *T*, *G* and *C* such that for any $u(\cdot) \in \mathcal{U}^t[t, t+\delta]$,

$$\begin{aligned} &|\mathbb{G}_{t,t+\delta}^{t,x,u}[V(t+\delta,X_{t+\delta}^{t,x,u})] - V(t+\delta,x)|^2 \\ &\leq C_1 \hat{\mathbb{E}}[|V(t+\delta,X_{t+\delta}^{t,x,u}) - V(t+\delta,x)|^2 + (\int_t^{t+\delta} (1+|X_s^{t,x,u}| + |V(t+\delta,x)|)ds)^2]. \end{aligned}$$

By Lemmas 18 and 19, we can get

$$\begin{split} &|\mathbb{G}_{t,t+\delta}^{t,x,u}[V(t+\delta,X_{t+\delta}^{t,x,u})] - V(t+\delta,x)|^2 \\ &\leq C_2 \hat{\mathbb{E}}[(1+|x|^2)\delta^2 + \delta \int_t^{t+\delta} |X_s^{t,x,u}|^2 ds + |X_{t+\delta}^{t,x,u} - x|^2], \end{split}$$

where C_2 depending on T, G and C. By Theorem 5, there exists a constant $C_3 > 0$ depending on T, n, U, G and C such that for any $u(\cdot) \in \mathcal{U}^t[t, t+\delta]$,

$$\hat{\mathbb{E}}[|X_s^{t,x,u}|^2] \le C_3(1+|x|^2), \ \hat{\mathbb{E}}[|X_{t+\delta}^{t,x,u}-x|^2] \le C_3(1+|x|^2)\delta.$$

Then we obtain

$$|\mathbb{G}_{t,t+\delta}^{t,x,u}[V(t+\delta, X_{t+\delta}^{t,x,u})] - V(t+\delta, x)|^2 \le C_4(1+|x|^2)\delta,$$

where C_4 depending on T, n, U, G and C. Thus

$$\begin{aligned} |V(t,x) - V(t+\delta,x)| &\leq \sup_{u(\cdot) \in \mathcal{U}^t[t,t+\delta]} |\mathbb{G}^{t,x,u}_{t,t+\delta}[V(t+\delta,X^{t,x,u}_{t+\delta})] - V(t+\delta,x)| \\ &\leq \sqrt{C_4(1+|x|^2)} \delta^{\frac{1}{2}}. \end{aligned}$$

The proof is complete.

6 The viscosity solution of HJB equation

The following theorem gives the relationship between the value function V and the second-order partial differential equation (13).

Theorem 26 Let Assumptions (A1) and (A2) hold. V is the value function defined by (7). Then V is the unique viscosity solution of the following second-order partial differential equation:

$$\partial_t V(t,x) + \inf_{u \in U} H(t,x,V,\partial_x V,\partial_{xx}^2 V,u) = 0,$$
(13)
$$V(T,x) = \Phi(x), \qquad x \in \mathbb{R}^n,$$

where

$$\begin{array}{ll} H(t,x,v,p,A,u) = & G(F(t,x,v,p,A,u)) + \langle p, b(t,x,u) \rangle + f(t,x,v,\sigma^{T}(t,x,u)p,u), \\ F_{ij}(t,x,v,p,A,u) = & (\sigma^{T}(t,x,u)A\sigma(t,x,u))_{ij} + 2\langle p, h_{ij}(t,x,u) \rangle \\ & + 2g_{ij}(t,x,v,\sigma^{T}(t,x,u)p,u), \end{array}$$

 $(t, x, v, p, A, u) \in [0, T] \times \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{S}_n \times U, G \text{ is defined by equation (3).}$

For simplicity, we only consider the case $h_{ij} = g_{ij} = 0$. Suppose $\varphi \in C^{2,3}_{b,Lip}([0,T] \times \mathbb{R}^n)$. Define

$$F_{1}(s, x, y, z, u) = \partial_{s}\varphi(s, x) + f(s, x, y + \varphi(s, x), z + \sigma^{T}(s, x, u)\partial_{x}\varphi(s, x), u) + \langle b(s, x, u), \partial_{x}\varphi(s, x) \rangle, F_{2}^{ij}(s, x, u) = \frac{1}{2} \langle \partial_{xx}^{2}\varphi(s, x)\sigma_{i}(s, x, u), \sigma_{j}(s, x, u) \rangle.$$
(14)

Consider the following G-BSDEs: $\forall s \in [t, t + \delta]$,

$$Y_{s}^{1,u} = \int_{s}^{t+\delta} F_{1}(r, X_{r}^{t,x,u}, Y_{r}^{1,u}, Z_{r}^{1,u}, u_{r}) dr + \int_{s}^{t+\delta} F_{2}^{ij}(r, X_{r}^{t,x,u}, u_{r}) d\langle B^{i}, B^{j} \rangle_{r} - \int_{s}^{t+\delta} Z_{r}^{1,u} dB_{r} - (K_{t+\delta}^{1,u} - K_{s}^{1,u}),$$
(15)

and

$$Y_{s}^{u} = \varphi(t+\delta, X_{t+\delta}^{t,x,u}) + \int_{s}^{t+\delta} f(r, X_{r}^{t,x,u}, Y_{r}^{u}, Z_{r}^{u}, u_{r}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - \int_{s}^{t+\delta} Z_{r}^{u} dB_{r} - (K_{t+\delta}^{u} - K_{s}^{u}) dr - (K_{t+\delta}^{u} - K_{t+\delta}^{u}) dr - (K_{t+\delta}^{$$

Lemma 27 For each $s \in [t, t + \delta]$, we have

$$Y_{s}^{1,u} = Y_{s}^{u} - \varphi(s, X_{s}^{t,x,u}).$$
(17)

Proof. Applying Itô's formula to $\varphi(s, X_s^{t,x,u})$, we have

$$d(Y_s^u - \varphi(s, X_s^{t,x,u})) = dY_s^{1,u}$$

Since $Y_{t+\delta}^u - \varphi(t+\delta, X_{t+\delta}^{t,x,u}) = Y_{t+\delta}^{1,u} = 0$, we obtain

$$Y_s^{1,u} = Y_s^u - \varphi(s, X_s^{t,x,u}), \quad \forall s \in [t, t+\delta].$$

The proof is completed. $\hfill\blacksquare$

Consider the G-BSDE: $\forall s \in [t, t + \delta]$,

$$Y_{s}^{2,u} = \int_{s}^{t+\delta} F_{1}(r, x, Y_{r}^{2,u}, Z_{r}^{2,u}, u_{r}) dr + \int_{s}^{t+\delta} F_{2}^{ij}(r, x, u_{r}) d\langle B^{i}, B^{j} \rangle_{r}$$

$$- \int_{s}^{t+\delta} Z_{r}^{2,u} dB_{r} - (K_{t+\delta}^{2,u} - K_{s}^{2,u}).$$
(18)

We have the following estimates.

Lemma 28 For each $u(\cdot) \in \mathcal{U}^t[t, t+\delta]$, we have

$$Y_t^{1,u} - Y_t^{2,u} \mid \le L_4 \delta^{3/2}, \tag{19}$$

where L_4 is a positive constant dependent on x and independent of $u(\cdot)$.

Proof. By Proposition 5.1 in [12], there exists a constant $C_1 > 0$ depending on T, G and C such that for any $u(\cdot) \in \mathcal{U}^t[t, t+\delta]$,

$$|Y_t^{1,u} - Y_t^{2,u}|^2 \le C_1 \hat{\mathbb{E}}[(\int_t^{t+\delta} \hat{F}_r dr)^2],$$

where

$$\hat{F}_{r} = |F_{1}(r, X_{r}^{t,x,u}, Y_{r}^{2,u}, Z_{r}^{2,u}, u_{r}) - F_{1}(r, x, Y_{r}^{2,u}, Z_{r}^{2,u}, u_{r})| + \sum_{i,j=1}^{d} |F_{2}^{ij}(r, X_{r}^{t,x,u}, u_{r}) - F_{2}^{ij}(r, x, u_{r})|.$$

Note that $\varphi \in C^{2,3}_{b,Lip}([0,T] \times \mathbb{R}^n)$, it is easy to verify that

$$\hat{F}_r \le C_2(|X_r^{t,x,u} - x| + |X_r^{t,x,u} - x|^2),$$

where C_2 is dependent on x and independent of $u(\cdot)$. By Theorem 5, we can obtain that for any $p \ge 2$,

$$\hat{\mathbb{E}}[\sup_{r \in [t,t+\delta]} | X_r^{t,x,u} - x |^p] \le C_3 (1 + |x|^p) \delta^{p/2},$$

where C_3 is independent of $u(\cdot)$. Then by Hölder's inequality we can deduce that $|Y_t^{1,u} - Y_t^{2,u}| \le L_4 \delta^{3/2}$, where L_4 is dependent on x and independent of $u(\cdot)$. This completes the proof. Now we compute $\inf_{u(\cdot) \in \mathcal{U}^t[t,t+\delta]} Y_t^{2,u}$.

Lemma 29 We have

$$\inf_{u(\cdot)\in\mathcal{U}^t[t,t+\delta]}\,Y^{2,u}_t=Y^0_t,$$

where Y^0 is the solution of the following ordinary differential equation

$$-dY_s^0 = F_0(s, x, Y_s^0, 0)ds, \quad Y_{t+\delta}^0 = 0, \ s \in [t, t+\delta]$$
(20)

and

$$F_0(s, x, y, z) := \inf_{u \in U} [F_1(s, x, y, z, u) + 2G(F_2(s, x, u))].$$

In order to prove Lemma 29, we need the following property of the decreasing G-martingale.

Lemma 30 Suppose that $(M_s)_{t \leq s \leq t+\delta}$ is a decreasing *G*-martingale. Then there exists a $Q \in \mathcal{P}$ such that

$$M_{t+\delta} = M_t, \ Q - a.s.$$

Proof. By the representation of *G*-expectation, we know that

$$\hat{\mathbb{E}}[M_{t+\delta} - M_t] = \sup_{P \in \mathcal{P}} E_P[M_{t+\delta} - M_t].$$

Thus there exist $Q_k \in \mathcal{P}, k = 1, 2, ...,$ such that

$$E_{Q_k}[M_{t+\delta} - M_t] \uparrow \hat{\mathbb{E}}[M_{t+\delta} - M_t] = \hat{\mathbb{E}}[\hat{\mathbb{E}}_t[M_{t+\delta}] - M_t] = 0.$$

Since \mathcal{P} is weakly compact, there exist $Q \in \mathcal{P}$ and a subsequence (Q_{k_i}) of (Q_k) such that Q_{k_i} converges weakly to Q. By Lemma 29 in [6], then we get

$$E_Q[M_{t+\delta} - M_t] = \lim_{i \to \infty} E_{Q_{k_i}}[M_{t+\delta} - M_t] = 0.$$

Note that $M_{t+\delta} - M_t \leq 0$, q.s.. Thus, we obtain that

$$M_{t+\delta} = M_t, \ Q - a.s..$$

This completes the proof. \blacksquare

Now we give the proof of Lemma 29. **Proof.** (1) We first prove that for any $u(\cdot) \in \mathcal{U}^t[t, t+\delta]$,

$$Y_t^{2,u} \ge Y_t^0.$$

Note that

$$\begin{split} Y^{2,u}_s = & \int_s^{t+\delta} [F_1(r,x,Y^{2,u}_r,Z^{2,u}_r,u_r) + 2G(F_2(r,x,u_r))] dr \\ & + [\int_s^{t+\delta} F_2^{ij}(r,x,u_r) d\langle B^i,B^j\rangle_r - \int_s^{t+\delta} 2G(F_2(r,x,u_r)) dr] \\ & - \int_s^{t+\delta} Z^{2,u}_r dB_r - (K^{2,u}_{t+\delta} - K^{2,u}_s), \text{ q.s..} \end{split}$$

It is easy to verify that

$$M_{s} = \int_{t}^{s} F_{2}^{ij}(r, x, u_{r}) d\langle B^{i}, B^{j} \rangle_{r} - \int_{t}^{s} 2G(F_{2}(r, x, u_{r})) dr$$

is a decreasing G-martingale.

By Lemma 30, there exists a $Q \in \mathcal{P}$ such that for $s \in [t, t + \delta]$,

$$M_s = 0, \ Q - a.s..$$

Then

$$\begin{split} Y^{2,u}_s = & \int_s^{t+\delta} [F_1(r,x,Y^{2,u}_r,Z^{2,u}_r,u_r) + 2G(F_2(r,x,u_r))] dr \\ & - \int_s^{t+\delta} Z^{2,u}_r dB_r - (K^{2,u}_{t+\delta} - K^{2,u}_s), \; Q-\text{a.s.}. \end{split}$$

Consider the following BSDE: for $s \in [t, t + \delta]$,

$$Y_s^0 = \int_s^{t+\delta} F_0(r, x, Y_r^0, Z_r^0) dr - \int_s^{t+\delta} Z_r^0 dB_r - (K_{t+\delta}^0 - K_s^0), \ Q - \text{a.s.}.$$

Since F_0 is a deterministic function, we obtain that $Z_s^0 = 0$, $K_s^0 = 0$ and Y_s^0 is just the solution of equation (20). Note that $(-K_s^{2,u})_{s \in [t,t+\delta]}$ is a increasing process and $F_1(r, x, y, z, u_r) + 2G(F_2(r, x, u_r)) \ge F_0(r, x, y, z)$, then by the comparison theorem of classical BSDE (under the reference probability measure Q), we deduce that

$$Y_t^{2,u} \ge Y_t^0$$

(2) We denote the class of all deterministic controls in $\mathcal{U}^t[t, t+\delta]$ by \mathcal{U}_1 . Then, for every $u(\cdot) \in \mathcal{U}_1$, $Y^{2,u}$ is the solution of the following ordinary differential equation:

$$-dY_s^{2,u} = [F_1(s, x, Y_s^{2,u}, 0, u_s) + 2G(F_2(s, x, u_s))]ds, \quad s \in [t, t+\delta],$$

$$Y_{t+\delta}^{2,u} = 0.$$

It is easy to check that

$$Y^0_t = \inf_{u(\cdot) \in \mathcal{U}_1} Y^{2,u}_t \geq \inf_{u(\cdot) \in \mathcal{U}^t[t,t+\delta]} Y^{2,u}_t.$$

This completes the proof. \blacksquare

Finally we give the proof of Theorem 26.

Proof. The uniqueness of viscosity solution of equation (13) can be proved similarly as in Theorem 6.1 in [1], we only prove that V is a viscosity solution of equation (13). By Lemmas 18 and 25, V is a continuous functions on $[0, T] \times \mathbb{R}^n$. We first prove that V is the subsolution of (13).

We first prove that V is the subsolution of (13). Given $t \leq T$ and $x \in \mathbb{R}^n$, suppose $\varphi \in C^{2,3}_{b,Lip}([0,T] \times \mathbb{R}^n)$ such that $\varphi(t,x) = V(t,x)$ and $\varphi \geq V$ on $[0,T] \times \mathbb{R}^n$. By Theorem 21, we have

$$V(t,x) = \inf_{u(\cdot) \in \mathcal{U}^t[t,t+\delta]} \mathbb{G}^{t,x,u}_{t,t+\delta}[V(t+\delta,X^{t,x,u}_{t+\delta})].$$

Note that $\varphi \geq V$ on $[0,T] \times \mathbb{R}^n$. Then by comparison theorem, we get

$$\inf_{\iota(\cdot)\in\mathcal{U}^t[t,t+\delta]} \left\{ \mathbb{G}^{t,x,u}_{t,t+\delta}[\varphi(t+\delta,X^{t,x,u}_{t+\delta})] - \varphi(t,x) \right\} \ge 0.$$

By equality (17), we have

$$\inf_{u(\cdot)\in\mathcal{U}^t[t,t+\delta]}Y_t^{1,u}\ge 0.$$

By inequality (19) and Lemma 29, we get

$$\inf_{u(\cdot)\in\mathcal{U}^t[t,t+\delta]}Y_t^{2,u} \ge -L_4\delta^{3/2}$$

and

$$Y_t^0 \ge -L_4 \delta^{3/2}$$

Thus

$$-L_4 \delta^{1/2} \le \delta^{-1} Y_t^0 = \delta^{-1} \int_t^{t+\delta} F_0(r, x, Y_r^0, 0) dr.$$

Letting $\delta \to 0$, we get $F_0(t, x, 0, 0) = \inf_{u \in U} (F_1(t, x, 0, 0, u) + G(F_2(t, x, u))) \ge 0$, which implies that V is a subsolution of (13). Using the same method, we can prove V is the supersolution of (13). This completes the proof.

Acknowledgments

We would like to thank S. Peng for many helpful discussions.

References

- R. Buckdahn and J. Li, Stochastic differential games and viscosity solutions for Hamilton-Jacobi-Bellman-Isaacs equations, SIAM J Control Optim., 47(2008), pp. 444–475.
- [2] R. Buckdahna, B. Labedb, C. Rainera and L. Tamer, Existence of an optimal control for stochastic control systems with nonlinear cost functional, Stochastics, An International Journal of Probability and Stochastics Processes, 82(2010), pp. 241–256.
- [3] Z. Chen and L. G. Epstein, Ambiguity, risk, and asset returns in continuous time, Econometrica 70(2002), pp. 1403–43.
- [4] L. Denis and M. Kervarec, Optimal investment under model uncertainty in non-dominated models, SIAM J Control Optim., 51(2013), pp. 1803-1822.
- [5] L. Denis and C. Martini, A theoretical framework for the pricing of contingent claims in the presence of model uncertainty, The Annals of Applied Probability, 16(2006), pp. 827-852.
- [6] L. Denis, M. Hu and S. Peng, Function spaces and capacity related to a sublinear expectation: application to G-Brownian motion paths, Potential Anal., 34(2011), pp. 139-161.
- [7] D. Duffie and L. Epstein, Stochastic differential utility, Econometrica, 60(1992), pp. 353–394.
- [8] N. El Karoui, S. Peng and M. C. Quenez, Backward stochastic differential equations in finance, Math. Finance, 7(1997), pp. 1-71.
- [9] L. Epstein and S. Ji, Ambiguous volatility, possibility and utility in continuous time, Journal of Mathematical Economics, 50(2014), pp. 269-282.
- [10] L. Epstein and S. Ji, Ambiguous volatility and asset pricing in continuous time, Rev. Finan. Stud., 26 (2013), pp. 1740-1786.

- [11] W. H. Fleming, and H. M. Soner, Control Markov processes and viscosity solutions (New York:Springer Verlag).
- [12] M. Hu, S. Ji, S. Peng and Y. Song, Backward stochastic differential equations driven by G-Brownian motion, Stochastic Processes and their Applications, 124(2014), pp. 759-784.
- [13] M. Hu, S. Ji, S. Peng and Y. Song, Comparison theorem, Feynman-Kac formula and Girsanov transformation for BSDEs driven by G-Brownian motion, Stochastic Processes and their Applications, 124(2014), pp. 1170-1195.
- [14] M. Hu and S. Peng, On representation theorem of G-expectations and paths of G-Brownian motion, Acta Math. Appl. Sin. Engl. Ser., 25(2009), pp. 539-546.
- [15] M. Hu, S. Ji and S. Yang, A stochastic recursive optimal control problem under the G-expectation framework, Appl. Math. Optim. 70(2014), pp. 253– 278.
- [16] M. Hu, F. Wang and G. Zheng, Quasi-continuous random variables and processes under the G-expectation framework, (2014), arXiv:1088060.
- [17] A. Matoussi, D. Possamai and C. Zhou, Robust utility maximization in nondominated models with 2BSDE: the uncertain volatility model, Mathematical Finance, Article first published online: 18 JUN 2013, DOI: 10.1111/mafi.12031.
- [18] S. Peng, Backward stochastic differential equation and application to optimal control, Applied Mathematics and Optimization, 27 (1993), pp. 125-144.
- [19] S. Peng, G-expectation, G-Brownian motion and related stochastic calculus of Itô type, Stochastic analysis and applications, Abel Symp., 2, Springer, Berlin, (2007), pp. 541-567.
- [20] S. Peng, G-Brownian motion and dynamic risk measure under volatility uncertainty, (2007), arXiv:0711.2834v1.
- [21] S. Peng, Multi-dimensional G-Brownian motion and related stochastic calculus under G-expectation, Stochastic Processes and their Applications, 118(2008), pp. 2223-2253.
- [22] S. Peng, Nonlinear expectations and stochastic calculus under uncertainty, (2010), arXiv:1002.4546v1.
- [23] S. Peng, A generalized dynamic programming principle and Hamilton-Jacobi-Bellmen equation, Stochastics Stochastics Rep., 38(1992), pp. 119– 134.

- [24] S. Peng, Backward stochastic differential equations—stochastic optimization theory and viscosity solutions of HJB equations, in Topics on Stochastic Analysis, J. Yan, S. Peng, S. Fang, and L. Wu, eds., Science Press, Beijing, 1997, pp. 85–138 (in Chinese).
- [25] T. Pham and J. Zhang, Two person zero-sum game in weak formulation and path dependent Bellman-Isaacs equation, SIAM J Control Optim., 52(2014), pp. 2090-2121.
- [26] H. M. Soner, N. Touzi and J. Zhang, Wellposedness of second order backward SDEs, Probability Theory and Related Fields, 153(2012), pp. 149-190.
- [27] R. Tevzadze, T. Toronjadze, and T. Uzunashvili, Robust utility maximization for a diffusion market model with misspecified coefficients, Finance and Stochastics, (2013), pp. 535-563.
- [28] J. Yong and X. Y. Zhou, Stochastic controls: Hamiltonian systems and HJB equations, 1999.