Optimal dividend payments under a time of ruin constraint: Exponential claims

Camilo Hernández^{*} Mauricio Junca[†]

March 2, 2022

Abstract

We consider the classical optimal dividends problem under the Cramér-Lundberg model with exponential claim sizes subject to a constraint on the time of ruin. We introduce the dual problem and show that the complementary slackness conditions are satisfied, thus there is no duality gap. Therefore the optimal value function can be obtained as the point-wise infimum of auxiliary value functions indexed by Lagrange multipliers. We also present a series of numerical examples.

1 Introduction

One of the most studied models in actuarial science to describe the reserves process of an insurance company is the Cramér-Lundberg model. In this model the company faces claims whose arrivals follow a compound Poisson process and a constant premium is paid by the insured clients.

After the model was introduced, the probability of ruin of such a portfolio was among the principal interests in this field, see [1]. Nowadays, results about minimizing ruin probability considering reinsurance and investment in risky assets are proved by [8]. In [9], similar results for a discrete time version of the model and a diffusion approximation are shown. However, a process that does not end in ruin in a model exceeds every finite level, this is, the company lives an infinite period of time and accumulates an infinite amount of money, which is quite unrealistic in practice.

This idea motivated the study of the performance, instead of the safety aspect, of such portfolio. In 1957, Bruno de Finetti was interested in finding a way of paying out dividends in order to optimise the expected present value of the total income of the shareholders from time zero until ruin. This problem is commonly referred as de Finetti's problem, see [4]. As a result, researchers addressed the optimality aspect under more general and realistic assumptions which has turned out to be an abundant and ambitious field of research. See [9] for results on this problem in the Cramér-Lundberg model and its diffusion approximation, and [2] when reinsurance is also considered.

Under exponential claims, see [9], the optimal dividends payment strategy to de Finetti's problem is known to be a *barrier strategy*, this is, there exists a value b such that the optimal strategy

^{*}Department of Mathematics, Universidad de los Andes, Bogota, Colombia. Email address: mc.hernandez131@uniandes.edu.co

[†]Department of Mathematics, Universidad de los Andes, Bogota, Colombia. Email address: mj.junca20@uniandes.edu.co

is to let $D_t = [\bar{X}_t - b] \vee 0$ where \bar{X}_t denotes the maximum of the reserves process X up to time t. However, the solution is not necessarily of this type in the general set up. In [2] the authors provided an example of claims distribution for which no barrier strategy is optimal.

However, there exists a trade-off between stability and profitability. Minimizing ruin probability means no dividends payment and profits tend to 0. On the contrary, maximizing the dividends leads to a situation for which ruin is certain regardless of the initial capital, see [3].

The idea of this work is to study a way to link two key concepts in the optimal dividends payment theory for the Cramér-Lundberg model: the profits and the time of ruin derived from a dividends payment strategy. Approaches in this direction have been already considered, see [6] for a solution to the optimal dividends payment problem under a ruin constraint in discrete time and state setting. In [7] the author addressed this matter introducing the concept of *solvency constraints* and studied how they affected the optimal level of the barrier. In [10] the authors consider the classical and diffusion approximation models and introduced a penalization on the time of ruin on the objective function, however an actual restriction on the time of ruin was not stated in this work. Finally, the model introduced in [5] took this link into account but in a different setting. More specifically, [5] proposed an iterative scheme to solve the problem of maximize the expected discounted consumption of an investor in finite time plus a penalisation on the level of the reserves at ruin for a given upper bound for the ruin probability.

This paper is organized as follows: Section 2 is dedicated to the formulation of the problem under consideration and to rewrite it using duality theory. In the following sections, under the exponential claim sizes assumption, we present the main results of this paper. We first focus on solving the dual problem in Section 3 and then show the absence of duality gap for this problem in Section 4. This paper's contribution relies on both the solution of the optimal dividend payment problem under a constrain on the time of ruin, and the tools developed in order to prove the duality gap of this problem to be zero. Section 5 is dedicated to the presentation of numerical examples that illustrate different scenarios of the solution. In the last section we give conclusions of this study and present directions in which this work can be continued.

2 Problem formulation

In this paper we consider the Cramér-Lundberg risk model, in which the surplus follows the process:

$$X_t = x_0 + ct - \sum_{i=1}^{N_t} Y_i,$$

where $N = (N_t)_{t\geq 0}$ represents a homogeneous Poisson process with rate $\lambda > 0$, modeling the claim occurrences. $\{Y_i\}$ models the sequence of claim amounts, $\{Y_i\} \stackrel{iid}{\sim} G(y)$, with G(.) is a continuous distribution function on $[0, \infty)$. $\{Y_i\}$ is assumed to be independent of the claim occurrences process N. The deterministic components are the premium rate c > 0 and the initial capital x_0 . All of the above are defined in the same filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$, with $(\mathcal{F}_t)_{t\geq 0}$ the filtration generated by the process X.

The insurance company is allowed to pay dividends which are model by the process $D = (D_t)_{t\geq 0}$ representing the cumulative payments up to time t. A dividends process is called admissible if it is a non negative, non decreasing càdlàg process adapted to the filtration $(\mathcal{F}_t)_{t\geq 0}$. Therefore, the surplus process under dividends process D reads as

$$X_t^D = x_0 + ct - \sum_{i=1}^{N_t} Y_i - D_t.$$
 (1)

Let τ^D denotes the time of ruin under dividends process D, i.e., $\tau^D = \inf\{t \ge 0 : X_t^D < 0\}$. We require the dividends process not to lead to ruin, i.e., $D_{t-} - D_t \le X_t^D$ for all t and $D_t = D_{\tau^D}$ for $t \ge \tau^D$, so no dividends are paid after ruin. We call Θ the set of such processes.

The company wants to maximise the expected value of the discounted flow of dividends payment along time, that is, to maximise

$$\mathcal{V}^D(x_0) := \mathbb{E}_{x_0} \Big[\int_0^{\tau^D} e^{-\delta t} dD_t \Big],$$

where the lifespan of the company will be determined by its ruin, see [10]. Also, δ is the discount factor.

Finally, we add a restriction on the dividends process D which we model by the equation:

$$\mathbb{E}_{x_0}\left[\int_0^{\tau^D} e^{-\delta s} ds\right] \ge \int_0^T e^{-\delta s} ds \quad T \ge 0 \text{ fixed.}$$
(2)

The motivation behind such a constraint is that it imposes a restriction on the time of ruin. For simplicity, let's denote the right hand side of the restriction by K_T , i.e., $K_T := \int_0^T e^{-\delta s} ds$. Note that $K_T \in [0, \frac{1}{\delta})$ and that the greater K_T the greater τ^D must be. Evidently there are other restrictions that capture this effect in a more direct way, e.g., $\mathbb{E}[\tau^D] \ge K$. However there is no, to the best of our knowledge, a satisfactory technique to introduce such type of constraint in the model; additionally, as it will be clear in the following sections, the chosen functional form of the constraint fits in with the model in a smooth way.

Combining all the above components we state the problem we aim to solve:

$$V(x_0) := \sup_{D \in \Theta} \quad \mathcal{V}^D(x_0) \quad \text{s.t.} \quad \mathbb{E}_{x_0} \left[\int_0^{\tau^D} e^{-\delta s} ds \right] \ge K_T \quad T \text{ fixed.} \tag{P1}$$

In order to solve this problem we use Lagrange multipliers to reformulate our problem. We first define the following for $\Lambda \ge 0$

$$\mathcal{V}_{\Lambda}^{D}(x_{0}) := \mathbb{E}_{x_{0}} \left[\int_{0}^{\tau^{D}} e^{-\delta t} dD_{t} + \Lambda \int_{0}^{\tau^{D}} e^{-\delta s} ds \right] - \Lambda K_{T}.$$
(3)

The following remark clears out the strategy we will follow in the remainder of the paper.

Remark 2.1. • Note that (P1) is equivalent to $\sup_{D \in \Theta} \inf_{\Lambda \ge 0} \mathcal{V}^D_{\Lambda}(x_0)$ since

$$\inf_{\Lambda \ge 0} \mathcal{V}^{D}_{\Lambda}(x_{0}) = \begin{cases} \mathcal{V}^{D}(x_{0}) & \text{if } \mathbb{E}_{x_{0}} \left[\int_{0}^{\tau^{D}} e^{-\delta s} ds \right] \ge K_{T} \\ -\infty & \text{otherwise} \end{cases}$$

• The dual problem of (P1), is defined as

$$\inf_{\Lambda \ge 0} \sup_{D \in \Theta} \mathcal{V}^{D}_{\Lambda}(x_{0}), \tag{D}$$

which is always an upper bound for the primal (P1). The main goal is to prove that $\sup_{D \in \Theta} \inf_{\Lambda \ge 0} \mathcal{V}^D_{\Lambda}(x_0) = \inf_{\Lambda \ge 0} \sup_{D \in \Theta} \mathcal{V}^D_{\Lambda}(x_0).$

To solve (D) we note the last term of (3) does not depend on D and is linear on Λ , therefore, we can focus on the first term on the right hand side of this equation and solve for fixed $\Lambda \geq 0$

$$V_{\Lambda}(x_0) := \sup_{D \in \Theta} \mathcal{V}^D_{\Lambda}(x_0).$$
(P2)

For this problem, it is known that its solution must satisfy the following HJB equation, see Proposition 11 in [10]

$$\max\{\Lambda + cV'(x) + \lambda \int_0^x V(x-y)dG(y) - (\lambda + \delta)V(x), 1 - V'(x)\} = 0.$$
 (4)

As a first approach to this problem, we restrict ourselves assuming an exponential distribution for the claim sizes. In this scenario, we succeed in solving (P1) via (D) proving there is no duality gap. However, we have not yet approached the general problem for an arbitrary claim sizes distribution.

From now on we assume $\{Y_i\} \stackrel{iid}{\sim} Exp(\alpha)$. As a result (4) converts into

$$\max\{\Lambda + cV'(x) + \lambda \int_0^x V(x-y)\alpha e^{-\alpha y} dy - (\lambda+\delta)V(x), 1 - V'(x)\} = 0,$$
(5)

3 Solution of (P2)

For this problem [10] proved that the optimal strategy corresponds to a barrier strategy and showed that the solution of (P2) is given by

Theorem 3.1 (Lemma 10 in [10]). Let $\Lambda \ge 0$. Then, if $\Lambda \le \frac{(\lambda+\delta)^2}{\alpha\lambda} - c$ the value function of (P2) is given by

$$V_{\Lambda}(x) = x + \frac{c + \Lambda}{\lambda + \delta} + \frac{\Lambda}{\delta}(e^{-\delta T} - 1).$$

Otherwise, there exists unique $b_{\Lambda} > 0$ such that

$$V_{\Lambda}(x) = \begin{cases} x - b_{\Lambda} + \frac{\alpha c - \lambda - \delta}{\alpha \delta} + \frac{\Lambda}{\delta} e^{-\delta T} & \text{if } x \ge b_{\Lambda} \\ C_1 e^{r_1 x} + C_2 e^{r_2 x} + \frac{\Lambda}{\delta} e^{-\delta T} & \text{if } x \le b_{\Lambda} \end{cases},$$
(6)

Where r_1, r_2 are roots of the polynomial

$$p(R) = cR^2 + (\alpha c - (\lambda + \delta))R - \alpha\delta.$$
(7)

 C_1, C_2 are given by

$$C_2 = -\frac{\alpha + r_2}{\alpha} \left\lfloor \frac{\alpha}{\alpha + r_1} C_1 + \frac{\Lambda}{\delta} \right\rfloor,$$
$$C_1 = \frac{(\alpha + r_1)(\alpha \delta + e^{r_2 b} \Lambda r_2(\alpha + r_2))}{\alpha \delta(e^{r_1 b} r_1(\alpha + r_1) - e^{r_2 b} r_2(\alpha + r_2))}.$$

and the value of the optimal barrier level b_{Λ} is given by the expression

$$(r_2 - r_1)(\alpha + r_1)(\alpha + r_2)\Lambda = -r_1 e^{-r_2 b} (r_1(\lambda + \delta) + \alpha \delta) + r_2 e^{-r_1 b} (r_2(\lambda + \delta) + \alpha \delta).$$
(8)

Let us define the critical value

$$\bar{\Lambda} = \frac{(\delta + \lambda)^2}{\alpha \lambda} - c$$

Remark 3.1. Note that for $0 \leq \Lambda \leq \overline{\Lambda}$ the optimal barrier strategy is b = 0.

The following proposition will be essential in the next section to prove the main result of this contribution.

Proposition 3.2. (i) If $\overline{\Lambda} \ge 0$ for each b > 0, there exists a unique $\Lambda > \overline{\Lambda}$ such that b is the optimal barrier dividend strategy for (P2) with Λ .

(ii) If $\overline{\Lambda} < 0$, there exists $b_0 > 0$ such that for each $b \ge b_0$, there exists a unique $\Lambda \ge 0$ such that b is the optimal barrier dividend strategy for (P2) with Λ .

Proof. Note that equation (8) defines a map $\Lambda : [0, \infty) \to \mathbb{R}$. Taking derivative with respect to b we obtain

$$\frac{d\Lambda(b)}{db} = \frac{r_1 r_2 [e^{-r_2 b} (r_1(\lambda + \delta) + \alpha \delta) - e^{-r_1 b} (r_2(\lambda + \delta) + \alpha \delta)]}{(r_2 - r_1)(\alpha + r_1)(\alpha + r_2)}$$
(9)

It can be easily shown that both roots of the characteristic polynomial are real and non zero, furthermore, one of them is positive and the other is negative. The first follows since $[\alpha c - (\lambda + \delta)]^2 + 4c\alpha\delta > 0$ and the second since $-[\alpha c - (\lambda + \delta)] + \sqrt{[\alpha c - (\lambda + \delta)]^2 + 4c\alpha\delta} > 0$, $-[\alpha c - (\lambda + \delta)] - \sqrt{[\alpha c - (\lambda + \delta)]^2 + 4c\alpha\delta} < 0$. Now, for the negative root, let's say r_2 , we have $\alpha + r_2 > 0$ (see Lemma A.1). This leaves us with the case $r_1 > 0 > r_2$, in which (9) is strictly positive as both the numerator and denominator are negative and therefore the map is injective. Furthermore, using the expressions for r_1 and r_2 in (8) we can show that

$$\Lambda(0) = \frac{(\lambda + \delta)^2 - \alpha \lambda c}{\alpha \lambda} = \bar{\Lambda}.$$
(10)

Hence, if $\Lambda(0) < 0$ and there exists $b_0 > 0$ (and solution of (??)) that satisfies (ii) since $\Lambda \to \infty$ when $b \to \infty$. (i) follows from (10).

Figure 1 shows values of b_{Λ} derived from equation (8) for $\lambda = 1$, c = 1.3, $Y_i \sim Exp(1)$, and $\delta = 0.1$. Note that this values fall in the second case of Proposition 3.2.

4 Solution of (P1)

Let X_t^b be the surplus process under dividend barrier strategy with level b denoted by D^b . Let τ^b be the time of ruin using such strategy, i.e., $\tau^b := \inf\{t : X_t^b < 0\}$. In order to find out the solution to (P1) we will need the following proposition.

Proposition 4.1. For each $x_0 \ge 0$ there exists $H_{x_0} \ge 0$ such that if $0 \le T < H_{x_0}$ there exists (Λ^*, b^*) that satisfies:

(i) $\Lambda^* \geq 0$ and b^* is the optimal barrier for (P2) with Λ^* and initial value x_0 ,



Figure 1: Optimal barrier

(*ii*)
$$\mathbb{E}_{x_0} \left[\int_0^{\tau^{b^*}} e^{-\delta s} ds \right] \ge K_T$$
 and
(*iii*) $\Lambda^* \left(\mathbb{E}_{x_0} \left[\int_0^{\tau^{b^*}} e^{-\delta s} ds \right] - K_T \right) = 0.$

Proof. Let $x_0 \ge 0$ fixed. Consider the following IDE problem:

$$\mathcal{A}(\Psi_b)(x) - \delta \Psi_b(x) = -1 \tag{A1}$$
$$\Psi'_b(b) = 0$$
$$\Psi_b \in C^1[0, b]$$

where $\mathcal{A}(f)(x)$ is $cf'_b(x) + \lambda \int_0^x f(x-y)\alpha e^{-\alpha y} dy - \lambda f(x)$, the infinitesimal generator of the surplus process X_t . It can be shown that for $0 \le x \le b$

$$\Psi_b(x) = \frac{1}{\delta} + C_1 e^{r_1 x} + C_2 e^{r_2 x}$$

with

$$C_1 = \frac{(\alpha + r_1)(\alpha + r_2)r_2e^{r_2b}}{\alpha\delta[r_1e^{r_1b}(\alpha + r_1) - r_2e^{r_2b}(\alpha + r_2)]}$$

and

$$C_2 = -\frac{(\alpha + r_2)^2 r_2 e^{r_2 b}}{\alpha \delta (e^{r_1 b} r_1 (\alpha + r_1) - e^{r_2 b} r_2 (\alpha + r_2))} - \frac{(\alpha + r_2)}{\alpha \delta}$$

is solution of (A1), where r_1, r_2 denote the roots of (7). Extend Ψ_b to \mathbb{R} so that $\Psi_b(x) = 0, x < 0$ and $\Psi_b(x) = \Psi_b(b), x \ge b$. Using Dynkin's formula and the Optional Stopping Theorem we obtain that for $0 \le x \le b$

$$\begin{split} \mathbb{E}_{x}[e^{-\delta\tau^{b}}\Psi_{b}(X^{b}_{\tau^{b}})] &= \Psi_{b}(x) + \mathbb{E}_{x}\bigg[\int_{0}^{\tau^{b}}e^{-\delta s}[\mathcal{A}(\Psi_{b}(X^{b}_{s})) - \delta\Psi_{b}(X^{b}_{s})]ds \\ &+ \sum_{0 \leq s \leq t, \bigtriangleup D^{b} \neq 0}e^{-\delta s}\left[\Psi_{b}(X^{b}_{s}) - \Psi_{b}(X^{b}_{s-})\right] + \int_{0}^{\tau^{b}}e^{-\delta s}\Psi_{b}'(X^{b}_{s})d\bar{D}^{b}_{s}\bigg] \\ &= \Psi_{b}(x) + \mathbb{E}_{x}\bigg[\int_{0}^{\tau^{b}}e^{-\delta s}[\mathcal{A}(\Psi_{b}(X_{s})) - \delta\Psi_{b}(X_{s})]ds\bigg] \\ &+ \mathbb{E}_{x}\bigg[\int_{0}^{\tau^{b}}e^{-\delta s}c\Psi_{b}'(b)\mathbf{1}_{\{X_{s}=b\}}ds\bigg], \end{split}$$

where \overline{D}^b denotes the continuous part of the control D^b . For the last equality we used that the continuous part of the control consists only of c at the moment at which $X_s^b = b$ and that the control has no jumps. Therefore, for $\Psi_b(x)$, the extended solution of (A1), we get that

$$\Psi_b(x) = \mathbb{E}_x \left[\int_0^{\tau^b} e^{-\delta s} ds \right].$$

Define,

$$\hat{\Psi}(x) := \lim_{b \to \infty} \Psi_b(x) = \frac{1}{\delta} - \frac{\alpha + r_2}{\alpha \delta} e^{r_2 x}.$$

Let $H_{x_0} := -\frac{1}{\delta} (\log(\frac{\alpha + r_2}{\alpha}) + r_2 x_0)$ and suppose $0 \le T < H_{x_0}$. Recall $K_T = \frac{1 - e^{-\delta T}}{\delta}$ and note that $K_{H_{x_0}} = \hat{\Psi}(x_0)$, so $K_T < \hat{\Psi}(x_0)$. Then we have the following to cases:

- 1. Suppose $\bar{\Lambda} \geq 0$. If $\Psi_0(x_0) \geq K_T$ then (ii) is satisfied and the barrier $b^* = 0$ is optimal for (P2) with $\Lambda^* = 0$ by Remark 3.1. On the other hand, if $\Psi_0(x_0) < K_T < \hat{\Psi}(x_0)$ Lemma A.2 guarantees the existence of a unique $b^* > 0$ such that $\Psi_{b^*}(x_0) = K_T$. In this later case, (i) of Proposition 3.2 guarantees the existence of a unique Λ^* for which b^* is optimal for (P2) with Λ^* . In both cases we have (iii).
- 2. Suppose $\overline{\Lambda} < 0$. If $\Psi_{b_0}(x_0) \ge K_T$ then (ii) is satisfied and the unconstrained problem satisfies the restriction. Therefore, $b^* = b_0$ is optimal for (P2) with $\Lambda^* = 0$. If $\Psi_{b_0}(x_0) < K_T < \hat{\Psi}(x_0)$ just as before we know there exists a unique b^* such that $\Psi_{b^*}(x_0) = K_T$. By (ii) of Proposition 3.2 there exists a unique Λ^* for which b^* is optimal for (P2) with Λ^* . In both cases we also have (iii).

As a consequence we have the main theorem:

Theorem 4.2. Let $x_0 \ge 0$, $T \ge 0$ and $V(x_0)$ be the optimal solution to (P1). Then

(i)
$$V(x_0) \leq \inf_{\Lambda > 0} V_{\Lambda}(x_0)$$
 and

(ii) $\inf_{\Lambda \ge 0} V_{\Lambda}(x_0) \le V(x_0).$

Therefore, $\inf_{\Lambda \ge 0} V_{\Lambda}(x_0) = V(x_0).$

Proof. Fix $x_0 \ge 0$. Condition (i) is satisfied since $\inf_{\Lambda \ge 0} V_{\Lambda}(x_0)$ is the dual problem of (P1). To verify condition (ii) we have the following cases:

(i) $T < H_{x_0}$: By Proposition 4.1 there is a pair (Λ^*, b^*) such that

$$\begin{split} \inf_{\Lambda \ge 0} V_{\Lambda}(x_0) &\leq V_{\Lambda^*}(x_0) \\ &= \mathbb{E}_{x_0} \left[\int_0^{\tau^{b^*}} e^{-\delta t} dD_t^{b^*} + \Lambda^* \int_0^{\tau^{b^*}} e^{-\delta t} dt \right] - \Lambda^* K_T \\ &= \mathbb{E}_{x_0} \left[\int_0^{\tau^{b^*}} e^{-\delta t} dD_t^{b^*} \right] \\ &\leq V(x_0), \end{split}$$

where the last inequality follows since the barrier strategy b^* satisfies (2).

(ii) $T = H_{x_0}$: In this case $K_T = \hat{\Psi}(x_0)$ and by Lemma A.3

$$\Lambda\left(\mathbb{E}_{x_0}\left[\int_0^{\tau^{b_\Lambda}} e^{-\delta s} ds\right] - K_T\right) \to 0 \quad \text{as} \quad \Lambda \to \infty.$$

Also $\mathbb{E}_{x_0}\left[\int_0^{\tau^{b_\Lambda}} e^{-\delta t} dD_t^{b_\Lambda}\right] \to 0$ as $\Lambda \to \infty$ since $b_\Lambda \to \infty$. Therefore, since $V_0(x_0) \ge 0$ and $V_\Lambda(x_0)$ is convex in Λ (it is the supremum of linear functions) we obtain that

$$\inf_{\Lambda \ge 0} V_{\Lambda}(x) = 0 \le V(x_0).$$

(iii) $\underline{T > H_{x_0}}$: In this case $K_T > \hat{\Psi}(x_0)$, therefore for all $b \ge 0$ it holds that $\mathbb{E}_{x_0} \left[\int_0^{\tau^b} e^{-\delta s} ds \right] < \hat{\Psi}(x_0) < K_T$. From this, one can deduce there exists $\epsilon > 0$ such that $\Lambda \left(\mathbb{E}_{x_0} [\int_0^{\tau^{b_\Lambda}} e^{-\delta s} ds] - K_T \right) < -\Lambda \epsilon$. Letting $\Lambda \to \infty$ we obtain $\inf_{\Lambda \ge 0} V_{\Lambda}(x) = -\infty \le V(x_0)$.

_			
г		ъ	
L		L	
		L	
_			

5 Numerical examples

In this section we illustrate with more detail the cases that came up in the proof of Proposition 4.1 and Theorem 4.2. As presented in the previous section to obtain the optimal value function of (P1) we showed a pair (b^*, Λ^*) that certified strong duality. To do so, we consider several cases depending on the initial value x_0 and T. We will continue to assume the following parameter values: $\lambda = 1, c = 1.3, Y_i \sim exp(1), \text{ and } \delta = 0.1$. In each case we will show two graphs. Graphs on the left show $\Psi_b(x)$ for different values of b and graphs on the right show $V_{\Lambda}(x_0)$ for different values of Λ .

For the first case, choose T and x_0 so that K_T lies below $\Psi_{b_0}(x_0)$. In this situation we know that the unconstrained solution satisfies the restriction. With such values the plot of $V_{\Lambda}(x_0)$ for



Figure 2: Inactive constraint.

different values of Λ illustrate that the minimum is attained at $\Lambda^* = 0$, see Figure 2. The optimal solution is $V_0(x_0)$ and the optimal barrier is $b^* = b_0 = 0.8$.

In the second case, let T and x_0 have values such that K_T lies between $\hat{\Psi}(x_0)$ and $\Psi_{b_0}(x_0)$. With such values the plot of $V_{\Lambda}(x_0)$ for different values of Λ reflects the existence of a minimum $\Lambda^* > 0$. To find it, find b^* that satisfies $\Psi_{b^*}(x_0) = K_T$ and use Proposition 3.2 to get Λ^* , see Figure 3. The optimal solution is $V_{\Lambda^*}(x_0)$ and the optimal barrier is b^* .



Figure 3: Active constraint.

Now, let T and x_0 have values such that $K_T = \hat{\Psi}(x_0)$. With such values the plot of $V_{\Lambda}(x_0)$ for different values of Λ shows that the minimum is attained at ∞ with a value of 0 see Figure 4. In this particular case we conclude that $V(x_0) = 0$ so that for (P1) the optimal strategy is to do nothing.



Figure 4: Do nothing.

In the last case, let T and x_0 have values such that K_T lies above $\hat{\Psi}(x_0)$. With such values the plot of $V_{\Lambda}(x_0)$ for different values of Λ reflects that the minimum is also attained at ∞ . This is due to the fact that there is no b such that $\Psi_b(x_0) = K_T$, see Figure 5. The problem is infeasible so its optimal value is $-\infty$.



Figure 5: Problem unfeasible.

Figure 6 shows the value functions of both the unconstrained (Solid line) and the constrained problem (Dashed line) for K_{20} . In this case for x < 4.23 the value function of the constrained problem equals $-\infty$.



Figure 6: $V(x_0)$ for both the unconstrained and the constrained problem.

Finally, Figure 7 shows how the solution to problem (P1) can be graphically characterized in terms of (x_0, K_T) , the horizontal line at level $\frac{1}{\delta}$, $\hat{\Psi}(x)$ and $\Psi_{b_0}(x)$.



Figure 7: Solution description.

6 Conclusions and future work

In the framework of the classical dividend problem there exists a trade-off between stability and profitability. Minimizing the ruin probability could lead to no dividend payment whereas maximizing the expected value of the discounted payments leads to a dividend payment trend for which ruin is certain regardless of the initial amount x_0 . In this work we study a way to link the profits and the time of ruin derived from a dividend payment strategy D. We introduced a restriction that imposes a constraint on the time of ruin. Under exponentially distributed claim sizes distribution we succeed in solving the constrained problem using Duality Theory. Consider the problem with general claims distribution is part of future research. Ongoing research also involves different type of restrictions and time-dependent optimal strategies as well.

A Auxiliary Lemmas

Lemma A.1. Let r_2 be the negative root of the characteristic polynomial (7). Then. $r_2 + \alpha > 0$.

Proof.

$$\begin{split} r_{2} + \alpha > 0 \\ \iff \frac{-\alpha c + (\lambda + \delta) - \sqrt{([\alpha c - (\lambda + \delta)]^{2} + 4c\alpha\delta)}}{2c} + \alpha > 0 \\ \iff 2c\alpha > \alpha c - (\lambda + \delta) + \sqrt{([\alpha c - (\lambda + \delta)]^{2} + 4c\alpha\delta)} \\ \iff c\alpha + (\lambda + \delta) > \sqrt{([\alpha c - (\lambda + \delta)]^{2} + 4c\alpha\delta)} \\ \iff c\alpha + (\lambda + \delta) > \sqrt{([\alpha c + (\lambda + \delta) - 2(\lambda + \delta)]^{2} + 4c\alpha\delta)} \\ \iff c\alpha + (\lambda + \delta) > \sqrt{([\alpha c + (\lambda + \delta)]^{2} - 4(\alpha c + (\lambda + \delta))(\lambda + \delta) + 4(\lambda + \delta)^{2} + 4c\alpha\delta)} \\ \iff -4(\alpha c + (\lambda + \delta))(\lambda + \delta) + 4(\lambda + \delta)^{2} + 4c\alpha\delta < 0 \\ \iff 4\alpha c(\lambda + \delta) > 4c\alpha\delta \\ \iff 4\alpha c(\lambda + \delta) > 4c\alpha\delta \\ \iff 4\alpha c\lambda > 0. \end{split}$$

Lemma A.2. Let $\Psi_b(x)$ be the solution of problem (A1). For $x \ge 0$ fixed, $\Psi_b(x)$ is increasing in b. *Proof.* The proof consists on calculate $\frac{d\Psi_b(x)}{db}$. For x < b

$$\begin{split} g(x) &\coloneqq \frac{d\Psi_b(x)}{db} \\ &= \frac{(\alpha + r_1)(\alpha + r_2)r_2^2 e^{r_2 b} \alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)]}{(\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)])^2} e^{r_1 x} \\ &- \frac{(\alpha + r_1)(\alpha + r_2)r_2 e^{r_2 b} \alpha \delta[r_1^2 e^{r_1 b}(\alpha + r_1) - r_2^2 e^{r_2 b}(\alpha + r_2)]}{(\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)])^2} e^{r_1 x} \\ &- \frac{(\alpha + r_2)^2 r_2^2 e^{r_2 b} \alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)])^2}{(\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)])^2} e^{r_2 x} \\ &+ \frac{(\alpha + r_2)^2 r_2 e^{r_2 b} \alpha \delta[r_1^2 e^{r_1 b}(\alpha + r_1) - r_2^2 e^{r_2 b}(\alpha + r_2)]}{(\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)])^2} e^{r_2 x}. \end{split}$$

The numerator of this expression can be reduced to

$$\begin{split} &\alpha \delta r_2(\alpha+r_1)(\alpha+r_2)e^{r_2(b+x)}[r_1r_2e^{r_1b}(\alpha+r_1)\\ &-r_2^2e^{r_2b}(\alpha+r_2)-r_1^2e^{r_1b}(\alpha+r_1)+r_2^2e^{r_2b}(\alpha+r_2)]\\ &-\alpha \delta r_2(\alpha+r_2)^2e^{r_2(b+x)}[r_1r_2e^{r_1b}(\alpha+r_1)\\ &-r_2^2e^{r_2b}(\alpha+r_2)-r_1^2e^{r_1b}(\alpha+r_1)+r_2^2e^{r_2b}(\alpha+r_2)]\\ &=\alpha \delta r_2r_1(\alpha+r_1)(\alpha+r_2)^2e^{r_2(b+x)}e^{r_1b}[r_1-r_2]\\ &-\alpha \delta r_2r_1(\alpha+r_1)^2(\alpha+r_2)e^{r_2(b+x)}e^{r_1b}[r_1-r_2]^2>0. \end{split}$$

Now, for $x \ge b$, $\Psi_b(x) = \Psi_b(b)$ and therefore we must calculate

$$\begin{split} \frac{d \varPsi_b(b)}{db} = & g(b) + \frac{(\alpha + r_1)(\alpha + r_2)r_2 e^{r_2 b}r_1 e^{r_1 b}}{\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)]} \\ & - \frac{(\alpha + r_2)^2 r_2 e^{r_2 b} r_2 e^{r_2 b}}{\alpha \delta(e^{r_1 b}r_1(\alpha + r_1) - e^{r_2 b}r_2(\alpha + r_2))} - \frac{(\alpha + r_2)r_2 e^{r_2 b}}{\alpha \delta} \\ & = & g(b) + \frac{(\alpha + r_1)(\alpha + r_2)r_1 r_2 e^{(r_1 + r_2) b}}{\alpha \delta[r_1 e^{r_1 b}(\alpha + r_1) - r_2 e^{r_2 b}(\alpha + r_2)]} \\ & - \frac{(\alpha + r_2)^2 r_2^2 e^{2r_2 b} - (\alpha + r_1)(\alpha + r_2)r_1 r_2 e^{(r_1 + r_2) b} + (\alpha + r_2)^2 r_2^2 e^{2r_2 b}}{\alpha \delta(e^{r_1 b}r_1(\alpha + r_1) - e^{r_2 b}r_2(\alpha + r_2))} \\ & = & g(b) > 0. \end{split}$$

Lemma A.3. Let $x \ge 0$ and $T \ge 0$. If $K_T = \hat{\Psi}(x)$ then $\Lambda \left[\mathbb{E}_x \left[\int_0^{\tau^{b_\Lambda}} e^{-\delta s} ds \right] - K_T \right] \to 0$ as $\Lambda \to \infty$.

Proof. From the proof of Proposition 4.1 we must calculate $\Lambda \left[\Psi_{b_{\Lambda}}(x) - K_T \right]$ as $\Lambda \to \infty$. Furthermore, from Proposition 3.2 and Lemma A.2 we know the second term goes to 0 as $\Lambda \to \infty$. Since $\Lambda \to \infty$ is equivalent to $b \to \infty$, we will calculate the limit of

$$-\frac{[\Psi_b(x) - K_T]^2}{\frac{d\Psi_b(x)}{d\Lambda}} = -\frac{[\Psi_b(x) - K_T]^2}{\frac{d\Psi_b(x)}{db}\frac{db}{d\Lambda}} \text{ as } b \to \infty.$$

Recall that

$$\Psi_b(x) = \frac{1}{\delta} + \frac{(\alpha + r_1)(\alpha + r_2)r_2e^{r_2b + r_1x} - (\alpha + r_2)(\alpha + r_1)r_1e^{r_1b + r_2x}}{\alpha\delta[r_1e^{r_1b}(\alpha + r_1) - r_2e^{r_2b}(\alpha + r_2)]}$$

and that $K_T = \hat{\Psi}(x) = \frac{1}{\delta} - \frac{\alpha + r_2}{\alpha \delta} e^{r_2 x}$. Therefore

$$(\Psi_b(x) - K_T)^2 = \left(\frac{(\alpha + r_2)r_2[e^{r_1x}(\alpha + r_1) - e^{r_2x}(\alpha + r_2)]}{(\alpha\delta[r_1e^{r_1b}(\alpha + r_1) - r_2e^{r_2b}(\alpha + r_2)]}e^{r_2b}\right)^2$$

From Proposition 3.2 and Lemma A.2 it can be derived that

$$\frac{d\Psi_b(x)}{db}\frac{db}{d\Lambda} = \frac{\delta\alpha(\alpha+r_1)^2(\alpha+r_2)^2[r_1-r_2]^3e^{r_1b}e^{r_2(b+x)}}{(\alpha\delta[r_1e^{r_1b}(\alpha+r_1)-r_2e^{r_2b}(\alpha+r_2)])^2} \cdot \frac{1}{e^{-r_2b}(r_1(\lambda+\delta)+\alpha\delta)-e^{-r_1b}(r_2(\lambda+\delta)+\alpha\delta)}$$

Now,

$$\begin{aligned} -\frac{(\Psi_b(x) - K_T)^2}{\frac{d\Psi_b(x)}{db}\frac{dh}{d\Lambda}} &= -\frac{\left((\alpha + r_2)r_2[e^{r_1x}(\alpha + r_1) - e^{r_2x}(\alpha + r_2)]\right)^2 e^{2r_2b}}{\alpha\delta(r_1 - r_2)^3(\alpha + r_1)^2(\alpha + r_2)^2 e^{r_1b}e^{r_2b}e^{r_2x}} \\ &\quad \cdot \left(e^{-r_2b}(r_1(\lambda + \delta) + \alpha\delta) - e^{-r_1b}(r_2(\lambda + \delta) + \alpha\delta)\right) \\ &= -\frac{(\alpha + r_2)r_2[e^{r_1x}(\alpha + r_1) - e^{r_2x}(\alpha + r_2)]\right)^2}{\alpha\delta(r_1 - r_2)^3(\alpha + r_1)^2(\alpha + r_2)^2 e^{r_2x}} \\ &\quad \cdot \left(e^{-r_1b}(r_1(\lambda + \delta) + \alpha\delta) - e^{(r_2 - 2r_1)b}(r_2(\lambda + \delta) + \alpha\delta)\right), \end{aligned}$$

from where one can conclude that the last expression goes to 0 as $b \to \infty$ since $r_2 < 0 < r_1$.

References

- S. Asmussen and H. Albrecher. *Ruin Probabilities*. Science and Applied Probability. Springer-Verlag, 2nd edition, 2010.
- [2] P. Azcue and N. Muler. Optimal reinsurance and dividend distribution policies in the cramérlundberg model. *Mathematical Finance*, 15(2):261308, 03 2005.
- [3] H. Bühlmann. Mathematical Methods of Risk Theory. Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1970.
- [4] B. De Finetti. Su unimpostazion alternativa dell teoria collectiva del rischio. In Transactions of the XVth International Congress of Actuaries, number 2, pages 433–443, 1957.
- [5] P. Grandits. An optimal consumption problem in finite time with a constraint on the ruin probability. *Preprint*, 2013.
- [6] C. Hipp. Optimal dividend payment under a ruin constraint: Discrete time and state space. Bltter der DGVFM, 26(2):255-264, 2003.
- [7] J. Paulsen. Optimal dividend payouts for diffusions with solvency constraints. Finance and Stochastics, 7(4):457-473, 09 2003.
- [8] H. Schmidli. On minimizing the ruin probability by investment and reinsurance. The Annals of Applied Probability, 12(3):890–907, 08 2002.
- [9] H. Schmidli. Stochastic Control in Insurance. Probability and Its Applications. Springer, 2008.
- [10] S. Thonhauser and H. Albrecher. Dividend maximization under consideration of the time value of ruin. *Insurance: Mathematics and Economics*, 41(1):163–184, 2007.