# Lattice Universe: examples and problems

## Maxim Brilenkov<sup>1</sup>, Maxim Eingorn<sup>2</sup>, and Alexander Zhuk<sup>3</sup>

<sup>1</sup>Department of Theoretical Physics, Odessa National University, Dvoryanskaya st. 2, Odessa 65082, Ukraine

<sup>2</sup>Physics Department, North Carolina Central University, Fayetteville st. 1801, Durham, North Carolina 27707, U.S.A.

<sup>3</sup>Astronomical Observatory, Odessa National University, Dvoryanskaya st. 2, Odessa 65082, Ukraine

E-mail: maxim.brilenkov@gmail.com, maxim.eingorn@gmail.com, ai.zhuk2@gmail.com

**Abstract.** We consider lattice Universes with spatial topologies  $T \times T \times T$ ,  $T \times T \times R$ and  $T \times R \times R$ . In the Newtonian limit of General Relativity, we solve the Poisson equation for the gravitational potential in the enumerated models. In the case of point-like massive sources in the  $T \times T \times T$  model, we demonstrate that the gravitational potential has no definite values on the edges joining identical masses in neighboring cells, i.e. at points where masses are absent. Clearly, this is a nonphysical result. The only way to avoid this problem and get a regular solution at any point of the cell is the smearing of these masses over some region. In the cases of  $T \times T \times R$  and  $T \times R \times R$  topologies, there is no way to get any physically reasonable and nontrivial solution. The only solutions we can get here are the ones which reduce these topologies to the  $T \times T \times T$  one.

#### Contents

1	Introduction	1
<b>2</b>	Topology $T \times T \times T$ . Point-like masses	3
3	Topology $T \times T \times T$ . Smeared masses	6
4	<b>Topology</b> $T \times T \times R$	9
5	<b>Topology</b> $T \times R \times R$	12
6	Conclusion	14

### 1 Introduction

Papers devoted to the lattice Universe can be divided into two groups. The first group includes articles (see, e.g., [1–11]) offering alternative cosmological models. Despite the great success of the standard ACDM model, it has some problematic aspects. The main one is the presence of dark energy and dark matter which constitute about 96% of the total energy density in the Universe. However, the nature of these components is still unknown. Another subtle point is that the conventional model is based on the Friedmann-Lemaitre-Robertson-Walker (FLRW) geometry with homogeneous and isotropic distribution of matter in the form of a perfect fluid. Observations show that such approximation works well on rather large scales. According to simple estimates made on the basis of statistical physics, these scales correspond to 190 Mpc [12] which is in good agreement with observations. This is the cell of uniformity size. Deep inside this cell, our Universe is highly inhomogeneous. Here, we clearly see galaxies, dwarf galaxies, groups and clusters of galaxies. Therefore, it makes sense to consider matter on such scales in the form of discrete gravitational sources. In this case, we arrive at the question how this discrete distribution influences global properties and dynamics of the Universe. This problem was investigated in the above mentioned papers (see also [13]). Here, gravitating masses are usually distributed in a very simplified and artificial way. They form either periodic structures of identical masses with proper boundary conditions or correspond to Einstein equation solutions (e.g., Schwarzschild or Schwarzschild-de Sitter solutions) matching with each other with the help of the Israel boundary conditions. Usually, such models do not rely on the  $\Lambda$ CDM background solution and do not include observable parameters (e.g., the average rest-mass density  $\bar{\rho}$  of matter in the Universe). As a result, these models have nothing common with the observable Universe. Their main task is to find new phenomena following from discretization and nontrivial topology.

Papers from the second class are devoted to numerical N-body simulations of the observable Universe. Here, the lattice is constructed as follows. In the spatially flat Universe, we choose a three-dimensional cell with N arbitrarily distributed gravitating masses  $m_i$  and suppose periodic boundary conditions for them on the boundary of the cell. Such models rely on background FLRW geometry with a scale factor a. It is supposed that the background solution is the  $\Lambda$ CDM model with the perfect fluid in the form of dust with the average rest-mass density  $\overline{\rho}$ . Discrete masses/inhomogeneities with the real rest-mass density  $\rho = \sum_{i=1}^{N} m_i \delta(\mathbf{r} - \mathbf{r}_i)$  perturb this background. The gravitational potential inside the cell is determined in the Newtonian limit by the following Poisson equation [14–16]:

$$\Delta \varphi(\mathbf{r}) = 4\pi G_N \left[ \sum_{i=1}^N m_i \delta(\mathbf{r} - \mathbf{r}_i) - \overline{\rho} \right], \qquad (1.1)$$

where  $G_N$  is the Newtonian gravitational constant, and  $\mathbf{r}, \mathbf{r}_i$  belong to the cell, e.g.,  $x_i \in [-l_1/2, l_1/2], y_i \in [-l_2/2, l_2/2], z_i \in [-l_3/2, l_3/2]$ . Here, the Laplace operator  $\Delta = \partial^2/\partial x^2 + \partial^2/\partial y^2 + \partial^2/\partial z^2$ , and the coordinates x, y and z, the gravitational potential  $\varphi$  and the restmass densities  $\rho$  and  $\overline{\rho}$  correspond to the comoving frame. All these quantities are connected with the corresponding physical ones as follows:  $\mathbf{R}_{\text{phys}} = a\mathbf{r}, \Phi_{\text{phys}} = \varphi/a$  and  $\overline{\rho}_{\text{phys}} = \overline{\rho}/a^3$ . The same equation can be also obtained in the Newtonian limit of General Relativity [12, 17, 18]. If we know the gravitational potential, then we can investigate dynamics of the inhomogeneities/galaxies taking into account both gravitational attraction between them and cosmological expansion of the Universe [17, 19, 20].

It can be easily seen that in the case of a finite volume (e.g., the volume of the cell) Eq. (1.1) satisfies the superposition principle. Here, for each gravitating mass  $m_i$  we can determine its contribution to the average rest-mass density:  $\overline{\rho}_i = m_i/(l_1 l_2 l_3), \ \overline{\rho} = \sum_{i=1}^N \overline{\rho}_i$ . Therefore, we can solve Eq. (1.1) for each mass  $m_i$  separately.

If we do not assume periodic boundary conditions, at least for one of directions, there is no lattice in these directions and space along them is not compact (in the sense of lack of the finite period of the lattice). Obviously, in infinite space the number of inhomogeneities must be also infinite:  $N \to \infty$ . This case has a number of potentially dangerous points. First, the superposition principle does not work here because we cannot determine  $\overline{\rho}_i$  for each of masses  $m_i$ . Second, it is known that the sum of an infinite number of Newtonian potentials diverges (the Neumann-Seeliger paradox [21]). Therefore, in general, the considered model can also suffer from this problem if we do not distribute masses in some specific way. Third, we can easily see from Eq. (1.1) that the presence of  $\overline{\rho}$  will result in quadratic (with respect to the noncompact distance) divergence. Hence, to avoid it, we should cut off gravitational potentials in these directions. This also may require a very specific distribution of the gravitating masses.

In the present paper, we investigate Eq. (1.1) for different topologies of space which imply different kinds of the lattice structure. First, in section 2, we consider the  $T \times T \times T$ topology with periodic boundary conditions in all three spatial dimensions. For point-like sources, we obtain a solution in the form of an infinite series. This series has the well known Newtonian type divergence in the positions of masses. However, we show that the sum of the series does not exist on the edges joining identical particles in neighboring cells. Therefore, there is no solution in points where masses are absent. This is a new result. To avoid this nonphysical property, in section 3, we smear point-like sources. We present them in the form of uniformly filled parallelepipeds. In this case, the infinite series has definite limits on the edges. Therefore, smearing of the gravitating masses in N-body simulations plays a dual role: first, this is the absence of the Newtonian divergence in the positions of masses, second, this is the regular behavior of the gravitational potential in all other points.

In sections 4 and 5 we consider a possibility to get reasonable solutions of Eq. (1.1) in the case of absence of periodicity in one or two spatial directions. In section 4, we investigate a model with the spatial topology  $T \times T \times R$ , i.e. with one noncompact dimension, let it be z. As we mentioned above, due to noncompactness, the gravitational potential may suffer from the Neumann-Seeliger paradox and additionally has a divergence of the form  $\overline{\rho}z^2 \to +\infty$ for  $|z| \to +\infty$ . In this section we try to resolve these problems with the help of a special arrangement of gravitating masses in the direction of z. Similar procedure in the flat Universe with topology  $R^3$  was performed in [17]. Unfortunately, in the case of topology  $T \times T \times R$ , there is no possibility to arrange masses in such a way that the gravitational potential is a smooth function in any point z. The same result takes place for the Universe with topology  $T \times R \times R$  which is considered in section 5. Here we also demonstrate impossibility of constructing a smooth potential. The main results are briefly summarized in concluding section 6.

#### 2 Topology $T \times T \times T$ . Point-like masses

Obviously, for topology  $T \times T \times T$  the space is covered by identical cells, and, instead of an infinite number of these cells, we may consider just one cell with periodic boundary conditions. As we mentioned in Introduction, due to the finite volume of the cell, we can apply the superposition principle. It means that we can solve Eq. (1.1) for one arbitrary gravitating mass, and the total gravitational potential in a point inside the cell is equal to a sum of gravitational potentials (in this point) of all N masses. Without loss of generality, we can put a gravitating mass m at the origin of coordinates. Then, the Poisson equation (1.1) for this mass reads

$$\Delta \varphi = 4\pi G_N \left( m\delta\left(\mathbf{r}\right) - \frac{m}{l_1 l_2 l_3} \right) \,. \tag{2.1}$$

Taking into account that delta-functions can be expressed as

$$\delta(x) = \frac{1}{l_1} \sum_{k_1 = -\infty}^{+\infty} \cos\left(\frac{2\pi k_1}{l_1}x\right) \,, \tag{2.2}$$

we get

$$\Delta \varphi = 4\pi G_N \left[ \frac{m}{l_1 l_2 l_3} \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \sum_{k_3 = -\infty}^{+\infty} \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) \cos\left(\frac{2\pi k_3}{l_3}z\right) - \frac{m}{l_1 l_2 l_3} \right]$$
(2.3)

Therefore, it makes sense to look for a solution of this equation in the form

$$\varphi = \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \sum_{k_3 = -\infty}^{+\infty} C_{k_1 k_2 k_3} \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) \cos\left(\frac{2\pi k_3}{l_3}z\right), \quad (2.4)$$

where unknown coefficients  $C_{k_1k_2k_3}$  can be easily found from Eq. (2.3):

$$C_{k_1k_2k_3} = -\frac{G_Nm}{\pi l_1 l_2 l_3} \frac{1}{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2} + \frac{k_3^2}{l_3^2}}, \quad k_1^2 + k_2^2 + k_3^2 \neq 0.$$
(2.5)

Hence, the desired gravitational potential is

$$\varphi = -\frac{G_N m}{\pi l_1 l_2 l_3} \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \sum_{k_3 = -\infty}^{+\infty} \frac{\cos\left(\frac{2\pi k_1}{l_1}x\right)\cos\left(\frac{2\pi k_2}{l_2}y\right)\cos\left(\frac{2\pi k_3}{l_3}z\right)}{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2} + \frac{k_3^2}{l_3^2}}, \quad (2.6)$$

where  $k_1^2 + k_2^2 + k_3^2 \neq 0$ . If x, y, z simultaneously tend to zero, then the gravitational potential (2.6) has the Newtonian limit:

$$\varphi \to -\frac{G_N m}{\pi} \iint_{-\infty}^{+\infty} \frac{\cos(2\pi x k_x) \cos(2\pi y k_y) \cos(2\pi z k_z)}{k^2} dk_x dk_y dk_z = -\frac{G_N m}{r}, \qquad (2.7)$$

where  $r = (x^2 + y^2 + z^2)^{1/2}$ , as it should be. A good feature of the potential (2.6) is that its average value (integral) over the cell is equal to zero:  $\overline{\varphi} = 0^1$ . This is a physically reasonable result because  $\overline{\rho - \overline{\rho}} = 0$ .

Clearly, in the case of a point-like gravitational source, we have usual divergence at the point of its location. Now, we want to demonstrate that there is also a problem at points where gravitating masses are absent. More precisely, we will show that the sum (2.6) is absent on edges which connect identical masses in neighboring cells. In our particular example, they are lines of intersection (pairwise) of the planes x = 0, y = 0 and z = 0. Let us consider the potential (2.6) on the edge y = 0, z = 0. The numerical calculation of the potential on this edge at the point  $x = l_1/2$  for different values of the limiting number n (being the maximum absolute value of the summation indexes:  $|k_{1,2,3}| \leq n$ ) is presented in the following table for the cubic cell case  $l_1 = l_2 = l_3 \equiv l$ . This table clearly demonstrates that the potential does not tend (with the growth of n) to any particular finite number.

n	$\frac{\varphi_n(l/2,0,0)}{G_N m/l}$	n	$\frac{\varphi_n(l/2,0,0)}{G_N m/l}$
40	-0.73371	41	0.89453
60	-0.72869	61	0.89969
80	-0.72614	81	0.90229

To understand the reason for this, let us analyze the structure of the expression (2.6) in more detail. For z = 0 the gravitational potential reads:

$$\varphi(x,y,0) = -\frac{G_N m}{\pi l_1 l_2 l_3} \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \sum_{k_3 = -\infty}^{+\infty} \frac{\cos\left(\frac{2\pi k_1}{l_1}x\right)\cos\left(\frac{2\pi k_2}{l_2}y\right)}{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2} + \frac{k_3^2}{l_3^2}} \\ = -\frac{4G_N m}{l_1 l_2} \sum_{k_1 = 1}^{+\infty} \sum_{k_2 = 1}^{+\infty} \frac{1}{\sqrt{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2}}} \cos\left(\frac{2\pi k_1}{l_1}x\right)\cos\left(\frac{2\pi k_2}{l_2}y\right) \coth\left(\pi\sqrt{\frac{l_3^2 k_1^2}{l_1^2} + \frac{l_3^2 k_2^2}{l_2^2}}\right) \\ - \frac{2G_N m}{l_2} \sum_{k_1 = 1}^{+\infty} \frac{1}{k_1}\cos\left(\frac{2\pi k_1}{l_1}x\right) \coth\left(\frac{\pi l_3 k_1}{l_1}\right) \\ - \frac{2G_N m}{l_1} \sum_{k_2 = 1}^{+\infty} \frac{1}{k_2}\cos\left(\frac{2\pi k_2}{l_2}y\right) \coth\left(\frac{\pi l_3 k_2}{l_2}\right) - \frac{G_N m \pi l_3}{3l_1 l_2}, \tag{2.8}$$

<sup>&</sup>lt;sup>1</sup>It is worth noting that in [12, 17, 22] the concrete mass distribution in the Universe with topology  $R^3$  is cited as an example of the case of nonzero average value  $\overline{\varphi} \neq 0$ . From the pure mathematical point of view this case is inadmissible in the framework of the first-order perturbation theory.

where we used the tabulated formulas for sums of series [23] (see, e.g., 5.1.25). All sums in this expression are potentially dangerous. To show it, we can drop the hyperbolic cotangents because  $\coth k \to 1$  for  $k \to +\infty$ . Two last sums are divergent depending on which edge we consider: x = 0 or y = 0, respectively. For example, on the edge y = 0, the sum  $\sum_{k_1=1}^{+\infty} \cos(2\pi k_1 x/l_1)/k_1 = -\ln[2\sin(\pi x/l_1)]$  (see 5.4.2 in [23]) is convergent for any ratio  $x/l_1 \neq 0, 1$  while  $\sum_{k_2=1}^{+\infty} (1/k_2) \sim \lim_{k_2 \to +\infty} \ln k_2$  is logarithmically divergent. The rough estimate of the double sum also leads to a divergent result. To be more precise, we investigate now finite sums<sup>2</sup> of the "suspicious" terms on the edge y = 0:

$$f_n(x) = 2\sum_{k_1=1}^n \sum_{k_2=1}^n \cos\left(\frac{2\pi k_1}{l_1}x\right) \frac{1}{\sqrt{\frac{l_2^2}{l_1^2}k_1^2 + k_2^2}} + \sum_{k_2=1}^n \frac{1}{k_2}.$$
 (2.9)

It is worth noting that in the case  $l_1 = l_2$  and  $x/l_1 = 1/2$  the logarithmically divergent terms exactly cancel each other. Really, it follows directly from the following estimates:

$$2\sum_{k_{1}=1}^{+\infty}\sum_{k_{2}=1}^{+\infty}\frac{\cos\left(\pi k_{1}\right)}{\sqrt{k_{1}^{2}+k_{2}^{2}}} = 2\sum_{k_{1}=1}^{+\infty}\sum_{k_{2}=1}^{+\infty}\frac{\left(-1\right)^{k_{1}}}{\sqrt{k_{1}^{2}+k_{2}^{2}}}$$
$$= 2\sum_{m=1}^{+\infty}\sum_{k_{2}=1}^{+\infty}\left[\frac{1}{\sqrt{(2m)^{2}+k_{2}^{2}}} - \frac{1}{\sqrt{(2m-1)^{2}+k_{2}^{2}}}\right]$$
$$\sim -2\sum_{m=1}^{+\infty}\sum_{k_{2}=1}^{+\infty}\frac{2m}{\left[(2m)^{2}+k_{2}^{2}\right]^{3/2}} \sim -2\int_{1}^{+\infty}\int_{1}^{+\infty}\frac{2x}{\left[(2x)^{2}+y^{2}\right]^{3/2}}dxdy$$
$$\sim -\lim_{R\to+\infty}\ln R = -\infty$$
(2.10)

and

$$\sum_{k_2=1}^{+\infty} \frac{1}{k_2} \sim \int_{1}^{+\infty} \frac{dx}{x} \sim \lim_{R \to +\infty} \ln R = +\infty.$$
 (2.11)

Therefore, both of these logarithmically divergent terms cancel each other. Nevertheless, the expression (2.9) does not have a definite limit for  $n \to +\infty$ . To demonstrate it, along with (2.9) let us introduce the function

$$f_{n+1}(x) = 2\sum_{k_1=1}^{n+1}\sum_{k_2=1}^{n+1}\cos\left(\frac{2\pi k_1}{l_1}x\right)\frac{1}{\sqrt{\frac{l_2^2}{l_1^2}k_1^2 + k_2^2}} + \sum_{k_2=1}^{n+1}\frac{1}{k_2}.$$
 (2.12)

Evidently, if the expression (2.9) is convergent for  $n \to +\infty$ , then in this limit the difference

 $<sup>^2 \</sup>rm Obviously,$  the inclusion of the hyperbolic cotangents does not effect the main results but makes calculations more complicated.

 $f_{n+1}(x) - f_n(x) \to 0$ . After some simple algebra we get (for  $l_1 = l_2$ )

$$f_{n+1}(x) - f_n(x) = 2 \sum_{k_1=1}^n \cos\left(\frac{2\pi k_1}{l_1}x\right) \frac{1}{\sqrt{k_1^2 + (n+1)^2}} + 2 \sum_{k_2=1}^n \cos\left(\frac{2\pi (n+1)}{l_1}x\right) \frac{1}{\sqrt{(n+1)^2 + k_2^2}} + \frac{1 + \sqrt{2}\cos\left(2\pi (n+1)x/l_1\right)}{n+1} \equiv \Delta f_n(x) + \frac{1 + \sqrt{2}\cos\left(2\pi (n+1)x/l_1\right)}{n+1}.$$
(2.13)

Here, the last term in the third line vanishes for  $n \to +\infty$ . Therefore, the problem of convergence of (2.9) is reduced now to the analysis of  $\Delta f_n(x)$ . In Figure 1, we show the graph of  $\Delta f_n(x)$  (for  $x/l_1 = 1/2$ ) as a function of n. Each point gives the value of  $\Delta f_n(x)$  for the corresponding number n. This picture clearly demonstrates that the difference  $f_{n+1}(x) - f_n(x)$  does not tend to zero for growing n. Even more, it does not go to any definite value. It can be also verified that a similar result takes place for any other point on any of the edges and holds also for  $l_1 \neq l_2$ . Therefore, we have proven that in the case of point-like gravitating masses in the considered lattice Universe the gravitational potential has no definite values on the edges joining identical masses in neighboring cells. Clearly, this is a nonphysical result.



**Figure 1**. The graph of  $\triangle f_n(l_1/2)$  as a function of the number *n*.

#### **3** Topology $T \times T \times T$ . Smeared masses

Can the smearing of gravitating masses resolve the problem found in the previous section? To answer this question, we present gravitating masses as uniformly filled parallelepipeds. This representation of masses looks a bit artificial. However, such form is the most appropriate for the considered cells, and the most important point is that the form of smearing does not matter for us at the moment. We just want to get a principal answer on a possibility to avoid the problem with the help of smearing. So, let the mass m be uniformly smeared over a parallelepiped (with the lengthes of edges a, b and c) which we put, without loss of generality, in the middle of the cell. It is convenient to introduce a function  $f_1(x)$  equal to 1 for  $x \in [-a/2, a/2]$  and 0 elsewhere inside  $[-l_1/2, l_1/2]$ . We can write down this function as

$$f_1(x) = \frac{a}{l_1} + \sum_{n=1}^{+\infty} \frac{2}{\pi n} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \,. \tag{3.1}$$

Similarly,

$$f_2(y) = \frac{b}{l_2} + \sum_{j=1}^{+\infty} \frac{2}{\pi j} \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right)$$
(3.2)

and

$$f_3(z) = \frac{c}{l_3} + \sum_{k=1}^{+\infty} \frac{2}{\pi k} \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \,. \tag{3.3}$$

Therefore, the rest-mass density of the mass under consideration is

$$\rho(\mathbf{r}) = \frac{m}{abc} f_1(x) f_2(y) f_3(z) \equiv \frac{m}{abc} f(\mathbf{r}) \,. \tag{3.4}$$

Then, Eq. (1.1) for this mass reads:

$$\begin{split} \Delta\varphi &= 4\pi G_N \left[ \frac{m}{abc} f(\mathbf{r}) - \frac{m}{l_1 l_2 l_3} \right] = 4\pi G_N m \left[ \frac{1}{l_1 l_2 c} \sum_{k=1}^{+\infty} \frac{2}{\pi k} \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \right. \\ &+ \frac{1}{l_1 l_3 b} \sum_{j=1}^{+\infty} \frac{2}{\pi j} \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) + \frac{1}{l_2 l_3 a} \sum_{n=1}^{+\infty} \frac{2}{\pi n} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \\ &+ \frac{1}{l_1 bc} \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} \frac{4}{\pi^2 j k} \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \\ &+ \frac{1}{l_2 ac} \sum_{n=1}^{+\infty} \sum_{k=1}^{+\infty} \frac{4}{\pi^2 n k} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \\ &+ \frac{1}{l_3 ab} \sum_{n=1}^{+\infty} \sum_{j=1}^{+\infty} \frac{4}{\pi^2 n j} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) \\ &+ \frac{1}{abc} \sum_{n=1}^{+\infty} \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} \frac{8}{\pi^3 n j k} \sin\left(\frac{a\pi n}{l_1}\right) \sin\left(\frac{b\pi j}{l_2}\right) \sin\left(\frac{c\pi k}{l_3}\right) \\ &\times \cos\left(\frac{2\pi n}{l_1}x\right) \cos\left(\frac{2\pi j}{l_2}y\right) \cos\left(\frac{2\pi k}{l_3}z\right) \right]. \end{split}$$

This equation implies that it makes sense to look for a solution in the following form:

$$\varphi(\mathbf{r}) = \frac{m}{l_1 l_2 c} \sum_{k=1}^{+\infty} C_k \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \\
+ \frac{m}{l_1 l_3 b} \sum_{j=1}^{+\infty} C'_j \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) + \frac{m}{l_2 l_3 a} \sum_{n=1}^{+\infty} C''_n \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \\
+ \frac{m}{l_1 b c} \sum_{j=1}^{+\infty} \sum_{k=1}^{+\infty} C_{jk} \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi k}{l_3}z\right) \\
+ \frac{m}{l_2 a c} \sum_{n=1}^{+\infty} \sum_{k=1}^{+\infty} C'_{nk} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \sin\left(\frac{c\pi k}{l_3}\right) \cos\left(\frac{2\pi j}{l_2}y\right) \\
+ \frac{m}{l_3 a b} \sum_{n=1}^{+\infty} \sum_{j=1}^{+\infty} C''_{nj} \sin\left(\frac{a\pi n}{l_1}\right) \cos\left(\frac{2\pi n}{l_1}x\right) \sin\left(\frac{b\pi j}{l_2}\right) \cos\left(\frac{2\pi j}{l_2}y\right) \\
+ \frac{m}{a b c} \sum_{n=1}^{+\infty} \sum_{j=1}^{+\infty} C_{njk} \sin\left(\frac{a\pi n}{l_1}\right) \sin\left(\frac{b\pi j}{l_2}\right) \sin\left(\frac{c\pi k}{l_3}\right) \\
\times \cos\left(\frac{2\pi n}{l_1}x\right) \cos\left(\frac{2\pi j}{l_2}y\right) \cos\left(\frac{2\pi k}{l_3}z\right)\right].$$
(3.6)

Substitution of this expression into the Poisson equation (3.5) gives

$$C_{k} = -\frac{2G_{N}}{\pi^{2}k} \left(\frac{l_{3}}{k}\right)^{2}, \quad C_{j}' = -\frac{2G_{N}}{\pi^{2}j} \left(\frac{l_{2}}{j}\right)^{2}, \quad C_{n}'' = -\frac{2G_{N}}{\pi^{2}n} \left(\frac{l_{1}}{n}\right)^{2},$$

$$C_{jk} = -\frac{4G_{N}}{\pi^{3}jk} \frac{1}{\left(\frac{j^{2}}{l_{2}} + \frac{k^{2}}{l_{3}^{2}}\right)}, \quad C_{nk}' = -\frac{4G_{N}}{\pi^{3}nk} \frac{1}{\left(\frac{n^{2}}{l_{1}^{2}} + \frac{k^{2}}{l_{3}^{2}}\right)},$$

$$C_{jn}'' = -\frac{4G_{N}}{\pi^{3}nj} \frac{1}{\left(\frac{n^{2}}{l_{1}^{2}} + \frac{j^{2}}{l_{2}^{2}}\right)}, \quad C_{njk} = -\frac{8G_{N}}{\pi^{4}njk} \frac{1}{\left(\frac{n^{2}}{l_{1}^{2}} + \frac{j^{2}}{l_{3}^{2}} + \frac{k^{2}}{l_{3}^{2}}\right)}.$$
(3.7)

Let us choose the same edge as in the previous section, that is y = 0, z = 0, and the same point  $x = l_1/2$ . The numerical calculation of the gravitational potential in this point for different values of the limiting number n is presented in the following table for the cubic cell case under the additional condition a = b = c = (3/7)l. In contrast to the previous case of a point-like source, here the potential apparently tends to a particular finite value. Therefore, in the case of smeared gravitating masses the gravitational potential has a regular behavior at any point inside the cell (including, e.g., the point x = y = z = 0).

n	$\frac{\varphi_n(l/2,0,0)}{G_N m/l}$	n	$\frac{\varphi_n(l/2,0,0)}{G_N m/l}$
$15 \\ 19 \\ 23$	$\begin{array}{c} 0.028717 \\ 0.028536 \\ 0.028368 \end{array}$	16 20 24	0.028443 0.028222 0.028223

### 4 Topology $T \times T \times R$

The  $T \times T \times R$  topology implies one noncompact dimension, let it be z. Therefore, there is a lattice structure in directions x and y and an irregular structure in the direction z. In a column  $x \in [-l_1/2, l_1/2], y \in [-l_2/2, l_2/2], z \in (-\infty, +\infty)$  there is an infinite number of gravitating masses. To obtain a "nice" regular solution, we will try to arrange masses in the z direction in such a way that in each point z the gravitational potential is determined by one mass only. There are two possibilities for that. Let this mass be at z = 0. In the first scenario, the potential and its first derivative (with respect to z) should vanish at some distance  $z_0$  (which we determine below). Then, the next mass should be at a distance (in the z direction) equal or greater than  $z_0 + z_1$ , where  $z_1$  is a distance at which the gravitational potential and its first derivative vanish for the second mass. Similarly, we should shift in the direction of z the third mass with respect to the second one and so on. In this scenario, we can arrange strips  $\Delta z$  between masses where the potential is absent. It occurs, e.g., between the first and second masses if the second mass is situated at distances greater that  $z_0 + z_1$ . In the strip, we place a uniform medium with the rest-mass density  $\overline{\rho}$ . Coordinates  $x \in [-l_1/2, l_1/2]$  and  $y \in [-l_2/2, l_2/2]$  of masses are arbitrary. In the second scenario, we should determine distances  $z_0, z_1, z_2, \ldots$  where potentials of neighboring (in the z direction) particles are smoothly matched to each other. This means that at these distances potentials are generally nonzero. Moreover, we suppose that their first derivatives are zero at the points of matching, i.e. potentials have extrema in these points. In this scenario, the neighboring (in the z direction) masses should have the same coordinates x and y. Now let us consider these scenarios in detail. For both of them, we need to look for a solution just for one particle. Let this particle be in the point x = y = z = 0. Then, Eq. (1.1) reads

$$\Delta \varphi = 4\pi G_N \left( m\delta(\mathbf{r}) - \bar{\rho} \right) \,. \tag{4.1}$$

Keeping in mind the regular structure in x and y directions, we can represent the delta functions  $\delta(x)$  and  $\delta(y)$  in the form (2.2). So, Eq. (4.1) is reduced to

$$\Delta \varphi = 4\pi G_N \left[ \frac{m}{l_1 l_2} \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \cos\left(\frac{2\pi k_1}{l_1} x\right) \cos\left(\frac{2\pi k_2}{l_2} y\right) \delta(z) - \bar{\rho} \right] . \tag{4.2}$$

Evidently, we can look for a solution of this equation in the form

$$\varphi = \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} C_{k_1 k_2}(z) \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) , \qquad (4.3)$$

and from the Poisson equation (4.2) we get

$$G_N \bar{\rho} = \sum_{k_1 = -\infty}^{+\infty} \sum_{k_2 = -\infty}^{+\infty} \pi \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) \left[\left(\frac{k_2^2}{l_2^2} + \frac{k_1^2}{l_1^2}\right) C_{k_1 k_2}(z) + \frac{m G_N}{l_1 l_2 \pi} \delta(z) - \frac{C_{k_1 k_2}''(z)}{4\pi^2}\right].$$
(4.4)

In this section, the prime denotes the derivative with respect to z. Now, we should determine unknown functions  $C_{k_1k_2}(z)$ . First, we find the zero mode  $C_{00}(z)$  which satisfies the equation

$$\frac{C_{00}''(z)}{4\pi} = \frac{mG_N}{l_1 l_2} \delta(z) - G_N \bar{\rho}.$$
(4.5)

This equation has the solution

$$C_{00}(z) = -2\pi G_N \bar{\rho} z^2 + \frac{2\pi}{l_1 l_2} m G_N |z| + B, \qquad (4.6)$$

where B is a constant of integration. This solution is a function growing with z. Therefore, we must cut off it at some distance  $z_0$ .

Let us consider the first scenario. From the condition  $C'_{00}(z_0) = 0$  we obtain

$$z_0 = \frac{m}{2\bar{\rho}l_1l_2} \quad \Leftrightarrow \quad \bar{\rho} = \frac{m}{2z_0l_1l_2}. \tag{4.7}$$

The second condition  $C_{00}(z_0) = 0$  provides the value of B:

$$B = -\frac{\pi m^2 G_N}{2\bar{\rho} l_1^2 l_2^2} \,. \tag{4.8}$$

Now, we want to determine the form of  $C_{k_1k_2}(z)$  when  $k_1^2 + k_2^2 \neq 0$ . In this case Eq. (4.4) is consistent only if the following condition holds true:

$$\left(\frac{k_2^2}{l_2^2} + \frac{k_1^2}{l_1^2}\right)C_{k_1k_2}(z) + \frac{mG_N}{l_1l_2\pi}\delta(z) - \frac{C_{k_1k_2}''(z)}{4\pi^2} = 0, \quad k_1^2 + k_2^2 \neq 0.$$
(4.9)

We look for a solution of this equation in the form

$$C_{k_1k_2}(z) = \widetilde{A}e^{-2\pi\beta|z|} + \widetilde{B}e^{2\pi\beta|z|}, \quad \beta \equiv \sqrt{\frac{k_2^2}{l_2^2} + \frac{k_1^2}{l_1^2}}, \quad (4.10)$$

where  $\widetilde{A}$  and  $\widetilde{B}$  are constants. The substitution of this function into Eq. (4.9) gives

$$\widetilde{A} - \widetilde{B} = -\frac{mG_N}{l_1 l_2 \beta}.$$
(4.11)

Therefore,

$$C_{k_1k_2}(z) = 2\widetilde{B}\cosh(2\pi\beta z) - \frac{mG_N}{l_1l_2\beta}e^{-2\pi\beta|z|}, \quad k_1^2 + k_2^2 \neq 0.$$
(4.12)

From the boundary condition  $C_{k_1k_2}(z_0) = 0$  we get

$$\widetilde{B} = \frac{mG_N}{2l_1 l_2 \beta} e^{-2\pi\beta z_0} \left[\cosh(2\pi\beta z_0)\right]^{-1} .$$
(4.13)

It can be easily verified that the function (4.12) (with  $\tilde{B}$  from (4.13)) does not satisfy the boundary condition  $C'_{k_1k_2}(z_0) = 0$ . Hence, we cannot determine the gravitational potential in accordance with the first scenario.

Now, we intend to demonstrate that there is a possibility to find the potential in the framework of the second scenario in the case of identical masses. However, this construction has a drawback inherent in the  $T \times T \times T$  model with the point-like source.

In the second scenario with identical masses m, all of them have the same coordinates x, y and are separated by the same distance  $2z_0 \equiv l_3$  in the direction of z. Here, the function  $C_{00}(z)$  still has the form (4.6). Since we require  $C'_{00}(z_0) = 0$ , the boundary  $z_0$  is determined by (4.7). However, the constant B is not given now by (4.8) because the condition  $C_{00}(z_0) = 0$ 

is absent. This constant can by found from the condition  $\overline{\varphi} = 0$  over the period  $l_3 = 2z_0$ . That is

$$\int_{-z_0}^{+z_0} C_{00}(z)dz = 0 \quad \Rightarrow \quad B = -\frac{2\pi G_N m z_0}{3l_1 l_2} = -\frac{\pi G_N m l_3}{3l_1 l_2}.$$
 (4.14)

The functions  $C_{k_1k_2}(z)$  for  $k_1^2 + k_2^2 \neq 0$  are given by Eq. (4.12), where the constant  $\widetilde{B}$  follows from the boundary condition  $C'_{k_1k_2}(z_0) = 0$ :

$$\widetilde{B} = -\frac{mG_N}{2l_1 l_2 \beta} e^{-2\pi\beta z_0} \left[\sinh(2\pi\beta z_0)\right]^{-1} .$$
(4.15)

It can be easily verified that  $C_{k_1k_2}(z)$  can be rewritten in the form

$$C_{k_1k_2}(z) = -\frac{G_N m}{l_1 l_2 \beta \sinh(2\pi\beta z_0)} \cosh\left[2\pi\beta(|z| - z_0)\right].$$
(4.16)

Therefore, in the second scenario the gravitational potential is

$$\begin{split} \varphi &= \sum_{k_1=-\infty}^{+\infty} \sum_{k_2=-\infty}^{+\infty} C_{k_1k_2}(z) \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) \\ &= C_{00}(z) + 2\sum_{k_1=1}^{+\infty} C_{k_10}(z) \cos\left(\frac{2\pi k_1}{l_1}x\right) + 2\sum_{k_2=1}^{+\infty} C_{0k_2}(z) \cos\left(\frac{2\pi k_2}{l_2}y\right) \\ &+ 4\sum_{k_1=1}^{+\infty} \sum_{k_2=1}^{+\infty} C_{k_1k_2}(z) \cos\left(\frac{2\pi k_1}{l_1}x\right) \cos\left(\frac{2\pi k_2}{l_2}y\right) \\ &= -G_N m \left\{ \frac{2\pi}{l_1 l_2 l_3} z^2 - \frac{2\pi}{l_1 l_2} |z| + \frac{\pi l_3}{3l_1 l_2} \right. \\ &+ \frac{2}{l_2} \sum_{k_1=1}^{+\infty} \frac{\cos\left(2\pi k_1 x/l_1\right)}{k_1} \frac{\cosh\left[2\pi k_1(|z|-z_0)/l_1\right]}{\sinh\left(2\pi k_1 z_0/l_1\right)} \\ &+ \frac{2}{l_1} \sum_{k_2=1}^{+\infty} \frac{\cos\left(2\pi k_2 y/l_2\right)}{k_2} \frac{\cosh\left[2\pi k_2(|z|-z_0)/l_2\right]}{\sinh\left(2\pi k_2 z_0/l_2\right)} \left. \begin{array}{c} \left(4.17\right) \\ &+ \frac{4}{l_1 l_2} \sum_{k_1=1}^{+\infty} \sum_{k_2=1}^{+\infty} \frac{\cos\left(2\pi k_1 x/l_1\right) \cos\left(2\pi k_2 y/l_2\right)}{\sqrt{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2}}} \frac{\cosh\left[2\pi \sqrt{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2}}(|z|-z_0)\right]}{\sinh\left(2\pi \sqrt{\frac{k_1^2}{l_1^2} + \frac{k_2^2}{l_2^2}}z_0\right)} \right\} \end{split}$$

When z = 0 and x, y simultaneously go to zero, the potential  $\varphi \to -G_N m/\sqrt{x^2 + y^2}$ , as it should be. From the physical point of view, it is clear that this scenario should coincide with the  $T \times T \times T$  case. Really, the triple sum (2.6) can be rewritten in the form (4.17) with the help of [23] (see 5.4.5). It can be also easily seen that on the plane z = 0 the expression (4.17) exactly coincides with Eq. (2.8). Therefore, in the second scenario we again arrive at the nonphysical result that the gravitational potential has no definite values on edges y = 0and x = 0.

### 5 Topology $T \times R \times R$

In this section we consider a model with a periodic boundary condition in one direction only. Two other spatial dimensions are noncompact. Here, in analogy with the previous section, we also suppose that the gravitational potential in the vicinity of a particle is determined by its mass only. The shape of such domain is dictated by the symmetry of the model and will be described below. On the boundary (in the direction of noncompact dimensions) of this domain the potential and its first derivative are equal to zero, and between such domains the potential is absent:  $\varphi = 0$ . Therefore, this model is similar to the first scenario in the previous section.

Let the mass be at the point x = y = z = 0 and the periodic (with the period  $l_1$ ) boundary condition be along the x coordinate. Then, the Poisson equation (4.1) for this topology can be written as follows:

$$\Delta \varphi = 4\pi G_N \frac{m}{l_1} \sum_{k_1 = -\infty}^{+\infty} \cos\left(\frac{2\pi k_1}{l_1} x\right) \delta(y) \delta(z) - 4\pi G_N \bar{\rho} \,. \tag{5.1}$$

We look for a solution of this equation in the form

$$\varphi = \sum_{k_1 = -\infty}^{+\infty} C_{k_1}(y, z) \cos\left(\frac{2\pi k_1}{l_1}x\right).$$
(5.2)

Then, from the Poisson equation (5.1) we get

$$G_N \bar{\rho} = \sum_{k_1 = -\infty}^{+\infty} \cos\left(\frac{2\pi k_1}{l_1}x\right) \left[ G_N \frac{m}{l_1} \delta(y) \delta(z) + \frac{\pi k_1^2}{l_1^2} C_{k_1}(y, z) - \frac{C_{k_1y}''(y, z)}{4\pi} - \frac{C_{k_1z}''(y, z)}{4\pi} \right],$$
(5.3)

where

$$C_{k_1y}''(y,z) \equiv \frac{\partial^2 C_{k_1}(y,z)}{\partial y^2}, \quad C_{k_1z}''(y,z) \equiv \frac{\partial^2 C_{k_1}(y,z)}{\partial z^2}.$$
 (5.4)

For the zero mode  $k_1 = 0$  this equation gives

$$G_N \bar{\rho} = G_N \frac{m}{l_1} \delta(y) \delta(z) - \frac{C_{0y}''(y,z)}{4\pi} - \frac{C_{0z}''(y,z)}{4\pi} \,.$$
(5.5)

Following the geometry of the model, it makes sense to turn to polar coordinates:

$$y = \xi \cos \phi, \quad z = \xi \sin \phi. \tag{5.6}$$

Then, the two-dimensional Laplace operator is

$$\Delta_{\xi\phi} = \frac{1}{\xi} \frac{\partial}{\partial \xi} \left( \xi \frac{\partial}{\partial \xi} \right) + \frac{1}{\xi^2} \frac{\partial^2}{\partial \phi^2} \equiv \Delta_{\xi} + \Delta_{\phi} \,. \tag{5.7}$$

It is clear that due to the symmetry of the problem the functions  $C_{k_1}$  do not depend on the azimuthal angle  $\phi$ . Therefore, Eq. (5.5) reads

$$G_N \bar{\rho} = G_N \frac{m}{l_1} \delta\left(\vec{\xi}\right) - \frac{\Delta_{\xi} C_0}{4\pi} \,. \tag{5.8}$$

This equation has the solution

$$C_0 = -\pi G_N \bar{\rho} \xi^2 + 2G_N \frac{m}{l_1} \ln \xi + \hat{B} , \qquad (5.9)$$

where we took into account the formula  $\triangle_{\xi} \ln \xi = 2\pi \delta\left(\vec{\xi}\right)$ . Similar to the previous model with one noncompact dimension, here the solution is also divergent in some directions. In the present case, it grows with the polar radius  $\xi$ . So, we must cut off this solution at some distance  $\xi_0$ . Clearly, this boundary represents the cylindrical surface  $\xi = \xi_0$ . The domain in which we put the mass  $m_0 = m$  is a cylinder with the radius  $\xi_0$  and the generator parallel to the axis x. The length of the cylinder along the x-axis is  $l_1$ . The mass m is in the center of the cylinder (with the coordinate x = 0 for the considered case). The next particle of the mass  $m_1$  is inside its own cylinder with the generator along the axis x and the radius  $\xi_1$ . This particle may have the coordinate x different from the first particle. All these cylinders have the periodic (with the period  $l_1$ ) boundary conditions along the x-axis. On the other hand, they should not overlap each other in the transverse (with respect to the x-axis) direction. Moreover, it is impossible to match them smoothly via cylindrical surfaces. Therefore, we demand that the gravitational potential outside the cylinders is equal to zero. Hence, on the boundaries of the cylinders ( $\xi = \xi_0$  for the first mass) the gravitational potential and its first derivative (with respect to  $\xi$ ) are equal to zero. These boundary conditions enable us to determine the radius  $\xi_0$  and the constant  $\hat{B}$  in Eq. (5.9). For example, from the condition  $dC_0(\xi_0)/d\xi = 0$  we get

$$\xi_0 = \sqrt{\frac{m}{\pi\bar{\rho}l_1}} \quad \Leftrightarrow \quad \bar{\rho} = \frac{m}{\pi\xi_0^2 l_1} \equiv \frac{m}{s_0 l_1}, \tag{5.10}$$

where  $s_0 = \pi \xi_0^2$  is the cross-sectional area of the cylinder. From the second boundary condition  $C_0(\xi_0) = 0$  we get the value of  $\hat{B}$ :

$$\hat{B} = \frac{G_N m}{l_1} \left( 1 - 2 \ln \xi_0 \right) \,. \tag{5.11}$$

Obviously, in the case  $k_1 \neq 0$ , Eq. (5.3) is consistent only if the following condition holds true:

$$G_N \frac{m}{l_1} \delta\left(\vec{\xi}\right) + \frac{\pi k_1^2}{l_1^2} C_{k_1}(y, z) - \frac{C_{k_1 y}'(y, z)}{4\pi} - \frac{C_{k_1 z}'(y, z)}{4\pi} = 0, \qquad (5.12)$$

which for  $\xi > 0$  can be rewritten in the form

$$\xi \frac{d^2 C_{k_1}}{d\xi^2} + \frac{d C_{k_1}}{d\xi} - \frac{4\pi^2 k_1^2}{l_1^2} C_{k_1} \xi = 0.$$
(5.13)

The solutions of this equation are the modified Bessel functions:

$$C_{k_1}(\xi) = C_1 I_0 \left(\frac{2\pi |k_1|}{l_1} \xi\right) - 2G_N \frac{m}{l_1} K_0 \left(\frac{2\pi |k_1|}{l_1} \xi\right) , \qquad (5.14)$$

where  $C_1$  is a constant of integration. We took into account that the function  $K_0(\xi) \to -\ln \xi$ for  $\xi \to 0$ , so the two-dimensional Laplacian acting on this function provides the necessary delta-function in Eq. (5.12). The function (5.14) should satisfy the same boundary conditions at  $\xi = \xi_0$  as the function  $C_0(\xi)$ . It can be easily verified that we cannot simultaneously reach both equalities  $C_{k_1}(\xi_0) = 0$  and  $dC_{k_1}(\xi_0)/d\xi = 0$ . Hence, we cannot determine the gravitational potential in accordance with the proposed scenario.

To conclude this section, it is worth noting that we can construct the potential in the scenario similar to the second one from the previous section. This is the case of identical masses distributed regularly in all directions. Obviously, this case is reduced to the  $T \times T \times T$  model from the section 2 with the drawback inherent in it.

Therefore, similar to the previous section, we also failed in determining a physically reasonable gravitational potential in the model with the topology  $T \times R \times R$ .

#### 6 Conclusion

Our paper was devoted to cosmological models with different spatial topology. According to the recent observations, our Universe is spatially flat with rather high accuracy. So, we restricted ourselves to this case. However, such spatially flat geometry may have different topology depending on a number of directions/dimensions with toroidal discrete symmetry. These topologies result in different kinds of the lattice Universe. There are a lot of articles exploring the lattice Universes (see, e.g., [1]-[11] and references therein). One of their main motivations is to provide an alternative (compared to the standard  $\Lambda$ CDM model) explanation of the late-time accelerated expansion of the Universe. Another important point is that our Universe is highly inhomogeneous inside the cell of uniformity with the size of the order of 190 Mpc [12]. Hence, it is quite natural to consider the Universe on such scales filled with discrete sources rather than a homogeneous isotropic perfect fluid.

On the other hand, N-body simulations of the evolution of structures in the Universe are based on dynamics of discrete sources in chosen cells. To perform such simulations, we should know gravitational potentials generated by these sources. Therefore, the main purpose of our paper was determination of gravitational potentials in the cases of three different spatial topologies:  $T \times T \times T$ ,  $T \times T \times R$  and  $T \times R \times R$ . The potential satisfies the corresponding Poisson equations of the form (1.1). These equations can be obtained as a Newtonian limit of General Relativity [12, 17]. So, to determine the potential, we should solve them. One of the main features of the analyzed Poisson equations is that they contain the average rest-mass density which represents a constant in the comoving frame. This results in two problems. First, we cannot, in general, apply the superposition principle. Second, the presence of such term leads to divergences in directions of noncompact dimensions. We tried to avoid these problems arranging masses in special ways. Our investigation has shown that the  $T \times T \times T$ model is the most physical one. Here, due to the discrete symmetry in all three directions, we can represent the infinite Universe as one finite cell with periodic boundary conditions in all dimensions. The finite volume of the cell enabled us to use the superposition principle and solve the Poisson equation for a single mass. The total potential in an arbitrary point of the cell is equal to the sum of potentials of all particles in the cell. Unfortunately, in the case of point-like gravitating sources the obtained solution has a very important drawback. Usually, it is expected that potentials diverge at the positions of masses. However, in the model under consideration the gravitational potential has no definite values on the edges joining identical masses in neighboring cells, i.e. at points where masses are absent. Clearly, this is a nonphysical result. Then, looking for a more physical solution, we smeared gravitating masses over some regions and showed that in this case the gravitational potential has regular behavior at any point inside the cell. Therefore, smearing represents the necessary condition of getting a regular gravitational potential in the lattice Universe. It is usually written that in N-body simulations some sort of smearing is used to avoid divergence at the positions of masses. Now, we have demonstrated that this procedure helps to avoid problems on the above mentioned edges as well. In the  $T \times T \times T$  model, particles in the cell may have different masses and be distributed arbitrarily. In the cases of topologies  $T \times T \times R$  and  $T \times R \times R$ , we cannot do this. We have shown that the only way to get a solution here is to suppose the periodic (in all dimensions) distribution of identical masses. However, such solution is reduced to the one obtained in the case of  $T \times T \times T$  topology. Therefore, first, this solution has a drawback inherent in this model, and, second, the distribution of masses looks very artificial.

#### Acknowledgements

The work of M. Eingorn was supported by NSF CREST award HRD-1345219 and NASA grant NNX09AV07A.

#### References

- R.W. Linquist and J.A. Wheeler, Dynamics of a Lattice Universe by the Schwarzschild-Cell methods, Rev. Mod. Phys. 29 (1957) 432.
- [2] D. Korotkin and H. Nicolai, A periodic analog of the Schwarzschild solution (1994); arXiv:gr-qc/9403029.
- [3] T. Clifton and P.G. Ferreira, Archipelagian Cosmology: dynamics and observables in a Universe with discretized matter content, Phys. Rev. D 80 (2009) 103503; arXiv:astro-ph/0907.4109.
- [4] T. Clifton and P.G. Ferreira, Errors in estimating  $\Omega_{\Lambda}$  due to the fluid approximation, JCAP 10 (2009) 026; arXiv:astro-ph/0908.4488.
- [5] T. Clifton, Cosmology without averaging, Class. Quant. Grav. 28 (2011) 164011; arXiv:gr-qc/1005.0788.
- [6] J.-P. Uzan, G.F.R. Ellis and J. Larena, A two-mass expanding exact space-time solution, Gen. Rel. Grav. 43 (2011) 191; arXiv:gr-qc/1005.1809.
- [7] T. Clifton, K. Rosquist and R. Tavakol, An exact quantification of backreaction in relativistic cosmology, Phys. Rev. D. 86 (2012) 043506; arXiv:gr-qc/1203.6478.
- [8] C.-M. Yoo, H. Abe, K.-I. Nakao and Y. Takamori, Black Hole Universe: construction and analysis of initial data, Phys. Rev. D. 86 (2012) 044027; arXiv:gr-qc/1204.2411.
- [9] E.G. Floratos and G.K. Leontaris, On topological modifications of Newton's law, JCAP 04 (2012) 024; arXiv:astro-ph/1202.6067.
- [10] J.-P. Bruneton and J. Larena, Dynamics of a lattice Universe, Class. Quant. Grav. 29 (2012) 155001; arXiv:gr-qc/1204.3433.
- [11] J.-P. Bruneton and J. Larena, Observables in a lattice Universe, Class. Quant. Grav. 30 (2013) 025002; arXiv:gr-qc/1208.1411.
- [12] M. Eingorn and A. Zhuk, Remarks on mechanical approach to observable Universe, JCAP 05 (2014) 024; arXiv:astro-ph/1309.4924.
- [13] C. Clarkson, G. Ellis, J. Larena and O. Umeh, Does the growth of structure affect our dynamical models of the universe? The averaging, backreaction and fitting problems in cosmology, Rept. Prog. Phys. 74 (2011) 112901; arXiv:astro-ph/1109.2314.
- [14] P.J.E. Peebles, The large-scale structure of the Universe, Princeton University Press, Princeton (1980).

- [15] J.S. Bagla, Cosmological N-Body simulation: techniques, scope and status, Curr. Sci. 88 (2005) 1088; arXiv:astro-ph/0411043.
- [16] V. Springel, The cosmological simulation code GADGET-2, Mon. Not. R. Astron. Soc. 364 (2005) 1105; arXiv:astro-ph/0505010.
- [17] M. Eingorn and A. Zhuk, Hubble flows and gravitational potentials in observable Universe, JCAP 09 (2012) 026; arXiv:astro-ph/1205.2384.
- [18] J. Adamek, R. Durrer and M. Kunz, N-body methods for relativistic cosmology (2014); arXiv:astro-ph/1408.3352.
- [19] M. Eingorn, A. Kudinova and A. Zhuk, Dynamics of astrophysical objects against the cosmological background, JCAP 04 (2013) 010; arXiv:astro-ph/1211.4045.
- [20] G.F.R. Ellis and G.W. Gibbons, Discrete Newtonian Cosmology: Perturbations (2014); arXiv:gr-qc/1409.0395.
- [21] J.D. Norton, The cosmological woes of Newtonian gravitation theory, in H. Goenner, J. Renn, J. Ritter and T. Sauer, eds., The expanding worlds of General Relativity: Einstein studies, volume 7, Boston: Birkhauser, pp. 271-323 (1999).
- [22] M. Eingorn, M. Brilenkov and B. Vlahovic, Zero average values of cosmological perturbations as an indispensable condition for the theory and simulations (2014); arXiv:astro-ph/1407.3244.
- [23] A.P. Prudnikov, Yu.A. Brychkov and O.I. Marichev, *Integrals and series*, Gordon & Breach Science Publishers, New York (1986).