

Dispersion of guided modes in two-dimensional split ring lattices

Per Lunnemann¹ and A. Femius Koenderink^{2,*}

¹*DTU Fotonik, Department of Photonics Engineering, Østedsplads 343, DK-2800, Denmark*

²*Center for Nanophotonics, FOM Institute AMOLF, Science Park 104, 1098 XG Amsterdam, The Netherlands*

(Dated: May 16, 2022)

We present a semi-analytical point-dipole method that uses Ewald lattice summation to find the dispersion relation of guided plasmonic and bianisotropic modes in metasurfaces composed of 2D periodic lattices of arbitrarily strongly scattering magneto-electric dipole scatterers. This method takes into account all retarded electrodynamic interactions as well as radiation damping selfconsistently. As illustration we analyze the dispersion of plasmon nanorod lattices, and of 2D split ring resonator lattices. Plasmon nanorod lattices support transverse and longitudinal in-plane electric modes. Scatterers that have an in-plane electric and out-of-plane magnetic polarizability, but without intrinsic magnetoelectric coupling, result in two bands that are mixtures of the bands of electric-only and magnetic-only lattices. Thereby bianisotropy through mutual coupling, in absence of building-block bianisotropy, is evident. Once strong bi-anisotropy is included in each building block, the Bloch modes become even more strongly magnetoelectric. Our results are important to understand spatial dispersion and bianisotropy of metasurface and metamaterial designs.

I. INTRODUCTION

Periodic structures of scatterers have a long standing history in photonics, traditionally in guise of diffraction gratings¹⁻³ and photonic crystals,^{4,5} and more recently in the context of plasmonics, metamaterials, and metasurfaces.⁶⁻⁸ In plasmonics with noble metal particles that support localized resonances, periodic chains of particles with subdiffraction pitch were already proposed in 1998⁹ and demonstrated in 2003¹⁰ as potential candidates for guiding signals in a deep subwavelength fashion through near-field dipole-dipole interaction.⁹⁻¹⁵ While transport in these systems is very lossy, the exact formalism to describe the guiding mechanism in presence of long range retarded dipole-dipole interactions has remained a topic of ongoing work.^{9,11-13,16} This is in part due to the associated mathematical intricacies¹⁷ and in part to the fact that transport along chains of Lorentzian dipole resonators transcends plasmonics in relevance. Two-dimensionally periodic systems of plasmon particles have traditionally been studied in case of diffractive lattices, in which case grating anomalies hybridize with localized surface plasmon resonances to give very sharp spectral features.¹⁸⁻²¹ These features have been pursued for field-enhanced spectroscopies,^{22,23} sensing,^{24,25} as well as improved solid-state light sources.²⁶⁻²⁸ Since the advent of 2D metamaterial arrays the response of subdiffraction pitch lattices of resonant scatterers has gained significantly in relevance.^{7,8,29}

While experimental studies of 2D metamaterials and metasurfaces usually probe transmission and reflection for some definite incident polarization and wave vector,³⁰⁻³² the fundamental underlying property of a lattice must be its dispersion relation or band structure, which summarizes the existence of guided as well as leaky modes. The spectrum of leaky modes supported by a lattice of split rings, for instance, would explain the origin of angle-dependent transmission and reflection features,³³

and would form an excellent basis to understand spatial dispersion in attributed effective material constants.³⁴⁻³⁶ Complementary to the leaky modes, the guided mode structure would also be relevant, for instance for the proposed ‘lasing spaser’²⁸ where a 2D metamaterial lattice is immersed in a gain medium, or when coupling a localized fluorescent source to a lattice in the near field. In this case, the modes subject to most gain, or the modes with strongest coupling to the source, need not correspond to resonances identified in normal incidence scattering experiments. Rather, any guided modes supported by the lattice could be excited in any experiment that does not a priori restrict or impose parallel wave vector. An understanding of the band structure of 2D lattices of magnetic and electric resonant scatterers is therefore important for metasurface research.

Previously, modal band structures of a 2D lattice of electric dipolar spherical scatterers have been theoretically treated,^{16,37} while only the leaky modes of SRR lattices have so far been assessed through transmission calculations and compared to experiments^{30,31,33} In this paper we present a method to calculate band structures for arbitrary lattices of arbitrary magneto-electric dipolar scatterers, and illustrate its properties for simple lattices of plasmon bars, as well as idealized split rings.

II. LATTICE RESPONSE

We consider a 2D lattice consisting of arbitrary magnetoelectric point scatterers in the dipole approximation, however, without making any electrostatic approximation. Each particle is described by a polarizability tensor, $\vec{\alpha}$ that relates the induced electric and magnetic dipole moment, \mathbf{p} and \mathbf{m} , to a driving electric and magnetic

field \mathbf{E} and \mathbf{H} according to^{38,39}

$$\begin{pmatrix} \mathbf{p} \\ \mathbf{m} \end{pmatrix} = \overleftrightarrow{\alpha} \begin{pmatrix} \mathbf{E} \\ \mathbf{H} \end{pmatrix}. \quad (1)$$

The magnetoelectric polarizability may be decomposed as

$$\overleftrightarrow{\alpha} = \begin{pmatrix} \overleftrightarrow{\alpha}_{EE} & \overleftrightarrow{\alpha}_{EH} \\ \overleftrightarrow{\alpha}_{HE} & \overleftrightarrow{\alpha}_{HH} \end{pmatrix}, \quad (2)$$

where $\overleftrightarrow{\alpha}_{EE}$ is the 3×3 electric polarizability tensor that quantifies the induced electric dipole moment in response to an electric field. Similarly, $\overleftrightarrow{\alpha}_{HH}$ describes the magnetic polarizability that quantifies the induced magnetic dipole in response to a magnetic driving field. Finally, $\overleftrightarrow{\alpha}_{EH} = -\overleftrightarrow{\alpha}_{HE}^T$ denotes the magnetoelectric coupling that describes the induced electric dipole moment in response to a magnetic field and vice versa. This element controls the bi-anisotropy³⁸ of the medium giving rise to chiral extinction under oblique incidence.^{39,40} We shall denote $\overleftrightarrow{\alpha}$ the bare polarizability, since it describes the induced dipole moments in the absence of neighbouring point scatterers. $\overleftrightarrow{\alpha}$ is subject to reciprocity and energy conservation constraints discussed in Ref. 39 and 41. We construct the electrodynamically consistent polarizability of a single scatterer, bound by the optical theorem, by addition of radiation damping

$$\overleftrightarrow{\alpha}^{-1} = \overleftrightarrow{\alpha}_0^{-1} - \frac{2}{3}k^3i\mathbb{I}, \quad (3)$$

to an electrostatic bare polarizability tensor $\overleftrightarrow{\alpha}_0$ which can for instance be derived from an LC model. Here k denotes the wave number, \mathbb{I} is the 6-dimensional identity tensor and \cdot^{-1} denotes matrix inversion. In this work we will illustrate band structure calculations by considering a specific group of scatterers representative of plasmon bars and of many metamaterial scatterers like split ring resonators (SRRs). In particular, we assume the only available responses to be electric along the x -direction (along the bar, or split, see fig. 1.) and/or magnetic along the z -direction (direction through the SRR loop), setting all other tensor elements to zero. I.e.

$$\overleftrightarrow{\alpha}_0 = \mathcal{L}(\omega) \begin{pmatrix} \eta_E & 0 & \dots & 0 & i\eta_C \\ 0 & 0 & & & 0 \\ \vdots & & \ddots & & \vdots \\ 0 & & & 0 & 0 \\ -i\eta_C & 0 & \dots & 0 & \eta_H \end{pmatrix}, \quad (4)$$

where $\mathcal{L}(\omega)$ is a Lorentzian pre-factor

$$\mathcal{L}(\omega) = V \frac{\omega_0^2}{\omega_0^2 - \omega^2 - i\omega\gamma}, \quad (5)$$

typical for a plasmon resonance or LC circuit model, where V is the physical volume of the scatterer, γ is the damping rate due to Ohmic losses, ω_0 denotes the




FIG. 1. Illustration of the considered lattice, here sketched for split ring resonators, with a plane wave incident with an in-plane wave vector k_{\parallel} .

resonance frequency, and $\eta_{E,H,C}$ are real dimensionless parameters that for LC circuits can be calculated from geometry. Recalling that the extinction cross section of a simple scatterer with scalar polarizability α is $\sigma_{\text{ext}} = 4\pi k \text{Im}(\alpha)$, we note that for tensorial polarizability, the extinction cross section varies with incidence condition, but is always a linear combination of the imaginary part of the tensor eigenvalues, α_1 and α_2 , of eq. (3). The corresponding eigenvectors may possess a magnetoelectric character, having both a component along p_x and m_z through intrinsic magneto-electric coupling when $\eta_C \neq 0$.³⁹ Eigenvectors with a phase-offset between p_x and m_z results in a ‘pseudochirality’, *i.e.*, a handedness-dependent extinction for some incidence angles.

The optical response of 2D periodic lattices of electric polarizabilities has been didactically reviewed by de Abajo⁴² and extended to full magneto-electric case in Refs. 30 and 43. For consistency we recapitulate the main findings. Consider a 2D periodic lattice of point scatterers placed at $\mathbf{R}_{mn} = m\mathbf{a}_1 + n\mathbf{a}_2$ (where m and n are integers, and $\mathbf{a}_{1,2}$ are the real space basis vectors, see Fig. 1). The response of a particle at position \mathbf{R}_{mn} is self-consistently set by the incident field, plus the field of all other dipoles in the lattice according to⁴²

$$\begin{pmatrix} \mathbf{p}_{mn} \\ \mathbf{m}_{mn} \end{pmatrix} = \overleftrightarrow{\alpha} \left[\begin{pmatrix} \mathbf{E}_{\text{in}}(\mathbf{R}_{mn}) \\ \mathbf{H}_{\text{in}}(\mathbf{R}_{mn}) \end{pmatrix} + \sum_{m' \neq m, n' \neq n} \overleftrightarrow{G}^0(\mathbf{R}_{mn} - \mathbf{R}_{m'n'}) \begin{pmatrix} \mathbf{p}_{m'n'} \\ \mathbf{m}_{m'n'} \end{pmatrix} \right] \quad (6)$$

where $\overleftrightarrow{G}^0(\mathbf{R}_{mn} - \mathbf{R}_{m'n'})$ is the 6×6 dyadic Green function of the medium surrounding the lattice. Plane wave incidence with parallel wave vector \mathbf{k}_{\parallel} allows a Bloch wave form $(\mathbf{p}_{mn}, \mathbf{m}_{mn})^T = e^{i\mathbf{k}_{\parallel} \cdot \mathbf{R}_{mn}} (\mathbf{p}_{00}, \mathbf{m}_{00})^T$ to obtain

$$\begin{pmatrix} \mathbf{p}_{00} \\ \mathbf{m}_{00} \end{pmatrix} = [\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{G}^{\neq}(\mathbf{k}_{\parallel}, 0)]^{-1} \begin{pmatrix} \mathbf{E}_{\text{in}}(\mathbf{R}_{00}) \\ \mathbf{H}_{\text{in}}(\mathbf{R}_{00}) \end{pmatrix} \quad (7)$$

Here, $\overleftrightarrow{\mathcal{G}}^\neq(\mathbf{k}_\parallel, 0)$ is a summation of the dyadic Green function \overleftrightarrow{G}^0 over all positions in the lattice barring the origin:

$$\overleftrightarrow{\mathcal{G}}^\neq(\mathbf{k}_\parallel, \mathbf{r}) = \sum_{m \neq 0, n \neq 0} \overleftrightarrow{G}^0(\mathbf{R}_{mn} - \mathbf{r}) e^{i\mathbf{k}_\parallel \cdot \mathbf{R}_{mn}} \quad (8)$$

In this work, we take the surrounding medium, that defines \overleftrightarrow{G}^0 , to be homogeneous. Implementation of the sum of \overleftrightarrow{G}^0 was carried out using the Ewald lattice summation technique,¹⁷ that consists of splitting a poorly convergent sum, like eq. (8), into two exponentially convergent sums as summarized in Refs. 30 and 43. The same techniques, can be extended to lattices in stratified dielectric systems, complex unit cells, and stacks of lattices.⁴³

The factor $[\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{\mathcal{G}}^\neq(\mathbf{k}_\parallel, 0)]^{-1}$ in Eq. (7) is identified as an *effective* polarizability tensor, $\overleftrightarrow{\alpha}_{\text{eff}}$, of the scatterer, renormalized by the scattering from all other lattice sites. In a lossless system the Bloch wave dispersion would correspond to those frequencies for which $\det(\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{\mathcal{G}}^\neq) = 0$, or equivalently, those frequencies for which α_{eff} has a pole. These two dimensional lattices in fact have radiative (if $k_\parallel \leq \omega/c$) and Ohmic loss. While one could in principle seek complex frequencies for which $\det(\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{\mathcal{G}}^\neq) = 0$,¹¹ we simply evaluate $\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{\mathcal{G}}^\neq$ for real ω and real k_\parallel . In particular we consider the imaginary part of the eigenvalues of $[\overleftrightarrow{\alpha}^{-1} - \overleftrightarrow{\mathcal{G}}^\neq]^{-1}$ as they directly relate to extinction. Distinct bands emerge, the width of which we identify as the damping rate³⁷ due to both Ohmic and radiation damping. For the effectively 2×2 form of the single-particle polarizability that we use in this paper (Eq. (4)), for each (ω, k_\parallel) -point we obtain at most two nontrivial eigenvectors and eigenvalues. To separate the dispersion in bands, we group eigenvalues by continuity of the projection of the corresponding eigenvectors with the eigenvectors of neighboring (ω, k_\parallel) -points.

III. RESULTS

As illustration we consider four types of scatterers, starting with plasmon bars and culminating at a realistic description for split rings. It has been experimentally demonstrated that SRRs are well described with a dipolar polarizability as in eqs. (4) and (3) provided one takes bianisotropy η_C at the upper limit $\sqrt{\eta_E \eta_H}$.^{30,39,40} To help understand what effect magnetoelectric coupling has, we consider three sub-cases prior to analyzing the SRR lattice with maximum intrinsic coupling. These consist of (1) plasmon bars along x (only $\eta_E \neq 0$), (2) magnetic dipolar antennas along z (only $\eta_H \neq 0$) and (3) uncoupled SRRs without bi-anisotropy (setting $\eta_E = \eta_H = 1$ and $\eta_C = 0$). Throughout the paper we use parameter values as stated in table I. To more clearly resolve the modes, we take an Ohmic damping rate $\gamma \sim 100$ times less than that of gold. The parameters yield

TABLE I. Chosen parameter values.

Parameter	Value	Description
V	$(80 \text{ nm})^3$	Physical volume of scatterer.
γ	$1 \cdot 10^{12} \text{ s}^{-1}$	Damping rate.
$a_1 = a_2$	300 nm	Lattice constant.
ω_0	$2\pi c/1.5 \text{ } \mu\text{m}$	Resonance frequency




FIG. 2. (Color online) Calculated effective polarizability, $\log[\text{Im}(\alpha)/V]$ as a function of k_\parallel and ω for a) an in-plane bar-type electric structure as illustrated in the inset with $\eta_H = \eta_C = 0$ and $\eta_E = 1$ in eq. (4) and b) a transverse bar-type magnetic structure as illustrated in the inset with $\eta_E = \eta_C = 0$ and $\eta_H = 1$ in eq. (4).

an extinction cross section per scatterer of $0.2 - 0.3 \text{ } \mu\text{m}^2$, comparable to measured values.^{30,44}

A. Scalar anisotropic scatterers

We first consider plasmon bars with bare polarizability given by eq. (4) and setting $\eta_H = \eta_C = 0$ and $\eta_E = 1$. In figure 2a) we present the calculated dispersion diagram sweeping (k_x, k_y) through the reduced Brillouin zone along the following path: $(\pi/d, \pi/d) \rightarrow (0, \pi/d) \rightarrow (0, 0) \rightarrow (\pi/d, 0) \rightarrow (\pi/d, \pi/d)$ also denoted $M \rightarrow Y \rightarrow \Gamma \rightarrow X \rightarrow M \rightarrow \Gamma$. The discontinuity on each side of Γ indicates the light line. The sharp resonances

below the light line are guided modes of the lattice. Two distinct modes are observed on either side of the Γ -point. Within the domain $\Gamma \rightarrow Y$, the dipole phase is constant along \hat{x} and varies along \hat{y} , transverse to the dipole moment orientation. Thus, this mode is a transverse in-plane electric (TIE) mode, with dipole moments perpendicular to k_{\parallel} . The heads to tail arrangement along \hat{x} and alternating direction when going along \hat{y} results in a redshift of the resonance as has previously been explained using simple hybridization models for electrostatic 1D and 2D systems.^{11–13,45,46} In such a hybridization model, resonance shifts of coupled dipoles can be understood from considering the quasi-static interaction energy, U , between two dipoles \mathbf{p}_1 and \mathbf{p}_2 separated by a vector \mathbf{r}

$$U \propto \frac{\mathbf{p}_1 \cdot \mathbf{p}_2 - 3[\mathbf{p}_1 \cdot \hat{\mathbf{r}}][\mathbf{p}_2 \cdot \hat{\mathbf{r}}]}{r^3}. \quad (9)$$

Accordingly, longitudinal coupling of parallel (antiparallel) dipoles leads to redshifts (blueshifts), while for transversely coupled dipoles the coupling strength is reduced and opposite in sign. As caveat we note that such hybridization models strictly apply only in electrostatics, whereas here we treat retarded interactions between lossy, resonant dipoles.

Analogously, within the domain $\Gamma \rightarrow X$ all dipoles point along the wave vector, and the mode is therefore longitudinal in-plane electric (LIE).³⁷ In this configuration, the head to head arrangement along \hat{x} and the fixed direction along \hat{y} results in a blueshift. For the regions $Y \rightarrow M$ and $X \rightarrow M$ the mode possesses a mixed transverse and longitudinal character. For wave vectors within the light cone, we notice a faint resonance. This resonance is very broad due to radiation damping, and is the resonance that is probed in farfield transmission spectra.^{30–32} Compared to the single particle radiative linewidth, the collective resonance linewidth is broader by more than an order of magnitude. This collective superradiant damping effect has been observed experimentally in density dependent studies of transmission at normal incidence for 2D lattices.³¹ Since the time-averaged far field flux from a single dipole pointing along \hat{x} is proportional to $\sin^2(\theta)$ with θ being the azimuthal angle along x ,⁴⁷ scattering out of the lattice plane is strong for in-plane modes. For longitudinal modes this damping monotonically reduces with increasing k_{\parallel} as dipoles do not radiate along their axis.^{11,13} For transverse modes the radiation damping is constant or increases with k_{\parallel} when approaching the light line. Finally, the modes far below the light line have a constant width comparable to the single particle Ohmic damping rate γ . However, very close to the light line, the damping rates drop well below the Ohmic damping rate indicating that the modes near the light line are very weakly confined. As is the case for, e.g., a thin dielectric slab in a symmetric host environment, even a plane of weakly polarizable particles binds a guided mode, however with a very large fraction of its energy density in air. We note the strong similarities to the calculated dispersion of 1D as well as 2D lattices of

spherical electric scatterers.^{11,13,37}

We now turn to the (hypothetical) case of out-of-plane magnetic scatterers with bare polarizability set by $\eta_E = \eta_C = 0$ and $\eta_H = 1$ in eq. (4). In fig. 2b) we present the calculated effective polarizability. The dispersion is symmetric about Γ owing to the rotational symmetry of this lattice. Obviously, the observed mode is a transverse magnetic (TM) mode. As opposed to the TIE and LIE modes in fig. 2, we can clearly resolve a resonance above the light line. For a single transverse dipole the radiated intensity perpendicular to the lattice is zero, while radiation in the plane is strong.⁴⁷ Hence for the array at $k_{\parallel} = 0$ there is no radiative loss. The increasing broadening when going from $k_{\parallel} = 0$ to the light line, was also claimed for 1D chains of plasmonic particles.^{11,13} As the wave vector sweeps to the light line, the radiative loss, which by momentum conservation has the same in-plane wave vector, has an increasingly good overlap with the single dipole radiation pattern, thereby causing the radiative loss to increase from 0 at $k_{\parallel} = 0$ to large values.

Comparing fig. 2a) and 2b), the TIE and TM modes are seen to converge asymptotically to the light line, while the LIE-mode crosses the light line. For the transverse modes all dipoles are perpendicular to the propagation vector and therefore couple strongly to free photons with an anti-crossing as a result. For the LIE mode all dipoles are parallel to the wave vector and thus hindered from coupling to the far field. Consequently no anti-crossing is observed in good agreement with previous results on 1D and 2D arrays of scatterers.^{11,13,37} Finally, we note that the modes of the out-of-plane magnetic antennas are much less dispersive than those of the electric in-plane antennas showing an almost flat band between Y-M and X-M. Inspecting the interaction energy in the magneto-static equivalent of eq. (9), we note that for the TM mode near Y (X) a blueshift is induced from adjacent parallel dipoles along \hat{x} (\hat{y}) while a redshift of equal magnitude is induced from adjacent anti-parallel dipoles along \hat{y} (\hat{x}), leading to net cancellation of the hybridization energies. This cancellation of nearest-neighbor contributions holds for any periodicity, i.e., also for retarded interactions. To the contrary, for the electric in-plane antennas near Y (X), the heads to tail arrangement along \hat{x} (\hat{y}) and anti-parallel arrangement along \hat{y} (\hat{x}) both contribute with a red (blue) shift. Hence any path connecting Y and X exhibits a larger variation in frequencies compared to the TM mode.

B. Split ring resonators

We now turn our attention to SRR-type scatterers. We describe the per-building block magnetoelectric coupling by the parameter η_C in eq. (4). Energy conservation dictates that η_C is bound by $|\eta_C| \leq \sqrt{\eta_E \eta_H}$,³⁹ where equality holds for a truly planar scatterer described as a single resonant circuit. It has been demonstrated experimentally, and by full-wave simulations^{30,39,40,48} that

real SRRs indeed possess a coupling strength η_C close to the upper limit $\sqrt{\eta_E\eta_H}$. We note, that for more complicated scatterers though, i.e. nested split rings, lower cross couplings can occur.^{40,48} For clarity we shall first consider the two cases of absent and partial coupling between the magnetic and electric dipole by setting $\eta_C = 0$ and $\eta_C = 0.5\sqrt{\eta_E\eta_H}$, respectively, in eq.(4) before finally considering the realistic case with full cross coupling $\eta_C = \sqrt{\eta_E\eta_H}$. The calculated dispersion diagrams are presented in figure 3.

1. SRR without cross coupling (anisotropic)

We start by considering the case of no electro-magnetic cross coupling, $\eta_C = 0$ and $\eta_E = \eta_H = 1$. In this case, for each (k_{\parallel}, ω) -pair two non-trivial eigenvalues exist. In fig. 3a) the calculated sum of imaginary part of eigenvalues, $\text{Im}\alpha_1 + \text{Im}\alpha_2$ is plotted. We immediately identify that the dispersion diagram resembles the superposition of those in fig. 2 for the purely electric, and purely magnetic objects. One mode traces the in-plane mode in fig. 2a) while we observe some differences between the other mode and the TM mode in fig. 2b), especially for the region M-Y- Γ . These differences arise from inter-particle coupling between electric dipoles, magnetic dipoles and between electric and magnetic dipoles.

To clarify how fig. 3a) and fig. 2b) (TM mode) differ we define an electromagnetic mixing ratio as

$$\zeta_j = |p_{j,x}| - |m_{j,z}|, \quad (10)$$

where $p_{j,x}$ ($m_{j,z}$) is the electric x (magnetic z) component of the j th normalized eigenvector. For $\zeta = -1$ the mode is purely TM while for $\zeta = 1$ the mode is purely in-plane electric. The mixing ratio of the bands in fig. 3a) is presented in fig. 3d). Generally the two modes are clearly identifiable as strongly electric resp. magnetic, as expected if the dispersion were that of an uncoupled electric lattice and magnetic lattice. In fact, the modes are *purely* in-plane electric, respectively TM, at all the symmetry point M, Y, Γ , and X. A strongly mixed character occurs close to the light line, in particular midway Γ and Y, and at the light line crossing between Γ and M, commensurate with the fact that there the mode must match a plane wave propagating in the array plane, which carries both out-of-plane H and in plane E .

Before we introduce cross coupling η_C , we consider the hybridization interaction between magnetic and electric dipoles. In fig. 4 we illustrate the spatial distribution of magnetic dipoles and their associated electric fields at the point in k_{\parallel} -space $Y = (0, \pi/2d)$ and the point midway between Γ and Y, denoted as $Y/2$. Considering the field lines at Y, depicted in fig. 4a), at the location of the central SRR the electric fields from adjacent magnetic moments cancel. We therefore conclude that at Y the lattices of magnetic and electric dipoles are essentially decoupled, in good agreement with the unit value of the mixing ratio in fig. 3d). A similar analysis holds for the

X-point. Considering the point $Y/2$ in fig. 4b), the electric field lines of magnetic dipoles adjacent to a central site add up along \hat{x} and from this we infer that the TIE and TM mode strongly mix at $Y/2$. This is in contrast to the point $X/2$, where the field lines add up along \hat{y} , along which the SRR is not polarizable. Hence no dipole moment is induced and one mode is therefore purely TM, while the other is purely electric in nature. The same results for the absence/presence of magnetoelectric cross-coupling is obtained by starting the hybridization analysis from magnetic fields due to in-plane electric dipoles rather than vice versa, as expected from reciprocity.

To conclude, even if one starts with building blocks that have no magnetoelectric coupling, once placed in a dense lattice, the collective modes have mixed electric and magnetic character except at symmetry points.

2. SRR with cross coupling (bi-anisotropic)

Most realized metamaterial lattices will consist of building blocks possessing an intrinsic magneto-electric coupling that couples the excitation of electric and magnetic dipoles in a single building block according to a definite amplitude and phase relation. We consider partial electromagnetic cross coupling setting $\eta_C = 0$ and $\eta_E = \eta_H = 1$ in fig. 3b). Comparing with the case $\eta_C = 0$ in fig. 3a), we immediately see a close resemblance, apart from a clearly resolved anti-crossing midway between M and Y. The calculated mixing ratio (see figure 3e)) evidences that the modes are no longer purely electric or purely magnetic at any of the symmetry points. Near the anti-crossing, the mixing ratio becomes 0 implying that the two modes carry equal electric and magnetic content. The associated complex phase difference defined as

$$\Delta\phi_j = \arg p_{j,x} - \arg m_{j,z} \quad (11)$$

is, at the anti-crossing point, found to be $\Delta\phi_1 = \pi/2$ for the lower band and $\Delta\phi_2 = -\pi/2$ for the upper band. This distinct phase difference between the two anticrossing bands signifies that one solution has the electric dipole a quarter cycle in advance of the magnetic one, while for the other solution the electric dipole lags the magnetic dipole by a quarter cycle. This distinction stems from the physics intrinsic to the single bi-anisotropic scatterer, wherein the polarizability tensor of a cross-coupled scatterer has two distinct eigenvalues, corresponding to high, respectively low scattering strength, with eigensolution corresponding to either an advanced or lagging electric dipole relative to the magnetic response. In LC -circuit terms, a strong difference in response to a driving field with either an advance or lag can be understood by noting that in the driving $E_x d + i\omega H_z A$ (d the capacitor gap, A the loop area) the electric term driving the capacitor and the electromotive force due to a changing flux, either add or cancel depending on phase. In a scattering experiment this results in a strongly handed response under oblique incidence, since




FIG. 3. (Color online) a-c) Calculated sum of eigenvalues, $\log[\text{Im}(\alpha_1 + \alpha_2)/V]$, as a function of wave vector, k_{\parallel} , and normalized frequency ω/ω_0 for a SRR lattice with a) cross coupling $\eta_C = 0$ and b) $\eta_C = 0.5$, and c) $\eta_C = 1$. d-f) calculated mixing ratio $\zeta = |p_x| - |m_z|$ of the eigenvectors for the extracted bands. Blue corresponds to a pure electric dipole while red corresponds to pure magnetic dipole. White is a balanced mix of magnetic and electric dipoles. Dashed black lines indicate the light lines.




FIG. 4. (Color online) Illustration of the distribution of magnetic dipoles and their associated electric fields in a SRR lattice with $\eta_C = 0$ for wave vectors at a) Y and b) Y/2 as indicated by a blue arrow in the dispersion map inset. Red circles with a dot (cross) indicate a magnetic dipole pointing out of (into) the paper.

oblique incidence circular polarization carries a phase difference between E_x and H_z .^{30,40} Returning to the physics of the lattice, given the Bloch wave vector $k_{\parallel} = (\text{M} + \text{Y})/2$ one can explicitly calculate the fields exerted on a central dipole by all its neighbors. We indeed find that when $p_x = im_z$ respectively $p_x = -im_z$, the overlap of the fields a central dipole scatterer receives from its neighbors in the lattice is very different in strength (addition resp. cancellation in terms of $E_x d + i\omega H_z A$).

Finally, we consider the maximally coupled case, $\eta_C = \eta_E = \eta_H = 1$, presented in figure 3c,f). The most remarkable aspect is that one of the two bands vanishes, leaving only one band. That this must happen is easily understood by noting that at maximum cross coupling $\eta_C = \sqrt{\eta_E \eta_H}$, one of the two eigenvalues of the single-object polarizability tensor vanishes. In other words, a

maximally crosscoupled SRR has just one mode of oscillation and not two, in which furthermore the relative phase and amplitude of p and m are locked. A didactic example is an LC circuit, which has just one resonant mode of oscillation where the same circulating charge gives rise to both p_x and m_z in a fixed phase and amplitude ratio. This intuition for the LC circuit is only reconcilable with a 2×2 polarizability tensor if one eigenvalue vanishes. By extension, this also means that a lattice of maximally cross-coupled scatterers presents only one band, not two. A remarkable observation is that the band structure is similar to that of the electric bar structure presented in fig. 2 as well as to the band of the uncoupled SRRs that is assigned as primarily electric in fig. 3(a,d). This similarity is strong for wave vectors well beyond the light line, while differences are large for wave vectors well within the light line. That the optical response within the light line must be very different between bars and split rings is well understood: bars do not show a handed response in transmission while split rings do,³⁰ directly evidencing large observable differences between electric only, and magnetoelectric objects. Well beyond the light line, the dispersion is surprisingly similar. This similarity only holds for the (ω, k_{\parallel}) -relation but not for the associated eigenfunctions, since the eigenfunctions necessarily show equally strong p and m with a quarter cycle phase slip at the maximum cross coupling condition, whereas $m_z = 0$ in the electric-only case. As regards the similarity in frequency dispersion, it is tempting to push the hybridization arguments based on electrostatic potential $U_E = -\langle \text{Re} \mathbf{E} \cdot \text{Re} \mathbf{p} \rangle$ to some magnetoelectric equivalent. If one takes energy as $U_{EH} = -\langle \text{Re}(\mathbf{p}, \mathbf{m}) \cdot \text{Re}(\mathbf{E}, \mathbf{H}) \rangle$ (where $\langle \cdot \rangle$ indicates cycle-averaging), evaluation shows that $U_E(\mathbf{k}_{\parallel})$ and $U_{EH}(\mathbf{k}_{\parallel})$ are qualitatively similar for

our parameters, with the same extrema at X and Y , though with deviations at Γ and M . While this correspondence is consistent with the similar dispersion in ω , in our view, this is fortuitous and a comparison through static dipole hybridization should not be too vigorously pursued. Generalizing potential energy arguments to complex fields and lossy, dispersive scatterers is firstly problematic, and secondly presents little benefit since ultimately our work shows that in lattices retarded, long-range interactions are so important, that leaving them out to obtain a hybridization model even simpler than our point-dipole theory is a perilous affair.

IV. CONCLUSION

To conclude, we discussed a method based on magneto-electric point-dipole interactions and Ewald lattice summation to approximate the dispersion relation of two dimensional lattices of bi-anisotropic scatterers, accounting for all retarded electrodynamic interparticle interactions. Our results show that simple square lattices of plasmon bars that are dense, i.e. of subdiffraction pitch, support a mode structure characterized by weakly confined guided modes with a dispersion very close to the light line for frequencies to the red of the single scatterer resonance, and tightly confined guided modes at wave vectors well away from the light line. These modes are dispersive in a manner similar to results obtained previously for 1D plasmon chains, with the added complication that modes can have a mixed transverse and longitudinal character. Lattices of scatterers that have a intrinsically decoupled electric and magnetic polarizability in each element, will have a dispersion in which modes have a mixed magneto-electric character. Furthermore, we reported how introduction of bi-anisotropy in each building block modifies the dispersion. For full cross coupling a single mode prevails with the electric and magnetic dipoles being interlocked with equal magnitude and a fixed $\pi/2$ phase.

As outlook, while we presented results for simple square lattices of split ring type resonators with just an in plane electric moment and out of plane magnetic moment, the method is easily generalized to deal with arbitrarily complex multi-element lattices of arbitrary

magneto-electric scatterers, provided that the dipole approximation is met. Thus our method is important for many structures, including metasurface designs with complex unit cells that comprise many elements. Since the method is fast, it should thus be possible to screen many different lattice symmetries and arrangements in the unit cell for desirable properties, such as minimized bi-anisotropy and spatial dispersion. As regards actual measurements of such dispersion relations, we note that measurements in the visible domain would likely be hampered by strong Ohmic damping, and the drawback that the Brillouin zone extends to very large wavevectors, rendering even near-field microscopy impractical. These are exactly the drawbacks that have made it impossible to verify the projected dispersion of 1D plasmon chains beyond the light line.^{14,15} However, in the radio frequency domain, the projected dispersion relations should be readily available. Finally we believe that the calculated dispersion relations should be important for lasing spaser experiments, in which lattices are studied in presence of gain to achieve lasing. The modes with the best tradeoff between loss and confinement should be found close to and just below the light line, rather than at the $k_{\parallel} = 0$ point, which was proposed as the lasing mode originally.²⁸ Furthermore, from work on Yagi-Uda antennas in the optical domain it is well known that directional scattering, and directional emission of embedded emitters is strongly linked to the modes of 1D particle chains just below the light line.^{49,50} Similarly, for 2D arrays our work points at design strategies for shaping directional emission.

ACKNOWLEDGMENTS

We are grateful to Yuntian Chen and Lutz Langguth for insightful comments. This work is part of the research program of the Stichting voor Fundamenteel Onderzoek der Materie (FOM), which is financially supported by the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO). AFK. gratefully acknowledges a NWO-VIDI grant for financial support. PL acknowledges support by the Carlsberg Foundation as well as the Danish Research Council for Independent Research (Grant No. FTP 11-116740).

* fkoenderink@amolf.nl

¹ R. Wood, *Phys. Rev.* **48**, 928 (1935).

² U. Fano, *J. Opt. Soc. Am.* **31**, 213 (1941).

³ E. G. Loewen and E. Popov, *Diffraction Gratings and Applications*, edited by B. J. Thompson (CRC Press, New York, USA, 1997).

⁴ E. Yablonovitch, *Phys. Rev. Lett.* **58**, 2059 (1987).

⁵ J. D. Joannopoulos, P. R. Villeneuve, and S. Fan, *Nature* **386**, 143 (1997).

⁶ S. Linden, J. Kuhl, and H. Giessen, *Phys. Rev. Lett.* **86**,

4688 (2001).

⁷ N. I. Zheludev and Y. S. Kivshar, *Nat. Mater.* **11**, 917 (2012).

⁸ A. V. Kildishev, A. Boltasseva, and V. M. Shalaev, *Science* **339**, 1232009 (2013).

⁹ M. Quinten, a. Leitner, J. R. Krenn, and F. R. Aussenegg, *Opt. Lett.* **23**, 1331 (1998).

¹⁰ S. A. Maier, P. G. Kik, H. A. Atwater, S. Meltzer, E. Harel, B. E. Koel, and A. A. G. Requicha, *Nat. Mater.* **2**, 229 (2003).

- ¹¹ A. F. Koenderink and A. Polman, *Phys. Rev. B* **74**, 033402 (2006).
- ¹² M. L. Brongersma, J. W. Hartman, and H. A. Atwater, *Phys. Rev. B* **62**, R16356 (2000).
- ¹³ W. H. Weber and G. W. Ford, *Phys. Rev. B* **70**, 125429 (2004).
- ¹⁴ R. de Waele, A. F. Koenderink, and A. Polman, *Nano Lett.* **7**, 2004 (2007).
- ¹⁵ T. Yang and K. B. Crozier, *Opt. Express* **16**, 8570 (2008).
- ¹⁶ A. L. Frutos, S. Campione, F. Capolino, and F. Mesa, *J. Opt. Soc. Am. B* **28**, 1446 (2011).
- ¹⁷ C. M. Linton, *SIAM Rev.* **52**, 630 (2010).
- ¹⁸ G. Vecchi, V. Giannini, and J. Gómez Rivas, *Phys. Rev. B* **80**, 201401 (2009).
- ¹⁹ G. Vecchi, V. Giannini, and J. Gómez Rivas, *Phys. Rev. Lett.* **102**, 146807 (2009).
- ²⁰ B. Auguié and W. L. Barnes, *Phys. Rev. Lett.* **101**, 143902 (2008).
- ²¹ V. G. Kravets, F. Schedin, and A. N. Grigorenko, *Phys. Rev. Lett.* **101**, 087403 (2008).
- ²² R. F. Aroca, *Phys. Chem. Chem. Phys.* **15**, 5355 (2013).
- ²³ L. Langguth and A. F. Koenderink, *Opt. Express* **22**, 15397 (2014).
- ²⁴ J. N. Anker, W. P. Hall, O. Lyandres, N. C. Shah, J. Zhao, and R. P. Van Duyne, *Nat. Mater.* **7**, 442 (2008).
- ²⁵ S. S. Aćimović, M. P. Kreuzer, M. U. González, and R. Quidant, *ACS Nano* **3**, 1231 (2009).
- ²⁶ S. R. K. Rodriguez, S. Murai, M. A. Verschuuren, and J. G. Rivas, *Phys. Rev. Lett.* **109**, 166803 (2012).
- ²⁷ G. Lozano, D. J. Louwers, S. R. Rodríguez, S. Murai, O. T. A. Jansen, M. A. Verschuuren, and J. Gómez Rivas, *Light Sci. Appl.* **2**, e66 (2013).
- ²⁸ N. I. Zheludev, S. L. Prosvirnin, N. Papasimakis, and V. A. Fedotov, *Nat. Photonics* **2**, 351 (2008).
- ²⁹ A. Polman, *Science* **322**, 868 (2008).
- ³⁰ P. Lunnemann, I. Sersic, and A. F. Koenderink, *Phys. Rev. B* **88**, 245109 (2013).
- ³¹ I. Sersic, M. Frimmer, E. Verhagen, and A. F. Koenderink, *Phys. Rev. Lett.* **103**, 213902 (2009).
- ³² C. Enkrich, M. Wegener, S. Linden, S. Burger, L. Zschiedrich, F. Schmidt, J. F. Zhou, T. Koschny, and C. M. Soukoulis, *Phys. Rev. Lett.* **95**, 203901 (2005).
- ³³ C. Rockstuhl, T. Zentgraf, H. Guo, N. Liu, C. Etrich, I. Loa, K. Syassen, J. Kuhl, F. Lederer, and H. Giessen, *Appl. Phys. B* **84**, 219 (2006).
- ³⁴ C. R. Simovski and S. A. Tretyakov, *Phys. Rev. B* **75**, 195111 (2007).
- ³⁵ C. Menzel, C. Rockstuhl, T. Paul, F. Lederer, and T. Pertsch, *Phys. Rev. B* **77**, 195328 (2008).
- ³⁶ A. Alù, *Phys. Rev. B* **83**, 081102 (2011).
- ³⁷ Y.-R. Zhen, K. H. Fung, and C. T. Chan, *Phys. Rev. B* **78**, 035419 (2008).
- ³⁸ I. Lindell, A. Sihvola, S. Tretyakov, and A. Viitanen, *Electromagnetic Waves in Chiral and Bi-Isotropic Media* (Artech Print on Demand, Norwood, USA, 1994) p. 352.
- ³⁹ I. Sersic, C. Tuambalangana, T. Kampfrath, and A. F. Koenderink, *Phys. Rev. B* **83**, 245102 (2011).
- ⁴⁰ I. Sersic, M. A. van de Haar, F. B. Arango, and A. F. Koenderink, *Phys. Rev. Lett.* **108**, 223903 (2012).
- ⁴¹ P. Belov, S. Maslovski, K. Simovski, and S. Tretyakov, *Tech. Phys. Lett.* **29**, 718 (2003).
- ⁴² F. J. García de Abajo, *Rev. Mod. Phys.* **79**, 1267 (2007).
- ⁴³ A. Kwadrin and A. F. Koenderink, *Phys. Rev. B* **89**, 045120 (2014).
- ⁴⁴ M. Husnik, M. W. Klein, N. Feth, J. Niegemann, K. Busch, S. Linden, and M. Wegener, *Nat. Photonics* **2**, 614 (2008).
- ⁴⁵ E. Prodan, C. Radloff, N. J. Halas, and P. Nordlander, *Science* **302**, 419 (2003).
- ⁴⁶ N. Liu, H. Liu, S. Zhu, and H. Giessen, *Nat. Photonics* **3**, 157 (2009).
- ⁴⁷ C. F. Bohren and D. R. Huffman, *Absorption and Scattering of Light by Small Particles* (John Wiley & Sons, New York, USA, 1983).
- ⁴⁸ F. Bernal Arango and A. F. Koenderink, *New J. Phys.* **15**, 1073023 (2013).
- ⁴⁹ A. F. Koenderink, *Nano Lett.* **9**, 4228 (2009).
- ⁵⁰ F. Bernal Arango, A. Kwadrin, and A. F. Koenderink, *ACS Nano* **6**, 10156 (2012).