

UNIVERSITY OF SOUTHAMPTON

Faculty of Social and Human Sciences School of Mathematics

Cohomological Finiteness Properties of Groups

Simon St. John-Green

Supervisor: Prof. Brita Nucinkis

Thesis for the degree of Doctor of Philosophy October 2014

UNIVERSITY OF SOUTHAMPTON <u>Abstract</u> Faculty of Social and Human Sciences School of Mathematics <u>Doctor of Philosophy</u> COHOMOLOGICAL FINITENESS PROPERTIES OF GROUPS by Simon St. John-Green

The main objects of study in this thesis are cohomological finiteness conditions of discrete groups. While most of the conditions we investigate are algebraic, they are inspired by topological invariants, particularly those concerning proper actions on CW-complexes.

The first two chapters contain preliminary material necessary for the remainder of the thesis. Chapter 2 concerns modules over a category with an emphasis on finiteness conditions. This material is well-known for (EI) categories, but we use a more general setup applicable to Mackey and cohomological Mackey functors, needed in Chapter 4. Chapter 3 specialises to Bredon cohomology, giving an overview of some results and detailing a few interesting examples.

In Chapter 4 we study finiteness conditions associated to Bredon cohomology with coefficients restricted to Mackey functors and cohomological Mackey functors, building again on the material in Chapter 2. In particular we characterise the corresponding FP_n conditions and prove that the Bredon cohomological dimension with coefficients restricted to cohomological Mackey functors is equal to the \mathfrak{F} -cohomological dimension for all groups.

We prove in Chapter 5 that for groups of finite \mathfrak{F} -cohomological dimension, the \mathfrak{F} -cohomological dimension equals the Gorenstein cohomological dimension, and give an application to the behaviour of the \mathfrak{F} -cohomological dimension under group extensions.

If a group G admits a closed manifold model for BG then G is a Poincaré duality group, in Chapter 6 we study Bredon–Poincaré duality groups, a generalisation of these. In particular if G admits a cocompact manifold model X for $E_{\mathcal{F}in}G$ (the classifying space for proper actions) with X^H a submanifold for any finite subgroup H of G, then G is a Bredon–Poincaré duality group. We give several sources of examples, including using the reflection group trick of Davis to produce examples where the dimensions of the submanifolds X^H are specified. We classify Bredon–Poincaré duality groups in low dimensions and examine their behaviour under group extensions.

In Chapter 7 we study Houghton's group H_n , calculating the centralisers of virtually cyclic subgroups and the Bredon cohomological dimension with respect to both the family of finite subgroups and the family of virtually cyclic subgroups.

Contents

List of Figures	ix
Declaration of Authorship	xi
Acknowledgements	xiii
Chapter 1. Introduction	1
1.1. Free actions and group cohomology	1
1.2. Proper actions and Bredon cohomology	2
1.3. Modules over a category	4
1.4. n_G and \mathfrak{F} -cohomological dimension	5
1.5. Mackey and cohomological Mackey functors	7
1.6. Gorenstein cohomological dimension	9
1.7. Bredon duality groups	11
1.8. Houghton's groups	14
Chapter 2. Modules over a category	17
2.1. Tensor products	21
2.1.1. Tensor product over \mathfrak{C}	21
2.1.2. Tensor product over R	22
2.2. Frees, projectives, injectives and flats	22
2.3. Restriction, induction and coinduction	24
2.4. Tor and Ext	26
2.5. Finiteness conditions	28
Chapter 3. Bredon modules	31
3.1. Free modules	31
3.2. Restriction, induction and coinduction	34
3.3. Bredon homology and cohomology of spaces	37
3.4. Homology and cohomology of groups	38
3.5. Cohomological dimension	38
3.5.1. Low dimensions	39
3.6. FP_n conditions	40
3.6.1. Quasi- $\mathcal{O}_{\mathcal{F}} FP_n$ conditions	41
3.7. Change of rings	42

3.8. Some interesting examples	44
3.9. Finitely generated projectives and duality	47
Chapter 4. Mackey and cohomological Mackey functors	55
4.1. Introduction	56
4.1.1. Mackey functors	56
4.1.1.1. Free modules	60
4.1.1.2. Induction	61
4.1.1.3. Homology and cohomology	62
4.1.2. Cohomological Mackey functors	63
4.1.2.1. Explicit description of π	66
4.1.2.2. Homology and cohomology	67
4.2. FP_n conditions for Mackey functors	68
4.3. Homology and cohomology of cohomological Mackey functors	72
4.4. FP_n conditions for cohomological Mackey functors	77
4.4.1. $\mathcal{H}_{\mathcal{F}} \mathrm{FP}_n$ implies $\mathcal{F} \mathrm{FP}_n$	79
4.4.2. $\mathcal{F}FP_n$ implies $\mathcal{H}_{\mathcal{F}}FP_n$	81
4.5. Cohomological dimension for cohomological Mackey functors	83
4.5.1. Closure properties	85
4.6. The family of <i>p</i> -subgroups	86
4.6.1. FP_n conditions over \mathbb{F}_p	90
Chapter 5. Gorenstein cohomology and \mathfrak{F} -cohomology	93
5.1. Preliminaries	93
5.1.1. Complete resolutions and complete cohomology	93
5.1.2. \mathfrak{F} -cohomology	94
5.1.3. Complete \mathfrak{F} -cohomology	95
5.1.4. Gorenstein cohomology	96
5.2. \mathfrak{F}_G -cohomology	97
5.2.1. Construction	97
5.2.2. Technical results	99
5.2.3. An Avramov–Martsinkovsky long exact sequence in \mathfrak{F} -cohomol	ogy 101
5.3. Group extensions	105
5.4. Rational cohomological dimension	105
Chapter 6. Bredon duality groups	107
6.1. Preliminary observations	108
6.2. Examples	109
6.2.1. Smooth actions on manifolds	110
6.2.2. A counterexample to the generalised PD^n conjecture	113
6.2.3. Actions on <i>R</i> -homology manifolds	114

×		7	1
1	٧		

CONTENTS

6.2.4. One relator groups	116
6.2.5. Discrete subgroups of Lie groups	118
6.2.6. Virtually soluble groups	118
6.2.7. Elementary amenable groups	119
6.3. Finite extensions of right-angled Coxeter groups	120
6.4. Low dimensions	124
6.5. Extensions	127
6.5.1. Direct products	127
6.5.2. Finite-by-duality groups	129
6.6. Graphs of groups	131
6.7. The wrong notion of Bredon duality	134
Chapter 7. Houghton's groups	137
7.1. Centralisers of finite subgroups in H_n	138
7.2. Centralisers of elements in H_n	142
7.3. Brown's model for $E_{\mathcal{F}in}H_n$	147
7.4. Finiteness conditions satisfied by H_n	149
7.4.1. The pushout of Lück and Weiermann	150
7.4.2. Calculation of $\mathcal{O}_{\mathcal{VCyc}} \operatorname{cd} H_n$	152
Bibliography	155
Index	163

vii

List of Figures

1 A graphical representation of the set S_a described in Section 7.1. 142

Declaration of Authorship

I, Simon St. John-Green, declare that this thesis and the work presented in it are my own and has been generated by me as the result of my own original research. Title: Cohomological Finiteness Properties of Groups

I confirm that:

- (1) This work was done wholly or mainly while in candidature for a research degree at this University;
- (2) Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated;
- (3) Where I have consulted the published work of others, this is always clearly attributed;
- (4) Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work;
- (5) I have acknowledged all main sources of help;
- (6) Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself;
- (7) Either none of this work has been published before submission, or parts of this work have been published as:
 - (a) Centralisers in Houghton's groups (2012, to appear Proc. Edinburgh Math. Soc.) [SJG12].
 - (b) Bredon–Poincaré duality groups (2013, to appear J. Group Theory) [SJG13a].
 - (c) Finiteness conditions for Mackey and cohomological Mackey functors (J. Algebra 411 (2014), no. 0, 225–258) [SJG14]
 - (d) On the Gorenstein and \$\vec{v}\$-cohomological dimensions (2013, to appear Bull. Lond. Math. Soc.) [SJG13b].

Signed:

Jina SI. Tche-Green

Date: October 2014.

Acknowledgements

First and foremost I'd like to thank my supervisor Brita Nucinkis, for introducing me to this beautiful area of mathematics and for being an endless source of motivation, support, and guidance throughout my studies.

I'd like to thank my advisor Ian Leary for his support and for many useful mathematical insights, including the examples in Section 6.3 and Example 6.2.11. I thank him as well for reading a first draft of this thesis.

For making the last three years such a wonderful experience, I thank all the PhD students and post-docs, past and present, from the Southampton mathematics department, including but not limited to: Alex Bailey, Matt Burfitt, Valerio Capraro, Chris Cave, Yuyen Chien, Charles Cox, Dennis Dreesen, Michal Ferov, Robin Frankhuizen, Tom Harris, Pippa Hiscock, Ana Khukhro, Dave Matthews, Ingrid Membrillo, Raffaele Rainone, Amin Saied, Conor Smyth, Rob Snocken, Joe Tait, Edwin Tye, and Mike West.

The material in Sections 7.4.1 and 7.4.2 is joint work with Nansen Petrosyan and I thank him for suggesting the topic and for sharing with me some fascinating mathematics.

Thanks to Dieter Degrijse, not only for pointing out to me an error in Lemma 3.9.5 while reading a preprint of [**SJG14**], but also for many interesting conversations concerning Mackey and cohomological Mackey functors.

I'd like to thank Conchita Martínez-Pérez for her very helpful comments concerning a preprint of [**SJG12**], material now contained in Chapter 7. Thanks to the referee of the Proceedings of the Edinburgh Mathematical Society who reviewed the same preprint for making useful comments, including suggesting the graph Γ used in Section 7.2. Thanks to Jim Davis for showing me that some groups constructed by Block and Weinberger, appearing now in Section 6.2.2, answered a question in an early preprint of [**SJG13a**]. Simplifications to the proofs of Lemmas 4.2.1 and 4.3.5 are due to the referee from the Journal of Algebra.

Finally but most importantly, I thank my family, especially my parents and my sister. I would dedicate this thesis to them, only then they might feel compelled to read it, and I would not want to force that upon anyone.

CHAPTER 1

Introduction

This introduction contains an overview of relevant background material and details the contributions of this thesis as they arise.

1.1. Free actions and group cohomology

For any group G there exists an aspherical CW-complex X with fundamental group G, this is called a *model for* BG or *Eilenberg-Mac Lane* space. Such a space is unique up to homotopy equivalence, a fact observed essentially by Hurewicz [**Hur36**], arguably kick-starting the field of group cohomology. The universal cover \tilde{X} , a contractible CW-complex with a free G-action, is called a model for EG or a classifying space for free actions. Equivalently one could define a model for EG as the terminal object in the G-homotopy category of free G-CW-complexes.

One can use invariants of these spaces to study the groups themselves, for example defining the group cohomology $H^*(G)$ to be $H^*(BG)$. Alternatively there is an algebraic definition of group cohomology, replacing the space BG with a resolution of \mathbb{Z} by projective $\mathbb{Z}G$ -modules.

An important invariant is the geometric dimension gd G, the minimal dimension of a model for EG. Its algebraic counterpart is the cohomological dimension cd G, the minimal length of a resolution of \mathbb{Z} by projective $\mathbb{Z}G$ -modules. It's easy to see that gd G = 0 if and only if cd G = 0 if and only if G is the trivial group. Also, by a theorem of Stallings and Swan, cd G = 1 if and only if gd G = 1 if and only if G is a free group [**Sta68, Swa69**]. Eilenberg and Ganea conjectured that cd G = gd G for all groups and, along with Stallings and Swan's result for the dimension one case, proved this conjecture for all cases, except for the possibility that cd G = 2 and gd G = 3 [**EG57**]. That this is impossible is still an open problem, known as the Eilenberg–Ganea conjecture.

A group G has type F_n if it admits a model for BG with finite n-skeleton, and on the algebraic side G has type FP_n if Z admits a resolution by projective ZGmodules, finitely generated up to dimension n. A group of type F_n is necessarily of type FP_n . All groups are of type F_0 , since there always exists a model for BG with a single 0-cell [**Geo08**, 7.1.5]. The conditions F_1 , FP_1 and finitely generated are all equivalent, but the situation is more complex for larger n. A group is F_2 if

1. INTRODUCTION

and only if it is finitely presented, and FP_n together with F_2 implies F_n [Geo08, 7.2.1][Bro94, VIII §7]. Bestvina and Brady use discrete Morse theory techniques to construct subgroups of right-angled Artin groups that are FP_n but not FP_{n+1} for all n, and groups of type FP_n which are not finitely presented for all n [BB97].

We say a group is type F if it is F_{∞} and $\operatorname{gd} G < \infty$, and type FP of it is $\operatorname{FP}_{\infty}$ and $\operatorname{cd} G < \infty$.

For a more detailed overview of these finiteness conditions see [**Bro94**, Chapter VIII], [**Bie81**] and [**Geo08**, Chapter II].

1.2. Proper actions and Bredon cohomology

Let \mathcal{F} be a family of subgroups of a group G, closed under conjugation and taking subgroups. A model for $E_{\mathcal{F}}G$, or classifying space for actions with isotropy in \mathcal{F} , is the terminal object in the G-homotopy category of G-CW complexes with isotropy in \mathcal{F} . A model for EG is thus the same as a model for $E_{Triv}G$, where Trivdenotes the family consisting of only the trivial subgroup.

Using the equivariant Whitehead theorem one can show that a G-CW-complex X is a model for $E_{\mathcal{F}}G$ if and only if X has isotropy in \mathcal{F} and X^H is contractible for all $H \in \mathcal{F}$ [Lüc05, Theorem 1.9]. Models for $E_{\mathcal{F}}G$ always exist—there are various standard constructions including the infinite join construction of Milnor [Mil56], Segals construction [Seg68], and a construction where one iteratively attaches equivariant cells to build a G-CW-complex with contractible fixed point sets [Lüc89, Proposition 2.3, p.35].

Let $\mathcal{F}in$ denote the family of all finite subgroups of a group G. There are many groups which admit natural models for $\mathbb{E}_{\mathcal{F}in}G$, for example mapping class groups, word-hyperbolic groups, and one-relator groups. A good survey is [Lüc05].

There has been recent interest in models for $E_{\mathcal{F}in}G$ and models for $E_{\mathcal{VCyc}}G$, where \mathcal{VCyc} denotes the family of virtually cyclic subgroups, because they appear on one side of the Baum–Connes and Farrell–Jones conjectures respectively [**LR05**]. These are deep conjectures with far reaching consequences in mathematics [**MV03**, **BLR08**].

We denote by $\operatorname{gd}_{\mathcal{F}} G$ the minimal dimension of a model for $\operatorname{E}_{\mathcal{F}} G$, if $\mathcal{F} = \mathcal{F}$ in then this is known as the proper geometric dimension of G. The cohomology theory most suited to the study of this geometric invariant is Bredon cohomology, introduced for finite groups by Bredon in [**Bre67**] to study equivariant obstructions and extended to the study of infinite groups by Lück [**Lüc89**].

Fixing G we consider the orbit category $\mathcal{O}_{\mathcal{F}}$. This is the small category whose objects are the transitive G-sets with stabilisers in \mathcal{F} and the morphisms between two such G-sets is the free abelian group on the G-maps between them. Bredon modules, or $\mathcal{O}_{\mathcal{F}}$ -modules, are contravariant additive functors from $\mathcal{O}_{\mathcal{F}}$ to the category of left *R*-modules, where *R* is some commutative ring. Our definition of $\mathcal{O}_{\mathcal{F}}$ is different from the usual definition given in, for example, [Lüc89], but the definitions lead to isomorphic $\mathcal{O}_{\mathcal{F}}$ -module categories (Remark 2.0.7).

The category of all Bredon modules is an abelian category with frees and projectives, so one can use techniques from homological algebra to study them. Let \underline{R} be the constant Bredon module, defined as $\underline{R}(G/H) = R$ for all $H \in \mathcal{F}$ and $\underline{R}(\alpha) = \operatorname{id}_R$ for any G-map $\alpha : G/H \to G/K$. As in ordinary group cohomology, using projective resolutions of \underline{R} one builds the Bredon cohomology of G. Analagously to $\operatorname{cd} G$, we denote by $\mathcal{O}_{\mathcal{F}}\operatorname{cd} G$ the Bredon cohomological dimension of G—the minimal length of a projective resolution of \underline{R} . We denote by $\mathcal{O}_{\mathcal{F}}\operatorname{FP}_n$ the Bredon cohomological analogue of the FP_n conditions of ordinary cohomology, so G is $\mathcal{O}_{\mathcal{F}}\operatorname{FP}_n$ if there exists a resolution of \underline{R} by projective Bredon modules which is finitely generated in all degrees $\leq n$.

That the Bredon cohomological dimension $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G$ is the correct algebraic invariant to mirror $\operatorname{gd}_{\mathcal{F}in} G$ is exemplified by the following theorem, an analogue of the classical results of Eilenberg–Ganea and Stallings–Swan.

THEOREM. [LM00, Theorem 0.1][Dun79] If $R = \mathbb{Z}$ then $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G = \operatorname{gd}_{\mathcal{F}in} G$, except for the possibility that $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G = 2$ and $\operatorname{gd}_{\mathcal{F}in} G = 3$.

Brady, Leary and Nucinkis construct groups with $\mathcal{O}_{\text{fin}} \operatorname{cd} G = 2$ and $\operatorname{gd}_{\text{fin}} G = 3$ [**BLN01**].

If G admits a model for $\mathbb{E}_{\mathcal{F}in}G$ with cocompact *n*-skeleton then G is $\mathcal{O}_{\mathcal{F}in}\mathrm{FP}_n$ over Z. In the other direction, if G is $\mathcal{O}_{\mathcal{F}in}\mathrm{FP}_n$ and the Weyl groups $WH = N_G H/H$ are finitely presented for all finite subgroups H of G, then G admits a model for $\mathbb{E}_{\mathcal{F}in}G$ with cocompact *n*-skeleton [**LM00**, Theorem 0.1].

PROPOSITION. [KMPN11b, Lemmas 3.1,3.2] A group G is $\mathcal{O}_{\text{fin}} \text{FP}_n$ if and only if G has finitely many conjugacy classes of finite subgroups and the Weyl groups $WH = N_G H/H$ are FP_n for all finite subgroups H.

We will discuss these conditions in more depth in Section 3.6.

Much of this thesis is concerned with the Bredon cohomological dimension and $\mathcal{O}_{\text{fin}} \text{FP}_n$ conditions, and how they interact with other cohomological finiteness conditions. This includes those obtained by restricting the coefficients of Bredon cohomology to Mackey functors or cohomological Mackey functors (Section 1.5 and Chapter 4) and those obtained via relative homological algebra, namely the Gorenstein cohomological dimension and \mathfrak{F} -cohomological dimension (Sections 1.4 and 1.6, and Chapter 5).

There are already many results giving bounds for the Bredon cohomological dimension in terms of other algebraic invariants. In [MP13a, MP13b], Martínez-Pérez uses the poset of finite subgroups of a group to provide bounds

1. INTRODUCTION

for $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G$ and in [**KMPN09**] Kropholler, Martínez-Pérez, and Nucinkis show that any elementary amenable group of type $\operatorname{FP}_{\infty}$ over \mathbb{Z} satisfies

$$\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{Z}} G = hG = \mathrm{cd}_{\mathbb{Q}} G,$$

where hG denotes the *Hirsch length* of *G*. See [Hil91] for a definition of Hirsch length for elementary amenable groups.

Finiteness conditions in Bredon cohomology are not well-behaved with respect to group extensions. This is exemplified by the constructions of Leary and Nucinkis [LN03] of

- (1) groups which are virtually-F (there exists a finite index subgroup of type F) and satisify $\operatorname{vcd} G < \mathcal{O}_{\operatorname{fin}} \operatorname{cd} G$, and
- (2) groups which are virtually-F with infinitely many conjugacy classes of finite subgroups (and hence not of type $\mathcal{O}_{\text{fin}} \text{FP}_0$ by Proposition 3.6.1).

Interestingly, a virtually-F group cannot contain infinitely many conjugacy classes of subgroups of prime power order [**Bro94**, IX.13.2], but may contain infinitely many conjugacy classes of subgroups isomorphic to some finite group H as long as H does not have prime power order [**Lea05**].

In Chapter 3 we look in detail at $\mathcal{O}_{\mathcal{F}}$ -modules and at some results concerning finiteness conditions in Bredon cohomology which will be needed later on in the thesis. We also give some interesting examples of groups whose Bredon cohomological dimension is not preserved under change of rings. Apart from these examples, this chapter contains mainly background material and straightforward extensions of known results.

1.3. Modules over a category

An **Ab** category (also called a pre-additive category) is a category \mathfrak{C} enriched over the category of abelian groups—for any two objects x and y in \mathfrak{C} the morphisms from x to y form an abelian group and morphism composition distributes over addition [**ML98**, p.28]. So for any $w, x, y, z \in \mathfrak{C}$ and morphisms

$$w \xrightarrow{f} x \xrightarrow{g} y \xrightarrow{k} z,$$

we have

$$k \circ (q+h) \circ f = k \circ q \circ f + k \circ h \circ f$$

If \mathfrak{C} is a small \mathbf{Ab} category then a \mathfrak{C} -module is a contravariant additive functor from \mathfrak{C} to the category of left R-modules. The theory of modules over a category specialises to Bredon cohomology by setting $\mathfrak{C} = \mathcal{O}_{\mathcal{F}}$. In Chapter 2 we study modules over an \mathbf{Ab} category \mathfrak{C} with the property that for all objects x and yin \mathfrak{C} , the set of morphisms from x to y forms a free abelian group. We describe standard constructions including tensor products; projective, injective, and flat modules; restriction, induction, and coinduction; the $\operatorname{Tor}^{\mathfrak{C}}_*$ and $\operatorname{Ext}^*_{\mathfrak{C}}$ functors; projective dimension and $\mathfrak{C}\operatorname{FP}_n$ conditions; and the Bieri–Eckmann criterion.

Let $[x, y]_{\mathfrak{C}}$ denote the morphisms from x to y in \mathfrak{C} . The category \mathfrak{C} is said to be an (EI) category if (see Remark 2.0.1):

(EI) For every $x \in \mathfrak{C}$ there is a distinguished basis of $[x, x]_{\mathfrak{C}}$, the elements of which are isomorphisms.

The material in this chapter is well-known in the case that \mathfrak{C} is an (EI) category. We study a more general case which can be specialised, not just to Bredon modules in Chapter 3, but also to Mackey and cohomological Mackey functors in Chapter 4—Mackey and cohomological Mackey functors may be described as modules over categories $\mathcal{M}_{\mathcal{F}}$ and $\mathcal{H}_{\mathcal{F}}$ respectively, categories which do not have (EI).

1.4. n_G and \mathfrak{F} -cohomological dimension

Let n_G denote the minimal dimension of a contractible proper *G*-CW complex. Nucinkis suggested \mathfrak{F} -cohomology in [**Nuc99**] as an algebraic analogue of n_G , it is a special case of the relative homology of Mac Lane [**ML95**] and Eilenberg– Moore [**EM65**]. Fix a subfamily \mathcal{F} of the family of finite subgroups, closed under conjugation and taking subgroups. Let Δ be the *G*-set $\coprod_{H \in \mathcal{F}} G/H$ and say that a module is \mathcal{F} -projective if it is a direct summand of a module of the form $N \otimes R\Delta$ where N is any RG-module. Short exact sequences are replaced with \mathcal{F} -split short exact sequences—short exact sequences which split when restricted to any subgroup in \mathcal{F} , or equivalently which split when tensored with $R\Delta$. The class of \mathcal{F} -split short exact sequences is allowable in the sense of Mac Lane, and the projective modules with respect to these sequences are exactly the \mathcal{F} -projectives. This means an RG-module P is \mathcal{F} -projective if and only if given any \mathcal{F} -split short exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

of RG-modules, applying $\operatorname{Hom}_{RG}(P, -)$ gives a short exact sequence

 $0 \longrightarrow \operatorname{Hom}_{RG}(P, A) \longrightarrow \operatorname{Hom}_{RG}(P, B) \longrightarrow \operatorname{Hom}_{RG}(P, C) \longrightarrow 0.$

There are enough \mathcal{F} -projectives and one can define Ext and Tor functors, denoted $\mathcal{F}\text{Ext}_{RG}^*$ and $\mathcal{F}\text{Tor}_*^{RG}$, for any RG-modules M and N,

$$\mathcal{F}\text{Ext}_{RG}^*(M,N) = H^* \operatorname{Hom}_{RG}(P_*,N)$$
$$\mathcal{F}\text{Tor}_*^{RG}(M,N) = H_*(P_* \otimes_{RG} N)$$

where P_* is a \mathcal{F} -split resolution of M by \mathcal{F} -projective modules. We define the \mathcal{F} -cohomology and \mathcal{F} -homology

$$\mathcal{F}H^*(G, M) = \mathcal{F}\text{Ext}^*_{RG}(R, M),$$

1. INTRODUCTION

$$\mathcal{F}H_*(G,M) = \mathcal{F}\mathrm{Tor}^{RG}_*(R,M).$$

The \mathcal{F} -cohomological dimension, denoted \mathcal{F} cd G, is the shortest length of an \mathcal{F} -split \mathcal{F} -projective resolution of R. A group G is \mathcal{F} FP_n if there exists an \mathcal{F} -split \mathcal{F} -projective resolution of R, finitely generated in all degrees $\leq n$.

NOTATION. When $\mathcal{F} = \mathcal{F}in$, we use the standard notation in the literature, writing $\mathfrak{F}cd$ instead of $\mathcal{F}incd$ and referring to the \mathfrak{F} -cohomological dimension.

A result of Bouc and Kropholler–Wall implies $\mathfrak{Fcd} G \leq n_G$ [Bou99, KW11], but it is unknown if $\mathfrak{Fcd} G < \infty$ implies $n_G < \infty$, or if there are any groups for which the two invariants differ. Unfortunately \mathcal{F} -cohomology can be very difficult to deal with, in particular it lacks some useful features such as free modules.

Since every model for $E_{\mathcal{F}in}G$ is a proper contractible *G*-CW complex, it is clear that $n_G \leq \operatorname{gd}_{\mathcal{F}in} G$.

CONJECTURE (Kropholler–Mislin Conjecture [**Gui08**, Conjecture 43.1]). If $n_G < \infty$ then $\operatorname{gd}_{\operatorname{fin}} G < \infty$.

Kropholler and Mislin verified their conjecture for groups of type FP_{∞} [KM98] and later Lück verified the conjecture for groups with $l(G) < \infty$ [Lüc00]. Here l(G) is the length of the longest chain

$$1 = H_0 \lneq H_1 \lneq H_2 \lneq \dots \lneq H_n \lneq G$$

of finite subgroups in G. Nucinkis posed an algebraic version of the conjecture, asking whether the finiteness of $\operatorname{\mathfrak{F}cd} G$ and $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G$ are equivalent and verifying this for groups with $l(G) < \infty$ [Nuc00, Conjecture on p.337, Corollary 4.5].

The class $\mathbf{H}\mathfrak{F}$ of hierarchically decomposable groups was introduced by Kropholler as the smallest class of groups such that if there exists a finite-dimensional contractible *G*-CW complex with stabilisers in $\mathbf{H}\mathfrak{F}$ then $G \in \mathbf{H}\mathfrak{F}$ [**Kro93**], he proves that every torsion-free group of type FP_{∞} in $\mathbf{H}\mathfrak{F}$ has finite cohomological dimension. The class $\mathbf{H}\mathfrak{F}$ is extremely large, containing for example all countable elementary amenable groups and all countable linear groups. The first known example of a group not in Kropholler's class $\mathbf{H}\mathfrak{F}$ was Thompson's group *F*, since *F* is torsion-free of type FP_{∞} but with infinite cohomological dimension [**BG84**]. Other examples have since been found [**Gan12b**, **ABJ+09**]. Gandini and Nucinkis have verified the Kropholler–Mislin conjecture for a class of groups containing many groups of unbounded torsion [**GN12**].

In [**MP13a**, Example 3.6] Martínez-Peréz modifies the Leary–Nucinkis construction [**LN03**] to produce an extension G of a torsion-free group by a cyclic group of order p, with $\operatorname{\mathfrak{F}cd} G = 3$ but $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G = 4$. Taking direct products of these groups and using [**DP12**, Theorem C] gives a family of virtually torsionfree groups G_n with $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G_n = \operatorname{\mathfrak{F}cd} G_n + n$ for all natural numbers n [**Deg13a**, Remark 3.6]. However one should note that in these examples $\operatorname{\mathfrak{F}cd} G_n$ is growing linearly with n.

Interestingly, it is still unknown if $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G = \operatorname{\mathfrak{F}cd} G$ when G is of type $\mathcal{O}_{\mathcal{F}in}\operatorname{FP}_{\infty}$, although Degrijse and Martínez-Pérez have obtained some results pertaining to this question in [**DMP13**]. They investigate groups admitting a cocompact model X for $\operatorname{E}_{\mathcal{F}in}G$, and find a description of $\mathcal{O}_{\mathcal{F}in}\operatorname{cd} G$ as largest nfor which $H_c^n(X^K, X_{\operatorname{sing}}^K)$ is non-trivial, where K runs over the finite subgroups of G, $X_{\operatorname{sing}}^K$ denotes the subspace of cells in X^K with isotropy containing Kbut not equal to K, and H_c^n denotes the cohomology with compact supports [**DMP13**, Corollary 2.5]. Using this they prove that if G acts properly and chamber-transitively on a building of type (W, S), where (W, S) is a finite Coxeter group, then $\mathcal{O}_{\mathcal{F}in}\operatorname{cd} G = \operatorname{\mathfrak{F}cd} G$ [**DMP13**, Theorem 5.4].

1.5. Mackey and cohomological Mackey functors

In [MPN06], Martínez-Peréz and Nucinkis studied cohomological finiteness conditions arising from taking the Bredon cohomology of a group G but restricting to Mackey functor coefficients. They showed that the associated Mackey cohomological dimension $\mathcal{M}_{\mathcal{F}in}$ cd G is always equal to both the virtual cohomological dimension and the \mathfrak{F} -cohomological dimension when G is virtually torsion-free. One can view Mackey functors as contravariant functors from a small category $\mathcal{M}_{\mathcal{F}in}$ into the category of left R-modules, and a crucial result in the paper of Martínez-Peréz and Nucinkis is that the Bredon cohomology with coefficients in a Mackey functor may be calculated using a projective resolution of Mackey functors. Specifically they prove that you can induce a resolution of \underline{R} by projective Bredon modules to a resolution of the Burnside functor B^G by projective Mackey functors. This is explained in more detail in Section 4.1.1.

Degripse showed that for groups with $l(G) < \infty$ the Mackey cohomological dimension is equal to the \mathfrak{F} -cohomological dimension [**Deg13a**, Theorem A]. He proves this via the study of Bredon cohomology with cohomological Mackey functor coefficients and the associated notion of cohomological dimension $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G$.

The main ingredient of Chapter 4 is a similar result to that of Martínez-Peréz and Nucinkis for Bredon cohomology with cohomological Mackey functor coefficients. Yoshida observed that a cohomological Mackey functor may be described as a contravariant functor from a small category $\mathcal{H}_{\mathcal{F}in}$ to the category of left *R*-modules [**Yos83**]. We use Yoshida's result to prove in Section 4.3 that the Bredon cohomology with coefficients in a cohomological Mackey functor may be calculated with a projective resolution of cohomological Mackey functors, by showing that a resolution of <u>R</u> by projective Bredon modules can be induced to a resolution of the fixed point functor R^- by projective cohomological Mackey functors.

Degripse also proves in $[\mathbf{Deg13a}]$ that $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G = \mathfrak{F}cd G$ when $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G < \infty$, and asks if they are always equal, we can verify this:

THEOREM 4.5.1. $\mathcal{H}_{\text{fin}} \operatorname{cd} G = \operatorname{\mathfrak{F}cd} G$ for all groups G.

Thus for an arbitrary group G we have a chain of inequalities

 $\mathfrak{F}\mathrm{cd}\,G = \mathcal{H}_{\operatorname{Fin}}\mathrm{cd}\,G \leq \mathcal{M}_{\operatorname{Fin}}\mathrm{cd}\,G \leq \mathcal{O}_{\operatorname{Fin}}\mathrm{cd}\,G.$

Since the new invariants $\mathcal{M}_{\mathcal{F}in}$ cd and $\mathcal{H}_{\mathcal{F}in}$ cd interpolate between $\mathcal{O}_{\mathcal{F}in}$ cd Gand $\mathfrak{F}cd G$, one might hope to use them to gain information about how the Kropholler–Mislin conjecture might fail. However, few of the inequalities above are well understood. The inequality $\mathfrak{F}cd G \leq \mathcal{O}_{\mathcal{F}in}$ cd G has already been discussed in Section 1.4. For groups with l(G) finite, $\mathcal{H}_{\mathcal{F}in}$ cd $G = \mathcal{M}_{\mathcal{F}in}$ cd G [**Deg13a**, Theorem 4.10], we don't know of any examples where they differ.

- QUESTION 1.5.1. (1) For an arbitrary group G, does the finiteness of $\mathcal{M}_{\mathcal{F}in} \operatorname{cd} G$ imply the finiteness of $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G$?
- (2) Is there any relation between $\mathcal{M}_{\text{fin}} \operatorname{cd} G$ and n_G ?

The $\mathcal{O}_{\mathcal{F}in} \mathrm{FP}_n$ conditions are well understood—see Section 3.6. We study the $\mathcal{M}_{\mathcal{F}in} \mathrm{FP}_n$ conditions corresponding to Mackey functors, the $\mathcal{H}_{\mathcal{F}in} \mathrm{FP}_n$ conditions corresponding to cohomological Mackey functors, and the \mathfrak{FP}_n conditions defined in the previous section.

COROLLARY 4.2.6. Over any ring R, a group is $\mathcal{M}_{\text{fin}} FP_n$ if and only if it is $\mathcal{O}_{\text{fin}} FP_n$.

THEOREM 4.4.1. If R is a commutative Noetherian ring, a group is $\mathcal{H}_{\text{fin}} FP_n$ if and only if it is $\mathfrak{F}P_n$.

In Section 4.6 we prove a result similar to that shown for \mathfrak{F} -cohomology in [**LN10**, §4], that depending on the coefficient ring, $\mathcal{H}_{\mathcal{F}in}$ cd may be calculated using a subfamily of the family of finite subgroups. For example when working over \mathbb{Z} we need consider only the family of finite subgroups of prime power order, and over either the finite field \mathbb{F}_p or over $\mathbb{Z}_{(p)}$ (the integers localised at p), we need consider only the family of subgroups of order a power of p.

THEOREM 4.6.1. Let R be either \mathbb{Z} , \mathbb{F}_p , or $\mathbb{Z}_{(p)}$. If $R = \mathbb{Z}$ then denote by \mathcal{P} the family of subgroups of prime-power order. If $R = \mathbb{F}_p$ or $\mathbb{Z}_{(p)}$ then let \mathcal{P} denote the family of subgroups of order a power of p.

For all $n \in \mathbb{N} \cup \{\infty\}$, the conditions $\mathcal{H}_{\text{fin}} \operatorname{cd} G = n$ and $\mathcal{H}_{\mathcal{P}} \operatorname{cd} G = n$ are equivalent, as are the conditions $\mathcal{H}_{\text{fin}} \operatorname{FP}_n$ and $\mathcal{H}_{\mathcal{P}} \operatorname{FP}_n$.

We also give a complete description of the condition $\mathcal{H}_{\text{fin}} FP_n$ over \mathbb{F}_p .

COROLLARY 4.6.11. G is $\mathcal{H}_{\text{fin}} \text{FP}_n$ over \mathbb{F}_p if and only if G has finitely many conjugacy classes of p-subgroups, and $WH = N_G H/H$ is FP_n over \mathbb{F}_p for all finite p-subgroups H.

1.6. Gorenstein cohomological dimension

An *RG*-module is *Gorenstein projective* if it is a cokernel in a strong complete resolution of *RG*-modules, these were first defined over an arbitrary ring by Enochs and Jenda [**EJ95**]. We give a full explanation of complete resolutions in Section 5.1.1. The *Gorenstein projective dimension* Gpd M of an *RG*-module M is the minimal length of a resolution of M by Gorenstein projective modules. Equivalently, Gpd $M \leq n$ if and only if M admits a complete resolution of coincidence index n [**BDT09**, p.864].

The Gorenstein cohomological dimension of a group G, denoted $\operatorname{Gcd} G$, is the Gorenstein projective dimension of R. If G is virtually torsion-free then $\operatorname{Gcd} G = \operatorname{vcd} G$ [**BDT09**, Remark 2.9(1)], indeed the Gorenstein cohomology can be seen as a generalisation of the virtual cohomological dimension.

Gcd G is closely related to the silp RG and spli RG invariants studied by Gedrich and Gruenberg [**GG87**] and recently shown to be equal by Emmanouil when $R = \mathbb{Z}$ [**Emm10**]. The invariants silp RG and spli RG are defined as the supremum of the injective lengths (injective dimensions) of the projective RGmodules and the supremum of the projective lengths (projective dimensions) of the injective RG-modules respectively. It is known that

$$\operatorname{Gcd} G \leq \operatorname{spli} RG \leq \operatorname{Gcd} G + 1,$$

and conjectured that $\operatorname{Gcd} G = \operatorname{spli} RG$ [**DT08**, Conjecture A]. In fact, Dembegioti and Talelli phrase this conjecture with the generalised cohomological dimension of Ikenaga [**Ike84**], but this is always equal to the Gorenstein cohomological dimension [**BDT09**, Theorem 2.5].

By [ABS09, Lemma 2.21], every permutation RG-module with finite stabilisers is Gorenstein projective, so combining with [Gan12b, Lemma 3.4] gives that $\operatorname{Gcd} G \leq \operatorname{\mathfrak{Fcd}} G$.

THEOREM 5.2.11. If $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G = \operatorname{Gcd} G$.

We don't know if $\operatorname{Gcd} G < \infty$ implies $\operatorname{\mathfrak{Fcd}} G < \infty$, although if $\operatorname{Gcd} G = 0$ or 1 then $\operatorname{Gcd} G = \operatorname{\mathfrak{Fcd}} G = \mathcal{O}_{\operatorname{fin}} \operatorname{cd} G$ [ABS09, Proposition 2.19][BDT09, Theorem 3.6]. Additionally if G is in Kropholler's class $\operatorname{H}\mathfrak{F}$ and has a bound on the orders of its finite subgroups then $\operatorname{\mathfrak{Fcd}} G = \operatorname{Gcd} G$ (see Example 5.2.12).

1. INTRODUCTION

It is conjectured by Talelli that $\operatorname{Gcd} G < \infty$ if and only if $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G < \infty$ (see for example, [**Tal07**, Conjecture A]). This is a stronger version of the conjecture of Nucinkis mentioned in Section 1.4, that $\operatorname{Fcd} G < \infty$ if and only if $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G < \infty$. If one could strengthen Theorem 5.2.11 to show that $\operatorname{Gcd} G = \operatorname{Fcd} G$ for all groups G, then the two conjectures would be equivalent. Using [**MP07**], Bahlekeh, Dembegioti, and Talelli show that for groups with $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G < \infty$, there is a bound $\mathcal{O}_{\operatorname{fin}} \operatorname{cd} G < l(G) + \operatorname{Gcd} G$ [**BDT09**, Theorem C].

Generalising a construction of Avramov–Martsinkovsky, it was shown by Asadollahi, Bahlekeh, and Salarian that if Gcd $G < \infty$ then there is a long exact sequence of cohomological functors relating group cohomology, complete cohomology and Gorenstein cohomology [**AM02**, §7][**ABS09**, §3]. Theorem 5.2.11 follows from constructing a similar long exact sequence relating \mathfrak{F} -cohomology, complete \mathfrak{F} -cohomology (defined in Section 5.1.3), and a new cohomology theory we call \mathfrak{F}_G -cohomology (defined in Section 5.2).

When they both exist, these two long exact sequences fit into the commutative diagram below, see Proposition 5.2.9,

where for conciseness we have written H^n for $H^n(G, -)$ etc. In the commutative diagram above, $\widehat{H}^n(G, -)$ is the complete cohomology, $GH^n(G, -)$ is the Gorenstein cohomology, $\widehat{\mathfrak{F}H}^n(G, -)$ is complete \mathfrak{F} -cohomology, and $\mathfrak{F}_G H^n(G, -)$ is the \mathfrak{F}_G -cohomology.

Since Theorem 5.2.11 is proved via this commutative diagram, it appears that the requirement that $\operatorname{\mathfrak{F}cd} G < \infty$ will be difficult to circumvent—without it we do not know how to construct the long exact sequence appearing on the top row.

In Section 5.3 we use that the Gorenstein cohomological dimension is subadditive to improve upon a result of Degrijse on the behaviour of the \mathfrak{F} -cohomological dimension under group extensions [**Deg13a**, Theorem B]. Degrijse phrased his result in terms of Bredon cohomological dimension of G with coefficients restricted to cohomological Mackey functors, but this invariant is equal to $\mathfrak{F}cd G$ by Theorem 4.5.1 (see previous section).

COROLLARY 5.3.2. Given a short exact sequence of groups

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1,$$

if $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G \leq \operatorname{\mathfrak{F}cd} N + \operatorname{\mathfrak{F}cd} Q$.

QUESTION 1.6.1. Is the \mathfrak{F} -cohomological dimension subadditive under group extensions?

In Section 5.4 we use the Avramov–Martsinkovsky long exact sequence to prove the following.

PROPOSITION 5.4.4. If $\operatorname{cd}_{\mathbb{O}} G < \infty$ then $\operatorname{cd}_{\mathbb{O}} G \leq \operatorname{Gcd}_{\mathbb{Z}} G$.

We know of no groups for which $\operatorname{cd}_{\mathbb{Q}} G \leq \operatorname{Gcd}_{\mathbb{Z}} G$ fails. If $\operatorname{cd}_{\mathbb{Z}} G < \infty$ then necessarily $\operatorname{cd}_{\mathbb{Z}} G = \operatorname{Gcd} G$ [ABS09, Corollary 2.25], but we cannot rule out the possibility that there exists a torsion-free group G with $\operatorname{cd}_{\mathbb{Z}} G = \infty$ but $\operatorname{Gcd} G < \infty$. In fact, the question below is still open even for torsion-free groups.

QUESTION 1.6.2. Do there exist groups G with $\operatorname{cd}_{\mathbb{Q}} G = \infty$ but $\operatorname{Gcd}_{\mathbb{Z}} G < \infty$?

1.7. Bredon duality groups

A duality group is a group G of type FP for which

$$H^{i}(G, \mathbb{Z}G) \cong \begin{cases} \mathbb{Z}\text{-flat} & \text{if } i = n, \\ 0 & \text{else,} \end{cases}$$

where n is necessarily the cohomological dimension of G. The name duality comes from the fact that this condition is equivalent to existence of a $\mathbb{Z}G$ -module D, giving an isomorphism

$$H^{i}(G, M) \cong H_{n-i}(G, D \otimes_{\mathbb{Z}} M)$$

for all *i* and all $\mathbb{Z}G$ -modules M. It can be proven that given such an isomorphism, the module D is necessarily $H^n(G, \mathbb{Z}G)$. A duality group G is called a *Poincaré duality group* if in addition

$$H^i(G, \mathbb{Z}G) \cong \begin{cases} \mathbb{Z} & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

These groups were first defined by Bieri [**Bie72**], and independently by Johnson–Wall [**JW72**]. Duality groups were first studied by Bieri and Eckmann in [**BE73**]. See [**Dav00**] and [**Bie81**, §III] for a thorough introduction.

If a group G has an oriented manifold model for BG then G is a Poincaré duality group [**Dav00**, p.1]. Wall asked if the converse is true [**Wal79**], the answer is no as Poincaré duality groups can be built which are not finitely presented [**Dav98**, Theorem C]. However the question remains a significant open problem if we include the requirement that G be finitely presented. The conjecture is known to hold only in dimensions at most 2 [**Eck87**].

Let R be a commutative ring. A group G is a *duality group over* R if G is FP over R and

$$H^{i}(G, RG) \cong \begin{cases} R\text{-flat} & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

G is Poincaré duality over R if

$$H^{i}(G, RG) \cong \begin{cases} R & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

An analogue of Wall's conjecture is whether every torsion-free finitely presented Poincaré duality group over R is the fundamental group of an aspherical closed R-homology manifold [**Dav00**, Question 3.5]. This is answered in the negative by Fowler for $R = \mathbb{Q}$ [**Fow12**], but remains open for $R = \mathbb{Z}$.

We study a generalisation of Poincaré duality groups, looking at the algebraic analogue of the condition that G admits a manifold model M for $\mathbb{E}_{\mathcal{F}in}G$ such that for any finite subgroup H the fixed point set M^H is a submanifold.

If G admits a cocompact manifold model M for $E_{\mathcal{F}in}G$ then G is $\mathcal{O}_{\mathcal{F}in}FP$. Also if for any finite subgroup H the fixed point set M^H is a submanifold, we have the following condition on the cohomology of the Weyl groups $WH = N_G H/H$:

$$H^{i}(WH, \mathbb{Z}[WH]) = \begin{cases} \mathbb{Z} & \text{if } i = \dim M^{H}, \\ 0 & \text{else,} \end{cases}$$

see [**DL03**, p.3] for a proof of the above. Building on this, in [**DL03**] and also in [**MP13a**, Definition 5.1] a *Bredon duality group over* R is defined as a group G of type $\mathcal{O}_{\mathcal{F}in}$ FP such that for every finite subgroup H of G there is an integer d_H with

$$H^{i}(WH, R[WH]) = \begin{cases} R\text{-flat} & \text{if } i = d_{H}, \\ 0 & \text{else.} \end{cases}$$

Furthermore, G is said to be *Bredon–Poincaré duality over* R if for all finite subgroups H,

$$H^{d_H}(WH, R[WH]) = R.$$

We say that a Bredon duality group G is dimension n if $\mathcal{O}_{\text{fin}} \operatorname{cd} G = n$. Note that for torsion-free groups these definitions reduce to the usual definitions of duality and Poincaré duality groups.

One might generalise Wall's conjecture: Let G be Bredon–Poincaré duality over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H. Does Gadmit a cocompact manifold model M for $\mathbb{E}_{\mathcal{F}in}G$? This is false by an example of Jonathan Block and Schmuel Weinberger, suggested to us by Jim Davis.

THEOREM 6.2.7. There exist examples of Bredon–Poincaré duality groups over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H but G doesn't admit a cocompact manifold model M for $\mathbb{E}_{\text{fin}}G$.

If G is Bredon–Poincaré duality and virtually torsion-free then G is virtually Poincaré duality. Thus an obvious question is whether all virtually Poincaré duality groups are Bredon–Poincaré duality, in [**DL03**] it is shown that this is not the case for $R = \mathbb{Z}$. An example is also given in [**MP13a**, §6] which fails for both $R = \mathbb{Z}$ and for $R = \mathbb{F}_p$, the finite field of p elements. One might also ask if every Bredon–Poincaré duality group is virtually torsion-free but this is also not the case, see for instance Examples 6.2.5 and 6.2.21.

In [Ham11, Theorems D,E] Hamilton shows that, over a field \mathbb{F} of characteristic p, given an extension Γ of a torsion-free group G of type $\operatorname{FP}_{\infty}$ by a finite pgroup, the group Γ will be of type $\mathcal{O}_{\operatorname{fin}}\operatorname{FP}_{\infty}$ (by examples of Leary and Nucinkis, an extension by an arbitrary finite group may not even be $\mathcal{O}_{\operatorname{fin}}\operatorname{FP}_0$ [LN03]). Martínez-Pérez builds on this result to show that if G is assumed Poincaré duality then Γ is Bredon–Poincaré duality over \mathbb{F} with $\mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{F}}\Gamma = \operatorname{cd}_{\mathbb{F}}G$ [MP13a, Theorem C]. However, her results do not extend to Bredon duality groups.

Given a Bredon duality group G we write $\mathcal{V}(G)$ for the set

$$\mathcal{V}(G) = \{d_F : F \text{ a non-trivial finite subgroup of } G\} \subseteq \{0, \dots, n\}.$$

In Example 6.6.8 we will build Bredon duality groups with arbitrary $\mathcal{V}(G)$. If G has a manifold model, or homology manifold model, for $\mathbb{E}_{\mathcal{F}in}G$ then there are some restrictions on $\mathcal{V}(G)$ —see Section 6.2.3 for this. In Section 6.3 we build Bredon–Poincaré duality groups for many choices of $\mathcal{V}(G)$, however the following question remains open:

QUESTION 1.7.1. Is it possible to construct Bredon–Poincaré duality groups with prescribed $\mathcal{V}(G)$?

It follows from Proposition 6.1.4 that for a Bredon–Poincaré duality group, $d_1 \leq \mathcal{O}_{\mathcal{F}in} \operatorname{cd} G$ (recall d_1 is the integer for which $H^{d_1}(G, RG) \cong R$) and also, if we are working over \mathbb{Z} , then $d_1 = \operatorname{cd}_{\mathbb{Q}} G$ (Lemma 6.1.2). Thus the following question is of interest.

QUESTION 1.7.2. Do there exist Bredon duality groups with $\mathcal{O}_{\text{fin}} \operatorname{cd} G \neq d_1$?

Examples of groups for which $\operatorname{cd}_{\mathbb{Q}} G \neq \mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Z}} G$ are known [**LN03**], but there are no known examples of type $\mathcal{O}_{\operatorname{fin}}\operatorname{FP}_{\infty}$. This question is also related to [**MP13a**, Question 5.8] where it is asked whether a virtually torsion-free Bredon duality group satisfies $\mathcal{O}_{\operatorname{fin}}\operatorname{cd} G = \operatorname{vcd} G$.

One might hope to give a definition of Bredon–Poincaré duality groups in terms of Bredon cohomology only, we do not know if this is possible but we show in Section 6.7 that the naïve idea of asking that a group be \mathcal{O}_{fin} FP with

$$H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G, R[?, -]) \cong \begin{cases} \underline{R} & \text{if } i = n, \\ 0 & \text{else,} \end{cases}$$

is not the correct definition, where in the above $H^i_{\mathcal{O}_{fin}}$ denotes the Bredon cohomology and <u>R</u> is the constant covariant Bredon module. Namely we show in Theorem 6.7.3 that any such group is necessarily a torsion-free Poincaré duality group over R.

1.8. Houghton's groups

Houghton's group H_n was introduced in [Hou79], as an example of a group acting on a set S with $H^1(H_n, A \otimes \mathbb{Z}[S]) = A^{n-1}$ for any abelian group A.

In [**Bro87**], Brown used an important new technique to show that the groups $F_{n,r}, T_{n,r}$, and $V_{n,r}$ of Thompson and Higman were FP_{∞} . In the same paper he showed that Houghton's group H_n is interesting from the viewpoint of cohomological finiteness conditions, namely H_n is FP_{n-1} but not FP_n . Brown proves this by studying the action of H_n on the geometric realisation $|\mathcal{M}|$ of a certain poset \mathcal{M} . More recently, Johnson gave a finite presentation for H_3 [Joh99], and later Lee did the same for H_n where $n \geq 3$ [Lee12].

Interestingly, H_n embeds in Thompson's group $V = V_{2,1}$ for all $n \ge 0$ [**Röv99**]. Antolín, Burillo, and Martino have shown that for $n \ge 2$, the group H_n has solvable conjugacy problem [**ABM13**] and Burillo, Cleary, Martino, and Röver have calculated the automorphism groups and abstract commensurators of H_n [**BCMR14**].

There has been recent interest in the structure of the centralisers of Thompson's groups and their generalisations [MPN13, BBG⁺11, MPMN13]. The results obtained here are similar to [MPMN13, 4.10,4.11] where it is shown that in the groups $V_r(\Sigma)$, generalisations of Thompson's V, the centralisers of finite subgroups are of type FP_{∞} whenever the groups $V_r(\Sigma)$ are of type FP_{∞}.

In Section 7.1 we completely describe centralisers of finite subgroups and prove the following.

COROLLARY 7.1.7. If Q is a finite subgroup of H_n then the centraliser $C_{H_n}Q$ is FP_{n-1} but not FP_n .

This contrasts with [**KMPN11a**] where examples are given of soluble groups of type FP_n with centralisers of finite subgroups that are not FP_n, although it is shown in [**MPN10**] that in virtually soluble groups of type FP_{∞} the centralisers of all finite subgroups are of type FP_{∞}.

In Section 7.2 our analysis is extended to arbitrary elements and virtually cyclic subgroups. Using this information elements in H_n are constructed whose centralisers are FP_i for any $0 \le i \le n-3$.

In Section 7.3 the space $|\mathcal{M}|$ mentioned previously is shown to be a model for $\mathbb{E}_{\text{fin}}H_n$.

Finally Section 7.4 contains a discussion of Bredon (co)homological finiteness conditions that are satisfied by Houghton's group. In particular we calculate the Bredon cohomological dimension with respect to the family of finite subgroups, and use a construction of Lück and Weiermann [LW12] to calculate the Bredon cohomological dimension with respect to the family of virtually cyclic subgroups.

PROPOSITION 7.4.3 AND THEOREM 7.4.4. $\mathcal{O}_{\text{fin}} \operatorname{cd} H_n = \mathcal{O}_{\mathcal{V}_{\text{fin}}} \operatorname{cd} H_n = n.$

CHAPTER 2

Modules over a category

Much of this chapter is based on [Lüc89]. Although we consider a slightly more general situation, as explained in Remark 2.0.1, the idea is the same. The material in this chapter is used in much of this thesis, especially in Chapters 3 and 4.

Let R be a commutative ring with unit and \mathfrak{C} a small \mathbf{Ab} category (sometimes called a preadditive category) with the condition below.

(A) For any two objects x and y in \mathfrak{C} , the set of morphisms, denoted $[x, y]_{\mathfrak{C}}$, between x and y is a free abelian group.

Recall that an **Ab** category is one where the morphisms between any two objects form an abelian group and where morphism composition distributes over this addition [**Wei94**, A.4].

REMARK 2.0.1. In [Lüc89, 9.2], categories \mathfrak{X} are considered with the property that every endomorphism in \mathfrak{X} is an isomorphism. However the approach to defining modules over a category in [Lüc89, 9.2] is different from that used here (see also Remark 2.0.7). One can translate between the different viewpoints in the following way:

 $[x, y]_{\mathfrak{X}} = \mathbb{Z}[\{\text{Morphisms } x \to y \text{ in the sense of Lück}\}],$

where $\mathbb{Z}[X]$ denotes the free abelian group on a set X.

The correct analogue of Lück's property with our definitions is the following:

(EI) For every $x \in \mathfrak{C}$, there is a distinguished basis of $[x, x]_{\mathfrak{C}}$, the elements of which are isomorphisms.

The main advantage of the (EI) property is that it allows objects in \mathfrak{C} to be given a partial order: setting $x \leq y$ if $[x, y]_{\mathfrak{C}}$ is non-empty. We choose not to ask for this property in this section, since we want everything discussed here to be relevant to the Mackey and Hecke categories, discussed in Chapter 4, which do not have (EI). The motivating example of a category with (EI) is the orbit category, see Example 2.0.6.

Throughout, the fraktur letters \mathfrak{C} , \mathfrak{D} , \mathfrak{E} etc. will always denote small \mathbf{Ab} categories with (A).

Define the category of covariant \mathfrak{C} -modules over R to be the category of additive covariant functors from \mathfrak{C} to **R-Mod**, the category of left R-modules. Similarly the category of contravariant \mathfrak{C} -modules over R is the category of additive contravariant functors from \mathfrak{C} to **R-Mod**.

If neither "covariant" or "contravariant" is specified in a statement about \mathfrak{C} -modules, the reader should assume the statement holds for both covariant and contravariant \mathfrak{C} -modules.

Since \mathfrak{C} -modules form a functor category and **R-Mod** is an abelian category, the category of \mathfrak{C} -modules is an abelian category [**Mur06**, 44]. In fact, it inherits all of Grothendieck's axioms for an abelian category which are satisfied by **R-Mod** [**Mur06**, 44,55], namely:

- (1) AB3 and AB4—Every small colimit exists and products of exact sequences are exact.
- (2) AB3* and AB4*—Every small limit exists and coproducts of exact sequences are exact.
- (3) AB5—Filtered colimits of exact sequences are exact.

Again because we are working in a functor category, a sequence of \mathfrak{C} -modules

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is exact if and only if it is exact when evaluated at every $x \in \mathfrak{C}$. Note that 0 denotes the zero functor, sending every object to the zero module. Similarly, using the fact that the category of \mathfrak{C} -modules is a functor category and the category of abelian groups is complete, limits and colimits are computed pointwise [**Mur06**, p.8].

Since $[x, y]_{\mathfrak{C}}$ is abelian for all x and y in \mathfrak{C} , for any $y \in \mathfrak{C}$ we can form a contravariant module $R[-, y]_{\mathfrak{C}}$ by

$$R[-,y]_{\mathfrak{C}}(x) = R \otimes_{\mathbb{Z}} [x,y]_{\mathfrak{C}}.$$

The analogous construction for covariant modules gives us

$$R[y,-]_{\mathfrak{C}}(x) = R \otimes_{\mathbb{Z}} [y,x]_{\mathfrak{C}}$$

In Section 2.2 we will show that these modules are analogues of free modules in the category of \mathfrak{C} -modules. Since $R[x, y]_{\mathfrak{C}}$ is a free *R*-module we write $r\alpha$ instead of $r \otimes \alpha$, for $r \in R$ and $\alpha \in [x, y]_{\mathfrak{C}}$.

If $f \in R[x, y]_{\mathfrak{C}}$, where $f = \sum_{i} r_i f_i$ for some $f_i \in [x, y]_{\mathfrak{C}}$, and Q is a \mathfrak{C} -module, then we will write Q(f) for the *R*-module homomorphism given by $\sum_{i} r_i Q(f_i)$.

Let A and B be any two covariant \mathfrak{C} -modules, or any two contravariant \mathfrak{C} -modules, then we denote by $\operatorname{Hom}_{\mathfrak{C}}(A, B)$ the \mathfrak{C} -module morphisms between A and B, i.e. the natural transformations from A to B.

LEMMA 2.0.2 (The Yoneda-type lemma). For any covariant functor A and $x \in \mathfrak{C}$, there is an isomorphism, natural in A:

Hom_c
$$(R[x, -]_{c}, A) \cong A(x)$$

 $f \mapsto f(x)(\mathrm{id}_{x})$

Similarly for any contravariant functor M and $x \in \mathfrak{C}$, there is an isomorphism, natural in M:

Hom_{$$\mathfrak{C}$$} $(R[-,x]_{\mathfrak{C}},M) \cong M(x)$
 $f \mapsto f(x)(\mathrm{id}_x)$

The proof is a generalisation of [MV03, p.9] into the setting of \mathfrak{C} -modules.

PROOF. We provide a proof only for covariant modules, that for contravariant modules is similar.

Let f be a morphism $f : R[x, -]_{\mathfrak{C}} \to A$, we claim f is completely determined by f(x). If $\alpha \in R[x, y]_{\mathfrak{C}}$ then we can view α as $\alpha = R[x, \alpha]_{\mathfrak{C}}(\mathrm{id}_x)$, thus

$$f(y)(\alpha) = f(y) \circ R[x, \alpha]_{\mathfrak{C}}(\mathrm{id}_x)$$
$$= A(\alpha) \circ f(x)(\mathrm{id}_x)$$

where we use that f is R-additive and that f is a morphism in the category of \mathfrak{C} -modules—so a natural transformation of functors—meaning the diagram below commutes.

Conversely, given an element $a \in A(x)$ we can define a morphism f, with $f(x)(\mathrm{id}_x) = a$, by

$$f(y): R[x, y]_{\mathfrak{C}} \longrightarrow A(y)$$
$$\alpha \longmapsto A(\alpha)(a).$$

The endomorphisms $[x, x]_{\mathfrak{C}}$ of an object $x \in \mathfrak{C}$ form an associative ring. This ring will appear often, so we write $\operatorname{End}(x)$ instead of $[x, x]_{\mathfrak{C}}$, and write $R \operatorname{End}(x)$ instead of $R \otimes_{\mathbb{Z}} \operatorname{End}(x)$.

REMARK 2.0.3. Given a covariant module A, evaluating A at x gives a left $R \operatorname{End}(x)$ -module, using the action

$$R \operatorname{End}(x) \times A(x) \longrightarrow A(x)$$
$$(f, a) \longmapsto A(f)(a)$$

This is a left $R \operatorname{End}(x)$ -module structure since given any two elements $g, f \in R \operatorname{End}(x)$,

$$(g \circ f) \cdot x = A(g \circ f)(x) = A(g) \circ A(f)(x) = g \cdot (f \cdot x).$$

Similarly, for a contravariant module M, M(x) has a right $R \operatorname{End}(x)$ -module structure.

REMARK 2.0.4. Let $\operatorname{End}(x)$ denote the category with one object and with morphisms the free abelian group $\operatorname{End}(x)$. Clearly $\operatorname{End}(x)$ has property (A) and it's possible to identify covariant $\operatorname{End}(x)$ -modules and left $R \operatorname{End}(x)$ -modules, similarly contravariant $\operatorname{End}(x)$ -modules and right $R \operatorname{End}(x)$ -modules.

There is often a need to consider bi-modules. A \mathfrak{C} - \mathfrak{D} bi-module (can be covariant or contravariant in either variable, although most of the bi-modules we shall use will be covariant in one variable in contravariant in the other), is a functor

$$Q(-,?): \mathfrak{C} \times \mathfrak{D} \to \mathbf{R}\text{-}\mathbf{Mod}$$
.

EXAMPLE 2.0.5. The \mathfrak{C} - \mathfrak{C} bi-module $R[-,?]_{\mathfrak{C}}$ is defined as

 $R[-,?]_{\mathfrak{C}}:(x,y)\mapsto R[x,y]_{\mathfrak{C}}.$

EXAMPLE 2.0.6 (The orbit category). The orbit category, denoted $\mathcal{O}_{\mathcal{F}}$, is the prototypical example of a category with property (A), and will be studied properly in Chapter 3. It was introduced for finite groups by Bredon [**Bre67**], who used the associated cohomology theory, Bredon cohomology, to develop equivariant obstruction theories. It was later generalised to arbitrary groups by Lück [**Lüc89**].

Fix a discrete group G and family \mathcal{F} of subgroups of G, closed under taking subgroups and conjugation. Commonly studied families are the family $\mathcal{F}in$ of all finite subgroups, and the family \mathcal{VCyc} of all virtually cyclic subgroups. The objects of the orbit category $\mathcal{O}_{\mathcal{F}}$ are all transitive G-sets with stabilisers in \mathcal{F} , ie. the G-sets G/H where H is a subgroup in \mathcal{F} . The morphism set $[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ is the free abelian group on the set of G-maps $G/H \to G/K$. A G-map

$$\alpha: G/H \longrightarrow G/K$$
$$H \longmapsto gK$$

is completely determined by the element $\alpha(H) = gK$, and such an element $gK \in G/K$ determines a *G*-map if and only if HgK = gK, usually written as $gK \in (G/K)^H$. Equivalently gK determines a *G*-map if and only if $g^{-1}Hg \leq K$. In particular if $\mathcal{F} \subseteq \mathcal{F}$ in then the orbit category has (EI), since any *G*-map $\alpha: G/K \to G/K$ is automatically an automorphism. The isomorphism classes of elements in $\mathcal{O}_{\mathcal{F}}$, denoted Iso $\mathcal{O}_{\mathcal{F}}$, are exactly the conjugacy classes of subgroups in \mathcal{F} .

REMARK 2.0.7. The morphisms from G/H to G/K in the orbit category are usually defined as just the *G*-maps $G/H \to G/K$. We show that this definition gives an isomorphic module category.

For this remark, let $\mathcal{O}_{\mathcal{F}}'$ denote the category with the same objects as $\mathcal{O}_{\mathcal{F}}$ but with morphisms from G/H to G/K just the *G*-maps $G/H \to G/K$. Let $\iota : \mathcal{O}_{\mathcal{F}}' \to \mathcal{O}_{\mathcal{F}}$ be the faithful inclusion and given an $\mathcal{O}_{\mathcal{F}}$ -module M define an $\mathcal{O}_{\mathcal{F}}'$ module $M' = M \circ \iota$. We claim that the functor $M \mapsto M'$ gives an isomorphism of categories between $\mathcal{O}_{\mathcal{F}}$ -modules and $\mathcal{O}_{\mathcal{F}}'$ modules.

Any $\mathcal{O}_{\mathcal{F}}'$ -module M' extends uniquely to an $\mathcal{O}_{\mathcal{F}}$ -module M by setting:

- (1) M(G/H) = M'(G/H) for all $H \in \mathcal{F}$.
- (2) $M(\sum_{i} z_{i}\alpha_{i}) = \sum_{i} z_{i}M'(\alpha_{i})$ for any $\mathcal{O}_{\mathcal{F}}$ -morphism $\sum_{i} z_{i}\alpha_{i}$ written as the sum of *G*-maps α_{i} .

This gives an inverse to the functor $M \mapsto M'$ described above.

2.1. Tensor products

2.1.1. Tensor product over \mathfrak{C}. We describe a construction, due to Lück **[Lüc89**, 9.12], of the categorical tensor product of **[Sch70**, 16.7]**[Fis68]** for the categories of \mathfrak{C} -modules over R.

For M contravariant and A covariant, the tensor product over \mathfrak{C} of M and A is

$$M \otimes_{\mathfrak{C}} A = \bigoplus_{x \in \mathfrak{C}} M(x) \otimes_R A(x) \Big/ \sim$$

where $M(\alpha)(m) \otimes a \sim m \otimes A(\alpha)(a)$ for all morphisms $\alpha \in [x, y]$ in \mathfrak{C} , elements $m \in M(y)$ and $a \in A(x)$, and objects $x, y \in \mathfrak{C}$. Since R is commutative, this construction yields an R-module. The tensor product is associative [**MP02**, Lemma 3.1], and commutes with direct sums.

EXAMPLE 2.1.1. If A is a left $R \operatorname{End}(x)$ -module and M is a right $R \operatorname{End}(x)$ module then, by Remark 2.0.4, A and M can be regarded as covariant and contravariant $\widehat{\operatorname{End}(x)}$ -modules. It's easy to check that

$$M \otimes_{\widehat{\operatorname{End}(x)}} A \cong M \otimes_{R \operatorname{End}(x)} A.$$

PROPOSITION 2.1.2. [Lüc89, p.166][MP02] There are adjoint natural isomorphisms of *R*-modules:

$$\operatorname{Hom}_{\mathfrak{D}}(M(?) \otimes_{\mathfrak{C}} Q(?, -), N(-)) \cong \operatorname{Hom}_{\mathfrak{C}}(M(?), \operatorname{Hom}_{\mathfrak{D}}(Q(?, -), N(-)))$$

$$\operatorname{Hom}_{\mathfrak{C}}(Q(?,-)\otimes_{\mathfrak{D}}A(-),B(?))\cong\operatorname{Hom}_{\mathfrak{D}}(A(-),\operatorname{Hom}_{\mathfrak{C}}(Q(?,-),B(?)))$$

Here M and N are contravariant modules, A and B are covariant modules, and Q(?, -) is an \mathfrak{D} - \mathfrak{C} -bi-module—a contravariant \mathfrak{D} -module in "-" and a covariant \mathfrak{C} -module in "?".

LEMMA 2.1.3. [**MV03**, p.14] There are natural isomorphisms of R-modules for any contravariant module M and covariant module A:

$$M \otimes_{\mathfrak{C}} R[x, -]_{\mathfrak{C}} \cong M(x)$$
$$R[-, x]_{\mathfrak{C}} \otimes_{\mathfrak{C}} A \cong A(x).$$

2.1.2. Tensor product over R. We describe the tensor product over R as in [Lüc89, 9.13]. If A and B are \mathfrak{C} -modules, either both covariant or both contravariant, then the tensor product over R is the \mathfrak{C} -module

$$(A \otimes_R B)(x) = A(x) \otimes_R B(x).$$

If $\alpha : x \to y$ is a morphism in \mathfrak{C} , then

$$(A \otimes_R B)(\alpha) = A(\alpha) \otimes_R B(\alpha).$$

2.2. Frees, projectives, injectives and flats

Free objects in a category are usually defined as left adjoint to some forgetful functor, often with codomain **Set**. For \mathfrak{C} -modules the necessary forgetful functor is

$$U: \{ \mathfrak{C}\text{-modules} \} \longrightarrow [\operatorname{Ob}(\mathfrak{C}), \mathbf{Set}]$$
$$UA: x \longmapsto A(x).$$

Here $[Ob(\mathfrak{C}), \mathbf{Set}]$ denotes the category of functors $Ob(\mathfrak{C}) \to \mathbf{Set}$, where $Ob(\mathfrak{C})$ is the category whose objects are the objects of \mathfrak{C} but with only the identity morphisms at each object. The functor F left adjoint to U is, for $X \in [Ob(\mathfrak{C}), \mathbf{Set}]$,

$$FX = \bigoplus_{x \in \mathfrak{C}} \bigoplus_{X(x)} R[x, -]_{\mathfrak{C}}.$$

Analagously, if we are working with contravariant functors,

$$FX = \bigoplus_{x \in \mathfrak{C}} \bigoplus_{X(x)} R[-, x]_{\mathfrak{C}}.$$
That (F, U) form an adjoint pair is a consequence of the Yoneda-type Lemma 2.0.2—for any covariant module A,

$$\operatorname{Hom}_{\mathfrak{C}}(FX, A) = \operatorname{Hom}_{\mathfrak{C}} \left(\bigoplus_{x \in \mathfrak{C}} \bigoplus_{X(x)} R [x, -]_{\mathfrak{C}}, A \right)$$
$$\cong \prod_{x \in \mathfrak{C}} \prod_{X(x)} \operatorname{Hom}_{\mathfrak{C}} \left(R [x, -]_{\mathfrak{C}}, A \right)$$
$$\cong \prod_{x \in \mathfrak{C}} \prod_{X(x)} A(x)$$
$$\cong \operatorname{Hom}_{[\operatorname{Ob}(\mathfrak{C}), \operatorname{Set}]}(X, UA).$$

The proof for contravariant functors is analogous.

Projective and injective modules are defined as in any abelian category—a \mathfrak{C} -module P is projective if $\operatorname{Hom}_{\mathfrak{C}}(P, -)$ is exact and a \mathfrak{C} -module I is injective if $\operatorname{Hom}_{\mathfrak{C}}(-, I)$ is exact [Wei94, §2.2]. Free modules are projective since if

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is an exact sequence of \mathfrak{C} -modules then, by the Yoneda-type Lemma (2.0.2), applying $\operatorname{Hom}_{\mathfrak{C}}(R[x,?]_{\mathfrak{C}},-)$ gives the exact sequence

$$0 \longrightarrow A(x) \longrightarrow B(x) \longrightarrow C(x) \longrightarrow 0.$$

Since direct sums of projectives are projective in any abelian category, this is enough to show the category of \mathfrak{C} -modules has enough projectives, in fact the counit of the adjunction between F and U,

$$\eta: (FUA) \longrightarrow A,$$

is always an epimorphism: By construction,

$$FUA = \bigoplus_{x \in \mathfrak{C}} \bigoplus_{a \in A(x)} F_a(x, -)$$

where $F_a(x, -) \cong R[x, -]_{\mathfrak{C}}$. The counit is the map defined on $F_a(x, -)$, via the Yoneda-type Lemma 2.0.2, by $\mathrm{id}_x \mapsto a$. It's clear that every $a \in A(x)$ is in the image of $\eta(x)$, and thus η is an epimorphism.

The category of \mathfrak{C} -modules also has enough injectives, see Remark 2.3.4 for a proof using coinduction.

A covariant (respectively contravariant) \mathfrak{C} -module F is flat if the functor $F \otimes_{\mathfrak{C}} \dagger$ (respectively $\dagger \otimes_{\mathfrak{C}} F$) is flat. Lemma 2.1.3 shows free modules are flat, and since the tensor product commutes with direct sums, projectives are flat also.

A covariant \mathfrak{C} -module M is said to be *finitely generated* if there exists an epimorphism

$$\bigoplus_{x \in I} R[x, -]_{\mathfrak{C}} \longrightarrow M,$$

for some finite indexing set I of objects in \mathfrak{C} . There is an analogous definition for contravariant \mathfrak{C} -modules.

2.3. Restriction, induction and coinduction

Given a functor $\iota : \mathfrak{C} \to \mathfrak{D}$, we define restriction, induction, and coinduction functors. Induction and restriction can be found in [Lüc89, §9.8] but with the names extension and restriction, he also defines an adjoint pair of functors called "splitting" and "inclusion". We don't include these here as the adjointness of these functors relies on the (EI) property which we are not assuming holds.

Restriction and induction are, for covariant modules:

 $\operatorname{Res}_{\iota} : \{ \operatorname{Covariant} \mathfrak{D}\operatorname{-modules} \} \longrightarrow \{ \operatorname{Covariant} \mathfrak{C}\operatorname{-modules} \}$ $\operatorname{Res}_{\iota} : A \longmapsto A \circ \iota$

$$\operatorname{Ind}_{\iota}: \{\operatorname{Covariant} \mathfrak{C}\operatorname{-modules}\} \longrightarrow \{\operatorname{Covariant} \mathfrak{D}\operatorname{-modules}\}$$

 $\operatorname{Ind}_{\iota}: A \longmapsto R[\iota(?), -]_{\mathfrak{D}} \otimes_{\mathfrak{C}} A(?).$

Where the notation $R[\iota(?), -]_{\mathfrak{D}}$ means that in the variable "?", this functor should be regarded as a \mathfrak{C} -module using ι . Coinduction is, for covariant modules:

$$\begin{split} \text{CoInd}_{\iota} : \{ \text{Covariant } \mathfrak{C}\text{-modules} \} &\longrightarrow \{ \text{Covariant } \mathfrak{D}\text{-modules} \} \\ & \text{CoInd}_{\iota} : A \longmapsto \text{Hom}_{\mathfrak{C}}(R[-,\iota(?)]_{\mathfrak{D}},A(?)). \end{split}$$

For contravariant functors, the definition of restriction is identical, and for induction and coinduction is nearly identical:

$$\operatorname{Ind}_{\iota}: \{\operatorname{Contravariant} \mathfrak{C}\operatorname{-modules}\} \longrightarrow \{\operatorname{Contravariant} \mathfrak{D}\operatorname{-modules}\}$$
$$\operatorname{Ind}_{\iota}: M \longmapsto M(?) \otimes_{\mathfrak{C}} R[-, \iota(?)]_{\mathfrak{D}}$$

 $\operatorname{CoInd}_{\iota} : \{\operatorname{Contravariant} \mathfrak{C}\operatorname{-modules}\} \longrightarrow \{\operatorname{Contravariant} \mathfrak{D}\operatorname{-modules}\}$

 $\operatorname{CoInd}_{\iota}: M(-) \longmapsto \operatorname{Hom}_{\mathfrak{C}}(R[\iota(?), -]_{\mathfrak{D}}, M(?)).$

Usually the functor ι will be implicit, and we will use the notation $\operatorname{Res}^{\mathfrak{D}}_{\mathfrak{C}}$ for $\operatorname{Res}_{\iota}$, and similarly for induction and coinduction. We will also write $\operatorname{Res}^{\mathfrak{C}}_{x}$ instead of $\operatorname{Res}^{\mathfrak{C}}_{\widehat{\operatorname{End}}(x)}$ and similarly for induction and coinduction.

Note that for any left $R \operatorname{End}(x)$ -module A,

$$\operatorname{Ind}_{x}^{\mathfrak{C}} A(x) = R[x, x] \otimes_{R \operatorname{End}(x)} A \cong A$$

 $\operatorname{CoInd}_{x}^{\mathfrak{C}} A(x) = \operatorname{Hom}_{R \operatorname{End}(x)}(R[x, x], A) \cong A.$

Similarly for contravariant induction and coinduction.

PROPOSITION 2.3.1. [MPN06, §2] Induction is left adjoint to restriction and coinduction is right adjoint to restriction.

The following proposition is almost entirely a consequence of this adjointness.

Proposition 2.3.2.

- (1) Restriction is exact.
- (2) Induction is right exact and preserves frees, projectives, flats and "finitely generated".
- (3) Coinduction preserves injectives.
- (4) Induction and restriction preserve colimits and coinduction and restriction preserve limits.
- PROOF. (1) Since a short exact sequence of modules over \mathfrak{C} is exact if and only if it's exact when evaluated at every element of \mathfrak{C} , restriction is always exact.
- (2) Since induction has an exact right adjoint it preserves projectives and is right-exact [Wei94, 2.3.10, 2.6.1].

That induction takes frees to frees is a consequence of Lemma 2.1.3,

 $\operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} R[\dagger, -]_{\mathfrak{C}} = R[\iota(?), -]_{\mathfrak{D}} \otimes_{\mathfrak{C}} R[\dagger, ?]_{\mathfrak{C}} \cong R[\dagger, -]_{\mathfrak{D}}$

and similarly for contravariant modules.

That induction takes flats to flats is a consequence of Lemma 2.3.3 below. In the covariant case, this implies the functor $? \otimes_{\mathfrak{D}} \operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} F$ is naturally isomorphic to the functor $(\operatorname{Res}_{\mathfrak{C}}^{\mathfrak{D}}?) \otimes_{\mathfrak{C}} F$. Thus if F is assumed flat then $? \otimes_{\mathfrak{D}} \operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} F$ is exact. An analogous proof holds for contravariant F.

If A is a finitely generated \mathfrak{C} -module then there is an epimorphism $F \longrightarrow A$ for some finitely generated free F. Induction is right exact so there is an epimorphism

$$\operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} F \longrightarrow \operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} A.$$

Induction takes finitely generated frees to finitely generated frees so $\operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}}A$ is finitely generated.

- (3) Since coinduction has an exact left adjoint it preserves injectives [Wei94, 2.3.10] and is left-exact [Wei94, 2.6.1]
- (4) This is another consequence of adjointness [ML98, p.118].

LEMMA 2.3.3. There exist natural isomorphisms for any contravariant \mathfrak{C} -module M and covariant \mathfrak{C} -module A:

$$M \otimes_{\mathfrak{D}} \operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} A \cong \operatorname{Res}_{\mathfrak{C}}^{\mathfrak{D}} M \otimes_{\mathfrak{C}} A$$
$$\operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} M \otimes_{\mathfrak{D}} A \cong M \otimes_{\mathfrak{C}} \operatorname{Res}_{\mathfrak{C}}^{\mathfrak{D}} A.$$

PROOF. We prove the first natural isomorphism, the second is analogous:

$$M \otimes_{\mathfrak{D}} \operatorname{Ind}_{\mathfrak{C}}^{\mathfrak{D}} A \cong M(-) \otimes_{\mathfrak{D}} \left(R[?,-]_{\mathfrak{D}} \otimes_{\mathfrak{C}} A(?) \right)$$
$$\cong \left(M(-) \otimes_{\mathfrak{D}} R[?,-]_{\mathfrak{D}} \right) \otimes_{\mathfrak{C}} A(?)$$
$$\cong \operatorname{Res}_{\mathfrak{C}}^{\mathfrak{D}} M \otimes_{\mathfrak{C}} A.$$

REMARK 2.3.4 (The category of \mathfrak{C} -modules has enough injectives). A consequence of Proposition 2.3.2(3) is that the category of \mathfrak{C} -modules has enough injectives. For any ring S and module M over S there always exists an injective module I and injection $M \hookrightarrow I$ [Wei94, 2.3.11]. Given a \mathfrak{C} -module M choose injective $R \operatorname{End}(x)$ -modules I_x such that M(x) injects into I_x for all $x \in \mathfrak{C}$, and consider the map

$$\prod_{x \in \mathfrak{C}} \eta_x : M \longrightarrow \prod_{x \in \mathfrak{C}} \operatorname{CoInd}_{R \operatorname{End}(x)}^{\mathfrak{C}} I_x$$

where η_x is chosen, via the adjointness of coinduction and restriction, such that $\eta_x(x)$ is the inclusion of M(x) into $\operatorname{CoInd}_x^{\mathfrak{C}} I_x(x) = I_x$.

Clearly the product of the η_x maps is an injection. The module on the right is injective by Proposition 2.3.2(3) and the fact that in any abelian category, products of injective modules are injective.

EXAMPLE 2.3.5. If A and B are covariant \mathfrak{C} -modules, we define a \mathfrak{C} - \mathfrak{C} bimodule:

$$A(?) \otimes_R B(-) : (x, y) \mapsto A(x) \otimes A(y).$$

Denote by $\Delta : \mathfrak{C} \to \mathfrak{C} \times \mathfrak{C}$ the diagonal functor $\Delta : x \to (x, x)$. The tensor product over R defined in Section 2.1.2 could be defined as

$$A \otimes_R B = \operatorname{Res}_{\Delta}(A(?) \otimes_R B(-)).$$

2.4. Tor and Ext

Since the categories of \mathfrak{C} -modules are abelian and have enough projectives, it is possible to use techniques from homological algebra to study them. For M a \mathfrak{C} -module, a projective resolution P_* of M is an exact chain complex of \mathfrak{C} -modules,

$$\cdots \longrightarrow P_i \longrightarrow P_{i-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow M \longrightarrow 0$$

where each P_i is projective.

If A is a covariant \mathfrak{C} -module and P_* a projective resolution of A then for any covariant module B and contravariant module M, we define $\operatorname{Ext}^*_{\mathfrak{C}}$ and $\operatorname{Tor}^{\mathfrak{C}}_*$ as

$$\operatorname{Ext}^{k}_{\mathfrak{C}}(A,B) = H^{k} \operatorname{Hom}_{\mathfrak{C}} \left(P_{*}, B \right)$$
$$\operatorname{Tor}^{\mathfrak{C}}_{k}(M,A) = H_{k} \left(M \otimes_{\mathfrak{C}} P_{*} \right).$$

We make the same definitions for contravariant modules, if M is a contravariant module, Q_* a projective resolution of M, A a covariant module and N a contravariant module then

$$\operatorname{Ext}_{\mathfrak{C}}^{k}(M,N) = H^{k}\operatorname{Hom}_{\mathfrak{C}}(Q_{*},N)$$

$$\operatorname{Tor}_{k}^{\mathfrak{C}}(M,A) = H_{k}(Q_{*} \otimes_{\mathfrak{C}} A)$$

A priori $\operatorname{Tor}_*^{\mathfrak{C}}$ has just been given two definitions, these are equivalent by Proposition 2.4.1 below, an analogue of the classical result that Tor for modules over a ring can be computed using a resolution in either variable.

PROPOSITION 2.4.1. If A is any covariant module, M is any contravariant module, P_* is a projective covariant resolution of A, and Q_* is a projective contravariant resolution of M then for all k,

$$H_k(M \otimes_{\mathfrak{C}} P_*) \cong H_k(Q_* \otimes_{\mathfrak{C}} A).$$

We need some notation for the proof: If (C_*, ∂_*) is an arbitrary chain complex of \mathfrak{C} -modules then we write C_{*+j} for the chain complex whose degree *i* term is C_{i+j} , and differential $(-1)^j \partial_{i+j}$. This change in the differential doesn't affect exactness, as the homology groups of the new complex are simply $H_n(C_{*+j}) =$ $H_{n+j}(C_*)$.

PROOF. The proof is a generalisation of [Wei94, Theorem 2.7.2, p.58] into the setting of \mathfrak{C} -modules. Form three double complexes, $M \otimes_{\mathfrak{C}} P_*$, $Q_* \otimes_{\mathfrak{C}} P_*$ and $Q_* \otimes_{\mathfrak{C}} A$. The augmentation maps $\varepsilon : P_* \longrightarrow A$ and $\eta : Q_* \longrightarrow M$ induce maps between the total complexes,

$$\operatorname{Tot}\left(Q_*\otimes_{\mathfrak{C}} P_*\right)\longrightarrow \operatorname{Tot}\left(M\otimes_{\mathfrak{C}} P\right)\cong M\otimes_{\mathfrak{C}} P_*$$

$$\operatorname{Tot}\left(Q_*\otimes_{\mathfrak{C}} P_*\right)\longrightarrow \operatorname{Tot}\left(Q_*\otimes_{\mathfrak{C}} A\right)\cong Q_*\otimes_{\mathfrak{C}} A_*$$

where Tot denotes the total complex of a bicomplex of R-modules (see [Wei94, 1.2.6] for the definition of total complex). We claim that these maps are weak equivalences. Define a new double complex C_{**} , by adding $A_* \otimes_{\mathfrak{C}} Q_{*-1}$ in the (-1) column of $P_* \otimes_{\mathfrak{C}} Q_*$, giving the following complex. Note that we need to shift Q_* so that the resulting complex is a bi-complex, without the shift the horizontal

and vertical differentials would not anti-commute.



The complex $\operatorname{Tot}(C_{**})_{*+1}$ is the mapping cone of $\varepsilon \otimes_{\mathfrak{C}} \operatorname{id}_Q$, so it suffices to show that it is acyclic (see [Wei94, §1.5]). But this follows from the Acyclic Assembly Lemma [Wei94, 2.7.3], since the flatness of Q_i means the functor $\dagger \otimes_{\mathfrak{C}} Q_i$ is exact for all *i* and hence the rows of C_{**} are exact.

Similarly, the mapping cone of $\operatorname{id}_P \otimes_{\mathfrak{C}} \eta$ is the complex $\operatorname{Tot}(D_{**})_{*+1}$, where D_{**} is the double complex obtained by adding $P_{*-1} \otimes_{\mathfrak{C}} B$ in row (-1) to the complex $P_* \otimes_{\mathfrak{C}} Q_*$. Since P_i is flat for all $i, P_i \otimes_{\mathfrak{C}} \dagger$ is exact, and the columns of D_{**} are exact. Thus $\operatorname{Tot}(D_{**})_{*+1}$ is acyclic, again by the Acyclic Assembly Lemma [Wei94, 2.7.3], showing $\operatorname{id}_P \otimes_{\mathfrak{C}} \eta$ is a weak equivalence.

 $\operatorname{Tor}_{*}^{\mathfrak{C}}$ could also be calculated using flat resolutions instead of projective resolutions. The standard proof of this in the case of modules over a ring goes through with almost no modification, see for example [Wei94, 3.2.8]. Similarly, we could calculate $\operatorname{Ext}_{\mathfrak{C}}^*$ using injective resolutions, again the proof is the standard one.

2.5. Finiteness conditions

We define projective and flat dimensions as one would expect, the *projective* dimension $\mathfrak{C}pdA$ of a contravariant \mathfrak{C} -module A is the minimal length of a projective resolution of A and the *flat dimension* $\mathfrak{C}fdA$ is the minimal length of a flat resolution. These can be characterised as the vanishing of the $\operatorname{Ext}^*_{\mathfrak{C}}$ and $\operatorname{Tor}^{\mathfrak{C}}_*$ groups as in ordinary homological algebra.

Recall that a \mathfrak{C} -module is finitely generated if it admits an epimorphism from a finite direct sum of modules of the form $R[x, -]_{\mathfrak{C}}$ for some $x \in \mathfrak{C}$. We say a \mathfrak{C} -module A is \mathfrak{CFP}_n if there is a projective resolution of A which is finitely generated up to degree n. Additionally we call \mathfrak{CFP}_0 modules *finitely generated* and \mathfrak{CFP}_1 modules *finitely presented*. There is an analogue of the Bieri-Eckmann criterion [**BE74**], see also [**Bie81**, Theorem 1.3]. A proof in the case that $\mathfrak{C} = \mathcal{O}_{\mathcal{F}}$ appears in [MPN13, Theorem 5.3] and no substantial change is required to prove for \mathfrak{C} -modules.

THEOREM 2.5.1 (Bieri–Eckmann Criterion). The following conditions on any contravariant \mathfrak{C} -module A are equivalent:

- (1) A is \mathfrak{CFP}_n .
- (2) If B_{λ} , for $\lambda \in \Lambda$, is a filtered system of \mathfrak{C} -modules then the natural map

$$\varinjlim_{\Lambda} \operatorname{Ext}^k_{\mathfrak{C}}(A, B_{\lambda}) \longrightarrow \operatorname{Ext}^k_{\mathfrak{C}}(A, \varinjlim_{\Lambda} B_{\lambda})$$

is an isomorphism for $k \leq n-1$ and a monomorphism for k = n.

(3) For any filtered system B_{λ} , for $\lambda \in \Lambda$, such that $\varinjlim_{\Lambda} B_{\lambda} = 0$,

$$\varinjlim_{\Lambda} \operatorname{Ext}^k_{\mathfrak{C}}(A, B_{\lambda}) = 0$$

for all $k \leq n$.

(4) For any collection of indexing sets Λ_x , for $x \in \mathfrak{C}$, the natural map

$$\operatorname{Tor}_{k}^{\mathfrak{C}}\left(M,\prod_{x\in\operatorname{Ob}\mathfrak{C}}\prod_{\Lambda_{x}}R[x,-]_{\mathfrak{C}}\right)\longrightarrow\prod_{x\in\operatorname{Ob}\mathfrak{C}}\prod_{\Lambda_{x}}\operatorname{Tor}_{k}^{\mathfrak{C}}\left(M,R[x,-]_{\mathfrak{C}}\right)$$

is an isomorphism for k < n and an epimorphism for k = n.

There is a similar result for covariant modules.

LEMMA 2.5.2. If

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is a short exact sequence of \mathfrak{C} -modules then

- (1) If A and B are \mathfrak{CFP}_n then C is \mathfrak{CFP}_n .
- (2) If A and C are \mathfrak{CFP}_n then B is \mathfrak{CFP}_n .
- (3) If B and C are \mathfrak{CFP}_n then A is \mathfrak{CFP}_{n-1} .

PROOF. This follows from the long exact sequence associated to $\text{Ext}^*_{\mathfrak{C}}$ and the Bieri–Eckmann criterion (Theorem 2.5.1).

CHAPTER 3

Bredon modules

Fix a family \mathcal{F} of subgroups of G, closed under subgroups and conjugation, and recall from Example 2.0.6 that the orbit category $\mathcal{O}_{\mathcal{F}}$ is the category whose objects are all transitive G-sets with stabilisers in \mathcal{F} and whose morphism set $[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ is the free abelian group on the set of G-maps $G/H \longrightarrow G/K$. Common families to study are the family $\mathcal{F}in$ of all finite subgroups and the family $\mathcal{V}Cyc$ of all virtually cyclic subgroups.

Contravariant $\mathcal{O}_{\mathcal{F}in}$ -modules and their associated finiteness conditions provide a good algebraic reflection of the geometric world of proper actions. This background has already been discussed in the introduction and we will discuss connections with geometry in Sections 3.3 and 3.5 also.

Sections 3.1 and 3.2 specialise information from Chapter 2 to modules over the orbit category, and the later sections discuss finiteness conditions for contravariant $\mathcal{O}_{\mathcal{F}}$ -modules.

Recall that a G-map $\alpha : G/H \longrightarrow G/K$ is completely determined by the element $\alpha(H) = gK$, and such an element $gK \in G/K$ determines a G-map if and only if HgK = gK, equivalently $g^{-1}Hg \leq K$.

3.1. Free modules

For this section we require that $\mathcal{F} \subseteq \mathcal{F}in$. In this section we describe the structure of free $\mathcal{O}_{\mathcal{F}}$ -modules. Throughout this section H and K will denote subgroups in \mathcal{F} and α_g will denote a G-map $\alpha_g : G/H \to G/K$ sending $H \mapsto gK$ for any H and K.

REMARK 3.1.1 (Structure of End(G/H)). If $\alpha_g : G/H \longrightarrow G/H$ is the *G*map sending $H \longmapsto gH$ then necessarily $g \in N_G H$ and two such g determine the same *G*-map if they are in the same left *H*-coset. Furthermore $\alpha_h \circ \alpha_g = \alpha_{gh}$ so, denoting by *WH* the Weyl group $N_G H/H$,

$$\operatorname{End}(G/H) = \widehat{\mathbb{Z}[WH]}^{\operatorname{op}}.$$

Here $\mathbb{Z}[\widehat{WH}]$ denotes the category of one element and morphisms given by $\mathbb{Z}[WH]$, and $\mathbb{Z}[\widehat{WH}]^{\text{op}}$ is the opposite of that category. As described in Remark 2.0.3, if Ais a covariant \mathfrak{C} -module then evaluating at x gives A(x) a left $R \operatorname{End}(x)$ -structure. Thus evaluating a covariant $\mathcal{O}_{\mathcal{F}}$ -module at G/H gives a left $\widehat{R[WH]}^{\text{op}}$ -structure, equivalently a right R[WH]-structure.

Similarly, if M is a contravariant $\mathcal{O}_{\mathcal{F}}$ -module then evaluating at G/H gives M(G/H) a left R[End(G/H)] structure, equivalently a left R[WH]-module structure.

Note that this description may fail when $\mathcal{F} \not\subseteq \mathcal{F}in$, as it is possible to have an infinite cyclic subgroup H of a group G along with an element $g \in G$ such that $g^{-1}Kg$ is a proper subgroup of K. This occurs for example in the Baumslag–Solitar group BS(1,2).

EXAMPLE 3.1.2 (Right action of R[WK] on $R[G/H, -]_{\mathcal{O}_{\mathcal{F}}}(G/K)$). The action of WK on $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ is as follows: If $f: G/H \to G/K$ with f(H) = gKand $w \in WK$ then

$$f \cdot w = R[G/H, \alpha_w]_{\mathcal{O}_{\mathcal{F}}}(f) = \alpha_w \circ f.$$

Since $(\alpha_w \circ f)(1) = gwK$, under the identification

$$R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} \cong R[(G/K)^H],$$

the action is given by $gK \cdot w = gwK$.

LEMMA 3.1.3. There is an isomorphism of right R[WK]-modules

$$R[G/H, -]_{\mathcal{O}_{\mathcal{F}}}(G/K) = R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} \cong \bigoplus_{\substack{gN_GK \in G/N_GK\\g^{-1}Hg \leq K}} R[WK].$$

PROOF. Firstly, $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} \cong R[(G/K)^H]$ is a free WK-module, since if $n \in N_G K$ is such that gnK = gK then nK = K and hence $n \in K$. Now, gK and g'K lie in the same WK orbit if and only if g(WK)K = g'(WK)K, equivalently $gN_GK = g'N_GK$, and gK determines an element of $R[(G/K)^K]$ if and only if $g^{-1}Hg \leq K$. Thus there is one R[WK] orbit for each element in the set

$$\{gN_GK \in G/N_GK : g^{-1}Hg \le K\}.$$

For contravariant modules the situation is more complex, evaluating at G/H doesn't always give a free R[WH]-module, although it does always give a R[WH]-module of type FP_{∞}. This is proved in the case $R = \mathbb{Z}$ in [**KMPN09**, Proof of 3.2], the proof for general rings R requires no substantial change, and is given in Corollary 3.1.6.

EXAMPLE 3.1.4 (Left action of R[WH] on $R[-, G/K]_{\mathcal{O}_{\mathcal{F}}}(G/H)$). A similar argument to the previous example shows that under the identification

$$R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} \cong R[(G/K)^H]$$

32

the action of R[WH] is given by $w \cdot gK = wgK$.

LEMMA 3.1.5. There is an isomorphism of left R[WH]-modules

$$R[-,G/K]_{\mathcal{O}_{\mathcal{F}}}(G/H) = R[G/H,G/K]_{\mathcal{O}_{\mathcal{F}}} = \bigoplus_{x} R[WH/WH_{xK}]$$

where x runs over a set of coset representatives of the subset of the set of N_GH -K double cosets

$$\{x \in N_G H \setminus G / K : x^{-1} H x \le K\},\$$

and the stabilisers are given by

$$WH_{xK} = \left(N_G H \cap xKx^{-1}\right)/H.$$

PROOF. Recall the identification $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} = R[(G/K)^H]$. The elements xK and yK are in the same WH-orbit if there exists some $nH \in WH$ (where $n \in N_GH$) with

$$nHxK = yK \Leftrightarrow nxK = yK \Leftrightarrow (N_GH)xK = (N_GH)yK.$$

Combining this with the fact that $xK \in (G/K)^H$ if and only if $x^{-1}Hx \leq K$ means there is a WH-orbit for each N_GH -K double coset N_GHxK such that $x^{-1}Hx \leq K$, i.e. coset representatives for

$$\{x \in N_G H \setminus G / K : x^{-1} H x \le K\}$$

are orbit representatives for the R[WH]-orbits in $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$.

The $N_G(H)$ -stabiliser of the point $xK \in (G/K)^H$ is the set

$$\{g \in N_G(H) : gxK = xK\} = \{g \in N_G(H) : g \in xKx^{-1}\} = N_G(H) \cap xKx^{-1}.$$

So the WH-stabiliser of $xK \in (G/K)^H$ is $WH_{xK} = (N_G(H) \cap xKx^{-1})/H.$

COROLLARY 3.1.6. The $\mathcal{O}_{\mathcal{F}}$ -module $R[-, G/K]_{\mathcal{O}_{\mathcal{F}}}(G/H) = R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ is a finite direct sum of projective R[WH]-permutation modules of type FP_{∞} with stabilisers in \mathcal{F} . In particular $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ is FP_{∞} .

PROOF. Since K is finite, the set $\{x \in N_G H \setminus G/K : x^{-1} H x \leq K\}$ is finite and $R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}$ can be written as a finite direct sum

$$R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} = \bigoplus_{x} R[WH/WH_{xK}]$$

where the WH_{xK} are finite groups. Since R is FP_{∞} as a $R[WH_{xK}]$ -module and

$$R[WH/WH_{xK}] = \operatorname{Ind}_{R[WH_{xK}]}^{R[WH]} R,$$

we can apply Lemma 3.1.7 below and deduce that $R[WH/WH_{xK}]$ is FP_{∞} as an *RG*-module. Finally, any finite direct sum of FP_{∞} modules is FP_{∞}.

LEMMA 3.1.7. If M is FP_{∞} as an RF-module for some subgroup $F \leq G$, then $Ind_{RF}^{RG}M = RG \otimes_{RF} M$ is FP_{∞} as an RG-module. **PROOF.** Let $\prod_i N_i$ be an arbitrary direct product of *RG*-modules, then

$$\operatorname{Tor}_{*}^{RG}\left(\operatorname{Ind}_{RF}^{RG}M,\prod_{i}N_{i}\right) = \operatorname{Tor}_{*}^{RF}\left(M,\prod_{i}N_{i}\right)$$
$$=\prod_{i}\operatorname{Tor}_{*}^{RF}\left(M,N_{i}\right)$$
$$=\prod_{i}\operatorname{Tor}_{*}^{RG}\left(\operatorname{Ind}_{RF}^{RG}M,N_{i}\right)$$

where the first and third equalities come from Shapiro's Lemma. This finishes the proof as $\operatorname{Ind}_{RF}^{RG} M$ is $\operatorname{FP}_{\infty}$ if and only if $\operatorname{Tor}_{*}^{RG}(\operatorname{Ind}_{RF}^{RG} M, -)$ commutes with direct products [**Bro94**, Theorem VIII.4.8].

3.2. Restriction, induction and coinduction

In this section we require that $\mathcal{F} \subseteq \mathcal{F}in$. We specialise the constructions of Section 2.3 to the categories of covariant and contravariant $\mathcal{O}_{\mathcal{F}}$ -modules. In order to match the literature, we write $\operatorname{Ind}_{WH}^{\mathcal{O}_{\mathcal{F}}} A$ instead of $\operatorname{Ind}_{G/H}^{\mathcal{O}_{\mathcal{F}}} A$ for induction with the inclusion functor

$$\iota: \operatorname{End}(\widehat{G/H}) \hookrightarrow \mathcal{O}_{\mathcal{F}}$$

and similarly for restriction and coinduction. Note that the notation for covariant and contravariant induction is the same, if neither covariant or contravariant is specified then contravariant should be assumed.

EXAMPLE 3.2.1. If R is the trivial RG module then inducing to a covariant $\mathcal{O}_{\mathcal{F}}$ -module gives

$$\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} R : G/H \longmapsto R \otimes_{RG} R[G/H] = R.$$

Checking the morphisms as well, $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} R = \underline{R}$, the constant covariant functor on *R*—sending every object to *R* and every *G*-map to the identity.

A group is said to *contain no R-torsion* if for every finite subgroup $F \leq G$, |F| is invertible in R. For example every group has no \mathbb{Q} -torsion. If

$$|F| = p_1^{n_1} \cdots p_m^{n_m}$$

is a prime factorisation of |F| then for each p_i there is an element of order p_i by Cauchy's Theorem [**Rob96**, 1.6.17]. Since the invertible elements R^* form a group, if all the p_i are invertible in R then so is |F|. Hence a group has no R-torsion if and only if the order of every finite-order element is invertible in R.

Recall from Proposition 2.3.2 that covariant and contravariant restriction are exact, in addition we have the following:

Proposition 3.2.2.

(1) Covariant restriction preserves projectives and flats.

- (2) Contravariant restriction preserves finite generation.
- (3) Contravariant restriction at H preserves projectives and flats if WH is R-torsion-free, if not then contravariant restriction takes projectives to FP_∞-modules.
- PROOF. (1) If P is a projective covariant $\mathcal{O}_{\mathcal{F}}$ -module and F a free covariant $\mathcal{O}_{\mathcal{F}}$ -module with a split epimorphism $F \longrightarrow P$ then restricting at G/H yields a split epimorphism $F(G/H) \longrightarrow P(G/H)$, by Lemma 3.1.3 F(G/H) is free and thus P(G/H) is projective.

If F is a flat covariant module and M any left R[WH]-module then,

$$F(G/H) \otimes_{R[WH]} M \cong (R[-, G/H]_{\mathcal{O}_{\mathcal{F}}} \otimes_{\mathcal{O}_{\mathcal{F}}} F) \otimes_{R[WH]} M$$
$$\cong (R[-, G/H]_{\mathcal{O}_{\mathcal{F}}} \otimes_{R[WH]} M) \otimes_{\mathcal{O}_{\mathcal{F}}} F$$

Thus for any short exact sequence of left R[WH]-modules

 $0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$

applying $F(G/H) \otimes_{R[WH]}$ – is equivalent to applying first the contravariant induction functor and then $\dagger \otimes_{\mathcal{O}_{\mathcal{F}}} F$. Since contravariant induction is exact (Proposition 3.2.5(2)) and F is assumed flat, exactness is preserved, and thus F(G/H) is flat as required.

- (2) Use the argument of the previous part, noting that Lemma 3.1.5 implies that for contravariant frees restricting at G/H preserves finite generation.
- (3) If WH is R-torsion-free then, using Lemma 3.1.5, restricting any free at G/H gives a projective module, and the result follows. To see that in this case, restriction preserves flats, let F be a contravariant flat module and consider a short exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

of left R[WH]-modules, thus by Proposition 3.2.5 below,

$$0 \longrightarrow \operatorname{Ind}_{WH}^{\mathcal{O}_{\mathcal{F}}} A \longrightarrow \operatorname{Ind}_{WH}^{\mathcal{O}_{\mathcal{F}}} B \longrightarrow \operatorname{Ind}_{WH}^{\mathcal{O}_{\mathcal{F}}} C \longrightarrow 0$$

is a short exact sequence of covariant modules. Since F is flat, the functor $\dagger \otimes_{\mathcal{O}_F} F$ is exact, applying this to the above and using Lemma 2.3.3 gives a short exact sequence

$$0 \longrightarrow A \otimes_R F(G/H) \longrightarrow B \otimes_R F(G/H) \longrightarrow C \otimes_R F(G/H) \longrightarrow 0$$

showing F(G/H) is flat.

If WH is not R-torsion free then the result is just Corollary 3.1.6.

3. BREDON MODULES

EXAMPLE 3.2.3. Unlike in the contravariant case, the covariant restriction functor does not preserve "finitely generated" in general: Take for example the infinite dihedral group $D_{\infty} = (\mathbb{Z}/2\mathbb{Z}) * (\mathbb{Z}/2\mathbb{Z})$ generated by the two elements aand b of order 2. The finite subgroup $\langle a \rangle$ is self-normalising, thus $R[W\langle a \rangle] = R$ and Lemma 3.1.3 implies that as R-modules,

$$R[D_{\infty}/1,D_{\infty}/\langle a
angle]_{\mathcal{O}_{\mathcal{F}\!in}} = igoplus_{g\langle a
angle\in D_{\infty}/\langle a
angle} R.$$

REMARK 3.2.4. The covariant restriction functor $\operatorname{Res}_{G}^{\mathcal{O}_{\mathcal{F}}}$ preserves "finitely generated". Recall that

$$R[G/K, G/1]_{\mathcal{O}_{\mathcal{F}}} \cong \begin{cases} RG & \text{if } K = 1\\ 0 & \text{else.} \end{cases}$$

So if A is an arbitrary finitely generated covariant $\mathcal{O}_{\mathcal{F}}$ -module and F a free covariant $\mathcal{O}_{\mathcal{F}}$ -module with an epimorphism onto A then F(G/1) is finitely generated as an RG-module and since $\operatorname{Res}_{RG}^{\mathcal{O}_{\mathcal{F}}}$ is exact there is a surjection $F(G/1) \longrightarrow A(G/1)$.

Recall from Proposition 2.3.2 that contravariant and covariant induction both preserve projectives, flats and finitely generation. In addition we have the following facts.

PROPOSITION 3.2.5.

- (1) If WH has no R-torsion the covariant induction functor $\operatorname{Ind}_{WH}^{\mathcal{O}_{\mathcal{F}}}$ is exact.
- (2) Contravariant induction is always exact.

PROOF. (1) Assume that WH has no R-torsion, we must check that the functor

$$A\longmapsto A\otimes_{R[WH]} R[G/H, -]_{\mathcal{O}_{\mathcal{F}}}$$

is exact, where A is an R[WH]-module. Equivalently that for any subgroup K in \mathcal{F} , the functor

$$-\otimes_{R[WH]} R[G/H,G/K]_{\mathcal{O}_{\mathcal{F}}}$$

is exact, but by Lemma 3.1.5

$$R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}} = \bigoplus_{x \in I} R\left[WH/WH_x\right]$$

for some finite indexing set I and WH_x finite subgroups of WH. By Maschke's Theorem, $R[WH/WH_x]$ is projective, and hence flat, as an R[WH]-module. Hence $- \otimes_{R[WH]} R[G/H, G/K]_{\mathcal{O}_F}$ is indeed exact.

(2) Similarly to the above, we must check the functor

$$R[G/K, G/H]_{\mathcal{O}_{\mathcal{F}}} \otimes_{R[WH]} -$$

is exact, but by Lemma 3.1.3, $R[G/K, G/H]_{\mathcal{O}_{\mathcal{F}}}$ is free as an R[WH]-module, so this is automatic.

3.3. Bredon homology and cohomology of spaces

Recall that a space X is a G-CW complex [tD87, §II.1] if there exists a filtration $\{X_i\}_{i\in\mathbb{Z}}$ of X such that

- (1) X has the colimit topology with respect to the filtration.
- (2) $X_{-1} = \emptyset$.
- (3) X_n is obtained from X_{n-1} via a pushout of G-spaces:



For any $j \in \Delta_n$ the image of $G/H_j \times D^n$ in X is called an *equivariant n-cell* with isotropy H_j . We say X has *isotropy in* \mathcal{F} if the subgroups H_j are elements of \mathcal{F} . For example a G-CW complex is proper if and only if it has isotropy in \mathcal{F} in and is free if and only if it has isotropy in $\mathcal{T}riv$ (the family consisting of only the trivial subgroup).

REMARK 3.3.1. [tD87, II.(1.15)] If X is a CW-complex with a G-action such that

- (1) For all $g \in G$, the map $x \mapsto gx$ takes cells to cells.
- (2) If $g \in G$ fixes a cell σ setwise then g fixes σ pointwise.

Then X is a G-CW-complex. Such an action is often called cellular or rigid.

X is finite-dimensional if $X = X_n$ for some integer n and the minimal such n is called the dimension, and X is finite-type if for all n, X_n is obtained from X_{n-1} by attaching finitely many equivariant n-cells (i.e. the set Δ_n is finite). X is finite (equivalently cocompact) if it is both finite-dimensional and finite-type.

Let \mathcal{F} be a family of subgroups such that X has isotropy in \mathcal{F} . Define the contravariant $\mathcal{O}_{\mathcal{F}}$ -module

$$C_n^{\mathcal{O}_{\mathcal{F}}}(X) = \bigoplus_{j \in \Delta_n} \mathbb{Z}[-, G/H_j]_{\mathcal{O}_{\mathcal{F}}}.$$

Denoting by $C_n(X)$ the ordinary cellular chain complex of X (see for example [Hat02, p.139])

$$C_n^{\mathcal{O}_F}(X)(G/K) = C_n(X^K).$$

The boundary maps from the cellular chain complexes $C_n(X^K)$ give boundary maps for $C_n^{\mathcal{O}_F}(X)$, including an augmentation map,

$$\varepsilon: C_0^{\mathcal{O}_{\mathcal{F}}}(X) \longrightarrow \underline{\mathbb{Z}},$$

which maps every 0-cell in $C_0(X^K)$ to $1 \in \mathbb{Z}(G/K) \cong \mathbb{Z}$. We obtain a free chain complex of contravariant $\mathcal{O}_{\mathcal{F}}$ -modules

$$\cdots \longrightarrow C_n^{\mathcal{O}_{\mathcal{F}}}(X) \longrightarrow C_{n-1}^{\mathcal{O}_{\mathcal{F}}}(X) \longrightarrow \cdots \longrightarrow C_0^{\mathcal{O}_{\mathcal{F}}}(X) \longrightarrow \underline{\mathbb{Z}} \longrightarrow 0.$$

The Bredon homology of X with coefficients in some covariant module A is

$$H^{\mathcal{O}_{\mathcal{F}}}_*(X,A) = H_*(C^{\mathcal{O}_{\mathcal{F}}}_*(X) \otimes_{\mathcal{O}_{\mathcal{F}}} A)$$

and similarly the *Bredon cohomology of* X with coefficients in some contravariant module M is

$$H^*_{\mathcal{O}_{\mathcal{F}}}(X, M) = H^* \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(C^{\mathcal{O}_{\mathcal{F}}}_*(X), M).$$

3.4. Homology and cohomology of groups

Recall the definitions of $\operatorname{Tor}_*^{\mathcal{O}_{\mathcal{F}}}$ and $\operatorname{Ext}_{\mathcal{O}_{\mathcal{F}}}^*$ from Section 2.4. For a group G, covariant module A, and contravariant module M we define

$$H^*_{\mathcal{O}_{\mathcal{F}}}(G, M) = \operatorname{Ext}^*_{\mathcal{O}_{\mathcal{F}}}(\underline{R}, M)$$
$$H^{\mathcal{O}_{\mathcal{F}}}_{*}(G, A) = \operatorname{Tor}^{\mathcal{O}_{\mathcal{F}}}_{*}(\underline{R}, A).$$

Note that in both statements above, \underline{R} denotes the contravariant constant functor on R.

If X is a model for $E_{\mathcal{F}}G$ then the chain complex $C^{\mathcal{O}_{\mathcal{F}}}_*$ is exact, since evaluating at G/H for any $H \in \mathcal{F}$ gives the cellular chain complex of the contractible space X^H . Thus there are isomorphisms for any covariant $\mathcal{O}_{\mathcal{F}}$ -module A and contravariant module M:

$$H^*_{\mathcal{O}_{\mathcal{F}}}(G,M) = H^*_{\mathcal{O}_{\mathcal{F}}}(X,M)$$
$$H^{\mathcal{O}_{\mathcal{F}}}_*(G,A) = H^{\mathcal{O}_{\mathcal{F}}}_*(X,A).$$

3.5. Cohomological dimension

Recall from Section 2.5 that the projective dimension of an $\mathcal{O}_{\mathcal{F}}$ -module M, denoted $\mathcal{O}_{\mathcal{F}}$ pd M, is the minimal length of a projective $\mathcal{O}_{\mathcal{F}}$ -module resolution of M. We say that G has Bredon cohomological dimension n, written $\mathcal{O}_{\mathcal{F}}$ cd G = n, if $\mathcal{O}_{\mathcal{F}}$ pd $\underline{R} = n$ where \underline{R} is the constant contravariant $\mathcal{O}_{\mathcal{F}}$ -module. If we want to emphasize the ring R we will write $\mathcal{O}_{\mathcal{F}}$ cd $_R$ instead of $\mathcal{O}_{\mathcal{F}}$ cd.

As mentioned in the previous section, if X is a model for $E_{\mathcal{F}}G$ then $C^{\mathcal{O}_{\mathcal{F}}}_*(X)$ is an exact resolution of $\underline{\mathbb{Z}}$ by free $\mathcal{O}_{\mathcal{F}}$ -modules, hence $\mathcal{O}_{\mathcal{F}}cd_{\mathbb{Z}}G \leq gd_{\mathcal{F}}G$ (recall $gd_{\mathcal{F}}$ is the minimal dimension of a model for $E_{\mathcal{F}}G$). A theorem of Lück and Meintrup in the high dimensional case and of Dunwoody in the dimension 1 case provides an inequality in the other direction when $\mathcal{F} = \mathcal{F}in$.

THEOREM 3.5.1. [LM00, Theorem 0.1][Dun79] Except for the possibility that $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G = 2$ and $\operatorname{gd}_{\text{fin}} G = 3$, $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G = \operatorname{gd}_{\text{fin}} G$.

In [**BLN01**], Brady, Leary and Nucinkis construct groups G with $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G = 2$, but $\operatorname{gd}_{\text{fin}} G = 3$, showing the bound is sharp.

3.5.1. Low dimensions. Recall that $cd_R G = 0$ if and only if G is finite with no *R*-torsion [**Bie81**, Proposition 4.12].

PROPOSITION 3.5.2. $\mathcal{O}_{\mathcal{F}} \mathrm{cd}_R G = 0$ if and only if there exists a subgroup $H \in \mathcal{F}$ with |G/H| invertible in R and every $K \in \mathcal{F}$ is subconjugate to H. In particular, $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_R G = 0$ if and only if G is finite and and $\mathcal{O}_{\mathcal{V}Cyc} \mathrm{cd}_R G = 0$ if and only if G is virtually cyclic.

A proof of this when $R = \mathbb{Z}$ is available in [Flu10, Prop 3.20], there are some minor modifications needed to generalise to arbitrary rings R.

PROOF. Using a more general definition of family of subgroups \mathcal{F} than we use here, Symonds proves that <u>R</u> is projective if and only if every component of \mathcal{F} has a unique maximal element M and $|N_G M : M|$ is finite and invertible in R, where he views \mathcal{F} as a poset with inclusion [**Sym10**, Lemma 2.5]. Since we assume \mathcal{F} is closed under intersection, for us \mathcal{F} may have only one component. Also, since we assume \mathcal{F} is closed under conjugation we must have $N_G M = G$ —if $g \in G \setminus N_G M$ then since M is maximal $M^g \leq M$ and thus $M \leq M^{g^{-1}}$ contradicting maximality of M. The proposition now follows immediately from Symonds' result and the fact that $\mathcal{O}_{\mathcal{F}}$ cd G = 0 if and only if <u>R</u> is projective.

Combining [**Bie81**, Proposition 4.12] and Proposition 3.5.2, $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G = 0$ if and only if $\operatorname{cd}_{\mathbb{Q}} G = 0$ if and only if G is finite.

Recall that $\operatorname{cd}_{\mathbb{Z}} G = 1$ if and only if G is a free group [Sta68, Swa69], $\operatorname{cd}_R G = 1$ if and only if G is R-torsion-free and acts properly on a tree, and $\operatorname{cd}_{\mathbb{Q}} G = 1$ if and only if G acts properly on a tree or equivalently G is virtuallyfree [Dun79].

LEMMA 3.5.3. For any group G, $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G = 1$ if and only if $\operatorname{cd}_{\mathbb{Q}} G = 1$.

PROOF. If $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_{\mathbb{Z}} G = 1$ then Lemma 3.7.1 implies $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_{\mathbb{Q}} G \leq 1$ and Lemma 3.7.2 implies $\operatorname{cd}_{\mathbb{Q}} G \leq 1$. Since G is not finite, $\operatorname{cd}_{\mathbb{Q}} G = 1$.

If $\operatorname{cd}_{\mathbb{Q}} G = 1$ then by [**Dun79**, Theorem 1.1], G acts properly and with finite stabilisers on a tree T. For any finite subgroup $H \leq G$, H acts on T, $T^H \neq \emptyset$ and in particular T^H is a sub-tree of T [**Ser03**, 6.1, 6.3.1]. T is thus a model for $\operatorname{E}_{\operatorname{fin}} G$ and $\mathcal{O}_{\operatorname{F}}\operatorname{cd}_{\mathbb{Z}} G = 1$.

COROLLARY 3.5.4. The following are equivalent for an infinite group G, and any ring R:

- (1) $\operatorname{cd}_R G = 1$.
- (2) G has no R-torsion and $\mathcal{O}_{\text{Fin}} \operatorname{cd}_{\mathbb{Z}} G = 1$.
- (3) G has no R-torsion and $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G = 1$.
- PROOF. $1 \Rightarrow 2$ If $\operatorname{cd}_R G = 1$ then G has no R-torsion [**Bie81**, Proposition 4.11] and G acts properly on a tree [**Dun79**, Theorem 1.1]. By the argument of Lemma 3.5.3 the tree is a model for $\operatorname{E}_{\operatorname{fin}} G$ and hence $\mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Z}} G = 1$.
- $2 \Rightarrow 3$ Lemma 3.7.1.
- $3 \Rightarrow 1$ Lemma 3.7.2.

QUESTION 3.5.5. What does the condition $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_R G = 1$ represent? Is it equivalent to $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_{\mathbb{Z}} G = 1$?

3.6. FP_n conditions

Recall from Section 2.5 that an $\mathcal{O}_{\mathcal{F}}$ -module M is $\mathcal{O}_{\mathcal{F}} FP_n$ if there is a resolution of M by projective $\mathcal{O}_{\mathcal{F}}$ -modules, finitely generated up to dimension n. We say G is $\mathcal{O}_{\mathcal{F}} FP_n$ if \underline{R} is $\mathcal{O}_{\mathcal{F}} FP_n$, if G is $\mathcal{O}_{\mathcal{F}} FP_\infty$ with finite Bredon cohomological dimension then we say G is $\mathcal{O}_{\mathcal{F}} FP$.

If G admits a model X for $E_{\mathcal{F}}G$ with cocompact *n*-skeleton then the chain complex $C^{\mathcal{O}_{\mathcal{F}}}_{*}(X)$ is finitely generated up to dimension *n* and so G is of type $\mathcal{O}_{\mathcal{F}}FP_n$ over Z. Conversely, if G is $\mathcal{O}_{\mathcal{F}}FP_n$ over Z and WH is finitely presented for every finite subgroup then G has a model for $E_{\mathcal{F}}G$ with cocompact *n*-skeleton [**LM00**, Theorem 0.1].

PROPOSITION 3.6.1. G is $\mathcal{O}_{\mathcal{F}} FP_0$ over R if and only if there exists a finite set $H_1, \ldots, H_m \in \mathcal{F}$ such that every $K \in \mathcal{F}$ is subconjugate to some H_i . In particular, if $\mathcal{F} \subseteq \mathcal{F}$ in, G is $\mathcal{O}_{\mathcal{F}} FP_0$ over R if and only if there are finitely many conjugacy classes of subgroups in \mathcal{F} .

The case $\mathcal{F} = \mathcal{F}$ in appears in [KMPN09, Lemma 3.1].

PROOF. If G is $\mathcal{O}_{\mathcal{F}} FP_0$ then there is an epimorphism,

$$\bigoplus_{i=1}^m R[-,G/H_i]_{\mathcal{O}_{\mathcal{F}}} \longrightarrow \underline{R},$$

where the indexing set I is finite. Let K be a subgroup in \mathcal{F} , evaluating at G/K gives a surjection

$$\bigoplus_{i=1}^m R[G/K, G/H_i]_{\mathcal{O}_{\mathcal{F}}} \longrightarrow R,$$

40

so for some *i* we have $R[G/K, G/H_i]_{\mathcal{O}_F} \neq 0$ and hence *K* is subconjugate to one of the H_i .

For the converse, one checks that the augmentation map

$$\bigoplus_{i=1}^m R[-, G/H_i]_{\mathcal{O}_{\mathcal{F}}} \longrightarrow \underline{R}$$

is a surjection.

If $\mathcal{F} \subseteq \mathcal{F}$ in then observe that each H_i has at most finitely many subconjugate subgroups, so the existence of such a collection H_1, \ldots, H_m is equivalent to \mathcal{F} having finitely many conjugacy classes.

PROPOSITION 3.6.2. Let G be $\mathcal{O}_{\mathcal{F}} FP_0$ and $\mathcal{F} \subseteq \mathcal{F}in$, then a contravariant module M is $\mathcal{O}_{\mathcal{F}} FP_n$ $(n \geq 1)$ over R if and only if M(G/K) is of type FP_n over R[WK] for all subgroups K in \mathcal{F} .

The proof in the case $R = \mathbb{Z}$ appears as [**KMPN09**, Lemma 3.2] and requires no substantial alteration to generalise to arbitrary rings R.

QUESTION 3.6.3. Is there an easy characterisation of the condition $\mathcal{O}_{\mathcal{F}} FP_n$ for arbitrary \mathcal{F} , or for $\mathcal{F} = \mathcal{VCyc}$?

COROLLARY 3.6.4. The following are equivalent for a group G and $\mathcal{F} \subseteq \mathcal{F}$ in,

- (1) G is $\mathcal{O}_{\mathcal{F}} FP_n$ over R.
- (2) G is $\mathcal{O}_{\mathcal{F}} FP_0$ and the Weyl groups WK are FP_n over R for all $K \in \mathcal{F}$.
- (3) G is $\mathcal{O}_{\mathcal{F}} FP_0$ and the centralisers $C_G K$ are FP_n over R for all $K \in \mathcal{F}$.

PROOF. By the previous Proposition (1) and (2) are equivalent. To see the equivalence of (2) and (3) consider the short exact sequence

$$0 \longrightarrow K \longrightarrow N_G K \longrightarrow W K \longrightarrow 0.$$

K is finite and hence FP_{∞} , so WK is FP_n over R if and only if N_GK is FP_n over R [**Bie81**, Proposition 2.7]. Since K is finite, so C_GK is finite index in N_GK [**Rob96**, 1.6.13] and so C_GK is FP_n over R if and only if N_GK is FP_n over R. Combining the last two results gives WK is FP_n over R if and only if C_GK is FP_n over R.

EXAMPLE 3.6.5. In [**BS80**], it's shown that Abels' group is FP₂ over \mathbb{Q} but not over \mathbb{Z} . The Bestvina Brady groups also provide examples of groups which are FP_n over some rings but not others [**BB97**].

3.6.1. Quasi- $\mathcal{O}_{\mathcal{F}} FP_n$ conditions. In [MPN13, §6], Martínez-Pérez and Nucinkis define the quasi- $\mathcal{O}_{\mathcal{F}} FP_n$ condition, a weakening of $\mathcal{O}_{\mathcal{F}} FP_n$, these are defined for all families $\mathcal{F} \subseteq \mathcal{F}$ in. We will need these conditions in Chapter 7. A

group G is quasi- $\mathcal{O}_{\mathcal{F}}FP_n$ if WK is FP_n for all $K \in \mathcal{F}$ and G has finitely many conjugacy classes of subgroups in \mathcal{F} isomorphic to a given finite subgroup.

For any positive integer k, define the module

$$\underline{R}_k(G/H) = \begin{cases} R & \text{if } |H| \le k, \\ 0 & \text{otherwise.} \end{cases}$$

Then G is quasi- $\mathcal{O}_{\mathcal{F}} FP_n$ if and only if \underline{R}_k is $\mathcal{O}_{\mathcal{F}} FP_n$ for all positive integers k [**MPN13**, Proposition 6.5].

Say G is quasi- $\mathcal{O}_{\mathcal{F}}\mathbf{F}_n$ if for all positive integers k, G has a finite type model for $E_{\mathcal{F}_k}G$, where \mathcal{F}_k is the subfamily of \mathcal{F} containing all subgroups of order less than k. If G is quasi- $\mathcal{O}_{\mathcal{F}}\mathbf{F}\mathbf{P}_n$ then G is quasi- $\mathcal{O}_{\mathcal{F}}\mathbf{F}_n$ if and only if the centralisers $C_G K$ are finitely presented for all $K \in \mathcal{F}$ [**MPN13**, Proposition 6.10].

We can give a geometric meaning to the quasi- $\mathcal{O}_{\mathcal{F}}\mathbf{F}_n$ conditions—a group G is quasi- $\mathcal{O}_{\mathcal{F}}\mathbf{F}_n$ if and only if G admits a model for $\mathbf{E}_{\mathcal{F}}G$ which is a mapping telescope of models for $E_{\mathcal{F}_k}G$ with cocompact *n*-skeleta [**MPN13**, Theorem 6.11].

3.7. Change of rings

If $\varphi : R_1 \to R_2$ is a ring homomorphism then we define the change of rings functor φ^* from $\mathcal{O}_{\mathcal{F}}$ -modules over R_2 to $\mathcal{O}_{\mathcal{F}}$ -modules over R_1 as follows,

$$\varphi^*A: G/H \mapsto A(G/H),$$

where we are viewing A(G/H) as an R_1 -module via φ . On morphisms:

$$\varphi^* A\left(\sum_i r_i \alpha_i\right) = \sum_i \varphi(r_i) A(\alpha_i)$$

where $r_i \in R_1$ and the α_i are morphisms $G/H \to G/K$ for some $G/H, G/K \in \mathcal{O}_F$.

We also define a functor $R_2 \otimes_{R_1}$ – from $\mathcal{O}_{\mathcal{F}}$ -modules over R_1 to $\mathcal{O}_{\mathcal{F}}$ -modules over R_2 by

$$R_2 \otimes_{R_1} A : G/H \mapsto R_2 \otimes_{R_1} A(G/H)$$

where we are using φ to view R_2 as an R_1 -module. Applying this to a free module gives

$$R_2 \otimes_{R_1} R_1[-, G/H]_{\mathcal{O}_{\mathcal{F}}} \cong R_2[-, G/H]_{\mathcal{O}_{\mathcal{F}}}.$$

Hence if P is a projective $\mathcal{O}_{\mathcal{F}}$ -module over R_1 then $R_2 \otimes_{R_1} P$ is a projective $\mathcal{O}_{\mathcal{F}}$ -module over R_2 .

LEMMA 3.7.1. If $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_{\mathbb{Z}} G \leq n$ then $\mathcal{O}_{\mathcal{F}} \operatorname{cd}_R G \leq n$ for all rings R. Similarly if G is $\mathcal{O}_{\mathcal{F}} \operatorname{FP}_n$ over \mathbb{Z} then G is $\mathcal{O}_{\mathcal{F}} \operatorname{FP}_n$ over R for all rings R.

PROOF. For the first part, take a projective resolution of $\underline{\mathbb{Z}}$ by contravariant $\mathcal{O}_{\mathcal{F}}$ -modules of length n and define a new resolution by $Q_n(G/H) = R \otimes_{\mathbb{Z}} P_n(G/H)$ for all $n \in \mathbb{N}$ and $G/H \in \mathcal{O}_{\mathcal{F}}$. Since for any $H \in \mathcal{F}$ the complex $P_*(G/H)$ is \mathbb{Z} -acyclic and hence \mathbb{Z} -split so $Q_*(G/H)$ is acyclic also. Finally each Q_n is projective.

The second part is similar—choose the projective $\mathcal{O}_{\mathcal{F}}$ -module resolution of $\underline{\mathbb{Z}}$ to be finitely generated in all degrees $i \leq n$ and use that if P_i is finitely generated then so is Q_i .

LEMMA 3.7.2. If G has no R-torsion then $\operatorname{cd}_R G \leq \mathcal{O}_{\mathcal{F}}\operatorname{cd}_R G$.

PROOF. Take a projective resolution of <u>R</u> by contravariant modules of length n and evaluate at G/1, since G is R-torsion-free, Proposition 3.2.2(3) implies that $P_*(G/1)$ is a length n projective resolution of R by RG-modules.

PROPOSITION 3.7.3. If $\varphi : R_1 \to R_2$ is a ring homomorphism and A is a $\mathcal{O}_{\mathcal{F}}$ -module over R_2 then

$$\operatorname{Tor}_{*}^{R_{1},\mathcal{O}_{\mathcal{F}}}(\underline{R}_{1},\varphi^{*}A) \cong \operatorname{Tor}_{*}^{R_{2},\mathcal{O}_{\mathcal{F}}}(\underline{R}_{2},A).$$

There are similar isomorphisms for contravariant modules and for $\operatorname{Ext}^*_{\mathcal{O}_{\mathcal{F}}}$.

PROOF. Firstly, consider the case $\varphi : \mathbb{Z} \to R$ for some ring R, we prove

$$\operatorname{Tor}_{*}^{\mathbb{Z},\mathcal{O}_{\mathcal{F}}}(\underline{\mathbb{Z}},\varphi^{*}A) = \operatorname{Tor}_{*}^{R,\mathcal{O}_{\mathcal{F}}}(\underline{R},A).$$

Choose a resolution P_* of $\underline{\mathbb{Z}}$ by contravariant projective $\mathcal{O}_{\mathcal{F}}$ -modules over \mathbb{Z} . For any G/H in $\mathcal{O}_{\mathcal{F}}$, $P_*(G/H)$ is a \mathbb{Z} -split resolution, so applying the functor $R \otimes_{\mathbb{Z}} -$ to P_* yields a projective resolution of \underline{R} by projective $\mathcal{O}_{\mathcal{F}}$ -modules over R. Observing that

$$P_* \otimes_{\mathcal{O}_{\mathcal{F}},\mathbb{Z}} \varphi^* A \cong (P_* \otimes_{\mathbb{Z}} R) \otimes_{\mathcal{O}_{\mathcal{F}},R} A$$

completes the proof.

For the general case, let $\varphi_1 : \mathbb{Z} \to R_1$ and $\varphi_2 : \mathbb{Z} \to R_2$ be (unique) ring homomorphisms, then $\varphi \circ \varphi_1 = \varphi_2$ and $\varphi_1^* \circ \varphi^* = \varphi_2^*$. Applying the previous part twice gives

$$\operatorname{Tor}_{*}^{R_{1},\mathcal{O}_{\mathcal{F}}}(\underline{R}_{1},\varphi^{*}A) \cong \operatorname{Tor}_{*}^{\mathbb{Z},\mathcal{O}_{\mathcal{F}}}(\underline{\mathbb{Z}},\varphi_{1}^{*}\circ\varphi^{*}A)$$
$$\cong \operatorname{Tor}_{*}^{R_{2},\mathcal{O}_{\mathcal{F}}}(\underline{R}_{2},A).$$

The next result is essentially [Ham08, 1.4.3], where it is proved for rings of prime characteristic in the setting of ordinary group cohomology.

PROPOSITION 3.7.4. Given some integer m > 0 and ring R with characteristic m, then G is $\mathcal{O}_{\mathcal{F}} FP_n$ over R if and only if G is $\mathcal{O}_{\mathcal{F}} FP_n$ over $\mathbb{Z}/m\mathbb{Z}$.

PROOF. The proof below is for contravariant modules, the proof for covariant modules is analogous.

Assume that G is $\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n$ over $\mathbb{Z}/m\mathbb{Z}$. If M_* is any directed system of contravariant $\mathcal{O}_{\mathcal{F}}$ -modules over R with $\varinjlim M_* = 0$, we necessarily have $\varinjlim \varphi^* M_* =$ 0. By the Bieri–Eckmann criterion (Theorem 2.5.1), and the fact that $\underline{\mathbb{Z}/m\mathbb{Z}}$ is assumed $\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n$ over $\mathbb{Z}/m\mathbb{Z}$, we have that for all $i \leq n$,

$$\varinjlim \operatorname{Ext}^{i}_{\mathcal{O}_{\mathcal{F}}, \mathbb{Z}/m\mathbb{Z}}(\mathbb{Z}/m\mathbb{Z}, \varphi^{*}M_{*}) = 0.$$

Thus by Proposition 3.7.3 applied to the canonical map $\mathbb{Z}/m\mathbb{Z} \to R$,

 $\lim_{\tau} \operatorname{Ext}^{i}_{\mathcal{O}_{\tau},R}(\underline{R}, M_{*}) = 0.$

The Bieri–Eckmann criterion (Theorem 2.5.1) gives that <u>R</u> is $\mathcal{O}_{\mathcal{F}}FP_n$ over R.

For the "only if" direction, suppose M_* is a directed system of $\mathcal{O}_{\mathcal{F}}$ -modules over $\mathbb{Z}/m\mathbb{Z}$, with $\varinjlim M_* = 0$ thus $\varinjlim M_* \otimes_{\mathbb{Z}/m\mathbb{Z}} R = 0$ and by Theorem 2.5.1 for all $i \leq n$,

$$\lim_{i \to \infty} \operatorname{Ext}^{i}_{\mathcal{O}_{\mathcal{F}},R}(\underline{R}, M_{*} \otimes_{\mathbb{Z}/m\mathbb{Z}} R) = 0.$$

Combining with Proposition 3.7.3

$$\varinjlim \operatorname{Ext}_{\mathbb{Z},\mathcal{O}_{\mathcal{F}}}^{i}(\underline{\mathbb{Z}/m\mathbb{Z}}, M_{*} \otimes_{\mathbb{Z}/m\mathbb{Z}} R) = \varinjlim \operatorname{Ext}_{R,\mathcal{O}_{\mathcal{F}}}^{i}(\underline{R}, M_{*} \otimes_{\mathbb{Z}/m\mathbb{Z}} R)$$
$$= 0.$$

Since $\mathbb{Z}/m\mathbb{Z}$ is self-injective [Lam99, Cor 3.13], R splits as a $\mathbb{Z}/m\mathbb{Z}$ module into $R \cong \mathbb{Z}/m\mathbb{Z} \oplus N$ where N is some $\mathbb{Z}/m\mathbb{Z}$ module. Thus we have

$$\varinjlim \left(\operatorname{Ext}^{i}_{\mathbb{Z}/m\mathbb{Z},\mathcal{O}_{\mathcal{F}}}(\underline{\mathbb{Z}/m\mathbb{Z}},M_{*}) \\ \oplus \operatorname{Ext}^{i}_{\mathbb{Z}/m\mathbb{Z},\mathcal{O}_{\mathcal{F}}}(\underline{\mathbb{Z}/m\mathbb{Z}},M_{*}\otimes_{\mathbb{Z}/m\mathbb{Z}}N) \right) = 0$$

In particular

$$\varinjlim \operatorname{Ext}^{i}_{\mathbb{Z}/m\mathbb{Z},\mathcal{O}_{\mathcal{F}}}(\underline{\mathbb{Z}/m\mathbb{Z}},M_{*}) = 0$$

so by the Bieri–Eckmann criterion (Theorem 2.5.1) $\underline{\mathbb{Z}/m\mathbb{Z}}$ is $\mathcal{O}_{\mathcal{F}}FP_n$ over $\mathbb{Z}/m\mathbb{Z}$, i.e. G is $\mathcal{O}_{\mathcal{F}}FP_n$ over $\mathbb{Z}/m\mathbb{Z}$.

REMARK 3.7.5. This proposition fails in characteristic zero as the ring \mathbb{Z} is not self-injective. For example \mathbb{Q} is not isomorphic, as a \mathbb{Z} -module, to $N \oplus \mathbb{Z}$ for any \mathbb{Z} -module N.

3.8. Some interesting examples

By the right-angled Coxeter group (W, S) corresponding to some flag complex L we mean the group W generated by a set S of involutions where S is in bijection with the vertices of L and two involutions commute if and only if they are adjacent in L. Given such a (W, S) we let S be the poset of spherical subsets of S (subsets

generating a finite subgroup of W) and form the geometric realisations $K = |\mathcal{S}|$ and $\partial K = |\mathcal{S}_{>\emptyset}|$.

We form the simplicial complexes $\mathcal{U}(W, \partial K)$ and $\mathcal{U}(W, K)$ as in [**Dav08**, 5.1.2], both admit W-actions and $\mathcal{U}(W, \partial K)$ is the singular set of $\mathcal{U}(W, K)$ (subcomplex with non-trivial isotropy). The complex $\mathcal{U}(W, K)$, often called the Davis complex, is known to be a model for $\mathbb{E}_{\text{fin}}W$ [**Dav08**, Theorem 12.3.4(ii)].

LEMMA 3.8.1. [Dav08, 8.2.8] $\mathcal{U}(W, \partial K)$ is *R*-acyclic if and only if $(\partial K)_T$ is *R*-acyclic for all spherical subsets $T \in S$, where

$$K_T = \bigcap_{s \in T} |\mathcal{S}_{\geq s}|.$$

The example below first appeared in [**Bes93**], see also [**Dav08**, 8.5.8], and much of the following argument appears in [**DL98**, proof of Theorem 2].

EXAMPLE 3.8.2 (A group W with $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{F}_3} W = 2$ but $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} W = 3$ which is not torsion-free). Consider the right-angled Coxeter group (W, S) corresponding to the barycentric subdivision L of the ordinary triangulation of \mathbb{RP}^2 .

Claim: $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_{\mathbb{Z}} W = 3$. Since $\mathcal{U}(W, K)$ is a model for $\operatorname{E}_{\mathcal{F}in} W$ and one can calculate that it is 3-dimensional, we conclude $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_{\mathbb{Z}} W \leq 3$. To see that $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_{\mathbb{Z}} W = 3$ we calculate $H^n_{\mathcal{O}_{\mathcal{F}in}}(W, \mathbb{Z}[-, W/1]_{\mathcal{O}_{\mathcal{F}in}})$ as in [**LN03**, p.147], using Lemma 3.8.5 at the end of this section,

$$H^{n}_{\mathcal{O}_{\mathcal{F}in}}(W, \mathbb{Z}[-, W/1]_{\mathcal{O}_{\mathcal{F}in}}) \cong H^{3}_{W}(\mathcal{U}(W, K), \mathcal{U}(W, K)^{\text{sing}}; \mathbb{Z}W)$$
$$\cong H^{3}_{W}(\mathcal{U}(W, K), \mathcal{U}(W, \partial K); \mathbb{Z}W).$$

Recall that $\mathcal{U}(W, K) = W \times K / \sim$ where the identification is only on $W \times \partial K$, that K is a fundamental domain for the W-action on $\mathcal{U}(W, K)$, and that $(K, \partial K) \simeq (\mathcal{C}\mathbb{R}\mathbb{P}^2, \mathbb{R}\mathbb{P}^2)$. Here $\mathcal{C}X$ denotes the cone on a space X. The action of W on $C_*(\mathcal{U}(W, K), \mathcal{U}(W, \partial K))$ is free, so

$$H^*_W(\mathcal{U}(W,K),\mathcal{U}(W,\partial K);\mathbb{Z}W) \cong H^*(K,\partial K;\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Z}W$$
$$\cong H^*(\mathcal{C}\mathbb{R}\mathbb{P}^2,\mathbb{R}\mathbb{P}^2;\mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Z}W.$$

In particular, in dimension 3,

$$H^{3}(\mathcal{C}\mathbb{RP}^{2},\mathbb{RP}^{2};\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{Z}W\cong H^{2}(\mathbb{RP}^{2};\mathbb{Z})\otimes_{\mathbb{Z}}\mathbb{Z}W\cong\mathbb{F}_{2}W.$$

We conclude $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_{\mathbb{Z}} W = 3$.

Claim: $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{F}_3} W = 2$. $\mathcal{U}(W, \partial K)$ is the singular set of $\mathcal{U}(W, K)$, so in particular the fixed point sets of finite subgroups (except for the trivial subgroup) agree. They are contractible and hence \mathbb{F}_3 -acyclic. We claim $\mathcal{U}(W, \partial K)$ is also \mathbb{F}_3 acyclic. We use Lemma 3.8.1—if $T \neq \emptyset$ then $(\partial K)_T = K_T$ which is contractible and hence \mathbb{F}_3 -acyclic and if $T = \emptyset$ then $(\partial K)_T = \partial K$ which is the barycentric subdivision of $L = \mathbb{RP}^2$ and hence \mathbb{F}_3 -acyclic. Taking the Bredon chain complex

$$P_* = C^{\mathcal{O}_{\mathcal{F}_{in}}}_*(\mathcal{U}(W, \partial K)) \otimes_{\mathbb{Z}} \mathbb{F}_3$$

gives that $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{F}_3} W \leq 2.$

W is a right-angled Coxeter group so every finite subgroup of W has order a power of 2, in particular W has no \mathbb{F}_3 -torsion. By Corollary 3.5.4, $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{F}_3} W = 1$ if and only if $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{Z}} W = 1$ but we have already shown that $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{Z}} W = 3$ proving that $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{F}_3} W \neq 1$ and in fact $\mathcal{O}_{\mathcal{F}in} \mathrm{cd}_{\mathbb{F}_3} W = 2$.

EXAMPLE 3.8.3 (A group with $\operatorname{cd}_{\mathbb{Q}} G \neq \mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Q}} G$). In [**LN03**], Leary and Nucinkis construct examples of groups with $\operatorname{vcd}_{\mathbb{Z}} G = nm$ and $\mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Z}} G = m(n+1)$ for various integers n and m. We show that these groups have $\mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Q}} G = m(n+1)$ as well, so since $\operatorname{cd}_{\mathbb{Q}} G \leq \operatorname{vcd}_{\mathbb{Z}} G$ this provides examples of groups with $\operatorname{cd}_{\mathbb{Q}} G \neq \mathcal{O}_{\operatorname{fin}}\operatorname{cd}_{\mathbb{Q}} G$.

We prove that groups G satisfying the assumptions of [**LN03**, Theorem 6] satisfy $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Q}} G \geq m(n+1)$ also, since combining this with the inequality $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Q}} G \leq \mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G$ gives $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Q}} G = m(n+1)$ as required.

Leary and Nucinkis show there exists a model X for $E_{\mathcal{F}in}G$ such that the cellular chain complex $C_*(X^{m(n+1)}, (X^{m(n+1)})^{\text{sing}})$ contains a copy of $\mathbb{Z}G$ in dimension m(n+1) as a direct summand. Here X^i denotes the *i* skeleton of some CW-complex X. Using Lemma 3.8.5 below,

$$H^{m(n+1)}_{\mathcal{O}_{\mathcal{F}in}}(G, \mathbb{Q}[-G/1]_{\mathcal{O}_{\mathcal{F}in}}) \cong H^{m(n+1)}_{G}(C_*(X, X^{\operatorname{sing}}); \mathbb{Q}G)$$
$$\cong H^{m(n+1)}_{G}(C_*(X^{m(n+1)}, (X^{m(n+1)})^{\operatorname{sing}}); \mathbb{Q}G)$$
$$\neq 0$$

showing $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_{\mathbb{Q}} G \ge m(n+1)$.

The examples constructed with the method above can never be of type $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$ [LN03, Question 2, p.154], so a natural question is:

QUESTION 3.8.4. Are there groups G with $\operatorname{cd}_{\mathbb{Q}} G \neq \mathcal{O}_{\operatorname{fin}} \operatorname{cd}_{\mathbb{Q}} G$ and type $\mathcal{O}_{\operatorname{fin}} \operatorname{FP}_{\infty}$?

LEMMA 3.8.5. For any group G and model X for $E_{\text{Fin}}G$

$$H^*_{\mathcal{O}_{\text{fin}}}(G, R[-, G/1]_{\mathcal{O}_{\text{fin}}}) \cong H^*_G(C_*(X, X^{\text{sing}}); RG)$$

where $C_*(X, X^{sing})$ denotes the cellular chain complex of the pair (X, X^{sing}) .

PROOF. Firstly,

$$H^*\left(\operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}\left(C^{\mathcal{O}_{\operatorname{fin}}}_*(X^{\operatorname{sing}}), R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}}\right)\right) = 0$$

since the G-orbits of cells in X^{sing} all give rise to contravariant modules of the form $R[-, G/H]_{\mathcal{O}_{\text{fin}}}$ for $H \neq 1$, and by the Yoneda-type Lemma 2.0.2,

$$\operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}(R[-,G/H]_{\mathcal{O}_{\operatorname{fin}}},R[-,G/1]_{\mathcal{O}_{\operatorname{fin}}})\cong R[G/H,G/1]_{\mathcal{O}_{\operatorname{fin}}}=0.$$

Using the long exact sequence in homology associated to the pair (X, X^{sing}) ,

$$\begin{aligned} H^*_{\mathcal{O}_{\mathcal{F}in}}(G, R[-, G/1]_{\mathcal{O}_{\mathcal{F}in}}) &\cong H^* \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}in}}(C^{\mathcal{O}_{\mathcal{F}in}}_*(X), R[-, G/1]_{\mathcal{O}_{\mathcal{F}in}}) \\ &\cong H^* \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}in}}(C^{\mathcal{O}_{\mathcal{F}in}}_*(X, X^{\operatorname{sing}}), R[-, G/1]_{\mathcal{O}_{\mathcal{F}in}}). \end{aligned}$$

Via the Yoneda-type Lemma 2.0.2, there is a chain of natural isomorphisms:

$$\operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}(C^{\mathcal{O}_{\operatorname{fin}}}_{*}(X, X^{\operatorname{sing}}), R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}})$$

$$\cong \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}\left(\bigoplus_{\substack{G\text{-orbits of }i\text{-cells}\\ \text{with trivial isotropy}}} R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}}, R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}}\right)$$

$$\cong \prod \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}\left(R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}}, R[-, G/1]_{\mathcal{O}_{\operatorname{fin}}}\right)$$

$$\cong \prod \operatorname{Hom}_{RG}(RG, RG)$$

$$\cong \operatorname{Hom}_{RG}\left(\bigoplus RG, RG\right)$$

$$\cong \operatorname{Hom}_{RG}(C^{\mathcal{O}_{\operatorname{fin}}}_{*}(X, X^{\operatorname{sing}}), RG).$$

Thus,

$$H^* \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}(C^{\mathcal{O}_{\operatorname{fin}}}_*(X, X^{\operatorname{sing}}) \cong H^* \operatorname{Hom}_{RG}(C_*(X, X^{\operatorname{sing}}), RG),$$

and combining this with the isomorphism (\star) completes the proof.

3.9. Finitely generated projectives and duality

In this section we require $\mathcal{F} \subseteq \mathcal{F}in$. This section contains a number of technical results concerning dual $\mathcal{O}_{\mathcal{F}}$ -modules, they are all analogs of results for modules over group rings that can be found in [**Bie81**]. The results in this section are built on in Section 6.7 and utilised in Section 4.3.

For M a contravariant module, denote by M^D the dual module

$$M^{D} = \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}}).$$

Similarly for A a covariant module,

$$A^{D} = \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}} \left(A(-), R[?, -]_{\mathcal{O}_{\mathcal{F}}} \right).$$

This definition should be compared with that of the dual of an RG-module M, namely $M^D = \operatorname{Hom}_{RG}(M, RG)$ [**Bie81**, §3.1].

3. BREDON MODULES

EXAMPLE 3.9.1. If G is an infinite group and <u>R</u> is the covariant constant functor on R then $\underline{R}^D = 0$, as

$$\underline{R}^{D} = \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(\underline{R}(?), R[-, ?]_{\mathcal{O}_{\mathcal{F}}})$$
$$\cong \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(\operatorname{Ind}_{G}^{\mathcal{O}_{\mathcal{F}}} R(?), R[-, ?]_{\mathcal{O}_{\mathcal{F}}})$$
$$\cong \operatorname{Hom}_{RG}(R, R[-, G/1]_{\mathcal{O}_{\mathcal{F}}}),$$

using Example 3.2.1 and the adjointness of induction and restriction. Finally, $\operatorname{Hom}_{RG}(R, R[-, G/1]_{\mathcal{O}_{\mathcal{T}}})$ is the zero module since G is infinite.

LEMMA 3.9.2. The dual functor takes projectives to projectives and the doubledual functor $-^{DD}$: { $\mathcal{O}_{\mathcal{F}}$ -modules} \rightarrow { $\mathcal{O}_{\mathcal{F}}$ -modules} is a natural isomorphism when restricted to the subcategory of finitely generated projective $\mathcal{O}_{\mathcal{F}}$ -modules.

PROOF. By the Yoneda-type Lemma 2.0.2,

$$R[-, G/H]^{D}_{\mathcal{O}_{\mathcal{F}}} \cong \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(R[?, G/H]_{\mathcal{O}_{\mathcal{F}}}, R[?, -]_{\mathcal{O}_{\mathcal{F}}}) \cong R[G/H, -]_{\mathcal{O}_{\mathcal{F}}}.$$

The proof for covariant frees is identical.

For any module M, there is a map $\zeta : M \longrightarrow M^{DD}$, given by $\zeta(m)(f) = f(m)$. If $M = R[-, G/H]_{\mathcal{O}_{\mathcal{F}}}$ then applying the Yoneda-type lemma twice shows $M^{DD} = M$. This generalises to projectives since the duality functor represents direct sums.

Naturality follows from naturality of the map ζ .

For M and N contravariant $\mathcal{O}_{\mathcal{F}}$ -modules, we construct an R-module homomorphism

$$\nu: N \otimes_{\mathcal{O}_{\mathcal{F}}} M^D \longrightarrow \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}} (M, N).$$

The main result of this section will be Lemma 3.9.4, that ν is an isomorphism when M is finitely generated projective and Proposition 3.9.8, that ν induces an isomorphism

$$N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} H^{i}_{\mathcal{O}_{\mathcal{F}}}(G, R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) \cong H^{i}_{\mathcal{O}_{\mathcal{F}}}(G, N)$$

for all $i \leq n$ when G is $\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n$.

Recall that elements of $N \otimes_{\mathcal{O}_{\mathcal{F}}} M^D$ are equivalence classes of finite sums of elements of the form

$$n_H \otimes \varphi_H \in \bigoplus_{G/H \in \mathcal{O}_F} N(G/H) \otimes_R \operatorname{Hom}_{\mathcal{O}_F} (M, R[-, G/H]_{\mathcal{O}_F}).$$

For any $G/L \in \mathcal{O}_{\mathcal{F}}$ and $m \in M(G/L)$ we define

$$\nu (n_H \otimes_R \varphi_H) (G/L) : M(G/L) \longrightarrow N(G/L)$$
$$m \longmapsto N (\varphi_H(G/L)(m)) (n_H).$$

48

This makes sense because $\varphi_H(G/L)(m) \in R[G/L, G/H]_{\mathcal{O}_F}$ and N is a contravariant module so

$$N(\varphi_H(G/L)(m)): N(G/H) \longrightarrow N(G/L).$$

We must check that $\nu(n_H \otimes_R \varphi_H)$ is a natural transformation, it's well defined including that it doesn't depend on the choice of equivalence class in $N(?) \otimes_{\mathcal{O}_F}$ $\operatorname{Hom}_{\mathcal{O}_F}(M(-), R[-, ?]_{\mathcal{O}_F})$, and that it is an *R*-module homomorphism. $\nu(n_H \otimes_R \varphi_H)$ is a natural transformation:

Let $\alpha : G/L_1 \mapsto G/L_2$ be a *G*-map and $G/L_i \in \mathcal{O}_F$, we must check the following diagram commutes.

$$M(G/L_{1}) \xrightarrow{\nu(n_{H} \otimes_{R} \varphi_{H})(G/L_{1}):m \longmapsto N(\varphi_{H}(G/L_{1})(m))(n_{H})} N(G/L_{1})}_{N(\alpha)} M(\alpha) \xrightarrow{\uparrow} N(\alpha) \xrightarrow{\uparrow} N(\alpha) \xrightarrow{\uparrow} N(G/L_{2}):m \longmapsto N(\varphi_{H}(G/L_{2})(m))(n_{H})} N(G/L_{2})$$

$$N(\alpha) \circ \left(\nu(n_H \otimes_R \varphi_H)(G/L_2)\right)(m) = N(\alpha) \circ N \left(\varphi_H(G/L_2)(m)\right)(n_H)$$

= $N\left(\varphi_H(G/L_2)(m) \circ \alpha\right)(n_H)$
= $N\left(\left(R[\alpha, G/H]_{\mathcal{O}_{\mathcal{F}}} \circ \varphi_H(G/L_2)\right)(m)\right)(n_H)$
= $N\left(\left(\varphi_H(G/L_1) \circ M(\alpha)\right)(m)\right)(n_H)$
= $\left(\nu(n_H \otimes_R \varphi_H)(G/L_2) \circ M(\alpha)\right)(m)$

Where the second equality is because N is a contravariant functor, the third is because by definition $\varphi_H(G/L_2)(m) \circ \alpha = (R[\alpha, G/H]_{\mathcal{O}_{\mathcal{F}}} \circ \varphi_H(G/L_2))(m)$, and the fourth is because φ_H is itself a natural transformation and hence the following diagram commutes.

$$(\dagger) \qquad \qquad M(G/L_1) \xrightarrow{\varphi_H(G/L_1)} R[G/L_1, G/H]_{\mathcal{O}_{\mathcal{F}}} \\ M(\alpha) \uparrow \qquad \qquad R[\alpha, G/H]_{\mathcal{O}_{\mathcal{F}}} \uparrow \\ M(G/L_2) \xrightarrow{\varphi_H(G/L_2)} R[G/L_2, G/H]_{\mathcal{O}_{\mathcal{F}}} \end{cases}$$

 ν is well-defined: Firstly,

$$\nu(rn_H \otimes \varphi_H) = \nu(n_H \otimes r\varphi_H)$$

this is because

$$\nu (n_H \cdot r \otimes_R \varphi_H) (G/L)(m) = N (\varphi_H(G/L)(m)) (rn_H)$$
$$= rN (\varphi_H(G/L)(m)) (n_H)$$
$$= N (r\varphi_H(G/L)(m)) (n_H)$$
$$= \nu (n_H \otimes r\varphi_H).$$

Secondly, we show ν doesn't depend on the choice of equivalence class in

$$N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}} (M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}}).$$

Choose $n_H \in N(G/H)$, $\varphi_M \in \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(M(-), R[-, G/M]_{\mathcal{O}_{\mathcal{F}}})$, $\alpha : G/H \to G/M$ a *G*-map and $G/H, G/M \in \mathcal{O}_{\mathcal{F}}$, we must show that

$$\nu(N(\alpha)(n_H) \otimes_R \varphi_M) = \nu\left(n_H \otimes_R \left(\operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(M(-), R[-, \alpha]_{\mathcal{O}_{\mathcal{F}}})\right)(\varphi_M)\right).$$

Let $G/L \in \mathcal{O}_{\mathcal{F}}$, then

$$\nu (N(\alpha)(n_H) \otimes_R \varphi_M) (G/L)(m) = N (\varphi_H(G/L_1)(m)) (N(\alpha)(n_H))$$

= $N (\alpha \circ \varphi_H(G/L)(m)) (n_H)$
= $N (R[G/L, \alpha]_{\mathcal{O}_F} (\varphi_H(G/L_1)(m))) (n_H)$
= $N (\operatorname{Hom}_{\mathcal{O}_F} (M(-), R[-, \alpha]_{\mathcal{O}_F}) (\varphi_H) (G/L_1)(m)) (n_H)$
= $\nu (n_H \otimes_R \operatorname{Hom}_{\mathcal{O}_F} (M(-), R[-, \alpha]_{\mathcal{O}_F}) (\varphi_M)) (G/L)(m).$

 ν is a map of *R*-modules: It's clear that ν is additive, and

$$\nu(rn_H\otimes\varphi_H)=r\nu(n_H\otimes\varphi_H)$$

since N being a module over R implies that $N(\varphi_H(G/L)(m))$ is an R-module homomorphism.

Lemma 3.9.3. ν is natural in N in M.

PROOF. We only prove naturality in N, the proof for M is similar. Let F be a morphism of contravariant modules $N \to N'$, we must show that the following diagram of R-modules commutes.

Let $n_H \otimes \varphi_H \in N(G/H) \otimes_{\mathcal{O}_F} \operatorname{Hom}_{\mathcal{O}_F}(M(-), R[-, G/H]_{\mathcal{O}_F})$ then moving about the top right of the diagram yields

$$(\operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(M(-), F(-)) \circ \nu_N(n_H \otimes \varphi_H))(G/L)(m)$$

= $F(G/L) \circ N(\varphi_H(G/L)(m))(n_H).$

Moving around the bottom left yields

$$(\nu_{N'} \circ F(?) \otimes \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}})(n_{H} \otimes \varphi_{H}))(G/L)(m)$$

= $\nu_{N'} (F(G/H)(n_{H}) \otimes \varphi_{H}))(G/L)(m)$
= $N' (\varphi_{H}(G/L)(m)) (F(G/H)(n_{H})).$

That these two are equivalent is because F is a natural transformation, so the diagram below commutes.

$$N(G/L) \xrightarrow{F(G/L)} N'(G/L)$$

$$N(\varphi_H(G/L)(m)) \uparrow \qquad \uparrow N'(\varphi_H(G/L)(m))$$

$$N(G/H) \xrightarrow{F(G/H)} N'(G/H)$$

The next lemma is an $\mathcal{O}_{\mathcal{F}}$ module version of [**Bie81**, Proposition 3.1].

LEMMA 3.9.4. If M is finitely generated projective then ν is an isomorphism.

PROOF. Consider first the case $M = R[-, G/H]_{\mathcal{O}_{\mathcal{F}}}$, then the map ν becomes $\nu : N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom} \left(R[-, G/H]_{\mathcal{O}_{\mathcal{F}}}, R[-, ?]_{\mathcal{O}_{\mathcal{F}}} \right) \longrightarrow \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}} \left(R[-, G/H]_{\mathcal{O}_{\mathcal{F}}}, N(-) \right).$ But, using Lemmas 2.0.2 and 2.1.3, the left hand side collapses to

$$N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom} \left(R[-, G/H]_{\mathcal{O}_{\mathcal{F}}}, R[-, ?]_{\mathcal{O}_{\mathcal{F}}} \right) \cong N(?) \otimes_{R} R[G/H, ?]_{\mathcal{O}_{\mathcal{F}}}$$

(*)
$$\cong N(G/H).$$

Under these isomorphisms $n_H \in N(G/H)$ maps to

$$n_H \otimes \mathrm{id}_H \in N(?) \otimes_R R[G/H, ?]_{\mathcal{O}_F}$$

and then to $n_H \otimes \varphi$ where φ is the unique natural transformation φ such that $\varphi(G/H)(\mathrm{id}_H) = \mathrm{id}_H$.

The right hand side collapses to

(†)
$$\operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(R[-,G/H]_{\mathcal{O}_{\mathcal{F}}},N(-)) \cong N(G/H)$$

again by the Yoneda-type Lemma 2.0.2, where n_H maps to the unique natural transformation ψ with $\psi(G/H)(id) = n_H$.

$$\nu(n_H \otimes \varphi)(G/H)(\mathrm{id}_H) = N(\varphi(G/H)(\mathrm{id}_H))(n_H) = N(\mathrm{id}_H)(n_H) = n_H$$

Precomposing ν with the isomorphism from (\star) and postcomposing with the isomorphism from (†) gives the identity map $N(G/H) \rightarrow N(G/H)$ and hence ν is an isomorphism.

The case for finitely generated free modules follows as all the necessary functors commute with finite direct sums, and for projectives from naturality of ν proved in Lemma 3.9.3.

The following result is an analog of [Bie81, 5.2(a,c)].

Lemma 3.9.5.

- (1) If M is finitely presented and N is flat then ν is an isomorphism.
- (2) If M is finitely generated and N is projective then ν is an isomorphism.

Proof. (1) If

 $F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$

is an exact sequence with F_i finitely generated free for i = 0, 1 then by the naturality of ν and flatness of N we have the following commutative diagram with exact rows (for brevity we write Hom for $\operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}$ and \otimes for $\otimes_{\mathcal{O}_{\mathcal{F}}}$).

The right hand and middle vertical maps are isomorphisms by Lemma 3.9.4, the result follows from the 5-Lemma.

(2) If F(?) is free then by Lemma 2.1.3 there is an isomorphism

$$F(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}(M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) \cong \operatorname{Hom}(M, F).$$

Checking the definition of this isomorphism shows it's induced by ν . If N(?) is projective and $i : N(?) \hookrightarrow F(?)$ is a split injection then by naturality of ν , the following diagram commutes:

$$\begin{array}{ccc} N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}(M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) & \longrightarrow & \operatorname{Hom}(M, N) \\ & & & & \downarrow \\ F(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}(M(-), R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) & \xrightarrow{\cong} & \operatorname{Hom}(M, F) \end{array}$$

Since *i* is a split injection, the left hand map is an injection and the top map must be an injection. Consider the commutative diagram in the proof of part 1, only F_0 is known to be projective so the middle vertical map is an isomorphism. Since N is projective the left and right hand vertical maps are monomorphisms and the Four Lemma completes the proof, implying that the left hand vertical map is an isomorphism.

LEMMA 3.9.6. If P_* is any chain complex of contravariant $\mathcal{O}_{\mathcal{F}}$ -modules and N is any contravariant $\mathcal{O}_{\mathcal{F}}$ -module, the following morphism is both well defined and natural in P_* and N:

$$\xi^{i}: N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} H^{i} P_{*}(?)^{D} \to H^{i} \big(N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} P_{*}(?)^{D} \big)$$

$$\xi^{i}: N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} H^{i} (\operatorname{Hom}(P_{*}(-), R[-, ?]) \to H^{i} \big(N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}(P_{*}(-), R[-, ?]) \big)$$

$$n_{H} \otimes [\varphi_{H}] \mapsto [n_{H} \otimes \varphi_{H}],$$

where $H^i P_*(?)^D : G/H \mapsto H^i P_*(G/H)^D$.

PROOF. If φ_H is a cocycle, $n_H \otimes \varphi_H$ is also a cocycle and similarly if φ_H is a coboundary then $n_H \otimes \varphi_H$ is a coboundary.

If $\alpha: G/L \to G/H$ is a G-map then by definition $\alpha_*[\varphi_H] = [\alpha_*\varphi_H]$ and

$$\begin{split} \xi^i \big(\alpha^* n_H \otimes [\varphi_H] - n_H \otimes \alpha_* [\varphi_H] \big) &= \xi^i \big(\alpha^* n_H \otimes [\varphi_H] - n_H \otimes [\alpha_* \varphi_H] \big) \\ &= [\alpha^* n_H \otimes \varphi_H - n_H \otimes \alpha_* \varphi_H] \\ &= 0. \end{split}$$

Finally naturality follows because the functors H^i and $\operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(-,?)$ are natural, and so is the process of taking tensor products.

Since ν is natural (Lemma 3.9.3), if P_* is a projective resolution of <u>R</u> by contravariant modules then ν induces chain homomorphisms

 $N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} P_*(?)^D \longrightarrow \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(P_*, N)$

which in turn induce maps on cohomology

$$H^i(N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} P_*(?)^D) \longrightarrow H^i_{\mathcal{O}_{\mathcal{F}}}(G,N).$$

Precomposing this with ξ^i gives a map

 $\nu^{i}: N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} H^{i}_{\mathcal{O}_{\mathcal{F}}}(G, R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) \longrightarrow H^{i}_{\mathcal{O}_{\mathcal{F}}}(G, N).$

PROPOSITION 3.9.7. If G is $\mathcal{O}_{\mathcal{F}} FP_n$ over R and N is projective then ν^i is an isomorphism for all $i \leq n$.

PROOF. Choose a projective resolution $P_* \longrightarrow \underline{R}$, finitely generated up to dimension n and write K_i for the i^{th} syzygy of P_* . Since N is projective it is also flat and we have the following commutative diagram with exact rows, where we omit the $\mathcal{O}_{\mathcal{F}}$ on \otimes , Hom, and H^i .

Since G is $\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n$, K_{i-1} and P_{i-1} are finitely generated, Lemma 3.9.5(2) implies the middle and left hand vertical maps are isomorphisms. The 5-Lemma completes the proof.

The following result is an analog of [Bie81, 9.1].

PROPOSITION 3.9.8. If G is $\mathcal{O}_{\mathcal{F}}FP$ over R, with $\mathcal{O}_{\mathcal{F}}cd_R G = n$, and N is any contravariant module then there is a natural isomorphism

$$\nu^n: N(?) \otimes_{\mathcal{O}_{\mathcal{F}}} H^n_{\mathcal{O}_{\mathcal{F}}}(G, R[-, ?]_{\mathcal{O}_{\mathcal{F}}}) \cong H^n_{\mathcal{O}_{\mathcal{F}}}(G, N).$$

PROOF. Let

$$0 \longrightarrow K \longrightarrow F \longrightarrow N \longrightarrow 0$$

be a short exact sequence of contravariant $\mathcal{O}_{\mathcal{F}}$ -modules over R with F free. By the naturality of ν^n we have the following commutative diagram with exact rows, we omit the $\mathcal{O}_{\mathcal{F}}$ decorations on \otimes , H^* , and R[-,?] for brevity.

$$\begin{split} K(?)\otimes H^n(G,R[-,?]) &\succ F(?)\otimes H^n(G,R[-,?]) &\succ N(?)\otimes H^n(G,R[-,?]) &\succ 0 \\ & \downarrow & \downarrow & \downarrow \\ H^n(G,K) & \longrightarrow H^n(G,F) & \longrightarrow H^n(G,N) &\longrightarrow 0 \end{split}$$

The middle vertical map is an isomorphism by Proposition 3.9.7, thus by the Four Lemma, the right hand vertical map is an epimorphism. Since there are no restrictions on N, we conclude that the left hand vertical map is an epimorphism and by the 5-Lemma that the right hand map is an isomorphism.

CHAPTER 4

Mackey and cohomological Mackey functors

This chapter contains material that has appeared in:

 Finiteness conditions for Mackey and cohomological Mackey functors (J. Algebra 411 (2014), no. 0, 225–258) [SJG14]

Throughout this section we will work over an arbitrary subfamily \mathcal{F} of $\mathcal{F}in$, closed under conjugation and taking subgroups. One could also work over larger families of subgroups such as \mathcal{VCyc} [**Deg13b**, p.101], however this necessitates a change in the construction of Mackey and cohomological Mackey functors and we shall not consider it.

In Section 4.1 we give an overview of Mackey functors and cohomological Mackey functors including the description due to Yoshida of cohomological Mackey functors as modules over the category $\mathcal{H}_{\mathcal{F}}$ [Yos83].

Section 4.2 contains a complete description of the condition $\mathcal{M}_{\mathcal{F}} FP_n$, the Mackey functor analogue of the $\mathcal{O}_{\mathcal{F}} FP_n$ conditions.

COROLLARY 4.2.6. Over any ring R, a group is $\mathcal{M}_{\mathcal{F}} FP_n$ if and only if it is $\mathcal{O}_{\mathcal{F}} FP_n$.

The main result of Section 4.3 is that the Bredon cohomology with coefficients in a cohomological Mackey functor may be calculated with a projective resolution of cohomological Mackey functors. We show in Proposition 4.3.6 that a projective resolution of <u>R</u> by Bredon modules can be induced to a projective resolution of the fixed point functor R^- by cohomological Mackey functors, this is an analogue of [**MPN06**, Theorem 3.8]—that one can induce a projective resolution of <u>R</u> by Bredon modules to a projective resolution of the Burnside functor B^G by Mackey functors.

Building on this, in Section 4.4 we study the $\mathcal{H}_{\mathcal{F}} FP_n$ conditions, the cohomological Mackey functor analogue of the $\mathcal{O}_{\mathcal{F}} FP_n$ conditions, relating them to the $\mathcal{F} FP_n$ conditions defined in Section 1.4.

THEOREM 4.4.1. If R is a commutative Noetherian ring, a group is $\mathcal{H}_{\mathcal{F}} FP_n$ if and only if it is $\mathcal{F} FP_n$.

In Section 4.5 the main result is the following.

THEOREM 4.5.1. $\mathcal{H}_{\mathcal{F}} \operatorname{cd} G = \mathcal{F} \operatorname{cd} G$ for all groups G.

In Section 4.6 we prove that, depending on the coefficient ring, $\mathcal{H}_{\mathcal{F}} cd G$ may be calculated using a proper subfamily of \mathcal{F} . When working over \mathbb{Z} we need consider only the family \mathcal{P} of subgroups in \mathcal{F} with prime power order, and over either the finite field \mathbb{F}_p or over $\mathbb{Z}_{(p)}$ (the integers localised at p), we need consider only the family \mathcal{P} of subgroups of \mathcal{F} with order a power of p. This is similar to a result of Leary and Nucinkis for \mathcal{F} -cohomology [**LN10**, §4].

THEOREM 4.6.1. For all $n \in \mathbb{N} \cup \{\infty\}$, the conditions $\mathcal{H}_{\mathcal{F}} cd G = n$ and $\mathcal{H}_{\mathcal{P}} cd G = n$ are equivalent, as are the conditions $\mathcal{H}_{\mathcal{F}} FP_n$ and $\mathcal{H}_{\mathcal{P}} FP_n$.

Over the finite field \mathbb{F}_p we can be even more precise.

COROLLARY 4.6.11. G is $\mathcal{H}_{\mathcal{F}} \mathrm{FP}_n$ over \mathbb{F}_p if and only if \mathcal{P} has finitely many conjugacy classes and WH is FP_n over \mathbb{F}_p for all $H \in \mathcal{P}$.

4.1. Introduction

4.1.1. Mackey functors. There are many constructions of Mackey functors, we use the construction coming from modules over a category, an approach due to Linder [Lin76]. Another construction is mentioned in Remark 4.1.6. We begin by building a small category $\mathcal{M}_{\mathcal{F}}$ then Mackey functors will be contravariant $\mathcal{M}_{\mathcal{F}}$ -modules. As in $\mathcal{O}_{\mathcal{F}}$, the objects of $\mathcal{M}_{\mathcal{F}}$ are the transitive *G*-sets with stabilisers in \mathcal{F} , the morphism set however is much larger. A *basic morphism* from G/H to G/K, where H and K are in \mathcal{F} , is an equivalence class of diagrams of the form

$$G/H \xleftarrow{\alpha} G/L \xrightarrow{\beta} G/K$$

where the maps are G-maps, and $L \in \mathcal{F}$. This basic morphism is equivalent to

$$G/H \xleftarrow{\alpha'} G/L' \xrightarrow{\beta'} G/K$$

if there is a bijective G-map $\sigma : G/L \to G/L'$, fitting into the commutative diagram below:

Form the free abelian monoid on these basic morphisms, and complete this free abelian monoid to a group, denoted $[G/H, G/K]_{\mathcal{M}_{\mathcal{F}}}$. This is the set of morphisms in $\mathcal{M}_{\mathcal{F}}$ from G/H to G/K.

REMARK 4.1.1. When building the Mackey category, we could instead have started with equivalence classes of diagrams

$$G/H \leftarrow \Delta \rightarrow G/K$$

where Δ is any finitely generated *G*-set with stabilisers in \mathcal{F} and the maps are *G*maps. This can be shown to be the free abelian monoid on the basic morphisms [**TW95**, Proposition 2.2]. Because of this alternative construction, we will pass freely between writing

$$(G/H \leftarrow G/L \to G/K) + (G/H \leftarrow G/L' \to G/K)$$

and

$$\left(G/H \leftarrow G/L \coprod G/L' \to G/K\right).$$

To complete the description of $\mathcal{M}_{\mathcal{F}}$, we must describe composition of morphisms. It's sufficient to describe composition of basic morphisms, and then use distributivity to extend this to all morphisms. If

$$G/H \leftarrow G/L \to G/K$$

and

$$G/K \leftarrow G/S \rightarrow G/Q$$

are two basic morphisms then their composition is the pullback diagram below in the category of G-sets.



LEMMA 4.1.2 (Composition of morphisms in $\mathcal{M}_{\mathcal{F}}$). [MPN06, §3] The diagram below is a pullback in the category of G-sets.



Notice that the subgroup $L^g \cap S^{g'x^{-1}}$ is both a subgroup of K via the maps on the left and subconjugated to K via the map α_x , which is the composition of the maps on the right.

If H is a subgroup of G the notation H^g means the conjugate $g^{-1}Hg$.

LEMMA 4.1.3 (Standard form for morphisms in $\mathcal{M}_{\mathcal{F}}$). [**TW95**, Lemma 2.1] Any basic morphism is equivalent to one in the standard form:



Recall that two such basic morphisms are equivalent if there is a commutative diagram of the form:

$$G/K \overset{\operatorname{id}}{\underset{\operatorname{id}}{\overset{\alpha_g}{\underset{\alpha_x}{\overset{\alpha_{g'}}}{\overset{\alpha_{g'}}{\overset{\alpha_{g'}}{\overset{\alpha_{g'}}{\overset{\alpha_{g'}}}{\overset{\alpha_{g'}}{\overset{\alpha_{g'}}}{\overset{\alpha_{g'}}{\overset{\alpha_{g'}}}}}}}}}}}}}}}}}}}}}$$

The commutativity of the left hand triangle ensures that $x \in K$, and that of the right hand diagram gives $\alpha_g = \alpha_{g'} \circ \alpha_x$, or more concisely gS = xg'S. This means KgS = Kg'S and $x = gS(g')^{-1} \cap K = gSg^{-1} \cap K$. Thus a basic morphism is determined by both an element of $K \setminus G/S$ and a subgroup L, subconjugate to K, unique up to conjugation by an element $x \in gSg^{-1} \cap K$. In summary,

(4.1)
$$[G/K, G/S]_{\mathcal{M}_{\mathcal{F}}} = \bigoplus_{g \in K \setminus G/S} \bigoplus_{\substack{L \le gSg^{-1} \cap K \\ \text{Up to } gSg^{-1} \cap K \text{-conjugacy}}} \mathbb{Z}_{L,g},$$

where $\mathbb{Z}_{L,g} \cong \mathbb{Z}$ for all L and g.

EXAMPLE 4.1.4. If S = 1 then (4.1) becomes

$$[G/K, G/1]_{\mathcal{M}_{\mathcal{F}}} = \bigoplus_{g \in K \setminus G} \mathbb{Z}_g \cong \mathbb{Z}[K \setminus G].$$

REMARK 4.1.5. The category $\mathcal{M}_{\mathcal{F}}$ has property (A) by construction, but it does not have property (EI). For example, given any non-trivial $H \in \mathcal{F}$, the endomorphism

$$e = \left(G/H \xleftarrow{\alpha_1} G/1 \xrightarrow{\alpha_1} G/H \right)$$

is not an isomorphism. If

$$m = \left(G/H \xleftarrow{\alpha_1} G/K \xrightarrow{\alpha_g} G/H \right)$$

is some other basic morphism then their composition is

$$m \circ e = \sum_{x \in H/K} \left(G/H \xleftarrow{\alpha_1} G/1 \xrightarrow{\alpha_{xg}} G/H \right).$$

So it's clear that e cannot be a sum of automorphisms of G/H.

Following [**MPN06**], we will mostly consider contravariant Mackey functors. From here on, whenever we write $\mathcal{M}_{\mathcal{F}}$ -module, we mean contravariant $\mathcal{M}_{\mathcal{F}}$ -module.

REMARK 4.1.6 (Green's alternative description of Mackey functors). There is an alternative description of Mackey functors, due to Green [**Gre71**], which we include here in full because when we later study cohomological Mackey functors we will need some of the language.
Green defined a Mackey functor M as a mapping,

$$M: \{G/H : H \in \mathcal{F}\} \longrightarrow \mathbf{R-Mod}$$

with morphisms for any finite subgroups $K \leq H$ in \mathcal{F} ,

$$M(I_K^H) : M(G/K) \longrightarrow M(G/H)$$
$$M(R_K^H) : M(G/H) \longrightarrow M(G/K)$$
$$M(c_g) : M(G/H) \longrightarrow M(G/H^{g^{-1}})$$

called *induction*, *restriction* and *conjugation* respectively. Induction is sometimes also called transfer. In the literature, $M(I_K^H)$, $M(R_K^H)$ and $M(c_g)$ are often written as just I_K^H , R_K^H and c_g , omitting the M entirely. We choose to use different notation so that we can identify I_K^H , R_K^H and c_g with specific morphisms in $\mathcal{M}_{\mathcal{F}}$ (see the end of this remark).

This mapping M must satisfy the following axioms,

- (0) $M(I_H^H)$, $M(R_H^H)$ and $M(c_h)$ are the identity morphism for all $h \in H$.
- (1) $M(R_J^K) \circ M(R_K^H) = M(R_J^H)$, where $J \leq K \leq H$ and $J, K, H \in \mathcal{F}$.
- (2) $M(I_K^H) \circ M(I_J^K) = M(I_J^H)$, where $J \leq K \leq H$ and $J, K, H \in \mathcal{F}$.
- (3) $M(c_g) \circ M(c_h) = M(c_{gh})$ for all $g, h \in G$.
- (4) $M(R_{Kg^{-1}}^{Hg^{-1}}) \circ M(c_g) = M(c_g) \circ M(R_K^H)$, where $K \leq H$ and $K, H \in \mathcal{F}$ and $g \in G$.
- (5) $M(I_{K^{g^{-1}}}^{H^{g^{-1}}}) \circ M(c_g) = M(c_g) \circ M(I_K^H)$, where $K \leq H$ and $K, H \in \mathcal{F}$ and $g \in G$.
- (6) $M(R_J^H) \circ M(I_K^H) = \sum_{x \in J \setminus H/K} M(I_{J \cap K^{x-1}}^J) \circ M(c_x) \circ M(R_{J^x \cap K}^K)$, where $J, K \leq H$ and $J, K, H \in \mathcal{F}$.

Axiom (6) is often called the Mackey axiom. Converting between this description and our previous description is done by rewriting induction, restriction and conjugation in terms of morphisms of $\mathcal{M}_{\mathcal{F}}$.

$$M(I_K^H) \longleftrightarrow M(G/H \xleftarrow{\alpha_1} G/K \xrightarrow{\operatorname{id}} G/K)$$
$$M(R_K^H) \longleftrightarrow M(G/K \xleftarrow{\operatorname{id}} G/K \xrightarrow{\alpha_1} G/H)$$
$$M(c_g) \longleftrightarrow M(G/H^{g^{-1}} \xleftarrow{\operatorname{id}} G/H^{g^{-1}} \xrightarrow{\alpha_g} G/H)$$

Because of the above, we make the following definitions,

$$I_K^H = \left(G/H \xleftarrow{\alpha_1} G/K \xrightarrow{\mathrm{id}} G/K\right)$$
$$R_K^H = \left(G/K \xleftarrow{\mathrm{id}} G/K \xrightarrow{\alpha_1} G/H\right)$$
$$c_g = \left(G/H^{g^{-1}} \xleftarrow{\mathrm{id}} G/H^{g^{-1}} \xrightarrow{\alpha_g} G/H\right)$$

It is possible to write any morphism in $\mathcal{M}_{\mathcal{F}}$ as a composition of the three types of morphisms above.

One can check that Green's axioms all follow from the description of the composition of morphisms in $\mathcal{M}_{\mathcal{F}}$ as pullbacks (Lemma 4.1.2), and vice versa. Complete proofs of the equivalence of this definition with our previous one can be found in [**TW95**, §2].

4.1.1.1. Free modules. In this section we describe the structure of $\operatorname{End}(G/H)$ and study free $\mathcal{M}_{\mathcal{F}}$ -modules.

REMARK 4.1.7 (Structure of $\operatorname{End}(G/H)$). As mentioned in Remark 4.1.5, $\mathcal{M}_{\mathcal{F}}$ doesn't have property (EI). Consider the endomorphisms of an object given by the diagrams of the form

$$a_q = \left(G/H \stackrel{\alpha_1}{\longleftarrow} G/H \stackrel{\alpha_g}{\longrightarrow} G/H \right).$$

Every $g \in WH$ uniquely determines a G-map $\alpha_g : G/H \to G/H$ and every Gmap comes from such a g. Finally, since $a_g \circ a_h = a_{hg}$, we determine that such endomorphisms give a copy of $\mathbb{Z}[WH]^{\mathrm{op}}$ inside $\mathrm{End}(G/H)$. This is similar to the situation over the orbit category, where $\mathrm{End}_{\mathcal{O}_F}(G/H) \cong \mathbb{Z}[WH^{\mathrm{op}}]$. Thus, as with \mathcal{O}_F -modules, if M is a Mackey functor, then M(G/H) is a right $R[WH^{\mathrm{op}}]$ module, equivalently a left R[WH]-module.

A basic morphism in $\operatorname{End}(G/H)$ is determined by a morphism in standard form

$$e_{L,g} = \left(G/H \xleftarrow{\alpha_1} G/L \xrightarrow{\alpha_g} G/H \right)$$

where L is some subgroup of G. As such we can filter $\operatorname{End}(G/H)$ via the poset \mathcal{F}/G of conjugacy classes of subgroups in \mathcal{F} . If L is a finite subgroup of G then we write $\operatorname{End}(G/H)_L$ for the basic morphisms $e_{L,g}$ for all $g \in G$. Note that in particular, $\operatorname{End}(G/H)_H \cong R[WH]$ by the paragraph above. Addition gives $\operatorname{End}(G/H)_L$ an abelian group structure. Composing two elements of $\operatorname{End}(G/H)_L$ doesn't necessarily give an element of $\operatorname{End}(G/H)_L$, but pre-composing an element of $\operatorname{End}(G/H)_L$ by some a_w does, since

$$e_{L,g} \circ a_w \cong e_{L,wg}.$$

Thus $R \operatorname{End}(G/H)_L$ is a left R[WH]-module. In summary, there is an R[WH]-module isomorphism

$$R \operatorname{End}(G/H) \cong \bigoplus_{L \in \mathcal{F}/G} R \operatorname{End}(G/H)_L$$

where $R \operatorname{End}(G/H)_H \cong R[WH]$.

EXAMPLE 4.1.8. Using (4.1),

$$R \operatorname{End}(G/H)_1 \cong \bigoplus_{H \setminus G/H} R_{1,g},$$

with left action of $w \in W_G H$ taking $g \mapsto wg$. In other words, $R \operatorname{End}(G/H)_1 \cong R[H \setminus G/H]$ with the canonical action of $W_G H$. This is not in general finitely generated—take for example $G = D_{\infty}$, the infinite dihedral group generated by the involutions a and b, and $H = \langle a \rangle$. Then $W_G H$ is the trivial group but $H \setminus G/H$ is an infinite set so $R[H \setminus G/H]$ is not a finitely generated R-module.

LEMMA 4.1.9. As a left $R[W_GS]$ -module, $R[G/S, G/K]_{\mathcal{M}_F}$ is an $R[W_GS]$ permutation module with finite stabilisers. In addition, $R[G/1, G/K]_{\mathcal{M}_F}$ is FP_{∞} over RG.

PROOF. The left action of $w \in W_G S$ on $[G/S, G/K]_{\mathcal{M}_{\mathcal{F}}}$ is the action given by pre-composing any basic morphism $G/S \stackrel{\text{id}}{\leftarrow} G/L \stackrel{\alpha_g}{\to} G/K$ with the morphism $G/S \stackrel{\text{id}}{\leftarrow} G/S \stackrel{\alpha_w}{\to} G/S$ to yield the morphism

$$G/S \stackrel{\alpha_1}{\leftarrow} G/L \stackrel{\alpha_{wg}}{\rightarrow} G/K.$$

To show this we calculate the pullback:



Under the identification (4.1), w maps $R_{L,g}$ onto $R_{L,wg}$, so the stabiliser of this action is the stabiliser of the action of $R[W_GS]$ on $R[S \setminus G/K]$, which is finite. In particular $R[G/S, G/K]_{\mathcal{M}_{\mathcal{F}}}$ is an $R[W_GS]$ -permutation module with finite stabilisers. If S = 1 then, using (4.1), $R[G/1, G/K]_{\mathcal{M}_{\mathcal{F}}} \cong R[G/K]$ with RGacting by multiplication on the left, thus $R[G/1, G/K]_{\mathcal{M}_{\mathcal{F}}}$ is FP $_{\infty}$ as a left RGmodule.

REMARK 4.1.10. $R[G/S, G/K]_{\mathcal{M}_{\mathcal{F}}}$ is not in general finitely generated as a left $R[W_GS]$ -module. For an example of this let \mathcal{F} be all finite subgroups and choose a group G with a finite subgroup S such that $S \setminus G$ has infinitely many W_GS -orbits. Then, by Example 4.1.4,

$$R[G/S, G/1]_{\mathcal{M}_{\mathcal{F}}} \cong R[S \setminus G]$$

which is not finitely generated as a left $R[W_GS]$ module.

4.1.1.2. Induction. Let $\sigma : \mathcal{O}_{\mathcal{F}} \to \mathcal{M}_{\mathcal{F}}$ be the covariant functor sending

$$\sigma(G/H) = G/H$$

$$\sigma(G/H \xrightarrow{\alpha} G/K) = (G/H \xleftarrow{\text{id}} G/H \xrightarrow{\alpha} G/K).$$

Thus σ induces restriction, induction, and coinduction between $\mathcal{O}_{\mathcal{F}}$ -modules and $\mathcal{M}_{\mathcal{F}}$ -modules.

LEMMA 4.1.11. [**MPN06**, Proposition 3.6] There is an $\mathcal{O}_{\mathcal{F}}$ -module isomorphism:

$$\operatorname{Res}_{\sigma} R[G/H, -]_{\mathcal{M}_{\mathcal{F}}} \cong \bigoplus_{L \leq H} R \otimes_{W_{H}L} R[G/L, -]_{\mathcal{O}_{\mathcal{F}}}$$

Let B^G denote the Burnside functor B^G which, by an abuse of notation since G/G is not an object of $\mathcal{M}_{\mathcal{F}}$, can be defined as

$$B^G = R[-, G/G]_{\mathcal{M}_{\mathcal{F}}}.$$

Upon evaluation at G/K for some $K \in \mathcal{F}$,

$$B^G(G/K) = \bigoplus_{\substack{L \le K \\ \text{Up to } K \text{-conjugacy}}} R_L.$$

This is not so dissimilar from the case of the orbit category $\mathcal{O}_{\mathcal{F}}$ where, using a similar abuse of notation, one could view <u>R</u> as $R[-, G/G]_{\mathcal{O}_{\mathcal{F}}}$.

EXAMPLE 4.1.12. If <u>R</u> is the constant contravariant $\mathcal{O}_{\mathcal{F}}$ -module then using Lemma 4.1.11,

$$\operatorname{Ind}_{\sigma} \underline{R}(G/H) \cong R[G/H, \sigma(-)]_{\mathcal{M}_{\mathcal{F}}} \otimes_{\mathcal{O}_{\mathcal{F}}} \underline{R}$$
$$\cong \bigoplus_{L \leq H} R \otimes_{W_{H}L} R[G/L, -]_{\mathcal{O}_{\mathcal{F}}} \otimes_{\mathcal{O}_{\mathcal{F}}} \underline{R}$$
$$\cong \bigoplus_{L \leq H} R.$$

Checking the morphisms as well, one sees that

$$\operatorname{Ind}_{\sigma} \underline{R} \cong B^G$$

PROPOSITION 4.1.13. [MPN06, Theorem 3.8] Although induction with σ is not exact in general, induction with σ takes contravariant resolutions of <u>R</u> by projective $\mathcal{O}_{\mathcal{F}}$ -modules to resolutions of B^G by projective $\mathcal{M}_{\mathcal{F}}$ -modules.

4.1.1.3. Homology and cohomology. We define the Mackey cohomology and Mackey homology for any contravariant $\mathcal{M}_{\mathcal{F}}$ -module M and covariant $\mathcal{M}_{\mathcal{F}}$ -module A as

$$H^*_{\mathcal{M}_{\mathcal{F}}}(G, M) = \operatorname{Ext}^*_{\mathcal{M}_{\mathcal{F}}}(B^G, M)$$
$$H^{\mathcal{M}_{\mathcal{F}}}_*(G, A) = \operatorname{Tor}^*_{\mathcal{M}_{\mathcal{F}}}(B^G, A).$$

A corollary of Proposition 4.1.13 is the following.

COROLLARY 4.1.14. [MPN06, Theorem 3.8]

$$H^n_{\mathcal{M}_{\mathcal{F}}}(G, M) \cong H^n_{\mathcal{O}_{\mathcal{F}}}(G, \operatorname{Res}_{\sigma} M).$$

G is said to be $\mathcal{M}_{\mathcal{F}} FP_n$ if there is a projective resolution of B^G , finitely generated up to degree n, and G has $\mathcal{M}_{\mathcal{F}} cd G \leq n$ if there is a length n projective resolution of B^G by $\mathcal{M}_{\mathcal{F}}$ -modules. **4.1.2. Cohomological Mackey functors.** A Mackey functor M is called cohomological if, using the language of Remark 4.1.6, it satisfies

(4.2)
$$M(I_K^H) \circ M(R_K^H) = (m \mapsto |H:K|m)$$

for all subgroups $K \leq H$ in \mathcal{F} . Recall from Remark 4.1.6 that to describe a Mackey functor M it is sufficient to describe it on objects and on the induction, restriction and conjugation morphisms in $\mathcal{M}_{\mathcal{F}}(I_K^H, R_K^H \text{ and } c_g)$, we use this in the examples below.

EXAMPLE 4.1.15 (Group cohomology). The group cohomology functor is cohomological Mackey, more precisely the functor

$$H^n(-,R): G/H \longmapsto H^n(H,R).$$

Where $H^n(-,R)(c_g)$ is induced by conjugation, $H^n(-,R)(R_K^H)$ is the usual restriction map and $H^n(-,R)(I_K^H)$ is the transfer (see for example [**Bro94**, §III.9]). That the group cohomology functor satisfies (4.2) is [**Bro94**, III.9.5(ii)].

EXAMPLE 4.1.16 (Fixed point and fixed quotient functors). If M is a RGmodule then we write M^- for the fixed point functor

$$M^-: G/H \longmapsto M^H$$

where $M^H = \operatorname{Hom}_{RH}(R, M)$. For any $K \leq H$ in \mathcal{F} , $M^-(R_K^H)$ is the inclusion, $M^-(I_K^H)$ is the trace $m \mapsto \sum_{h \in H/K} hm$, and $M^-(c_g)$ is the map $m \mapsto gm$.

We write M_{-} for the fixed quotient functor

$$M_{-}: G/H \longmapsto M_H$$

where $M_H = R \otimes_{RH} M$. For any $K \leq H$ in \mathcal{F} , $M_-(R_K^H)$ is the trace $1 \otimes m \mapsto 1 \otimes \sum_{h \in H/K} hm$, $M_-(I_K^H)$ is the inclusion, and $M_-(c_g)$ is the map $m \mapsto gm$.

LEMMA 4.1.17. [**MPN06**, Lemma 4.2][**TW90**, 6.1] There are Mackey functor isomorphisms for any RG-module M,

$$\operatorname{CoInd}_{RG}^{\mathcal{M}_{\mathcal{F}}} M \cong M^{-}$$
$$\operatorname{Ind}_{RG}^{\mathcal{M}_{\mathcal{F}}} M \cong M_{-}$$

where induction and coinduction are with the functor $\widehat{\mathbb{Z}G} \to \mathcal{M}_{\mathcal{F}}$ given by composition of the usual inclusion functor $\widehat{\mathbb{Z}G} \to \mathcal{O}_{\mathcal{F}}$ and the functor $\sigma : \mathcal{O}_{\mathcal{F}} \to \mathcal{M}_{\mathcal{F}}$. Thus there are also adjoint isomorphisms, for any Mackey functor N.

$$\operatorname{Hom}_{RG}(N(G/1), M) \cong \operatorname{Hom}_{\mathcal{M}_{\mathcal{F}}}(N, M^{-})$$
$$\operatorname{Hom}_{RG}(M, N(G/1)) \cong \operatorname{Hom}_{\mathcal{M}_{\mathcal{F}}}(M^{-}, N)$$

4. MACKEY AND COHOMOLOGICAL MACKEY FUNCTORS

As observed by Thévenaz and Webb in [**TW95**, §16], in [**Yos83**] Yoshida proves that the category of cohomological Mackey modules is isomorphic to the category of modules over the Hecke category $\mathcal{H}_{\mathcal{F}}$, which we shall describe below. Yoshida concentrates mainly on finite groups but observes in [**Yos83**, §5, Theorem 4.3'] that this isomorphism will hold for $\mathcal{M}_{\mathcal{F}}$ -modules, where \mathcal{F} is any subfamily of the family of finite groups.

The Hecke category $\mathcal{H}_{\mathcal{F}}$ has for objects the transitive *G*-sets with stabilisers in \mathcal{F} . The morphisms between the objects G/H and G/K are exactly the $\mathbb{Z}G$ module homomorphisms, $\operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}[G/H], \mathbb{Z}[G/K])$.

REMARK 4.1.18. In [Yos83], Yoshida actually uses the category $\mathcal{H}_{\mathcal{F}}'$, this category has the same objects, but the morphisms between G/H and G/K are the RG-module homomorphisms $\operatorname{Hom}_{RG}(R[G/H], R[G/K])$. He then studies Radditive functors from $\mathcal{H}_{\mathcal{F}}'$ into the category of left R-modules and proves these are exactly the cohomological Mackey functors. We claim that the category of R-additive functors $\mathcal{H}_{\mathcal{F}}' \to \mathbf{R}$ -Mod and the category of additive functors $\mathcal{H}_{\mathcal{F}} \to \mathbf{R}$ -Mod are isomorphic, where the isomorphism preserves the values the functors take on objects.

Since $\mathbb{Z}[G/H]$ is finitely presented as a $\mathbb{Z}G$ -module and R is flat as a \mathbb{Z} -module there is an isomorphism for all $H, K \in \mathcal{F}$ (the proof is essentially the proof of [Wei94, 3.3.8])

$$\operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}[G/H], \mathbb{Z}[G/K]) \otimes_{\mathbb{Z}} R \cong \operatorname{Hom}_{RG}(R[G/H], R[G/K]).$$

Using the above and that $\operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}[G/H], \mathbb{Z}[G/K])$ is free as a \mathbb{Z} -module, there is a natural isomorphism for any *R*-module *A*

$$\operatorname{Hom}_{\mathbb{Z}}(\operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}[G/H],\mathbb{Z}[G/K]), A)$$
$$\cong \operatorname{Hom}_{R}(\operatorname{Hom}_{\mathbb{Z}G}(\mathbb{Z}[G/H],\mathbb{Z}[G/K]) \otimes_{\mathbb{Z}} R, A)$$
$$\cong \operatorname{Hom}_{R}(\operatorname{Hom}_{RG}(R[G/H], R[G/K]), A).$$

The claim follows from this isomorphism.

REMARK 4.1.19. In [**Deg13a**] Degripse considers the categories $\operatorname{Mack}_{\mathcal{F}}G$ and $\operatorname{coMack}_{\mathcal{F}}G$. In the notation used here $\operatorname{Mack}_{\mathcal{F}}G$ is the category of $\mathcal{M}_{\mathcal{F}}$ -modules and $\operatorname{coMack}_{\mathcal{F}}G$ is the subcategory of cohomological Mackey functors, Degripse doesn't study modules over $\mathcal{H}_{\mathcal{F}}$.

LEMMA 4.1.20 (Free and projective $\mathcal{H}_{\mathcal{F}}$ -modules). [**TW95**, Theorem 16.5(ii)] The free $\mathcal{H}_{\mathcal{F}}$ -modules are exactly the fixed point functors of permutation modules with stabilisers in \mathcal{F} , and the projective $\mathcal{H}_{\mathcal{F}}$ -modules are exactly the fixed point functors of direct summands of permutation modules with stabilisers in \mathcal{F} .

4.1. INTRODUCTION

Thévenaz and Webb describe a map $\pi : \mathcal{M}_{\mathcal{F}} \to \mathcal{H}_{\mathcal{F}}$ (they call this map α), taking objects G/H in $\mathcal{M}_{\mathcal{F}}$ to G/H in $\mathcal{H}_{\mathcal{F}}$ and on morphisms as follows, for any $K \leq H$,

- $\pi(R_K^H)$ is the natural projection map $\mathbb{Z}[G/K] \to \mathbb{Z}[G/H]$.
- $\pi(I_K^H)$ takes $gH \mapsto \sum_{h \in H/K} ghK$.
- $\pi(c_x)$ takes $gH \mapsto gxH^x$.

If M is an $\mathcal{H}_{\mathcal{F}}$ -module then it is straightforward to check that $M \circ \pi$ is a $\mathcal{M}_{\mathcal{F}}$ module, see for example [**Tam89**, p.809] for a proof. Moreover, every cohomological Mackey functor $M : \mathcal{M}_{\mathcal{F}} \to \mathbf{R}$ -Mod factors through the map π , this is the main result in [**Yos83**], see also [**Web00**, §7]. Thus we may pass freely between cohomological Mackey functors and modules over $\mathcal{H}_{\mathcal{F}}$.

LEMMA 4.1.21. [Yos83, Lemma 3.1'] There is an isomorphism for any finite subgroups H and K of G,

$$R[H\backslash G/K] \cong R[G/H, G/K]_{\mathcal{H}_{\mathcal{F}}}.$$

Under this identification, morphism composition is given by

$$(HxK) \cdot (KyL) = \sum_{z \in H \setminus G/L} |(HxK \cap zLy^{-1}K)/K| (HzL).$$

REMARK 4.1.22. The identification in the lemma above relates to the usual definition of $R[G/H, G/K]_{\mathcal{H}_{\mathcal{F}}}$ as $\operatorname{Hom}_{RG}(R[G/H], R[G/H])$ with the isomorphism

$$\begin{split} \psi: R[H\backslash G/K] & \stackrel{\cong}{\longrightarrow} \operatorname{Hom}_{RG}(R[G/H], R[G/K]) \\ HxK & \longmapsto \left(gH \longmapsto \sum_{u \in H/(H \cap xKx^{-1})} guxK\right) \end{split}$$

Notice that ψ satisfies

$$\psi((HxK) \cdot (KxL)) = \psi(KxL) \circ \psi(HxK).$$

LEMMA 4.1.23. If $\alpha: G/L \longrightarrow G/K$ is the G-map $L \mapsto xK$ then the induced map α_* on $R[G/H, -]_{\mathcal{H}_F}$ can be written as

$$\alpha_* : R[H \setminus G/L] \longrightarrow R[H \setminus G/K]$$
$$(HzL) \longmapsto \sum_{y \in H \cap K^{(zx)^{-1}}/H \cap L^{z^{-1}}} (HyzxK).$$

PROOF. We calculate

$$\psi\left(\sum_{y\in H\cap K^{(zx)^{-1}}/H\cap L^{z^{-1}}}HyzxK\right)$$
$$=\left(H\longmapsto\sum_{y\in H\cap K^{(zx)^{-1}}/H\cap L^{z^{-1}}}\sum_{u\in H/H\cap K^{(yzx)^{-1}}}uyzxK\right)$$
$$=\left(H\longmapsto\sum_{u\in H/H\cap L^{z^{-1}}}uzxL\right)$$

which is exactly $\alpha_*(\psi(HzL))$. The final equality comes from the fact that $y \in K^{(zx)^{-1}}$ so $K^{(yzx)^{-1}} = K^{(zx)^{-1}}$.

4.1.2.1. Explicit description of π . Using the identification of Lemma 4.1.21, for any $K \leq H$, we can describe π as follows.

- $\pi(R_K^H) = KH$, since according to Lemma 4.1.21, KH corresponds to the morphism $gK \mapsto gH$, which is exactly Thévenaz and Webb's description of $\pi(R_K^H)$.
- $\pi(I_K^H) = HK$, as according to Lemma 4.1.21, HK corresponds to the morphism $gH \mapsto \sum_{u \in H/K} uK$, which is Thévenaz and Webb's description of $\pi(I_K^H)$.
- $\pi(c_x) = HxH^x$, similarly to the above because HxH^x corresponds to the morphism $gH \mapsto gxH^x$.

It is interesting to write down the effect of π on a basic morphism

$$m = \begin{pmatrix} G/L \\ \swarrow & \swarrow \\ G/H & G/K \end{pmatrix}$$

This morphism may be rewritten as

$$\begin{pmatrix} \alpha_1 & G/L \\ & \swarrow & & \\ G/H & & G/L \end{pmatrix} \circ \begin{pmatrix} \alpha_1 & G/L & \alpha_x \\ & \swarrow & & \\ G/L & & G/L^x \end{pmatrix} \circ \begin{pmatrix} \alpha_1 & G/L^x & \alpha_1 \\ & \swarrow & & \\ G/L^x & & G/K \end{pmatrix}$$

So,

$$m = R_{L^x}^K \circ c_x \circ I_L^H.$$

Using the definition of π in [**TW95**, §16], $\pi(m)$ maps

π

$$(m) = \pi (R_{L^x}^K \circ c_x \circ I_L^H)$$

$$= \left(H \mapsto \sum_{h \in H/L} hxK \right)$$

$$= \left(H \mapsto \sum_{h \in H/H \cap K^{x^{-1}}} \sum_{y \in H \cap K^{x^{-1}}/L} hyxK \right)$$

$$= \left(H \mapsto \sum_{y \in H \cap K^{x^{-1}}/L} \sum_{h \in H/H \cap K^{(yx)^{-1}}} hyxK \right)$$

$$= \sum_{y \in H \cap K^{x^{-1}}/L} (HyxK)$$

$$= |H \cap K^{x^{-1}} : L|(HxK).$$

In summary,

$$\pi \begin{pmatrix} G/L \\ \swarrow \\ G/H \\ G/K \end{pmatrix} = |H \cap K^{x^{-1}} : L|(HxK).$$

4.1.2.2. Homology and cohomology. In Section 4.3 we will prove results similar to Proposition 4.1.13 and Corollary 4.1.14, showing that inducing a projective resolution of <u>R</u> by projective $\mathcal{O}_{\mathcal{F}}$ -modules yields a projective resolution of R^- by projective $\mathcal{H}_{\mathcal{F}}$ -modules. For any group G, we define the cohomology and homology functors $H^*_{\mathcal{H}_{\mathcal{F}}}(G, -)$ and $H^{\mathcal{H}_{\mathcal{F}}}_*(G, -)$ as

$$H^*_{\mathcal{H}_{\mathcal{F}}}(G, M) = \operatorname{Ext}^*_{\mathcal{H}_{\mathcal{F}}}(R^-, M)$$
$$H^{\mathcal{H}_{\mathcal{F}}}_*(G, A) = \operatorname{Tor}^*_{\mathcal{H}_{\mathcal{F}}}(R^-, A)$$

where M is any contravariant $\mathcal{H}_{\mathcal{F}}$ -module and A is any covariant $\mathcal{H}_{\mathcal{F}}$ -module. In Proposition 4.3.8 we show that there is an isomorphism

$$H^n_{\mathcal{H}_{\mathcal{F}}}(G, M) \cong H^n_{\mathcal{O}_{\mathcal{F}}}(G, \operatorname{Res}_{\pi \circ \sigma} M).$$

The $\mathcal{H}_{\mathcal{F}}$ cohomological dimension of a group G, denoted $\mathcal{H}_{\mathcal{F}} cd G$, is defined to be the length of the shortest projective resolution of R^- by $\mathcal{H}_{\mathcal{F}}$ -modules, or equivalently

$$\mathcal{H}_{\mathcal{F}}$$
cd $G = \sup\{n : H^n_{\mathcal{H}_{\mathcal{F}}}(G, M) \neq 0\}, M$ some $\mathcal{H}_{\mathcal{F}}$ -module.}

Note that in [Deg13a] the $\mathcal{H}_{\mathcal{F}}$ cohomological dimension is defined as

$$\mathcal{H}_{\mathcal{F}} \operatorname{cd} G = \sup\{n : H^n_{\mathcal{O}_{\mathcal{F}}}(G, \operatorname{Res}_{\pi \circ \sigma} M) \neq 0, M \text{ some } \mathcal{H}_{\mathcal{F}} \operatorname{-module.}\}$$

These two definitions are equivalent by the isomorphism of Proposition 4.3.8 mentioned above.

We say G is $\mathcal{H}_{\mathcal{F}} FP_n$ if there exists a projective $\mathcal{H}_{\mathcal{F}}$ -module resolution of R^- , finitely generated up to degree n.

4.2. FP_n conditions for Mackey functors

As far as we are aware, there are no results in the literature on the conditions $\mathcal{M}_{\mathcal{F}} FP_n$. We show in this section that the conditions $\mathcal{M}_{\mathcal{F}} FP_n$ and $\mathcal{O}_{\mathcal{F}} FP_n$ are equivalent. From this point on, unless otherwise stated, all results are valid over any commutative ring R.

PROPOSITION 4.2.1. If G is $\mathcal{O}_{\mathcal{F}} FP_n$ then G is $\mathcal{M}_{\mathcal{F}} FP_n$.

PROOF. Combine Proposition 4.1.13 with the fact that induction preserves finite generation (Proposition 2.3.2(2)). \Box

Recall that G is $\mathcal{O}_{\mathcal{F}} FP_0$ if and only if \mathcal{F} has finitely many conjugacy classes (Corollary 3.6.4). In the lemmas below \mathcal{F}/G denotes the poset of conjugacy classes in \mathcal{F} , ordered by subconjugation. We write $H \leq_G K$ if H is subconjugate to K.

LEMMA 4.2.2. G is $\mathcal{M}_{\mathcal{F}}FP_0$ if and only if G is $\mathcal{O}_{\mathcal{F}}FP_0$.

PROOF. We prove first that if G is $\mathcal{M}_{\mathcal{F}}FP_0$ then \mathcal{F}/G has a finite cofinal subset, since \mathcal{F} is a subfamily of the family of finite subgroups this implies that \mathcal{F}/G is finite.

Let f be an $\mathcal{M}_{\mathcal{F}}$ -module morphism

$$f: R[-, G/K]_{\mathcal{M}_{\mathcal{F}}} \longrightarrow B^G \cong R[-, G/G]_{\mathcal{M}_{\mathcal{F}}}.$$

Firstly, we claim that the element m of $R[G/S, G/G]_{\mathcal{M}_{\mathcal{F}}}$ given by

$$m = \left(G/S \xleftarrow{\text{id}} G/S \longrightarrow G/G\right)$$

cannot be in the image of f(G/S) unless S is subconjugate to K. Assume for a contradiction that S is not subconjugate to K and assume m is in the image of f(G/S). Thus $m = f(G/S)\varphi$ for some $\varphi \in [G/S, G/K]_{\mathcal{M}_{\mathcal{F}}}$. Thinking of f as a natural transformation gives the commutative diagram below

$$R[G/S, G/K]_{\mathcal{M}_{\mathcal{F}}} \xrightarrow{f(G/S)} R[G/S, G/G]_{\mathcal{M}_{\mathcal{F}}}$$

$$\uparrow^{\varphi^{*}} \qquad \uparrow^{\varphi^{*}}$$

$$R[G/K, G/K]_{\mathcal{M}_{\mathcal{F}}} \xrightarrow{f(G/K)} R[G/K, G/G]_{\mathcal{M}_{\mathcal{F}}}$$

where

$$\begin{split} m &= f(G/S)\varphi \\ &= f(G/S) \circ \varphi^* \operatorname{id}_{[G/K, G/K]_{\mathcal{M}_{\mathcal{F}}}} \\ &= \big(\varphi^* \circ f(G/K)\big) (\operatorname{id}_{[G/K, G/K]_{\mathcal{M}_{\mathcal{F}}}}). \end{split}$$

Let $f(G/K)(\operatorname{id}_{[G/K,G/K]_{\mathcal{M}_{\mathcal{F}}}}) = \sum_{i} r_{i}x_{i}$, where $r_{i} \in R$ and the x_{i} are basic morphisms in $R[G/K, G/G]_{\mathcal{M}_{\mathcal{F}}}$. Similarly, let $\varphi = \sum_{j} s_{j}y_{j}$ for $s_{j} \in R$ and where the y_{j} are basic morphisms in $R[G/S, G/K]_{\mathcal{M}_{\mathcal{F}}}$. By assumption we have that

$$m = \varphi^* \sum_i r_i x_i$$
$$= \sum_i r_i x_i \circ \sum_j s_j y_j$$
$$= \sum_{i,j} (r_i s_j) x_i \circ y_j.$$

There must exist some i and j for which $x_i \circ y_j$ is a morphism which, when written as a sum of basic morphisms, has one component some multiple of m. We calculate $x_i \circ y_j$ for this i and j. Write x_i and y_j in their standard forms as below,

$$x_i = \left(G/K \longleftarrow G/L_i \longrightarrow G/G\right)$$
$$y_j = \left(G/S \longleftarrow G/J_j \longrightarrow G/K\right).$$

Their composition is (see Lemma 4.1.2)

$$x_i \circ y_j = \sum_k \begin{pmatrix} G/X_k \\ \swarrow & \searrow \\ G/J_j & G/L_i \\ \swarrow & \swarrow & \swarrow \\ G/S & G/K & G/G \end{pmatrix}$$

where X_k is some finite subgroup of G which is subconjugate to both L_i and J_j .

We claim $|J_j|$ is strictly smaller than |S|. Since J_j is subconjugate to S we have $|J_j| \leq |S|$. If the cardinalities were equal then S and J_j would be conjugate, but J_j is subconjugate to K whereas by assumption S is not subconjugate to K.

Since $|X_k| \leq |J_j| \leq |S|$, the subgroup X_k cannot be conjugate to S. This contradicts our earlier assertion that $x_i \circ y_j$ when written as a sum of basic morphisms, has one component some multiple of m. Thus, for m to be in the image of f(G/S), S must be subconjugate to K.

Now, if G is $\mathcal{M}_{\mathcal{F}} FP_0$ then B^G admits an epimorphism from some finitely generated free

$$\bigoplus_{i\in I} R[-,G/K_i]_{\mathcal{M}_{\mathcal{F}}} \longrightarrow B^G.$$

As this set I is finite, the argument above implies that all the subgroups in \mathcal{F} are subconjugate to one of a finite collection of subgroups in \mathcal{F} . Thus there is a finite cofinal subset of \mathcal{F}/G and \mathcal{F}/G is finite.

For the converse, use Proposition 4.2.1.

This remainder of this section is devoted to a proof that for any n, $\mathcal{M}_{\mathcal{F}} FP_n$ implies $\mathcal{O}_{\mathcal{F}} FP_n$. We will assume G is $\mathcal{M}_{\mathcal{F}} FP_0$, equivalently \mathcal{F} contains finitely many conjugacy classes.

In [**HPY13**, 4.9, 4.10], there are the following definitions, for M an $\mathcal{O}_{\mathcal{F}}$ -module

$$D_H M = \operatorname{CoInd}_{R[WH]}^{\mathcal{O}_F} M(G/H)$$

 $j_H : M \longrightarrow D_H M$

where $\operatorname{CoInd}_{R[WH]}^{\mathcal{O}_{\mathcal{F}}}$ denotes coinduction (see Section 2.3 for the definition of coinduction) with the functor $\iota : \mathbb{Z}[WH] \longrightarrow \mathcal{O}_{\mathcal{F}}$. Here we view $\mathbb{Z}[WH]$ as a category with one object and morphisms elements of $\mathbb{Z}[WH]$, then $\mathbb{Z}[WH]$ has property (A) and ι maps the one object to G/H and morphisms to the free abelian group on the automorphisms of G/H in $\mathcal{O}_{\mathcal{F}}$. Equivalently,

$$\operatorname{CoInd}_{R[WH]}^{\mathcal{O}_{\mathcal{F}}} M(G/H) \cong \operatorname{Hom}_{R[WH]}(R[G/H, -]_{\mathcal{O}_{\mathcal{F}}}, M(G/H)).$$

The map j_H is the counit of the adjunction between coinduction and restriction. Since evaluation, coinduction, and counits are all natural constructions, D_H and j_H are natural. Crucially the $\mathcal{O}_{\mathcal{F}}$ -module $D_H M$ extends to a Mackey functor **[HPY13**, Example 4.8]. Also defined are:

$$DM = \prod_{H \in \mathcal{F}/G} D_H M$$
$$CM = \operatorname{CoKer} \left(C \xrightarrow{\prod j_H} DM \right)$$

Again all the constructions are natural and DM extends to a Mackey functor. Naturality means that if M_{λ} , for $\lambda \in \Lambda$, is a directed system of $\mathcal{O}_{\mathcal{F}}$ -modules then DM_{λ} and CM_{λ} form directed systems also.

LEMMA 4.2.3. If M_{λ} is a directed system of $\mathcal{O}_{\mathcal{F}}$ -modules with $\varinjlim M_{\lambda} = 0$ then $\lim DM_{\lambda} = 0$.

PROOF. Since the colimit of M_{λ} is zero, so is the colimit of $M_{\lambda}(G/H)$, and for any $K \in \mathcal{F}$,

$$\underbrace{\lim D_H M_\lambda(G/K)}_{K} = \underbrace{\lim \operatorname{CoInd}}_{R[WH]}^{\mathcal{O}_{\mathcal{F}}} M_\lambda(G/H)(G/K)$$

$$= \underbrace{\lim \operatorname{Hom}}_{R[WH]}(R[G/H, G/K]_{\mathcal{O}_{\mathcal{F}}}, M_\lambda(G/H))$$

$$= \underbrace{\lim \operatorname{Hom}}_{R[WH]} \left(\bigoplus_{i \in I} R[WH/WH_i], M_\lambda(G/H) \right)$$

where the last line is Lemma 3.1.5, the indexing set I is finite and WH_i is a finite subgroup of WH. Hence

$$\varinjlim \operatorname{Hom}_{R[WH]} \left(\bigoplus_{i \in I} R[WH/WH_i], M_{\lambda}(G/H) \right) \cong \bigoplus_{i \in I} \varinjlim \operatorname{Hom}_{R[WH]} (R[WH/WH_i], M_{\lambda}(G/H)) \cong \bigoplus_{i \in I} \varinjlim \operatorname{Hom}_{R[WH_i]} (R, M_{\lambda}(G/H)) = 0$$

where the final zero is by the Bieri–Eckmann criterion (Theorem 2.5.1), since R is $R[WH_i]$ -finitely generated. Thus

$$\varinjlim DM_{\lambda}(G/K) = \varinjlim \prod_{H \in \mathcal{F}/G} D_H M_{\lambda}(G/K)$$
$$= \prod_{H \in \mathcal{F}/G} \varinjlim D_H M_{\lambda}(G/K)$$
$$= 0$$

where the commuting of the product and the colimit is because the product is finite $(\mathcal{F}/G \text{ is assumed finite})$.

LEMMA 4.2.4. If M_{λ} is a directed system of $\mathcal{O}_{\mathcal{F}}$ -modules with $\varinjlim M_{\lambda} = 0$ then $\varinjlim CM_{\lambda} = 0$.

PROOF. There is a natural short exact sequence for each λ

 $0 \longrightarrow M_{\lambda} \longrightarrow DM_{\lambda} \longrightarrow CM_{\lambda} \longrightarrow 0.$

Since the colimit of the left hand and centre term are zero (Lemma 4.2.3), and colimits are exact in the category of $\mathcal{O}_{\mathcal{F}}$ -modules, so $\lim CM_{\lambda} = 0$ also.

PROPOSITION 4.2.5. If G is $\mathcal{M}_{\mathcal{F}} FP_n$ then G is $\mathcal{O}_{\mathcal{F}} FP_n$.

PROOF. Let G be of type $\mathcal{M}_{\mathcal{F}} FP_n$ and let M_{λ} , for $\lambda \in \Lambda$, be a directed system of $\mathcal{O}_{\mathcal{F}}$ -modules with colimit zero. Following the notation in [**Deg13a**], we define

$$C^{0}M_{\lambda} = M_{\lambda}$$
$$C^{i}M_{\lambda} = CC^{i-1}M_{\lambda}$$

for all natural numbers $i \ge 0$ and all $\lambda \in \Lambda$. There are short exact sequences of directed systems,

$$0 \longrightarrow C^i M_{\lambda} \longrightarrow DC^i M_{\lambda} \longrightarrow C^{i+1} M_{\lambda} \longrightarrow 0$$

all the terms of which have colimit zero.

As G is assumed $\mathcal{M}_{\mathcal{F}} FP_n$ and $DC^i M_{\lambda}$ extends to a Mackey functor for all i, the Bieri–Eckmann criterion (Theorem 2.5.1) gives that for all $m \leq n$,

$$\varinjlim H^m_{\mathcal{O}_{\mathcal{F}}}(G, DC^i M_\lambda) = 0$$

and thus using exactness of colimits and the long exact sequence associated to cohomology gives that for all non-negative integers m and i,

$$\varinjlim H^m_{\mathcal{O}_{\mathcal{F}}}(G, C^{i+1}M_{\lambda}) = \varinjlim H^{m+1}_{\mathcal{O}_{\mathcal{F}}}(G, C^iM_{\lambda}).$$

So,

$$\varinjlim H^m_{\mathcal{O}_{\mathcal{F}}}(G, M_{\lambda}) = \varinjlim H^{m-1}_{\mathcal{O}_{\mathcal{F}}}(G, C^1 M_{\lambda}) \cong \cdots \cong \varinjlim H^0_{\mathcal{O}_{\mathcal{F}}}(G, C^m M_{\lambda}) \cong 0$$

where the zero is from the Bieri–Eckmann criterion (Theorem 2.5.1), because G is assumed $\mathcal{M}_{\mathcal{F}}FP_0$ hence $\mathcal{O}_{\mathcal{F}}FP_0$ by Lemma 4.2.2. Using the Bieri–Eckmann criterion again, G is $\mathcal{O}_{\mathcal{F}}FP_n$.

COROLLARY 4.2.6. The conditions $\mathcal{O}_{\mathcal{F}} FP_n$ and $\mathcal{M}_{\mathcal{F}} FP_n$ are equivalent.

PROOF. Combine Propositions 4.2.1 and 4.2.5.

4.3. Homology and cohomology of cohomological Mackey functors

The main result of this section is Proposition 4.3.6, that we may induce projective $\mathcal{O}_{\mathcal{F}}$ -module resolutions of <u>R</u> to projective $\mathcal{H}_{\mathcal{F}}$ -module resolutions of R^- . The following diagram shows the relationship between the different induction functors we will be using (for a small category \mathfrak{C} , we denote by \mathfrak{C} -Mod the category of contravariant \mathfrak{C} -modules).



LEMMA 4.3.1. For any $L \in \mathcal{F}$, there is an isomorphism of covariant $\mathcal{O}_{\mathcal{F}}$ -modules

$$\operatorname{Res}_{\pi\circ\sigma} R[G/L, -]_{\mathcal{H}_{\mathcal{F}}} \cong \operatorname{Hom}_{RL}(R, R[G/1, -]_{\mathcal{O}_{\mathcal{F}}}).$$

PROOF. If H is a subgroup in \mathcal{F} , then evaluating the left hand side at G/H yields $R[G/L, G/H]_{\mathcal{H}_{\mathcal{F}}}$ while evaluating the right hand side at G/H yields

$$\operatorname{Hom}_{RL}(R, R[G/H]) \cong \operatorname{Hom}_{RG}(RG \otimes_{RL} R, R[G/H])$$
$$\cong \operatorname{Hom}_{RG}(R[G/L], R[G/H])$$
$$\cong R[G/L, G/H]_{\mathcal{H}_{\mathcal{F}}}$$

where the first isomorphism is [**Bro94**, p.63 (3.3)]. If $\alpha_x : G/H \to G/K$ is the *G*-map $H \mapsto xK$ then, looking at the left hand side,

$$\operatorname{Res}_{\pi \circ \sigma} R[G/L, -]_{\mathcal{H}_{\mathcal{F}}}(\alpha_x) = R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}(c_x \circ R_H^{K^{x^{-1}}})$$
$$\cong R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}(c_x) \circ R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}(R_H^{K^{x^{-1}}}).$$

But $R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}(R_H^{K^{x^{-1}}})$ is post-composition with the *G*-map

 $\alpha_1: G/H \to G/K^{x^{-1}}$

and $R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}(c_x)$ is post-composition with the G-map

$$\alpha_x: G/K^{x^{-1}} \to G/K.$$

In summary, $\operatorname{Res}_{\pi\circ\sigma} R[G/L, -]_{\mathcal{H}_{\mathcal{F}}}(\alpha_x)$ is the map:

$$\operatorname{Hom}_{RG}(R[G/L], R[G/H]) \longrightarrow \operatorname{Hom}_{RG}(R[G/L], R[G/K])$$
$$f \longmapsto \alpha_x \circ f$$

Now, the right hand side, recall that

$$R[G/L, -]_{\mathcal{O}_{\mathcal{F}}}(\alpha_x) : f \longmapsto \alpha_x \circ f$$

so $\operatorname{Hom}_{RL}(R, R[G/1, -]_{\mathcal{O}_{\mathcal{F}}})(\alpha_x)$ is the map:

$$\begin{split} \operatorname{Hom}_{RL}(R,R[G/H]) &\longrightarrow \operatorname{Hom}_{RL}(R,R[G/K]) \\ f &\longmapsto \alpha_x \circ f \end{split}$$

Showing the left and right hand sides agree on morphisms.

Recall that $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}}$ denotes induction with the functor $\iota : \mathbb{Z}G \longrightarrow \mathcal{O}_{\mathcal{F}}$, where we view $\mathbb{Z}G$ as the single object category whose morphisms are elements of $\mathbb{Z}G$ and ι maps the single object to G/1. Equivalently for an RG-module M,

$$\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} M \cong R[-, G/1]_{\mathcal{O}_{\mathcal{F}}} \otimes_{RG} M.$$

LEMMA 4.3.2. The functor $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}}$ is exact.

PROOF. This is because for any $H \in \mathcal{F}$,

$$\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} M(G/H) = \begin{cases} M & \text{if } H = 1\\ 0 & \text{else.} \end{cases}$$

LEMMA 4.3.3. For any finite subgroup H of G, the $\mathcal{O}_{\mathcal{F}}$ -module $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} \operatorname{Ind}_{RH}^{RG} R$ is of type $\mathcal{O}_{\mathcal{F}} \operatorname{FP}_{\infty}$.

PROOF. Since R is $\operatorname{FP}_{\infty}$ as a RH module, $\operatorname{Ind}_{RH}^{RG} R$ is of type $\operatorname{FP}_{\infty}$ over RG. Choose a finite type free resolution F_* of $\operatorname{Ind}_{RH}^{RG} R$ by RG-modules, by Lemma 4.3.2, $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} F_*$ is a finite type free resolution of $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} \operatorname{Ind}_{RH}^{RG} R$ by $\mathcal{O}_{\mathcal{F}}$ -modules.

LEMMA 4.3.4. If N is a projective $\mathcal{O}_{\mathcal{F}}$ -module and $H \in \mathcal{F}$, there is an isomorphism

$$N \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Res}_{\pi \circ \sigma} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}} \cong \operatorname{Hom}_{RH}(R, N(G/1)).$$

PROOF. The adjointness of induction and restriction gives an isomorphism of $\mathcal{O}_{\mathcal{F}}$ -modules, for any $\mathcal{O}_{\mathcal{F}}$ -module N,

$$\operatorname{Hom}_{RH}(R, N(G/1)) \cong \operatorname{Hom}_{RG}(\operatorname{Ind}_{RH}^{RG} R, N(G/1))$$
$$\cong \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} \operatorname{Ind}_{RH}^{RG} R, N).$$

There is a chain of isomorphisms,

$$N \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Res}_{\pi \circ \sigma} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$$

$$\cong N \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}_{RH}(R, R[G/1, -]_{\mathcal{O}_{\mathcal{F}}})$$

$$\cong N(-) \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} \operatorname{Ind}_{RH}^{RG} R(?), R[?, -]_{\mathcal{O}_{\mathcal{F}}})$$

$$\cong \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}} \operatorname{Ind}_{RH}^{RG} R(?), N(?))$$

$$\cong \operatorname{Hom}_{RH}(R, N(G/1))$$

where the first isomorphism is Lemma 4.3.1 and the second and fourth are the adjoint isomorphism mentioned above. The third isomorphism is from Lemma 3.9.5 for which we need that $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}}}\operatorname{Ind}_{RH}^{RG}R$ is finitely generated, but this is implied by Lemma 4.3.3.

Recall from Example 4.1.16 the definition of the fixed point functor. For the constant RG-module R the fixed point functor R^- can be described explicitly as $R^H = R$ for all $H \in \mathcal{F}$, and on morphisms,

$$R^{-}(R_{K}^{H}) = \mathrm{id}_{R}$$
$$R^{-}(I_{K}^{H}) = (r \mapsto |H:K|r)$$
$$R^{-}(c_{g}) = \mathrm{id}_{R}.$$

LEMMA 4.3.5. $\operatorname{Ind}_{\pi \circ \sigma} \underline{R} \cong R^-$.

PROOF. The proof is split into two parts, first we check that the two functors agree on objects, then we check they agree on morphisms. Throughout the proof

 $H, K \text{ and } L \text{ are elements of } \mathcal{F}.$ If $\alpha : G/L \to G/K$ is a G-map then we will write α_* for the induced map

$$\alpha_* : \operatorname{Hom}_{RG}(R[G/H], R[G/L]) \longrightarrow \operatorname{Hom}_{RG}(R[G/H], R[G/K])$$

and also for the induced map

$$\alpha_*: R[H \backslash G/L] \longrightarrow R[H \backslash G/K]$$

where $R[H \setminus G/L]$ is identified with $\operatorname{Hom}_{RG}(R[G/H], R[G/L])$ using the isomorphism ψ of Remark 4.1.22.

The functors $\operatorname{Ind}_{\pi\circ\sigma} \underline{R}$ and R^- agree on objects:

For any subgroup $H \in \mathcal{F}$,

$$\operatorname{Ind}_{\pi\circ\sigma}\underline{R}(G/H) = \underline{R} \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Res}_{\pi\circ\sigma} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$$
$$\cong \underline{R} \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Hom}_{RG}(R[G/H], R[G/1, -]_{\mathcal{O}_{\mathcal{F}}})$$
$$\cong \bigoplus_{K \in \mathcal{F}} \operatorname{Hom}_{RG}(R[G/H], R[G/K]) \Big/ \stackrel{\alpha : G/L \to G/K \text{ any } G \text{ map}}{\underset{x_{L} \in \operatorname{Hom}_{RG}(R[G/H], R[G/K])}{\underset{x_{L} \in \operatorname{Hom}_{RG}(R[G/H], R[G/L])}}$$
$$\cong \bigoplus_{K \in \mathcal{F}} R[H \backslash G/K] \Big/ \stackrel{(HxK) \sim \alpha_{*}(HxL)}{\alpha : G/L \to G/K \text{ any } G \text{ map}}$$

where the first isomorphism is Lemma 4.3.1 and the last is Lemma 4.1.21. Let $(HxK) \in R[H \setminus G/K]$ be an arbitrary element, and consider the *G*-map

$$\alpha_x : G/(H \cap K^{x^{-1}}) \longrightarrow G/K$$
$$(H \cap K^{x^{-1}}) \longmapsto xK.$$

Then, using Lemma 4.1.23 we calculate

$$(\alpha_x)_*\left(H1(H\cap K^{x^{-1}})\right) = (HxK)$$

so in $\operatorname{Ind}_{\pi\circ\sigma} \underline{R}(G/H)$, the elements $[H \cdot x \cdot K]$ and $[H \cdot 1 \cdot H \cap K^{x^{-1}}]$ are equal, where [-] denotes an equivalence class of elements under the relation \sim . Similarly if $K \leq H$ then $[H \cdot 1 \cdot K] = |H : K|[H \cdot 1 \cdot H]$ since if $\alpha_1 : G/K \to G/H$ is the projection, then using Lemma 4.1.23 again,

$$\alpha_{1*}(H1K) = |H:K|(H1H).$$

Combining the two facts proved above,

$$(\star) \qquad \qquad [H \cdot x \cdot K] = |H : H \cap K^{x^{-1}}|[H \cdot 1 \cdot H].$$

In particular, any element $[H \cdot x \cdot K]$ is equal to some multiple of $[H \cdot 1 \cdot H]$, so

Ind_{$$\pi \circ \sigma$$} R(G/H) \cong R.

The functors $\operatorname{Ind}_{\pi\circ\sigma} \underline{R}$ and R^- agree on morphisms:

Recall from Remark 4.1.6 that we must only check this for the morphisms R_K^H , I_K^H and c_x .

Following the generator $[H \cdot 1 \cdot H]$ up the chain of isomorphisms at the beginning of the proof shows the element

$$1 \otimes \operatorname{id}_{R[G/H]} \in R \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Res}_{\pi \circ \sigma} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$$

generates $\operatorname{Ind}_{\pi \circ \sigma} \underline{R}(G/H) \cong R$, where

$$\operatorname{id}_{R[G/H]} \in \operatorname{Hom}_{RG}(R[G/H], R[G/H]) \cong R[G/H, G/H]_{\mathcal{H}_{\mathcal{F}}}.$$

For some subgroup $K \in \mathcal{F}$ with $K \leq H$,

$$\operatorname{Ind}_{\pi\circ\sigma}\underline{R}(R_K^H): 1\otimes \operatorname{id}_{R[G/H]} \mapsto 1\otimes \pi$$

where $\pi : R[G/K] \mapsto R[G/H]$ is the projection map. Following this back down the chain of isomorphisms at the beginning of the proof, gives the element $[K \cdot 1 \cdot H]$. Using $(\star), [K \cdot 1 \cdot H] = [K \cdot 1 \cdot K]$, so $\operatorname{Ind}_{\pi \circ \sigma} \underline{R}(R_K^H)$ is the identity on R, as required.

Similarly, for some $L \in \mathcal{F}$ with $H \leq L$, we calculate

$$\operatorname{Ind}_{\pi\circ\sigma}\underline{R}(I_H^L): 1\otimes \operatorname{id}_{R[G/H]} \longmapsto 1\otimes t_{L/H}$$

where $t_{L/H} \in \operatorname{Hom}_{RG}(R[G/L], R[G/H])$ denotes the map $L \mapsto \sum_{l \in L/H} lH$. Following this element back down the chain of isomorphisms we get the element $[L \cdot 1 \cdot H]$, which by (\star) is equal to $|L : H|[H \cdot 1 \cdot H]$. Thus $\operatorname{Ind}_{\pi \circ \sigma} \underline{R}(I_H^L)$ acts as multiplication by |L : H| on R, as required.

For any element $x \in G$, we calculate

 $\operatorname{Ind}_{\pi \circ \sigma} \underline{R}(c_x) : 1 \otimes \operatorname{id}_{R[G/H]} \longmapsto 1 \otimes \gamma_x$

where $\gamma_x \in \operatorname{Hom}_{RG}(R[G/H^{x^{-1}}], R[G/H])$ is the map $H^{x^{-1}} \mapsto xH$. Following this down the chain of isomorphisms we get the element $[H^{x^{-1}} \cdot x \cdot H]$, which by (\star) is equal to $[H^{x^{-1}} \cdot 1 \cdot H^{x^{-1}}]$. Thus $\operatorname{Ind}_{\pi \circ \sigma} \underline{R}(c_x)$ acts as the identity on R, as required.

The next proposition should be compared with Proposition 4.1.13. Recall that a chain complex is \mathcal{F} -split if it splits when restricted to RH for all $H \in \mathcal{F}$.

PROPOSITION 4.3.6. Induction with $\pi \circ \sigma$ takes projective $\mathcal{O}_{\mathcal{F}}$ -module resolutions of <u>R</u> to projective $\mathcal{H}_{\mathcal{F}}$ -module resolutions of R^- .

PROOF. Let P_* be a projective resolution of <u>R</u> by $\mathcal{O}_{\mathcal{F}}$ -modules, then by Lemma 4.3.4,

$$\operatorname{Ind}_{\pi\circ\sigma} P_*(G/H) = P_* \otimes_{\mathcal{O}_{\mathcal{F}}} \operatorname{Res}_{\pi\circ\sigma} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$$
$$\cong \operatorname{Hom}_{RH}(R, P_*(G/1)).$$

76

$$\operatorname{Ind}_{\pi\circ\sigma} P_* \longrightarrow R^-$$

Induction preserves projectives, so we must show only that the above is exact. Since induction is right exact, it is necessarily exact at degree -1 and degree 0. Evaluating at G/H gives the resolution

$$\operatorname{Hom}_{RH}(R, P_*(G/1)) \longrightarrow R.$$

By [**Nuc00**, Theorem 3.2], the resolution $P_*(G/1)$ is \mathcal{F} -split. Since $\operatorname{Hom}_{RH}(R, -)$ preserves the exactness of RH-split complexes, $\operatorname{Hom}_{RH}(R, P_*(G/1))$ is exact at position i for all $i \geq 1$, completing the proof.

REMARK 4.3.7. The proposition above may not hold with <u>R</u> replaced by an arbitrary $\mathcal{O}_{\mathcal{F}}$ -module M, as a resolution of M by projective $\mathcal{O}_{\mathcal{F}}$ -modules will not in general split when evaluated at G/1.

Recall that in Section 4.1.2.2 we defined, for any $\mathcal{H}_{\mathcal{F}}$ -module M,

$$H^*_{\mathcal{H}_{\mathcal{T}}}(G, M) \cong \operatorname{Ext}^*_{\mathcal{H}_{\mathcal{T}}}(R^-, M).$$

There is an analogue of Corollary 4.1.14:

PROPOSITION 4.3.8. For any $\mathcal{H}_{\mathcal{F}}$ -module M and any natural number n,

$$H^n_{\mathcal{H}_{\mathcal{F}}}(G, M) = H^n_{\mathcal{O}_{\mathcal{F}}}(G, \operatorname{Res}_{\pi \circ \sigma} M).$$

PROOF. Let P_* be a projective $\mathcal{O}_{\mathcal{F}}$ -module resolution of <u>R</u>, then

$$H^{n}_{\mathcal{O}_{\mathcal{F}}}(G, \operatorname{Res}_{\pi \circ \sigma} M) = H^{n} \operatorname{Hom}_{\mathcal{O}_{\mathcal{F}}}(P_{*}, \operatorname{Res}_{\pi \circ \sigma} M)$$
$$\cong H^{n} \operatorname{Hom}_{\mathcal{H}_{\mathcal{F}}}(\operatorname{Ind}_{\pi \circ \sigma} P_{*}, M)$$
$$= H^{n}_{\mathcal{H}_{\mathcal{F}}}(G, M)$$

where the isomorphism is adjoint isomorphism between induction and restriction and $\operatorname{Ind}_{\pi\circ\sigma} P_*$ is a projective $\mathcal{H}_{\mathcal{F}}$ -module resolution of R^- by Proposition 4.3.6.

4.4. FP_n conditions for cohomological Mackey functors

The main result of this section is Theorem 4.4.1 below. For a detailed construction of \mathcal{F} -cohomology and the condition $\mathcal{F}FP_n$ see [**Nuc99**], for an overview see Section 4.1.

THEOREM 4.4.1. For any ring R, if G is $\mathcal{H}_{\mathcal{F}}FP_n$ then G is $\mathcal{F}FP_n$. If R is Noetherian and G is $\mathcal{F}FP_n$ then G is $\mathcal{H}_{\mathcal{F}}FP_n$.

The proof is contained in Sections 4.4.1 and 4.4.2.

PROPOSITION 4.4.2. If G is $\mathcal{M}_{\mathcal{F}}FP_n$ then G is $\mathcal{H}_{\mathcal{F}}FP_n$.

PROOF. Combining Corollary 4.1.14 and Proposition 4.3.8 shows that for all groups G, and all non-negative integers i,

$$H^i_{\mathcal{H}_{\mathcal{F}}}(G,-) \cong H^i_{\mathcal{M}_{\mathcal{F}}}(G,\operatorname{Res}_{\pi}-).$$

Let G be a group of type $\mathcal{M}_{\mathcal{F}} \mathrm{FP}_n$ and let M_{λ} , for $\lambda \in \Lambda$, be a directed system of $\mathcal{H}_{\mathcal{F}}$ -modules with colimit zero. Then the colimit of $\mathrm{Res}_{\pi} M_{\lambda}$ is zero also and by the Bieri–Eckmann criterion (Theorem 2.5.1), for any $i \leq n$,

$$\varinjlim_{\Lambda} H^{i}_{\mathcal{H}_{\mathcal{F}}}(G, M_{\lambda}) \cong \varinjlim_{\Lambda} H^{i}_{\mathcal{M}_{\mathcal{F}}}(G, \operatorname{Res}_{\pi} M_{\lambda})$$
$$\cong 0.$$

Applying the Bieri–Eckmann criterion again shows G is of type $\mathcal{H}_{\mathcal{F}}FP_n$.

PROPOSITION 4.4.3. If G is $\mathcal{H}_{\mathcal{F}}FP_n$ then G is FP_n .

PROOF. Let $P_* \longrightarrow R^-$ be a resolution of R^- by free $\mathcal{H}_{\mathcal{F}}$ -modules, finitely generated up to degree n. Since the finitely generated free $\mathcal{H}_{\mathcal{F}}$ -modules are fixed point functors of finitely generated permutation modules with stabilisers in \mathcal{F} , evaluating at G/1 gives a resolution of R by RG-modules of type FP_{∞} and a standard dimension shifting argument completes the proof.

So there is a chain of implications:

$$\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n \Leftrightarrow \mathcal{M}_{\mathcal{F}} \mathrm{FP}_n \Rightarrow \mathcal{H}_{\mathcal{F}} \mathrm{FP}_n \Rightarrow \mathrm{FP}_n + \left\{ \begin{smallmatrix} G \text{ has finitely many} \\ \mathrm{conjugacy \ classes \ of} \\ \mathrm{finite} \ p \text{-subgroups in } \mathcal{F} \end{smallmatrix} \right\}$$

Where the final implication is [**LN10**, Proposition 4.2], where it is proved that G is $\mathcal{F}FP_0$ if and only if G has finitely many conjugacy classes of finite p-subgroups in \mathcal{F} , for all primes p. It is conjectured in the same paper that a group G of type FP_{∞} with finitely many conjugacy classes of finite p-subgroups in \mathcal{F} is $\mathcal{F}FP_{\infty}$ [**LN10**, Conjecture 4.3].

Since G is $\mathcal{M}_{\mathcal{F}} FP_0$ if and only if G has finitely many conjugacy classes in \mathcal{F} (Lemma 4.2.2), the implication $\mathcal{M}_{\mathcal{F}} FP_n \Rightarrow \mathcal{H}_{\mathcal{F}} FP_n$ is not reversible.

There are examples due to Leary and Nucinkis of groups which act properly and cocompactly on contractible *G*-CW-complexes but which are not of type $\mathcal{O}_{\mathcal{F}}FP_0$ [**LN03**, Example 3, p.149]. By Remark 4.5.6, these groups are of type $\mathcal{H}_{\mathcal{F}}FP_{\infty}$ showing that $\mathcal{H}_{\mathcal{F}}FP_{\infty} \not\Rightarrow \mathcal{O}_{\mathcal{F}}FP_0$. Leary and Nucinkis also give examples of groups which act properly and cocompactly on contractible *G*-CWcomplexes, are of type $\mathcal{O}_{\mathcal{F}}FP_0$ but which are not $\mathcal{O}_{\mathcal{F}}FP_{\infty}$ [**LN03**, Example 4, p.150]. Hence there can be no implication $\mathcal{H}_{\mathcal{F}}FP_n + \mathcal{O}_{\mathcal{F}}FP_0 \Rightarrow \mathcal{O}_{\mathcal{F}}FP_n$. **4.4.1.** $\mathcal{H}_{\mathcal{F}} FP_n$ implies $\mathcal{F} FP_n$. This section comprises a series of lemmas, building to the proof of Proposition 4.4.11, that for any commutative ring R the condition $\mathcal{H}_{\mathcal{F}} FP_n$ implies the condition $\mathcal{F} FP_n$. Throughout, H and K are arbitrary subgroups in \mathcal{F} .

We say a short exact sequence of RG-modules

$$(\star) \qquad \qquad 0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is H-good if

$$0 \longrightarrow A^H \longrightarrow B^H \longrightarrow C^H \longrightarrow 0$$

is exact. Similarly an exact chain complex C_* is *H*-good if C_*^H is exact. If an exact chain complex is *H*-good for all $H \in \mathcal{F}$ we say it is \mathcal{F} -good. Note that an \mathcal{F} -split exact chain complex is automatically \mathcal{F} -good.

REMARK 4.4.4. If C_*^- is an exact chain complex of fixed point functors then C_* is \mathcal{F} -good.

REMARK 4.4.5. In general being *H*-good is a weaker property than being *RH*-split: Applying $\operatorname{Hom}_{RH}(R, -)$ to (\star) gives

$$0 \to \operatorname{Hom}_{RH}(R, A) \to \operatorname{Hom}_{RH}(R, B) \to \operatorname{Hom}_{RH}(R, C) \to H^{1}(H, A) \to \cdots$$

So to find an example of an *H*-good short exact sequence which is not *RH*-split it is sufficient to find modules *C* and *A* such that $H^1(H, A) = 0$ and $\operatorname{Ext}^1_{RH}(C, A) \neq 0$. For example if *H* is any finite group we may set $R = \mathbb{Z}$, $A = \mathbb{Z}H$ and $C = (\mathbb{Z}/2\mathbb{Z})H$.

Additionally, we say that an RH-module M has property (P_H) if for any \mathcal{F} -good short exact sequence (\star) , $\operatorname{Hom}_{RH}(M, -)$ preserves the exactness of (\star) . Since $\operatorname{Hom}_{RH}(M, -)$ is always left exact, having (P_H) is equivalent to asking that for any \mathcal{F} -good short exact sequence (\star) and any RH-module homomorphism $f: M \to C$, there is a RH-module homomorphism $l: M \to B$ such that the diagram below commutes.



Note that the trivial RG-module R has property (P_H) .

LEMMA 4.4.6. If M has (P_H) then any direct summand of M, as an RH-module, has (P_H) .

PROOF. This is, with a minor alteration, the proof of [**Rot09**, Theorem 3.5(ii)]. Let N be a direct summand of M and consider the diagram with exact bottom row. Assume the bottom row is \mathcal{F} -good.



Here f is some arbitrary homomorphism, and π and ι are the projection and inclusion maps respectively. Since M has P_H , there is a map $l: M \to B$ such that $g \circ l = f \circ \pi$, the composition $l \circ \iota$ is the required map.

LEMMA 4.4.7. For any $K \in \mathcal{F}$, the permutation module R[G/K] has (P_H) .

PROOF. Let L be any subgroup of H, then using the natural isomorphism

$$\operatorname{Hom}_{RH}(R[H/L], -) \cong \operatorname{Hom}_{RL}(R, -)$$

we see that R[H/L] has (P_H) . Now use [**Bro94**, Proof of §III.9.5(ii) on p.83] to rewrite R[G/K] (as an *RH*-module), as

$$R[G/K] \cong \bigoplus_{g \in H \setminus G/K} R[H/K_g]$$

where $K_g = \{h \in H : g^{-1}hg \le K\}$. Thus:

$$\operatorname{Hom}_{RH}(R[G/K],-)\cong\prod_{g\in H\backslash G/K}\operatorname{Hom}_{RH}(R[H/K_g],-).$$

Now use that R[H/L] has (P_H) and that direct products of exact sequences are exact.

LEMMA 4.4.8. If

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is an H-good short exact sequence and both B and C have (P_H) then the short exact sequence is H-split and A has (P_H) .

PROOF. Apply $\operatorname{Hom}_{RH}(C, -)$ to see that the short exact sequence is *H*-split. Then since, as *RH*-modules, *B* is the direct sum of *C* and *A*, *A* necessarily has (P_H) by Lemma 4.4.6.

LEMMA 4.4.9. If P_* is an \mathcal{F} -good resolution of R by permutation RG-modules with stabilisers in \mathcal{F} , then P_* is \mathcal{F} -split.

PROOF. Use induction with Lemmas 4.4.7 and 4.4.8. \Box

REMARK 4.4.10. Similarly to Proposition 4.3.6, the above lemma may fail for \mathcal{F} -good resolutions of arbitrary modules.

PROPOSITION 4.4.11. If G is $\mathcal{H}_{\mathcal{F}}FP_n$ then G is $\mathcal{F}FP_n$.

PROOF. Find a free $\mathcal{H}_{\mathcal{F}}$ -module resolution P_* of R^- , finitely generated up to dimension n. By Remark 4.4.4, $P_*(G/1)$ is an \mathcal{F} -good resolution of R by permutation RG-modules with stabilisers in \mathcal{F} . By Lemma 4.4.9 P_* is \mathcal{F} -split, and by [**Nuc99**, Definition 2.2] permutation RG-modules with stabilisers in \mathcal{F} are \mathcal{F} -projective.

4.4.2. $\mathcal{F}FP_n$ implies $\mathcal{H}_{\mathcal{F}}FP_n$. This section comprises a series of lemmas, building to the proof of Proposition 4.4.17, that if R is commutative Noetherian and G is $\mathcal{F}FP_n$ then G is $\mathcal{H}_{\mathcal{F}}FP_n$.

LEMMA 4.4.12. For any $H \in \mathcal{F}$, inducing R[G/H] to a covariant $\mathcal{H}_{\mathcal{F}}$ -module gives the free module $R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$.

PROOF. On objects the two functors are equal:

$$\operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} R[G/H](G/K) = R[G/H] \otimes_{RG} R[G/1, -]_{\mathcal{H}_{\mathcal{F}}}(G/K)$$
$$= R[G/H] \otimes_{RG} \operatorname{Hom}_{RG}(RG, R[G/K])$$
$$= R[G/H] \otimes_{RG} R[G/K]$$
$$= R[H \setminus G/K]$$
$$= \operatorname{Hom}_{RG}(R[G/H], R[G/K]).$$

If $L \leq K$ are in \mathcal{F} , and $\sum_{I} g_{i}L \in R[G/L]^{H}$ then

$$R[G/1, -]_{\mathcal{H}_{\mathcal{F}}}(R_L^K) : R[G/1, G/L]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow R[G/1, G/K]_{\mathcal{H}_{\mathcal{F}}}$$
$$\sum_I g_i L \longmapsto \sum_I g_i K.$$

Following this down the chain of isomorphisms, then

$$\operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} R[G/H](R_{L}^{K}) : \operatorname{Hom}_{RG}(R[G/H], R[G/L]) \longrightarrow \operatorname{Hom}_{RG}(R[G/H], R[G/K])$$
$$\sum_{I} g_{i}L \longmapsto \sum_{I} g_{i}K$$

as required. Similarly, if $\sum_I g_i K \in R[G/K]^H$ then

$$R[G/1, -]_{\mathcal{H}_{\mathcal{F}}}(I_L^K) : R[G/1, G/K]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow R[G/1, G/L]_{\mathcal{H}_{\mathcal{F}}}$$
$$\sum_I g_i K \longmapsto \sum_{k \in K/L} \sum_I g_i kL.$$

Following this down the chain of isomorphisms,

$$\operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} R[G/H](I_L^K) : \operatorname{Hom}_{RG}(R[G/H], R[G/K]) \longrightarrow \operatorname{Hom}_{RG}(R[G/H], R[G/L])$$
$$\sum_I g_i K \longmapsto \sum_{k \in K/L} \sum_I g_i kL$$

again as required.

The proof for the conjugation morphisms c_g is similar to the above.

Lemma 4.4.13.

$$\operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} \prod_{H \in \mathcal{F}/G} \prod_{\Lambda_H} R[G/H] = \prod_{H \in \mathcal{F}/G} \prod_{\Lambda_H} R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}$$

where for each $H \in \mathcal{F}/G$, Λ_H is any indexing set and we are using covariant induction.

PROOF. In this proof, we use \prod as a shorthand for $\prod_{H \in \mathcal{F}/G} \prod_{\Lambda_H}$. On objects, the two functors are equal:

$$\operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} \prod R[G/H](G/K) = \left(\prod R[G/H] \right) \otimes_{RG} R[G/1, -]_{\mathcal{H}_{\mathcal{F}}}(G/K)$$
$$= \left(\prod R[G/H] \right) \otimes_{RG} \operatorname{Hom}_{RG}(RG, R[G/K])$$
$$= \left(\prod R[G/H] \right) \otimes_{RG} R[G/K]$$
$$\cong \prod (R[G/H] \otimes_{RG} R[G/K])$$
$$= \prod R[H \setminus G/K]$$
$$= \prod \operatorname{Hom}_{RG}(R[G/H], R[G/K]).$$

Where the isomorphism marked (\star) is the Bieri–Eckmann criterion [**Bie81**, Theorem 1.3], which is valid because R[G/K] is FP_{∞}. That the morphisms are equal can be checked as in the previous lemma.

LEMMA 4.4.14.

$$\mathcal{F}H_*\left(G, \prod_{H\in\mathcal{F}/G}\prod_{\Lambda_H}R[G/H]\right) = H_*^{\mathcal{H}_{\mathcal{F}}}\left(G, \prod_{H\in\mathcal{F}/G}\prod_{\Lambda_H}R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}\right)$$

where for each $H \in \mathcal{F}/G$, Λ_H is any indexing set.

PROOF. Again we use \prod to stand for $\prod_{H \in \mathcal{F}/G} \prod_{\Lambda_H}$. Let P_* be a free $\mathcal{H}_{\mathcal{F}}$ -module resolution of R^- , then $P_*(G/1)$ is an \mathcal{F} -split resolution of R by \mathcal{F} -projective modules by Lemma 4.4.9, so

$$\mathcal{F}H_*(G, R[G/H]) \cong H_*\left(P_*(G/1) \otimes_{RG} \prod R[G/H]\right)$$
$$\cong H_*\left(P_* \otimes_{\mathcal{H}_{\mathcal{F}}} \operatorname{Ind}_{RG}^{\mathcal{H}_{\mathcal{F}}} \prod R[G/H]\right)$$
$$\cong H_*(P_* \otimes_{\mathcal{H}_{\mathcal{F}}} \prod R[G/H, -]_{\mathcal{H}_{\mathcal{F}}})$$
$$\cong H_*^{\mathcal{H}_{\mathcal{F}}}(G, \prod R[G/H, -]_{\mathcal{H}_{\mathcal{F}}})$$

where the second isomorphism is the adjoint isomorphism between induction and restriction and the third is Lemma 4.4.13

LEMMA 4.4.15. For any group G, any commutative Noetherian ring R, any RG-module A of type $\mathcal{F}FP_n$, and any exact limit, the natural map

$$\mathcal{F}\operatorname{Tor}_{i}^{RG}(A, \varprojlim_{\lambda \in \Lambda} M_{\lambda}) \longrightarrow \varprojlim_{\lambda \in \Lambda} \mathcal{F}\operatorname{Tor}_{i}^{RG}(A, M_{\lambda})$$

is an isomorphism for i < n and an epimorphism for i = n.

PROOF. The proof is analogous to [**Bie81**, Theorem 1.3] and [**Nuc99**, Theorem 7.1], using [**Nuc99**, Proposition 6.3] which states that for R commutative Noetherian, finitely generated \mathcal{F} -projective modules over RG are of type FP_{∞}.

Specialising the previous lemma to M = R:

COROLLARY 4.4.16. If R is commutative Noetherian and G is $\mathcal{F}FP_n$ over R, then for any exact limit, the natural map

$$\mathcal{F}H_i(G, \varprojlim_{\lambda \in \Lambda} M_\lambda) \longrightarrow \varprojlim_{\lambda \in \Lambda} \mathcal{F}H_i(G, M_\lambda)$$

is an isomorphism for i < n and an epimorphism for i = n.

PROPOSITION 4.4.17. If R is commutative Noetherian and G is $\mathcal{F}FP_n$ over R then G is $\mathcal{H}_{\mathcal{F}}FP_n$ over R.

PROOF. In this proof, we write \prod for $\prod_{H \in \mathcal{F}/G} \prod_{\Lambda_H}$ where Λ_H is any indexing set. Using Lemmas 4.4.14 and 4.4.16, for any i < n:

$$H_i^{\mathcal{H}_{\mathcal{F}}}\left(G, \prod R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}\right) = \mathcal{F}H_i\left(G, \prod R[G/H]\right)$$
$$= \prod \mathcal{F}H_i\left(G, R[G/H]\right)$$
$$= \prod H_i^{\mathcal{H}_{\mathcal{F}}}\left(G, R[G/H, -]_{\mathcal{H}_{\mathcal{F}}}\right)$$

Thus G is $\mathcal{H}_{\mathcal{F}} FP_n$ by the Bieri–Eckmann criterion (Theorem 2.5.1).

REMARK 4.4.18. The requirement that R be Noetherian was needed only for Lemma 4.4.15, where we need that finitely generated \mathcal{F} -projectives are FP_{∞}. Nucinkis has given an example of a finitely generated \mathcal{F} -projective module which is not FP_{∞} [Nuc99, Remark on p.167], but the following question is still open.

QUESTION 4.4.19. Does Proposition 4.4.17 remain true if R is not Noetherian?

4.5. Cohomological dimension for cohomological Mackey functors

In [**Deg13a**], Degrijse shows that for all groups G with $\mathcal{H}_{\mathcal{F}} \operatorname{cd} G < \infty$,

$$\mathcal{F}\mathrm{cd}\,G=\mathcal{H}_{\mathcal{F}}\mathrm{cd}\,G.$$

We can improve this.

THEOREM 4.5.1. For all groups G,

$$\mathcal{F}\mathrm{cd}\,G=\mathcal{H}_{\mathcal{F}}\mathrm{cd}\,G.$$

PROOF. Remark 4.4.4 and Lemma 4.4.9 imply $\mathcal{F}cd G \leq \mathcal{H}_{\mathcal{F}}cd G$.

For the opposite inequality, we first use [Gan12b, Lemma 3.4] which states that for a group G with $\mathcal{F}cdG \leq n$ there is an \mathcal{F} -projective resolution P_* of Rof length n, where each P_i is a permutation module with stabilisers in \mathcal{F} . Given such a P_* , we take fixed points of P_* to get the $\mathcal{H}_{\mathcal{F}}$ resolution P_*^- . Since P_* is \mathcal{F} -split, P_*^- is exact.

Recall that $\mathcal{F}in$ denotes the family of finite subgroups of G and n_G denotes the minimal dimension of a proper contractible G-CW complex.

PROPOSITION 4.5.2. For all groups G,

$$\mathcal{H}_{\mathcal{F}in} \mathrm{cd}\, G \leq n_G.$$

This fact is well-known for \mathfrak{F} cd instead of $\mathcal{H}_{\mathcal{F}in}$ cd, but since a direct proof for $\mathcal{H}_{\mathcal{F}in}$ cd is both interesting and short we provide one.

PROOF. Let P_* denote the cellular chain complex for a contractible *G*-CWcomplex *X* of dimension *n* and take fixed points to get the complex $P_*^- \longrightarrow R^$ of $\mathcal{H}_{\mathcal{F}in}$ -modules. Since the action of *G* on *X* is proper the modules comprising P_* are permutation modules with finite stabilisers and so P_*^- is a chain complex of free $\mathcal{H}_{\mathcal{F}in}$ -modules. By a result of Bouc [**Bou99**] and Kropholler–Wall [**KW11**] this chain complex splits when restricted to a complex of *RH*-modules for any finite subgroup *H* of *G*. In other words, P_* is \mathcal{F} -good, thus $P_*^H \longrightarrow R$ is exact for any finite subgroup *H* by Remark 4.4.4.

This leads naturally to the question:

QUESTION 4.5.3. Does $\mathcal{H}_{\text{Fin}} \operatorname{cd} G < \infty$ imply $n_G < \infty$?

We know of no group for which n_G and $\mathcal{H}_{\mathcal{F}}$ cd G differ. Brown has asked the following:

QUESTION 4.5.4. [**Bro94**, VIII.11 p.226] If G is virtually torsion-free with $\operatorname{vcd} G < \infty$, then is $n_G = \operatorname{vcd} G$?

If G is virtually torsion free then $\operatorname{vcd} G = \mathcal{H}_{\operatorname{fin}} \operatorname{cd} G$ [MPN06], so a constructive answer to Question 4.5.3 would give information about Question 4.5.4 as well.

Related to this is the following question, posed using \mathfrak{F} cd instead of $\mathcal{H}_{\mathcal{F}in}$ cd by Nucinkis.

QUESTION 4.5.5. [Nuc00, p.337] Does $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G < \infty$ imply that $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G < \infty$?

REMARK 4.5.6. If G acts properly and cocompactly on a finite-dimensional contractible G-CW-complex then, by a modification of the argument of the proof of Lemma 4.5.2, G is $\mathcal{H}_{\mathcal{F}in} FP_{\infty}$ also. However, if G acts properly on a finite type but infinite dimensional contractible complex X, then the theorem of Bouc and Kropholler–Wall doesn't apply, and the cellular chain complex of X may not be \mathfrak{F} -split, thus we cannot deduce G is $\mathcal{H}_{\mathcal{F}in} FP_{\infty}$.

For an example of a group G acting properly on a finite-type but infinite dimensional contractible CW-complex with a cellular chain complex which is not \mathfrak{F} -split, take the cyclic group $K \cong C_2$ acting antipodally on the infinite sphere S_{∞} , with the usual CW structure of 2 cells in each dimension. One calculates that $C_*(S^{\infty})^K$ is not exact and hence that $C_*(S_{\infty})$ is not $\mathbb{Z}K$ -split.

QUESTION 4.5.7. If G acts properly on a contractible G-CW-complex of finite type, but not necessarily finite dimension, then is G of type $\mathcal{H}_{\mathcal{F}in} FP_{\infty}$?

4.5.1. Closure properties. The class of groups G with $\mathcal{H}_{\mathcal{F}} cd G < \infty$ is closed under subgroups, free products with amalgamation, HNN extensions [**Nuc00**, Corollary 2.7], direct products [**Gan12b**, Corollary 3.9] and extensions of finite groups by groups with $\mathcal{H}_{\mathcal{F}} cd$ finite [**Deg13a**, Lemma 5.1].

Section 5.3 contains a proof, via the Gorenstein cohomological dimension, that for a group extension

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1$$

where $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G < \infty$, we have $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} N + \mathcal{H}_{\mathcal{F}in} \operatorname{cd} Q \leq \mathcal{H}_{\mathcal{F}in} \operatorname{cd} G$.

PROPOSITION 4.5.8. [Gan12b, 3.8,3.10] Let

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1$$

be a group extension such that for any finite extension H of N where H/N has prime power order, $\mathcal{H}_{\text{fin}} \operatorname{cd} H \leq m$, then $\mathcal{H}_{\text{fin}} \operatorname{cd} G \leq n + m$.

LEMMA 4.5.9. Let N be any group and p any prime. If for any extension

 $1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1$

we have that $\mathcal{H}_{\text{fin}} \operatorname{cd} G = \mathcal{H}_{\text{fin}} \operatorname{cd} N$ where Q is the cyclic group of order p, then $\mathcal{H}_{\text{fin}} \operatorname{cd} G = \mathcal{H}_{\text{fin}} \operatorname{cd} N$, where Q is any finite p-group.

PROOF. We prove by induction on the order of Q, the case |Q| = p is by assumption. Let Q' be a normal subgroup of index p in Q (such a subgroup exists by [Rot95, Theorem 4.6(ii)]) and consider the diagram below.



Since Q' is normal in Q, the preimage $\pi^{-1}(Q')$ is normal in G, with quotient group $G/\pi^{-1}(Q')$ of order p so $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G = \mathcal{H}_{\mathcal{F}in} \operatorname{cd} \pi^{-1}(Q')$. Finally by the induction assumption $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} \pi^{-1}(Q') = \mathcal{H}_{\mathcal{F}in} \operatorname{cd} N$.

Combining the results above, if $\mathcal{H}_{\text{fin}} \operatorname{cd} G$ fails to be subadditive there must exist a finite cyclic group Q, group N with $\mathcal{H}_{\text{fin}} \operatorname{cd} N < \infty$, and an extension Gof Q by N with $\mathcal{H}_{\text{fin}} \operatorname{cd} G = \infty$.

QUESTION 4.5.10. If N is a group with $\mathcal{H}_{\text{fin}} \operatorname{cd} N < \infty$ then does every extension G of a cyclic group of order p by N satisfy $\mathcal{H}_{\text{fin}} \operatorname{cd} G < \infty$?

Any counterexample cannot be virtually torsion-free, since $\mathcal{H}_{\mathcal{F}in} \operatorname{cd} G = \operatorname{vcd} G$ for all virtually torsion-free groups [**MPN06**], and neither can it be elementary amenable [**Gan12b**, Proposition 3.13].

4.6. The family of *p*-subgroups

Throughout this section q is an arbitrary fixed prime and R will denote one of the following rings: the integers \mathbb{Z} , the finite field \mathbb{F}_q , or the integers localised at q denoted $\mathbb{Z}_{(q)}$. If $R = \mathbb{F}_q$ or $\mathbb{Z}_{(q)}$ then let \mathcal{P} denote the subfamily of \mathcal{F} consisting of all finite q-subgroups of groups in \mathcal{F} . If $R = \mathbb{Z}$ then let \mathcal{P} denote the subfamily of finite p-subgroups of groups in \mathcal{F} for all primes p.

We will always treat the cases $R = \mathbb{F}_q$ and $R = \mathbb{Z}_{(q)}$ together, in fact the only property of these rings that we use is that for any integer *i* coprime to *q*, the image of *i* under the map $\mathbb{Z} \to R$ is invertible in *R*. Hence the arguments in this section generalise to any other rings with this property, for example any ring with characteristic *q*. The argument used for $R = \mathbb{Z}$ will go through for any ring *R*.

For $R = \mathbb{Z}$ and $\mathcal{F} = \mathcal{F}in$, Leary and Nucinkis prove that the conditions $\mathcal{F}FP_n$ and $\mathcal{P}FP_n$ are equivalent, and that $\mathcal{F}cd G = \mathcal{P}cd G$ [LN10, Theorem 4.1]. We use an averaging method similar to theirs to show that, for $R = \mathbb{Z}$, \mathbb{F}_q , or $\mathbb{Z}_{(q)}$:

THEOREM 4.6.1. For $n \in \mathbb{N} \cup \{\infty\}$, the conditions $\mathcal{H}_{\mathcal{F}} cd G = n$ and $\mathcal{H}_{\mathcal{P}} cd G = n$ n are equivalent, as are the conditions $\mathcal{H}_{\mathcal{F}} FP_n$ and $\mathcal{H}_{\mathcal{P}} FP_n$.

REMARK 4.6.2. If $R = \mathbb{Z}_{(q)}$ or \mathbb{F}_q and all subgroups of G have order coprime to q then \mathcal{P} contains only the trivial subgroup. Thus $\mathcal{H}_{\mathcal{F}} \operatorname{cd}_R G = \operatorname{cd}_R G$ and the conditions $\mathcal{H}_{\mathcal{F}} \operatorname{FP}_n$ and FP_n are equivalent. At the end of the section we will look at the case that $R = \mathbb{K}$ is a field of characteristic zero, and prove that in this case $\mathcal{H}_{\mathcal{F}} \operatorname{cd} G = \operatorname{cd} G$ and that the conditions $\mathcal{H}_{\mathcal{F}} \operatorname{FP}_n$ and FP_n are equivalent.

The argument relies on two maps ι_H and ρ_H defined for any subgroup H in $\mathcal{F} \setminus \mathcal{P}$. These maps have different definitions depending on the ring R.

We treat the case that $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$ first. Let $H \in \mathcal{F} \setminus \mathcal{P}$ and let Q be a Sylow q-subgroup of H, define

$$\rho_H = R_Q^H \in R[G/Q, G/H]_{\mathcal{H}_F}$$
$$\iota_H = (1/|H:Q|) I_Q^H \in R[G/H, G/Q]_{\mathcal{H}_F}.$$

The map ι_H is well defined since |H : Q| contains no powers of q and hence is invertible in R.

If $R = \mathbb{Z}$ and $H \in \mathcal{F} \setminus \mathcal{P}$ then let $\{P_i\}_{i \in I}$ run over the non-trivial Sylow *p*-subgroups of H (choosing one subgroup for each p). We necessarily have that $gcd\{|H:P_i|: i \in I\} = 1$ so, by Bézout's identity, we may choose integers z_i such that $\sum_{i \in I} z_i |H:P_i| = 1$. Define, with a slight abuse of notation,

$$\rho_H = \bigoplus_{i \in I} R_{P_i}^H.$$

By which we mean that for any $\mathcal{H}_{\mathcal{F}}$ -module M,

$$M(\rho_H) : M(G/H) \longrightarrow \bigoplus_{i \in I} M(G/P_i)$$
$$m \longmapsto \bigoplus_{i \in I} M(R_{P_i}^H)(m).$$

With a similar abuse of notation we define

$$\iota_H = \sum_{i \in I} z_i I_{P_i}^H.$$

By which we mean that for any $\mathcal{H}_{\mathcal{F}}$ -module M,

$$M(\iota_H) : \bigoplus_{i \in I} M(G/P_i) \longrightarrow M(G/H)$$
$$(m_i)_{i \in I} \longmapsto \sum_{i \in I} z_i M(I_{P_i}^H)(m_i).$$

The next couple of lemmas catalogue properties of the maps ι_H and ρ_H which are needed for the proof of Theorem 4.6.1.

LEMMA 4.6.3. For any $\mathcal{H}_{\mathcal{F}}$ -module M and subgroup $H \in \mathcal{F} \setminus \mathcal{P}$,

$$M(\iota_H) \circ M(\rho_H) = \mathrm{id}_{M(G/H)}.$$

PROOF. In the case $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$, this follows from the fact that $M(R_Q^H \circ I_Q^H)$ is multiplication by |H:Q|. For $R = \mathbb{Z}$,

$$M(\iota_{H}) \circ M(\rho_{H}) = \sum_{i} z_{i} M(R_{P_{i}}^{H} \circ I_{P_{i}}^{H}) = \sum_{i} z_{i} |H:P_{i}| = 1.$$

LEMMA 4.6.4. If $H \in \mathcal{F}$ then $\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} R[-, G/H]_{\mathcal{H}_{\mathcal{F}}}$ is a finitely generated projective $\mathcal{H}_{\mathcal{P}}$ -module.

PROOF. If H is an element of \mathcal{P} then this is obvious so assume that $H \notin \mathcal{P}$. First, the case $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$. The projection

$$s: R[-, G/Q]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow R[-, G/H]_{\mathcal{H}_{\mathcal{F}}}$$

corresponding to ι_H under the Yoneda-type lemma (2.0.2) is split by the map

$$i: R[-, G/H]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow R[-, G/Q]_{\mathcal{H}_{\mathcal{F}}}$$

corresponding to ρ_H under the Yoneda-type lemma: It is sufficient to calculate

$$s \circ i(G/H)(\mathrm{id}_H) = \rho_H \circ \iota_H = \mathrm{id}_H$$

Applying $\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}}$ gives a split surjection

$$\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} s : \operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} R[-, G/Q]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow \operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} R[-, G/H]_{\mathcal{H}_{\mathcal{F}}}.$$

Since $\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} R[-, G/Q]_{\mathcal{H}_{\mathcal{F}}} = R[-, G/Q]_{\mathcal{H}_{\mathcal{P}}}$ this completes the proof.

Now the case $R = \mathbb{Z}$, this time we construct a split surjection

$$s: \bigoplus_{i \in I} R[-, G/P_i]_{\mathcal{H}_{\mathcal{F}}} \longrightarrow R[-, G/H]_{\mathcal{H}_{\mathcal{F}}}$$

using the maps corresponding to ι_H and ρ_H under the Yoneda-type lemma. The rest of the proof is identical to the case $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$.

LEMMA 4.6.5. A chain complex C_* of $\mathcal{H}_{\mathcal{F}}$ -modules is exact if and only if it is exact at G/P for all subgroups $P \in \mathcal{P}$.

PROOF. The "only if" direction is obvious so assume C_* is a chain complex of $\mathcal{H}_{\mathcal{F}}$ -modules, exact at all $P \in \mathcal{P}$ and let $H \in \mathcal{F} \setminus \mathcal{P}$.

We claim that the maps $C_*(\iota_H)$ and $C_*(\rho_H)$ are chain complex maps, we show this below for $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$, the proof for $R = \mathbb{Z}$ is analogous. The only non-obvious part of this claim is that the maps commute with the boundary maps ∂_i of C_* , in other words the diagrams below commute:

$$\begin{array}{ccc} C_i(G/H) \xrightarrow{\partial_i(G/H)} C_{i-1}(G/H) \\ \xrightarrow{C_i(\rho_H)} & & & \downarrow C_{i-1}(\rho_H) \\ C_i(G/Q) \xrightarrow{\partial_i(G/Q)} C_{i-1}(G/Q) \end{array}$$

a (a (a)

$$\begin{array}{ccc} C_i(G/Q) \xrightarrow{\partial_i(G/Q)} C_{i-1}(G/Q) \\ \xrightarrow{C_i(\iota_H)} & & & \downarrow C_{i-1}(\iota_H) \\ C_i(G/H) \xrightarrow{\partial_i(G/H)} C_{i-1}(G/H) \end{array}$$

This follows from the fact that ∂_i is an $\mathcal{H}_{\mathcal{F}}$ -module map.

Lemma 4.6.3 gives that $C_*(\iota_H) \circ C_*(\rho_H)$ is the identity on the chain complex $C_*(G/H)$. The induced maps ι_H^* and ρ_H^* on homology satisfy

$$\iota_H^* \circ \rho_H^* = \mathrm{id} : H_*(C_*(G/H)) \longrightarrow H_*(C_*(G/H))$$

so ρ_H^* is injective. The image of ρ_H^* lies in $H_*(C_*(G/Q)) = 0$ if $R = \mathbb{F}_q$ or $R = \mathbb{Z}_{(q)}$, or $\oplus_i H_*(C_*(G/P_i)) = 0$ if $R = \mathbb{Z}$, hence $H_*(C_*(G/H))$ is zero. \Box

LEMMA 4.6.6. If P is a projective (respectively finitely generated projective) $\mathcal{H}_{\mathcal{P}}$ -module then there exists a $\mathcal{H}_{\mathcal{F}}$ -module Q such that

$$\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}}Q = P$$

and Q is projective (resp. finitely generated projective).

PROOF. Recall from Lemma 4.1.20 that the projective $\mathcal{H}_{\mathcal{P}}$ -modules are exactly those of the form V^- for V some direct summand of a permutation RG-module whose stabilisers lie in \mathcal{P} . The required module is just V^- regarded as a $\mathcal{H}_{\mathcal{F}}$ -module.

PROOF OF THEOREM 4.6.1. Assume that $\mathcal{H}_{\mathcal{F}} \mathrm{cd} G \leq n$ and let P_* be a length *n* projective resolution of R^- by $\mathcal{H}_{\mathcal{F}}$ -modules, then restricting to the family \mathcal{P} and using Lemma 4.6.4 gives a length *n* projective resolution by $\mathcal{H}_{\mathcal{P}}$ modules. A similar argument shows that $\mathcal{H}_{\mathcal{F}} \mathrm{FP}_n$ implies $\mathcal{H}_{\mathcal{P}} \mathrm{FP}_n$.

For the converse, let P_* be a length n projective resolution of R^- by $\mathcal{H}_{\mathcal{P}^-}$ modules. Lemma 4.6.6 gives projective $\mathcal{H}_{\mathcal{F}^-}$ -modules Q_i such that $\operatorname{Res}_{\mathcal{H}_{\mathcal{P}}}^{\mathcal{H}_{\mathcal{F}}} Q_i = P_i$ for each i. Denoting by d_i the boundary maps in P_* , define boundary maps of Q_* as $\partial_i(G/P) = d_i(G/P)$ if $P \in \mathcal{P}$ and if $H \notin \mathcal{P}$ then

$$\partial_i(G/H) = P_{i-1}(\iota_H) \circ d_i(G/H) \circ P_i(\rho_H).$$

One can check that these maps are indeed $\mathcal{H}_{\mathcal{F}}$ -module maps and that this makes Q_* a chain complex:

$$\begin{aligned} \partial_i(G/H) &\circ \partial_{i+1}(G/H) \\ &= P_{i-1}(\iota_H) \circ d_i(G/H) \circ P_i(\rho_H) \circ P_i(\iota_H) \circ d_{i+1}(G/H) \circ P_{i+1}(\rho_H) \\ &= P_{i-1}(\iota_H) \circ d_i(G/H) \circ d_{i+1}(G/H) \circ P_{i+1}(\rho_H) \\ &= 0. \end{aligned}$$

Finally P_* is exact by Lemma 4.6.5.

Since at all stages of the argument above finite generation is preserved, we get that $\mathcal{H}_{\mathcal{P}}FP_n$ implies $\mathcal{H}_{\mathcal{F}}FP_n$ too.

For the remainder of this section $R = \mathbb{K}$ is a field of characteristic zero, in this case we can reduce to the family *Triv* containing only the trivial subgroup. For any $H \in \mathcal{F}$, let

$$\rho_H = R_1^H$$
$$\iota_H = (1/|H|)I_1^H$$

All the arguments of the section go through with no alteration, showing:

PROPOSITION 4.6.7. $\mathcal{H}_{\mathcal{F}} \operatorname{cd}_{\mathbb{K}} G = \mathcal{H}_{\operatorname{Triv}} \operatorname{cd}_{\mathbb{K}} G$ and the conditions $\mathcal{H}_{\mathcal{F}} \operatorname{FP}_n$ over \mathbb{K} and $\mathcal{H}_{\operatorname{Triv}} \operatorname{FP}_n$ over \mathbb{K} are equivalent for any $n \in \mathbb{N} \cup \{\infty\}$.

COROLLARY 4.6.8. $\mathcal{H}_{\mathcal{F}} \operatorname{cd}_{\mathbb{K}} G = \operatorname{cd}_{\mathbb{K}} G$ and the conditions $\mathcal{H}_{\mathcal{F}} \operatorname{FP}_n$ over \mathbb{K} and FP_n over $\mathbb{K} G$ are equivalent for any $n \in \mathbb{N} \cup \{\infty\}$.

PROOF. The category of \mathcal{H}_{Triv} -modules is isomorphic to the category of $\mathbb{K}G$ -modules.

4.6.1. FP_n conditions over \mathbb{F}_p . Throughout this section, we fix a prime p and work over the ring \mathbb{F}_p with the family \mathcal{P} of all finite *p*-subgroups of groups in \mathcal{F} .

LEMMA 4.6.9. [HPY13, Lemma 5.3] For any finite subgroup $H \in \mathcal{P}$ and $\mathcal{H}_{\mathcal{P}}$ -module M, $D_H M$ extends to a cohomological Mackey functor.

PROPOSITION 4.6.10. G is $\mathcal{H}_{\mathcal{P}} FP_n$ over \mathbb{F}_p if and only if G is $\mathcal{O}_{\mathcal{P}} FP_n$ over \mathbb{F}_p .

The proof is basically that of Proposition 4.2.5 combined with the lemma above.

PROOF. We know already from Proposition 4.4.2 and Corollary 4.2.6 that $\mathcal{O}_{\mathcal{P}}FP_n$ implies $\mathcal{H}_{\mathcal{P}}FP_n$.

Let M_{λ} , for $\lambda \in \Lambda$, be a directed system of $\mathcal{O}_{\mathcal{P}}$ -modules with colimit zero. Using the notation of Proposition 4.2.5 there is an exact sequence of directed systems for each $i \geq 0$,

$$0 \longrightarrow C^{i} M_{\lambda} \longrightarrow DC^{i} M_{\lambda} \longrightarrow C^{i+1} M_{\lambda} \longrightarrow 0,$$

each of which has colimit zero. Moreover, DC^iM_{λ} extends to a cohomological Mackey functor so using the Bieri–Eckmann criterion (Theorem 2.5.1), if $m \leq n$ then for all $i \geq 0$,

$$\lim_{\Lambda} H^m_{\mathcal{O}_{\mathcal{P}}}(G, DC^i M_{\lambda}) = 0.$$

Thus,

$$\varinjlim H^m_{\mathcal{O}_{\mathcal{P}}}(G, M_{\lambda}) = \varinjlim H^m_{\mathcal{O}_{\mathcal{P}}}(G, C^0 M_{\lambda})$$

$$= \varinjlim H^{m-1}_{\mathcal{O}_{\mathcal{P}}}(G, C^1 M_{\lambda})$$

$$= \cdots$$

$$= \varinjlim H^0_{\mathcal{O}_{\mathcal{P}}}(G, C^m M_{\lambda})$$

$$= 0.$$

Where the final zero is because G is $\mathcal{O}_{\mathcal{P}} FP_0$ (by [**LN10**, Proposition 4.2] and Theorem 4.4.1).

COROLLARY 4.6.11. G is $\mathcal{H}_{\mathcal{F}} FP_n$ over \mathbb{F}_p if and only if \mathcal{P} contains finitely many conjugacy classes, and WH is FP_n over \mathbb{F}_p for all $H \in \mathcal{P}$.

PROOF. G is $\mathcal{H}_{\mathcal{F}} FP_n$ if and only if G is $\mathcal{O}_{\mathcal{P}} FP_n$ by Theorem 4.6.1 and Proposition 4.6.10. Now use that G is $\mathcal{O}_{\mathcal{P}} FP_n$ if and only if \mathcal{P} contains finitely many conjugacy classes, and WH is FP_n for all $H \in \mathcal{P}$ (Corollary 3.6.4).

PROPOSITION 4.6.12. If G is virtually torsion-free then the conditions virtually FP over \mathbb{F}_p and \mathcal{H}_F FP over \mathbb{F}_p are equivalent.

PROOF. If G is virtually FP over \mathbb{F}_p then G has finitely many conjugacy classes of finite p-subgroups [**Bro94**, IX.(13.2)]. A result of Hamilton gives that for any finite p-subgroup H of G, WH is virtually FP over \mathbb{F}_p , in particular WH is FP_{∞} over \mathbb{F}_p [**Ham11**, Theorem 7]. Finally, [**Ham11**, Proposition 34] gives that G acts properly on a finite-dimensional \mathbb{F}_p -acyclic space, thus in particular $\mathcal{H}_{\mathcal{F}}cd_{\mathbb{F}_p} G < \infty$. The other direction is obvious.

In [LN10] it is conjectured that, if $\mathcal{F} = \mathcal{F}in$, G is $\mathcal{F}FP_{\infty}$ if and only if G is FP_{∞} and has finitely many conjugacy classes of finite p-subgroups for all primes p. One could generalise this and ask:

QUESTION 4.6.13. Let $\mathcal{F} = \mathcal{F}in$ and $n \in \mathbb{N} \cup \{\infty\}$.

- (1) If G is FP_n over \mathbb{Z} with finitely many conjugacy classes of finite psubgroups for all primes p, then is G of type $\mathcal{H}_{\mathcal{F}}FP_n$ over \mathbb{Z} ?
- (2) Fixing a prime p, if G is FP_n over \mathbb{F}_p with finitely many conjugacy classes of finite p-subgroups then is G of type $\mathcal{H}_{\mathcal{F}}FP_n$ over \mathbb{F}_p ?

A problem with finding a counterexample to Question 4.6.13(2) is that if G admits a cocompact action on a finite-dimensional \mathbb{F}_p -acyclic space X then, via Smith theory, X^P is \mathbb{F}_p -acyclic for any finite p-subgroup P and thus WP is FP_n over \mathbb{F}_p . For this reason one cannot use the examples of Leary and Nucinkis in [**LN03**], their construction requires actions of finite groups on finite dimensional \mathbb{F}_p -acyclic flag complexes with fixed point sets that are not \mathbb{F}_p -acyclic.

CHAPTER 5

Gorenstein cohomology and *F*-cohomology

This chapter contains material that has appeared in:

• On the Gorenstein and \mathfrak{F} -cohomological dimensions (2013, to appear Bull. Lond. Math. Soc.) [SJG13b].

We study the Gorenstein cohomological dimension $\operatorname{Gcd} G$ and prove the following result.

THEOREM 5.2.11. If $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G = \operatorname{Gcd} G$.

The proof is via the construction in Theorem 5.2.7 of a long exact sequence relating the \mathfrak{F} -cohomology, the complete \mathfrak{F} -cohomology, and a new cohomology theory we call the \mathfrak{F}_G -cohomology. The construction is analogous to the construction of the long exact sequence of Avramov–Martsinkovsky relating the group cohomology, complete cohomology, and Gorenstein cohomology [**AM02**, §7][**ABS09**, Theorem 3.11].

In Section 5.3 we use Theorem 5.2.11 and subadditivity of the Gorenstein cohomological dimension to study the behaviour of the \mathfrak{F} -cohomological dimension under group extensions.

COROLLARY 5.3.2. Given a short exact sequence of groups

 $1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1,$

if $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G \leq \operatorname{\mathfrak{F}cd} N + \operatorname{\mathfrak{F}cd} Q$.

Finally, in Section 5.4 we use the Avramov–Martsinkovsky long exact sequence to prove the following.

PROPOSITION 5.4.4. If $\operatorname{Gcd} G < \infty$ and $\operatorname{cd}_{\mathbb{Q}} G < \infty$ then $\operatorname{cd}_{\mathbb{Q}} G \leq \operatorname{Gcd} G$.

5.1. Preliminaries

5.1.1. Complete resolutions and complete cohomology. A weak complete resolution of a module M is an acyclic complex T_* of projective modules which coincides with an ordinary projective resolution P_* of M in sufficiently high degree. The degree in which the two coincide is called the *coincidence index*. A weak complete resolution is called a *strong complete resolution* if $\text{Hom}_{RG}(T_*, Q)$ is acyclic for every projective module Q. We avoid the term "complete resolution" since some authors use it to refer to a weak complete resolution and others to a strong complete resolution.

PROPOSITION 5.1.1. [ABS09, Proposition 2.8] A group G admits a strong complete resolution if and only if $\operatorname{Gcd} G < \infty$.

The advantage of strong complete resolutions is that given strong complete resolutions T_* and S_* of modules M and N, any module homomorphism $M \to N$ lifts to a morphism of strong complete resolutions $T_* \to S_*$ [**CK97**, Lemma 2.4]. Thus they can be used to define a cohomology theory: given a strong complete resolution T_* of M we define

$$\widehat{\operatorname{Ext}}_{RG}^*(M,-) \cong H^* \operatorname{Hom}_{RG}(T_*,-).$$

We also set $\widehat{H}^*(G, -) = \widehat{\operatorname{Ext}}_{RG}^*(R, -)$. This coincides with the complete cohomology of Mislin [**Mis94**], Vogel [**Goi92**], and Benson–Carlson [**BC92**] (see [**CK97**, Theorem 1.2] for a proof). Recall that the complete cohomology is itself a generalisation of the Farrell–Tate Cohomology, defined only for groups with finite virtual cohomological dimension [**Br094**, §X].

Even weak complete resolutions do not always exist, for example a free Abelian group of infinite rank cannot admit a weak complete resolution [**MT00**, Corollary 2.10]. It is conjectured by Dembegioti and Talelli that a $\mathbb{Z}G$ -module admits a weak complete resolution if and only if it admits a strong complete resolution [**DT10**, Conjecture B].

5.1.2. *F*-cohomology. This section contains two technical lemmas we will need later.

If M is any RG-module and $F_i = R\Delta^i$ is the standard \mathfrak{F} -split resolution of R[**Nuc00**, p.342], then $F_* \otimes_R M$ is an \mathfrak{F} -split \mathfrak{F} -projective resolution of M. Thus we've shown:

LEMMA 5.1.2. \mathfrak{F} -split \mathfrak{F} -projective resolutions exist for all RG-modules M.

There is also a version of the Horseshoe lemma.

LEMMA 5.1.3 (Horseshoe lemma). If

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is an \mathfrak{F} -split short exact sequence and P_* and Q_* are \mathfrak{F} -split \mathfrak{F} -projective resolutions of A and C respectively then there is an \mathfrak{F} -split \mathfrak{F} -projective resolution S_* of B such that $S_i = P_i \oplus Q_i$ and there is an \mathfrak{F} -split short exact sequence of augmented complexes

$$0 \longrightarrow \tilde{P}_* \longrightarrow \tilde{S}_* \longrightarrow \tilde{Q}_* \longrightarrow 0.$$
The proof is similar to [EJ11, Lemma 8.2.1].

PROOF. First build the diagram below as in, for example, [**Rot09**, Proposition 6.24] where it is shown to commute and have exact rows and columns. Here, K_A , K_B and K_C are the kernels of the maps $P_0 \longrightarrow A$, $P_0 \oplus Q_0 \longrightarrow B$, and $Q_0 \longrightarrow C$ respectively.



Since P_0 and Q_0 are both \mathfrak{F} -projective, $P_0 \oplus Q_0$ is \mathfrak{F} -projective, and since the middle row is split, it is \mathfrak{F} -split.

Let Δ be the *G*-set $\coprod_{H \in \mathcal{F}in} G/H$ and apply $- \otimes R\Delta$ to the commutative diagram to obtain a new commutative diagram with exact left column, right column, bottom row, and central row.

$$(P_0 \oplus Q_0) \otimes R\Delta \longrightarrow B \otimes R\Delta$$

is surjective is because the tensor product is right exact, and an application of the 5-Lemma [**Rot09**, Lemma 2.72] shows

$$K_B \otimes R\Delta \longrightarrow (P_0 \oplus Q_0) \otimes R\Delta$$

is injective. Hence the central column of our new commutative diagram is exact. The 3×3 -Lemma provides that the top row is exact too [**Rot09**, Ex 2.32], thus all rows and columns of the first commutative diagram are \mathfrak{F} -split.

Now repeat this process, but starting with the \mathfrak{F} -split short exact sequence

$$0 \longrightarrow K_A \longrightarrow K_B \longrightarrow K_C \longrightarrow 0.$$

5.1.3. Complete \mathfrak{F} -cohomology. In [Nuc99], Nucinkis constructs a complete \mathfrak{F} -cohomology, we give a brief outline here. An \mathfrak{F} -complete resolution T_* of M is an acyclic \mathfrak{F} -split complex of \mathfrak{F} -projectives which coincides with an \mathfrak{F} -split \mathfrak{F} -projective resolution of M in high enough dimensions. An \mathfrak{F} -strong \mathfrak{F} -complete

resolution T_* has $\operatorname{Hom}_{RG}(T_*, Q)$ exact for all \mathfrak{F} -projectives Q. Given such a T_* we define

$$\widehat{\mathfrak{F}}\operatorname{Ext}_{RG}^{*}(M,-) = H^{*}\operatorname{Hom}_{RG}(T_{*},-)$$
$$\widehat{\mathfrak{F}}H^{*}(G,-) = \widehat{\mathfrak{F}\operatorname{Ext}}_{RG}^{*}(R,-).$$

Nucinkis also describes a Mislin style construction and a Benson–Carlson construction of complete \mathfrak{F} -cohomology defined for all groups, proves they are equivalent, and proves that whenever there exists an \mathfrak{F} -complete resolution they agree with the definition above.

5.1.4. Gorenstein cohomology. The Gorenstein cohomology is, like the \mathfrak{F} -cohomology, a special case of the relative homology of Mac Lane [ML95, §IX] and Eilenberg–Moore [EM65].

Recall that a module is Gorenstein projective if it is a cokernel in a strong complete resolution. An acyclic complex C_* of Gorenstein projective modules is *G*-proper if $\operatorname{Hom}_{RG}(Q, C_*)$ is exact for every Gorenstein projective Q. The class of G-proper short exact sequences is allowable in the sense of Mac Lane [**ML95**, §IX.4]. The projectives objects with respect to G-proper short exact sequences are exactly the Gorenstein projectives (for the definition of a projective object with respect to a class of short exact sequences see [**ML95**, p.261]). For M and N any RG-modules, we define

$$\operatorname{GExt}_{RG}^*(M,N) = H^* \operatorname{Hom}_{RG}(P_*,N)$$
$$\operatorname{GH}^*(G,N) = \operatorname{GExt}_{RG}^*(R,N)$$

where P_* is a G-proper resolution of M by Gorenstein projectives.

The usual method of producing a "Gorenstein projective dimension" of a module M in this setting would be to look at the shortest length of a G-proper resolution of M by Gorenstein projectives. A priori this could be larger than the Gorenstein projective dimension defined in the introduction, where the G-proper condition is not required. Fortunately there is the following theorem of Holm:

THEOREM 5.1.4. [Hol04, Theorem 2.10] If M has finite Gorenstein projective dimension then M admits a G-proper Gorenstein projective resolution of length Gpd M.

Generalising an argument of Avramov and Martsinkovsky in [AM02, §7] Asadollahi, Bahlekeh, and Salarian construct a long exact sequence:

THEOREM 5.1.5 (Avramov–Martsinkovsky long exact sequence). [ABS09, Theorem 3.11] For a group G with $\operatorname{Gcd} G < \infty$, there is a long exact sequence of cohomology functors

$$0 \longrightarrow \mathrm{G}H^1(G, -) \longrightarrow H^1(G, -) \longrightarrow \cdots$$

$$\cdots \longrightarrow \mathrm{G}H^n(G,-) \longrightarrow H^n(G,-) \longrightarrow \widehat{H}^n(G,-) \longrightarrow \mathrm{G}H^{n+1}(G,-) \longrightarrow \cdots$$

The construction relies on the complete cohomology being calculable via a complete resolution, hence the requirement that $\operatorname{Gcd} G < \infty$.

We will need the following lemma later:

LEMMA 5.1.6. Any G-proper resolution of R is \mathfrak{F} -split.

PROOF. If P_* is a G-proper resolution of R then since R[G/H] is a Gorenstein projective [**ABS09**, Lemma 2.21],

$$\operatorname{Hom}_{RG}(R[G/H], P_*) \cong \operatorname{Hom}_{RH}(R, P_*) \cong P_*^H$$

is exact, thus by the argument of Proposition 4.4.11 P_* is \mathfrak{F} -split.

5.2. \mathfrak{F}_G -cohomology

5.2.1. Construction. We define another special case of relative homology, which we call the \mathfrak{F}_G -cohomology. It enables us to build an Avramov–Martsinkovsky long exact sequence of cohomology functors containing $\mathfrak{F}H^*(G, -)$ and $\widehat{\mathfrak{F}H}^*(G, -)$.

We define an \mathfrak{F}_G -projective to be the cokernel in a \mathfrak{F} -complete \mathfrak{F} -strong resolution and say a complex C_* of RG-modules is \mathfrak{F}_G -proper if $\operatorname{Hom}_{RG}(Q, C_*)$ is exact for any \mathfrak{F}_G -projective Q. The \mathfrak{F}_G -proper short exact sequences form an allowable class in the sense of Mac Lane, whose projective objects are the \mathfrak{F}_G projectives—to check the class of \mathfrak{F}_G -proper short exact sequences is allowable we need only check that given a \mathfrak{F}_G -proper short exact sequence, any isomorphic short exact sequence is \mathfrak{F}_G -proper and that for any RG-module A the short exact sequences

$$0 \longrightarrow A \stackrel{\mathrm{id}}{\longrightarrow} A \longrightarrow 0 \longrightarrow 0$$

and

$$0 \longrightarrow 0 \longrightarrow A \xrightarrow{\mathrm{id}} A \longrightarrow 0$$

are \mathfrak{F}_G -proper.

We don't know if the class of \mathfrak{F}_G -projectives is precovering (see [**EJ11**, §8]), so we don't know if there always exists an \mathfrak{F}_G -proper \mathfrak{F}_G -projective resolution. However, if A and B admit \mathfrak{F}_G -proper \mathfrak{F}_G -resolutions P_* and Q_* , respectively, then any map $A \longrightarrow B$ induces a map of resolutions $P_* \longrightarrow Q_*$ which is unique up to chain homotopy equivalence [**ML95**, IX.4.3] and we have a slightly weaker form of the Horseshoe lemma.

LEMMA 5.2.1 (Horseshoe lemma). Suppose

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is a \mathfrak{F}_G -proper short exact sequence of RG-modules and both A and C admit \mathfrak{F}_G proper \mathfrak{F}_G -projective resolutions P_* and Q_* then there is an \mathfrak{F}_G -proper resolution S_* of B such that $S_i = P_i \oplus Q_i$ and there is an \mathfrak{F}_G -proper short exact sequence of augmented complexes

$$0 \longrightarrow \tilde{P}_* \longrightarrow \tilde{S}_* \longrightarrow \tilde{Q}_* \longrightarrow 0.$$

The proof is similar to that of [EJ11, 8.2.1] and Lemma 5.1.3.

PROOF. First build the same commutative diagram as in the proof of Lemma 5.1.3. Since P_0 and Q_0 are both \mathfrak{F}_G -projective, $P_0 \oplus Q_0$ is \mathfrak{F}_G -projective, and since the middle row is split, it is \mathfrak{F}_G -proper.

Let T be an \mathfrak{F}_G -projective and apply $\operatorname{Hom}_{RG}(T, -)$ to obtain a new commutative diagram with exact left column, right column, bottom row, and central row. That

$$\operatorname{Hom}_{RG}(T, P_0 \oplus Q_0) \longrightarrow \operatorname{Hom}_{RG}(T, B)$$

is surjective is an application of the 5-Lemma [**Rot09**, Lemma 2.72], and another application of the same lemma shows

$$\operatorname{Hom}_{RG}(T, K_B) \longrightarrow \operatorname{Hom}_{RG}(T, P_0 \oplus Q_0)$$

is injective. Hence the central column of our new commutative diagram is exact, and an application of the 3×3 -Lemma shows the top row is exact [**Rot09**, Ex 2.32], thus the original commutative diagram is \mathfrak{F} -split. The rest of the proof is the same as that of Lemma 5.1.3.

For any module M which admits an \mathfrak{F}_G -proper resolution P_* by \mathfrak{F}_G -projectives we define

$$\mathfrak{F}_{G}Ext^*_{RG}(M,N) = H^* \operatorname{Hom}_{RG}(P_*,N).$$

We define also

$$\mathfrak{F}_{\mathrm{G}}H^*(G,-) = \mathfrak{F}_{\mathrm{G}}\mathrm{Ext}^*_{RG}(R,-)$$

The next lemma follows from Lemma 5.2.1, see [EJ11, 8.2.3].

LEMMA 5.2.2. Suppose

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

is a \mathfrak{F}_G -proper short exact sequence of RG-modules and both A and C admit \mathfrak{F}_G -proper \mathfrak{F}_G -projective resolutions, then there is an \mathfrak{F}_G Ext $^*_{RG}(-, M)$ long exact sequence for any RG-module M.

For any RG-module M the \mathfrak{F}_G projective dimension of G denoted $\mathfrak{F}_G \text{pd} M$ is the minimal length of an \mathfrak{F}_G -proper resolution of M by \mathfrak{F}_G -projectives. We set $\mathfrak{F}_G \text{cd} G = \mathfrak{F}_G \text{pd} R$. Note that these finiteness conditions will not be defined unless R admits an \mathfrak{F}_G -proper resolution by \mathfrak{F}_G -projectives. One could think of \mathfrak{F}_G -cohomology as the "Gorenstein cohomology relative \mathfrak{F} ".

5.2.2. Technical results. We need some results for the \mathfrak{F}_G -cohomology whose analogues are well-known for Gorenstein cohomology [Hol04].

We say an RG-module M admits a *right* resolution by \mathfrak{F} -projectives if there exists an exact chain complex

$$0 \longrightarrow M \longrightarrow T_{-1} \longrightarrow T_{-2} \longrightarrow \cdots$$

where the T_i are \mathfrak{F} -projectives. \mathfrak{F} -strong right resolutions and \mathfrak{F} -split right resolutions are defined as for any chain complex.

LEMMA 5.2.3. An RG-module M is \mathfrak{F}_{G} -projective if and only if M satisfies

(*)
$$\mathfrak{FExt}^{i}_{RG}(M,Q) \cong 0 \text{ for all } \mathfrak{F}\text{-projective } Q$$

for all $i \geq 1$ and M admits a right \mathfrak{F} -strong \mathfrak{F} -split resolution by \mathfrak{F} -projectives.

PROOF. If M is the cokernel of a \mathfrak{F} -strong \mathfrak{F} -complete resolution T_* then for all $i \geq 1$ and any \mathfrak{F} -projective Q,

$$\mathfrak{F}Ext^i_{RG}(M,Q) \cong H^i \operatorname{Hom}_{RG}(T^+_*,Q)$$

where T_*^+ denotes the resolution $T_i^+ = T_i$ if $i \ge 0$ and $T_i^+ = 0$ for i < 0. Then (\star) follows because T_* is \mathfrak{F} -strong.

Conversely given (\star) and an \mathfrak{F} -strong right resolution T^-_* then let T^+_* be the standard \mathfrak{F} -split resolution for M (Lemma 5.1.2), (\star) ensures that T^+_* is \mathfrak{F} -strong and splicing together T^+_* and T^-_* gives the required resolution.

LEMMA 5.2.4. If $\mathfrak{F}pd N < \infty$ and M is \mathfrak{F}_G -projective then $\mathfrak{F}Ext^i_{RG}(M, N) = 0$ for all $i \geq 1$.

PROOF. Let $P_* \longrightarrow N$ be a \mathfrak{F} -split \mathfrak{F} -projective resolution then by a standard dimension shifting argument

$$\mathfrak{F}Ext^i(M,N) \cong \mathfrak{F}Ext^{i+j}(M,K_j)$$

where K_j is the j^{th} syzygy of P_* . Since K_j is \mathfrak{F} -projective for $j \ge n$ the result follows from Lemma 5.2.3.

PROPOSITION 5.2.5. Let A be any RG-module and $P_* \longrightarrow A$ a length n \mathfrak{F} -split resolution of A with $P_i \mathfrak{F}$ -projective for $i \geq 1$, then P_* is \mathfrak{F}_G -proper.

PROOF. The case n = 0 is obvious. If n = 1 then for any \mathfrak{F}_G -projective Q, there is a long exact sequence

$$0 \longrightarrow \operatorname{Hom}_{RG}(Q, P_1) \longrightarrow \operatorname{Hom}_{RG}(Q, P_0) \longrightarrow \operatorname{Hom}_{RG}(Q, A)$$
$$\longrightarrow \mathfrak{F}\operatorname{Ext}^1_{RG}(Q, P_1) \longrightarrow \cdots$$

but $\mathfrak{F}Ext^1_{RG}(Q, P_1) = 0$ by Lemma 5.2.4.

Assume $n \geq 2$ and let K_* be the syzygies of P_* , then there is an \mathfrak{F} -split resolution

$$0 \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_{i+1} \longrightarrow K_i \longrightarrow 0$$

so $\mathfrak{F}pd K_i < \infty$ for all $i \geq 0$. Thus every short exact sequence

$$0 \longrightarrow K_i \longrightarrow P_i \longrightarrow K_{i-1}$$

is \mathfrak{F}_G -proper by Lemma 5.2.4, so P_* is \mathfrak{F}_G -proper.

LEMMA 5.2.6 (Comparison Lemma). Let A and B be two RG-modules with \mathfrak{F} -strong \mathfrak{F} -split right resolutions by \mathfrak{F} -projectives called S^* and T^* respectively, then any map $f: A \to B$ lifts to a map f_* of complexes as shown below:

$$0 \longrightarrow A \longrightarrow S^{1} \longrightarrow S^{2} \longrightarrow \cdots$$
$$\left| \begin{array}{c} f \\ f \end{array} \right| f_{1} \\ 0 \longrightarrow B \longrightarrow T^{1} \longrightarrow T^{2} \longrightarrow \cdots$$

The map of complexes is unique up to chain homotopy and if f is \mathfrak{F} -split then so is f_* .

PROOF. The lemma without the \mathfrak{F} -splitting comes from dualising [EJ11, p.169], see also [Hol04, Proposition 1.8].

Assume f is \mathfrak{F} -split and consider the map of complexes restricted to RH for some finite subgroup H of G. Let ι_*^T and ι_*^S denote the splittings of the top and bottom rows and s_* the splitting of f_* , constructed only up to degree i - 1. The base case of the induction, when i = 0, holds because f is \mathfrak{F} -split.

$$\cdots \underbrace{ \overset{\partial_{i-2}^S}{\overset{i_{i-2}}{\underset{s_{i-1}}{\overset{s_{i-1}}{\underset{i_{i-1}}{\overset{s_{i-1}}{\underset{s_{i-1}}}{\underset{s_{i-1}}{\underset{s_{i-1}}}{\underset{s_{i-1}}{\underset{s_{i-1}}}{\underset{s_{i-1}}{\underset{s_{i-1}}}{\underset{s_{i-1}}{\underset{s_{i-1}}{\underset{s_{i-1}}{\underset{s_{i-1}}{\underset{s_{i-1}}{\underset{s_{i-1}}}{\underset{s_{i-1}}}{\underset{s_{i-1}}{\underset{s_{i$$

Let $s_i = \partial_{i-1}^S \circ s_{i-1} \circ \iota_{i-1}^T$. Then,

$$f_i \circ s_i = f_i \circ \partial_{i-1}^S \circ s_{i-1} \circ \iota_{i-1}^T$$
$$= \partial_{i-1}^T \circ f_{i-1} \circ s_{i-1} \circ \iota_{i-1}^T$$
$$= \partial_{i-1}^T \circ \iota_{i-1}^T$$
$$= \mathrm{id}_{T^i}$$

where the second equality is the commutativity condition coming from the fact that f_* is a chain map.

100

5.2.3. An Avramov–Martsinkovsky long exact sequence in \mathfrak{F} -cohomology.

THEOREM 5.2.7. Given an \mathfrak{F} -strong \mathfrak{F} -complete resolution of R there is a long exact sequence

$$0 \longrightarrow \mathfrak{F}_{G}H^{1}(G, -) \longrightarrow \cdots$$
$$\cdots \longrightarrow \widehat{\mathfrak{F}H}^{n-1}(G, -) \longrightarrow \mathfrak{F}_{G}H^{n}(G, -) \longrightarrow \mathfrak{F}H^{n}(G, -)$$
$$\longrightarrow \widehat{\mathfrak{F}H}^{n}(G, -) \longrightarrow \mathfrak{F}_{G}H^{n+1}(G, -) \longrightarrow \cdots$$

PROOF. We follow the proof in [ABS09, §3]. Let T_* be an \mathfrak{F} -strong \mathfrak{F} complete resolution coinciding with an \mathfrak{F} -projective \mathfrak{F} -split resolution P_* in degrees n and above. We may choose $\theta_* : T_* \longrightarrow P_*$ to be \mathfrak{F} -split by Lemma 5.2.6 and without loss of generality we may also assume that θ_i is surjective for all i.

Truncating at position 0 and adding cokernels gives the bottom two rows of the diagram below, the row above is the row of kernels. Note that the map $A \to R$ is necessarily surjective since the maps $T_0 \to P_0$ and $P_0 \to R$ are surjective.



We make some observations about the diagram: Firstly, since the module A is the cokernel of a \mathfrak{F} -strong \mathfrak{F} -complete resolution, A is \mathfrak{F}_G -projective. Secondly, in degree $i \geq 0$ the columns are \mathfrak{F} -split and the P_i are \mathfrak{F} -projective, thus the K_i are \mathfrak{F} -projective for all $i \geq 0$. Thirdly the far right vertical short exact sequence is \mathfrak{F} -split since the degree 0 column and the rows are \mathfrak{F} -split. Finally the top row is exact and \mathfrak{F} -split since the other two rows are.

Apply the functor $\operatorname{Hom}_{RG}(-, M)$ for an arbitrary RG-module M and take homology. This gives a long exact sequence

$$\cdots \longrightarrow \mathfrak{F}H^i(G,M) \longrightarrow \widehat{\mathfrak{F}H}^i(G,M) \longrightarrow H^i \operatorname{Hom}_{RG}(K_*,M) \longrightarrow \cdots$$

We can simplify the right-hand term:

$$H^{i}\operatorname{Hom}_{RG}(K_{*}, M) \cong \mathfrak{F}_{G}\operatorname{Ext}_{RG}^{i}(K, M)$$

 $\cong \mathfrak{F}_{G}H^{i+1}(G, M)$

where the first isomorphism is because, by Proposition 5.2.5, the top row is \mathfrak{F}_{G} -proper. For the second isomorphism note that the short exact sequence

$$0 \longrightarrow K \longrightarrow A \longrightarrow R \longrightarrow 0$$

is \mathfrak{F}_G -proper by Proposition 5.2.5, so

$$0 \longrightarrow K_{n-1} \longrightarrow \cdots \longrightarrow K_0 \longrightarrow A \longrightarrow R \longrightarrow 0$$

is an \mathfrak{F}_G -proper \mathfrak{F}_G -projective resolution of R. Thus the second isomorphism follows from the short exact sequence and Lemma 5.2.2.

COROLLARY 5.2.8. If there exists an \mathfrak{F} -strong \mathfrak{F} -complete resolution of R then $\mathfrak{F}_{G} \operatorname{cd} G < \infty$.

PROOF. In the proof of the theorem we assumed an \mathfrak{F} -strong \mathfrak{F} -complete resolution of R and built a finite length \mathfrak{F}_G -proper resolution of R by \mathfrak{F}_G -projectives.

PROPOSITION 5.2.9. If the Avramov–Martsinkovsky long exact sequence and the long exact sequence of Theorem 5.2.7 both exist, then there is a commutative diagram:

Where for conciseness we have written H^n for $H^n(G, -)$ etc.

PROOF. The Avramov-Martsinkovsky long exact sequence is constructed analogously to in the proof of Theorem 5.1.5, we give a quick sketch below as we will need the notation. Take a strong complete resolution T'_* of R coinciding with a projective resolution P'_* in high dimensions and let A' be the zeroth cokernel of T'_* . Thus A' is Gorenstein projective. Again, the map $T'_* \to P'_*$ is assumed surjective and the kernel K'_* is a projective resolution of K', the kernel of the map $A' \longrightarrow R$. Applying $\operatorname{Hom}_{RG}(-, M)$, for some RG-module M, to the short exact sequence of complexes

$$0 \longrightarrow K'_* \longrightarrow T'_* \longrightarrow P'_* \longrightarrow 0$$

gives the Avramov–Martsinkovsky long exact sequence.

Let T_* , P_* , K_* , K and A be as defined in the proof of Theorem 5.1.5. There is a commutative diagram of chain complexes

where the maps β exists by the comparison theorem for projective resolutions and γ exists by the comparison theorem for strong complete resolutions [CK97, Lemma 2.4]. The map α is the induced map on the kernels. Applying the functor $\operatorname{Hom}_{RG}(-, M)$ for some RG-module M, and taking homology, the maps α , β and γ induce the maps α_* , β_* and γ_* .

Finally we construct the map $\eta_n : GH^n(G, -) \longrightarrow \mathfrak{F}H^n(G, -)$. Let B_* be a G-proper Gorenstein projective resolution and recall P_* is an \mathfrak{F} -split resolution by \mathfrak{F} -projectives. Then B_* is \mathfrak{F} -split (Lemma 5.1.6) so there is a chain map $P_* \to B_*$ inducing η_* on cohomology.

Commutativity is obvious for the diagram with the maps η_i removed, leaving us with two relations to prove. Let

$$\varepsilon_n^G: \mathrm{G}H^n(G,-) \longrightarrow H^n(G,-)$$

denote the map from the commutative diagram. This is the map induced by comparison of a resolution of Gorenstein projectives and ordinary projectives [**ABS09**, 3.2,3.11]. We get $\beta_* \circ \eta_* = \varepsilon_*^G$, since all the maps are induced by comparison of resolutions, and such maps are unique up to chain homotopy equivalence.

The final commutativity relation, that $\eta_* \circ \alpha_* = \varepsilon_*^{\mathfrak{F}_G}$, is the most difficult to show. Here

$$\varepsilon_n^{\mathfrak{F}_G}:\mathfrak{F}_GH^n(G,-)\longrightarrow\mathfrak{F}_H^n(G,-)$$

denotes the map from the commutative diagram, it is induced by comparison of resolutions.

Here is a commutative diagram showing the resolutions involved:

$$0 \longrightarrow K \longrightarrow A \longrightarrow R \longrightarrow 0$$

$$0 \longrightarrow K_{*} \xrightarrow{\checkmark} T_{*} \xrightarrow{\checkmark} P_{*} \xrightarrow{\checkmark} 0$$

$$0 \longrightarrow K'_{*} \longrightarrow T'_{*} \longrightarrow P_{*} \xrightarrow{\checkmark} 0$$

$$0 \longrightarrow K'_{*} \longrightarrow T'_{*} \longrightarrow P'_{*} \longrightarrow 0$$

Let L_* be the chain complex defined by $L_i = K_{i-1}$ for all $i \ge 1$ and $L_0 = A$, with boundary map at i = 1 the composition of the maps $K_0 \to K$ and $K \to A$. Thus L_* is acyclic except at degree zero where $H_0L_* = R$. Similarly, let L'_* denote chain complex with $L'_i = K'_{i-1}$ for all $i \ge 1$ and $L'_0 = A'$ augmented by A', so L'_* is acyclic except at degree zero where $H_0L'_* = R$. Note that L_* is an \mathfrak{F}_G -proper resolution of R by Proposition 5.2.5 and L'_* is a G-proper resolution of R by the Gorenstein cohomology version of the same proposition.

Recall that the maps $\varepsilon_*^{\mathfrak{F}_G}$ and η_* are induced by comparison of resolutions: $\varepsilon_*^{\mathfrak{F}_G}$ is induced by a map $P_* \to L_*$ and η_* is induced by a map $P_* \to L'_*$. The map

$$\mathfrak{F}_{\mathrm{G}}\mathrm{Ext}^{i}_{RG}(K,-)\longrightarrow \mathrm{GExt}^{i}_{RG}(K',-)$$

is induced by $\alpha: K'_* \longrightarrow K_*$. Thus the map

$$\alpha_*:\mathfrak{F}_{\mathbf{G}}H^n(G,-)\longrightarrow \mathbf{G}H^n(G,-)$$

is induced by $L'_* \longrightarrow L_*$. The diagram below is the one we must show commutes.

$$\mathfrak{F}_{\mathbf{G}}H^{n}(G,-)\cong H^{n}\operatorname{Hom}_{RG}(L_{*},-)\xrightarrow{\varepsilon_{n}^{\mathfrak{F}_{G}}}\mathfrak{F}^{H^{n}}(G,-)\cong H^{n}\operatorname{Hom}_{RG}(P_{*},-)$$

$$\downarrow^{\alpha_{n}}$$

$$\mathfrak{G}H^{n}(G,-)\cong H^{n}\operatorname{Hom}_{RG}(L'_{*},-)$$

Since the composition P_* to L'_* to L_* is a map of resolutions from P_* to L_* , and such maps are unique up to chain homotopy equivalence, this completes the proof.

COROLLARY 5.2.10. Given an \mathfrak{F} -strong \mathfrak{F} -complete resolution of R, $\operatorname{Gcd} G = n < \infty$ implies $\mathfrak{F}H^i(G, -)$ injects into $\widehat{\mathfrak{F}H}^i(G, -)$ for all $i \ge n+1$.

PROOF. Gcd $G < \infty$ implies the Avramov–Martsinkovsky long exact sequence exists (Theorem 5.1.5). Consider the the commutative diagram of Proposition 5.2.9. The map

$$\mathfrak{F}_{\mathbf{G}}H^{i}(G,-)\longrightarrow \mathfrak{F}H^{i}(G,-)$$

factors as $\eta_i \circ \alpha_i = 0$, so since $GH^i(G, -) = 0$ for all $i \ge n+1$, $\mathfrak{F}H^i(G, -)$ injects into $\widehat{\mathfrak{F}H}^i(G, -)$ for all $i \ge n+1$.

THEOREM 5.2.11. If $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G = \operatorname{Gcd} G$.

PROOF. We know already that $\operatorname{Gcd} G \leq \operatorname{\mathfrak{F}cd} G$ (see Section 1.6). If $\operatorname{\mathfrak{F}cd} G < \infty$ then it is trivially true that \mathfrak{F} admits an \mathfrak{F} -strong \mathfrak{F} -complete resolution, thus $\mathfrak{F}H^i(G,-)$ injects into $\operatorname{\mathfrak{F}H}^i(G,-)$ for all $i \geq \operatorname{Gcd} G + 1$, but $\operatorname{\mathfrak{F}H}^i(G,-)$ is always zero since $\operatorname{\mathfrak{F}cd} G < \infty$ [**Kro93**, 4.1(i)].

EXAMPLE 5.2.12. Let $R = \mathbb{Z}$ for this example. Kropholler introduced the class $\mathbf{H}\mathfrak{F}$ of hierarchically decomposable groups in [**Kro93**] as the smallest class of groups such that if there exists a finite-dimensional contractible *G*-CW complex with stabilisers in $\mathbf{H}\mathfrak{F}$ then $G \in \mathbf{H}\mathfrak{F}$. Let $\mathbf{H}\mathfrak{F}_b$ denote the subclass of $\mathbf{H}\mathfrak{F}$ containing groups with a bound on the orders of their finite subgroups.

The $\mathbb{Z}G$ -module $B(G,\mathbb{Z})$ of bounded functions from G to \mathbb{Z} was first studied in [**KT91**], Kropholler and Mislin proved that if G is $\mathbf{H}\mathfrak{F}$ with a bound on lengths of chains of finite subgroups and $\operatorname{pd}_{\mathbb{Z}G}B(G,\mathbb{Z}) < \infty$ then $\mathcal{O}_{\operatorname{fin}}\operatorname{cd} G < \infty$, in particular $\operatorname{\mathfrak{F}cd} G < \infty$ [**KM98**, Theorem B]. If $\operatorname{Gcd} G < \infty$ then $\operatorname{pd}_{\mathbb{Z}G}B(G,\mathbb{Z}) < \infty$ ∞ [**ABS09**, 2.10][**CK98**, Theorem C]. Thus if $G \in \operatorname{H}\mathfrak{F}_b$ then $\operatorname{Gcd} G = \operatorname{\mathfrak{F}cd} G$.

5.3. Group extensions

Recall Theorem 4.5.1, that $\mathfrak{F}cd G = \mathcal{H}_{\mathcal{F}in}cd G$ for all groups. The invariant $\mathcal{H}_{\mathcal{F}in}cd G$ was studied by Degrijse in [**Deg13a**] where he proves the following (though stated for $\mathcal{H}_{\mathcal{F}in}cd G$ not $\mathfrak{F}cd G$):

THEOREM 5.3.1. [Deg13a, Theorem B] Let

$$1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1$$

be a short exact sequence of groups such that every finite index overgroup of N in G has a bound on the orders of the finite subgroups not contained in N. If $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G \leq \operatorname{\mathfrak{F}cd} N + \operatorname{\mathfrak{F}cd} Q$.

Since Gorenstein cohomological dimension is subadditive under extensions [BDT09, Remark 2.9(2)], an application of Theorem 5.2.11 removes the condition on the orders of finite subgroups:

COROLLARY 5.3.2. Given a short exact sequence of groups

 $1 \longrightarrow N \longrightarrow G \longrightarrow Q \longrightarrow 1$

if $\operatorname{\mathfrak{F}cd} G < \infty$ then $\operatorname{\mathfrak{F}cd} G \leq \operatorname{\mathfrak{F}cd} N + \operatorname{\mathfrak{F}cd} Q$.

For further discussion on the behaviour of $\operatorname{\mathfrak{F}cd} G$ (equivalently $\mathcal{H}_{\operatorname{fin}} \operatorname{cd} G$) under group extensions and other standard constructions see Section 4.5.1.

5.4. Rational cohomological dimension

For this section, let $R = \mathbb{Z}$. Gandini has shown that for groups in $\mathbf{H}\mathfrak{F}$, $\mathrm{cd}_{\mathbb{Q}} G \leq \mathrm{Gcd} G$ [Gan12b, Remark 4.14] and this is the only result we are aware of relating $\mathrm{cd}_{\mathbb{Q}} G$ and $\mathrm{Gcd} G$. In Proposition 5.4.4 we show that $\mathrm{cd}_{\mathbb{Q}} G \leq \mathrm{Gcd} G$ for all groups with $\mathrm{cd}_{\mathbb{Q}} G < \infty$. Recall there are examples of torsion-free groups with $\mathrm{cd}_{\mathbb{Q}} G < \mathrm{cd}_{\mathbb{Z}} G$ [Dav08, Example 8.5.8] and $\mathrm{Gcd} G = \mathrm{cd}_{\mathbb{Z}} G$ whenever $\mathrm{cd}_{\mathbb{Z}} G < \infty$ [ABS09, Corollary 2.9], so we cannot hope for equality of $\mathrm{cd}_{\mathbb{Q}} G$ and $\mathrm{Gcd} G$ in general.

QUESTION 5.4.1. Are there groups G with $\operatorname{Gcd} G < \infty$ but $\operatorname{cd}_{\mathbb{Q}} G = \infty$?

Recall from Section 1.6 that silp RG denotes the supremum of the injective lengths (injective dimensions) of all projective RG-modules and spli RG denotes the infimum of the projective lengths (projective dimensions) of all injective RGmodules.

LEMMA 5.4.2. For any group G, $\operatorname{silp} \mathbb{Q}G \leq \operatorname{silp} \mathbb{Z}G$.

PROOF. Using [**Emm10**, Theorem 4.4], $\operatorname{silp} \mathbb{Q}G = \operatorname{spli} \mathbb{Q}G$ and $\operatorname{silp} \mathbb{Z}G = \operatorname{spli} \mathbb{Z}G$. Combining with [**GG87**, Lemma 6.4] that $\operatorname{spli} \mathbb{Q}G \leq \operatorname{spli} \mathbb{Z}G$ gives the result.

LEMMA 5.4.3. If $\operatorname{Gcd} G < \infty$ then for any $\mathbb{Q}G$ -module M there is a natural isomorphism

$$\widehat{H}^*(G,M)\otimes \mathbb{Q}\cong \widehat{\operatorname{Ext}}^*_{\mathbb{O}G}(\mathbb{Q},M).$$

PROOF. Let T_* be a strong complete resolution of \mathbb{Z} by $\mathbb{Z}G$ -modules, then $T_* \otimes \mathbb{Q}$ is a strong complete resolution of \mathbb{Q} by $\mathbb{Q}G$ -modules. By an obvious generalisation of [**MT00**, Lemma 2.2], if silp $\mathbb{Q}G \leq \infty$ then any complete $\mathbb{Q}G$ -module resolution is a strong complete $\mathbb{Q}G$ -module resolution, so since silp $\mathbb{Q}G < \operatorname{silp} \mathbb{Z}G \leq \infty$, $T_* \otimes \mathbb{Q}$ is a strong complete resolution. This gives a chain of isomorphisms for any $\mathbb{Q}G$ -module M:

$$H^*(G, M) \otimes \mathbb{Q} \cong H^* \operatorname{Hom}_{\mathbb{Z}G}(T_*, M) \otimes \mathbb{Q}$$
$$\cong H^* \operatorname{Hom}_{\mathbb{Q}G}(T_* \otimes \mathbb{Q}, M)$$
$$\cong \widehat{\operatorname{Ext}}^*_{\mathbb{Q}G}(\mathbb{Q}, M).$$

PROPOSITION 5.4.4. If $\operatorname{cd}_{\mathbb{Q}} G < \infty$ then $\operatorname{cd}_{\mathbb{Q}} G \leq \operatorname{Gcd} G$.

PROOF. There is nothing to show if $\operatorname{Gcd} G = \infty$ so assume that $\operatorname{Gcd} G < \infty$. Since \mathbb{Q} is flat over \mathbb{Z} , tensoring the Avramov–Martsinkovsky long exact sequence with \mathbb{Q} preserves exactness. Combining this with Lemma 5.4.3 and the wellknown fact that for any $\mathbb{Q}G$ -module M there is a natural isomorphism [**Bie81**, p.2]

$$H^*(G, M) \otimes \mathbb{Q} \cong \operatorname{Ext}^*_{\mathbb{O}G}(\mathbb{Q}, M)$$

gives the long exact sequence

$$\cdots \longrightarrow \mathrm{G}H^i(G,M) \otimes \mathbb{Q} \longrightarrow \mathrm{Ext}^i_{\mathbb{Q}G}(\mathbb{Q},M) \longrightarrow \widehat{\mathrm{Ext}}^i_{\mathbb{Q}G}(\mathbb{Q},M) \longrightarrow \cdots$$

Since $\operatorname{cd}_{\mathbb{Q}} G < \infty$, we have that $\widehat{\operatorname{Ext}}_{\mathbb{Q}G}^{i}(\mathbb{Q}, M) = 0$ [**Kro93**, 4.1(i)]. Thus there is an isomorphism for all i,

$$GH^i(G,M) \otimes \mathbb{Q} \cong \operatorname{Ext}^i_{\mathbb{O}G}(\mathbb{Q},M)$$

and the result follows.

CHAPTER 6

Bredon duality groups

This chapter contains material that has appeared in:

• Bredon–Poincaré duality groups (2013, to appear J. Group Theory) [SJG13a].

In this chapter we study Bredon duality and Bredon–Poincaré duality groups. Recall that a *Bredon duality group over* R is a group G of type $\mathcal{O}_{\mathcal{F}in}$ FP over Rsuch that for every finite subgroup H of G there is an integer d_H with

$$H^{i}(WH, R[WH]) = \begin{cases} R\text{-flat} & \text{if } i = d_{H}, \\ 0 & \text{else.} \end{cases}$$

Furthermore, G is said to be *Bredon–Poincaré duality* over R if for all finite subgroups H,

$$H^{d_H}(WH, R[WH]) = R.$$

In Section 6.2 we give several sources of examples of both Bredon duality and Bredon–Poincaré duality groups, including the example below of Jonathan Block and Schmuel Weinberger, suggested to us by Jim Davis.

THEOREM 6.2.7. There exist examples of Bredon–Poincaré duality groups over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H but G doesn't admit a cocompact manifold model M for $\mathbb{E}_{\text{fin}}G$.

This is a counterexample to a possible generalisation of Wall's conjecture, which asks if a finitely presented Poincaré duality group admits a cocompact manifold model for EG: Let G be Bredon–Poincaré duality over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H, does G admit a cocompact manifold model M for $\mathbb{E}_{\text{fin}}G$?

Section 6.4 contains an analysis of Bredon duality and Bredon–Poincaré duality groups of low dimension and Section 6.5 looks at when these properties are preserved under group extensions.

Recall that given a Bredon duality group G of dimension n we write $\mathcal{V}(G)$ for the set

 $\mathcal{V}(G) = \{ d_F : F \text{ a non-trivial finite subgroup of } G \} \subseteq \{0, \dots, n\}.$

In Example 6.6.8 we build Bredon duality groups with arbitrary $\mathcal{V}(G)$ and in Section 6.3 we build Bredon–Poincaré duality groups with a large selection of $\mathcal{V}(G)$, although we are unable to produce arbitrary $\mathcal{V}(G)$.

One might hope to give a definition of Bredon–Poincaré duality groups in terms of Bredon cohomology only, we show in Section 6.7 that the naïve idea of asking that a group be \mathcal{O}_{fin} FP with

$$H^{i}_{\mathcal{O}_{\mathcal{F}_{in}}}(G, R[?, -]_{\mathcal{O}_{\mathcal{F}_{in}}}) \cong \begin{cases} \underline{R} & \text{if } i = n, \\ 0 & \text{else,} \end{cases}$$

is not the correct definition, namely we show in Theorem 6.7.3 that any such group is necessarily a torsion-free Poincaré duality group over R.

6.1. Preliminary observations

Recall that a group G is *R*-torsion-free if the order of every finite subgroup of G is invertible in R, equivalently the order of every finite order element is invertible in R (see page 34).

Recall that a Bredon duality group is said to be dimension n if $\mathcal{O}_{\text{fin}} \operatorname{cd} G = n$.

LEMMA 6.1.1. If G is Bredon duality of dimension n over \mathbb{Z} then G is Bredon duality of dimension n over any ring R, with the same values of d_H for all finite subgroups H.

PROOF. Since G is $\mathcal{O}_{\mathcal{f}in}$ FP over Z, G is $\mathcal{O}_{\mathcal{f}in}$ FP over R (Lemma 3.7.1). Also because G is $\mathcal{O}_{\mathcal{f}in}$ FP over Z, WH is FP_{∞} over Z for all finite subgroups H (Corollary 3.6.4) and we may apply [**Bie81**, Corollary 3.6] to get a short exact sequence

$$0 \longrightarrow H^{q}(WH, \mathbb{Z}[WH]) \otimes_{\mathbb{Z}} R \longrightarrow H^{q}(WH, R \otimes_{\mathbb{Z}} \mathbb{Z}[WH])$$
$$\longrightarrow \operatorname{Tor}_{1}^{\mathbb{Z}}(H^{q+1}(WH, \mathbb{Z}[WH]), R) \longrightarrow 0.$$

 $H^{q+1}(WH, \mathbb{Z}[WH])$ is \mathbb{Z} -flat for all q giving an isomorphism

$$H^q(WH, \mathbb{Z}[WH]) \otimes_{\mathbb{Z}} R \cong H^q(WH, R[WH]).$$

Observing that if an Abelian group M is \mathbb{Z} -flat then $M \otimes R$ is R-flat completes the proof.

LEMMA 6.1.2. If G is R-torsion-free and Bredon duality of dimension n over R then $d_H = \operatorname{cd}_R WH$ and $d_1 \leq n$.

To prove the Lemma we need the following proposition, an analogue of [Bro94, VIII.6.7] for arbitrary rings R and proved in exactly the same way.

PROPOSITION 6.1.3. If G is FP over R then

 $\operatorname{cd}_R G = \max\{n : H^n(G, RG) \neq 0\}.$

PROOF OF LEMMA 6.1.2. If G is R-torsion-free then for any finite subgroup H,

$$\operatorname{cd}_R N_G H \le \operatorname{cd}_R G \le \mathcal{O}_{\operatorname{Fin}} \operatorname{cd}_R G$$

and $N_G H$ is FP_{∞} over R by Corollary 3.6.4 and Lemma 3.7.2. The short exact sequence

 $1 \longrightarrow H \longrightarrow N_G H \longrightarrow W H \longrightarrow 1$

and Lemma 6.5.4 implies that

$$H^i(N_GH, R[N_GH]) \cong H^i(WH, R[WH]).$$

Thus Proposition 6.1.3 shows $d_H = \operatorname{cd}_R N_G H = \operatorname{cd}_R W H$. Finally, $d_1 \leq n$ because $\operatorname{cd}_R G \leq \mathcal{O}_{\operatorname{fin}} \operatorname{cd}_R G$ (Lemma 3.7.2).

In the proposition below $\mathfrak{F}cd G$ denotes the \mathfrak{F} -cohomological dimension (see Section 1.4) and $\operatorname{Gcd} G$ denotes the Gorenstein cohomological dimension (see Section 1.6).

This proposition implies that if G is Bredon–Poincaré duality over R then $\operatorname{Gcd} G = \operatorname{\mathfrak{Fcd}} G = d_1$ and if G is also virtually torsion-free then $\operatorname{vcd} G = d_1$ also.

PROPOSITION 6.1.4. If G is $\operatorname{FP}_{\infty}$ with $\mathcal{O}_{\operatorname{Fin}} \operatorname{cd} G < \infty$ then

$$\operatorname{Gcd} G = \operatorname{\mathfrak{Fcd}} G = \sup\{n : H^n(G, RG) \neq 0\},\$$

and if G is also virtually torsion-free then $\operatorname{vcd} G = \operatorname{Gcd} G$ also.

PROOF. This proof uses an argument due to Degrijse and Martínez-Pérez in [**DMP13**]. By [**Hol04**, Theorem 2.20] the Gorenstein cohomological dimension can be characterised as

 $\operatorname{Gcd} G = \sup\{n : H^n(G, P) \neq 0 \text{ for } P \text{ any projective } RG\text{-module}\}.$

As G is $\operatorname{FP}_{\infty}$ we need only check when P = RG. Since $\operatorname{\mathfrak{Fcd}} G \leq \mathcal{O}_{\operatorname{fin}} \operatorname{cd} G < \infty$, we can conclude that $\operatorname{\mathfrak{Fcd}} G = \operatorname{Gcd} G$ (Theorem 5.2.11) and finally for virtually torsion-free groups $\operatorname{\mathfrak{Fcd}} G = \operatorname{vcd} G$ [**MPN06**, Theorem 5.1].

6.2. Examples

In this section we provide several sources of examples of Bredon duality and Bredon–Poincaré duality groups, showing that these properties are not too rare. **6.2.1.** Smooth actions on manifolds. Recall from the introduction that if G has a cocompact manifold model M for $\mathbb{E}_{\mathcal{F}in}G$ such that M^H is a submanifold for all finite subgroups H then G is Bredon–Poincaré duality. The following lemma gives a condition which guarantees that M^H is a submanifold of M.

LEMMA 6.2.1. [**Dav08**, 10.1 p.177] If G is a discrete group acting properly and locally linearly on a manifold M then the fixed points subsets of finite subgroups of G are submanifolds of M.

Locally linear is a technical condition, the definition of which can be found in [**Dav08**, Definition 10.1.1], for our purposes it is enough to know that if Mis a smooth manifold and G acts by diffeomorphisms then the action is locally linear. The locally linear condition is necessary however—in [**DL03**] examples are given of virtually torsion-free groups acting as a discrete cocompact group of isometries of a CAT(0) manifold which are not Bredon duality.

EXAMPLE 6.2.2. Let p be a prime, let K be a cyclic group of order p, and let G be the wreath product

$$G = \mathbb{Z} \wr K = \left(\bigoplus_{K} \mathbb{Z}\right) \rtimes K.$$

G acts properly and by diffeomorphisms on \mathbb{R}^p : The copies of \mathbb{Z} act by translation along the axes, and the *K* permutes the axes. The action is cocompact with fundamental domain the quotient of the *p*-torus by the action of *K*. The finite subgroup *K* is a representative of the only conjugacy class of finite subgroups in *G*, and has fixed point set the line $\{(\lambda, \dots, \lambda) : \lambda \in \mathbb{R}\}$. If $z = (z_1, \dots, z_p) \in \mathbb{Z}^p$ then

$$(\mathbb{R}^p)^{K^z} = \{ (\lambda + z_1, \dots, \lambda + z_p) : \lambda \in \mathbb{R} \}.$$

Hence \mathbb{R}^p is a model for $\mathbb{E}_{\text{fin}}G$ and, invoking Lemma 6.2.1, G is a Bredon–Poincaré duality group of dimension p.

We can explicitly calculate the Weyl group WK: let L denote the copy of \mathbb{Z} inside \mathbb{Z}^p generated by $(1, 1, \ldots, 1)$, then the normaliser $N_G K$ is $L \rtimes K$ and thus the Weyl group WK is isomorphic to \mathbb{Z} . Since K is a representative of the only non-trivial conjugacy class of finite subgroups it provides the only element of $\mathcal{V}(G)$, thus $\mathcal{V}(G) = \{1\}$.

EXAMPLE 6.2.3. Fixing positive integers $m \leq n$, if $G = \mathbb{Z}^n \rtimes C_2$ where C_2 , the cyclic group of order 2, acts as the antipodal map on $\mathbb{Z}^{n-m} \leq \mathbb{Z}^n$ then

$$N_G C_2 = C_G C_2 = \{ g \in G : gz = zg \}.$$

6.2. EXAMPLES

But this is exactly the fixed points of the action of C_2 on G, hence $N_G C_2 = \mathbb{Z}^m \rtimes C_2$ and

$$H^{i}(W_{G}C_{2}, R[W_{G}C_{2}]) \cong \begin{cases} R & \text{if } i = m, \\ 0 & \text{else.} \end{cases}$$

G embeds as a discrete subgroup of $\text{Isom}(\mathbb{R}^n) = \mathbb{R}^n \rtimes GL_n(\mathbb{R})$ and acts properly and cocompactly on \mathbb{R}^n . It follows that *G* is \mathcal{O}_{fin} FP and \mathcal{O}_{fin} cd G = n so *G* is Bredon–Poincaré duality of dimension *n* over any ring *R* with $\mathcal{V} = \{m\}$.

EXAMPLE 6.2.4. Similarly to the previous example we can take

$$G = \mathbb{Z}^n \rtimes \bigoplus_{i=1}^n C_2$$

where the j^{th} copy of C_2 acts antipodally on the j^{th} copy of \mathbb{Z} in \mathbb{Z}^n . Note that G is isomorphic to $(D_{\infty})^n$ where D_{∞} denotes the infinite dihedral group. As before G embeds as a discrete subgroup of $\text{Isom}(\mathbb{R}^n) = \mathbb{R}^n \rtimes GL_n(\mathbb{R})$ and acts properly and cocompactly on \mathbb{R}^n . Thus G is \mathcal{O}_{fin} FP and \mathcal{O}_{fin} cd G = n, so G is Bredon–Poincaré duality of dimension n over any ring R with $\mathcal{V}(G) = \{0, \ldots, n\}$.

More generally, we could take a subgroup $\bigoplus_{i=1}^{m} C_2 \hookrightarrow \bigoplus_{i=1}^{n} C_2$ and form the semi-direct product of \mathbb{Z}^n with this subgroup. Although this gives us a range of possible values for $\mathcal{V}(G)$ it is impossible to produce a full range of values. Consider the case m = 2, so we have a group

$$G = \mathbb{Z}^n \rtimes (A \times B)$$

where $A \cong B \cong C_2$, and both A and B act either trivially or antipodally on each coordinate of \mathbb{Z}^n . We can describe the normaliser $N_G A$ by an element $(a_1, \ldots, a_n) \in \{0, 1\}^n$, so A acts trivially on the i^{th} copy of \mathbb{Z} if $a_i = 1$ and acts antipodally otherwise. Thus,

$$N_G A = \left(\bigoplus_{i=1}^n \left\{ \begin{array}{cc} \mathbb{Z} & \text{if } a_i = 1 \\ 0 & \text{else.} \end{array} \right\} \right) \rtimes (A \times B).$$

Similarly we can describe $N_G B$ by an element $(b_1, \ldots, b_n) \in \{0, 1\}^n$. One calculates that the normaliser $N_G(A \times B)$ is described by the element

$$(a_1 \wedge b_1, \ldots, a_n \wedge b_n)$$

where \wedge denotes the boolean AND function. This is because the i^{th} copy of \mathbb{Z} is normalised by $A \times B$ if and only if it is normalised by A and also by B.

If C denotes the subgroup of $A \times B$ generated by the element (1, 1) then the normaliser of $N_G C$ is described by the element

$$(\neg (a_1 \oplus b_1), \ldots, \neg (a_n \oplus b_n))$$

where \oplus denotes the boolean XOR function, and \neg the unary negation operator.

Now, using the above it can be shown that, for example, a Bredon–Poincaré duality group of dimension 4 with the form

$$G = \mathbb{Z}^4 \rtimes \bigoplus_{i=1}^m C_2$$

cannot have $\mathcal{V}(G) = \{1, 3\}$. Assume that such a G exists, clearly $m \geq 2$, let A and B denote two of the C_2 summands of $\bigoplus_{1=1}^m C_2$. Without loss of generality we can assume that A and B don't have the same action on \mathbb{Z}^4 . If $d_A = d_B = 1$ then by the description of the normaliser of $A \times B$ above, $d_{A \times B} = 0$, a contradiction. If $d_A = d_B = 3$ then in order for A and B not to have the same action on \mathbb{Z}^4 , we must have (up to some reordering of the coordinates)

$$(a_1, \dots, a_4) = (1, 1, 1, 0)$$

 $(b_1, \dots, b_4) = (0, 1, 1, 1).$

So $d_{A \times B} = 2$, a contradiction. Finally, if $d_A = 1$ and $d_B = 3$ then let C be the subgroup of $A \times B$ generated by (1,1). There are two possibilities, up to reordering of the coordinates, either

$$(a_1, \dots, a_4) = (1, 1, 1, 0)$$

 $(b_1, \dots, b_4) = (1, 0, 0, 0)$

or

$$(a_1, \dots, a_4) = (1, 1, 1, 0)$$

 $(b_1, \dots, b_4) = (0, 0, 0, 1).$

In the first case, $d_C = 2$, and in the second case $d_{A \times B} = 0$, both contradictions.

EXAMPLE 6.2.5. In [**FW08**, Theorem 6.1], Farb and Weinberger construct a Bredon–Poincaré duality group G arising from a proper cocompact action on \mathbb{R}^n by diffeomorphisms, however G is not virtually torsion-free.

REMARK 6.2.6. Restrictions on the dimensions of the fixed point sets. Suppose G is a group acting smoothly on an m-dimensional manifold M, and suppose furthermore that G contains a finite cyclic subgroup C_p fixing a point $x \in M$. There is an induced linear action of C_p on the tangent space $T_x M \cong \mathbb{R}^m$, equivalently a representation of C_p into the orthogonal group O(m). We can use this to give some small restrictions on the possible dimensions of the submanifold M^{C_p} , and hence on the values of d_{C_p} .

A representation of C_p in O(m) is simply a matrix N with $N^p = 1$. Using the Jordan–Chevalley decomposition (expressing a matrix as the product of its semi-simple and nilpotent parts), we see that N is semi-simple, so viewing N as a matrix over \mathbb{C} it is diagonalisable. However, since $N^p = 1$ and the characteristic

6.2. EXAMPLES

polynomial has coefficients in \mathbb{R} , all the eigenvalues come in pairs ω, ω^{-1} , where ω is a p^{th} root of unity. Thus N is conjugate via complex matrices to

$$\begin{pmatrix} \omega_1 & & & \\ & \omega_1^{-1} & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \frac{\omega_m}{2} & \\ & & & & \frac{\omega_m^{-1}}{2} & \\ & & & & \frac{\omega_{m-1}}{2} & \\ & & & & \frac{\omega_{m-1}}{2} & \\ & & & & \frac{\omega_{m-1}}{2} & \\ & & & &$$

depending on whether m is even or odd. The blank space in the matrices should be filled with zeros. Note that the ± 1 term can only be a -1 if p = 2. The matrix

$$\begin{pmatrix} \omega & 0 \\ 0 & \omega^{-1} \end{pmatrix}$$

is conjugate via complex matrices to

$$R_{\theta} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

Thus N is conjugate via complex matrices to $R_{\theta_1} \oplus \cdots \oplus R_{\theta_{m/2}}$ or $R_{\theta_1} \oplus \cdots \oplus R_{\theta_{(m-1)/2}} \oplus (\pm 1)$, and by [**Zha11**, 5.11], they are conjugate via real matrices as well. Hence the dimensions of the fixed point sets are the same. Noting that the rotation matrix R_{θ} fixes only the origin when $\theta \neq 0$, we conclude that for $p \neq 2$, the fixed point set M^{C_p} must be even dimensional if m is even, and odd dimensional otherwise.

Consider the case that G is a Bredon–Poincaré duality group, arising from a smooth cocompact action on an m-dimensional manifold M, and C_p for $p \neq 2$ is some finite subgroup of G. Then d_{C_p} is exactly the dimension of the submanifold M^{C_p} , and by the discussion above d_{C_p} is odd dimensional if m is odd dimensional, even dimensional otherwise. As demonstrated by Example 6.2.3, there are no restrictions when p = 2.

6.2.2. A counterexample to the generalised PD^n conjecture. Let G be Bredon–Poincaré duality over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H. One might ask if G admits a cocompact manifold model M for $E_{\mathcal{F}in}G$. This is generalisation of the famous PD^n -conjecture, due to Wall [Wal79]. This example is due to Jonathan Block and Schmuel Weinberger and was suggested to us by Jim Davis.

THEOREM 6.2.7. There exist examples of Bredon–Poincaré duality groups over \mathbb{Z} , such that WH is finitely presented for all finite subgroups H, but there doesn't exist a cocompact manifold model M for $\mathbb{E}_{\text{fin}}G$. Combining Theorems 1.5 and 1.8 of [**BW08**] gives the following example.

THEOREM 6.2.8 (Block–Weinberger). There exists a short exact sequence of groups

$$1 \longrightarrow K \longrightarrow G \longrightarrow Q \longrightarrow 1$$

with Q finite, such that

- (1) All torsion in G is contained in K.
- (2) There exists a cocompact manifold model for $E_{\text{Fin}}K$.
- (3) $\operatorname{gd}_{\operatorname{Fin}} G < \infty$.
- (4) There exists no manifold model for $E_{\mathcal{F}in}G$.

PROOF OF THEOREM 6.2.7. Let G be one of the groups constructed by Block and Weinberger in the theorem above. Since K has a cocompact model for $E_{\mathcal{f}in}K$ it has finitely many conjugacy classes of finite subgroups hence G has finitely many conjugacy classes of finite subgroups, since all torsion in G is contained in K. Let H be a finite subgroup of G, so H is necessarily a subgroup of Kand the normaliser $N_K H$ is finite index in $N_G H$. Since there is a cocompact model for $E_{\mathcal{f}in}K$, the normaliser $N_K H$ is FP_{∞} and finitely presented [LM00, Theorem 0.1] hence $N_G H$ and $W_G H$ are FP_{∞} and finitely presented too [Bro94, VIII.5.1][Rob96, 2.2.5]. Using Corollary 3.6.4, G is of type $\mathcal{O}_{\mathcal{f}in}FP$.

Finally, using [**Bro94**, III.(6.5)], there is a chain of isomorphisms for all natural numbers i,

$$H^{i}(W_{G}H, R[W_{G}H]) \cong H^{i}(N_{G}H, R[N_{G}H])$$
$$\cong H^{i}(N_{K}H, R[N_{K}H])$$
$$\cong H^{i}(W_{K}H, R[W_{K}H])$$

proving that the Weyl groups of finite subgroups have the correct cohomology. \Box

REMARK 6.2.9. Although it doesn't appear in the statements of [**BW08**, Theorems 1.5, 1.8], Block and Weinberger do prove that there is a cocompact model for $E_{\mathcal{F}in}G$, in their notation this is the space \widetilde{X} .

6.2.3. Actions on *R*-homology manifolds. Following [DL98] we define an *R*-homology *n*-manifold to be a locally finite simplicial complex *M* such that the link σ of every *i*-simplex of *M* satisfies

$$H_j(\sigma, R) = \begin{cases} R & \text{if } j = n - i - 1 \text{ or } j = 0, \\ 0 & \text{else,} \end{cases}$$

for all *i* such that $n - i - 1 \ge 0$ and the link is empty if n - i - 1 < 0. In particular *M* is an *n*-dimensional simplicial complex. *M* is called *orientable* if we can choose an orientation for each *n*-simplex which is consistent along the (n-1)-simplices and we say that M is R-orientable if either M is orientable or if R has characteristic 2.

A topological space X is called R-acyclic if the reduced homology $\hat{H}_*(X, R)$ is trivial.

THEOREM 6.2.10. If G is a group acting properly and cocompactly on an R-acyclic R-orientable R-homology n-manifold M then

$$H^{i}(G, RG) \cong \begin{cases} R & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

PROOF. By [**Dav08**, Lemma F.2.2] $H^i(G, RG) \cong H^i_c(M, R)$, where H^i_c denotes cohomology with compact supports. By Poincaré duality for *R*-orientable *R*-homology manifolds (see for example [**DL98**, Theorem 5]), there is a duality isomorphism $H^i_c(M, R) \cong H_{n-i}(M, R)$. Finally, since *M* is assumed *R*-acyclic,

$$H_{n-i}(M,R) \cong \begin{cases} R & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

EXAMPLE 6.2.11. In [**DL98**, Example 3], Dicks and Leary construct a group which is Poincaré duality over R, arising from an action on an R-orientable Racyclic R-homology manifold, but which is not Poincaré duality over \mathbb{Z} . Here Rmay be any ring for which a fixed prime q is invertible, for example $R = \mathbb{F}_p$ for $p \neq q$ or $R = \mathbb{Q}$.

COROLLARY 6.2.12. Let G be a group which admits a cocompact model X for $E_{\text{fin}}G$ such that for every finite subgroup H of G, X^H is an R-orientable R-acyclic R-homology manifold, then G is Bredon–Poincaré duality over R.

REMARK 6.2.13. In the case $R = \mathbb{Z}$ we can drop the condition that M be orientable since this is implied by being acyclic. This is because if M is acyclic then $\pi_1(M)$ is perfect, thus $\pi_1(M)$ has no normal subgroups of prime index, in particular M has no index 2 subgroups. But if M were non-orientable then the existence of an orientable double cover (see for example [Hat02, p.234]) would imply that $\pi_1(M)$ has a subgroup of index 2.

Let p be a prime and \mathbb{F}_p the field of p elements. A consequence of Smith theory is the following theorem, for background on Smith theory see [Bre72, §III]

THEOREM 6.2.14 ([**Bor60**, §5 Theorem 2.2][**Dav08**, 10.4.3]). If G is a finite p-group acting properly on an \mathbb{F}_p -homology manifold M then the fixed point set M^G is also an \mathbb{F}_p -homology manifold. If $p \neq 2$ then M^G has even codimension in M.

COROLLARY 6.2.15 (Actions on homology manifolds).

- (1) Let G have an n-dimensional F_p-homology manifold model M for E_{fin}G. If H is a finite p-subgroup of G then M^H is an F_p-homology manifold. In particular if all finite subgroups of G are p-groups and M is cocompact then G is Bredon-Poincaré duality over F_p. If p ≠ 2 and H is a finite p-subgroup of G then n - d_H is even.
- (2) Let G have an n-dimensional Z-homology manifold model M for E_{fin}G.
 If p ≠ 2 is a prime and H is a finite p-subgroup of G such that M^H is a Z-homology manifold then n − dim M^H is even.

REMARK 6.2.16. Given a group G with subgroup H which is not of prime power order, looking at the Sylow *p*-subgroups can give further restrictions. For example if P_i for $i \in I$ is a set of Sylow *p*-subgroups of H, one for each prime p, then G is generated by the P_i [**Rot95**, Ex. 4.10]. Thus if G acts on an Rhomology manifold then the fixed points of H are exactly the intersection of the fixed points of the P_i .

6.2.4. One relator groups. The following lemma is adapted from [BE73, 5.2].

LEMMA 6.2.17. If G is FP₂ with $\operatorname{cd} G = 2$ and $H^1(G, \mathbb{Z}G) = 0$ then G is a duality group.

PROOF. We must show that $H^2(G, \mathbb{Z}G)$ is a flat \mathbb{Z} -module. Consider the short exact sequence of $\mathbb{Z}G$ modules

$$0 \longrightarrow \mathbb{Z}G \xrightarrow{\times p} \mathbb{Z}G \longrightarrow \mathbb{F}_pG \longrightarrow 0.$$

This yields a long exact sequence

$$\cdots \longrightarrow H^1(G, \mathbb{F}_pG) \longrightarrow H^2(G, \mathbb{Z}G) \xrightarrow{\times p} H^2(G, \mathbb{Z}G) \longrightarrow \cdots$$

By [**Bie81**, Corollary 3.6], $H^1(G, \mathbb{F}_p G) \cong H^1(G, \mathbb{Z}G) \otimes_{\mathbb{Z}} \mathbb{F}_p = 0$. Hence the map $H^2(G, \mathbb{Z}G) \xrightarrow{\times p} H^2(G, \mathbb{Z}G)$ must have zero kernel for all p, in other words $H^2(G, \mathbb{Z}G)$ is torsion-free, but the torsion-free \mathbb{Z} -modules are exactly the flat \mathbb{Z} -modules.

Let G be a one-relator group—a group admitting a presentation of finitely many generators and one relator (see [LS01, \S 5] for background), then:

- (1) G is $\mathcal{O}_{\mathcal{F}in}$ FP and $\mathcal{O}_{\mathcal{F}in}$ cd_{\mathbb{Z}} G = 2 [Lüc05, 4.12].
- (2) G contains a torsion-free subgroup Q of finite index [FKS72].
- (3) The normaliser of every non-trivial finite subgroup F is finite. In fact, every such F is subconjugate to a finite cyclic self-normalising subgroup

C of G, and furthermore the normaliser $N_G F$ is subconjugate to C [LS01, II.5.17, II.5.19]. Thus,

$$H^{i}(N_{G}F, \mathbb{Z}[N_{G}F]) = \begin{cases} 0 & \text{if } i > 0, \\ \mathbb{Z} & \text{if } i = 0. \end{cases}$$

Assume further that $H^1(G, \mathbb{Z}G) = 0$.

If $\operatorname{cd}_{\mathbb{Z}} Q \leq 1$ then Q is either trivial or a finitely generated free group so G is either finite or virtually finitely generated free. Thus G is Bredon duality over \mathbb{Z} by 6.4.1, 6.4.3, and 6.4.2. Assume therefore that $\operatorname{cd}_{\mathbb{Z}} Q = 2$. Being finite index in G, Q is also FP₂ and $H^1(Q, \mathbb{Z}Q) = H^1(G, \mathbb{Z}G) = 0$ [**Bro94**, III.(6.5)], thus by Lemma 6.2.17 Q is a duality group and G is virtual duality. Combining with (iii) above, G is also Bredon duality of dimension 2.

In summary:

PROPOSITION 6.2.18. If G is a one relator group with $H^1(G, \mathbb{Z}G) = 0$ then G is Bredon duality over any ring R.

REMARK 6.2.19. If G is a one relator group with $H^1(G, \mathbb{Z}G) \neq 0$ then, since G is $\mathcal{O}_{\mathcal{F}in} \operatorname{FP}_0$, G has bounded orders of finite subgroups by Proposition 3.6.1. By a result of Linnell, G admits a decomposition as the fundamental group of a finite graph of groups with finite edge groups and vertex groups G_v satisfying $H^1(G_v, \mathbb{Z}G_v) = 0$ [Lin83]. These vertex groups are subgroups of virtually torsion-free groups so in particular virtually torsion-free with $\mathcal{O}_{\mathcal{F}in}\operatorname{cd}_{\mathbb{Z}} G_v \leq 2$. Lemma 6.2.20 below gives that the vertex groups are FP₂ and Lemma 6.2.17 shows that the edge groups are virtually duality.

LEMMA 6.2.20. Let G be a group which splits as a finite graph of groups with finite edge groups G_e , indexed by E, and vertex groups G_v , indexed by V. Then if G is FP₂, so are the vertex groups G_v .

PROOF. Fix a vertex group G_v . Let M_λ , for $\lambda \in \Lambda$, be a directed system of $\mathbb{Z}G_v$ modules with $\varinjlim M_\lambda = 0$. To use the Bieri–Eckmann criterion [**Bie81**, Theorem 1.3], we must show that $\varinjlim H^i(G_v, M_\lambda) = 0$ for i = 1, 2.

The Mayer–Vietoris sequence associated to the graph of groups is

$$\cdots \longrightarrow H^i(G, -) \longrightarrow \bigoplus_{u \in V} H^i(G_u, -) \longrightarrow \bigoplus_{e \in E} H^i(G_e, -) \longrightarrow \cdots$$

Now $\varinjlim M_{\lambda} = 0$, so $\varinjlim \operatorname{Ind}_{\mathbb{Z}G_v}^{\mathbb{Z}G} M_{\lambda} = 0$ as well. Evaluating the Mayer–Vietoris sequence at $\operatorname{Ind}_{\mathbb{Z}G_v}^{\mathbb{Z}G} M_{\lambda}$, taking the limit, and using the Bieri–Eckmann criterion, implies

$$\lim_{\Lambda} \bigoplus_{u \in V} H^i(G_u, \operatorname{Ind}_{\mathbb{Z}G_v}^{\mathbb{Z}G} M_{\lambda}) = 0.$$

In particular $\varinjlim_{\mathcal{Z}G_v} H^i(G_v, \operatorname{Ind}_{\mathbb{Z}G_v}^{\mathbb{Z}G} M_\lambda) = 0$ and because, as $\mathbb{Z}G_v$ -module, M_λ is a direct summand of $\operatorname{Ind}_{\mathbb{Z}G_v}^{\mathbb{Z}G} M_\lambda$ [**Bro94**, VII.5.1], this implies $\varinjlim_{\mathcal{H}^i} H^i(G_v, M_\lambda) = 0$.

6.2.5. Discrete subgroups of Lie groups. If L is a Lie group with finitely many path components, K a maximal compact subgroup and G a discrete subgroup then L/K is a model for $E_{\mathcal{F}in}G$. The space L/K is a manifold and the action of G on L/K is smooth so the fixed point subsets of finite groups are submanifolds of L/K, using Lemma 6.2.1. If we assume that the action is co-compact then G is seen to be of type $\mathcal{O}_{\mathcal{F}in}$ FP, $\mathcal{O}_{\mathcal{F}in}$ cd $G = \dim L/K$ and G is a Bredon–Poincaré duality group. See [Lüc05, Theorem 5.24] for a statement of these results.

EXAMPLE 6.2.21. In [**Rag84**, **Rag95**], examples of cocompact lattices in finite covers of the Lie group Spin(2, n) are given which are not virtually torsion-free.

6.2.6. Virtually soluble groups. For G a soluble group the *Hirsch length* hG is the sum of the torsion-free ranks of the factors in an abelian series [**Rob96**, p.422]. Hillman later extended this definition to elementary amenable groups [**Hil91**].

Much of the observation below appears in [MP13a, Example 5.6].

Torsion-free soluble groups of type FP_{∞} are duality [**Kro86**]. We combine this with [**MPN10**], that virtually soluble groups of type FP_{∞} are $\mathcal{O}_{\mathcal{f}in}FP$ with $\mathcal{O}_{\mathcal{f}in}cd G = hG$, and deduce that if G is a virtually soluble duality group (equivalently virtually soluble of type FP_{∞}) then G virtually duality of type $\mathcal{O}_{\mathcal{f}in}FP$ with $\mathcal{O}_{\mathcal{f}in}cd G = hG$. We claim G is also Bredon duality, so we must check the cohomology condition on the Weyl groups. Since G is $\mathcal{O}_{\mathcal{f}in}FP$, the normalisers N_GF of any finite subgroup F of G are FP_{∞} (Corollary 3.6.4). Subgroups of virtually-soluble groups are virtually-soluble [**Rob96**, 5.1.1], so the normalisers N_GF are virtually-soluble FP_{∞} and hence virtually duality, and so the Weyl groups satisfy the required condition on cohomology.

Additionally, if G is a virtually soluble Poincaré duality group then we claim G is Bredon–Poincaré duality. By [**Bie81**, Theorem 9.23], G is virtually-polycyclic. Subgroups of virtually-polycyclic groups are virtually-polycyclic [**Rob96**, p.52], so N_GF is polycyclic FP_{∞} for all finite subgroups F and, since polycylic groups are Poincaré duality,

$$H^{d_F}(N_G F, \mathbb{Z}[N_G F]) = \mathbb{Z}.$$

PROPOSITION 6.2.22. The following conditions on a virtually-soluble group G are equivalent:

- (1) G is FP_{∞} .
- (2) G is virtually duality.
- (3) G is virtually torsion-free and $\operatorname{vcd} G = hG < \infty$.
- (4) G is Bredon duality.

Additionally, if G is Bredon duality then G is virtually Poincaré duality if and only if G is virtually-polycyclic if and only if G is Bredon–Poincaré duality.

PROOF. The equivalence of the first three is [Kro86] and [Kro93]. The rest is the discussion above.

6.2.7. Elementary amenable groups. If G is an elementary amenable group of type FP_{∞} then G is virtually soluble [**KMPN09**, p.4], in particular Bredon duality over \mathbb{Z} of dimension hG. The converse, that every elementary amenable Bredon duality group is FP_{∞} , is obvious.

If G is elementary amenable FP_{∞} then the condition $H^n(G, \mathbb{Z}G) \cong \mathbb{Z}$ implies that G is Bredon–Poincaré duality, so for all finite subgroups,

$$H^{d_F}(N_G F, \mathbb{Z}[N_G F]) \cong \mathbb{Z}.$$

A natural question is whether

$$H^{d_F}(N_G F, \mathbb{Z}[N_G F]) = \mathbb{Z}$$

can ever occur for an elementary amenable, or indeed a soluble Bredon duality, but not Bredon–Poincaré duality group. An example of this behaviour is given below.

EXAMPLE 6.2.23. We construct a finite index extension of the Baumslag–Solitar group BS(1, p), for p a prime.

$$BS(1,p) = \langle x, y : y^{-1}xy = x^p \rangle$$

This has a normal series [LR04, p.60],

$$1 \trianglelefteq \langle x \rangle \trianglelefteq \langle \langle x \rangle \rangle \trianglelefteq BS(1,p),$$

where $\langle \langle x \rangle \rangle$ denotes the normal closure of x. The quotients of this normal series are $\langle x \rangle /1 \cong \mathbb{Z}, \langle \langle x \rangle \rangle / \langle x \rangle \cong C_{p^{\infty}}$ and $BS(1,p)/\langle \langle x \rangle \rangle \cong \mathbb{Z}$, where $C_{p^{\infty}}$ denotes the Prüfer group (see [**Rob96**, p.94]). Clearly BS(1,p) is finitely generated torsionfree soluble with hBS(1,p) = 2, but not polycyclic, since $C_{p^{\infty}}$ does not have the maximal condition on subgroups [**Rob96**, 5.4.12], thus BS(1,p) is not Poincaré duality. Also since BS(1,p) is an HNN extension of $\langle x \rangle \cong \mathbb{Z}$ it has cohomological dimension 2 [**Bie81**, Proposition 6.12] and thus cd BS(1,p) = hBS(1,p). By Proposition 6.2.22, BS(1,p) is a Bredon duality group.

Recall that elements of BS(1,p) can be put in a normal form: $y^i x^k y^{-j}$ where $i, j \ge 0$ and if i, j > 0 then $n \nmid k$. Consider the automorphism φ of BS(1,p),

sending $x \mapsto x^{-1}$ and $y \mapsto y$, an automorphism since it is its own inverse and because the relation $y^{-1}xy = x^p$ in BS(1,p) implies the relation $y^{-1}x^{-1}y = x^{-p}$. Let $y^i x^k, y^{-j}$ be an element in normal form, then

$$\varphi: y^i x^k y^{-j} \longmapsto y^i x^{-k} y^{-j}.$$

So the only fixed points of φ are in the subgroup $\langle y \rangle \cong \mathbb{Z}$. Form the extension

$$1 \longrightarrow BS(1,p) \longrightarrow G \longrightarrow C_2 \longrightarrow 1$$

where C_2 acts by the automorphism φ . The property of being soluble is extension closed [**Rob96**, 5.1.1], so *G* is soluble virtual duality and Bredon duality by Proposition 6.2.22. The normaliser

$$N_G C_2 = C_G C_2 = \{g \in G : gz = zg \text{ for the generator } z \in C_2\}$$

is exactly $\langle y \rangle \times C_2 \cong \mathbb{Z} \times C_2$. Thus $W_G C_2 \cong \mathbb{Z}$ and $H^1(W_G C_2, \mathbb{Z}[W_G C_2]) \cong \mathbb{Z}$. Since BS(1, p) is finite index in G, by [**Bro94**, III.(6.5)]

$$H^2(G, \mathbb{Z}G) \cong H^2(BS(1, p), \mathbb{Z}[BS(1, p)]).$$

However since BS(1,p) is not Poincaré duality, $H^n(BS(1,p),\mathbb{Z}[BS(1,p)])$ is \mathbb{Z} -flat but not isomorphic to \mathbb{Z} .

REMARK 6.2.24. Baues [**Bau04**] and Dekimpe [**Dek03**] proved independently that any virtually polycyclic group G can be realised as a NIL affine crystallographic group—G acts properly, cocompactly, and by diffeomorphisms on a simply connected nilpotent Lie group of dimension hG. Any connected, simply connected nilpotent Lie group is diffeomorphic to some Euclidean space [**Kna02**, §I.16] and hence contractible, so any elementary amenable Bredon–Poincaré duality group has a cocompact manifold model for $E_{\text{fin}}G$.

6.3. Finite extensions of right-angled Coxeter groups

Recall Corollary 6.2.15, that if G has a cocompact n-dimensional Z-homology manifold model M for $\mathbb{E}_{\mathcal{F}in}G$ such that all fixed point sets M^H are Z-homology manifolds, and if H is a finite p-subgroup of G with $p \neq 2$ then $n - d_H$ is even. In this section we construct Bredon–Poincaré duality groups G over Z of arbitrary dimension such that, for any fixed prime $p \neq 2$:

- (1) All of the finite subgroups of G are p-groups.
- (2) $\mathcal{V}(G)$ is any set with $n d_H$ even for all finite subgroups H.

The method of constructing these examples was recommended to us by Ian Leary and utilises methods from [**DL03**, §2] and [**Dav08**, §11]. We write " Γ " instead of "W", as is used in [**DL03**], to denote a Coxeter group so the notation can't be confused with our use of W_GH for the Weyl group.

Let M be an compact contractible *n*-manifold with boundary ∂M , such that ∂M is triangulated as a flag complex. Let G be a group acting on M such that the induced action on the boundary is by simplicial automorphisms. Let Γ be the right angled Coxeter group corresponding to the flag complex ∂M , the group G acts by automorphisms on Γ and we can form the semi-direct product $\Gamma \rtimes G$. Moreover there is a space $\mathcal{U} = \mathcal{U}(M, \partial M, G)$ such that:

- (1) \mathcal{U} is a contractible *n*-manifold without boundary.
- (2) $\Gamma \rtimes G$ acts properly and cocompactly on \mathcal{U} .
- (3) For any finite subgroup H of G, we have $\mathcal{U}^H = \mathcal{U}(M^H, (\partial M)^H, W_G H)$, in particular dim $\mathcal{U}^H = \dim M^H$.
- (4) $W_{\Gamma \rtimes G}H = \Gamma^H \rtimes W_G H$, where Γ^H is the right-angled Coxeter group associated to the flag complex $(\partial M)^H$.
- (5) If M is the cone on a finite complex then \mathcal{U} has a CAT(0) cubical structure (1-connected cubical complex whose links are simplicial flag complexes) such that the action of $\Gamma \rtimes G$ is by isometries.

Every Coxeter group contains a finite-index torsion-free subgroup [**Dav08**, Corollary D.1.4], let Γ' denote such a subgroup of Γ and assume that Γ' is normal. Then $\Gamma' \rtimes G$ is finite index in $\Gamma \rtimes G$ and so acts properly and cocompactly on \mathcal{U} also.

LEMMA 6.3.1. If M is the cone on a finite complex then \mathcal{U} is a cocompact model for $\mathbb{E}_{\text{fin}}(\Gamma \rtimes G)$ and for $\mathbb{E}_{\text{fin}}(\Gamma' \rtimes G)$. In particular, $\Gamma' \rtimes G$ is of type \mathcal{O}_{fin} FP.

PROOF. A CAT(0) cubical complex has a CAT(0) metric [Wis12, Remark 2.1] and any contractible CAT(0) space on which a group acts properly is a model for the classifying space for proper actions [BH99, Corollary II.2.8] (see also [Lüc05, Theorem 4.6]).

LEMMA 6.3.2. Let M be the cone on the finite complex ∂M . If K is a finite subgroup of $\Gamma' \rtimes G$ then K is subconjugate in $\Gamma \rtimes G$ to G.

PROOF. Since, by Lemma 6.3.1, \mathcal{U} is a model for $\mathbb{E}_{\text{fin}}(\Gamma \rtimes G)$, the finite subgroup K necessarily fixes a vertex v of \mathcal{U} and hence is a subgroup of the stabiliser of v.

Recall from $[Dav08, \S5]$ and Section 3.8 that

$$\mathcal{U} = \Gamma \times M / \sim$$

where the identification is along $\Gamma \times \partial M$ only and the action of $\Gamma \rtimes G$ on \mathcal{U} is given by

$$(\gamma', g) \cdot (\gamma, m) = (\gamma'\gamma, gm).$$

A fundamental domain for the Γ -action is the copy of M inside \mathcal{U} given by $1 \times M$ and as such the stabiliser of any vertex is conjugate via an element of Γ

to the stabiliser of a vertex v' in $1 \times M$. Finally, the only elements from $\Gamma' \rtimes G$ stabilising $v' \in 1 \times M$ are contained in G (Γ' moves M about \mathcal{U} freely, whereas G stabilises M setwise).

THEOREM 6.3.3. Let G be a finite group with real representation $\rho : G \hookrightarrow GL_n\mathbb{R}$ and, for all subgroups H of G, let d_H denote the dimension of the subspace of \mathbb{R}^n fixed by H. Then there exists a Bredon-Poincaré duality group $\Gamma' \rtimes G$ of dimension n such that

$$\mathcal{V}(\Gamma' \rtimes G) = \{ d_H : H \le G \}.$$

PROOF. Restrict ρ to an action on (D^n, S^{n-1}) and choose a *G*-equivariant flag triangulation of S^{n-1} (use, for example, [**III78**]). We obtain a Coxeter group Γ , normal finite-index torsion-free subgroup Γ' , and space \mathcal{U} . Lemma 6.3.1 gives that $\Gamma' \rtimes G$ is of type \mathcal{O}_{fin} FP.

Since $\Gamma' \rtimes G$ has an *n*-dimensional model for $\mathbb{E}_{\mathcal{F}in}\Gamma' \rtimes G$ we have that $\mathrm{gd}_{\mathcal{F}in}\Gamma' \rtimes G \leq n$, by Lemma 6.1.2(1) $\mathrm{cd}_{\mathbb{Q}}\Gamma' \rtimes G = d_1$, and by Theorem 6.2.10 and Corollary 6.2.12 $d_1 = n$. Using the chain of inequalities

$$n = \operatorname{cd}_{\mathbb{Q}} \Gamma' \rtimes G \leq \mathcal{O}_{\operatorname{Fin}} \operatorname{cd} \Gamma' \rtimes G \leq \operatorname{gd}_{\operatorname{Fin}} \Gamma' \rtimes G \leq n,$$

shows that $\mathcal{O}_{\text{fin}} \operatorname{cd} \Gamma' \rtimes G = n$. It remains only to check the condition on the cohomology of the Weyl groups of the finite subgroups.

For any finite subgroup H of G, the Weyl group $W_{\Gamma' \rtimes G}H$ acts properly and cocompactly on $\mathcal{U}(M^H, (\partial M)^H, W_G H)$ which is a contractible d_H -manifold without boundary. By Theorem 6.2.10,

$$H^{n}(W_{\Gamma' \rtimes G}H, \mathbb{Z}[W_{\Gamma' \rtimes G}H]) = \begin{cases} \mathbb{Z} & \text{if } i = d_{H}, \\ 0 & \text{else.} \end{cases}$$

If K is any finite subgroup of $\Gamma' \rtimes G$ then, by Lemma 6.3.2, K is conjugate in $\Gamma \rtimes G$ to some $H \leq G$. In particular the normalisers of H and K in $\Gamma \rtimes G$ are isomorphic. Also, since $\Gamma' \rtimes G$ is finite index in $\Gamma \rtimes G$, the normaliser $N_{\Gamma' \rtimes G}K$ is finite index in $N_{\Gamma \rtimes G}K$, thus:

$$H^{n}(N_{\Gamma' \rtimes G}K, \mathbb{Z}[N_{\Gamma' \rtimes G}K]) \cong H^{n}(N_{\Gamma \rtimes G}K, \mathbb{Z}[N_{\Gamma \rtimes G}K])$$
$$\cong H^{n}(N_{\Gamma \rtimes G}H, \mathbb{Z}[N_{\Gamma \rtimes G}H])$$
$$\cong H^{n}(N_{\Gamma' \rtimes G}H, \mathbb{Z}[N_{\Gamma' \rtimes G}H]).$$

From the short exact sequence

$$1 \longrightarrow K \longrightarrow N_{\Gamma' \rtimes G} K \longrightarrow W_{\Gamma' \rtimes G} K \longrightarrow 1$$

and Lemma 6.5.4,

$$H^{n}(N_{\Gamma' \rtimes G}H, \mathbb{Z}[N_{\Gamma' \rtimes G}H]) \cong H^{n}(W_{\Gamma' \rtimes G}K, \mathbb{Z}[W_{\Gamma' \rtimes G}K]).$$

Thus,

$$H^{n}(W_{\Gamma' \rtimes G}K, \mathbb{Z}[W_{\Gamma' \rtimes G}K]) = \begin{cases} \mathbb{Z} & \text{if } i = d_{H}, \\ 0 & \text{else.} \end{cases}$$

EXAMPLE 6.3.4. We construct a group using Theorem 6.3.3 with the properties mentioned at the beginning of this section. It will be of the form $\Gamma' \rtimes C_{p^m}$, where C_{p^m} is the cyclic group of order p^m .

For *i* between 1 and *m* let w_i be any collection of positive integers and let $n = \sum_i 2w_i$. If *c* is a generator of the cyclic group C_{p^m} , then C_{p^m} embeds into the orthogonal group O(n) via the real representation

$$\rho: C_{p^m} \longrightarrow O(n)$$
$$c \longmapsto (R_{2\pi/p})^{\oplus w_1} \oplus (R_{2\pi/p^2})^{\oplus w_2} \oplus \dots \oplus (R_{2\pi/p^m})^{\oplus w_m}$$

where R_{θ} is the 2-dimensional rotation matrix of angle θ . The image is in O(n) since we chose n such that $2w_1 + \cdots + 2w_n = n$.

If *i* is some integer between 1 and *m* then there is a unique subgroup $C_{p^{m-i+1}}$ of C_{p^m} with generator c^{p^i} , in fact this enumerates all subgroups of C_{p^m} except the trivial subgroup. Under ρ , this generator maps to

$$\rho: c^{p^i} \longmapsto R_0^{\oplus w_1} \oplus \cdots \oplus R_0^{\oplus w_i} \oplus \left(R_{p^i 2\pi/p^{i+1}} \right)^{\oplus w_{i+1}} \oplus \cdots \oplus \left(R_{p^i 2\pi/p^m} \right)^{\oplus w_n}$$

In other words, the fixed point set corresponding to $C_{p^{m-i+1}}$ is $\mathbb{R}^{2w_1+\cdots+2w_i}$. Thus the set of dimensions of the fixed point subspaces of non-trivial finite subgroups of C_{p^m} are

 $\{2w_1, 2(w_1+w_2), \ldots, 2(w_1+w_2+\ldots+w_{m-1})\}.$

Applying Theorem 6.3.3 gives a group $\Gamma' \rtimes C_{p^m}$ of type $\mathcal{O}_{\text{fin}} \text{FP}$ with

$$\mathcal{O}_{\mathcal{F}in} \operatorname{cd} G = n = \sum_{i=1}^{m} 2w_i$$

and such that

$$\mathcal{V}(\Gamma' \rtimes C_{p^m}) = \{2w_1, 2(w_1 + w_2), \dots, 2(w_1 + w_2 + \dots + w_{m-1})\}$$

Since there were no restrictions on the integers w_i , using this technique we can build an even dimensional Bredon–Poincaré duality group with any $\mathcal{V}(G)$, as long as all the integers d_H are even.

The case *n* is odd reduces to the case *n* is even. Proposition 6.5.3 shows that if a group *G* is Bredon–Poincaré duality then taking the direct product with \mathbb{Z} gives a Bredon–Poincaré duality group $G \times \mathbb{Z}$ where

$$\mathcal{V}(G \times \mathbb{Z}) \cong \{ v + 1 : v \in \mathcal{V}(G) \}.$$

Thus we can build a group with odd n_H and \mathcal{V} containing only odd elements by building a group with even n_H and then taking a direct product with \mathbb{Z} .

6.4. Low dimensions

This section is devoted to the study of Bredon duality groups and Bredon– Poincaré duality groups of low dimension. We completely classify those of dimension 0 in Lemma 6.4.1. We partially classify those of dimension 1—see Propositions 6.4.2 and 6.4.5, and Question 6.4.4. There is a discussion of the dimension 2 case.

Recall that a group G is duality of dimension 0 over R if and only if |G| is finite and invertible in R, and any such group is necessarily Poincaré duality [**Bie81**, Proposition 9.17(a)].

LEMMA 6.4.1. G is Bredon duality of dimension 0 over R if and only if |G| is finite. Any such group is necessarily Bredon–Poincaré duality.

PROOF. By [Geo08, 13.2.11],

$$H^{0}(G, RG) = \begin{cases} R & \text{if } |G| \text{ is finite,} \\ 0 & \text{else.} \end{cases}$$

Hence if G is Bredon duality of dimension 0 then G is finite and moreover G is Bredon–Poincaré duality.

Conversely, if G is finite then $\mathcal{O}_{\text{fin}} \text{cd}_R G = 0$ and G is $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$ over R (Propositions 3.5.2 and 3.6.1). Finally the Weyl groups of any finite subgroup will be finite so by [**Geo08**, 13.2.11, 13.3.1],

$$H^{n}(WH, R[WH]) = \begin{cases} R & \text{if } n = 0, \\ 0 & \text{if } n > 0. \end{cases}$$

Thus G is Bredon–Poincaré duality of dimension 0.

The duality groups of dimension 1 over R are exactly the groups of type FP₁ over R (equivalently finitely generated groups [**Bie81**, Proposition 2.1]) with $cd_R G = 1$ [**Bie81**, Proposition 9.17(b)].

PROPOSITION 6.4.2. If G is infinite R-torsion free, then the following are equivalent:

- (1) G is Bredon duality over R, of dimension 1.
- (2) G is finitely generated and virtually-free.
- (3) G is virtually duality over R, of dimension 1.

PROOF. That $2 \Rightarrow 3$ is [**Bie81**, Proposition 9.17(b)]. For $3 \Rightarrow 2$, let G be virtually duality over R of dimension 1, then $\operatorname{cd}_R G \leq 1$ so by [**Dun79**] G acts

properly on a tree. Since G is assumed finitely generated, G is virtually-free [Ant11, Theorem 3.3].

For $1 \Rightarrow 2$, if G is Bredon duality over R of dimension 1, then G is automatically finitely generated and $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G = 1$. By Lemma 3.7.2 $\operatorname{cd}_R G = 1$ so, as above, by [**Dun79**] and [**Ant11**, Theorem 3.3], G is virtually-free.

For $2 \Rightarrow 1$, if G is virtually finitely generated free then G acts properly and cocompactly on a tree [Ant11, Theorem 3.3], so G is $\mathcal{O}_{\mathcal{F}in}$ FP over R with $\mathcal{O}_{\mathcal{F}in} cd_R G = 1$. As G is $\mathcal{O}_{\mathcal{F}in}$ FP, for any finite subgroup K, the normaliser $N_G K$ is finitely generated. Subgroups of virtually-free groups are virtually-free, so $N_G K$ is virtually finitely generated free, in particular a virtual duality group [**Bie81**, Proposition 9.17(b)], so

$$H^{i}(WK, \mathbb{Z}[WK]) = H^{i}(N_{G}K, \mathbb{Z}[N_{G}K]) = \begin{cases} \mathbb{Z}\text{-flat} & \text{for } i = d_{K} \\ 0 & \text{else,} \end{cases}$$

where $d_K = 0$ or 1. Thus G is Bredon duality over \mathbb{Z} and hence also over R. \Box

REMARK 6.4.3. The only place that the condition G be R-torsion-free was used was in the implication $1 \Rightarrow 2$, the problem for groups which are not Rtorsion-free is that the condition $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G \leq 1$ is not known to imply that Gacts properly on a tree.

If we take $R = \mathbb{Z}$ then $\mathcal{O}_{\text{fin}} \operatorname{cd}_{\mathbb{Z}} G \leq 1$ implies G acts properly on a tree by a result of Dunwoody [**Dun79**]. Thus over \mathbb{Z} , G is Bredon duality of dimension 1 if and only G is finitely generated virtually free, if and only if G is virtually duality of dimension 1.

QUESTION 6.4.4. What characterises Bredon duality groups of dimension 1 over R?

PROPOSITION 6.4.5. If G is infinite then the following are equivalent:

- (1) G is Bredon-Poincaré duality over R, of dimension 1.
- (2) G is virtually infinite cyclic.
- (3) G is virtually Poincaré duality over R, of dimension 1.

PROOF. The equivalence follows from the fact that for G a finitely generated group, G is virtually infinite cyclic if and only if $H^1(G, RG) \cong R$ [Geo08, 13.5.5,13.5.9].

In dimension 2 we can only classify Bredon–Poincaré duality groups over \mathbb{Z} . The following result appears in [**MP13a**, Example 5.7], but a proof is not given there. Recall that a surface group is the fundamental group of a compact surface without boundary.

LEMMA 6.4.6. If G is virtually a surface group then G is Bredon–Poincaré duality.

PROOF. As G is a virtual surface group, G has finite index subgroup H with H the fundamental group of some compact surface without boundary. Firstly, assume $H = \pi_1(S_g)$ where S_g is the orientable surface of genus g. If g = 0 then S_g is the 2-sphere and G is a finite group, thus G is Bredon–Poincaré duality by Lemma 6.4.1. If g > 0 then by [Mis10, Lemma 4.4(b)] G is \mathcal{O}_{fin} FP over \mathbb{Z} with \mathcal{O}_{fin} cd_{\mathbb{Z}} $G \leq 2$.

We now treat the cases g = 1 and g > 1 separately. If g > 1 then, in the same lemma, Mislin shows that the upper half-plane is a model for $E_{\mathcal{F}in}G$ with Gacting by hyperbolic isometries. Thus [**Dav08**, §10.1] gives that the fixed point sets are all submanifolds, hence G is Bredon–Poincaré duality of dimension 2. If g = 1 then by [**Mis10**, Lemma 4.3], G acts by affine maps on \mathbb{R}^2 so again \mathbb{R}^2 is a model for $E_{\mathcal{F}in}G$ whose fixed point sets are submanifolds, and thus G is Bredon–Poincaré duality of dimension 2.

Now we treat the non-orientable case, so $H = \pi_1(T_k)$ where T_k is a closed nonorientable surface of genus k. In particular T_k has Euler characteristic $\chi(T_k) =$ 2 - k. H has an index 2 subgroup H' isomorphic to the fundamental group of the closed orientable surface of Euler characteristic $2\chi(T_k)$, thus $H' = \pi_1(S_{k-1})$. If k = 1 then $H = \mathbb{Z}/2$ and G is a finite group, thus Bredon–Poincaré duality by Lemma 6.4.1. Assume then that k > 1, we are now back in the situation above where G is virtually $\pi_1(S_g)$ for g > 0 and as such G is Bredon–Poincaré duality of dimension n, by the previous part of the proof.

PROPOSITION 6.4.7. The following conditions are equivalent:

- (1) G is virtually Poincaré duality of dimension 2 over \mathbb{Z} .
- (2) G is virtually surface.
- (3) G is Bredon–Poincaré duality of dimension 2 over \mathbb{Z} .

PROOF. That $1 \Leftrightarrow 2$ is [Eck87] and that $2 \Rightarrow 3$ is Lemma 6.4.6. The implication $3 \Rightarrow 1$ is provided by [Bow04, Theorem 0.1] which states that any FP₂ group with $H^2(G, \mathbb{Q}G) = \mathbb{Q}$ is a virtual surface group and hence a virtual Poincaré duality group. If G is Bredon–Poincaré duality of dimension 2 then $H^i(G, \mathbb{Q}G) = H^i(G, \mathbb{Z}G) \otimes \mathbb{Q} = \mathbb{Q}$ (see proof of Lemma 6.1.2(1)) and G is FP₂ so we may apply the aforementioned theorem.

The above proposition doesn't extend from Poincaré duality to just duality, as demonstrated by [Sch78, p.163] where an example, based on Higman's group, is given of a Bredon duality group of dimension 2 over \mathbb{Z} which is not virtual duality. This example is extension of a finite group by a torsion-free duality group of dimension 2. Schneebeli proves that the group is not virtually torsion-free, that it is Bredon duality follows from Proposition 6.5.8.

6.5. EXTENSIONS

QUESTION 6.4.8. Is there an easy characterisation of Bredon duality, or Bredon–Poincaré duality groups, of dimension 2 over R?

6.5. Extensions

In the classical case, extensions of duality groups by duality groups are always duality [**Bie81**, 9.10]. In the Bredon case the situation is more complex, for example semi-direct products of torsion-free groups by finite groups may not even be $\mathcal{O}_{\text{fin}} \text{FP}_0$ [**LN03**]. Davis and Leary build examples of finite index extensions of Poincaré duality groups which are not Bredon duality, although they are $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$ [**DL03**, Theorem 2], and examples of virtual duality groups which are not of type $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$ [**DL03**, Theorem 1]. In [**FL04**], Farrell and Lafont give examples of prime index extensions of δ -hyperbolic Poincaré duality groups which are not Bredon–Poincaré duality. In [**MP13a**, §5], Martínez-Pérez considers *p*-power extensions of duality groups over fields of characteristic *p*, showing that if *Q* is a *p*-group and *G* is Poincaré duality of dimension *n* over a field of characteristic *p* then then $G \rtimes Q$ is Bredon–Poincaré duality groups to duality groups however [**MP13a**, §6].

We study direct products of Bredon duality groups and extensions of the form finite-by-Bredon duality.

6.5.1. Direct products.

LEMMA 6.5.1. For all groups G_1 and G_2 ,

- (1) If G_1 and G_2 are \mathcal{O}_{fin} FP over R then $G_1 \times G_2$ is \mathcal{O}_{fin} FP over R.
- (2) $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G_1 \times G_2 \leq \mathcal{O}_{\text{fin}} \operatorname{cd}_R G_1 + \mathcal{O}_{\text{fin}} \operatorname{cd}_R G_2.$

PROOF. That $\mathcal{O}_{\mathcal{F}in} \operatorname{cd}_R G_1 \times G_2 \leq \mathcal{O}_{\mathcal{F}in} \operatorname{cd}_R G_1 + \mathcal{O}_{\mathcal{F}in} \operatorname{cd}_R G_2$ is a special case of [**Flu10**, 3.62], where Fluch proves that given projective resolutions P_* of \underline{R} by $\mathcal{O}_{\mathcal{F}in}$ -modules for G_1 and Q_* of \underline{R} by $\mathcal{O}_{\mathcal{F}in}$ -modules for G_2 , the total complex of the tensor product double complex is a projective resolution of \underline{R} by projective $\mathcal{O}_{\mathcal{F}in}$ -modules for $G_1 \times G_2$. So to prove that $G_1 \times G_2$ is $\mathcal{O}_{\mathcal{F}in}$ FP it is sufficient to show that if P_* and Q_* are finite type resolutions, then so is the total complex, but this follows from [**Flu10**, 3.52].

LEMMA 6.5.2. If L is a finite subgroup of $G_1 \times G_2$ then the normaliser $N_{G_1 \times G_2}L$ is finite index in $N_{G_1}\pi_1L \times N_{G_2}\pi_2L$, where π_1 and π_2 are the projection maps from $G_1 \times G_2$ onto the factors G_1 and G_2 .

PROOF. It's straightforward to check that

$$N_{G_1 \times G_2} L \le N_{G_1} \pi_1 L \times N_{G_2} \pi_2 L.$$

Next, observe that $N_{G_1}\pi_1L \times N_{G_2}\pi_2L$ acts by conjugation on $\pi_1L \times \pi_2L$ and the setwise stabiliser of $L \leq (\pi_1L \times \pi_2L)$ is exactly $N_{G_1 \times G_2}L$. Since $\pi_1L \times \pi_2L$ is finite, any stabiliser of a subset is necessarily finite-index (via the orbit-stabiliser theorem), thus $N_{G_1 \times G_2}L$ is finite index in $N_{G_1}\pi_1L \times N_{G_2}\pi_2L$.

PROPOSITION 6.5.3. If G_1 and G_2 are Bredon duality (resp. Bredon-Poincaré duality), then $G = G_1 \times G_2$ is Bredon duality (resp. Bredon-Poincaré duality). Furthermore,

$$\mathcal{V}(G_1 \times G_2) = \{ v_1 + v_2 : v_i \in \mathcal{V}(G_i) \} \cup \{ v_1 + d_1(G_2) : v_1 \in \mathcal{V}(G_1) \}$$
$$\cup \{ d_1(G_1) + v_2 : v_2 \in \mathcal{V}(G_2) \}.$$

PROOF. By Lemma 6.5.1, $G_1 \times G_2$ is \mathcal{O}_{fin} FP. If L is some finite subgroup of G, then, via Lemma 6.5.2, the normaliser $N_G L$ is finite index in $N_{G_1} \pi_1 L \times N_{G_2} \pi_2 L$ so an application of Shapiro's Lemma [**Bro94**, III.(6.5) p.73] gives that for all i,

 $H^{i}(N_{G}L, R[N_{G}L]) \cong H^{i}(N_{G_{1}}\pi_{1}L \times N_{G_{2}}\pi_{2}L, R[N_{G_{1}}\pi_{1}L \times N_{G_{2}}\pi_{2}L]).$

Noting the isomorphism of RG-modules

$$R[N_{G_1}\pi_1L \times N_{G_2}\pi_2L] \cong R[N_{G_1}\pi_1L] \otimes R[N_{G_2}\pi_2L],$$

the Künneth formula for group cohomology (see [Bro94, p.109]) is:

$$\begin{array}{c}
0 \\
\downarrow \\
\bigoplus_{i+j=k} \left(H^{i}(N_{G_{1}}\pi_{1}L, R[N_{G_{1}}\pi_{1}L]) \otimes H^{j}(N_{G_{1}}\pi_{1}L, R[N_{G_{1}}\pi_{1}L]) \right) \\
\downarrow \\
H^{k}(N_{G_{1}}\pi_{1}L \times N_{G_{2}}\pi_{2}L, R[N_{G_{1}}\pi_{1}L \times N_{G_{2}}\pi_{2}L]) \\
\downarrow \\
\bigoplus_{i+j=k+1} \operatorname{Tor}_{1}^{R}(H^{i}(N_{G_{1}}\pi_{1}L, R[N_{G_{1}}\pi_{1}L]), H^{j}(N_{G_{2}}\pi_{2}L, R[N_{G_{2}}\pi_{2}L])) \\
\downarrow \\
0
\end{array}$$

Here we are using that the $R[N_{G_i}\pi_i L]$ are *R*-free. Since $H^i(N_{G_1}\pi_1 L, R[N_{G_1}\pi_1 L])$ is *R*-flat the Tor₁ term is zero. Hence the central term is non-zero only when $i = d_{\pi_1 L}$ and $j = d_{\pi_2 L}$, in which case it is *R*-flat. Furthermore, $d_L = d_{\pi_1 L} + d_{\pi_2 L}$.

If G_1 and G_2 are Bredon–Poincaré duality then the central term in this case is R.

Since if L is non-trivial one of $\pi_1 L$ and $\pi_2 L$ must be non-trivial, the argument above implies that

$$\mathcal{V}(G_1 \times G_2) \subseteq \{v_1 + v_2 : v_i \in \mathcal{V}(G_i)\} \cup \{v_1 + d_1(G_2) : v_1 \in \mathcal{V}(G_1)\}$$
$$\cup \{d_1(G_1) + v_2 : v_2 \in \mathcal{V}(G_2)\}.$$

For the other inclusion let

$$v \in \{v_1 + v_2 : v_i \in \mathcal{V}(G_i)\} \cup \{v_1 + d_1(G_2) : v_1 \in \mathcal{V}(G)\}$$
$$\cup \{d_1(G_1) + v_2 : v_2 \in \mathcal{V}(G)\}.$$

Thus there exist finite finite subgroups L_1 of G_1 and L_2 of G_2 such that $d_{L_1} = v_1$, $d_{L_2} = v_2$, and one of the L_i is non-trivial. Using the Künneth formula again, one calculates that $d_{L_1 \times L_2} = v$.

6.5.2. Finite-by-duality groups. Throughout this section, F, G and Q will denote groups in a short exact sequence

$$1 \longrightarrow F \longrightarrow G \xrightarrow{\pi} Q \longrightarrow 1,$$

where F is finite. This section builds up to the proof of Proposition 6.5.8 that if Q is Bredon duality of dimension n over R, then G is also.

LEMMA 6.5.4. $H^i(G, RG) \cong H^i(Q, RQ)$ for all *i*.

PROOF. The Lyndon–Hochschild–Serre spectral sequence associated to the extension is [**Bro94**, VII§6]

$$H^p(Q, H^q(F, RG)) \Rightarrow H^{p+q}(G, RG).$$

RG is projective as a RF-module so by [Bie81, Proposition 5.3, Lemma 5.7],

$$H^{q}(F, RG) \cong H^{q}(F, RF) \otimes_{RF} RG = \begin{cases} R \otimes_{RF} RG = RQ & \text{if } q = 0, \\ 0 & \text{else.} \end{cases}$$

The spectral sequence collapses to $H^i(G, RG) \cong H^i(Q, RQ)$.

LEMMA 6.5.5. If Q is $\mathcal{O}_{\text{fin}} FP_0$, then G is $\mathcal{O}_{\text{fin}} FP_0$.

PROOF. Let B_i for i = 0, ..., n be a collection of conjugacy class representatives of all finite subgroups in Q. For each i, let B_i^j be a collection of conjugacy class representatives of finite subgroups in G which project onto B_i . Since F is finite $\pi^{-1}(B_i)$ is finite and there are only finitely many j for each i, we claim that these B_i^j are conjugacy class representatives for all finite subgroups in G.

Let K be some finite subgroup of G, we need to check it is conjugate to some B_i^j . $A = \pi(K)$ is conjugate to B_i , let $q \in Q$ be such that $q^{-1}Aq = B_i$ and let $g \in G$ be such that $\pi(g) = q$.

 $\pi(g^{-1}Kg) = q^{-1}Aq = B_i$ so $g^{-1}Kg$ is conjugate to some B_i^j and hence K is conjugate to some B_i^j . Since we have already observed that for each *i*, the set $\{B_i^j\}_j$ is finite, G has finitely many conjugacy classes of finite subgroups. \Box

LEMMA 6.5.6. If K is a finite subgroup of G then N_GK is finite index in $N_G(\pi^{-1} \circ \pi(K))$.

PROOF. $N_G K$ is a subgroup of $N_G(\pi^{-1} \circ \pi(K))$ since if $g^{-1}Kg = K$ then

$$(\pi^{-1} \circ \pi(g)) (\pi^{-1} \circ \pi(K)) (\pi^{-1} \circ \pi(g))^{-1} = \pi^{-1} \circ \pi(K),$$

but $g \in \pi^{-1} \circ \pi(g)$ so $g(\pi^{-1} \circ \pi(K))g^{-1} = \pi^{-1} \circ \pi(K)$.

Consider the action of $N_G(\pi^{-1} \circ \pi(K))$ on $\pi^{-1} \circ \pi(K)$ by conjugation, the setwise stabiliser of K is exactly $N_G K$. Since $\pi^{-1} \circ \pi(K)$ is finite, any stabiliser is finite index via the orbit-stabiliser theorem. We conclude that $N_G K$ is finite index in $N_G(\pi^{-1} \circ \pi(K))$.

LEMMA 6.5.7. If L is a subgroup of Q then $N_G \pi^{-1}(L) = \pi^{-1} N_Q L$.

PROOF. If $g \in N_G \pi^{-1}(L)$ then $g^{-1}\pi^{-1}(L)g = \pi^{-1}(L)$ so applying π gives that $\pi(g)^{-1}L\pi(g) = L$. Thus $\pi(g) \in N_Q L$, equivalently $g \in \pi^{-1}N_Q L$.

Conversely if $g \in \pi^{-1}(N_Q L)$ then $\pi(g)^{-1}L\pi(g) = L$ so

$$\left(\pi^{-1} \circ \pi(g)\right)^{-1} \pi^{-1}(L) \left(\pi^{-1} \circ \pi(g)\right) = \pi^{-1}(L).$$

Since $g \in \pi^{-1} \circ \pi(g)$, we have that $g^{-1}\pi^{-1}(L)g = \pi^{-1}(L)$.

PROPOSITION 6.5.8. Q is Bredon duality of dimension n over R if and only if G is Bredon duality of dimension n over R. Moreover, $\mathcal{V}(G) = \mathcal{V}(Q)$.

PROOF. Assume that Q is Bredon duality of dimension n of R. Let K be a finite subgroup of G. We combine Lemma 6.5.6 and Lemma 6.5.7 to see that $N_G K$ is finite index in $N_G(\pi^{-1} \circ \pi(K)) = \pi^{-1}(N_Q \pi(K))$. Hence

$$H^{i}(W_{G}K, R[W_{G}K]) \cong H^{i}(N_{G}K, R[N_{G}K])$$
$$\cong H^{i}(\pi^{-1}(N_{Q}\pi(K)), R[\pi^{-1}(N_{Q}\pi(K))])$$
$$\cong H^{i}(N_{Q}\pi(K), R[N_{Q}\pi(K)])$$
$$\cong H^{i}(W_{Q}\pi(K), R[W_{Q}\pi(K)])$$

where the first isomorphism is from the short exact sequence

$$1 \longrightarrow K \longrightarrow N_G K \longrightarrow W_G K \longrightarrow 1$$

and Lemma 6.5.4, the fourth isomorphism is from the same lemma and a similar short exact sequence containing $N_Q K$, and the third isomorphism follows from Lemma 6.5.4 and the short exact sequence

$$1 \longrightarrow F \longrightarrow \pi^{-1}(N_Q \pi(K)) \longrightarrow N_Q \pi(K) \longrightarrow 1.$$

Since Q is Bredon duality of dimension n this gives the condition on the cohomology of the Weyl groups.

G is $\mathcal{O}_{\text{fin}} \text{FP}_0$ by Lemma 6.5.5, and $\mathcal{O}_{\text{fin}} \text{cd} G = \mathcal{O}_{\text{fin}} \text{cd} Q = n$ by [Nuc04, Theorem 5.5]. So by Corollary 3.6.4, it remains to show that the Weyl groups of the finite subgroups are FP_{∞} . For any finite subgroup K of G, the short
exact sequence above and [**Bie81**, Proposition 1.4] gives that $\pi^{-1}(N_Q\pi(K))$ is $\operatorname{FP}_{\infty}$. But, as discussed at the beginning of the proof, $N_G K$ is finite index in $N_G(\pi^{-1} \circ \pi(K)) = \pi^{-1}(N_Q\pi(K))$, so $N_G K$ is $\operatorname{FP}_{\infty}$ also.

For the converse, assume that G is Bredon duality of dimension n over R. Let K be a finite subgroup of Q then

$$H^{i}(W_{Q}K, R[W_{Q}K]) \cong H^{i}(N_{Q}K, R[N_{Q}K])$$
$$\cong H^{i}(\pi^{-1}(N_{Q}K), R[\pi^{-1}(N_{Q}K)])$$
$$\cong H^{i}(N_{G}\pi^{-1}K, R[N_{G}\pi^{-1}K])$$
$$\cong H^{i}(W_{G}\pi^{-1}K, R[W_{G}\pi^{-1}K]),$$

where the first isomorphism is from the short exact sequence

$$1 \longrightarrow K \longrightarrow N_Q K \longrightarrow W_Q K \longrightarrow 1$$

and Lemma 6.5.4, the fourth isomorphism is from the same lemma and a similar short exact sequence containing $N_G \pi^{-1} K$, the second isomorphism follows from Lemma 6.5.4 and the short exact sequence

$$1 \longrightarrow F \longrightarrow \pi^{-1}(N_Q K) \longrightarrow N_Q K \longrightarrow 1,$$

and the third isomorphism is from Lemma 6.5.7.

Since G is Bredon duality of dimension n this gives the condition on the cohomology of the Weyl groups. Finally, since G is $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$, thus also Q is $\mathcal{O}_{\text{fin}} \text{FP}_{\infty}$.

6.6. Graphs of groups

An amalgamated free product of two duality groups of dimension n over a duality group of dimension n-1 is duality of dimension n, similarly an HNN extension of a duality group of dimension n relative to a duality group of dimension n-1 is duality of dimension n [Bie81, Proposition 9.15]. Unfortunately we know of no such result for Bredon–Poincaré duality groups: the problem is how to obtain the correct condition on the cohomology of the Weyl groups of the finite subgroups. However by putting some restrictions on the graph of groups, we can obtain some useful examples. For instance using graphs of groups of Bredon duality groups we will be able to build Bredon duality groups G with arbitrary $\mathcal{V}(G)$.

Throughout this section, G is the fundamental group of a finite graph of groups. Let T = (V, E) denote the associated Bass–Serre tree, we denote by G_v the stabiliser of the vertex $v \in V$ and we denote by G_e the stabiliser of the edge $e \in E$. See [Ser03] for the necessary background on Bass–Serre trees and graphs of groups.

We need some preliminary results, showing that a graph of groups is $\mathcal{O}_{\mathcal{F}in}$ FP if all groups involved are $\mathcal{O}_{\mathcal{F}in}$ FP. See [Ser03] for the necessary background on Bass–Serre trees and graphs of groups.

LEMMA 6.6.1. [GN12, Lemma 3.2] There is an exact sequence, arising from the Bass–Serre tree.

$$\cdots \longrightarrow H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G,-) \longrightarrow \bigoplus_{v \in V} H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G_{v}, \operatorname{Res}_{G_{v}}^{G}-)$$
$$\longrightarrow \bigoplus_{e \in E} H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G_{e}, \operatorname{Res}_{G_{e}}^{G}-) \longrightarrow \cdots$$

LEMMA 6.6.2. If all vertex groups G_v are of type $\mathcal{O}_{\text{fin}} \text{FP}_n$ and all edge groups G_e are of type $\mathcal{O}_{\text{fin}} \text{FP}_{n-1}$ over R then G is of type $\mathcal{O}_{\text{fin}} \text{FP}_n$ over R.

PROOF. Let M_{λ} , for $\lambda \in \Lambda$, be a directed system of \mathcal{O}_{fin} -modules with colimit zero. For any subgroup H of G, the directed system $\operatorname{Res}_{H}^{G} M_{\lambda}$ also has colimit zero. The long exact sequence of Lemma 6.6.1, and the exactness of colimits gives that for all i, there is an exact sequence

$$\cdots \longrightarrow \lim_{\lambda \in \Lambda} H^{i-1}_{\mathcal{O}_{\mathcal{F}_{in}}}(G, M_{\lambda}) \longrightarrow \bigoplus_{v \in V} \lim_{\lambda \in \Lambda} H^{i}_{\mathcal{O}_{\mathcal{F}_{in}}}\left(G_{v}, \operatorname{Res}_{G_{v}}^{G} M_{\lambda}\right)$$
$$\longrightarrow \bigoplus_{e \in E} \lim_{\lambda \in \Lambda} H^{i}_{\mathcal{O}_{\mathcal{F}_{in}}}\left(G_{e}, \operatorname{Res}_{G_{e}}^{G} M_{\lambda}\right) \longrightarrow \cdots .$$

If $i \leq n$ then by the Bieri–Eckmann criterion (Theorem 2.5.1), the left and right hand terms vanish, thus the central term vanishes. Another application of the Bieri–Eckmann criterion gives that G is $\mathcal{O}_{\text{fin}} \operatorname{FP}_n$.

LEMMA 6.6.3. If $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G_v \leq n$ for all vertex groups G_v and $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G_e \leq n-1$ for all edge groups G_e then $\mathcal{O}_{\text{fin}} \operatorname{cd}_R G \leq n$.

PROOF. Use the long exact sequence of Lemma 6.6.1.

LEMMA 6.6.4. If there exists a positive integer n such that:

(1) For every $v \in V$, $H^i(G_v, RG_v)$ is R-flat if i = n and 0 otherwise.

(2) For every $e \in E$, $H^i(G_e, RG_e)$ is R-flat if i = n - 1 and 0 otherwise. Then $H^i(G, RG)$ is R-flat if i = n and 0 else.

PROOF. The Mayer–Vietoris sequence associated to the graph of groups is

$$\cdots \longrightarrow H^q(G, RG) \longrightarrow \bigoplus_{v \in V} H^q(G_v, RG) \longrightarrow \bigoplus_{e \in E} H^q(G_e, RG) \longrightarrow \cdots$$

 $H^{q}(G_{v}, RG) = H^{q}(G_{v}, RG_{v}) \otimes_{RG_{v}} RG$ by [**Bie81**, Proposition 5.4] so we have

$$H^q(G, RG) = 0$$
 for $q \neq n$,

and a short exact sequence

$$0 \longrightarrow \bigoplus_{e \in E} H^{n-1}(G_e, RG_e) \otimes_{RG_e} RG \longrightarrow H^n(G, RG)$$
$$\longrightarrow \bigoplus_{v \in V} H^n(G_v, RG_v) \otimes_{RG_v} RG \longrightarrow 0.$$

Finally, extensions of flat modules by flat modules are flat (use, for example, the long exact sequence associated to $\operatorname{Tor}_*^{RG}$).

REMARK 6.6.5. In the above, if $H^n(G, RG_v) \cong R$ and $H^{n-1}(G_e, RG_e) \cong R$ for all vertex and edge groups then $H^n(G, RG)$ will not be isomorphic to R.

LEMMA 6.6.6. If K is a subgroup of the vertex group G_v and K is not subconjugate to any edge group then $N_G K = N_{G_v} K$.

PROOF. Let T be the Bass–Serre tree, then the normaliser $N_G K$ fixes T^K setwise, but T^K is the single vertex v (if $w \neq v$ was also fixed by K then K would fix all edges on the path from v to w, but it is assumed that K is not subconjugate to any edge stabiliser). Thus, $N_G K \leq G_v$.

EXAMPLE 6.6.7. Let S_n denote the star graph of n + 1 vertices—a single central vertex v_0 , and a single edge connecting every other vertex v_i to the central vertex. Let G be the fundamental group of a graph of groups on S_n , where the central vertex group G_0 is torsion-free duality of dimension n, the edge groups are torsion-free duality of dimension n - 1 and the remaining vertex groups G_i are Bredon duality of dimension n with $n_1 = n$.

By Lemmas 6.6.2 and 6.6.3, G is $\mathcal{O}_{\mathcal{F}in}$ FP of dimension n, so to prove it is Bredon duality it suffices to check the cohomology of the Weyl groups of the finite subgroups. Any non-trivial finite subgroup is subconjugate to a unique vertex group G_i , and cannot be subconjugate to an edge group since they are assumed torsion-free. If K is a subgroup of G_i then by Lemma 6.6.6, $H^i(N_GK, R[N_GK]) \cong$ $H^i(N_{G_i}K, R[N_{G_i}K])$ and the condition follows as G_i was assumed to be Bredon duality. Finally, for the trivial subgroup we must calculate $H^i(G, RG)$, which is Lemma 6.6.4.

 $\mathcal{V}(G)$ is easily calculable too,

$$\mathcal{V}(G) = \{ v : v \in \mathcal{V}(G_i) \text{ for some } i \in \{1, \dots, n\} \}.$$

EXAMPLE 6.6.8 (A Bredon duality group with prescribed $\mathcal{V}(G)$). We specialise the above example. Let $\mathcal{V} = \{v_1, \ldots, v_t\} \subset \{0, 1, \ldots, n\}$ be given. Choosing $G_i = \mathbb{Z}^n \rtimes \mathbb{Z}_2$ as in Example 6.2.3 so that $\mathcal{V}(G_i) = v_i$, let $G_0 = \mathbb{Z}^n$, let the edge groups be \mathbb{Z}^{n-1} , and choose injections $\mathbb{Z}^{n-1} \to \mathbb{Z}^n$ and $\mathbb{Z}^{n-1} \to \mathbb{Z}^n \rtimes \mathbb{Z}_2$ from the edge groups into the vertex groups. Then form the graph of groups as in the previous example to get, for G the fundamental group of the graph of groups,

$$\mathcal{V}(G) = \{v_1, \dots, v_t\}.$$

Because of Remark 6.6.5 the groups constructed in the example above will not be Bredon–Poincaré duality groups.

6.7. The wrong notion of Bredon duality

This section grew out of an investigation into which groups were \mathcal{O}_{fin} FP over some ring R with

$$H^{i}_{\mathcal{O}_{\mathcal{F}_{in}}}(G, R[-, ?]_{\mathcal{O}_{\mathcal{F}_{in}}}) \cong \begin{cases} \underline{R}(?) & \text{if } i = n, \\ 0 & \text{else.} \end{cases}$$

One might hope that this naïve definition would give a duality similar to Poincaré duality, we show this is not the case. Namely we prove in Theorem 6.7.3 that the only groups satisfying this property are torsion-free, and hence torsion-free Poincaré duality groups over R. We need a couple of technical results before we can prove the theorem.

Recall from Section 3.9 that for M a contravariant \mathcal{O}_{fin} -module we denote by M^D the dual module

$$M^{D} = \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}} \left(M(-), R[-, ?]_{\mathcal{O}_{\operatorname{fin}}} \right).$$

Note that M^D is a covariant $\mathcal{O}_{\mathcal{F}in}$ -module. Similarly for A a covariant $\mathcal{O}_{\mathcal{F}in}$ -module,

$$A^D = \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}} \left(A(-), R[?, -]_{\mathcal{O}_{\operatorname{fin}}} \right)$$

LEMMA 6.7.1. If there exists a length n resolution of the constant covariant module \underline{R} by projective covariant $\mathcal{O}_{\mathcal{F}in}$ -modules then G is R-torsion free and $\operatorname{cd}_R G \leq n$.

PROOF. Let $P_* \longrightarrow \underline{R}$ be a length n projective covariant resolution of \underline{R} , evaluating at G/1 gives a length n resolution of R by projective RG-modules (Propositions 2.3.2(1) and 3.2.2)(1). Thus $\operatorname{cd}_R G \leq n$ and it follows that G is R-torsion free.

Let M be an RG-module and recall from Section 3.2 that inducing M to a covariant $\mathcal{O}_{\mathcal{F}in}$ -module gives $\operatorname{Ind}_{RG}^{\mathcal{O}_{\mathcal{F}in}} M = M \otimes_{RG} R[G/1, -]_{\mathcal{O}_{\mathcal{F}in}}$. The covariant induction functor maps projective modules to projective modules and satisfies the following adjoint isomorphism for any covariant $\mathcal{O}_{\mathcal{F}in}$ -module A (Propositions 2.3.2 and 2.3.1),

$$\operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}}(\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}}M, A) \cong \operatorname{Hom}_{RG}(M, A(G/1)).$$

LEMMA 6.7.2. If $\operatorname{cd}_R G \leq n$ then there exists a length n projective covariant resolution of R.

PROOF. Let P_* be a length n projective RG-module resolution of R, then we claim $\operatorname{Ind}_{RG}^{\mathcal{O}_{\overline{fin}}} P_*$ is a projective covariant resolution of \underline{R} . One can easily check that $\operatorname{Ind}_{RG}^{\mathcal{O}_{\overline{fin}}} R = \underline{R}$ (Example 3.2.1) and since G is necessarily R-torsion-free, $\operatorname{Ind}_{RG}^{\mathcal{O}_{\overline{fin}}} P_*$ is exact (Proposition 3.2.5).

THEOREM 6.7.3. If G is $\mathcal{O}_{\text{fin}} \text{FP}$ with $\mathcal{O}_{\text{fin}} \text{cd}_R G = n$ and

$$H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G, R[-, ?]_{\mathcal{O}_{\mathcal{F}in}}) \cong \begin{cases} \underline{R}(?) & \text{if } i = n, \\ 0 & \text{else,} \end{cases}$$

then G is torsion-free. Note that in the above, <u>R</u> denotes the constant covariant \mathcal{O}_{Fin} -module.

PROOF. Choose a length n finite type projective $\mathcal{O}_{\mathcal{F}in}$ -module resolution P_* of <u>R</u> then by the assumption on $H^n_{\mathcal{O}_{\mathcal{F}in}}(G, R[-,?]_{\mathcal{O}_{\mathcal{F}in}})$, we know that P^D_* is a covariant resolution by finitely generated projectives of <u>R</u>:

$$0 \to P_0^D(-) \stackrel{\partial_1^D}{\to} P_1^D(-) \stackrel{\partial_2^D}{\to} \cdots \stackrel{\partial_n^D}{\to} P_n^D(-) \to H^n_{\mathcal{O}_{\text{fin}}}(G, R[?, -]_{\mathcal{O}_{\text{fin}}}) \cong \underline{R}(-) \to 0.$$

By Lemma 6.7.1 G is R-torsion-free and $\operatorname{cd}_R G \leq n$. Since G is $\mathcal{O}_{\operatorname{fin}}\operatorname{FP}_{\infty}$, G is $\operatorname{FP}_{\infty}$ (Corollary 3.6.4) and we may choose a length n finite type projective RG-module resolution Q_* of R. Lemma 6.7.2 gives that $\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_* \longrightarrow \underline{R}$ is a projective covariant resolution.

By the $\mathcal{O}_{\mathcal{F}in}$ -module analogue of the comparison theorem [Wei94, 2.2.6], the two projective covariant resolutions of <u>R</u> are chain homotopy equivalent. Any additive functor preserves chain homotopy equivalences, so applying the dual functor to both complexes gives a chain homotopy equivalence between

$$0 \longrightarrow \underline{R}^{D} \cong 0 \longrightarrow (\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_{0})^{D} \longrightarrow \cdots \longrightarrow (\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_{n})^{D}$$

and

$$0 \longrightarrow \underline{R}^D \cong 0 \longrightarrow P_n^{DD} \longrightarrow P_{n-1}^{DD} \longrightarrow \cdots \longrightarrow P_0^{DD},$$

(that $\underline{R}^D \cong 0$ is Example 3.9.1). Since $\operatorname{Hom}_{\mathcal{O}_{fin}}$ is left exact we know both complexes above are left exact. Lemma 3.9.2 gives the commutative diagram below.

The lower complex, P_* , satisfies $H_0P_* \cong \underline{R}$ and $H_iP_* = 0$ for all $i \neq 0$. Thus the same is true for the top complex, and also the complex $\operatorname{Ind}_{RG}^{\mathcal{O}_{\text{fin}}} Q_*^D$, since this is homotopy equivalent to it. In particular, there is an epimorphism of \mathcal{O}_{fin} -modules,

$$\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_n^D \longrightarrow \underline{R}.$$

The left hand side simplifies, using the adjoint isomorphism

$$\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_n^D = \operatorname{Hom}_{\mathcal{O}_{\operatorname{fin}}} \left(\operatorname{Ind}_{RG}^{\mathcal{O}_{\operatorname{fin}}} Q_n, R[?, -]_{\mathcal{O}_{\operatorname{fin}}} \right) \cong \operatorname{Hom}_{RG}(Q_n, R[?, G/1]_{\mathcal{O}_{\operatorname{fin}}}).$$

Since $\operatorname{Hom}_{RG}(Q_n, R[?, G/1]_{\mathcal{O}_{\operatorname{fin}}})(G/H) = 0$ if $H \neq 1$, this module cannot surject onto <u>R</u> unless G is torsion-free.

CHAPTER 7

Houghton's groups

This chapter, with the exception of Sections 7.4.1 and 7.4.2, contains material that has appeared in:

• Centralisers in Houghton's Groups (2012, to appear Proc. Edinburgh Math. Soc.) [SJG12].

Sections 7.4.1 and 7.4.2 contain unpublished joint work with Nansen Petrosyan.

Section 7.1 contains an analysis of the centralisers of finite subgroups in Houghton's group. As Corollary 7.1.7 we obtain that centralisers of finite subgroups are FP_{n-1} but not FP_n . In Section 7.2 our analysis is extended to arbitrary elements and virtually cyclic subgroups. Using this information elements in H_n are constructed whose centralisers are FP_i for any $0 \le i \le n-3$.

In Section 7.3 the space that Brown constructed in [**Bro87**], in order to prove that H_n is FP_{n-1} but not FP_n, is shown to be a model for $E_{\mathcal{F}in}H_n$, the classifying space for proper actions of H_n . Finally Section 7.4 contains a discussion of Bredon (co)homological finiteness conditions satisfied by Houghton's group, namely we show in Proposition 7.4.1 that H_n is not quasi- $\mathcal{O}_{\mathcal{F}in}$ FP₀ and in Proposition 7.4.3 that the Bredon cohomological dimension with respect to the family of finite subgroups and virtually cyclic subgroups are both equal to n. See Section 3.6.1 for the definition of quasi- $\mathcal{O}_{\mathcal{F}in}$ FP_n.

Fixing a natural number n > 1, define Houghton's group H_n to be the group of permutations of $S = \mathbb{N} \times \{1, \ldots, n\}$ which are "eventually translations", ie. for any given permutation $h \in H_n$ there are collections $\{z_1, \ldots, z_n\} \in \mathbb{N}^n$ and $\{m_1, \ldots, m_n\} \in \mathbb{Z}^n$ with

(7.1)
$$h(i,x) = (i+m_x,x)$$
 for all $x \in \{1,\ldots,n\}$ and all $i \ge z_x$.

Define a map ϕ as follows:

(7.2) $\phi: H_n \to \{(m_1, \dots, m_n) \in \mathbb{Z}^n : \sum m_i = 0\} \cong \mathbb{Z}^{n-1}$

(7.3) $\phi: h \mapsto (m_1, \dots, m_n).$

Its kernel is exactly the permutations which are "eventually zero" on S, i.e. the infinite symmetric group Sym_{∞} (the finite support permutations of the countable set S).

7.1. Centralisers of finite subgroups in H_n

First we recall some properties of group actions on sets, before specialising to Houghton's group.

PROPOSITION 7.1.1. If G is a group acting on a countable set X and H is any subgroup of G then

- (1) If x and y are in the same G-orbit then their isotropy subgroups G_x and G_y are G-conjugate.
- (2) If $g \in C_G(H)$ then $H_{qx} = H_x$ for all $x \in X$.
- (3) Partition X into {X_a}^t_{a=1}, where t ∈ N ∪ {∞}, via the equivalence relation x ~ y if and only if H_x is H-conjugate to H_y. Any two points in the same H-orbit will lie in the same partition and any c ∈ C_G(H) maps X_a onto X_a for all a.
- (4) Let G act faithfully on X, with the property that for all $g \in G$ and $X_a \subseteq X$ as in the previous section, there exists a group element $g_a \in G$ which fixes $X \setminus X_a$ and acts as g does on X_a . Then $C_G(H) = C_1 \times \cdots \times C_t$ where C_a is the subgroup of $C_G(H)$ acting trivially on $X \setminus X_a$.

PROOF. (1) and (2) are standard results.

- (3) This follows immediately from (1) and (2).
- (4) This follows from (3) and our new assumption on G: Let $c \in C_G(H)$ and c_a be the element given by the assumption. Since the action of Gon X is faithful, c_a is necessarily unique. That the action is faithful also implies $c = c_1 \cdots c_t$ and that any two c_a and c_b commute in G because they act non-trivially only on distinct X_a . Thus we have the necessary isomorphism $C_G(H) \longrightarrow C_1 \times \cdots \times C_t$.

Let $Q \leq H_n$ be a finite subgroup of Houghton's group H_n and $S_Q = S \setminus S^Q$ the set of points of S which are *not fixed* by Q. Q being finite implies $\phi(Q) = 0$ as any element q with $\phi(q) \neq 0$ necessarily has infinite order. For every $q \in Q$ there exists $\{z_1, \ldots, z_n\} \in \mathbb{N}^n$ such that

$$q(i,x) = (i,x)$$
 if $i \ge z_x$.

Taking z'_x to be the maximum of these z_x over all elements in Q, then Q must fix the set $\{(i, x) : i \ge z'_x\}$ and in particular $S_Q \subseteq \{(i, x) : i < z'_x\}$ is finite.

We need to see that the subgroup $Q \leq H_n$ acting on the set S satisfies the conditions of Proposition 7.1.1(4). We give the following lemma in more generality than is needed here, as it will come in useful later on. That the action is faithful is automatic as an element $h \in H_n$ is uniquely determined by its action on the set S. LEMMA 7.1.2. Let $Q \leq H_n$ be a subgroup, which is either finite or of the form $F \rtimes \mathbb{Z}$ for F a finite subgroup of H_n . Partition S with respect to Q into sets $\{S_a\}_{a=1}^t$ as in Proposition 7.1.1(3) applied to the action of H_n on S and the subgroup Q of H_n . Then the conditions of Proposition 7.1.1(4) are satisfied.

PROOF. Fix $a \in \{1, \ldots, t\}$ and let h_a denote the permutation of S which fixes $S \setminus S_a$ and acts as h does on S_a . We wish to show that h_a is an element of H_n .

There are only finitely many elements in Q with finite order so as in the argument just before this lemma we may choose integers z_x for $x \in \{1, \ldots, n\}$ such that if q is a finite order element of Q then q(i, x) = (i, x) whenever $i \ge z_x$. If Q is a finite group then either:

• S_a is fixed by Q, in which case

$$\{(i,x) : i \ge z_x, x \in \{1,\ldots,n\}\} \subseteq S_a$$

so $h_a(i,x) = h(i,x)$ for all $i \ge z_x$. In particular for large enough i, h_a acts as a translation on (i,x) and is hence an element of H_n .

Or

• S_a is not fixed by Q, in which case

$$S_a \subseteq \{(i, x) : i < z_x, x \in \{1, \dots, n\}\}.$$

In particular S_a is finite and $h_a(i, x) = (i, x)$ for all $i \ge z_x$. Hence h_a is an element of H_n .

It remains to treat the case where $Q = F \rtimes \mathbb{Z}$. Write w for a generator of \mathbb{Z} in $F \rtimes \mathbb{Z}$. By choosing a larger z_x if needed we may assume w acts either trivially or as a translation on (i, x) whenever $i \ge z_x$. Hence for any $x \in \{1, \ldots, n\}$, the isotropy group in Q of $\{(i, x) : i \ge z_x\}$ is either F or Q.

If S_a has isotropy group Q or F then for some $x \in \{1, \ldots, n\}$, either

٠

$$S_a \cap \{(i,x) : i \ge z_x\} = \{(i,x) : i \ge z_x\}$$

in which case $h_a(i, x) = h(i, x)$ for $i \ge z_x$. In particular for large enough i, h_a acts as a translation on (i, x) and hence is an element of H_n .

Or

$$S_a \cap \{(i, x) : i \ge z_x\} = \emptyset$$

in which case $h_a(i, x) = (i, x)$ for $i \ge z_x$. In particular for large enough i, h_a fixes (i, x) and hence is an element of H_n .

If S_a is the set corresponding to an isotropy group not equal to F or Q then

 $S_a \subseteq \{(i,x) : i \ge z_x, x \in \{1,\ldots,n\}\}.$

So h_a fixes (i, x) for $i \ge z_x$ and hence h_a is an element of H_n .

Partition S into disjoint sets according to the Q-conjugacy classes of the stabilisers, as in Proposition 7.1.1(3). The set with isotropy in Q equal to Q is S^Q and since S_Q is finite the partition is finite, thus

$$S = S^Q \cup S_1 \cup \dots \cup S_t.$$

Proposition 7.1.1(4) gives that

$$C_{H_n}(Q) = H_n|_{S^Q} \times C_1 \times \ldots \times C_t$$

where each C_a acts only on S_a and leaves S^Q and S_b fixed for $a \neq b$ (where $a, b \in \{1, \ldots, t\}$). The first element of the direct product decomposition is the subgroup of $C_{H_n}(Q)$ acting only on S^Q and leaving $S \setminus S^Q$ fixed. This is $H_n|_{S^Q}$ (H_n restricted to S^Q) because, as the action of Q on S^Q is trivial, any permutation of S^Q will centralise Q. Choose a bijection $S^Q \to S$ such that for all x, $(i, x) \mapsto (i + m_x, x)$ for large enough i and some $m_x \in \mathbb{Z}$, this induces an isomorphism between $H_n|_{S^Q}$ and H_n .

To give an explicit definition of the group C_a we need three lemmas.

LEMMA 7.1.3. C_a is isomorphic to the group T of Q-set automorphisms of S_a .

PROOF. An element $c \in C_a$ determines a Q-set automorphism of S_a , giving a map $C_a \to T$. Since the action of C_a on S_a is faithful this map is injective. Any Q-set automorphism α of S_a may be extended to a Q-set automorphism of S, where α acts trivially on $S \setminus S_a$. Since S_a is a finite set, α acts trivially on (i, x) for large enough i and any $x \in \{1, \ldots, n\}$, and hence α is an element of H_n . Finally, since α is a Q-set automorphism $q\alpha s = \alpha qs$, equivalently $\alpha^{-1}q\alpha s = s$, for all $s \in S$ and $q \in Q$, showing that $\alpha \in C_a$ and so the map $C_a \to T$ is surjective. \Box

LEMMA 7.1.4. S_a is Q-set isomorphic to the disjoint union of r copies of Q/Q_a , where Q_a is an isotropy group of S_a and $r = |S_a|/|Q:Q_a|$.

PROOF. S_a is finite and so splits as a disjoint union of finitely many Q-orbits. Choose orbit representatives $\{s_1, \ldots, s_r\} \subset S_a$ for these orbits, these s_k may be chosen to have the same Q-stabilisers: If $Q_{s_1} \neq Q_{s_2}$ then there is some $q \in Q$ such that $Q_{qs_2} = qQ_{s_2}q^{-1} = Q_{s_1}$ (the partitions S_a were chosen to have this property by Proposition 7.1.1), iterating this procedure we get a set of representatives who all have isotropy group Q_{s_1} . Now set $Q_a = Q_{s_1}$ and note that there are $|Q:Q_a|$ elements in each of the Q-orbits so $r|Q:Q_a| = |S_a|$. Recall that if G is any group and $r \ge 1$ is some natural number then the wreath product $G \wr \operatorname{Sym}_r$ is the semi-direct product

$$G\wr \operatorname{Sym}_r = \prod_{k=1}^r G \rtimes \operatorname{Sym}_r$$

where the symmetric group Sym_r acts by permuting the factors in the direct product.

Recall also that for any subgroup H of a group G, the Weyl group W_GH is defined to be $W_GH = N_GH/H$.

LEMMA 7.1.5. The group C_a is isomorphic to the wreath product $W_Q Q_a \wr \operatorname{Sym}_r$, where Q_a is some isotropy group of S_a and $r = |S_a|/|Q : Q_a|$.

PROOF. Using Lemmas 7.1.3 and 7.1.4, C_a is isomorphic to the group of Q-set automorphisms of the disjoint union of r copies of Q/Q_a .

To begin, we show the group of automorphisms of the Q-set Q/Q_a is isomorphic to W_QQ_a . An automorphism $\alpha : Q/Q_a \to Q/Q_a$ is determined by the image $\alpha(Q_a) = qQ_a$ of the identity coset and such an element determines an automorphism if and only if $q^{-1}Q_aq \leq Q_a$, equivalently $q \in N_QQ_a$. Since two elements $q_1, q_2 \in Q$ will determine the same automorphism if and only if $q_1Q_a = q_2Q_a$, the group of Q-set automorphisms of Q/Q_a is the Weyl group W_QQ_a .

For the general case, note that if $c \in C_a$ then c permutes the Q-orbits $\{Qs_1, \ldots, Qs_r\}$, so there is a map $\pi : C_a \to \text{Sym}_r$. Assume that the representatives $\{s_1, \ldots, s_r\}$ have been chosen, as in the proof of Lemma 7.1.4, to have the same Q-stabilisers. The map π is split by the map

$$\begin{split} \iota: \operatorname{Sym}_r &\to C_a \\ & \sigma \mapsto \left(\iota(\sigma): qs_k \mapsto qs_{\sigma(k)} \text{ for all } q \in Q \right). \end{split}$$

Each $\iota(\sigma)$ is a well defined element of H_n since

$$qs_k = \tilde{q}s_k \Leftrightarrow \tilde{q}^{-1}q \in Q_{s_k} = Q_{s_{\sigma(k)}} \Leftrightarrow qs_{\sigma(k)} = \tilde{q}s_{\sigma(k)}.$$

The kernel of the map π is exactly the elements of C_a which fix each Q-orbit but may permute the elements inside the Q-orbits, by the previous part this is exactly $\prod_{k=1}^{r} W_Q Q_a$. For any $\sigma \in \text{Sym}_r$, the element $\iota(\sigma)$ acts on $\prod_{k=1}^{r} W_Q Q_a$ by permuting the factors, so the group C_a is indeed isomorphic to the wreath product.

The centraliser $C_{H_n}Q$ can now be completely described.

PROPOSITION 7.1.6. The centraliser $C_{H_n}(Q)$ of any finite subgroup $Q \leq H_n$ splits as a direct product

$$C_{H_n}(Q) \cong H_n|_{S^Q} \times C_1 \times \cdots \times C_t,$$



FIGURE 1. A representation of S_a . The large circles are the sets $\{Qs_1 \ldots, Qs_r\}$ (in this figure r = 3). Elements of Sym_r permute only the large circles, while elements of $\prod_{k=1}^r W_Q Q_a$ leave the large circles fixed and permute only elements inside them.

where $H_n|_{S^Q} \cong H_n$ is Houghton's group restricted to S^Q and for all $a \in \{1, \ldots, t\}$,

$$C_a \cong W_Q Q_a \wr \operatorname{Sym}_r$$

for Q_a is an isotropy group of S_a and $r = |S_a|/|Q : Q_a|$. In particular H_n is finite index in $C_{H_n}(Q)$.

PROOF. We have already proven that

$$C_{H_n}(Q) \cong H_n|_{S^Q} \times C_1 \times \cdots \times C_t$$

and Lemma 7.1.5 gives the required description of C_a .

COROLLARY 7.1.7. If Q is a finite subgroup of H_n then the centraliser $C_{H_n}(Q)$ is FP_{n-1} but not FP_n .

PROOF. H_n is finite index in the centraliser $C_{H_n}(Q)$ by Proposition 7.1.6. Appealing to Brown's result [**Bro87**, 5.1] that H_n is FP_{n-1} but not FP_n, and that a group is FP_n if and only if a finite index subgroup is FP_n [**Bro94**, VIII.5.5.1] we can deduce $C_{H_n}(Q)$ is FP_{n-1} but not FP_n.

7.2. Centralisers of elements in H_n

If $q \in H_n$ is an element of finite order then the subgroup $Q = \langle q \rangle$ is a finite subgroup and the previous section may be used to describe the centraliser $C_{H_n}(q) = C_{H_n}(Q)$. Thus for an element q of finite order $C_{H_n}(q) \cong C \times H_n$ for some finite group C.

If $q \in H_n$ is an element of infinite order and $Q = \langle q \rangle$ then we may apply Proposition 7.1.1(3) to split up S into a disjoint collection $\{S_a : a \in A \subseteq \mathbb{N}\} \cup S^Q$ (S^Q) is the element of the collection associated to the isotropy group Q). Assume that S_0 is the set associated to the trivial isotropy group. Since q is a translation on $(i, x) \in S = \mathbb{N} \times \{1, \ldots, n\}$ for large enough i and points acted on by such a translation have trivial isotropy, there are only finitely many elements of S whose isotropy group is neither the trivial group nor Q. Hence S_a is finite for $a \neq 0$ and the set A is finite. From now on let $A = \{0, \ldots, t\}$. We now use Lemma 7.1.2 and Proposition 7.1.1(4) as in the previous section: $C_{H_n}(Q)$ splits as

$$C_{H_n}(Q) \cong C_0 \times C_1 \times \cdots \times C_t \times H_n|_{S^Q}$$

where C_a acts only on S_a and $H_n|_{S^Q}$ is Houghton's group restricted to S^Q . Unlike in the last section, $H_n|_{S^Q}$ may not be isomorphic to H_n . Let $J \subseteq \{1, \ldots, n\}$ satisfy

 $x \in J$ if and only if $(i, x) \in S^Q$ for all $i \ge z_x$, some $z_x \in \mathbb{N}$.

If $x \notin J$ then for large enough i, q must act as a non-trivial translation on (i, x), and the set $(\mathbb{N} \times \{x\}) \cap S^Q$ is finite. Clearly $|J| \leq n-2$, but different elements q may give values $0 \leq |J| \leq n-2$. In the case |J| = 0, S^Q is necessarily finite and so $H_n|_{S^Q}$ is isomorphic to a finite symmetric group on S^Q . It is also possible that $S^Q = \emptyset$, in which case $H_n|_{S^Q}$ is just the trivial group. If $|J| \neq 0$ then the argument proceeds by choosing a bijection

$$S^Q \to \mathbb{N} \times J$$

such that $(i, x) \mapsto (i + m_x, x)$ for some $m_x \in \mathbb{Z}$ whenever *i* is large enough and $x \in J$. This set map induces a group isomorphism between $H_n|_{S^Q}$ and $H_{|J|}$ (Houghton's group on the set $J \times \mathbb{N}$).

Lemma 7.1.5 describes the groups C_a for $a \neq 0$, so it remains only to treat the case a = 0. We cannot use the arguments used for $a \neq 0$ here as the set S_0 is not finite, in particular Lemma 7.1.3 doesn't apply: Every *Q*-set isomorphism of S_0 is realised by an element of the infinite support permutation group on S_0 , but there are *Q*-set isomorphisms of S_0 which are not realised by an element of H_n .

The next three lemmas are needed to describe C_0 , this description will use the graph Γ which we now describe. The vertices of Γ are those $x \in \{1, \ldots, n\}$ for which q acts non-trivially on infinitely many elements of $\mathbb{N} \times \{x\}$. Equivalently, the vertices are the elements of $\{1, \ldots, n\} \setminus J$. There is an edge from x to y in Γ if there exists $s \in S_0$ and $N \in \mathbb{N}$ such that for all $m \geq N$ we have $q^{-m}s \in \mathbb{N} \times \{x\}$ and $q^m s \in \mathbb{N} \times \{y\}$. Let $\pi_0 \Gamma$ denote the path components of Γ , and for any vertex x of Γ denote by [x] the element of $\pi_0 \Gamma$ corresponding to that vertex.

Let $z \in \mathbb{N}$ be some integer such that for all $i \geq z$, q acts trivially or as a translation on (i, x) for all $x \in \{1, \ldots, n\}$. Fix z for the remainder of this section.

For each path component [x] in $\pi_0\Gamma$, let $S_0^{[x]}$ denote the smallest *Q*-subset of S_0 containing the set $\{(i, y) : i \geq z, y \in [x]\}$. Note that $(i, y) \notin S_0^{[x]}$ for any $y \notin [x]$ and $i \geq z$, since if (i, x) and (j, y) are two elements of S_0 in the same

Q-orbit with $i \ge z$ and $j \ge z$ then there is an edge between x and y in Γ : If $(i, x) = q^k(j, y)$ and q acts as a positive translation on the element (i, x) then let N = k and s = (i, x), similarly for when q acts as a negative translation. This gives a Q-set decomposition of S_0 as

$$S_0 = \coprod_{[x]\in\pi_0\Gamma} S_0^{[x]},$$

where \coprod denotes disjoint union.

LEMMA 7.2.1. Let $[x] \in \pi_0 \Gamma$, if $C_0^{[x]}$ denotes the subgroup of C_0 which acts non-trivially only on $S_0^{[x]}$ then there is an isomorphism

$$C_0 \cong C_0^{[x_1]} \times \cdots \times C_0^{[x_r]},$$

where $[x_1], [x_2], \ldots, [x_r]$ are all elements of $\pi_0 \Gamma$.

PROOF. If $c \in C_0$ and $[x] \in \pi_0 \Gamma$ then let $c_{[x]}$ denote the permutation of S such that $c_{[x]}$ acts as c does on $S_0^{[x]}$, and acts trivially on $S \setminus S_0^{[x]}$. We will show that $c_{[x]}$ is an element of C_0 . Since the action of C_0 on S_0 is faithful it follows that the elements $c_{[x]}$ and $c_{[y]}$ commute and

$$c = c_{[x_1]} c_{[x_2]} \cdots c_{[x_r]},$$

which suffices to prove the lemma.

Let $y \in \{1, \ldots, n\}$. The element $c_{[x]}$ acts trivially on (i, y) for $i \ge z$ if $y \notin [x]$ and acts as as c does on (i, y) for $i \ge z$ if $y \in [x]$, thus $c_{[x]}$ is an element of H_n . Since $c_{[x]}$ is also a Q-set automorphism of S, $c_{[x]}$ is a member of C_0 .

LEMMA 7.2.2. Let $[x] \in \pi_0 \Gamma$, let $c \in C_0$, and let $z' \in \mathbb{N}$ be such that c acts either trivially or as a translation on (i, x) for all $x \in \{1, \ldots, n\}$ and $i \geq z'$. Then the action of c on some element $(i, x) \in S$ for $i \geq z'$ completely determines the action of c on $S_0^{[x]}$.

PROOF. Firstly, note that knowing the action of c on some element (i, x) for $i \ge z'$ determines the action of c on the set $\{(i, x) : i \ge z'\}$, since we chose z' in order to have this property.

Let $y \in [x]$ such that there is an edge from x to y, so there is a natural number N and element $s \in S_0^{[x]}$ such that $q^N s = (i, x)$ and $q^{-N} s = (j, y)$ for some natural numbers i and j. By choosing N larger if necessary we can take $i, j \ge z'$. The action of c on (j, y) is now completely determined by the action on (i, x), since

$$c(j,y) = cq^{-2N}(i,x) = q^{-2N}c(i,x)$$

For any $y \in [x]$ there is a path from x to y in Γ , so we've determined the action of c on the set $X = \{(j, y) : j \geq z', y \in [x]\}$. If $s \in S_0^{[x]} \setminus X$ then, since $S_0^{[x]} \setminus X$

is finite, there is some integer m with $q^m s = x \in X$. So $cs = cq^{-m}x = q^{-m}cx$, which completely determines the action of c on s.

LEMMA 7.2.3. For any $[x] \in \pi_0 \Gamma$, there is an isomorphism

$$C_0^{[x]} \cong \mathbb{Z}$$

PROOF. By Lemma 7.2.2 the action is completely determined by the action on some element (i, x) for large enough i, and the action on this element is necessarily by translation by some element $m_x(c)$. This defines an injective homomorphism $C_0^{[x]} \to \mathbb{Z}$, sending $c \mapsto m_x(c)$. Let $q_{[x]}$ be the element of $C_0^{[x]}$ described in the proof of Lemma 7.2.1, $q_{[x]}$ is a non-trivial element of $C_0^{[x]}$ so $C_0^{[x]}$ is mapped isomorphically onto a non-trivial subgroup of \mathbb{Z} .

Combining Lemmas 7.2.1 and 7.2.3 shows $C_0 \cong \mathbb{Z}^r$ where $r = |\pi_0 \Gamma|$.

Recall that the vertices of Γ are indexed by the set $\{1, \ldots, n\} \setminus J$. Since there are no isolated vertices in Γ , $|\pi_0\Gamma| \leq \lfloor (n-|J|)/2 \rfloor$ (where $\lfloor - \rfloor$ denotes the integer floor function). Recalling that $0 \leq |J| \leq n-2$, the set $\{1, \ldots, n\} \setminus J$ is necessarily non-empty so $1 \leq |\pi_0\Gamma|$, combining these gives

$$1 \le |\pi_0 \Gamma| \le \lfloor (n - |J|)/2 \rfloor.$$

We can now completely describe the centraliser $C_{H_n}(q)$.

THEOREM 7.2.4.

(1) If $q \in H_n$ is an element of finite order then

$$C_{H_n}(q) \cong H_n|_{S^Q} \times C_1 \times \cdots \times C_t$$

where $H_n|_{S^Q} \cong H_n$ is Houghton's group restricted to S^Q and for all $a \in \{1, \ldots, t\},\$

$$C_a \cong W_Q Q_a \wr \operatorname{Sym}_r$$

for Q_a an isotropy group of S_a and $r = |S_a|/|Q : Q_a|$. In particular H_n is finite index in $C_{H_n}Q$.

(2) If $q \in H_n$ is an element of infinite order then either

$$C_{H_n}(q) \cong H_k \times \mathbb{Z}^r \times C_1 \times \cdots \times C_t$$

or

$$C_{H_n}(q) \cong F \times \mathbb{Z}^r \times C_1 \times \cdots \times C_t$$

where F is some finite symmetric group, H_k is Houghton's group with $0 \le k \le n-2$, and the groups C_a are as in the previous part. In the first case $1 \le r \le \lfloor (n-k)/2 \rfloor$, and in the second case $1 \le r \le \lfloor n/2 \rfloor$.

7. HOUGHTON'S GROUPS

In Corollary 7.1.7 it was proved that for an element q of finite order, $C_{H_n}(q)$ is FP_{n-1} but not FP_n . The situation is much worse for elements q of infinite order, in which case the centraliser may not even be finitely generated, for example when n is odd and q is the element acting on $S = \mathbb{N} \times \{1, \ldots, n\}$ as

$$q: \begin{cases} (i,x) \mapsto (i+1,x) & \text{if } x \le (n-1)/2\\ (i,x) \mapsto (i-1,x) & \text{if } (n+1)/2 \le x \le n-1 \text{ and } i \ne 0\\ (0,x) \mapsto (0,x-((n-1)/2)) & \text{if } (n+1)/2 \le x \le n-1\\ (i,n) \mapsto (i,n) \end{cases}$$

then the only fixed points are on the ray $\mathbb{N} \times \{n\}$. The argument leading up to Theorem 7.2.4 shows that the centraliser is a direct product of groups, one of which is Houghton's group H_1 which is isomorphic to the infinite symmetric group and hence not finitely generated. In particular for this q, the centraliser $C_{H_n}(q)$ is not even FP₁. A similar example can easily be constructed when n is even.

All the groups in the direct product decomposition from Theorem 7.2.4 except H_k are FP_{∞}, being built by extensions from finite groups and free Abelian groups. By choosing various infinite order elements q, for example by modifying the example of the previous paragraph, the centralisers can be chosen to be FP_k for $0 \le k \le n-3$. The upper bound of n-3 arises because any infinite order element q must necessarily be "eventually a translation" (in the sense of (7.1)) on $\mathbb{N} \times \{x\}$ for at least two x. As such the copy of Houghton's group in the centraliser can act on at most n-2 rays and is thus at largest H_{n-2} , which is FP_{n-3}.

COROLLARY 7.2.5. If Q is an infinite virtually cyclic subgroup of H_n then either

$$C_{H_n}(Q) \cong H_k \times \mathbb{Z}^r \times C_1 \times \cdots \times C_t$$

or

$$C_{H_n}(Q) \cong F \times \mathbb{Z}^r \times C_1 \times \cdots \times C_t$$

where the elements in the decomposition are all as in Theorem 7.2.4.

This corollary can be proved by reducing to the case of Theorem 7.2.4, but before that we require the following lemma.

LEMMA 7.2.6. Every infinite virtually cyclic subgroup Q of H_n is finite-by- \mathbb{Z} .

PROOF. By [**JPL06**, Proposition 4], Q is either finite-by- \mathbb{Z} or finite-by- D_{∞} where D_{∞} denotes the infinite dihedral group, we show the latter cannot occur. Assume that there is a short exact sequence of groups

$$1 \longrightarrow F \longmapsto Q \xrightarrow{\pi} \mathcal{D}_{\infty} \longrightarrow 1,$$

regarding F as a subgroup of Q. Let a, b generate D_{∞} , so that

$$\mathbf{D}_{\infty} = \langle a, b \mid a^2 = b^2 = 1 \rangle.$$

Let $p, q \in Q$ be lifts of a, b, such that $\pi(p) = a, \pi(q) = b$, then $p^2 \in F$. Since F is finite, p^2 has finite order and hence p has finite order. The same argument shows that q has finite order. The element $pq \in Q$ necessarily has infinite order as $\pi(pq)$ is infinite order in D_{∞} .

However, since p and q are finite order elements of H_n , by the argument at the beginning of Section 7.1 they both permute only a finite subset of S. Thus pq permutes a finite subset of S and is of finite order, but this contradicts the previous paragraph.

PROOF OF COROLLARY 7.2.5. Using the previous lemma, write Q as $Q = F \rtimes \mathbb{Z}$ where F is a finite group. As F is finite, the set S_F of points not fixed by F is finite (see the argument at the beginning of Section 7.1). Let $z \in \mathbb{N}$ be such that for $i \ge z$, F acts trivially on (i, x) for all x, and \mathbb{Z} acts on (i, x)either trivially or as a translation. Applying Lemma 7.1.2 and Proposition 7.1.1, S splits as a disjoint union

$$S = S^Q \cup S_0 \cup S_1 \cup \dots \cup S_t$$

where S^Q is the fixed point set, S_0 is the set with isotropy group F and the S_a for $1 \le a \le t$ are subsets of $\{(i, x) : i \le z\}$, and hence all finite. By Proposition 7.1.1, $C_{H_n}(Q)$ splits as a direct product

$$C = H_n|_{S^Q} \times C_0 \times C_1 \times \ldots \times C_t$$

where $H_n|_{S^Q}$ denotes Houghton's group restricted to S^Q . The argument of Theorem 7.2.4 showing that $H_n|_{S^Q}$ is isomorphic to either a finite symmetric group or to H_k for some $0 \le k \le n-2$ goes through with no change, as does the proof of the structure of the groups C_a for $1 \le a \le t$. It remains to observe that because every element in S_0 is fixed by F, any element of H_n centralising \mathbb{Z} and fixing $S \setminus S_0$ necessarily also centralises Q and is thus a member of C_0 . This reduces us again to the case of Theorem 7.2.4 showing that $C_0 \cong \mathbb{Z}^r$ for some natural number $1 \le r \le \lfloor (n-k)/2 \rfloor$, or $1 \le r \le \lfloor n/2 \rfloor$ if $H_n|_{S^Q}$ is a finite symmetric group.

7.3. Brown's model for $E_{\text{fin}}H_n$

The main result of this section will be Corollary 7.3.4, where the construction of Brown [**Bro87**] used to prove that H_n is FP_{n-1} but not FP_n is shown to be a model for $E_{\mathcal{F}in}H_n$.

In this section, maps are written from left to right.

Write \mathcal{M} for the monoid of injective maps $S \to S$ with the property that every permutation is "eventually a translation" (in the sense of (7.1)), and write T for the free monoid generated by $\{t_1, \ldots, t_n\}$ where

$$(i,x)t_y = \begin{cases} (i+1,x) & \text{if } x = y, \\ (i,x) & \text{if } x \neq y. \end{cases}$$

The elements of T will be called *translations*. The map $\phi : H_n \to \mathbb{Z}^n$, defined in (7.2), extends naturally to a map $\phi : \mathcal{M} \to \mathbb{Z}^n$. Give \mathcal{M} a poset structure by setting $\alpha \leq \beta$ if $\beta = t\alpha$ for some $t \in T$. The monoid \mathcal{M} can be given the obvious action on the right by H_n , which in turn gives an action of H_n on the poset (\mathcal{M}, \leq) since $\beta = t\alpha$ implies $\beta h = t\alpha h$ for all $h \in H_n$. Let $|\mathcal{M}|$ be the geometric realisation of this poset, namely simplicies in $|\mathcal{M}|$ are finite ordered collections of elements in \mathcal{M} with the obvious face maps. An element $h \in H_n$ fixes a vertex $\{\alpha\} \in |\mathcal{M}|$ if and only if $s\alpha h = s\alpha$ for all $s \in S$ if and only if hfixes $S\alpha$, so the stabiliser $(H_n)_{\alpha}$ may only permute the finite set $S \setminus S\alpha$ and we may deduce:

PROPOSITION 7.3.1. Stabilisers of simplicies in $|\mathcal{M}|$ are finite.

We now build up to the the proof that $|\mathcal{M}|$ is a model for $\mathbb{E}_{\mathcal{F}_{in}}H_n$ with a few lemmas.

PROPOSITION 7.3.2. If $Q \leq H_n$ is a finite group then the fixed point set $|\mathcal{M}|^Q$ is non-empty and contractible.

PROOF. For all $q \in Q$, choose $\{z_0(q), \ldots, z_n(q)\}$ to be an *n*-tuple of natural numbers such that (i, x)q = (i, x) whenever $i \geq z_x(q)$ for all *i*. *Q* then fixes all elements $(i, x) \in S$ with $i \geq \max_Q z_x(q)$. Define a translation $t = t_1^{\max_Q z_1(q)} \cdots t_n^{\max_Q z_n(q)}, t \in \mathcal{M}^Q$ so $\{t\}$ is a vertex of $|\mathcal{M}|^Q$ and $|\mathcal{M}|^Q \neq \emptyset$.

If $\{m\}, \{n\} \in |\mathcal{M}|^Q$ then let $a, b \in T$ be two translations such that

$$\phi(m) - \phi(n) = \phi(b) - \phi(a)$$

(recall that for a translation t, $\phi(t)$ must be an *n*-tuple of positive numbers). Thus $\phi(am) = \phi(bn)$, and since $am, bn \in \mathcal{M}$ there exist *n*-tuples $\{z_1, \ldots, z_n\}$ and $\{z'_1, \ldots, z'_n\}$ such that am acts as a translation for all $(i, x) \in S$ with $i \geq z_x$ and bn acts as a translation for all $(i, x) \in S$ with $i \geq z'_x$. Let

$$c = t_1^{\max\{z_1, z_1'\}} \dots t_n^{\max\{z_n, z_n'\}}$$

so that cam = cbn, further pre-composing c with a large translation (for example that from the first section of this proof) we can assume that $cam = cbn \in \mathcal{M}^Q$, and $\{cam = cbn\} \in |\mathcal{M}|^Q$. This shows the poset \mathcal{M}^Q is directed and hence the simplicial realisation $|\mathcal{M}^Q| = |\mathcal{M}|^Q$ is contractible.

PROPOSITION 7.3.3. If $Q \leq H_n$ is an infinite group then $|\mathcal{M}|^Q = \emptyset$.

PROOF. Consider an infinite subgroup $Q \leq H_n$ with $|\mathcal{M}|^Q \neq \emptyset$ and choose some vertex $\{m\} \in |\mathcal{M}|^Q$. For any $q \in Q$, since mq = m it must be that $\phi(m) + \phi(q) = \phi(m)$ and so $\phi(q) = 0$, hence Q is a subgroup of $\operatorname{Sym}_{\infty} \leq H_n$. Furthermore Q must permute an infinite subset of S (if it permuted just a finite set it would be a finite subgroup). That mq = m implies that this infinite subset is a subset of $S \setminus Sm$ but this is finite by construction. So the fixed point subset $|\mathcal{M}|^Q$ for any infinite subgroup Q is empty. \Box

COROLLARY 7.3.4. $|\mathcal{M}|$ is a model for $\mathbb{E}_{\text{Fin}}H_n$.

PROOF. Combine Propositions 7.3.1, 7.3.2 and 7.3.3.

7.4. Finiteness conditions satisfied by H_n

Recall from Proposition 3.6.1 that a group G is $\mathcal{O}_{\text{fin}} \text{FP}_0$ if and only if it has finitely many conjugacy classes of finite subgroups. G satisfies the weaker quasi- $\mathcal{O}_{\text{fin}} \text{FP}_0$ condition if and only if it has finitely many conjugacy classes of subgroups isomorphic to a given finite subgroup (see Section 3.6.1).

PROPOSITION 7.4.1. H_n is not quasi- $\mathcal{O}_{\text{Fin}} \text{FP}_0$.

Before the above proposition is proved, we need a lemma. In the infinite symmetric group $\operatorname{Sym}_{\infty}$ acting on the set S, elements can be represented by products of disjoint cycles. We use the standard notation for a cycle: (s_1, s_2, \ldots, s_m) represents the element of $\operatorname{Sym}_{\infty}$ sending $s_i \mapsto s_{i+1}$ for i < n and $s_n \mapsto s_1$. Any element of finite order in H_n is contained in the infinite symmetric group $\operatorname{Sym}_{\infty}$ by the argument at the beginning of Section 7.1. We say two elements of $\operatorname{Sym}_{\infty}$ have the same cycle type if they have the same number of cycles of length m for each $m \in \mathbb{N}$.

LEMMA 7.4.2. If q is a finite order element of H_n and h is an arbitrary element of H_n , then hqh^{-1} is the permutation given in the disjoint cycle notation by applying h to each element in each disjoint cycle of q. In particular, if q is represented by the single cycle (s_1, \ldots, s_m) , then hqh^{-1} is represented by (hs_1, \ldots, hs_m) .

Furthermore, two finite order elements of H_n are conjugate if and only if they have the same cycle type.

PROOF. The proof of the first part is analogous to [**Rot95**, Lemma 3.4]. Let q be an element of finite order and h an arbitrary element of H_n . If q fixes $s \in S$ then hqh^{-1} fixes hs. If q(i) = j, h(i) = k and h(j) = l, for $i, j, k, l \in S$, then $hqh^{-1}(k) = l$ exactly as required.

By the above, conjugate elements have the same cycle type. For the converse, notice any two finite order elements with the same cycle type necessarily lie in Sym_r for some $r \in \mathbb{N}$ so by [**Rot95**, Theorem 3.5] they are conjugated by an element of Sym_r .

PROOF OF PROPOSITION 7.4.1. Choosing a collection of elements q_i for each $i \in \mathbb{N}_{\geq 1}$, so that q_i has *i* disjoint 2-cycles gives a collection of isomorphic subgroups which are all non-conjugate by Lemma 7.4.2.

PROPOSITION 7.4.3. $\mathcal{O}_{\text{fin}} \operatorname{cd} H_n = \operatorname{gd}_{\text{fin}} H_n = n.$

PROOF. As described in the introduction, H_n can be written as

$$\operatorname{Sym}_{\infty} \hookrightarrow H_n \longrightarrow \mathbb{Z}^{n-1}$$

Now, $\operatorname{gd}_{\operatorname{fin}} \mathbb{Z}^{n-1} = \operatorname{gd} \mathbb{Z}^{n-1} = n-1$ and $\operatorname{gd}_{\operatorname{fin}} \operatorname{Sym}_{\infty} = 1$ by [**LW12**, Theorem 4.3], as it is the colimit of its finite subgroups each of which have proper geometric dimension 0, and the directed category over which the colimit is taken has homotopy dimension 1 [**LW12**, Lemma 4.2]. \mathbb{Z}^{n-1} is torsion free and so has a bound of 1 on the orders of its finite subgroups and we deduce from [**Lüc00**, Theorem 3.1] that $\operatorname{gd}_{\operatorname{fin}} H_n \leq n-1+1=n$.

To deduce the other bound, we use an argument due to Gandini [Gan12a]. Assume that $\mathcal{O}_{\text{fin}} \operatorname{cd} H_n \leq n-1$. By [BLN01, Theorem 2] we have

$$\operatorname{cd}_{\mathbb{Q}} H_n \leq \mathcal{O}_{\operatorname{Fin}} \operatorname{cd} H_n = n - 1.$$

In [**Bro87**, Theorem 5.1], it is proved that H_n is FP_{n-1} (but not FP_n), combining this with [**LN01**, Proposition 1] we deduce that there is a bound on the orders of the finite subgroups of H_n , but this is obviously a contradiction, thus

$$n \leq \mathcal{O}_{\mathcal{F}in} \operatorname{cd} H_n \leq \operatorname{gd}_{\mathcal{F}in} H_n \leq n.$$

The remainder of this section is devoted to proving the following.

THEOREM 7.4.4. $\mathcal{O}_{\mathcal{VCyc}} \operatorname{cd} H_n = n.$

The proof is based on a pushout of Lück and Weiermann [**LW12**], described below.

7.4.1. The pushout of Lück and Weiermann. For any group G, we say two infinite virtually cyclic subgroups K and K' of G are commensurate, written $K \sim K'$, if $|K \cap K'| = \infty$. Commensurability is an equivalence relation and we write $[\mathcal{V}Cyc \setminus \mathcal{F}in]$ for the set of equivalence classes. The normaliser of

an equivalence class [K] is defined to be the stabiliser of the action of G on $[\mathcal{VCyc} \setminus \mathcal{Fin}]$ by conjugation, namely

$$N_G[K] = \{ x \in G : K^x \sim K \}.$$

Associated to each infinite virtually cyclic subgroup K we define the subfamily $\mathcal{VCyc}[K]$ of \mathcal{VCyc} by

$$\mathcal{V}\!\mathit{Cyc}[K] = \{L \in \mathit{V}\!\mathit{Cyc} \setminus \mathit{Fin} : L \sim K\} \cup (\mathit{Fin} \cap K) \, .$$

If X is a right G-space and Y is a left G-space then we denote by $X \times_G Y$ the *twisted product* of X and Y, defined to be the quotient space of $X \times Y$ under the action $g \cdot (x, y) = (xg^{-1}, gy)$ [**Bre72**, §II.2]. If Y is a left H-space for some subgroup H of G then $G \times_H Y$ is a left G-space via the usual left action of G on itself.

PROPOSITION 7.4.5 ([**tD87**, Proposition I.(4.3)]). Let H be a subgroup of G, let Y be a left H-space, and let X be a left G-space. There is an adjoint isomorphism

$$[G \times_H Y, X]_G = [Y, X]_H,$$

where $[Z, X]_G$ denotes the set of G-homotopy classes of G-equivariant maps between two G-spaces Z and X.

THEOREM 7.4.6. [LW12, Theorem 2.3, Remark 2.5] Let I denote a complete set of representatives of the G-orbits in $[\mathcal{VCyc} \setminus \mathcal{Fin}]$. Choosing arbitrary $N_G[K]$ -CW-models for $E_{\mathcal{Fin}}N_G[K]$ and $E_{\mathcal{VCyc}[K]}N_G[K]$ and an arbitrary G-CW-model for $E_{\mathcal{Fin}}G$, the cellular G-pushout described below may be constructed with the maps i and $f_{[K]}$ equivariant cellular maps, and either with i an inclusion of G-CW-complexes or with every $f_{[K]}$ an inclusion of $N_G[K]$ -CW-complexes and i a cellular G-map.

Moreover the space X defined by the pushout is a model for $E_{\mathcal{VCyc}}G$.

We can describe explicitly the G-homotopy classes of the maps i and $f_{[K]}$ in the pushout above: By restricting the G-action, any model for $E_{\mathcal{F}in}G$ is a model for $E_{\mathcal{F}in}N_G[K]$ so there is a $N_G[K]$ -map $E_{\mathcal{F}in}N_G[K] \to E_{\mathcal{F}in}G$, and using the adjoint isomorphism of Proposition 7.4.5 there is a G-map $G \times_{N_G[K]} E_{\mathcal{F}in}N_G[K] \to$ $E_{\mathcal{F}in}G$. The coproduct of these maps, one for each $[K] \in I$, is the map i. Since $E_{\mathcal{F}in}N_G[K]$ is an $N_G[K]$ -space with finite isotropy, it is a priori an $N_G[K]$ -space

7. HOUGHTON'S GROUPS

with isotropy in $\mathcal{VCyc}[K]$, there is a map $E_{\mathcal{Fin}}N_G[K] \to E_{\mathcal{VCyc}[K]}N_G[K]$. This is the map $f_{[K]}$.

This pushout gives a long exact sequence in Bredon cohomology [Lüc89, Lemma 13.7]:

$$\cdots \longrightarrow H^{i}_{\mathcal{O}_{\mathcal{VCyc}}}(G,-) \longrightarrow$$

$$\left(\prod_{[K]\in I} H^{i}_{\mathcal{O}_{\mathcal{H}\mathcal{Y}\mathcal{G}}[K]}(N_{G}[K], \operatorname{Res}^{\mathcal{O}_{\mathcal{H}\mathcal{Y}\mathcal{G}}G}_{\mathcal{H}\mathcal{Y}\mathcal{G}[K]}N_{G}[K] -)\right) \oplus H^{i}_{\mathcal{O}_{\mathcal{F}in}}(G, \operatorname{Res}^{\mathcal{O}_{\mathcal{H}\mathcal{Y}\mathcal{G}}G}_{\mathcal{O}_{\mathcal{F}in}G} -)$$
$$\longrightarrow \prod_{[K]\in I} H^{i}_{\mathcal{O}_{\mathcal{F}in}}(N_{G}[K], \operatorname{Res}^{\mathcal{O}_{\mathcal{H}\mathcal{Y}\mathcal{G}}G}_{\mathcal{O}_{\mathcal{F}in}N_{G}[K]} -) \longrightarrow \cdots.$$

For brevity we will usually omit the restriction maps from now on.

Given an infinite virtually cyclic subgroup K of G, let $\pi^K : N_G K \to W K$ denote the projection map and for any $\mathcal{O}_{\mathcal{VCyc}[K]}N_G K$ -module let π^K_*M denote the $\mathcal{O}_{\mathcal{Fin}}WK$ -module given by

$$\pi^K_*M: WK/L \mapsto M(N_GK/\pi^{-1}(L)),$$

for any finite subgroup L of WK.

LEMMA 7.4.7. [**DP12**, Lemma 4.2] If K is an infinite virtually cyclic subgroup and $N_G[K] = N_G K$ then there is an isomorphism,

$$H^{i}_{\mathcal{O}_{\mathscr{V}_{\mathcal{C}\mathcal{V}^{c}}[K]}}(N_{G}K,-)\cong H^{i}_{\mathcal{O}_{\mathcal{F}in}}(W_{G}K,\pi^{K}_{*}-).$$

Combining this lemma with the long exact sequence gives the following.

PROPOSITION 7.4.8. If every G-orbit in $[\mathcal{VCyc} \setminus \mathcal{Fin}]$ contains a K such that $N_G[K] = N_G K$ then, letting A be a set of representatives with that property, there is a long exact sequence:

$$\cdots \longrightarrow H^{i}_{\mathcal{O}_{\mathscr{V}_{\mathcal{G}_{\mathscr{V}}}}}(G,-) \longrightarrow \left(\prod_{K \in A} H^{i}_{\mathcal{O}_{\mathscr{F}_{in}}}(W_{G}K,\pi^{K}_{*}-)\right) \oplus H^{i}_{\mathcal{O}_{\mathscr{F}_{in}}}(G,-)$$
$$\longrightarrow \prod_{K \in A} H^{i}_{\mathcal{O}_{\mathscr{F}_{in}}}(N_{G}K,-) \longrightarrow \cdots.$$

7.4.2. Calculation of $\mathcal{O}_{\mathcal{VCyc}} \operatorname{cd} H_n$.

LEMMA 7.4.9. For every infinite virtually cyclic subgroup K of H_n there exists L commensurate to K with

$$N_G[K] = N_G[L] = N_GL.$$

Moreover, we may assume $L \cong \mathbb{Z}$.

PROOF. Firstly replace K with a finite-index subgroup isomorphic to \mathbb{Z} and let k generate K. Consider the action of K on S. There are finitely many finite K-orbits (see Section 7.2), so for large enough m, the subgroup $\langle k^m \rangle$ acts semifreely (freely away from the fixed point set), let $L = \langle k^m \rangle$.

If $S^L \neq \emptyset$ then let $s \in S^L$ and pick any $n \in N_{H_n}[L]$, so the stabiliser of ns is L^n . Since L acts semi-freely $L^n = L$ or $L^n = 1$ but since $L^n \sim L$ this forces $L^n = L$, thus $N_{H_n}[L] = N_{H_n}L$.

If $S^L = \emptyset$ then let $n \in N_{H_n}[L]$, let $l \in L$, and let $i \in \{1, \ldots, n\}$. So for x large enough,

$$l: (i, x) \mapsto (i, x + t_l)$$
$$n: (i, x) \mapsto (i, x + t_n).$$

For some $t_l, t_n \in \mathbb{Z}$ with $t_l \neq 0$ (*L* acts non-trivially everywhere),

$$n^{-1}ln: (i, x) \mapsto (i, x + t_n + t_l - t_n) = (i, x + t_l)$$

so $n^{-1}ln$ acts as L does on all (i, x) for large enough x, in particular for all but finitely many elements of S. Since all orbits are infinite this means $n^{-1}ln$ acts as L does on all of S. Hence $n^{-1}ln = l$, in particular $n \in N_G L$.

Let A denote a set of representatives of H_n -orbits in $[\mathcal{VCyc} \setminus \mathcal{Fin}]$ such that for all $L \in A$, we have $N_G[L] = N_G L$ and $L \cong \mathbb{Z}$.

LEMMA 7.4.10. For any $K \in A$ we have $N_{H_n}K \cong C_{H_n}K$.

PROOF. Recall that there is a short exact sequence

$$1 \longrightarrow C_{H_n} K \longrightarrow N_{H_n} K \longrightarrow Q \longrightarrow 1$$

where Q is a subgroup of Aut(K) [**Rob96**, 1.6.13].

Let $n \in N_{H_n}K$ and choose some $k \in K$ generating K. Assume that K acts non-trivially on the i^{th} ray, so for x large enough,

$$k: (i, x) \mapsto (i, x + t_k)$$
$$n: (i, x) \mapsto (i, x + t_n)$$

for some $t_k, t_n \in \mathbb{Z}$ with $t_k \neq 0$. Let $a \in \mathbb{Z}$ be such that $n^{-1}kn = k^a$, then

$$n^{-1}kn: (i, x) \mapsto (i, x + t_k)$$

but

$$k^a: (i, x) \mapsto (i, x + at_k).$$

Thus a = 1 and n acts as the trivial automorphism on K, thus Q = 1, proving the lemma.

LEMMA 7.4.11. If $K \in A$ then $\mathcal{O}_{\text{fin}} \operatorname{cd} N_{H_n} K \leq n-1$ and $\mathcal{O}_{\text{fin}} \operatorname{cd} W_{H_n} K \leq n-2$.

PROOF. Recall from Corollary 7.2.5 that

$$C_{H_n}K \cong H_k \times \mathbb{Z}^r \times F,$$

where F is a finite group, $0 \le k \le n-2$, and $1 \le r \le \lfloor (n-k)/2 \rfloor$. Thus,

$$\mathcal{O}_{\mathcal{F}in} \operatorname{cd} N_{H_n} K = \mathcal{O}_{\mathcal{F}in} \operatorname{cd} C_{H_n} K$$

= $\mathcal{O}_{\mathcal{F}in} \operatorname{cd} (H_k \times \mathbb{Z}^r \times F)$
 $\leq \mathcal{O}_{\mathcal{F}in} \operatorname{cd} H_k + \mathcal{O}_{\mathcal{F}in} \operatorname{cd} \mathbb{Z}^r + \mathcal{O}_{\mathcal{F}in} \operatorname{cd} F$
= $k + r$
 $\leq \max_{1 \leq k \leq n-2} \left(k + \lfloor (n-k)/2 \rfloor \right),$

where we've used Proposition 7.4.3 that $\mathcal{O}_{\text{fin}} \operatorname{cd} H_k = k$, Lemma 7.4.10, and Lemma 6.5.1. We claim that $\max_{1 \le k \le n-2} k + \lfloor (n-k)/2 \rfloor = n-1$, indeed we can always achieve n-1 by choosing k = n-2 and $k + \lfloor (n-k)/2 \rfloor$ is an increasing function of k.

Examining the proof of Corollary 7.2.5 that $C_{H_n}K \cong H_k \times \mathbb{Z}^r$, we see that K is a subgroup of \mathbb{Z}^r , so

$$W_G H \cong H_k \times \mathbb{Z}^{r-1} \times F',$$

for some finite subgroup F', which gives the second inequality.

PROOF OF 7.4.4. Via Lemma 7.4.9 we have the long exact sequence of Proposition 7.4.8,

$$\cdots \longrightarrow \prod_{K \in A} H^{i-1}_{\mathcal{O}_{\mathcal{F}_{in}}}(N_{H_n}K, -) \longrightarrow H^i_{\mathcal{O}_{\mathcal{V}_{\mathcal{G}_{in}}}}(H_n, -)$$
$$\longrightarrow \left(\prod_{K \in A} H^i_{\mathcal{O}_{\mathcal{F}_{in}}}(W_{H_n}K, \pi^K_* -)\right) \oplus H^i_{\mathcal{O}_{\mathcal{F}_{in}}}(H_n, -) \longrightarrow \cdots .$$

Let i = n + 1 then using Lemma 7.4.11, the left and right hand terms vanish. Thus the central term vanishes proving $\mathcal{O}_{\mathcal{W}_{\mathcal{W}}} \operatorname{cd} H_n \leq n$.

Let i = n then, using Lemma 7.4.11 again, there is a long exact sequence which terminates as

$$\cdots \longrightarrow H^n_{\mathcal{O}_{\mathcal{V}_{\mathcal{U}}}}(H_n,-) \longrightarrow H^n_{\mathcal{O}_{\mathcal{F}_{in}}}(H_n,-) \longrightarrow 0.$$

Let M be an $\mathcal{O}_{\mathcal{F}in}$ -module such that $H^n_{\mathcal{O}_{\mathcal{F}in}}(H_n, M) \neq 0$ then we may extend M to an $\mathcal{O}_{\mathcal{V}Cyc}$ -module by setting M(G/K) = 0 for all virtually cyclic subgroups K and thus $H^n_{\mathcal{O}_{\mathcal{V}Cyc}}(H_n, M) \neq 0$. In particular, $\mathcal{O}_{\mathcal{V}Cyc} \operatorname{cd} H_n \geq n$. \Box

154

Bibliography

 $[ABJ^+09]$ Goulnara Arzhantseva, Martin R. Bridson, Tadeusz Januszkiewicz, Ian J. Leary, Ashot Minasyan, and Jacek Świątkowski, Infinite groups with fixed point properties, Geom. Topol. 13 (2009), no. 3, 1229-1263. MR 2496045 (2010b:20069) Yago Antolín, José Burillo, and Armando Martino, Conjugacy in Houghton's [ABM13] groups, preprint, arXiv:1305.2044 (2013). [ABS09] Javad Asadollahi, Abdolnaser Bahlekeh, and Shokrollah Salarian, On the hierarchy of cohomological dimensions of groups, J. Pure Appl. Algebra 213 (2009), no. 9, 1795-1803. MR 2518178 (2010i:20062) Luchezar L. Avramov and Alex Martsinkovsky, Absolute, relative, and Tate coho-[AM02] mology of modules of finite Gorenstein dimension, Proc. London Math. Soc. (3) 85 (2002), no. 2, 393-440. MR 1912056 (2003g:16009) [Ant11] Yago Antolín, On Cayley graphs of virtually free groups, Groups Complex. Cryptol. 3 (2011), no. 2, 301-327. MR 2898895 [Bau04] Oliver Baues, Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups, Topology 43 (2004), no. 4, 903-924. MR 2061212 (2005c:57048) [BB97] Mladen Bestvina and Noel Brady, Morse theory and finiteness properties of groups, Invent. Math. 129 (1997), no. 3, 445-470. MR 1465330 (98i:20039) $[BBG^+11]$ Collin Bleak, Hannah Bowman, Alison Gordon, Garrett Graham, Jacob Hughes, Francesco Matucci, and Jenya Sapir, Centralizers in R. Thompson's group V_n , preprint, arXiv:1107.0672 (2011). [BC92] D. J. Benson and Jon F. Carlson, Products in negative cohomology, J. Pure Appl. Algebra 82 (1992), no. 2, 107-129. MR 1182934 (93i:20058) [BCMR14] José Burillo, Sean Cleary, Armando Martino, and Class E. Röver, Commensurators and metric properties of Houghton's groups, preprint, arXiv:1403.0026 (2014). [BDT09] Abdolnaser Bahlekeh, Fotini Dembegioti, and Olympia Talelli, Gorenstein dimension and proper actions, Bull. Lond. Math. Soc. 41 (2009), no. 5, 859-871. MR 2557467 (2010k:20084) [BE73] Robert Bieri and Beno Eckmann, Groups with homological duality generalizing Poincaré duality, Invent. Math. 20 (1973), 103-124. MR 0340449 (49 #5204) , Finiteness properties of duality groups, Comment. Math. Helv. 49 (1974), [BE74] 74-83. MR 0340450 (49 #5205) [Bes93] Mladen Bestvina, The virtual cohomological dimension of Coxeter groups, Geometric group theory, Vol. 1 (Sussex, 1991), London Math. Soc. Lecture Note Ser., vol. 181, Cambridge Univ. Press, Cambridge, 1993, pp. 19–23. MR 1238512 (94g:20056) Kenneth S. Brown and Ross Geoghegan, An infinite-dimensional torsion-free FP_{∞} [BG84] group, Invent. Math. 77 (1984), no. 2, 367-381. MR 752825 (85m:20073) [BH99] Martin R. Bridson and André Haefliger, Metric spaces of non-positive curvature, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles

of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999. MR 1744486 (2000k:53038)

- [Bie72] Robert Bieri, Gruppen mit Poincaré-Dualität, Comment. Math. Helv. 47 (1972), 373–396. MR 0352290 (50 #4777)
- [Bie81] _____, Homological dimension of discrete groups, second ed., Queen Mary College Mathematical Notes, Queen Mary College Department of Pure Mathematics, London, 1981. MR 715779 (84h:20047)
- [BLN01] Noel Brady, Ian J. Leary, and Brita E. A. Nucinkis, On algebraic and geometric dimensions for groups with torsion, J. London Math. Soc. (2) 64 (2001), no. 2, 489–500. MR 1853466 (2002h:57007)
- [BLR08] Arthur Bartels, Wolfgang Lück, and Holger Reich, On the Farrell-Jones conjecture and its applications, J. Topol. 1 (2008), no. 1, 57–86. MR 2365652 (2008m:19001)
- [Bor60] Armand Borel, Seminar on transformation groups, With contributions by G. Bredon, E. E. Floyd, D. Montgomery, R. Palais. Annals of Mathematics Studies, No. 46, Princeton University Press, Princeton, N.J., 1960. MR 0116341 (22 #7129)
- [Bou99] Serge Bouc, Le complexe de chaînes d'un G-complexe simplicial acyclique, J. Algebra 220 (1999), no. 2, 415–436. MR 1717349 (2000k:18009)
- [Bow04] Brian H. Bowditch, Planar groups and the Seifert conjecture, J. Reine Angew. Math. 576 (2004), 11–62. MR 2099199 (2005h:57033)
- [Bre67] Glen E. Bredon, Equivariant cohomology theories, Lecture Notes in Mathematics, No. 34, Springer-Verlag, Berlin-New York, 1967. MR 0214062 (35 #4914)
- [Bre72] _____, Introduction to compact transformation groups, Academic Press, New York, 1972, Pure and Applied Mathematics, Vol. 46. MR 0413144 (54 #1265)
- [Bro87] Kenneth S. Brown, Finiteness properties of groups, Proceedings of the Northwestern conference on cohomology of groups (Evanston, Ill., 1985), vol. 44, 1987, pp. 45– 75. MR 885095 (88m:20110)
- [Bro94] _____, Cohomology of groups, Graduate Texts in Mathematics, vol. 87, Springer-Verlag, New York, 1994, Corrected reprint of the 1982 original. MR 1324339 (96a:20072)
- [BS80] Robert Bieri and Ralph Strebel, Valuations and finitely presented metabelian groups, Proc. London Math. Soc. (3) 41 (1980), no. 3, 439–464. MR 591649 (81j:20080)
- [BW08] Jonathan Block and Shmuel Weinberger, On the generalized Nielsen realization problem, Comment. Math. Helv. 83 (2008), no. 1, 21–33. MR 2365406 (2009a:57037)
- [CK97] Jonathan Cornick and Peter H. Kropholler, On complete resolutions, Topology Appl. 78 (1997), no. 3, 235–250. MR 1454602 (98k:20087)
- [CK98] _____, Homological finiteness conditions for modules over group algebras, J. London Math. Soc. (2) 58 (1998), no. 1, 49–62. MR 1666074 (99k:20105)
- [Dav98] Michael W. Davis, The cohomology of a Coxeter group with group ring coefficients, Duke Math. J. 91 (1998), no. 2, 297–314. MR 1600586 (99b:20067)
- [Dav00] _____, Poincaré duality groups, Surveys on surgery theory, Vol. 1, Ann. of Math. Stud., vol. 145, Princeton Univ. Press, Princeton, NJ, 2000, pp. 167–193. MR 1747535 (2001b:57001)

- [Dav08] _____, The geometry and topology of Coxeter groups, London Mathematical Society Monographs Series, vol. 32, Princeton University Press, Princeton, NJ, 2008. MR 2360474 (2008k:20091)
- [Deg13a] Dieter Degrijse, Bredon cohomological dimensions for proper actions and Mackey functors, preprint, arXiv:1305.1227 (2013).
- [Deg13b] _____, Cohomology of split extensions and classifying spaces for families of subgroups, Ph.D. thesis, 2013.
- [Dek03] Karel Dekimpe, Any virtually polycyclic group admits a NIL-affine crystallographic action, Topology 42 (2003), no. 4, 821–832. MR 1958530 (2004i:20096)
- [DL98] Warren Dicks and Ian J. Leary, On subgroups of Coxeter groups, Geometry and cohomology in group theory (Durham, 1994), London Math. Soc. Lecture Note Ser., vol. 252, Cambridge Univ. Press, Cambridge, 1998, pp. 124–160. MR 1709956 (2000g:20067)
- [DL03] M. W. Davis and I. J. Leary, Some examples of discrete group actions on aspherical manifolds, High-dimensional manifold topology, World Sci. Publ., River Edge, NJ, 2003, pp. 139–150. MR 2048719 (2005f:57050)
- [DMP13] Dieter Degrijse and Conchita Martínez-Pérez, Dimension invariants for groups admitting a cocompact model for proper actions, preprint, arXiv:1311.6419 (2013).
- [DP12] Dieter Degrijse and Nansen Petrosyan, Geometric dimension of groups for the family of virtually cyclic subgroups, preprint, arXiv:1204.3482 (2012).
- [DT08] Fotini Dembegioti and Olympia Talelli, On a relation between certain cohomological invariants, J. Pure Appl. Algebra 212 (2008), no. 6, 1432–1437. MR 2391658 (2009d:20122)
- [DT10] _____, A note on complete resolutions, Proc. Amer. Math. Soc. 138 (2010), no. 11, 3815–3820. MR 2679604 (2011h:20114)
- [Dun79] M. J. Dunwoody, Accessibility and groups of cohomological dimension one, Proc.
 London Math. Soc. (3) 38 (1979), no. 2, 193–215. MR 531159 (80i:20024)
- [Eck87] Beno Eckmann, Poincaré duality groups of dimension two are surface groups, Combinatorial group theory and topology (Alta, Utah, 1984), Ann. of Math. Stud., vol. 111, Princeton Univ. Press, Princeton, NJ, 1987, pp. 35–51. MR 895608 (88e:57001)
- [EG57] Samuel Eilenberg and Tudor Ganea, On the Lusternik-Schnirelmann category of abstract groups, Ann. of Math. (2) 65 (1957), 517–518. MR 0085510 (19,52d)
- [EJ95] Edgar E. Enochs and Overtoun M. G. Jenda, Gorenstein injective and projective modules, Math. Z. 220 (1995), no. 4, 611–633. MR 1363858 (97c:16011)
- [EJ11] _____, Relative homological algebra. Volume 1, extended ed., de Gruyter Expositions in Mathematics, vol. 30, Walter de Gruyter GmbH & Co. KG, Berlin, 2011. MR 2857612 (2012g:16029)
- [EM65] Samuel Eilenberg and J. C. Moore, Foundations of relative homological algebra, Mem. Amer. Math. Soc. No. 55 (1965), 39. MR 0178036 (31 #2294)
- [Emm10] Ioannis Emmanouil, On certain cohomological invariants of groups, Adv. Math.
 225 (2010), no. 6, 3446–3462. MR 2729012 (2011m:20122)
- [Fis68] Janet L. Fisher, The tensor product of functors; satellites; and derived functors, J.
 Algebra 8 (1968), 277–294. MR 0237593 (38 #5874)
- [FKS72] J. Fischer, A. Karrass, and D. Solitar, On one-relator groups having elements of finite order, Proc. Amer. Math. Soc. 33 (1972), 297–301. MR 0311780 (47 #342)

158	BIBLIOGRAPHY
[FL04]	F. T. Farrell and JF. Lafont, <i>Finite automorphisms of negatively curved Poincaré duality groups</i> , Geom. Funct. Anal. 14 (2004), no. 2, 283–294. MR 2060195 (2005e:20062)
[Flu10]	Martin Fluch, On Bredon (co-)homological dimensions of groups, Ph.D. thesis, University of Southampton, UK, September 2010.
[Fow12]	Jim Fowler, <i>Finiteness properties for some rational Poincaré duality groups</i> , Illinois J. Math. 56 (2012), no. 2, 281–299. MR 3161324
[FW08]	Benson Farb and Shmuel Weinberger, <i>Isometries, rigidity and universal covers</i> , Ann. of Math. (2) 168 (2008), no. 3, 915–940. MR 2456886 (2009k:53094)
[Gan12a]	Giovanni Gandini, Bounding the homological finiteness length, Bull. Lond. Math. Soc. 44 (2012), no. 6, 1209–1214. MR 3007653
[Gan12b]	, Cohomological invariants and the classifying space for proper actions, Groups Geom. Dvn. 6 (2012), no. 4, 659–675. MR 2996406
[Geo08]	Ross Geoghegan, <i>Topological methods in group theory</i> , Graduate Texts in Mathematics, vol. 243, Springer, New York, 2008, MR 2365352 (2008):57002)
[GG87]	T. V. Gedrich and K. W. Gruenberg, <i>Complete cohomological functors on groups</i> , Topology Appl. 25 (1987), no. 2, 203–223, Singapore topology conference (Singapore, 1985). MR 884544 (89h:20073)
[GN12]	Giovanni Gandini and Brita Nucinkis, Some $\mathbf{H}_1\mathcal{F}$ -groups with unbounded torsion and a conjecture of Kropholler and Mislin, preprint, to appear Forum Mathe- maticum (2012).
[Goi92]	François Goichot, <i>Homologie de Tate-Vogel équivariante</i> , J. Pure Appl. Algebra 82 (1992), no. 1, 39–64. MR 1181092 (94d:55014)
[Gre71]	J. A. Green, Axiomatic representation theory for finite groups, J. Pure Appl. Al- gebra 1 (1971), no. 1, 41–77, MR 0279208 (43 #4931)
[Gui08]	Guido's book of conjectures, Monographies de L'Enseignement Mathématique [Monographs of L'Enseignement Mathématique], vol. 40, L'Enseignement Mathématique, Geneva, 2008, A gift to Guido Mislin on the occasion of his re- tirement from ETHZ June 2006, Collected by Indira Chatterji. MR 2499538 (2010d:00003)
[Ham 08]	Martin Hamilton, Finiteness conditions in group cohomology, Ph.D. thesis, 2008.
[Ham11]	, When is group cohomology finitary?, J. Algebra 330 (2011), 1–21. MR 2774614 (2012e:20122)
[Hat02]	Allen Hatcher, <i>Algebraic topology</i> , Cambridge University Press, Cambridge, 2002. MR 1867354 (2002k:55001)
[Hil91]	Jonathan A. Hillman, Elementary amenable groups and 4-manifolds with Eu- ler characteristic 0, J. Austral. Math. Soc. Ser. A 50 (1991), no. 1, 160–170. MR 1094067 (92g:20057)
[Hol04]	Henrik Holm, <i>Gorenstein homological dimensions</i> , J. Pure Appl. Algebra 189 (2004), no. 1-3, 167–193. MR 2038564 (2004k:16013)
[Hou79]	C. H. Houghton, The first cohomology of a group with permutation module coefficients, Arch. Math. (Basel) 31 (1978/79), no. 3, 254–258. MR 521478 (80c:20073)
[HPY13]	Ian Hambleton, Semra Pamuk, and Ergün Yalçın, <i>Equivariant CW-complexes and the orbit category</i> , Comment. Math. Hely. 88 (2013). no. 2, 369–425. MR 3048191
[Hur36]	Witold Hurewicz, <i>Beiträge zur Topologie der Deformationen. IV. Asphärische Räume</i> , Nederl. Akad. Wetensch. Proc. 39 (1936), 215–224.

- [Ike84] Bruce M. Ikenaga, Homological dimension and Farrell cohomology, J. Algebra 87 (1984), no. 2, 422–457. MR 739945 (85k:20152)
- [III78] Sören Illman, Smooth equivariant triangulations of G-manifolds for G a finite group, Math. Ann. 233 (1978), no. 3, 199–220. MR 0500993 (58 #18474)
- [Joh99] D. L. Johnson, Embedding some recursively presented groups, Groups St. Andrews
 1997 in Bath, II, London Math. Soc. Lecture Note Ser., vol. 261, Cambridge Univ.
 Press, Cambridge, 1999, pp. 410–416. MR 1676637 (2000h:20057)
- [JPL06] Daniel Juan-Pineda and Ian J. Leary, On classifying spaces for the family of virtually cyclic subgroups, Recent developments in algebraic topology, Contemp. Math., vol. 407, Amer. Math. Soc., Providence, RI, 2006, pp. 135–145. MR 2248975 (2007d:19001)
- [JW72] F. E. A. Johnson and C. T. C. Wall, On groups satisfying Poincaré duality, Ann. of Math. (2) 96 (1972), 592–598. MR 0311796 (47 #358)
- [KM98] Peter H. Kropholler and Guido Mislin, Groups acting on finite-dimensional spaces with finite stabilizers, Comment. Math. Helv. 73 (1998), no. 1, 122–136. MR 1610595 (99f:20086)
- [KMPN09] P. H. Kropholler, C. Martínez-Pérez, and B. E. A. Nucinkis, Cohomological finiteness conditions for elementary amenable groups, J. Reine Angew. Math. 637 (2009), 49–62. MR 2599081 (2011e:20074)
- [KMPN11a] Dessislava H. Kochloukova, Conchita Martínez-Pérez, and Brita E. A. Nucinkis, Centralisers of finite subgroups in soluble groups of type FP_n, Forum Math. 23 (2011), no. 1, 99–115. MR 2769866 (2012b:20125)
- [KMPN11b] _____, Cohomological finiteness conditions in Bredon cohomology, Bull. Lond. Math. Soc. 43 (2011), no. 1, 124–136. MR 2765556 (2012b:20124)
- [Kna02] Anthony W. Knapp, Lie groups beyond an introduction, second ed., Progress in Mathematics, vol. 140, Birkhäuser Boston Inc., Boston, MA, 2002. MR 1920389 (2003c:22001)
- [Kro86] Peter H. Kropholler, Cohomological dimension of soluble groups, J. Pure Appl. Algebra 43 (1986), no. 3, 281–287. MR 868988 (88h:20063)
- [Kro93] _____, On groups of type $(FP)_{\infty}$, J. Pure Appl. Algebra **90** (1993), no. 1, 55–67. MR 1246274 (94j:20051b)
- [KT91] P. H. Kropholler and Olympia Talelli, On a property of fundamental groups of graphs of finite groups, J. Pure Appl. Algebra 74 (1991), no. 1, 57–59. MR 1129129 (92h:57003)
- [KW11] Peter H. Kropholler and Charles T. C. Wall, Group actions on algebraic cell complexes, Publ. Mat. 55 (2011), no. 1, 3–18. MR 2779573 (2012e:20121)
- [Lam99] T. Y. Lam, Lectures on modules and rings, Graduate Texts in Mathematics, vol. 189, Springer-Verlag, New York, 1999. MR 1653294 (99i:16001)
- [Lea05] Ian J. Leary, On finite subgroups of groups of type VF, Geom. Topol. 9 (2005), 1953–1976 (electronic). MR 2175161 (2006j:20058)
- [Lee12] Sang Rae Lee, *Geometry of Houghton's groups*, preprint, arXiv:1212.0257 (2012).
- [Lin76] Harald Lindner, A remark on Mackey-functors, Manuscripta Math. 18 (1976), no. 3, 273–278. MR 0401864 (53 #5691)
- [Lin83] P. A. Linnell, On accessibility of groups, J. Pure Appl. Algebra 30 (1983), no. 1, 39–46. MR 716233 (85c:20021)

- [LM00] Wolfgang Lück and David Meintrup, On the universal space for group actions with compact isotropy, Geometry and topology: Aarhus (1998), Contemp. Math., vol. 258, Amer. Math. Soc., Providence, RI, 2000, pp. 293–305. MR 1778113 (2001e:55023)
- [LN01] Ian J. Leary and Brita E. A. Nucinkis, Bounding the orders of finite subgroups, Publ. Mat. 45 (2001), no. 1, 259–264. MR 1829588 (2002b:20074)
- [LN03] _____, Some groups of type VF, Invent. Math. 151 (2003), no. 1, 135–165. MR 1943744 (2003k:20086)
- [LN10] _____, On groups acting on contractible spaces with stabilizers of prime-power order, J. Group Theory 13 (2010), no. 5, 769–778. MR 2720203 (2012b:20035)
- [LR04] John C. Lennox and Derek J. S. Robinson, The theory of infinite soluble groups, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, Oxford, 2004. MR 2093872 (2006b:20047)
- [LR05] Wolfgang Lück and Holger Reich, The Baum-Connes and the Farrell-Jones conjectures in K- and L-theory, Handbook of K-theory. Vol. 1, 2, Springer, Berlin, 2005, pp. 703–842. MR 2181833 (2006k:19012)
- [LS01] Roger C. Lyndon and Paul E. Schupp, Combinatorial group theory, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1977 edition. MR 1812024 (2001i:20064)
- [Lüc89] Wolfgang Lück, Transformation groups and algebraic K-theory, Lecture Notes in Mathematics, vol. 1408, Springer-Verlag, Berlin, 1989, Mathematica Gottingensis. MR 1027600 (91g:57036)

[Lüc00] _____, The type of the classifying space for a family of subgroups, J. Pure Appl.
 Algebra 149 (2000), no. 2, 177–203. MR 1757730 (2001i:55018)

- [Lüc05] _____, Survey on classifying spaces for families of subgroups, Infinite groups: geometric, combinatorial and dynamical aspects, Progr. Math., vol. 248, Birkhäuser, Basel, 2005, pp. 269–322. MR 2195456 (2006m:55036)
- [LW12] Wolfgang Lück and Michael Weiermann, On the classifying space of the family of virtually cyclic subgroups, Pure and Applied Mathematics Quaterly (Special Issue: In honour of Thomas Farrell and Lowell E. Jones Part 2) 8 (2012), no. 2, 497–555.
 [Mil56] John Milnor, Construction of universal bundles. I, Ann. of Math. (2) 63 (1956),
- [Win50] 5000 Wintor, Construction of universal bundles. 1, Ann. of Wath. (2) 05 (1950), 272–284. MR 0077122 (17,994b)
- [Mis94] Guido Mislin, Tate cohomology for arbitrary groups via satellites, Topology Appl. 56 (1994), no. 3, 293–300. MR 1269317 (95c:20072)
- [Mis10] _____, Classifying spaces for proper actions of mapping class groups, Münster J.
 Math. 3 (2010), 263–272. MR 2775365 (2012c:55015)
- [ML95] Saunders Mac Lane, Homology, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1975 edition. MR 1344215 (96d:18001)
- [ML98] _____, Categories for the working mathematician, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998. MR 1712872 (2001j:18001)
- [MP02] Conchita Martínez-Pérez, A spectral sequence in Bredon (co)homology, J. Pure Appl. Algebra 176 (2002), no. 2-3, 161–173. MR 1933713 (2003h:20095)
- [MP07] _____, A bound for the Bredon cohomological dimension, J. Group Theory 10 (2007), no. 6, 731–747. MR 2364823 (2008j:20159)
- [MP13a] _____, Euler classes and Bredon cohomology for groups with restricted families of finite subgroups, Math. Z. 275 (2013), no. 3-4, 761–780. MR 3127036

[MP13b] _, Subgroup posets, Bredon cohomology and equivariant Euler characteristics, Trans. Amer. Math. Soc. 365 (2013), no. 8, 4351-4370. MR 3055698 Conchita Martínez-Pérez, Francesco Matucci, and Brita E. A. Nucinkis, Cohomo-[MPMN13] logical finiteness conditions and centralisers in generalisations of Thompson's group V, preprint, arXiv:1309.7858 (2013). [MPN06] Conchita Martínez-Pérez and Brita E. A. Nucinkis, Cohomological dimension of Mackey functors for infinite groups, J. London Math. Soc. (2) 74 (2006), no. 2, 379-396. MR 2269585 (2007h:20053) [MPN10] , Virtually soluble groups of type FP_{∞} , Comment. Math. Helv. 85 (2010), no. 1, 135-150. MR 2563683 (2011e:20075) _____, Bredon cohomological finiteness conditions for generalisations of Thomp-[MPN13] son groups, Groups Geom. Dyn. 7 (2013), no. 4, 931-959. MR 3134031 [MT00] Guido Mislin and Olympia Talelli, On groups which act freely and properly on finite dimensional homotopy spheres, Computational and geometric aspects of modern algebra (Edinburgh, 1998), London Math. Soc. Lecture Note Ser., vol. 275, Cambridge Univ. Press, Cambridge, 2000, pp. 208-228. MR 1776776 (2001i:20110) [Mur06] Daniel Murfet, Abelian categories, unpublished (2006). [MV03] Guido Mislin and Alain Valette, Proper group actions and the Baum-Connes conjecture, Advanced Courses in Mathematics. CRM Barcelona, Birkhäuser Verlag, Basel, 2003. MR 2027168 (2005d:19007) [Nuc99] Brita E. A. Nucinkis, Cohomology relative to a G-set and finiteness conditions, Topology Appl. 92 (1999), no. 2, 153-171. MR 1669819 (2000b:20072) [Nuc00] ____, Is there an easy algebraic characterisation of universal proper G-spaces?, Manuscripta Math. 102 (2000), no. 3, 335-345. MR 1777524 (2001h:20075) [Nuc04] _, On dimensions in Bredon homology, Homology Homotopy Appl. 6 (2004), no. 1, 33-47. MR 2061566 (2005c:20092) M. S. Raghunathan, Torsion in cocompact lattices in coverings of Spin(2, n), Math. [Rag84] Ann. 266 (1984), no. 4, 403-419. MR 735524 (86k:22027) _____, Corrigendum: "Torsion in cocompact lattices in coverings of Spin(2,n)" [Rag95] [Math. Ann. 266 (1984), no. 4, 403-419; MR0735524 (86k:22027)], Math. Ann. **303** (1995), no. 3, 575–578. MR 1355004 (96g:22019) [Rob96] Derek J. S. Robinson, A course in the theory of groups, second ed., Graduate Texts in Mathematics, vol. 80, Springer-Verlag, New York, 1996. MR 1357169 (96f:20001) Joseph J. Rotman, An introduction to the theory of groups, fourth ed., Graduate [Rot95] Texts in Mathematics, vol. 148, Springer-Verlag, New York, 1995. MR 1307623 (95m:20001)[Rot09] _, An introduction to homological algebra, second ed., Universitext, Springer, New York, 2009. MR 2455920 (2009i:18011) [Röv99] Class H. E. W. Röver, Subgroups of finitely presented simple groups, Ph.D. thesis, 1999. [Sch70] H. Schubert, Kategorien II, Springer, Berlin, 1970. [Sch78] Hans Rudolf Schneebeli, On virtual properties and group extensions, Math. Z. 159 (1978), no. 2, 159–167. MR 0473038 (57 #12717) [Seg68]Graeme Segal, Classifying spaces and spectral sequences, Inst. Hautes Études Sci.

Publ. Math. (1968), no. 34, 105-112. MR 0232393 (38 #718)

162	BIBLIOGRAPHY
[Ser03]	Jean-Pierre Serre, <i>Trees</i> , Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003, Translated from the French original by John Stillwell, Corrected 2nd printing of the 1980 English translation, MR 1954121 (2003m:20032)
[SJG12]	Simon St. John-Green, <i>Centralisers in Houghton's groups</i> , preprint, to appear Proc. Edinburgh Math. Soc. (2012)
[SJG13a]	, Bredon–Poincaré duality groups, preprint, to appear J. Group Theory (2013)
[SJG13b]	(2013). \dots , On the Gorenstein and \mathfrak{F} -cohomological dimensions, preprint, to appear Bull Lond Math. Soc. (2013)
[SJG14]	, Cohomological finiteness conditions for Mackey and cohomological Mackey functors. I. Algebra 411 (2014) 225–258. MR 3210927
[Sta68]	John R. Stallings, On torsion-free groups with infinitely many ends, Annals of Mathematics 88 (1968), no. 2, pp. 312–334
[Swa69]	Richard G. Swan, <i>Groups of cohomological dimension one</i> , J. Algebra 12 (1969), 585–610
[Sym10]	Peter Symonds, <i>The Bredon cohomology of subgroup complexes</i> , Journal of Pure and Applied Algebra 199 (2010), 261 – 208
[Tal07]	Olympia Talelli, On groups of type Φ , Arch. Math. (Basel) 89 (2007), no. 1, 24–32. MR 2322776 (2009c;20094)
[Tam89]	Daisuke Tambara, Homological properties of the endomorphism rings of certain permutation modules, Osaka J. Math. 26 (1989), no. 4, 807–828. MR 1040426 (91a:20060)
[tD87]	Tammo tom Dieck, <i>Transformation groups</i> , de Gruyter Studies in Mathematics, vol 8 Walter de Gruyter & Co. Berlin 1987 MB 889050 (89c:57048)
[TW90]	Jacques Thévenaz and Peter J. Webb, <i>Simple Mackey functors</i> , Proceedings of the Second International Group Theory Conference (Bressanone, 1989), no. 23, 1990, pp. 299–319. MR 1068370 (91g:20011)
[TW95]	, The structure of Mackey functors, Trans. Amer. Math. Soc. 347 (1995), no. 6, 1865–1961. MR 1261590 (95i:20018)
[Wal79]	C. T. C. Wall (ed.), <i>Homological group theory</i> , London Mathematical Society Lecture Note Series, vol. 36, Cambridge, Cambridge University Press, 1979. MR 564417 (80m:20001)
[Web00]	Peter Webb, A guide to Mackey functors, Handbook of algebra, Vol. 2, North-Holland, Amsterdam, 2000, pp. 805–836. MR 1759612 (2001f:20031)
[Wei94]	Charles A. Weibel, <i>An introduction to homological algebra</i> , Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR 1269324 (95f:18001)
[Wis12]	Daniel T. Wise, From riches to raags: 3-manifolds, right-angled Artin groups, and cubical geometry, CBMS Regional Conference Series in Mathematics, vol. 117, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2012. MR 2986461
[Yos83]	Tomoyuki Yoshida, On G-functors. II. Hecke operators and G-functors, J. Math. Soc. Japan 35 (1983), no. 1, 179–190. MR 679083 (84b:20010)
[Zha11]	Fuzhen Zhang, <i>Matrix theory</i> , second ed., Universitext, Springer, New York, 2011, Basic results and techniques. MR 2857760 (2012h:15001)

Index

 ϕ , group homomorphism $H_n \longrightarrow \mathbb{Z}^{n-1}$, 137 Avramov–Martsinkovsky long exact sequence, 96 Bieri–Eckmann criterion, 29 Bredon (co)homology $H^*_{\mathcal{O}_{\mathcal{F}}}(G, -)$ and $H_*^{\mathcal{O}_{\mathcal{F}}}(G, -), 38$ Bredon cohomological dimension $\mathcal{O}_{\mathcal{F}} \operatorname{cd} G$, 38 Bredon duality group, 12 Bredon–Poincaré duality group, 12 Burnside functor B^G , 62 C-module, 18 CAT(0) cubical complex, 121 Cellular chain complex $C_*^{\mathcal{O}_{\mathcal{F}}}$, 37 $\mathfrak{C}\mathrm{FP}_n$ condition, 28 c_g , Conjugation morphism in $\mathcal{M}_{\mathcal{F}}$, 59 Cohomological dimension $\operatorname{cd} G$, 1 Cohomological Mackey (co)homology $H^*_{\mathcal{H}_{\mathcal{F}}}(G,-)$ and $H^{\mathcal{H}_{\mathcal{F}}}_*(G,-), 67$ Cohomological Mackey cohomological dimension $\mathcal{H}_{\mathcal{F}}$ cd G, 67 Coinducton CoInd, 24 Complete cohomology $\widehat{H}^*(G, -), 94$ Complete Ext functor $\widehat{\operatorname{Ext}}_{RG}^*(M, -)$, 94 Constant $\mathcal{O}_{\mathcal{F}}$ -module R, 3Davis complex \mathcal{U} , 45, 121 d_H , integer such that $H^{d_H}(WH, R[WH]) \neq 0, \ 12$

 $\operatorname{Ext}_{\mathfrak{C}}^*, 26$

Dual module M^D , 47

Duality group, 11

 \mathcal{F} -cohomological dimension \mathcal{F} cd, 6

 \mathcal{F} -cohomology $\mathcal{F}H^*(G, -), 6$ \mathfrak{F} -complete cohomology $\widehat{\mathfrak{F}H}^*(G,-), 96$ \mathfrak{F} -complete Ext functor $\widehat{\mathfrak{FExt}}_{RG}^*(M, -), 96$ \mathfrak{F} -complete resolution, 95 \mathcal{F} -Ext functor \mathcal{F} Ext $^*_{BG}(M, -), 5$ \mathcal{F} -good, 79 \mathcal{F} -homology $\mathcal{F}H_*(G, -), 6$ \mathcal{F} -projective, 5 \mathcal{F} -split, 5 \mathfrak{F} -strong resolution, 96 \mathcal{F} -Tor functor \mathcal{F} Tor $^{RG}_*(M, -), 5$ Family of subgroups \mathcal{F} , 2 $\mathcal{F}FP_n$ condition, 6 \mathfrak{F}_G cohomological dimension \mathfrak{F}_G cd G, 98 \mathfrak{F}_G cohomology $\mathfrak{F}_G H^*(G, -), 98$ \mathfrak{F}_G Ext functor \mathfrak{F}_G Ext $^*_{RG}(M, -)$, 98 \mathfrak{F}_G projective dimension \mathfrak{F}_G pd, 98 \mathfrak{F}_G -proper chain complex, 97 \mathfrak{F}_G -projective, 97 Fin, family of all finite subgroups, 2 Fixed point functor M^- , 63 Fixed quotient functor M_{-} , 63 Free \mathfrak{C} -module $R[-, x]_{\mathfrak{C}}$, 18

G-CW-complex, 37 G-proper chain complex, 96 Geometric dimension gd G, 1 Gorenstein cohomological dimension Gcd, 9 Gorenstein cohomology $GH^*(G, -)$, 96 Gorenstein Ext functor $GExt^*_{RG}(M, -)$, 96 Gorenstein projective, 9 Gorenstein projective dimension Gpd, 9

H-good, 79 Hecke category $\mathcal{H}_{\mathcal{F}}$, 64 $\mathcal{H}_{\mathcal{F}} FP_n$ condition, 68

INDEX

164 hG, Hirsch length, 118 H_n , Houghton's group, 137 Home, 18 I_K^H , Induction morphism in $\mathcal{M}_{\mathcal{F}}$, 59 Induction Ind_{ι} , 24 Kropholler's class of hierarchically decomposable groups H3, 6 l(G), length of a group, 6 Mackey (co)homology $H^*_{\mathcal{M}_{\mathcal{F}}}(G, -)$ and $H^{\mathcal{M}_{\mathcal{F}}}_{*}(G,-), 62$ Mackey category $\mathcal{M}_{\mathcal{F}}$, 56 Mackey cohomological dimension $\mathcal{M}_{\mathcal{F}}$ cd, 62 $\mathcal{M}_{\mathcal{F}} \mathrm{FP}_n$ condition, 62 Model for BG, 1 Model for EG, 1 Model for $E_{\mathcal{F}}G$, 2 n_G , minimal dimension of a proper contractible G-CW complex, 5 $\mathcal{O}_{\mathcal{F}} \mathrm{FP}_n$ condition, 40 Orbit category $\mathcal{O}_{\mathcal{F}}$, 20 Poincaré duality group, 11 Projective dimension Cpd, 28 Property (P_H) , 79 Property (A), 17 Property (EI), 17 Quasi- $\mathcal{O}_{\mathcal{F}} FP_n$ condition, 41 R_K^H , Restriction morphism in \mathcal{M}_F , 59 R-acyclic space, 115 R-homology manifold, 114 R-orientable space, 115 R-torsion, 34 Reflection group trick, 120 Restriction $\operatorname{Res}_{\iota}$, 24 Strong complete resolution, 94 Surface group, 126 Tensor product over a category, $\otimes_{\mathfrak{C}}$, 21 $\operatorname{Tor}_*^{\mathfrak{C}}, 26$

Triv, family of only the trivial subgroup, 2 \mathcal{VCyc} , family of all virtually cyclic subgroups, 2 $\mathcal{V}(G), 13$ Weak complete resolution, 94 Weyl group WH, 31 Yoneda-type lemma, 19