

Covariant description of isothermic surfaces

J. Tafel

Institute of Theoretical Physics, University of Warsaw,
Hoza 69, 00-681 Warsaw, Poland, email: tafel@fuw.edu.pl

Abstract

We present a covariant formulation of the Gauss-Weingarten equations and the Gauss-Mainardi-Codazzi equations for surfaces in 3-dimensional curved spaces. We derive a coordinate invariant condition on the first and second fundamental form which is necessary and sufficient for the surface to be isothermic.

Keywords: isothermic surfaces, the Gauss-Weingarten equations

MCS: 53C42, 53A35

1 Introduction

Studies of isothermic surfaces were originated in 1837 by Lamé [1] who considered surfaces of constant temperature in a solid body. Bertrand [2] is an author of the present definition of isothermic surfaces as surfaces admitting coordinates in which the first fundamental form is conformally flat and the second one is diagonal. Their basic properties were found by Bour [3], Darboux [4], Calapso [5] and Bianchi [6] (see [7] for a review). A transformation given by Darboux and developed by Bianchi allowed to obtain a family of new isothermic surfaces from a given one. This property was crucial for an interpretation of isothermic surfaces as soliton surfaces by Cieśliński, Goldstein and Sym [5]. They recognized the Gauss-Mainardi-Codazzi equations for the isothermic surfaces as a completely integrable system. These results initiated a new interest in isothermic surfaces (see e.g. [9]).

Most studies on isothermic surfaces refer to surfaces in a flat or conformally flat 3-dimensional space. Moreover, their definition is coordinate dependent. The main aim of this paper is to replace this definition by a coordinate invariant condition which is also valid in a general curved 3-dimensional space. We begin with a covariant formulation of the generalized Gauss-Weingarten (GW) equations and their integrability conditions known as the Gauss-Mainardi-Codazzi (GMC) equations. Unfortunately, the latter equations are not a good tool to study isothermic surfaces in spaces which are not conformally flat. In section 3 we analyze coordinate transformations on surfaces and we find a covariant condition which assures the existence of special coordinates assumed in the definition of isothermic surfaces.

2 The Gauss-Weingarten equations

Let S with coordinates x^a , where $a \in \{1, 2\}$, be a 2-dimensional surface in a 3-dimensional manifold M with metric \tilde{g} and coordinates v^i , $i \in \{1, 2, 3\}$. The first and the second fundamental form on S are given by the induced metric g_{ab} and the exterior curvature tensor K_{ab} . Derivatives of v^i over x^a on S define the matrix

$$v^i_{,a} = v^i_{,a} \quad (1)$$

which represents the operator of a pullback of forms on M to forms on S . The induced metric on S is given by

$$g_{ab} = \tilde{g}_{ij} v^i_{,a} v^j_{,b} . \quad (2)$$

Vectors $v_a = v^i_{,a} \partial_i$ are tangent to S . Together with the unit normal vector $n = n^i \partial_i$ they compose a basis such that

$$n_i v^i_{,a} = 0 , \quad n_i n^i = 1 . \quad (3)$$

The field $v^i_{,a}$ transforms as a vector under transformations of coordinates v^i and as a covector under a change of x^a . Thus, it makes sense to define its covariant derivative on S as

$$Dv^i_{,a} = dv^i_{,a} + \tilde{\omega}^i_{,j} v^j_{,a} - \omega^b_{,a} v^i_{,b} , \quad (4)$$

where $\tilde{\omega}^i_{,j}$ and $\omega^a_{,b}$ are the Levi-Civita connections related, respectively, to metric \tilde{g} and g . The tensorial 1-form $Dv^i_{,a}$ transforms as $v^i_{,a}$ under a change of coordinates. Its decomposition in the basis dx^a yields

$$Dv^i_{,a} = v^i_{,ab} dx^b , \quad (5)$$

where

$$v^i_{,ab} = v^i_{,a,b} + \tilde{\Gamma}^i_{,jk} v^j_{,a} v^k_{,b} - \Gamma^c_{,ab} v^i_{,c} \quad (6)$$

and $\tilde{\Gamma}^i_{,jk}$ and $\Gamma^c_{,ab}$ are, respectively, the Christoffel symbols of \tilde{g} and g . Note that $v^i_{,ab}$ is symmetric in the lower indices if (1) is satisfied.

For any pair (a, b) we can decompose vector $v^i_{,ab} \partial_i$ in the basis $\{v_a, n\}$

$$v^i_{,ab} = n^i f_{ab} + v^i_{,d} f^d_{,ab} . \quad (7)$$

Unknown coefficients f_{ab} and $f^c_{,ab}$ are symmetric in indices a, b . The contraction of (7) with n_i yields $f_{ab} = -K_{ab}$, where K_{ab} are components of the exterior curvature (the second fundamental form) of S given by

$$K = K_{ab} dx^a dx^b = Dn_i dv^i = (n_{i,j} - \tilde{\Gamma}^k_{,ij} n_k) v^j_{,a} v^i_{,b} dx^a dx^b . \quad (8)$$

A contraction of (7) with v_{ic} shows that $f_{abc} = -f_{bac}$. This together with $f_{abc} = f_{acb}$ yields $f_{abc} = 0$. Hence,

$$Dv_a^i = -n^i K_{ab} dx^b. \quad (9)$$

Equation (9) is a part of the Gauss-Weingarten equations [9]. In order to find the remaining part we calculate the covariant derivative of n^i written in the form

$$n^i = \frac{1}{2} \tilde{\eta}^i{}_{jk} v_a^j v_b^k \eta^{ab}. \quad (10)$$

Here $\tilde{\eta}^{ijk}$ and η^{ab} are the Levi-Civita completely antisymmetric tensors corresponding to \tilde{g} and g , respectively. Taking into account (9) and $D\eta^{ijk} = D\eta^{ab} = 0$ one obtains

$$Dn^i = v_a^i K^a{}_b dx^b. \quad (11)$$

Equations (9) and (11) generalize the GW equations.

Assume that a 3-dimensional metric \tilde{g}_{ij} on M and tensors g_{ab} and K_{ab} on a 2-dimensional manifold S are given. If equations (1), (9) and (11), together with constraints (2) and (3), can be solved with respect to v^i , v_a^i and n^i then S can be immersed in M in such a way that g_{ab} and K_{ab} are, respectively, the first and the second fundamental form of S . Let us consider this overdetermined system of equations. Covariant derivatives of constraints (2) and (3) vanish thanks to the other equations. Thus, it is sufficient to assume them in a single point. Integrability of (1) is assured by the symmetry of v_{ab}^i in the lower indices. Integrability conditions of (9) and (11) yield the generalized GMC equations. They can be written in the following form

$$\tilde{G}_{ij} n^i n^j = \frac{1}{2} (H^2 - K_{ab} K^{ab} - R) \quad (12)$$

$$\tilde{G}_{ij} n^i v_a^j = (H \delta_a^b - K^b{}_a) |_{|b}. \quad (13)$$

Here \tilde{G}_{ij} is the Einstein tensor of \tilde{g} , R is the Ricci scalar of g , $H = K^a{}_a$ and $|_b$ denotes the covariant derivative corresponding to metric g . These equations can be obtained directly from (9) and (11), but they can be easier deduced from the Arnowitt-Deser-Misner decomposition [10] of the Einstein tensor of \tilde{g} .

For general \tilde{g} equations (12) and (13) are extra constraints for n^i and v_a^i and they cannot be easily solved. They reduce to equations for g and K if \tilde{g} satisfies the Einstein equations with a cosmological constant $\tilde{G}_{ij} = \Lambda \tilde{g}_{ij}$ [11]. However, then the Cotton-York tensor of \tilde{g} vanishes and \tilde{g} must be conformally flat. In this case one can transform (12) and (13) to the standard GMC equations in a flat space.

3 Isothermic surfaces

The conformal transformation

$$\tilde{g} \rightarrow \Omega^2 \tilde{g} \quad (14)$$

implies

$$g \rightarrow \Omega^2 g, \quad K_{ab} \rightarrow \Omega K_{ab} + n(\Omega) g_{ab}. \quad (15)$$

Fundamental properties of isothermic surfaces (conformally flat g and diagonal K in some coordinates x'^a) are preserved by these transformations. If \tilde{g} is conformally flat then in order to find isothermic surfaces one can first look for solutions of the flat GMC equations written in these special coordinates. In particular, one can reduce the problem to the Calapso equation [12]. This approach fails if \tilde{g} is not conformally flat. For this reason, in this section we will characterize isothermic surfaces independly of the GW and GMC equations. Our aim is to find a coordinate invariant condition which is necessary and sufficient for the existence of isothermic coordinates. An interpretation of K is not relevant for this condition.

Theorem 3.1. *Let g be a metric and K be a symmetric tensor on a 2-dimensional manifold. There exist coordinates such that g is conformally flat and K is diagonal iff*

$$\eta^{ab}(u^c u_{a|c} - u^c_{|c} u_a)_{|b} = 0 \quad (16)$$

or

$$(k^{ab} k_{bc|d} \eta^{cd})_{|a} = 0, \quad (17)$$

where u^a is an unit eigenvector of K^a_b ,

$$k_{ab} = \alpha(K_{ab} - \frac{1}{2} H g_{ab}) \quad (18)$$

and α is a normalization factor such that $k_{ab} k^{ab} = 1$.

Proof. Let x'^a be the isothermic coordinates in which $g'_{12} = 0$, $g'_{11} = g'_{22}$ and $K'_{12} = 0$. A transformation from general coordinates x^a to x'^a has to satisfy the conditions

$$L^a_1 g_{ab} L^b_2 = 0 \quad (19)$$

$$L^a_1 g_{ab} L^b_1 = L^a_2 g_{ab} L^b_2 \quad (20)$$

$$L^a_1 K_{ab} L^b_2 = 0, \quad (21)$$

where $L^a_b = \frac{\partial x'^a}{\partial x^b}$. Let β be an eigenvalue of $K^a_b = g^{ac} K_{cb}$. It must be real since any 2-dimensional metric g_{ab} admits conformally flat coordinates and in these coordinates matrix K^a_b is symmetric. Hence, K_{ab} can be written in the form

$$K_{ab} = \alpha u_a u_b + \beta g_{ab}, \quad (22)$$

where $u^a u_a = 1$. If $\alpha = 0$ then conformally flat coordinates of g_{ab} are also isothermic since g_{ab} and K_{ab} are proportional. In this case any vector is an eigenvector of K^a_b . Among them we can certainly find one which satisfies (16).

A non-trivial case occurs when $\alpha \neq 0$. Then eigenvectors of K^a_b are given by u^a and $\eta^a_b u^b$. By virtue of (22) equation (21) leads to $L^a_1 u_a = 0$ (or $L^a_2 u_a = 0$ but then we can interchange indices 1 and 2). Hence

$$L^a_1 = \gamma \eta^{ab} u_b, \quad (23)$$

where γ is a function. Substituting (23) into (19) yields $L^a_2 = \gamma' u^a$. Equation (20) shows that $\gamma'^2 = \gamma^2$. Thus, equations (19)-(21) are equivalent to (23) and

$$L^a_2 = \pm \gamma u^a. \quad (24)$$

From (23) and (24) we obtain the following expressions for the matrix L^{-1}

$$\frac{\partial x'^1}{\partial x^a} = \lambda \eta_{ab} u^b \quad (25)$$

$$\frac{\partial x'^2}{\partial x^a} = \pm \lambda u_a, \quad (26)$$

where λ is a nonvanishing function. Conditions (25) and (26), considered as equations for coordinates x'^a , are integrable provided

$$(\ln \lambda)_{,a} = (\eta^{cd} u_{c|d}) \eta_{ab} u^b - u^b_{|b} u_a. \quad (27)$$

It is easy to check that equation (27) defines λ iff (16) is satisfied.

In order to express equation (16) directly in terms of K_{ab} let us write it in the form

$$\eta^{ab} (4u^c u_{[a|c]} - (u^c u_a)_{|c})_{|b} = 0. \quad (28)$$

Due to the identity

$$u_{[a|c]} = u^d (u_d u_{[a|c]}) = \frac{1}{2} \eta_{ac} u^d (u_d u_e)_{|f} \eta^{ef} \quad (29)$$

equation (28) can be transformed to

$$2\eta^{ac} (u^b u^d (u_d u_a)_{|c})_{|b} = \eta^{ab} (u^c u_a)_{|cb}. \quad (30)$$

One can change order of derivatives on the rhs of (30) in virtue of the identities

$$\eta^{ab} R_{bc} u^c u_a = 0, \quad \eta^{ab} R^c_{abd} u^d u_c = 0 \quad (31)$$

satisfied by the Riemann tensor of a 2-dimensional metric (note that $R_{abcd} \sim \eta_{ab}\eta_{cd}$ and $R_{ab} \sim g_{ab}$). Hence,

$$2\eta^{ac}(u^b u^d (u_d u_a)_{|c})_{|b} = \eta^{ab}(u^c u_a)_{|bc} \quad (32)$$

and we obtain equation (17) with

$$k_{ab} = \sqrt{2}(u_a u_b - \frac{1}{2}g_{ab}) . \quad (33)$$

It follows from (33) that $k_{ab}k^{ab} = 1$ and k_{ab} is proportional to the traceless part of K_{ab} . Hence, one obtains relation (18) with α given by

$$\alpha = (K_{ab}K^{ab} - \frac{1}{2}H^2)^{-\frac{1}{2}} . \quad (34)$$

□

There exist other formulations of equation (16) or (17). For instance, equation (16) can be written in the form

$$d((*d\omega)^*\omega - *(d^*\omega)\omega) = 0 , \quad (35)$$

where $\omega = u_a dx^a$ and the star denotes the Hodge dual corresponding to metric g_{ab} . Equation (35) is transparently invariant under an interchange of ω and $*\omega$. This transformation corresponds to the replacement of the eigenvector u^a by $\eta^{ab}u_b$.

Since k^a_b is traceless and normalized it satisfies

$$k^a_b k^b_c = -\frac{1}{2}\delta^a_c . \quad (36)$$

Substituting (18) and (36) into equation (17) yields

$$(\alpha^2 \hat{K}^{ab} \hat{K}_{bc|d} \eta^{cd})_{|a} = 0 , \quad (37)$$

where $\hat{K}_{ab} = K_{ab} - \frac{1}{2}H g_{ab}$ is the traceless part of K_{ab} . If we insert $\delta^a_b = \eta^{ac}\eta_{bc}$ after \hat{K}^{ab} in (37) and use the identity $\eta^{ab}\eta_{cd} = \delta^a_c \delta^b_d - \delta^a_d \delta^b_c$ we obtain still another form of condition (17)

$$\hat{K}^{ab} \eta_{bc} (\alpha^2 \hat{K}^{cd})_{|d} = 0 . \quad (38)$$

It is known that surfaces in flat space with a constant mean curvature are isothermic. This property can be easily obtained from (38) since substituting (13) to (38) yields the equation

$$\hat{K}^{ab} \eta_{bc} (\alpha^2 H^{|c} - 2\tilde{G}_{ij} n^i v^{jc})_{|a} = 0 \quad (39)$$

which is satisfied if $\tilde{C}_{ij} = 0$ and $H = \text{const}$.

Equation (38), as well as (16) and (17), is preserved by transformations of coordinates and conformal transformation (14). This is an advantage with respect to the standard description of isothermic surfaces. However, equation (38) is rather complicated. Locally, every surface $S \subset M$ corresponds to a function f of 2 variables, e.g. S is defined by $z = f(x^a)$, where x^a and z are coordinates of M . The induced metric and the normal vector of S depend on f and its first derivatives. Hence, the exterior curvature tensor depends on second derivatives of f and equation (38) is a fourth order nonlinear equation for f .

4 Summary

We presented a covariant formulation of the Gauss-Weingarten and the Gauss-Mainardi-Codazzi equations for surfaces in curved 3-dimensional spaces. We found a covariant condition on the first and second fundamental form of a surface which assures existence of isothermic coordinates. It is given by (16), (17) or (38).

References

- [1] G. Lamé, Mémoire sur les surfaces isothermes dans les corps solides homogènes en équilibre de température, J. Math. Pur. Appl. (Liouville J.) 2 (1837) 147-183
- [2] J. Bertrand, Mémoire sur les surfaces isothermes orthogonales, J. Math. Pur. Appl. (Liouville J.) 9 (1844) 117-130
- [3] E. Bour, Théorie de la déformation des surfaces, J. l'École Imperiale Polytech. 19, Cahier 39 (1862) 1-48
- [4] G. Darboux, Sur les surfaces isothermiques, C. R. Acad. Sci. Paris 128 (1899) 1299-1305
- [5] P. Calapso, Sulla superficie a linee di curvatura isoterme, Rend. Circ. Mat. Palermo 17 (1903) 275-286
- [6] L. Bianchi, Ricerche sulle superficie isoterme e sulle deformazione delle quadriche, Ann. Matem. 11 (1905) 93-157
- [7] P. Klimczewski, M. Nieszporski and A. Sym, Luigi Bianchi, Pasquale Calapso and solitons, Rend. Sem. Mat. Messina, Atti del Congresso Internazionale in onore di Pasquale Calapso, Messina 1998 (2000) 223-240

- [8] J. Cieřliński, P Goldstein and A. Sym, Isothermic surfaces in E^3 as soliton surfaces, *Phys. Lett. A* 205 (1995) 37-43
- [9] U. Hertrich-Jeromin, *Introduction to Möbius Differential Geometry*, Cambridge University Press, 2003
- [10] N. Straumann, *General Relativity (Second Edition)*, Graduate Text in Physics, Springer, 2013
- [11] A. Garcia, F.W. Hehl, C. Heinicke and A. Macias, The Cotton tensor in Riemannian spacetimes, *Class. Quantum Grav.* 21 (2004) 1099-1118
- [12] C. Rogers and W.K. Schief, *Bäcklund and Darboux Transformations. Geometry and Modern Applications in Soliton Theory*, Cambridge Texts in Applied Mathematics, Cambridge University Press, 2002