

# Operator inequalities among arithmetic mean, geometric mean and harmonic mean

Shigeru Furuichi\*

Department of Information Science,  
College of Humanities and Sciences, Nihon University,  
3-25-40, Sakurajyousui, Setagaya-ku, Tokyo, 156-8550, Japan

**Abstract.** We give an upper bound for the weighted geometric mean using the weighted arithmetic mean and the weighted harmonic mean. We also give a lower bound for the weighted geometric mean. These inequalities are proven for two invertible positive operators.

**Keywords :** Operator inequality and operator mean

**2010 Mathematics Subject Classification :** 15A39 and 15A45

## 1 Introduction

Let  $\mathcal{H}$  be a complex Hilbert space. We represent the set of all bounded operators on  $\mathcal{H}$  by  $B(\mathcal{H})$ . If  $A \in B(\mathcal{H})$  satisfies  $A^* = A$ , then  $A$  is called a self-adjoint operator. If a self-adjoint operator  $A$  satisfies  $\langle x|A|x \rangle \geq 0$  for any  $|x \rangle \in \mathcal{H}$ , then  $A$  is called a positive operator. For two self-adjoint operators  $A$  and  $B$ ,  $A \geq B$  means  $A - B \geq 0$ . The notation  $A > 0$  means  $A$  is an invertible positive operator.

It is well-known that we have the following Young inequalities for invertible positive operators  $A$  and  $B$ :

$$(1 - \nu)A + \nu B \geq A\#_{\nu}B \geq \{(1 - \nu)A^{-1} + \nu B^{-1}\}^{-1}, \quad (1)$$

where  $A\#_{\nu}B \equiv A^{1/2}(A^{-1/2}BA^{-1/2})^{\nu}A^{1/2}$  represents the geometric mean for two positive operators  $A$  and  $B$  and a weighted parameter  $\nu \in [0, 1]$  [1]. (In this paper, we use the notation  $A\#B$  instead of  $A\#_{1/2}B$  for the simplicity.)  $(1 - \nu)A + \nu B$  and  $\{(1 - \nu)A^{-1} + \nu B^{-1}\}^{-1}$  are called weighted arithmetic mean and harmonic mean for two positive operators, respectively. The simplified and elegant proof for the inequalities (1) was given in [2]. Recently, refinements of the inequalities (1) were given in our papers [3, 4]. It is also notable that improvements of [4] have been given in the paper [5]. And further improvements have been given in quite recent papers [6] and [7]. In this short note, we consider the relations among operator means for two positive operators.

We start from the following proposition.

**Proposition 1.1** *Let  $A, B$  be invertible positive operators and  $r$  be a real number. Then we have the following inequalities.*

$$(i) \text{ If } r \geq 2, \text{ then } rA\#B + (1 - r)\frac{A+B}{2} \leq \left(\frac{A^{-1}+B^{-1}}{2}\right)^{-1}.$$

---

\*E-mail:furuichi@chs.nihon-u.ac.jp

(ii) If  $r \leq 1$ , then  $rA\#B + (1-r)\frac{A+B}{2} \geq \left(\frac{A^{-1}+B^{-1}}{2}\right)^{-1}$ .

*Proof:* In general, by using the notion of the representing function  $f_m(x) = 1mx$  for operator mean  $m$ , it is well-known [1] that  $f_m(x) \leq f_n(x)$  holds for  $x > 0$  if and only if  $AmB \leq AnB$  holds for all positive operators  $A$  and  $B$ . Thus we can prove this proposition from the following scalar inequalities for  $t > 0$ .

$$(i) \quad r\sqrt{t} + (1-r)\frac{t+1}{2} \leq \frac{2t}{t+1}, \quad (r \geq 2).$$

$$(ii) \quad r\sqrt{t} + (1-r)\frac{t+1}{2} \geq \frac{2t}{t+1}, \quad (r \leq 1).$$

Actually (i) above can be proven in the following way. We set  $f_r(t) \equiv \frac{2t}{t+1} - r\sqrt{t} - (1-r)\frac{t+1}{2}$ , then  $\frac{df_r(t)}{dr} = -\sqrt{t} + \frac{t+1}{2} \geq 0$  implies  $f_r(t) \geq f_2(t)$  for  $r \geq 2$ . From the relation  $\frac{2t}{t+1} + \frac{t+1}{2} \geq 2\sqrt{t}$ , we have  $f_2(t) \geq 0$ . We also give the proof for (ii) above. We set  $g_r(t) \equiv r\sqrt{t} + (1-r)\frac{t+1}{2} - \frac{2t}{t+1}$ , then  $\frac{dg_r(t)}{dr} = \sqrt{t} - \frac{t+1}{2} \leq 0$  implies  $g_r(t) \geq g_1(t)$  for  $r \leq 1$ . From the relation  $\frac{2t}{t+1} \leq \sqrt{t}$ , we have  $g_1(t) \geq 0$ . ■

**Remark 1.2** We have counter-examples of both inequalities (i) and (ii) in Proposition 1.1 for  $1 < r < 2$ . For example, we take  $r = 1.5$ . Then we have the following computations.  $\frac{2t}{t+1} - r\sqrt{t} - (1-r)\frac{t+1}{2} \simeq 0.122302$  when  $t = 0.01$  and  $\frac{2t}{t+1} - r\sqrt{t} - (1-r)\frac{t+1}{2} \simeq -0.037987$  when  $t = 2$ .

## 2 Main results

Proposition 1.1 can be generalized by means of weighted parameter  $\nu \in [0, 1]$ , as the second inequality in (2) below.

**Theorem 2.1** If (i)  $0 \leq \nu \leq 1/2$  and  $0 < A \leq B$  or (ii)  $1/2 \leq \nu \leq 1$  and  $0 < B \leq A$ , then the following inequalities hold

$$A\#B + \left(\nu - \frac{1}{2}\right)(B - A) \leq A\#_\nu B \leq \frac{1}{2}\{(1-\nu)A + \nu B\} + \frac{1}{2}\{(1-\nu)A^{-1} + \nu B^{-1}\}^{-1}. \quad (2)$$

**Remark 2.2** Under the same conditions as in Theorem 2.1, we have  $A\#B \geq A\#_\nu B$ .

In order to prove Theorem 2.1, we firstly prove the corresponding scalar inequalities, as it was similarly done in Proposition 1.1.

**Lemma 2.3** If (i)  $0 \leq \nu \leq 1/2$  and  $t \geq 1$  or (ii)  $1/2 \leq \nu \leq 1$  and  $0 < t \leq 1$ , then the following inequalities hold

$$2\sqrt{t} + (2\nu - 1)(t - 1) \leq 2t^\nu \leq (1 - \nu) + \nu t + \left\{(1 - \nu) + \frac{\nu}{t}\right\}^{-1}. \quad (3)$$

*Proof:* It is trivial for the case  $t = 1$ . For the cases  $\nu = 0, 1/2$  or  $1$ , the inequalities (3) hold. So we assume  $t \neq 1$  and  $\nu \neq 0, 1/2, 1$ . We firstly prove the first inequality of the inequalities (3), under the condition (i)  $0 < \nu < 1/2$  and  $t > 1$  or (ii)  $1/2 < \nu < 1$  and  $0 < t < 1$ . Here we put  $f_\nu(t) \equiv t^\nu - \sqrt{t} - (\nu - \frac{1}{2})(t - 1)$ . Then we have  $f'_\nu(t) = \nu t^{\nu-1} - \frac{1}{2}\frac{1}{\sqrt{t}} - (\nu - \frac{1}{2})$  and  $f'_\nu(1) = 0$ . We also have  $f''_\nu(t) = -\nu(1 - \nu)t^{\nu-2} + \frac{1}{4}t^{-3/2}$ . Thus we have  $f''_\nu(t) = 0 \Leftrightarrow t = t_\nu \equiv \{4\nu(1 - \nu)\}^{\frac{2}{1-2\nu}}$ . We find  $t_\nu < 1$  in the case  $0 < \nu < 1/2$  and  $t > 1$ . Then we find

$f''_\nu(t) \geq 0$  for  $t > 1 (> t_\nu)$ . So  $f'_\nu(t)$  is monotone increasing for  $t > 1$  and we have  $f'_\nu(1) = 0$ . Thus we find  $f'_\nu(t) \geq 0$  for  $t > 1$ . So  $f_\nu(t)$  is monotone increasing for  $t > 1$ . Therefore we have  $f_\nu(t) \geq f_\nu(1) = 0$ . We also find  $t_\nu > 1$  in the case  $1/2 < \nu < 1$  and  $0 < t < 1$ . Then we find  $f''_\nu(t) \geq 0$  for  $0 < t < 1 (< t_\nu)$ . So  $f'_\nu(t)$  is monotone increasing for  $0 < t < 1$  and we have  $f'_\nu(1) = 0$ . Thus we find  $f'_\nu(t) \leq 0$  for  $0 < t < 1$ . So  $f_\nu(t)$  is monotone decreasing for  $0 < t < 1$ . Therefore we have  $f_\nu(t) \geq f_\nu(1) = 0$ . Thus the proof for the first inequality of the inequalities (3) is done.

We prove the second inequality of the inequalities (3). We put  $g_\nu(t) \equiv (1 - \nu) + \nu t + \frac{1}{1 - \nu + \frac{\nu}{t}} - 2t^\nu$ . Then we have  $g_\nu(t) = (1 - \nu) + \nu t + \frac{t}{(1 - \nu)t + \nu} - 2t^\nu \geq 2\sqrt{\frac{\{(1 - \nu) + \nu t\}t}{(1 - \nu)t + \nu}} - 2t^\nu$ . Since  $g_\nu(t) \geq 0$  is equivalent to  $\frac{(1 - \nu) + \nu t}{(1 - \nu)t + \nu} \geq t^{2\nu - 1}$ , we put again  $h_\nu(t) \equiv (1 - \nu) + \nu t - \{(1 - \nu)t + \nu\}t^{2\nu - 1}$ . Then we prove  $h_\nu(t) > 0$  under the condition (i)  $0 < \nu < 1/2$  and  $t > 1$  or (ii)  $1/2 < \nu < 1$  and  $0 < t < 1$ . By the elementary calculations, we have  $h'_\nu(t) = \nu - 2\nu(1 - \nu)t^{2\nu - 1} - \nu(2\nu - 1)t^{2\nu - 2}$ ,  $h'_\nu(1) = 0$  and  $h''_\nu(t) = -2\nu(1 - \nu)(2\nu - 1)t^{2\nu - 3}(t - 1)$ . Then we find  $h''_\nu(t) = 0 \Leftrightarrow t = 1$ . In the case  $t > 1$ , we have  $h''_\nu(t) \geq 0$ . So  $h'_\nu(t)$  is monotone increasing for  $t > 1$  and we have  $h'_\nu(1) = 0$ . Thus we have  $h'_\nu(t) \geq 0$  for  $t > 1$ . So  $h_\nu(t)$  is monotone increasing for  $t > 1$ . Thus we have  $h_\nu(t) \geq h_\nu(1) = 0$ . In the case  $0 < t < 1$ , we also have  $h''_\nu(t) \geq 0$ . So  $h'_\nu(t)$  is monotone increasing for  $0 < t < 1$  and we have  $h'_\nu(1) = 0$ . Thus we have  $h'_\nu(t) \leq 0$  for  $0 < t < 1$ . So  $h_\nu(t)$  is monotone decreasing for  $0 < t < 1$ . Thus we have  $h_\nu(t) \geq h_\nu(1) = 0$ . Thus the proof for the second inequality of the inequalities (3) is done. ■

**Lemma 2.4** *Let  $r \in \mathbb{R}$ . Then the function  $k_{r,\nu}(t) \equiv rt^\nu + (1 - r)\{(1 - \nu) + \nu t\}$ , ( $0 \leq \nu \leq 1, t > 0$ ) is monotone decreasing with respect to  $r$ . Therefore,  $k_{r,\nu}(t) \leq k_{2,\nu}(t)$  for  $r \geq 2$  and  $k_{r,\nu}(t) \geq k_{1,\nu}(t)$  for  $r \leq 1$ .*

*Proof* : The proof is done by  $\frac{\partial k_{r,\nu}(t)}{\partial r} = t^\nu - \{(1 - \nu) + \nu t\} \leq 0$ , for  $\nu \in [0, 1]$  and  $t > 0$ . ■

Lemma 2.4 provides the following results.

**Lemma 2.5** *Let  $r \geq 2$ . If (i)  $0 \leq \nu \leq 1/2$  and  $t \geq 1$  or (ii)  $1/2 \leq \nu \leq 1$  and  $0 < t \leq 1$ , then*

$$rt^\nu + (1 - r)\{(1 - \nu) + \nu t\} \leq \left\{(1 - \nu) + \frac{\nu}{t}\right\}^{-1}.$$

*Proof* : The proof follows directly from Lemma 2.3 and Lemma 2.4. ■

**Lemma 2.6** *Let  $r \leq 1$ . For  $0 < \nu \leq 1$  and  $t > 0$ , we have*

$$rt^\nu + (1 - r)\{(1 - \nu) + \nu t\} \geq \left\{(1 - \nu) + \frac{\nu}{t}\right\}^{-1}.$$

*Proof*: For  $r \leq 1$ , it follows from Lemma 2.4 that  $rt^\nu + (1 - r)\{(1 - \nu) + \nu t\} \geq t^\nu$ . Since we have  $t^\nu \geq \left\{(1 - \nu) + \frac{\nu}{t}\right\}^{-1}$ , the proof is done. ■

Finally we have the following corollary.

**Corollary 2.7** *Let  $r \geq 2$ . If (i)  $0 < \nu \leq 1/2$  and  $0 < A \leq B$  or (ii)  $1/2 \leq \nu \leq 1$  and  $0 < B \leq A$ , then*

$$rA\#_\nu B + (1 - r)\{(1 - \nu)A + \nu B\} \leq \{(1 - \nu)A^{-1} + \nu B^{-1}\}^{-1}.$$

Let  $r \leq 1$ . For  $0 < \nu < 1$  and  $t > 0$ , we have

$$rA\#_{\nu}B + (1-r)\{(1-\nu)A + \nu B\} \geq \{(1-\nu)A^{-1} + \nu B^{-1}\}^{-1}.$$

*Proof:* The proof can be done applying Lemma 2.5, Lemma 2.6 and Theorem 2.1. ■

## Acknowledgement

The author was partially supported by JSPS KAKENHI Grant Number 24540146.

## References

- [1] F.Kubo and T.Ando, Means of positive operators, Math. Ann.,Vol.264(1980),pp.205-224.
- [2] T.Furuta and M.Yanagida, Generalized means and convexity of inversion for positive operators, Amer.Math.Monthly, Vol.105 (1998),pp.258-259.
- [3] S.Furuichi, On refined Young inequalities and reverse inequalities, J. Math. Ineq., Vol.5(2011), pp.21-31.
- [4] S.Furuichi, Refined Young inequalities with Specht's ratio, J. Egypt. Math. Soc., Vol.20(2012), pp.46-49.
- [5] H.Zuo, G.Shi and M.Fujii, Refined Young inequality with Kantorovich constant, J. Math. Ineq., Vol.5(2011), pp.551-556.
- [6] M.Krnić, N.Lovcićević and J.Pečarić, Jensen's operator and applications to mean inequalities for operators in Hilbert space, Bull. Malays. Math. Sci. Soc., Vol.35 (2012), pp.1-14.
- [7] F.Kittanech, M.Krnić, N.Lovcićević and J.Pečarić, Improved arithmetic-geometric and Heinz means inequalities for Hilbert space operators, Publ. Math. Debrecen, Vol.80(2012), pp.465-478.