# Space periodic Jacobi elliptic solution for triad modified Schrödinger equations

Usama Kadri

Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

#### Abstract

We present an analytical solution for triad nonlinear evolution equations with modified Schrödinger terms. An example for application in compressible water waves is presented.

#### 1 Introduction

We consider a two-dimensional problem of an interacting wave triad of the form

$$\Psi_{j,t}(x,t) = i\alpha_j \left[ \Psi_{j,xx}(x,t) + \delta_j^2 \Psi_j(x,t) \right] + \gamma_j V_j(t) \Psi_j(x,t) \qquad j = 1, 2, 3$$
(1)

with  $\alpha_j$ ,  $\delta_j$ , and  $\gamma$  are parameters of the physical problem. We also assume that the relations

$$V_1\Psi_1 = \Psi_2\Psi_3, \quad V_2\Psi_2 = \Psi_1\Psi_3^*, \quad V_3\Psi_3 = \Psi_1\Psi_2^*,$$
 (2)

are satisfied,  $\delta_2 = \delta_3 = \delta_1/2 \equiv \delta/2$ , and asterisks denote complex conjugates. The objective is to derive a periodic analytical solution, for application in long and short wave triad interactions.

Email address: ukadri@mit.edu (Usama Kadri).

## 2 Solution

Redefine  $\Psi_j(x,t) = g_j(x)f_j(t)$ , so that equations (1) can be written as

$$g_1 f_{1,t} = i\alpha_1 \left[ g_{1,xx} + \delta_1^2 g_1 \right] f_1 + \gamma_1 g_2 g_3 f_2 f_3 \tag{3}$$

$$g_2 f_{2,t} = i\alpha_2 \left[ g_{2,xx} + \delta_2^2 g_2 \right] f_2 + \gamma_2 g_1 g_3^* f_1 f_3^* \tag{4}$$

$$g_3 f_{3,t} = i\alpha_3 \left[ g_{3,xx} + \delta_3^2 g_3 \right] f_3 + \gamma_3 g_1 g_2^* f_1 f_2^* \tag{5}$$

Based on (2),  $g_1 = g_2 g_3$ ,  $g_2 = g_1 g_3^*$ , and  $g_3 = g_1 g_2^*$ , so that equations (3), (4), and (5), can be rewritten as

$$f_{1,t} = \gamma_1 f_2 f_3 + i\alpha_1 \left[ g_{1,xx} + \delta_1^2 g_1 \right] f_1 g_1^{-1} \tag{6}$$

$$f_{2,t} = \gamma_2 f_1 f_3^* + i\alpha_2 \left[ g_{2,xx} + \delta_2^2 g_2 \right] f_2 g_2^{-1} \tag{7}$$

$$f_{3,t} = \gamma_3 f_1 f_2^* + i\alpha_3 \left[ g_{3,xx} + \delta_3^2 g_3 \right] f_3 g_3^{-1} \tag{8}$$

Now we can seek a solution in two parts. The first part requires that

$$g_{j,xx} + \delta_j^2 g_j = 0, \qquad j = 1, 2, 3.$$
 (9)

A general solution of (9) is given by

$$g_j = a_j e^{i\delta_j x} + b_j e^{-i\delta_j x}.$$
 (10)

For the second part of the solution we need to solve the following simplified system of three ordinary differential equations that amplitudes satisfy

$$f_{1,t} = \gamma_1 f_2 f_3, \qquad f_{2,t} = \gamma_2 f_1 f_3^*, \qquad f_{3,t} = \gamma_3 f_1 f_2^*$$
(11)

Multiplying (11) by  $f_j^*$  and adding its conjugate multiplied by  $f_j$ , for j=1,2,3 respectively, we obtain the following set of equations

$$f_1^* f_{1,t} + f_1 f_{1,t}^* = \gamma_1 \left( f_1^* f_2 f_3 + f_1 f_2^* f_3^* \right) \tag{12}$$

$$f_2^* f_{2,t} + f_2 f_{2,t}^* = \gamma_2 \left( f_1^* f_2 f_3 + f_1 f_2^* f_3^* \right) \tag{13}$$

$$f_3^* f_{3,t} + f_3 f_{3,t}^* = \gamma_3 \left( f_1^* f_2 f_3 + f_1 f_2^* f_3^* \right) \tag{14}$$

More compactly we can write  $|f_j|_{t}^2 = 2\gamma_j \Im\{f_1^* f_2 f_3\}$ , where the Hamiltonian  $\Im\{f_1^* f_2 f_3\}$  is a constant of the motion (*Holm and Lynch* (2002)). Now define  $\mathbb{Z}_{t} = \Im\{f_1^* f_2 f_3\}$  gives  $|f_j|^2 = 2\gamma_j \mathbb{Z} + \psi_{0j}^2$ . In order to carry on with the

solution the signs of  $\gamma_j$  have to be determined. Note that for a resonating triad,  $\gamma_1 + \gamma_2 + \gamma_3 = 0$ , (Lynch et al. (2003)), thus one has a different sign than the others. Assume, with no loss of generality, that  $\gamma_1$  is negative, that  $\Psi_1(x, t = 0) = \psi_{01} = 0$ , and that  $|\psi_{03}| < |\psi_{02}|$  we obtain

$$\mathbb{Z}_{,t} = \sqrt{-8|\gamma_1|\gamma_2\gamma_3\mathbb{Z}\left(\mathbb{Z} + \frac{\psi_{02}^2}{\gamma_2}\right)\left(\mathbb{Z} + \frac{\psi_{03}^2}{\gamma_3}\right)} \tag{15}$$

This is an elliptic function with a solution given by (see *Byrd and Friedman* (1971), equation 236.00, p.79)

$$\mathbb{Z} = -\frac{\psi_{03}^2}{\gamma_3} \mathrm{sn}^2(u,k)$$
(16)

where  $\operatorname{sn}(u, k)$  is the sine amplitude Jacobian elliptic function of argument u, and modulus k given by

$$u = \sqrt{2|\gamma_1|\gamma_3|}\psi_{02}|t, \qquad k = \frac{|\psi_{03}|}{|\psi_{02}|}\sqrt{\frac{\gamma_2}{\gamma_3}}$$
(17)

and the expression for  $|f_j|$  are

$$|f_j|^2 = |\psi_{0j}|^2 - 2\gamma_j \frac{|\psi_{03}|^2}{\gamma_3} \operatorname{sn}^2(u,k)$$
(18)

Finally, the analytical solution is given by

$$\Psi_1(x,t)|^2 = -2\gamma_1 \frac{|\psi_{03}|^2}{\gamma_3} \operatorname{sn}^2(u,k) \left[ \exp\left(2i\delta x\right) + \exp\left(-2i\delta x\right) \right]$$
(19)

$$|\Psi_2(x,t)|^2 = \left[|\psi_{02}|^2 - 2\gamma_2 \frac{|\psi_{03}|^2}{\gamma_3} \operatorname{sn}^2(u,k)\right] \left[\exp\left(i\delta x\right) + \exp\left(-i\delta x\right)\right]$$
(20)

$$|\Psi_3(x,t)|^2 = |\psi_{03}|^2 \left[1 - 2\mathrm{sn}^2(u,k)\right] \left[\exp\left(i\delta x\right) + \exp\left(-i\delta x\right)\right]$$
(21)

# 3 Application

The solution presented here can be applied in various long-short wave triad interactions, such as Rossby-type waves (see *Pedlosky* (1987); *Charney* (1948)), or wave motion in an inhomogeneous plasma (*Hasegawa and Mima* (1977)). Nevertheless, the following example considers the interaction of two surface gravity waves with an acoustic wave in a mechanism similar to that proposed by *Longuet–Higgins* (1950), and more recently by *Kadri and Stiassnie* (2013) and Kadri (2015).

Given an acoustic wave and two gravity waves with potential amplitudes  $\phi_a$ ,  $\phi_{g1}$ , and  $\phi_{g2}$ , satisfying the following evolution equations

$$\phi_{a,t} = -\frac{ic^2\delta}{2\omega h} \left(\phi_{a,xx} + \frac{4\omega^2}{c^2}\phi_a\right) - \frac{2\omega}{hc}\phi_{g1}\phi_{g2} \tag{22}$$

$$\phi_{g1,t} = \frac{2\omega^3}{qc} \phi_a \phi_{g2}^*; \quad \phi_{g2,t} = \frac{2\omega^3}{qc} \phi_a \phi_{g1}^* \tag{23}$$

where c = 1500 m/s, is the speed of sound in water,  $\omega$  is the frequency of the gravity waves, h is the water depth. The solution of evolution equations is then given by

$$|\Phi_a(x,t)|^2 = \frac{2g|\phi_{0(g2)}|^2}{\omega h} \operatorname{sn}^2(u,k) \left(e^{4i\omega x/c} + e^{-4i\omega x/c}\right)$$
(24)

$$|\Phi_{g1}(x,t)|^2 = \left[ |\phi_{0(g1)}|^2 - 2|\phi_{0(g2)}|^2 \operatorname{sn}^2(u,k) \right] \left( e^{2i\omega x/c} + e^{-2i\omega x/c} \right)$$
(25)

$$|\Phi_{g2}(x,t)|^2 = |\phi_{0(g2)}|^2 \left[1 - 2\mathrm{sn}^2(u,k)\right] \left(e^{2i\omega x/c} + e^{-2i\omega x/c}\right)$$
(26)

with  $u = 2\sqrt{2/gh}\omega^2/c$ , and  $k = |\phi_{0(g2)}|/|\phi_{0(g1)}|$ .

### References

- Byrd, P. F. and Friedman M. D. (1971), Handbook of elliptic integrals for engineers and scientists, *Springer-Verlag*, pp. 358.
- Charney, J.G. (1948) On the Scale of Atmospheric Motions. *Geofys. Publ.*, 17, No. 2, 17pp.
- Hasegawa, A. and Mima, K. (1977) Pseudo-three-dimensional turbulence in magnetized nonuniform plasmas. *Phys. Fluids*, 21(1), 8792.
- Holm, D.D. and Lynch, P., (2002) Stepwise Precession of the Resonant Swinging Spring. SIAM J. Appl. Dy- nam. Systems, 1, 4464.
- Kadri, U. (2015) Wave motion in a heavy compressible fluid: revisited, European Journal of Mechanics -B/Fluids, 49(A), 50–57, 10.1016/j.euromechflu.2014.07.008.
- Kadri, U. and Stiassnie, M. (2013) Generation of an acoustic-gravity wave by two gravity waves, and their mutual interaction. J. Fluid Mech., 735 R6, doi:10.1017/jfm.2013.539.
- Longuet-Higgins, M. S. (1950) A theory of the origin of microseisms, *Philos. Trans. R. Soc. London, Ser. A*, 243, 1–35, doi:10.1098/rsta.1950.0012.
- Pedlosky, J. (1987) Geophysical Fluid Dynamics. Second edition, Springer, New York. 710pp.
- Lynch, P., Eireann, M., Hill, H. (2003) Resonant Rossby Wave Triads and the Swinging Spring, Bull. Amer. Met. Soc., DOI: 10.1175/BAMS-84-5-605.