

# Conditional Preference Orders and their Numerical Representations

Samuel Drapeau<sup>a,1</sup>, Asgar Jamneshan<sup>b,2</sup>

April 27, 2022

---

## ABSTRACT

This work provides an axiomatic framework to the concept of conditional preference orders based on conditional sets. Conditional numerical representations of such preference orders are introduced and a conditional version of the theorems of Debreu about the existence of such numerical representations is given. The continuous representations follow from a conditional version of Debreu's Gap Lemma the proof of which is free of any measurable selection arguments but is derived from the existence of a conditional axiom of choice. As an example, a conditional version of the classical von Neumann and Morgenstern representation is provided.

KEYWORDS: Conditional Preferences, Utility Theory, Gap Lemma, von Neumann and Morgenstern

---

## AUTHORS INFO

<sup>a</sup> SAIF/(CAFR) and Mathematics Departement, Shanghai Jiao Tong University, 211 Huaihai Road, Shanghai, China 200030

<sup>b</sup> Konstanz University, Universitätsstraße 10, 78464 Konstanz, Germany

<sup>1</sup> [sdrapeau@saif.sjtu.edu.cn](mailto:sdrapeau@saif.sjtu.edu.cn)

<sup>2</sup> [asgar.jamneshan@uni-konstanz.de](mailto:asgar.jamneshan@uni-konstanz.de)

---

## PAPER INFO

AMS CLASSIFICATION: 91B06

JEL CLASSIFICATION: C60, D81

---

## 1 Introduction

In decision theory, the normative framework of preference ordering classically requires the completeness axiom. Yet, there are good reasons to question this requirement as famously pointed out by Aumann [2]:

Of all the axioms of utility theory, the completeness axiom is perhaps the most questionable. [...] For example, certain decisions that an individual is asked to make might involve highly hypothetical situations, which he will never face in real life. He might feel that he cannot reach an “honest” decision in such cases. Other decision problems might be extremely complex, too complex for intuitive “insight”, and our individual might prefer to make no decision at all in these problems. Is it “rational” to force decision in such cases?

Aumann's remark, supported by empirical evidence, resulted into a consequent literature on general incomplete preferences in terms of interpretation, axiomatization, and representation, see [3, 11, 13, 15, 29, 31] and the references therein. Yet, Aumann's quote as well as a correspondence with Savage [32], where he exposes the idea of state-dependent preferences, also suggest that the information underlying a decision process is a natural source of incompleteness. Indeed, consider for instance the situation of a person facing a decision between visiting a museum or going for a walk in a park. One month ahead, she cannot express an unequivocal preference between these two prospective situations, since it depends on the knowledge of too many uncertain factors: weather, person accompanying, etc. However, the nature of this information-based incompleteness suggests a contingent form of completeness: in the previous situation, conditioned under the additional information ‘*sunny and warm day*’, she doubtlessly opts for a walk outside. This example illustrates that a complex decision problem, conditioned on sufficient information, leads to an “honest” decision. The present work provides a framework formalizing this notion of a contingent decision making and its quantification.

Numerous quantification instruments in finance and economics entail a conditional dimension by mapping prospective outcomes to random variables. For instance, this is the case of conditional/dynamic monetary risk measures [1, 5, 6, 9], conditional expected utilities/certainty equivalents or dynamic assessment indices [4, 19], or recursive utilities [12, 14]. However, few papers address the axiomatization of conditional preferences which corresponds to these conditional quantitative instruments. In this direction is the work of Luce and Krantz [26] where they consider an event-dependent preference ordering. This approach is further refined and extended in Wakker [36] and Karni [22, 23]. State-wise dependency is also used in Kreps and Porteus [24, 25] and Maccheroni et al. [27] in order to study intertemporal preferences and a dynamic version of variational preferences, respectively. Also remarkable, is the abstract approach towards conditionality by Skiadas [33, 34]. He gives an axiomatic analysis of conditional preferences on random variables which admit a conditional Savage representation of the form

$$U(x) = E_Q[u(x) | \mathcal{F}],$$

where  $\mathcal{F}$  is a  $\sigma$ -algebra representing the information,  $Q$  is a subjective probability measure and  $u$  a utility index. As in the previous works, its decision-theoretical foundation relies on a whole family of total pre-orders  $\succsim^A$ , one for each event  $A \in \mathcal{F}$ , and a consistent aggregation property in order to obtain the conditional representation. However, the decision maker is assumed to implicitly take into account a large number<sup>1</sup> of complete pre-orders. From this viewpoint, this approach does not address Aumann's observation on incompleteness.

Our axiomatic approach to conditional preferences differs in the following regard: Instead of a family of complete preference orders indexed by the contingent information, we consider a *single* but possibly *incomplete* preference order  $\succsim$ . Even though for two prospective outcomes  $x$  and  $y$  one cannot a priori decide whether  $x \succsim y$  or  $y \succsim x$ , there may exist a contingent information  $a$  conditioned on which  $x$  is preferable to  $y$ . In this case we formally write  $ax \succsim ay$ . The set of contingent information is modeled as an algebra  $\mathcal{A} = (\mathcal{A}, \vee, \wedge, \overset{c}{}, 0, 1)$  of conditions where the operations describe the disjunction  $\vee$ , coincidence  $\wedge$ , and absence  $\overset{c}{}{}$  of conditions while 0 represents the empty condition and 1 the full information. A typical example is an algebra of events. To describe the conditional nature of the preference, we require that  $\succsim$  interacts consistently with the information, that is

- *consistency*: if either  $ax \succsim ay$  or  $bx \succsim by$ , then  $(a \wedge b)x \succsim (a \wedge b)y$ ;
- *stability*: if  $ax \succsim ay$  and  $bx \succsim by$ , then  $(a \vee b)x \succsim (a \vee b)y$ ;
- *local completeness*: for every  $x, y$  there exists a non-trivial condition  $a$  such that either  $ax \succsim ay$  or  $ay \succsim ax$ .

The first two assumptions bear a certain normative appeal in view of the conditional approach that we are aiming at. Indeed, if our person prefers to go out for a walk rather than visit a museum if it is 'sunny', then a fortiori she prefers to go out if it is 'sunny' and 'warm'. The second assumption tells that if she prefers to go out if it is 'sunny' but also if it is 'not sunny', then on any day where at least one of these conditions are met, here always, she goes for a walk. In contrast to classical preferences, we only assume a local completeness: for any two situations, she is able to meet a decision provided that she is given enough – possibly extremely precise – information. Coming back to our example of going out in one month, there exists a rather unlikely but still non-trivial coincidence of conditions 'sunny', and 'humidity between 15 and 20%' and 'wind between 0 and 10km/h' under which she prefers going out rather than to the museum.

<sup>1</sup>In a five steps binary tree, 4.294.967.296 is the cardinality of the family of total pre-orders  $\succsim^A$ . Even in a simple finite state space, this large number is also disputable from a normative viewpoint.

Unlike classical completeness, the information necessary to decide between  $x$  and  $y$  depends on the pair  $(x, y)$ . Finally, if the set of contingent information reduces to the trivial information  $\mathcal{A} = \{0, 1\}$ , then, as expected, a conditional preference is a classical complete preference order. In particular, classical decision theory is a special case of the conditional one.

Although being intuitive, it is mathematically not obvious what is meant by the contingent prospective element  $ax$ . The formalization of which corresponds to the notion of a conditional set introduced recently by Drapeau et al. [10]. An heuristic introduction to conditional sets is given in Section 2, for an exhaustive mathematical presentation we refer to [10]. The formalization and properties of conditional preferences are given in Section 3. In Section 4, we address the notion of conditional numerical representation by introducing the conditional real numbers. While the proof techniques differ, the classical statements in decision theory translate to the conditional case. An example of which is the derivation of a conditional version of the classical representation of von Neumann and Morgenstern [35]. We show a conditional version of Debreu's existence result of continuous numerical representations. They rely on the Gap Lemma of Debreu [7, 8], the adaptation of which to the conditional case does not involve any measurable selection arguments but relies on the existence of a conditional axiom of choice as proven in [10]. Section 5 is dedicated to the formulation and proof of this conditional Gap Lemma.

## 2 Conditional Sets

As mentioned in the introduction, we model the contingent information, conditioned on which a decision maker ranks prospective outcomes, by a complete<sup>2</sup> Boolean algebra  $\mathcal{A}$ . The order  $a \leq b$  stays for  $a \wedge b = a$ , which for an algebra of events corresponds to the set inclusion. A set  $X$  is *conditional* on  $\mathcal{A}$  if it allows for actions  $a : X \rightarrow aX$  where  $a \in \mathcal{A}$  which satisfy a consistency and aggregation property:

**Consistency:** For any two elements  $x, y \in X$  and conditions  $a \leq b$ , if  $x$  and  $y$  coincide conditioned on  $b$ , that is  $bx = by$ , then they also coincide conditioned on  $a$ , that is  $ax = ay$ .

**Stability:** For any two elements  $x_1, x_2 \in X$  and  $a \in \mathcal{A}$ , there is  $x \in X$  such that  $x$  coincides with  $x_1$  conditioned on  $a$  and with  $x_2$  otherwise. We denote this element  $x = ax_1 + a^c x_2$ .<sup>3</sup>

Intuitively, the action  $x \mapsto ax$  tells how the elements of  $X$  are conditioned on the information  $a$  and  $aX$  represents the elements of  $X$  conditioned to  $a$ .

**Example 2.1.** Following the example from the introduction, there are two unconditional alternatives

$$x = \text{'going for a walk'} \quad \text{and} \quad y = \text{'going to the museum'},$$

and the information is reduced to a single condition  $a = \text{'sunny'}$  which yields the algebra

$$\mathcal{A} = \{0, a, a^c, 1\} = \{\text{'no information'}, \text{'sunny'}, \text{'not sunny'}, \text{'full information'}\}.$$

<sup>2</sup>The completeness is a technical but central assumption. Though, from an economical viewpoint it entails most standard frameworks, such as any finite event algebra, or any  $\sigma$ -algebra where sets are identified in the almost sure sense with respect to some probability measure, see [10, 20].

<sup>3</sup>Since  $\mathcal{A}$  is complete, the concatenation property is required for any partition  $(a_i) \subseteq \mathcal{A}$  and  $(x_i) \subseteq X$ , and we denote  $x = \sum a_i x_i$  the unique element such that  $x$  coincides with  $x_i$  conditioned on  $a_i$ .

The corresponding conditional set is then given by

$$X = \{x, y, ax + a^c y, ay + a^c x\}.$$

For instance, the conditional element  $ax + a^c y$  stays for going for a walk provided it is sunny and going to the museum otherwise.  $\diamond$

**Example 2.2.** The conditional rational numbers are defined as follows: given two rational numbers  $q_1, q_2 \in \mathbb{Q}$  and  $a \in \mathcal{A}$ , let  $q := aq_1 + a^c q_2$  be the conditional rational number which is  $q_1$  conditioned on  $a$  and  $q_2$  otherwise.<sup>4</sup> The set of conditional rational numbers, denoted by  $\mathbf{Q}$  is a conditional set whereby the conditioning action is given by  $bq = (b \wedge a)q_1 + (b \wedge a^c)q_2 \in b\mathbf{Q}$ . The conditional natural numbers  $\mathbf{N}$  are defined analogously.  $\diamond$

**Example 2.3.** Another example is the collection of random variables on a probability space  $(\Omega, \mathcal{A}, P)$  where  $\mathcal{A}$  is a  $\sigma$ -algebra of events. Events  $A \in \mathcal{A}$  act on random variables by the restriction operator on  $A$ . In the case of two random variables  $x^1, x^2$ , consistency means that if  $x^1 = x^2$  restricted to  $B \in \mathcal{A}$ , then  $x^1 = x^2$  restricted to  $A$  for any event  $A \subseteq B$ . As for the stability, if we are given  $A \in \mathcal{A}$  and two random variables  $x_1, x_2$ , then

$$x = 1_A x_1 + 1_{A^c} x_2$$

is the unique random variable  $x$  coinciding with  $x_1$  restricted to  $A$  and  $x_2$  otherwise.

Given another  $\sigma$ -algebra  $\mathcal{B}$  such that  $\mathcal{A} \subseteq \mathcal{B}$ , we can also define the conditional set  $L^1(\mathcal{B})$  of those  $\mathcal{B}$ -measurable random variable  $x$  with finite conditional expectation, that is

$$E[|x| \mid \mathcal{A}] < \infty.$$

It is in fact an  $L^0$ -module as studied and introduced in [16, 17].  $\diamond$

The relation between conditional sets is described by the *conditional inclusion* which is characterized by two dimensions, a classical inclusion and a conditioning:

- On the one hand, every set  $Y \subseteq X$  which is *stable*, that is  $ax + a^c y \in Y$  for every  $x, y \in Y$  and  $a \in \mathcal{A}$ , is a conditional subset of  $X$ .
- On the other hand,  $aX$  is a conditional set but on the relative algebra  $\mathcal{A}_a := \{b \in \mathcal{A} : b \leq a\}$  and a subset of  $X$  but conditioned on  $a$ .

Combining the two dimensions, a conditional set  $Y$  is said to be *conditionally included* in  $X$ , denoted  $Y \sqsubseteq X$ , if  $Y = aZ$  for some stable  $Z \subseteq X$  and condition  $a \in \mathcal{A}$ , as illustrated in Figure 1a. In this case  $Y$  is a conditional subset “*living*” on  $a$ . If  $a = 0$ , then  $Y = 0Z$  lives nowhere and in particular is conditionally contained in any conditional set, and thus is conditionally the emptyset. In order to distinguish between both the conditional emptyset is denoted by  $\mathbf{0}$ . The *conditional power-set*


$$\mathcal{P}(X) := \{Y : Y \sqsubseteq X\} = \{Y : Y = aZ \text{ for some stable } Z \subseteq X \text{ and } a \in \mathcal{A}\}$$

consists of the collection of all conditional subsets of  $X$ .


<sup>4</sup>More generally, given a partition of information  $(a_i) \subseteq \mathcal{A}$  and a corresponding family of rationals  $(q_i) \subseteq \mathbb{Q}$ , define the conditional rational number  $q := \sum a_i q_i$  as the conditional element valued  $q_i$  conditioned on  $a_i$ .

**Example 2.4.** In Example 2.1, the set  $\{x, y\} \subseteq X$  is not stable since  $ax + a^c y \notin \{x, y\}$ . Hence  $\{x, y\}$  is not a conditional set whereas  $Z = \{x, ay + a^c x\}$  is a conditional subset of  $X$  living on 1. However,  $Y = \{ax, ay\}$  is a conditional subset of  $X$  living on  $a$ . Indeed,  $Y = a\{x, ay + a^c x\} = aZ$ .  $\diamond$

The conditional intersection of two conditional sets  $X, Y$  is the intersection on the largest condition  $a^*$  on which  $X$  and  $Y$  have a non-empty classical intersection as illustrated in Figure 1b. The conditional



(a) Illustration of the conditional inclusion.




(b) Illustration of the conditional intersection.


union of two conditional subsets  $X, Y$  is the collection of all elements which can be concatenated such that each piece of the concatenation conditionally falls either in  $X$  or in  $Y$ . Formally, the conditional union is

$$X \sqcup Y := \{ax + by : ax \in aX, by \in bY, (a, b) \text{ partition}\},$$

which is illustrated in Figure 1c. Finally, the conditional complement in  $Z$  of a conditional set  $X \sqsubseteq Z$  is the collection of all those elements  $x$  which nowhere fall into  $X$ , as illustrated in Figure 1d.



(c) Illustration of the conditional union..



(d) Illustration of the conditional complement.

One of the central results in [10] is that the conditional power set together with these operations

$$(\mathcal{P}(X), \sqcup, \sqcap, \sqsupseteq, \mathbf{0}, X)$$

forms a complete Boolean algebra. Following the classical constructions, the conditional power set allows to define conditional relations, functions, topologies, etc. [10].

### 3 Conditional Preference Orders

For the remainder of the paper  $X$  denotes a conditional set. A conditional binary relation  $\succsim$  is a conditional subset  $G \sqsubseteq X \times X$  and we write  $x \succsim y$  if and only if  $(x, y) \in G$ . In particular, a conditional binary relation is at first a classical binary relation. However, due to the fact that the graph  $G$  is a conditional set the following additional properties hold<sup>5</sup>

- *consistency*: if either  $ax \succsim ay$  or  $bx \succsim by$ , then  $(a \wedge b)x \succsim (a \wedge b)y$ ;
- *stability*: if  $ax \succsim ay$  and  $bx \succsim by$ , then  $(a \vee b)x \succsim (a \vee b)y$ ;

corresponding to two of the normative properties mentioned in the introduction. Given a conditional binary relation,  $\sim$  denotes the symmetric part of the binary relation and we use the notation

$$x \succ y \quad \text{if and only if} \quad x \succsim y \text{ and } ay \not\sucsim ax \text{ for every non-trivial condition } a \in \mathcal{A}.$$

In other words,  $x \succ y$  means that  $x$  is strictly preferred to  $y$  on any non-trivial condition. Both  $\sim$  and  $\succ$  are also conditional binary relations.

**Definition 3.1.** A conditional binary relation  $\succsim$  on  $X$  is called a *conditional preference order* if  $\succsim$  is

- *reflexive*:  $x \succsim x$  for every  $x$ ;
- *transitive*: From  $x \succsim y$  and  $y \succsim z$  it follows that  $x \succsim z$ ;
- *locally complete*: for every  $x \not\sucsim y$ , there exists a non-trivial condition  $a \neq 0$  such that either  $ax \succ ay$  or  $ay \succ ax$ .

Even if a conditional preference is not total, the following lemma shows that local completeness allows to derive for every two elements a partition on which a comparison can be achieved.

**Lemma 3.2.** *Let  $\succsim$  be a conditional preference order on  $X$  and  $x, y \in X$ . There is a disjoint family of conditions  $a, b, c$  such that  $a \vee b \vee c = 1$  and*

$$ax \sim ay, \quad bx \succ by \quad \text{and} \quad cy \succ cx.$$

*Proof.* Let  $x, y \in X$  and define

$$a = \vee\{\tilde{a} \in \mathcal{A} : \tilde{a}x \sim \tilde{a}y\}, \quad b = \vee\{\tilde{b} \in \mathcal{A} : \tilde{b}x \succ \tilde{b}y\} \quad \text{and} \quad c = \vee\{\tilde{c} \in \mathcal{A} : \tilde{c}y \succ \tilde{c}x\}$$

which are the largest conditions on which  $x$  is conditionally equivalent, strictly better or worse than  $y$ , respectively. Due to the consistency property of conditional relations, it follows that these conditions

<sup>5</sup>We write  $ax \succ ay$  for  $a(x, y) = (ax, ay) \in aG$ .

are mutually disjoint. For the sake of contradiction, suppose that  $d := a \vee b \vee c < 1$ . It follows by conditionality that outside  $d$ , that is, conditioned on  $d^c$ , the element  $x$  is nowhere either equivalent, strictly better or worse than  $y$ . Indeed, this contradicts otherwise the definition of  $a$ ,  $b$  and  $c$ . Define  $\tilde{y} = dx + d^c y$  being  $x$  conditioned on  $d$  and  $y$  conditioned on  $d^c$ . Since  $x \not\sim y$  and  $\sim$  is a conditional equivalence relation, it follows that  $x \not\sim \tilde{y}$ , otherwise  $d^c x \sim d^c y$  contradicting the definition of  $a$ . By local completeness, there exists a non-trivial condition  $e > 0$  such that either  $ex \succ e\tilde{y}$  or  $e\tilde{y} \succ ex$ . Without loss of generality, suppose that  $ex \succ e\tilde{y}$ . Since  $d\tilde{y} = dx \sim dx$  by reflexivity and consistency, it follows that  $e$  is disjoint from  $d$ , in other words  $e \leq d^c$ . In particular  $e\tilde{y} = ey$  implying that  $ex \succ ey$ , which together with  $0 < e \leq d^c$  contradict the maximality of  $b$ . Thus  $d = 1$  which ends the proof.  $\square$

**Example 3.3.** Let us give a complete formal description of the example in the introduction. Recall from Example 2.1 that  $\mathcal{A} = \{\text{'no information'}, \text{'sunny'}, \text{'not sunny'}, \text{'full information'}\}$ , and that the conditional set generated from the two unconditional choices  $x = \text{'going for a walk'}$  and  $y = \text{'going to the museum'}$  is given by  $X = \{x, y, ax + a^c y, ay + a^c x\}$ . The conditional preference being reflexive, it trivially holds

$$x \succcurlyeq x, \quad y \succcurlyeq y, \quad ax + a^c y \succcurlyeq ax + a^c y, \quad ay + a^c x \succcurlyeq ay + a^c x.$$

Further, the individual prefers going out if it is sunny and go to the museum otherwise. This translate into

$$ax \succ ay \quad \text{and} \quad a^c y \succ a^c x.$$

Since the preference is assumed to be conditional, it also holds

$$ax + a^c y \succcurlyeq ay + a^c x, \quad x \succcurlyeq ay + a^c x, \quad ax + a^c y \succcurlyeq y.$$

For instance, the relation  $x \succcurlyeq ay + a^c x$  states that going out is in any case better than going to the museum if it is sunny and going out otherwise. Inspection shows that the conditional preference is indeed a transitive and reflexive conditional relation. As mentioned, this relation does not tell though whether  $x$  is preferred to  $y$ , that is, whether she wants to go for a walk or to the museum. There exists however a condition  $a = \text{'sunny'}$ , such that  $ax \succ ay$  which shows that it is locally complete. In particular, the partitioning given in Lemma 3.2 corresponds to

$$0x \sim 0y, \quad ax \succ ay, \quad a^c y \succ a^c x.$$

Note also that conditional sets allow to solve the following puzzle: Define  $Y = \{z \succ y\}$ , the set of elements which are strictly preferred to  $y$ . In a classical setting this set is empty. Indeed, there exists no alternative which is strictly preferred to going to the museum since conditioned on  $a^c$ ,  $y$  is maximal for the preference order. However, as our intuition suggests, this set should not be empty and indeed it is conditionally non-empty since  $Y = \{ax\}$ . The importance of this fact is observed in the proof of Debreu's Theorem 4.5 with the definitions of  $Z^\pm$  towards the construction of a conditional numerical representation.  $\diamond$

**Example 3.4.** In the unconditional case  $(\mathbb{Q}, \leq)$  is a classical example of a total order. We extend this order to the conditional rational numbers  $\mathbb{Q}$ :

$$q = aq_1 + a^c q_2 \leq r = br_1 + b^c r_2 \quad \text{if and only if} \quad \begin{cases} q_1 \leq r_1 & \text{if } a \wedge b > 0 \\ q_1 \leq r_2 & \text{if } a \wedge b^c > 0 \\ q_2 \leq r_1 & \text{if } a^c \wedge b > 0 \\ q_2 \leq r_2 & \text{if } a^c \wedge b^c > 0 \end{cases}$$

Its extension to the general case makes this partial order locally complete, see [10]. In particular, this allows to define on  $\mathbf{Q}$  the conditional variant of the Euclidean topology on  $\mathbb{Q}$  by the conditional balls:

$$B_\varepsilon(q) := \{r \in \mathbf{Q} : |q - r| \leq \varepsilon\},$$

for  $q \in \mathbf{Q}$  and  $\varepsilon \in \mathbf{Q}_{++} := \{r \in \mathbf{Q} : r > 0\}$ . It behaves like the standard topology on  $\mathbb{Q}$  with the additional local property:

$$aB_{\varepsilon_1}(q) + a^cB_{\varepsilon_2}(r) := B_{a\varepsilon_1 + a^c\varepsilon_2}(aq + a^cr),$$

for every  $\varepsilon_1, \varepsilon_2 \in \mathbf{Q}_{++}$  and  $q, r \in \mathbf{Q}_{++}$ . In other words, a conditional neighborhood of 3 conditioned on  $a$  and  $2/5$  on  $a^c$ , is itself a conditional neighborhood of the conditional rational  $a3 + a^c2/5$ .  $\diamond$

*Remark 3.5.* In general a conditional preference can be an equivalence relation conditioned on  $a$  and strictly non-trivial on  $a^c$ . However, the case of interest lives on  $a^c$ . Therefore, throughout this paper, we assume that a conditional preference is conditionally non-trivial, that is, there exists a pair  $x, y \in X$  such that  $x \succ y$   $\diamond$

## 4 Conditional Numerical Representations

Next we address the quantification of such a conditional ranking. First, we need the notion of a *conditional function*. A conditional function  $f : X \rightarrow Y$  between two conditional sets is a classical function with the additional property of stability:

$$f(ax + a^cy) = af(x) + a^cf(y).$$

**Example 4.1.** The  $\mathcal{A}$ -conditional expectation of elements of  $L^1(\mathcal{B})$  introduced in Example 2.3, is a conditional function. Indeed, for every  $x, y \in L^1(\mathcal{B})$  and  $A \in \mathcal{A}$  it holds

$$f(1_Ax + 1_{A^c}y) := E[1_Ax + 1_{A^c}y | \mathcal{A}] = 1_AE[1_Ax | \mathcal{A}] + 1_{A^c}E[y | \mathcal{A}] = 1_Af(x) + 1_{A^c}f(y)$$

since  $1_A$  is  $\mathcal{A}$ -measurable.  $\diamond$

**Example 4.2.** For  $q = aq_1 + a^cq_2$  and  $r = br_1 + b^cr_2$ , define the conditional addition and conditional absolute value on  $\mathbf{Q}$  as

$$q + r := \begin{cases} q_1 + r_1 & \text{on } a \wedge b \\ q_1 + r_2 & \text{on } a \wedge b^c \\ q_2 + r_1 & \text{on } a^c \wedge b \\ q_2 + r_2 & \text{on } a^c \wedge b^c \end{cases} \quad \text{and} \quad |q| := a|q_1| + a^c|q_2|.$$

Together with an analogous definition for conditional multiplication these operations make  $\mathbf{Q}$  a conditional totally ordered field in the sense given in [10].  $\diamond$

For the quantification, we secondly need a conditional analogue of the real line which allows to represent the conditional preferences. The conditional real numbers, denoted by  $\mathbf{R}$ , are obtained from the



conditional rational numbers by adapting Cantor's construction. As in the standard theory, the conditional real numbers can be characterized as a conditional field where every bounded subset has an infimum and a supremum and which is topologically conditionally separable. In particular,  $\mathbf{Q}$  is conditionally dense in  $\mathbf{R}$ . For further details, we refer the interested reader to [10].

**Definition 4.3.** A conditional numerical representation of a conditional preference order  $\succsim$  on  $X$  is a conditional function  $U : X \rightarrow \mathbf{R}$  such that

$$x \succsim y \quad \text{if and only if} \quad U(x) \geq U(y). \quad (4.1)$$

Note that every conditional function  $U : X \rightarrow \mathbf{R}$  defines a conditional preference order by means of (4.1). Furthermore, if  $U : X \rightarrow \mathbf{R}$  is a conditional numerical representation, then  $\varphi \circ U$  is a conditional numerical representation for every conditionally strictly increasing function  $\varphi : \mathbf{R} \rightarrow \mathbf{R}$ .

*Remark 4.4.* In the case where  $\mathcal{A}$  is an algebra of events on a probability space,  $\mathbf{R}$  is isometric to the conditional set of random variables for the  $L^0$ -topology introduced in [16], as shown in [10]. Hence, the conditional entropic monetary utility function studied for instance in [19] as a special case of a conditional certainty equivalent and given by

$$U(x) = \ln (E [e^x | \mathcal{A}]), \quad x \in L^1(\mathcal{B}),$$

is a representation of a conditional preference. Indeed, this function is local since for every  $A \in \mathcal{A}$  it holds

$$\begin{aligned} U(1_A x + 1_{A^c} y) &= \ln (E [e^{1_A x + 1_{A^c} y} | \mathcal{A}]) = \ln (1_A E [e^x | \mathcal{A}] + 1_{A^c} E [e^y | \mathcal{A}]) \\ &= 1_A \ln (E [e^x | \mathcal{A}]) + 1_{A^c} \ln (E [e^y | \mathcal{A}]) = 1_A U(x) + 1_{A^c} U(y). \end{aligned}$$

The same argumentation holds for all conditional certainty equivalents, conditional/dynamic risk measures or acceptability indices mentioned in the introduction.  $\blacklozenge$

Given a conditional preference order, we address the necessary and sufficient conditions under which a conditional numerical representation exists. The first result is a conditional version of Debreu [7] and necessitates the notion of conditionally order dense. A conditional subset  $Z \sqsubseteq X$  is *conditionally order dense* if for every  $x, y \in X$  with  $x \succ y$ , there exists  $z \in Z$  such that  $x \succ z \succ y$ . The case of interest is when  $Z$  is conditionally countable, that is, there exists a conditional injection  $\varphi : Z \rightarrow \mathbf{Q}$ . Equivalently,  $Z$  is conditionally countable if it is a conditional sequence  $Z = (z_n)_{n \in \mathbf{N}}$  where  $\mathbf{N}$  is the conditional natural numbers. There exists a difference between a conditional sequence and a standard sequence: Analogous to the classical case, a conditional sequence  $(z_n)$  in  $Z$  is a conditional function  $f : \mathbf{N} \rightarrow Z$ ,  $n \mapsto f(n) = z_n$ . However stability yields  $az_n + a^c z_m = af(n) + a^c f(m) = f(an + a^c m) = z_{an + a^c m}$ . In other words, the sequence step  $n$  conditioned on  $a$  and the sequence step  $m$  conditioned on  $b$  result into the sequence step  $an + a^c m$ .

**Theorem 4.5.** A conditional preference order  $\succsim$  on  $X$  admits a conditional numerical representation if and only if  $X$  has a conditionally countable order dense subset.

*Proof. The if-part:* Without loss of generality assume  $Z = (z_n)_{n \in \mathbf{N}}$  is a conditionally countable order dense subset of  $X$  which is not conditionally finite. Consider now

$$Z^+(x) := \{z \in Z : z \succ x\} \quad \text{and} \quad Z^-(x) := \{z \in Z : x \succ z\}.$$

Since  $\succsim$  is a conditional binary relation,  $Z^+(x)$  and  $Z^-(x)$  are conditional subsets of  $Z$  for every  $x \in X$ . However, as mentioned in Example 3.3,  $Z^\pm(x)$  may both live on some conditions smaller than 1.<sup>6</sup> Further,  $(Z^\pm(x))_{x \in X}$  is a conditional family in the conditional power-set  $\mathcal{P}(Z)$ , that is  $Z^\pm(x) = aZ^\pm(x_1) + a^cZ^\pm(x_2)$  for every  $x = ax_1 + a^cx_2 \in X$ . Due to transitivity,

$$x \succsim y \text{ implies } Z^+(x) \sqsubseteq Z^+(y) \text{ and } Z^-(y) \sqsubseteq Z^-(x). \quad (4.2)$$

It follows from conditional order denseness that  $x \succ y$  implies that there is  $z \in Z$  such that  $x \succ z \succsim y$  on some condition  $a$  and  $x \succ z \succ y$  on  $a^c$ . Thus,

$$az \in a[Z^-(x) \setminus Z^-(y)] \quad \text{and} \quad a^cz \in a^c[Z^+(y) \setminus Z^+(x)] \text{ for some } z \in Z \text{ and } a \in \mathcal{A}. \quad (4.3)$$

Let now  $\mu$  be a strictly positive conditional measure on  $Z$ <sup>7</sup>, that is  $\mu(\{z_n\}) > 0$  for every  $n \in \mathbf{N}$ . Define then  $U(x) = \mu(Z^-(x)) - \mu(Z^+(x))$  for every  $x \in X$ . Then  $U$  is a conditional function since  $(Z^\pm(x))_{x \in X}$  is a conditional family and  $\mu$  a conditional function. On the one hand, from (4.2) and  $\mu$  being conditionally increasing it follows that  $x \succsim y$  implies  $U(x) \geq U(y)$ . On the other hand, assume that  $x \succ y$  on some condition  $a > 0$ . Without loss of generality  $a = 1$ . Then from (4.3) and  $\mu$  being strictly positive, it follows that  $x \succ y$  yields

$$\begin{aligned} U(x) &= \mu(Z^-(x)) - \mu(Z^+(x)) \\ &\geq a[\mu(\{z\}) + \mu(Z^-(y))] - a^c[\mu(Z^+(y)) - \mu(\{z\})] \\ &= \mu(\{z\}) + U(y) > U(y). \end{aligned}$$

From the conditional completeness of  $\succsim$  it follows that  $U$  is a conditional numerical representation.

**The only if-part:** A conditional preference order which admits a conditional numerical representation is conditionally complete since the conditional reals are so. It holds that  $Y := \text{Im}(U)$  is a conditional subset of  $\mathbf{R}$ . Choose a conditionally countable order dense subset  $I \sqsubseteq Y$  by Lemma 5.1. Then  $Z := U^{-1}(I)$  is conditionally countable and since  $U$  is a conditional numerical representation, it is conditionally order dense.  $\square$

The existence of a conditionally countable order dense subset is rather of technical nature. In the classical case, Debreu [7, 8] and then Rader [30] showed that under some topological assumptions a numerical representation exists. And even more, by means of Debreu's Gap Lemma, the existence of an upper semi-continuous or continuous representation is guaranteed. The conditional counterparts of these results also hold, based on a conditional adaptation of Debreu's Gap Lemma in Section 5.

**Definition 4.6.** Let  $\succsim$  be a conditional preference order on a conditional topological space  $X$ .<sup>8</sup> We say that  $\succsim$  is *conditionally upper semi-continuous* if  $\mathcal{U}(x) := \{y \in X : y \succsim x\}$  is conditionally closed for every  $x \in X$ . A conditional numerical representation  $U : X \rightarrow \mathbf{R}$  is said to be *conditionally upper semi-continuous* if  $\{x \in X : U(x) \geq m\}$  is conditionally closed for every  $m \in \mathbf{R}$ .

**Theorem 4.7.** *Let  $\succsim$  be a conditionally upper semi-continuous preference order on a conditionally second countable<sup>9</sup> topological space  $X$ . Then  $\succsim$  admits a conditionally upper semi-continuous numerical*

<sup>6</sup>Let  $a$  be the condition on which  $Z^+(x)$  lives. It means that there exists no  $z \in Z$  such that  $z$  is strictly preferred to  $x$  conditioned on  $a^c$ . Since  $Z$  is conditionally order dense, it follows that  $x$  is a maximal element conditioned on  $a^c$ .

<sup>7</sup>For instance, define  $\mu(\{z_n\}) = 2^{-n} := \sum a_k 2^{-n_k}$  for every  $n = \sum a_k n_k \in \mathbf{N}$ .

<sup>8</sup>A conditional topology is the counterpart to a classical topology but with respect to the conditional operations of union and intersection, see [10].

<sup>9</sup>The conditional topology of which is generated by a conditionally countable neighborhood base.

representation. In particular, if  $\succsim$  is conditionally continuous<sup>10</sup>, then it admits a conditionally continuous numerical representation.

*Proof.* Let  $\mathcal{O} = (O_n)_{n \in \mathbf{N}}$  be a conditionally countable topological base of  $X$  and  $\mu$  be a strictly positive measure on  $\mathbf{N}$ . We know that  $Z(x) := \mathcal{U}(x)^\square$  is conditionally open for every  $x \in X$ . Fix some  $x \in X$  and let  $a$  be the condition on which lives  $Z(x)$ . Then  $\{n \in \mathbf{N} : aO_n \sqsubseteq Z(x)\}$  is a conditional subset of  $\mathbf{N}$ . Next define

$$U(x) = a\mu(\{n \in \mathbf{N} : aO_n \sqsubseteq Z(x)\}) + a^c0.$$

If  $x \succsim y$ , then  $U(x) \geq U(y)$  since  $Z(y) \sqsubseteq Z(x)$ . Otherwise if  $x \succ y$ , then  $y \in Z(x)$  and since  $Z(x)$  is conditionally open, there exists a neighborhood  $O_{i_0}$  of  $y$  such that  $O_{i_0} \sqsubseteq Z(x)$ . However, since  $y \in Z(y)^\square \cap O_{i_0}$ , it follows that  $O_{i_0}$  is nowhere a subset of  $Z(y)$ . Hence,  $U(x) \geq U(y) + \mu(\{i_0\}) > U(y)$ . By Theorem 5.3 we can choose  $U$  to be conditionally upper semi-continuous which ends the proof.  $\square$

A classical class of preferences are the affine one, due to von Neumann and Morgenstern [35]. This representation can be carried over to the conditional case as follows. Let  $X$  be a conditionally convex subset living on 1 of some conditional vector space. We say that a conditional preference order on  $X$  satisfies the

- **conditional independence axiom:** if  $x \succ y$  then  $\alpha x + (1 - \alpha)z \succ \alpha y + (1 - \alpha)z$  for every  $z \in X$  and  $\alpha \in ]0, 1[$ ;
- **conditional Archimedean axiom:** if  $x \succ y \succ z$  then  $\alpha x + (1 - \alpha)z \succ y \succ \beta x + (1 - \beta)z$  for some  $\alpha, \beta \in ]0, 1[$ .

A conditional real-valued numerical representation  $U$  of  $\succsim$  is *conditionally affine*, if

$$U(\alpha x + (1 - \alpha)y) = \alpha U(x) + (1 - \alpha)U(y),$$

for every  $x, y \in X$  and  $\alpha \in [0, 1]$ .

**Theorem 4.8.** *Let  $\succsim$  be a conditionally complete preference order satisfying both the conditional Archimedean and independence axioms. Then  $\succsim$  admits a conditionally affine representation  $U$ . Moreover, if  $\hat{U}$  is another conditionally affine representation, then  $\hat{U} = \alpha U + \beta$ , where  $\alpha > 0$  and  $\beta \in \mathbf{R}$ .*

We will follow the classical proof adapted to the conditional setting.

**Lemma 4.9.** *Let  $\succsim$  be a conditionally complete preference order satisfying both the conditional independence and Archimedean axioms. Then the following assertions hold:*

- (i) *If  $x \succ y$ , then  $\beta x + (1 - \beta)y \succ \alpha x + (1 - \alpha)y$  for all  $0 \leq \alpha < \beta \leq 1$ .*
- (ii) *If  $x \succ z$  and  $x \succ y \succ z$ , then there exists a unique  $\alpha \in [0, 1]$  with  $y \sim \alpha x + (1 - \alpha)z$ .*
- (iii) *If  $x \sim y$ , then  $\alpha x + (1 - \alpha)z \sim \alpha y + (1 - \alpha)z$  for all  $\alpha \in [0, 1]$  and all  $z \in \mathcal{M}$ .*

*Proof.* (i) Strictly analogous to the classical proof, e.g. [18, p. 54].

<sup>10</sup>That is  $\mathcal{U}(x) = \{y \in X : y \succ x\}$  and  $\mathcal{U}(y) = \{y \in X : x \succ y\}$  are conditionally closed for every  $x \in X$ .

(ii) Up to conditioning, we may assume that  $x \succ y \succ z$ .<sup>11</sup> The candidate is

$$\alpha := \sup\{\beta \in [0, 1] : y \succ \beta x + (1 - \beta)z\}$$

We obtain a partition  $a, b, c$  of 1 such that  $y \sim \alpha x + (1 - \alpha)z$  on  $a$ ,  $y \succ \alpha x + (1 - \alpha)z$  on  $b$  and  $\alpha x + (1 - \alpha)z \succ y$  on  $c$ . Conditioned on  $b$  and  $c$  respectively, we may apply the classical argumentation, e.g. [18, p. 54] yielding a contradiction showing that  $b = c = 0$  and therefore  $a = 1$  which ends the proof. As for the uniqueness, this is a consequence of the first point.

(iii) The same conditioning technique holds also here. Let  $\alpha \in [0, 1]$  and  $z \in X$ . There exists a partition of  $a, b, c$ , of 1 such that  $\alpha x + (1 - \alpha)z \sim \alpha y + (1 - \alpha)z$  on  $a$ ,  $\alpha x + (1 - \alpha)z \succ \alpha y + (1 - \alpha)z$  on  $b$  and  $\alpha x + (1 - \alpha)z \succ \alpha y + (1 - \alpha)z$  on  $c$ . The same contradiction argumentation as in the classical case, [18, p. 54–55], conditioned on  $b$  and  $c$ , respectively, shows that  $b = c = 0$  and therefore  $a = 1$ .  $\square$

*Proof (Theorem 4.8).* Let  $x, y \in X$  be such that  $x \succ y$  and define the conditional convex subset  $N_{x,y} := \{z \in X : x \succ z \succ y\}$ . For  $z \in N_{x,y}$ , part (ii) of Lemma 4.9 yields a unique  $\alpha \in [0, 1]$  such that  $z \sim \alpha x + (1 - \alpha)y$ . Setting  $U(z) := \alpha$ ,  $z \in N_{x,y}$  provides a well defined conditional function from  $N_{x,y}$  to  $[0, 1]$ . Indeed, let  $[a_i, z_i] \subseteq \mathcal{A} \times N_{x,y}$ , and denote  $\alpha_i = U(z_i)$  and  $\alpha = U(\sum a_i z_i)$ . There exists a partition  $a, b, c$  of 1 such that  $\alpha = \sum a_i \alpha_i$  on  $a$ ,  $\alpha > \sum a_i \alpha_i$  on  $b$  and  $\sum a_i \alpha_i > \alpha$  on  $c$ . It means in particular that  $\alpha > \alpha_i$  on  $b \wedge a_i$  and  $\alpha_i > \alpha$  on  $c \wedge a_i$  for every  $i$ . Hence, if either  $b$  or  $c$  were strictly positive conditions, this would contradict the uniqueness of some  $\alpha_i$  on  $b \wedge a_i > 0$  or  $c \wedge a_i > 0$ . Hence  $b = c = 0$  showing that  $a = 1$ . The extension to  $X$  follows exactly the same argumentation as the classical case, see [18, p. 55].  $\square$

The result of Neumann and Morgenstern goes a step forward by providing a utility index against which lotteries are ranked according to expectation. In our context, lotteries become conditional lotteries

$$X := \{\mu : \text{is a conditional probability distribution on } \mathbf{R}\}$$

see Jamneshan et al. [21] for the construction and definition of such conditional probability distributions. Conditional probability distributions can be identified with stochastic kernels with respect to  $\mathcal{A}$  or Anscombe and Aumann acts which scenario dependence is described by  $\mathcal{A}$ . In this framework, the classical representation Theorem of von Neumann and Morgenstern carries over

**Theorem 4.10.** *Let  $\succ$  be a conditional preference order on the conditional convex set of lotteries  $X$ . Suppose that  $\succ$  fulfills the conditional independence and Archimedean axioms. Suppose finally that  $\succ$  is  $\sigma(X, C)$ -continuous<sup>12</sup>, then there exists a unique, up to strictly positive affine transformation, conditionally continuous utility function  $u : \mathbf{R} \rightarrow \mathbf{R}$  such that*

$$\mu \succ \nu \quad \text{if and only if} \quad \int_{\mathbf{R}} u(x) \mu(dx) \geq \int_{\mathbf{R}} u(x) \nu(dx)$$

*Proof.* This is a consequence of Theorem 4.8 together with the conditional version of Riez Theorem, see [21].  $\square$

<sup>11</sup>On the condition where  $x \sim y$  set  $\alpha = 1$  and on the condition where  $y \sim z$ , set  $\alpha = 0$ .

<sup>12</sup>For the conditional weak topology generated by the set of conditionally continuous functions  $f : \mathbf{R} \rightarrow \mathbf{R}$ .

*Remark 4.11.* In the case where  $\mathcal{A}$  is an algebra of events on a probability space  $(\Omega, \mathcal{A}, P)$ , it follows that conditional lotteries on  $\mathbb{R}$  can be identified with stochastic kernels  $\mu : \Omega \otimes \mathcal{B}(\mathbb{R}) \rightarrow [0, 1]$  where  $\mathcal{B}(\mathbb{R})$  is the Borel  $\sigma$ -algebra on the real line. The conditional utility function can be identified with a function  $u : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  where  $u(\omega, \cdot)$  is a state-wise utility function. It follows that the von Neuman and Morgenstern conditional preference order can be characterized by

$$\mu \succcurlyeq \nu \quad \text{if and only if} \quad \int_R u(\omega, x) \mu(\omega, dx) \geq \int_R u(\omega, x) \nu(\omega, dx) \quad \text{for } P\text{-almost all } \omega \in \Omega. \quad \blacklozenge$$

## 5 Conditional Gap Lemma

For  $s, t \in \mathbf{R}$  with  $s \leq t$  we denote  $[s, t] = \{u \in \mathbf{R} : s \leq u \leq t\}$  and if  $s < t$  we denote  $[s, t[ = \{u \in \mathbf{R} : s \leq u < t\}$ ,  $]s, t] = \{u \in \mathbf{R} : s < u \leq t\}$ , and  $]s, t[ = \{u \in \mathbf{R} : s < u < t\}$ , all conditionally convex subsets of  $\mathbf{R}$  which live on 1. In the conditional topology of  $\mathbf{R}$  the conditionally convex subset  $[s, t]$  is conditionally closed whereas  $]s, t[$  is conditionally open. A conditionally convex subset  $I \sqsubseteq \mathbf{R}$  is an *interval*. Denoting by  $s = \inf I$  and  $t = \sup I$ , an interval is generically denoted by  $(s, t)$ .<sup>13</sup> Up to conditioning, all conditionally convex subsets of  $\mathbf{R}$  are characterized as conditional intervals. If we assume that an interval lives on 1, convexity yields

$$(s, t) = a[s, t] + b[s, t[ + c]s, t] + d]s, t[ \quad (5.1)$$

where  $a = e \wedge f$ ,  $b = e \wedge f^c$ ,  $c = e^c \wedge f$ ,  $d = e^c \vee f^c$  and  $e = \vee \{\tilde{e} : \tilde{e}s \in \tilde{e}I\}$  and  $f = \vee \{\tilde{f} : \tilde{f}t \in \tilde{f}I\}$ , as illustrated in Figure 1.

Let  $S \sqsubseteq \mathbf{R}$  and  $(s, t)$  be an interval<sup>14</sup> such that  $(s, t) \sqsubseteq S^\square$ . Inspection shows that there exists a unique maximal interval  $(s^*, t^*)$  with respect to conditional inclusion such that  $(s, t) \sqsubseteq (s^*, t^*) \sqsubseteq S^\square$ . Such a maximal interval with  $\inf S^\square < s^* \leq t^* < \sup S^\square$  on the conditions where  $s^*, t^*$  are living is called a *conditional gap* of  $S$ . Note that any conditional gap  $(s, t)$  of  $S$  can be decomposed into

$$(s, t) = a\{s\} + b(s, t)$$

where  $a \wedge b = 0$  and  $s < t$  on  $b$ , that is the conditional interior of  $(s, t)$  lives on  $b$ . Moreover, the family of conditional gaps of  $S$  is itself stable and therefore each of the conditional gaps of  $S$  lives on the same condition. Indeed, suppose that two conditional gaps  $(s_1, t_1)$  and  $(s_2, t_2)$  live on  $a$  and  $b$ , respectively, and  $a < b$ . Then it follows that

$$(s_1, t_1) = a(s_1, t_1) \sqsubseteq a(s_1, t_1) + (b \wedge a^c)(s_2, t_2) \sqsubseteq S^\square$$

contradicting the maximality of  $(s_1, t_1)$ . Hence  $a = b$ .

**Lemma 5.1.** *The conditionally complete order  $\succcurlyeq$  restricted to any  $S \sqsubseteq \mathbf{R}$  living on 1 admits a conditionally countable order dense subset.*

*Proof.* Analogous to conditional gaps, we define a *predecessor-successor* as a maximal interval  $(s, t) \sqsubseteq S^\square$  but under the additional requirement that  $s < t$ . In other words, these are conditionally maximal non-trivial conditional gaps. Alike conditional gaps, predecessor-successor pairs of  $S$  form a conditional

<sup>13</sup>It is possible that  $s, t$  attain  $\pm\infty$  on some positive condition.

<sup>14</sup>May live on a condition  $a < 1$ .




Figure 1: Illustration of a Gap.

family and therefore all live on the same condition. This condition is per definition smaller than the one on which the conditional gaps live.

Now, up to conditioning, we may assume that the conditional gaps of  $S$  are all living on 1.<sup>15</sup> Since  $B_{1/m}(q)$  for  $m \in \mathbf{N}$  and  $q \in \mathbf{Q}$  is a conditionally countable base of the topology, the family of those intersections  $B_{1/m}(q) \cap S$  living on 1 is a conditionally countable family which we denote  $(U_n)$ . By means of the conditional axiom of choice, see [10, Theorem 2.17], there exists a conditionally countable family  $(u_n)$  such that  $u_n \in U_n$  for all  $n$ . Let further  $a$  be the condition on which the conditional family  $(s_i, t_i)$  of the predecessors-successors of  $S$  is living. It follows that  $U = (u_n) \sqcup (s_i)$  is a conditionally countable order dense subset of  $S$  living on 1. Indeed, let  $s < t$  for  $s, t \in S$  and  $b$  the condition on which  $(s, t)$  is a predecessor-successor pair, that is the maximal condition such that  $s$  is an element of  $(s_i)$ . It follows that there exists  $v \in S$  such that  $s < v < t$  on  $b^c$ . Hence, we may find  $q \in \mathbf{Q}$  and  $n \in \mathbf{N}$  such that  $s < q - 1/m < v < q + 1/m < t$  on  $b^c$  which ensures the existence of some  $u_n$  in the family  $(u_n)$  such that  $s < u_n < t$  on  $b^c$ . It follows that  $u = bs + b^c u_n \in U$  and  $s \leq u \leq t$ .

We are then left to show that  $U$  is conditionally countable. Since  $(u_n)$  is conditionally countable, according to [10, Lemma 2.33], it is enough to show that the conditional family of open sets  $]s_i, t_i[ = ]s_i, t_i[_{i \in I}$ , where  $(s_i, t_i)$  is the conditional collection of predecessor-successors, is conditionally countable. Without loss of generality, suppose that this family lives on 1. For any two  $]s_i, t_i[_$  and  $]s_j, t_j[_$  such that  $s_i \neq s_j$  on any condition, it follows that  $]s_i, t_i[_ \cap ]s_j, t_j[_ = \mathbf{0}$ . This provides a conditionally mutually disjoint family of conditionally open sets on 1. By means of the conditional axiom of choice, [10, Theorem 2.17], we choose a conditional family  $(q_i)$  of elements of  $\mathbf{Q}$  such that  $q_i \in ]s_i, t_i[_$  for every  $i$ . For  $P = \sqcup \{q_i\} \subseteq \mathbf{Q}$ , define  $f : I \rightarrow P$ ,  $i \mapsto q_i$ . This function is a well-defined conditional function. Indeed, for  $q_i = q_j$ , it follows that  $q_i \in ]s_i, t_i[_ \cap ]s_j, t_j[_$ . Both being conditional gaps of  $S$ , this implies that

<sup>15</sup>On the condition where none of the gaps lives, it holds  $S = \mathbf{R}$  for which  $\mathbf{Q}$  is a conditionally countable order dense subset.

$]s_i, t_i[ = ]s_j, t_j[$  and therefore  $i = j$ . This also shows that  $f$  is a conditional injection, thus  $I$  is at most conditionally countable.  $\square$

**Theorem 5.2 (Debreu's Gap Lemma).** *For every  $S \sqsubseteq \mathbf{R}$  there exists a conditionally strictly increasing function  $g : S \rightarrow \mathbf{R}$  such that all the conditional gaps  $(s, t)$  of  $g(S)$  are of the form*

$$(s, t) = a\{s\} + b]s, t[.$$

This theorem says that there exists a strictly increasing transformation of  $S$  such that any conditional gap which is of the form (5.1) is transformed in a gap which conditionally is either empty, a singleton or an open set. The following argumentation follows the proof idea in [28].

*Proof. Step 1:* According to Lemma 5.1, let  $U = (u_n)_{n \in \mathbf{N}}$  be a conditionally countable order dense subset of  $S$ . We construct a conditionally increasing function  $f : U \rightarrow [0, 1]$ . Let  $H$  be the set of conditional functions  $f : V \rightarrow [0, 1]$ , where  $V = \{u_k : 1 \leq k \leq n\}$ ,  $n \in \mathbf{N}$  or  $V = U$  and such that<sup>16</sup>

$$\begin{cases} f(u_1) = 1/2, \\ f(u_k) = \frac{\sup_{l \leq k-1} \{f(u_l) : u_l < u_k\} + \inf_{l \leq k-1} \{f(u_l) : u_k < u_l\}}{2}, \quad k \geq 2. \end{cases}$$

By definition, any  $f \in H$  is conditionally strictly increasing on its domain and  $H$  is a conditional set. Furthermore  $f : \{u_1\} \rightarrow [0, 1]$  with  $f(u_1) = 1/2$  is an element of  $H$  so that  $H$  lives on 1. We show that there exists a function  $f \in H$  with domain  $U$ . For  $f : V \rightarrow [0, 1]$  and  $g : W \rightarrow [0, 1]$  in  $H$ , define  $f \preceq g$  if  $V \sqsubseteq W$  and  $f = g$  restricted on  $V$ . Let now  $(f_i)$  be a chain in  $H$  and define  $f : V = \sqcup V_i \rightarrow [0, 1]$ ,  $u = \sum a_j u_j \mapsto f(u) = \sum a_j f_j(u_j)$  where  $u_j \in V_j$  for every  $j$  is a well-defined conditional function in  $H$ . Indeed,  $H$  is a conditional set, the  $f_i$  are restrictions of each others and  $V_i \sqsubseteq V_j$  if  $f_i \preceq f_j$ . Hence by Zorn's Lemma there exists a maximal function  $f \in H$ ,  $f : V \rightarrow [0, 1]$ . Next we show that  $V = U$ . For the sake of contradiction, suppose that  $V = \{u_k : 1 \leq k \leq n\}$  for some  $n \in \mathbf{N}$  on some non-trivial condition  $a$ . Without loss of generality, assume that  $a = 1$ . Define  $g : \{u_k : 1 \leq k \leq n+1\} \rightarrow [0, 1]$  by setting  $g = f$  on  $\{u_k : 1 \leq k \leq n\}$  and

$$g(u_{n+1}) = \frac{\sup_{l \leq n} \{f(u_l) : u_l < u_{n+1}\} + \inf_{l \leq n} \{f(u_l) : u_{n+1} < u_l\}}{2}$$

As for those  $n \leq m \leq n+1$ , it follows that  $m = an + a^c(n+1)$  and we set  $g(u_m) = ag(u_n) + a^c g(u_{n+1})$ . By construction  $g : \{u_k : 1 \leq k \leq n+1\} \rightarrow [0, 1]$  is an element of  $H$  which coincides on  $V$  with  $f$ . Since  $g$  is defined on  $V \sqcup \{u_{n+1}\}$  it contradicts the maximality of  $f$ . Thus the domain of the maximal function  $f \in H$  is  $U$ .

*Step 2:* Let  $U = (u_n)$  and  $f : U \rightarrow [0, 1]$  as defined in the previous step. Suppose that  $V, W \sqsubseteq U$  living on 1 satisfy

(a)  $V \sqcup W = U$ ,

<sup>16</sup>With the classical convention that the infimum and supremum over emptyset is equal to 1 and 0, respectively. That is,  $f(u_k) = \inf_{l \leq k-1} f(u_l)/2$  on the condition where  $u_k < u_l$  for every  $l \leq k-1$  and  $f(u_k) = (\sup_{l \leq k-1} f(u_l) + 1)/2$  on the condition where  $u_l < u_k$  for every  $k-1 \leq l$ .

(b)  $V \leq W$ ,<sup>17</sup>

(c) 0 is the unique condition on which  $V$  and  $W$  have both a maximum and a minimum, respectively.

Then

$$\sup_{s \in V} f(s) = \inf_{t \in W} f(t).$$

By (b) it holds  $\sup_V f(s) \leq \inf_W f(t)$ . In order to show the reverse inequality, according to (c), it is sufficient to suppose that  $V$  and  $W$  have both nowhere a maximum and a minimum, respectively since then the gap is the largest. For the sake of contradiction, up to conditioning, suppose that

$$\sup_V f(s) + \varepsilon < \inf_W f(t)$$

for some  $\varepsilon > 0$ . Choose  $s_0 = u_m \in V$  and  $t_0 = u_n \in W$  such that

$$\sup_V f(s) - \varepsilon \leq f(s_0) \leq \sup_V f(s) \quad \text{and} \quad \inf_W f(t) \leq f(t_0) \leq \inf_W f(t) + \varepsilon. \quad (5.2)$$

Since  $V$  has nowhere a maximum, there exists  $u_k \in V$  such that  $u_n < u_k < u_m$  and  $k > n, m$ . Let  $k' = \min\{k > n, m : u_n < u_k < u_m\}$ . By construction of  $f$  and since (a) holds, it follows that

$$f(u_{k'}) = \frac{f(u_n) + f(u_m)}{2} = \frac{f(s_0) + f(t_0)}{2}.$$

Adding both inequalities in (5.2) yields

$$\frac{\sup_V f(s) + \inf_W f(t)}{2} - \frac{\varepsilon}{2} \leq f(u_{k'}) \leq \frac{\sup_V f(s) + \inf_W f(t)}{2} + \frac{\varepsilon}{2},$$

and therefore  $\sup_V f(s) < f(u_{k'}) < \inf_W f(t)$  contradicting  $u_{k'} \in V$ .

*Step 3:* Define  $g : S \rightarrow \mathbf{R}$  by  $g(s) = \sup_{u \in U, u \leq s} f(u)$ . By construction  $g$  is a conditionally strictly increasing extension of  $f$  since  $U$  is a conditionally countable order dense subset of  $S$ . Let  $(s, t)$  be a conditional gap of  $g(S)$  and  $a$  be the maximal condition such that  $s < t$ , that is  $(s, t) = a(s, t) + b\{s\}$ . Without loss of generality, suppose that  $a = 1$  and  $b = 0$ . Define  $V = \{u \in U : f(u) \leq s\}$  and  $W = \{u \in U : f(u) \geq t\}$ . Since  $s < t$  and  $V, W$  satisfy (a) and (b) of the previous step, (c) has to be violated. Hence  $V$  and  $W$  have both a conditional maximum and minimum respectively on some maximal condition  $c > 0$ , that is  $s = f(u_n)$  and  $t = f(u_m)$  on  $c$  for some  $n, m$ . Thus  $cs, ct \in cg(S)$  showing that  $c(s, t) = c]s, t[$ . If  $c < 1$ , we follow the same argumentation but conditioned on  $c^c$  which yields a contradiction with the maximality of  $c$ . Therefore,  $(s, t) = ]s, t[$  which ends the proof.  $\square$

**Theorem 5.3.** *Any numerically representable conditionally upper semi-continuous preference order admits a conditionally upper semi-continuous numerical representation.*

*Proof.* Let  $\tilde{U}$  be a numerical representation of a conditionally upper semicontinuous preference order  $\succsim$ . According to Debreu's Gap Lemma 5.2 there exists a conditional function  $g : Im(\tilde{U}) \rightarrow \mathbf{R}$  such that all the conditional gaps  $(s, t)$  of  $g(Im(\tilde{U}))$  are of the following form

$$(s, t) = as + b]s, t[, \quad \text{for } s < t.$$

<sup>17</sup>In the sense that for all  $s \in V$  and for all  $t \in W$  it holds  $s \leq t$ .



Since  $g$  is strictly increasing, it follows that  $U = g \circ \tilde{U}$  is a conditional numerical representation of  $\succsim$  as well. Clearly  $Im(U) = g(Im(\tilde{U}))$ . In order to verify the upper semi-continuity, pick  $m \in \mathbf{R}$ . We distinguish between the following cases:

- If  $m = U(y)$ , then  $\{x \in X : U(x) \geq m\} = \{x \in X : U(x) \geq U(y)\} = \{x \in X : x \succsim y\}$  which is conditionally closed by assumption.
- If  $m \in ]s, t[$  where  $]s, t[$  is a conditional gap of  $Im(U)$ , then  $t = U(y)$  for some  $y \in X$ , and thus  $\{x \in X : U(x) \geq m\} = \{x \in X : U(x) \geq t\} = \{x \in X : U(x) \geq U(y)\} = \{x \in X : x \succsim y\}$  which is also conditionally closed by assumption.
- If  $m = s$  where  $\{s\}$  is a conditional gap of  $Im(U)$ , then let  $(s_n) = (U(y_n)) \sqsubseteq Im(U)$  be a conditional sequence such that  $s_n \nearrow s$ . It holds  $\{x \in X : U(x) \geq s\} = \bigcap_n \{x \in X : U(x) \geq s_n\} = \bigcap_n \{x \in X : U(x) \geq U(y_n)\} = \bigcap_n \{x \in X : x \succsim y_n\}$  which is conditionally closed as the conditional intersection of closed sets.

Since  $\mathbf{R} = Im(U) \sqcup [Im(U)]^\square$  and  $[Im(U)]^\square$  is made of gaps of the form  $(s, t) = a\{s\} + b]s, t[$ , it follows that any  $m \in \mathbf{R}$  belongs conditionally to one of the previous three cases. Thus  $U$  is conditionally upper semi-continuous.  $\square$

## References

- [1] B. Acciaio, H. Föllmer, and I. Penner. Risk assessment for uncertain cash flows: Model ambiguity, discounting ambiguity, and the role of bubbles. *Forthcoming in Finance and Stochastics*, 2011.
- [2] R. J. Aumann. Utility theory without the completeness axiom. *Econometrica*, 30(3):445–462, 1962.
- [3] T. F. Bewley. Knightian decision theory. Part I. *Decisions in Economics and Finance*, 2001.
- [4] T. R. Bielecki, I. Cialenco, S. Drapeau, and M. Karliczek. Dynamic assessment indices. *Preprint*, 2014.
- [5] P. Cheridito and M. Kupper. Recursivity of indifference prices and translation-invariant preferences. *Mathematics and Financial Economics*, 2:173–188, 2009.
- [6] P. Cheridito, F. Delbaen, and M. Kupper. Dynamic monetary risk measures for bounded discrete-time processes. *Electronic Journal of Probability*, 11(3):57–106, 2006.
- [7] G. Debreu. Representation of a preference ordering by a numerical function. In C. C. Thrall, R.M. and R. Davis, editors, *Decision Process*, pages 159–165. John Wiley, New York, 1954.
- [8] G. Debreu. Continuity properties of Paretian utility. *International Economic Review*, 5(3):285–293, 1964.
- [9] K. Detlefsen and G. Scandolo. Conditional and dynamic convex risk measures. *Finance and Stochastics*, 9: 539–561, 2005.
- [10] S. Drapeau, A. Jamneshan, M. Karliczek, and M. Kupper. The algebra of conditional sets and the concepts of conditional topology and compactness. *Preprint*, 2013.
- [11] J. Dubra, F. Maccheroni, and E. A. Ok. Expected utility theory without the completeness axiom. *Journal of Economic Theory*, 115(1):118–133, 2004.
- [12] D. Duffie and L. G. Epstein. Stochastic differential utility. *Econometrica*, 60(2):353–94, 1992.
- [13] K. Eliaz and E. A. Ok. Indifference or indecisiveness? Choice-theoretic foundations of incomplete preferences. *Games and Economic Behavior*, 56(1):61 – 86, 2006.
- [14] L. G. Epstein and S. E. Zin. Substitution, risk aversion, and the temporal behavior of consumption and asset returns: A theoretical framework. *Econometrica*, 57(4):937–69, 1989.
- [15] O. Evren and E. A. Ok. On the multi-utility representation of preference relations. *Journal of Mathematical Economics*, 47(4-5):554–563, 2011.

- [16] D. Filipovic, M. Kupper, and N. Vogelpoth. Separation and duality in locally  $L^0$ -convex modules. *Journal of Functional Analysis*, 256:3996–4029, 2009.
- [17] D. Filipovic, M. Kupper, and N. Vogelpoth. Approaches to conditional risk. *SIAM Journal of Financial Mathematics*, 2011.
- [18] H. Föllmer and A. Schied. *Stochastic Finance. An Introduction in Discrete Time*. de Gruyter Studies in Mathematics. Walter de Gruyter, Berlin, New York, 2 edition, 2004.
- [19] M. Frittelli and M. Maggis. Conditional certainty equivalent. *International Journal of Theoretical and Applied Finance*, 14(1):41–59, 2011.
- [20] S. Givant and P. Halmos. *Introduction to Boolean algebras*. Springer, 2009.
- [21] A. Jamneshan, M. Kupper, and M. Streckfuss. Conditional Lebesgue integration. *In preparation*, 2015.
- [22] E. Karni. Subjective expected utility theory with state-dependent preferences. *Journal of Economic Theory*, 60(2):428–438, 1993.
- [23] E. Karni. A definition of subjective probabilities with state-dependent preferences. *Econometrica*, 61(1):pp. 187–198, 1993.
- [24] D. M. Kreps and E. L. Porteus. Temporal resolution of uncertainty and dynamic choice theory. *Econometrica*, 46(1):185–200, 1978.
- [25] D. M. Kreps and E. L. Porteus. Temporal von Neumann-Morgenstern and induced preferences. *Journal of Economic Theory*, 20(1):81–109, 1979.
- [26] R. D. Luce and D. H. Krantz. Conditional expected utility. *Econometrica*, 39(2):pp. 253–271, 1971.
- [27] F. Maccheroni, M. Marinacci, and A. Rustichini. Dynamic variational preferences. *Journal of Economic Theory*, 127(1):4–44, 2006.
- [28] P. Ouwehand. A simple proof of Debreu’s Gap Lemma. *ORiON*, 2010.
- [29] B. Peleg. Utility functions for partially ordered topological spaces. *Econometrica*, 38(1):93–96, 1970.
- [30] T. Rader. The existence of a utility function to represent preferences. *The Review of Economic Studies*, 30(3): 229–232, 1963.
- [31] M. K. Richter. Revealed preference theory. *Econometrica*, 34(3):635–645, 1966.
- [32] A. Robert. Letter from Robert Aumann to Leonard Savage. In J. Drèze, editor, *Essays on Economic Decisions under Uncertainty*. Cambridge University Press, 1987.
- [33] C. Skiadas. Conditioning and aggregation of preferences. *Econometrica*, 65(2):347–368, 1997.
- [34] C. Skiadas. Subjective probability under additive aggregation of conditional preferences. *Journal of Economic Theory*, 76(2):242–271, 1997.
- [35] J. von Neumann and O. Morgenstern. *Theory of Games and Economics Behavior*. Princeton University Press, 2nd edition, 1947.
- [36] P. Wakker. Subjective probabilities for state dependent continuous utility. *Mathematical Social Sciences*, 14(3):289–298, 1987.