

# SINGULAR TWISTED SUMS GENERATED BY COMPLEX INTERPOLATION

JESUS M. F. CASTILLO, VALENTIN FERENCZI AND MANUEL GONZÁLEZ

ABSTRACT. We present new methods to obtain singular twisted sums  $X \oplus_{\Omega} X$  (i.e., exact sequences  $0 \rightarrow X \rightarrow X \oplus_{\Omega} X \rightarrow X \rightarrow 0$  in which the quotient map is strictly singular), in which  $X$  is the interpolation space arising from a complex interpolation scheme and  $\Omega$  is the induced centralizer.

Although our methods are quite general, in our applications we are mainly concerned with the choice of  $X$  as either a Hilbert space, or Ferenczi's uniformly convex Hereditarily Indecomposable space. In the first case, we construct new singular twisted Hilbert spaces, including the only known example so far: the Kalton-Peck space  $Z_2$ . In the second case we obtain the first example of an H.I. twisted sum of an H.I. space. We then use Rochberg's description of iterated twisted sums to show that there is a sequence  $\mathcal{F}_n$  of H.I. spaces so that  $\mathcal{F}_{m+n}$  is a singular twisted sum of  $\mathcal{F}_m$  and  $\mathcal{F}_n$ , while for  $l > n$  the direct sum  $\mathcal{F}_n \oplus \mathcal{F}_{l+m}$  is a nontrivial twisted sum of  $\mathcal{F}_l$  and  $\mathcal{F}_{m+n}$ .

We also introduce and study the notion of disjoint singular twisted sum of Köthe function spaces and construct several examples involving reflexive  $p$ -convex Köthe function spaces, which include the function version of the Kalton-Peck space  $Z_2$ .

## 1. INTRODUCTION

For all unexplained notation see Sections 2 (background on exact sequences and quasi-linear maps) and 3 (background and preliminary results on complex interpolation and centralizers).

This paper focuses on the study of the existence and properties of exact sequences

$$(1) \quad 0 \longrightarrow X \xrightarrow{j} E \xrightarrow{q} X \longrightarrow 0,$$

in which the Banach space  $X$  has been obtained by complex interpolation. The exact sequence is called *nontrivial* when  $j(X)$  is not complemented in the middle space  $E$ , which is then called a (nontrivial) twisted sum of  $X$  (or a twisting of  $X$ , or even a twisted  $X$ ). The exact sequence is called *singular* (and  $E$  is called a *singular twisted sum*) when the operator  $q$  is strictly singular. The key example on which all the theory is modeled is the Kalton-Peck twisted Hilbert space  $Z_2$  obtained in [32], which provides the first and only known singular sequence

$$0 \longrightarrow \ell_2 \xrightarrow{j} Z_2 \xrightarrow{q} \ell_2 \longrightarrow 0.$$

Singular sequences correspond to twisted sums which are, in some sense, as far as possible from being direct sums. For example, in Kalton-Peck example  $Z_2$ , the natural copy of  $\ell_2$  does not even admit a "relative" summand, i.e. there is no infinite dimensional subspace  $Z$  of  $Z_2$  forming a topological direct sum  $j(\ell_2) \oplus Z$  inside  $Z_2$ .

In [26] Kalton showed that exact sequences (1) are in correspondence with certain non-linear maps  $F : X \rightarrow X$ , called quasi-linear maps. So, twisted sum spaces, and in particular

---

This research was supported by Project MTM2013-45643, D.G.I. Spain. The second author was supported by Fapesp project 2013/11390-4, including for visits of the first and third author to the University of São Paulo.

exact sequences, can be written in the form

$$(2) \quad 0 \longrightarrow X \longrightarrow X \oplus_F X \longrightarrow X \longrightarrow 0.$$

Following [10, 14], we say that a quasi-linear map  $F$  is *singular* if the associated exact sequence (2) is singular. In [32] Kalton and Peck presented a method to show an explicit construction of quasi-linear maps on Banach spaces with unconditional basis. This method was refined by Kalton [28] and extended to Köthe function spaces. The special type of quasi-linear maps obtained by this method were called *centralizers*. The main examples are the so called *Kalton-Peck maps*:

$$\mathcal{K}_\phi(x) = x\phi\left(-\log\frac{|x|}{\|x\|}\right)$$

where  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  is a certain Lipschitz map. The choice of the function  $\phi_r(t) = t^r$  (when  $t \geq 1$ ), and  $\phi_r(t) = t$  (when  $0 \leq t \leq 1$ ); with  $0 < r \leq 1$  will have a especial interest for us. We simply write  $\mathcal{K}$  for Kalton-Peck space instead of  $\mathcal{K}_{\phi_1}$ . In [32] it is shown that  $\mathcal{K}$  is singular on  $\ell_p$  spaces for  $1 < p < \infty$ ; in [14] for  $p = 1$ ; and in [10] for the whole range  $0 < p < \infty$ . It was soon observed that the Kalton-Peck map  $\mathcal{K}$  on  $\ell_2$  could be generated from the interpolation scale of  $\ell_p$  spaces. Taking this as starting point, Kalton unfolds in [28, 29] the existence of a correspondence between centralizers defined on Köthe function spaces and interpolation scales of Köthe function spaces. This opens the door to the possibility of obtaining nontrivial quasi-linear maps in Banach spaces generated by an interpolation scale, even when no unconditional structure is present.

Such is the point of view we adopt in this paper to tackle the study of singular centralizers and singular quasi-linear maps on Banach spaces obtained by complex interpolation. In the case of centralizers this leads us to obtain new singularity results for Kalton-Peck sums of sequence spaces as well as of function spaces; and, in particular, new singular twisted Hilbert spaces. We introduce a new concept of singularity, that we call *disjoint singularity*, which is relevant to the study of interpolation schemes of function spaces. In the case of general quasi-linear maps, not just centralizers, we “localize” the techniques developed and apply them to spaces with monotone basis and obtain the first H.I. twisted sum of an H.I. space.

A description of the contents of the paper is in order: after this introduction and a preliminary Section 2 on basic facts about exact sequences and quasi-linear maps, Section 3 takes root in Kalton’s work and so it contains an analysis of centralizers arising from an interpolation scheme; the analysis is centered on an interpolation couple  $(X_0, X_1)$  and the centralizer  $\Omega_\theta$  obtained at the interpolation space  $X_\theta = (X_0, X_1)_\theta$ ; although the results extend (see subsection 5.4) to cover the case of a measurable family of spaces. We observe, and derive a few consequences from it, the fact that such centralizers admit an overall form as  $\Omega_\theta(x) = x \log \frac{a_0(x)}{a_1(x)}$ , where  $a_0(x)^{1-\theta} a_1(x)^\theta$  is a Lozanovskii factorization of  $|x|$  with respect to the couple  $(X_0, X_1)$ . Section 4 contains the two fundamental estimates we use through the paper: Lemma 4.3 (estimate for non-singular maps) and Lemma 4.5 (general estimate for centralizers arising from an interpolation scheme). Section 5 contains several criteria for singularity based on the previous two lemmata. The first two subsections treat the Banach lattice case: we recover the singularity of Kalton-Peck maps associated to the interpolation scale of  $\ell_p$  spaces, as a particular case of a general criterion for singularity, and we prove the disjoint singularity of Kalton-Peck maps associated to the interpolation scale of  $L_p$  spaces, for which it was known that they were not singular. We also prove the disjoint singularity of Kalton-Peck maps on more general  $p$ -convex Köthe spaces, by means of the interpolation

formula  $X = (L_\infty, X^{(p)})_{1/p}$ . In the third subsection, we give conditions implying the singularity in the conditional case (spaces admitting a basis not necessarily unconditional) which will be needed to cover the case of H.I. spaces. In Section 6 we obtain new singular twisted Hilbert spaces; we also complete previous results by showing that centralizers  $\mathcal{K}_\phi$  is singular under rather mild conditions on  $\phi$ , satisfied in particular by the complex versions [30] of  $\mathcal{K}$ . In Section 7 we connect the results about singular sequences with the twisting of H.I. spaces: a twisted sum of two H.I. spaces is H.I. if and only if it is singular. One of the difficulties for such construction is, as we show, that a nontrivial twisted sum of two H.I. spaces can be decomposable; note that it was known [23, Theorem 1] that such twisted sums should be at most 2-decomposable. Section 8 applies the previous techniques to the quasi-linear map associated to the construction of Ferenczi's H.I. space  $\mathcal{F}$  [21] by complex interpolation of a suitable family of Banach spaces. In Section 9 we complete and improve the results in Sections 7 and 8 by showing new natural H.I. and decomposable twistings; precisely, that there is a sequence  $(\mathcal{F}_n)$  of H.I. spaces so that:

- (i) For each  $m, n \geq 1$  there is a singular exact sequence

$$0 \longrightarrow \mathcal{F}_m \longrightarrow \mathcal{F}_{m+n} \longrightarrow \mathcal{F}_n \longrightarrow 0.$$

- (ii) For each  $l, m, n \geq 1$  with  $l > n$  there is a nontrivial exact sequence

$$0 \longrightarrow \mathcal{F}_l \longrightarrow \mathcal{F}_n \oplus \mathcal{F}_{l+m} \longrightarrow \mathcal{F}_{m+n} \longrightarrow 0.$$

## 2. EXACT SEQUENCES, TWISTED SUMS AND QUASI-LINEAR MAPS

A *twisted sum* of two Banach spaces  $Y$  and  $Z$  is a space  $X$  which has a subspace  $M$  isomorphic to  $Y$  with the quotient  $X/M$  isomorphic to  $Z$ . The space  $X$  is a quasi-Banach space in general [32]. Recall that a Banach space is *B-convex* when it does not contain  $\ell_1^n$  uniformly. Theorem 2.6 of [26] implies that a twisted sum of two B-convex Banach spaces is isomorphic to a Banach space.

An exact sequence  $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$ , where  $Y, Z$  are Banach spaces and the arrows are (bounded) operators is a diagram in which the kernel of each arrow coincides with the image of the preceding one. By the open mapping theorem this means that the middle space  $X$  is a twisted sum of  $Y$  and  $Z$ .

Two exact sequences  $0 \rightarrow Y \rightarrow X_1 \rightarrow Z \rightarrow 0$  and  $0 \rightarrow Y \rightarrow X_2 \rightarrow Z \rightarrow 0$  are *equivalent* if there exists an operator  $T : X_1 \rightarrow X_2$  such that the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & Y & \xrightarrow{i} & X_1 & \xrightarrow{q} & Z \longrightarrow 0 \\ & & \parallel & & \downarrow T & & \parallel \\ 0 & \longrightarrow & Y & \xrightarrow{j} & X_2 & \xrightarrow{p} & Z \longrightarrow 0 \end{array}$$

The classical 3-lemma (see [13, p. 3]) shows that  $T$  must be an isomorphism. An exact sequence is trivial if and only if it is equivalent to  $0 \rightarrow Y \rightarrow Y \times Z \rightarrow Z \rightarrow 0$ , where  $Y \times Z$  is endowed with the product norm. In this case we say that the exact sequence *splits*.

A map  $F : Z \rightarrow Y$  is called *quasi-linear* if it is homogeneous and there is a constant  $M$  such that  $\|F(u+v) - F(u) - F(v)\| \leq M\|u+v\|$  for all  $u, v \in Z$ . There is a correspondence (see [13, Theorem 1.5.c, Section 1.6]) between exact sequences  $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$  of Banach spaces and a special kind of quasi-linear maps  $\omega : Z \rightarrow Y$ , called *z-linear maps*, which satisfy  $\|\omega(\sum_{i=1}^n u_i) - \sum_{i=1}^n \omega(u_i)\| \leq M \sum_{i=1}^n \|u_i\|$  for all finite sets  $u_1, \dots, u_n \in Z$ . A quasi-linear map  $F : Z \rightarrow Y$  induces the exact sequence  $0 \rightarrow Y \xrightarrow{j} Y \oplus_F Z \xrightarrow{p} Z \rightarrow 0$  in which  $Y \oplus_F Z$

denotes the vector space  $Y \times Z$  endowed with the quasi-norm  $\|(y, z)\|_F = \|y - F(z)\| + \|z\|$ . The embedding is  $j(y) = (y, 0)$  while the quotient map is  $p(y, z) = z$ . When  $F$  is  $z$ -linear, this quasi-norm is equivalent to a norm [13, Chapter 1]. On the other hand, the process to obtain a  $z$ -linear map out from an exact sequence  $0 \rightarrow Y \xrightarrow{i} X \xrightarrow{q} Z \rightarrow 0$  of Banach spaces is the following one: get a homogeneous bounded selection  $b : Z \rightarrow X$  for the quotient map  $q$ , and then a linear  $\ell : Z \rightarrow X$  selection for the quotient map. Then  $\omega = i^{-1}(b - \ell)$  is a  $z$ -linear map from  $Z$  to  $Y$ . The commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & Y & \xrightarrow{i} & X & \xrightarrow{q} & Z & \longrightarrow & 0 \\ & & \parallel & & \downarrow T & & \parallel & & \\ 0 & \longrightarrow & Y & \xrightarrow{j} & Y \oplus_{\omega} Z & \xrightarrow{p} & Z & \longrightarrow & 0 \end{array}$$

obtained by taking as  $T : X \rightarrow Y \oplus_{\omega} Z$  the operator  $T(x) = (x - \ell qx, qx)$  shows that the upper and lower exact sequences are equivalent. Two quasi-linear maps  $F, F' : Z \rightarrow Y$  are said to be equivalent, denoted  $F \equiv G$ , if the difference  $F - F'$  can be written as  $B + L$ , where  $B : Z \rightarrow Y$  is a homogeneous bounded map (not necessarily linear) and  $L : Z \rightarrow Y$  is a linear map (not necessarily bounded). Of course two quasi-linear maps are equivalent if and only if the associated exact sequences are equivalent. Thus, two exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & Y & \longrightarrow & Y \oplus_{\Omega} Z & \longrightarrow & Z & \longrightarrow & 0 \\ & & \parallel & & & & \parallel & & \\ 0 & \longrightarrow & Y & \longrightarrow & Y \oplus_{\Psi} Z & \longrightarrow & Z & \longrightarrow & 0 \end{array}$$

(or two quasi-linear maps  $\Omega, \Psi$ ) are equivalent (i.e.,  $\Omega \equiv \Psi$ ) if there exists a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & Y & \longrightarrow & Y \oplus_{\Omega} Z & \longrightarrow & Z & \longrightarrow & 0 \\ & & \alpha \downarrow & & \beta \downarrow & & \downarrow \gamma & & \\ 0 & \longrightarrow & Y & \longrightarrow & Y \oplus_{\Psi} Z & \longrightarrow & Z & \longrightarrow & 0 \end{array}$$

with  $\alpha = id_Y$  and  $\gamma = id_Z$ . Imposing other conditions on the maps  $\alpha, \beta, \gamma$  yields other notions of equivalence appearing in the literature. From the most restrictive to the more general they are:

- (1) Bounded equivalence [28, 29] (see also Section 3 below): asking that  $\Omega - \Psi$  is bounded.
- (2) Projective equivalence [32]: asking  $\alpha, \gamma$  to be scalar multiples of the identity. Equivalently,  $\Omega \equiv \mu\Psi$  for some scalar  $\mu$ .
- (3) We will need in this paper “permutative projective equivalence”: when  $Y$  and  $Z$  have unconditional bases  $(e_n)$ , asking  $T_{\sigma}\Omega \equiv \mu\Psi T_{\sigma}$  for some scalar  $\mu$  and some operator  $T_{\sigma}(\sum_i x_i e_i) = \sum_i x_i e_{\sigma(i)}$  induced by a permutation  $\sigma$  of the integers. When  $\mu = 1$  we will just say that  $\Omega$  and  $\Lambda$  are permutatively equivalent.
- (4) Isomorphic equivalence [7, 15]: asking  $\alpha, \beta, \gamma$  to be isomorphisms. In quasi-linear terms, this means that  $\alpha\Omega \equiv \Psi\gamma$ .

Obviously, equivalence takes place between (1) and (2). In conclusion, each of (1), (2), (3), (4) yields a “natural” isomorphism  $\beta$  between  $Y \oplus_{\Omega} Z$  and  $Y \oplus_{\Psi} Z$  of a specific form prescribed by the forms of the maps  $\alpha$  and  $\gamma$ .

A few facts about the connections between quasi-linear maps and the associated exact sequences will be needed in this paper, and can be explicitly found in [16, Section 1]. Given

an exact sequence  $0 \rightarrow Y \rightarrow X \rightarrow Z \rightarrow 0$  with associated quasi-linear map  $F$  and an operator  $\alpha : Y \rightarrow Y'$ , there is a commutative diagram

$$(3) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & Y & \xrightarrow{i} & X & \xrightarrow{q} & Z & \longrightarrow & 0 \\ & & \alpha \downarrow & & T \downarrow & & \parallel & & \\ 0 & \longrightarrow & Y' & \xrightarrow{i'} & \text{PO} & \xrightarrow{q'} & Z & \longrightarrow & 0. \end{array}$$

The lower sequence is called the *push-out sequence*, its associated quasi-linear map is equivalent to  $\alpha \circ F$ , and the space PO is called the *push-out space*. When  $F$  is  $z$ -linear, so is  $\alpha \circ F$ . Given a commutative diagram like (3) the *diagonal push-out sequence* is the exact sequence generated by the quasi-linear map  $F \circ q'$ , and is equivalent to the exact sequence

$$0 \longrightarrow Y \xrightarrow{d} Y' \oplus X \xrightarrow{m} X' \longrightarrow 0$$

where  $d(y) = (-\alpha y, iy)$  and  $m(y', x) = i'y' + Tx$ .

### 3. COMPLEX INTERPOLATION AND CENTRALIZERS

Here we explain the connections between complex interpolation, twisted sums and quasi-linear maps that we use throughout the paper.

**3.1. Complex interpolation and twisted sums.** We describe the complex interpolation method for a pair of spaces following [5]. Other general references are [17, 29, 31, 36].

Let  $\mathbb{S}$  denote the closed strip  $\{z \in \mathbb{C} : 0 \leq \text{Re}(z) \leq 1\}$  in the complex plane, and let  $\mathbb{S}^\circ$  be its interior and  $\partial\mathbb{S}$  be its boundary. Given an admissible pair  $(X_0, X_1)$  of complex Banach spaces, we denote by  $\mathcal{H} = \mathcal{H}(X_0, X_1)$  the space of functions  $g : \mathbb{S} \rightarrow \Sigma := X_0 + X_1$  satisfying the following conditions:

- (1)  $g$  is  $\|\cdot\|_\Sigma$ -bounded and  $\|\cdot\|_\Sigma$ -continuous on  $\mathbb{S}$ , and  $\|\cdot\|_\Sigma$ -analytic on  $\mathbb{S}^\circ$ ;
- (2)  $g(it) \in X_0$  for each  $t \in \mathbb{R}$ , and the map  $t \in \mathbb{R} \mapsto g(it) \in X_0$  is bounded and continuous;
- (3)  $g(it+1) \in X_1$  for each  $t \in \mathbb{R}$ , and the map  $t \in \mathbb{R} \mapsto g(it+1) \in X_1$  is bounded and continuous;

The space  $\mathcal{H}$  is a Banach space under the norm  $\|g\|_{\mathcal{H}} = \sup\{\|g(j+it)\|_j : j = 0, 1; t \in \mathbb{R}\}$ . For  $\theta \in [0, 1]$ , define the interpolation space

$$X_\theta = (X_0, X_1)_\theta = \{x \in \Sigma : x = g(\theta) \text{ for some } g \in \mathcal{H}\}$$

with the norm  $\|x\|_\theta = \inf\{\|g\|_{\mathcal{H}} : x = g(\theta)\}$ . So  $(X_0, X_1)_\theta$  is the quotient of  $\mathcal{H}$  by  $\ker \delta_\theta$ , and thus it is a Banach space.

For  $0 < \theta < 1$ , we will consider the maps  $\delta_\theta^n : \mathcal{H} \rightarrow \Sigma$  –evaluation of the  $n^{\text{th}}$ -derivative at  $\theta$ – that appear in Schechter's version of the complex method of interpolation [37]. Note that  $\delta_\theta \equiv \delta_\theta^0$  is bounded by the definition of  $\mathcal{H}$ , and this fact and the Cauchy integral formula imply the boundedness of  $\delta_\theta^n$  for  $n \geq 1$  (see also [9]). We will also need the following result (see [12, Theorem 4.1]):

**Lemma 3.1.**  $\delta'_\theta : \ker \delta_\theta \rightarrow X_\theta$  is bounded and onto for  $0 < \theta < 1$ .

For future use, note that given  $G \in \ker \delta_\theta$ , the function  $H$  defined by  $H(z) = G(z)/(z - \theta)$  belongs to  $\mathcal{H}$  and satisfies  $\delta'_\theta(G) = H(\theta)$ , which implies the estimate

$$\|\delta'_\theta|_{\ker \delta_\theta}\| \leq \text{dist}(\theta, \partial\mathbb{S})^{-1}.$$

Lemma 3.1 provides the connection with exact sequences and twisted sums through the following push-out diagram:

$$(4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \ker \delta_\theta & \xrightarrow{i_\theta} & \mathcal{H} & \xrightarrow{\delta_\theta} & X_\theta \longrightarrow 0 \\ & & \delta'_\theta \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & X_\theta & \longrightarrow & \text{PO} & \longrightarrow & X_\theta \longrightarrow 0 \end{array}$$

whose lower row is obviously a twisted sum of  $X_\theta$ .

Apart from the obvious description as a push-out space, PO can be represented as:

(1) *A twisted sum space.* Let  $B_\theta : X_\theta \rightarrow \mathcal{H}$  be a bounded homogeneous selection for  $\delta_\theta$ , and let  $L_\theta : X_\theta \rightarrow \mathcal{H}$  be a linear selection. The map  $\omega_\theta = B_\theta - L_\theta : X_\theta \rightarrow \ker \delta_\theta$  is an associated quasi-linear map for the upper sequence in diagram (4). The lower push-out sequence then comes defined by the quasi-linear map  $\delta'_\theta \omega_\theta$ . Hence,  $\text{PO} \simeq X_\theta \oplus_{\delta'_\theta \omega_\theta} X_\theta$ .

(2) *A derived space.* With the same notation as above, set

$$d_{\delta'_\theta B_\theta}(X_\theta) = \{(y, z) \in \Sigma \times X_\theta : y - \delta'_\theta B_\theta z \in X_\theta\}$$

endowed with the quasi-norm  $\|(y, z)\|_d = \|y - \delta'_\theta B_\theta z\|_{X_\theta} + \|z\|_{X_\theta}$ . This is a twisted sum of  $X_\theta$  since the embedding  $y \rightarrow (y, 0)$  and quotient map  $(y, z) \rightarrow z$  yield an exact sequence

$$0 \longrightarrow X_\theta \longrightarrow d_{\delta'_\theta B_\theta}(X_\theta) \longrightarrow X_\theta \longrightarrow 0.$$

Moreover, the two exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & X_\theta & \longrightarrow & X_\theta \oplus_{\delta'_\theta \omega_\theta} X_\theta & \longrightarrow & X_\theta \longrightarrow 0 \\ & & \parallel & & \downarrow T & & \parallel \\ 0 & \longrightarrow & X_\theta & \longrightarrow & d_{\delta'_\theta B_\theta}(X_\theta) & \longrightarrow & X_\theta \longrightarrow 0. \end{array}$$

are isometrically equivalent via the isometry  $T(y, z) = (y + \delta'_\theta L_\theta z, z)$ .

Thus, we can pretend that the quasi-linear map associated to the push-out sequence is  $\delta'_\theta B_\theta$ , usually more intuitive than the true quasi-linear map  $\delta'_\theta(B_\theta - L_\theta)$ . Such map has been sometimes called “the  $\Omega$ -operator”. Needless to say, the  $\Omega$ -operator depends on the choice of  $B_\theta$ . However the difference between two associated  $\Omega$ -operators must be bounded:

$$\|\delta'_\theta(\tilde{B}_\theta - B_\theta)x\|_{X_\theta} \leq \|\delta'_\theta|_{\ker \delta_\theta}\|(\|\tilde{B}_\theta\| + \|B_\theta\|)\|x\|_{X_\theta}.$$

The derived space admits the following useful representation, see [36, p.323] for an embryonic finite-dimensional version; also quoted in [18, p.218]; see [12, Prop.7.1] for a general version involving two compatible interpolators, and [9] for a rather complete exposition, variations and applications of that representation.

**Proposition 3.2.**  $d_{\delta'_\theta B_\theta}(X_\theta) = \{(f'(\theta), f(\theta)) : f \in \mathcal{H}\}$ , and the quotient norm of  $\mathcal{H}/(\ker \delta_\theta \cap \ker \delta'_\theta)$  is equivalent to the quasi-norm  $\|(\cdot, \cdot)\|_d$ .

*Proof.* Given  $f \in \mathcal{H}$ , since  $f - B_\theta(f(\theta)) \in \ker \delta_\theta$ , by Lemma 3.1 one has

$$f'(\theta) - \delta'_\theta B_\theta(f(\theta)) = \delta'_\theta(f - B_\theta(f(\theta))) \in X_\theta,$$

hence  $(f'(\theta), f(\theta)) \in d_{\delta'_\theta B_\theta}(X_\theta)$ .

Conversely, let  $(y, z) \in d_{\delta'_\theta B_\theta}(X)$ . We have  $z \in X_\theta$ , so  $B_\theta z \in \mathcal{H}$ . Since  $y - \delta'_\theta B_\theta z \in X_\theta$ , there exists  $g \in \ker \delta_\theta$  such that  $y - \delta'_\theta B_\theta z = g'(\theta)$ . Thus taking  $f = B_\theta z + g$  we have  $f(\theta) = z$  and  $f'(\theta) = y$ , and the equality is proved.

For the equivalence, given  $(y, z) \in d_{\delta'_\theta B_\theta}(X)$ , take  $f \in \mathcal{H}$  with  $\|f\| \leq 2 \operatorname{dist}(f, \ker \delta_\theta \cap \ker \delta'_\theta)$  such that  $y = f'(\theta)$  and  $z = f(\theta)$ . Then  $\|z\|_{X_\theta} = \operatorname{dist}(f, \ker \delta_\theta)$  and

$$\|y - \delta'_\theta B_\theta z\|_{X_\theta} = \|\delta'_\theta(f - B_\theta z)\|_{X_\theta}.$$

Since  $f - B_\theta z \in \ker \delta_\theta$ , we get

$$\|(y, z)\|_d \leq \|\delta'_\theta|_{\ker \delta_\theta}\| (1 + \|B_\theta\|) \|f\| + \|f\| \leq 2(\|\delta'_\theta|_{\ker \delta_\theta}\| (1 + \|B_\theta\|) + 1) \operatorname{dist}(f, \ker \delta_\theta \cap \ker \delta'_\theta),$$

and there exists a constant  $C$  so that  $\operatorname{dist}(f, \ker \delta_\theta \cap \ker \delta'_\theta) \leq C\|(y, z)\|_d$  by the open-mapping theorem.  $\square$

The results mentioned so far remain valid in the wider context of the general method of interpolation considered in [31, Section 10]. In Section 8 we will need to work with the complex interpolation method associated to a family  $(X_{(0,t)}, X_{(1,t)})_{t \in \mathbb{R}}$  of complex Banach spaces as described in [17]; which is a special case of the general method mentioned above.

**3.2. Centralizers.** Here we consider Köthe function spaces  $X$  over a measure space  $(\Sigma, \mu)$  with their  $L_\infty$ -module structure. As a particular case, we have Banach spaces with a 1-unconditional basis with their associated  $\ell_\infty$ -structure. We denote by  $L_0$  the space of all  $\mu$ -measurable functions, and given  $g \in L_0$ , we understand that  $\|g\|_X < \infty$  implies  $g \in X$ .

**Definition 1.** A centralizer on a Köthe function space  $X$  is a homogeneous map  $\Omega : X \curvearrowright L_0$  such that  $\|\Omega(ax) - a\Omega(x)\|_X \leq C\|x\|_X\|a\|_\infty$  for all  $a \in L_\infty$  and  $x \in X$ .

A centralizer on  $X$  will be denoted by  $\Omega : X \curvearrowright X$ . We use this notation to stress the fact that a centralizer on  $X$  is not a map  $X \rightarrow X$ , but only a map  $X \rightarrow L_0$  so that the differences  $\Omega(ax) - a\Omega(x)$  belong to  $X$ . This notion coincides with that of Kalton's "strong centralizer" introduced in [28]. Centralizers arise naturally in a complex interpolation scheme in which the interpolation scale of spaces share a common  $L_\infty$ -module structure: in such case, the space  $\mathcal{H}$  also enjoys the same  $L_\infty$ -module structure in the form  $(u \cdot f)(z) = u \cdot f(z)$ . In this way, the fundamental sequence of the interpolation scheme  $0 \rightarrow \ker \delta_\theta \rightarrow \mathcal{H} \rightarrow X_\theta \rightarrow 0$  is an exact sequence in the category of  $L_\infty$ -modules. In an interpolation scheme starting with a couple  $(X_0, X_1)$  of Köthe function spaces, the map  $\Omega_\theta = \delta_\theta' B_\theta$  is a centralizer on  $X_\theta$ .

For a centralizer  $\Omega : X \curvearrowright X$  on a Köthe function space  $X$ , it was proved in [28, Lemma 4.2] that there exists  $M > 0$  such that  $\|\Omega(u + v) - \Omega(u) - \Omega(v)\|_X \leq M(\|u\|_X + \|v\|_X)$ . So we can assume that  $\Omega$  is a quasi-linear map. The smallest of the constants  $M$  satisfying the above inequality is denoted  $\rho(\Omega)$ . Note that  $\Omega : X \curvearrowright X$  induces an exact sequence in the category of (quasi-)Banach  $L_\infty$ -modules  $0 \rightarrow X \rightarrow d_\Omega(X) \rightarrow X \rightarrow 0$ , where

$$d_\Omega(X) = \{(w, z) \in L_0 \times X : w - \Omega z \in X\}$$

endowed with the quasi-norm  $\|(w, z)\|_\Omega = \|w - \Omega z\|_X + \|z\|_X$ ; with embedding  $y \rightarrow (y, 0)$  and quotient map  $(w, z) \rightarrow z$ . The derived space  $d_\Omega(X)$  admits a  $L_\infty$ -module structure defined by  $a(w, z) = (aw, az)$ . Kalton proved in [28, Section 4] that every self-extension of a Köthe function space  $X$  is (equivalent to) the extension induced by a centralizer on  $X$ . Sometimes we will take the restriction of  $\Omega$  to a closed subspace  $Y$  of  $X$ , and consider  $d_\Omega(X, Y)$  defined in the same way as a subspace of  $L_0 \times Y$ .

A centralizer  $\Omega : X \curvearrowright X$  is said to be *bounded* when there exists a constant  $C > 0$  so that  $\|\Omega(x)\|_X \leq C\|x\|_X$  for all  $x \in X$ ; which in particular means that  $\Omega(x) \in X$  for all  $x \in X$ . Two centralizers  $\Omega_1 : X \curvearrowright X$  and  $\Omega_2 : X \curvearrowright X$  are *equivalent* if and only if the induced exact sequences are equivalent, which happens if and only if there exists a linear map  $L : X \rightarrow L_0$

so that  $\Omega_1 - \Omega_2 - L$  is bounded. Two centralizers  $\Omega_1 : X \curvearrowright X$  and  $\Omega_2 : X \curvearrowright X$  are said to be *boundedly equivalent* when  $\Omega_1 - \Omega_2$  is bounded. The interest in this notion (which, to some extent, plays the role of triviality for quasi-linear maps) stems from the following outstanding result of Kalton [29, Theorem 7.6]:

**Theorem 3.3.** *Let  $X$  be a separable superreflexive Köthe function space. Then there exists a constant  $c$  (depending on the concavity of a  $q$ -concave renorming of  $X$ ) such that if  $\Omega : X \curvearrowright X$  is a real centralizer on  $X$  with  $\rho(\Omega) \leq c$ , then*

- (1) *There is a pair of Köthe function spaces  $X_0, X_1$  such that  $X = (X_0, X_1)_{1/2}$  and  $\Omega - \Omega_{1/2}$  is bounded.*
- (2) *The spaces  $X_0, X_1$  are uniquely determined up to equivalent renorming.*

An example is in order: taking the couple  $(\ell_1, \ell_\infty)$ , the map  $B(x) = x^{2(1-z)}$  is a homogeneous bounded selection for the evaluation map  $\delta_{1/2} : \mathcal{H} \rightarrow \ell_2$ ; hence the interpolation procedure yields the centralizer  $-2\mathcal{K}$ ; while the couple  $(\ell_p, \ell_{p^*})$  yields  $-2(\frac{1}{p} - \frac{1}{p^*})\mathcal{K}$ . As we see the two centralizers are the same up to the scalar factor. Theorem 3.3 shows however that the scalar factor cannot be overlooked since it actually determines the end points  $X_0, X_1$  in the interpolation scale. See the general situation in Proposition 3.7.

We note for future use that the condition on  $\rho(\Omega)$ , which is necessary for existence, is not necessary for uniqueness; thus, uniqueness may be stated as follows:

**Proposition 3.4.** *Let  $X$  be a separable superreflexive Köthe function space. Assume that  $X = (X_0, X_1)_\theta = (Y_0, Y_1)_\theta$ , where  $0 < \theta < 1$  and  $X_i, Y_i$  are Köthe function spaces, and that the associated maps  $\Omega_X$  and  $\Omega_Y$  are boundedly equivalent. Then  $X_0 = Y_0$  and  $X_1 = Y_1$ .*

*Proof.* We follow Kalton's notation and the first steps of the proof of uniqueness in Kalton's theorem [29, Theorem 7.6], which is written in the case  $\theta = 1/2$ . Since  $\Omega_X$  and  $\Omega_Y$  are boundedly equivalent,  $\Omega_X^{[1]}$  and  $\Omega_Y^{[1]}$  are boundedly equivalent. Hence on a suitable strict semi-ideal,  $\Phi^{\Omega_X}$  is equivalent to  $\Phi_{Y_1} - \Phi_{Y_0}$ , while  $(1 - \theta)\Phi_{Y_0} + \theta\Phi_{Y_1}$  is equivalent to  $\Phi_X$ . Thus, up to equivalence  $\Phi_{Y_0}$  and  $\Phi_{Y_1}$  are uniquely determined. [29, Proposition 4.5] shows then that the spaces  $Y_0$  and  $Y_1$  are unique up to equivalence of norm.  $\square$

### 3.3. Centralizers and Lozanovskii's decomposition.

Here we follow Kalton (see [29, formula (3.2)]) to obtain a formula for the centralizer  $\Omega_\theta$  corresponding to the interpolation of a couple of Köthe function spaces  $(X_0, X_1)$ . Let  $0 < \theta < 1$ , and suppose that one of the spaces  $X_0, X_1$  has the Radon-Nikodym property. The Lozanovskii decomposition formula allows us to show (see [31, Theorem 4.6]) that the complex interpolation space  $X_\theta$  is isometric to the space  $X_0^{1-\theta}X_1^\theta$ , with

$$\|x\|_\theta = \inf\{\|y\|_0^{1-\theta}\|z\|_1^\theta : y \in X_0, z \in X_1, |x| = |y|^{1-\theta}|z|^\theta\}.$$

By homogeneity we may always assume that  $\|y\|_0 = \|z\|_1$  for  $y, z$  in this infimum. When  $\|y\|_0, \|z\|_1 \leq K\|x\|_\theta$  we shall say that  $|x| = |y|^{1-\theta}|z|^\theta$  is a *K-optimal decomposition* for  $x$ . When  $x$  is finitely supported or  $X$  is uniformly convex a 1-optimal (or simply, optimal) decomposition may be achieved. A simple choice of  $B_\theta(x)$  can be made for positive  $x$  as follows: Let  $a_0(x), a_1(x)$  be a  $(1+\epsilon)$ -optimal (or optimal when possible) Lozanovskii decomposition for  $x$ . Since  $\|x\|_\theta = \|a_0(x)\|_0 = \|a_1(x)\|_1$ , set  $B_\theta(x) \in \mathcal{H}$  given by  $B_\theta(x)(z) = |a_0(x)|^{1-z}|a_1(x)|^z$ . One thus gets for positive  $x$  the formula:

$$\Omega_\theta(x) = \delta'_\theta B_\theta(x) = |a_0(x)|^{1-\theta}|a_1(x)|^\theta \log \frac{|a_1(x)|}{|a_0(x)|} x = x \log \frac{|a_1(x)|}{|a_0(x)|}.$$



Using  $B_\theta(x) = (\text{sgn } x)B_\theta(|x|)$  for general  $x$  one still gets

$$(5) \quad \Omega_\theta(x) = x \log \frac{|a_1(x)|}{|a_0(x)|}.$$

Recall that a *unit*  $u$  in  $L_\infty$  is an element which only takes the values  $\pm 1$ . Thus one has:

**Lemma 3.5.** *The centralizer  $\Omega_\theta = \delta'_\theta B_\theta$  on  $X_\theta = (X_0, X_1)_\theta$  verifies:*

- (1)  $\Omega_\theta(ux) = u\Omega_\theta(x)$  for every unit  $u$  and  $x \in X_\theta$ ;
- (2)  $\text{supp } \Omega_\theta(x) \subset \text{supp } x$  for every  $x \in X_\theta$ ;
- (3) when  $X_0$  and  $X_1$  are spaces with a normalized 1-unconditional basis  $(e_n)$ ,  $\Omega_\theta(e_n) = 0$  for all  $n$ .

The Lozanovskii approach can be used to make explicit the Kalton correspondence between centralizers and interpolation scales in some special cases. Recall that the  $p$ -convexification of a Köthe function space  $X$  is defined by the norm  $\|x\| = \| |x|^p \|^{1/p}$ . Conversely, when  $X$  is  $p$ -convex, the  $p$ -concavification of  $X$  is given by  $\|x\| = \| |x|^{1/p} \|^p$ . Modulo the fact that every uniformly convex space may be renormed to be  $p$ -convex for some  $p > 1$ , the following proposition interprets Kalton-Peck maps defined on uniformly convex spaces as induced by specific interpolation schemes.

**Proposition 3.6.** *Let  $0 < \theta < 1 < p < \infty$ , and let  $X$  be a Banach space with 1-unconditional basis (respectively a Köthe function space). Then  $X_\theta = (\ell_\infty, X)_\theta$  (respectively  $(L_\infty(\mu), X)_\theta$ ) is the  $\theta^{-1}$ -convexification of  $X$ , and the induced centralizer on  $X_\theta$  is*

$$\Omega(x) = \theta^{-1} x \log(|x|/\|x\|_\theta).$$

*Conversely if  $X$  is  $p$ -convex and  $X^p$  is the  $p$ -concavification of  $X$  then  $X = (\ell_\infty, X^p)_{1/p}$  (respectively  $X = (L_\infty(\mu), X^p)_{1/p}$ ), and the induced centralizer is defined on  $X$  by*

$$\Omega(x) = p x \log(|x|/\|x\|).$$

*Proof.* We write down the proof for unconditional basis, the other being analogous. For normalized positive  $x$  in  $X_\theta$ , write  $x = a_0(x)^{1-\theta} a_1(x)^\theta$  and look for such a (normalized) decomposition which is optimal. Since  $a_0(x) \in \ell_\infty$ , we may assume that  $a_0(x)$  has constant coefficients equal to 1 on the support of  $x$ : otherwise, we may increase the non 1 coordinates of  $a_0(x)$  to 1, therefore diminishing the corresponding coordinates of  $a_1(x)$  and non-increasing the norm of  $a_1(x)$  by 1-unconditionality, and still get something optimal. So  $a_0(x) = \mathbf{1}_{\text{supp}(x)}$  and  $x = a_1(x)^\theta$ . Therefore  $\|x\|_\theta = \|a_1(x)\|^\theta = \|x^{1/\theta}\|^\theta$ . So  $X_\theta$  is the  $\theta^{-1}$ -convexification of  $X$  and

$$\Omega_\theta(x) = x \log(a_1(x)/a_0(x)) = \frac{1}{\theta} x \log(x).$$

As for the converse, note that when we interpolate  $\ell_\infty$  and some  $Y$  we have  $|a_1(x)| = |x|^p$  for  $x$  normalized in  $Y_\theta$ , so if we interpolate  $\ell_\infty$  and  $Y = X^{(p)}$  we obtain for such  $x$

$$\|x\|_{Y_\theta} = 1 = \|a_1(x)\|_Y = \| |x|^p \|_Y = \| (|x|^p)^\theta \|_X^p = \|x\|_X^p,$$

therefore  $X = Y_\theta = (\ell_\infty, X^{(p)})_\theta$ .

The remaining part of the converse is an immediate consequence of the first part of the proposition.  $\square$

As we announced before Theorem 3.3, we show now the dependence of the scalar factor with respect to different choices of endpoints in a given interpolation scale:

**Proposition 3.7.** *Let  $(X_0, X_1)$  be an admissible pair of Köthe function spaces and for some  $0 < \alpha < \beta < 1$ , consider also the admissible pair  $(X_\alpha, X_\beta)$ . Let  $\alpha < \theta < \beta$  so that one has  $(X_0, X_1)_\theta = (X_\alpha, X_\beta)_\rho$  for some  $0 < \rho < 1$ . Let  $\Omega$  (resp.  $\Omega'$ ) denote the centralizers generated by the couple  $(X_0, X_1)$  (resp.  $(X_\alpha, X_\beta)$ ). Then  $\Omega'_\rho = (\beta - \alpha)\Omega_\theta$ .*

*Proof.* It is easy to check (see [31, Theorem 4.5]) that  $\rho$  is given by  $\alpha(1 - \rho) + \beta\rho = \theta$ . Let us consider the centralizers

$$\Omega_\theta(x) = x \log \frac{|a_1(x)|}{|a_0(x)|} \quad \text{and} \quad \Omega'_\rho(x) = x \log \frac{|a_\beta(x)|}{|a_\alpha(x)|}.$$

Since  $x = a_0(x)^{1-\theta} a_1(x)^\theta$ ,  $1 - \theta = (1 - \alpha)(1 - \rho) + (1 - \beta)\rho$  and  $\theta = \alpha(1 - \rho) + \beta\rho$  we get

$$x = (a_0(x)^{1-\alpha} a_1(x)^\alpha)^{1-\rho} (a_0(x)^{1-\beta} a_1(x)^\beta)^\rho.$$

Thus taking  $a_\alpha(x) = a_0(x)^{1-\alpha} a_1(x)^\alpha$  and  $a_\beta(x) = a_0(x)^{1-\beta} a_1(x)^\beta$  it is not difficult to check that the minimality of  $x = a_0(x)^{1-\theta} a_1(x)^\theta$  implies the minimality of  $x = a_\alpha(x)^{1-\rho} a_\beta(x)^\rho$ , and the equality  $\Omega'_\rho(x) = (\beta - \alpha)\Omega_\theta(x)$  follows from the properties of the logarithm function.  $\square$

**3.4. The case of Orlicz function spaces.** We now describe the centralizers associated to Orlicz function spaces over a measure space  $(\Sigma, \mu)$ . Recall that an  $N$ -function is a map  $\varphi : [0, \infty) \rightarrow [0, \infty)$  which is strictly increasing, continuous,  $\varphi(0) = 0$ ,  $\varphi(t)/t \rightarrow 0$  as  $t \rightarrow 0$ , and  $\varphi(t)/t \rightarrow \infty$  as  $t \rightarrow \infty$ . An  $N$ -function  $\varphi$  satisfies the  $\Delta_2$ -property if there exists a number  $C > 0$  such that  $\varphi(2t) \leq C\varphi(t)$  for all  $t \geq 0$ . For  $1 < p < \infty$ ,  $\varphi(t) = t^p$  is  $N$ -function satisfying the  $\Delta_2$ -property.

When an  $N$ -function  $\varphi$  satisfies the  $\Delta_2$ -property, the Orlicz space  $L_\varphi(\mu)$  is given by

$$L_\varphi(\mu) = \{f \in L_0(\mu) : \varphi(|f|) \in L_1(\mu)\}.$$

with the norm

$$\|f\| = \inf\{r > 0 : \int \varphi(|f|/r) d\mu \leq 1\}$$

The following result was proved in [25], and a clear exposition can be found in [11].

**Proposition 3.8.** *Let  $\varphi_0$  and  $\varphi_1$  be two  $N$ -functions satisfying the  $\Delta_2$ -property, and let  $0 < \theta < 1$ . Then the formula  $\varphi^{-1} = (\varphi_0^{-1})^{1-\theta} (\varphi_1^{-1})^\theta$  defines an  $N$ -function  $\varphi$  satisfying the  $\Delta_2$ -property, and  $(L_{\varphi_0}(\mu), L_{\varphi_1}(\mu))_\theta = L_\varphi(\mu)$ .*

Next we give an expression for the centralizer associated to a Hilbert space obtained by complex interpolation of Orlicz spaces. Note that once we have defined a centralizer  $\Omega$  for normalized  $0 \leq f \in X$ , we can define  $\Omega(0) = 0$  and  $\Omega(g) = g \cdot \Omega(|g|/\|g\|)$  for  $0 \neq g \in X$ .

**Proposition 3.9.** *Let  $\varphi_0$  and  $\varphi_1$  be two  $N$ -functions satisfying the  $\Delta_2$ -property and such that  $t = \varphi_0^{-1}(t) \cdot \varphi_1^{-1}(t)$ . Then  $(L_{\varphi_0}(\mu), L_{\varphi_1}(\mu))_{1/2} = L_2(\mu)$  and the induced centralizer is*

$$\Omega_{1/2}(f) = f \log \frac{\varphi_1^{-1}(f^2)}{\varphi_0^{-1}(f^2)} = 2f \log \frac{\varphi_1^{-1}(f^2)}{f} \quad (0 \leq f \in L_2(\mu), \|f\|_2 = 1).$$

*Proof.* First we consider the general case  $\varphi^{-1} := (\varphi_0^{-1})^{1-\theta} (\varphi_1^{-1})^\theta$ , as in Proposition 3.8. For  $0 \leq f$  normalized in  $L_\varphi(\mu)$  we can write  $f = (\varphi_0^{-1}\varphi(f))^{1-\theta} (\varphi_1^{-1}\varphi(f))^\theta$ . Thus a selection of the quotient map  $\mathcal{H} \rightarrow L_\varphi(\mu)$  is given by  $B_\theta(f)(z) = (\varphi_0^{-1}\varphi(f))^{1-z} (\varphi_1^{-1}\varphi(f))^z$ . Differentiating  $B_\theta(f)'(z) = B_\theta(f)(z) \log \frac{|\varphi_1^{-1}(\varphi(f))|}{|\varphi_0^{-1}\varphi(f)|}$ , hence  $\Omega_{1/2}(f) = B_{1/2}(f)'(1/2) = f \log \frac{|\varphi_1^{-1}(\varphi(f))|}{|\varphi_0^{-1}\varphi(f)|}$ , which gives the desired result when  $\varphi(t) = t^2$ .  $\square$

**3.5. Additional properties.** The properties of  $\Omega_\theta$  obtained in Lemma 3.5 will turn out essential for our estimates, so they deserve a definition.

**Definition 2.** Let  $X$  be a Köthe function space. A centralizer  $\Omega : X \curvearrowright X$  is called exact if for each  $x \in X$  and every unit  $u$  one has  $\Omega(ux) = u\Omega x$ . It is called contractive if  $\text{supp } \Omega(x) \subset \text{supp } x$  for every  $x \in X$ .

One has:

**Lemma 3.10.** Let  $X$  be a Köthe function space.

- (1) Every exact quasi-linear map on  $X$  is contractive.
- (2) If  $X$  is reflexive, then every exact trivial centralizer  $\Omega$  on  $X$  admits an exact linear map  $\Lambda$  such that  $\Omega - \Lambda$  is bounded.
- (3) If  $X$  has unconditional basis  $(e_n)$  and is reflexive, and if  $\Omega$  is exact and trivial on  $X$ , and satisfies  $\Omega(e_n) = 0$  for all  $n$ , then  $\Omega$  is bounded.

*Proof.* (1) Let  $u \in L_\infty$  be the function with value 1 on the support of  $x$  and  $-1$  elsewhere, then  $ux = x$ , therefore  $u\Omega(x) = \Omega(ux) = \Omega(x)$  which means that  $\text{supp } \Omega(x)$  is included in the support of  $x$ .

(2) Let  $\Omega$  be a centralizer with constant  $C$  and assume that it is trivial. So some linear map  $\ell : X \rightarrow L_0$  exists such that  $B := \Omega - \ell$  is bounded. Let  $U$  denote the abelian group of units in  $L_\infty$ . Then  $U$  is amenable, so there exists a left invariant finitely additive mean  $m$  on  $U$  allowing to define for any bounded  $f : U \rightarrow \mathbb{R}$  an integral  $\int_U f(u)dm$ . Since  $X$  is reflexive we may then define for any bounded  $f : U \rightarrow X$  an element  $x = \int_U f(u)dm \in X$  in the natural way, i.e.

$$\phi(x) = \int_U \phi(f(u))dm$$

for every  $\phi \in X^*$ . One can therefore define a map  $\Lambda : X \rightarrow L_0$  as follows:

$$\Lambda(x) = \Omega(x) - \int_U uB(ux)dm.$$

Then the homogeneous map  $x \mapsto \int_U uB(ux)dm$  is bounded, and by exactness of  $\Omega$  and invariance of  $m$ , we have that  $\Lambda$  is exact. It is also easy to check that  $\Lambda$  is linear. Indeed, denoting by  $\Delta(x, y)$  the element  $\Omega(x+y) - \Omega x - \Omega y = B(x+y) - Bx - By \in X$ , and observing that  $\Delta(ux, uy) = u\Delta(x, y)$ , we obtain

$$\begin{aligned} \Omega(x+y) - \Lambda(x+y) &= \int_U uB(ux+uy)dm \\ &= \int_U u\Delta(ux, uy)dm + \int_U uBuxdm + \int_U uBuydm \\ &= \Delta(x, y) + \Omega(x) - \Lambda(x) + \Omega(y) - \Lambda(y) \\ &= \Omega(x+y) - \Lambda(x) - \Lambda(y). \end{aligned}$$

- (3) We claim that  $\Lambda(x) = ax$  for all  $x \in X$ , where  $\Lambda(e_n) = a_n e_n$ . Indeed

$$\Lambda(x) = \Lambda(x - x_n e_n) + \Lambda(x_n e_n) = \Lambda(x - x_n e_n) + a_n x_n e_n$$

which, since  $\Lambda(x - x_n e_n)$  has support disjoint from  $n$ , implies that the  $n$ -th entry of  $\Lambda(x)$  is  $a_n x_n$ . Since  $\Omega(e_n) = 0$ ,  $a_n e_n = -B(e_n)$ , and therefore  $(a_n)_n$  is a bounded sequence. So unconditionality applies to make  $\Lambda$  bounded. Since  $\Omega - \Lambda$  is also bounded,  $\Omega$  is bounded.  $\square$

A reformulation of (3) will provide us in due time with a criterion to distinguish between permutatively projectively equivalent centralizers:

**Corollary 3.11.** *Let  $\Omega$  and  $\Psi$  be exact centralizers on a reflexive space  $X$  with 1-unconditional basis  $(e_n)$ , and such that  $\Omega(e_n) = \Psi(e_n) = 0$  for all  $n \in \mathbb{N}$ . If  $\Omega$  and  $\Lambda$  are equivalent then they are boundedly equivalent.*

*Proof.*  $\Omega - \Lambda$  is still an exact centralizer vanishing on the  $e_n$ . Thus, if it is trivial then it is bounded.  $\square$

Lemma 3.10 can be generalized for maps between two different modules. We are interested in the particular case in which one has to combine two related actions: let  $X$  be an  $L_\infty$ -Banach module and let  $W \subset X$  be a subspace generated by disjointly supported elements  $W = [u_n]$ . Consider in this case the subspace  $L_\infty^W \subset L_\infty$  formed by the elements which are constant on the supports of all the  $u_n$ . Let  $U_W$  be its group of units. We say that a map  $\Omega : W \rightarrow X$  is *relatively exact* if  $\Omega(ux) = u\Omega(x)$  for all  $u \in U_W$  and  $x \in W$ , and we say that  $\Omega$  is *relatively contractive* if  $\text{supp}_X \Omega(x) \subset \text{supp}_X x$ , for all  $x \in W$ . One has:

**Lemma 3.12.** *Let  $X$  be a Köthe function space, and let  $W$  be a subspace of  $X$  generated by disjointly supported elements. Then:*

- (1) *If  $\Omega : X \curvearrowright X$  is a exact centralizer then the restriction  $\Omega|_W$  is relatively exact.*
- (2) *Every relatively exact map  $W \curvearrowright X$  is relatively contractive.*
- (3) *Assume  $X$  is reflexive. If some relatively exact  $\Omega : W \curvearrowright X$  is trivial then there exists a relatively exact linear map  $\Lambda : W \rightarrow X$  such that  $\Omega - \Lambda$  is bounded.*

*Proof.* Assertion (1) is obvious, (2) has the same proof as before. For (3), assuming  $\Omega = B + \ell$ , where  $B : W \rightarrow X$  is bounded and  $\ell : W \rightarrow L_0$  is linear, define for  $x \in W$ ,

$$\Lambda(x) = \Omega(x) - \int_{U_W} uB(ux)dm,$$

where  $m$  is a left invariant finitely additive mean on  $U_W$ .  $\square$

**Lemma 3.13.**

- (1) *Every centralizer  $\Omega$  on a Köthe function space admits a exact centralizer  $\omega$  such that  $\Omega - \omega$  is bounded.*
- (2) *Every exact centralizer (resp. quasi-linear map)  $\Omega$  between Banach spaces with unconditional basis admits a exact centralizer (resp. quasi-linear map)  $\omega$  such that  $\omega(e_n) = 0$  and  $\Omega - \omega$  is linear and exact.*
- (3) *Every contractive centralizer (resp. quasi-linear map)  $\Omega$  between Köthe function spaces admits, for every sequence  $(f_n)$  of disjointly supported vectors, a contractive centralizer (resp. quasi-linear map)  $\omega$  such that  $\omega(f_n) = 0$  and  $\Omega - \omega$  is linear and contractive.*

*Proof.* Assertion (1) is in [28, Prop. 4.1]. In fact,  $\omega(x) = \|x\| \text{sgn}(x)\Omega(|x|/\|x\|)$  for  $x \neq 0$ . To prove (2), note that since  $\Omega$  is contractive,  $\Omega(e_n) = \mu_n e_n$ , and we may define the multiplication linear map  $\ell(x) = \mu x$ , where  $\mu = (\mu_n)_n$ . Thus  $\omega = \Omega - \ell$  is the desired map. To prove (3), define as above a linear map by  $\ell(f_n) = \Omega(f_n)$ . If  $\Omega$  is contractive, so is  $\ell$  and thus  $\omega = \Omega - \ell$  is the desired map.  $\square$

#### 4. SINGULARITY AND ESTIMATES FOR EXACT CENTRALIZERS

Recall that an operator between Banach spaces is said to be *strictly singular* if no restriction to an infinite dimensional closed subspace is an isomorphism.

**Definition 3.** A quasi-linear map (in particular, a centralizer) is said to be singular if its restriction to every infinite dimensional closed subspace is never trivial. An exact sequence induced by a singular quasi-linear map is called a singular sequence.

It is well known [14] that a quasi-linear map is singular if and only if the associated exact sequence has strictly singular quotient map. So singular quasi-linear maps induce twisted sums which are, in some sense, as nontrivial as is possible. The following notion is perhaps more suitable to work with Köthe function spaces.

**Definition 4.** A quasi-linear map on a Köthe function space is called disjointly singular if its restriction to every subspace generated by a disjoint sequence is never trivial.

One can show that a quasi-linear map  $F : Z \rightarrow Y$  is disjointly singular if and only if the quotient map of the induced exact sequence  $0 \rightarrow Y \rightarrow Y \oplus_F Z \xrightarrow{q} Z \rightarrow 0$  is never an isomorphism on a subspace  $X$  of  $Y \oplus_F Z$  such that  $q(X)$  is generated by disjoint vectors in  $Z$ . Observe that when  $Z$  has an unconditional basis and the lattice structure one considers is the one induced by the basis then the two notions coincide since saying that  $q$  is an isomorphism on some subspace is the same that saying that it is an isomorphism on some subspace whose image is generated by blocks of the basis. Thus:

**Lemma 4.1.** A quasi-linear map  $F : Z \rightarrow Y$  on a Banach space  $Z$  with unconditional basis is singular if and only if it is disjointly singular with respect to the induced lattice structure.

Singularity implies disjoint singularity and, as we shall see, the reverse implication does not hold in general. Of course, a disjointly singular quasi-linear map is nontrivial. The following “transfer principle” ([14], [10]) will be essential for us.

**Lemma 4.2.** If a quasi-linear map defined on a Banach space  $X$  with basis is trivial on some infinite dimensional subspace of  $X$  then it is also trivial on some subspace  $W = [w_n]$  of  $X$  spanned by normalized blocks of the basis.

Observe that if  $F$  is a quasi-linear map on a Köthe space  $X$ , and if for some sequence  $(u_n)$  of disjointly supported vectors and some constant  $K$  one has

$$\left\| F\left(\sum \lambda_j u_j\right) - \sum \lambda_j F(u_j) \right\| \leq K \left\| \sum \lambda_j u_j \right\|$$

for all choices of scalars  $(\lambda_j)$  then  $F$  is not singular: indeed, the estimate above means that the linear map  $[u_j] \rightarrow X \oplus_F [u_j]$  given by  $u_j \rightarrow (0, u_j)$  is continuous. Under exactness conditions we can get a partial converse.

**Lemma 4.3.** Let  $\Omega : X \curvearrowright X$  be an exact centralizer on a reflexive Köthe function space. If  $\Omega$  is not disjointly singular, then there exists a subspace  $W$  of  $X$  generated by a disjoint sequence and a constant  $K$  such that given vectors  $u_1, \dots, u_n$  in  $W$  there are vectors  $z_1, \dots, z_n$  in  $X$  with  $\text{supp} z_i \subset \text{supp} u_i$  and  $\|z_i\| \leq K \|u_i\|$  such that for all scalars  $\lambda_1, \dots, \lambda_n$  one has

$$(6) \quad \left\| \Omega\left(\sum_{i=1}^n \lambda_i u_i\right) - \sum_{i=1}^n \lambda_i \Omega(u_i) \right\| \leq K \left( \left\| \sum_{i=1}^n \lambda_i u_i \right\| + \left\| \sum_{i=1}^n \lambda_i z_i \right\| \right).$$

*Proof.* Since  $\Omega$  is not disjointly singular, it is trivial on some subspace  $W = [u_n]$  spanned by disjointly supported vectors. Then by Lemma 3.12 there exists a linear relatively exact map  $\Lambda : W \rightarrow X$  so that  $\Omega|_W - \Lambda$  is bounded. Since both  $\Omega$  and  $\Lambda$  (by Lemma 3.12 (2)) are relatively contractive, so is  $\Omega - \Lambda$ . Set  $z_i = (\Omega - \Lambda)(u_i)$  and  $K = \|\Omega|_W - \Lambda\|$ .  $\square$

The preceding estimate can be considered as a subtler version of the “upper  $p$ -estimates” argument for non-splitting, which can be quickly described as: if the space  $X$  verifies some type of upper  $p$ -estimate and the twisted sum  $X \oplus_{\Omega} X$  splits then the space  $X \oplus_{\Omega} X$  must also verify the upper  $p$ -estimate (the key here is the  $p$  since, in general, if  $X$  has type  $p$  then  $X \oplus_{\Omega} X$  only needs to have type  $p + \varepsilon$  for every  $\varepsilon$  (see [27])). Therefore, given suitable vectors  $(u_n)$  in  $X$  the elements  $(0, u_n)$  in  $X \oplus_{\Omega} X$  should verify an upper  $p$ -estimate; which amounts to

$$\left\| \Omega\left(\sum_{i=1}^n u_i\right) - \sum_{i=1}^n \Omega(u_i) \right\| \leq C \sqrt[p]{p}.$$

We now introduce the notion of standard class of finite families of elements of Köthe spaces to simplify the exposition.

**Definition 5.** A standard class  $\mathcal{S}$  is a class of finite families ( $n$ -tuples) of elements of Köthe function spaces (respect. spaces with 1-unconditional bases)  $X$  satisfying

- (i) whenever  $(x_i) \in \mathcal{S}$  and  $\text{supp} z_i \subset \text{supp} x_i$  for all  $i$  then  $(z_i) \in \mathcal{S}$ ;
- (ii) assume that  $W$  is a subspace generated by disjoint vectors (resp. generated by successive vectors) of  $X$ , and  $(x_i)$  is  $n$ -tuple of elements of  $W$ ; if  $(x_i)$  belongs to  $\mathcal{S}$  as a family in  $W$ , then it also belongs to  $\mathcal{S}$  as a family in  $X$ .

The two standard classes we shall use in this paper are disjointly supported vectors in Köthe spaces and “Schreier” successive vectors on 1-unconditional bases, i.e. families  $(x_1, \dots, x_n)$  such that  $n < \text{supp } x_1 < \dots < \text{supp } x_n$ , but some other examples like successive vectors on 1-unconditional bases could also be of interest for other applications.

Given a standard class  $\mathcal{S}$  and a space  $X$ , we consider the following indicator function:

$$M_{X, \mathcal{S}}(n) := \sup\{\|x_1 + \dots + x_n\| : (x_j) \in \mathcal{S}, \|x_j\| \leq 1\}.$$

Lemma 4.3 can be rewritten as:

**Lemma 4.4.** Let  $\mathcal{S}$  be a standard class, and let  $\Omega : X \curvearrowright X$  be an exact centralizer on a reflexive Köthe function space. If  $\Omega$  is not disjointly singular, then there exists a subspace  $W$  of  $X$  generated by a disjoint sequence and a constant  $K$  such that given any  $n$ -tuple  $(u_i) \in \mathcal{S}$  belonging to the unit ball of  $W$ , one has

$$\left\| \Omega\left(\sum_{i=1}^n u_i\right) - \sum_{i=1}^n \Omega(u_i) \right\| \leq K M_{X, \mathcal{S}}(n).$$

We arrive now to the core of our method:

**Lemma 4.5.** Let  $(X_0, X_1)$  be an admissible couple of Köthe function spaces, fix  $0 < \theta < 1$ , and let  $\Omega_{\theta}$  be the induced centralizer on  $X_{\theta}$ . If  $(x_i) \in \mathcal{S}$  is a  $n$ -tuple in the unit ball of  $X_{\theta}$ , then

$$(7) \quad \left\| \Omega_{\theta}\left(\sum_{i=1}^n x_i\right) - \sum_{i=1}^n \Omega_{\theta}(x_i) - \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \left(\sum_{i=1}^n x_i\right) \right\| \leq 3 \frac{M_{X_0, \mathcal{S}}(n)^{1-\theta} M_{X_1, \mathcal{S}}(n)^{\theta}}{\text{dist}(\theta, \partial \mathbb{S})}.$$

*Proof.* To simplify, let us write  $M(n, z) = M_{X_0, \mathcal{S}}(n)^{1-z} M_{X_1, \mathcal{S}}(n)^z$ . Given  $0 < \epsilon < 1/4$ , let  $(x_i) \in \mathcal{S}$  be a  $n$ -tuple in the unit ball of  $X_{\theta}$ . Let  $B_{\theta}$  be a  $(1 + \epsilon)$ -bounded selection  $X_{\theta} \rightarrow \mathcal{H}$  such that  $\text{supp } B_{\theta}(x) \subset \text{supp } x$  for all  $x$ . Let  $F_i = B_{\theta}(x_i)$  for each  $i$ . Note that  $(F_i(z))$  is a  $n$ -tuple in  $\mathcal{S}$  for any  $z$  in the strip. Let  $F$  be the function

$$F(z) = \frac{F_1(z) + \dots + F_n(z)}{M(n, z)}$$

for  $z \in \mathbb{S}$ . We know that  $\|F\| \leq 1 + \epsilon$  and

$$F(\theta) = \frac{1}{M(n, \theta)}(x_1 + \dots + x_n).$$

Set  $k = \|F(\theta)\|^{-1}$ . The map  $\Phi : F(\theta) \mapsto F$  defines a linear bounded selection on the one-dimensional subspace  $[F(\theta)]$  having norm at most  $k$ . Therefore

$$\|B_{\theta|[F(\theta)]} - \Phi\| \leq 1 + \epsilon + k \leq k(1 + \epsilon) + \epsilon + k.$$

Thus, if  $x \in [F(\theta)]$ , and denoting  $\delta' = \delta'_\theta$ ,

$$\|(\delta' B_\theta - \delta' \Phi)(x)\|_\theta \leq (2k + k\epsilon + \epsilon) \|\delta'_{\ker \delta_\theta}\| \|x\|_\theta.$$

In particular

$$\left\| (\delta' B_\theta - \delta' \Phi) \left( \sum_{i=1}^n x_i \right) \right\|_\theta \leq (2k + k\epsilon + \epsilon) \operatorname{dist}(\theta, \partial\mathbb{S})^{-1} \left\| \sum_{i=1}^n x_i \right\|_\theta,$$

or equivalently

$$\left\| (\delta' B_\theta - \delta' \Phi) \left( \sum_{i=1}^n x_i \right) \right\|_\theta \leq \operatorname{dist}(\theta, \partial\mathbb{S})^{-1} (2 + \epsilon + \frac{\epsilon}{k}) M(n, \theta) \leq 3 \operatorname{dist}(\theta, \partial\mathbb{S})^{-1} M(n, \theta).$$

On the other hand,

$$F'(\theta) = F(\theta) \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} + \frac{1}{M(n, \theta)} \sum_i B_\theta(x_i)'(\theta),$$

which means

$$\delta' \Phi \left( \sum_i x_i \right) = \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \left( \sum_i x_i \right) + \sum_i \delta' B_\theta(x_i).$$

Therefore

$$\delta' \Phi \left( \sum_i x_i \right) - \delta' B_\theta \left( \sum_i x_i \right) = \sum_i \delta' B_\theta(x_i) - \delta' B_\theta \left( \sum_i x_i \right) + \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \left( \sum_i x_i \right)$$

which yields

$$\left\| \sum_i \delta' B_\theta(x_i) - \delta' B_\theta \left( \sum_i x_i \right) + \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \left( \sum_i x_i \right) \right\|_\theta \leq 3 \operatorname{dist}(\theta, \partial\mathbb{S})^{-1} M(n, \theta),$$

hence

$$(8) \quad \left\| \Omega_\theta \left( \sum_{i=1}^n x_i \right) - \sum_{i=1}^n \Omega_\theta(x_i) - \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \left( \sum_{i=1}^n x_i \right) \right\| \leq 3 \operatorname{dist}(\theta, \partial\mathbb{S})^{-1} M(n, \theta)$$

as desired.  $\square$

Observe that the estimate above applies (after suitable normalization) to all real centralizers; it is not equally clear the form such estimate should adopt for complex centralizers or for centralizers generated by arbitrary families. We show now that the function  $\theta \mapsto M_{X_\theta, \mathcal{S}}(n)$  is log-convex:

**Lemma 4.6.** *Given an interpolation scale  $(X_\theta)$  of Köthe function spaces associated to a pair  $(X_0, X_1)$  one has*

$$M_{X_\theta, \mathcal{S}}(n) \leq M_{X_0, \mathcal{S}}(n)^{1-\theta} M_{X_1, \mathcal{S}}(n)^\theta.$$

*Proof.* Let  $F(z) = (F_1(z) + \cdots + F_n(z))/M(n, z)$  be the function in the proof of Lemma 4.5. The inequalities  $\|F(\theta)\|_\theta \leq \|F\| \leq 1 + \epsilon$  imply  $\|x_1 + \cdots + x_n\|_\theta \leq (1 + \epsilon)M(n, \theta)$ , from where the conclusion follows.  $\square$

## 5. CRITERIA FOR SINGULARITY

Here we give some results that will allow us to recognize nontrivial exact sequences by showing that the quasi-linear map is singular in some sense.

**5.1. A general criterion in Köthe function spaces.** We set now the core of our criterion to obtain disjointly singular sequences: to combine Lemma 4.4, Lemma 4.5 and Lemma 4.6 to get the following result.

**Proposition 5.1.** *Let  $\mathcal{S}$  be a standard class. Let  $(X_0, X_1)$  be an interpolation couple of Köthe function spaces generating the interpolation scale  $(X_\theta)$ ; assume  $X_\theta$  is reflexive and let  $\Omega_\theta$  be the induced centralizer on  $X_\theta$ ,  $0 < \theta < 1$ . If  $\Omega_\theta$  is not disjointly singular then there exists a subspace  $W \subset X_\theta$  spanned by disjointly supported vectors and a constant  $K$  such that*

$$(9) \quad \left| \log \frac{M_{X_0, \mathcal{S}}(n)}{M_{X_1, \mathcal{S}}(n)} \right| M_{W, \mathcal{S}}(n) \leq K M_{X_0, \mathcal{S}}(n)^{1-\theta} M_{X_1, \mathcal{S}}(n)^\theta.$$

An even more general criterion could be obtained by using in the definition of  $M_X$  sequences of vectors whose norms are at most some prescribed varying values, instead of vectors of norm at most 1. We shall not write it since it will not be needed to deal with the applications we are interested in.

We consider first the standard class  $\mathcal{D}$  of all disjointly supported sequences in a Köthe function space  $X$ , and simplify notation to:

$$M_X(n) = M_{X, \mathcal{D}}(n) = \sup\{\|x_1 + \cdots + x_n\| : x_1, \dots, x_k \text{ disjoint in the unit ball of } X\}.$$

Recall that two functions  $f, g : \mathbb{N} \rightarrow \mathbb{R}$  are called equivalent, and denoted  $f \sim g$ , if  $0 < \liminf f(n)/g(n) \leq \limsup f(n)/g(n) < +\infty$ . As a direct application of the criterion in Proposition 5.1 we have:

**Proposition 5.2.** *Let  $(X_0, X_1)$  be an interpolation couple of two Köthe function spaces so that  $M_{X_0}$  and  $M_{X_1}$  are not equivalent. Let  $0 < \theta < 1$ . Assume that  $X_\theta$  is reflexive, "self-similar" in the sense that  $M_W \sim M_{X_\theta}$  for every infinite-dimensional subspace generated by a disjoint sequence  $W \subset X_\theta$ , and  $M_{X_\theta} \sim M_{X_0}^{1-\theta} M_{X_1}^\theta$ . Then  $\Omega_\theta$  is disjointly singular.*

*Proof.* Otherwise, the estimate (9) yields that, on some subspace  $W$ , one gets

$$\left| \log \frac{M_{X_0}(n)}{M_{X_1}(n)} \right| M_W(n) = O(M(n, \theta)) = O(M_{X_\theta}(n)) = O(M_W(n)),$$

which is impossible unless  $M_{X_0}$  and  $M_{X_1}$  are equivalent.  $\square$

Let us see these criteria at work. The simplest case of course concerns the scale of  $\ell_p$  spaces,  $1 < p < +\infty$ . These spaces are self similar with  $M_{\ell_p}(n) = n^{1/p}$ , while re-iteration theorems allow one to fix  $X_0$  and  $X_1$  at any two different values  $p, q$  so that  $\lim \left| \log \frac{M_{X_0}(n)}{M_{X_1}(n)} \right| = \lim \left| \log n^{1/p-1/q} \right| = +\infty$ . Thus, the induced centralizer, which is actually (projectively equivalent to) the Kalton-Peck  $\ell_\infty$ -centralizer  $\mathcal{K}$ , is disjointly singular, hence



singular on  $\ell_p$ . The case of  $L_p$  spaces,  $1 < p < +\infty$  is also simple: Proposition 5.1 yields that if the twisted sum fails to be disjointly singular then

$$\left| \log \frac{M_{L_\infty}(n)}{M_{L_1}(n)} \right| M_{\ell_p}(n) \leq K M_{L_\infty}^{1-\frac{1}{p}}(n) M_{L_1}^{\frac{1}{p}}(n).$$

Therefore  $(\log n)n^{1/p} \leq Kn^{1/p}$ , which is impossible. So the induced centralizer, which is actually (projectively equivalent to) the Kalton-Peck  $L_\infty$ -centralizer  $\mathcal{K}$ , in  $L_p$  is disjointly singular.

**Proposition 5.3.** *For  $1 < p < +\infty$ , the Kalton-Peck  $L_\infty$ -centralizer  $\mathcal{K}(f) = f \log \frac{|f|}{\|f\|}$  is disjointly singular on  $L_p$ ,*

In [6] it was shown that no  $L_\infty$ -centralizer on  $L_p$  is singular for  $0 < p < \infty$ ; previously, it had been shown in [38] that the Kalton-Peck  $L_\infty$ -centralizer  $\Omega(f) = f \log |f|/\|f\|$  on  $L_p$  is not singular since it becomes trivial on the Rademacher copy of  $\ell_2$ . Proposition 5.3 tells us that it is not trivial on any subspace generated by disjointly supported vectors. In [10, Theorem 2(b)] it was shown that the Kalton-Peck centralizer on  $\ell_p$  is singular for  $0 < p < \infty$ . Cabello [6] remarks that it would be interesting to know whether there exist singular quasi-linear maps  $L_p \rightarrow L_p$  for  $p < 2$ .

A tricky question is what occurs with the scale of  $L_p$ -spaces in their  $\ell_\infty$ -module structure generated by the Haar basis. Is the associated  $\ell_\infty$ -centralizer  $\Omega_\theta$  singular? Khintchine's inequality makes possible to define  $B_\theta(r) = f_r$  (the constant function  $f_r(z) = r$  on the subspace  $\ell_2^R$  generated by the Rademacher functions, so  $\Omega_\theta(r) = \delta'_\theta B_\theta(r) = 0$  on  $\ell_2^R$  and thus  $\Omega_\theta$  is not singular. Since the Haar basis is unconditional, this means that it is not disjointly singular either.

In sharp contrast with this, it was shown in [10] that the Kalton-Peck centralizer  $f \rightarrow f \log \frac{|f|}{\|f\|}$  (relative to the Haar basis) is singular for  $2 \leq p < \infty$ . This means, in particular, that the Kalton-Peck  $\ell_\infty$ -centralizer relative to the Haar basis is not the  $\ell_\infty$ -centralizer induced by the interpolation scale of  $L_p$  spaces in their  $\ell_\infty$ -module structure.

We may use Proposition 5.2 together with Proposition 3.6 to prove singularity of Kalton-Peck maps on more general classes of Banach lattices.

**Theorem 5.4.** *Let  $X$  be a reflexive,  $p$ -convex Köthe function space,  $p > 1$ . Assume  $M_X(n) \sim M_Y(n)$  for every subspace  $Y$  of  $X$  generated by a sequence of disjointly supported vectors. Then the Kalton-Peck map  $\mathcal{K}(x) = x \log \frac{|x|}{\|x\|}$  is disjointly singular on  $X$ .*

*Proof.* Since  $X$  is  $p$ -convex we may by Proposition 3.6 write  $X = (L_\infty, X^p)_{1/p}$ . Furthermore the centralizer induced by this interpolation scheme is a multiple of the Kalton-Peck map. In particular, the two twisted sums are projectively equivalent in the sense of Section 2. Thus one is singular if and only if the other is. Since the norm on  $X^p$  is defined as  $\|x\| = \||x|^{1/p}\|_X^p$ , we have immediately that  $M_{X^p}(n) = M_X(n)^p$ . Since  $X$  is  $p$ -convex,  $M_X(n)$  is not bounded and so  $M_X(n)^p$  is not equivalent to  $M_{L_\infty}(n) = 1$ . Furthermore

$$M_{L_\infty}(n)^{1-\frac{1}{p}} M_{X^p}(n)^{\frac{1}{p}} = (M_X(n)^p)^{1/p} = M_X(n),$$

and by Proposition 5.2 the centralizer (hence the Kalton-Peck map) is disjointly singular.  $\square$

**5.2. The criterion in spaces with unconditional bases.** We consider now the following asymptotic variation of  $M_X$  with its associated standard class:

$$A_X(n) = \sup\{\|x_1 + \dots + x_n\| : \|x_i\| \leq 1, n < x_1 < \dots < x_n\},$$

when  $X$  has a 1-monotone basis. Then Proposition 5.1 can be reformulated as follows:

**Proposition 5.5.** *Let  $(X_0, X_1)$  be an admissible pair of Banach spaces with a common 1-unconditional basis, and  $0 < \theta < 1$ .*

- a) *If the associated centralizer  $\Omega_\theta$  is not singular then there exists a block subspace  $W \subset X_\theta$  and a constant  $K$  such that:*

$$\left| \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \right| A_W(n) \leq K A_{X_0}^{1-\theta}(n) A_{X_1}^\theta(n).$$

- b) *If  $A_{X_0} \not\sim A_{X_1}$  and  $A_{X_0}^{1-\theta} A_{X_1}^\theta \sim A_{X_\theta} \sim A_Y$  for all subspaces  $Y \subset X_\theta$  then  $\Omega_\theta$  is singular.*

Recall that a Banach space with a basis is said to be *asymptotically  $\ell_p$*  if there exists  $C \geq 1$  such that for all  $n$  and normalized  $n < x_1 < \dots < x_n$  in  $X$ , the sequence  $(x_i)_{i=1}^n$  is  $C$ -equivalent to the basis of  $\ell_p^n$ . Apart from the  $\ell_p$  spaces, Tsirelson's space is asymptotically  $\ell_1$  as well as a class of H.I. spaces (this one without unconditional basic sequences) defined by Argyros and Delyanii [2]. One has:

**Corollary 5.6.** *Let  $(X_0, X_1)$  be an interpolation pair of Banach spaces with a common 1-unconditional basis. Let  $p_0 \neq p_1$  and  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ . The induced centralizer  $\Omega_\theta : X_\theta \curvearrowright X_\theta$  is singular in any of the following cases:*

- (1) *The spaces  $X_j$ ,  $j = 0, 1$  are reflexive asymptotically  $\ell_{p_j}$ .*
- (2) *Successive vectors in  $X_j$ ,  $j = 0, 1$  satisfy an asymptotic upper  $\ell_{p_j}$ -estimate; and for every block-subspace  $W$  of  $X_\theta$ , there exist a constant  $C$  and, for each  $n$ , a finite block-sequence  $n < y_1 < \dots < y_n$  in  $B_W$  such that  $\|y_1 + \dots + y_n\| \geq C^{-1}n^{1/p}$ .*

We also obtain as immediate corollary, with the same method as in Theorem 5.4:

**Corollary 5.7.** *Let  $X$  be a  $p$ -convex reflexive space with 1-unconditional basis, such that  $A_X(n) \sim A_Y(n)$  for every block-subspace  $Y$  of  $X$ . Then the Kalton-Peck map  $\mathcal{K}(x) = x \log \frac{\|x\|}{\|x\|}$  is singular on  $X$ .*

Spaces to which Corollary 5.7 apply include, for example, the  $p$ -convexified Tsirelson spaces  $T^{(p)}$ ,  $p > 1$ ; since then  $A_Y(n) \sim n^{1/p}$  for any block subspace  $Y$ . Thus, the Kalton-Peck map on  $T^{(p)}$  is singular.

**5.3. The criterion in spaces with monotone bases.** Let  $\Omega : X \rightarrow X$  be a quasi-linear map acting on a space with 1-monotone basis. This case does not fit under the umbrella of Kalton theorem, so it could well occur that  $\Omega$  could not be recovered from an interpolation scheme. Without the lattice structure, supports cannot be used in the same way as before, although successive vectors and asymptoticity still makes sense, so that the function  $A_X$  still may be defined. In this context one uses the range of vectors ( $\text{ran } x$  is the minimal interval of integers containing its support) instead of their supports. In the general case of 1-monotone bases the maps  $\Omega_\theta$  appearing in an interpolation process are not  $\ell_\infty$ -centralizers or contractive. However, the maps can be chosen to be ‘‘range’’ contractive, in the sense of verifying  $\text{ran } \Omega_\theta(x) \subset \text{ran } x$ . Indeed if for  $x \in c_{00}$ ,  $b_\theta(x)$  is an almost optimal selection, then  $B_\theta(x) = 1_{\text{ran } x} b_\theta(x)$  will also be almost optimal and range contractive, so  $\delta'_\theta B_\theta$  will be the

required map. The transfer principle still works and thus a non-singular  $\Omega : X \rightarrow X$  must be trivial on some subspace  $W$  generated by blocks of the basis.

Note that the lattice structure was not used in the proof of Lemma 4.5, apart from the use of supports, which are here replaced by ranges. So a proof entirely similar to that of Lemma 4.5, using instead the function

$$F(z) = \frac{1}{A_{X_0}(n)^{1-z} A_{X_1}(n)^z} (B_\theta(y_1) + \cdots + B_\theta(y_n))(z),$$

immediately yields the estimate

$$(10) \quad \left\| \Omega_\theta \left( \sum_{i=1}^n y_i \right) - \sum_{i=1}^n \Omega_\theta(y_i) - \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \sum_{i=1}^n y_i \right\| \leq k_\theta A_{X_0}^{1-\theta} A_{X_1}^\theta(n),$$

for all  $n < y_1 < \cdots < y_n$  in the unit ball of  $X_\theta$ , in an interpolation scale  $(X_0, X_1)$  of spaces with common 1-monotone basis (here  $k_\theta = 3 \operatorname{dist}(\theta, \partial\mathbb{S})^{-1}$ ). One can also prove that the function  $\theta \mapsto A_{X_\theta}(n)$  is log-convex working as in Lemma 4.6. On the other hand the estimate in Lemma 4.3 requires lattice structure in a deep way, and so something new is needed in the conditional case: we shall now see how the lattice structure may be replaced by hypotheses of local unconditionality and complementation.

**Proposition 5.8.** *Assume we have a complex interpolation scheme of two spaces  $X_0, X_1$  with a common 1-monotone basis. Assume that for every block-subspace  $W$  of  $X_\theta$ , there exists for every  $n$  a finite successive sequence  $n < y_1 < \cdots < y_n$  with  $\|y_i\| \leq 1 \forall i = 1, \dots, n$ , and constants  $\varepsilon_n, \lambda_n, M_n$  satisfying*

- (i) *The block sequence is  $\varepsilon_n$ -optimal, in the sense that  $\|\sum_{i=1}^n y_i\| \geq \varepsilon_n A_{X_0}(n)^{1-\theta} A_{X_1}(n)^\theta$ ;*
- (ii) *The block sequence  $\{y_1, \dots, y_n\}$  is  $\lambda_n$ -unconditional;*
- (iii) *the space  $[y_1, \dots, y_n]$  is  $M_n$ -complemented in  $X_\theta$ ;*

and so that

$$\liminf_{n \rightarrow +\infty} \frac{\lambda_n^3 M_n}{\varepsilon_n \left| \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \right|} = 0.$$

Then  $\Omega_\theta$  is singular.

*Proof.* Suppose that the restriction of  $\Omega_\theta$  to some subspace of  $X$  is trivial. By the hypothesis  $\Omega_\theta$  is trivial on some block subspace  $Y_\theta$  of  $X_\theta$ , and we can pick for any  $n$  a  $\lambda_n$ -unconditional,  $\varepsilon_n$ -optimal, finite sequence  $[y_i]_{i=1}^n$  of blocks in  $B_{Y_\theta}$  that is  $M_n$ -complemented in  $X_\theta$  by a projection  $P_n$ .

Then a local version of the proof of Lemma 3.12 (3) can be made. Let  $\ell : Y_\theta \rightarrow L_0$  be a linear map so that  $\|\Omega_{|Y_\theta} - \ell\| \leq K$ . Let then  $G_n \simeq \{-1, 1\}^n$  be the group of units of  $\ell_\infty^n$  acting on  $Y_n = [y_1, \dots, y_n]$  in the natural way by change of signs of the coordinates on the  $y_i$ 's, and let, for  $y \in Y_n$ ,  $\psi_n(y)$  be the finite average

$$\psi_n(y) = \operatorname{Ave}_{u \in G_n} u P_n (\Omega_{|Y_\theta} - \ell)(uy).$$

Note that  $\psi_n$  takes values in  $Y_n$ , and that this homogeneous map is bounded by  $K M_n \lambda_n^2$ . It is also an exact  $\ell_\infty^n$ -centralizer in the sense that  $\psi_n(uy) = u \psi_n(y)$  for  $u \in G_n$ , so  $\operatorname{supp} \psi_n(y) \subset \operatorname{supp} y$  for  $y \in Y_n$ . This implies that  $\psi_n(y_i) = \mu_i y_i$  for some scalars  $\mu_i$  with  $|\mu_i| \leq K M_n \lambda_n^2$ . Thus

$$(11) \quad \begin{aligned} \|\psi_n(\sum_{i=1}^n y_i) - \sum_{i=1}^n \psi_n(y_i)\| &\leq K M_n \lambda_n^2 \|\sum_{i=1}^n y_i\| + \|\sum_{i=1}^n \mu_i y_i\| \\ &\leq K M \lambda_n^2 (1 + \lambda_n) \|\sum_{i=1}^n y_i\|. \end{aligned}$$

Consider the estimate (10), and observe that replacing  $\Omega_\theta$  by  $\Omega_\theta - \ell$  with  $\ell$  linear changes nothing, and projecting and averaging on  $\pm$  signs as in the definition of  $\psi_n$  only changes the estimate by  $\lambda_n \|P_n\| \leq \lambda_n M_n$ ; so one gets

$$\left\| \psi_n \left( \sum_{i=1}^n y_i \right) - \sum_{i=1}^n \psi_n(y_i) - \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \sum_{i=1}^n y_i \right\| \leq k_\theta M_n \lambda_n A_{X_0}(n)^{1-\theta} A_{X_1}(n)^\theta.$$

On the other hand by log-convexity of  $A_{X_\theta}$  we can rewrite (11) as

$$(12) \quad \left\| \psi_n \left( \sum_i y_i \right) - \sum_i \psi_n(y_i) \right\| \leq K M_n \lambda_n^2 (1 + \lambda_n) A_{X_0}^{1-\theta}(n) A_{X_1}^\theta(n).$$

Putting both estimates together we get

$$\left| \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \right| \cdot \left\| \sum_{i=1}^n y_i \right\| \leq (K \lambda_n (1 + \lambda_n) + k_\theta) M_n \lambda_n A_{X_0}^{1-\theta}(n) A_{X_1}^\theta(n).$$

Condition (i) yields that

$$\varepsilon_n \left| \log \frac{A_{X_0}(n)}{A_{X_1}(n)} \right| \leq (K \lambda_n (1 + \lambda_n) + k_\theta) M_n \lambda_n$$

in contradiction with the hypothesis.  $\square$

**Corollary 5.9.** *Assume we have an interpolation scheme of two spaces  $X_0$  and  $X_1$  with a common 1-monotone basis. Let  $1 \leq p_0 \neq p_1 \leq +\infty$ ,  $0 < \theta < 1$ , and  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$  and assume that the spaces  $X_j$ ,  $j = 0, 1$  satisfy an asymptotic upper  $\ell_{p_j}$ -estimate; and that for every block-subspace  $W$  of  $X_\theta$ , there exist a constant  $C$  and for each  $n$ , a  $C$ -unconditional finite block-sequence  $n < y_1 < \dots < y_n$  in  $B_W$  such that  $\|y_1 + \dots + y_n\| \geq C^{-1} n^{1/p}$  and  $[y_1, \dots, y_n]$  is  $C$ -complemented in  $X_\theta$ . Then  $\Omega_\theta$  is singular.*

It was proved by Pisier [34] that a B-convex Banach space contains  $\ell_2^n$  uniformly complemented. Condition (ii) in Proposition 5.8 could suggest to apply this result to B-convex Banach spaces. Proposition 7.2 below states that when  $X$  is B-convex, nontrivial twisted sums  $X \oplus_F X$  always exist.

**5.4. Interpolation of families of spaces.** Here we apply the preceding criteria to spaces induced by complex interpolation of a family of spaces (see [17]), as will be necessary in Section 8. We thus take a family of compatible Banach spaces  $\{X_{(j,t)} : j = 0, 1; t \in \mathbb{R}\}$  with index in the boundary of  $\mathbb{S}$ , and denote by  $\Sigma(X_{j,t})$  the algebraic sum of these spaces with the norm

$$\|x\|_\Sigma = \inf \{ \|x_1\|_{(j_1,t_1)} + \dots + \|x_n\|_{(j_n,t_n)} : x = x_1 + \dots + x_n \}.$$

Let  $\mathcal{H}(X_{j,t})$  denote the space of functions  $g : \mathbb{S} \rightarrow \Sigma := \Sigma(X_{j,t})$  which are  $\|\cdot\|_\Sigma$ -bounded,  $\|\cdot\|_\Sigma$ -continuous on  $\mathbb{S}$  and  $\|\cdot\|_\Sigma$ -analytic on  $\mathbb{S}^\circ$ ; and satisfy  $g(it) \in X_{(0,t)}$  and  $g(it+1) \in X_{(1,t)}$  for each  $t \in \mathbb{R}$ . Note that  $\mathcal{H}(X_{j,t})$  is a Banach space under the norm

$$\|g\|_{\mathcal{H}} = \sup \{ \|g(j+it)\|_{(j,t)} : j = 0, 1; t \in \mathbb{R} \}.$$

For each  $\theta \in (0, 1)$ , or even  $\theta \in \mathbb{S}$ , we define

$$X_\theta := \{x \in \Sigma(X_{j,t}) : x = g(\theta) \text{ for some } g \in \mathcal{H}(X_{j,t})\}$$

with the norm  $\|x\|_\theta = \inf \{ \|g\|_{\mathcal{H}} : x = g(\theta) \}$ . Clearly  $X_\theta$  is the quotient of  $\mathcal{H}(X_{j,t})$  by the kernel of the evaluation map  $\ker \delta_\theta$ , and thus it is a Banach space.

All the ingredients of our constructions straightforwardly adapt to this context, and the only relevant modification is to set  $A_j(n) = \text{ess sup}_{t \in \mathbb{R}} A_{X_{j+it}}(n)$  instead of  $A_{X_j}(n)$ ,  $j = 0, 1$ .

**Proposition 5.10.** *Consider an interpolation scheme given by a family  $\{X_{(j,t)} : j = 0, 1; t \in \mathbb{R}\}$  of spaces with a common 1-monotone basis. Let  $1 \leq p_0 \neq p_1 \leq +\infty$ ,  $0 < \theta < 1$ , and  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ . Assume that all the spaces  $X_{j,t}$  satisfy an asymptotic upper  $\ell_{p_j}$ -estimate with uniform constant; and for every block-subspace  $W$  of  $X_\theta$ , there exist a constant  $C$  and for each  $n$ , a  $C$ -unconditional finite block-sequence  $n < y_1 < \dots < y_n$  in  $B_W$  such that  $\|y_1 + \dots + y_n\| \geq C^{-1}n^{1/p}$  and  $[y_1, \dots, y_n]$  is  $C$ -complemented in  $X_\theta$ . Then  $\Omega_\theta$  is singular.*

*Proof.* It is similar to those of Proposition 5.8 and Corollary 5.9.  $\square$

## 6. SINGULAR TWISTED HILBERT SPACES

In many cases, complex interpolation between a Banach space and its dual gives  $(X, X^*)_{1/2} = \ell_2$ . See e.g., the comments at [35, around Theorem 3.1]. Also Watbled [39] claims that her results cover the case of spaces with a 1-unconditional basis  $X$ . We do not know whether there could be counterexamples with monotone basis. So, for the sake of clarity, let us briefly explain the situation.

Given a Banach space  $X$  with a normalized basis  $(e_n)$ , we denote by  $(e_n^*)$  the corresponding sequence of biorthogonal functionals. We identify  $X$  with  $\{(e_n^*(x)) : x \in X\}$ , and its antidual space  $\hat{X}^*$  with  $\{\overline{(x^*(e_n))} : x^* \in X\}$ , both linear subspaces of  $\ell_\infty$ , in such a way that  $X \cap \hat{X}^*$  is continuously embedded in  $\ell_2$ . Indeed,  $x = (a_n) \in X \cap \hat{X}^*$  implies  $x(x) = \sum |a_n|^2 \leq \|x\|_X \cdot \|x\|_{\hat{X}^*}$ .

**Proposition 6.1.** *Let  $X$  be a Banach space with a monotone shrinking basis. Then  $(X, \hat{X}^*)_{1/2} = \ell_2$  with equality of norms.*

*Proof.* It is enough to show that  $\ell_2$  is continuously embedded in  $X + \hat{X}^*$  and apply [39, Corollary 4]. Let  $T : X \cap \hat{X}^* \rightarrow \ell_2$  be the embedding. Since the basis is shrinking,  $X \cap \hat{X}^*$  is dense in both  $X$  and  $\hat{X}^*$ . Thus the dual of  $X \cap \hat{X}^*$  is  $X^* + (\hat{X}^*)^* = X^{**} + \hat{X}^*$  [5, 2.7.1 Theorem], and the conjugate operator  $T^*$  embeds  $\ell_2$  into  $X + \hat{X}^*$ , which is a closed subspace of  $X^{**} + \hat{X}^*$  by the arguments in [39, p. 204].  $\square$

We have a similar result for Köthe function spaces  $X$ . Observe that in this case  $X^*$  and  $\hat{X}^*$  coincide as sets.

**Proposition 6.2.** [39, Corollary 5] *Let  $X$  be a Köthe function space on a complete  $\sigma$ -finite measurable space  $S$ . Suppose that  $X \cap X^*$  is dense in  $X$  and*

$$L_1(S) \cap L_\infty(S) \subset X \cap X^* \subset L_2(S) \subset X + X^* \subset L_1(S) + L_\infty(S).$$

*Then  $(X, X^*)_{1/2} = L_2(S)$ .*

Arguing like in Proposition 6.1, we can show that the conditions  $X$  and  $X^*$  intermediate spaces between  $L_1(S)$  and  $L_\infty(S)$ , and  $X \cap X^*$  dense in both  $X$  and  $X^*$  imply the hypothesis of Proposition 6.2.

In all the previous situations the twisted sum space induced by the interpolation of a space and its antidual is a twisted Hilbert space. Proposition 5.2 fits appropriately in this situation since  $\ell_2$  is “asymptotically self-similar” in the sense that  $A_W(n) = n^{1/2}$  for all infinite dimensional block subspaces. Thus, we are ready to construct singular exact sequences

$$0 \longrightarrow \ell_2 \longrightarrow E \longrightarrow \ell_2 \longrightarrow 0.$$

The first consequence of Corollary 5.9 is:

**Proposition 6.3.** *The interpolation of a reflexive asymptotically  $\ell_p$  space,  $p \neq 2$ , with its antidual induces a singular twisted Hilbert space.*

Thus interpolation of Tsirelson's space  $\mathcal{T}$  with its dual  $\mathcal{T}^*$ ; or interpolation of Argyros-Deliyanni's H.I. asymptotically  $\ell_1$ -space [2] with its antidual produce new singular exact sequences

$$0 \longrightarrow \ell_2 \longrightarrow X \longrightarrow \ell_2 \longrightarrow 0.$$

By uniqueness in Kalton's theorem (Proposition 3.4), the singular sequence induced by interpolation of  $\mathcal{T}$  with  $\mathcal{T}^*$  is not boundedly equivalent to

$$0 \longrightarrow \ell_2 \longrightarrow Z_2 \longrightarrow \ell_2 \longrightarrow 0.$$

Thus, by Corollary 3.11, they cannot be even equivalent. In favorable situations this can be improved to be non-permutatively projectively equivalent. Indeed, given a reflexive Banach space  $X$  with normalized subsymmetric basis  $(e_n)$ , we denote as usual [33]

$$\lambda_X(n) := \left\| \sum_{i=1}^n e_i \right\|_X.$$

Then  $\lambda_{X^*}(n) \simeq n/\lambda_X(n)$  (see [33, Proposition 3.a.6]). One has

**Proposition 6.4.** *Let  $\ell_M$  be the symmetric Orlicz space with function  $M_\alpha(t) = e^{-t^{-\alpha}}$ ,  $\alpha > 0$ . The induced centralizers at  $\ell_2 = (\ell_M, \ell_M^*)_{1/2}$  for different values of  $\alpha$  are not permutatively projectively equivalent.*

*Proof.* Let  $X$  and  $Y$  be reflexive spaces with normalized 1-unconditional and 1-subsymmetric bases, and let  $\Omega$  (resp.  $\Psi$ ) be the induced centralizers at  $\ell_2$  defined on terms of the Lozanovskii decompositions associated to  $(X, X^*)_{1/2}$  (resp.  $(Y, Y^*)_{1/2}$ ). Then

$$(\Omega - \mu\Psi)(x) = \left( \log \frac{|a_0(x)|}{|a_1(x)|} - \mu \log \frac{|a'_0(x)|}{|a'_1(x)|} \right) x.$$

Pick  $x = \sum_{i=1}^n x_i e_i$  with  $x_i = 1/\sqrt{n}$  and apply the above formula with

$$\begin{aligned} |a_0(x)| &= \lambda_X(n)^{-1} 1_{[1,n]}, & |a_1(x)| &= \frac{\lambda_X(n)}{n} 1_{[1,n]}, & \text{and} \\ |a'_0(x)| &= \lambda_Y(n)^{-1} 1_{[1,n]}, & |a'_1(x)| &= \frac{\lambda_Y(n)}{n} 1_{[1,n]}. \end{aligned}$$

If  $\Omega - \mu\Psi$  is trivial then it is bounded by Corollary 3.11, so the function  $\log(n\lambda_X(n)^{-2}) - \mu \log(n\lambda_Y(n)^{-2})$  on  $\mathbb{N}$  is bounded, which implies that the functions  $n\lambda_X(n)^{-2}$  and  $(n\lambda_Y(n)^{-2})^\mu$  are equivalent. It is not difficult to check that that is impossible for different  $\alpha, \beta > 0$  since the choice of  $M_\alpha$  in the statement yields  $\lambda_{\ell_{M_\alpha}}(n) \simeq (\log n)^{1/\alpha}$ . Since the symmetric Orlicz spaces have symmetric bases, the corresponding induced centralizers are not even permutatively projectively equivalent.  $\square$

We have found no specific criterion to show when twisted Hilbert sums induced by interpolation of spaces with subsymmetric bases are singular. Let us move our attention back to asymptotically  $\ell_p$  spaces.

**Proposition 6.5.** *Let  $X, Y$  be spaces with asymptotically  $\ell_p$  normalized 1-unconditional bases,  $1 \leq p \leq +\infty$ . Then the singular twisted Hilbert sums induced by the interpolation couples  $(X, X^*)$  and  $(Y, Y^*)$  at  $1/2$  are (permutatively) projectively equivalent if and only if the bases of  $X$  and  $Y$  are (permutatively) equivalent.*

*Proof.* The key is to show that projective equivalence actually implies equivalence, hence bounded equivalence; which implies, by Kalton's result (Proposition 3.4), that the bases of  $X$  and  $Y$  are equivalent.

Assume thus that the induced centralizers are  $\lambda$ -projectively equivalent. By Lemma 3.5 (3) and Corollary 3.11

$$\sum_i a_i^2 \left( \log \frac{\mu_i}{\nu_i} - \lambda \log \frac{\mu'_i}{\nu'_i} \right)^2 \leq K,$$

whenever  $x = \sum_i a_i e_i$  in  $\ell_2$  is normalized, and  $a_i^2 = \nu_i \mu_i = \nu'_i \mu'_i$  with

$$1 \leq \left\| \sum_i \nu_i e_i \right\|_X, \left\| \sum_i \mu_i e_i \right\|_{X^*}, \left\| \sum_i \nu'_i e_i \right\|_Y, \left\| \sum_i \mu'_i e_i \right\|_{Y^*} \leq c.$$

Taking  $x$  with support far enough on the basis, we may choose  $a_i = n^{-1/2}$  and  $\nu_i = \nu'_i \simeq n^{-1/p}$ ,  $\mu_i = \mu'_i \simeq n^{-1/p'}$ . Then  $|(1 - \lambda) \log n|^2 \leq K'$ , which means that  $\lambda = 1$ . Therefore we have equivalence, and even bounded equivalence by Corollary 3.11.

To deduce the permutative projective equivalence case from the projective equivalence case just note that if a basis  $(e_n)$  is asymptotically  $\ell_p$  then any permutation of  $(e_n)$  is again asymptotically  $\ell_p$  "in the long distance", in the sense that there exists  $C \geq 1$  and a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  such that for all  $n$  and normalized  $f(n) < x_1 < \dots < x_n$  in  $X$ , the sequence  $(x_i)_{i=1}^n$  is  $C$ -equivalent to the basis of  $\ell_p^n$ .  $\square$

From the purely Banach space theory it is interesting to decide whether the twisted Hilbert spaces thus obtained are isomorphic. We can obtain non-isomorphic singular twisted Hilbert spaces as follows.

**Definition 6.** A Lipschitz function  $\phi : [0 + \infty) \rightarrow \mathbb{C}$  with  $\phi(0) = 0$  is called *expansive* if for every  $M$  there exists  $N$  such that  $|s - t| \geq N \Rightarrow |\phi(s) - \phi(t)| \geq M$ .

Observe that Lipschitz functions for which  $\lim_{t \rightarrow \infty} \phi'(t) = 0$  are not expansive. In particular the functions  $\phi_r$  for  $0 < r < 1$  are not expansive, while  $\phi_1$  is expansive.

**Proposition 6.6.** Let  $X$  be a space with a normalized 1-unconditional basis that is self-similar, in the sense that  $M_X \sim M_Y$  for all subspaces  $Y \subset X$  generated by a disjoint sequence, and such that  $\lim_{n \rightarrow \infty} M_X(n) = \infty$ . Assume  $\phi : [0 + \infty) \rightarrow \mathbb{C}$  is an expansive Lipschitz function. Then the Kalton-Peck map  $\mathcal{K}_\phi(x) = x \phi \left( -\log \frac{\|x\|}{\|x\|} \right)$  is singular.

*Proof.* To simplify notation we write  $\Omega = \mathcal{K}_\phi$ . Observe that  $\Omega$  is a contractive centralizer. Assume that  $Y$  is a sublattice of  $X$  such that  $\Omega|_Y$  is trivial. Let  $M$  be arbitrary positive,  $N$  be such that  $|s - t| \geq N \Rightarrow |\phi(s) - \phi(t)| \geq M$ , and  $n$  be such that  $M_Y(n) \geq 2e^N$ . We may consider disjoint vectors  $y_1, \dots, y_n$  in  $Y$  of norm at most 1 such that  $\|y_1 + \dots + y_n\| \geq M_Y(n)/2$ . An easy calculation shows that

$$\Omega \left( \sum_i y_i \right) - \sum_i \Omega(y_i) = \sum_i y_i \left( \phi \left( -\log \left( \frac{\sum_i y_i}{K} \right) \right) - \phi \left( -\log \left( \sum_i y_i \right) \right) \right),$$

where  $K = \left\| \sum_{i=1}^n y_i \right\|$ . Each coordinate of the vector  $\log(\sum_i y_i) - \log(\sum_i y_i/K)$  is  $\log K$  which is larger than  $\log(M_Y(n)/2) \geq N$ . Therefore each coordinate of the vector  $\phi(-\log(\sum_i y_i)) - \phi(-\log(\sum_i y_i/K))$  is larger than  $M$  in modulus. We deduce that

$$\left\| \Omega \left( \sum_i y_i \right) - \sum_i \Omega(y_i) \right\| \geq M \left\| \sum_i y_i \right\| \geq M M_Y(n)/2.$$

By Lemma 4.4, this implies for some fixed constant  $k$  that  $k M_X(n) \geq M M_Y(n)/2$ , therefore  $M_X \not\sim M_Y$ , a contradiction which proves that  $\Omega$  is singular.  $\square$

Observe that the condition  $\lim_{n \rightarrow \infty} M_X(n) = \infty$  can be obtained assuming that  $X$  is self-similar and does not contain  $c_0$ .

In [30] Kalton obtained a family  $Z_2(\alpha)$  of complex twisted Hilbert spaces induced by the centralizers

$$\mathcal{K}_{i\alpha}(x) = x \left( -\log \frac{|x|}{\|x\|} \right)^{1+i\alpha}$$

for  $-\infty < \alpha < \infty$  (see also [28]). Since these are not real centralizers they appear, according to [29], as induced by the interpolation of three spaces. They are singular because:

**Lemma 6.7.** *The Lipschitz function  $\phi(t) = t^{1+i\alpha}$  is expansive.*

*Proof.*  $|\phi(s) - \phi(t)| = |se^{i\alpha \log(s)} - te^{i\alpha \log(t)}| \geq ||s| - |t|| = |s - t|$ . □

Thus, according to Proposition 6.6 [30] we get:

**Proposition 6.8.** *Given  $\alpha \in \mathbb{R}$ , the exact sequences*

$$0 \longrightarrow \ell_2 \longrightarrow Z_2(\alpha) \longrightarrow \ell_2 \longrightarrow 0$$

*are singular and for  $\alpha \neq \beta$  the spaces  $Z_2(\alpha)$  and  $Z_2(\beta)$  are not isomorphic.*

We consider now the Kalton-Peck centralizers  $\mathcal{K}_{\phi_r}(x) = x\phi_r(-\log(|x|/\|x\|_2))$  induced by the Lipschitz functions  $\phi_r(t) = t$  for  $0 \leq t \leq 1$ , and  $\phi_r(t) = t^r$  for  $1 < t < \infty$ , and the twisted Hilbert spaces  $\ell_2(\phi_r) = \ell_2 \oplus_{\mathcal{K}_{\phi_r}} \ell_2$  they generate, introduced by Kalton and Peck in [32]. Note that  $\ell_2(\phi_1) = Z_2$ . It follows from Kalton's theorem 3.3 ([29, Theorem 7.6]) that  $\ell_2(\phi_r)$  comes generated by some interpolation scale, and we show now that it is a scale of Orlicz spaces.

**Proposition 6.9.** *Let  $0 < r < 1$  and  $\varphi_0, \varphi_1$  be the maps  $[0, \infty) \rightarrow [0, \infty)$  defined by*

$$\varphi_0^{-1}(t) = t^{\frac{1}{2} + \frac{1}{4}(-\log t)^{r-1}}, \quad \varphi_1^{-1}(t) = t^{\frac{1}{2} - \frac{1}{4}(-\log t)^{r-1}},$$

*on a neighborhood of 0, and extended to  $[0, \infty)$  to be  $N$ -functions with the  $\Delta_2$ -property. Then*

$$\ell_2(\phi_r) \simeq (\ell_{\varphi_0}, \ell_{\varphi_1})_{1/2}.$$

*Proof.* We note that everything here is well defined since by choice of  $r$  and after an easy calculation,  $t^{3/4} \leq \varphi_0^{-1}(t) \leq t^{1/4}$ ,  $t^{3/4} \leq \varphi_1^{-1}(t) \leq t^{1/4}$  and  $\varphi_1^{-1}(t)$  and  $\varphi_0^{-1}(t)$  are increasing, for  $t$  in some neighborhood of 0. This is enough to make sure that  $\varphi_1$  and  $\varphi_0$  define  $N$ -function Orlicz spaces. The  $\Delta_2$ -property is also satisfied on a neighborhood of 0. Indeed

$$\begin{aligned} \varphi_0^{-1}(9t) &= 3t^{\frac{1}{2} + \frac{1}{4}(-\log 9t)^{r-1}} = 3\varphi_0^{-1}(t)t^{\frac{1}{4}[(-\log 9 - \log t)^{r-1} - (-\log t)^{r-1}]} \\ &= 3\varphi_0^{-1}(t) \exp\left(-\frac{1}{4}(-\log t)^r \left[\left(1 + \frac{\log 9}{\log t}\right)^{r-1} - 1\right]\right). \end{aligned}$$

The exponential in this expression is easily seen to tend to 1 when  $t$  tends to 0, so close enough to 0,  $\varphi_0^{-1}(9t) \geq 2\varphi_0^{-1}(t)$ , and  $\varphi_0$  satisfies the  $\Delta_2$  condition  $\varphi_0(2s) \leq 9\varphi_0(s)$  for  $s$  in a neighborhood of 0. The same holds for  $\varphi_1$ . Since  $\varphi_0^{-1}(t)\varphi_1^{-1}(t) = t$  on a neighborhood of 0, the equality  $(\ell_{\varphi_0}, \ell_{\varphi_1})_{1/2} = \ell_2$  holds up to equivalence of bases.

Let  $\psi$  be the map so that

$$\varphi_1^{-1}(t) = t^{\frac{1}{2} - \frac{1}{4}\psi(-\log(t))}.$$

Note that  $\psi$  is continuous,  $\psi(s) = s^{r-1}$  for  $s$  on a neighborhood  $V$  of  $+\infty$ , and only the value of  $\psi(s)$  for  $s \geq 0$  is relevant here. Suppose that  $\|x\|_2 = 1$ . Then the centralizer  $\Omega$  associated



to  $(\ell_{\varphi_0}, \ell_{\varphi_1})_{1/2} = \ell_2$  (see Proposition 3.9), is given by

$$\Omega(x) = 2x \log \frac{\varphi_1^{-1}(|x|^2)}{|x|} = 2x \log |x|^{-\frac{1}{2}\psi(-\log|x|)} = x\psi(-\log|x|)(-\log|x|),$$

while  $\mathcal{K}_{\phi_r}(x)_n = x_n \cdot (-\log|x_n|)^r$  whenever  $|x_n|$  is less than some constant  $c$  depending on  $V$ . So we deduce that

$$\begin{aligned} \|\Omega(x) - \mathcal{K}_{\phi_r}(x)\|^2 &\leq \sum_{|x_n| \geq c} 2(\Omega(x))_n^2 + (\mathcal{K}_{\phi_r}(x))_n^2 \\ &\leq 2((-\log c)^2 \sup_{[0, -\log c]} |\psi| + (-\log c)^{2r}). \end{aligned}$$

Since  $\Omega$  and  $\mathcal{K}_{\phi_r}$  are homogeneous, they are boundedly equivalent. Hence  $\ell_2 \oplus_{\Omega} \ell_2$  and  $\ell_2(\phi_r)$  are isomorphic.  $\square$

Recall from [32, Corollary 5.5] that the spaces  $\ell_2(\phi_r)$  are mutually non-isomorphic for different values of  $0 < r \leq 1$ . We already know [32, Corollary 5.5] that  $\mathcal{K} = \mathcal{K}_{\phi_1}$  is singular but, since the function  $\phi_r$  is not expansive for  $r < 1$ , we do not know if also  $\mathcal{K}_{\phi_r}$  is singular for  $0 < r < 1$ .

## 7. THE TWISTING OF H.I. SPACES

A Banach space  $X$  is said to be *indecomposable* if it cannot be decomposed as  $A \oplus B$  for two infinite dimensional subspaces  $A, B$ . An infinite dimensional space  $X$  is said to be *hereditarily indecomposable* (H.I., in short) if all subspaces are indecomposable [24]. It is said to be *quotient hereditarily indecomposable* (Q.H.I., in short) if all its quotients of subspaces are indecomposable [22]. In particular, Q.H.I. spaces are H.I. The existence of Q.H.I. Banach spaces was proved in [22]. The simplest connection between H.I. spaces and the theory of singular exact sequences is described in the following folklore proposition; we present its proof for the sake of completeness.

**Lemma 7.1.** *Given an exact sequence of Banach spaces*

$$0 \longrightarrow Y \longrightarrow X \xrightarrow{q} Z \longrightarrow 0,$$

*the space  $X$  is H.I. if and only if  $Y$  is H.I. and  $q$  is strictly singular.*

*Proof.* Suppose  $X$  is H.I. Then clearly  $Y$  is H.I., and if  $q$  is not strictly singular,  $q|_V$  is an isomorphism for some (infinite dimensional) subspace  $V$  of  $X$ , hence  $Y \oplus V$  is a subspace of  $X$  and thus  $X$  cannot be H.I. Conversely, suppose that  $q$  is strictly singular. If  $X$  is not H.I. we can find a decomposable subspace  $X_1 \oplus X_2$  of  $X$ , and  $q$  has compact (even nuclear) restrictions on some subspaces  $Y_1 \subset X_1$  and  $Y_2 \subset X_2$ . Thus we can assume that there exists a bijective isomorphism  $U : X \rightarrow X$  such that  $U(Y_1)$  and  $U(Y_2)$  are contained in  $Y$ . Since  $U(Y_1) \oplus U(Y_2)$  is closed, we conclude that  $Y$  is not H.I.  $\square$

The basic question we tackle in this section is whether it is possible to obtain nontrivial twisted sums of H.I. spaces. The existence of a nontrivial twisted sum of  $A$  and  $B$  will be denoted  $\text{Ext}(B, A) \neq 0$ . On one hand, if  $X$  is a given example of a Q.H.I. space and  $Y$  is a subspace of  $X$  with  $\dim Y = \dim X/Y = \infty$ , then  $X$  is a nontrivial twisted sum of the two H.I. spaces  $Y$  and  $X/Y$ . However, what one is looking for is to obtain methods to twist two specified H.I. spaces. Recall that the Kalton-Peck method [32] to twist spaces only works, in principle, under unconditionality assumptions. A second method is to use the local theory of exact sequences as developed in [8]. The following result is a good example; we could not find it explicitly in the literature, but it is certainly known:

**Proposition 7.2.** *If  $X$  is a  $B$ -convex Banach space then  $\text{Ext}(X, X) \neq 0$ .*

*Proof.* If  $X$  contains  $\ell_2^n$  uniformly complemented, as it is the case of  $B$ -convex Banach spaces, then  $\text{Ext}(X, \ell_2) \neq 0$  [8]. And if  $\text{Ext}(X, X) = 0$  then  $\text{Ext}(X, \ell_2) = 0$  [8].  $\square$

The only currently known  $B$ -convex H.I. space is the one constructed by Ferenczi in [21]. So, calling this space  $\mathcal{F}$  one gets  $\text{Ext}(\mathcal{F}, \mathcal{F}) \neq 0$ . However this is not entirely satisfactory since this twisting does not provide any information about the twisted sum space, apart from its existence. So we formulate the following question:

**Problem 1.** *Let  $X$  be an H.I. space. Does there exist an H.I. twisted sum of  $X$ ?*

Focusing again on Ferenczi's space  $\mathcal{F}$ , since it is a space obtained via an interpolation scheme, i.e.,  $\mathcal{F} = X_\theta$  for a certain configuration of spaces, the induced centralizer  $\Omega_\theta$  provides a natural twisted sum of  $\mathcal{F}$  with itself that we call  $\mathcal{F}_2$ :

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F} \longrightarrow 0.$$

We will show in Section 8 that this sequence is singular, which implies that  $\mathcal{F}_2$  is H.I.

By the characterization in Lemma 7.1 it is tempting to believe that a twisted sum of two H.I. spaces is H.I. whenever is not trivial. However, this is not the case:

**Proposition 7.3.** *There exists a nontrivial twisted sum of two H.I. spaces which is indecomposable but not H.I.*

*Proof.* Recall that two Banach spaces  $A, B$  are said to be totally incomparable if no infinite dimensional subspace of  $A$  is isomorphic to a subspace of  $B$ . It was proved in [22, Prop. 25] that there exist two reflexive Q.H.I. spaces  $X_1, X_2$  admitting infinite dimensional subspaces  $Y_1 \subset X_1$  and  $Y_2 \subset X_2$  such that  $Y_1$  is isometric to  $Y_2$  and  $X_1/Y_1$  and  $X_2/Y_2$  are infinite dimensional and totally incomparable. Note that  $X_1^*$  and  $X_2^*$  are Q.H.I.

Given a bijective isometry  $U : Y_1 \rightarrow Y_2$ , we consider the subspace  $\hat{Y} := \{(y, Uy) : y \in Y_1\}$  of  $X_1 \times X_2$ , the quotient  $\hat{X} := (X_1 \times X_2)/\hat{Y}$ , and the quotient map  $Q : X_1 \times X_2 \rightarrow \hat{X}$ . Note that  $\hat{X}_1 := Q(X_1 \times \{0\})$  and  $\hat{X}_2 := Q(\{0\} \times X_2)$  are subspaces of  $\hat{X}$  isometric to  $X_1$  and  $X_2$  respectively, and  $\hat{Z} := \hat{X}_1 \cap \hat{X}_2 = Q(Y_1 \times \{0\}) = Q(\{0\} \times Y_2)$ . Thus  $\hat{X}/\hat{Z}$  is isomorphic to  $\hat{X}_1/\hat{Z} \oplus \hat{X}_2/\hat{Z}$ , hence  $\hat{Z}^\perp$  is decomposable and  $\hat{X}^*$  is not H.I. Let us see that  $\hat{X}^*$  is a nontrivial twisted sum of two H.I. spaces: Since  $\hat{X}$  is reflexive and H.I. [22, Proposition 23], the dual space  $\hat{X}^*$  is indecomposable, hence the exact sequence

$$0 \longrightarrow \hat{X}_1^\perp \longrightarrow \hat{X}^* \longrightarrow \hat{X}^*/\hat{X}_1^\perp \longrightarrow 0$$

is nontrivial. Moreover,  $\hat{X}_1^\perp$  and  $\hat{X}^*/\hat{X}_1^\perp$  are H.I. because  $\hat{X}_1 \simeq X_1$  and  $\hat{X}/\hat{X}_1 \simeq X_2/Y_2$  are Q.H.I. and reflexive.  $\square$

We can present an alternative construction of nontrivial and non H.I. twisted sums of H.I. spaces. Let us say that a Banach space  $X$  admits a singular extension if there exists a singular exact sequence

$$0 \longrightarrow X \longrightarrow Y \xrightarrow{q} Z \longrightarrow 0;$$

i.e., an exact sequence with  $q$  strictly singular and  $Z$  infinite dimensional. By Lemma 7.1, a HI space admits a singular extension if and only if it admits a non trivial extension which is a HI space.

**Proposition 7.4.** *Every separable H.I. space  $X$  which admits a singular extension is a complemented subspace of a nontrivial twisted sum of two H.I. spaces.*

*Proof.* Let  $0 \rightarrow X \xrightarrow{i} Y \xrightarrow{q} Z \rightarrow 0$  be a singular extension of  $X$  with  $Y$  separable. It follows from Proposition 7.1 that  $Y$  is H.I. By [3, Theorems 14.5 and 14.8] there exists a separable H.I. space  $W$  and a surjective operator  $p : W \rightarrow Y$  with infinite dimensional kernel. Note that  $p$  is strictly singular by Proposition 7.1. We consider the closed subspace  $\text{PB} := \{(w, x) \in W \oplus X : p(w) = i(x)\}$  of  $W \oplus X$  and the projection operators  $\alpha : \text{PB} \rightarrow W$  and  $\beta : \text{PB} \rightarrow X$ . Note that  $\beta$  is strictly singular because  $i\beta = q\alpha$ , and that  $\beta$  is surjective with  $\ker(\beta) = \ker(p)$  an H.I. space. Hence  $\text{PB}$  is H.I.

Since the operator  $U : (w, x) \in Z \oplus X \rightarrow i(x) - p(w) \in Y$  is surjective, we have a twisted sum of two H.I. spaces

$$(13) \quad 0 \longrightarrow \text{PB} \longrightarrow W \oplus X \xrightarrow{U} Y \longrightarrow 0.$$

To finish the proof it is enough to show that this twisted sum is nontrivial. Indeed, otherwise  $U$  would be in the class  $\Phi_r$  of operators with complemented kernel and finite codimensional closed range. By the stability of  $\Phi_r$  under strictly singular perturbations [1, Theorem 7.23], the operator  $T(w, x) \in Z \oplus W \rightarrow i(x) \in Y$  would define an isomorphism of  $X$  onto a finite codimensional subspace of  $Y$ , which is not possible.  $\square$

We do not know if every separable H.I. space admit a singular extension. On the other hand, the exact sequence (13) also shows that there are nontrivial twisted sums of H.I. spaces which are decomposable (“two” is the maximum number of summands by [23, Theorem 1]). In Section 9 we will give other examples of this kind.

To conclude this section, we formulate the general problem about twisting H.I.:

**Problem 2.** *Does there exists an H.I. space  $X$  so that  $\text{Ext}(X, X) = 0$ ?*

Note (see [4]) that there are only a few known solutions to the equation  $\text{Ext}(X, X) = 0$ : the spaces  $L_1(\mu)$ ,  $c_0$ ,  $\ell_\infty(\Gamma)$  and  $\ell_\infty/c_0$ .

## 8. AN H.I. TWISTED SUM OF $\mathcal{F}$

Ferenci’s H.I. uniformly convex space  $\mathcal{F}$  [21] comes induced by a complex interpolation scheme associated to a family of Banach spaces (briefly described in Subsection 5.4) setting  $X_{(1,t)} = \ell_q$ ,  $q > 1, t \in \mathbb{R}$ , and as  $X_{(0,t)}$  certain Gowers-Maurey-like spaces with 1-monotone basis. We fix  $\theta \in (0, 1)$ , and define  $\mathcal{F} = \{x \in \Sigma(X_{j,t}) : x = g(\theta) \text{ for some } g \in \mathcal{H}(X_{j,t})\}$  with the quotient norm of  $\mathcal{H}(X_{j,t})/\ker \delta_\theta$ , given by  $\|x\|_\theta = \inf\{\|g\|_{\mathcal{H}} : x = g(\theta)\}$ . In this section we will show that the space  $\mathcal{F}$  satisfies the hypotheses of Proposition 5.10 with  $C = 1 + \epsilon$  for any  $\epsilon > 0$  and thus:

**Theorem 8.1.** *The induced exact sequence*

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F} \longrightarrow 0.$$

*is singular. Therefore  $\mathcal{F}_2$  is H.I.*

We have trivial upper  $\ell_1$ -estimates in spaces  $X_{(0,t)}$  and upper  $\ell_q$ -estimates in spaces  $X_{(1,t)}$ . So we only need to check the  $\ell_p$ -condition of Proposition 5.10 in the middle space  $X_\theta$ , for  $\frac{1}{p} = 1 - \theta + \frac{\theta}{q}$ . Let  $f(x) := \log_2(1+x)$ . We first state estimates relative to successive vectors in the space  $\mathcal{F}$  [21, Proposition 1], as well as estimates for successive functionals in  $\mathcal{F}^*$  obtained by standard duality arguments:

**Lemma 8.2.** *For all successive vectors  $x_1 < \dots < x_n$  in  $\mathcal{F}$ ,*

$$\frac{1}{f(n)^{1-\theta}} \left( \sum_{i=1}^n \|x_i\|^p \right)^{1/p} \leq \left\| \sum_{i=1}^n x_i \right\| \leq \left( \sum_{i=1}^n \|x_i\|^p \right)^{1/p},$$

*and for all successive functionals  $\phi_1 < \dots < \phi_n$  in  $\mathcal{F}^*$ ,*

$$\left( \sum_{i=1}^n \|\phi_i\|^{p'} \right)^{1/p'} \leq \left\| \sum_{i=1}^n \phi_i \right\| \leq f(n)^{1-\theta} \left( \sum_{i=1}^n \|\phi_i\|^{p'} \right)^{1/p'}.$$

In [21],  $\ell_{p+}^n$ -averages are defined as normalized vectors of the form  $\sum_{i=1}^n x_i$ , where the  $x_i$ 's are successive of norm at most  $(1+\epsilon)n^{-1/p}$ , and may be found in any block-subspace of  $\mathcal{F}$  (see [21, Lemma 2]). However here we need to control not only the norm of  $\sum_{i=1}^n x_i$  but also of  $\sum_{i=1}^n \pm x_i$  for any choice of signs  $\pm$ , so [21, Lemma 2] is not quite enough. To this end we shall use RIS sequences as defined in [21, Definition 3].

RIS sequences with constant  $C > 1$  are successive sequences of  $\ell_{p+}^{n_k}$ -averages with a technical "rapidly" increasing condition on the  $n_k$ 's and therefore are also present in every block subspace of  $\mathcal{F}$ . Every subsequence of a RIS sequence is again a RIS sequence. In what follows  $L$  is some lacunary infinite subset of  $\mathbb{N}$  whose exact definition may be found in [21]. As a consequence of Lemma 8.2, [21, Lemma 10] and standard duality arguments we have:

**Lemma 8.3.** *Let  $y_1 < \dots < y_n$  be a RIS sequence in  $\mathcal{F}$ , with constant  $1 + \epsilon^2/100$ , where  $n \in [\log N, \exp N]$  for some  $N$  in  $L$ , and  $0 < \epsilon < 1/16$ . Then*

$$\frac{n^{1/p}}{f(n)^{1-\theta}} \leq \left\| \sum_{i=1}^n y_i \right\| \leq (1+\epsilon) \frac{n^{1/p}}{f(n)^{1-\theta}}.$$

*Furthermore if for all  $i$ ,  $\phi_i \in \mathcal{F}^*$  satisfies  $\|\phi_i\| = \phi_i(y_i) = 1$  and  $\text{ran } \phi_i \subset \text{ran } y_i$ , then*

$$(1+\epsilon)^{-1} f(n)^{1-\theta} n^{1/p'} \leq \left\| \sum_{i=1}^n \phi_i \right\| \leq f(n)^{1-\theta} n^{1/p'}.$$

We deduce the existence of sequences satisfying the condition of Proposition 5.10 in any block-subspace of  $\mathcal{F}$ :

**Proposition 8.4.** *Let  $Y$  be a block sequence of  $\mathcal{F}$ ,  $n \in \mathbb{N}$ , and  $\epsilon > 0$ . Then there exists a block-sequence  $y_1 < \dots < y_n$  in  $Y$  and a block-sequence  $\psi_1 < \dots < \psi_n$  in  $\mathcal{F}^*$  such that:*

- (1)  $(1+\epsilon)^{-1} \leq \|\psi_i\| \leq 1 \leq \|y_j\| \leq 1+\epsilon$  and  $\psi_i(y_j) = \delta_{ij}$  for  $i, j = 1, \dots, n$ ,
- (2) for any complex  $\alpha_1, \dots, \alpha_n$ ,  $\left\| \sum_{i=1}^n \alpha_i y_i \right\| \geq (1+\epsilon)^{-1} (\sum_{i=1}^n |\alpha_i|^p)^{1/p}$
- (3) for any complex  $\alpha_1, \dots, \alpha_n$ ,  $\left\| \sum_{i=1}^n \alpha_i \psi_i \right\| \leq (1+\epsilon) (\sum_{i=1}^n |\alpha_i|^{p'})^{1/p'}$

*In particular the block sequence  $y_1 < \dots < y_n$  of  $Y$  is  $(1+\epsilon)$ -equivalent to the unit vector basis of  $\ell_p^n$  and  $[y_1, \dots, y_n]$  is  $(1+\epsilon)$ -complemented in  $Y$ .*

*Proof.* Assuming  $\epsilon \leq 1/16$ , pick  $m$  such that  $\text{dist}(mn, N) < n$  for some  $N \in L$  and big enough to ensure that  $m$  and  $mn$  belong to  $[\log N, \exp N]$ , and that  $f(mn)/f(m) < 1+\epsilon$ . Denote  $M = mn$ . Let  $x_1, \dots, x_M$  be a RIS in  $Y$  with constant  $1 + \epsilon^2/100$  and  $\phi_1, \dots, \phi_M$  be a sequence of successive norming functionals in  $X^*$  for  $x_1, \dots, x_M$ .

Now for  $j = 1, \dots, n$ , let

$$y_j = \frac{f(m)^{1-\theta}}{m^{1/p}} \sum_{i=(j-1)m+1}^{jm} x_i, \quad \text{and} \quad \psi_j = \frac{1}{f(m)^{1-\theta} m^{1/p'}} \sum_{i=(j-1)m+1}^{jm} \phi_i.$$

Since  $x_{(j-1)m+1}, \dots, x_{jm}$  is a RIS with constant  $1 + \epsilon^2/100$ , we have by Lemma 8.3 that for  $j = 1, \dots, n$ ,

$$1 \leq \|y_j\| \leq (1 + \epsilon), \quad (1 + \epsilon)^{-1} \leq \|\psi_j\| \leq 1,$$

and clearly  $\psi_j(y_k) = \delta_{j,k}$ . For any complex  $\alpha_1, \dots, \alpha_n$ , Lemma 8.2 implies

$$\frac{m^{1/p}}{f(m)^{1-\theta}} \left\| \sum_{j=1}^n \alpha_j y_j \right\| \geq \frac{(\sum_{j=1}^n m |\alpha_j|^p)^{1/p}}{f(M)^{1-\theta}},$$

so

$$\left\| \sum_{j=1}^n \alpha_j y_j \right\| \geq \left( \sum_{j=1}^n |\alpha_j|^p \right)^{1/p} \left( \frac{f(m)}{f(M)} \right)^{1-\theta} \geq \left( \sum_{j=1}^n |\alpha_j|^p \right)^{1/p} (1 + \epsilon)^{-1}.$$

Lemma 8.2 also implies  $f(m)^{1-\theta} m^{1/p'} \|\sum_{j=1}^n \alpha_j \psi_j\| \leq f(M)^{1-\theta} (\sum_{j=1}^n m |\alpha_j|^p)^{1/p'}$ , so  $\|\sum_{j=1}^n \alpha_j \psi_j\| \leq (1 + \epsilon) (\sum_{j=1}^n |\alpha_j|^{p'})^{1/p'}$ .

Clearly  $(y_i)_{i=1}^n$  is  $(1 + \epsilon)$ -equivalent to the unit basis of  $\ell_p^n$ . We claim that  $Px = \sum_{i=1}^n \psi_i(x) y_i$  defines a projection from  $\mathcal{F}$  onto  $[y_1, \dots, y_n]$  of norm at most  $(1 + \epsilon)^{2p}$ . Indeed for  $x \in \mathcal{F}$ ,

$$\|Px\|^p \leq (1 + \epsilon)^p \left( \sum_{i=1}^n |\psi_i(x)|^p \right) = (1 + \epsilon)^p \left( \sum_{i=1}^n \alpha_i |\psi_i(x)|^{p-1} \psi_i(x) \right)$$

for some  $\alpha_1, \dots, \alpha_n$  of modulus 1. So

$$\|Px\|^p \leq (1 + \epsilon)^p \|x\| \left\| \sum_{i=1}^n \alpha_i |\psi_i(x)|^{p-1} \psi_i \right\| \leq (1 + \epsilon)^{p+1} \|x\| \left( \sum_{i=1}^n |\psi_i(x)|^{p-1} \right)^{1/p'}.$$

Since

$$\sum_{i=1}^n |\psi_i(x)|^{p-1} \right)^{p'} = \sum_{i=1}^n |\psi_i(x)|^p \leq (1 + \epsilon)^p \|Px\|^p,$$

we deduce  $\|Px\|^p \leq (1 + \epsilon)^{p+1+p/p'} \|x\| \|Px\|^{p/p'}$ , therefore  $\|Px\| \leq (1 + \epsilon)^{2p} \|x\|$ . This concludes the proof of the claim, and up to appropriate choice of  $\epsilon$ , that of the proposition.  $\square$

## 9. ITERATED TWISTING OF $\mathcal{F}$

The results in this section are the particular cases of [9, Cor. 2 and Prop.3] for the admissible families yielding Ferenczi's space. For the sake of completeness we include a rather complete sketch with somewhat different proofs. To unify the notation, let us set  $\mathcal{F}_1 = \mathcal{F}$ . As above,  $\mathcal{F}_2$  denote the self-extension of  $\mathcal{F}_1$  obtained in Section 8. As it is showed in Proposition 3.2,

$$\mathcal{F}_2 = \{(g'(\theta), g(\theta)) : g \in \mathcal{H}(X_{j,t})\},$$

endowed with the quotient norm of  $\mathcal{H}(X_{j,t})/(\ker \delta_\theta \cap \ker \delta'_\theta)$ . Let us show that the twisting process can be iterated obtaining a sequence  $(\mathcal{F}_n)$  of H.I. spaces such that  $\mathcal{F}_{n+m}$  is a twisted sum of  $\mathcal{F}_n$  and  $\mathcal{F}_m$ .

Given a function  $g \in \mathcal{H}(X_{j,t})$  and an integer  $k \in \mathbb{N}$ , we denote  $\hat{g}[k] := g^{(k-1)}(\theta)/(k-1)!$ , the  $(k)$ -th coefficient of the Taylor series of  $g$  at  $\theta$ . Following the constructions in [9], we define for  $n \geq 3$ :

$$\mathcal{F}_n := \{(\hat{g}[n], \dots, \hat{g}[2], \hat{g}[1]) : g \in \mathcal{H}(X_{j,t})\}$$

endowed with the quotient norm of  $\mathcal{H}(X_{j,t})/\bigcap_{k=0}^{n-1} \ker \delta_\theta^{(k)}$ .

**Proposition 9.1.** *Let  $m, n \in \mathbb{N}$  with  $m > n$ .*

- (1) The expression  $\pi_{m,n}(x_m, \dots, x_n, \dots, x_1) = (x_n, \dots, x_1)$  defines a surjective operator  $\pi_{m,n} : \mathcal{F}_m \rightarrow \mathcal{F}_n$ .
- (2) The expression  $i_{n,m}(x_n, \dots, x_1) := (x_n, \dots, x_1, 0, \dots, 0)$  defines a isomorphic embedding  $i_{n,m} : \mathcal{F}_n \rightarrow \mathcal{F}_m$  with  $\text{ran}(i_{n,m}) = \ker(\pi_{m,m-n})$ .
- (3) The operator  $\pi_{m,n}$  is strictly singular.

*Proof.* (1) Since  $\text{dist}(g, \bigcap_{k=0}^{n-1} \ker \delta_\theta^{(k)}) \leq \text{dist}(g, \bigcap_{k=0}^{m-1} \ker \delta_\theta^{(k)})$ , we have  $\|\pi_{m,n}\| \leq 1$ . And it is obvious that  $\pi_{m,n}$  is surjective.

(2) Let  $\phi \in H^\infty(\mathbb{S})$  be a scalar function such that  $\hat{\phi}[k] = \delta_{k,m-n}$  for  $1 \leq k \leq m$ . For the existence of  $\phi$ , we consider a conformal equivalence  $\varphi : \mathbb{S} \rightarrow \mathbb{D}$  satisfying  $\varphi(\theta) = 0$ , and the polynomial  $p(z) := (z - \theta)^{m-n}$ . The function  $p \circ \varphi^{-1} \in H(\mathbb{D})$  admits a representation  $p \circ \varphi^{-1}(\omega) = \sum_{l=0}^{\infty} a_l \omega^l$ , and it is not difficult to check that  $\phi(z) := \sum_{l=0}^m a_l \varphi(z)^l$  defines a function that satisfies the required conditions.

Given  $(x_n, \dots, x_1) \in \mathcal{F}_n$ , we take  $g \in \mathcal{H}(X_{j,t})$  such that  $\hat{g}[k] = x_k$  for  $k = 1, \dots, n$ . Then  $f := \phi \cdot g \in \mathcal{H}(X_{j,t})$  with  $\|f\| \leq \|\phi\|_\infty \cdot \|g\|$  and, by the Leibnitz rule,

$$\hat{f}[k] = \sum_{l=1}^k \hat{\phi}[l] \hat{g}[k-l].$$

Thus  $\hat{f}[k] = 0$  for  $1 \leq k \leq m-n$  and  $\hat{f}[k] = \hat{g}[k-m+n]$  for  $m-n < k \leq m$ ; i.e.,  $(\hat{f}[m], \dots, \hat{f}[1]) = (x_n, \dots, x_1, 0, \dots, 0)$ . Hence  $i_{n,m}$  is well-defined and  $\|i_{n,m}\| \leq \|\phi\|_\infty$ .

Clearly  $i_{n,m}$  is injective and  $\text{ran}(i_{n,m}) \subset \ker(\pi_{m,m-n})$ . Let  $(y_n, \dots, y_1, 0, \dots, 0) \in \ker(\pi_{m,m-n})$ . Then there exists  $g \in \mathcal{H}(X_{j,t})$  such that  $\hat{g}[k] = 0$  for  $1 \leq k \leq m-n$  and  $\hat{g}[k] = y_{k-m+n}$  for  $m-n < k \leq m$ . Since  $g$  has a zero of order  $m-n$  at  $\theta$ , there exists  $f \in \mathcal{H}(X_{j,t})$  such that  $g(z) = f(z)(z - \theta)^{m-n}$ , and it is not difficult to check that  $i_{n,m}(\hat{f}[n], \dots, \hat{f}[1]) = (y_n, \dots, y_1, 0, \dots, 0)$ .

(3) Since  $\pi_{m,n} = \pi_{m-1,n} \pi_{m,m-1}$  for  $m > n+1$ , it is enough to prove that  $\pi_{m,m-1}$  is strictly singular. We will do it by induction:

We proved in Theorem 8.1 that  $\pi_{2,1}$  is strictly singular. Let  $m > 2$  and assume that  $\pi_{m-1,m-2}$  is strictly singular. Note that  $\pi_{m,1} = \pi_{m,2} \pi_{2,1}$ ; hence  $\pi_{m,1}$  is also strictly singular.

We consider the following commuting diagram:

$$(14) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}_{m-1} & \xrightarrow{i_{m-1,m}} & \mathcal{F}_m & \xrightarrow{\pi_{m,1}} & \mathcal{F}_1 \longrightarrow 0 \\ & & \pi_{m-1,m-2} \downarrow & & \downarrow \pi_{m,m-1} & & \parallel \\ 0 & \longrightarrow & \mathcal{F}_{m-2} & \xrightarrow{i_{m-2,m-1}} & \mathcal{F}_{m-1} & \xrightarrow{\pi_{m-1,1}} & \mathcal{F}_1 \longrightarrow 0. \end{array}$$

By (1) and (2), the two rows are exact. Suppose that  $M$  is an infinite dimensional closed subspace of  $\mathcal{F}_m$  such that  $\pi_{m,m-1}|_M$  is an isomorphism. Since  $\pi_{m,m-1} i_{m-1,m}$  is strictly singular and  $\text{ran}(i_{m-1,m}) = \ker(\pi_{m,1})$ ,  $M \cap \ker(\pi_{m,1})$  is finite dimensional and  $M + \ker(\pi_{m,1})$  is closed. But this is impossible, because  $\pi_{m,1}$  is strictly singular.  $\square$

Since uniform convexity is a 3-space property [13], as an immediate consequence we get:

**Corollary 9.2.** *Let  $m, n \in \mathbb{N}$ . Then the sequence*

$$0 \longrightarrow \mathcal{F}_m \xrightarrow{i_{m,m+n}} \mathcal{F}_{m+n} \xrightarrow{\pi_{m+n,n}} \mathcal{F}_n \longrightarrow 0$$

*is exact and singular. Therefore, all the spaces  $\mathcal{F}_n$  are uniformly convex H.I.*

Next we show that there are natural nontrivial twisted sums of spaces  $\mathcal{F}_n$  which are not H.I. Let  $l, m, n \in \mathbb{N}$  with  $l > n$ . We consider the following push-out diagram:

$$(15) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}_l & \xrightarrow{i_{l,l+m}} & \mathcal{F}_{l+m} & \xrightarrow{\pi_{l+m,m}} & \mathcal{F}_m \longrightarrow 0 \\ & & \pi_{l,n} \downarrow & & \downarrow \pi_{l+m,n+m} & & \parallel \\ 0 & \longrightarrow & \mathcal{F}_n & \xrightarrow{i_{n,n+m}} & \mathcal{F}_{n+m} & \xrightarrow{\pi_{n+m,m}} & \mathcal{F}_m \longrightarrow 0. \end{array}$$

**Proposition 9.3.** *Let  $l, m, n \in \mathbb{N}$  with  $l > n$ . Then the diagonal push-out sequence*

$$(16) \quad 0 \longrightarrow \mathcal{F}_l \xrightarrow{i} \mathcal{F}_n \oplus \mathcal{F}_{l+m} \xrightarrow{\pi} \mathcal{F}_{m+n} \longrightarrow 0$$

*obtained from diagram (15) is a nontrivial exact sequence.*

*Proof.* As we saw in Section 2, the maps  $i$  and  $\pi$  are given by

$$i(x) = (-\pi_{l,n} x, i_{l,l+m} x) \quad \text{and} \quad \pi(y, z) = i_{n,n+m} y + \pi_{l+m,n+m} z,$$

and it is easy to check that the sequence (16) is exact. Since  $l > n$ , every operator from  $\mathcal{F}_l$  or  $\mathcal{F}_{m+n}$  into  $\mathcal{F}_n$  is strictly singular. Thus  $\mathcal{F}_l \oplus \mathcal{F}_{m+n}$  is not isomorphic to  $\mathcal{F}_n \oplus \mathcal{F}_{l+m}$ , and the exact sequence (16) is nontrivial.  $\square$

## REFERENCES

- [1] P. Aiena, *Fredholm and local spectral theory with applications to multipliers*, Kluwer Acad. Publ., Dordrecht, 2004.
- [2] S.A. Argyros and Deliyanni, *Examples of asymptotically  $\ell_1$  Banach spaces*, Trans. Amer. Math. Soc. 349 (1997) 973–995.
- [3] S.A. Argyros and A. Tolas, *Methods in the theory of hereditarily indecomposable Banach spaces*, Mem. A.M.S. 806, 2004.
- [4] A. Avilés, F. Cabello Sánchez, J.M.F. Castillo, M. González and Y. Moreno, *On separably injective Banach spaces*. Advances in Math. 234 (2013) 192–216.
- [5] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Springer-Verlag, 1976.
- [6] F. Cabello Sánchez, *There is no strictly singular centralizer on  $L_p$* , Proc. Amer. Math. Soc. 142 (2014) 949–955.
- [7] F. Cabello Sánchez and J.M.F. Castillo, *Duality and twisted sums of Banach spaces*, J. Funct. Anal. 175 (2000) 1–16.
- [8] F. Cabello Sánchez and J.M.F. Castillo, *Uniform boundedness and twisted sums of Banach spaces*, Houston J. Math. 30 (2004) 523–536.
- [9] F. Cabello Sánchez, J.M.F. Castillo and N.J. Kalton, *Complex interpolation and twisted Hilbert spaces*, To appear in Pacific J. Math.
- [10] F. Cabello Sánchez, J.M.F. Castillo and J. Suárez, *On strictly singular exact sequences*, Nonlinear Anal. 75 (2012) 3313–3321.
- [11] R. del Campo, A. Fernández, A. Manzano, F. Mayoral and F. Naranjo, *Complex interpolation of Orlicz spaces with respect to a vector measure*, Math. Nachr. 287 (2014) 23–31.
- [12] M.J. Carro, J. Cerdá and J. Soria, *Commutators and interpolation methods*, Ark. Mat. 33 (1995) 199–216.
- [13] J.M.F. Castillo and M. González, *Three-space problems in Banach space theory*, Lecture Notes in Math. 1667, Springer-Verlag, 1997.
- [14] J.M.F. Castillo and Y. Moreno, *Strictly singular quasi-linear maps*, Nonlinear Analysis - TMA 49 (2002) 897–904.
- [15] J.M.F. Castillo and Y. Moreno, *On the Lindenstrauss-Rosenthal theorem*, Israel J. Math. 140 (2004) 253–270.

- [16] J.M.F. Castillo and Y. Moreno, *On the bounded approximation property in Banach spaces*, Israel J. Math. 198 (2013) 243–59.
- [17] R.R. Coifman, M. Cwikel, R. Rochberg, Y. Sagher and G. Weiss, *A theory of complex interpolation for families of Banach spaces*, Advances in Math. 43 (1982) 203–229.
- [18] M. Cwikel, B. Jawerth, M. Milman and R. Rochberg, *Differential estimates and commutators in interpolation theory*, Analysis at Urbana II, London Math. Soc. L.N. 138 (1989), 170–220.
- [19] M. Cwikel, N. Kalton, M. Milman and R. Rochberg, *A unified theory of commutator estimates for a class of interpolation methods*, Advances in Math. 169 (2002) 241–312.
- [20] P. Enflo, J. Lindenstrauss and G. Pisier, *On the "three-space" problem for Hilbert spaces*, Math. Scand. 36 (1975), 199–210.
- [21] V. Ferenczi, *A uniformly convex hereditarily indecomposable Banach space*, Israel J. Math. 102 (1997), 199–225.
- [22] V. Ferenczi, *Quotient Hereditarily Indecomposable Banach spaces*, Canad. J. Math. 51 (1999) 566–584.
- [23] M. González and J.M. Herrera, *Finitely decomposable Banach spaces and the three-space property*, Archiv Math. 80 (2003) 647–654.
- [24] W.T. Gowers and B. Maurey, *The unconditional basic sequence problem*, J. Amer. Math. Soc. 6 (1993) 851–874.
- [25] J. Gustavsson and J. Peetre, *Interpolation of Orlicz spaces*. Studia Math. 60 (1977) 33–59.
- [26] N.J. Kalton, *The three-space problem for locally bounded  $F$ -spaces*, Compositio Math. 37 (1978) 243–276.
- [27] N.J. Kalton, *Convexity, type and the three space problem*, Studia Math. 69 (1980/81), 247–287.
- [28] N.J. Kalton, *Nonlinear commutators in interpolation theory*, Memoirs of the A.M.S. 385, 1988.
- [29] N.J. Kalton, *Differentials of complex interpolation processes for Köthe function spaces*, Trans. Amer. Math. Soc. 333 (1992) 479–529.
- [30] N.J. Kalton, *An elementary example of a Banach space not isomorphic to its complex conjugate*, Canad. Math. Bull. 38 (1995) 218–222.
- [31] N.J. Kalton and S. Montgomery-Smith, *Interpolation of Banach spaces*, Chapter 26 in “Handbook of the geometry of Banach spaces, vol. 2; W. B. Johnson and J. Lindenstrauss (eds.) Elsevier 2003, pp. 1131–1175.
- [32] N.J. Kalton and N.T. Peck, *Twisted sums of sequence spaces and the three-space problem*, Trans. Amer. Math. Soc. 255 (1979) 1–30.
- [33] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I*, Springer-Verlag, 1977.
- [34] G. Pisier, *Holomorphic semi-groups and the geometry of Banach spaces*, Ann. of Math. 115 (1982) 375–392.
- [35] G. Pisier and H. Xu, *Non-commutative  $L^p$ -spaces*, Chapter 34 in “Handbook in the geometry of Banach spaces vol. 2”, W.B. Johnson and J. Lindenstrauss (eds.) Elsevier 2003, pp. 1459–1518.
- [36] R. Rochberg and G. Weiss, *Derivatives of analytic families of Banach spaces*, Ann. of Math. 118 (1983) 315–347.
- [37] M. Schechter, *Complex interpolation*, Compositio Math. 18 (1967) 117–147.
- [38] J. Suárez de la Fuente, *The Kalton centralizer on  $L_p[0, 1]$  is not strictly singular*, Proc. Amer. Math. Soc. 141 (2013) 3447–3451.
- [39] F. Watbled, *Complex interpolation of a Banach space with its dual*, Math. Scand. 87 (2000) 200–210.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, AVENIDA DE ELVAS S/N, 06011 BADAJOZ, SPAIN

*E-mail address:* castillo@unex.es

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, RUA DO MATÃO 1010, 05508-090 SÃO PAULO SP, BRAZIL, AND  
EQUIPE D’ANALYSE FONCTIONNELLE, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UNIVERSITÉ PIERRE ET MARIE CURIE - PARIS 6, CASE 247, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE.

*E-mail address:* ferenczi@ime.usp.br

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE CANTABRIA, AVENIDA DE LOS CASTROS S/N, 39071 SANTANDER, SPAIN

*E-mail address:* manuel.gonzalez@unican.es