

AVERAGE MIXED VOLUME UNDER PROJECTION

GREGORIO MALAJOVICH

ABSTRACT. The average mixed volume of a random projection of d convex bodies in \mathbb{R}^n is bounded above in terms of a quermassintegral.

1. INTRODUCTION

The *mixed volume* of the n -tuple of convex bodies $(\mathcal{A}_1, \dots, \mathcal{A}_n)$, $\mathcal{A}_i \subset \mathbb{R}^n$ is defined as

$$V(\mathcal{A}_1, \dots, \mathcal{A}_n) \stackrel{\text{def}}{=} \frac{1}{n!} \frac{\partial^n}{\partial t_1 \partial t_2 \dots \partial t_n} \text{Vol}(t_1 \mathcal{A}_1 + \dots + t_n \mathcal{A}_n)$$

where $t_1, \dots, t_n \geq 0$ and the derivative is taken at $t = 0$. The normalization factor $1/n!$ ensures that $V(\mathcal{A}, \dots, \mathcal{A}) = \text{Vol}(\mathcal{A})$. In that sense the mixed volume generalizes the usual volume.

The definition above is still valid if we allow the \mathcal{A}_i to have empty volume, so in this paper we will only require the \mathcal{A}_i to be closed convex sets.

The mixed volume was introduced by Minkowski (1901) for $n = 3$. If $\mathcal{A} \subset \mathbb{R}^3$ is a convex body, the Steiner formula

$$\text{Vol}(\mathcal{A} + \epsilon B^3) = \text{Vol}(\mathcal{A}) + S\epsilon + \pi B\epsilon^2 + \frac{4}{3}\pi\epsilon^3$$

implies that

$$V(\mathcal{A}, \mathcal{A}, B^3) = 3S \quad \text{and} \quad V(\mathcal{A}, B^3, B^3) = 3\pi B$$

with S is the total area of $\partial\mathcal{A}$ and B its total mean curvature (assuming $\partial\mathcal{A}$ is smooth). The quantities $V(\mathcal{A}, \dots, \mathcal{A}, B^n, \dots, B^n)$ are known as *quermassintegrals*. There is a vast literature on recovering quantities such as the area of $\partial\mathcal{A}$ from the area of projections of \mathcal{A} .

However, most of the literature deals with mixed volumes for two different convex bodies, one being the ball B^n . Mixed volumes with many possibly different convex bodies seem to arise mostly in connection with counting zeros of sparse polynomial systems or obtaining a starting system for path-following. The main result in this note arised during the complexity analysis of such an algorithm.

Let $G(d, n)$ denote the Grassmannian manifold of d -dimensional subspaces of \mathbb{R}^n . It is endowed with the unique probability measure invariant under the orthogonal group $O(n)$. In this note we prove the following result:

Date: July 5, 2018.

2010 Mathematics Subject Classification. Primary 52A39.

Key words and phrases. mixed volume.

This paper was written while visiting the Simons Institute for the Theory of Computing in the University of California at Berkeley. This visit was funded by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil. Proc. BEX 2388/14-6).

Theorem 1. *Let $\mathcal{A}_1, \dots, \mathcal{A}_d$ be compact convex sets in \mathbb{R}^n . Then,*

$$\operatorname{Avg}_{P \in G(d,n)} V(P(\mathcal{A}_1), \dots, P(\mathcal{A}_d)) \leq \frac{\operatorname{Vol}(B^d)}{\operatorname{Vol}(B^n)} V(\mathcal{A}_1, \dots, \mathcal{A}_d, B^n, \dots, B^n).$$

By replacing \mathcal{A}_i with B^n and $P(\mathcal{A}_i)$ by B^d , one quickly checks that the bound above is sharp.

2. PROOF OF THE THEOREM

Before proving theorem 1, we will restate it in a more convenient form. If $Q \in O(n)$, let q_j denote the j -th row of Q . If Q is uniformly distributed in $O(n)$ with respect to the Haar measure, the projection $P : \mathbb{R}^n \rightarrow \mathbb{R}^d$,

$$P : x \mapsto \begin{bmatrix} q_1 x \\ \vdots \\ q_d x \end{bmatrix}$$

is uniformly distributed in $G(d, n)$. We have

$$d! V(P(\mathcal{A}_1), \dots, P(\mathcal{A}_d)) = n! V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_n]).$$

Therefore, what we need to prove is the following restatement of Theorem 1:

Theorem 1-A. *Let $\mathcal{A}_1, \dots, \mathcal{A}_d$ be compact convex sets in \mathbb{R}^n . Then,*

$$\begin{aligned} \operatorname{Avg}_{Q \in O(n)} V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_{d+n}]) &\leq \\ &\leq \frac{d! \operatorname{Vol}(B^d)}{n! \operatorname{Vol}(B^n)} V(\mathcal{A}_1, \dots, \mathcal{A}_d, B^n, \dots, B^n). \end{aligned}$$

Remark 2. Explicitly,

$$\frac{d! \operatorname{Vol}(B^d)}{n! \operatorname{Vol}(B^n)} = \frac{1}{(2\sqrt{\pi})^{n-d}} \frac{\Gamma\left(\frac{d+1}{2}\right)}{\Gamma\left(\frac{n+1}{2}\right)}.$$

This is an immediate consequence of the duplication formula (Abramowitz and Stegun, 1964, formula 6.1.18),

$$\Gamma(2z) = \frac{1}{\sqrt{2\pi}} 2^{2z-1/2} \Gamma(z) \Gamma(z + 1/2).$$

Let CONV_n denote the set of all compact convex subsets of \mathbb{R}^n . This set is closed under Minkowski sum and under multiplication by non-negative real numbers. In that sense, CONV_n is an ‘algebra’ over the non-negative real numbers. We recall the following definitions:

Definition 3. A function $F : \operatorname{CONV}_n \rightarrow \mathbb{R}^+$ is linear if for all $t_1, t_2 \in \mathbb{R}^+$ and for all $\mathcal{A}_1, \mathcal{A}_2 \in \operatorname{CONV}_n$, $F(t_1\mathcal{A}_1 + t_2\mathcal{A}_2) = t_1F(\mathcal{A}_1) + t_2F(\mathcal{A}_2)$.

Definition 4. A function $F : \operatorname{CONV}_n \rightarrow \mathbb{R}^+$ is monotone if $\mathcal{A} \subseteq \mathcal{B}$ implies that $F(\mathcal{A}) \leq F(\mathcal{B})$.

The mixed volume $V(\mathcal{A}_1, \dots, \mathcal{A}_n)$ is linear and monotone in each of the variables. The key for proving theorem 1 is the following Lemma, that I could not find it in the literature.

Lemma 5. *Let $F : \text{CONV}_k \rightarrow \mathbb{R}^+$ be linear and monotone. Then*

$$\text{Avg}_{c \in S^{k-1}} (F([0, c])) \leq r_k F(B^k)$$

with $r_k = \frac{\Gamma(k/2)}{\sqrt{\pi}(k-1)\Gamma(\frac{k-1}{2})}$. This constant r_k is the smallest with such a property.

Proof. For every $\epsilon > 0$, there is a closed covering of S^{k-1} into measurable subsets of radius ϵ , say $S^{k-1} = \cup_{\lambda \in \Lambda} V_\lambda$, and such that $V_\lambda \cap V_{\lambda'}$ has measure zero for $\lambda \neq \lambda'$. The sets $W_\lambda = \text{Conv}(\{0\} \cup V_\lambda)$ are a closed covering of B^k with the same property.

$$\begin{aligned} \text{Avg}_{c \in S^{k-1}} (F([0, c])) &= \frac{1}{\text{Vol}S^{k-1}} \int_{S^{k-1}} F([0, c]) \, dS^{k-1}(c) && \text{by definition,} \\ &= \frac{1}{\text{Vol}S^{k-1}} \sum_{\lambda \in \Lambda} \int_{V_\lambda} F([0, c]) \, dS^{k-1}(c) && \text{trivially,} \\ &\leq \frac{1}{\text{Vol}S^{k-1}} \sum_{\lambda \in \Lambda} \int_{V_\lambda} F(W_\lambda) \, dS^{k-1}(c) && \text{by monotonicity,} \\ &\leq \frac{1}{\text{Vol}S^{k-1}} \sum_{\lambda \in \Lambda} (\text{Vol}V_\lambda) F(W_\lambda) && \text{trivially,} \\ &\leq \frac{1}{\text{Vol}S^{k-1}} F \left(\sum_{\lambda \in \Lambda} (\text{Vol}V_\lambda) W_\lambda \right) && \text{by linearity} \end{aligned}$$

After an orthogonal change of coordinates, assume that $\rho(\epsilon)e_1$ is a point of largest norm in $\sum_{\lambda \in \Lambda} (\text{Vol}V_\lambda) W_\lambda$. In particular, there are $x_\lambda \in W_\lambda$ such that $\rho(\epsilon)e_1 = \sum_{\lambda \in \Lambda} (\text{Vol}V_\lambda)x_\lambda$. So,

$$\rho(\epsilon) = \sum_{\lambda \in \Lambda} (\text{Vol}V_\lambda)(x_\lambda)_1 \leq \sum_{(x_\lambda)_1 \geq 0} \text{Vol}V_\lambda \frac{(x_\lambda)_1}{\|x_\lambda\|}.$$

Passing to the limit,

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \rho(\epsilon) &= \int_{S^{k-1} \cap [y_1 \geq 0]} y_1 \, dS^{k-1}(y) \\ &= \int_{S^{k-1} \cap [y_1 \geq 0]} e_1 \cdot dn(y) && \text{interpreting as a flow,} \\ &= \int_{B^k \cap [y_1 \geq 0]} \text{div}(e_1) \, dB^k(y) - \int_{B^{k-1}} e_1 \cdot dn(y) && \text{by Gauss theorem,} \\ &= 0 - \int_{B^{k-2}} -1 \, dB^{k-1}(y) \\ &= \text{Vol}B^{k-1} \end{aligned}$$

By monotonicity of F and then linearity, we conclude that

$$\text{Avg}_{c \in S^{k-1}} (F([0, c])) \leq \frac{1}{\text{Vol}S^{k-1}} F(\lim_{\epsilon \rightarrow 0} \rho(\epsilon) B^k) \leq r_k F(B^k)$$

with $r_k = \text{Vol}B^{k-1}/\text{Vol}S^{k-1}$. Using the well-known formulas

$$\text{Vol}B^k = \frac{2\pi^{k/2}}{k \Gamma\left(\frac{k}{2}\right)} \quad \text{and} \quad \text{Vol}S^{k-1} = \frac{2\pi^{k/2}}{\Gamma\left(\frac{k}{2}\right)}$$

we deduce that

$$r_k = \frac{\Gamma\left(\frac{k}{2}\right)}{\sqrt{\pi}(k-1) \Gamma\left(\frac{k-1}{2}\right)}.$$

To establish sharpness, we consider the operator $F(X) = V(B^k, \dots, B^k, X)$. Because of the properties of the mixed volume,

$$F([0, c]) = V(B^k \cap c^\perp, \dots, B^k \cap c^\perp, [0, c]).$$

Thus,

$$\text{Avg}_{c \in S^{k-1}} F([0, c]) = F([0, e_k]) = V(B^{k-1} \times \{0\}, \dots, B^{k-1} \times \{0\}, [0, e_k]).$$

This last value is by definition $\frac{1}{k!}$ times the coefficient in $t_1 t_2 \cdots t_k$ of

$$\text{Vol}(t_1 + \cdots + t_{k-1})B^{k-1} + t_k[0, e_k] = (t_1 + \cdots + t_{k-1})^{k-1} t_k \text{Vol}B^{k-1}.$$

Thus,

$$\text{Avg}_{c \in S^{k-1}} F([0, c]) = \frac{1}{k} \text{Vol}B^{k-1}$$

while the predicted value is $r_k \text{Vol}B^k = \frac{\text{Vol}B^{k-1}}{\text{Vol}S^{k-1}} \text{Vol}B^k = \frac{1}{k} \text{Vol}B^{k-1}$. \square

Proof of Theorem 1. The last row q_n of Q is uniformly distributed in S^{n-1} , so

$$\begin{aligned} \text{Avg}_{Q \in SO(n)} (V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_n])) &= \\ &= \text{Avg}_{q_n \in S^{n-1}} \left(\text{Avg} (V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_n])) \right) \end{aligned}$$

where the inner average is taken over orthonormal frames $(q_{d+1}, \dots, q_{n-1})$ that are orthogonal to q_n . By Lemma 5,

$$\begin{aligned} \text{Avg}_{Q \in SO(n)} (V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_n])) &= \\ &= r_n \text{Avg} (V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_{n-1}], B^n)). \end{aligned}$$

Repeating the same argument,

$$\begin{aligned} \text{Avg}_{Q \in SO(n)} V(\mathcal{A}_1, \dots, \mathcal{A}_d, [0, q_{d+1}], \dots, [0, q_n]) & \\ &\leq r_{d+1} r_{d+2} \cdots r_n V(\mathcal{A}_1, \dots, \mathcal{A}_d, B^n, \dots, B^n). \end{aligned}$$

Finally,

$$r_{d+1} r_{d+2} \cdots r_n = \frac{\Gamma(n/2) \Gamma(d)}{\pi^{\frac{n-d}{2}} \Gamma(d/2) \Gamma(n)}$$

\square

REFERENCES

- Abramowitz, Milton and Irene A. Stegun. 1964. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, vol. 55, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. MR0167642 (29 #4914)
- Minkowski, Hermann. 1901. *Sur les surfaces convexes fermées*, C.R. Acad.Sci., Paris **132**, 21–24.

DEPARTAMENTO DE MATEMÁTICA APLICADA, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO. CAIXA POSTAL 68530, RIO DE JANEIRO RJ 21941-909, BRASIL.

E-mail address: `gregorio.malajovich@gmail.com`