

Superheavy Dark Matter in Light of Dark Radiation

Jong-Chul Park* and Seong Chan Park†

Department of Physics, Sungkyunkwan University, Suwon 440-746, Korea

Abstract

Superheavy dark matter can satisfy the observed dark matter abundance if the stability condition is fulfilled. Here, we propose a new Abelian gauge symmetry $U(1)_H$ for the stability of superheavy dark matter as the electromagnetic gauge symmetry to the electron. The new gauge boson associated with $U(1)_H$ contributes to the effective number of relativistic degrees of freedom in the universe as dark radiation, which has been recently measured by several experiments, *e.g.*, PLANCK. We calculate the contribution to dark radiation from the decay of a scalar particle via the superheavy dark matter in the loop. Interestingly enough, this scenario will be probed by a future LHC run in the invisible decay signatures of the Higgs boson.

PACS numbers: 11.30.-j, 12.90.+b, 95.35.+d, 98.80.Cq

Keywords: Superheavy dark matter, Dark radiation, Higgs invisible decay, Planck, LHC

I. INTRODUCTION

Although dark matter (DM) is one of the most important constituent of the universe [1], its nature is still almost unknown. Indeed, we still have no concrete evidence for the weakly interacting massive particle (WIMP) from any experiments including the Large Hadron Collider (LHC) and DM direct searches below $\mathcal{O}(\text{TeV})$ even though the WIMP or “heavy neutrino” [2] has been regarded as a promising candidate since the late 1970s. On the other hand, heavier DM

particles, $\gg \mathcal{O}(\text{TeV})$, have been relatively less studied due to a mass limit from the so-called *unitarity bound*. If this bound is applied on the DM relic density, for a Majorana fermion, one can obtain the following relation [3]:

$$\Omega_{\text{DM}} h^2 \geq 1.7 \times 10^{-6} \sqrt{x_f} \left[\frac{m_{\text{DM}}}{1\text{TeV}} \right]^2, \quad (1)$$

where $x_f = m_{\text{DM}}/T_f$ and T_f is the freeze-out temperature. The result is similar for a scalar, while $\Omega_{\text{DM}} h^2$ is a factor of 2 larger for a Dirac fermion. To avoid the overclosure of the universe, we should set $\Omega_{\text{DM}} h^2 \leq 1$. Thus, we find the upper limit on the DM mass, $m_{\text{DM}} \lesssim 340\text{TeV}$ for $x_f \approx 28$. If the current bound on the DM abundance is

*Electronic address: log1079@gmail.com

†Electronic address: s.park@skku.edu

used, $\Omega_{\text{DM}}h^2 \approx 0.12$ [4], the constraint can be stronger, $m_{\text{DM}} \lesssim 120\text{TeV}$. In addition, other crucial challenges exist for heavy DM particles:

- *Instability of DM:* A heavier particle is generally more unstable if no symmetry principle precludes its decay.
- *Lack of testability:* In currently ongoing or future experiments, heavier DM well above $\mathcal{O}(\text{TeV})$ is difficult to test [1].

One can evade the unitarity bound in the case of superheavy DM with mass $m_{\text{DM}} \approx 10^{12-14}$ GeV, which is generically called as *WIMPZILLA*. Independently of the nongravitational interactions, *WIMPZILLA* particles can be produced in various ways and satisfy the observed DM abundance: *i.e.*, from the expansion of the background spacetime at the end of the inflation or in the bubble collision in a first-order phase transition. For more details, see, *e.g.*, Refs. [5] and [6].

On the other hand, for *WIMPZILLA*, the instability problem gets worse due to the extremely high mass. In the presence of gravity, a global symmetry does not work, so all the stable particles need gauge symmetries [7, 8]. In this work, for the stability of *WIMPZILLA*, we introduce an Abelian group $U(1)_H$, which similarly acts as $U(1)_{em}$ for the electron in the standard model (SM).

Moreover, a new massless gauge boson or hidden photon associated with $U(1)_H$ could provide a possibility to test *WIMPZILLA*. The new boson does not directly interact with the SM sector, thus, we regard this new boson as dark radiation. Dark radiation could have left evidence of its presence in various stages of cosmological history. Recently, observations show that additional relativistic particles may have existed at the Big Bang nucleosynthesis (BBN) time and at the recombination era shown in cosmic microwave background radiation (CMBR). In the rest of this work, we will introduce a model with *WIMPZILLA* and study the testability of this *WIMPZILLA* via the associated dark radiation in BBN and CMBR [9, 10] and the possible signatures at the LHC [11].

II. THE MODEL

We propose the minimal model including two hidden sector particles: a Dirac fermion ψ and a scalar ϕ . The fermion ψ , the DM candidate, is only charged under the ‘hidden’ Abelian gauge symmetry $U(1)_H$. The scalar particle ϕ is neutral under $U(1)_H$ as well as the SM gauge symmetries, but the late time decay of ϕ into hidden photons is responsible for dark radiation. The gauge invariant




FIG. 1: Scalar particle ϕ decaying into two hidden photons through the WIMPZILLA ψ loop.

Lagrangian[25] is given by

$$\mathcal{L} \supset \mathcal{L}_{SM} + \mathcal{L}_\phi + \lambda_{\phi H} \phi^2 (H^\dagger H) - \frac{1}{4} F_{H\mu\nu} F_H^{\mu\nu} - y_\psi \phi \bar{\psi} \psi + i \bar{\psi} \gamma^\mu (\partial_\mu - i g_H A_\mu^H) \psi - m_\psi \bar{\psi} \psi, \quad (2)$$

where $\mathcal{L}_\phi = \frac{1}{2}[(\partial\phi)^2 - m_\phi^2\phi^2]$ and the ‘Higgs portal’ interaction [14] is allowed. The new fermion can be naturally heavy enough due to the vectorlike mass term and, thus, is consistent with the WIMPZILLA picture. For the late time decay, however the new scalar is required to be much lighter than the WIMPZILLA, which provides a similar hierarchy problem as the SM Higgs boson. If a (super)symmetry is allowed in the hidden sector, the scalar boson can be protected from the radiative correction. In addition, unnecessary terms such as $\phi H^\dagger H$ are automatically forbidden. The details will be given in Ref. [15].

The scalar particle decays into two hidden photons through a diagram with the virtual ψ in the loop, as shown in Fig. 1. The decay rate of ϕ is strongly suppressed by the

superheavy mass of WIMPZILLA ψ :

$$\frac{1}{\tau_{\phi \rightarrow 2\gamma_H}} = \Gamma_{\phi \rightarrow 2\gamma_H} \approx \frac{(\alpha_H y_\psi)^2 m_\phi^3}{144\pi^3 m_\psi^2}, \quad (3)$$

where $\alpha_H = g_H^2/4\pi$. With appropriate parameters, the life time of ϕ can lie in an interesting epoch: the BBN epoch, $t_{\text{BBN}} \approx \mathcal{O}(0.1 - 1000)\text{s}$, or the CMB epoch, $t_{\text{CMB}} \approx 1.2 \times 10^{13}\text{s}$. Consequently, we could test this WIMPZILLA model by observing dark radiation in these epochs.

III. OBSERVATIONAL LIMITS

Now, we will consider the contributions to dark radiation observed in BBN and CMB data and the invisible decay of the Higgs at the LHC.

A. Dark Radiation

The recent observational results on the effective number of relativistic degrees of freedom, N_{eff} , are as follows:

$$\begin{aligned} N_{\text{eff}}^{\text{CMB}} &= 3.30 \pm 0.27 \quad (\text{Planck 2013 [4]}), \\ N_{\text{eff}}^{\text{CMB}} &= 3.84 \pm 0.40 \quad (\text{WMAP9 [16]}), \quad (4) \\ N_{\text{eff}}^{\text{BBN}} &= 3.71_{-0.45}^{+0.47} \quad (\text{BBN [17]}), \end{aligned}$$

which show some deviations, in particular, in WMAP9 and the BBN results compared to the SM expectation, $N_{\text{eff}}^{\text{SM}} = 3.046$ [18]. One should note that adding the independent H_0 measurement to the Planck CMB data provides $N_{\text{eff}} = 3.62 \pm 0.25$, which corresponds to

a 2.3σ deviation from the SM expectation [4]. Indeed, a new relativistic degree of freedom, the so-called *Dark radiation*, alleviates the tension between the CMB data and H_0 .

In this model, the late time decay of ϕ , in contact with the SM sector through the Higgs portal interaction, provides sizable dark radiation contributions at the BBN and the CMB epochs. If ϕ has never dominated the expansion of the universe, the extra contribution to N_{eff} , ΔN_{eff} , by its decay [19] is computed with $Y_\phi = n_\phi/s$, the actual number of particles per comoving volume and with the mass and life time of ϕ by using a simple relation [9, 20]:

$$\Delta N_{\phi \text{ decay}}^{\text{CMB}} \simeq 8.3(Y_\phi m_\phi/\text{MeV})(\tau_\phi/\text{s})^{1/2}. \quad (5)$$

Besides the dark radiation component coming from the ϕ decay, the primordial hidden photon γ_H can contribute to ΔN_{eff} . In this case, the primordial contribution is, however, quite suppressed as $\Delta N_{\text{primo } \gamma_H} \approx 0.053$, which is well explained in Ref. [21].

Finally, the total contribution of this WIMPZILLA model to N_{eff} is given by

$$\begin{aligned} \Delta N_{\text{eff}}^{\text{WZ}} &= N_{\text{eff}} - N_{\text{eff}}^{\text{SM}} \\ &= \Delta N_{\phi \text{ decay}}^{\text{CMB}} + \Delta N_{\text{primo } \gamma_H}, \end{aligned} \quad (6)$$

which should be compared with the observational results in Eq. (4), in particular, the most stringent Planck 2013 result [4]. The lower region below thick-solid line in Fig. 2,




FIG. 2: Contour plots for $\text{BR}_{h \rightarrow 2\phi}$ in the $m_\phi - \lambda_{\phi H}$ plane. The shaded region is constrained by the conservative invisible Higgs decay width limit $\text{BR}_{\text{inv}} < 0.34$ at the 95% C.L.. The lower region below thick-solid line is excluded by the Planck limit on N_{eff} at the 95% C.L. when $\tau_\phi = 1\text{s}$.

where $\tau_\phi = 1\text{ s}$ is chosen as a reference value, is excluded by the Planck result at the 95% confidence level (C.L.) due to the large contribution to N_{eff} . The dip around $m_\phi = m_h/2$ appears because of the resonant s -channel annihilation of ϕ into SM particles mediated by the Higgs boson. $\Delta N_{\text{eff}}^{\text{WZ}}$ at the CMB and BBN epochs is discussed in detail in Ref. [21].

B. Invisible Decay of the Higgs

In our scenario, we have another important possibility to test the model in collider experiments because the hidden scalar ϕ interacts with the SM Higgs boson h . The new decay channel of h to ϕ is open due to the non-vanishing ‘Higgs-portal’ interaction when kinematically allowed. The decay rate is given by

$$\Gamma_{h \rightarrow 2\phi} \simeq \frac{\lambda_{\phi H}^2}{32\pi} \frac{v^2}{m_h} \sqrt{1 - \frac{4m_\phi^2}{m_h^2}}. \quad (7)$$

Recently, the invisible branching fraction of the Higgs was measured by ATLAS [22] and CMS [23]. A global fit analysis based on all the Higgs search data provides a constraint on the invisible branching fraction of the Higgs: $\text{BR}_{\text{inv}} = \Gamma_{h \rightarrow \text{invisible}}/\Gamma_{h \rightarrow \text{all}} < 0.24$ at the 95% C.L. assuming the SM decay rates for the other decay modes and $\text{BR}_{\text{inv}} < 0.34$ at the 95% C.L. allowing non-standard values for $h \rightarrow \gamma\gamma$ and $h \leftrightarrow gg$ [24]. We finally obtained a limit on the Higgs portal coupling which is shown in Fig. 2.

IV. CONCLUSION

WIMPZILLA with $m_{DM} \approx 10^{12-14}$ GeV can satisfy the required DM relic density when a new gauge symmetry $U(1)_H$ protects WIMPZILLA from decay as $U(1)_{em}$ for the electron. The new gauge boson of $U(1)_H$ can provide a possibility of testing this simple WIMPZILLA model by tracing dark radiation in the BBN and the CMBR data. In addition, we may (dis)prove this model by measuring the invisible decay of the Higgs at collider experiments.

Acknowledgments

This work is supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology NRF-2013R1A1A2061561 (JC), NRF-2013R1A1A2064120 (SC). We appreciate APCTP for its hospitality during completion of this work.

[1] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267**, 195 (1996); G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005); J. L. Feng, Ann. Rev. Astron. Astrophys. **48**, 495 (2010).

[2] B. W. Lee and S. Weinberg, Phys. Rev.

Lett. **39**, 165 (1977).

[3] K. Griest and M. Kamionkowski, Phys. Rev. Lett. **64**, 615 (1990).

[4] P. A. R. Ade *et al.* [Planck Collaboration], [arXiv:1303.5076 [astro-ph.CO]].

[5] D. J. H. Chung, E. W. Kolb and A. Ri-

- otto, Phys. Rev. D **59**, 023501 (1999); D. J. H. Chung, E. W. Kolb and A. Riotto, Phys. Rev. Lett. **81**, 4048 (1998); D. J. H. Chung, E. W. Kolb and A. Riotto, Phys. Rev. D **60**, 063504 (1999).
- [6] D. J. H. Chung, P. Crotty, E. W. Kolb and A. Riotto, Phys. Rev. D **64**, 043503 (2001).
- [7] R. Kallosh, A. D. Linde, D. A. Linde and L. Susskind, Phys. Rev. D **52**, 912 (1995).
- [8] T. Banks and N. Seiberg, Phys. Rev. D **83**, 084019 (2011).
- [9] J. L. Menestrina and R. J. Scherrer, Phys. Rev. D **85**, 047301 (2012).
- [10] W. Fischler and J. Meyers, Phys. Rev. D **83**, 063520 (2011); M. C. Gonzalez-Garcia, V. Niro and J. Salvado, JHEP **1304**, 052 (2013); J. Hasenkamp and J. Kersten, JCAP **1308**, 024 (2013).
- [11] C. S. Kim, S. C. Park, K. Wang and G. Zhu, Phys. Rev. D **81**, 054004 (2010).
- [12] B. Holdom, Phys. Lett. B **166**, 196 (1986); L. B. Okun, Sov. Phys. JETP **56**, 502 (1982) [Zh. Eksp. Teor. Fiz. **83**, 892 (1982)].
- [13] K. R. Dienes, C. F. Kolda and J. March-Russell, Nucl. Phys. B **492**, 104 (1997); J. -H. Huh, J. E. Kim, J. -C. Park and S. C. Park, Phys. Rev. D **77**, 123503 (2008); E. J. Chun and J. -C. Park, JCAP **0902**, 026 (2009); E. J. Chun, J. -C. Park and S. Scopel, JHEP **1102**, 100 (2011); G. Belanger and J. -C. Park, JCAP **1203**, 038 (2012); J. -C. Park and S. C. Park, Phys. Lett. B **718**, 1401 (2013).
- [14] T. Binoth and J. J. van der Bij, Z. Phys. C **75**, 17 (1997); R. Schabinger and J. D. Wells, Phys. Rev. D **72**, 093007 (2005).
- [15] J. C. Park and S. C. Park, in preparation.
- [16] G. Hinshaw *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **208**, 19 (2013).
- [17] G. Steigman, Adv. High Energy Phys. **2012**, 268321 (2012).
- [18] G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti and P. D. Serpico, Nucl. Phys. B **729**, 221 (2005).
- [19] D. Hooper, F. S. Queiroz and N. Y. Gnedin, Phys. Rev. D **85**, 063513 (2012); K. Choi, K. -Y. Choi and C. S. Shin, Phys. Rev. D **86**, 083529 (2012); P. Di Bari, S. F. King and A. Merle, Phys. Lett. B **724**, 77 (2013); C. Kelso, S. Profumo and F. S. Queiroz, Phys. Rev. D **88**, 023511 (2013).
- [20] R. J. Scherrer and M. S. Turner, Astrophys. J. **331**, 33 (1988).
- [21] J. -C. Park and S. C. Park, arXiv:1305.5013 [hep-ph].
- [22] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012).
- [23] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).
- [24] P. P. Giardino, K. Kannike, I. Masina, M. Raidal and A. Strumia,

[arXiv:1303.3570 [hep-ph]].

[25] The kinetic mixing term ($\sim F_{\mu\nu}F_H^{\mu\nu}$) can be another source of interactions between the hidden and the SM sectors [12, 13] if there

exists a bi-charged particle under the gauge symmetries of both sectors. However, such a particle is absent in this model.