

Purely affine gravity

Aureliano Skirzewski^{1,*} and Oscar Castillo-Felisola^{2,3,†}

¹*Centro de Física Fundamental, Universidad de los Andes,
5101 Mérida, Venezuela.*

²*Centro Científico Tecnológico de Valparaíso,
Casilla 110-V, Valparaíso, Chile.*

³*Departamento de Física, Universidad Técnica Federico Santa María,
Casilla 110-V, Valparaíso, Chile.*

We present a purely affine gravitational model in four dimensions built up entirely on the bases of full diffeomorphism invariance, and power-counting renormalizability. We show that its non-relativistic limit around a homogeneous and isotropic spacetime yields to a Newtonian gravity.

PACS numbers: 02.40.Ma,04.50.Kd,04.90.+e

Keywords: Affine Gravity, Torsion, Generalised Gravity.

I. INTRODUCTION

In a critique to Newtonian mechanics, Mach proposed that inertial forces should have a dynamical rather than a kinematical origin (for a deeper discussion on the subject of Mach's principle see Ref. [1] and references therein). Any locally Minkowskian metric in the kinematics of the description of spacetime will introduce a notion of inertial forces at a microscopic level [2]. Hence, we will explore the dynamical origin of inertial forces, studying the dynamics of the affine connection of a non-metric manifold with torsion. To this end, we use the most general power-counting renormalizable action that includes only the gauge connection associated to diffeomorphisms invariance.

During the last years an increasing amount of alternative theories of gravity have being built and tested. Yet, General Relativity (GR) has proven to be the most successful theory of gravity [3]. However, we consider as a problem that the standard quantization procedure cannot be applied properly on GR. Moreover, not only it is not renormalizable, but there are problems with the choice of variables to be quantized and the choice of the Hilbert space to be used. Additionally, to sum over all possible field configurations of the metric seems to be wrong, as this would imply summing Euclidean and Minkowski like contributions to the transition amplitudes on equal terms. We might also consider the difficulties of in the quantization of non-polynomial field theories, and more specifically the square root of the metric that appears in the Hamiltonian in an ADM formulation of GR.

In order to bypass some of these issues, several approaches have been designed that use the connection as a fundamental field. For instance, a well-known example comes from the context of Cartan formulations of gravity, using the relation between the Weitzenböck and

Levi-Civita connections it is possible to obtain an equivalent Lagrangian to the one by Einstein and Hilbert, as a function of the torsion field. This approach is known as Teleparallel Gravity (see Ref. [4–6] and references within).

Furthermore, another alternative description of GR developed initially by Ashtekar uses the spin connection as the fundamental field and the frame field turns out to be its canonically conjugated momentum. In the context of Loop Quantum Gravity (LQG), using Ashtekar connection, a successful quantization program has been achieved [7, 8]. Originally, this approach towards quantum gravity addressed the concerns of the quantization of non-polynomial functions of the gravitational field, but later on it turned out that diffeomorphisms symmetry would not show up when the quantum operators were not of the correct density weight, which forces one to reintroduce the squared root [9]. Some strengths of this quantization program lies within a theorem by H. Sahlmann *et al.* in Ref. [10] that states the only diffeomorphisms invariant Hilbert space that supports the Heisenberg algebra, for the connection and its associated momentum, is the one of LQG. In spite of its success, LQG has not advanced enough to conclude that its low energy effective description is GR. Currently, there is no good description for the LQG effective model at other scales, nor its continuum spacetime limit either. Therefore, we cannot conclude that the search for a fundamental theory of gravitational interactions has ended. On the contrary, there are increasingly many alternatives to the usual metric description of gravity and they all must be tested against experiments and observations.

The earliest model that argues a description of gravitational interaction in terms of connections as fundamental fields was presented by Eddington [11], for a spacetime with positive cosmological constant. He proposed the square root of the determinant of the Ricci tensor as the gravitational Lagrangian. Authors like N. Popławski [12] and K. Krasnov [13] have advanced the road towards a pure connection gravity theory. In this paper we study a power-counting renormalizable, diffeomorphism invariant model consisting solely of an affine connection with

* askirz@gmail.com

† o.castillo.felisola@gmail.com

torsion. We expect this model may overcome the uniqueness theorem about diffeomorphism invariant theories of connections, since we have no fundamental metric field to quantize. Additionally, all terms in the action will involve the torsion field, therefore three- and four-point graviton amplitudes will all include torsion too. This might be considered relevant to avoid the results pointing that there are no renormalizable three-point vertices for graviton interactions in a model independent way, since in a Lorentz invariant set up kinematics alone determines this result, as reviewed in Refs. [14, 15].

The article is organized as follows: In Sec. II we analyse the three-dimensional theory built with the affine connection and power-counting renormalizable. In Sec. III we study the four-dimensional model, built under the same precepts than before. Additionally, we find solutions to the equations of motion assuming a static, homogeneous and isotropic background, and show that in the non-relativistic limit of the theory the gravitational potential is Newtonian. Finally, in Sec. IV we briefly discuss the reaching consequences of the model.

With these ingredients, in a three-dimensional space we write an action

$$\begin{aligned}
S[\hat{\Gamma}] = \int d^3x \left\{ \hat{R}_{\mu_1\mu_2}{}^\rho{}_{\mu_3} T^\sigma{}_{\mu_4\mu_5} \sum_{\pi \in Z_5} C_\pi \delta_\rho^{\mu_\pi(1)} \delta_\sigma^{\mu_\pi(2)} \epsilon^{\mu_\pi(3)\mu_\pi(4)\mu_\pi(5)} \right. \\
+ T^\rho{}_{\mu_1\mu_2} T^\sigma{}_{\mu_3\mu_4} T^\tau{}_{\mu_5\mu_6} \sum_{\pi \in Z_6} D_\pi \delta_\rho^{\mu_\pi(1)} \delta_\sigma^{\mu_\pi(2)} \delta_\tau^{\mu_\pi(3)} \epsilon^{\mu_\pi(4)\mu_\pi(5)\mu_\pi(6)} \\
\left. + T^\rho{}_{\mu_1\mu_2} \hat{\nabla}_{\mu_3} T^\sigma{}_{\mu_4\mu_5} \sum_{\pi \in Z_5} E_\pi \delta_\rho^{\mu_\pi(1)} \delta_\sigma^{\mu_\pi(2)} \epsilon^{\mu_\pi(3)\mu_\pi(4)\mu_\pi(5)} \right\}, \quad (2)
\end{aligned}$$

where all possible permutations of n elements $\pi \in Z_n$ have been included in the sums with different constants C_π , D_π and E_π for permutation.

The torsion field can be decomposed into invariant tensors respecting the symmetry,

$$T^\sigma{}_{\mu\nu} = \epsilon_{\mu\nu\rho} T^{\sigma\rho} + A_{[\mu} \delta^\sigma{}_{\nu]}, \quad (3)$$

with a symmetric $T^{\sigma\rho}$ of density weight $w = 1$, and $A_\mu = T^\nu{}_{\mu\nu}$ is the trace part of the more arbitrary $T^\sigma{}_{\mu\nu}$. In the action in Eq. (2) an extra Chern–Simons term, which in three dimensions is invariant under diffeomorphisms, with an independent coefficient.

The affine connection can be decomposed into its symmetric and antisymmetric parts,

$$\hat{\Gamma}^\lambda{}_{\mu\rho} = \Gamma^\lambda{}_{(\mu\rho)} + \epsilon_{\mu\rho\sigma} T^{\lambda\sigma} + A_{[\mu} \delta^\lambda{}_{\rho]}, \quad (4)$$

where $\epsilon_{\mu\rho\sigma}$ has been introduced, and it is related to the skew-symmetric $\epsilon^{\mu\rho\sigma}$ through the identity $\epsilon^{\lambda\mu\nu} \epsilon_{\rho\sigma\tau} = 3! \delta^\lambda{}_{[\rho} \delta^\mu{}_{\sigma} \delta^\nu{}_{\tau]}$. Therefore, the curvature ten-

II. WARMING UP: THE THREE-DIMENSIONAL CASE

Formally, the curvature of a manifold is defined through the commutator of covariant derivatives under diffeomorphisms, $\hat{\nabla}_\mu$, but for general choice of the connection, $\hat{\Gamma}^\mu{}_{\nu\lambda}$, there is an extra contribution given by its antisymmetric part in the lower indices, $T^\mu{}_{\nu\lambda} = 2\hat{\Gamma}^\mu{}_{[\nu\lambda]}$. Therefore, the commutator of the covariant derivatives yields,

$$[\hat{\nabla}_\mu, \hat{\nabla}_\nu] V^\rho = \hat{R}_{\mu\nu}{}^\rho{}_\lambda V^\lambda - T^\rho{}_{\mu\nu} \hat{\nabla}_\rho V^\rho. \quad (1)$$

Note that $T^\rho{}_{\mu\nu}$ is a nine-dimensional tensor representation under diffeomorphisms.

In order to build topological invariants of density one, we can use the skew-symmetric Levi-Civita tensor $\epsilon^{\mu_1\mu_2\cdots\mu_n}$ in n -dimensional space(-time).

sor can be expressed as

$$\begin{aligned}
\hat{R}_{\mu\nu}{}^\sigma{}_\rho = R_{\mu\nu}{}^\sigma{}_\rho - 2\epsilon_{\rho\alpha[\mu} \nabla_{\nu]} T^{\sigma\alpha} + \partial_{[\mu} A_{\nu]} \delta^\sigma{}_\rho \\
+ \delta^\sigma{}_{[\mu} \nabla_{\nu]} A_\rho + \epsilon_{\mu\nu\kappa} T^{\kappa\sigma} A_\rho - \delta^\sigma{}_{[\mu} \epsilon_{\nu]\rho\alpha} T^{\alpha\beta} A_\beta \\
+ \frac{1}{2} \delta^\sigma{}_{[\mu} A_{\nu]} A_\rho - 2\epsilon_{\alpha\beta[\mu} \epsilon_{\nu]\rho\delta} T^{\sigma\alpha} T^{\beta\delta}, \quad (5)
\end{aligned}$$

where ∇_ρ and $R_{\mu\nu}{}^\lambda{}_\rho$ are the covariant derivative and curvature associated to the symmetric part of the connection. Notice that Bianchi identity for the Riemann curvature, obtained as $\epsilon^{\mu\nu\lambda} R_{\mu\nu}{}^\rho{}_\lambda = 0$, leads us to

$$\epsilon^{\mu\nu\rho} \hat{R}_{\mu\nu}{}^\lambda{}_\rho = 4\nabla_\rho T^{\rho\lambda} + 2\epsilon^{\mu\nu\lambda} \partial_\mu A_\nu - 4T^{\lambda\rho} A_\rho. \quad (6)$$

Using the Eqs. (5) and (6) one can rewrite the action (up

to a boundary term) as

$$\begin{aligned}
S[\Gamma, T, A] = & \int d^3x \left(A_1 R_{\mu\nu}{}^\mu{}_\rho T^{\nu\rho} + A_2 \epsilon^{\mu\nu\rho} R_{\mu\nu}{}^\sigma{}_\sigma A_\rho \right. \\
& + A_3 \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho + A_4 T^{\mu\nu} \nabla_\mu A_\nu \\
& + A_5 T^{\mu\nu} A_\mu A_\nu + A_6 \det(T^{\mu\nu}) \\
& + A_7 \epsilon^{\mu\nu\lambda} \left(\Gamma^\sigma{}_{\mu\rho} \partial_\nu \Gamma^\rho{}_{\lambda\sigma} + \frac{2}{3} \Gamma^\tau{}_{\mu\rho} \Gamma^\rho{}_{\nu\sigma} \Gamma^\sigma{}_{\lambda\tau} \right) \\
& \left. + A_8 \epsilon^{\mu\nu\rho} \Gamma^\sigma{}_{\mu\sigma} \partial_\nu \Gamma^\tau{}_{\rho\tau} \right). \tag{7}
\end{aligned}$$

At this point, it is useful to introduce what we have called the ‘‘Eddington’s trick’’ [11]. First of all, notice that in the usual Einstein–Hilbert action the variation of the action with respect to the Ricci tensor yields an inverse metric density. Thus, a sort of dual theory could be obtained by identifying the tensor density obtained from the variation of the action with respect to the symmetric part of Ricci tensor with the inverse metric density (see Ref. [11, 12])

$$\frac{\delta}{\delta R_{(\mu\nu)}} S[\hat{\Gamma}] \implies \sqrt{\mathfrak{g}} \mathfrak{g}^{\mu\nu} \equiv \bar{g}^{\mu\nu}. \tag{8}$$

Noticing that in the first term, the variation respect to the Ricci tensor yields to $T^{\mu\nu}$, it can be argued that in a standard theory of gravity this tensor density corresponds to $\bar{g}^{\mu\nu}$. Therefore, Eq. (7) reveals a one-to-one correspondence with general relativity nonminimally coupled to the A_μ field,

$$\begin{aligned}
S[\mathfrak{g}, \Gamma, A] = & \int d^3x \left(\sqrt{\mathfrak{g}} \left(A_1 R + A_4 \nabla^\mu A_\mu \right. \right. \\
& + A_5 A_\mu A^\mu + A_6 \left. \right) + A_2 \epsilon^{\mu\nu\rho} R_{\mu\nu}{}^\sigma{}_\sigma A_\rho \\
& + A_3 \epsilon^{\mu\nu\rho} A_\mu \partial_\nu A_\rho + A_7 \epsilon^{\mu\nu\lambda} \left(\Gamma^\sigma{}_{\mu\rho} \partial_\nu \Gamma^\rho{}_{\lambda\sigma} \right. \\
& \left. + \frac{2}{3} \Gamma^\tau{}_{\mu\rho} \Gamma^\rho{}_{\nu\sigma} \Gamma^\sigma{}_{\lambda\tau} \right) + A_8 \epsilon^{\mu\nu\rho} \Gamma^\sigma{}_{\mu\sigma} \partial_\nu \Gamma^\tau{}_{\rho\tau} \left. \right). \tag{9}
\end{aligned}$$

Thus, an interesting sector of the theory corresponds to the space of non-degenerated $T^{\mu\nu}$.

For the degrees of freedom count we have used Ref. [16] where a lagrangian approach to the counting has been developed. We part from the $N = 27$ dimensional affine connection. For the action (7), we compute the equations of motion and obtain only first order equations that figure as 27 lagrangian constraints. Only 3 of these equations admit an additional time derivative without becoming of second order in time, i.e.,

$$\frac{\delta S}{\delta \Gamma^i{}_{00}} \text{ and } A_4 \frac{\delta S}{\delta \Gamma^0{}_{00}} - 2A_1 \frac{\delta S}{\delta A_0}.$$

Thus, the total number of Lagrangian constraints is $l = 30$. As for gauge transformations, we know this model to be invariant under diffeomorphisms, described through three local parameters which appear with first order time derivatives in the change of $T^{\mu\nu}$ and A_0 and with second order time derivatives in the transformation rule of $\Gamma^{\mu}{}_{00}$, hence we have $e = 9$ effective gauge parameters (because each time derivative count as independent) and $g = 3$ gauge identities. Finally, we conclude that $N - \frac{1}{2}(l + g + e) = 6$ is the total number of degrees of freedom in this model.

III. FOUR-DIMENSIONAL METRICLESS (AND TORSIONFUL) ACTION

Following the precepts already stated, we start by defining an irreducible representation decomposition for the full connection field

$$\hat{\Gamma}^\mu{}_{\rho\sigma} = \Gamma^\mu{}_{\rho\sigma} + T^\mu{}_{\rho\sigma} = \Gamma^\mu{}_{\rho\sigma} + \epsilon_{\rho\sigma\lambda\kappa} T^{\mu,\lambda\kappa} + A_{[\rho} \delta^\mu_{\nu]}, \tag{10}$$

where $\Gamma^\mu{}_{\rho\sigma}$ denotes a forty-dimensional symmetric connection, A_μ is a four-dimensional vector field that gives trace to the antisymmetric part of the full connection, and $T^{\mu,\lambda\kappa}$ is a twenty-dimensional Curtright field (see Ref. [17]) that is defined through the symmetry of its indices: antisymmetric in the last two indices, and it has a cyclic property $T^{\mu,\lambda\kappa} + T^{\lambda,\kappa\mu} + T^{\kappa,\mu\lambda} = 0$. In other words that $T^{[\mu,\lambda]\kappa} = \frac{1}{2} T^{\kappa,\lambda\mu}$, just as for the Riemann tensor $R_{\mu[\nu}{}^\alpha{}_{\lambda]} = \frac{1}{2} R_{\lambda\nu}{}^\alpha{}_\mu$. Notice that due to its symmetries, the contraction $\epsilon_{\rho\sigma\lambda\kappa} T^{\mu,\lambda\kappa}$ is traceless.

Additionally, since no metric is present the epsilon symbols are not related by lowering or raising their indices, but instead one demands that $\epsilon^{\delta\eta\lambda\kappa} \epsilon_{\mu\nu\rho\sigma} = 4! \delta^\delta{}_{[\mu} \delta^\eta{}_\nu \delta^\lambda{}_\rho \delta^\kappa{}_{\sigma]}$.

One can write all the combinations of fields that would presumably be renormalizable with these three independent

fields — up to a boundary term —,

$$\begin{aligned}
S[\Gamma, T, A] = \int d^4x \left[B_1 R_{\mu\nu}{}^\mu{}_\rho T^{\nu,\alpha\beta} T^{\rho,\gamma\delta} \epsilon_{\alpha\beta\gamma\delta} + B_2 \left(R_{\mu\nu}{}^\sigma{}_\rho + \frac{2}{3} \delta^\sigma{}_{[\mu} R_{\nu]\lambda}{}^\lambda{}_\rho \right) T^{\beta,\mu\nu} T^{\rho,\gamma\delta} \epsilon_{\sigma\beta\gamma\delta} \right. \\
+ B_3 R_{\mu\nu}{}^\mu{}_\rho T^{(\nu,\rho)\sigma} A_\sigma + B_4 \left(R_{\mu\nu}{}^\sigma{}_\rho + \frac{2}{3} \delta^\sigma{}_{[\mu} R_{\nu]\lambda}{}^\lambda{}_\rho \right) \left(T^{\rho,\mu\nu} A_\sigma - \frac{1}{4} \delta_\sigma^\rho T^{\kappa,\mu\nu} A_\kappa \right) + B_5 R_{\mu\nu}{}^\rho{}_\sigma T^{\sigma,\mu\nu} A_\sigma \\
+ C_1 R_{\mu\nu}{}^\mu{}_\rho \nabla_\sigma T^{(\nu,\rho)\sigma} + C_2 R_{\mu\nu}{}^\rho{}_\sigma \nabla_\sigma T^{\sigma,\mu\nu} + D_1 T^{\alpha,\mu\nu} T^{\beta,\rho\sigma} \nabla_\gamma T^{(\lambda,\kappa)\gamma} \epsilon_{\beta\mu\nu\lambda} \epsilon_{\alpha\rho\sigma\kappa} \\
+ D_2 T^{\alpha,\mu\nu} T^{\lambda,\beta\gamma} \nabla_\lambda T^{\delta,\rho\sigma} \epsilon_{\alpha\beta\gamma\delta} \epsilon_{\mu\nu\rho\sigma} + D_3 T^{\mu,\alpha\beta} T^{\lambda,\nu\gamma} \nabla_\lambda T^{\delta,\rho\sigma} \epsilon_{\alpha\beta\gamma\delta} \epsilon_{\mu\nu\rho\sigma} \\
+ D_4 T^{\lambda,\mu\nu} T^{\kappa,\rho\sigma} \nabla_{(\lambda} A_{\kappa)} \epsilon_{\mu\nu\rho\sigma} + D_5 T^{\lambda,\mu\nu} \nabla_{[\lambda} T^{\kappa,\rho\sigma} A_{\kappa]} \epsilon_{\mu\nu\rho\sigma} + D_6 T^{\lambda,\mu\nu} A_\nu \nabla_{(\lambda} A_{\mu)} + D_7 T^{\lambda,\mu\nu} A_\lambda \nabla_{[\mu} A_{\nu]} \\
+ E_1 \nabla_{(\rho} T^{\rho,\mu\nu} \nabla_{\sigma)} T^{\sigma,\lambda\kappa} \epsilon_{\mu\nu\lambda\kappa} + E_2 \nabla_{(\lambda} T^{\lambda,\mu\nu} \nabla_{\mu)} A_\nu + T^{\alpha,\beta\gamma} T^{\delta,\eta\kappa} T^{\lambda,\mu\nu} T^{\rho,\sigma\tau} (\Lambda_1 \epsilon_{\beta\gamma\eta\kappa} \epsilon_{\alpha\rho\mu\nu} \epsilon_{\delta\lambda\sigma\tau} \\
+ \Lambda_2 \epsilon_{\beta\lambda\eta\kappa} \epsilon_{\gamma\rho\mu\nu} \epsilon_{\alpha\delta\sigma\tau}) + \Lambda_3 T^{\rho,\alpha\beta} T^{\gamma,\mu\nu} T^{\lambda,\sigma\tau} A_\tau \epsilon_{\alpha\beta\gamma\lambda} \epsilon_{\mu\nu\rho\sigma} + \Lambda_4 T^{\eta,\alpha\beta} T^{\kappa,\gamma\delta} A_\eta A_\kappa \epsilon_{\alpha\beta\gamma\delta} \left. \right], \tag{11}
\end{aligned}$$

where the terms B_2 and B_4 contain a traceless contribution of the curvature.

In this case, the Eddington's "inverse metric density" [see Eq. (8)] is

$$\bar{g}^{\mu\nu} = B_1 T^{\mu,\lambda\kappa} T^{\nu,\rho\sigma} \epsilon_{\lambda\kappa\rho\sigma} + B_3 T^{(\mu,\nu)\lambda} A_\lambda + C_1 \nabla_\lambda T^{(\mu,\nu)\lambda}. \tag{12}$$

Symmetric solution to the equations of motion

In four dimensions there is no obvious equivalence of Eq. (11) with GR, specially due to the lack of a fundamental metric field in the given model. However, both models are explicitly invariant under diffeomorphisms, and even if their structures and number of degrees of freedom differ, the action in Eq. (11) provides a context where parallel transport of particle's velocities on a purely torsional background is nontrivial.

Here, we wish to establish the model's non-relativistic (Newtonian) limit for the "geodesic" deviation of "inertial" observers at rest with respect to a static, isotropic, homogeneous and spatially flat background within the context provided by Eq. (11). In order to properly analyse the model, we propose the following expansion of the fields

$$A_\mu = \delta_\mu^0 A + a_\mu, \tag{13}$$

$$T^{\mu,\nu\rho} = \delta_m^\mu \delta_{m0}^{\nu\rho} T + t^{\mu,\nu\rho}, \tag{14}$$

and

$$\Gamma^\lambda{}_{\mu\nu} = E \delta_0^\lambda \delta_\mu^m \delta_\nu^m + F \delta_m^\lambda \delta_{(\mu}^m \delta_{\nu)}^0 + G \delta_0^\lambda \delta_\mu^0 \delta_\nu^0 + \gamma^\lambda{}_{\mu\nu}, \tag{15}$$

where $\delta_{\lambda\kappa}^{\mu\nu} = \delta_\lambda^\mu \delta_\kappa^\nu - \delta_\kappa^\mu \delta_\lambda^\nu$.

The Eddington's metric density in Eq. (12) on the background is

$$\bar{g}^{\mu\nu} = \left(B_3 A + \frac{1}{2} C_1 F \right) T \delta_m^\mu \delta_m^\nu - 3 C_1 E T \delta_0^\mu \delta_0^\nu, \tag{16}$$

while the Ricci curvature tensor calculated from Eq. (15) is

$$R_{\mu\nu} = \frac{1}{2} E F \delta_\mu^m \delta_\nu^m - \frac{3}{4} F^2 \delta_\mu^0 \delta_\nu^0. \tag{17}$$

Therefore, whether the four-dimensional metric structure is Riemannian or pseudo-Riemannian will depend exclusively on the values of the parameters of the action in Eq. (11) and the signs of the components of the connection field.

The first order perturbations of the action yields

$$\begin{aligned}
\delta S = & \left((B_3 + \frac{8}{3} B_4 + \frac{1}{2} E_2) A + 4 C_1 F - 2 C_1 G \right) E + 8 (-D_1 + 2 D_2 + D_3) T^2 \Big) T \delta \Gamma^m{}_{0m} \\
& + \left((\frac{1}{2} B_3 + \frac{4}{3} B_4 + \frac{1}{4} E_2) A F + (B_3 - \frac{4}{3} B_4 - \frac{1}{2} E_2) A G + C_1 F^2 - C_1 F G - D_6 A^2 \right) T \delta \Gamma^{0m}{}_m \\
& + \left((-\frac{1}{2} B_3 + \frac{4}{3} B_4 + \frac{1}{4} E_2) A F + (-B_3 + \frac{4}{3} B_4 + \frac{1}{2} E_2) A G - C_1 F^2 + C_1 F G + D_6 A^2 \right) E \\
& + \left(12 (D_1 - 2 D_2 - D_3) F + 24 \Lambda_3 A \right) T^2 \Big) \delta T_m{}^{0m} + \left((3 B_3 - 4 B_4 - \frac{3}{2} E_2) A - 3 C_1 F \right) E T \delta \Gamma^0{}_{00} \\
& + \left(3 \left(-2 D_6 A + (\frac{1}{2} B_3 + \frac{4}{3} B_4 + \frac{1}{4} E_2) F + (B_3 - \frac{4}{3} B_4 - \frac{1}{2} E_2) G \right) E - 24 \Lambda_3 T^2 \right) T \delta A_0 = 0,
\end{aligned} \tag{18}$$

and we are most interested in solutions to the connection field whose contribution to the parallel transport equation of a test particle's velocity is that of a free particle, at least at the low velocity regime

$$\ddot{x}^i + 2F \dot{x}^0 \dot{x}^i = 0, \quad \text{and} \quad \ddot{x}^0 + E (\dot{x}^i)^2 + G (\dot{x}^0)^2 = 0, \tag{19}$$

which we can achieve by setting $F = G = 0$ and $E \neq 0$ since $(\dot{x}^i)^2$ is already second order in the velocities. Thus, looking again at the equations of motion we can find a nontrivial solution if we set all coupling constants to zero but $B_3 \neq 0$, $B_4 = -\frac{3}{2} B_3$, $C_1 \neq 0$ and $E_2 = 6B_3$.

Additionally, we can incorporate perturbative inhomogeneous sources to the connection field equations and check on how these fluctuations affect motion. For this, we consider a matter's action, whose dependence on the affine connection can be almost arbitrary. However, we will presume that it will depend only on the barred metric in Eq. (12)

$$S_{\text{Matter}} = S_{\text{Matter}}[\bar{g}^{\mu\nu}].$$

Thus, a non-moving matter point particle at the origin of the reference frame will contribute to the equations of motion for the gravitational field through the component \bar{g}^{00} following the symmetries of the matter source

$$\begin{aligned}
\delta S_{\text{Matter}} = & C_1 \left(-\frac{1}{2} (\delta \Gamma^0{}_{mn}) T \delta^{mn} - \frac{1}{2} (\delta T^{00m}) i p_m \right. \\
& \left. + \frac{1}{2} (\delta T^{m0n}) E \delta_{mn} \right) \frac{\partial \mathcal{L}_{\text{Matter}}}{\partial \bar{g}^{00}}.
\end{aligned} \tag{20}$$

Scalar modes and Newtonian limit

In order to obtain the non-relativistic limit, *i.e.*, the Newtonian potential, one performs the scalar mode per-

turbative expansion. One proceeds by substituting the connection and torsion components by their scalar perturbation decomposition,

$$a_\mu \rightarrow \delta_\mu^0 a + \delta_\mu^m \partial_m a_L, \tag{21}$$

$$\begin{aligned}
t^{\mu,\nu\rho} \rightarrow & \delta_m^\mu \delta_{n0}^{\nu\rho} \left(t \delta^{mn} + \partial^m \partial^n t_L \right) + \delta_0^\mu \delta_{m0}^{\nu\rho} \partial^m c_L \\
& + \left(\delta_0^\mu \delta_{mn}^{\nu\rho} - \delta_m^\mu \delta_{n0}^{\nu\rho} \right) \epsilon^{mnp} \partial_p b \\
& + \delta_m^\mu \delta_n^\nu \delta_p^\rho \left(\epsilon^{npq} \partial_q \partial^m d_1 + (\delta^{mn} \partial^p - \delta^{mp} \partial^n) d_2 \right)
\end{aligned} \tag{22}$$

and

$$\begin{aligned}
\gamma_{\mu\nu}^\lambda \rightarrow & \delta_0^\lambda \delta_\mu^0 \delta_\nu^0 u + \delta_m^\lambda \delta_\mu^0 \delta_\nu^0 \partial^m v_L + 2 \delta_0^\lambda \delta_{(\mu}^0 \delta_{\nu)}^m \partial_m w_L \\
& + \delta_0^\lambda \delta_\mu^m \delta_\nu^n \left(x \delta_{mn} + \partial_m \partial_n x_L \right) \\
& + 2 \delta_m^\lambda \delta_{(\mu}^0 \delta_{\nu)}^n \left(y_1 \delta^m{}_n + \epsilon^{mp}{}_n \partial_p y_2 + \partial^m \partial_n y_L \right) \\
& + \delta_m^\lambda \delta_\mu^n \delta_\nu^p \left(\delta_{np} \partial^m z_1 + (\delta^m{}_n \partial_p + \delta^m{}_p \partial_n) z_2 \right. \\
& \left. + (\epsilon^{mq}{}_n \partial_p + \epsilon^{mq}{}_p \partial_n) \partial_q z_3 + \partial^m \partial_n \partial_p z_L \right),
\end{aligned} \tag{23}$$

where the scalar fields identified with the sub-index ‘‘L’’ correspond to longitudinal degrees of freedom. Vector and tensor perturbations are left for further investigations of the dynamical structure of the model.

The first order perturbative expansion of the equations of motion around the already described background in momentum space with $p_0 = 0$ is given by

$$\begin{aligned}
\delta S = & \left(-2Ed_2 + t - p^2 t_L + Tw_L + 3Tz_1 + 2Tz_2 - Tp^2 z_L \right) 6B_3 p^2 \delta A_0 + \left(-Ep^2 d_2 + \frac{1}{2}p^2 t - \frac{1}{2}p^4 t_L - \frac{1}{2}Tp^2 w_L \right. \\
& - 6ETy_1 + 2ETp^2 y_L + \frac{3}{2}Tp^2 z_1 + Tp^2 z_2 - \frac{1}{2}Tp^2 p^2 z_L \left. \right) C_1 \delta \Gamma^0_{00} + \left(6B_3 a - \frac{1}{2}C_1 u + 2C_1 E v_L + \frac{1}{2}C_1 y_1 \right. \\
& - \frac{3}{2}C_1 p^2 y_L \left. \right) T v^m \delta \Gamma^0_{0m} + C_1 T p^2 v_L \delta^{mn} \delta \Gamma^0_{mn} + \left(-p^2 c_L + 2E^2 d_2 + 4Et - 2Ep^2 t_L + 2ETw_L \right. \\
& + 3Tx - Tp^2 x_L - 10ETz_2 + 2ETp^2 z_L \left. \right) C_1 v p_m \delta \Gamma^m_{00} + \left(Ep^2 d_2 + \frac{1}{2}p^2 t - \frac{1}{2}p^4 t_L - 2ETu - \frac{1}{2}Tp^2 w_L \right. \\
& + 8ETy_1 - 2ETp^2 y_L + \frac{1}{2}Tp^2 z_1 + Tp^2 z_2 - \frac{1}{2}Tp^4 z_L \left. \right) C_1 \delta^m_n \delta \Gamma^n_{0m} + \left(-4Ed_2 - t + p^2 t_L + 2Tw_L \right. \\
& - 2ETy_L - 2Tz_1 + Tp^2 z_L \left. \right) C_1 p_n p^m \delta \Gamma^n_{0m} - 6ETy_2 C_1 v^p \epsilon_{np}{}^m \delta \Gamma^n_{0m} + \left(6B_3 Ta + C_1 p^2 d_2 + \frac{1}{2}C_1 Tu \right. \\
& + C_1 ETv_L - \frac{1}{2}C_1 Ty_1 + \frac{1}{2}C_1 Tp^2 y_L \left. \right) v p_m \delta^{np} \delta \Gamma^m_{np} + \left(-3Ev_L + y_1 - p^2 y_L \right) C_1 T v^p \delta^m_p \delta \Gamma^m_{np} + \left(-d_2 \right. \\
& + Ty_L \left. \right) C_1 v p_m p^n p^p \delta \Gamma^m_{np} + v_L C_1 p^2 v p_m \delta T^{00m} - v_L C_1 E p^2 \delta_{mn} \delta T^{m0n} + \left(-6B_3 a - \frac{1}{2}C_1 u - C_1 E v_L - \frac{1}{2}C_1 y_1 \right. \\
& - \frac{1}{2}C_1 p^2 y_L \left. \right) p_m p_n \delta T^{m0n} + \left(6B_3 Ea + \frac{1}{2}C_1 E u - C_1 E^2 v_L + \frac{1}{2}C_1 E y_1 - \frac{3}{2}C_1 E p^2 y_L - C_1 p^2 z_1 \right) v p_m \delta_{np} \delta T^{mnp}.
\end{aligned} \tag{24}$$

which we will add to the variations of the action of the matter from Eq. (20) and set $\delta S_{\text{total}} = 0$.

Solutions to this set of equations are in general a highly difficult problem that concerns twenty equations of motion with twenty scalar fields to be fixed. Yet, knowledge of the value of some of these scalars does not necessarily help to determine how geodesics are affected. For this, we only need $\gamma^i{}_{00}$ and $\gamma^0{}_{00}$ as these provide the first order contributions to the equations

$$\ddot{x}^i + \gamma^i{}_{00}(\dot{x}^0)^2 = 0, \quad \text{and} \quad \ddot{x}^0 + \gamma^0{}_{00}(\dot{x}^0)^2 = 0. \tag{25}$$

From Eq. (25) we can conclude that we need only know $\gamma^i{}_{00} = \partial^i v_L$, which we obtain in Fourier space to be

$$v_L = \frac{1}{2} \frac{\partial \mathcal{L}_{\text{Matter}}}{\partial \bar{g}^{00}} \frac{1}{p^2}. \tag{26}$$

In position space,

$$v_L = \frac{1}{8\pi} \frac{\partial \mathcal{L}_{\text{Matter}}}{\partial \bar{g}^{00}} \frac{1}{|\vec{x}|} \tag{27}$$

is the usual Newtonian potential for a massive far off source.

IV. DISCUSSION

In this paper we have proposed novel model of gravitational interactions with full diffeomorphisms invariance as the main guiding principle, whose fundamental field is an affine connection and no metric field is assumed (nor needed). Surprisingly, in four dimensions, it upholds the

correct Newtonian limit, supporting the suspicions that it may describe some aspects of gravitational physics that have not been exposed yet. Still, the model is alien for anyone accustomed to metric spacetimes or their extensions, specially since no local Lorentz structure is present. We also argue that in the absence of fundamental inertial structure, it becomes a natural playground to test the full reaches of Mach's Principle.

Since GR is the best gravitational theory we have so far, we are currently working in obtaining the relativistic limit of our model, in the limit of vanishing torsion [18]. Interestingly enough, in this limit we get a generalization of Einstein's equations which support the principal result of this paper, i.e., our model has a well-defined Newtonian limit — even in the non-vanishing torsion case —. Also, we couple scalar matter to the gravity sector which, at the level of equation of motions, is equivalent to the usual Einstein–scalar-field theory with minimal coupling [18].

Additionally, within the model, all coupling constants turn out to be dimensionless, a property that has been related to scale invariance and conformally invariant theories (see Refs. [19, 20]). In fact, renormalization is intimately related to the scaling properties of a model and it may be worth to study the quantization and renormalizability of this model. In doing so, we believe the lack of metric may allow the model to bypass the uniqueness of the diffeomorphisms invariant Hilbert space stated in Ref. [10].

Sticking to the classical theory, the analysis of the cosmological implications is needed. In cosmology there are important aspects with unsatisfactory explanations, such as those related to the matter content of the Universe

(in particular the dark energy sector), or the large scale structure formation [21]. Additionally, other formal aspects of the model remain unknown, such as the proper number of propagating degrees of freedom, and whether or not a duality exists between our model and one of the well-known metric models (say for example Hořava-Lifshitz gravity [22]).

We also believe that almost every aspect related to coupling gravity to matter fields can be extrapolated to couplings with the affine connection without making reference to a fundamental metric or a local Lorentz symmetry. This could be done rewriting the affine connection in terms of a local $GL(4)$ connection $\omega_\mu{}^a{}_b$, relating the two of them through use of a frame field e_μ^a and its inverse e_a^μ

$$\omega_\mu{}^a{}_b = e_\lambda^a e_b^\nu \Gamma^\lambda{}_{\mu\nu} - e_b^\nu \partial_\mu e_\nu^a, \quad (28)$$

where a and b are indices in the defining representation of the local group $GL(4)$. In particular, $GL(4) = \mathbb{R}_+ \times SL(4)$ and using $SL(4) \simeq SO(3,3)$, we can define spinorial representations for the diffeomorphisms group, eventually we will be able to define an action for $SO(3,3)$ spinors in four dimensions (check Appendix A).

ACKNOWLEDGMENTS

We thank to J. Zanelli for fruitful discussions, and also to K. Peeters for helpful advises in the manipulation of the software CADABRA [23], which was used extensively to achieve the results presented in this paper. Additionally, we thank the developers of the mathematical software SAGE [24], used to achieve several manipulations. This work was partially supported by CONICYT (Chile) under grant No. 79140040.

Appendix A: Matter Fields

The Dirac equation relies on the local $SO(3,1)$ Lorentz symmetry for everything. The aim of this section is to describe the inclusion of Dirac spinor without a local Lorentz symmetry in four dimensions.

Dirac spinors are the fields that transform under representation of the local symmetry group that corresponds to the double cover of the original symmetry. We are interested in the representations of the diffeomorphisms

group, that could be associated to the local symmetry generated by the semi-simple Lie group $SL(4, \mathbb{R})$ in four dimensions. This notion is unusual because typically one would think of $SO(3,1)$ as the local symmetry, and its double cover would generally be called the spin group $Spin(3,1)$. Instead, we realize that the local symmetry we have got, $SL(4, \mathbb{R})$ is equivalent to $SO(3,3)$ and the fundamental six dimensional representation of this group corresponds to the space of 4×4 antisymmetric matrix representation of $SL(4, \mathbb{R})$. In terms of their invariant tensors, it is easy to see their correspondence. Consider a vector with components $v^A = \frac{1}{\sqrt{2}}(F^{01} - F^{23}, F^{02} - F^{31}, F^{03} - F^{12}, F^{01} + F^{23}, F^{02} + F^{31}, F^{03} + F^{12})$, where F^{ab} is an antisymmetric tensor representation of $SL(4)$. A diagonal metric $\eta_{AB} = \text{diag}(-, -, -, +, +, +)$ allows us to compute inner products, and $\vec{v} \cdot \vec{v}$ is

$$\begin{aligned} \vec{v} \cdot \vec{v} &= 2 \left(F^{01} F^{23} + F^{02} F^{31} + F^{03} F^{12} \right) \\ &= \frac{1}{4} \epsilon_{abcd} F^{ab} F^{cd}. \end{aligned} \quad (A1)$$

Thus, in order to consider $SL(4, \mathbb{R})$ double cover we will use the Clifford algebra $\mathcal{Cl}_{3,3}$ defined by

$$[\Gamma_A, \Gamma_B] = 2\eta_{AB} \quad (A2)$$

and redefine

$$\begin{aligned} \Gamma_A \rightarrow \frac{1}{\sqrt{2}} (\gamma_{01} - \gamma_{23}, \gamma_{02} - \gamma_{31}, \gamma_{03} - \gamma_{12}, \gamma_{01} \\ + \gamma_{23}, \gamma_{02} + \gamma_{31}, \gamma_{03} + \gamma_{12}) \end{aligned} \quad (A3)$$

to rewrite the Clifford algebra in Eq. (A2) as

$$\{\gamma_{ab}, \gamma_{cd}\} = 2\epsilon_{abcd}, \quad (A4)$$

where γ_{ab} are antisymmetric in $a \leftrightarrow b$, 8×8 complex matrices and η_{AB} is basically ϵ_{abcd} in a different basis.

Thus, several topological scalars can be added to the Lagrangian density

$$\begin{aligned} \mathcal{L}_\Psi &= g_1 \bar{\Psi} \gamma_{ab} e_\mu^a e_\nu^b T^{\lambda, \mu\nu} \nabla_\lambda \Psi + g_2 \bar{\Psi} \gamma_{ab} e_\mu^a e_\nu^b \epsilon^{\lambda\kappa\mu\nu} A_\kappa \nabla_\lambda \Psi \\ &+ g_3 \bar{\Psi} \gamma_{ab} e_\mu^a e_\nu^b \nabla_\lambda T^{\lambda, \mu\nu} \Psi + g_4 \bar{\Psi} \gamma_{ab} e_\mu^a e_\nu^b \epsilon^{\lambda\kappa\mu\nu} \nabla_\lambda A_\kappa \Psi. \end{aligned} \quad (A5)$$

-
- [1] H. Lichtenegger and B. Mashhoon, (2004), [arXiv:physics/0407078](https://arxiv.org/abs/physics/0407078) [physics.hist-ph].
- [2] D. W. Sciama, *Rev. Mod. Phys.* **36**, 463 (1964).
- [3] E. Berti, E. Barausse, V. Cardoso, L. Gualtieri, P. Pani, *et al.*, (2015), [arXiv:1501.07274](https://arxiv.org/abs/1501.07274) [gr-qc].
- [4] J. W. Maluf, *Annalen Phys.* **525**, 339 (2013), [arXiv:1303.3897](https://arxiv.org/abs/1303.3897) [gr-qc].
- [5] R. Aldrovandi and J. G. Pereira, *Teleparallel Gravity*, Fundamental Theories of Physics, Vol. 173 (Springer, 2013).
- [6] J. C. Baez and D. K. Wise, (2012), [arXiv:1204.4339](https://arxiv.org/abs/1204.4339) [gr-qc].
- [7] A. Ashtekar and J. Lewandowski, *Class. Quant. Grav.* **21**, R53 (2004), [arXiv:gr-qc/0404018](https://arxiv.org/abs/gr-qc/0404018) [gr-qc].
- [8] T. Thiemann, in *Approaches to Fundamental Physics*

- (Springer, 2007) pp. 185–263.
- [9] T. Thiemann, *Class. Quant. Grav.* **15**, 839 (1998), [arXiv:gr-qc/9606089 \[gr-qc\]](#).
- [10] J. Lewandowski, A. Okolow, H. Sahlmann, and T. Thiemann, *Commun. Math. Phys.* **267**, 703 (2006), [arXiv:gr-qc/0504147 \[gr-qc\]](#).
- [11] A. S. Eddington, *The mathematical theory of relativity* (CUP, 1923).
- [12] N. J. Popławski, *Gen. Rel. Grav.* **46**, 1625 (2014), [arXiv:1203.0294 \[gr-qc\]](#).
- [13] K. Krasnov, *Phys. Rev. Lett.* **106**, 251103 (2011), [arXiv:1103.4498 \[gr-qc\]](#).
- [14] D. A. McGady and L. Rodina, *Phys. Rev. D* **90**, 084048 (2014), [arXiv:1311.2938 \[hep-th\]](#).
- [15] X. O. Camanho, J. D. Edelstein, J. Maldacena, and A. Zhiboedov, (2014), [arXiv:1407.5597 \[hep-th\]](#).
- [16] B. Diaz, D. Higuera, and M. Montesinos, *J. Math. Phys.* **55**, 122901 (2014), [arXiv:1406.1156 \[hep-th\]](#).
- [17] T. Curtright, *Phys. Lett. B* **165**, 304 (1985).
- [18] O. Castillo-Felisola and A. Skirzewski, “Einstein’s gravity from an affine model,” (in preparation).
- [19] D. Buchholz and K. Fredenhagen, *J. Math. Phys.* **18**, 1107 (1977).
- [20] J. Maldacena, (2011), [arXiv:1105.5632 \[hep-th\]](#).
- [21] There is a controversy between the experimental results obtained by BICEP2 [25] and Planck [26] in the respect of their interpretation associated with inflation.
- [22] P. Hořava, *Phys. Rev. D* **79**, 084008 (2009), [arXiv:0901.3775 \[hep-th\]](#); T. P. Sotiriou, *J. Phys. Conf. Ser.* **283**, 012034 (2011), [arXiv:1010.3218 \[hep-th\]](#).
- [23] K. Peeters, (2007), [arXiv:hep-th/0701238 \[HEP-TH\]](#); *Computeralgebra Rundbrief* **41**, 16 (2007); *Comput. Phys. Commun.* **176**, 550 (2007).
- [24] W. A. Stein *et al.*, *Sage Mathematics Software (Version 6.4.1)*, The Sage Development Team (2015).
- [25] P. A. R. Ade *et al.* (BICEP2 Collaboration), *Phys. Rev. Lett.* **112**, 241101 (2014), [arXiv:1403.3985 \[astro-ph.CO\]](#).
- [26] R. Adam *et al.* (Planck Collaboration), (2014), [arXiv:1409.5738 \[astro-ph.CO\]](#).