

# Finite $p$ -groups all of whose subgroups of index $p^3$ are abelian \*

Qinhai Zhang<sup>†</sup>, Libo Zhao, Miaomiao Li and Yiqun Shen  
 Department of Mathematics, Shanxi Normal University, Linfen, Shanxi 041004  
 People's Republic of China

September 4, 2018

## Abstract

Suppose that  $G$  is a finite  $p$ -group. If all subgroups of index  $p^t$  of  $G$  are abelian and at least one subgroup of index  $p^{t-1}$  of  $G$  is not abelian, then  $G$  is called an  $\mathcal{A}_t$ -group. In this paper, some information about  $\mathcal{A}_t$ -groups are obtained and  $\mathcal{A}_3$ -groups are completely classified. This solves an *old problem* proposed by Berkovich and Janko in their book [8]. Abundant information about  $\mathcal{A}_3$ -groups are given.

**Keywords** finite  $p$ -groups, minimal non-abelian  $p$ -groups,  $\mathcal{A}_t$ -groups.

**2010MSC** 20D15

## 1 Introduction

Finite  $p$ -groups are an important class of finite groups. After the classification of finite simple groups was finally completed, the study of finite  $p$ -groups becomes more and more active. Many leading group theorists, for example, Glauberman and Janko have turned their attentions to the study of finite  $p$ -groups. Although finite simple groups are classified, it is impossible to classify finite  $p$ -groups in the classical sense. The reason is that a finite  $p$ -group has “too many” normal subgroups and consequently there is an extremely large number of non-isomorphic  $p$ -groups of a given fixed order. In fact, Higman[15, 16] gave a formula for the number  $f(n, p)$  of non-isomorphic  $p$ -groups of order  $p^n$ :

$$\text{when } n \rightarrow \infty, f(n, p) = p^{n^3(2/27 + O(n^{-1/3}))}.$$

It is easy to see that when  $n$  becomes large, the number of non-isomorphic  $p$ -groups of order  $p^n$  becomes large in exponent speed. For example, up to now, the known results about the classification of 2-groups [11] are:

$n$	2	$2^2$	$2^3$	$2^4$	$2^5$	$2^6$	$2^7$	$2^8$	$2^9$	$2^{10}$
$f(n, 2)$	1	2	5	14	51	267	2328	56092	10494213	49487365422

\*This work was supported by NSFC (no.11371232) and by NSF of Shanxi Province (no.2012011001).

<sup>†</sup>Corresponding author. e-mail: zhangqh@dns.sxnu.edu.cn zhangqhc@hotmail.com

For  $p > 2$ ,  $p$ -groups of order  $p^7$  are classified by [21]. The result is  $f(7, 3) = 9310$ ,  $f(7, 5) = 34297$ . For  $p > 5$ ,  
 $f(7, p) = 3p^5 + 12p^4 + 44p^3 + 170p^2 + 707p + 2455 + (4p^2 + 44p + 291) \gcd(p - 1, 3) + (p^2 + 19p + 135) \gcd(p - 1, 4) + (3p + 31) \gcd(p - 1, 5) + 4 \gcd(p - 1, 7) + 5 \gcd(p - 1, 8) + \gcd(p - 1, 9)$ .

Because of the difficulty of the classification of finite  $p$ -groups in the classical sense, Janko and Berkovich sponsored and led a research project that aims at classifying certain classes of finite  $p$ -groups defined by their subgroup structure. As Janko mentioned in the Foreword of [7], to study  $p$ -groups with “large” abelian subgroups is another approach to finite  $p$ -groups. We know that non-abelian  $p$ -groups with “largest” abelian subgroups are minimal non-abelian groups. A non-abelian group  $G$  is said to be *minimal non-abelian* if every proper subgroup of  $G$  is abelian. Minimal non-abelian groups were classified in [20], and in more detail for finite  $p$ -groups in [25]. Berkovich and Janko in [6] introduced a new concept,  $\mathcal{A}_t$ -groups, which is a more general concept than that of minimal non-abelian  $p$ -groups. For a positive integer  $t$ , a finite  $p$ -group  $G$  is called an  $\mathcal{A}_t$ -group if all subgroups of index  $p^t$  of  $G$  are abelian, and at least one subgroup of index  $p^{t-1}$  of  $G$  is not abelian. In this paper, an  $\mathcal{A}_0$ -group is an abelian group. Obviously,  $\mathcal{A}_1$ -groups are exactly the minimal non-abelian  $p$ -groups. For small  $t$ ,  $\mathcal{A}_t$ -groups can be considered as groups having “large” abelian subgroups. Many scholars studied and classified  $\mathcal{A}_2$ -groups, see [6, 8, 14, 18, 26, 31]. Obviously, classifying  $\mathcal{A}_3$ -groups is a fascinating problem. In fact, this is a problem proposed by Berkovich and Janko in their joint book [8].

**Problem 1278.** (*Old problem*) *Classify  $\mathcal{A}_3$ -groups.*

In this paper, we completely classify  $\mathcal{A}_3$ -groups in classical sense. The groups described in the title are the totality of abelian groups,  $\mathcal{A}_1$ -groups,  $\mathcal{A}_2$ -groups and  $\mathcal{A}_3$ -groups. Thus the groups described in the title are completely classified.

Related to  $\mathcal{A}_3$ -groups, Berkovich and Janko proposed the following



**Problem 893([8]).** *Describe the set  $\{\alpha_1(G) \mid G \text{ is an } \mathcal{A}_3\text{-group}\}$ .*

**Problem 1595([9]).** *Classify the  $\mathcal{A}_3$ -groups  $G$  such that  $\alpha_1(G) < p^2 + p + 1$ , where  $\alpha_1(G)$  denotes the number of  $\mathcal{A}_1$ -subgroups in a  $p$ -group  $G$ .*

**Problem 2829([10]).** *Find  $\max\{\alpha_1(G)\}$ , where a  $p$ -group  $G$  runs over all  $\mathcal{A}_3$ -groups.*

As a corollary of the classification of  $\mathcal{A}_3$ -groups, we get all  $\alpha_1(G)$  for  $\mathcal{A}_3$ -groups  $G$ . Hence these problems mentioned above are also solved. Moreover, we give the triple  $(\mu_0, \mu_1, \mu_2)$ , where  $\mu_i$  denotes the number of  $\mathcal{A}_i$ -subgroups of index  $p$  in  $\mathcal{A}_3$ -groups.

We classify  $\mathcal{A}_3$ -groups  $G$  in two parts:  $G$  has an  $\mathcal{A}_1$ -subgroup of index  $p$ , and  $G$  has no  $\mathcal{A}_1$ -subgroup of index  $p$ . The sketch of the classification of  $\mathcal{A}_3$ -groups are as follows.



## 2 Preliminaries

In this paper,  $p$  is always a prime. We use  $F_p$  to denote the finite field containing  $p$  elements.  $F_p^*$  is the multiplicative group of  $F_p$ .  $(F_p^*)^2 = \{a^2 \mid a \in F_p^*\}$  is a subgroup of  $F_p^*$ .

Let  $G$  be a finite group. We use  $c(G)$ ,  $\exp(G)$  and  $d(G)$  to denote the nilpotency class, the exponent and the minimal number of generators of  $G$  respectively. We use  $C_{p^m}$ ,  $C_{p^m}^n$  and  $H * K$  to denote the cyclic group of order  $p^m$ , the direct product of  $n$  cyclic groups of order  $p^m$ , and a central product of  $H$  and  $K$  respectively. We use  $M \triangleleft G$  to denote  $M$  is a maximal subgroup of  $G$  and

$$G > G' = G_2 > G_3 > \cdots > G_{c+1} = 1$$

denote the lower central series of  $G$ , where  $c = c(G)$ .

Let  $G$  be a finite  $p$ -group. We use  $G \in \mathcal{A}_t$  to denote  $G$  is an  $\mathcal{A}_t$ -group. For any positive integer  $s$ , we define

$$\Omega_s(G) = \langle a \in G \mid a^{p^s} = 1 \rangle \text{ and } \mathcal{U}_s(G) = \langle a^{p^s} \mid a \in G \rangle.$$

We use  $M_p(n, m)$  to denote the  $p$ -groups

$$\langle a, b \mid a^{p^n} = b^{p^m} = 1, a^b = a^{1+p^{n-1}} \rangle, \text{ where } n \geq 2.$$

We use  $M_p(n, m, 1)$  to denote the  $p$ -groups

$$\langle a, b, c \mid a^{p^n} = b^{p^m} = c^p = 1, [a, b] = c, [c, a] = [c, b] = 1 \rangle,$$

where

$$n \geq m, \text{ and if } p = 2, \text{ then } m + n \geq 3.$$

For other notation and terminology the reader is referred to [17].

The following results are used in this paper, we gather them together.

**Lemma 2.1.** ([12, Theorem 2.3]) *Suppose that  $G$  is a finite  $p$ -group. Then  $G$  is metacyclic if and only if  $G/\Phi(G')G_3$  is metacyclic.*

**Lemma 2.2.** ([29, Lemma 2.2]) *Suppose that  $G$  is a finite non-abelian  $p$ -group. Then the following conditions are equivalent:*

- (1)  $G$  is minimal non-abelian;
- (2)  $d(G) = 2$  and  $|G'| = p$ ;
- (3)  $d(G) = 2$  and  $\Phi(G) = Z(G)$ .

**Lemma 2.3.** ([25]) *Suppose that  $G$  is an  $\mathcal{A}_1$ -group. Then  $G$  is one of the following groups:  $Q_8$ ,  $M_p(n, m)$  or  $M_p(n, m, 1)$ .*

**Lemma 2.4.** ([7, §9, Exercise 10]) *Let  $G$  be a 3-group of maximal class. Then the fundamental subgroup  $G_1$  of  $G$  is either abelian or minimal nonabelian.*

**Lemma 2.5.** ([31]) *A finite  $p$ -group  $G$  is an  $\mathcal{A}_2$ -group if and only if  $G$  is one of the following pairwise non-isomorphic groups:*

- (I)  $d(G) = 2$  and  $G$  has an abelian subgroup of index  $p$ . In this case,  $\alpha_1(G) = p$ .
- (1)  $\langle a, b \mid a^8 = b^{2^m} = 1, [a, b] = a^{-2} \rangle$ , where  $m \geq 1$ ;
  - (2)  $\langle a, b \mid a^8 = b^{2^m} = 1, [a, b] = a^2 \rangle$ , where  $m \geq 1$ ;
  - (3)  $\langle a, b \mid a^8 = 1, b^{2^m} = a^4, [a, b] = a^{-2} \rangle$ , where  $m \geq 1$ ;
  - (4)  $\langle a_1, b; a_2, a_3 \mid a_1^p = a_2^p = a_3^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = 1, [a_i, a_j] = 1 \rangle$ , where  $p \geq 5$  for  $m = 1$ ,  $p \geq 3$  and  $1 \leq i, j \leq 3$ ;
  - (5)  $\langle a_1, b; a_2 \mid a_1^p = a_2^p = b^{p^{m+1}} = 1, [a_1, b] = a_2, [a_2, b] = b^{p^m}, [a_1, a_2] = 1 \rangle$ , where  $p \geq 3$ ;
  - (6)  $\langle a_1, b; a_2 \mid a_1^{p^2} = a_2^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_1^{\nu p}, [a_1, a_2] = 1 \rangle$ , where  $p \geq 3$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .
  - (7)  $\langle a_1, b; a_2 \mid a_1^9 = a_2^3 = 1, b^3 = a_1^3, [a_1, b] = a_2, [a_2, b] = a_1^{-3}, [a_2, a_1] = 1 \rangle$ .
- (II)  $d(G) = 3$ ,  $|G'| = p$  and  $G$  has an abelian subgroup of index  $p$ . In this case,  $\alpha_1(G) = p^2$ .
- (8)  $\langle a, b, x \mid a^4 = x^2 = 1, b^2 = a^2 = [a, b], [x, a] = [x, b] = 1 \rangle \cong Q_8 \times C_2$ ;
  - (9)  $\langle a, b, x \mid a^{p^{n+1}} = b^{p^m} = x^p = 1, [a, b] = a^{p^n}, [x, a] = [x, b] = 1 \rangle \cong M_p(n+1, m) \times C_p$ ;
  - (10)  $\langle a, b, x; c \mid a^{p^n} = b^{p^m} = c^p = x^p = 1, [a, b] = c, [c, a] = [c, b] = [x, a] = [x, b] = 1 \rangle \cong M_p(n, m, 1) \times C_p$ , where  $n \geq m$ , and  $n \geq 2$  if  $p = 2$ ;
  - (11)  $\langle a, b, x \mid a^4 = 1, b^2 = x^2 = a^2 = [a, b], [x, a] = [x, b] = 1 \rangle \cong Q_8 * C_4$ ;
  - (12)  $\langle a, b, x \mid a^{p^n} = b^{p^m} = x^{p^2} = 1, [a, b] = x^p, [x, a] = [x, b] = 1 \rangle \cong M_p(n, m, 1) * C_{p^2}$ , where  $n \geq 2$  if  $p = 2$  and  $n \geq m$ .
- (III)  $d(G) = 3$ ,  $|G'| = p^2$  and  $G$  has an abelian subgroup of index  $p$ . In this case,  $\alpha_1(G) = p^2 + p$ .
- (13)  $\langle a, b, c \mid a^4 = b^4 = 1, c^2 = a^2b^2, [a, b] = b^2, [c, a] = a^2, [c, b] = 1 \rangle$ ;
  - (14)  $\langle a, b, d \mid a^{p^m} = b^{p^2} = d^p = 1, [a, b] = a^{p^{m-1}}, [d, a] = b^p, [d, b] = 1 \rangle$ , where  $m \geq 3$  if  $p = 2$ ;
  - (15)  $\langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}, [d, b] = 1 \rangle$ , where  $(j, p) = 1$ ,  $p > 2$ ,  $j$  is a fixed quadratic non-residue modulo  $p$ , and  $-4j$  is a quadratic non-residue modulo  $p$ ;

(16)  $\langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}d^p, [d, b] = 1 \rangle$ , where if  $p$  is odd, then  $4j = 1 - \rho^{2r+1}$  with  $1 \leq r \leq \frac{p-1}{2}$  and  $\rho$  the smallest positive integer which is a primitive root (mod  $p$ ); if  $p = 2$ , then  $j = 1$ .

(IV)  $d(G) = 2$  and  $G$  has no abelian subgroup of index  $p$ . In this case,  $\alpha_1(G) = 1 + p$ .

(17)  $\langle a, b \mid a^{p^{r+2}} = 1, b^{p^{r+s+t}} = a^{p^{r+s}}, [a, b] = a^{p^r} \rangle$ , where  $r \geq 2$  for  $p = 2$ ,  $r \geq 1$  for  $p \geq 3$ ,  $t \geq 0$ ,  $0 \leq s \leq 2$  and  $r + s \geq 2$ ;

(18)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = a^p \rangle$ , where  $p \geq 5$ ,  $\nu$  is a fixed quadratic non-residue modulo  $p$ ;

(19)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = a^{-p}b^{-lp}, [c, b] = a^{-p} \rangle$ , where  $p \geq 5$ ,  $4l = \rho^{2r+1} - 1$ ,  $r = 1, 2, \dots, \frac{1}{2}(p-1)$ ,  $\rho$  is the smallest positive integer which is a primitive root modulo  $p$ ;

(20)  $\langle a, b, c \mid a^9 = b^9 = c^3 = 1, [a, b] = c, [c, a] = b^{-3}, [c, b] = a^3 \rangle$ ;

(21)  $\langle a, b, c \mid a^9 = b^9 = c^3 = 1, [a, b] = c, [c, a] = b^{-3}, [c, b] = a^{-3} \rangle$ .

(V)  $d(G) = 3$  and  $G$  has no abelian subgroup of index  $p$ . In this case,  $\alpha_1(G) = 1 + p + p^2$ .

(22)  $\langle a, b, d \mid a^4 = b^4 = d^4 = 1, [a, b] = d^2, [d, a] = b^2d^2, [d, b] = a^2b^2, [a^2, b] = [b^2, a] = 1 \rangle$ .

By analyzing the groups in Lemma 2.5, we have following lemma.

**Lemma 2.6.** *Suppose that  $G$  is an  $\mathcal{A}_2$ -group of order  $p^n$ . Then*

- (1)  $d(G) \leq 3$  and  $c(G) \leq 3$ ;
- (2) if  $d(G) = 3$ , then  $c(G) = 2$  and  $G' \leq C_p^3$ . Moreover, if  $G' = C_p^3$ , then  $|G| = 2^6$ ,  $G' = \Omega_1(G) = Z(G) \cong C_2^3$  and  $G$  has no abelian subgroup of index  $p$ ;
- (3) if  $d(G) = 3$  and  $|G'| = p$ , then  $|G : Z(G)| = p^2$  and the type of  $G/G'$  is  $(p^u, p^v, p)$  where  $u + v = n - 2$ ;
- (4) if  $d(G) = 3$  and  $|G'| = p^2$ , then  $c(G) = 2$ ,  $\Phi(G) = \mathcal{U}_1(G) = Z(G)$ ,  $G$  has a unique abelian subgroup  $A$  of index  $p$  and  $\exp(A) > p$ ,  $G/G'$  has the type  $(p^{n-4}, p, p)$  and  $A/G'$  has the type  $(p^{n-5}, p, p)$ .
- (5) if  $G$  has an abelian subgroup of index  $p$ , then  $G$  is metacyclic if and only if  $p = 2$ ;
- (6) if  $p = 2$ , then  $G$  is metacyclic if and only if  $d(G) = 2$ ;
- (7) if  $|G'| = p$ , then  $\alpha_1(G) = p^2$ ;
- (8) if  $d(G) = 2$  and  $|G'| = p^3$ , then  $\alpha_1(G) = 1 + p$ ;

(9) if  $d(G) = 2$  and  $G' \cong C_p^2$ , then  $\alpha_1(G) = p$ .

**Lemma 2.7.** ([7, p27, Exercise 6]) *Let  $G$  be a non-abelian  $p$ -group. Then the number of abelian subgroups of index  $p$  in  $G$  is  $0, 1$  or  $p + 1$ .*

**Lemma 2.8.** [17, p259, Aufgabe 2]) *Suppose that a finite non-abelian  $p$ -group  $G$  has an abelian normal subgroup  $A$ , and  $G/A = \langle bA \rangle$  is cyclic. Then the map  $a \mapsto [a, b]$ ,  $a \in A$ , is an epimorphism from  $A$  to  $G'$ , and  $G' \cong A/A \cap Z(G)$ . In particular, if a non-abelian  $p$ -group  $G$  has an abelian subgroup of index  $p$ , then  $|G| = p|G'| |Z(G)|$ .*

**Lemma 2.9.** *Let  $p$  be an odd prime. Then the equation  $x^2 + ry^2 - u = 0$  about  $x, y$  over  $F_p$  has a solution, and the following conclusions hold:*

(1) *If  $-r \in (F_p^*)^2$  and  $u \in F_p^*$ , then the equation  $x^2 + ry^2 - u = 0$  has exactly  $p - 1$  solutions;*

(2) *If  $-r \notin (F_p^*)^2$  and  $u \in F_p^*$ , then the equation  $x^2 + ry^2 - u = 0$  has exactly  $p + 1$  solutions.*

**Proof** Let  $A = \{a^2 \mid a \in F_p\}$  and  $B = \{u - rb^2 \mid b \in F_p\}$ . Then  $|A| = |B| = \frac{p+1}{2}$ . It follows that  $A \cap B \neq \emptyset$ . Hence there exist  $a, b \in F_p$  such that  $a^2 = u - rb^2$ .

(1) Let  $-r = \alpha^2$ . Then the equation  $x^2 + ry^2 - u = 0$  is  $x^2 - \alpha^2 y^2 = u$ , and hence is equivalent to

$$\begin{cases} x + \alpha y = k \\ x - \alpha y = k^{-1}u \end{cases}$$

where  $k = 1, 2, \dots, p - 1$ . Hence the equation  $x^2 + ry^2 - u = 0$  has exactly  $p - 1$  solutions.

(2) Let  $n(u)$  be the number of solutions of the equation  $x^2 + ry^2 - u = 0$ . Since  $-r \notin (F_p^*)^2$ ,  $n(0) = 1$ . It is obvious that  $n(u_1) = n(u_2)$  for  $u_1 u_2^{-1} \in (F_p^*)^2$ . Let  $\nu \notin (F_p^*)^2$ . Since  $F_p \times F_p$  has a partition

$$\{(x, y) \in F_p \times F_p \mid x^2 + ry^2 \in (F_p^*)^2\} \cup \{(x, y) \in F_p \times F_p \mid x^2 + ry^2 \notin (F_p^*)^2\},$$

$p^2 = n(0) + \frac{p-1}{2}n(1) + \frac{p-1}{2}n(\nu)$ . It follows that  $n(1) + n(\nu) = 2p + 2$ . If we prove that  $n(1) = p + 1$ , then  $n(u) = p + 1$  for all  $u \in F_p^*$ .

Now we calculate  $n(1)$ . If  $y = 0$ , then the equation  $x^2 + ry^2 - 1 = 0$  has two solutions  $(1, 0)$  and  $(-1, 0)$ . If  $y \neq 0$ , then the equation  $x^2 + ry^2 - 1 = 0$  is equivalent to  $(xy^{-1})^2 - y^{-2} + r = 0$ . By (1), the later has  $p - 1$  solutions. Hence  $n(1) = p + 1$ .  $\square$

**Lemma 2.10.** *Let  $p$  be an odd prime. If  $s^2 - 4r \not\equiv 0 \pmod{p}$ , then the equation  $x^2 + sxy + ry^2 + wx + vy + u = 0$  about  $x, y$  over  $F_p$  has a solution.*

**Proof** Let  $x_1 = x + 2^{-1}sy$ ,  $y_1 = y$ ,  $r_1 = r - 4^{-1}s^2$  and  $v_1 = v - 2^{-1}ws$ . Then the equation is turned to  $x_1^2 + r_1 y_1^2 + w x_1 + v_1 y_1 + u = 0$ . Let  $x_2 = x_1 + 2^{-1}w$ ,  $y_2 = y_1 + 2^{-1}r_1^{-1}v_1$  and  $u_2 = u - 4^{-1}w^2 - 4^{-1}r_1^{-1}v_1^2$ . Then the equation is turned to  $x_2^2 + r_1 y_2^2 + u_2 = 0$ . By Lemma 2.9, the last equation has a solution  $(a, b)$ . Thus  $(a - 2^{-1}w - 2^{-1}sb + 4^{-1}sr_1^{-1}v_1, b - 2^{-1}r_1^{-1}v_1)$  is a solution of the first equation.  $\square$

**Lemma 2.11.** ([29, Lemma 3.1]) *Let  $G$  be a non-abelian two-generator  $p$ -group having an abelian subgroup  $A$  of index  $p$ . Assume that  $|G/G'| = p^{m+1}$  and  $c(G) = c$ . Then  $m \geq 1$ ,  $c \geq 2$  and*

- (1)  $G$  has the lower central complexion  $(m+1, \underbrace{1, \dots, 1}_{c-1})$  and hence  $|G'| = p^{c-1}$ ,  $|G| = p^{m+c}$ ;
- (2)  $|Z(G)| = p^m$  and  $G/Z(G)$  is of maximal class;
- (3)  $Z(G) \leq \Phi(G)$ ,  $\Phi(G) = G'Z(G)$  and  $G' \cap Z(G) = G_c$ ;
- (4) Let  $M$  be a non-abelian subgroup of index  $p$  of  $G$ . Then  $Z(M) = Z(G)$  and

$$M' = G_3, M_3 = G_4, \dots, M_{c-1} = G_c.$$

**Lemma 2.12.** ([29, Corollary 3.8]) *Suppose that  $G$  is a finite  $p$ -group having an abelian subgroup of index  $p$ , and all non-abelian subgroups of  $G$  are generated by two elements. Then*

- (1)  $G \in \mathcal{A}_2$  if and only if  $c(G) = 3$ ;
- (2) If  $p = 2$ , then  $G$  is metacyclic;
- (3) If  $c(G) \leq p$ , then  $d(G') = c(G) - 1$ . If  $c(G) \geq p + 1$ , then  $d(G') = p - 1$ .

**Lemma 2.13.** ([29, Theorem 4.1]) *Suppose that  $G$  is a three-generator non-abelian  $p$ -group, and all non-abelian proper subgroups of  $G$  are generated by two elements. If  $G$  has an abelian subgroup of index  $p$ , then  $G \in \mathcal{A}_2$ .*

**Lemma 2.14.** ([13, Theorem 4]) *Let  $G$  be a  $p$ -group. If both  $G$  and  $G'$  can be generated by two elements, then  $G'$  is abelian.*

**Lemma 2.15.** *Suppose that  $G$  is a finite  $p$ -group.  $M_1$  and  $M_2$  are two distinct maximal subgroups of  $G$ . Then  $|G'| \leq p|M'_1M'_2|$ .*

**Proof** Let  $\bar{G} = G/M'_1M'_2$ . Then  $\bar{G}$  has abelian subgroups  $\bar{M}_1$  and  $\bar{M}_2$  of index  $p$ . Hence  $Z(\bar{G}) \geq \bar{M}_1 \cap \bar{M}_2$ . By Lemma 2.8,  $|\bar{G}'| \leq p$ . Thus  $|G'| \leq p|M'_1M'_2|$ .  $\square$

**Lemma 2.16.** ([1, Theorem 5.6]) *If  $G$  is a finite non-abelian  $p$ -group and  $|G'| = p^k$ , then  $G$  has a subgroup  $K$  such that  $d(K) \leq k + 1$  and  $K_n = G_n$  for all  $2 \leq n \leq c(G)$ .*

**Proposition 2.17.** ([27]) *Let  $G$  be a metabelian group and  $a, b \in G$ . For any positive integers  $i$  and  $j$ , let*

$$[ia, jb] = [a, \underbrace{b, a, \dots, a}_{i-1}, \underbrace{b, \dots, b}_{j-1}].$$

*Then, for any positive integers  $m$  and  $n$ ,*

$$[a^m, b^n] = \prod_{i=1}^m \prod_{j=1}^n [ia, jb] \binom{m}{i} \binom{n}{j}.$$



The following lemma is equivalent to [19, Theorem 3.4]

**Lemma 2.18.** *Assume  $G$  is a finite  $p$ -group. If  $c(H) \leq 2$  for all  $H < G$ , then  $c(G) \leq 3$ .*

Let  $G$  be a group of order  $p^m$ ,  $|G : \Phi(G)| = p^d$ ,

$$\Gamma_i = \{H < G \mid \Phi(G) \leq H, |G : H| = p^i\}$$

and  $\mathfrak{M}$  be a set of proper subgroups of  $G$ . For  $K \leq G$ , we denote by  $\alpha(K)$  the number of members of the set  $\mathfrak{M}$  that are subgroups of  $K$ . Obviously,  $\alpha(G) = |\mathfrak{M}|$ .

**Theorem 2.19.** (*Hall's enumeration principle*) *In the above notation,*

$$\alpha(G) = \sum_{i=1}^d \sum_{H \in \Gamma_i} (-1)^{i-1} p^{\binom{i}{2}} \alpha(H).$$

### 3 Some general properties of $\mathcal{A}_t$ -groups with $t \geq 3$

Y. Berkovich was the first where  $\mathcal{A}_t$ -groups were investigated. He obtained many significant results. The following are some related to  $\mathcal{A}_3$ -groups.

**Lemma 3.1.** ([5, Lemma J(i)]) *Let  $G$  be a metacyclic  $p$ -group. Then  $G$  is an  $\mathcal{A}_t$ -group if and only if  $|G'| = p^t$ .*

**Lemma 3.2.** ([8, Proposition 72.2]) *If a  $p$ -group  $G$  is an  $\mathcal{A}_3$ -group, then  $|G'| \leq p^4$ . For  $p = 5$ , there exists an  $\mathcal{A}_3$ -group  $G$  with  $|G'| = 5^4$ .*

**Lemma 3.3.** ([8, Proposition 72.3]) *Suppose that a  $p$ -group  $G$  is an  $\mathcal{A}_3$ -group with  $|G'| = p^4$ . Then*

- (a)  $G'$  is abelian.
- (b) If  $\exp(G') = p$ , then  $p > 2$ ,  $G/G'$  is abelian of type  $(p^n, p)$  ( $n \geq 1$ ) and  $G' \cong C_p^m$ .  
If, in addition,  $n > 1$ , then  $\Omega_1(G) = G'$ .
- (c)  $G'$  is not metacyclic.

**Theorem 3.4.** ([8, Theorem 72.6]) *If a  $p$ -group  $G$  is an  $\mathcal{A}_4$ -group, then  $|G'| \leq p^6$ .*

In following we give more information about  $\mathcal{A}_t$ -groups.

**Lemma 3.5.** ([2, Corollary 2.4]) (1) *Let  $M$  be an  $\mathcal{A}_t$ -group, and  $A$  be an abelian group of order  $p^k$ . Then  $G = M \times A$  is an  $\mathcal{A}_{t+k}$ -group.*

(2) *Let  $M$  be an  $\mathcal{A}_t$ -group with  $|M'| = p$ ,  $G = M * A$ , where  $A$  is an abelian group of order  $p^{k+1}$  and  $M \cap A = G'$ . Then  $G$  is an  $\mathcal{A}_{t+k}$ -group.*

**Lemma 3.6.** ([24, Theorem 5.4]) *Assume  $G$  is a finite 2-group having a unique  $\mathcal{A}_1$ -subgroup  $A$  of index  $p$ . If  $|G| = 2^t \geq 2^9$  and  $G/A'$  has a unique abelian subgroup of index  $p$ , then one of the following is true:*

- (1)  $G$  is metacyclic;
- (2)  $G$  is an  $\mathcal{A}_{t-2}$ -group or an  $\mathcal{A}_{t-3}$ -group.

**Lemma 3.7.** ([24, Corollary 5.5]) *Assume that  $G$  is a finite 2-group having a unique  $\mathcal{A}_1$ -subgroup  $A$  of index  $p$ . If  $G$  is an  $\mathcal{A}_3$ -group and  $|G| \geq 2^9$ , then  $G$  is metacyclic.*

**Lemma 3.8.** *Suppose that  $G$  is a finite  $p$ -group having an abelian subgroup of index  $p$ . If all non-abelian subgroups of  $G$  are generated by two elements, then  $G$  is an  $\mathcal{A}_t$ -group if and only if  $c(G) = t + 1$ .*

**Proof** Suppose that  $c(G) = t + 1$  and  $M$  is a non-abelian subgroup of index  $p$ . By Lemma 2.11(4),  $c(M) = t$ . By hypothesis,  $d(M) = 2$ . By induction on  $t$ ,  $M$  is an  $\mathcal{A}_{t-1}$ -group. Hence  $G$  is an  $\mathcal{A}_t$ -group.  $\square$

**Lemma 3.9.** *If  $G$  is a  $p$ -group of maximal class of order  $p^n$  with  $n \geq 3$ , then  $G$  is an  $\mathcal{A}_{n-2}$ -group.*

**Proof** By [7, Theorem 9.6(f)], there exists a subgroup  $A$  of order  $p^2$  of  $G$  such that  $C_G(A) = A$ . Let  $B$  be a subgroup of  $G$  with  $B \geq A$  and  $|B| = p^3$ . Then  $B$  is not abelian. It follows that  $G \in \mathcal{A}_{n-2}$ .  $\square$

**Lemma 3.10.** *If a  $p$ -group  $G$  is an  $\mathcal{A}_t$ -group and  $G'$  is regular, then  $\exp(G') \leq p^t$ .*

**Proof** By induction on  $t$  we have  $\exp(M') \leq p^{t-1}$  for all  $M \triangleleft G$ . Let  $N = \prod_{M \triangleleft G} M'$ . Since  $G'$  is regular,  $\exp(N) \leq p^{t-1}$  by [7, Theorem 7.2(b)]. It is easy to see that  $G/N$  is abelian or an  $\mathcal{A}_1$ -group. It follows by Lemma 2.2(2) that  $|G'N/N| = |(G/N)'| \leq p$ . Thus  $\exp(G') \leq p \cdot p^{t-1} = p^t$ .  $\square$

**Corollary 3.11.** *Assume  $G$  is a finite  $p$ -group.*

- (1) *If  $G$  is an  $\mathcal{A}_3$ -group, then  $\exp(G') \leq p^3$ .*
- (2) *If  $p \geq 7$  and  $G$  is an  $\mathcal{A}_4$ -group, then  $\exp(G') \leq p^4$ .*

**Proof** By Lemma 3.3(a), Theorem 3.4 and [7, Theorem 7.1(b)] we get  $G'$  is regular. Thus the conclusion follows by Lemma 3.10.  $\square$

**Theorem 3.12.** *If a  $p$ -group  $G$  is an  $\mathcal{A}_3$ -group, then  $c(G) \leq 4$ .*

**Proof** Let  $N = \prod_{M \triangleleft G} M_3$ . Since  $M_3$  is characteristic in  $M$ ,  $M_3 \triangleleft G$ . Thus  $N \triangleleft G$ . By Lemma 2.18,  $c(G/N) \leq 3$ . If  $N \leq Z(G)$ , then  $c(G) \leq 4$ . Assume  $N \not\leq Z(G)$ . Then there exists  $M \triangleleft G$  such that  $M_3 \not\leq Z(G)$ . Thus  $|M_3| \geq p^2$ . Notice that  $M$  is an  $\mathcal{A}_2$ -group. Hence  $|M| = p^5$  by checking the list of groups in Lemma 2.5. Thus  $|G| = p^6$ . If  $c(G) = 5$ , then  $G$  is of maximal class. It follows by Lemma 3.9 that  $G$  an  $\mathcal{A}_4$ -group. This is a contradiction.  $\square$

**Remark 3.13.** *There exist  $\mathcal{A}_3$ -groups of class 4. In fact, those groups listed in Theorem 5.1 are all  $\mathcal{A}_3$ -groups of class 4.*

**Lemma 3.14.** *Assume a  $p$ -group  $G$  is an  $\mathcal{A}_3$ -group.*

- (1) *If  $\Phi(G)$  is non-abelian, then  $\Phi(G)$  is metacyclic.*
- (2) *If  $p > 2$ , then  $\Phi(G)$  is non-abelian if and only if  $G$  is metacyclic.*

**Proof** (1) Since  $G$  is an  $\mathcal{A}_3$ -group,  $d(G) = 2$  and  $\Phi(G) \in \mathcal{A}_1$ . By Lemma 2.2,  $d(\Phi(G)) = 2$ . It follows from [7, Theorem 44.12] that  $\Phi(G)$  is metacyclic.

(2)  $\Leftarrow$  Let  $G = \langle a, b \rangle$ . Then  $o([a, b]) = p^3$  by Lemma 3.1. Since  $G$  is regular,  $[a^p, b^p] = 1$  if and only if  $[a, b]^{p^2} = 1$  by [7, Theorem 7.2(e)]. Since  $o([a, b]) = p^3$ ,  $[a^p, b^p] \neq 1$ . It follows that  $\Phi(G)$  is non-abelian.

$\Rightarrow$  Assume  $G$  is not metacyclic. It follows by Lemma 3.1 and Lemma 3.3(a) and (c) that  $|G' : \Omega_1(G')| \leq p$ . Thus  $G_3 \leq \Omega_1(G')$ . It follows that  $\exp(G_3) \leq p$ . on the other hand,  $c(G) \leq 4$  by Lemma 3.12. Since  $G'$  is abelian,  $G$  is metabelian. By lemma 2.17 we have

$$[x^p, y^p] = \prod_{i=1}^p \prod_{j=1}^p [ix, jy] \binom{p}{i} \binom{p}{j}, \quad [x^p, z] = \prod_{i=1}^p [ix, z] \binom{p}{i}.$$

where  $x, y \in G, z \in G'$ .

It follows from  $\exp(G') \leq p^2$ ,  $\exp(G_3) \leq p$ ,  $c(G) \leq 4$  and  $p > 2$  that  $[x^p, y^p] = 1$  and  $[x^p, z] = 1$ . Thus  $\Phi(G)$  is abelian. This is a contradiction.  $\square$

**Remark 3.15.** *In Lemma 3.14, if  $p = 2$ , then (2) is not true. For example,  $D_{32}$  is a metacyclic  $\mathcal{A}_3$ -group, but  $\Phi(D_{32}) \cong C_8$ .*

**Lemma 3.16.** *Suppose that  $G$  is an  $\mathcal{A}_3$ -group. If  $G$  has a maximal subgroup  $M$  such that  $d(M) = 3$  and  $M' \not\leq Z(G)$ , then  $G$  has a three-generator maximal subgroup  $L$  such that  $d(L) = 3$ ,  $|L'| \leq p^2$  and  $L' \not\leq Z(G)$ .*

**Proof** Otherwise, for every three-generator maximal subgroup  $L$  with  $d(L) = 3$  and  $|L'| \leq p^2$ , we have  $L' \leq Z(G)$ . Hence  $|M'| \geq p^3$ . It follows from Lemma 2.6(2) that

$$p = 2, |M| = 2^6 \text{ and } M' = \Omega_1(M) = Z(M) \cong C_2^3.$$

Let  $d(G) = r$  where  $2 \leq r \leq 4$ . Then  $G$  has  $s = 1 + 2 + \dots + 2^{r-1}$  maximal subgroups. Let  $H_1, H_2, \dots, H_{s-1}$  and  $M$  be all maximal subgroups of  $G$ . Since  $M' \leq H_i$ ,  $H_i$  is not metacyclic. It follows from Lemma 2.6(6) that  $d(H_i) = 3$ . If  $|H'| = 8$ , then we also have  $H' = \Omega_1(H) = Z(H)$ . Since  $M' \cong C_2^3 \leq H$ ,  $M' = H'$ . It follows that  $[M', G] = [M', HM] = 1$  and hence  $M' \leq Z(G)$ , a contradiction. Hence  $|H'_i| \leq 4$  and  $H'_i \leq Z(G)$ . Let  $N = H'_1 H'_2 \dots H'_{s-1}$ . Then  $G/N$  has  $s - 1$  abelian subgroups of index  $p$ . Since  $s - 1 \geq 2$ , we have, by Lemma 2.7, the number of abelian subgroups of index  $p$  of  $G/N$  is 3. Hence  $s - 1 \leq 3$ . It follows that  $d(G) = 2$  and  $M/N$  is also abelian. Thus  $M' \leq N \leq Z(G)$ , a contradiction.  $\square$

**Lemma 3.17.** *Suppose that  $G$  is a four-generator  $\mathcal{A}_3$ -group. Then  $c(G) = 2$ ,  $\Phi(G) \leq Z(G)$ ,  $G' \leq C_p^3$  and all  $\mathcal{A}_1$ -subgroups of  $G$  contain  $\Phi(G)$ .*

**Proof** Let  $M$  be an  $\mathcal{A}_1$ -subgroup of  $G$  and  $L$  be a maximal subgroup containing  $M$ . Since  $d(G) = 4$ , we have  $d(L) \geq 3$ . Hence  $L \in \mathcal{A}_2$ . By Lemma 2.6(2), we get  $d(L) = 3$ ,  $c(L) = 2$  and  $L' \leq C_p^3$ . By comparing the indexes of  $\Phi(M)$ ,  $\Phi(L)$  and  $\Phi(G)$ , we get

$$Z(M) = \Phi(M) = \Phi(L) = \Phi(G).$$

By the arbitrariness of  $M$  we get  $\Phi(G) \leq Z(G)$ . Hence  $c(G) = 2$ . Let  $N = \prod_{L < G} L'$ . Then  $N \leq Z(G)$  and  $\exp(N) = p$ . Since all subgroups of  $G/N$  are abelian, we have  $G/N$  is also abelian. It follows that  $G' = N$ . Hence  $\exp(G') = p$ . Since  $G' \leq \Omega_1(\Phi(G)) = \Omega_1(\Phi(M))$ ,  $G' \leq C_p^3$ .  $\square$

**Lemma 3.18.** *Let  $G$  be an  $\mathcal{A}_3$ -group such that  $d(G) = 3$  and  $\Phi(G) \leq Z(G)$ . Then  $\alpha_1(G) = \mu_1 + p^2\mu_2$ .*

**Proof** Groups satisfying  $d(G) = 3$  and  $\Phi(G) \leq Z(G)$  were classified in [2, 23]. The derived groups of non-abelian maximal subgroups of  $G$  are of order  $p$ .

Let  $H \in \Gamma_2$ . Then, by  $\Phi(G) \leq Z(G)$ ,  $H$  is abelian and hence  $\alpha_1(H) = 0$ . Let  $M \in \Gamma_1$ . If  $M \in \mathcal{A}_2$ , then, by Lemma 2.6(7),  $\alpha_1(M) = p^2$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p^2\mu_2$ .  $\square$

## 4 $\mathcal{A}_3$ -groups with an $\mathcal{A}_1$ -subgroup of index $p$

In this section, we classify  $\mathcal{A}_3$ -groups with an  $\mathcal{A}_1$ -subgroup of index  $p$ . Since finite  $p$ -groups with an  $\mathcal{A}_1$ -subgroup of index  $p$  are classified by [2, 3, 22, 23, 24], it is enough to check whether those groups in [2, 3, 22, 23, 24] are  $\mathcal{A}_3$ -groups or not. In this case,  $\mathcal{A}_3$ -groups have 72 non-isomorphic types.

For convenience, in this section, assume  $G$  is an  $\mathcal{A}_3$ -group with an  $\mathcal{A}_1$ -subgroup of index  $p$  in Theorem 4.1, 4.2, 4.4, 4.6 and 4.7. It is easy to see that  $d(G) \leq 3$ .

**Theorem 4.1.**  *$G$  has an abelian subgroup of index  $p$  and  $d(G) = 2$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Ai)  $c(G) = 3$  and  $G' \cong C_4$ .

(A1)  $\langle a, b, c \mid a^4 = b^2 = c^4 = 1, [a, b] = c, [c, a] = [c, b] = c^2 \rangle$ . Moreover  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_4 \times C_2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2, c^2 \rangle \cong C_2^2$ .

(A2)  $\langle a, b, c \mid a^4 = b^4 = 1, c^2 = a^2, [a, b] = c, [c, a] = [c, b] = c^2 \rangle$ . Moreover,  $|G| = 2^5$ ,  $\Phi(G) = \langle b^2, c \rangle \cong C_4 \times C_2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle b^2, c^2 \rangle \cong C_2^2$ .

(A3)  $\langle a, b, c \mid a^8 = b^2 = 1, c^2 = a^4, [a, b] = c, [c, a] = [c, b] = c^2 \rangle$ . Moreover,  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_4 \times C_2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2 \rangle \cong C_4$ .

(Aii)  $c(G) = 3$  and  $G' \cong C_p^2$  where  $p > 2$ .

- (A4)  $\langle a, b, c \mid a^{p^3} = b^p = c^p = 1, [a, b] = c, [c, a] = 1, [c, b] = a^{\nu p^2} \rangle$ , where  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ . Moreover,  $|G| = p^5$ ,  $\Phi(G) = \langle a^p, c \rangle \cong C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^p \rangle \cong C_{p^2}$ .
- (A5)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = 1, [c, b] = b^p \rangle$ . Moreover,  $|G| = p^5$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_p^3$ ,  $G' = \langle b^p, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_p^2$ .
- (A6)  $\langle a, b, c, d \mid a^{p^2} = b^p = c^p = d^p = 1, [a, b] = c, [c, a] = 1, [c, b] = d, [d, a] = [d, b] = 1 \rangle$ . Moreover,  $|G| = p^5$ ,  $\Phi(G) = \langle a^p, c, d \rangle \cong C_p^3$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle a^p, d \rangle \cong C_p^2$ .

Moreover,  $(\mu_0, \mu_1, \mu_2) = (1, p-1, 1)$  and  $\alpha_1(G) = p^2 + p - 1$ .

**Proof** Let  $A$  be an abelian subgroup of index  $p$  and  $B$  be an  $\mathcal{A}_1$ -subgroup of index  $p$ . By Lemma 2.11,  $G_3 = B'$ ,  $c(G) = 3$  and  $|G'| = p^2$ . Let  $\bar{G} = G/G_3$ . Then  $|\bar{G}'| = p$ . Since  $d(\bar{G}) = d(G) = 2$ ,  $\bar{G} \in \mathcal{A}_1$  by Lemma 2.2. Obviously,  $G_3 = \Phi(G')G_3$ . If  $\bar{G}$  is metacyclic, then  $G$  is also metacyclic by Lemma 2.1. Hence  $G$  is an  $\mathcal{A}_2$ -group by Lemma 3.1. This contradicts that  $G$  is an  $\mathcal{A}_3$ -group. Thus  $\bar{G}$  is a non-metacyclic  $\mathcal{A}_1$ -group. By Lemma 2.3,  $\bar{G} \cong M_p(n, m, 1)$ . Now we have  $\Phi(G')G_3 \leq C_p^2$ ,  $\Phi(G')G_3 \leq Z(G)$  and  $G/\Phi(G')G_3 \cong M_p(n, m, 1)$ . Thus  $G$  is one of the groups classified in [3]. Since  $|G'| = p^2$ ,  $G$  is either one of the groups listed in [3, Theorem 3.5] or [3, Theorem 4.6]. By hypothesis, we only need pick out those groups satisfying the following three conditions:

- (1) the minimal index of  $\mathcal{A}_1$ -subgroups is 1;
- (2) the maximal index of  $\mathcal{A}_1$ -subgroups is 2;
- (3)  $G$  has an abelian subgroup of index  $p$ .

If  $G' \cong C_{p^2}$ , then  $G$  is one of the groups listed in [3, Theorem 3.5]. [3, Theorem 3.1 & 3.6] tell us those groups in [3, Theorem 3.5] satisfy the conditions (1) and (2). Moreover, it is easy to check which one satisfies the condition (3) in these groups. Thus we get the groups (A1)–(A3). If  $G' \cong C_p^2$ , then  $G$  is one of the groups listed in [3, Theorem 4.6]. [3, Theorem 4.1 & 4.7] tell us those groups in [3, Theorem 4.6] satisfy the conditions (1) and (2). Moreover, in these groups we check which one satisfies the condition (3). Thus we get the groups (A4)–(A6). For convenience, in Table 1 we give the relationship between the groups in Theorem 4.1 and those groups in paper [3].

Groups in Theorem 4.1	Groups in [3, Theorem 3.5]	Groups in Theorem 4.1	Groups in [3, Theorem 4.6]
(A1)	(A1)	(A4)	(J1) where $n = 2$ and $m = 1$
(A2)	(A2)	(A5)	(J3) where $n = 2$ and $m = 1$
(A3)	(A3)	(A6)	(J5) where $n = 2$ and $m = 1$

Table 1: The correspondence from Theorem 4.1 to [3, Theorem 3.5 & 4.6]

In following, we calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$ .

Since  $d(G) = 2$ ,  $G$  has  $1 + p$  maximal subgroups. For any  $\mathcal{A}_3$ -group, we always have  $\mu_2 \geq 1$ . By the hypothesis,  $\mu_0 \geq 1$  and  $\mu_1 \geq 1$ .

If  $G$  is one of the groups (A1)–(A3), then  $p = 2$ . Hence  $\mu_0 + \mu_1 + \mu_2 = 1 + p = 3$ . It follows that  $(\mu_0, \mu_1, \mu_2) = (1, 1, 1)$ .

If  $G$  is one of the groups (A4)–(A6), then it is easy to see that  $\langle c, a, \Phi(G) \rangle$  and  $\langle c, b, \Phi(G) \rangle$  are the unique abelian subgroup of index  $p$  and the  $\mathcal{A}_2$ -subgroup of index  $p$  respectively. Hence  $(\mu_0, \mu_1, \mu_2) = (1, p-1, 1)$ .

Now we calculate  $\alpha_1(G)$ . It is obvious that  $\alpha_1(\Phi(G)) = 0$ . Let  $M \in \Gamma_1$ . If  $M$  is the  $\mathcal{A}_2$ -subgroup of index  $p$ , then  $|M'| = p$ . By Lemma 2.6 (7),  $\alpha_1(M) = p^2$ .

By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p\alpha_1(\Phi(G)) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p^2\mu_2 = p^2 + p - 1.$$

□

**Theorem 4.2.**  $G$  has an abelian subgroup of index  $p$  and  $d(G) = 3$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

(Bi)  $c(G) = 2$  and  $G' \cong C_p$ .

(B1)  $\langle a, b, c \mid a^4 = c^4 = 1, b^2 = a^2 = [a, b], [c, a] = [c, b] = 1 \rangle \cong Q_8 \times C_4$ ; where  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c^2 \rangle \cong C_2 \times C_2$ ,  $G' = \langle a^2 \rangle$  and  $Z(G) = \langle a^2, c \rangle \cong C_4 \times C_2$ .

(B2)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^m} = c^{p^2} = 1, [a, b] = a^{p^n}, [c, a] = [c, b] = 1 \rangle \cong M_p(n+1, m) \times C_{p^2}$ , where  $\min\{n, m\} = 1$ ; where  $|G| = p^{m+n+3}$ ,  $\Phi(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_p$  if  $m > 1$ ,  $\Phi(G) \cong C_{p^n} \times C_p$  if  $m = 1$ ,  $G' = \langle a^{p^n} \rangle$ ,  $Z(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_{p^2}$  if  $m > 1$  and  $Z(G) \cong C_{p^n} \times C_{p^2}$  if  $m = 1$ .

(B3)  $\langle a, b, c, d \mid a^{p^n} = b^p = c^{p^2} = d^p = 1, [a, b] = d, [c, a] = [c, b] = 1 \rangle \cong M_p(n, 1, 1) \times C_{p^2}$ , where  $n \geq 2$  for  $p = 2$ ; where  $|G| = p^{n+4}$ ,  $\Phi(G) = \langle a^p, c^p, d \rangle \cong C_{p^{n-1}} \times C_p \times C_p$  if  $n > 1$ ,  $\Phi(G) \cong C_p \times C_p$  if  $n = 1$ ,  $G' = \langle d \rangle$ ,  $Z(G) = \langle a^p, c, d \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_p$  if  $n > 1$ ,  $Z(G) \cong C_{p^2} \times C_p$  if  $n = 1$ .

(B4)  $\langle a, b, c \mid a^4 = 1, b^2 = c^4 = a^2 = [a, b], [c, a] = [c, b] = 1 \rangle \cong Q_8 * C_8$ ; where  $|G| = 2^5$ ,  $\Phi(G) = \langle c^2 \rangle \cong C_4$ ,  $G' = \langle c^4 \rangle$  and  $Z(G) = \langle c \rangle \cong C_8$ .

(B5)  $\langle a, b, c \mid a^{p^n} = b^p = c^{p^3} = 1, [a, b] = c^{p^2}, [c, a] = [c, b] = 1 \rangle \cong M_p(n, 1, 1) * C_{p^3}$ , where  $n \geq 2$  for  $p = 2$ . Moreover,  $|G| = p^{n+4}$ ,  $\Phi(G) = \langle a^p, c^p \rangle \cong C_{p^{n-1}} \times C_{p^2}$  if  $n > 1$ ,  $\Phi(G) \cong C_{p^2}$  if  $n = 1$ ,  $G' = \langle c^{p^2} \rangle$ ,  $Z(G) = \langle a^p, c \rangle \cong C_{p^{n-1}} \times C_{p^3}$  if  $n > 1$ ,  $Z(G) \cong C_{p^3}$  if  $n = 1$ .

(Bii)  $c(G) = 2$  and  $G' \cong C_p^2$ .

(B6)  $\langle a, b, c \mid a^p = b^{p^2} = c^{p^2} = 1, [b, c] = 1, [c, a] = c^p, [a, b] = b^{-p} \rangle$ , where  $p$  is odd; and  $|G| = p^5$ ,  $\Phi(G) = G' = Z(G) = \langle b^p, c^p \rangle \cong C_p^2$ .

- (B7)  $\langle a, b, c \mid a^{p^l} = b^{p^2} = c^{p^2} = 1, [b, c] = 1, [c, a] = b^p c^p, [a, b] = b^{-p} \rangle$ , where  $p$  is odd; and  $|G| = p^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{l-1}} \times C_p \times C_p$  if  $l > 1$ ,  $\Phi(G) = Z(G) \cong C_p \times C_p$  if  $l = 1$ ,  $G' = \langle b^p, c^p \rangle$ .
- (B8)  $\langle a, b, c \mid a^{p^l} = b^{p^2} = c^{p^2} = 1, [b, c] = 1, [c, a] = b^p c^{tp}, [a, b] = b^{-tp} c^{\nu p} \rangle$ , where  $p$  is odd,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ,  $-\nu \in (F_p^*)^2$ , and  $t \in \{0, 1, \dots, \frac{p-1}{2}\}$  such that  $t^2 \neq -\nu$ ; and  $|G| = p^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{l-1}} \times C_p \times C_p$  if  $l > 1$ ,  $\Phi(G) = Z(G) \cong C_p \times C_p$  if  $l = 1$ ,  $G' = \langle b^p, c^p \rangle$ .
- (B9)  $\langle a, b, c \mid a^p = b^{p^3} = c^{p^3} = 1, [b, c] = 1, [c, a] = b^{p^2} c^{tp^2}, [a, b] = b^{-tp^2} c^{\nu p^2} \rangle$ , where  $p$  is odd,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ,  $-\nu \notin (F_p^*)^2$ , and  $t \in \{0, 1, \dots, \frac{p-1}{2}\}$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle b^p, c^p \rangle \cong C_{p^2} \times C_{p^2}$ ,  $G' = \langle b^{p^2}, c^{p^2} \rangle$ .
- (B10)  $\langle a, b, c \mid a^{2^l} = b^4 = c^4 = 1, [b, c] = 1, [c, a] = b^2, [a, b] = c^2 \rangle$ ; where  $|G| = 2^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_{2^{l-1}} \times C_2 \times C_2$  if  $l > 1$ ,  $\Phi(G) = Z(G) \cong C_2 \times C_2$  if  $l = 1$ ,  $G' = \langle b^2, c^2 \rangle$ .
- (B11)  $\langle a, b, c \mid a^2 = b^8 = c^8 = 1, [b, c] = 1, [c, a] = b^4, [a, b] = b^4 c^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = Z(G) = \langle b^2, c^2 \rangle \cong C_4 \times C_4$ ,  $G' = \langle b^4, c^4 \rangle$ .
- (B12)  $\langle a, b, c \mid a^{p^l} = b^{p^3} = c^{p^2} = 1, [b, c] = 1, [a, b] = c^{\nu p}, [c, a] = b^{p^2} \rangle$ , where  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; and  $|G| = p^{l+5}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{l-1}} \times C_{p^2} \times C_p$  if  $l > 1$ ,  $\Phi(G) = Z(G) \cong C_{p^2} \times C_p$  if  $l = 1$ ,  $G' = \langle b^{p^2}, c^p \rangle$ .
- (B13)  $\langle a, b, c \mid a^{p^{l+1}} = b^{p^m} = c^{p^2} = 1, [b, c] = 1, [c, a] = c^p, [a, b] = a^{p^l} \rangle$ , where  $m \leq 2$ ; moreover,  $|G| = p^{l+m+3}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^l} \times C_p^2$  if  $m = 2$ ,  $\Phi(G) = Z(G) \cong C_{p^l} \times C_p$  if  $m = 1$ ,  $G' = \langle a^{p^l}, c^p \rangle$ .
- (B14)  $\langle a, b, c \mid a^{p^{l+1}} = b^{p^{m+1}} = c^{p^n} = 1, [b, c] = 1, [c, a] = b^{p^m}, [a, b] = a^{p^l} \rangle$ , where  $\min\{l, m, n\} = 1$  and  $\max\{m, n\} = 2$ ; moreover,  $|G| = p^{l+m+n+2}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^l} \times C_{p^m} \times C_{p^{n-1}}$  if  $n > 1$ ,  $\Phi(G) = Z(G) \cong C_{p^l} \times C_{p^m}$  if  $n = 1$ ,  $G' = \langle a^{p^l}, b^{p^m} \rangle$ .
- (B15)  $\langle a, b, c \mid a^4 = b^4 = c^4 = 1, [b, c] = 1, [c, a] = a^2 c^2, [a, b] = c^2 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_2^3$ ,  $G' = \langle a^2, c^2 \rangle$ .
- (B16)  $\langle a, b, c \mid a^4 = b^4 = c^4 = 1, [b, c] = 1, [c, a] = a^2 = c^2, [a, b] = b^2 \rangle$ ; where  $|G| = 2^5$ ,  $\Phi(G) = Z(G) = G' = \langle a^2, b^2 \rangle \cong C_2^2$ .
- (B17)  $\langle a, b, c; x \mid a^{p^l} = b^p = c^{p^2} = x^p = 1, [a, b] = x, [a, c] = c^p, [b, c] = [x, a] = [x, b] = [x, c] = 1 \rangle$ , where  $l \geq 2$  if  $p = 2$ ; moreover,  $|G| = p^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^p, c^p, x \rangle \cong C_{p^{l-1}} \times C_p \times C_p$  if  $l > 1$ ,  $\Phi(G) = Z(G) \cong C_p \times C_p$  if  $l = 1$ ,  $G' = \langle c^p, x \rangle$ .
- (B18)  $\langle a, b, c; x \mid a^{p^l} = b^{p^m} = c^{p^2} = x^p = 1, [a, b] = c^p, [a, c] = x, [b, c] = [x, a] = [x, b] = [x, c] = 1 \rangle$ , where  $l \geq 2$  if  $p = 2$ ,  $m \leq 2$  and  $\min\{l, m\} = 1$ ; moreover,

$|G| = p^{l+m+3}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p, x \rangle \cong C_{p^{l-1}} \times C_p \times C_p$  if  $m = 1$ ,  
 $\Phi(G) = Z(G) \cong C_{p^{m-1}} \times C_p \times C_p$  if  $l = 1$ ,  $G' = \langle c^p, x \rangle$ .

(B19)  $\langle a, b, c; x \mid a^{p^{l+1}} = b^{p^m} = c^p = x^p = 1, [a, b] = a^{p^l}, [a, c] = x, [b, c] = [x, a] = [x, b] = [x, c] = 1 \rangle$ , where  $m \leq 2$ ; moreover,  $|G| = p^{l+m+3}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, x \rangle \cong C_{p^l} \times C_{p^{m-1}} \times C_p$  if  $m > 1$ ,  $\Phi(G) = Z(G) \cong C_{p^l} \times C_p$  if  $m = 1$ ,  $G' = \langle a^{p^l}, x \rangle$ .

(B20)  $\langle a, b, c; x \mid a^4 = b^2 = c^4 = x^2 = 1, [a, b] = x, [a, c] = a^2 = c^2, [b, c] = [x, a] = [x, b] = [x, c] = 1 \rangle$ ; where  $|G| = 2^5$ ,  $\Phi(G) = Z(G) = G' = \langle a^2, x \rangle \cong C_2^2$ .

Moreover, Table 2 gives  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  for the groups (B1)–(B20).

$(\mu_0, \mu_1, \mu_2)$	$\alpha_1(G)$	types of $\mathcal{A}_3$ groups
$(p+1, p^2-1, 1)$	$2p^2-1$	(B1); (B2) where $m = n = 1$ ; (B3) where $n = 1$ ; (B4); (B5) where $n = 1$
$(p+1, p^2-p, p)$	$p^3+p^2-p$	(B2) where $m > 1 = n$ or $n > 1 = m$ ; (B3) where $n > 1$ ; (B5) where $n > 1$
$(1, p^2-1, p+1)$	$p^3+2p^2-1$	(B6); (B9); (B11); (B13) where $p = 2$ and $m = l = 1$ (B17) where $l = 1$ ; (B19) where $p = 2$ and $m = l = 1$
$(1, p^2, p)$	$p^3+p^2$	(B7) where $l > 1$ ; (B10); (B12) where $l > 1$ ; (B13) where $m = 1$ and $l > 1$ (B14) where $n = 1$ or $m = 1$ ; (B16); (B18) where $m = 1$ (B19) where $p = 2$ and $l > 1 = m$ ; (B19) where $p > 2$ and $m = 1$
$(1, p^2+p-1, 1)$	$2p^2+p-1$	(B7) where $l = 1$ ; (B12) where $l = 1$ ; (B13) where $p > 2$ and $m = l = 1$ ; (B20)
$(1, p^2+p-2, 2)$	$3p^2+p-2$	(B8) where $l = 1$
$(1, p^2-p, 2p)$	$2p^3+p^2-p$	(B8) where $l > 1$ ; (B13) where $m = 2$ ; (B14) where $n = m = 2$ (B15); (B17) where $l > 1$ ; (B18) where $m = 2$ ; (B19) where $m = 2$

Table 2: The enumeration of (B1)–(B20)

**Proof** Let  $A$  be an abelian subgroup of index  $p$  and  $B$  be an  $\mathcal{A}_1$ -subgroup of index  $p$ . By Lemma 2.15,  $|G'| \leq p|A'||B'| = p^2$ . It is easy to see that  $\Phi(B) = \Phi(G)$ . By Lemma 2.2,  $\Phi(B) = Z(B)$ . Since  $[\Phi(G), A] = [\Phi(G), B] = 1$  and  $G = AB$ , we have  $\Phi(G) \leq Z(G)$ . Moreover, it is easy to prove that  $G' \leq C_p^2$ . If  $|G'| = p$ , then  $G$  is one of the groups listed in [2, Theorem 3.1]. Since  $G$  has an  $\mathcal{A}_1$ -subgroup of index  $p$ , they are the groups (B1)–(B5). If  $|G'| = p^2$ , then  $G$  is one of the groups listed in [2, Theorem 4.7]. By checking [2, Table 4] we get the groups (B6)–(B20). The correspondence shows as Table 3.

Groups	Groups in [2, Theorem 4.8]	Groups	Groups in [2, Theorem 4.8]
(B6)	(A1) where $l = m = 1$	(B14)	(A10) where $\min\{l, m, n\} = 1$ and $\max\{m, n\} = 2$
(B7)	(A2) where $m = 1$	(B15)	(A11) where $n = 2$
(B8)	(A3) where $m = 1$ and $-\nu \in (F_p^*)^2$	(B16)	(A12) where $h = 1$
(B9)	(A3) where $l = 1, m = 2$ and $-\nu \notin (F_p^*)^2$	(B17)	(B1) where $m = n = 2 = 1$
(B10)	(A4) where $m = 1$	(B18)	(B2) where $n = 1, m \leq 2$ and $\min\{l, m\} = 1$
(B11)	(A6) where $m = 2$	(B19)	(B3) where $n = 1$ and $m \leq 2$
(B12)	(A8) where $m = 2$ and $n = 1$	(B20)	(B4) where $h = 1$
(B13)	(A9) where $n = 1$ and $m \leq 2$		

Table 3: The correspondence from Theorem 4.2 to [2, Theorem 4.7]

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 4.2 as follows.

By Lemma 3.18,  $\alpha_1(G) = \mu_1 + p^2\mu_2$ . Hence we only need to calculate  $(\mu_0, \mu_1, \mu_2)$ . Since  $d(G) = 3$ ,  $G$  has  $1 + p + p^2$  maximal subgroups. They are respectively

$$M = \langle b, c, \Phi(G) \rangle;$$

$$M_i = \langle ab^i, c, \Phi(G) \rangle, \text{ where } 0 \leq i \leq p-1;$$



$M_{ij} = \langle ac^i, bc^j, \Phi(G) \rangle$ , where  $0 \leq i, j \leq p-1$ .

**Case 1.**  $G' \cong C_p$ . That is,  $G$  is one of the groups (B1)–(B5).

In this case,  $M$  and  $M_i$  are abelian. Notice that  $|M'_{ij}| = p$ . If  $d(M_{ij}) = 2$ , then, by Lemma 2.2,  $M_{ij} \in \mathcal{A}_1$ . Since  $\Phi(M_{ij}) \leq \Phi(G)$ , we get that  $M_{ij} \in \mathcal{A}_1$  if and only if  $\Phi(G) = \Phi(M_{ij})$ .

By calculations,  $\Phi(G) = \langle a^p, b^p, c^p, G' \rangle$  and  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, G' \rangle$ . Since  $\Phi(G)/G' = \langle \bar{a}^p \rangle \times \langle \bar{b}^p \rangle \times \langle \bar{c}^p \rangle$  where  $\bar{a}^p = 1$  or  $\bar{b}^p = 1$ , and  $\langle \bar{c}^p \rangle \cong C_p$ , the following conclusions hold:

- (i) If  $\bar{a}^p = 1$  and  $\bar{b}^p \neq 1$ , then  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$ ;
- (ii) If  $\bar{a}^p \neq 1$  and  $\bar{b}^p = 1$ , then  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$ ;
- (iii) If  $\bar{a}^p = \bar{b}^p = 1$ , then  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$  or  $(j, p) = 1$ .

If  $|G| = p^5$ , then case (iii) happens. In this case,  $M_{00}$  is the unique  $\mathcal{A}_2$ -subgroup of  $G$ . Hence  $(\mu_0, \mu_1, \mu_2) = (p+1, p^2-1, 1)$ .

If  $|G| > p^5$ , then case (i) or (ii) happens. If (i) happens, then  $M_{ij} \in \mathcal{A}_2$  if and only if  $i = 0$ . If (ii) happens, then  $M_{ij} \in \mathcal{A}_2$  if and only if  $j = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (p+1, p^2-p, p)$ .

**Case 2.**  $G' \cong C_p^2$ . That is,  $G$  is one of the groups (B6)–(B20).

In this case,  $M$  is the unique abelian subgroup of index  $p$ . Notice that  $|M'_i| = p$ . If  $d(M_i) = 2$ , then, by Lemma 2.2,  $M_i \in \mathcal{A}_1$ . Since  $\Phi(M_i) \leq \Phi(G)$ , we get that  $M_i \in \mathcal{A}_1$  if and only if  $\Phi(G) = \Phi(M_{ij})$ . Similar argument gives that  $M_{ij} \in \mathcal{A}_1$  if and only if  $\Phi(G) = \Phi(M_{ij})$ .

Subcase 1.  $G$  is the group (B6).

In this case,  $M_0 = \langle a, c \rangle \times \langle b^p \rangle \in \mathcal{A}_2$ . For  $i \neq 0$ ,  $M_i = \langle ab^i, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle b^p, c^p \rangle$  and  $\Phi(M_{ij}) = \langle c^{ip}, b^p c^{jp} \rangle$ . Hence  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$ . In Summary,  $G$  has  $p+1$   $\mathcal{A}_2$ -subgroups. They are  $M_0$  and  $M_{0j}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2-1, p+1)$ .

Subcase 2.  $G$  is the group (B7).

In this case,  $M_i = \langle ab^i, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle a^p, b^p, c^p \rangle$  and  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, b^{-p} b^{-jp} c^{-jp} \rangle = \langle a^p c^{ip}, b^p c^{jp}, b^{-jp} \rangle$ .

If  $l = 1$ , then  $a^p = 1$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$  or  $(j, p) = 1$ . That is,  $G$  has a unique  $\mathcal{A}_2$ -subgroup  $M_{00}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2+p-1, 1)$ .

If  $l > 1$ , then  $a^p \neq 1$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$ . That is,  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 3.  $G$  is the group (B8).

In this case,  $M_i = \langle ab^i, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle a^p, b^p, c^p \rangle$  and  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, b^{-tp} c^{\nu p} b^{-jp} c^{-tjp} \rangle = \langle a^p c^{ip}, b^p c^{jp}, c^{\nu p} b^{-jp} \rangle$ .

If  $l = 1$ , then  $a^p = 1$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$  or  $\begin{vmatrix} 1 & j \\ -j & \nu \end{vmatrix} \neq 0$ . That is,  $G$  has  $2$   $\mathcal{A}_2$ -subgroups  $M_{0j}$  where  $j^2 = -\nu$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 + p - 2, 2)$ .

If  $l > 1$ , then  $a^p \neq 1$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $\begin{vmatrix} 1 & j \\ -j & \nu \end{vmatrix} \neq 0$ . That is,  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_{ij}$  where  $j^2 = -\nu$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

Subcase 4.  $G$  is the group (B9).

In this case,  $M_0 = \langle a, c \rangle * \langle b^p c^{tp} \rangle \in \mathcal{A}_2$ . For  $i \neq 0$ ,  $M_i = \langle ab^i, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle b^p, c^p \rangle$  and  $\Phi(M_{ij}) = \langle c^{ip}, b^p c^{jp}, G' \rangle$ . Hence  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$ . In summary,  $G$  has  $p$   $\mathcal{A}_2$ -subgroups. They are  $M_0$  and  $M_{0j}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - 1, p + 1)$ .

Subcase 5.  $G$  is the group (B10).

In this case,  $M_i = \langle ab^i, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle a^2, b^2, c^2 \rangle$  and  $\Phi(M_{ij}) = \langle a^2 c^{2i} b^{2i}, b^2 c^{2j}, c^2 b^{2j} \rangle$ . Hence  $\Phi(G) = \Phi(M_{ij})$  if and only if  $j = 0$ . In summary,  $G$  has  $p$   $\mathcal{A}_2$ -subgroups. They are  $M_{i1}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 6.  $G$  is the group (B11).

In this case,  $M_0 = \langle a, c \rangle * \langle b^2 \rangle \in \mathcal{A}_2$  and  $M_1 = \langle ab, c \rangle \in \mathcal{A}_1$ . By calculations,  $\Phi(G) = \langle b^2, c^2 \rangle$  and  $\Phi(M_{ij}) = \langle c^{2i}, b^2 c^{2j}, G' \rangle$ . Hence  $\Phi(G) = \Phi(M_{ij})$  if and only if  $i = 1$ . In summary,  $G$  has  $p + 1$   $\mathcal{A}_2$ -subgroups. They are  $M_0$  and  $M_{0j}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - 1, p + 1)$ .

Subcase 7.  $G$  is the group (B12).

By calculations,  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, c^{\nu p} b^{-jp^2} \rangle = \langle a^p, b^p, c^p \rangle = \Phi(G)$ . Hence  $M_{ij} \in \mathcal{A}_1$ . Notice that  $\Phi(M_i) = \langle a^p b^{ip}, c^p, b^{p^2} \rangle$ .

If  $l = 1$ , then  $a^p = 1$ . In this case,  $\Phi(G) = \Phi(M_i)$  if and only if  $(i, p) = 1$ . That is,  $G$  has a unique  $\mathcal{A}_2$ -subgroup  $M_0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 + p - 1, 1)$ .

If  $l > 1$ , then  $a^p \neq 1$  and  $\Phi(G) > \Phi(M_i)$ . Hence  $M_i \in \mathcal{A}_2$  and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 8.  $G$  is the group (B13).

By calculations,  $\Phi(G) = \langle a^p, b^p, c^p \rangle$ . If  $p > 2$ , then  $\Phi(M_i) = \langle a^p b^{ip}, c^p \rangle$ . Hence  $M_i \in \mathcal{A}_2$  for  $m = 2$  and  $M_i \in \mathcal{A}_1$  for  $m = 1$ . If  $p = 2$ , then  $\Phi(M_i) = \langle a^2 b^{2i} a^{i^2}, c^2 \rangle$ . Hence  $M_i \in \mathcal{A}_2$  for  $m = 2$  and  $M_i \in \mathcal{A}_1$  for  $m = 1 \leq l$ . If  $l = m = 1$  and  $p = 2$ , then  $M_0 \in \mathcal{A}_1$  and  $M_1 \in \mathcal{A}_2$ .

If  $p > 2$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, a^{p^l} c^{-jp} \rangle$ . We discuss the value of  $m$  and  $l$ .

If  $m = 2$  and  $l > 1$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, c^{-jp} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$  and  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $m = 2$  and  $l = 1$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, c^{(i+j)p} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i + j, p) = 1$  and  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{ij}$  where

$i + j = p$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $m = 1$  and  $l > 1$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, c^{jp} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$  and  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $l = m = 1$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, c^{jp}, a^p c^{-jp} \rangle = \langle a^p, c^{ip}, c^{jp} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$  or  $(j, p) = 1$ , and  $G$  has a unique  $\mathcal{A}_2$ -subgroup  $M_{00}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 + p - 1, 1)$ .

If  $p = 2$ , then  $\Phi(M_{ij}) = \langle a^2, b^2 c^{2j}, a^{2l} c^{-2j} \rangle = \langle a^2, b^2, c^{-2j} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, 2) = 1$  and  $M_{i0} \in \mathcal{A}_2$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$  for  $m = 2$ ,  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$  for  $m = 1 < l$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - 1, p + 1)$  for  $l = m = 1$ .

Subcase 9.  $G$  is the group (B14).

By calculations,  $\Phi(G) = \langle a^p, b^p, c^p \rangle$ . If  $p > 2$ , then  $\Phi(M_i) = \langle a^p b^{ip}, c^p, b^{p^m} \rangle$ . Hence  $M_i \in \mathcal{A}_2$  for  $m = 2$  and  $M_i \in \mathcal{A}_1$  for  $m = 1$ . If  $p = 2$ , then  $\Phi(M_i) = \langle a^2 b^{2i} a^{i2^l}, c^2, b^{2^m} \rangle$ . Hence  $M_i \in \mathcal{A}_2$  for  $m = 2$  and  $M_i \in \mathcal{A}_1$  for  $m = 1 \leq l$ . If  $l = m = 1$  and  $p = 2$ , then  $M_0 \in \mathcal{A}_1$  and  $M_1 \in \mathcal{A}_2$ .

If  $p > 2$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, a^{p^l} b^{-jp^m} \rangle$ . We discuss the value of  $m$  and  $l$ .

If  $n = 1$ , then  $\Phi(M_{ij}) = \langle a^p, b^p \rangle = \Phi(G)$  and hence  $\Phi(M_{ij}) \in \mathcal{A}_1$ . Since  $\max\{m, n\} = 2$ ,  $m = 2$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $m = n = 2$ , then  $l = 1$  since  $\min\{l, m, n\} = 1$ . Notice that  $\Phi(M_{ij}) = \langle c^{ip}, b^p c^{jp}, a^p \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, p) = 1$  and  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{0j}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $n = 2$  and  $l = m = 1$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, a^p b^{-jp} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $j^2 - i \neq 0$  and  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{ij}$  where  $i = j^2$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $n = 2$  and  $m = 1 < l$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, b^p c^{jp}, b^{-jp} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$  and  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $p = 2$ , then  $\Phi(M_{ij}) = \langle a^2 c^{2i} b^{i2^m}, b^2 c^{2j}, a^{2^l} b^{-j2^m} \rangle$ . We discuss the value of  $m$  and  $l$ .

If  $n = 1$ , then  $\Phi(M_{ij}) = \langle a^2, b^2 \rangle = \Phi(G)$  and hence  $\Phi(M_{ij}) \in \mathcal{A}_1$ . Since  $\max\{m, n\} = 2$ ,  $m = 2$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $m = n = 2$ , then  $l = 1$  since  $\min\{l, m, n\} = 1$ . Notice that  $\Phi(M_{ij}) = \langle c^{2i}, b^2 c^{2j}, a^2 \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(i, 2) = 1$  and  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{0j}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $n = 2$  and  $l = m = 1$ , then  $\Phi(M_{ij}) = \langle a^2 c^{2i} b^{2i}, b^2 c^{2j}, a^2 b^{2j} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $i = 1$  or  $j = 1$ , and  $G$  has 2  $\mathcal{A}_2$ -subgroups  $M_1$  and  $M_{00}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $n = 2$  and  $m = 1 < l$ , then  $\Phi(M_{ij}) = \langle a^2 c^{2i} b^{2i}, b^2 c^{2j}, b^{-2j} \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, 2) = 1$  and  $G$  has 2  $\mathcal{A}_2$ -subgroups  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 10.  $G$  is the group (B15).

By calculations,  $\Phi(G) = \langle a^2, b^2, c^2 \rangle$ ,  $M_0 = \langle c, a \rangle \times \langle b^2 \rangle \in \mathcal{A}_2$  and  $M_1 = \langle c, ab \rangle \in \mathcal{A}_1$ . Since  $\Phi(M_{ij}) = \langle a^{2(1+i)}, b^2 c^{2j}, c^{2(1+j)} a^{2j} \rangle$ ,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $i = j = 0$ . Hence  $G$  has 4  $\mathcal{A}_2$ -subgroups  $M_0$  and  $M_{ij}$  where  $i \neq 0$  or  $j \neq 0$ , and  $(\mu_0, \mu_1, \mu_2) = (1, 2, 4)$ .

Subcase 11.  $G$  is the group (B16).

By calculations,  $\Phi(G) = \langle a^2, b^2 \rangle$ ,  $M_i = \langle c, ab^i \rangle \times \langle b^2 \rangle \in \mathcal{A}_2$ . Since  $\Phi(M_{ij}) = \langle a^2 = c^2, b^2 c^{2j}, b^2 a^{2j} \rangle = \Phi(G)$ ,  $M_{ij} \in \mathcal{A}_1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 12.  $G$  is the group (B17).

By calculations,  $\Phi(G) = \langle a^p, c^p, x \rangle$  and  $M_i = \langle ab^i, c \rangle \times \langle x \rangle \in \mathcal{A}_2$ .

If  $p > 2$ , then  $\Phi(M_{ij}) = \langle a^p c^{ip}, c^{jp}, x c^{-jp} \rangle = \langle a^p c^{ip}, c^{jp}, x \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, p) = 1$  and  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $p = 2$ , then  $\Phi(M_{ij}) = \langle a^2, c^{2j}, x c^{2j} \rangle = \langle a^2, c^{2j}, x \rangle$ . In this case,  $\Phi(G) = \Phi(M_{ij})$  if and only if  $(j, 2) = 1$  and  $G$  has 4  $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{i0}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

Subcase 13.  $G$  is the group (B18).

If  $l = 1$  and  $m = 2$ , then  $p > 2$  and  $\Phi(G) = \langle b^p, c^p, x \rangle$ . In addition,  $\Phi(M_i) = \langle b^{ip}, c^p, x \rangle$  and  $\Phi(M_{ij}) = \langle c^{ip}, b^p c^{jp}, c^p x^{-j} \rangle$ . In this case,  $\Phi(M_i) = \Phi(G)$  if and only if  $(i, p) = 1$ , and  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(i, p) = 1$  and  $(j, p) = 1$ . Hence  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_0, M_{0j}$  and  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $l = m = 1$ , then  $p > 2$  and  $\Phi(G) = \langle c^p, x \rangle$ . Since  $\Phi(M_i) = \langle c^p, x \rangle = \Phi(G)$ ,  $M_i \in \mathcal{A}_1$ . Since  $\Phi(M_{ij}) = \langle c^{ip}, c^{jp}, c^p x^{-j} \rangle$ ,  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(j, p) = 1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

If  $l > 1$ , then  $m = 1$  since  $\min\{l, m\} = 1$ . In this case,  $\Phi(G) = \langle a^p, c^p, x \rangle$ . In addition, if  $p > 2$ , then  $\Phi(M_i) = \langle a^p, c^p, x \rangle$  and  $\Phi(M_{ij}) = \langle a^p c^{ip}, c^{jp}, c^p x^{-j} \rangle$ . In this case,  $M_i \in \mathcal{A}_1$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(j, p) = 1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$  and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ ; if  $p = 2$ , then  $\Phi(M_i) = \langle a^2 c^{2i}, c^2, x \rangle$  and  $\Phi(M_{ij}) = \langle a^2 c^{2i} x^i, c^{2j}, c^2 x^j \rangle$ . In this case,  $M_i \in \mathcal{A}_1$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(j, 2) = 1$ . Hence  $G$  has 2  $\mathcal{A}_2$ -subgroups  $M_{i0}$  and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 14.  $G$  is the group (B19).

If  $m = 2$ , then  $\Phi(G) = \langle a^p, b^p, x \rangle$ . In addition, if  $p > 2$ , then  $\Phi(M_i) = \langle a^p b^{ip}, x \rangle$  and  $\Phi(M_{ij}) = \langle a^p, b^p, a^{p^l} x^{-j} \rangle$ . In this case,  $\Phi(M_i) \in \mathcal{A}_2$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(j, p) = 1$ . Hence  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ ; if  $p = 2$ , then  $\Phi(M_i) = \langle a^2 b^{2i} a^{i^{2^l}}, x \rangle$  and  $\Phi(M_{ij}) = \langle a^2 x^i, b^2, a^{2^l} x^{-j} \rangle$ . In this case,  $\Phi(M_i) \in \mathcal{A}_2$ .  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $(j, 2) = 1$  for  $l > 1$ . Hence  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{i0}$  for  $l > 1$ .  $\Phi(M_{ij}) = \Phi(G)$  if and only if  $i \neq j$  for  $l = 1$ .

Hence  $G$  has  $2p$   $\mathcal{A}_2$ -subgroups  $M_i$  and  $M_{ij}$  where  $i \neq j$  for  $l > 1$ . In summary,  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - p, 2p)$ .

If  $l = m = 1$ , then  $\Phi(G) = \langle a^p, x \rangle$ . In addition, if  $p > 2$ , then  $\Phi(M_i) = \langle a^p, x \rangle$  and  $\Phi(M_{ij}) = \langle a^p, a^p x^{-j} \rangle$ . In this case,  $\Phi(M_i) \in \mathcal{A}_1$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only  $(j, p) = 1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ ; if  $p = 2$ , then  $\Phi(M_i) = \langle a^2 a^{i^2}, x \rangle$  and  $\Phi(M_{ij}) = \langle a^2 x^i, a^2 x^{-j} \rangle$ . In this case,  $\Phi(M_0) \in \mathcal{A}_1$ ,  $\Phi(M_1) \in \mathcal{A}_2$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only  $i \neq j$ . Hence  $G$  has  $p + 1$   $\mathcal{A}_2$ -subgroups  $M_1$  and  $M_{ij}$  where  $i \neq j$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2 - 1, p + 1)$ .

If  $l > 1 = m$ , then  $\Phi(G) = \langle a^p, x \rangle$ . In addition, if  $p > 2$ , then  $\Phi(M_i) = \langle a^p, x \rangle$  and  $\Phi(M_{ij}) = \langle a^p, a^{p^l} x^{-j} \rangle$ . In this case,  $\Phi(M_i) \in \mathcal{A}_1$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only  $(j, p) = 1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ ; if  $p = 2$ , then  $\Phi(M_i) = \langle a^2 a^{i2^l}, x \rangle$  and  $\Phi(M_{ij}) = \langle a^2 x^i, a^{2^l} x^{-j} \rangle$ . In this case,  $\Phi(M_i) \in \mathcal{A}_1$  and  $\Phi(M_{ij}) = \Phi(G)$  if and only  $(j, 2) = 1$ . Hence  $G$  has  $p$   $\mathcal{A}_2$ -subgroups  $M_{i0}$ , and  $(\mu_0, \mu_1, \mu_2) = (1, p^2, p)$ .

Subcase 15.  $G$  is the group (B20).

By calculations,  $\Phi(G) = \langle a^2 = c^2, x \rangle$ ,  $M_0 = \langle a, c \rangle \times \langle x \rangle \in \mathcal{A}_2$ , and  $M_1 = \langle ab, c \rangle \in \mathcal{A}_1$ . Since  $\Phi(M_{ij}) = \langle a^2, c^{2j}, xc^{2j} \rangle = \langle a^2, x \rangle = \Phi(G)$ ,  $M_{ij} \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, p^2 + p - 1, 1)$ .

To sum up, we get the Table 3. □

**Lemma 4.3.** *Suppose that  $G$  is a finite  $p$ -group such that  $\Phi(G') = 1$ ,  $C_p^2 \cong G_3 \leq Z(G)$  and  $G/G_3 \cong M_p(1, 1, 1)$ , where  $p \geq 3$ . Let  $G = \langle a, b, c, x, y \rangle$  such that*

$$a^p = x^{w_{11}} y^{w_{12}}, b^p = x^{w_{21}} y^{w_{22}}, c^p = x^p = y^p = 1, [a, b] = c, [c, a] = x, [c, b] = y,$$

$N = \langle c, b, x, y \rangle$  and  $M_i = \langle ab^i, c, x, y \rangle$ . Then  $N$  and  $M_i$  are all maximal subgroups of  $G$ , and the following conclusions hold:

- (1)  $N \in \mathcal{A}_2$  if and only if  $w_{21} = 0$ ;
- (2) If  $p > 3$ , then  $M_i \in \mathcal{A}_2$  if and only if  $w_{12} + i(w_{22} - w_{11}) - i^2 w_{21} = 0$ ;
- (3) If  $p = 3$ , then  $M_i \in \mathcal{A}_2$  if and only if  $w_{12} + i(w_{22} - w_{11}) - i^2 w_{21} - i^2 = 0$ ;
- (4)  $\alpha_1(G) = \mu_1 + \mu_2 p^2$ .

**Proof** If  $w_{21} = 0$ , then  $N = \langle c, b \rangle \times \langle x \rangle \in \mathcal{A}_2$ . If  $w_{21} \neq 0$ , then  $N = \langle c, b \rangle \in \mathcal{A}_1$ . Hence (1) holds.

It is easy to see that  $M_i \in \mathcal{A}_2$  if and only if  $\langle (ab^i)^p \rangle = \langle [c, ab^i] \rangle = \langle xy^i \rangle$ .

If  $p > 3$ , then  $(ab^i)^p = a^p b^{ip} = x^{w_{11} + iw_{21}} y^{w_{12} + iw_{22}}$ . Hence  $M_i \in \mathcal{A}_2$  if and only if  $w_{12} + iw_{22} = i(w_{11} + iw_{21})$ . The later is  $w_{12} + i(w_{22} - w_{11}) - i^2 w_{21} = 0$ . Hence (2) holds.

If  $p = 3$ , then  $(ab^i)^3 = a^3 b^{3i} [a, b^{-i}, a][a, b^{-i}, b^{-i}] = x^{w_{11} + iw_{21} - i} y^{w_{12} + iw_{22} + i^2}$ . Hence  $M_i \in \mathcal{A}_2$  if and only if  $w_{12} + iw_{22} + i^2 = i(w_{11} + iw_{21} - i)$ . The later is  $w_{12} + i(w_{22} - w_{11}) - i^2 w_{21} - i^2 = 0$ . Hence (3) holds.

Let  $M \in \Gamma_1$ . Then, by above argument,  $|M'| = p$ . If  $M \in \mathcal{A}_2$ , then, by Lemma 2.6 (7),  $\alpha_1(M) = p^2$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p^2\mu_2$ . Hence (4) holds.  $\square$

**Theorem 4.4.**  *$G$  has no abelian subgroup of index  $p$ ,  $G$  has at least two distinct  $\mathcal{A}_1$ -subgroups of index  $p$  and  $d(G) = 2$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Ci)  $c(G) = 3$  and  $G' \cong C_2^2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 2, 1)$  and  $\alpha_1(G) = 6$ .

(C1)  $\langle a, b, c, d \mid a^4 = b^2 = c^2 = d^2 = 1, [a, b] = c, [c, a] = d, [c, b] = [d, a] = [d, b] = 1 \rangle$ ; where  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c, d \rangle \cong C_2^3$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle d \rangle \cong C_2$ .

(C2)  $\langle a, b, c \mid a^8 = b^2 = c^2 = 1, [a, b] = c, [c, a] = a^4, [c, b] = 1 \rangle$ ; where  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_4 \times C_2$ ,  $G' = \langle c, a^4 \rangle$  and  $Z(G) = \langle a^4 \rangle \cong C_2$ .

(C3)  $\langle a, b, c \mid a^8 = c^2 = 1, b^2 = a^4, [a, b] = c, [c, a] = [a^2, b] = b^2, [c, b] = 1 \rangle$ ; where  $|G| = 2^5$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_4 \times C_2$ ,  $G' = \langle c, a^4 \rangle$  and  $Z(G) = \langle a^4 \rangle \cong C_2$ .

(C4)  $\langle a, b, c \mid a^{2^{n+1}} = b^2 = c^2 = 1, [a, b] = c, [c, a] = a^{2^n}, [c, b] = 1 \rangle$ , where  $n \geq 3$ ; moreover,  $|G| = 2^{n+3}$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_{2^n} \times C_2$ ,  $G' = \langle c, a^{2^n} \rangle$  and  $Z(G) = \langle a^4 \rangle \cong C_{2^{n-1}}$ .

(C5)  $\langle a, b, c \mid a^{2^n} = b^4 = c^2 = 1, [a, b] = c, [c, a] = b^2, [c, b] = 1 \rangle$ , where  $n \geq 3$ ; moreover,  $|G| = 2^{n+3}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^{n-1}} \times C_2 \times C_2$ ,  $G' = \langle c, b^2 \rangle$  and  $Z(G) = \langle a^4, b^2 \rangle \cong C_{2^{n-2}} \times C_2$ .

(C6)  $\langle a, b, c, d \mid a^{2^n} = b^2 = c^2 = d^2 = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle$ , where  $n \geq 3$ ; moreover,  $|G| = 2^{n+3}$ ,  $\Phi(G) = \langle a^2, c, d \rangle \cong C_{2^{n-1}} \times C_2 \times C_2$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle a^4, d \rangle \cong C_{2^{n-2}} \times C_2$ .

(Cii)  $\Phi(G') \leq G_3 \cong C_p^2$ .

(C7)  $\langle a, b, c, d, e \mid a^3 = b^3 = c^3 = d^3 = e^3 = 1, [a, b] = c, [c, a] = d, [c, b] = e, [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle$ , where  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle c, d, e \rangle$  and  $Z(G) = \langle d, e \rangle \cong C_3^2$ .

(C8)  $\langle a, b, c, d \mid a^9 = c^3 = d^3 = 1, b^3 = a^3, [a, b] = c, [c, a] = d, [c, b] = a^3, [d, a] = [d, b] = 1 \rangle$ ; where  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle a^3, c, d \rangle$ ,  $Z(G) = \langle a^3, d \rangle \cong C_3^2$ .

(C9)  $\langle a, b, c, d \mid a^9 = b^3 = c^3 = d^3 = 1, [a, b] = c, [c, a] = d, [c, b] = a^{-3}, [d, a] = [d, b] = 1 \rangle$ ; where  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle a^3, c, d \rangle$  and  $Z(G) = \langle a^3, d \rangle \cong C_3^2$ .

(C10)  $\langle a, b, c \mid a^9 = b^9 = c^3 = 1, [a, b] = c, [c, a] = a^3, [c, b] = b^3 \rangle$ ; where  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle a^3, b^3, c \rangle$  and  $Z(G) = \langle a^3, b^3 \rangle \cong C_3^2$ .

(C11)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = a^p b^{\nu p}, [c, b] = b^p \rangle$ , where  $p > 3$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_p^2$ .

- (C12)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = a^{-p}, [a^p, b] = 1 \rangle$ , where  $p > 3$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  such that  $-\nu \in F_p^2$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_p^2$ .
- (C13)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a]^{1+r} = a^p b^p, [c, b]^{1+r} = a^{-r p} b^p, [a^p, b] = 1 \rangle$ , where  $p > 3$ ,  $r \neq 0, -1$  and  $-r \in (F_p^*)^2$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_p^2$ .
- (C14)  $\langle a, b, c, d \mid a^{p^2} = b^p = c^p = d^p = 1, [a, b] = c, [c, a] = a^p, [c, b] = d \rangle$ , where  $p > 3$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a^p, c, d \rangle$  and  $Z(G) = \langle a^p, d \rangle \cong C_p^2$ .
- (C15)  $\langle a, b, c, d \mid a^p = b^{p^2} = c^p = d^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = d, [d, a] = [d, b] = 1 \rangle$ , where  $p > 3$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle b^p, c, d \rangle$  and  $Z(G) = \langle b^p, d \rangle \cong C_p^2$ .
- (C16)  $\langle a, b, c \mid a^8 = c^4 = 1, b^2 = a^4, [a, b] = c, [c, a] = a^4, [c, b] = c^2 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, c \rangle \cong C_4^2$ ,  $G' = \langle c, a^4 \rangle$  and  $Z(G) = \langle a^4, c^2 \rangle \cong C_2^2$ .
- (C17)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^{\nu_1 p}, [c, b] = a^{-\nu_2 p^2} \rangle$ , where  $p \geq 3$ , and  $\nu_1, \nu_2 = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p \times C_p$ ,  $G' = \langle c, a^{p^2}, b^p \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_p$ .

Moreover, Table 4 gives  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  for the groups (C7)–(C17).

$(\mu_0, \mu_1, \mu_2)$	$\alpha_1(G)$	types of $\mathcal{A}_3$ groups
$(0, p-1, 2)$	$2p^2 + p - 1$	(C7); (C10); (C12)–(C14)
$(0, p, 1)$	$p^2 + p$	(C8)–(C9); (C11); (C15); (C17)
$(0, p, 1)$	$p^2 + 2p$	(C16)

Table 4: The enumeration of (C7)–(C17)

**Proof** Let  $A$  and  $B$  be two distinct  $\mathcal{A}_1$ -subgroups of index  $p$ . Then  $|A'| = |B'| = p$  and hence  $A'B' \leq Z(G)$ . Let  $\bar{G} = G/A'B'$ . Then  $\bar{G}$  has two abelian subgroups  $\bar{A}$  and  $\bar{B}$  of index  $p$ . By Lemma 2.7,  $\bar{G}$  has  $1 + p$  abelian subgroups. Since  $d(\bar{G}) = 2$ ,  $\bar{G}$  is an  $\mathcal{A}_1$ -group. It is easy to see that  $A'B' = \Phi(G')G_3$ . If  $\bar{G}$  is metacyclic, then  $G$  is also metacyclic by Lemma 2.1 and hence  $A' = B'$ . It follows that  $|G'| = p^2$ . by Lemma 3.1,  $G$  is an  $\mathcal{A}_2$ -group. This contradicts that  $G$  is an  $\mathcal{A}_3$ -group. Thus  $\bar{G}$  is a non-metacyclic  $\mathcal{A}_1$ -group.

If  $G' \cong C_{p^2}$ , then  $G_3 \leq \Phi(G') \cong C_p$ . Thus  $G$  is one of the groups listed in [3, Theorem 3.5]. By [3, Theorem 3.6] we get no group.

If  $G' \cong C_p^2$ , then  $G$  is one of the groups listed in [3, Theorem 4.6]. By [3, Theorem 4.7] we get groups (C1)–(C6).

If  $G_3 \cong C_p$  and  $\Phi(G')G_3 \cong C_p^2$ , then, by [3, Theorem 5.1, 5.2, 5.5 & 5.6], we get no group.

If  $\Phi(G') \leq G_3 \cong C_p^2$ , then  $G$  is one of the groups listed in [3, Theorem 6.5]. By [3, Theorem 6.1], we get the groups (C7)–(C17).

Table 5 gives the correspondence from Theorem 4.4 to [3, Theorem 4.6 & 6.5].

Groups	Groups in [3, Theorem 4.6]	Groups	Groups in [3, Theorem 6.5]
(C1)	(H1)	(C7)	(O1)
(C2)	(H2)	(C8)	(O3)
(C3)	(H3)	(C9)	(O4)
(C4)	(I1)	(C10)	(O5)
(C5)	(I2)	(C11)	(P2) where $n = 1$
(C6)	(I3)	(C12)	(P3) where $n = 1$ and $-\nu \in F_p^2$
		(C13)	(P4) where $n = 1$ and $-r \in (F_p^*)^2$
		(C14)	(P8) where $n = 1$
		(C15)	(P9) where $n = 1$
		(C16)	(R4)
		(C17)	(S2) where $n = 2$

Table 5: The correspondence from Theorem 4.4 to [3, Theorem 4.6 & 6.5]

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 4.4 as follows.

Since  $d(G) = 2$ ,  $G$  has  $1 + p$  maximal subgroups. For any  $\mathcal{A}_3$ -group, we always have  $\mu_2 \geq 1$ . By the hypothesis of Theorem 4.4,  $\mu_0 = 0$  and  $\mu_1 \geq 2$ .

**Case 1.**  $c(G) = 3$  and  $G' \cong C_2^2$ . That is,  $G$  is one of the groups (C1)–(C6).

Obviously,  $(\mu_0, \mu_1, \mu_2) = (0, 2, 1)$ . Let  $M \in \Gamma_1$ . If  $M$  is the unique  $\mathcal{A}_2$ -subgroup, then  $|M'| = p$ . By Lemma 2.6,  $\alpha_1(M) = p^2 = 4$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + 4\mu_2 = 6$ .

**Case 2.**  $G$  is one of the groups (C7)–(C15).

In this case,  $p \geq 3$ ,  $\Phi(G') = 1$ ,  $G_3 \cong C_p^2$ ,  $G_3 \leq Z(G)$  and  $G/G_3 \cong M_p(1, 1, 1)$ .  $G = \langle a, b, c, x, y \rangle$  such that

$$a^p = x^{w_{11}}y^{w_{12}}, b^p = x^{w_{21}}y^{w_{22}}, c^p = x^p = y^p = 1, [a, b] = c, [c, a] = x, [c, b] = y.$$

Let  $N = \langle c, b, x, y \rangle$  and  $M_i = \langle ab^i, c, x, y \rangle$ . Then  $N$  and  $M_i$  are all maximal subgroups of  $G$ . Let  $w(G) = \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix}$ .

If  $G$  is the group (C7), then  $p = 3$  and  $w(G) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p - 1, 2)$  and  $\alpha_1(G) = 2p^2 + p - 1$ .

If  $G$  is the group (C8), then  $p = 3$  and  $w(G) = \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p, 1)$  and  $\alpha_1(G) = p^2 + p$ .

If  $G$  is the group (C9), then  $p = 3$  and  $w(G) = \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p, 1)$  and  $\alpha_1(G) = p^2 + p$ .



If  $G$  is the group (C10), then  $p = 3$  and  $w(G) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p - 1, 2)$  and  $\alpha_1(G) = 2p^2 + p - 1$ .

If  $G$  is the group (C11), then  $p > 3$  and  $w(G) = \begin{pmatrix} 1 & -\nu \\ 0 & 1 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p, 1)$  and  $\alpha_1(G) = p^2 + p$ .

If  $G$  is the group (C12), then  $p > 3$  and  $w(G) = \begin{pmatrix} 0 & -1 \\ \nu^{-1} & 0 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_1$  and  $M_i \in \mathcal{A}_2$  if and only if  $i^2 = -\nu$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p - 1, 2)$  and  $\alpha_1(G) = 2p^2 + p - 1$ .

If  $G$  is the group (C13), then  $p > 3$  and  $w(G) = \begin{pmatrix} 1 & -1 \\ r & 1 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_1$  and  $M_i \in \mathcal{A}_2$  if and only if  $i^2 = -r^{-1}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p - 1, 2)$  and  $\alpha_1(G) = 2p^2 + p - 1$ .

If  $G$  is the group (C14), then  $p > 3$  and  $w(G) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p - 1, 2)$  and  $\alpha_1(G) = 2p^2 + p - 1$ .

If  $G$  is the group (C15), then  $p > 3$  and  $w(G) = \begin{pmatrix} 0 & 0 \\ \nu^{-1} & 0 \end{pmatrix}$ . By Lemma 4.3,  $N \in \mathcal{A}_1$  and  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p, 1)$  and  $\alpha_1(G) = p^2 + p$ .

**Case 3.**  $G$  is the group (C16).

It is obvious that  $(\mu_0, \mu_1, \mu_2) = (0, 2, 1) = (0, p, 1)$ . The unique  $\mathcal{A}_2$ -subgroup of index  $p$  is  $M = \langle c, b, a^2 \rangle$ , where  $d(M) = 3$  and  $|M'| = 4$ . Notice that  $\alpha_1(M) = p^2 + p = 6$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + (p^2 + p)\mu_2 = p^2 + 2p = 8.$$

**Case 4.**  $G$  is the group (C17).

Let  $N = \langle c, b, a^p, b^p \rangle$  and  $M_i = \langle c, ab^i, a^p, b^p \rangle$ . Then  $N$  and  $M_i$  are all maximal subgroups of  $G$ . Since  $N = \langle c, b \rangle * \langle a^p \rangle$ ,  $N \in \mathcal{A}_2$  with  $d(N) = 3$  and  $|N'| = p$ . By Lemma 2.6 (7),  $\alpha_1(N) = p^2$ . Since  $M_i = \langle c, ab^i \rangle$  and  $|M'_i| = p$ ,  $M_i \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p, 1)$  and, by Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p^2\mu_2 = p^2 + p.$$

To sum up, we get the Table 4. □

**Lemma 4.5.** *Suppose that  $G$  is an  $\mathcal{A}_3$ -group with  $d(G) = 3$ ,  $\Phi(G) = Z(G)$  and  $G' \cong C_p^3$ . Then we may assume that  $G = \langle a_1, a_2, a_3; x, y, z \rangle$  with*

$$x = [a_2, a_3], y = [a_3, a_1], z = [a_1, a_2], x^p = y^p = z^p = 1, a_i^{p^{m_i}} = x^{w_{i1}} y^{w_{i2}} z^{w_{i3}}, i = 1, 2, 3.$$

Let  $M = \langle a_2, a_3, \Phi(G) \rangle$ ,  $M_i = \langle a_1 a_2^i, a_3, \Phi(G) \rangle$  and  $M_{ij} = \langle a_1 a_3^i, a_2 a_3^j, \Phi(G) \rangle$ , where  $0 \leq i, j \leq p-1$ . Then  $M$ ,  $M_i$  and  $M_{ij}$  are all maximal subgroups of  $G$ , and the following conclusions hold:

(1) If  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $a_1^{p^2} = x$ , then  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  if and only if  $w_{33} = 0$ , and  $M_{ij} \in \mathcal{A}_2$  if and only if  $w_{22} + j(w_{23} + w_{32}) + j^2 w_{33} = 0$ ;

(2) If  $m_1 = m_2 = 2$ ,  $m_3 = 1$ ,  $a_3^p = z$  and  $w_{13} = w_{23} = 0$ , then  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$ , and  $M_{ij} \in \mathcal{A}_2$  if and only if  $w_{11} w_{22} - w_{12} w_{21} = 0$ ;

(3) If  $m_1 = m_2 = m_3 = 1$  and  $p > 2$ , then  $M \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{vmatrix} = 0$ ,  $M_i \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} i & -1 & 0 \\ w_{11} + i w_{21} & w_{12} + i w_{22} & w_{13} + i w_{23} \\ w_{31} & w_{32} & w_{33} \end{vmatrix} = 0$ , and  $M_{ij} \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} -i & -j & 1 \\ w_{11} + i w_{31} & w_{12} + i w_{32} & w_{13} + i w_{33} \\ w_{21} + j w_{31} & w_{22} + j w_{32} & w_{23} + j w_{33} \end{vmatrix} = 0$ ;

(4) If  $m_1 = m_2 = m_3 = 1$  and  $p = 2$ , then  $M \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{vmatrix} = 0$ ,  $M_i \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} i & 1 & 0 \\ w_{11} + i w_{21} & w_{12} + i w_{22} & w_{13} + i w_{23} + i \\ w_{31} & w_{32} & w_{33} \end{vmatrix} = 0$ , and  $M_{ij} \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} i & j & 1 \\ w_{11} + i w_{31} & w_{12} + i w_{32} + i & w_{13} + i w_{33} \\ w_{21} + j w_{31} + j & w_{22} + j w_{32} & w_{23} + j w_{33} \end{vmatrix} = 0$ .

**Proof** Since  $G \in \mathcal{A}_3$ , any maximal subgroup is either  $\mathcal{A}_1$ -group or  $\mathcal{A}_2$ -group. Let  $H \in \Gamma_1(G)$ . Then  $|H'| = p$ . If  $d(H) = 2$ , then, by Lemma 2.2,  $H \in \mathcal{A}_1$ . Since  $\Phi(H) \leq \Phi(G)$ , we get that  $H \in \mathcal{A}_1$  if and only if  $\Phi(G) = \Phi(H)$ .

(1) If  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $a_1^{p^2} = x$ , then it is obvious that  $M = \langle a_2, a_3 \rangle * \langle a_1 \rangle \in \mathcal{A}_2$ . Since  $x = (a_1 a_2^i)^{p^2} \in \Phi(M_i)$  and  $y = [a_3, a_1 a_2^i] x^i \in \Phi(M_i)$ ,  $\Phi(M_i) = \Phi(G)$  if and only if  $a_3^p \notin \langle x, y \rangle$ . Hence  $M_i \in \mathcal{A}_2$  if and only if  $w_{33} = 0$ . Similar reason gives that  $M_{ij} \in \mathcal{A}_1$  if and only if  $(a_2 a_3^j)^p \notin \langle x, [a_1 a_3^i, a_2 a_3^j] \rangle = \langle x, y^{-j} z \rangle$ . Hence  $M_{ij} \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} 1 & 0 & 0 \\ 0 & -j & 1 \\ * & w_{22} + j w_{32} & w_{23} + j w_{33} \end{vmatrix} = 0$ . The later is,  $w_{22} + j(w_{23} + w_{32}) + j^2 w_{33} = 0$ .

(2) If  $m_1 = m_2 = 2$ ,  $m_3 = 1$ ,  $a_3^p = z$  and  $w_{13} = w_{23} = 0$ , then it is obvious that  $a_1^p \notin \langle a_2, a_3 \rangle$  and  $\langle a_1^p, a_2^p \rangle \not\leq \langle a_1 a_2^i, a_3 \rangle$ . Hence  $M \in \mathcal{A}_2$  and  $M_i \in \mathcal{A}_2$ . Since  $(a_1 a_3^i)^{p^2} = a_1^{p^2} = x^{w_{11}} y^{w_{12}}$  and  $(a_2 a_3^j)^{p^2} = a_2^{p^2} = x^{w_{21}} y^{w_{22}}$  and  $[a_1 a_2^i, a_2 a_3^j] \notin \langle x, y \rangle$   $M_{ij} \in \mathcal{A}_1$  if and only if  $\begin{vmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{vmatrix} = 0$ . The later is,  $w_{11} w_{22} - w_{12} w_{21} = 0$ .

(3) If  $m_1 = m_2 = m_3 = 1$  and  $p > 2$ , then  $M \in \mathcal{A}_1$  if and only if  $\langle a_2^p, a_3^p, [a_2, a_3] \rangle = G'$ . The later is  $\begin{vmatrix} w_{21} & w_{22} & w_{23} \\ w_{31} & w_{32} & w_{33} \\ 1 & 0 & 0 \end{vmatrix} \neq 0$ . Hence  $M \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{vmatrix} = 0$ .

Similar arguments give that  $M_i \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} i & -1 & 0 \\ w_{11} + i w_{21} & w_{12} + i w_{22} & w_{13} + i w_{23} \\ w_{31} & w_{32} & w_{33} \end{vmatrix} = 0$ , and  $M_{ij} \in \mathcal{A}_2$  if and only if  $\begin{vmatrix} -i & -j & 1 \\ w_{11} + i w_{31} & w_{12} + i w_{32} & w_{13} + i w_{33} \\ w_{21} + j w_{31} & w_{22} + j w_{32} & w_{23} + j w_{33} \end{vmatrix} = 0$ .

(4) It follows from similar arguments as (3). The details are omitted.  $\square$

**Theorem 4.6.**  *$G$  has no abelian subgroup of index  $p$ ,  $G$  has at least two distinct  $\mathcal{A}_1$ -subgroups of index  $p$  and  $d(G) = 3$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Di) *The type of  $G/G'$  is  $(p^2, p, p)$ .*

- (D1)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [b, c] = a^{p^2}, [c, a] = c^{-p}, [a, b] = b^p c^{\nu p} \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_p \times C_p$  and  $G' = \langle a^{p^2}, b^p, c^p \rangle \cong C_p^3$ .
- (D2)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [b, c] = a^{p^2}, [c, a] = b^p, [a, b] = c^{\nu p} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_p \times C_p$  and  $G' = \langle a^{p^2}, b^p, c^p \rangle \cong C_p^3$ .
- (D3)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [b, c] = a^{p^2}, [c, a]^{1+r} = b^{r p} c^{-p}, [a, b]^{1+r} = b^p c^p \rangle$ , where  $p > 2$ ,  $r = 1, 2, \dots, p-2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_p \times C_p$  and  $G' = \langle a^{p^2}, b^p, c^p \rangle \cong C_p^3$ .
- (D4)  $\langle a, b, c \mid a^8 = b^4 = c^4 = 1, [b, c] = a^4, [c, a] = b^2, [a, b] = c^2 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_4 \times C_2 \times C_2$  and  $G' = \langle a^4, b^2, c^2 \rangle \cong C_2^3$ .
- (D5)  $\langle a, b, c \mid a^8 = b^4 = c^4 = 1, [b, c] = a^4, [c, a] = b^2, [a, b] = b^2 c^2 \rangle$ , where  $|G| = 2^7$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_4 \times C_2 \times C_2$  and  $G' = \langle a^4, b^2, c^2 \rangle \cong C_2^3$ .

(Dii) *The type of  $G/G'$  is  $(p^2, p^2, p)$ .*

- (D6)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = c^{p^2} = 1, [b, c] = a^{p^2}, [c, a] = b^{\nu p^2}, [a, b] = c^p \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  such that  $-\nu \notin (F_p)^2$ ; moreover,  $|G| = p^8$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_{p^2} \times C_p$  and  $G' = \langle a^{p^2}, b^{p^2}, c^p \rangle \cong C_p^3$ .
- (D7)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = c^{p^2} = 1, [b, c]^{1+r} = a^{r p^2} b^{p^2}, [c, a]^{1+r} = a^{-r p^2} b^{p^2}, [a, b] = c^p \rangle$ , where  $p > 2$ ,  $r = 1, 2, \dots, p-2$  such that  $-r \notin (F_p)^2$ ; moreover,  $|G| = p^8$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_{p^2} \times C_p$  and  $G' = \langle a^{p^2}, b^{p^2}, c^p \rangle \cong C_p^3$ .
- (D8)  $\langle a, b, c \mid a^8 = b^8 = c^4 = 1, [b, c] = a^4 b^4, [c, a] = b^4, [a, b] = c^2 \rangle$ ; where  $|G| = 2^8$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_4 \times C_4 \times C_2$  and  $G' = \langle a^4, b^4, c^2 \rangle \cong C_2^3$ .

(Diii) *The type of  $G/G'$  is  $(p, p, p)$  where  $p > 2$ .*

- (D9)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [b, c] = a^p, [c, a] = b^p, [a, b] = c^p \rangle$ ; where  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, c^p \rangle \cong C_p^3$ .
- (D10)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [b, c] = a^p, [c, a] = c^{-p}, [a, b] = b^p \rangle$ ; where  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, c^p \rangle \cong C_p^3$ .
- (D11)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [b, c] = a^p, [c, a] = c^{-p}, [a, b] = b^p c^{\nu p} \rangle$ , where  $\nu = 1$  or is a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, c^p \rangle \cong C_p^3$ .

- (D12)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [b, c] = a^p, [c, a]^{1+r} = b^{rp}c^{-p}, [a, b]^{1+r} = b^p c^p \rangle$ ,  
where  $r = 1, 2, \dots, p-2$ . Moreover,  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, c^p \rangle \cong C_p^3$ .
- (D13)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [b, c] = a^{-p}b^p c^p, [c, a] = a^{-p}b^p, [a, b] = a^p \rangle$ ;  
where  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, c^p \rangle \cong C_p^3$ .
- (D14)  $\langle a, b, c; d \mid a^{p^2} = b^{p^2} = c^p = d^p = 1, [b, c] = a^p, [c, a] = b^{\nu p}, [a, b] = d, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  such that  $-\nu \notin (F_p)^2$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, d \rangle \cong C_p^3$ .
- (D15)  $\langle a, b, c; d \mid a^{p^2} = b^{p^2} = c^p = d^p = 1, [b, c] = a^p, [c, a] = d, [a, b] = b^p, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^p, b^p, d \rangle \cong C_p^3$ .
- (D16)  $\langle a, b, c; d \mid a^p = b^{p^2} = c^{p^2} = d^p = 1, [b, c] = d, [c, a]^{1+r} = b^{rp}c^{-p}, [a, b]^{1+r} = b^p c^p, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $p > 2$ ,  $r = 1, 2, \dots, p-2$  such that  $-r \notin (F_p)^2$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = Z(G) = G' = \langle b^p, c^p, d \rangle \cong C_p^3$ .

(Div) The type of  $G/G'$  is  $(2, 2, 2)$ .

- (D17)  $\langle a, b, c; d \mid a^4 = b^2 = c^4 = d^2 = 1, [b, c] = d, [c, a] = a^2, [a, b] = c^2, [d, a] = [d, b] = [d, c] = 1 \rangle$ ; where  $\Phi(G) = Z(G) = G' = \langle a^2, c^2, d \rangle \cong C_2^3$ .
- (D18)  $\langle a, b, c; d \mid a^4 = b^4 = c^4 = d^2 = 1, [b, c] = d, [c, a] = a^2, [a, b] = b^2 = c^2, [d, a] = [d, b] = [d, c] = 1 \rangle$ ; where  $\Phi(G) = Z(G) = G' = \langle a^2, b^2, d \rangle \cong C_2^3$ .
- (D19)  $\langle a, b, c; d \mid a^4 = b^4 = c^4 = d^2 = 1, [b, c] = d, [c, a] = a^2 b^2, [a, b] = a^2 = c^2, [d, a] = [d, b] = [d, c] = 1 \rangle$ ;  $\Phi(G) = Z(G) = G' = \langle a^2, b^2, d \rangle \cong C_2^3$ .

Moreover, Table 6 gives  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  for the groups (D1)–(D19).

$(\mu_0, \mu_1, \mu_2)$	$\alpha_1(G)$	types of $\mathcal{A}_3$ groups
$(0, p^2, p+1)$	$p^3 + 2p^2$	(D1), (D4), (D6)–(D10), (D12)–(D14), (D16)
$(0, p^2 + p, 1)$	$2p^2 + p$	(D2) where $-\nu \notin (F_p^*)^2$ ; (D3) where $-r \notin (F_p^*)^2$ ; (D5); (D11) where $-\nu \notin (F_p^*)^2$ ;
$(0, p^2 - p, 2p+1)$	$2p^3 + 2p^2 - p$	(D2) where $-\nu \in (F_p^*)^2$ ; (D3) where $-\nu \in (F_p^*)^2$ ; (D11) where $-\nu \in (F_p^*)^2$ ; (D15)
$(0, p^2 - 1, p+2)$	$p^3 + 3p^2 - 1$	(D17), (D18)
$(0, p^2 + 1, p)$	$p^3 + p^2 + 1$	(D19)

Table 6: The enumeration of (D1)–(D19)

**Proof** Let  $A$  and  $B$  be two distinct  $\mathcal{A}_1$ -subgroups of index  $p$ . By Lemma 2.15,  $|G'| \leq p|A'||B'| \leq p^3$ . It is easy to see that  $\Phi(A) = \Phi(B) = \Phi(G)$ . By Lemma 2.2, we have  $\Phi(A) = Z(A)$  and  $\Phi(B) = Z(B)$ . Since  $[\Phi(G), A] = [\Phi(G), B] = 1$  and  $G = AB$ , we have  $\Phi(G) \leq Z(G)$ . Moreover,  $G' \leq C_p^3$ . If  $|G'| \leq p^2$ , then  $G$  has an abelian subgroup of index  $p$ , a contradiction. Now we have  $d(G) = 3$ ,  $\Phi(G) \leq Z(G)$  and  $G' \cong C_p^3$ . Thus  $G$  is one of the groups classified in [23]. Let the type of  $G/G'$  be  $(p^{m_1}, p^{m_2}, p^{m_3})$ , where  $m_1 \geq m_2 \geq m_3$ . By [23, Theorem 2.7],  $m_3 = 1$ . By [23, Theorem 2.8],  $m_1 \leq 2$ .

If  $m_1 = 2$  and  $m_2 = 1$ , then  $G$  is either one of the groups determined by [23, Theorem 4.1] for  $p > 2$  or one of the groups determined by [23, Theorem 7.1] for  $p = 2$ . By [23, Theorem 4.3 & 7.4], we get the groups (D1)–(D5).

If  $m_1 = m_2 = 2$ , then  $G$  is one of the groups determined by [23, Theorem 5.1]. By [23, Theorem 5.2], we get the groups (D6)–(D8).

If  $m_1 = 1$  and  $p > 2$ , then  $G$  is one of the groups determined by [23, Theorem 6.1]. By [23, Theorem 6.3], we get the groups (D9)–(D16).

If  $m_1 = 1$  and  $p = 2$ , then  $G$  is one of the groups listed in [23, Theorem 7.6]. By [23, Theorem 7.7], we get the groups (D17)–(D19). Table 7 gives the correspondence.

Groups	Groups in [23, Theorem 4.1 & 5.1 & 6.1 & 7.1 & 7.6]	Groups	Groups in [23, Theorem 4.1 & 5.1 & 6.1 & 7.1 & 7.6]
(D1)	(D2) where $m_1 = 2$ and $m_2 = 1$	(D11)	(J3) where $m_1 = m_2 = m_3 = 1$
(D2)	(D3) where $m_1 = 2$ and $m_2 = 1$	(D12)	(J4) where $m_1 = m_2 = m_3 = 1$
(D3)	(D4) where $m_1 = 2$ and $m_2 = 1$	(D13)	(J5) where $m_1 = m_2 = m_3 = 1$
(D4)	(M2) where $m_1 = 2$	(D14)	(K1) where $m_1 = m_2 = m_3 = 1$ and $-\nu \notin F_p^2$
(D5)	(M3) where $m_1 = 2$	(D15)	(K2)
(D6)	(G3) where $-\nu \notin F_p^2$ , $m_1 = m_2 = 2$ and $m_3 = 1$	(D16)	(K6) where $m_1 = m_2 = m_3 = 1$ and $-r \notin F_p^2$
(D7)	(G4) where $-r \notin F_p^2$ , $m_1 = m_2 = 2$ and $m_3 = 1$	(D17)	(S7)
(D8)	(G7) where $m_1 = m_2 = 2$ and $m_3 = 1$	(D18)	(S8)
(D9)	(J1) where $m_1 = m_2 = m_3 = 1$	(D19)	(S9)
(D10)	(J2) where $m_1 = m_2 = m_3 = 1$		

Table 7: The correspondence from Theorem 4.6 to [23, Theorem 4.1,5.1,6.1,7.1 & 7.6]

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 4.6 as follows.

By Lemma 3.18,  $\alpha_1(G) = \mu_1 + p^2\mu_2$ . Hence we only need to calculate  $(\mu_0, \mu_1, \mu_2)$ . we may assume that  $G = \langle a_1, a_2, a_3; x, y, z \rangle$  with

$$x = [a_2, a_3], y = [a_3, a_1], z = [a_1, a_2], x^p = y^p = z^p = 1, a_i^{p^{m_i}} = x^{w_{i1}} y^{w_{i2}} z^{w_{i3}}, i = 1, 2, 3.$$

Let  $M = \langle a_2, a_3, \Phi(G) \rangle$ ,  $M_i = \langle a_1 a_2^i, a_3, \Phi(G) \rangle$  and  $M_{ij} = \langle a_1 a_3^i, a_2 a_3^j, \Phi(G) \rangle$ , where  $0 \leq i, j \leq p-1$ . Then  $M$ ,  $M_i$  and  $M_{ij}$  are all maximal subgroups of  $G$ . Let  $w(G) = (w_{ij})$ .

If  $G$  is the group (D1), then  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \nu & 1 \\ 0 & -1 & 0 \end{pmatrix}$ . By Lemma 4.5 (1),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  and  $M_{ij} \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

If  $G$  is the group (D2), then  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \nu^{-1} \end{pmatrix}$ . By Lemma 4.5 (1),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_1$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $j^2 = -\nu$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2 + p, 1)$  for  $-\nu \notin (F_p^*)^2$  and  $(\mu_0, \mu_1, \mu_2) = (0, p^2 - p, 2p + 1)$  for  $-\nu \in (F_p^*)^2$ .

If  $G$  is the group (D3), then  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & r \end{pmatrix}$ . By Lemma 4.5 (1),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_1$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $j^2 = -r^{-1}$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2 + p, 1)$  for  $-r \notin (F_p^*)^2$  and  $(\mu_0, \mu_1, \mu_2) = (0, p^2 - p, 2p + 1)$  for  $-r \in (F_p^*)^2$ .

If  $G$  is the group (D4), then  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By Lemma 4.5 (1),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_1$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $j = 1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

If  $G$  is the group (D5), then  $m_1 = 2$ ,  $m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}$ . By Lemma 4.5 (1),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_1$  and  $M_{ij} \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2 + p, 1)$  and  $\alpha_1(G) = 2p^2 + p$ .

If  $G$  is one of the groups (D6)–(D8), then  $m_1 = m_2 = 2$ ,  $m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \nu & 0 \\ 0 & 0 & 1 \end{pmatrix}$ ,  $\begin{pmatrix} 1 & -1 & 0 \\ 1 & r & 0 \\ 0 & 0 & 1 \end{pmatrix}$ , or  $\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By Lemma 4.5 (2),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  and  $M_{ij} \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

If  $G$  is the group (D9), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i^2 + 1 = 0$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $i^2 + j^2 + 1 = 0$ .

If  $-1 \in (F_p^*)^2$ , then  $i^2 + 1 = 0$  has two solutions, and by Lemma 2.9,  $i^2 + j^2 + 1 = 0$  has  $p - 1$  solutions. Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ . If  $-1 \notin (F_p^*)^2$ , then  $i^2 + 1 = 0$  has no solution, and by Lemma 2.9,  $i^2 + j^2 + 1 = 0$  has  $p + 1$  solutions. Hence we also have  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ .

If  $G$  is the group (D10), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ .

If  $G$  is the group (D11), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \nu & 1 \\ 0 & -1 & 0 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $i^2 = -\nu$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2 + p, 1)$  for  $-\nu \notin (F_p^*)^2$  and  $(\mu_0, \mu_1, \mu_2) = (0, p^2 - p, 2p + 1)$  for  $-\nu \in (F_p^*)^2$ .

If  $G$  is the group (D12), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & r \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i^2(r+1) + r = 0$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $i^2(r+1) + rj^2 + 1 = 0$ .

If  $-(r+1)^{-1}r \in (F_p^*)^2$ , then  $i^2(r+1) + r = 0$  has two solutions, and by Lemma 2.9,  $i^2(r+1) + j^2r + 1 = 0$  has  $p - 1$  solutions. Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ . If  $-(r+1)^{-1}r \notin (F_p^*)^2$ , then  $i^2(r+1) + r = 0$  has no solution, and by Lemma 2.9,  $i^2(r+1) + j^2r + 1 = 0$  has  $p + 1$  solutions. Hence we also have  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ .

If  $G$  is the group (D13), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & -1 & 0 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i^2 = 1$  and  $M_{ij} \in \mathcal{A}_2$  if and only if  $(i+1)^2 - j^2 - 1 = 0$ . By Lemma 2.9,  $(i+1)^2 - j^2 - 1 = 0$  has  $p - 1$  solutions. Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p+1)$ .

If  $G$  is the group (D14), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \nu & 0 \\ 0 & 0 & 0 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  and  $M_{ij} \in \mathcal{A}_1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p + 1)$ .

If  $G$  is the group (D15), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  and  $M_{ij} \in \mathcal{A}_1$  if and only if  $j = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2 - p, 2p + 1)$ .

If  $G$  is the group (D16), then  $p > 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & r \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_1$  if and only if  $i = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, p^2, p + 1)$ .

If  $G$  is the group (D17), then  $p = 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_1$  if and only if  $j = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, 3, 4)$ .

If  $G$  is the group (D18), then  $p = 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_2$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_1$  if and only if  $j = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, 3, 4)$ .

If  $G$  is the group (D19), then  $p = 2$  and  $m_1 = m_2 = m_3 = 1$  and  $w(G) = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ . By Theorem 4.5 (3),  $M \in \mathcal{A}_1$ ,  $M_i \in \mathcal{A}_2$  if and only if  $i = 0$  and  $M_{ij} \in \mathcal{A}_1$  if and only if  $i = j = 0$ . Hence  $(\mu_0, \mu_1, \mu_2) = (0, 5, 2)$ .  $\square$

**Theorem 4.7.**  *$G$  has no abelian subgroup of index  $p$  and  $G$  has a unique  $\mathcal{A}_1$ -subgroups of index  $p$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Ei)  $p > 2$ . In this case  $p = 3$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 1, 3)$  and  $\alpha_1(G) = 10$  except (E2), in which  $\alpha_1(G) = 28$ .

(E1)  $\langle s_1, s; s_2 \mid s_1^9 = s_2^9 = 1, s^3 = s_2^{3\delta}, [s_1, s] = s_2, [s_2, s] = s_2^{-3}s_1^{-3}, [s_2, s_1] = s_2^3 \rangle$  where  $\delta = 0, 1, 2$ ; moreover,  $|G| = 3^5$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle s_1^3, s_2 \rangle \cong C_3 \times C_9$  and  $Z(G) = \langle s_2^3 \rangle \cong C_3$ .

(E2)  $\langle a, b; c \mid a^{3^2} = b^{3^2} = c^3 = 1, [b, a] = c, [c, a] = a^3, [c, b] = b^{-3} \rangle$ ; where  $|G| = 3^5$ ,  $c(G) = 3$ ,  $\Phi(G) = G' = \langle a^3, b^3, c \rangle \cong C_3^3$  and  $Z(G) = \langle a^3, b^3 \rangle \cong C_3^2$ .

(E3)  $\langle s_1, \beta; s_2, x \mid s_1^9 = s_2^9 = x^3 = 1, \beta^3 = x, [s_1, \beta] = s_2, [s_2, \beta] = s_2^{-3}s_1^{-3}, [s_1, s_2] = x, [x, s_1] = [x, \beta] = 1 \rangle$ ; where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle s_1^3, s_2, x \rangle \cong C_9 \times C_3 \times C_3$  and  $Z(G) = \langle s_2^3, x \rangle \cong C_3^2$ .

- (E4)  $\langle s_1, \beta; s_2, x \mid s_1^9 = s_2^9 = x^3 = 1, \beta^3 = s_2^3 x, [s_1, \beta] = s_2, [s_2, \beta] = s_2^{-3} s_1^{-3}, [s_1, s_2] = x, [x, s_1] = [x, \beta] = 1 \rangle$ ; where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle s_1^3, s_2, x \rangle \cong C_9 \times C_3 \times C_3$  and  $Z(G) = \langle s_2^3, x \rangle \cong C_3^2$ .
- (E5)  $\langle \alpha, \beta; s_1, s_2, x \mid s_1^9 = s_2^3 = x^3 = 1, \beta^3 = x^2, \alpha^3 = s_2^{-1}, [\alpha, \beta] = s_1, [s_1, \alpha] = x, [s_1, \beta] = s_2, [s_2, \beta] = s_1^{-3}, [s_1, s_2] = [x, \alpha] = [x, \beta] = 1 \rangle$ ; where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle s_1, s_2, x \rangle \cong C_9 \times C_3 \times C_3$  and  $Z(G) = \langle s_1^3, x \rangle \cong C_3^2$ .
- (E6)  $\langle \alpha, \beta; s_1, s_2, x \mid s_1^9 = s_2^3 = x^3 = 1, \beta^3 = x, \alpha^3 = s_2^{-1} x, [\alpha, \beta] = s_1, [s_1, \alpha] = x, [s_1, \beta] = s_2, [s_2, \beta] = s_1^{-3}, [s_1, s_2] = [x, \alpha] = [x, \beta] = 1 \rangle$ , where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle s_1, s_2, x \rangle \cong C_9 \times C_3 \times C_3$  and  $Z(G) = \langle s_1^3, x \rangle \cong C_3^2$ .

(Eii)  $p = 2$ .

- (E7)  $\langle a, b \mid a^{16} = 1, b^{2^{s+t+2}} = a^{2^{s+t'+2}}, [a, b] = a^2 \rangle$ , where  $s, t, t'$  are non-negative integers with  $1 \leq s + t' \leq 2$ ,  $tt' = 0$  and if  $t' \neq 0$ , then  $s + t' = 2$ ; moreover,  $|G| = 2^{s+t+6}$ ,  $c(G) = 4$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_8 \times C_{2^{t-t'+3}}$ ,  $G' = \langle a^2 \rangle \cong C_8$ ,  $Z(G) = \langle a^8, b^4 \rangle \cong C_2 \times C_{2^{t-t'+2}}$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 1, 2)$  and  $\alpha_1(G) = 5$ ;
- (E8)  $\langle a, b, c \mid a^2 = b^8 = 1, c^2 = b^{4t}, [a, b] = b^4, [a, c] = 1, [c, b] = b^2 \rangle$  where  $t = 0, 1$ ; moreover,  $|G| = 2^5$ ,  $c(G) = 3$ ,  $\Phi(G) = G' = \langle b^2 \rangle \cong C_4$ ,  $Z(G) = \langle b^4 \rangle \cong C_2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 1, 6)$  and  $\alpha_1(G) = 9$ ;
- (E9)  $\langle a, b, c; d \mid a^4 = b^4 = c^2 = d^2 = 1, [a, b] = b^2, [c, a] = a^2 b^2, [c, b] = d, [d, a] = [d, b] = [d, c] = 1 \rangle$ ; moreover  $|G| = 2^6$ ,  $c(G) = 2$ ,  $\Phi(G) = Z(G) = G' = \langle a^2, b^2, d \rangle \cong C_2^3$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 1, 6)$  and  $\alpha_1(G) = 25$ ;
- (E10)  $\langle a, b, c; d \mid a^4 = b^4 = c^2 = d^2 = 1, [a, b] = a^2, [c, a] = a^2 b^2, [c, b] = d, [d, a] = [d, b] = [d, c] = 1 \rangle$ , moreover  $|G| = 2^6$ ,  $c(G) = 2$ ,  $\Phi(G) = Z(G) = G' = \langle a^2, b^2, d \rangle \cong C_2^3$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 1, 6)$  and  $\alpha_1(G) = 25$ .

**Proof** Finite  $p$ -groups with a unique  $\mathcal{A}_1$ -subgroup were classified in [22] for  $p > 2$  and [24] for  $p = 2$ . If  $p > 2$ , then, by using the classification in [22] and calculating the maximal index of  $\mathcal{A}_1$ -subgroups for these groups, we get the groups (E1)–(E6). The details are omitted. Table 8 gives the correspondence.

Groups	Groups in [22, Lemma 2.12]	Groups	Groups in [22, Corollary 3.5]	Groups	Groups in [22, Theorem 3.8]
(E1)	(2) where $e = 2$	(E2)	(2)	(E3)	(1) where $e = 2$ and $k = 1$
				(E4)	(2) where $e = 2$ and $k = 1$
				(E5)	(3) where $e = 2$ and $k = 2$
				(E6)	(4) where $e = 2$ and $k = 1$

Table 8: The correspondence from Theorem 4.6 to [22, Lemma 2.12, Corollary 3.5, Theorem 3.8]

If  $p = 2$ , then, by lemma 3.7,  $G$  is either metacyclic or  $|G| \leq 2^8$ . If  $G$  is metacyclic, then, by Lemma 3.1,  $|G'| = p^3$ . By using the classification of metacyclic 2-groups in [30] and checking the number of their  $\mathcal{A}_1$ -subgroups of index  $p$ , we get the group (E7). The details is omitted. If  $G$  is not metacyclic, then  $|G| \leq 2^8$ . By using Magma to check the SmallGroup database, we get the groups (E8)–(E10).



We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 4.7 as follows.

By hypothesis,  $\mu_0 = 0$  and  $\mu_1 = 1$ . Hence  $\mu_2 = p$  for  $d(G) = 2$  and  $\mu_2 = p + p^2$  for  $d(G) = 3$ . In the following, we calculate  $\alpha_1(G)$ .

If  $G$  is the group (E2), then  $|G| = 3^5$ , and  $|H'| = p$  for any maximal subgroup  $H$ . By Lemma 2.6 (7),  $\alpha_1(H) = p^2$  for  $H \in \mathcal{A}_2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p^2 \mu_2 = p^3 + 1 = 28.$$

If  $G$  is one of the group (E1) and (E3)–(E6), then  $c(G) = 4$ , and  $c(H) = 3$  for any  $\mathcal{A}_2$ -subgroup  $H$ . Since  $d(H) = 2$ ,  $\alpha_1(H) = p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p \mu_2 = p^2 + 1 = 10.$$

If  $G$  is the group (E7), then  $G$  is metacyclic. Hence  $H$  is metacyclic for any  $\mathcal{A}_2$ -subgroup  $H$ . Since  $d(H) = 2$ ,  $\alpha_1(H) = p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_1 + p \mu_2 = p^2 + 1 = 5.$$

If  $G$  is the group (E8), then  $d(G) = 3$  and  $c(G) = 3$ . All maximal subgroups of  $G$  are  $M = \langle b, a \rangle$ ,  $M_0 = \langle c, a, b^2 \rangle$ ,  $M_1 = \langle cb, a, b^2 \rangle$  and  $M_{ij} = \langle ca^i, ba^j \rangle$  where  $i, j = 0, 1$ . Here  $M$  is the unique  $\mathcal{A}_1$ -subgroup of index 2,  $|M'_i| = 2$  and  $c(M_{ij}) = 3$ . By Lemma 2.6,  $\alpha_1(M_i) = 4$  and  $\alpha_1(M_{ij}) = 2$ . Let  $H \in \Gamma_2(G)$ . Then  $H = \langle a^i b^j c^k, b^2 \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $k = 1$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - 2 \sum_{H \in \Gamma_2} \alpha_1(H) = 1 + 4 \times 2 + 2 \times 4 - 2 \times 4 = 9.$$

If  $G$  is the group (E9) or (E10), then  $d(G) = 3$  and  $\Phi(G) = Z(G)$ . By Theorem 3.18,  $\alpha_1(G) = \mu_1 + 4\mu_2 = 25$ .  $\square$

## 5 $\mathcal{A}_3$ -groups without $\mathcal{A}_1$ -subgroup of index $p$

Assume  $G$  is an  $\mathcal{A}_3$ -group without an  $\mathcal{A}_1$ -subgroup of index  $p$ . We discuss according to  $G$  has an abelian subgroup of index  $p$  or not. In this case,  $\mathcal{A}_3$ -groups have 151 non-isomorphic types. Theorem 5.1, 5.2, 5.3, 5.4 and 5.5 give the classification of  $\mathcal{A}_3$ -groups without an  $\mathcal{A}_1$ -subgroup of index  $p$  and with an abelian subgroup of index  $p$ . They are the groups (F1)–(F8), (G1)–(G12), (H1)–(H10), (I1)–(I11) and (J1)–(J9). Theorem 5.6, 5.7, 5.13, 5.15 and 5.16 give the classification of  $\mathcal{A}_3$ -groups without an  $\mathcal{A}_1$ -subgroup of index  $p$  and an abelian subgroup of index  $p$ . They are the groups (K1)–(K5), (L1)–(L2), (M1)–(M62), (N1)–(N26) and (O1)–(O6).

## 5.1 $G$ has an abelian subgroup of index $p$

In this section assume  $G$  is an  $\mathcal{A}_3$ -group with an abelian subgroup of index  $p$ , and  $G$  without an  $\mathcal{A}_1$ -subgroup of index  $p$  in Theorem 5.1, 5.2, 5.3, 5.4 and 5.5.

**Theorem 5.1.**  $d(G) = 2$  and  $c(G) = 4$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

- (F1)  $\langle a, b \mid a^{16} = b^{2^m} = 1, [a, b] = a^{-2} \rangle$ ; where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_8 \times C_{2^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_8$  if  $m = 1$ ,  $G' = \langle a^2 \rangle \cong C_8$ ,  $Z(G) = \langle a^8, b^2 \rangle \cong C_2 \times C_{2^{m-1}}$  if  $m > 1$ ,  $Z(G) \cong C_2$  if  $m = 1$ .
- (F2)  $\langle a, b \mid a^{16} = b^{2^m} = 1, [a, b] = a^6 \rangle$ ; where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_8 \times C_{2^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_8$  if  $m = 1$ ,  $G' = \langle a^2 \rangle \cong C_8$ ,  $Z(G) = \langle a^8, b^2 \rangle \cong C_2 \times C_{2^{m-1}}$  if  $m > 1$ ,  $Z(G) \cong C_2$  if  $m = 1$ .
- (F3)  $\langle a, b \mid a^{16} = 1, b^{2^m} = a^8, [a, b] = a^{-2} \rangle$ ; where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_2^3$  if  $m = 1$ ,  $\Phi(G) \cong C_2^3 \times C_2$  if  $m = 2$ ,  $\Phi(G) \cong C_{2^m} \times C_{2^2}$  if  $m > 2$ ,  $G' = \langle a^2 \rangle \cong C_8$  and  $Z(G) = \langle b^2 \rangle \cong C_{2^m}$ .
- (F4)  $\langle a_1, b; a_2 \mid a_1^9 = a_2^9 = b^{3^m} = 1, [a_1, b] = a_2, [a_2, b] = a_1^{-3} a_2^{3t}, [a_1, a_2] = 1 \rangle$ , where  $t = 1, 2$ ; moreover,  $|G| = 3^{m+4}$ ,  $\Phi(G) = \langle a_2, a_1^3, b^3 \rangle \cong C_3 \times C_9 \times C_{3^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_3 \times C_9$  if  $m = 1$ ,  $G' = \langle a_2, a_1^3 \rangle \cong C_3 \times C_9$ ,  $Z(G) = \langle a_2^3, b^3 \rangle \cong C_3 \times C_{3^{m-1}}$  if  $m > 1$ ,  $Z(G) \cong C_3$  if  $m = 1$ .
- (F5)  $\langle a_1, b; a_2 \mid a_1^9 = a_2^9 = 1, b^{3^m} = a_2^{-3}, [a_1, b] = a_2, [a_2, b] = a_1^{-3} a_2^{-3}, [a_1, a_2] = 1 \rangle$ ; where  $|G| = 3^{m+4}$ ,  $\Phi(G) = \langle a_2, a_1^3, b^3 \rangle \cong C_9 \times C_3$  if  $m = 1$ ,  $\Phi(G) \cong C_{3^m} \times C_3 \times C_3$  if  $m > 1$ ,  $G' = \langle a_2, a_1^3 \rangle \cong C_9 \times C_3$  and  $Z(G) = \langle b^3 \rangle \cong C_{3^m}$ .
- (F6)  $\langle a_1, b; a_2, a_3, a_4 \mid a_i^p = b^{p^m} = 1, [a_j, b] = a_{j+1}, [a_4, b] = 1, [a_i, a_j] = 1 \rangle$ , where  $p \geq 5$ ,  $1 \leq i \leq 4$ ,  $1 \leq j \leq 3$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_3, a_4, b^p \rangle \cong C_p^3 \times C_{p^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_p^3$  if  $m = 1$ ,  $G' = \langle a_2, a_3, a_4 \rangle \cong C_p^3$ ,  $Z(G) = \langle a_4, b^p \rangle \cong C_p \times C_{p^{m-1}}$  if  $m > 1$ ,  $Z(G) \cong C_p$  if  $m = 1$ .
- (F7)  $\langle a_1, b; a_2, a_3 \mid a_i^p = b^{p^{m+1}} = 1, [a_j, b] = a_{j+1}, [a_3, b] = b^{p^m}, [a_i, a_j] = 1 \rangle$ , where  $p \geq 5$ ,  $1 \leq i \leq 3$ ,  $1 \leq j \leq 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_3, b^p \rangle \cong C_p^2 \times C_{p^m}$ ,  $G' = \langle a_2, a_3, b^{p^m} \rangle \cong C_p^3$  and  $Z(G) = \langle b^p \rangle \cong C_{p^m}$ .
- (F8)  $\langle a_1, b; a_2, a_3 \mid a_1^{p^2} = a_i^p = b^{p^m} = 1, [a_j, b] = a_{j+1}, [a_3, b] = a_1^{tp}, [a_i, a_j] = 1 \rangle$ , where  $2 \leq i \leq 3$ ,  $1 \leq j \leq 2$ , and  $t = t_1, t_2, \dots, t_{(3,p-1)}$ , where  $p \geq 5$ ,  $t_1, t_2, \dots, t_{(3,p-1)}$  are the coset representatives of the subgroup  $(F_p^*)^3$  in  $F_p^*$ . Moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_3, a_1^p, b^p \rangle \cong C_p^3 \times C_{p^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_p^3$  if  $m = 1$ ,  $G' = \langle a_2, a_3, a_1^p \rangle \cong C_p^3$ ,  $Z(G) = \langle a_1^p, b^p \rangle \cong C_p \times C_{p^{m-1}}$  if  $m > 1$ ,  $Z(G) \cong C_p$  if  $m = 1$ .

Moreover,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p)$  and  $\alpha_1(G) = p^2$ .

**Proof** Let  $A$  be an abelian subgroup of index  $p$  and  $B$  be a non-abelian subgroup of index  $p$ . By Lemma 2.11,  $G_3 = B'$ ,  $G_4 = B_3$ ,  $|G_3/G_4| = p$  and  $|G'| = p^{c(G)-1} = p^3$ . Since  $c(G) = 4$ ,  $c(B) \geq 3$ . Hence Lemma 2.6 gives that  $d(B) = 2$  and  $c(B) = 3$ . By arbitrariness of  $B$ , all non-abelian subgroups of  $G$  are generated by two elements. Such groups were classified by [29]. It follows from [29, Main Theorem] that  $G$  is either a  $p$ -groups of maximal class or one of the groups listed in [29, Theorem 3.12-3.13]. Since  $c(G) = 4$ , they are the groups (F1)–(F8).

Since  $d(G) = 2$ ,  $G$  has  $1+p$  maximal subgroups. Hence  $(\mu_0, \mu_1, \mu_2) = (1, 0, p)$ . Since  $d(H) = 2$  for any non-abelian maximal subgroup  $H$ ,  $\alpha_1(H) = p$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p\mu_2 = p^2$ .  $\square$

**Theorem 5.2.**  $d(G) = 2$  and  $c(G) = 3$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

(Gi)  $G' \cong C_{p^2}$ .

(G1)  $\langle a, b, c \mid a^8 = 1, c^2 = a^4 = b^4, [a, b] = c, [c, a] = 1, [c, b] = c^2 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_2 \times C_2 \times C_4$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle ca^2, b^2 \rangle \cong C_2 \times C_4$ .

(G2)  $\langle a, b, c \mid a^8 = b^4 = 1, c^2 = a^4, [a, b] = c, [c, a] = 1, [c, b] = c^2 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_2 \times C_2 \times C_4$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle ca^2, b^2 \rangle \cong C_2^2$ .

(G3)  $\langle a, b, c \mid a^8 = b^4 = 1, c^2 = a^4, [a, b] = c, [c, a] = c^2, [c, b] = 1 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_2^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2, cb^2 \rangle \cong C_4 \times C_2$ .

(G4)  $\langle a, b, c \mid a^{2^{n+1}} = b^4 = 1, c^2 = a^{2^n}, [a, b] = c, [c, a] = c^2, [c, b] = 1 \rangle$ , where  $n > 2$ ; moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^n} \times C_2^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2, cb^2 \rangle \cong C_{2^n} \times C_2$ .

(G5)  $\langle a, b, c \mid a^{2^n} = b^8 = 1, c^2 = b^4, [a, b] = c, [c, a] = c^2, [c, b] = 1 \rangle$ , where  $n > 2$ ; moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^{n-1}} \times C_2 \times C_4$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2, cb^2 \rangle \cong C_{2^{n-1}} \times C_2$ .

(G6)  $\langle a, b, c \mid a^{2^n} = b^4 = c^4 = 1, [a, b] = c, [c, a] = c^2, [c, b] = 1 \rangle$ , where  $n > 2$ . Moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^{n-1}} \times C_2 \times C_4$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^2, cb^2 \rangle \cong C_{2^{n-1}} \times C_4$ .

(Gii)  $G' \cong C_p^2$ .

(G7)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = 1, [c, b] = a^{\nu p^2} \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p^2$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_p$ .

(G8)  $\langle a, b, c, d \mid a^{p^2} = b^{p^2} = c^p = d^p = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle$ , where  $p > 2$ . Moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p, c, d \rangle \cong C_p^4$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle a^p, b^p, d \rangle \cong C_p^3$ .

- (G9)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = a^{p^2}, [c, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p^2$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_p$ .
- (G10)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = a^{p^n}, [c, b] = 1 \rangle$ , where  $p > 2$  and  $n > 2$ ; moreover,  $|G| = p^{n+4}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_p^2$ ,  $G' = \langle a^{p^n}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^n} \times C_p$ .
- (G11)  $\langle a, b, c \mid a^{p^n} = b^{p^3} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p^2}, [c, b] = 1 \rangle$ , where  $p > 2$ ,  $n > 2$ , and  $\nu = 1$  or or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^{n+4}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^{n-1}} \times C_{p^2}$ .
- (G12)  $\langle a, b, c, d \mid a^{p^n} = b^{p^2} = c^p = d^p = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle$ , where  $p > 2$  and  $n > 2$ . Moreover,  $|G| = p^{n+4}$ ,  $\Phi(G) = \langle a^p, b^p, c, d \rangle \cong C_{p^{n-1}} \times C_p^3$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle a^p, b^p, d \rangle \cong C_{p^{n-1}} \times C_p^2$ .

Moreover,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p)$  and  $\alpha_1(G) = p^3$ .

**Proof** Let  $A$  be an abelian subgroup of index  $p$  and  $B$  be a non-abelian subgroup of index  $p$ . By Lemma 2.11,  $G_3 = B'$ ,  $G_4 = B_3$ ,  $|G_3/G_4| = p$  and  $|G'| = p^{c(G)-1}$ .

Since  $c(G) = 3$ ,  $|B'| = |G_3| = p$  and  $|G'| = p^2$ . Let  $\bar{G} = G/G_3$ . Then  $|\bar{G}'| = p$  and  $d(\bar{G}) = 2$ . Hence  $\bar{G}$  is an  $\mathcal{A}_1$ -group by Lemma 2.2. If  $\bar{G}$  is metacyclic, then  $G$  is also metacyclic by Lemma 2.1 and hence  $G$  is an  $\mathcal{A}_2$ -group by Lemma 3.1. This contradicts that  $G$  is an  $\mathcal{A}_3$ -group. Thus  $\bar{G}$  is a non-metacyclic  $\mathcal{A}_1$ -group. If  $G' \cong C_{p^2}$ , then  $G$  is one of the groups listed in [3, Theorem 3.5]. By [3, Theorem 3.1 & 3.6] we get groups (G1)–(G6). If  $G' \cong C_p^2$ , then  $G$  is one of the groups listed in [3, Theorem 4.6]. By [3, Theorem 4.1 & 4.7] we get groups (G7)–(G12).

Groups	Groups in [3, Theorem 3.5]	Groups	Groups in [3, Theorem 4.6]
(G1)	(C3)	(G7)	(G1) where $m = 2$ and $p > 2$
(G2)	(C4)	(G8)	(G2) where $m = 2$ and $p > 2$
(G3)	(C5)	(G9)	(G3) where $m = 2$ and $p > 2$
(G4)	(E2) where $p = m = 2$	(G10)	(J2) where $m = 2$ and $p > 2$
(G5)	(E5) where $p = m = 2$	(G11)	(J4) where $m = 2$ and $p > 2$
(G6)	(E8) where $p = m = 2$	(G12)	(J6) where $m = 2$ and $p > 2$

Table 9: The correspondence from Theorem 5.2 to [3, Theorem 3.5 & 4.6]

Now we calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$ . Since  $d(G) = 2$ ,  $G$  has  $1 + p$  maximal subgroups. Hence  $(\mu_0, \mu_1, \mu_2) = (1, 0, p)$ . Since  $|H'| = p$  for any non-abelian maximal subgroup  $H$ , by Lemma 2.6 (7),  $\alpha_1(H) = p^2$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p^2 \mu_2 = p^3$ .  $\square$

**Theorem 5.3.**  $d(G) = 3$  and  $\Phi(G) \leq Z(G)$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

(Hi)  $\Phi(G) < Z(G)$ ,  $G' \cong C_p$  and  $c(G) = 2$ . Moreover,  $(\mu_0, \mu_1, \mu_2) = (1 + p, 0, p^2)$  and  $\alpha_1(G) = p^4$ .

(H1)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^m} = c^{p^2} = 1, [a, b] = a^{p^n}, [c, a] = [c, b] = 1 \rangle \cong M_p(n + 1, m) \times C_{p^2}$ , where  $\min\{n, m\} \geq 2$ ; moreover,  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_p$ ,  $G' = \langle a^{p^n} \rangle$  and  $Z(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_{p^2}$ .

(H2)  $\langle a, b, c; d \mid a^{p^n} = b^{p^m} = c^{p^2} = d^p = 1, [a, b] = d, [c, a] = [c, b] = 1 \rangle \cong M_p(n, m, 1) \times C_{p^2}$ , where  $n \geq m \geq 2$ ; moreover,  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p, c^p, d \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_p \times C_p$ ,  $G' = \langle d \rangle$  and  $Z(G) = \langle a^p, b^p, c, d \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_{p^2} \times C_p$ .

(H3)  $\langle a, b, c \mid a^{p^n} = b^{p^m} = c^{p^3} = 1, [a, b] = c^{p^2}, [c, a] = [c, b] = 1 \rangle \cong M_p(n, m, 1) * C_{p^3}$ , where  $n \geq m \geq 2$ . Moreover,  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_{p^2}$ ,  $G' = \langle c^{p^2} \rangle$  and  $Z(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_{p^3}$ .

(Hii)  $\Phi(G) = Z(G)$ ,  $G' \cong C_p^2$  and  $c(G) = 2$ . Moreover,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p + p^2)$  and  $\alpha_1(G) = p^4 + p^3$ .

(H4)  $\langle a, b, c \mid a^{p^l} = b^{p^2} = c^{p^2} = 1, [b, c] = 1, [c, a] = c^p, [a, b] = b^{-p} \rangle$ , where  $p > 2$  and  $l \geq 2$ ; moreover,  $|G| = p^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{l-1}} \times C_p^2$ ,  $G' = \langle b^p, c^p \rangle$ .

(H5)  $\langle a, b, c \mid a^{p^l} = b^{p^3} = c^{p^3} = 1, [b, c] = 1, [c, a] = b^{p^2} c^{tp^2}, [a, b] = b^{-tp^2} c^{\nu p^2} \rangle$ , where  $p > 2$ ,  $l \geq 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  such that  $-\nu \notin (F_p^*)^2$  and  $t \in \{0, 1, \dots, \frac{p-1}{2}\}$ ; moreover,  $|G| = p^{l+6}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^{l-1}} \times C_{p^2} \times C_{p^2}$ ,  $G' = \langle b^{p^2}, c^{p^2} \rangle$ .

(H6)  $\langle a, b, c \mid a^{2^l} = b^4 = c^4 = 1, [b, c] = 1, [c, a] = c^2, [a, b] = b^2 \rangle$ ; moreover,  $|G| = 2^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_{2^{l-1}} \times C_2 \times C_2$  if  $l > 1$ ,  $G' = \langle b^2, c^2 \rangle$ ,  $\Phi(G) = Z(G) \cong C_2 \times C_2$  if  $l = 1$ .

(H7)  $\langle a, b, c \mid a^{2^l} = b^8 = c^8 = 1, [b, c] = 1, [c, a] = b^4, [a, b] = b^4 c^4 \rangle$ , where  $l \geq 2$ ; moreover,  $|G| = 2^{l+6}$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_{2^{l-1}} \times C_4 \times C_4$ ,  $G' = \langle b^4, c^4 \rangle$ .

(H8)  $\langle a, b, c \mid a^{p^{l+1}} = b^{p^3} = c^{p^2} = 1, [b, c] = 1, [c, a] = b^{p^2}, [a, b] = a^{p^l} \rangle$ , where  $l \geq 2$ ; moreover,  $|G| = p^{l+6}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^l} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^l}, b^{p^2} \rangle$ .

(H9)  $\langle a, b, c; x \mid a^{p^l} = b^{p^2} = c^{p^2} = x^p = 1, [a, b] = c^p, [a, c] = x, [b, c] = [x, a] = [x, b] = [x, c] = 1 \rangle$ , where  $l \geq 2$ ; moreover,  $|G| = p^{l+5}$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p, x \rangle \cong C_{p^{l-1}} \times C_p^3$ ,  $G' = \langle c^p, x \rangle$ .

(H10)  $\langle a, b, c; x, y \mid a^{p^l} = b^p = c^p = x^p = y^p = 1, [a, b] = x, [a, c] = y, [b, c] = [x, a] = [x, b] = [x, c] = [y, a] = [y, b] = [y, c] = 1 \rangle$ , where  $l \geq 2$  if  $p = 2$ ; moreover,  $|G| = p^{l+4}$ ,  $\Phi(G) = Z(G) = \langle a^p, x, y \rangle \cong C_{p^{l-1}} \times C_p^2$ ,  $G' = \langle x, y \rangle$ .

**Proof** By Lemma 2.8,  $|G'| \leq p^2$ . If  $|G'| = p$ , then  $G$  is one of the groups listed in [2, Theorem 3.1]. Since  $G$  has no  $\mathcal{A}_1$ -subgroup of index  $p$ , they are the groups (H1)–(H3). If  $|G'| = p^2$ , then  $G$  is one of the groups listed in [2, Theorem 4.8]. By checking [2, Table 4] we get groups (H4)–(H10).

Groups	Groups in [2, Theorem 4.8]	Groups	Groups in [2, Theorem 4.8]
(H4)	(A1) where $l \geq 2$ and $m = 1$	(H8)	(A10) where $l \geq 2 = m = n$
(H5)	(A3) where $l \geq m = 2$ and $-\nu \notin F_p^2$	(H9)	(B2) where $m = 2$ and $n = 1$
(H6)	(A5) where $m = 1$	(H10)	(C) where $m = n = 1$
(H7)	(A6) where $l \geq m = 2$ and $-\nu \notin (F_p^*)^2$		

Table 10: The correspondence from Theorem 5.3 to [2, Theorem 4.8]

Now we calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$ . If  $G$  is one of the groups (H1)–(H3), then  $\mu_0 = 1 + p$ . Since  $\mu_1 = 0$ ,  $\mu_2(G) = p^2$ . By Lemma 3.18,  $\alpha_1(G) = p^2\mu_2 = p^4$ .

If  $G$  is one of the groups (H4)–(H10), then  $d(G) = 3$  and  $G' \cong C_p^2$ . Since  $\Phi(G) = Z(G)$ ,  $\mu_0 = 1$ . Hence  $(\mu_0, \mu_1, \mu_2) = (1, 0, p + p^2)$ . By Lemma 3.18,  $\alpha_1(G) = p^2\mu_2 = p^4 + p^3$ .  $\square$

**Theorem 5.4.**  $d(G) = 3$  and  $\Phi(G) \not\subseteq Z(G)$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

- (Ii)  $|G'| = p^2$  and  $c(G) = 3$ . In this case,  $Z(G) \not\subseteq \Phi(G)$ ,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p^2 + p)$  and  $\alpha_1(G) = p^3$ .
- (II)  $\langle a, b, x \mid a^8 = b^{2^m} = x^2 = 1, [a, b] = a^{-2}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ ;  
where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_{2^{m-1}} \times C_4$  if  $m > 1$ ,  $\Phi(G) \cong C_4$  if  $m = 1$ ,  $G' = \langle a^2 \rangle \cong C_4$ ,  $Z(G) = \langle a^4, b^2, x \rangle \cong C_{2^{m-1}} \times C_2^2$  if  $m > 1$ ,  $Z(G) \cong C_2^2$  if  $m = 1$ .
- (I2)  $\langle a, b, x \mid a^8 = b^{2^m} = 1, x^2 = a^4, [a, b] = a^{-2}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle * \langle x \rangle$ ;  
where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_{2^{m-1}} \times C_4$  if  $m > 1$ ,  $\Phi(G) \cong C_4$  if  $m = 1$ ,  $G' = \langle a^2 \rangle \cong C_4$ ,  $Z(G) = \langle b^2, x \rangle \cong C_{2^{m-1}} \times C_4$  if  $m > 1$ ,  $Z(G) \cong C_4$  if  $m = 1$ .
- (I3)  $\langle a, b, x \mid a^8 = b^{2^m} = x^2 = 1, [a, b] = a^2, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ ;  
where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_{2^{m-1}} \times C_4$  if  $m > 1$ ,  $\Phi(G) \cong C_4$  if  $m = 1$ ,  $G' = \langle a^2 \rangle \cong C_4$ ,  $Z(G) = \langle a^4, b^2, x \rangle \cong C_{2^{m-1}} \times C_2^2$  if  $m > 1$ ,  $Z(G) \cong C_2^2$  if  $m = 1$ .
- (I4)  $\langle a, b, x \mid a^8 = x^2 = 1, b^{2^m} = a^4, [a, b] = a^{-2}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ ;  
where  $|G| = 2^{m+4}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_4$  if  $m = 1$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_{2^m} \times C_2$  if  $m > 1$ ,  $G' = \langle a^2 \rangle \cong C_4$  and  $Z(G) = \langle b^2, x \rangle \cong C_{2^m} \times C_2$ .
- (I5)  $\langle a_1, b, x; a_2, a_3 \mid a_1^p = a_2^p = a_3^p = b^{p^m} = x^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = [a_3, a_1] = [a_3, a_2] = [x, a_1] = [x, b] = 1 \rangle = \langle a_1, b \rangle \times \langle x \rangle$ ,  
where  $p > 3$  for  $m = 1$  and  $p > 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) =$

- $\langle a_2, a_3, b^p \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $\Phi(G) \cong C_p^2$  if  $m = 1$ ,  $G' = \langle a_2, a_3 \rangle \cong C_p^2$ ,  $Z(G) = \langle a_3, b^p, x \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $Z(G) \cong C_p^2$  if  $m = 1$ .
- (I6)  $\langle a_1, x, b; a_2 \mid a_1^p = a_2^p = b^{p^m} = x^{p^2} = 1, [a_1, b] = a_2, [a_2, b] = x^p, [a_2, a_1] = [x, a_1] = [x, b] = 1 \rangle = \langle a_1, b \rangle * \langle x \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, b^p, x^p \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $\Phi(G) \cong C_p^2$  if  $m = 1$ ,  $G' = \langle a_2, x^p \rangle \cong C_p^2$ ,  $Z(G) = \langle x, b^p \rangle \cong C_{p^{m-1}} \times C_{p^2}$  if  $m > 1$ ,  $Z(G) \cong C_{p^2}$  if  $m = 1$ .
- (I7)  $\langle a_1, x, b; a_2 \mid a_1^p = a_2^p = b^{p^{m+1}} = x^p = 1, [a_1, b] = a_2, [a_2, b] = b^{p^m}, [a_2, a_1] = [x, a_1] = [x, b] = 1 \rangle = \langle a_1, b \rangle \times \langle x \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, b^p \rangle \cong C_{p^m} \times C_p$ ,  $G' = \langle a_2, b^{p^m} \rangle \cong C_p^2$  and  $Z(G) = \langle x, b^p \rangle \cong C_{p^m} \times C_p$ .
- (I8)  $\langle a_1, x, b; a_2 \mid a_1^{p^2} = a_2^p = b^{p^m} = x^p = 1, [a_1, b] = a_2, [a_2, b] = a_1^{\nu p}, [a_2, a_1] = [x, a_1] = [x, b] = 1 \rangle = \langle a_1, b \rangle \times \langle x \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; Moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_1^p, b^p \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $\Phi(G) \cong C_p^2$  if  $m = 1$ ,  $G' = \langle a_2, a_1^p \rangle \cong C_p^2$ ,  $Z(G) = \langle a_1^p, b^p, x \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $Z(G) \cong C_p^2$  if  $m = 1$ .
- (I9)  $\langle a_1, x, b; a_2 \mid a_1^9 = a_2^3 = x^3 = 1, b^3 = a_1^3, [a_1, b] = a_2, [a_2, b] = a_1^{-3}, [x, a_1] = [x, a_2] = 1 \rangle = \langle a_1, b \rangle \times \langle x \rangle$ ; where  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle a_2, a_1^3 \rangle \cong C_3^2$  and  $Z(G) = \langle a_1^3, x \rangle \cong C_3^2$ .
- (Iii)  $|G'| = p^3$  and  $c(G) = 3$ . In this case,  $Z(G) < \Phi(G)$ ,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p^2 + p)$  and  $\alpha_1(G) = p^3 + p^2$ .
- (I10)  $\langle a, b, c \mid a^8 = b^{2^{m+1}} = 1, c^2 = a^4 b^{2^m}, [a, b] = a^2, [c, a] = 1, [c, b] = b^{2^m} \rangle$ ; where  $|G| = 2^{m+5}$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_{2^m} \times C_4$ ,  $G' = \langle a^2, b^{2^m} \rangle \cong C_4 \times C_2$  and  $Z(G) = \langle a^4, b^2 \rangle \cong C_{2^m} \times C_2$ .
- (I11)  $\langle a, b, c; d \mid a^{p^{n+1}} = b^p = c^{p^2} = d^p = 1, [a, b] = d, [c, a] = a^{p^n}, [d, a] = c^p, [c, b] = [d, b] = [d, c] = 1 \rangle$ , where  $p > 2$ . Moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, c^p, d \rangle \cong C_{p^n} \times C_p^2$ ,  $G' = \langle a^{p^n}, c^p, d \rangle \cong C_p^3$  and  $Z(G) = \langle c^p, a^p \rangle \cong C_{p^n} \times C_p$ .

**Proof** It is easy to see that  $Z(G) \not\leq \Phi(G)$  for groups (I1)–(I9) and  $Z(G) < \Phi(G)$  for (I10)–(I11). By checking maximal subgroups of groups (I1)–(I11), we know that they are all  $\mathcal{A}_3$ -groups and are pairwise non-isomorphic. Conversely, in the following, we prove that  $G$  is one of the groups (I1)–(I11).

Let  $A$  be an abelian subgroup of index  $p$  of  $G$ , and  $B$  be a non-abelian subgroup of index  $p$  of  $G$ . Then  $A \cap B$  is an abelian subgroup of index  $p$  of  $B$ . It follows that  $Z(B) \leq A \cap B$  and hence  $Z(B) \leq Z(G)$ .

We claim that there exists a non-abelian subgroup  $B$  of index  $p$  such that  $d(B) = 3$ . If not, then all non-abelian proper subgroups of  $G$  are generated by two elements. By Lemma 2.13,  $G \in \mathcal{A}_2$ , a contradiction.

By Lemma 2.6 (2),  $B' \leq C_p^2$  and  $B' \leq Z(B) \leq Z(G)$ . If  $G' = B'$ , then we have  $\Phi(G) \leq Z(G)$ , a contradiction. Hence  $G' > B'$ . By Lemma 2.15,  $|G'| \leq p|A'||B'| = p|B'|$ . Thus  $|G'| = p|B'|$ . Let  $D$  be another non-abelian subgroup of index  $p$  of  $G$  such that  $d(D) = 3$ . Similarly we have  $D' \leq C_p^2$  and  $D' \leq Z(D) \leq Z(G)$ . If  $D' \not\leq B'$ , then  $G' = B'D' \leq Z(G)$  and hence  $\Phi(G) \leq Z(G)$ , a contradiction. Thus we have  $D' \leq B'$ . The same reason gives that  $B' \leq D'$  and hence  $B' = D'$ . By arbitrariness of  $D$ , there exists a non-abelian subgroup  $M$  of index  $p$  such that  $d(M) = 2$  and all non-abelian subgroups of  $G/B'$  are generated by two elements.

Case 1.  $|G'| = p^2$

By Lemma 2.8,  $|G : Z(G)| = p^3$ . Since  $|G : \Phi(G)| = p^3$  and  $\Phi(G) \neq Z(G)$ , there exists a non-abelian subgroup  $M$  of index  $p$  such that  $Z(G) \not\leq M$ . Let  $x \in Z(G) \setminus M$ . Then  $G = \langle M, x \rangle$  and  $G' = M'$ . If  $d(M) = 3$ , then, by Lemma 2.6 (2),  $M' \leq C_p^3$  and  $M' \leq Z(M)$ . It follows that  $G' \leq C_p^3$  and  $G' \leq Z(G)$ . By calculation we get  $\Phi(G) \leq Z(G)$ , a contradiction. Hence  $d(M) = 2$ . It follows that  $\Phi(M) = \Phi(G)$  and  $M$  is one of the groups (1)–(7) in Lemma 2.5.

Subcase 1.1.  $M$  is the group (1) in Lemma 2.5. That is,  $M = \langle a, b \mid a^8 = b^{2^m} = 1, a^b = a^{-1} \rangle$ .

Since  $Z(M) = \langle a^4, b^2 \rangle$  and  $x^2 \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^2 = a^{4i}b^{2j}$ . Thus  $(xb^{-j})^2 = a^{4i}$ . If  $(2, j) = 1$ , then  $|\langle a^2, xb^{-j} \rangle| = 8$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq 2^5$  and  $|M| \leq 2^4$ . Hence  $m = 1$ . It follows that  $x^2 = a^{4i}$ . If  $2 \mid j$ , then, replacing  $x$  with  $xb^{-j}$ , we also have  $x^2 = a^{4i}$ . Hence we get the groups (I1) for  $2 \mid i$  and (I2) for  $(2, i) = 1$ , respectively.

Subcase 1.2.  $M$  is the group (2) in Lemma 2.5. That is,  $M = \langle a, b \mid a^8 = b^{2^m} = 1, a^b = a^3 \rangle$ .

Since  $Z(M) = \langle a^4, b^2 \rangle$  and  $x^2 \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^2 = a^{4i}b^{2j}$ . Thus  $(xb^{-j})^2 = a^{4i}$ . If  $(2, j) = 1$ , then  $|\langle a^2, xb^{-j} \rangle| = 8$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq 2^5$  and  $|M| \leq 2^4$ . Hence  $m = 1$ . It follows that  $x^2 = a^{4i}$ . If  $2 \mid j$ , then, replacing  $x$  with  $xb^{-j}$ , we also have  $x^2 = a^{4i}$ . If  $x^2 = 1$ , then we get the group (I3). If  $x^2 = a^4$ , then, replacing  $a$  with  $ax$ , we get the group (I2).

Subcase 1.3.  $M$  is the group (3) in Lemma 2.5. That is,  $M = \langle a, b \mid a^8 = 1, b^{2^m} = a^4, a^b = a^{-1} \rangle$ .

Since  $Z(M) = \langle a^4, b^2 \rangle$  and  $x^2 \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^2 = a^{4i}b^{2j}$ . Thus  $(xb^{-j})^2 = a^{4i}$ . If  $(2, j) = 1$ , then  $|\langle a^2, xb^{-j} \rangle| = 8$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq 2^5$  and  $|M| \leq 2^4$ . Hence  $m = 1$ . It follows that  $x^2 = a^{4(i+1)}$ . If  $2 \mid j$ , then, replacing  $x$  with  $xb^{-j}$ , we also have  $x^2 = a^{4i}$ . If  $x^2 = 1$ , then we get the group (I4). In the following, we may assume that  $x^2 = a^4$ . If  $m = 1$ , then, replacing  $b$  with  $xb$ , we get the group (I2). If  $m \geq 2$ , then, replacing  $x$  with  $xb^{2^{m-1}}$ , we get the group (I4).



Subcase 1.4.  $M$  is the group (4) in Lemma 2.5. That is,  $M = \langle a_1, b; a_2, a_3 \mid a_1^p = a_2^p = a_3^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = 1, [a_i, a_j] = 1 \rangle$ , where  $p > 3$  for  $m = 1$ ,  $p > 2$  and  $1 \leq i, j \leq 3$ .

Since  $Z(M) = \langle a_3, b^p \rangle$  and  $x^p \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^p = a_3^i b^{jp}$ . Thus  $(xb^{-j})^p = a_3^i$ . If  $(j, p) = 1$ , then  $|\langle a_2, xb^{-j} \rangle| = p^3$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq p^5$  and  $|M| \leq p^4$ . Hence  $m = 1$ . It follows that  $x^p = a_3^i$ . If  $p \mid j$ , then, replacing  $x$  with  $xb^{-j}$ , we also have  $x^p = a_3^i$ . If  $p \mid i$ , then we get the group (I5). If  $(i, p) = 1$ , then, replacing  $x$  with  $x^{i^{-1}}$ , we get the group (I6).

Subcase 1.5.  $M$  is the group (5) in Lemma 2.5. That is,  $M = \langle a_1, b; a_2 \mid a_1^p = a_2^p = b^{p^{m+1}} = 1, [a_1, b] = a_2, [a_2, b] = b^{p^m}, [a_1, a_2] = 1 \rangle$ , where  $p > 2$ .

Since  $Z(M) = \langle b^p \rangle$  and  $x^p \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^p = b^{jp}$ . Thus  $(x^{-j^{-1}}b)^p = 1$ . If  $p \mid j$ , then, replacing  $x$  with  $x^{-j^{-1}}b$ , we get the group (I7). If  $(j, p) = 1$ , then  $|\langle a_2, x^{-j^{-1}}b \rangle| = p^3$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq p^5$  and  $|M| \leq p^4$ . Hence  $m = 1$ . By replacing  $b$  with  $x^{-j^{-1}}b$  we get the group (I6).

Subcase 1.6.  $M$  is the group (6) in Lemma 2.5. That is,  $M = \langle a_1, b; a_2 \mid a_1^{p^2} = a_2^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_1^{\nu p}, [a_1, a_2] = 1 \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .

Since  $Z(M) = \langle a_1^p, b^p \rangle$  and  $x^p \in Z(G) \cap \Phi(G) = Z(M)$ , we may assume that  $x^p = a_1^{ip} b^{jp}$ . Thus  $(x^{-j^{-1}}b)^p = a_1^{ip}$ . If  $(j, p) = 1$ , then  $|\langle a_2, x^{-j^{-1}}b \rangle| = p^3$ . Since  $G$  is an  $\mathcal{A}_3$ -group, we have  $|G| \leq p^5$  and  $|M| \leq p^4$ . Hence  $m = 1$ . It follows that  $x^p = a_1^{ip}$ . If  $p \mid j$ , then, replacing  $x$  with  $xb^{-j}$ , we also have  $x^p = a_1^{ip}$ . If  $p \mid i$ , then we get the group (I8). If  $(i, p) = 1$ , then, replacing  $a_1$  and  $x$  with  $a_1 x^{-i^{-1}}$  and  $x^{i^{-1}\nu}$  respectively, we get the group (I6).

Subcase 1.7.  $M$  is the group (7) in Lemma 2.5. That is,  $M = \langle a_1, b; a_2 \mid a_1^9 = a_2^3 = 1, b^3 = a_1^3, [a_1, b] = a_2, [a_2, b] = a_1^{-3} \rangle$ .

If  $x^3 = 1$ , then  $G$  is the group (I9). If  $x^3 \neq 1$ , then we may assume that  $a_1^3 = b^3 = x^{-3}$ . By replacing  $a_1$  and  $b$  with  $a_1 x$  and  $bx$  respectively, we get the group (I6).

Case 2.  $|G'| = p^3$

In this case,  $B' \cong C_p^2$  and  $Z(B) = \Phi(B)$ . By Lemma 2.8,  $|G : Z(G)| = p^4$ . Since  $|G : Z(B)| = p^4$  and  $Z(B) \leq Z(G)$ , we have  $Z(G) = Z(B) = \Phi(B) < \Phi(G)$ . By Lemma 2.13,  $G/B' \in \mathcal{A}_2$ . Hence  $G/B'$  is one of the groups (8)–(12) in Lemma 2.5.

Subcase 2.1.  $G/B'$  is the group (8) or (11) in Lemma 2.5. That is,  $G/B' = \langle \bar{a}, \bar{b}, \bar{c} \mid \bar{a}^4 = \bar{c}^2 = 1, \bar{b}^2 = \bar{a}^2 = [\bar{a}, \bar{b}], [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle \cong Q_8 \times C_2$  or  $G/B' = \langle \bar{a}, \bar{b}, \bar{c} \mid \bar{a}^4 = 1, \bar{b}^2 = \bar{c}^2 = \bar{a}^2 = [\bar{a}, \bar{b}], [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle \cong Q_8 * C_4$ .

In this subcase,  $\Phi(G) = G' = \langle a^2, B' \rangle$ . Since  $a^2 = b^2 z$  where  $z \in B' \leq Z(G)$ , we have  $[a^2, b] = 1$ . It follows that  $\Phi(G) \leq Z(G)$ , a contradiction.

Subcase 2.2.  $G/B'$  is the group (9) in Lemma 2.5. That is,  $G/B' = \langle \bar{a}, \bar{b}, \bar{c} \mid \bar{a}^{p^{n+1}} = \bar{b}^{p^m} = \bar{c}^p = 1, [\bar{a}, \bar{b}] = \bar{a}^{p^n}, [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle \cong M(n+1, m) \times C_p$ .

If  $o(a) = p^{n+1}$ , then, by calculation, we have  $\Phi(G) \leq Z(G)$ , a contradiction. Hence  $o(a) = p^{n+2}$ . If  $p > 2$ , then  $\langle a, b \rangle$  has no abelian subgroup of index  $p$ , a contradiction. Hence  $p = 2$ . If  $b^{2^m} \in \langle a^{2^{n+1}} \rangle$ , then  $\langle a^2, b \rangle$  is non-abelian and is of order  $2^{n+m+1}$ , which is contrary to  $G \in \mathcal{A}_3$ . Hence  $o(b) = 2^{m+1}$  and  $B' = \langle a^{2^{n+1}}, b^{2^m} \rangle$ . Since  $\langle a, b \rangle$  has an abelian subgroup of index  $p$ , by calculation we have  $n = 1$  and  $A \cap \langle a, b \rangle = \langle a, b^2 \rangle$ , where  $A$  is the abelian subgroup of index  $p$  of  $G$ . Hence  $A = \langle a, b^2, c \rangle$ . It follows that  $[c, a] = 1$ . Assume that  $[c, b] = a^{4w}b^{u2^m}$ . By replacing  $c$  with  $ca^{2w}$  we have  $[c, b] = b^{u2^m}$ . Since  $Z(G) < \Phi(G)$ , we have  $[c, b] = b^{2^m}$ . If  $c^2 \in \langle b^{2^m} \rangle$ , then  $|\langle c, b \rangle| = 2^{m+2}$ , which is contrary to  $G \in \mathcal{A}_3$ . Thus we may assume that

$$G = \langle a, b, c \mid a^8 = b^{2^{m+1}} = 1, c^2 = a^4 b^{i2^m}, [a, b] = a^2 a^{4j} b^{k2^m}, [c, a] = 1, [c, b] = b^{2^m} \rangle.$$

By replacing  $a$  with  $ac^j$  we may assume that  $[a, b] = a^2 b^{k'2^m}$ . If  $m \geq 2$ , then, replacing  $a$  and  $c$  with  $ab^{k'2^{m-1}}$  and  $cb^{(i+1)2^{m-1}}$  respectively, we have  $[a, b] = a^2$  and  $c^2 = a^4 b^{2^m}$ . Hence we get the group (I10). If  $m = 1$ , then  $(ba)^2 = b^2 a^2 [a, b] = a^4 b^{2(k'+1)}$ . Since  $G \in \mathcal{A}_3$ , we have  $|\langle a^2, ba \rangle| = 2^4$ . It follows that  $(ba)^2 = a^4 b^2$  and  $[a, b] = a^2$ . Again since  $G \in \mathcal{A}_3$ , we have  $|\langle ca^2, ba \rangle| = 2^4$ . It follows that  $(ca)^2 = b^2$  and  $c^2 = a^4 b^2$ . Hence we also get the group (I10).

Subcase 2.3.  $G/B'$  is the group (10) in Lemma 2.5. That is,  $G/B' = \langle \bar{a}, \bar{b}, \bar{c}; \bar{d} \mid \bar{a}^{p^n} = \bar{b}^{p^m} = \bar{c}^p = \bar{d}^p = 1, [\bar{a}, \bar{b}] = \bar{d}, [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle \cong M(n, m, 1) \times C_p$ , where  $n \geq m$ , and  $n \geq 2$  if  $p = 2$ .

Since  $\langle \bar{a}, \bar{b} \rangle$  is not metacyclic,  $\langle a, b \rangle$  is not metacyclic. Since  $\langle a, b \rangle$  is a two-generator  $\mathcal{A}_2$ -subgroup, by Lemma 2.6 (6) and (7) we have  $p > 2$  and  $\exp(\langle a, b \rangle) = p$ . Hence  $d^p = 1$  and we may assume  $[a, b] = d$ . If  $m \geq 2$ , then

$$|G : \langle d, a \rangle| \geq p^{m+1} \geq p^3 \text{ and } |G : \langle d, b \rangle| \geq p^{n+1} \geq p^3.$$

Since  $G \in \mathcal{A}_3$ ,  $[d, a] = [d, b] = 1$ . It follows that  $\langle a, b \rangle \in \mathcal{A}_1$ , a contradiction. Hence we have  $m = 1$ . Since the index of  $\langle b, c \rangle$  is at least  $p^{n+1}$ , we have  $[b, c] = 1$  for  $n \geq 2$ . If  $n = 1$ , then, without loss of generality, we may assume  $[b, c] = 1$ . Since  $Z(G) < \Phi(G)$ , we have  $[c, a] \neq 1$ . Hence  $A = \langle b, c, \Phi(G) \rangle$ . It follows that  $[d, b] = 1$  and hence  $[d, a] \neq 1$ . Since  $G \in \mathcal{A}_3$  and  $|G| = p^{n+5}$ , we have  $|\langle a, d \rangle| = p^{n+3}$ . Hence

$$o(a) = p^{n+1}, [d, a] \notin \langle a^{p^n} \rangle \text{ and } B' = \langle a^{p^n}, [d, a] \rangle.$$

Since  $Z(G) < \Phi(G)$ , we have  $[c, a] \notin \langle [d, a] \rangle$ . Hence we may assume  $[c, a] = a^{ip^n} [d, a]^j$  where  $(i, p) = 1$ . By replacing  $c$  with  $c^{i^{-1}} d^{-i^{-1}j}$  we have  $[c, a] = a^{p^n}$ . Hence  $|\langle c, a \rangle| = p^{n+3}$  and  $c^p \notin \langle a^{p^n} \rangle$ . It follows that  $G' = \langle d, a^{p^n}, c^p \rangle$ . Hence we may assume  $[d, a] = c^{up} a^{vp^n}$  where  $(u, p) = 1$ . Since  $|\langle d, c^u a^{vp^{n-1}} \rangle| = p^3 \leq p^{n+3}$ ,  $[d, c^u a^{vp^{n-1}}] = 1$  and hence

$n \geq 2$  or  $v = 0$ . By replacing  $b$ ,  $c$  and  $d$  with  $b^{u^{-1}}$ ,  $ca^{u^{-1}vp^{n-1}}$  and  $d^{u^{-1}}$ , respectively, we get  $[d, a] = c^p$ . We may assume  $b^p = a^{sp^n} \pmod{\langle c^p \rangle}$ . By calculation we have  $(ba^{-sp^{n-1}})^p \in \langle c^p \rangle$ . Since  $|\langle c, ba^{-sp^{n-1}} \rangle| \leq p^3$ , we have  $[c, ba^{-sp^{n-1}}] = 1$ . Assume  $(ba^{-sp^{n-1}})^p = c^{tp}$ . Since  $|\langle d, ba^{-sp^{n-1}}c^{-t} \rangle| \leq p^3$ , we have  $[d, ba^{-sp^{n-1}}c^{-t}] = 1$ . By replacing  $b$  with  $ba^{-sp^{n-1}}c^{-t}$  we have  $b^p = 1$ . Hence  $G$  is the group (I11).

Subcase 2.4.  $G/B'$  is the group (12) in Lemma 2.5. That is,  $G/B' = \langle \bar{a}, \bar{b}, \bar{c} \mid \bar{a}^{p^n} = \bar{b}^{p^m} = \bar{c}^{p^2} = 1, [\bar{a}, \bar{b}] = \bar{c}^p, [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle \cong M(n, m, 1) * C_{p^2}$ , where  $n \geq m$ , and  $n \geq 2$  if  $p = 2$ .

Since  $\langle \bar{a}, \bar{b} \rangle$  is not metacyclic,  $\langle a, b \rangle$  is not metacyclic. Since  $\langle a, b \rangle$  is a two-generator  $\mathcal{A}_2$ -group, by Lemma 2.6 (6) and (7) we have  $p > 2$  and  $\exp(\langle a, b \rangle) = p$ . Hence  $c^{p^2} = 1$  and  $\exp(G') = p$ . It follows that  $[c^p, a] = [c^p, b] = 1$ . Thus  $\langle a, b \rangle \in \mathcal{A}_1$ , a contradiction.

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 5.4 as follows. It is easy to see that  $(\mu_0, \mu_1, \mu_2) = (1, 0, p^2 + p)$ . We only need to calculate  $\alpha_1(G)$ .

**Case 1.**  $G$  is one of the groups (I1)–(I4).

All maximal subgroups of  $G$  are:

$$N = \langle a, x, \Phi(G) \rangle;$$

$$N_i = \langle ba^i, x, \Phi(G) \rangle, \text{ where } i = 0, 1;$$

$$N_{ij} = \langle ax^i, bx^j, \Phi(G) \rangle, \text{ where } i, j = 0, 1.$$

It is easy to see that  $N = \langle a, x, \Phi(G) \rangle$  is the unique abelian maximal subgroup of  $G$ . Since  $|N'_i| = p$  and  $d(N_i) = 3$ , by Lemma 2.6 (7),  $\alpha_1(N_i) = p^2 = 4$ . Since  $N_{ij} = \langle ax^i, bx^j \rangle$ ,  $\alpha_1(N_{ij}) = p = 2$ .

Let  $H \in \Gamma_2(G)$ . Then  $H = \langle a^i b^j x^k, \Phi(G) \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $j \neq 0$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p \times p^2 + p^2 \times p - p \times p^2 = p^3.$$

**Case 2.**  $G$  is one of the groups (I5)–(I9).

All maximal subgroups of  $G$  are:

$$N = \langle a_1, x, \Phi(G) \rangle;$$

$$N_i = \langle ba_1^i, x, \Phi(G) \rangle, \text{ where } 0 \leq i \leq p-1;$$

$$N_{ij} = \langle a_1 x^i, b x^j, \Phi(G) \rangle, \text{ where } 0 \leq i, j \leq p-1.$$

It is easy to see that  $N = \langle a_1, x, \Phi(G) \rangle$  is the unique abelian maximal subgroup of  $G$ . Since  $|N'_i| = p$  and  $d(N_i) = 3$ , by Lemma 2.6,  $\alpha_1(N_i) = p^2$ . Since  $N_{ij} = \langle a_1 x^i, b x^j \rangle$ ,  $\alpha_1(N_{ij}) = p$ .

Let  $H \in \Gamma_2(G)$ . Then  $H = \langle a_1^i b^j x^k, \Phi(G) \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $j \neq 0$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p \times p^2 + p^2 \times p - p \times p^2 = p^3.$$

**Case 3.**  $G$  is the group (I10).

All maximal subgroups of  $G$  are:

$$N = \langle a, c, \Phi(G) \rangle;$$

$$N_i = \langle ba^i, c, \Phi(G) \rangle, \text{ where } i = 0, 1;$$

$$N_{ij} = \langle ac^i, bc^j, \Phi(G) \rangle, \text{ where } i, j = 0, 1.$$

It is easy to see that  $N = \langle a, c, \Phi(G) \rangle$  is the unique abelian maximal subgroup of  $G$ . Since  $|N'_i| = p^2 = 4$  and  $d(N_i) = 3$ , by Lemma 2.6,  $\alpha_1(N_i) = p^2 + p$ . By calculation,  $N_{ij} = \langle ac^i, bc^j \rangle$ . Hence  $\alpha_1(N_{ij}) = p$ .

Let  $H \in \Gamma_2(G)$ . Then  $H = \langle a^i b^j c^k, \Phi(G) \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $j \neq 0$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p \times (p^2 + p) + p^2 \times p - p \times p^2 = p^3 + p^2.$$

**Case 4.**  $G$  is the group (I11).

All maximal subgroups of  $G$  are:

$$N = \langle b, c, \Phi(G) \rangle;$$

$$N_i = \langle ab^i, c, \Phi(G) \rangle, \text{ where } 0 \leq i \leq p-1;$$

$$N_{ij} = \langle ac^i, bc^j, \Phi(G) \rangle, \text{ where } 0 \leq i, j \leq p-1.$$

It is easy to see that  $N = \langle b, c, \Phi(G) \rangle$  is the unique abelian maximal subgroup of  $G$ . Since  $|N'_i| = p^2$  and  $d(N_i) = 3$ , by Lemma 2.6,  $\alpha_1(N_i) = p^2 + p$ . By calculation,  $N_{ij} = \langle ac^i, bc^j \rangle$ . Hence  $\alpha_1(N_{ij}) = p$ .

Let  $H \in \Gamma_2$ . Then  $H = \langle a^i b^j c^k, \Phi(G) \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $i \neq 0$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p \times (p^2 + p) + p^2 \times p - p \times p^2 = p^3 + p^2.$$

□

**Theorem 5.5.**  $d(G) = 4$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:

(Ji)  $G' \cong C_p$  and  $c(G) = 2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (1 + p, 0, p^2 + p^3)$  and  $\alpha_1(G) = p^4$ .

(J1)  $\langle a, b, x, y \mid a^4 = x^2 = y^2 = 1, b^2 = a^2 = [a, b], [x, a] = [x, b] = [y, a] = [y, b] = [x, y] = 1 \rangle \cong Q_8 \times C_2 \times C_2$ ; where  $|G| = 2^5$ ,  $\Phi(G) = G' = \langle a^2 \rangle$  and  $Z(G) = \langle a^2, x, y \rangle \cong C_2^3$ .

(J2)  $\langle a, b, x, y \mid a^{p^{n+1}} = b^{p^m} = x^p = y^p = 1, [a, b] = a^{p^n}, [x, a] = [x, b] = [y, a] = [y, b] = [x, y] = 1 \rangle \cong M_p(n+1, m) \times C_p \times C_p$ ; where  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^n} \times C_{p^{m-1}}$  if  $m > 1$ ,  $\Phi(G) \cong C_{p^n}$  if  $m = 1$ ,  $G' = \langle a^{p^n} \rangle$ ,  $Z(G) = \langle a^p, b^p, x, y \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $Z(G) \cong C_{p^n} \times C_p^2$  if  $m = 1$ .

- (J3)  $\langle a, b, x, y; c \mid a^{p^n} = b^{p^m} = c^p = x^p = y^p = 1, [a, b] = c, [c, a] = [c, b] = [x, a] = [x, b] = [y, a] = [y, b] = [x, y] = 1 \rangle \cong M_p(n, m, 1) \times C_p \times C_p$ , where  $n \geq m$ , and  $n \geq 2$  if  $p = 2$ ; moreover,  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_p$  if  $m > 1$ ,  $\Phi(G) \cong C_{p^{n-1}} \times C_p$  if  $m = 1$  and  $n > 1$ ,  $\Phi(G) \cong C_p$  if  $m = n = 1$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle a^p, b^p, c, x, y \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_p^3$  if  $m > 1$ ,  $Z(G) \cong C_{p^{n-1}} \times C_p^3$  if  $m = 1$  and  $n > 1$ ,  $Z(G) \cong C_p^3$  if  $m = n = 1$ .
- (J4)  $\langle a, b, x, y \mid a^4 = y^2 = 1, b^2 = x^2 = a^2 = [a, b], [x, a] = [x, b] = [y, a] = [y, b] = [x, y] = 1 \rangle \cong Q_8 * C_4 \times C_2$ ; where  $|G| = 2^5$ ,  $\Phi(G) = G' = \langle a^2 \rangle$  and  $Z(G) = \langle x, y \rangle \cong C_4 \times C_2$ .
- (J5)  $\langle a, b, x, y \mid a^{p^n} = b^{p^m} = x^{p^2} = y^p = 1, [a, b] = x^p, [x, a] = [x, b] = [y, a] = [y, b] = [x, y] = 1 \rangle \cong M_p(n, m, 1) * C_{p^2} \times C_p$ , where  $n \geq 2$  if  $p = 2$  and  $n \geq m$ . Moreover,  $|G| = p^{n+m+3}$ ,  $\Phi(G) = \langle a^p, b^p, x^p \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_p$  if  $m > 1$ ,  $\Phi(G) \cong C_{p^{n-1}} \times C_p$  if  $m = 1$  and  $n > 1$ ,  $\Phi(G) \cong C_p$  if  $m = n = 1$ ,  $G' = \langle x^p \rangle$ ,  $Z(G) = \langle a^p, b^p, x, y \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_p \times C_{p^2}$  if  $m > 1$ ,  $Z(G) \cong C_{p^{n-1}} \times C_p \times C_{p^2}$  if  $m = 1$  and  $n > 1$ ,  $Z(G) \cong C_p \times C_{p^2}$  if  $m = n = 1$ .
- (Jii)  $G' \cong C_p^2$  and  $c(G) = 2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (1, 0, p + p^2 + p^3)$  and  $\alpha_1(G) = p^4 + p^3$ .
- (J6)  $K \times C_2$ , where  $K = \langle a, b, c \mid a^4 = b^4 = 1, c^2 = a^2b^2, [a, b] = b^2, [c, a] = a^2, [c, b] = 1 \rangle$ ; and  $|G| = 2^6$ ,  $\Phi(G) = G' = \langle a^2, b^2 \rangle \cong C_2^2$ ,  $Z(G) \cong C_2^3$ .
- (J7)  $K \times C_p$ , where  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^p = 1, [a, b] = a^{p^{m-1}}, [d, a] = b^p, [d, b] = 1 \rangle$ , where  $m \geq 3$  if  $p = 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^{m-1}} \times C_p$ ,  $G' = \langle a^{p^{m-1}}, b^p \rangle$ ,  $Z(G) \cong C_{p^{m-1}} \times C_p^2$ .
- (J8)  $K \times C_p$ , where  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}, [d, b] = 1 \rangle$ , where  $(j, p) = 1$ ,  $p > 2$ ,  $j$  is a fixed quadratic non-residue modulo  $p$ , and  $-4j$  is a quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^{m+5}$ ,  $\Phi(G) = \langle a^p, b^p, d^p \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $\Phi(G) \cong C_p^2$  if  $m = 1$ ,  $G' = \langle b^p, d^p \rangle$ ,  $Z(G) \cong C_{p^{m-1}} \times C_p^3$  if  $m > 1$ ,  $Z(G) \cong C_p^3$  if  $m = 1$ .
- (J9)  $K \times C_p$ , where  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}d^p, [d, b] = 1 \rangle$ , where if  $p$  is odd, then  $4j = 1 - \rho^{2r+1}$  with  $1 \leq r \leq \frac{p-1}{2}$  and  $\rho$  the smallest positive integer which is a primitive root (mod  $p$ ); if  $p = 2$ , then  $j = 1$ . Moreover,  $|G| = p^{m+5}$ ,  $\Phi(G) = \langle a^p, b^p, d^p \rangle \cong C_{p^{m-1}} \times C_p^2$  if  $m > 1$ ,  $\Phi(G) \cong C_p^2$  if  $m = 1$ ,  $G' = \langle b^p, d^p \rangle$ ,  $Z(G) \cong C_{p^{m-1}} \times C_p^3$  if  $m > 1$ ,  $Z(G) \cong C_p^3$  if  $m = 1$ .

**Proof** By Lemma 3.5, groups (J1)–(J9) are  $\mathcal{A}_3$ -groups. By checking maximal subgroups of groups (J1)–(J9), we know that they are pairwise non-isomorphic. Conversely, in the following, we prove that  $G$  is one of the groups (J1)–(J9).

By Lemma 3.17, we have  $c(G) = 2$ ,  $\Phi(G) \leq Z(G)$ , all  $\mathcal{A}_1$ -subgroups of  $G$  contain  $\Phi(G)$  and  $G' \leq C_p^3$ . Let  $S$  be an abelian subgroup of index  $p$  of  $G$  and  $t \in G \setminus A$ . By Lemma 2.8,  $G' = \langle [s, t] \mid s \in S \rangle$ .

We claim  $G' \leq C_p^2$ . Otherwise,  $G' \cong C_p^3$ . Let  $M = \langle s, t \rangle$  be an  $\mathcal{A}_1$ -subgroup of  $G$ . Since  $G' \leq M$ ,  $M$  is not metacyclic. Hence we may assume that

$$M = \langle s, t \mid s^{p^m} = t^{p^n} = r^p = 1, [s, t] = r, [r, s] = [r, t] = 1 \rangle$$

and

$$G' = \langle s^{p^{m-1}}, t^{p^{n-1}}, r \rangle.$$

By Lemma 2.8, there exists  $a \in S$  such that  $[a, t] = t^{p^{n-1}}$ . Hence  $\langle a, t \rangle$  is metacyclic, which is contrary to  $G' \leq \langle a, t \rangle$ . Hence  $G' \leq C_p^2$ .

By Lemma 2.16, there exists  $K \leq G$  such that  $K' = G'$  and  $d(K) = 3$ . Since  $K \in \mathcal{A}_2$ ,  $K$  is maximal in  $G$ . By Lemma 2.8,

$$|Z(G)| = \frac{|G|}{p|G'|} \text{ and } |Z(K)| = \frac{|K|}{p|K'|}.$$

It follows that  $Z(G) \not\leq K$ . Let  $y \in Z(G) \setminus K$ . Then  $G = \langle K, y \rangle$ . By Lemma 2.5,  $K$  is one of the groups (8)–(16) in Lemma 2.5.

Case 1.  $K$  is one of the groups (8)–(10) in Lemma 2.5. That is,  $K = \langle a, b \rangle \times \langle x \rangle$ .

Let  $L = \langle a, b, y \rangle$ . Then  $G = L \times \langle x \rangle$  and  $L$  is a group of Type (8)–(12) in Lemma 2.5. Hence we get groups (J1)–(J5).

Case 2.  $K$  is the groups (11) in Lemma 2.5. That is,  $K = \langle a, b, x \mid a^4 = 1, b^2 = x^2 = a^2 = [a, b], [x, a] = [x, b] = 1 \rangle \cong Q_8 * C_4$ .

Since  $y^2 \in \Phi(G) = \Phi(K) = \langle x^2 \rangle$ , we may assume  $y^2 = x^{2i}$ . By replacing  $y$  with  $yx^i$  we have  $y^2 = 1$ . Hence we get the group (J4).

Case 3.  $K$  is the group (12) in Lemma 2.5. That is,  $K = \langle a, b, x \mid a^{p^n} = b^{p^m} = x^{p^2} = 1, [a, b] = x^p, [x, a] = [x, b] = 1 \rangle \cong M_p(n, m, 1) * C_{p^2}$ , where  $n \geq 2$  if  $p = 2$  and  $n \geq m$ .

Let  $L = \langle a, b, y \rangle$ . Then  $G = L * \langle x \rangle$  and  $L$  is one of the groups (8)–(12) in Lemma 2.5. If  $L$  is one of the groups (8)–(11) in Lemma 2.5, then it is reduced to Case 1 or 2. Hence we may assume that  $L$  is also a group of Type (12) in Lemma 2.5. Let

$$L = \langle a, b, y \mid a^{p^n} = b^{p^m} = y^{p^2} = 1, [a, b] = y^p, [y, a] = [y, b] = 1 \rangle.$$

Since  $x^p \in G' = K'$ , we may assume that  $x^p = y^{ip}$ . By replacing  $x$  with  $xy^{-i}$  we have  $x^p = 1$ . Hence we get the group (J5).

Case 4.  $K$  is the group (13) in Lemma 2.5. That is,  $K = \langle a, b, c \mid a^4 = b^4 = 1, c^2 = a^2 b^2, [a, b] = b^2, [c, a] = a^2, [c, b] = 1 \rangle$ .

Since  $y^2 \in \Phi(G) = \Phi(K) = \langle a^2, b^2 \rangle$ , we may assume  $y^2 = a^{2i}b^{2j}$ . Since  $|\langle ya^i, b \rangle| \leq 8$ , we have  $[ya^i, b] = 1$  and hence  $y^2 = b^{2j} = c^{2j}a^{2j}$ . Since  $|\langle yc^j, a \rangle| \leq 8$ , we have  $[yc^j, a] = 1$  and hence  $y^2 = 1$ . Thus we get the group (J6).

Case 5.  $K$  is the group (14) in Lemma 2.5. That is,  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^p = 1, [a, b] = a^{p^{m-1}}, [d, a] = b^p, [d, b] = 1 \rangle$ , where  $m \geq 3$  if  $p = 2$ .

Since  $y^p \in \Phi(G) = \Phi(K) = \langle a^p, b^p \rangle$ , we may assume  $y^p = a^{ip}b^{jp}$ . Since  $|\langle yb^{-j}, a \rangle| \leq p^{m+1}$ , we have  $[yb^{-j}, a] = 1$  and hence  $y^p = a^{ip}$ . Since  $|\langle ya^{-i}, d \rangle| \leq p^3$ , we have  $[ya^{-i}, d] = 1$  and hence  $a^i \in Z(K)$ . By replacing  $y$  with  $ya^{-i}$  we have  $y^p = 1$ . Hence we get the group (J7).

Case 6.  $K$  is the group (15) in Lemma 2.5. That is,  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}, [d, b] = 1 \rangle$ , where  $(j, p) = 1$ ,  $p > 2$ ,  $j$  is a fixed quadratic non-residue modulo  $p$ , and  $-4j$  is a quadratic non-residue modulo  $p$ .

Since  $y^p \in \Phi(G) = \Phi(K) = \langle b^p, d^p \rangle$ , we may assume that  $y^p = b^{rp}d^{sp}$ . Since  $|\langle yb^{-r}, a \rangle| \leq p^{m+2}$ , we have  $[yb^{-r}, a] = 1$  and hence  $y^p = d^{sp}$ . Since  $|\langle yd^{-s}, a \rangle| \leq p^{m+2}$ , we have  $[yd^{-s}, a] = 1$  and hence  $y^p = 1$ . Thus we get the group (J8).

Case 7.  $K$  is the group (16) in Lemma 2.5. That is,  $K = \langle a, b, d \mid a^{p^m} = b^{p^2} = d^{p^2} = 1, [a, b] = d^p, [d, a] = b^{jp}d^p, [d, b] = 1 \rangle$ , where if  $p$  is odd, then  $4j = 1 - \rho^{2r+1}$  with  $1 \leq r \leq \frac{p-1}{2}$  and  $\rho$  the smallest positive integer which is a primitive root (mod  $p$ ); if  $p = 2$ , then  $j = 1$ .

Since  $y^p \in \Phi(G) = \Phi(K) = \langle b^p, d^p \rangle$ , we may assume  $y^p = b^{rp}d^{sp}$ . Since  $|\langle yb^{-r}, a \rangle| \leq p^{m+2}$ , we have  $[yb^{-r}, a] = 1$  and hence  $y^p = d^{sp}$ . Since  $|\langle yd^{-s}, a \rangle| \leq p^{m+2}$ , we have  $[yd^{-s}, a] = 1$  and hence  $y^p = 1$ . Thus we get the group (J9).

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 5.5 as follows.

Since  $d(G) = 4$  and  $\mu_1 = 0$ ,  $\mu_0 + \mu_2 = 1 + p + p^2 + p^3$ . Let  $N$  be an  $\mathcal{A}_1$ -subgroup of  $G$ . Since  $G \in \mathcal{A}_3$ ,  $N\Phi(G) \in \Gamma_1$  or  $\Gamma_2$ . If  $N\Phi(G) \in \Gamma_1$ , then  $d(G) \leq 3$ , a contradiction. Hence  $N\Phi(G) \in \Gamma_2$ . Since  $G \in \mathcal{A}_3$ ,  $N\Phi(G) \in \mathcal{A}_1$ . That is,  $N = N\Phi(G) \in \Gamma_2$ . Hence  $\sum_{H \in \Gamma_2} \alpha_1(H) = \alpha_1(G)$ . By Hall's enumeration principle,  $\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = \sum_{H \in \Gamma_1} \alpha_1(H) - p\alpha_1(G)$ . Hence  $\alpha_1(G) = \frac{1}{1+p} \sum_{H \in \Gamma_1} \alpha_1(H)$ .

**Case 1.**  $G$  is one of the groups (J1)–(J5).

Since  $Z(G)$  is of index  $p^2$ ,  $\mu_0 = 1 + p$  and hence  $\mu_2(G) = p^2 + p^3$ . Let  $H \in \Gamma_1$ . If  $H$  is not abelian, then  $|H'| = p$ . By Lemma 2.6,  $\alpha_1(H) = p^2$ . Hence

$$\alpha_1(G) = \frac{1}{1+p} \sum_{H \in \Gamma_1} \alpha_1(H) = \frac{1}{1+p} \mu_2 p^2 = p^4.$$

**Case 2.**  $G$  is one of the groups (J6)–(J9).

Let  $A$  be the unique abelian maximal subgroup of  $K$ . Then  $A \times \langle x \rangle$  is the unique abelian maximal subgroup of  $G$ . Other maximal subgroups of  $G$  are:

$N_i = M_i \times \langle x \rangle$ , where  $M_i$  are non-abelian maximal subgroups of  $K$ ;

$N_{ijk} = \langle ax^i, bx^j, dx^k \rangle$  (or  $\langle ax^i, bx^j, dx^k \rangle$  for (J6)), where  $0 \leq i, j, k \leq p-1$ .

It is easy to see that  $|N'_i| = p$ . By Lemma 2.6,  $\alpha_1(N_i) = p^2$ . Since  $N_{ijk} \cong K$ ,  $\alpha_1(N_{ijk}) = p + p^2$ . Hence

$$\alpha_1(G) = \frac{1}{1+p} \sum_{H \in \Gamma_1} \alpha_1(H) = \frac{1}{1+p} ((p+p^2)p^2 + p^3(p+p^2)) = p^3 + p^4.$$

□

## 5.2 $G$ has no abelian subgroup of index $p$

In this section assume  $G$  is an  $\mathcal{A}_3$ -group without an abelian subgroup of index  $p$  and an  $\mathcal{A}_1$ -subgroup of index  $p$  in Theorem 5.6, 5.7, 5.13, 5.15 and 5.16.

**Theorem 5.6.** *Suppose that  $G$  is a  $p$ -group all of whose maximal subgroups are  $\mathcal{A}_2$ -groups, and every maximal subgroup of  $G$  is generated by two elements. Then  $G$  is an  $\mathcal{A}_3$ -group if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Ki)  $G$  is metacyclic. In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = 1 + p + p^2$ .

(K1)  $\langle a, b \mid a^{p^{r+3}} = 1, b^{p^{r+s+t}} = a^{p^{r+s}}, [a, b] = a^{p^r} \rangle$ , where  $p > 2$  and  $r, s, t$  are non-negative integers with  $r \geq 1$  and  $r + s \geq 3$ ;  $|G| = p^{2r+s+t+3}$ ,  $c(G) = 4$  for  $r = 1$ ;  $c(G) = 3$  for  $r = 2$  and  $c(G) = 2$  for  $r > 2$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong M_p(r+2, r+s+t-1)$  if  $s \geq 3$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong M_p(r+t+2, r+s-1)$  if  $s < 3$ ; moreover,  $G' = \langle a^{p^r} \rangle \cong C_{p^3}$ ,  $Z(G) = \langle a^{p^3}, b^{p^3} \rangle \cong C_{p^r} \times C_{p^{r+t+s-3}}$  if  $s \geq 3$ ,  $Z(G) = \langle a^{p^3}, b^{p^3} \rangle \cong C_{p^{r+s-3}} \times C_{p^{r+t}}$  if  $s < 3$ .

(K2)  $\langle a, b \mid a^{2^{r+3}} = 1, b^{2^{r+s+t}} = a^{2^{r+s}}, [a, b] = a^{2^r} \rangle$ , where  $r, s, t$  are non-negative integers with  $r \geq 2$  and  $r + s \geq 3$ . Moreover,  $|G| = 2^{2r+s+t+3}$ ,  $c(G) = 3$  for  $r = 2$  and  $c(G) = 2$  for  $r > 2$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong M_2(r+2, r+s+t-1)$  if  $s \geq 3$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong M_2(r+t+2, r+s-1)$  if  $s < 3$ ;  $G' = \langle a^{2^r} \rangle \cong C_{2^3}$ ,  $Z(G) = \langle a^8, b^8 \rangle \cong C_{2^r} \times C_{2^{r+t+s-3}}$  if  $s \geq 3$ ,  $Z(G) = \langle a^{2^3}, b^{2^3} \rangle \cong C_{2^{r+s-3}} \times C_{2^{r+t}}$  if  $s < 3$ .

(Kii)  $G$  is not metacyclic. In this case,  $p > 2$ ,  $|G| = p^6$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = p + p^2$ .

(K3)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, b] = a^p c^{mp}, [c, a] = b^{\nu p} c^{np}, [a, b^p] = [a^p, b] = c^p, [c, a^p] = [c, b^p] = [c^p, a] = [c^p, b] = 1 \rangle$ , where  $\nu$  is a fixed quadratic non-residue modulo  $p$ . The parameters  $m, n$  are the smallest positive integers satisfying  $(m-1)^2 - \nu^{-1}(n+\nu)^2 \equiv r \pmod{p}$ , for  $r = 0, 1, \dots, p-1$ ; moreover,  $|G| = p^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle \cong C_p \times C_p \times C_{p^2}$  and  $Z(G) = \langle c^p \rangle \cong C_p$ .



(K4)  $\langle a, b, c, d \mid a^9 = b^9 = c^3 = d^3 = 1, [a, b] = c, [c, b] = a^3, [c, a] = b^{-3}, [a^3, b] = [a, b^3] = d, [d, a] = [d, b] = 1 \rangle$ ; where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle a^3, b^3, c, d \rangle \cong C_3^4$  and  $Z(G) = \langle d \rangle \cong C_3$ .

(K5)  $\langle a, b, c, d \mid a^9 = b^9 = c^3 = d^3 = 1, [a, b] = c, [c, b] = a^3d, [c, a] = b^{-3}d, [a^3, b] = [a, b^3] = d, [d, a] = [d, b] = 1 \rangle$ , where  $|G| = 3^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle a^3, b^3, c, d \rangle \cong C_3^4$  and  $Z(G) = \langle d \rangle \cong C_3$ .

**Proof** Since all non-abelian proper subgroups of  $G$  are generated by two elements,  $G$  is one of the groups classified in [29]. It follows from [29, Main Theorem],  $G$  is either a metacyclic group or a 3-group of maximal class or the group in [29, Theorem 5.5 & 5.6].

If  $G$  is metacyclic, then  $|G'| = p^3$  by Lemma 3.1. By using the classification of metacyclic  $p$ -groups in [28, 30] and checking their maximal subgroups, we get the groups (K1)–(K2). The details is omitted.

If  $G$  is of maximal class, then, by Lemma 2.4,  $G$  does not satisfy the hypothesis.

If  $G$  is the group in [29, Theorem 5.5], then  $G$  is the group (K3). If  $G$  is the group in [29, Theorem 5.6], then  $G$  is the group (K4)–(K5).

Since  $d(G) = 2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$ . If  $G$  is one of the groups (K1)–(K2), the  $G$  is metacyclic and  $\Phi(G) \in \mathcal{A}_1$ . Let  $H \in \Gamma_1$ . Then  $H$  is also metacyclic and  $\alpha_1(H) = 1 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = \mu_2(1 + p) - p = p^2 + p + 1.$$

If  $G$  is one of the groups (K3)–(K5), then  $\Phi(G)$  is abelian. Let  $H \in \Gamma_1$ . Then  $\alpha_1(H) = p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = \mu_2 p = p^2 + p.$$

□

**Theorem 5.7.** *Suppose that  $|G| = p^n$  and  $G$  is an  $\mathcal{A}_3$ -group. If  $G$  has a three-generator maximal subgroup  $M$  such that  $d(M) = 3$  and  $M' \not\leq Z(G)$ , then  $p \geq 5$ ,  $n = 6$ , all maximal subgroup of  $G$  are  $\mathcal{A}_2$ -groups and  $G$  is one of the following non-isomorphic groups:*

(L1)  $\langle x, m, a \mid x^{p^2} = m^{p^2} = a^{p^2} = 1, [x, m] = a, [a, x] = x^p, [a, m] = m^{-p} \rangle$ ; where  $|G| = p^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle a, x^p, m^p \rangle \cong C_p^2 \times C_{p^2}$  and  $Z(G) = \langle a^p \rangle \cong C_p$ .

(L2)  $\langle x, m, a \mid x^{p^2} = m^{p^2} = a^{p^2} = 1, [x, m] = a, [a, x] = x^p a^p, [a, m] = m^{-p} a^{vp} \rangle$ , where  $v \in F_p$ . Moreover,  $|G| = p^6$ ,  $c(G) = 4$ ,  $\Phi(G) = G' = \langle a, x^p, m^p \rangle \cong C_p^2 \times C_{p^2}$  and  $Z(G) = \langle a^p \rangle \cong C_p$ .

Moreover,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = 3p^2 + p$ .

**Proof** By Lemma 3.16, we may assume  $|M'| \leq p^2$ . If  $|M'| = p$ , then  $M' \leq Z(G)$ . Hence  $|M'| = p^2$ . It follows from Lemma 2.6(4) that  $c(M) = 2$ ,  $M' \cong C_p^2$ ,  $\Phi(M) = \mathcal{U}_1(M) = Z(M)$ ,  $M$  has a unique abelian subgroup  $A$  of index  $p$  and  $A/M'$  has type  $(p^{n-6}, p, p)$ . Since  $A$  is characteristic in  $M$  and  $M \trianglelefteq G$ , we have  $A \trianglelefteq G$ . We take  $x \in G \setminus M$  and let  $H = \langle x, M' \rangle$ . Then  $H$  is not abelian since  $M' \not\leq Z(G)$ . Moreover,  $|H'| = p$ ,  $H' \leq M'$  and  $H' \leq Z(G)$ . By Lemma 2.2 we have  $H$  is an  $\mathcal{A}_1$ -group. It follows that  $|H| \geq p^{n-2}$ . Hence  $o(xM') \geq p^{n-4}$ . We will prove that

$$(1) \Phi(G) \leq A.$$

Otherwise,  $G/A$  is cyclic,  $G = \langle x, A \rangle$  and  $M = \langle x^p, A \rangle$ . Since  $x^{p^2} \in \mathcal{U}_1(M) = Z(M)$ , we have  $x^{p^2} \in Z(G)$ . It follows that

$$|Z(G)/H'| = |Z(G)/Z(G) \cap M'| = |Z(G)M'/M'| \geq o(x^{p^2}M') \geq p^{n-6}.$$

Hence  $|Z(G)| \geq p^{n-5}$ . By Lemma 2.8,  $|G'| = |A/Z(G)| \leq p^3$ . Then  $G_3 \leq M'$ ,  $c(G) \leq 4$  and  $G_4 \leq Z(G)$ . Since  $G' \geq M' \cong C_p^2$ ,  $d(G') \geq 2$  and hence  $|\Phi(G')| \leq p$ . It follows that  $\Phi(G') \leq H'$ .

We claim that  $[x, w, x]^{(2)} \in Z(G)$  for all  $w \in G$ . Otherwise, we may assume that  $p = 2$  since  $[x, w, x]^{(2)} \in \Phi(G') \leq Z(G)$  for  $p > 2$ . In this case,  $[x, w, x, x] \neq 1$ . Let  $L = \langle [x, w], x \rangle$ . Then  $c(L) = 3$ . Since  $G$  is an  $\mathcal{A}_3$ -group,  $L$  is an  $\mathcal{A}_2$ -group. By Lemma 2.6 (6),  $L' \cong C_4$ . On the other hand,  $L' \leq G_3 \leq M'$ , a contradiction.

Since  $[x^p, w] \equiv [x, w]^p[x, w, x]^{(2)} \pmod{G_4}$  and  $\Phi(G') \leq Z(G)$ ,  $[x^p, w] \in Z(G)$ . By Lemma 2.8,  $M' = \{[x^p, a] \mid a \in A\} \leq Z(G)$ , a contradiction.

$$(2) |G| = p^6, o(xM') = p^2 \text{ and } x^p \in A \setminus M'.$$

By (1),  $x^p \in A$ . Since  $o(xM') \geq p^{n-4}$ , we have  $o(x^pM') \geq p^{n-5}$  and hence  $\exp(A/M') \geq p^{n-5}$ . On the other hand,  $A/M'$  has type  $(p^{n-6}, p, p)$ . If  $n \geq 7$ , then  $\exp(A/M') = p^{n-6}$ , a contradiction. Hence  $n = 6$ ,  $\exp(A/M') = p$ ,  $o(xM') = p^2$  and  $x^p \in A \setminus M'$ .

(3)  $G = \langle x, m; a \mid x^{p^2} = m^{p^2} = a^{p^2} = 1, [x, m] = a, [a, x] = x^p a^{up}, [a, m] = m^{-p} a^{vp} \rangle$ , where  $u, v \in F_p$ .

Let  $A = \langle x^p, a, M' \rangle$  and  $M = \langle m, A \rangle$ . Then  $M' = \langle [m, x^p] \rangle \times \langle [m, a] \rangle$ . Since  $[x^p, x, m] = 1$  and  $[x, m, x^p] = 1$ , we have  $[m, x^p, x] = 1$  by Witt's formula. Hence  $Z = \langle [m, x^p] \rangle$ . Since  $M' \not\leq Z(G)$ ,  $[m, a, x] \neq 1$ . Let  $[m, a, x] = [m, x^p]^i$ , where  $(i, p) = 1$ . By calculation we have

$$[[a, x]x^{-ip}, m] = [a, x, m][x^{-ip}, m] = [a, x, m][m, a, x] = 1.$$

It follows that  $[a, x]x^{-ip} \in Z(M) = M'$ .

We claim that  $x^{p^2} = 1$ . Otherwise,  $[a, x]^p \neq 1$ . Hence  $\langle a, x \rangle \in \mathcal{A}_2$ . By Lemma 2.6 (7),  $\langle a, x \rangle$  is metacyclic. If  $p > 2$ , then  $[a, x]^p = [a, x]^p \neq 1$ , a contradiction. If

$p = 2$ , then, by calculation, we have  $1 = [a, x^2] = [a, x]^2[a, x, x]$ . Hence  $[a, x, x] = x^4$ . By calculation,  $[a^2, x] = [a, x]^2[a, x, a] = x^4$ . Hence  $a^2 \notin Z(G)$ . It follows that  $M' = \langle a^2, x^4 \rangle$ . Since  $[a, x, x] \neq 1$ , we have  $[a, x] \equiv x^2 a^2 \pmod{Z(G)}$ . By calculation we have  $\langle a^2, ax \rangle \cong D_8$ , which is contrary to that  $G$  is an  $\mathcal{A}_3$ -group. Hence  $x^{p^2} = 1$ .

By Lemma 2.6 (4),  $\exp(A) > p$ . It follows that  $a^p \neq 1$ . Since  $[a^p, x] = [a, x]^p = 1$ ,  $Z(G) = \langle a^p \rangle$  and hence  $M' = \langle a^p, [a, m] \rangle$ .

We claim  $p > 2$ . Otherwise, since  $[a, x^2] = 1$ , we have  $[a, x, x] = 1$ . Hence we may assume  $[a, x] = x^2 a^{2u}$ . By calculation we have  $\langle [a, m], ax \rangle \cong D_8$ , which is contrary to that  $G$  is an  $\mathcal{A}_3$ -group. Hence  $p \geq 3$ .

Since  $[a, x, x] \in H'$ , we may assume  $[a, x, x] = a^{sp}$ . Since  $1 \neq [a, m, x] \in H'$ , we may assume  $[a, m, x] = a^{tp}$  where  $(t, p) = 1$ . Let  $j = -2^{-1}t^{-1}s$ . Then

$$[a, m^j x, m^j x] = [a, x, x][a, x, m]^j [a, m, x]^j = a^{(s+2jtp)} = 1.$$

By replacing  $x$  with  $xm^j$  we have  $[a, x, x] = 1$ .

Since  $[x^p, m] \neq 1$ , we have  $[x, m]^p \neq 1$  and  $[x, m^p] \neq 1$ . Hence  $M' = \langle a^p, m^p \rangle$  and we may assume  $[x, m] \equiv a^v \pmod{\langle x^p, M' \rangle}$ , where  $(v, p) = 1$ . By replacing  $m$  with  $m^{v^{-1}}$  we have  $[x, m] \equiv a \pmod{\langle x^p, M' \rangle}$ . Without loss of generality assume that  $[x, m] = a$ . Hence  $G = \langle x, m \rangle$  satisfy the following relations:

$$x^{p^2} = m^{p^2} = a^{p^2} = 1, [x, m] = a, [a, x] \equiv x^{ip} \pmod{\langle a^p \rangle}, [a, m] \equiv m^{jp} \pmod{\langle a^p \rangle}.$$

By replacing  $m$  and  $a$  with  $m^{i^{-1}}$  and  $[x, m^{i^{-1}}]$  respectively, we have  $[a, x] \equiv x^p \pmod{\langle a^p \rangle}$ . Since  $G$  is metabelian, we have  $[a, x, m] = [a, m, x]$ . It follows that  $j = -1$ . If  $p = 3$ , then  $\langle x^3, xm \rangle$  is not abelian and is of order  $3^3$ , which is contrary to that  $G$  is an  $\mathcal{A}_3$ -group. Hence  $p \geq 5$  and we may assume that

$$G = \langle x, m; a \mid x^{p^2} = m^{p^2} = a^{p^2} = 1, [x, m] = a, [a, x] = x^p a^{up}, [a, m] = m^{-p} a^{vp} \rangle,$$

where  $u, v \in F_p$ .

(4) Two groups with parameters  $(u, v)$  and  $(u', v')$  are pairwise isomorphic if and only if there exists  $t \in F_p^*$  such that  $(u', v') = (u, v) \begin{bmatrix} t & \\ & t^{-1} \end{bmatrix}$  or  $(u', v') = (v, u) \begin{bmatrix} t & \\ & t^{-1} \end{bmatrix}$ .

Suppose that  $G = \langle x, m, a \rangle$  and  $\bar{G} = \langle \bar{x}, \bar{m}, \bar{a} \rangle$  with parameters  $(u, v)$  and  $(u', v')$  respectively. Let  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Assume that

$$\bar{x}^\theta = x^{s_{11}} m^{s_{12}} a^{s_{13}} r_1, \bar{m}^\theta = x^{s_{21}} m^{s_{22}} a^{s_{23}} r_2,$$

where  $s_{ij} \in F_p$  such that  $s := s_{11}s_{22} - s_{12}s_{21} \in F_p^*$  and  $r_1, r_2 \in G_3$ .

By calculation,  $\bar{a}^\theta = [\bar{x}, \bar{m}]^\theta \equiv a^s \pmod{G_3}$ . Since  $[\bar{a}^\theta, \bar{x}^\theta] \equiv (\bar{x}^\theta)^\theta \equiv (\bar{x}^\theta)^p \pmod{G_4}$ , we have  $[a^s, x^{s_{11}} m^{s_{12}}] \equiv (x^{s_{11}} m^{s_{12}})^p \pmod{G_4}$ . Comparing indexes of  $x^p$  and  $m^p$  in two sides, we have  $ss_{11} = s_{11}$  and  $ss_{12} = -s_{12}$ . Since  $[\bar{a}^\theta, \bar{m}^\theta] = (\bar{m}^{-p})^\theta = (\bar{m}^\theta)^{-p}$ , we

have  $[a^s, x^{s_{21}}m^{s_{22}}] \equiv (x^{s_{21}}m^{s_{22}})^{-p} \pmod{G_4}$ . Comparing indexes of  $x^p$  and  $m^p$  in two sides, we have  $ss_{21} = -s_{21}$  and  $ss_{22} = s_{22}$ . It follows that

$$s_{11}s_{22} = 1, s_{12} = s_{21} = 0 \text{ or } s_{11} = s_{22} = 0, s_{12}s_{21} = 1.$$

If  $s_{11}s_{22} = 1$  and  $s_{12} = s_{21} = 0$ , then, by calculation,

$$\bar{a}^\theta = [\bar{x}, \bar{m}]^\theta = [x^{s_{11}}a^{s_{13}}, m^{s_{22}}a^{s_{23}}] \equiv ax^{-s_{23}s_{11}p}m^{-s_{13}s_{22}p} \pmod{G_4}.$$

Since

$$[\bar{a}^\theta, \bar{x}^\theta] = (\bar{x}^p\bar{a}^{u'p})^\theta = (\bar{x}^\theta)^p(\bar{a}^\theta)^{u'p},$$

we have

$$[ax^{-s_{23}s_{11}p}m^{-s_{13}s_{22}p}, x^{s_{11}}] = (x^{s_{11}}a^{s_{13}})^p a^{u'p}.$$

Comparing index of  $a^p$  in two sides, we have

$$s_{11}u = u'. \tag{1}$$

Since

$$[\bar{a}^\theta, \bar{m}^\theta] = (\bar{m}^{-p}\bar{a}^{v'p})^\theta = (\bar{m}^\theta)^{-p}(\bar{a}^\theta)^{v'p},$$

we have

$$[ax^{-s_{23}s_{11}p}, m^{s_{22}}] = (m^{s_{22}}a^{s_{23}})^{-p} a^{v'p}.$$

Comparing index of  $a^p$  in two sides, we have

$$s_{22}v = v'. \tag{2}$$

Let  $t = s_{11}$ . Then  $(u', v') = (u, v) \begin{bmatrix} t \\ t^{-1} \end{bmatrix}$ .

On the other hand, if there exists  $t \in F_p^*$  such that  $(u', v') = (u, v) \begin{bmatrix} t \\ t^{-1} \end{bmatrix}$ , then  $\theta : \bar{x} \mapsto x^t, \bar{m} \mapsto m^{t^{-1}}$  is an isomorphism from  $\bar{G}$  to  $G$ .

If  $s_{11} = s_{22} = 0$  and  $s_{12}s_{21} = 1$ , then, by calculation,

$$\bar{a}^\theta = [\bar{x}, \bar{m}]^\theta = [m^{s_{12}}a^{s_{13}}, x^{s_{21}}a^{s_{23}}] \equiv a^{-1}x^{s_{13}s_{21}p}m^{s_{12}s_{23}p} \pmod{G_4}.$$

Since

$$[\bar{a}^\theta, \bar{x}^\theta] = (\bar{x}^p\bar{a}^{u'p})^\theta = (\bar{x}^\theta)^p(\bar{a}^\theta)^{u'p},$$

we deduce that

$$[a^{-1}x^{s_{13}s_{21}p}m^{s_{12}s_{23}p}, m^{s_{12}}] = (m^{s_{12}}a^{s_{13}})^p a^{u'p}.$$

Comparing index of  $a^p$  in two sides, we have

$$-s_{12}v = u'. \tag{3}$$

Since

$$[\bar{a}^\theta, \bar{m}^\theta] = (\bar{m}^{-p} \bar{a}^{v'p})^\theta = (\bar{m}^\theta)^{-p} (\bar{a}^\theta)^{v'p},$$

we deduce that

$$[a^{-1} x^{s_{13} s_{21} p} m^{s_{12} s_{23} p}, x^{s_{21}}] = (x^{s_{21}} a^{s_{23}})^{-p} a^{v'p}.$$

Comparing index of  $a^p$  in two sides, we have

$$-s_{21}u = v'. \quad (4)$$

Let  $t = -s_{12}$ . Then  $(u', v') = (v, u) \begin{bmatrix} t \\ t^{-1} \end{bmatrix}$ .

On the other hand, if there exists  $t \in F_p^*$  such that  $(u', v') = (v, u) \begin{bmatrix} t \\ t^{-1} \end{bmatrix}$ , then  $\theta : \bar{x} \mapsto m^{-t}, \bar{m} \mapsto x^{-t^{-1}}$  is an isomorphism from  $\bar{G}$  to  $G$ .

(5)  $G$  is one of the groups in the theorem. That is, we may assume that  $u = v = 0$  or  $u = 1$ .

If  $u = v = 0$ , then  $G$  is the group (L1). If  $u \neq 0$ , then, replacing  $x$  and  $m$  with  $x^{u^{-1}}$  and  $m^u$  respectively,  $G$  is the group (L2). If  $v \neq 0$ , then, replacing  $x$  and  $m$  with  $m^{v^{-1}}$  and  $x^v$  respectively,  $G$  is the group (L2).

By (4), groups in Theorem 5.7 are pairwise non-isomorphic.

Now we calculate  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$ .

Since  $d(G) = 2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$ . All maximal subgroups of  $G$  are:

$$M = \langle a, x, \Phi(G) \rangle;$$

$$M_i = \langle a, mx^i, \Phi(G) \rangle, \text{ where } 0 \leq i \leq p-1.$$

It is easy to see that  $M = \langle a, x, m^p \rangle$  such that  $|M'| = p^2$  and  $d(M) = 3$ . Hence  $\alpha_1(M) = p^2 + p$  and  $\Phi(G)$  is the unique abelian maximal subgroup of  $M$ . Similarly,  $M_0 = \langle a, m, x^p \rangle$  and  $\alpha(M_0) = p^2 + p$ . By calculation,  $d(M_i) = 2$  for  $i \neq 0$ . Hence  $\alpha_1(M_i) = p$  for  $i \neq 0$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = 2 \times (p + p^2) + (p-1) \times p = 3p^2 + p.$$

□

**Lemma 5.8.** *Suppose that  $G(i, j) = \langle b, a_1; a_2, a_3 \mid b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = b^{ip}, [a_2, a_1] = b^{jp}, [a_3, a_1] = 1 \rangle$ , where  $i, j \in F_p^*$  and  $p \geq 5$ .*

- (1) *If  $G = G(i, j)$ , then  $|G| = p^5$ ,  $G' = \langle a_2, a_3, b^p \rangle$ ,  $G_3 = \langle a_3, b^p \rangle$ ,  $G_4 = \langle b^p \rangle$ ,  $G \in \mathcal{A}_3$ , and  $M = \langle b^p, a_1, a_2, a_3 \rangle$  is the unique three-generator maximal subgroup of  $G$ ;*
- (2)  *$G(i, j) \cong G(i', j')$  if and only if there exist  $r, s \in F_p^*$  such that  $j' = s^2 j$  and  $i' = r^2 s i$ .*

**Proof** (1) Let

$$K = \langle a_2, a_1; d \mid a_2^p = a_1^p = d^p = 1, [a_2, a_1] = d^j, [d, a_1] = [d, a_2] = 1 \rangle \times \langle a_3 \rangle.$$

We define an automorphism  $\beta$  of  $K$  as follows:

$$a_2^\beta = a_2 a_3, \quad a_1^\beta = a_1 a_2 \quad \text{and} \quad a_3^\beta = a_3 d^i.$$

Then  $o(\beta) = p$ . It follows from the cyclic extension theory that  $G = \langle K, b \rangle$  is a cyclic extension of  $K$  by  $C_p$ . Hence  $|G| = p^5$ . It is easy to verify that  $G' = \langle a_2, a_3, b^p \rangle$ ,  $G_3 = \langle a_3, b^p \rangle$ ,  $G_4 = \langle b^p \rangle$ ,  $G \in \mathcal{A}_3$ , and  $M = \langle b^p, a_1, a_2, a_3 \rangle$  is the unique three-generator maximal subgroup of  $G$ .

(2) For convenience, let  $G = \langle b, a_1, a_2, a_3 \rangle \cong G(i, j)$ ,  $\bar{G} = \langle \bar{b}, \bar{a}_1, \bar{a}_2, \bar{a}_3 \rangle \cong G(i', j')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle b^p, a_1, a_2, a_3 \rangle$  and  $\bar{M} = \langle \bar{b}^p, \bar{a}_1, \bar{a}_2, \bar{a}_3 \rangle$  are the unique three-generator maximal subgroups of  $G$  and  $\bar{G}$  respectively, we have  $\bar{M}^\theta = M$ . Hence we may assume that  $\bar{b}^\theta = b^r x$  and  $\bar{a}_1^\theta = a_1^s y$ , where  $r, s \in F_p^*$  and  $x \in M$ ,  $y \in \Phi(G)$ .

By calculation we have  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta \equiv a_2^{r^2 s} \pmod{G_3}$ ,  $\bar{a}_3^\theta = [\bar{a}_2, \bar{b}]^\theta \equiv a_3^{r^2 s} \pmod{G_4}$ .

Since  $[\bar{a}_2, \bar{a}_1^\theta] = (\bar{b}^{j'p})^\theta = (\bar{b}^\theta)^{j'p}$ , we have  $[a_2^{r^2 s}, a_1^s] = b^{r^2 j'p}$ . Left side of the equation is  $b^{r^2 s^2 j'p}$ . By comparing index of  $b^p$  in two sides we have  $j' = s^2 j$ .

Since  $[\bar{a}_3, \bar{b}^\theta] = (\bar{b}^{i'p})^\theta = (\bar{b}^\theta)^{i'p}$ , we have  $[a_3^{r^2 s}, b^r] = b^{r i'p}$ . Left side of the equation is  $b^{r^3 s i'p}$ . By comparing index of  $b^p$  in two sides we have  $i' = r^2 s i$ .

On the other hand, if there exist  $r, s \in F_p^*$  such that  $j' = s^2 j$  and  $i' = r^2 s i$ , then,  $\theta : \bar{a}_1 \rightarrow a_1^s, \bar{b} \rightarrow b^r$  is an isomorphism from  $\bar{G}$  to  $G$ .  $\square$

**Lemma 5.9.** *Suppose that  $G(i, j) = \langle b, a_1; a_2, a_3 \mid b^p = a_1^{p^2} = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = a_1^{ip}, [a_2, a_1] = a_1^{jp}, [a_3, a_1] = 1 \rangle$ , where  $i, j \in F_p^*$  and  $p \geq 5$ .*

(1) *If  $G = G(i, j)$ , then  $|G| = p^5$ ,  $G' = \langle a_2, a_3, a_1^p \rangle$ ,  $G_3 = \langle a_3, a_1^p \rangle$ ,  $G_4 = \langle a_1^p \rangle$ ,  $G \in \mathcal{A}_3$ , and  $M = \langle a_1, a_2, a_3 \rangle$  is the unique three-generator maximal subgroup of  $G$ ;*

(2)  *$G(i, j) \cong G(i', j')$  if and only if there exist  $r, s \in F_p^*$  such that  $j' = rsj$  and  $i' = r^3 i$ .*

**Proof** (1) Let

$$K = \langle a_2, a_1 \mid a_2^p = a_1^{p^2} = 1, [a_2, a_1] = a_1^{ip} \rangle \times \langle a_3 \rangle.$$

We define an automorphism  $\beta$  of  $K$  as follows:

$$a_2^\beta = a_2 a_3, \quad a_1^\beta = a_1 a_2 \quad \text{and} \quad a_3^\beta = a_3 a_1^{ip}.$$

Then  $o(\beta) = p$ . Hence  $G = K \rtimes \langle b \rangle$  and  $|G| = p^5$ . It is easy to verify that  $G' = \langle a_2, a_3, a_1^p \rangle$ ,  $G_3 = \langle a_3, a_1^p \rangle$ ,  $G_4 = \langle a_1^p \rangle$ ,  $G \in \mathcal{A}_3$ , and  $M = \langle a_1, a_2, a_3 \rangle$  is the unique three-generator maximal subgroup of  $G$ .

(2) For convenience, let  $G = \langle b, a_1, a_2, a_3 \rangle \cong G(i, j)$ ,  $\bar{G} = \langle \bar{b}, \bar{a}_1, \bar{a}_2, \bar{a}_3 \rangle \cong G(i', j')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle a_1, a_2, a_3 \rangle$  and  $\bar{M} = \langle \bar{a}_1, \bar{a}_2, \bar{a}_3 \rangle$  are the unique three-generator maximal subgroups of  $G$  and  $\bar{G}$  respectively, we have  $\bar{M}^\theta = M$ . Hence we may assume that  $\bar{b}^\theta = b^r x$ ,  $\bar{a}_1^\theta = a_1^s y$ , where  $r, s \in F_p^*$  and  $x \in \Omega_1(M)$ ,  $y \in \Phi(G)$ .

By calculation we have  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta \equiv a_2^{rs} \pmod{G_3}$ .  $\bar{a}_3^\theta = [\bar{a}_2, \bar{b}]^\theta \equiv a_3^{r^2s} \pmod{G_4}$ .

Since  $[\bar{a}_2^\theta, \bar{a}_1^\theta] = (\bar{a}_1^{j'p})^\theta = (\bar{a}_1^\theta)^{j'p}$ , we have  $[a_2^{rs}, a_1^s] = a_1^{sj'p}$ . Left side of the equation is  $a_1^{rs^2jp}$ . By comparing index of  $a_1^p$  in two sides we have  $j' = rsj$ .

Since  $[\bar{a}_3^\theta, \bar{b}^\theta] = (\bar{a}_1^{i'p})^\theta = (\bar{a}_1^\theta)^{i'p}$ , we have  $[a_3^{r^2s}, b^r] = a_1^{si'p}$ . Left side of the equation is  $a_1^{r^3sip}$ . By comparing index of  $a_1^p$  in two sides we have  $i' = r^3i$ .

On the other hand, if there exist  $r, s \in F_p^*$  such that  $j' = rsj$  and  $i' = r^3i$ , then  $\theta : \bar{a}_1 \rightarrow a_1^s, \bar{b} \rightarrow b^r$  is an isomorphism from  $\bar{G}$  to  $G$ .  $\square$

**Lemma 5.10.** *Suppose that  $p \geq 3$  and  $G = G(r, s) = \langle b, a_1, a_2 \mid b^{p^3} = a_1^{p^2} = a_2^{p^2} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2} a_2^{rp}, [a_2, b] = a_1^{\nu_2 p} a_2^{sp} \rangle$ , where  $r, s \in F_p$  and  $\nu_1, \nu_2 = 1$  or a fixed quadratic non-residue modula  $p$ . Then*

(1)  $|G| = p^7$ ,  $G' = \langle a_1^p, a_2, b^{p^2} \rangle$ ,  $G_3 = \langle a_1^p, a_2^p, b^{p^2} \rangle$ ,  $G_4 = \langle a_2^p \rangle$ ,  $Z(G) \cap G' = \langle a_2^p, b^{p^2} \rangle$  and  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ ;

(2)  $G$  is one of the following non-isomorphic groups:

- (i)  $\langle b, a_1, a_2 \mid b^{p^3} = a_1^{p^2} = a_2^{p^2} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2}, [a_2, b] = a_1^{\nu_2 p} a_2^{sp} \rangle$ , where  $\nu_1, \nu_2 = 1$  or a fixed quadratic non-residue modula  $p$  and  $s = \nu_2, \nu_2 + 1, \dots, \nu_2 + \frac{p-1}{2}$ ;
- (ii)  $\langle b, a_1, a_2 \mid b^{p^3} = a_1^{p^2} = a_2^{p^2} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2} a_2^{rp}, [a_2, b] = a_1^{\nu_2 p} \rangle$ , where  $\nu_1, \nu_2 = 1$  or a fixed quadratic non-residue modula  $p$  and  $r = 1, 2, \dots, \frac{p-1}{2}$ .

(3)  $[a_1^p, b] = a_2^p$ ;

(4) all maximal subgroups of  $G$ , except  $M$ , are  $\mathcal{A}_2$ -groups;

(5) If  $p = 3$  and  $\nu_2 = -1$ , then  $|M'| = 3$  and  $G \in \mathcal{A}_3$ ;

(6) If  $p = 3$  and  $\nu_2 = 1$ , then  $|M'| = 9$ , and  $G \in \mathcal{A}_3$  if and only if  $r^2 + 4\nu_1$  is not a square;

(7) If  $p \geq 5$ , then  $|M'| = p^2$ , and  $G \in \mathcal{A}_3$  if and only if  $r^2 - 4\nu_1$  is not a square.

**Proof** (1) Let

$$K = \langle a_2, a_1, d \mid a_2^{p^2} = a_1^{p^2} = d^p = 1, [a_2, a_1] = d \rangle.$$

We define an automorphism  $\beta$  of  $K$  as follows:

$$a_2^\beta = a_2 a_1^{\nu_2 p} a_2^{sp}, \quad a_1^\beta = a_1 a_2.$$

Then

$$a_2^{\beta p} = a_2, a_1^{\beta p^2} = 1, o(\beta) \leq p^2.$$

It follows from the cyclic extension theory that  $G = \langle K, b \rangle$  is a cyclic extension of  $K$  by  $C_{p^2}$ . Hence  $|G| = p^7$ . It is easy to check that  $G' = \langle a_1^p, a_2, b^{p^2} \rangle$ ,  $G_3 = \langle a_1^p, a_2^p, b^{p^2} \rangle$ ,  $G_4 = \langle a_2^p \rangle$ ,  $Z(G) \cap G' = \langle a_2^p, b^{p^2} \rangle$  and  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ .

(2) For convenience, let  $G = \langle b, a_1, a_2 \rangle \cong G(r, s)$ ,  $\bar{G} = \langle \bar{b}, \bar{a}_1, \bar{a}_2 \rangle \cong G(r', s')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle b^p, a_1, a_2 \rangle$  and  $\bar{M} = \langle \bar{b}^p, \bar{a}_1, \bar{a}_2 \rangle$  are the unique three-generator maximal subgroups of  $G$  and  $\bar{G}$  respectively, we have  $\bar{M}^\theta = M$ . Hence we may assume that  $\bar{b}^\theta = b^l a_1^m x$ ,  $\bar{a}_1^\theta = a_1^i a_2^j b^{kp} y$ , where  $l, i \in F_p^*$  and  $x \in \Phi(G)$ ,  $y \in \mathcal{U}_1(M)$ .

By calculation,  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta \equiv a_2^{il} \pmod{G_3}$ .

Since  $[\bar{a}_2^\theta, \bar{a}_1^\theta] = (\bar{b}^{\nu_1 p^2} \bar{a}_2^{r' p})^\theta = (\bar{b}^\theta)^{\nu_1 p^2} (\bar{a}_2^\theta)^{r' p}$ , we have  $[a_2^{il}, a_1^i] = b^{\nu_1 p^2} a_2^{il r' p}$ . Left side of above equation is  $b^{i^2 \nu_1 p^2} a_2^{i^2 l r p}$ . Comparing indexes of  $b^{p^2}$  and  $a_2^p$  in two sides, we have  $i^2 = 1$  and  $r' = ir$ .

Since  $[\bar{a}_2^\theta, \bar{b}^\theta, \bar{b}^\theta] = (\bar{a}_2^{\nu_2 p})^\theta = (\bar{a}_2^\theta)^{\nu_2 p}$ , we have  $[a_2^{il}, b^l, b^l] = a_2^{il \nu_2 p}$ . Left side of above equation is  $a_2^{i^3 \nu_2 p}$ . Comparing index of  $a_2^p$  in two sides, we have  $l^2 = 1$ .

By calculation we have

$$\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta = [a_1^i a_2^j b^{kp} y, b^l a_1^m] \equiv [a_1^i a_2^j, b^l] \pmod{Z(G) \cap G'}.$$

Moreover, by calculation we have

$$\bar{a}_2^\theta \equiv a_2^i a_1^{j \nu_2 p} \pmod{Z(G) \cap G'} \text{ for } l = 1$$

and

$$\bar{a}_2^\theta \equiv a_2^{-i} a_1^{(i-j) \nu_2 p} \pmod{Z(G) \cap G'} \text{ for } l = -1.$$

We have  $[\bar{a}_2^\theta, \bar{b}^\theta] = (\bar{a}_1^{\nu_2 p} \bar{a}_2^{s' p})^\theta = (\bar{a}_1^\theta)^{\nu_2 p} (\bar{a}_2^\theta)^{s' p}$ .

If  $l = 1$ , then we have  $[a_2^i a_1^{j \nu_2 p}, b a_1^m] = (a_1^i a_2^j b^{kp})^{\nu_2 p} a_2^{i s' p}$ . Left side of above equation is  $b^{i m \nu_1 p^2} a_2^{i m r p} a_1^{i \nu_2 p} a_2^{i s p} a_2^{j \nu_2 p}$ . Comparing indexes of  $b^{p^2}$ ,  $a_1^p$  and  $a_2^p$  in two sides, we have  $s' = s + i k \nu_2 \nu_1^{-1} r$ .

If  $l = -1$ , then we have  $[a_2^{-i} a_1^{(i-j) \nu_2 p}, b^{-1} a_1^m] = (a_1^i a_2^j b^{kp})^{\nu_2 p} a_2^{-i s' p}$ . Left side of above equation is  $b^{-i m \nu_1 p^2} a_2^{-i m r p} a_1^{i \nu_2 p} a_2^{i s p} a_2^{-i \nu_2 p} a_2^{(j-i) \nu_2 p}$ . Comparing indexes of  $b^{p^2}$ ,  $a_1^p$  and  $a_2^p$  in two sides, we have  $s' + s = 2 \nu_2 + i k \nu_2 \nu_1^{-1} r$ .

On the other hand, if there exists  $i, k$  such that  $i^2 = 1$ ,  $r' = ir$  and  $s' + s = 2 \nu_2 + i k \nu_2 \nu_1^{-1} r$ , then,  $\theta : \bar{a}_1 \rightarrow a_1^i b^{kp}$ ,  $\bar{b} \rightarrow b^{-1} a_1^{-i k \nu_2 \nu_1^{-1}}$  is an isomorphism from  $\bar{G}$  to  $G$ .

By above argument, if  $r = 0$ , then  $G$  is a group of Type (i), and if  $r \neq 0$ , then  $G$  is a group of Type (ii).

(3) By calculation we have  $[a_1^p, b] = [a_1, b]^p [a_1, b, a_1]^{\binom{p}{2}} [a_1, b, a_1, a_1]^{\binom{p}{3}} = a_2^p$ .



(4) Let  $N$  be a maximal subgroup of  $G$  such that  $N \neq M$ . Then we may assume  $N = \langle ba_1^i, \Phi(G) \rangle$ . By calculation we have  $[a_2, ba_1^i, ba_1^i] = [a_1^{\nu_1 2^p}, ba_1^i] = a_2^{\nu_1 2^p}$ . Hence  $N = \langle ba_1^i, a_2 \rangle$  is isomorphic to a group of Type (6) of Lemma 2.5. Thus  $N \in \mathcal{A}_2$ .

(5) By calculation,  $[a_1, b^3] = [a_1, b]^3 [a_1, b, b]^3 [a_1, b, b, b] = a_2^3 [a_1^{-3}, b] = 1$ . Since  $M' = \langle [a_1, b^3], [a_2, a_1] \rangle$ ,  $|M'| = 3$ . Moreover,  $M = \langle a_2, a_1 \rangle * \langle b^{3\nu_1} a_2^r \rangle$ . By Lemma 3.5 (2) we get  $M \in \mathcal{A}_2$ . By (4) we get  $G \in \mathcal{A}_3$ .

(6) By calculation,  $[a_1, b^3] = [a_1, b]^3 [a_1, b, b]^3 [a_1, b, b, b] = a_2^3 [a_1^3, b] = a_2^{-3}$ . Since  $M' = \langle [a_1, b^3], [a_2, a_1] \rangle$ ,  $|M'| = 9$ . By (4) we get  $G \in \mathcal{A}_3$  if and only if  $M \in \mathcal{A}_2$ . Moreover, we claim that  $M \in \mathcal{A}_2$  if and only if  $r^2 + 4\nu_1$  is not a square.

If  $M \in \mathcal{A}_2$ , then  $\langle a_2 a_1^x, b^3 a_1^y, \Phi(M) \rangle$  is either abelian or minimal non-abelian, where  $x, y \in F_3$ . For  $x \neq 0$  or  $y \neq 0$ , since  $[a_2 a_1^x, b^3 a_1^y] = b^{9y\nu_1} a_2^{3yr} a_2^{-3x} \neq 1$ , we have  $\langle a_2 a_1^x, b^3 a_1^y \rangle$  is minimal non-abelian. Hence  $\Phi(M) \leq \langle a_2 a_1^x, b^3 a_1^y \rangle$ . Moreover,  $\Phi(M) =$

$$\langle a_1^{3x} a_2^3, a_1^{3y} b^9, a_2^{3(yr-x)} b^{9y\nu_1} \rangle. \text{ It follows that the equation } \begin{vmatrix} x & 1 & 0 \\ y & 0 & 1 \\ 0 & yr-x & y\nu_1 \end{vmatrix} = 0 \text{ only}$$

has solution  $x = y = 0$ . That is, the equation  $x^2 - xyr - y^2\nu_1 = 0$  only has solution  $x = y = 0$ . Hence  $r^2 + 4\nu_1$  is not a square. Conversely, if  $r^2 + 4\nu_1$  is not a square, then, by above argument,  $\langle a_2 a_1^x, b^3 a_1^y, \Phi(M) \rangle$  is either abelian or minimal non-abelian. It is easy to see that  $\langle a_1, b^3, \Phi(M) \rangle$  and  $\langle a_1, a_2 b^{3z}, \Phi(M) \rangle$  are minimal non-abelian. Hence all maximal subgroups of  $M$  are abelian or minimal non-abelian. That is,  $M \in \mathcal{A}_2$ .

(7) By calculation,  $[a_1, b^p] = [a_1, b]^p = a_2^p$ . Since  $M' = \langle [a_1, b^p], [a_2, a_1] \rangle$ ,  $|M'| = p^2$ . By (4) we get  $G \in \mathcal{A}_3$  if and only if  $M \in \mathcal{A}_2$ . Moreover, we claim that  $M \in \mathcal{A}_2$  if and only if  $r^2 - 4\nu_1$  is not a square.

If  $M \in \mathcal{A}_2$ , then  $\langle a_2 a_1^x, b^p a_1^y, \Phi(M) \rangle$  is either abelian or minimal non-abelian, where  $x, y \in F_p$ . For  $x \neq 0$  or  $y \neq 0$ , since  $[a_2 a_1^x, b^p a_1^y] = b^{y\nu_1 p^2} a_2^{yrp} a_2^{xp} \neq 1$ , we have  $\langle a_2 a_1^x, b^p a_1^y \rangle$  is minimal non-abelian. Hence  $\Phi(M) \leq \langle a_2 a_1^x, b^p a_1^y \rangle$ . Moreover,  $\Phi(M) =$

$$\langle a_1^{xp} a_2^p, a_1^{yp} b^{p^2}, a_2^{(yr+x)p} b^{y\nu_1 p} \rangle. \text{ It follows that the equation } \begin{vmatrix} x & 1 & 0 \\ y & 0 & 1 \\ 0 & yr+x & y\nu_1 \end{vmatrix} = 0 \text{ only}$$

has solution  $x = y = 0$ . That is, the equation  $x^2 + xyr + y^2\nu_1 = 0$  only has solution  $x = y = 0$ . Hence  $r^2 - 4\nu_1$  is not a square. Conversely, if  $r^2 - 4\nu_1$  is not a square, then, by above argument,  $\langle a_2 a_1^x, b^p a_1^y, \Phi(M) \rangle$  is either abelian or minimal non-abelian. It is easy to see that  $\langle a_1, b^p, \Phi(M) \rangle$  and  $\langle a_1, a_2 b^{zp}, \Phi(M) \rangle$  are minimal non-abelian. Hence all maximal subgroups of  $M$  are abelian or minimal non-abelian. That is,  $M \in \mathcal{A}_2$ .  $\square$

**Lemma 5.11.** *Suppose that  $G(\nu, t) = \langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^2} = a_2^{tp}, [a_1, b] = a_2, [a_2, a_1] = a_2^p, [a_2, b] = a_1^{\nu p} \rangle$ , where  $t \in F_p$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .*

- (1) *If  $G = G(\nu, t)$ , then  $|G| = p^6$ ,  $G' = \langle a_1^p, a_2 \rangle$ ,  $G_3 = \langle a_1^p, a_2^p \rangle$ ,  $G_4 = \langle a_2^p \rangle$ ,  $G \in \mathcal{A}_3$ , and  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ ;*

(2)  $G(\nu, t) \cong G(\nu', t')$  if and only if  $(\nu', t') = (\nu, t)$ .

**Proof** (1) Let

$$K = \langle a_2, a_1 \mid a_2^{p^2} = a_1^{p^2} = 1, [a_2, a_1] = a_2^p \rangle.$$

We define an automorphism  $\beta$  of  $K$  as follows:

$$a_2^\beta = a_2 a_1^{\nu p}, \quad a_1^\beta = a_1 a_2.$$

Then  $a_2^{\beta p} = a_2$ ,  $a_1^{\beta p^2} = a_1$ ,  $o(\beta) \leq p^2$ . It follows from the cyclic extension theory that  $G = \langle K, b \rangle$  is a cyclic extension of  $K$  by  $C_{p^2}$ . Hence  $|G| = p^6$ . It is easy to check that  $G' = \langle a_1^p, a_2 \rangle$ ,  $G_3 = \langle a_1^p, a_2^p \rangle$ ,  $G_4 = \langle a_2^p \rangle$  and  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ .

(2) For convenience, let  $G = \langle b, a_1, a_2 \rangle \cong G(\nu, t)$ ,  $\bar{G} = \langle \bar{b}, \bar{a}_1, \bar{a}_2 \rangle \cong G(\nu', t')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle b^p, a_1, a_2 \rangle$  and  $\bar{M} = \langle \bar{b}^p, \bar{a}_1, \bar{a}_2 \rangle$  are the unique three-generator maximal subgroups of  $G$  and  $\bar{G}$  respectively, we have  $\bar{M}^\theta = M$ . Hence we may assume that  $\bar{b}^\theta = b^l x$  and  $\bar{a}_1^\theta = a_1^i y$ , where  $l, i \in F_p^*$  and  $x \in M$ ,  $y \in \Phi(G)$ .

By calculation,  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta \equiv a_2^{il} \pmod{G_3}$ .

Since  $[\bar{a}_2^\theta, \bar{a}_1^\theta] = (\bar{a}_2^\theta)^\theta = (\bar{a}_2^\theta)^p$ , we have  $[a_2^{il}, a_1^i] = a_2^{ilp}$ . Left side of above equation is  $a_2^{i^2 lp}$ . Comparing index of  $a_2^p$  in two sides, we have  $i = 1$ .

Since  $(\bar{b}^\theta)^{p^2} = (\bar{b}^{p^2})^\theta = (\bar{a}_2^{t'p})^\theta = (\bar{a}_2^\theta)^{t'p}$ , we have  $(b^l)^{p^2} = a_2^{t'lp}$ . Left side of above equation is  $a_2^{ltp}$ . Comparing index of  $a_2^p$  in two sides, we have  $t' = t$ .

Since  $[\bar{a}_2^\theta, \bar{b}^\theta, \bar{b}^\theta] = (\bar{a}_2^{\nu'p})^\theta = (\bar{a}_2^\theta)^{\nu'p}$ , we have  $[a_2^l, b^l, b^l] = a_2^{\nu'lp}$ . Left side of above equation is  $a_2^{l^3 \nu p}$ . Comparing index of  $a_2^p$  in two sides, we have  $\nu' = \nu$ .  $\square$

**Lemma 5.12.** *Suppose that  $G(\nu, s, t) = \langle b, a_1, a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^m} = a_2^{tp}, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} a_2^{sp} \rangle$ , where  $m \geq 2$ ,  $t \in F_p$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .*

(1) *If  $G = G(\nu, s, t)$ , then  $|G| = p^{m+4}$ ,  $G' = \langle a_1^p, a_2 \rangle$ ,  $G_3 = \langle a_1^p, a_2^p \rangle$ ,  $G_4 = \langle a_2^p \rangle$ ,  $G \in \mathcal{A}_3$ ,  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ ;*

(2)  *$G$  is one of the following non-isomorphic groups:*

(i)  $\langle b, a_1, a_2 \mid a_1^{p^2} = a_2^{p^2} = b^{p^m} = 1, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} a_2^{sp} \rangle$ , where  $m \geq 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ,  $s = \nu, \nu + 1, \dots, \nu + \frac{p-1}{2}$ ;

(ii)  $\langle b, a_1, a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^m} = a_2^p, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} \rangle$ , where  $m \geq 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .

(3) *If  $p = 3$  and  $\nu = -1$ , then  $M$  is abelian. If  $\nu = 1$  or  $p \geq 5$ , then  $|M'| = p$ .*

**Proof** (1) Let

$$K = \langle a_2, a_1 \mid a_2^{p^2} = a_1^{p^2} = 1, [a_2, a_1] = a_2^p \rangle.$$

We define an automorphism  $\beta$  of  $K$  as follows:

$$a_2^\beta = a_2 a_1^{\nu p} a_2^{s p}, \quad a_1^\beta = a_1 a_2.$$

Then  $a_2^{\beta^p} = a_2$ ,  $a_1^{\beta^{p^2}} = a_1$ ,  $o(\beta) \leq p^2$ . It follows from the cyclic extension theory that  $G = \langle K, b \rangle$  is a cyclic extension of  $K$  by  $C_{p^m}$ . Hence  $|G| = p^{m+4}$ . It is easy to verify that  $G' = \langle a_1^p, a_2 \rangle$ ,  $G_3 = \langle a_1^p, a_2^p \rangle$ ,  $G_4 = \langle a_2^p \rangle$ ,  $G \in \mathcal{A}_3$  and  $M = \langle b^p, a_1, a_2 \rangle$  is the unique three-generator maximal subgroup of  $G$ .

(2) For convenience, let  $G = \langle b, a_1, a_2 \rangle \cong G(\nu, s, t)$ ,  $\bar{G} = \langle \bar{b}, \bar{a}_1, \bar{a}_2 \rangle \cong G(\nu', s', t')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle b^p, a_1, a_2 \rangle$  and  $\bar{M} = \langle \bar{b}^p, \bar{a}_1, \bar{a}_2 \rangle$  are the unique three-generator maximal subgroups of  $G$  and  $\bar{G}$  respectively, we have  $\bar{M}^\theta = M$ . Hence we may assume that  $\bar{b}^\theta = b^l x$  and  $\bar{a}_1^\theta = a_1^i a_2^j b^{k p^{m-1}} y$ , where  $l, i \in F_p^*$  and  $x \in M, y \in G_3$ .

By calculation we have  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta \equiv a_2^{il} \pmod{G_3}$ .

Since  $(\bar{b}^\theta)^{p^m} = (\bar{b}^{p^m})^\theta = (\bar{a}_2^{t'p})^\theta = (\bar{a}_2^\theta)^{t'p}$ , we have  $(b^l x)^{p^m} = a_2^{t'ip}$ . Left side of the equation is  $a_2^{ltp}$ . By comparing index of  $a_2^3$  in two sides we have  $t'i = t$ .

Since  $[\bar{a}_2^\theta, \bar{b}^\theta, \bar{b}^\theta] = (\bar{a}_2^{\nu'p})^\theta = (\bar{a}_2^\theta)^{\nu'p}$ , we have  $[a_2^l, b^l, b^l] = a_2^{\nu'lp}$ . Left side of the equation is  $a_2^{l^3 \nu p}$ . By comparing index of  $a_2^2$  in two sides we have  $\nu' = \nu$  and  $l^2 = 1$ .

By calculation we have  $\bar{a}_2^\theta = [\bar{a}_1, \bar{b}]^\theta = [a_1^i a_2^j b^{k p^{m-1}} y, b^l] \equiv [a_1^i a_2^j, b^l] \pmod{G_4}$ . Moreover, by calculation we have  $\bar{a}_2^\theta \equiv a_2^i a_1^{j \nu p} \pmod{G_4}$  for  $l = 1$  and  $\bar{a}_2^\theta \equiv a_2^{-i} a_1^{(i-j) \nu p} \pmod{G_4}$  for  $l = -1$ .

If  $l = 1$ , then

$$[a_2^i a_1^{j \nu p}, b] = [\bar{a}_2^\theta, \bar{b}^\theta] = (\bar{a}_1^{\nu p} \bar{a}_2^{s'p})^\theta = (\bar{a}_1^\theta)^{\nu p} (\bar{a}_2^\theta)^{s'p} = (a_1^i a_2^j b^{k p^{m-1}})^{\nu p} a_2^{is'p}.$$

Left side of the equation is  $a_1^{i \nu p} a_2^{i s p} a_2^{j \nu p}$ . By comparing indexes of  $a_1^p$  and  $a_2^p$  in two sides we have  $s' = s - i^{-1} t k \nu$ . On the other hand, if there exists  $i, k$  such that  $t'i = t$  and  $s' = s - i^{-1} t k \nu$ , then,  $\theta: \bar{a}_1 \rightarrow a^i b^{k p^{m-1}}, \bar{b} \rightarrow b$  is an isomorphism from  $\bar{G}$  to  $G$ .

If  $l = -1$ , then we have

$$[a_2^{-i} a_1^{(i-j) \nu p}, b^{-1}] = [\bar{a}_2^\theta, \bar{b}^\theta] = (\bar{a}_1^{\nu p} \bar{a}_2^{s'p})^\theta = (\bar{a}_1^\theta)^{\nu p} (\bar{a}_2^\theta)^{s'p} = (a_1^i a_2^j b^{k p^{m-1}})^{\nu p} a_2^{-is'p}.$$

Left side of the equation is  $a_1^{i \nu p} a_2^{i s p} a_2^{-i \nu p} a_2^{(j-i) \nu p}$ . By comparing indexes of  $a_1^p$  and  $a_2^p$  in two sides we have  $s' + s = 2\nu + i^{-1} t k \nu$ . On the other hand, if there exists  $i, k$  such that  $t'i = t$  and  $s' + s = 2\nu + i^{-1} t k \nu$ , then,  $\theta: \bar{a}_1 \rightarrow a^i b^{k p^{m-1}}, \bar{b} \rightarrow b^{-1}$  is an isomorphism from  $\bar{G}$  to  $G$ .

By above argument we get that  $G(\nu, s, 0) \cong G(\nu', s', t')$  if and only if  $\nu = \nu', t' = 0$  and  $s = s'$  or  $s + s' = 2\nu$ . It follows that  $G$  is a group of Type (i) for  $t = 0$ . If  $t \neq 0$ , then, by above argument,  $G(\nu, s, t) \cong G(\nu, 0, 1)$ . Hence  $G$  is a group of Type (ii).

(3) By calculation,  $[a_1^{\nu p}, b] = a_2^{\nu p}$ . Hence

$$[a_1, b^3] = [a_1, b]^3 [a_1, b, b]^3 [a_1, b, b, b] = a_2^3 [a_1^{3\nu}, b] = a_2^{3(1+\nu)}.$$

If  $p \geq 5$ , then  $[a_1, b^p] = a_2^p$ . Since  $M' = \langle [a_1, b^p] \rangle$ , the result is proved.  $\square$

**Theorem 5.13.** *Suppose that  $G$  is an  $\mathcal{A}_3$ -group all of whose maximal subgroups are  $\mathcal{A}_2$ -groups and there exists a three-generator maximal subgroup in  $G$ , and  $M' \leq Z(G)$  for every three-generator maximal subgroup  $M$ . Then  $d(G) = 2$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Mi)  $G' \cong C_{p^2}$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = p^3 + p^2$ .

(M1)  $\langle a, b, c \mid a^8 = 1, c^2 = a^4 = b^4, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ ; where  $|G| = 2^6$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_2^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle c \rangle \cong C_4$ .

(M2)  $\langle a, b, c \mid a^8 = b^4 = 1, c^2 = a^4, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ ; moreover,  $|G| = 2^6$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_2^2$ ,  $G' = Z(G) = \langle c \rangle \cong C_4$ .

(M3)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = 1, c^p = a^{p^2}, [a, b] = c, [c, a] = 1, [c, b] = c^{tp} \rangle$ , where  $p > 2$  and  $t \in F_p^*$ ; moreover,  $|G| = p^6$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle c^p \rangle \cong C_p$ .

(M4)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = 1, c^p = a^{p^2}, [a, b] = c, [c, a] = c^p, [c, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^6$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^{p^2} \rangle \cong C_p$ .

(M5)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^n} = 1, c^p = a^{p^n}, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ , where  $n \geq 3$  for  $p = 2$  and  $n \geq 2$ ; moreover,  $|G| = p^{2n+2}$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^{n-1}} \times C_p$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-1}} \times C_{p^{n-2}} \times C_p$  if  $n > 2$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_p^2$  if  $n = 2$ .

(M6)  $\langle a, b, c \mid a^{p^2} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, a] = c^p, [c, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^6$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_p^2$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle c^p \rangle \cong C_p$ .

(M7)  $\langle a, b, c \mid a^{p^n} = b^{p^n} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ , where  $n \geq 3$  for  $p = 2$  and  $n \geq 2$ ; moreover,  $|G| = p^{2n+2}$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}}^2 \times C_{p^2}$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-2}}^2 \times C_{p^2}$  if  $n > 2$ ,  $Z(G) \cong C_{p^2}$  if  $n = 2$ .

(M8)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^2} = 1, c^p = a^{p^n}, [a, b] = c, [c, a] = c^p, [c, b] = 1 \rangle$ , where  $p > 2$  and  $n > 2$ ; moreover,  $|G| = p^{n+4}$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_p^2$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^{p^2} \rangle \cong C_{p^{n-1}}$ .

(M9)  $\langle a, b, c \mid a^{p^n} = b^{p^3} = 1, c^p = b^{p^2}, [a, b] = c, [c, a] = c^{tp}, [c, b] = 1 \rangle$ , where  $p > 2$ ,  $n > 2$  and  $t \in F_p^*$ ; moreover,  $|G| = p^{n+4}$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_p$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_{p^{n-2}} \times C_p$ .

- (M10)  $\langle a, b; c \mid a^{p^n} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, a] = c^p, [c, b] = 1 \rangle$ , where  $p > 2$  and  $n > 2$ . Moreover,  $|G| = p^{n+4}$ ,  $c(G) = 3$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_p$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_{p^{n-2}} \times C_p$ .
- (M11)  $\langle a, b; c \mid a^{p^{n+1}} = b^{p^m} = 1, c^p = a^{p^n}, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ , where  $n > m \geq 2$ ; moreover,  $|G| = p^{n+m+2}$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^{m-1}} \times C_p$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-1}} \times C_{p^{m-2}} \times C_p$  if  $m > 2$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-1}} \times C_p$  if  $m = 2$ .
- (M12)  $\langle a, b; c \mid a^{p^n} = b^{p^{m+1}} = 1, c^p = b^{p^m}, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ , where  $n > m \geq 2$ ;  $|G| = p^{n+m+2}$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^m} \times C_p$ ,  $G' = \langle c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-2}} \times C_{p^{m-1}} \times C_p$ .
- (M13)  $\langle a, b; c \mid a^{p^n} = b^{p^m} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [c, b] = 1 \rangle$ , where  $n > m \geq 2$ ; moreover,  $|G| = p^{n+m+2}$ ,  $c(G) = 2$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^{m-1}} \times C_{p^2}$ ,  $G' = \langle c \rangle$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, c \rangle \cong C_{p^{n-2}} \times C_{p^{m-2}} \times C_{p^2}$  if  $m > 2$ ,  $Z(G) \cong C_{p^{n-2}} \times C_{p^2}$  if  $m = 2$ .
- (Mii)  $c(G) = 3$  and  $G' \cong C_p^2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = p^3 + p^2$ .
- (M14)  $\langle a, b; c \mid a^8 = b^4 = c^2 = 1, [a, b] = c, [c, a] = 1, [c, b] = a^4 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_2^2$ ,  $G' = \langle a^4, c \rangle$  and  $Z(G) = \langle a^2 \rangle \cong C_4$ .
- (M15)  $\langle a, b; c, d \mid a^4 = b^4 = c^2 = d^2 = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle$ ; moreover,  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c, d \rangle \cong C_4^2$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle b^2, d \rangle \cong C_2^2$ .
- (M16)  $\langle a, b; c \mid a^8 = b^4 = c^2 = 1, [a, b] = c, [c, a] = a^4, [c, b] = 1 \rangle$ , where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_2^2$ ,  $G' = \langle a^4, c \rangle$  and  $Z(G) = \langle a^4, b^2 \rangle \cong C_2^2$ .
- (M17)  $\langle a, b; c \mid a^{2^{n+1}} = b^4 = c^2 = 1, [a, b] = c, [c, a] = a^{2^n}, [c, b] = 1 \rangle$ , where  $n > 2$ ; moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^n} \times C_2^2$ ,  $G' = \langle a^{2^n}, c \rangle$  and  $Z(G) = \langle a^4, b^2 \rangle \cong C_{2^{n-1}} \times C_2$ .
- (M18)  $\langle a, b; c \mid a^{2^n} = b^8 = c^2 = 1, [a, b] = c, [c, a] = b^4, [c, b] = 1 \rangle$ , where  $n > 2$ ; moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_{2^{n-1}} \times C_4 \times C_2$ ,  $G' = \langle b^4, c \rangle$  and  $Z(G) = \langle a^4, b^2 \rangle \cong C_{2^{n-2}} \times C_4$ .
- (M19)  $\langle a, b; c, d \mid a^{2^n} = b^4 = c^2 = d^2 = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle$ , where  $n > 2$ ; moreover,  $|G| = 2^{n+4}$ ,  $\Phi(G) = \langle a^2, b^2, c, d \rangle \cong C_{2^{n-1}} \times C_2^3$ ,  $G' = \langle c, d \rangle$  and  $Z(G) = \langle a^4, b^2, d \rangle \cong C_{2^{n-2}} \times C_2^2$ .
- (Miii)  $G_3 \cong C_p$  and  $\Phi(G')G_3 \cong C_p^2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = p^3 + 2p^2$ .
- (M20)  $\langle a, b; c \mid a^8 = b^8 = 1, c^2 = b^4, [a, b] = c, [c, a] = 1, [c, b] = a^4b^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle a^4, c \rangle$  and  $Z(G) = \langle a^4, c^2 \rangle \cong C_2^2$ .

- (M21)  $\langle a, b, c \mid a^8 = c^4 = 1, b^4 = c^2, [a, b] = c, [c, a] = a^4, [c, b] = 1 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle a^4, c \rangle$  and  $Z(G) = \langle a^4, c^2 \rangle \cong C_2^2$ .
- (M22)  $\langle a, b, c \mid a^8 = c^4 = 1, b^4 = c^2, [a, b] = c, [c, a] = 1, [c, b] = a^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle a^4, c \rangle$ ,  $Z(G) = \langle a^4, c^2 \rangle \cong C_2^2$ .
- (M23)  $\langle a, b, c \mid a^4 = b^8 = c^4 = 1, [a, b] = c, [c, a] = 1, [c, b] = b^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle b^4, c \rangle$  and  $Z(G) = \langle b^4, c^2 \rangle \cong C_2^2$ .
- (M24)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = 1, c^p = a^{p^2} b^{sp^2}, [a, b] = c, [c, a] = 1, [b, c] = b^{p^2} \rangle$ , where  $p > 2$ ,  $s \in F_p$  and  $1 + 4s \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p^2$ .
- (M25)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [c, b] = c^{tp} a^{-tp^2} \rangle$ , where  $p > 2$  and  $t^2 - 4t \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_p^2$ .
- (M26)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = 1, [a, b] = c, c^p = b^{p^2}, [c, a] = 1, [b, c] = a^{\nu p^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ , and  $-\nu \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p^2$ .
- (M27)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [b, c] = a^{\nu p^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ , and  $-\nu \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_p^2$ .
- (M28)  $\langle a, b, c \mid a^{p^2} = b^{p^3} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [c, b] = c^p b^{-p^2}, [b^{p^2}, a] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle c^p, b^{p^2} \rangle \cong C_p^2$ .
- (M29)  $\langle a, b, c \mid a^{p^2} = b^{p^3} = c^{p^2} = 1, [a, b] = c, [c, a] = 1, [b, c] = b^{p^2} \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle c^p, b^{p^2} \rangle \cong C_p^2$ .
- (M30)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^3} = 1, [a, b] = c, c^p = a^{p^n}, [a, c] = b^{\nu p^2}, [b, c] = 1 \rangle$ , where  $n > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_{p^{n-1}} \times C_p$ .
- (M31)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^3} = 1, [a, b] = c, c^p = a^{p^n} b^{s\eta p^2}, [a, c] = b^{\eta p^2}, [b, c] = 1 \rangle$ , where  $p > 2$ ,  $n > 2$ ,  $s = 1, 2, \dots, \frac{p-1}{2}$ , and  $\eta$  is a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^2} \times C_p$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_{p^{n-1}} \times C_p$ .
- (M32)  $\langle a, b, c \mid a^{p^{n+1}} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, b] = 1, [c, a] = c^p a^{-p^n}, [a^{p^n}, b] = 1 \rangle$ , where  $p > 2$  and  $n > 2$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^n}, c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_{p^{n-1}} \times C_p$ .

- (M33)  $\langle a, b; c \mid a^{p^{n+1}} = b^{p^3} = 1, [a, b] = c, c^p = b^{p^2}, [b, c] = 1, [a, c] = a^{p^n} \rangle$ , where  $n > 2$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^n}, c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_{p^{n-1}} \times C_p$ .
- (M34)  $\langle a, b; c \mid a^{p^{n+1}} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [b, c] = 1, [a, c] = a^{p^n} \rangle$ , where  $n > 2$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^n} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^n}, c \rangle$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_{p^{n-1}} \times C_p$ .
- (M35)  $\langle a, b; c \mid a^{p^n} = b^{p^3} = c^{p^2} = 1, [a, b] = c, [c, b] = 1, [c, a] = c^{tp}b^{-tp^2}, [b^{p^2}, a] = 1 \rangle$ , where  $p > 2$ ,  $n > 2$  and  $t^2 + 4t \notin F_p^2$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_{p^2}$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2}, c^p \rangle \cong C_{p^{n-2}} \times C_p \times C_p$ .
- (M36)  $\langle a, b; c \mid a^{p^n} = b^{p^3} = c^{p^2} = 1, [a, b] = c, [c, b] = 1, [a, c] = b^{\eta p^2} \rangle$ , where  $p > 2$ ,  $n > 2$ ,  $\eta$  is a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^{n+5}$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^{n-1}} \times C_{p^2} \times C_{p^2}$ ,  $G' = \langle b^{p^2}, c \rangle$  and  $Z(G) = \langle a^{p^2}, b^{p^2}, c^p \rangle \cong C_{p^{n-2}} \times C_p \times C_p$ .
- (Miv)  $\Phi(G') \leq G_3 \cong C_p^2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = p^3 + p^2$  except for (M37)–(M39).
- (M37)  $\langle a, b; c \mid a^8 = b^8 = c^2 = 1, [a, b] = c, [c, a] = a^4b^4, [c, b] = a^4, [a^4, b] = 1 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle a^4, b^4, c \rangle \cong C_2^3$ ,  $Z(G) = \langle a^4, b^4 \rangle \cong C_2^2$ , and  $\alpha_1(G) = 18$ ;
- (M38)  $\langle a, b; c \mid a^8 = b^8 = 1, [a, b] = c, [c, a] = c^2 = b^4, [c, b] = a^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4^2 \times C_2$ ,  $G' = \langle a^4, c \rangle \cong C_4 \times C_2$ ,  $Z(G) = \langle a^4, b^4 \rangle \cong C_2^2$ , and  $\alpha_1(G) = 14$ ;
- (M39)  $\langle a, b; c \mid a^8 = c^4 = 1, [a, b] = c, [c, a] = c^2, [c, b] = a^4 = b^4 \rangle$ ; where  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2, c \rangle \cong C_4 \times C_4 \times C_2$ ,  $G' = \langle a^4, c \rangle \cong C_4 \times C_2$ ,  $Z(G) = \langle a^4, c^2 \rangle \cong C_2^2$ , and  $\alpha_1(G) = 18$ ;
- (M40)  $\langle a, b; c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = a^p, [c, b] = b^p \rangle$ , where  $p > 3$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle \cong C_p^3$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_p^2$ ;
- (M41)  $\langle a, b; c \mid a^{p^3} = b^{p^3} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p^2}, [c, b] = a^{-p^2}, [a^{p^2}, b] = 1 \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$  such that  $-\nu \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, b^{p^2}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_{p^2}$ ;
- (M42)  $\langle a, b; c \mid a^{p^3} = b^{p^3} = c^p = 1, [a, b] = c, [c, a]^{1+r} = a^{p^2}b^{p^2}, [c, b]^{1+r} = a^{-rp^2}b^{p^2}, [a^{p^2}, b] = 1 \rangle$ , where  $p > 3$  and  $-r \notin (F_p)^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, b^{p^2}, c \rangle$  and  $Z(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_{p^2}$ ;
- (M43)  $\langle a, b; c, d, e \mid a^p = b^p = c^p = d^p = e^p = 1, [a, b] = c, [c, a] = d, [c, b] = e, [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle$ , where  $p > 3$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle c, d, e \rangle \cong C_p^3$  and  $Z(G) = \langle d, e \rangle \cong C_p^2$ ;

- (M44)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = 1, [a, b] = c, [c, a] = c^p = b^{sp^2}, [c, b] = a^{-\nu p^2} b^{st\nu p^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ,  $s \in F_p^*$ ,  $t = 0, 1, \dots, \frac{p-1}{2}$  such that  $(st\nu)^2 - 4\nu(s+1) \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle \cong C_{p^2} \times C_p$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p^2$ ;
- (M45)  $\langle a, b, c \mid a^{p^3} = b^{p^3} = 1, [a, b] = c, [c, a] = c^p = b^{-p^2}, [b, c] = a^{\nu p^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle \cong C_{p^2} \times C_p$  and  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p^2$ ;
- (M46)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [a, b] = c, [c, a] = c^p, [c, b] = a^{-\nu p^2} c^{t\nu p} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ,  $t = 0, 1, \dots, \frac{p-1}{2}$  such that  $(t\nu)^2 - 4\nu \notin F_p^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p, c \rangle \cong C_{p^2} \times C_{p^2} \times C_p$ ,  $G' = \langle a^{p^2}, c \rangle \cong C_{p^2} \times C_p$  and  $Z(G) = \langle a^{p^2}, c^p \rangle \cong C_p^2$ ;
- (M47)  $\langle a, b, c, d \mid a^{p^2} = b^{p^2} = c^p = d^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = d, [d, a] = [d, b] = 1 \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modular  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p, c, d \rangle \cong C_p^4$ ,  $G' = \langle b^p, c, d \rangle \cong C_p^3$  and  $Z(G) = \langle a^p, b^p, d \rangle \cong C_p^3$ .
- (Mv)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ , and  $G/M'$  is a group of Type (4) in Lemma 2.5. In this case,  $p \geq 5$ ,  $|M'| = p$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = 2p^2$ .
- (M48)  $\langle b, a_1, a_2, a_3 \mid b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = b^p, [a_3, b] = b^p, [a_3, a_1] = 1 \rangle$ ; where  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3, b^p \rangle \cong C_p^3$  and  $Z(G) = \langle b^p \rangle \cong C_p$ .
- (M49)  $\langle b, a_1, a_2, a_3 \mid b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = b^{\eta p}, [a_3, b] = b^p, [a_3, a_1] = 1 \rangle$ , where  $\eta$  is a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3, b^p \rangle \cong C_p^3$  and  $Z(G) = \langle b^p \rangle \cong C_p$ .
- (M50)  $\langle b, a_1, a_2, a_3 \mid b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = b^p, [a_3, b] = b^{\eta p}, [a_3, a_1] = 1 \rangle$ , where  $p \equiv 1 \pmod{4}$  and  $\eta$  is a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3, b^p \rangle \cong C_p^3$  and  $Z(G) = \langle b^p \rangle \cong C_p$ .
- (M51)  $\langle b, a_1, a_2, a_3 \mid b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = b^{\eta p}, [a_3, b] = b^{\eta p}, [a_3, a_1] = 1 \rangle$ , where  $p \equiv 1 \pmod{4}$  and  $\eta$  is a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3, b^p \rangle \cong C_p^3$  and  $Z(G) = \langle b^p \rangle \cong C_p$ .
- (M52)  $\langle b, a_1, a_2, a_3 \mid b^p = a_1^{\nu p^2} = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = a_1^p, [a_3, b] = a_1^{\nu p}, [a_3, a_1] = 1 \rangle$ , where  $\nu = 1, \eta_1$  or  $\eta_2$ ,  $\{1, \eta_1, \eta_2\}$  is a transversal



sal for  $(F_p^*)^3$  in  $F_p^*$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3, a_1^p \rangle \cong C_p^3$  and  $Z(G) = \langle a_1^p \rangle \cong C_p$ .

$$(M53) \langle b, a_1; a_2, a_3, a_4 \mid b^p = a_1^p = a_2^p = a_3^p = a_4^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_2, a_1] = a_4, [a_3, b] = a_4, [a_3, a_1] = [a_4, a_1] = [a_4, b] = 1 \rangle, \text{ where } |G| = p^5, \\ \Phi(G) = G' = \langle a_2, a_3, a_4 \rangle \cong C_p^3 \text{ and } Z(G) = \langle a_4 \rangle \cong C_p.$$

(Mvi)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ ,  $|M'| = 9$  and  $G/M'$  is a group of Type (6) in Lemma 2.5. In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = 2p^2 + p$ .

$$(M54) \langle b, a_1; a_2 \mid b^{27} = a_1^9 = a_2^9 = 1, [a_1, b] = a_2, [a_2, a_1] = b^{-9}, [a_2, b] = a_1^3 a_2^{3s} \rangle, \\ \text{where } s = 0, 2; \text{ moreover, } |G| = 3^7, \Phi(G) = \langle a_2, a_1^3, b^3 \rangle \cong C_3 \times C_9 \times C_9, \\ G' = \langle a_2, a_1^3, b^9 \rangle \cong C_3^2 \times C_9 \text{ and } Z(G) = \langle a_2^3, b^9 \rangle \cong C_3^2.$$

$$(M55) \langle b, a_1; a_2 \mid b^{27} = a_1^9 = a_2^9 = 1, [a_1, b] = a_2, [a_2, a_1] = b^9 a_2^3, [a_2, b] = a_1^3 \rangle, \text{ where } \\ |G| = 3^7, \Phi(G) = \langle a_2, a_1^3, b^3 \rangle \cong C_3 \times C_9 \times C_9, G' = \langle a_2, a_1^3, b^9 \rangle \cong C_3^2 \times C_9 \\ \text{and } Z(G) = \langle a_2^3, b^9 \rangle \cong C_3^2.$$

(Mvii)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ ,  $|M'| = p^2$  where  $p \geq 5$  and  $G/M'$  is a group of Type (6) in Lemma 2.5. In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = 2p^2 + p$ .

$$(M56) \langle b, a_1; a_2 \mid b^{p^3} = a_1^{p^2} = a_2^{p^2} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2}, [a_2, b] = a_1^{\nu_2 p} a_2^{s p} \rangle, \\ \text{where } \nu_1, \nu_2 = 1 \text{ or a fixed quadratic non-residue modulo } p \text{ such that } -\nu_1 \text{ is} \\ \text{not a square, and } s = 2^{-1} \nu_2, 2^{-1} \nu_2 + 1, \dots, 2^{-1} \nu_2 + \frac{p-1}{2}; \text{ moreover, } |G| = p^7, \\ \Phi(G) = \langle a_2, a_1^p, b^p \rangle \cong C_{p^2} \times C_{p^2} \times C_p, G' = \langle a_2, a_1^p, b^{p^2} \rangle \cong C_p^2 \times C_{p^2} \text{ and} \\ Z(G) = \langle a_2^p, b^{p^2} \rangle \cong C_p^2.$$

$$(M57) \langle b, a_1; a_2 \mid b^{p^3} = a_1^{p^2} = a_2^{p^2} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2} a_2^{r p}, [a_2, b] = \\ a_1^{\nu_2 p} \rangle, \text{ where } \nu_1, \nu_2 = 1 \text{ or a fixed quadratic non-residue modulo } p \text{ and } r = \\ 1, 2, \dots, \frac{p-1}{2} \text{ such that } r^2 - 4\nu_1 \text{ is not a square. Moreover, } |G| = p^7, \Phi(G) = \\ \langle a_2, a_1^p, b^p \rangle \cong C_{p^2} \times C_{p^2} \times C_p, G' = \langle a_2, a_1^p, b^{p^2} \rangle \cong C_p^2 \times C_{p^2} \text{ and } Z(G) = \\ \langle a_2^p, b^{p^2} \rangle \cong C_p^2.$$

(Mviii)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ ,  $|M'| = p$  where  $p \geq 3$  and  $G/M'$  is a group of Type (6) in Lemma 2.5. In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p)$  and  $\alpha_1(G) = 2p^2$ .

$$(M58) \langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^2} = a_2^{t p}, [a_1, b] = a_2, [a_2, a_1] = a_2^p, [a_2, b] = a_1^{\nu p} \rangle, \\ \text{where } t \in F_p, \nu = 1 \text{ or a fixed quadratic non-residue modulo } p; \text{ moreover, } \\ |G| = p^6, \Phi(G) = \langle a_2, a_1^p, b^p \rangle \cong C_{p^2} \times C_p^2, G' = \langle a_2, a_1^p \rangle \cong C_{p^2} \times C_p, Z(G) = \\ \langle a_2^p, b^{p^2} \rangle \cong C_p \text{ except for } p = 3 \text{ and } \nu = -1, \text{ In case of } p = 3 \text{ and } \nu = -1, \\ \text{we have } Z(G) = \langle a_2^p, b^p \rangle \cong C_p^2 \text{ if } t \equiv 0 \pmod{p} \text{ or } Z(G) = \langle a_2^p, b^p \rangle \cong C_{p^2} \text{ if} \\ t \not\equiv 0 \pmod{p}.$$

(M59)  $\langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = b^{p^m} = 1, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} a_2^{s p} \rangle$ , where  $m \geq 2$ ,  $\nu = 1$  for  $p = 3$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  for  $p \geq 5$ ,  $s = \nu, \nu + 1, \dots, \nu + \frac{p-1}{2}$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_1^p, b^p \rangle \cong C_p \times C_{p^{m-1}} \times C_{p^2}$ ,  $G' = \langle a_2, a_1^p \rangle \cong C_p \times C_{p^2}$ ,  $Z(G) = \langle a_2^p, b^{p^2} \rangle \cong C_p \times C_{p^{m-2}}$  for  $m > 2$ ,  $Z(G) = \langle a_2^p, b^{p^2} \rangle \cong C_p$  for  $m = 2$ .

(M60)  $\langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^m} = a_2^p, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} \rangle$ , where  $m \geq 2$ ,  $\nu = 1$  for  $p = 3$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  for  $p \geq 5$ . Moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a_2, a_1^p, b^p \rangle \cong C_{p^m} \times C_p^2$ ,  $G' = \langle a_2, a_1^p \rangle \cong C_p \times C_{p^2}$ ,  $Z(G) = \langle a_2^p, b^{p^2} \rangle \cong C_{p^{m-2}}$  for  $m > 2$ ,  $Z(G) = \langle a_2^p, b^{p^2} \rangle \cong C_p$  for  $m = 2$ .

(Mix)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ ,  $|M'| = 3$  and  $G/M' \in \mathcal{A}_3$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = 2p^2$ .

(M61)  $\langle b, a_1; a_2 \mid a_1^9 = a_2^9 = b^{27} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{9s}, [a_2, b] = a_1^{-3} a_2^{3t} \rangle$ , where  $s, t = 1, 2$ . Moreover,  $|G| = 3^7$ ,  $\Phi(G) = \langle a_2, a_1^3, b^3 \rangle \cong C_3 \times C_9 \times C_9$ ,  $G' = \langle a_2, a_1^3, b^9 \rangle \cong C_9 \times C_3 \times C_3$  and  $Z(G) = \langle a_2^3, b^3 \rangle \cong C_3 \times C_9$ .

(Mx)  $c(G) = 4$ ,  $G$  has a unique three-generator maximal subgroup  $M$ ,  $|M'| = p$  where  $p \geq 5$  and  $G/M' \in \mathcal{A}_3$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$  and  $\alpha_1(G) = 2p^2$ .

(M62)  $\langle b, a_1; a_2 \mid b^{p^2} = a_1^{p^2} = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu p}, [a_2, b] = a_3, [a_3, b] = a_1^{t p}, [a_3, a_1] = 1 \rangle$ , where  $p \geq 5$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ,  $t = t_1, t_2, \dots, t_{(3, p-1)}$ , where  $t_1, t_2, \dots, t_{(3, p-1)}$  are the coset representatives of the subgroup  $(F_p^*)^3$  in  $F_p^*$ . Moreover,  $|G| = p^6$ ,  $\Phi(G) = G' = \langle a_2, a_3, a_1^p, b^p \rangle \cong C_p^4$  and  $Z(G) = \langle a_1^p, b^p \rangle \cong C_p^2$ .

**Proof** If  $\Phi(G')G_3 \leq Z(G)$  and  $\Phi(G')G_3 \leq C_p^2$ , then  $G$  is one of the groups listed in [3, Theorem 3.5, 4.6, 5.5, 5.8 and 6.5]. By [3, Theorem 3.6, 4.7, 5.1-5.2, 5.5-5.6 and 6.1], we get the groups (M1)-(M47). For convenience, in Table 11 we give the correspondence from those groups (M1)-(M47) to [3, Theorem 3.5, 4.6, 5.5, 5.8 and 6.5]. In the following, we may assume that  $\Phi(G')G_3 \not\leq Z(G)$  or  $\Phi(G')G_3 \not\leq C_p^2$ .

If  $G$  has two distinct three-generator maximal subgroup  $M_1$  and  $M_2$ , then, by Lemma 2.6 (2),  $M'_1 \leq C_p^3$  and  $M'_2 \leq C_p^3$ . By hypothesis,  $M'_1 \leq Z(G)$  and  $M'_2 \leq Z(G)$ . Let  $N = M'_1 M'_2$ . Then  $\exp(N) = p$  and  $N \leq Z(G)$ . Let  $\bar{G} = G/N$ . Then  $\bar{G}$  has two distinct abelian subgroups  $\bar{M}_1$  and  $\bar{M}_2$  of index  $p$ . By Lemma 2.2 we have  $\bar{G} \in \mathcal{A}_1$ . It follows that  $\Phi(G')G_3 = N$ . By above assumption,  $|N| \geq p^3$ . Hence  $G$  is not metacyclic. By Lemma 2.1,  $\bar{G}$  is not metacyclic. Let

$$\bar{G} = \langle \bar{a}, \bar{b}; \bar{c} \mid \bar{a}^{p^n} = \bar{b}^{p^m} = \bar{c}^p = 1, [\bar{a}, \bar{b}] = \bar{c}, [\bar{c}, \bar{a}] = [\bar{c}, \bar{b}] = 1 \rangle,$$

Groups	Groups in [3, Theorem 3.5, 4.6, 5.5, 5.8 and 6.5]	Groups	Groups in [3, Theorem 3.5, 4.6, 5.5, 5.8 and 6.5]
(M1)	(C1)	(M25)	(L2) where $n = 2$ and $t^2 - 4t \notin F_p^2$
(M2)	(C2)	(M26)	(L4) where $n = 2$ and $-\nu \notin F_p^2$
(M3)	(D1) where $p > 2 = n$	(M27)	(L5) where $n = 2$ and $-\nu \notin F_p^2$
(M4)	(D2) where $p > 2 = n$	(M28)	(L6) where $n = 2$
(M5)	(D3)	(M29)	(L8) where $n = 2$
(M6)	(I4) where $p > 2 = n$	(M30)	(M1) where $m = 2$ and $s = 0$
(M7)	(D5)	(M31)	(M1) where $m = 2$ , $s = 0$ and $\nu = \eta$
(M8)	(E2) where $p > 2 = m$	(M32)	(M2) where $m = 2$
(M9)	(E5) where $p > 2 = m$	(M33)	(M4) where $m = 2$
(M10)	(E8) where $p > 2 = m$	(M34)	(M5) where $m = 2$
(M11)	(E3)	(M35)	(M6) where $m = 2$ and $t^2 + 4t \notin F_p^2$
(M12)	(E6)	(M36)	(M8) where $m = 2$ and $\nu = \eta$
(M13)	(E9)	(M37)	(N3)
(M14)	(G1) where $p = m = 2$	(M38)	(N8)
(M15)	(G2) where $p = m = 2$	(M39)	(N9)
(M16)	(G3) where $p = m = 2$	(M40)	(P1) where $n = 1$
(M17)	(J2) where $p = m = 2$	(M41)	(P3) where $n = 2$ and $-\nu \notin F_p^2$
(M18)	(J4) where $p = m = 2$	(M42)	(p4) where $n = 2$ and $-r \notin F_p^2$
(M19)	(J6) where $p = m = 2$	(M43)	(P10) where $n = 1$
(M20)	(K2)	(M44)	(Q1) where $n = 2$ and $(st\nu)^2 - 4\nu(s+1) \notin F_p^2$
(M21)	(K4)	(M45)	(Q1) where $n = 2$ , $s = -1$ and $t = 0$
(M22)	(K8)	(M46)	(Q2) where $n = 2$
(M23)	(K10)	(M47)	(S4) where $n = 2$
(M24)	(L1) where $n = 2$ and $1 + 4s \notin F_p^2$		

Table 11: The correspondence from Theorem 5.13 to [3, Theorem 3.5, 4.6, 5.5, 5.8 and 6.5]

where  $n \geq m$ . Then  $N = \langle c^p, [c, a], [c, b] \rangle$ ,  $|N| = p^3$  and  $|G| = p^{n+m+4}$ . Since  $c^p \neq 1$ , by calculation we have  $[a, b^p] \neq 1$ . It follows that  $m \geq 2$ . Since  $\langle c, b \rangle \in \mathcal{A}_1$ , we have  $|\langle c, b \rangle| = p^{n+m+2}$ . It follows that  $n \leq 2$  and hence  $n = m = 2$ . Moreover,

$$N = \langle c^p, a^{p^2}, b^{p^2} \rangle, \text{ where } [c, a] \notin \langle c^p, a^{p^2} \rangle \text{ and } [c, b] \notin \langle c^p, b^{p^2} \rangle.$$

By suitable replacement, we may assume that  $[c, a] = b^{rp^2} c^{sp}$  and  $[c, b] = a^{up^2} b^{vp^2} c^{wp}$ , where  $r, s, u, v, w \in F_p$ ,  $r \neq 0$  and  $u \neq 0$ . If  $p = 2$ , then we can prove that  $G \in \mathcal{A}_4$ . The details are omitted. Hence  $p \geq 3$ . If  $s^2 - 4r$  is a square, then the equation  $x^2 - sx + r = 0$  has a solution  $x_1$ . By calculation we have  $\langle a, cb^{x_1 p} \rangle$  is not abelian and of order  $p^5$ , which contradicts that  $G \in \mathcal{A}_3$ . Hence  $s^2 - 4r$  is not a square. By Lemma 2.10, the equation  $x^2 + sxy + ry^2 + wx + vy - u = 0$  has a solution  $(x_0, y_0)$ . By calculation,  $\langle ca^{x_0 p}, ba^{y_0} \rangle$  is not abelian and of order  $p^5$ , which contradicts that  $G \in \mathcal{A}_3$  again.

By above argument,  $G$  has a unique three-generator maximal subgroup  $M$ . Let  $\bar{G} = G/M'$ . Then  $\bar{G}$  has a three-generator abelian subgroup  $M/M'$  of index  $p$ , and every non-abelian subgroup of  $\bar{G}$  is generated by two elements. By Lemma 2.12 (2),  $p \geq 3$ . Since  $\bar{G} \in \mathcal{A}_2$  or  $\mathcal{A}_3$ ,  $\bar{G}$  is either one of the groups (4)–(7) in Lemma 2.5 or, by Theorem 5.1, one of the groups (F4)–(F8).

We claim that  $G'$  is abelian. Otherwise,  $|G : G'| = p^2$  and  $G'$  is minimal non-abelian. By Lemma 2.14,  $G'$  is abelian, a contradiction.

Case 1:  $\bar{G}$  is the group (4) in Lemma 2.5. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2, \bar{a}_3 \mid \bar{a}_1^p = \bar{a}_2^p = \bar{a}_3^p = \bar{b}^{p^m} = 1, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{a}_3, [\bar{a}_3, \bar{b}] = 1, [\bar{a}_i, \bar{a}_j] = 1 \rangle$ , where  $1 \leq i, j \leq 3$ .

Since  $d(\bar{M}) = 3$ , we have  $m = 1$  and  $M = \langle a_1, a_2, a_3 \rangle$ . Hence  $p \geq 5$ . Since  $G'$  is abelian,  $[a_2, a_3] = 1$ . By calculation we have

$$[a_3, a_1] = [a_2, b, a_1] = [a_2, a_1, b] = 1.$$

It follows that

$$[a_1, a_2] \neq 1, |M'| = p \text{ and } |G| = p^5.$$

By calculation we have

$$a_3^p = [a_2, b]^p = [a_2^p, b] = 1 \text{ and } a_2^p = [a_1^p, b] = 1.$$

Since  $\langle a_2, b \rangle \in \mathcal{A}_2$ ,  $[a_3, b] \neq 1$  and hence  $G$  is of maximal class.

If  $\exp(G) = p^2$  and  $\exp(M) = p$ , then  $a_1^p = 1$  and  $b^{p^2} = 1$ . Hence  $G = \langle b, a_1, a_2, a_3 \rangle$  has following relations:

$$b^{p^2} = a_1^p = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = b^{ip}, [a_2, a_1] = b^{jp}$$

where  $i, j \in F_p^*$ . By Lemma 5.8 we get groups (M48)–(M51).

If  $\exp(G) = p^2$  and  $\exp(M) = p^2$ , then  $a_1^{p^2} = 1$ . By suitable replacement, we may assume that  $b^p = 1$ . Hence  $G = \langle b, a_1, a_2, a_3 \rangle$  has following relations:

$$b^p = a_1^{p^2} = a_2^p = a_3^p = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = a_1^{ip}, [a_2, a_1] = a_1^{jp}$$

where  $i, j \in F_p^*$ . By Lemma 5.9 we get the group (M52).

If  $\exp(G) = p$ , then we get the group (M53).

Case 2:  $\bar{G}$  is the group (5) in Lemma 2.5. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2 \mid \bar{a}_1^p = \bar{a}_2^p = \bar{b}^{p^{m+1}} = 1, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{b}^{p^m}, [\bar{a}_1, \bar{a}_2] = 1 \rangle$ .

Since  $d(\bar{M}) = 3$ ,  $M = \langle a_1, a_2, b^p \rangle$ . Let  $N = \langle a_2, b, M' \rangle$ . Then  $N$  is maximal in  $G$  and  $d(N) = 2$ . It follows that  $N \in \mathcal{A}_2$ . Since  $G'$  is abelian, we have  $[b^{p^m}, a_2] = 1$  and hence  $c(N) = 2$ . By calculation we get  $b^{p^{m+1}} = [a_2, b]^p = [a_2^p, b] = 1$  and hence  $|N'| = p$ . By Lemma 2.2 we have  $N \in \mathcal{A}_1$ , a contradiction.

Case 3:  $\bar{G}$  is the group (6) in Lemma 2.5. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2 \mid \bar{a}_1^p = \bar{a}_2^p = \bar{b}^{p^m} = 1, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{a}_1^{\nu p}, [\bar{a}_1, \bar{a}_2] = 1 \rangle$ , where  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .

Since  $d(\bar{M}) = 3$ , we have  $m \geq 2$  and  $M = \langle a_1, a_2, b^p \rangle$ . Let  $N = \langle a_2, b, M' \rangle$ . Then  $N$  is maximal in  $G$  and  $d(N) = 2$ . It follows that  $N \in \mathcal{A}_2$ . By calculation,  $a_1^{\nu p^2} = [a_2, b]^p = [a_2^p, b] = 1$ . Hence  $a_1^{p^2} = 1$  and  $\exp(N') = p$ . If  $c(N) = 2$ , then  $G \in \mathcal{A}_1$ , a contradiction. Hence  $c(N) = 3$ . By calculation,  $[a_2, b^p] = 1$ . Since  $[a_1^p, a_2] = 1$ , we have  $[a_1^p, b] = a_2^p \neq 1$ .

If  $|M'| = p^2$ , then  $M' = \langle [a_1, a_2] \rangle \times \langle [a_1, b^p] \rangle$ . Since  $G \in \mathcal{A}_3$ , we have  $|\langle [a_1, a_2] \rangle| = |\langle [a_1, b^p] \rangle| = p^{m+3}$ . It follows that  $m = 2$ ,  $|G| = p^7$  and  $M' = \langle a_2^p, b^{p^2} \rangle$ . By suitable replacement, we may assume that

$$G = \langle b, a_1; a_2 \mid b^{p^3} = a_1^{p^2} = a_2^p = 1, [a_1, b] = a_2, [a_2, a_1] = b^{\nu_1 p^2} a_2^{rp}, [a_2, b] = a_1^{\nu_2 p} a_2^{sp} \rangle,$$

where  $\nu_1, \nu_2 = 1$  or a fixed quadratic non-residue modulo  $p$ . By Lemma 5.10 we get groups (M54)–(M57).

In the following, we may assume that  $|M'| = p$ . Hence  $M' = \langle a_2^2 \rangle$  and  $|G| = p^{m+4}$ .

If  $[a_2, a_1] \neq 1$ , then  $|\langle a_1, a_2 \rangle| = p^{m+2}$  since  $G \in \mathcal{A}_3$ . It follows that  $m = 2$  and  $|G| = p^6$ . By suitable replacement, we may assume that

$$G = \langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^2} = a_2^{tp}, [a_1, b] = a_2, [a_2, a_1] = a_2^p, [a_2, b] = a_1^{\nu p} \rangle,$$

where  $t \in F_p$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ . By Lemma 5.11 we get the group (M58).

If  $[a_2, a_1] = 1$ , then, by suitable replacement, we may assume that

$$G = \langle b, a_1; a_2 \mid a_1^{p^2} = a_2^{p^2} = 1, b^{p^m} = a_2^{tp}, [a_1, b] = a_2, [a_2, a_1] = 1, [a_2, b] = a_1^{\nu p} a_2^{sp} \rangle,$$

where  $s, t \in F_p$ . By Lemma 5.12 we get groups (M59)–(M60).

Case 4:  $\bar{G}$  is the group (7) in Lemma 2.5. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2 \mid \bar{a}_1^9 = \bar{a}_2^3 = 1, \bar{b}^3 = \bar{a}_1^3, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{a}_1^{-3}, [\bar{a}_2, \bar{a}_1] = 1 \rangle$ .

In this case, we have  $d(\bar{M}) = d(\langle \bar{a}_2, \bar{a}_1 \rangle) = 2$ , a contradiction.

Case 5:  $\bar{G}$  is the group (F4) in Main Theorem. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2 \mid \bar{a}_1^9 = \bar{a}_2^9 = \bar{b}^{3m} = 1, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{a}_1^{-3} \bar{a}_2^{3t}, [\bar{a}_1, \bar{a}_2] = 1 \rangle$ , where  $t = 1, 2$ .

Since  $d(\bar{M}) = 3$ , we have  $m \geq 2$  and  $M = \langle a_1, a_2, b^3 \rangle$ . By Lemma 2.6 (4),  $|M'| = 3$ . By Lemma 2.6 (3),  $m = 2$  and hence  $|G| = 3^7$ . Let  $N = \langle a_2, b, M' \rangle$ . Then  $N$  is maximal in  $G$  and  $d(N) = 2$ . It follows that  $N \in \mathcal{A}_2$ . By calculation,  $a_2^9 = [a_1^3, b]^3 = [a_1^9, b] = 1$ . Since  $|\langle a_2^3, b \rangle| \leq 3^4$ , we have  $[a_2^3, b] = 1$ . By calculation,  $a_1^9 = [b, a_2]^3 = [b, a_2^3] = 1$ . Hence  $\exp(N') = 3$ . Let  $L = \langle a_1^3, b \rangle$ . Then  $L \in \mathcal{A}_1$ . It follows that  $|L| = 3^5$ . Hence  $b^9 \neq 1$ . That is,  $M' = \langle b^9 \rangle$ . By calculation,  $[a_2, b^3] = 1$  and  $[a_1, b^3] = 1$ . Hence  $[a_2, a_1] \neq 1$ . Then we may assume

$$G = \langle b, a_1; a_2 \mid a_1^9 = a_2^9 = b^{27} = 1, [a_1, b] = a_2, [a_2, a_1] = b^{9s}, [a_2, b] = a_1^{-3} a_2^{3t} b^{9r} \rangle,$$

where  $s, t = 1, 2$ . By replacing  $b$  with  $a_1^{-rs}$  we get the group (M61).

Case 6:  $\bar{G}$  is the group (F5) in Main Theorem. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2 \mid \bar{a}_1^9 = \bar{a}_2^9 = 1, \bar{b}^{3m} = \bar{a}_2^{-3}, [\bar{a}_1, \bar{b}] = \bar{a}_2, [\bar{a}_2, \bar{b}] = \bar{a}_1^{-3} \bar{a}_2^{-3}, [\bar{a}_1, \bar{a}_2] = 1 \rangle$ .

Since  $d(\bar{M}) = 3$ , we have  $m \geq 2$  and  $M = \langle a_1, a_2, b^3 \rangle$ . By Lemma 2.6 (4),  $|M'| = 3$ . By Lemma 2.6 (3),  $m = 2$  and hence  $|G| = 3^7$ . Let  $N = \langle a_2, b, M' \rangle$ . Then  $N$  is maximal in  $G$  and  $d(N) = 2$ . It follows that  $N \in \mathcal{A}_2$ . By calculation we have

$$a_2^9 = [a_1^3, b]^3 = [a_1^9, b] = 1 \text{ and } a_1^9 = [b, a_2]^3 = [b, a_2^3] = 1.$$

Hence  $\exp(N') = 3$ . Let  $L = \langle a_1^3, b \rangle$ . Then  $L \in \mathcal{A}_1$ . It follows that  $|L| = 3^5$ . Hence  $b^9 \neq a_2^{-3}$ . That is,  $M' = \langle a_2^3 b^9 \rangle$ . By calculation,  $[a_2, b^3] = 1$  and  $[a_1, b^3] = 1$ . Hence  $[a_2, a_1] \neq 1$ . Then we may assume  $[a_2, a_1] = a_2^{3s} b^{9s}$ , where  $s = 1, 2$ . By calculation,  $\langle a_2 b^3, a_1 \rangle \in \mathcal{A}_1$  and is of order  $3^4$ , which contradicts that  $G \in \mathcal{A}_3$ .

Case 7:  $\bar{G}$  is the group (F6) in Main Theorem. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2, \bar{a}_3, \bar{a}_4 \mid \bar{a}_i^p = \bar{b}^{p^m} = 1, [\bar{a}_j, \bar{b}] = \bar{a}_{j+1}, [\bar{a}_4, \bar{b}] = 1, [\bar{a}_i, \bar{a}_j] = 1 \rangle$ , where  $p \geq 5$ ,  $1 \leq i \leq 4$ ,  $1 \leq j \leq 3$ .

In this case, we have  $d(\bar{M}) = d(\langle \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{a}_4, \bar{b}^p \rangle) = 5$ , a contradiction.

Case 8:  $\bar{G}$  is the group (F7) in Main Theorem. That is,  $\bar{G} = \langle \bar{a}_1, \bar{b}; \bar{a}_2, \bar{a}_3 \mid \bar{a}_i^p = \bar{b}^{p^{m+1}} = 1, [\bar{a}_j, \bar{b}] = \bar{a}_{j+1}, [\bar{a}_3, \bar{b}] = \bar{b}^{p^m}, [\bar{a}_i, \bar{a}_j] = 1 \rangle$ , where  $p \geq 5$ ,  $1 \leq i \leq 3$ ,  $1 \leq j \leq 2$ .

In this case, we have  $d(\bar{M}) = d(\langle \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{b}^p \rangle) = 4$ , a contradiction.

Case 9:  $\bar{G}$  is the group (F8) in Main Theorem. That is,  $\bar{G} = \langle a_1, b; \bar{a}_2, \bar{a}_3 \mid \bar{a}_1^{p^2} = \bar{a}_i^p = \bar{b}^{p^m} = 1, [\bar{a}_j, b] = \bar{a}_{j+1}, [\bar{a}_3, \bar{b}] = \bar{a}_1^{tp}, [\bar{a}_i, \bar{a}_j] = 1 \rangle$ , where  $2 \leq i \leq 3$ ,  $1 \leq j \leq 2$ , and  $t = t_1, t_2, \dots, t_{(3,p-1)}$ , where  $p \geq 5$ ,  $t_1, t_2, \dots, t_{(3,p-1)}$  are the coset representatives of the subgroup  $(F_p^*)^3$  in  $F_p^*$ .

Since  $d(\bar{M}) = 3$ , we have  $m = 1$  and  $M = \langle a_1, a_2, a_3 \rangle$ . Since  $G'$  is abelian,  $[a_2, a_3] = 1$ . By calculation we have

$$[a_3, a_1] = [a_2, b, a_1] = [a_2, a_1, b] = 1.$$

It follows that

$$[a_1, a_2] \neq 1, |M'| = p \text{ and } |G| = p^6.$$

By calculation we have

$$a_3^p = [a_2, b]^p = [a_2^p, b] = 1, a_2^p = [a_1^p, b] = 1 \text{ and } a_1^{tp^2} = [a_3^p, b] = 1.$$

Let  $[a_2, a_1] = d$ . Then we may assume

$$G = \langle b, a_1; a_2, a_3, d \mid a_1^{p^2} = a_2^p = a_3^p = 1, b^p = d^s, [a_1, b] = a_2, [a_2, a_1] = d, [a_2, b] = a_3, [a_3, b] = a_1^{tp} d^r \rangle.$$

Since  $[a_3, ba_1^x] = a_1^{tp} d^r \neq 1$ , we have  $|\langle a_3, ba_1^x \rangle| = p^4$  for any  $x$ . It follows that the equation  $st - rx \equiv 0 \pmod{p}$  about  $x$  has no solution. Hence we have  $r = 0$  and  $s \neq 0$ . By suitable replacement we get the group (M62).

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 5.13 as follows. Since  $d(G) = 2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p)$ . In the following, we calculate  $\alpha_1(G)$ .

**Case 1.**  $G$  is one of the groups (M1)–(M19).

Since  $|G'| = p^2$ ,  $|M'| = p$  for any  $M \in \Gamma_1$ . By Lemma 2.6 (7),  $\alpha_1(M) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (1 + p)p^2 = p^2 + p^3.$$

**Case 2.**  $G$  is one of the groups (M20)–(M36).

In this case,  $G = \langle a, b \rangle$  such that  $[a, b] = c$ ,  $c^{p^2} = 1$ ,  $[c, a] = 1$  and  $[c, b] \notin \langle c^p \rangle$ . All maximal subgroups of  $G$  are:

$$M = \langle c, a, \Phi(G) \rangle;$$

$$M_i = \langle c, ba^i, \Phi(G) \rangle, \text{ where } 0 \leq i \leq p-1.$$

It is easy to see that  $|M'| = p$ . By Lemma 2.6 (7),  $\alpha_1(M) = p^2$ . By calculation,  $d(M_i) = 3$  and  $|M'_i| = p^2$ . Hence  $\alpha_1(M_i) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p^2 + p \times (p^2 + p) = p^3 + 2p^2.$$

**Case 3.**  $G$  is one of the groups (M37).

All maximal subgroups of  $G$  are:

$$M = \langle c, a, \Phi(G) \rangle;$$

$$M_i = \langle c, ba^i, \Phi(G) \rangle, \text{ where } i = 0, 1.$$

By calculation,  $d(M) = 3$  and  $|M'| = p^2$ . Hence  $\alpha_1(M) = p^2 + p$ . Similarly,  $\alpha_1(M_i) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (1+p)(p^2+p) = p^3 + 2p^2 + p.$$

**Case 4.**  $G$  is one of the groups (M38)–(M39).

In this case,  $G = \langle a, b \rangle$  such that  $[a, b] = c, c^4 = 1, [c, a] = c^2$  and  $[c, b] \notin \langle c^2 \rangle$ . All maximal subgroups of  $G$  are:

$$M = \langle c, a, \Phi(G) \rangle;$$

$$M_i = \langle c, ba^i, \Phi(G) \rangle, \text{ where } i = 0, 1.$$

By calculation,  $M = \langle c, a, b^2 \rangle$  such that  $d(M) = 3$  and  $|M'| = p^2 = 4$ ,  $M_i = \langle c, ba^i, a^2 \rangle$  such that  $d(M_i) = 3$  and  $|M'_i| = p = 2$ . By Lemma 2.6,  $\alpha_1(M) = p^2 + p$  and  $\alpha_1(M_i) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (p^2 + p) + p \times p^2 = p^3 + p^2 + p.$$

**Case 5.**  $G$  is one of the groups (M40)–(M43).

Let  $H \in \Gamma_1$ . Then  $|H'| = p$ . By Lemma 2.6,  $\alpha_1(H) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (1+p) \times p^2 = p^3 + p^2.$$

**Case 6.**  $G$  is one of the groups (M44)–(M46).

In this case,  $G = \langle a, b \rangle$  such that  $[a, b] = c, c^{p^2} = 1, [c, a] = c^p$  and  $[c, b] \notin \langle c^p \rangle$  where  $p > 2$ . All maximal subgroups of  $G$  are:

$$M = \langle c, a, \Phi(G) \rangle;$$

$$M_i = \langle c, ba^i, \Phi(G) \rangle, \text{ where } i = 0, 1.$$

By calculation,  $M = \langle c, a, b^2 \rangle$  such that  $d(M) = 3$  and  $|M'| = p$ ,  $M_i = \langle c, ba^i, a^2 \rangle$  such that  $d(M_i) = 3$  and  $|M'_i| = p^2$ . By Lemma 2.6,  $\alpha_1(M) = p^2$  and  $\alpha_1(M_i) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p^2 + p \times (p^2 + p) = p^3 + 2p^2.$$

**Case 7.**  $G$  is the group (M47).

Let  $H \in \Gamma_1$ . Then  $|H'| = p$ . By Lemma 2.6,  $\alpha_1(H) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (1 + p) \times p^2 = p^3 + p^2.$$

**Case 8.**  $G$  is one of the groups (M48)–(M53).

In this case,  $\Phi(G)$  is abelian and  $|M'| = p$ . Let  $H \in \Gamma_1 \setminus \{M\}$ . Then  $d(H) = 2$ . Hence  $\alpha_1(H) = p$  and  $\alpha_1(M) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p^2 + p \times p = 2p^2.$$

**Case 9.**  $G$  is one of the groups (M54)–(M57).

In this case,  $\Phi(G)$  is abelian and  $|M'| = p^2$ . Let  $H \in \Gamma_1 \setminus \{M\}$ . Then  $d(H) = 2$ . Hence  $\alpha_1(H) = p$  and  $\alpha_1(M) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = (p^2 + p) + p \times p = 2p^2 + p.$$

**Case 10.**  $G$  is one of the groups (M58)–(M62).

In this case,  $\Phi(G)$  is abelian and  $|M'| = p$ . Let  $H \in \Gamma_1 \setminus \{M\}$ . Then  $d(H) = 2$ . Hence  $\alpha_1(H) = p$  and  $\alpha_1(M) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) = p^2 + p \times p = 2p^2.$$

□

**Lemma 5.14.** *Suppose that  $p \geq 3$  and  $G(\nu, k) = \langle a, b, x \mid a^{p^3} = b^{p^3} = x^p = 1, [a, b] = a^p, [x, b] = a^{p^2}, [x, a] = b^{\nu p^2} a^{k p^2} \rangle$ , where  $k \in F_p$  and  $\nu = 1$  or a fixed quadratic non-residue modula  $p$ .*

- (1) *If  $G = G(\nu, k)$ , then  $|G| = p^7$ ,  $G' = \langle a^p, b^{p^2} \rangle$ ,  $G_3 = \langle a^{p^2} \rangle$ ;*
- (2)  *$G(\nu', k') \cong G(\nu, k)$  if and only if  $\nu' = \nu$  and  $k' = \pm k$ .*
- (3)  *$G(\nu, k) \in A_3$  if and only if  $k^2 + 4\nu$  is not a square.*



**Proof** (1) Let

$$K = \langle a, b \mid a^{p^3} = b^{p^3} = 1, [a, b] = a^p \rangle.$$

We define an automorphism  $\gamma$  of  $K$  as follows:

$$a^\gamma = aa^{-kp^2}b^{-\nu p^2}, \quad b^\gamma = ba^{-p^2}.$$

Then

$$a^{\gamma^p} = 1, \quad b^{\gamma^p} = 1, \quad o(\gamma) = p.$$

It follows from the cyclic extension theory that  $G = \langle K, x \rangle$  is a cyclic extension of  $K$  by  $C_p$ . Hence  $|G| = p^7$ . It is easy to check that  $G' = \langle a^p, b^{p^2} \rangle$ ,  $G_3 = \langle a^{p^2} \rangle$ .

(2) For convenience, let  $G = \langle a, b, x \rangle \cong G(\nu, k)$ ,  $\bar{G} = \langle \bar{a}, \bar{b}, \bar{x} \rangle \cong G(\nu', k')$  and  $\theta$  be an isomorphism from  $\bar{G}$  to  $G$ . Since  $M = \langle x, b, a^p \rangle$  is the unique three-generator maximal subgroups such that  $M' \cong C_p^2$ ,  $M \text{ char } G$ . Similarly,  $\bar{M} = \langle \bar{x}, \bar{b}, \bar{a}^p \rangle \text{ char } \bar{G}$ . Since  $M, \Phi(G), \Omega_1(G) = \langle x, a^{p^2}, b^{p^2} \rangle$  and  $\bar{M}, \Phi(\bar{G}), \Omega_1(\bar{G}) = \langle \bar{x}, \bar{a}^{p^2}, \bar{b}^{p^2} \rangle$  are characteristic in  $G$  and  $\bar{G}$  respectively, we may assume that  $\bar{a}^\theta = a^l x^n w$ ,  $\bar{b}^\theta = b^i x^j y$ ,  $\bar{x}^\theta = x^r z$  where  $l, i, r \in F_p^*$  and  $w, y \in \Phi(G)$ ,  $z \in \Omega_1(M)$ .

Since  $[a^l, b^{ip}] = [\bar{a}^\theta, (\bar{b}^p)^\theta] = [\bar{a}, \bar{b}^p]^\theta = (\bar{a}^{p^2})^\theta$ , we have  $a^{lip^2} = a^{lp^2}$ . Comparing index of  $a^{p^2}$  in two sides, we have  $i = 1$ .

Since  $[x^r, b] = [\bar{x}^\theta, \bar{b}^\theta] = (\bar{a}^{p^2})^\theta = a^{lp^2}$ , we have  $a^{rp^2} = a^{lp^2}$ . Comparing index of  $a^{p^2}$  in two sides, we have  $r = l$ .

Since  $[x^r, a^r] = [\bar{x}^\theta, \bar{a}^\theta] = (\bar{b}^{\nu' p^2} \bar{a}^{k' p^2})^\theta = b^{\nu' p^2} a^{rk' p^2}$ , we have  $b^{r^2 \nu p^2} a^{r^2 k p^2} = b^{\nu' p^2} a^{rk' p^2}$ . Comparing indexes of  $a^{p^2}$  and  $b^{p^2}$  in two sides, we have  $\nu' = r^2 \nu$  and  $k' = rk$ .

Since  $\nu' = r^2 \nu$ ,  $\nu = \nu' = 1$  or  $\nu' = \nu$  is a fixed quadratic non-residue modula  $p$ . Moreover,  $r^2 = 1$  and hence  $r = \pm 1$ . It follows that  $k' = \pm k$ .

On the other hand, if  $\nu' = \nu$  and  $k' = -k$ , then,  $\theta : \bar{a} \rightarrow a^{-1}, \bar{b} \rightarrow b, \bar{x} \rightarrow x^{-1}$  is an isomorphism from  $\bar{G}$  to  $G$ .

(3) All maximal subgroups of  $G(\nu, k)$  are:

$$N = \langle b, x, a^p \rangle;$$

$$N_i = \langle ab^i, x, b^p \rangle \text{ where } 0 \leq i, j \leq p-1.$$

$$N_{ij} = \langle ax^i, bx^j \rangle \text{ where } 0 \leq i, j \leq p-1.$$

It is easy to see that  $N = \langle x, b \rangle * \langle a^p \rangle \in \mathcal{A}_2$ . Since  $N_{ij} \cong \langle a, b \rangle$ ,  $N_{ij} \in \mathcal{A}_2$ . Hence  $G(\nu, k) \in \mathcal{A}_3$  if and only if  $N_i \in \mathcal{A}_2$  for any  $i$ . Let  $a_1 = ab^i$  and  $b_1 = b^p$ . Then  $N_i = \langle a_1, b_1, x \rangle$  such that

$$a_1^{p^3} = b_1^{p^2} = x^p = 1, [a_1, b_1] = a_1^{p^2} b_1^{-ip}, [x, a_1] = a_1^{(k+i)p^2} b_1^{(\nu-ik-i^2)p}, [x, b_1] = 1).$$

By calculation,  $[x, a_1] \neq 1$  for any  $i$ . Hence  $\langle x, a_1 \rangle \in \mathcal{A}_1$ . Since  $G(\nu, k) \in \mathcal{A}_3$ ,  $|\langle x, a_1 \rangle| = p^5$ . It follows that  $\nu - ik - i^2 \neq 0$  for any  $i$ . That is, equation  $i^2 + ik - \nu = 0$  about  $i$  has no solution. Hence  $k^2 + 4\nu$  is not a square.

On the other hand, if  $k^2 + 4\nu$  is not a square, then it can be proved that  $N_i \in \mathcal{A}_2$  for any  $i$ . Hence  $G(\nu, k) \in \mathcal{A}_3$  if and only if  $k^2 + 4\nu$  is not a square.  $\square$

**Theorem 5.15.** *Suppose that  $G$  is an  $A_3$ -group all of whose maximal subgroups are  $A_2$ -groups and there exists a three-generator maximal subgroup in  $G$ , and  $M' \leq Z(G)$  for every three-generator maximal subgroup  $M$ . Then  $d(G) = 3$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Ni)  $\Phi(G) \leq Z(G)$  and  $c(G) = 2$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^4 + p^3 + p^2$ .

(N1)  $\langle a, b, c \mid a^{p^3} = b^{p^2} = c^{p^2} = 1, [b, c] = a^{p^2}, [c, a] = c^p, [a, b] = b^{-p} \rangle$ , where  $p$  is odd; moreover,  $|G| = p^7$ ,  $G' = \langle a^{p^2}, b^p, c^p \rangle \cong C_p^3$  and  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p \rangle \cong C_{p^2} \times C_p \times C_p$ .

(N2)  $\langle a, b, c; d \mid a^{p^2} = b^{p^2} = c^{p^2} = d^p = 1, [b, c] = d, [c, a] = b^p, [a, b] = c^{\nu p}, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $p$  is odd,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$  such that  $-\nu \notin (F_p^*)^2$ ; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p, d \rangle \cong C_p^4$  and  $G' = \langle b^p, c^p, d \rangle \cong C_p^3$ .

(N3)  $\langle a, b, c; d \mid a^{p^2} = b^{p^2} = c^{p^2} = d^p = 1, [b, c] = d, [c, a]^{1+r} = b^{r p} c^{-p}, [a, b]^{1+r} = b^p c^p, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $p$  is odd,  $-r \in F_p$  is not a square; moreover,  $|G| = p^7$ ,  $\Phi(G) = Z(G) = \langle a^p, b^p, c^p, d \rangle \cong C_p^4$  and  $G' = \langle b^p, c^p, d \rangle \cong C_p^3$ .

(N4)  $\langle a, b, c \mid a^8 = b^4 = c^4 = 1, [b, c] = a^4, [c, a] = c^2, [a, b] = b^2 \rangle$ ; where  $|G| = 2^7$ ,  $G' = \langle a^4, b^2, c^2 \rangle \cong C_2^3$  and  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2 \rangle \cong C_4 \times C_2 \times C_2$ .

(N5)  $\langle a, b, c; d \mid a^4 = b^4 = c^4 = d^2 = 1, [b, c] = d, [c, a] = b^2, [a, b] = b^2 c^2, [d, a] = [d, b] = [d, c] = 1 \rangle$ , where  $|G| = 2^7$ ,  $\Phi(G) = Z(G) = \langle a^2, b^2, c^2, d \rangle \cong C_2^4$  and  $G' = \langle b^2, c^2, d \rangle \cong C_2^3$ .

(Nii)  $Z(G) \not\leq \Phi(G)$  and  $c(G) = 3$ . In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^3 + p^2$ .

(N6)  $\langle a, b, x \mid a^{p^{r+t}} = x^p = 1, b^{p^{r+s+t}} = a^{p^{r+s}}, [a, b] = a^{p^r}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ , where  $r \geq 2$  for  $p = 2$ ,  $r \geq 1$  for  $p \geq 3$ ,  $t \geq 0$ ,  $0 \leq s \leq 2$  and  $r + s \geq 2$ ; moreover,  $|G| = p^{2r+s+t+3}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^{r+t+1}} \times C_{p^{r+s-1}}$ ,  $G' = \langle a^{p^r} \rangle \cong C_{p^2}$  and  $Z(G) = \langle a^{p^2}, b^{p^2}, x \rangle \cong C_{p^{r+t}} \times C_{p^{r+s-2}} \times C_p$ .

(N7)  $\langle a, b, x \mid a^{p^3} = b^{p^{t+3}} = 1, x^p = a^{p^2}, [a, b] = a^p, [x, a] = [x, b] = 1 \rangle$ , where  $p \geq 3$  and  $t \geq 0$ ; moreover,  $|G| = p^{t+7}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^{t+2}} \times C_{p^2}$ ,  $G' = \langle a^p \rangle \cong C_{p^2}$ ,  $Z(G) = \langle a^{p^2}, b^{p^2}, x \rangle \cong C_{p^{t+1}} \times C_p^2$ .

(N8)  $\langle a, b, x \mid a^{p^3} = 1, b^{p^2} = x^p = a^{p^2}, [a, b] = a^p, [x, a] = [x, b] = 1 \rangle$ , where  $p \geq 3$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_p$ ,  $G' = \langle a^p \rangle \cong C_{p^2}$  and  $Z(G) = \langle x \rangle \cong C_{p^2}$ .

(N9)  $\langle a, b, x; c \mid a^{p^2} = b^{p^2} = c^p = x^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = a^p, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ , where  $p \geq 5$ ,  $\nu$  is a fixed quadratic non-residue

modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle \cong C_p^3$  and  $Z(G) = \langle a^p, b^p, x \rangle \cong C_p^3$ .

(N10)  $\langle a, b, x; c \mid a^{p^2} = b^{p^2} = c^p = x^p = 1, [a, b] = c, [c, a] = a^{-p}b^{-lp}, [c, b] = a^{-p}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ , where  $p \geq 5$ ,  $4l = \rho^{2r+1} - 1$ ,  $r = 1, 2, \dots, \frac{1}{2}(p-1)$ ,  $\rho$  is the smallest positive integer which is a primitive root modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = G' = \langle a^p, b^p, c \rangle \cong C_p^3$  and  $Z(G) = \langle a^p, b^p, x \rangle \cong C_p^3$ .

(N11)  $\langle a, b, x; c \mid a^9 = b^9 = c^3 = x^3 = 1, [a, b] = c, [c, a] = b^{-3}, [c, b] = a^3, [a^3, b] = [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ ; moreover,  $|G| = 3^6$ ,  $\Phi(G) = G' = \langle a^3, b^3, c \rangle \cong C_3^3$  and  $Z(G) = \langle a^3, b^3, x \rangle \cong C_3^3$ .

(N12)  $\langle a, b, x; c \mid a^9 = b^9 = c^3 = x^3 = 1, [a, b] = c, [c, a] = b^{-3}, [c, b] = a^{-3}, [x, a] = [x, b] = 1 \rangle = \langle a, b \rangle \times \langle x \rangle$ ; moreover,  $|G| = 3^6$ ,  $\Phi(G) = G' = \langle a^3, b^3, c \rangle \cong C_3^3$  and  $Z(G) = \langle a^3, b^3, x \rangle \cong C_3^3$ .

(Niii)  $Z(G) < \Phi(G)$  and  $G$  has at least two three-generator maximal subgroups, in this case,  $c(G) = 2$  for (N24) and  $c(G) = 3$  for else. Moreover,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^3 + p^2$  except for (N14), (N18) and (N22)–(N24).

(N13)  $\langle a, b, x \mid a^8 = b^4 = x^2 = 1, [a, b] = a^{-2}, [x, a] = a^4, [x, b] = 1 \rangle$ ; moreover,  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_4 \times C_2$ ,  $G' = \langle a^2 \rangle \cong C_4$ ,  $Z(G) = \langle a^4, b^2 \rangle \cong C_2^2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 7)$  and  $\alpha_1(G) = 12$ ;

(N14)  $\langle a, b, x \mid a^8 = b^8 = x^2 = 1, [a, b] = a^{-2}, [x, a] = b^4, [x, b] = a^4 \rangle$ ; moreover,  $|G| = 2^7$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_4 \times C_4$ ,  $G' = \langle a^2, b^4 \rangle \cong C_4 \times C_2$ ,  $Z(G) = \langle a^4, b^2 \rangle \cong C_4 \times C_2$  and  $\alpha_1(G) = 14$ ;

(N15)  $\langle a, b, x \mid a^8 = x^2 = 1, b^4 = a^4, [a, b] = a^{-2}, [x, a] = a^4, [x, b] = 1 \rangle$ ; where  $|G| = 2^6$ ,  $\Phi(G) = \langle a^2, b^2 \rangle \cong C_4 \times C_2$ ,  $G' = \langle a^2 \rangle \cong C_4$  and  $Z(G) = \langle b^2 \rangle \cong C_4$ .

(N16)  $\langle a_1, b, x; a_2, a_3 \mid a_1^p = a_2^p = a_3^p = b^p = x^p = 1, [a_1, b] = a_2, [a_2, b] = [x, a_1] = a_3, [a_3, b] = 1, [x, b] = [a_i, a_j] = 1 \rangle$ , where  $p > 3$  and  $1 \leq i, j \leq 3$ ; where  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_3 \rangle \cong C_p^2$  and  $Z(G) = \langle a_3 \rangle \cong C_p$ .

(N17)  $\langle a_1, b, x; a_2 \mid a_1^p = a_2^p = b^p = x^{p^2} = 1, [a_1, b] = a_2, [a_2, b] = [x, a_1] = x^p, [a_2, a_1] = [a_2, x] = [x, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, x^p \rangle \cong C_p^2$  and  $Z(G) = \langle x^p \rangle \cong C_p$ .

(N18)  $\langle a_1, b, x; a_2 \mid a_1^p = a_2^p = b^{p^2} = x^{p^2} = 1, [a_1, b] = a_2, [a_2, b] = x^p, [x, a_1] = b^p, [a_2, a_1] = [a_2, x] = [x, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = G' = \langle a_2, x^p, b^p \rangle \cong C_p^3$ ,  $Z(G) = \langle b^p, x^p \rangle \cong C_p \times C_p$  and  $\alpha_1(G) = p^2 + 2p^2 - p$ .

(N19)  $\langle a_1, b, x; a_2 \mid a_1^p = a_2^p = b^{p^2} = x^p = 1, [a_1, b] = a_2, [a_2, b] = [x, a_1] = b^p, [a_2, a_1] = [a_2, x] = [x, b] = 1 \rangle$ , where  $p > 2$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, b^p \rangle \cong C_p^2$  and  $Z(G) = \langle b^p \rangle \cong C_p$ .

(N20)  $\langle a_1, b, x; a_2 \mid a_1^{p^2} = a_2^p = b^p = x^p = 1, [a_1, b] = a_2, [a_2, b] = a_1^{\nu p}, [x, a_1] = a_1^p, [a_2, a_1] = [a_2, x] = [x, b] = 1 \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^5$ ,  $\Phi(G) = G' = \langle a_2, a_1^p \rangle \cong C_p^2$  and  $Z(G) = \langle a_1^p \rangle \cong C_p$ .

(N21)  $\langle a_1, b, x; a_2 \mid a_1^9 = a_2^3 = x^3 = 1, b^3 = a_1^3, [a_1, b] = a_2, [a_2, b] = a_1^{-3}, [x, a_1] = a_1^3, [a_2, a_1] = [a_2, x] = [x, b] = 1 \rangle$ ; moreover,  $|G| = 3^5$ ,  $\Phi(G) = G' = \langle a_2, a_1^3 \rangle \cong C_3^2$  and  $Z(G) = \langle a_1^3 \rangle \cong C_3$ .

(N22)  $\langle a, b, x \mid a^{p^3} = b^{p^3} = x^p = 1, [a, b] = a^p, [x, b] = a^{p^2}, [x, a] = b^{\nu p^2} a^{kp^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ,  $0 \leq k \leq \frac{p-1}{2}$  such that  $k^2 + 4\nu$  is not a square; moreover,  $|G| = p^7$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_{p^2}$ ,  $G' = \langle a^p, b^{p^2} \rangle \cong C_{p^2} \times C_p$ ,  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p^2$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

(N23)  $\langle a, b, x \mid a^{p^3} = b^{p^{t+3}} = x^p = 1, [a, b] = a^p, [x, b] = a^{p^2} b^{p^{t+2}}, [x, a] = 1 \rangle$ , where  $p > 2$  and  $t \geq 1$ ; moreover,  $|G| = p^{t+7}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_{p^{t+2}}$ ,  $G' = \langle a^p, b^{p^{t+2}} \rangle \cong C_{p^2} \times C_p$ ,  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p \times C_{p^{t+1}}$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

(N24)  $\langle a, b, x \mid a^{p^{t+4}} = b^{p^3} = x^p = 1, [a, b] = a^{p^{t+2}}, [x, a] = b^{p^2}, [x, b] = 1 \rangle$ , where  $t \geq 0$ ; moreover,  $|G| = p^{t+8}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^2} \times C_{p^{t+3}}$ ,  $G' = \langle a^{p^{t+2}}, b^{p^2} \rangle \cong C_{p^2} \times C_p$ ,  $Z(G) = \langle a^{p^2}, b^{p^2} \rangle \cong C_p \times C_{p^{t+2}}$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = p^3 + 2p^2$ .

(Niv)  $Z(G) < \Phi(G)$ ,  $c(G) = 3$  and  $G$  has a unique three-generator maximal subgroup. In this case,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2)$  and  $\alpha_1(G) = 2p^2$ .

(N25)  $\langle a, b, d \mid a^{p^{m+1}} = b^{p^2} = d^p = 1, [a, b] = a^{p^{m-1}}, [d, a] = b^p, [d, b] = 1 \rangle$ , where  $p > 2$  and  $m \geq 2$ ; moreover,  $|G| = p^{m+4}$ ,  $\Phi(G) = \langle a^p, b^p \rangle \cong C_{p^m} \times C_p$ ,  $G' = \langle a^{p^{m-1}}, b^p \rangle \cong C_{p^2} \times C_p$  and  $Z(G) = \langle a^{p^2} \rangle \cong C_{p^{m-1}}$ .

(N26)  $\langle a, b, d \mid a^{p^3} = b^{p^2} = d^p = 1, [a, b] = a^p, [d, a] = b^p, [d, b] = a^{\nu p^2} \rangle$ , where  $p > 2$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ ; moreover,  $|G| = p^6$ ,  $\Phi(G) = G' = \langle a^p, b^p \rangle \cong C_p \times C_{p^2}$  and  $Z(G) = \langle a^{p^2} \rangle \cong C_p$ .

**Proof** If  $\Phi(G) \leq Z(G)$ , then, since  $G$  has no abelian maximal subgroup,  $\Phi(G) = Z(G)$ . Hence  $G$  is one of the groups listed in [23, Theorem 3.1,4.1,5.1,6.1,7.1-7.2,7.6]. By hypothesis, the minimal index of  $\mathcal{A}_1$ -subgroups and the maximal index of  $\mathcal{A}_1$ -subgroups are 2. By checking [23, Theorem 3.3, 4.3,5.2,6.3,7.4-7.5,7.7], we get groups (N1)–(N5).

In the following, we may assume  $\Phi(G) \not\leq Z(G)$ .

**Case 1:**  $Z(G) \not\leq \Phi(G)$ .

Then there exists a maximal subgroup  $M$  such that  $Z(G) \not\leq M$ . Let  $x \in Z(G) \setminus M$ . Then  $G = \langle M, x \rangle$  and  $x^p \in Z(M)$ . If  $d(M) = 3$ , then  $G' = M' \leq Z(G)$  and  $\exp(G') =$

Groups	Groups in [23, Theorem 4.1 & 7.1]	Groups	Groups in [23, Theorem 4.1 & 7.1]
(N1)	(D1) where $m_1 = 2$	(N4)	(M1) where $m_1 = 2$
(N2)	(E3) where $m_1 = 2$ and $-\nu \notin F_p^2$	(N5)	(N3) where $m_1 = 2$
(N3)	(E4) where $m_1 = 2$ and $-\nu \notin F_p^2$		

Table 12: The correspondence from (N1)–(N5) to [23, Theorem 4.1 & 7.1]

$p$ . It follows that  $\Phi(G) \leq Z(G)$ , a contradiction. Hence  $d(M) = 2$ . If  $M$  has an abelian subgroup of index  $p$ , then  $G$  also has an abelian subgroup of index  $p$ , a contradiction. Hence  $\alpha_1(M) = 1 + p$ .

Since any maximal subgroup of  $M$  is an  $\mathcal{A}_1$ -group,  $M$  is one of the groups (17)–(21) in Lemma 2.5.

Subcase 1.1:  $M$  is the group of Type (17) in Lemma 2.5. That is,  $M = \langle a, b \mid a^{p^{r+2}} = 1, b^{p^{r+s+t}} = a^{p^{r+s}}, [a, b] = a^{p^r} \rangle$ , where  $r \geq 2$  for  $p = 2$ ,  $r \geq 1$  for  $p \geq 3$ ,  $t \geq 0$ ,  $0 \leq s \leq 2$  and  $r + s \geq 2$ .

If  $s = 2$ , then

$$Z(M) = \langle a^{p^2} \rangle \times \langle b^{p^2} \rangle, \quad |M| = p^{2r+t+4} \quad \text{and} \quad |G| = p^{2r+t+5}.$$

Since  $x^p \in Z(G)$ , we may assume  $x^p = a^{ip^2} b^{jp^2}$ . Since

$$|\langle xb^{-jp}, a \rangle| = p^{r+3} < p^{2r+t+3} = \frac{|G|}{p^2},$$

we have  $[xb^{-jp}, a] = 1$  and hence  $xb^{-jp} \in Z(G)$ . By replacing  $x$  with  $xb^{-jp}$  we get  $x^p = a^{ip^2}$ .

If  $[xa^{-ip}, b] = 1$ , then, replacing  $x$  with  $xa^{-ip}$ , we get  $x^p = 1$  and  $G$  is a group of Type (N6).

If  $[xa^{-ip}, b] \neq 1$ , then  $(i, p) = 1$ . Without loss of generality, we may assume  $x^p = a^{p^2}$ . Since  $G \in \mathcal{A}_3$ , we have

$$|\langle xa^{-ip}, b \rangle| = p^{r+t+4} = \frac{|G|}{p^2} = p^{2r+t+3}.$$

It follows that  $r = 1$ ,  $p \geq 3$  and  $G$  is a group of Type (N7).

If  $s = 1$  and  $r \geq 2$ , then

$$Z(M) = \langle a^{p^2} b^{-p^{t+2}} \rangle \times \langle b^{p^2} \rangle, \quad |M| = p^{2r+t+3} \quad \text{and} \quad |G| = p^{2r+t+4}.$$

Since  $x^p \in Z(G)$ , we may assume  $x^p = (a^{p^2} b^{-p^{t+2}})^i b^{jp^2}$ . Since

$$|\langle xa^{-ip}, b \rangle| = p^{r+t+3} < p^{2r+t+2} = \frac{|G|}{p^2},$$

we have  $[xa^{-ip}, b] = 1$  and hence  $xa^{-ip} \in Z(G)$ . By replacing  $x$  with  $xa^{-ip}$  we get  $x^p = b^{j'p^2}$ . Since

$$|\langle xb^{-j'p}, a \rangle| = p^{r+3} < p^{2r+t+2} = \frac{|G|}{p^2},$$

we have  $[xb^{-j'p}, a] = 1$  and hence  $xb^{-j'p} \in Z(G)$ . By replacing  $x$  with  $xb^{-j'p}$  we get  $x^p = 1$  and  $G$  is a group of Type (N6).

If  $s = 1$  and  $r = 1$ , then

$$p \geq 3 \text{ and } Z(M) = \langle b^{p^2} \rangle, |M| = p^{t+5} \text{ and } |G| = p^{t+6}.$$

Since  $x^p \in Z(G)$ , we may assume  $x^p = b^{jp^2}$ .

If  $[xb^{-jp}, a] = 1$ , then, replacing  $x$  with  $xb^{-jp}$ , we get  $x^p = 1$  and  $G$  is a group of Type (N6).

If  $[xb^{-jp}, a] \neq 1$ , then  $(j, p) = 1$ . Without loss of generality, we may assume  $x^p = b^{p^2}$ . Since  $G \in \mathcal{A}_3$ , we have

$$|\langle xb^{-p}, a \rangle| = p^4 = \frac{|G|}{p^2} = p^{t+4}.$$

It follows that  $t = 0$  and  $G$  is a group of Type (N8).

If  $s = 0$  and  $r \geq 3$ , then

$$Z(M) = \langle a^{p^2} b^{-p^{t+2}} \rangle \times \langle b^{p^2} \rangle, |M| = p^{2r+t+2} \text{ and } |G| = p^{2r+t+3}.$$

Since  $x^p \in Z(G)$ , we may assume  $x^p = (a^{p^2} b^{-p^{t+2}})^i b^{jp^2}$ . Since

$$|\langle xa^{-ip}, b \rangle| = p^{r+t+3} < p^{2r+t+1} = \frac{|G|}{p^2},$$

we have  $[xa^{-ip}, b] = 1$  and hence  $xa^{-ip} \in Z(G)$ . Replacing  $x$  with  $xa^{-ip}$ , we get  $x^p = b^{j'p^2}$ . Since

$$|\langle xb^{-j'p}, a \rangle| = p^{r+3} < p^{2r+t+1} = \frac{|G|}{p^2},$$

we have  $[xb^{-j'p}, a] = 1$  and hence  $xb^{-j'p} \in Z(G)$ . Replacing  $x$  with  $xb^{-j'p}$ , we get  $x^p = 1$  and  $G$  is a group of Type (N6).

If  $s = 1$  and  $r \leq 2$ , then

$$p \geq 3, r = 2, Z(M) = \langle b^{p^2} \rangle, |M| = p^{t+6} \text{ and } |G| = p^{t+7}.$$

Since  $x^p \in Z(G)$ , we may assume  $x^p = b^{jp^2}$ . Since

$$|\langle xb^{-jp}, ab^{-p^t} \rangle| \leq p^4 < p^{t+5} = \frac{|G|}{p^2},$$

we have  $[xb^{-jp}, a] = 1$  and hence  $xb^{-jp} \in Z(G)$ . By replacing  $x$  with  $xb^{-jp}$  we get  $x^p = 1$  and  $G$  is a group of Type (N6).

Subcase 1.2:  $M$  is the group of Type (18) in Lemma 2.5. That is,  $M = \langle a, b, c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = a^p \rangle$ , where  $p \geq 5$ ,  $\nu$  is a fixed quadratic non-residue modulo  $p$ .

In this subcase,  $Z(M) = \langle a^p, b^p \rangle$ ,  $|M| = p^5$  and  $|G| = p^6$ . Since  $x^p \in Z(G)$ , we may assume  $x^p = a^{ip} b^{jp}$ . Since  $|\langle c, xa^{-i} b^{-j} \rangle| \leq p^3$ , we have  $[c, xa^{-i} b^{-j}] = 1$ . It follows that  $x^p = 1$ . Hence we get the group (N9).

Subcase 1.3:  $M$  is one of the groups (19)–(21) in Lemma 2.5.

By an argument similar to that of subcase 1.2, we get groups (N10)–(N12) respectively. The details is omitted.

**Case 2:**  $Z(G) < \Phi(G)$  and  $G$  has at least two three-generator maximal subgroups.

Let  $M_1, \dots, M_s$  be all three-generator maximal subgroups of  $G$ ,  $N = M'_1 M'_2 \dots M'_s$  and  $\bar{G} = G/N$ . Then  $\bar{G}$  has at least two abelian subgroups of index  $p$  and every non-abelian subgroup of  $\bar{G}$  is generated by two elements. Since  $\Phi(G) \not\leq Z(G)$ ,  $\bar{G}$  is not abelian. By Lemma 2.13 we have  $\bar{G} \in \mathcal{A}_2$ . Hence  $\bar{G}$  is a group of Type (8)–(12) in Lemma 2.5. Let  $\bar{G} = \langle \bar{a}, \bar{b}, \bar{x} \rangle$  and  $K = \langle a, b, \Phi(G) \rangle$ . Then  $d(K) = 2$  since  $\bar{K}$  is not abelian. Since  $d(G) = 3$ , we have  $\Phi(K) = \Phi(G)$ . It follows that  $K = \langle a, b \rangle$  is maximal in  $G$  and  $K \in \mathcal{A}_2$ . Hence  $K$  is a group of Type (1)–(7) or (17)–(21) in Lemma 2.5.

Subcase 2.1:  $K$  is a group of Type (1) in Lemma 2.5. That is,  $K = \langle a, b \mid a^8 = b^{2^m} = 1, a^b = a^{-1} \rangle$ .

If  $m = 1$ , then  $\Phi(G) = \Phi(K) = \langle a^2 \rangle$  and  $N = \langle a^4 \rangle$ . By calculation, a maximal subgroup  $\langle a, c \rangle$  of  $G$  is abelian or minimal non-abelian, a contradiction. Hence

$$m \geq 2, \Phi(G) = \Phi(K) = \langle a^2, b^2 \rangle, \langle a^4 \rangle \leq N \leq \langle a^4, b^{2^{m-1}} \rangle \text{ and } |G| = 2^{m+4}.$$

Let  $M = \langle x, a, b^2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $x^2 \in Z(K) \cap \Phi(M)$ , we may assume that  $x^2 = a^{4i} b^{4j}$ . By replacing  $x$  with  $xa^{2i} b^{-2j}$  we get  $x^2 = 1$ . Since  $[x, b^2] = [a, b^2] = 1$ ,  $[x, a] \neq 1$ . Since  $G \in \mathcal{A}_3$ ,  $|\langle x, a \rangle| = 2^{m+2}$ . Since  $|\langle x, a \rangle| \leq 2^5$ ,  $m \leq 3$ .

If  $m = 2$ , then  $N = \langle a^4 \rangle$ . We have  $[x, a] = a^4$  and  $[x, b] = a^{4k}$ . By replacing  $b$  with  $ba^k$  we get  $[x, b] = 1$ . Hence  $G$  is the group (N13).

If  $m = 3$ , then  $|\langle x, a \rangle| = 2^5$ . It follows that  $N = \langle a^4, b^4 \rangle$  and we may assume that  $[x, a] = b^4 a^{4i}$ . If  $[x, a] = b^4 a^4$ , then  $[x, b] = b^4$  or  $[x, ba] = b^4$ . It follows that  $|\langle x, b \rangle| = 2^4$  or  $|\langle x, ba \rangle| = 2^4$ , a contradiction. Hence  $[x, a] = b^4$ . By suitable replacement we get  $[x, b] = a^4$ . Hence  $G$  is the group (N14).

Subcase 2.2:  $K$  is a group of Type (2) in Lemma 2.5. That is,  $K = \langle a, b \mid a^8 = b^{2^m} = 1, a^b = a^3 \rangle$ .

If  $m = 1$ , then  $\Phi(G) = \Phi(K) = \langle a^2 \rangle$  and  $N = \langle a^4 \rangle$ . By calculation, a maximal subgroup  $\langle a, c \rangle$  of  $G$  is abelian or minimal non-abelian, a contradiction. Hence

$$m \geq 2, \Phi(G) = \Phi(K) = \langle a^2, b^2 \rangle, \langle a^4 \rangle \leq N \leq \langle a^4, b^{2^{m-1}} \rangle \text{ and } |G| = 2^{m+4}.$$

Let  $M = \langle x, a, b^2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $x^2 \in Z(K) \cap \Phi(M)$ , we may assume  $x^2 = a^{4i} b^{4j}$ . By replacing  $x$  with  $xa^{2i} b^{-2j}$  we get  $x^2 = 1$ . Since  $[x, b^2] = [a, b^2] = 1$ ,  $[x, a] \neq 1$ . If  $[x, a] = a^4$ , then, replacing  $b$  with  $bx$ , it is reduced to Subcase 2.1. Hence  $|\langle x, a \rangle| = 2^5$ ,  $m = 3$ ,  $N = \langle a^4, b^4 \rangle$  and we may assume  $[x, a] = b^4 a^{4i}$ . If  $[x, a] = b^4 a^4$ , then  $[x, b] = b^4$  or  $[x, ba] = b^4$ . It follows that  $|\langle x, b \rangle| = 2^4$  or  $|\langle x, ba \rangle| = 2^4$ ,

a contradiction. Hence  $[x, a] = b^4$ . By suitable replacement we get  $[x, b] = a^4$ . By replacing  $a$  and  $b$  with  $ax$  and  $bx$ , respectively, we get the group (N14) again.

Subcase 2.3:  $K$  is a group of Type (3) in Lemma 2.5. That is,  $K = \langle a, b \mid a^8 = 1, b^{2^m} = a^4, a^b = a^{-1} \rangle$ .

In this subcase,  $\Phi(G) = \Phi(K) = \langle a^2, b^2 \rangle$  and  $N = \langle a^4 \rangle$ . Let  $M = \langle x, a, b^2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $x^2 \in Z(K) \cap \Phi(M)$ , we may assume that  $x^2 = b^{4i}$ . By replacing  $x$  with  $xb^{-2j}$  we get  $x^2 = 1$ . Since  $[x, b^2] = [a, b^2] = 1$ ,  $[x, a] = a^4$ . It follows that  $m = 2$ . By suitable replacement we get  $[x, b] = 1$ . Hence  $G$  is the group (N15).

Subcase 2.4:  $K$  is a group of Type (4) in Lemma 2.5. That is,  $K = \langle a_1, b; a_2, a_3 \mid a_1^p = a_2^p = a_3^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_3, [a_3, b] = 1, [a_i, a_j] = 1 \rangle$ , where  $p \geq 5$  for  $m = 1$ ,  $p \geq 3$  and  $1 \leq i, j \leq 3$ .

If  $m = 1$ , then  $p \geq 5$  and  $N = \langle a_3 \rangle$ . Let  $M = \langle x, a_1, a_2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $[x, a_2] = [a_1, a_2] = 1$ ,  $[x, a_1] \neq 1$ . Without loss of generality assume  $[x, a_1] = a_3$ . By suitable replacement we get  $[x, b] = 1$ .

If  $x^p = 1$ , then  $G$  is the group (N16).

If  $x^p \neq 1$ , then we may assume that  $a_3 = x^{ip}$ , where  $(i, p) = 1$ . By replacing  $a_1$  and  $a_2$  with  $a_1^{i-1}$  and  $[a_1^{i-1}, b]$ , respectively, we get  $[a_2, b] = [x, a_1] = x^p$ . Hence  $G$  is the group (N17).

If  $m \geq 2$ , then

$$\Phi(G) = \Phi(K) = \langle a_2, a_3, b^p \rangle \text{ and } \langle a_3 \rangle \leq N \leq \langle a_3, b^{p^{m-1}} \rangle.$$

Let  $M = \langle x, a_1, a_2, a_3, b^p \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $x^p \in Z(K) \cap \Phi(M)$ , we may assume  $x^p = a_3^i b^{jp}$ . If  $(j, p) = 1$ , then  $\langle a_1, xb^{-j} \rangle \in \mathcal{A}_2$  and is of order  $p^4$ , a contradiction. Hence  $p \mid j$ . By replacing  $x$  with  $xb^{-j}$  we get  $x^p = a_3^i$ . Since  $[x, a_2] = [a_1, a_2] = 1$ ,  $[x, a_1] \neq 1$ . Since  $|\langle x, a_1 \rangle| \leq p^4$ , we have  $m = 2$ ,  $(i, p) = 1$  and  $[x, a_1] \notin \langle a_3 \rangle$ . It follows that  $N = \langle a_3, b^p \rangle$ . Without loss of generality assume that  $[a_2, b] = x^p$  and  $[x, a_1] = b^{jp} x^{kp}$ , where  $(j, p) = 1$ . Let  $[x, b] = b^{sp} x^{tp}$ . By replacing  $x$  and  $b$  with  $xa_2^{j^{-1}s-t}$  and  $ba_1^{-j^{-1}s}$ , respectively, we get  $[x, b] = 1$ . By replacing  $b$  with  $bx^{j^{-1}k}$  we get  $[x, a_1] = b^{jp}$ . By replacing  $b$  and  $x$  with  $b^{j^{-1}}$  and  $x^{j^{-2}}$ , respectively, we get  $[x, a_1] = b^p$ . Hence  $G$  is the group (N18).

Subcase 2.5:  $K$  is a group of Type (5) in Lemma 2.5. That is,  $K = \langle a_1, b; a_2 \mid a_1^p = a_2^p = b^{p^{m+1}} = 1, [a_1, b] = a_2, [a_2, b] = b^{p^m}, [a_1, a_2] = 1 \rangle$ , where  $p > 2$ .

In this subcase,  $N = \langle b^{p^m} \rangle$ . Let  $M = \langle x, a_1, a_2, b^p \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $[x, a_2] = [a_1, a_2] = 1$ ,  $[x, a_1] \neq 1$ . Without loss of generality assume  $[x, a_1] = b^{p^m}$ . Since  $x^p \in Z(K)$ , we may assume  $x^p = b^{ip}$ . If  $(i, p) = 1$ , then  $\langle a_1, xb^{-i} \rangle \in \mathcal{A}_2$  and is of order  $p^4$ . If  $p \mid i$ , then  $\langle a_1, xb^{-i} \rangle \in \mathcal{A}_1$  and is of order  $p^3$ . Hence  $m = 1$  and  $|G| = p^5$ . By suitable replacement we get  $[x, b] = 1$ .

If  $x^p = 1$ , then  $G$  is the group of Type (N19).



If  $x^p = b^{ip} \neq 1$ , then, replacing  $a_1$  and  $b$  with  $a_1^i$  and  $bx^{-i-1}$ , respectively, we get  $b^p = 1$  and  $[a_2, b] = [x, a_1] = x^p$ . Hence  $G$  is the group (N17).

Subcase 2.6:  $K$  is a group of Type (6) in Lemma 2.5. That is,  $K = \langle a_1, b \mid a_1^{p^2} = a_2^p = b^{p^m} = 1, [a_1, b] = a_2, [a_2, b] = a_1^{\nu p}, [a_1, a_2] = 1 \rangle$ , where  $p > 2$  and  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ .

If  $m = 1$ , then  $N = \langle a_1^p \rangle$ . Let  $M = \langle x, a_1, a_2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $[x, a_2] = [a_1, a_2] = 1$ ,  $[x, a_1] \neq 1$ . Without loss of generality assume  $[x, a_1] = a_1^p$ . By suitable replacement we get  $[x, b] = 1$ .

If  $x^p = 1$ , then  $G$  is the group (N20).

If  $x^p = a_1^{ip} \neq 1$ , then, by suitable replacement, we get the group (N17).

If  $m \geq 2$ , then

$$\Phi(G) = \Phi(K) = \langle a_2, a_1^p, b^p \rangle \text{ and } \langle a_1^p \rangle \leq N \leq \langle a_3, b^{p^{m-1}} \rangle.$$

Let  $M = \langle x, a_1, a_2, b^p \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $x^p \in Z(K) \cap \Phi(M)$ , we may assume that  $x^p = a_1^{ip} b^{jp}$ . If  $(j, p) = 1$ , then  $\langle a_1, x b^{-j} \rangle \in \mathcal{A}_2$  and is of order  $p^4$ , a contradiction. Hence  $p \mid j$ . By replacing  $x$  with  $x b^{-j}$  we get  $x^p = a_1^{ip}$ . If  $x^p \neq 1$ , then, replacing  $a_1$  with  $a_1 x^{-i-1}$ , it is reduced to Subcase 2.4. Hence we may assume  $x^p = 1$ . Since  $[x, a_2] = [a_1, a_2] = 1$ , we have  $[x, a_1] \neq 1$ . Since  $|\langle x, a_1 \rangle| \leq p^4$ , we have  $m = 2$  and  $[x, a_1] \notin \langle a_1^p \rangle$ . It follows that  $N = \langle a_1^p, b^p \rangle$  and  $[x, a_1] = b^{jp} a_1^{kp}$ , where  $(j, p) = 1$ . Let  $[x, b] = b^{sp} a_1^{tp}$ . By replacing  $x$  with  $x a_2^{-\nu-1t}$  we get  $[x, b] = b^{sp}$ . Since  $|\langle x, b \rangle| = p^3$ ,  $[x, b] = 1$ . Assume  $(ba_1)^p = b^p a_1^{rp}$ . By calculation we have  $[x^{j-1} a_2^{r-j-1k}, ba_1] = (ba_1)^p$ . It follows that  $\langle x^{j-1} a_2^{r-j-1k}, ba_1 \rangle \in \mathcal{A}_1$  and is of order  $p^3$ , a contradiction.

Subcase 2.7:  $K$  is a group of Type (7) in Lemma 2.5. That is,  $K = \langle a_1, b; a_2 \mid a_1^9 = a_2^3 = 1, b^3 = a_1^3, [a_1, b] = a_2, [a_2, b] = a_1^{-3} \rangle$ .

In this subcase,  $N = \langle a^3 \rangle = \langle b^3 \rangle$ . If  $x^p = b^{3i} \neq 1$ , then, replacing  $b$  with  $bx^{-i}$ , it is reduced to Subcase 2.6. Hence we may assume  $x^3 = 1$ . Let  $M = \langle x, a_1, a_2 \rangle$ . Then  $M \in \mathcal{A}_2$ . Since  $[x, a_2] = [a_1, a_2] = 1$ ,  $[x, a_1] \neq 1$ . Without loss of generality assume  $[x, a_1] = a_1^3$ . By suitable replacement we get  $[x, b] = 1$ . Hence  $G$  is the group (N21).

Subcase 2.8:  $K$  is a group of Type (17) in Lemma 2.5. That is,  $K = \langle a, b \mid a^{p^{r+2}} = 1, b^{p^{r+s+t}} = a^{p^{r+s}}, [a, b] = a^{p^r} \rangle$ , where  $r \geq 2$  for  $p = 2$ ,  $r \geq 1$  for  $p \geq 3$ ,  $t \geq 0$ ,  $0 \leq s \leq 2$  and  $r + s \geq 2$ .

In this subcase,

$$Z(K) = \langle a^{p^2}, b^{p^2} \rangle, |K| = p^{2r+s+t+2} \text{ and } |G| = p^{2r+s+t+3}.$$

If  $N = \langle a^{p^{r+1}} \rangle$ , then, assuming that  $[x, a] = a^{ip^{r+1}}$  and  $[x, b] = a^{jp^{r+1}}$ , we get  $x a^{-jp} b^{ip} \in Z(G)$ , a contradiction. Hence  $|N| = p^2$ . Since  $x^p \in Z(G)$ , we may assume that  $x^p = a^{ip^2} b^{jp^2}$ . By replacing  $x$  with  $x a^{-ip} b^{-jp}$  we get  $x^p = 1$ .

If  $s = 2$ , then

$$Z(K) = \langle a^{p^2} \rangle \times \langle b^{p^2} \rangle, |K| = p^{2r+t+4}, |G| = p^{2r+t+5} \text{ and } N = \langle a^{p^{r+1}}, b^{p^{r+t+1}} \rangle.$$

Without loss of generality assume  $[x, b] \neq 1$ . Since  $|\langle x, b \rangle| \leq p^{r+t+4}$ ,  $r+t+4 \geq 2r+t+3$ . It follows that  $r = 1$ ,  $p \geq 3$  and  $[x, b] \notin \langle b^{p^{t+2}} \rangle$ . We may assume  $[x, b] = a^{p^2} b^{kp^{t+2}}$ .

If  $[x, a] \neq 1$ , then  $t = 0$ ,  $|G| = p^7$  and  $[x, a] \notin \langle a^{p^2} \rangle$ . By suitable replacement we may assume that  $[x, b] = a^{p^2}$  and  $[x, a] = b^\nu p^2 a^{kp^2}$ , where  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ . By Lemma 5.14 we have  $G$  is the group (N22).

If  $[x, a] = 1$ , then  $(k, p) = 1$  since  $xa^{-p} \notin Z(G)$ . By replacing  $a$  and  $x$  with  $a^{k^{-1}}$  and  $x^{k^{-1}}$ , respectively, we have  $[x, b] = a^{p^2} b^{p^{t+2}}$ . If  $t = 0$ , then  $\langle x, ba \rangle$  is not abelian and is of order  $p^4$ , a contradiction. Hence  $t \geq 1$  and  $G$  is the group (N23).

If  $s = 1$ , then

$$|K| = p^{2r+t+3}, |G| = p^{2r+t+4}, r \geq 2 \text{ and } N = \langle a^{p^{r+1}}, a^{-p^r} b^{p^{r+t}} \rangle.$$

Since  $|\langle x, a^{-1} b^{p^t} \rangle| \leq p^{r+3} < p^{2r+t+2}$ ,  $[x, a^{-1} b^{p^t}] = 1$ . Since  $Z(G) < \Phi(G)$ ,  $[x, b] \neq 1$ . Since  $|\langle x, b \rangle| = p^{r+t+4} = p^{2r+t+2} = \frac{|G|}{p^2}$ ,  $r = 2$ . Without loss of generality assume  $[x, b] = a^{-p^2} b^{p^{t+2}} b^{ip^{t+3}}$ . By replacing  $a$  with  $axa^{-ip}$  we have  $[a, b] = b^{p^{t+2}}$ . By replacing  $a$  and  $b$  with  $b$  and  $a^{-1} b^{p^t} b^{ip^{t+1}}$ , respectively, we get

$$G = \langle a, b, x \mid a^{p^{t+4}} = b^{p^3} = x^p = 1, [a, b] = a^{p^{t+2}}, [x, b] = 1, [x, a] = b^{p^2} \rangle.$$

Hence  $G$  is a group of Type (N24).

If  $s = 0$ , then  $r \geq 2$ . By suitable replacement we have

$$K = \langle a, b \mid a^{p^{r+t+2}} = b^{p^r} = 1, [a, b] = a^{p^{r+t}} \rangle.$$

By calculation we have

$$Z(K) = \langle a^{p^2} \rangle \times \langle b^{p^2} \rangle, |K| = p^{2r+t+2}, |G| = p^{2r+t+3}, r \geq 3 \text{ and } N = \langle a^{p^{r+t+1}}, b^{p^{r-1}} \rangle.$$

Since  $|\langle x, a \rangle| = p^{r+t+4} = \frac{|G|}{p^2}$ ,  $r = 3$ . Without loss of generality assume  $[x, a] = b^{p^2} a^{kp^{t+4}}$ . By replacing  $b$  with  $ba^{kp^{t+2}}$  we get  $[x, a] = b^{p^2}$ . Hence  $G$  is a group of Type (N24).

Subcase 2.9:  $K$  is a group of Type (18) in Lemma 2.5. That is,  $K = \langle a, b \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^\nu p, [c, b] = a^p \rangle$ , where  $p \geq 5$ ,  $\nu$  is a fixed quadratic non-residue modulo  $p$ .

In this subcase,  $|G| = p^6$  and  $N = Z(K) = \langle a^p, b^p \rangle$ . We claim that  $x^p = 1$ . Otherwise, since  $x^p \in Z(K)$ , we may assume that  $x^p = a^{ip} b^{jp}$ , where  $(i, j) \neq (0, 0)$ . If  $i \neq 0$ , then  $\langle xa^{-i}, c \rangle$  is not abelian and is of order  $p^3$ , a contradiction. If  $j \neq 0$ , then  $\langle xb^{-j}, c \rangle$  is not abelian and is of order  $p^3$ , a contradiction again. Hence  $x^p = 1$ . Let  $[x, a] = a^{sp} b^{tp}$ . Since  $|\langle xc^{-\nu^{-1}t}, a \rangle| = p^3$ , we have  $[xc^{-\nu^{-1}t}, a] = 1$  and hence  $[x, a] = b^{tp}$ .

By replacing  $x$  with  $xc^{-\nu^{-1}t}$  we get  $[x, a] = 1$ . The same reason gives that  $[x, b] = a^{up}$ . Since  $Z(G) < \Phi(G)$ ,  $(u, p) = 1$ . Without loss of generality assume  $[x, b] = a^p$ . Then  $\langle cx^{\nu-1}, ab \rangle$  is not abelian and is of order  $p^3$ , a contradiction.

Subcase 2.10:  $K$  is a group of Type (19)–(21) in Lemma 2.5. That is,  $K = \langle a, b; c \mid a^{p^2} = b^{p^2} = c^p = 1, [a, b] = c, [c, a] = b^{\nu p}, [c, b] = a^p \rangle$ , where  $p \geq 5$ ,  $\nu$  is a fixed quadratic non-residue modulo  $p$ .

By an argument similar to that of subcase 2.9, we get a contradiction, respectively. The details is omitted.

**Case 3:**  $Z(G) < \Phi(G)$  and  $M$  is the unique three-generator maximal subgroup of  $G$ .

Then  $G/M'$  has a unique abelian subgroup  $M/M'$  of index  $p$ , and all non-abelian subgroups of  $G/M'$  are generated by two elements. By Lemma 2.13,  $G/M' \in \mathcal{A}_2$ . Hence  $G$  a group of Type (13)–(16) in Lemma 2.5.

Subcase 3.1:  $G/M'$  is a group of Type (13) in Lemma 2.5. That is,  $G/M' = \langle \bar{a}, \bar{b}, \bar{c} \mid \bar{a}^4 = \bar{b}^4 = 1, \bar{c}^2 = \bar{a}^2\bar{b}^2, [\bar{a}, \bar{b}] = \bar{b}^2, [\bar{c}, \bar{a}] = \bar{a}^2, [\bar{c}, \bar{b}] = 1 \rangle$ .

Let  $K = \langle a, b, \Phi(G) \rangle$ . Since  $K/M'$  is not abelian, we have  $d(K) = 2$  and  $K \in \mathcal{A}_2$ . Hence  $\Phi(K) = \Phi(G)$  and  $K = \langle a, b \rangle$ . By calculation we get

$$[c, a^2] = [c, a]^2[c, a, a] = a^4 \text{ and } [c^2, a] = [c, a]^2[c, a, c] = a^8 = 1.$$

Since  $c^2 \equiv a^2b^2 \pmod{M'}$ ,  $[a, b^2] = 1$ . It follows that  $b^4 = [a, b^2] = 1$  and hence  $\langle a, b \rangle \in \mathcal{A}_1$ , a contradiction.

Subcase 3.2:  $G/M'$  is a group of Type (14) in Lemma 2.5. That is,  $G/M' = \langle \bar{a}, \bar{b}, \bar{d} \mid \bar{a}^{p^m} = \bar{b}^{p^2} = \bar{d}^p = 1, [\bar{a}, \bar{b}] = \bar{a}^{p^{m-1}}, [\bar{d}, \bar{a}] = \bar{b}^p, [\bar{d}, \bar{b}] = 1 \rangle$ , where  $m \geq 3$  if  $p = 2$ .

Let  $K = \langle a, b, \Phi(G) \rangle$  and  $L = \langle d, a, \Phi(G) \rangle$ . Since  $K/M'$  and  $L/M'$  is not abelian, we have

$$d(K) = 2, d(L) = 2, K \in \mathcal{A}_2 \text{ and } L \in \mathcal{A}_2.$$

Hence

$$\Phi(K) = \Phi(L) = \Phi(G), K = \langle a, b \rangle \text{ and } L = \langle d, a \rangle.$$

By calculation,  $b^{p^2} = [d^p, a] = 1$ . Since  $L \in \mathcal{A}_2$ ,  $[a, b^p] \neq 1$ . It follows that  $a^{p^m} = [a, b^p] \neq 1$  and hence  $o(a) = p^{m+1}$ . Let  $N = \langle a^p, b \rangle$ . Then  $N \in \mathcal{A}_1$ . Since  $|N| = p^{m+2}$ , we get  $|G| = p^{m+4}$  and hence  $M' = \langle a^{p^m} \rangle$ . We claim that  $p > 2$ . Otherwise,  $\langle d, a^2 \rangle$  is not abelian and is of order  $2^{m+1}$ , a contradiction. Hence  $p > 2$ . Let  $d^p = a^{ip^m}$ . By replacing  $d$  with  $da^{-ip^{m-1}}$  we get  $d^p = 1$ . By suitable replacement we get  $[a, b] = a^{p^{m-1}}$  and  $[d, a] = b^p$ .

If  $[d, b] = 1$  then  $G$  is the group (N25).

If  $[d, b] \neq 1$ , then, since  $|\langle d, b \rangle| = p^4$ , we have  $m = 2$ . Assume that  $[d, b] = a^{jp^2}$ , where  $j = k^2\nu$ ,  $\nu = 1$  or a fixed quadratic non-residue modulo  $p$ . By replacing  $a$  and  $d$  with  $a^k$  and  $d^{k^{-1}}$ , respectively, we get  $[d, b] = a^{\nu p^2}$ . Hence  $G$  is the group (N26).

Subcase 3.3:  $G/M'$  is a group of Type (15) in Lemma 2.5. That is,  $\langle \bar{a}, \bar{b}, \bar{d} \mid \bar{a}^{p^m} = \bar{b}^{p^2} = \bar{d}^{p^2} = 1, [\bar{a}, \bar{b}] = \bar{d}^p, [\bar{d}, \bar{a}] = \bar{b}^{jp}, [\bar{d}, \bar{b}] = 1 \rangle$ , where  $(j, p) = 1$ ,  $p > 2$ ,  $j$  is a fixed quadratic non-residue modulo  $p$ , and  $-4j$  is a quadratic non-residue modulo  $p$ .

Let  $K = \langle a, b, \Phi(G) \rangle$  and  $L = \langle d, a, \Phi(G) \rangle$ . Since  $K/M'$  and  $L/M'$  is not abelian, we have

$$d(K) = 2, \quad d(L) = 2, \quad K \in \mathcal{A}_2 \text{ and } L \in \mathcal{A}_2.$$

Hence

$$\Phi(K) = \Phi(L) = \Phi(G), \quad K = \langle a, b \rangle \text{ and } L = \langle d, a \rangle.$$

By Lemma 2.6 (7),  $\exp(K') = \exp(L') = p$ . It follows that  $d^{p^2} = b^{p^2} = 1$ . By calculation we have  $[d^p, b] = [d, b]^p = 1$  and  $[d^p, a] = [d, a]^p = 1$ . Hence  $K \in \mathcal{A}_1$ , a contradiction.

Subcase 3.4:  $G/M'$  is a group of Type (16) in Lemma 2.5. That is,  $\langle \bar{a}, \bar{b}, \bar{d} \mid \bar{a}^{p^m} = \bar{b}^{p^2} = \bar{d}^{p^2} = 1, [\bar{a}, \bar{b}] = \bar{d}^p, [\bar{d}, \bar{a}] = \bar{b}^{jp}\bar{d}^p, [\bar{d}, \bar{b}] = 1 \rangle$ , where if  $p$  is odd, then  $4j = 1 - \rho^{2r+1}$  with  $1 \leq r \leq \frac{p-1}{2}$  and  $\rho$  the smallest positive integer which is a primitive root (mod  $p$ ); if  $p = 2$ , then  $j = 1$ .

Let  $K = \langle a, b, \Phi(G) \rangle$  and  $L = \langle d, a, \Phi(G) \rangle$ . Since  $K/M'$  and  $L/M'$  is not abelian, we have

$$d(K) = 2, \quad d(L) = 2, \quad K \in \mathcal{A}_2 \text{ and } L \in \mathcal{A}_2.$$

Hence

$$\Phi(K) = \Phi(L) = \Phi(G), \quad K = \langle a, b \rangle \text{ and } L = \langle d, a \rangle.$$

By Lemma 2.6 (7),  $\exp(K') = \exp(L') = p$ . It follows that  $d^{p^2} = b^{p^2} = 1$ . By calculation we have  $[d^p, b] = [d, b]^p = 1$  and  $[d^p, a] = [d, a]^p = 1$ . Hence  $K \in \mathcal{A}_1$ , a contradiction.

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 5.15 as follows. Since  $d(G) = 3$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1+p+p^2)$ . In the following, we calculate  $\alpha_1(G)$ .

**Case 1.**  $G$  is one of the groups (N1)–(N5).

Since  $\Phi(G) \leq Z(G)$ , by Theorem 3.18,  $\alpha_1(G) = \mu_1 + \mu_2 p^2 = p^4 + p^3 + p^2$ .

**Case 2.**  $G$  is one of the groups (N6)–(N12).

In this case,  $G = M * \langle x \rangle$  where  $M = \langle a, b \rangle$  such that  $\alpha_1(M) = 1+p$  and  $x^p \in Z(M)$ .

Other maximal subgroups of  $G$  are:

$N_i = M_i \langle x \rangle$  where  $M_i < M$ ;

$N_{ij} = \langle ax^i, bx^j, \Phi(M) \rangle$  where  $0 \leq i, j \leq p-1$ .

Since  $d(N_i) = 3$  and  $N'_i = M'_i$ , by Lemma 2.6 (7),  $\alpha_1(N_i) = p^2$ . Notice that  $G = N_{ij} * \langle x \rangle$ . If  $d(N_{ij}) = 3$ , then  $G' = N'_{ij} \leq Z(G)$  and  $\exp(G') = p$ . It follows that  $\Phi(G) \leq Z(G)$ , a contradiction. Hence  $d(N_{ij}) = 2$ . If  $N_{ij}$  has an abelian subgroup

of index  $p$ , then  $G$  also has an abelian subgroup of index  $p$ , a contradiction. Hence  $\alpha_1(N_{ij}) = 1 + p$ . Let  $H \in \Gamma_2$ . Then  $H = \langle xm, \Phi(G) \rangle$  where  $m \in M$  or  $H = \langle m, \Phi(G) \rangle$  where  $m \in M \setminus \Phi(G)$ . It is obvious that  $H' = 1$  if and only if  $H = \langle x, \Phi(G) \rangle$ . Thus  $\sum_{H \in \Gamma_2} \alpha_1(H) = p + p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = (1+p) \times p^2 + p^2 \times (1+p) - p \times (p+p^2) = p^3 + p^2.$$

**Case 3.**  $G$  is one of the groups (N13)–(N21) except for (N14) and (N18).

In this case,  $G = M * \langle x \rangle$  where  $\langle x \rangle \cong C_p$  and  $M = \langle a, b \rangle$  such that  $d(M) = 2$ ,  $|M'| = p^2$ ,  $[x, b] = 1$ ,  $[x, a] \in M_3$  and  $\langle a, \Phi(M) \rangle$  is the unique abelian maximal subgroup of  $G$ . Hence all maximal subgroups are:

$$N_i = K_i * \langle x \rangle \text{ where } K_i \text{ are maximal subgroups of } M;$$

$$N_{ij} = \langle ax^i, bx^j \rangle \text{ where } 0 \leq i, j \leq p-1.$$

It is easy to see that  $K'_i = M_3$  and hence  $|N'_i| = p$ . By Lemma 2.6,  $\alpha_1(N_i) = p^2$ . Since  $N_{ij} \cong M$ ,  $\alpha_1(N_{ij}) = p$ . Let  $H \in \Gamma_2$ . Then  $H = \langle a^i b^j x^k, \Phi(G) \rangle$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $j \neq 0$ . Hence  $\sum_{H \in \Gamma_2} \alpha_1(H) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = (1+p) \times p^2 + p^2 \times p - p \times p^2 = p^3 + p^2.$$

**Case 4.**  $G$  is either the group (N14) or (N18).

In this case,  $G = M * \langle x \rangle$  where  $\langle x \rangle \cong C_p$  and  $M = \langle a, b \rangle$  such that  $d(M) = 2$ ,  $|M'| = p^2$ ,  $[x, b] \in M_3$ ,  $[x, a] \notin M'$  and  $\langle a, \Phi(M) \rangle$  is the unique abelian maximal subgroup of  $G$ . Hence all maximal subgroups are:

$$N = \langle b, x, \Phi(M) \rangle;$$

$$N_i = \langle ab^i, x, \Phi(M) \rangle \text{ where } 0 \leq i, j \leq p-1.$$

$$N_{ij} = \langle ax^i, bx^j \rangle \text{ where } 0 \leq i, j \leq p-1.$$

It is easy to see that  $N' = M_3$ . By Lemma 2.6 (7),  $\alpha_1(N) = p^2$ . By calculation,  $|N'_0| = p$  and  $|N'_i| = p^2$  for  $i \neq 0$ . Hence  $\alpha_1(N_0) = p^2$  and  $\alpha_1(N_i) = p^2 + p$  for  $i \neq 0$ . Since  $N_{ij} \cong M$ ,  $\alpha_1(N_{ij}) = p$ . Let  $H \in \Gamma_2$ . Then  $H = \langle a^i b^j x^k, \Phi(G) \rangle$ . It is obvious that  $H \in \mathcal{A}_1$  if and only if  $j \neq 0$ . Hence  $\sum_{H \in \Gamma_2} \alpha_1(H) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = 2 \times p^2 + (p-1) \times (p^2 + p) + p^2 \times p - p \times p^2 = p^3 + 2p^2 - p.$$

**Case 5.**  $G$  is one of the groups (N22)–(N24).

In this case,  $p > 2$ ,  $G = M * \langle x \rangle$  where  $\langle x \rangle \cong C_p$  and  $M = \langle a, b \rangle$  such that  $M$  is metacyclic,  $[x, b] \in \Phi(M')$ ,  $[x, a] \notin M'$ . Hence all maximal subgroups are:

$$N = \langle b, x, \Phi(M) \rangle;$$

$$N_i = \langle ab^i, x, \Phi(M) \rangle \text{ where } 0 \leq i, j \leq p-1.$$

$N_{ij} = \langle ax^i, bx^j \rangle$  where  $0 \leq i, j \leq p-1$ .

It is easy to see that  $N' = \Phi(M')$ . By Lemma 2.6 (7),  $\alpha_1(N) = p^2$ . By calculation,  $|N'_i| = p^2$ . Hence  $\alpha_1(N_i) = p^2 + p$ . Since  $N_{ij} \cong M$ ,  $\alpha_1(N_{ij}) = 1 + p$ . Let  $H \in \Gamma_2$ . Then  $H = \langle a^i b^j x^k, \Phi(G) \rangle$ . It is obvious that  $H \notin \mathcal{A}_1$  if and only if  $i = j = 0$ . Hence  $\sum_{H \in \Gamma_2} \alpha_1(H) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p^2 + p \times (p^2 + p) + p^2 \times (1 + p) - p \times (p^2 + p) = p^3 + 2p^2.$$

**Case 6.**  $G$  is one of the groups (N25)–(N26).

It is easy to verify that  $M = \langle b, d, a^p \rangle$  is the unique three-generator maximal subgroup of  $G$ . Since  $|M'| = p$ , by Lemma 2.6,  $\alpha_1(N) = p^2$ . Other maximal subgroups are:

$N_i = \langle ab^i, d \rangle$  where  $0 \leq i \leq p-1$ .

$N_{ij} = \langle ad^i, bd^j \rangle$  where  $0 \leq i, j \leq p-1$ .

It is easy to see that  $N_i$  has a unique abelian maximal subgroup  $\langle d, a^p, b^p \rangle$  and  $N_{ij}$  has no abelian maximal subgroup. Hence  $\alpha_1(N_i) = p$  and  $\alpha_1(N_{ij}) = 1 + p$ . Let  $H \in \Gamma_2$ . Then  $H = \langle a^i b^j d^k, \Phi(G) \rangle$  where  $(i, j, k) \neq (0, 0, 0)$ . It is obvious that  $H \notin \mathcal{A}_1$  if and only if  $i = j = 0$ , hence if and only if  $H = \langle d, \Phi(G) \rangle$ . Thus  $\sum_{H \in \Gamma_2} \alpha_1(H) = p^2 + p$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = p^2 + p \times p + p^2 \times (1 + p) - p \times (p^2 + p) = 2p^2.$$

□

**Theorem 5.16.** *Suppose that  $G$  is an  $\mathcal{A}_3$ -group having no abelian subgroup of index  $p$ . Then  $d(G) = 4$  if and only if  $G$  is isomorphic to one of the following pairwise non-isomorphic groups:*

(Oi)  $G' \cong C_p$  and  $c(G) = 2$ . In this case,  $\Phi(G) = Z(G) = G'$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2 + p^3)$  and  $\alpha_1(G) = p^2 + p^4$ .

(O1)  $D_8 * Q_8$ ;

(O2)  $Q_8 * Q_8$ ;

(O3)  $M_p(1, 1, 1) * M_p(2, 1)$ , where  $p > 2$ ;

(O4)  $M_p(1, 1, 1) * M_p(1, 1, 1)$ , where  $p > 2$ .

(Oii)  $G' \cong C_p^2$  and  $c(G) = 2$ .

(O5)  $G = \langle a, b, c, d \mid a^4 = b^4 = 1, c^2 = a^2, d^2 = b^2, [a, b] = 1, [a, c] = b^2, [b, c] = a^2, [a, d] = a^2, [b, d] = a^2 b^2, [c, d] = 1 \rangle$ ; here  $|G| = 2^6$ ,  $\Phi(G) = Z(G) = G' = \langle a^2, b^2 \rangle \cong C_2^2$ , any two noncommutative elements of  $G$  generate  $M_2(2, 2)$ .  $(\mu_0, \mu_1, \mu_2) = (0, 0, 15)$  and  $\alpha_1(G) = 30$ .

(Oiii)  $G' \cong C_p^3$  and  $c(G) = 2$ .

(O6)  $G = K \times \langle a_4 \rangle$  where  $K = \langle a_1, a_2, a_3 \mid a_1^4 = a_2^4 = a_3^4 = 1, [a_1, a_2] = a_3^2, [a_1, a_3] = a_2^2 a_3^2, [a_2, a_3] = a_1^2 a_2^2, [a_1^2, a_2] = [a_2^2, a_1] = 1 \rangle$  and  $\langle a_4 \rangle \cong C_2$ ; here  $|G| = 2^7$ ,  $\Phi(G) = G' = \langle a_1^2, a_2^2, a_3^2 \rangle$ ,  $Z(G) = \langle a_1^2, a_2^2, a_3^2, a_4 \rangle \cong C_2^4$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 15)$  and  $\alpha_1(G) = 30$ .

**Proof** Let the type of  $G/G'$  be  $(p^{m_1}, p^{m_2}, p^{m_3}, p^{m_4})$ , where  $m_1 \geq m_2 \geq m_3 \geq m_4$ , and  $G/G' = \langle a_1 G' \rangle \times \langle a_2 G' \rangle \times \langle a_3 G' \rangle \times \langle a_4 G' \rangle$ , where  $o(a_i G') = p^{m_i}$ ,  $i = 1, 2, 3, 4$ . Then  $G = \langle a_1, a_2, a_3, a_4 \rangle$ . By Lemma 3.17,  $c(G) = 2$ ,  $\Phi(G) \leq Z(G)$ ,  $G' \leq C_p^3$  and all  $\mathcal{A}_1$ -subgroups of  $G$  contain  $\Phi(G)$ . The last property gives that  $\sum_{H \in \Gamma_2} \alpha_1(H) = \alpha_1(G)$ .

We claim that  $m_1 = 1$ . Otherwise,  $m_1 \geq 2$ . Let  $B = \langle a_2, a_3, a_4 \rangle$ . Since  $|G : B| \geq p^2$ , we deduce that  $B' = 1$ . Hence  $A = \langle B, a_1^p \rangle$  is an abelian subgroup of index  $p$  of  $G$ , a contradiction. Hence  $G/G'$  is elementary abelian.

**Case 1:**  $G' \cong C_p$ .

In this case,  $|G| = p^5$ . We claim that  $Z(G) = G'$ . Otherwise, without loss of generality assume  $a_1 \in Z(G)$ . Let  $B = \langle a_2, a_3, a_4 \rangle$ . Then  $B' \cong C_p$ . By Lemma 2.6 (3),  $|Z(B)| = p^2$ . Since  $Z(G) \geq \langle a_1, Z(B) \rangle$ ,  $|Z(G)| \geq p^3$ . Hence  $G$  has an abelian subgroup of index  $p$  of  $G$ , a contradiction.

By above argument,  $G$  is an extraspecial  $p$ -group. Hence we get groups (O1)–(O4).

**Case 2:**  $G' \cong C_p^2$ .

In this case,  $|G| = p^6$  and any two noncommutative elements generate an  $\mathcal{A}_1$ -group of order  $p^4$ . Such groups were classified by [4]. By checking those groups listed in [4] we get the group (O5).

**Case 3:**  $G' \cong C_p^3$ .

In this case,  $|G| = p^7$  and any  $\mathcal{A}_1$ -subgroup is  $M_p(2, 2, 1)$ . It follows that  $\Omega_1(G) \leq Z(G)$ . Let  $N \leq G'$  such that  $|N| = p$  and  $\bar{G} = G/N$ . Then  $|\bar{G}'| = p^2$ . By Lemma 2.16, there exists  $\bar{K} \leq \bar{G}$  such that  $d(\bar{K}) = 3$  and  $\bar{K}' = \bar{G}'$ . Without loss of generality assume  $K = \langle a_1, a_2, a_3 \rangle$ . It follows that  $|K'| \geq p^2$  and  $G' = \langle a_1^p, a_2^p, a_3^p \rangle$ .

Subcase 3.1:  $K' \cong C_p^3$ .

By Lemma 2.5,

$$K = \langle a_1, a_2, a_3 \mid a_1^4 = a_2^4 = a_3^4 = 1, [a_1, a_2] = a_3^2, [a_1, a_3] = a_2^2 a_3^2, [a_2, a_3] = a_1^2 a_2^2 \rangle.$$

Since  $a_4^2 \in G' = K'$ , there exists  $c \in K$  such that  $a_4^2 = c^2$ . Since  $|\langle a_4, c \rangle| \leq 16$ ,  $[a_4, c] = 1$ . By replacing  $a_4$  with  $a_4 c$  we get  $a_4^2 = 1$ . Hence  $a_4 \in \Omega_1(G) \leq Z(G)$  and we get the group (O6).

Subcase 3.2:  $K' \cong C_p^2$ .

Without loss of generality assume  $K' = \langle [a_1, a_2], [a_1, a_3] \rangle$  and  $[a_2, a_3] = 1$ . If  $G' = \langle K', [a_1, a_4] \rangle$ , then there exists  $b \in \langle a_2, a_3, a_4 \rangle$  such that  $[a_1, b] = a_1^p$ , which contradicts that  $G' \leq \langle a_1, b \rangle$ . Hence  $[a_1, a_4] \in K'$ . By suitable replacement we get  $[a_1, a_4] = 1$ . Without loss of generality assume  $G' = \langle K', [a_2, a_4] \rangle$ . By replacing  $a_3$  with  $a_3a_4$  we have  $K' = G'$ . It is reduced to subcase 3.1.

We calculate the  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  of those groups in Theorem 5.16 as follows.

Since  $d(G) = 4$ ,  $(\mu_0, \mu_1, \mu_2) = (0, 0, 1 + p + p^2 + p^3)$ . In the following, we calculate  $\alpha_1(G)$ .

**Case 1.**  $G$  is one of the groups (O1)–(O4).

Let  $H \in \Gamma_1$ . Then  $|H'| = p$ . By Lemma 2.6 (7),  $\alpha_1(H) = p^2$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = (1 + p + p^2 + p^3) \times p^2 - p\alpha_1(G).$$

Hence

$$\alpha_1(G) = \frac{(1 + p + p^2 + p^3)p^2}{1 + p} = p^2 + p^4.$$

**Case 2.**  $G$  is the group (O5).

By calculation,  $d(H) = 3$  and  $H' \cong C_2^2$  for any  $H \in \Gamma_1$ . Hence  $\alpha_1(H) = 6$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = 15 \times 6 - 2\alpha_1(G).$$

Hence  $\alpha_1(G) = 30$ .

**Case 3.**  $G$  is the group (O6).

Let  $H \in \Gamma_1$ . Then  $H$  is one of the following types: (1)  $H = M \times \langle a_4 \rangle$  where  $M \triangleleft K$ ; (2)  $H = \langle a_1a_4^i, a_2a_4^j, a_3a_4^k \rangle$  where  $i, j, k = 0, 1$ . If  $H$  is of Type (1), then  $|H'| = 2$  and hence  $\alpha_1(H) = 4$  by Lemma 2.6. If  $H$  is of Type (2), then  $H \cong K$  and hence  $\alpha_1(H) = 7$ . By Hall's enumeration principle,

$$\alpha_1(G) = \sum_{H \in \Gamma_1} \alpha_1(H) - p \sum_{H \in \Gamma_2} \alpha_1(H) = 7 \times 4 + 8 \times 7 - 2\alpha_1(G).$$

Hence  $\alpha_1(G) = 28$ . □

Now we list the triple  $(\mu_0, \mu_1, \mu_2)$  and  $\alpha_1(G)$  for  $\mathcal{A}_3$ -groups in Table 13 and 14 respectively. This solves Problem 893 in [8], Problem 1595 in [9] and Problem 2829 in [10] respectively.

At the end of this paper, we list some enumeration properties of  $\mathcal{A}_3$ -groups, which can easily be obtained from Table 14.



**Theorem 5.17.** *Let  $G$  be an  $\mathcal{A}_3$ -group. Then  $p^2 \leq \alpha_1(G) \leq p^4 + p^3 + p^2 + p$ , and the following conclusions hold.*

(1) *If  $\alpha_1(G) = p^2$ , then  $d(G) = 2$ ,  $c(G) = 4$  and  $G$  has an abelian maximal subgroup. In this case, non-abelian subgroups of  $G$  are generated by two elements;*

(2) *If  $\alpha_1(G) = p^4 + p^3 + p^2 + p$ , then  $p = 2$ ,  $c(G) = 2$ ,  $d(G) = 4$  and  $G = K \times C_2$  where  $K$  is the smallest Suzuki 2-group;*

(3) *If  $p > 2$ , then  $\alpha_1(G) \leq p^4 + p^3 + p^2$ ;*

(4) *If  $d(G) = 2$ , then  $\alpha_1(G) \leq p^3 + 2p^2 + p$ ;*

(5) *If  $d(G) = 2$  and  $\alpha_1(G) \leq p^3 + 2p^2 + p$ , then  $p = 2$  and  $G$  is the group (M37) in Theorem 5.13;*

(6) *If  $d(G) \geq 3$ , then  $\alpha_1(G) \geq 2p^2 - 1$ .*

**Acknowledgments** During writing the paper, Professor Y. Berkovich gave us many correspondences, we cordially express thanks for his encouragements and many valuable comments. All these helped us to improve the whole paper considerably.

$(\mu_0, \mu_1, \mu_2)$	types of $\mathcal{A}_3$ groups
$(1, p-1, 1)$	(A1)–(A6)
$(p+1, p^2-1, 1)$	(B1); (B2) where $m = n = 1$ ; (B3) where $n = 1$ ; (B4); (B5) where $n = 1$
$(p+1, p^2-p, p)$	(B2) where $m > 1 = n$ or $n > 1 = m$ ; (B3) where $n > 1$ ; (B5) where $n > 1$
$(1, p^2-1, p+1)$	(B6); (B9); (B11); (B13) where $p = 2$ and $m = l = 1$ (B17) where $l = 1$ ; (B19) where $p = 2$ and $m = l = 1$
$(1, p^2, p)$	(B7) where $l > 1$ ; (B10); (B12) where $l > 1$ ; (B13) where $m = 1$ and $l > 1$ (B14) where $n = 1$ or $m = 1$ ; (B16); (B18) where $m = 1$ (B19) where $p = 2$ and $l > 1 = m$ ; (B19) where $p > 2$ and $m = 1$
$(1, p^2+p-1, 1)$	(B7) where $l = 1$ ; (B12) where $l = 1$ ; (B13) where $p > 2$ and $m = l = 1$ ; (B20)
$(1, p^2+p-2, 2)$	(B8) where $l = 1$
$(1, p^2-p, 2p)$	(B8) where $l > 1$ ; (B13) where $m = 2$ ; (B14) where $n = m = 2$ (B15); (B17) where $l > 1$ ; (B18) where $m = 2$ ; (B19) where $m = 2$
$(0, p, 1)$	(C1)–(C6), (C8), (C9), (C11), (C15)–(C17)
$(0, p-1, 2)$	(C7), (C10), (C12)–(C14)
$(0, p^2, p+1)$	(D1), (D4), (D6)–(D10), (D12)–(D14), (D16)
$(0, p^2+p, 1)$	(D2) where $-\nu \notin (F_p^*)^2$ ; (D3) where $-r \notin (F_p^*)^2$ ; (D5); (D11) where $-\nu \notin (F_p^*)^2$ ;
$(0, p^2-p, 2p+1)$	(D2) where $-\nu \in (F_p^*)^2$ ; (D3) where $-\nu \in (F_p^*)^2$ ; (D11) where $-\nu \in (F_p^*)^2$ ; (D15)
$(0, p^2-1, p+2)$	(D17), (D18)
$(0, p^2+1, p)$	(D19)
$(0, 1, p)$	(E1)–(E7)
$(0, 1, p^2+p)$	(E8)–(E10)
$(1, 0, p)$	(F1)–(F8); (G1)–(G12)
$(p+1, 0, p^2)$	(H1)–(H3)
$(1, 0, p^2+p)$	(H4)–(H10), (I1)–(I11)
$(p+1, 0, p^3+p^2)$	(J1)–(J5)
$(1, 0, p^3+p^2+p)$	(J6)–(J9)
$(0, 0, p+1)$	(K1)–(K5); (L1)–(L2); (M1)–(M62)
$(0, 0, p^2+p+1)$	(N1)–(N26)
$(0, 0, p^3+p^2+p+1)$	(O1)–(O6)

Table 13: The number of  $\mathcal{A}_0$ -,  $\mathcal{A}_1$ -,  $\mathcal{A}_2$ -subgroups of index  $p$  in  $\mathcal{A}_3$ -groups

$\alpha_1(G)$	types of $\mathcal{A}_3$ -groups $G$
$p^2$	(F1)–(F8)
$p^2 + 1$	(E1), (E3)–(E7)
$p^2 + p - 1$	(A1)–(A6)
$p^2 + p$	(C1)–(C6), (C8)–(C9), (C11), (C15), (C17), (K3)–(K5)
$p^2 + p + 1$	(K1)–(K2)
$p^2 + 2p$	(C16)
$2p^2 - 1$	(B1), (B2) where $n = m = 1$ , (B3) where $n = 1$ , (B4), (B5) where $n = 1$
$2p^2$	(M48)–(M53), (M58)–(M62), (N25)–(N26)
$2p^2 + p - 1$	(B7) where $l = 1$ , (B12) where $l = 1$ , (B13) where $p > 2$ and $l = m = 1$ , (B20) (C7), (C10), (C12)–(C14)
$2p^2 + p$	(D2) where $-\nu \notin (F_p^*)^2$ ; (D3) where $-r \notin (F_p^*)^2$ ; (D5); (D11) where $-\nu \notin (F_p^*)^2$ ; (M54)–(M57)
$3p^2 + 1$	(D19)
$3p^2 + p - 2$	(B8) where $l = 1$
$3p^2 + p$	(L1)–(L2)
$p^3$	(G1)–(G12), (I1)–(I9)
$p^3 + 1$	(E2), (E8)
$p^3 + p^2 - p$	(B2) where $m > 1 = n$ or $n > 1 = m$ ; (B3) where $n \geq 2$ , (B5) where $n \geq 2$ (B7) where $l \geq 2$ , (B10), (B12) where $l \geq 2$ , (B13) where $l \geq 2$ and $m = 1$
$p^3 + p^2$	(B14) where $m = 1$ or $n = 1$ , (B16), (B18), (B18) where $m = 1$ , (B19) where $m = 1$ and $p > 2$ (B19) where $p = 2$ and $m = 1$ and $l > 1$ , (I10)–(I11), (M1)–(M19), (M40)–(M43), (M47) (N6)–(N13), (N15)–(N17), (N19)–(N21)
$p^3 + p^2 + p$	(M38)–(M39)
$p^3 + 2p^2 - p$	(N14), (N18)
$p^3 + 2p^2 - 1$	(B6), (B9), (B11), (B13) where $p = 2$ and $l = m = 1$ (B17) where $l = 1$ , (B19) where $p = 2$ and $l = m = 1$
$p^3 + 2p^2$	(D1), (D4), (D6)–(D10), (D12)–(D14), (D16), (M20)–(M36), (M44)–(M46), (N22)–(N24)
$p^3 + 2p^2 + p$	(M37)
$p^3 + 3p^2 - 1$	(D17)–(D18)
$2p^3 + p^2 - p$	(B8) where $l \geq 2$ , (B13) where $m = 2$ , (B14) where $m = n = 2$ , (B15) (B17) where $l \geq 2$ , (B18) where $m = 2$ , (B19) where $m = 2$
$2p^3 + 2p^2 - p$	(D2) where $-\nu \in (F_p^*)^2$ ; (D3) where $-\nu \in (F_p^*)^2$ ; (D11) where $-\nu \in (F_p^*)^2$ ; (D15)
$p^4$	(H1)–(H3), (J1)–(J5)
$p^4 + p^2$	(O1)–(O4)
$p^4 + p^3 + 1$	(E9)–(E10)
$p^4 + p^3$	(H4)–(H10), (J6)–(J9)
$p^4 + p^3 + p^2$	(N1)–(N5), (O6)
$p^4 + p^3 + p^2 + p$	(O5)

Table 14: The number of  $\mathcal{A}_1$ -subgroups in  $\mathcal{A}_3$ -groups

## References

- [1] L.J. An, J.F. Ding and Q.H. Zhang, Finite self dual group, *J. Algebra*, 341(2011), 35–44.
- [2] L.J. An, L.L. Li, H.P. Qu and Q.H. Zhang, Finite  $p$ -groups with a minimal non-abelian subgroup of index  $p$  (II), *Sci. China Ser.A*, 57:4(2014), 737–753.
- [3] L.J. An, R.F. Hu and Q.H. Zhang, Finite  $p$ -groups with a minimal non-abelian subgroup of index  $p$  (IV), *J. Algebra Appl.*, accepted. DOI: 10.1142/S0219498815500206. <http://arxiv.org/pdf/1310.5503v2.pdf>
- [4] L.J. An and J. Peng, Finite  $p$ -groups in which any two noncommutative elements generate an inner abelian group of order  $p^4$ , *Algebra Colloq.*, 20:2(2013), 215–226.

- [5] Y. Berkovich, Finite  $p$ -groups with few minimal nonabelian subgroups, *J. Algebra*, 297 (2006), 62–100.
- [6] Y. Berkovich and Z. Janko, Structure of finite  $p$ -groups with given subgroups, *Contemp. Math.* 402, Amer. Math. Soc., Providence, RI. (2006), 13–93.
- [7] Y. Berkovich, Groups of Prime Power Order Vol. 1, *Walter de Gruyter · Berlin · New York*, 2008.
- [8] Y. Berkovich and Z. Janko, Groups of Prime Power Order Vol. 2, *Walter de Gruyter · Berlin · New York*, 2008.
- [9] Y. Berkovich and Z. Janko, Groups of Prime Power Order Vol. 3, *Walter de Gruyter · Berlin · New York*, 2011.
- [10] Y. Berkovich and Z. Janko, Groups of Prime Power Order Vol. 4, *in preparation*.
- [11] Hans Ulrich Besche, Bettina Eick and E.A. O'Brien, A millennium project: constructing small groups, *International J. of Algebra and Computation*, 12:5(2002), 623–644.
- [12] N. Blackburn, On prime power groups with two generators, *Proc. Cambridge Phil. Soc.*, **54**(1957), 327–337.
- [13] N. Blackburn, On prime-power groups in which the derived group has two generators, *Proc. Cambridge Phil. Soc.* **53**(1957), 19–27.
- [14] S.V. Draganyuk, On the structure of finite primary groups all 2-maximal subgroups of which are abelian (Russian), in *Complex analysis, Algebra and topology*, Akad. Nauk Ukrain. SSR, Inst. Mat., Kiev.(1990), 42–51.
- [15] G. Higman, Enumerating  $p$ -groups I: Problems whose solution is PORC, *Proc. London Math.Soc.*, **10:3**(1960), 24–30.
- [16] G. Higman, Enumerating  $p$ -groups II: Inequalities, *Proc. London Math.Soc.*, **10:3**(1960), 566–582.
- [17] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, 1967.
- [18] L.S. Kazarin, On certain classes of finite groups, *Dokl. Akad. Nauk SSSR (Russian)*, **197**(1971), 773–776.
- [19] P.J. Li, H.P. Qu and J.W.Zeng, Finite  $p$ -groups whose proper subgroups are of class  $\leq n$ , *submitted*.
- [20] G.A. Miller, H.C. Moreno, Non-abelian groups in which every subgroup is abelian, *Trans. Amer. Math. Soc.*, **4**(1903), 398–404.

- [21] E.A.O'Brien and M.Vaughan-Lee, The groups with order  $p^7$  for odd prime  $p$ , *J. Algebra*, 292(2005), 243–258.
- [22] H.P. Qu, S.S. Yang, M.Y. Xu and L.J. An, Finite  $p$ -groups with a minimal non-abelian subgroup of index  $p$  (I), *J. Algebra*, **358**(2012), 178–188.
- [23] H.P. Qu, M.Y. Xu and L.J. An, Finite  $p$ -groups with a minimal non-abelian subgroup of index  $p$  (III), *Sci. China Ser.A*, accepted. <http://arxiv.org/pdf/1310.5496v2.pdf>
- [24] H.P. Qu, L.P. Zhao, J. Gao and L.J. An, Finite  $p$ -groups with a minimal non-abelian subgroup of index  $p$  (V), *J. Algebra Appl.*, 13:7 (2014), 1450032(35 pages).
- [25] L.Rédei, Das schiefe Produkt in der Gruppen theorie, *Comment. Math. Helvet.*, **20**(1947), 225–267.
- [26] V.A. Sheriev, A description of the class of finite  $p$ -groups whose 2-maximal subgroups are all abelian II, in Primary groups, *Proc. Sem. Algebraic Systems*, **2**(1970), 54–76.
- [27] M.Y. Xu, A theorem on metabelian  $p$ -groups and some consequences, *Chin. Ann. Math.* **5B** (1984), 1–6.
- [28] M.Y. Xu, A complete classification of metacyclic  $p$ -groups of odd order (Chinese), *Adv. in Math. (Beijing)* **12**(1983), 72-73.
- [29] M.Y. Xu, L.J. An and Q.H. Zhang, Finite  $p$ -groups all of whose non-abelian proper subgroups are generated by two elements, *J. Algebra*, 319(2008), 3603–3620.
- [30] M.Y. Xu and Q.H. Zhang, A classification of metacyclic 2-groups, *Algebra Colloq.*, 131(2006), 25–34.
- [31] Q.H. Zhang, X.J. Sun, L.J. An and M.Y. Xu, Finite  $p$ -groups all of whose subgroups of index  $p^2$  are abelian, *Algebra Colloq.*, **15:1**(2008), 167–180.