

The hadron structure and the description of the elastic scattering in a wide region of t and s

O.V. Selyugin

BLTP, JINR, Dubna, Russia

Abstract. On the basis of the new high energy general structure (HEGS) model, which takes into account the different moments of the General Parton Distributions (GPDs) of the hadron, the quantitative descriptions of all existing experimental data from $\sqrt{s} = 9.8$ GeV to 7 TeV, including the Coulomb range and large momentum transfers up to $-t = 15$ GeV², are obtained with only 3 free fitting high energy parameters. The real part of the hadronic amplitude is determined only through complex s satisfying the dispersion relations. The negligible contributions of the hard Pomeron and non-small contributions of the maximal Odderon is shown. Predictions for LHC energies are made.

Keywords: Elastic hadron scattering, hadron form factors, general parton distributions

PACS: 13.40.Gp, 14.20.Dh, 12.38.Lg

INTRODUCTION

Now we present a model based on the assumption that the hadron interaction is sensitive to the generalized parton distributions, whose moments can be represented in the form of two different distributions over momentum transfer: charge and matter separately. Hence, this model uses the exact electromagnetic and matter form-factors determined by one function - generalized parton distributions (GPDs). So both form factors are independent of the fitting procedure of the differential elastic cross sections. Note that the form of the GPDs is determined, on the one hand, by the deep-inelastic processes and, on the other hand, by the measure of the electromagnetic form factor from the electron-nucleon elastic scattering. This picture is supported by a good description of the experimental data in the Coulomb-hadron interference region and a large momentum transfer at high energies by one amplitude with a few free parameters. The differential cross sections of the nucleon-nucleon elastic scattering can be written as a sum of different helicity amplitudes [1]:

$$\frac{d\sigma}{dt} = \frac{2\pi}{s^2} (|\Phi_1|^2 + |\Phi_2|^2 + |\Phi_3|^2 + |\Phi_4|^2 + 4|\Phi_5|^2). \quad (1)$$

The total helicity amplitudes can be written as $\Phi_i(s,t) = F_i^h(s,t) + F_i^{\text{em}}(s,t)e^{\varphi(s,t)}$, where $F_i^h(s,t)$ comes from the strong interactions, $F_i^{\text{em}}(s,t)$ from the electromagnetic interactions and $\varphi(s,t)$ is the interference phase factor between the electromagnetic and strong interactions [2]. For the hadron part the amplitude with spin-flip is neglected in this approximation, as usual at high energy.

THE HIGH ENERGY GENERAL STRUCTURE (HEGS) MODEL

We suppose that at high energies a hadron interaction in the non-perturbative regime is determined by the reggenized-gluon exchange. The cross-even part of this amplitude can have 2 non-perturbative parts, possible standard pomeron (P_{2np}) and cross-even part of the 3-non-perturbative gluons (P_{3np}). The interaction of these two objects is proportional to two different form factors of the hadron. This is the main assumption of the model. The second important assumption is that we chose the slope of the second term 4 times smaller than the slope of the first term, by analogy with the two pomeron cut. Both terms have the same intercept. The electromagnetic form factors can be represented as the first moments of GPDs

$$F_1(t) = \int_0^1 dx \sum_{u,d} \mathcal{H}^q(x,t); \quad F_2(t) = \int_0^1 dx \sum_{u,d} \mathcal{E}^q(x,t), \quad (2)$$

In [3] the t -dependence of GPDs was chosen in the form

$$\mathcal{H}^q(x,t) = q(x)_{nf} \exp[a_+ \frac{(1-x)^2}{x^m} t]; \quad \mathcal{E}^q(x,t) = q(x)_{sf} \exp[a_- \frac{(1-x)^2}{x^m} t]; \quad (3)$$

The function $q(x)$ was taken at the scale $\mu^2 = 1$ and was based on the MRST2002 global fit [4]. For $\xi = 0$ one has

$$\int_0^1 dx x \sum_{u,d} [\mathcal{H}(x,t) \pm \mathcal{E}(x,t)] = A(t) \pm B(t). \quad (4)$$

The integration of the second momentum of GPDs over x gave the momentum-transfer representation of the form factor. It was approximated by the dipole form [5] $A(t) = L_2^4 / (L_2^2 - t)^2$. Hence, the Born term of the elastic hadron amplitude can be written as

$$F_h^{Born}(s,t) = h_1 G^2(t) F_a(s,t) (1 + r_1/s^{0.5}) + h_2 A^2(t) F_b(s,t) (1 + r_2/s^{0.5}) \quad (5)$$

where $F_a(s,t)$ and $F_b(s,t)$ has the standard Regge form

$$F_a(s,t) = \hat{s}^{\epsilon_1} e^{B(s)t}; \quad F_b(s,t) = \hat{s}^{\epsilon_1} e^{B(s)/4 t}, \quad (6)$$

The slope of the scattering amplitude has the standard logarithmic dependence on the energy. $B(s) = \alpha_1 \ln(\hat{s})$, with $\alpha_1 = 0.24 \text{ GeV}^{-2}$ and $\hat{s} = s e^{-i\pi/2}/s_0$; $s_0 = 1 \text{ GeV}^2$. The final elastic hadron scattering amplitude is obtained after unitarization of the Born term. So, first, we have to calculate the eikonal phase

$$\chi(s,b) = \frac{1}{2\pi} \int d^2q e^{i\vec{b}\cdot\vec{q}} F_h^{Born}(s,q^2), \quad (7)$$

and then obtain the final hadron scattering amplitude

$$F_h(s,t) = is \int b J_0(bq) \Gamma(s,b) db; \quad \Gamma(s,b) = 1 - \exp[-\chi(s,b)]. \quad (8)$$

All these calculations are carried out by the FORTRAN program. The model has only 3 high energy fitting parameters and 2 low energy parameters, which reflect some small contribution coming from the different low energy terms. We take all existing experimental data in the energy range $52.8 \leq \sqrt{s} \leq 1960 \text{ GeV}$ and the region of the momentum transfer $0.0008 \leq -t \leq 9.75 \text{ GeV}^2$ of the elastic differential cross sections of proton-proton and proton-antiproton data [6]. As a result, one obtains $\sum \chi_i^2 / N \simeq 1.8$ where $N = 975$ is the number of experimental points. Note that the parameters of the model are energy-independent. The energy dependence of the scattering amplitude is determined only by the single intercept and the logarithmic dependence on s of the slope.

EXTENSION OF THE HEGS MODEL

Further development of the model requires careful analysis of the t form of GPDs and a properly chosen form of PDFs. In [7], the analysis of more than 20 different PDFs was made. We slightly complicated the form of GPDs

$$\mathcal{H}^u(x,t) = q(x)_{nf} \exp[2a_+ \frac{(1-x)^{2+\epsilon_1}}{x^m} t]; \quad \mathcal{H}_1^d(x,t) = q(x)_{nf} \exp[2a_+ (\frac{(1-x)^{2+\epsilon_2} + dx(1-x)}{x^m}) t]. \quad (9)$$

$$\mathcal{E}^u(x,t) = q(x)_{nf} \exp[2a_+ \frac{(1-x)^{2+\epsilon_1}}{x^m} t]; \quad \mathcal{E}_1^d(x,t) = q(x)_{nf} \exp[2a_+ (\frac{(1-x)^{2+\epsilon_2} + dx(1-x)}{x^m}) t]. \quad (10)$$

On the basis of this analysis we calculated the nucleon electromagnetic form factor and the matter form factors. The fit of these calculations gave the parameters of the electromagnetic $G(t)$ and matter form factors $A(t)$.

Then, as we intend to describe sufficiently low energies, the possible odderon contributions were taken into account.

$$F_{odd}^{Born}(s,t) = \pm i h_{odd} A^2(t) \frac{t}{1 - r_0^2 t} F_b(s,t). \quad (11)$$




FIGURE 1. $d\sigma/dt$ a) for pp (left panel) at $\sqrt{s} = 9.8$ GeV and b) $p\bar{p}$ (central panel) at $\sqrt{s} = 11.3$ GeV and c) pp (right panel) at $\sqrt{s} = 7$ TeV [cross - TOTEM data [10], circles - ATLAS data [11]].

As we supposed in the previous variant of the HEGS model that the $F_b(s,t)A(t)^2$ correspond to the cross-even part of the three gluon exchange, our odderon contribution is also connected with the matter form factor $A(t)$. Our ansatz for the Odderon slightly differs from the cross-even part by some kinematic factor. The form of the odderon works in all t and has the same behavior at non-small t as the cross-even part, of course, with changing sign for proton-proton and proton-antiproton reactions.

We added a small contribution of the asymptotic energy independent part of the spin flip amplitude in the form similar to the proposed in [?]. $F_{sf}(s,t) = h_{sf}q^3G^2(t)$. It has only one additional free parameter.

In our new description of the experimental data of the proton-proton and proton-antiproton we took the energy region from $\sqrt{s} > 9.8$ GeV. It gave us many new high precision experimental data especially at small t in the CNI region. Hence, we need to more careful by examine the slope of the scattering amplitude. Taking into account the work [8] we take some small nonlinear form of the slope of the scattering amplitude with the standard logarithmic dependence on the energy. $B(s) = (\alpha_1 + qk_0e^{k_0t}) \ln(\hat{s})$, with $\alpha_1 = 0.24$ GeV $^{-2}$ and $\Delta = 0.11$; $s_0 = 4m_p^2$ - fixed. Our complete fit of 3088 experimental data in the energy range $9.8 \leq \sqrt{s} \leq 7000$ GeV and the region of the momentum transfer $0.00045 \leq -t \leq 14.75$ GeV 2 gave $\sum_{i=1}^N \chi_i^2/N = 1.33$ with the parameters $h_1 = 0.82$; $h_2 = 0.31$; $h_{odd} = 0.15$; $k_0 = 0.17$; $r_0^2 = 3.9$ and the low energy parameters $h_{sf} = 0.05$; $R_1 = 51$; $R_2 = 4.4$.

The model reproduces $d\sigma/dt$ at very small and large t and provides a qualitative description of the dip region at $-t \approx 1.4$ GeV 2 , for $\sqrt{s} = 53$ GeV 2 and for $\sqrt{s} = 62.1$ GeV 2 . Noteworthy is a good description of the CNI region in a very wide energy region, approximately 3 orders, with the same slope of the scattering amplitude. The differential cross sections of the elastic scattering pp and $p\bar{p}$ at small t and different s are presented in Fig.1. for $\sqrt{s} = 9.8$ GeV and 7 TeV (pp) and $\sqrt{s} = 11$. GeV ($p\bar{p}$). The model quantitatively reproduces the differential cross sections in the whole examined energy region in spite of the size of the slope changing essentially in this region (due to the standard Regge behavior $\log(\hat{s})$) and of the real part of the scattering amplitude having the different behavior for pp and $p\bar{p}$.

In Fig. 2, the description of the differential cross sections of the elastic scattering pp and $p\bar{p}$ at large t and different s is presented. It should be noted that the calculations of our integrals with complex oscillation functions at large t is a difficult task and require high precision of the calculations. In any case, we obtain a quantitatively good description of the differential cross sections at large t . In this region of t the contribution of the spin-flip amplitude is visible. We take into account only the asymptotic part of this amplitude with the simplest and energy independent forms. Although it has a small size, its constant is determined sufficiently well $h_{sf} = 0.05 \pm 0.002$.

CONCLUSIONS

We present the new model of the hadron-hadron interaction at high energies. The model is very simple from the viewpoint of the number of parameters and functions. There are no any artificial functions or some cuts which bound the separate parts of the amplitude by some region of momentum transfer or energy. One of the most remarkable properties is that the real part of the hadron scattering amplitude is determined only by complex energy \hat{s} that satisfies the crossing-symmetries. The new HEGS model gives the quantitatively description of the elastic nucleon scattering at high energy with only 6 fitting high energy parameters. Our model of GPDs leads to the well description of the proton and neutron electromagnetic form factors and its elastic scattering simultaneously. The successful description




FIGURE 2. $d\sigma/dt$ at large t a) for pp (left panel) at $\sqrt{s} = 27.43$ GeV and $p\bar{p}$ (central panel) at $\sqrt{s} = 1.8$ TeV and the model predictions at $\sqrt{s} = 7$ and 14 TeV.

of the existing experimental data by the model show that the elastic scattering determined by the generalized structure of the hadron. The model leads to a good coincidence of the model calculations with the preliminary data at 7 TeV. We find that the standard eikonal approximation [13] works perfectly well from $\sqrt{s} = 9$ GeV up to 7 TeV. The extending variant of the model shows the contribution of the "maximal" odderon with specific kinematic properties and does not show a visible contribution of the hard pomeron [14].

The slope of the differential cross sections at small t has a small peculiarity and has the same properties in the whole energy region examined. Such a uniform picture for the slope gives a possibility to further research the small peculiarity of the different cross sections like some possible oscillations [1, 15, 16]. Note that we do not see contributions of the second Reggions with a large slope and intercept above 0.5.

Including the data of the TOTEM Collaboration in our fit increases σ_{tot} at $\sqrt{s} = 7$ TeV from 95 mb [5] to 97 mb now.

ACKNOWLEDGMENTS

O.S. would like to thank the organizers R. Fiore and A. Papa for the invitation and support of his participation in the conference.

REFERENCES

1. R. Fiore, L. Jenkovszky, R. Orava, E. Predazzi, A. Prokudin, O. Selyugin, Mod.Phys., A24: 2551-2559 (2009).
2. O.V. Selyugin, Phys. Rev. D **60** 074028 (1999).
3. O.V. Selyugin, O.V. Teryaev, Phys.Rev. **D 79** 033003 (2009).
4. A.D. Martin *et al.*, Phys. Lett. B **531** 216 (2002).
5. O.V. Selyugin, Eur. Phys. J., **C 72** 2073-2081 (2012).
6. Spires Durham data base, (<http://durpdg.dur.ac.uk/hepdata>).
7. O.V. Selyugin, Phys. Rev. **D 89** 093007 (2014).
8. L.L. Jenkovszky, Phys.Part.Nucl. **34** 619 (2003); Fiz.Elem.Chas.Atom.Yadra **34** 1196 (2003).
9. The TOTEM Collaboration (G. Antchev *et al.*) EPL, **95** 41001 (2011).
10. The ATLAS Collaboration, arXiv:1408.5778.
11. O.V. Selyugin, Particles & Nuclei, **45** 39 (2014).
12. O.V. Selyugin, Nucl.Phys. A **922** 180 (2014).
13. J.-R. Cudell, E. Predazzi, O. V. Selyugin, Phys.Rev. **D 79** , 034033 (2009).
14. O. V. Selyugin, Nucl.Phys., **100** (2013). Nucl.Phys. A **903** 54 (2013).
15. P. Gauron, B. Nicolescu, J.V. Selyugin, Phys.Lett. **B397** 305 (1997).
16. O. V. Selyugin, J.-R. Cudell, talk presented at the International Workshop on Diffraction in High-Energy Physics, Otranto (Lecce, Italy), September 10-15, (2010); hep-ph/1011.4177.