

R-matrix and Mickelsson algebras for orthosymplectic quantum groups

Thomas Ashton and Andrey Mudrov

Department of Mathematics,

University of Leicester,

University Road, LE1 7RH Leicester, UK

e-mail: am405@le.ac.uk

Abstract

Let \mathfrak{g} be a complex orthogonal or symplectic Lie algebra and $\mathfrak{g}' \subset \mathfrak{g}$ the Lie subalgebra of rank $\text{rk } \mathfrak{g}' = \text{rk } \mathfrak{g} - 1$ of the same type. We give an explicit construction of generators of the Mickelsson algebra $Z_q(\mathfrak{g}, \mathfrak{g}')$ in terms of Chevalley generators via the R-matrix of $U_q(\mathfrak{g})$.

Mathematics Subject Classifications: 81R50, 81R60, 17B37.

Key words: Mickelsson algebras, quantum groups, R-matrix, lowering/raising operators.

1 Introduction

In the mathematics literature, lowering and raising operators are known as generators of step algebras, which were originally introduced by Mickelsson [1] for reductive pairs of Lie algebras, $\mathfrak{g}' \subset \mathfrak{g}$. These algebras naturally act on \mathfrak{g}' -singular vectors in $U(\mathfrak{g})$ -modules and are important in representation theory, [2, 3].

The general theory of step algebras for classical universal enveloping algebras was developed in [2, 4] and was extended to the special linear and orthogonal quantum groups in [5]. They admit a natural description in terms of extremal projectors, [4], introduced for classical groups in [6, 7] and extended to the quantum group case in [8]. It is known that the step algebra $Z(\mathfrak{g}, \mathfrak{g}')$ is generated by the image of the orthogonal complement $\mathfrak{g} \ominus \mathfrak{g}'$ under the extremal projector of the \mathfrak{g}' . Another description of lowering/raising operators for classical groups was obtained in [9, 10, 11, 12] in an explicit form of polynomials in \mathfrak{g} .

A generalization of the results of [9, 10] to quantum $\mathfrak{gl}(n)$ can be found in [13]. In this special case, the lowering operators can be also conveniently expressed through "modified commutators" in the Chevalley generators of $U(\mathfrak{g})$ with coefficients in the field of fractions of $U(\mathfrak{h})$. Extending [11, 12] to a general quantum group is not straightforward, since there are no immediate candidates for the nilpotent triangular Lie subalgebras \mathfrak{g}_\pm in $U_q(\mathfrak{g})$. We suggest such a generalization, where the lack of \mathfrak{g}_\pm is compensated by the entries of the universal R-matrix with one leg projected to the natural representation. Those entries are nicely expressed through modified commutators in the Chevalley generators turning into elements of \mathfrak{g}_\pm in the quasi-classical limit. Their commutation relation with the Chevalley generators modify the classical commutation relations with \mathfrak{g}_\pm in a tractable way. This enabled us to generalize the results of [9, 10, 11, 12] and construct generators of Mickelsson algebras for the non-exceptional quantum groups.

1.1 Quantized universal enveloping algebra

In this paper, \mathfrak{g} is a complex simple Lie algebra of type B , C or D . The case of $\mathfrak{gl}(n)$ can be easily derived from here due to the natural inclusion $U_q(\mathfrak{gl}(n)) \subset U_q(\mathfrak{g})$, so we do not pay special attention to it. We choose a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ with the canonical inner product (\cdot, \cdot) on \mathfrak{h}^* . By R we denote the root system of \mathfrak{g} with a fixed subsystem of positive roots $R^+ \subset R$ and the basis of simple roots $\Pi^+ \subset R^+$. For every $\lambda \in \mathfrak{h}^*$ we denote by h_λ its image under the isomorphism $\mathfrak{h}^* \simeq \mathfrak{h}$, that is $(\lambda, \beta) = \beta(h_\lambda)$ for all $\beta \in \mathfrak{h}^*$. We put $\rho = \frac{1}{2} \sum_{\alpha \in R^+} \alpha$ for the Weyl vector.

Suppose that $q \in \mathbb{C}$ is not a root of unity. Denote by $U_q(\mathfrak{g}_\pm)$ the \mathbb{C} -algebra generated by $e_{\pm\alpha}$, $\alpha \in \Pi^+$, subject to the q -Serre relations

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_{\alpha_i}} e_{\pm\alpha_i}^{1-a_{ij}-k} e_{\pm\alpha_j} e_{\pm\alpha_i}^k = 0,$$

where $a_{ij} = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$, $i, j = 1, \dots, n = \text{rk } \mathfrak{g}$, is the Cartan matrix, $q_\alpha = q^{\frac{(\alpha, \alpha)}{2}}$, and

$$\begin{bmatrix} m \\ k \end{bmatrix}_q = \frac{[m]_q!}{[k]_q! [m-k]_q!}, \quad [m]_q! = [1]_q \cdot [2]_q \cdots [m]_q.$$

Here and further on, $[z]_q = \frac{q^z - q^{-z}}{q - q^{-1}}$ whenever $q^{\pm z}$ make sense.

Denote by $U_q(\mathfrak{h})$ the commutative \mathbb{C} -algebra generated by $q^{\pm h_\alpha}$, $\alpha \in \Pi^+$. The quantum group $U_q(\mathfrak{g})$ is a \mathbb{C} -algebra generated by $U_q(\mathfrak{g}_\pm)$ and $U_q(\mathfrak{h})$ subject to the relations

$$q^{h_\alpha} e_{\pm\beta} q^{-h_\alpha} = q^{\pm(\alpha, \beta)} e_{\pm\beta}, \quad [e_\alpha, e_{-\beta}] = \delta_{\alpha, \beta} \frac{q^{h_\alpha} - q^{-h_\alpha}}{q_\alpha - q_\alpha^{-1}}.$$

Remark that \mathfrak{h} is not contained in $U_q(\mathfrak{g})$, still it is convenient for us to keep reference to \mathfrak{h} .

Fix the comultiplication in $U_q(\mathfrak{g})$ as in [14]:

$$\begin{aligned} \Delta(e_\alpha) &= e_\alpha \otimes q^{h_\alpha} + 1 \otimes e_\alpha, & \Delta(e_{-\alpha}) &= e_{-\alpha} \otimes 1 + q^{-h_\alpha} \otimes e_{-\alpha}, \\ \Delta(q^{\pm h_\alpha}) &= q^{\pm h_\alpha} \otimes q^{\pm h_\alpha}, \end{aligned}$$

for all $\alpha \in \Pi^+$.

The subalgebras $U_q(\mathfrak{b}_\pm) \subset U_q(\mathfrak{g})$ generated by $U_q(\mathfrak{g}_\pm)$ over $U_q(\mathfrak{h})$ are quantized universal enveloping algebras of the Borel subalgebras $\mathfrak{b}_\pm = \mathfrak{h} + \mathfrak{g}_\pm \subset \mathfrak{g}$.

The Chevalley generators e_α can be extended to a set of higher root vectors e_β for all $\beta \in R$. A normally ordered set of root vectors generate a Poincaré-Birkhoff-Witt (PBW) basis of $U_q(\mathfrak{g})$ over $U_q(\mathfrak{h})$, [14]. We will use \mathfrak{g}_\pm to denote the vector space spanned by $\{e_{\pm\beta}\}_{\beta \in R^+}$.

The universal R -matrix is an element of a certain extension of $U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$. We heavily use the intertwining relation

$$\mathcal{R}\Delta(x) = \Delta^{op}(x)\mathcal{R}, \tag{1.1}$$

between the coproduct and its opposite for all $x \in U_q(\mathfrak{g})$. Let $\{\varepsilon_i\}_{i=1}^n \subset \mathfrak{h}^*$ be the standard orthonormal basis and $\{h_{\varepsilon_i}\}_{i=1}^n$ the corresponding dual basis in \mathfrak{h} . The exact expression for \mathcal{R} can be extracted from [14], Theorem 8.3.9, as the ordered product

$$\mathcal{R} = q^{\sum_{i=1}^n h_{\varepsilon_i} \otimes h_{\varepsilon_i}} \prod_{\beta} \exp_{q_\beta} \{(1 - q_\beta^{-2})(e_\beta \otimes e_{-\beta})\} \in U_q(\mathfrak{b}_+) \hat{\otimes} U_q(\mathfrak{b}_-), \tag{1.2}$$

where $\exp_q(x) = \sum_{k=0}^{\infty} q^{\frac{1}{2}k(k+1)} \frac{x^k}{[k]_q!}$.

We use the notation $e_i = e_{\alpha_i}$ and $f_i = e_{-\alpha_i}$ for $\alpha_i \in \Pi^+$, in all cases apart from $i = n$, $\mathfrak{g} = \mathfrak{so}(2n+1)$, where we set $f_n = [\frac{1}{2}]_q e_{-\alpha_n}$. The reason for this is two-fold. Firstly, the natural representation can be defined through the classical assignment on the generators, as given below. Secondly, we get rid of $q_{\alpha_n} = q^{\frac{1}{2}}$ and can work over $\mathbb{C}[q]$, as the relations involved turn into

$$[e_n, f_n] = \frac{q^{h_{\alpha_n}} - q^{-h_{\alpha_n}}}{q - q^{-1}},$$

$$f_n^3 f_{n-1} - (q+1+q^{-1})f_n^2 f_{n-1} f_n + (q+1+q^{-1})f_n f_{n-1} f_n^2 - f_{n-1} f_n^3 = 0.$$

It is easy to see that the square root of q disappears from the corresponding factor in the presentation (1.2).

In what follows, we regard $\mathfrak{gl}(n) \subset \mathfrak{g}$ to be the Lie subalgebra with the simple roots $\{\alpha_i\}_{i=1}^{n-1}$ and $U_q(\mathfrak{gl}(n))$ the corresponding quantum subgroup in $U_q(\mathfrak{g})$.

Consider the natural representation of \mathfrak{g} in the vector space \mathbb{C}^N . We use the notation $i' = N+1-i$ for all integers $i = 1, \dots, N$. The assignment

$$\pi(e_i) = e_{i,i+1} \pm e_{i',i'-1}, \quad \pi(f_i) = e_{i+1,i} \pm e_{i',i'-1}, \quad \pi(h_{\alpha_i}) = e_{ii} - e_{i+1,i+1} + e_{i',i'-1} - e_{i'i'},$$

for $i = 1, \dots, n-1$, defines a direct sum of two representations of $\mathfrak{gl}(n)$ for each sign. It extends to the natural representation of the whole \mathfrak{g} by

$$\pi(e_n) = e_{n,n+1} \pm e_{n',n'}, \quad \pi(f_n) = e_{n+1,n} \pm e_{n',n'-1}, \quad \pi(h_{\alpha_n}) = e_{nn} - e_{n'n'},$$

$$\pi(e_n) = e_{nn'}, \quad \pi(f_n) = e_{n'n}, \quad \pi(h_{\alpha_n}) = 2e_{nn} - 2e_{n'n'},$$

$$\pi(e_n) = e_{n-1,n'} \pm e_{n,n'+1}, \quad \pi(f_n) = e_{n',n-1} \pm e_{n'+1,n}, \quad \pi(h_{\alpha_n}) = e_{n-1,n-1} + e_{nn} - e_{n'n'} - e_{n'+1,n'+1},$$

respectively, for $\mathfrak{g} = \mathfrak{so}(2n+1)$, $\mathfrak{g} = \mathfrak{sp}(2n)$, and $\mathfrak{g} = \mathfrak{so}(2n)$.

Two values of the sign give equivalent representations. The choice of minus corresponds to the standard representation that preserves the bilinear form with entries $C_{ij} = \delta_{i'j}$, for $\mathfrak{g} = \mathfrak{so}(N)$, and $C_{ij} = \text{sign}(i' - i)\delta_{i'j}$, for $\mathfrak{g} = \mathfrak{sp}(N)$. However, we fix the sign to $+$ in order to simplify calculations. The above assignment also defines representations of $U_q(\mathfrak{g})$.

2 R -matrix of non-exceptional quantum groups

Define $\check{\mathcal{R}} = q^{-\sum_{i=1}^n h_{\varepsilon_i} \otimes h_{\varepsilon_i}} \mathcal{R}$. Denote by $\check{R}^- = (\pi \otimes \text{id})(\check{\mathcal{R}}) \in \text{End}(\mathbb{C}^N) \otimes U_q(\mathfrak{g}_-)$ and by $\check{R}^+ = (\pi \otimes \text{id})(\check{\mathcal{R}}_{21}) \in \text{End}(\mathbb{C}^N) \otimes U_q(\mathfrak{g}_+)$. In this section, we deal only with \check{R}^- and suppress the label "–" for simplicity, $\check{R} = \check{R}^-$.

Denote by N_+ the ring of all upper triangular matrices in $\text{End}(\mathbb{C}^N)$ and by N'_+ its ideal spanned by e_{ij} , $i < j + 1$.

Lemma 2.1. *One has*

$$\check{R} = 1 \otimes 1 + (q^{1+\delta_{1n}} - q^{-1-\delta_{1n}}) \sum_{i=1}^n \pi(e_i) \otimes f_i \pmod{N'_+ \otimes U_q(\mathfrak{g}_-),}$$

where δ_{1n} is present only for $\mathfrak{g} = \mathfrak{sp}(2n)$.

Proof. For all positive roots α, β the matrix $\pi(e_\alpha e_\beta)$ belongs to N'_+ . Also, $\pi(e_\beta) \in N'_+$ for all $\beta \in R^+ \setminus \Pi^+$. Therefore, the only terms that contribute to $\text{Span}_{\varepsilon_i - \varepsilon_j \in \Pi^+} \{e_{ij} \otimes U_q(\mathfrak{g}_-)\}$ are those of degree 1 from the series $\exp_{q_\alpha}(1 - q_\alpha^{-2})(e_\alpha \otimes e_{-\alpha})$ with $\alpha \in \Pi^+$. \square

Write $\check{R} = \sum_{i,j=1}^N e_{ij} \otimes \check{R}_{ij}$, where $\check{R}_{ij} = 0$ for $i > j$. Due to the \mathfrak{h} -invariance of \check{R} , the entry $\check{R}_{ij} \in U_q(\mathfrak{g}_-)$ carries weight $\varepsilon_j - \varepsilon_i$.

For all \mathfrak{g} , we have $f_{k,k+1} = f_k = f_{k'-1,k'}$ once $k < n$ and $f_{n,n+1} = f_n = f_{n+1,n'}$ for $\mathfrak{g} = \mathfrak{so}(2n+1)$, $f_{n-1,n'} = f_n = f_{n,n'+1}$ for $\mathfrak{g} = \mathfrak{so}(2n)$, and $f_{nn'} = [2]_q f_n$ for $\mathfrak{g} = \mathfrak{sp}(2n)$. We present explicit expressions for the entries f_{ij} in terms of modified commutators in Chevalley generators, $[x, y]_a = xy - ayx$, where a is a scalar; we also put $\bar{q} = q^{-1}$.

Proposition 2.2. *Suppose that $\varepsilon_i - \varepsilon_j \in R^+ \setminus \Pi^+$. Then the elements f_{ij} are given by the following formulas:*

For all \mathfrak{g} and $i + 1 < j \leq \frac{N+1}{2}$:

$$f_{ij} = [f_{j-1}, \dots [f_{i+1}, f_i]_{\bar{q}} \dots]_{\bar{q}}, \quad f_{j'i'} = [\dots [f_i, f_{i+1}]_{\bar{q}}, \dots f_{j-1}]_{\bar{q}}. \quad (2.3)$$

Furthermore,

- For $\mathfrak{g} = \mathfrak{so}(2n+1)$: $f_{nn'} = (q-1)f_n^2$ and

$$f_{i,n+1} = [f_n, f_{i,n}]_{\bar{q}}, \quad f_{n+1,i'} = [f_{n',i'}, f_n]_{\bar{q}}, \quad i < n,$$

$$f_{ij'} = q^{\delta_{ij}} [f_{n+1,j'}, f_{i,n+1}]_{\bar{q}^{\delta_{ij}}}, \quad i, j < n.$$

- For $\mathfrak{g} = \mathfrak{sp}(2n)$: $f_{nn'} = [2]_q f_n$ and

$$f_{in'} = [f_n, f_{in}]_{\bar{q}^2}, \quad f_{n'i'} = [f_{n',i'}, f_n]_{\bar{q}^2}, \quad i < n,$$

$$f_{ij'} = q^{\delta_{ij}} [f_{n,j'}, f_{in}]_{\bar{q}^{1+\delta_{ij}}}, \quad i, j < n.$$

- For $\mathfrak{g} = \mathfrak{so}(2n)$: $f_{nn'} = 0$ and

$$f_{in'} = [f_n, f_{i,n-1}]_{\bar{q}}, \quad f_{ni'} = [f_{n'+1,i'}, f_n]_{\bar{q}}, \quad i < n - 2,$$

$$f_{ji'} = q^{\delta_{ij}} [f_{ni'}, f_{j,n}]_{\bar{q}^{1+\delta_{ij}}}, \quad i, j \leq n - 1.$$

Proof. The proof is a direct calculation with the use of the identity

$$(f_\alpha \otimes 1)\check{\mathcal{R}} - \check{\mathcal{R}}(f_\alpha \otimes 1) = \check{\mathcal{R}}(q^{-h_\alpha} \otimes f_\alpha) - (q^{h_\alpha} \otimes f_\alpha)\check{\mathcal{R}},$$

which follows from the intertwining axiom (1.1) for $x = f_\alpha$. This allows us to construct the elements f_{ij} by induction starting from f_α , $\alpha \in \Pi^+$. \square

For each $\alpha \in \Pi^+$, denote by $P(\alpha)$ the set of ordered pairs $l, r = 1, \dots, N$, with $\varepsilon_l - \varepsilon_r = \alpha$. We call such pairs simple.

Proposition 2.3. *The matrix entries $f_{i,j} \in U_q(\mathfrak{g}_-)$ such that $\varepsilon_i - \varepsilon_j \notin \Pi^+$ satisfy the identity*

$$[e_\alpha, f_{ij}] = \sum_{(l,r) \in P(\alpha)} (f_{il} \delta_{jr} q^{h_\alpha} - q^{-h_\alpha} \delta_{il} f_{rj}),$$

for all simple positive roots α .

Proof. The proof is a straightforward calculation based on the intertwining relation (1.1), which is equivalent to

$$(1 \otimes e_\alpha)\check{\mathcal{R}} - \check{\mathcal{R}}(1 \otimes e_\alpha) = \check{\mathcal{R}}(e_\alpha \otimes q^{h_\alpha}) - (e_\alpha \otimes q^{-h_\alpha})\check{\mathcal{R}},$$

for $x = e_\alpha$, $\alpha \in \Pi^+$. Alternatively, one can use the expressions for f_{ij} from Proposition 2.2. \square

3 Mickelsson algebras

Consider the Lie subalgebra $\mathfrak{g}' \subset \mathfrak{g}$ corresponding to the root subsystem $R_{\mathfrak{g}'} \subset R_{\mathfrak{g}}$ generated by $\alpha_i, i > 1$, and let $\mathfrak{h}' \subset \mathfrak{g}'$ denote its Cartan subalgebra. Let the triangular decomposition $\mathfrak{g}'_- \oplus \mathfrak{h}' \oplus \mathfrak{g}'_+$ be compatible with the triangular decomposition of \mathfrak{g} . Recall the definition of step algebra $Z_q(\mathfrak{g}, \mathfrak{g}')$ of the pair $(\mathfrak{g}, \mathfrak{g}')$. Consider the left ideal $J = U_q(\mathfrak{g})\mathfrak{g}'_+$ and its normalizer $\mathcal{N} = \{x \in U_q(\mathfrak{g}) : e_\alpha x \subset J, \forall \alpha \in \Pi_{\mathfrak{g}'}^+\}$. By construction, J is a two-sided ideal in the algebra \mathcal{N} . Then $Z_q(\mathfrak{g}, \mathfrak{g}')$ is the quotient \mathcal{N}/J .

For all $\beta_i \in R_{\mathfrak{g}}^+ \setminus R_{\mathfrak{g}'}^+$ let e_{β_i} be the corresponding PBW generators and let Z be the vector space spanned by $e_{-\beta_l}^{k_l} \dots e_{-\beta_1}^{k_1} e_0^{k_0} e_{\beta_1}^{m_1} \dots e_{\beta_l}^{m_l}$, where $e_0 = q^{h_{\alpha_1}}$, $k_i \in \mathbb{Z}_+$, and $k_0 \in \mathbb{Z}$. The PBW factorization $U_q(\mathfrak{g}) = U_q(\mathfrak{g}'_-) Z U_q(\mathfrak{h}') U_q(\mathfrak{g}'_+)$ gives rise to the decomposition

$$U_q(\mathfrak{g}) = Z U_q(\mathfrak{h}') \oplus (\mathfrak{g}'_- U_q(\mathfrak{g}) + U_q(\mathfrak{g}) \mathfrak{g}'_+).$$

Proposition 3.1 ([5], Theorem 1). *The projection $U_q(\mathfrak{g}) \rightarrow Z U_q(\mathfrak{h}')$ implements an embedding of $Z_q(\mathfrak{g}, \mathfrak{g}')$ in $Z U_q(\mathfrak{h}')$.*

Proof. The statement is proved in [5] for the orthogonal and special linear quantum groups but the arguments apply to symplectic groups too. \square

It is proved within the theory of extremal projectors that generators of $Z_q(\mathfrak{g}, \mathfrak{g}')$ are labeled by the roots $\beta \in R_{\mathfrak{g}} \setminus R_{\mathfrak{g}'}$ plus $z_0 = q^{h_{\alpha_1}}$. We calculate them in the subsequent sections, cf. Propositions 3.5 and 3.9.

3.1 Lowering operators

In what follows, we extend $U_q(\mathfrak{g})$ along with its subalgebras containing $U_q(\mathfrak{h})$ over the field of fractions of $U_q(\mathfrak{h})$ and denote such an extension by hat, e.g. $\hat{U}_q(\mathfrak{g})$. In this section we calculate representatives of the negative generators of $Z_q(\mathfrak{g}, \mathfrak{g}')$ in $\hat{U}_q(\mathfrak{b}_-)$.

Set $h_i = h_{\varepsilon_i} \in \mathfrak{h}$ for all $i = 1, \dots, N$ and introduce $\eta_{ij} \in \mathfrak{h} + \mathbb{C}$ for $i, j = 1, \dots, N$, by

$$\eta_{ij} = h_i - h_j + (\varepsilon_i - \varepsilon_j, \rho) - \frac{1}{2} \|\varepsilon_i - \varepsilon_j\|^2. \quad (3.4)$$

Here $\|\mu\|$ is the Euclidean norm on \mathfrak{h}^* .

Lemma 3.2. *Suppose that $(l, r) \in P(\alpha)$ for some $\alpha \in \Pi^+$. Then*

- i) if $l < r < j$, then $\eta_{lj} - \eta_{rj} = h_{\alpha} + (\alpha, \varepsilon_j - \varepsilon_r)$,
- ii) if $i < l < r$, then $\eta_{li} - \eta_{ri} = h_{\alpha} + (\alpha, \varepsilon_i - \varepsilon_r)$,
- iii) $\eta_{lr} = h_{\alpha}$.

Proof. We have $(\alpha, \rho) = \frac{1}{2} \|\alpha\|^2$ for all $\alpha \in \Pi^+$. This proves iii). Further, for $\varepsilon_l - \varepsilon_r = \alpha$:

$$\begin{aligned} \eta_{lj} - \eta_{rj} &= h_{\alpha} + \frac{1}{2} \|\alpha\|^2 + \frac{1}{2} \|\varepsilon_j - \varepsilon_r\|^2 - \frac{1}{2} \|\varepsilon_j - \varepsilon_r - \alpha\|^2 = h_{\alpha} + (\alpha, \varepsilon_j - \varepsilon_r), \quad r < j, \\ \eta_{li} - \eta_{ri} &= h_{\alpha} + \frac{1}{2} \|\alpha\|^2 + \frac{1}{2} \|\varepsilon_i - \varepsilon_r\|^2 - \frac{1}{2} \|\varepsilon_i - \varepsilon_r - \alpha\|^2 = h_{\alpha} + (\alpha, \varepsilon_i - \varepsilon_r), \quad i < l, \end{aligned}$$

which proves i) and ii). \square

We call a strictly ascending sequence $\vec{m} = (m_1, \dots, m_s)$ of integers a route from m_1 to m_s . We write $m < \vec{m}$ and $\vec{m} < m$ for $m \in \mathbb{Z}$ if, respectively, $m < \min \vec{m}$ and $\max \vec{m} < m$. More generally, we write $\vec{m} < \vec{k}$ if $\max \vec{m} < \min \vec{k}$. In this case, a sequence (\vec{m}, \vec{k}) is a route from $\min \vec{m}$ to $\max \vec{k}$.

Given a route $\vec{m} = (m_1, \dots, m_s)$, define the product $f_{\vec{m}} = f_{m_1, m_2} \cdots f_{m_{s-1}, m_s} \in U_q(\mathfrak{g}_-)$. Consider a free right $\hat{U}_q(\mathfrak{h})$ -module $\Phi_{1\vec{m}}$ generated by $f_{\vec{m}}$ with $1 \leq \vec{m} \leq j$ and define an operation $\partial_{lr}: \Phi_{1j} \rightarrow \hat{U}_q(\mathfrak{b}_-)$ for $(l, r) \in P(\alpha)$ as follows. Assuming $1 \leq \vec{\ell} < l < r < \vec{\rho} < j$, set

$$\begin{aligned} \partial_{lr} f_{(\vec{\ell}, l)} f_{(l, r)} f_{(r, \vec{\rho})} &= f_{(\vec{\ell}, l)} f_{(r, \vec{\rho})} [\eta_{lj} - \eta_{rj}]_q, \\ \partial_{lr} f_{(\vec{\ell}, l)} f_{(l, \vec{\rho})} &= -f_{(\vec{\ell}, l)} f_{(r, \vec{\rho})} q^{-\eta_{lj} + \eta_{rj}}, \\ \partial_{lr} f_{(\vec{\ell}, r)} f_{(r, \vec{\rho})} &= f_{(\vec{\ell}, l)} f_{(r, \vec{\rho})} q^{\eta_{lj} - \eta_{rj}}, \\ \partial_{lr} f_{\vec{m}} &= 0, \quad l \notin \vec{m}, \quad r \notin \vec{m}. \end{aligned}$$

Extend ∂_{lr} to entire Φ_{1j} by $\hat{U}_q(\mathfrak{h})$ -linearity. Let $p: \Phi_{1j} \rightarrow \hat{U}(\mathfrak{g})$ denote the natural homomorphism of $\hat{U}_q(\mathfrak{h})$ -modules.

Lemma 3.3. *For all $\alpha \in \Pi^+$ and all $x \in \Phi_{1j}$, $e_\alpha \circ p(x) = \sum_{(l, r) \in P(\alpha)} \partial_{lr} x \pmod{\hat{U}_q(\mathfrak{g})e_\alpha}$.*

Proof. A straightforward analysis based on Proposition 2.3 and Lemma 3.2. \square

To simplify the presentation, we suppress the symbol of projection p in what follows.

Introduce elements $A_r^j \in \hat{U}_q(\mathfrak{h})$ by

$$A_r^j = \frac{q - q^{-1}}{q^{-2\eta_{rj}} - 1}, \quad (3.5)$$

for all $r, j \in [1, N]$ subject to $r < j$. For each simple pair (l, r) we define (l, r) -chains as

$$f_{(\vec{\ell}, l)} f_{(l, \vec{\rho})} A_l^j + f_{(\vec{\ell}, l)} f_{(l, r)} f_{(r, \vec{\rho})} A_l^j A_r^j + f_{(\vec{\ell}, r)} f_{(r, \vec{\rho})} A_r^j, \quad f_{(\vec{\ell}, l)} f_{l, j} A_l^j + f_{(\vec{\ell}, j)}, \quad (3.6)$$

where $1 \leq \vec{\ell} < l$ and $r < \vec{\rho} \leq j$. Remark that $f_{(l, r)} = \left[\frac{(\alpha, \alpha)}{2} \right]_q e_{-\alpha}$, where $\alpha = \varepsilon_l - \varepsilon_r$.

Lemma 3.4. *The operator ∂_{lr} annihilates (l, r) -chains.*

Proof. Applying ∂_{lr} to the 3-chain in (3.6), we get

$$f_{(\vec{\ell}, l)} f_{(r, \vec{\rho})} (-q^{-\eta_{lj} + \eta_{rj}} A_l^j + [\eta_{lj} - \eta_{rj}]_q A_l^j A_r^j + q^{\eta_{lj} - \eta_{rj}} A_r^j).$$

The factor in the brackets turns zero on substitution of 3.5.

Now apply ∂_{lj} to the right expression in (3.6) and get

$$f_{(\vec{\ell}, l)} ([h_\alpha]_q A_l^j + q^{h_\alpha}) = f_{(\vec{\ell}, l)} \left(\frac{q^{h_\alpha} - q^{-h_\alpha}}{q^{-2\eta_{lj}} - 1} + q^{h_\alpha} \right) = f_{(\vec{\ell}, l)} \frac{[h_\alpha - \eta_{lj}]_q}{[-\eta_{lj}]_q} = 0,$$

so long as $\eta_{lj} = h_\alpha$ by Lemma 3.2. \square

Given a route $\vec{m} = (m_1, \dots, m_s)$, put $A_{\vec{m}}^j = A_{m_1}^j \cdots A_{m_s}^j \in \hat{U}_q(\mathfrak{h})$ (and $A_{\vec{m}}^j = 1$ for the empty route) and define

$$z_{-j+1} = \sum_{1 < \vec{m} < j} f_{(1, \vec{m}, j)} A_{\vec{m}}^j \in \hat{U}_q(\mathfrak{b}_-), \quad j = 2, \dots, N, \quad (3.7)$$

where the summation is taken over all possible \vec{m} subject to the specified inequalities plus the empty route.

Proposition 3.5. $e_\alpha z_{-j} = 0 \pmod{\hat{U}_q(\mathfrak{g})e_\alpha}$ for all $\alpha \in \Pi_{\mathfrak{g}'}^+$ and $j = 1, \dots, N-1$.

Proof. Thanks to Lemma 3.3, we can reduce consideration to the action of operators ∂_{lr} , with $(l, r) \in P(\alpha)$. According to the definition of ∂_{lr} the summands in (3.7) that survive the action of ∂_{lr} can be organized into a linear combination of (l, r) -chains with coefficients in $\hat{U}_q(\mathfrak{h})$. By Lemma 3.4 they are killed by ∂_{lr} . \square

The elements z_{-i} , $i = 1, \dots, N-1$, belong to the normalizer \mathcal{N} and form the set of negative generators of $Z_q(\mathfrak{g}, \mathfrak{g}')$ for symplectic \mathfrak{g} . In the orthogonal case, the negative part of $Z_q(\mathfrak{g}, \mathfrak{g}')$ is generated by z_{-i} , $i = 1, \dots, N-2$.

3.2 Raising operators

In this section we construct positive generators of $Z_q(\mathfrak{g}, \mathfrak{g}')$, which are called raising operators. Consider an algebra automorphism $\omega: U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g})$ defined on the generators by $f_\alpha \leftrightarrow e_\alpha$, $q^{\pm h_\alpha} \mapsto q^{\mp h_\alpha}$. For $i < j$, let g_{ji} be the image of f_{ij} under this isomorphism. The natural representation restricted to $U_q(\mathfrak{g}_\pm)$ intertwines ω and matrix transposition. Since $(\omega \otimes \omega)(\check{\mathcal{R}}) = \check{\mathcal{R}}_{21}$, the matrix $\check{R}^+ = (\pi \otimes \text{id})(\check{\mathcal{R}}_{21})$ is equal to $1 \otimes 1 + (q - q^{-1}) \sum_{i < j} e_{ji} \otimes g_{ji}$.

Lemma 3.6. For all $\alpha \in \Pi_{\mathfrak{g}'}^+$ and all $i > 1$, $e_\alpha g_{i1} = \sum_{(l,r) \in P(\alpha)} \delta_{il} g_{r1} \pmod{\hat{U}_q(\mathfrak{g})e_\alpha}$.

Proof. Follows from the intertwining property of the R-matrix. \square

Consider the right $\hat{U}_q(\mathfrak{h})$ -module Ψ_{i1} freely generated by $f_{(\vec{m}, k)} g_{k1}$ with $i \leq \vec{m} < k$. We define operators $\partial_{lr}: \Psi_{i1} \rightarrow \hat{U}_q(\mathfrak{g})$ similarly as we did it for Φ_{1j} . For a simple pair $(l, r) \in P(\alpha)$, put

$$\partial_{l,r} f_{(\vec{m}, k)} g_{k1} = \begin{cases} f_{(\vec{m}, l)} g_{r1}, & l = k, \\ (\partial_{l,r} f_{(\vec{m}, k)}) g_{k1}, & l \neq k, \end{cases} \quad i \leq \vec{m} < r.$$

The Cartan factors appearing in $\partial_{lr} f_{(\vec{m}, k)}$ depend on h_α . When pushed to the right-most position, h_α is shifted by $(\alpha, \varepsilon_1 - \varepsilon_r)$. We extend ∂_{lr} to an action on Ψ_{i1} by the requirement

that ∂_{lr} commutes with the right action of $\hat{U}_q(\mathfrak{h})$. Let p denote the natural homomorphism of $\hat{U}_q(\mathfrak{h})$ -modules, $p: \Psi_{i1} \rightarrow \hat{U}_q(\mathfrak{g})$. One can prove the following analog of Lemma 3.3.

Lemma 3.7. *For all $\alpha \in \Pi_{\mathfrak{g}'}^+$ and all $x \in \Psi_{i1}$, $e_\alpha \circ p(x) = \sum_{(l,r) \in P(\alpha)} \partial_{lr} x \pmod{\hat{U}_q(\mathfrak{g})e_\alpha}$.*

Proof. Straightforward. \square

We suppress the symbol of projection p to simplify the formulas.

Define σ_i for all $i = 1, \dots, N$ as follows. For $i < j$ let $|i - j|$ be the number of simple positive roots entering $\varepsilon_i - \varepsilon_j$. For all $i, k = 2, \dots, N$, $i < k$, put

$$A_k^i = \frac{q^{\eta_{k1} - \eta_{i1}}}{[\eta_{i1} - \eta_{k1}]_q}, \quad B_k^i = \frac{(-1)^{|i-k|}}{[\eta_{i1} - \eta_{k1}]_q},$$

For each $(l, r) \in P(\alpha)$, where $\alpha \in \Pi_{\mathfrak{g}'}^+$, define 3-chains as

$$f_{(i, \vec{m}, l)} g_{l1} B_l^i + f_{(i, \vec{m}, l)} f_{(l, r)} g_{r1} A_l^i B_r^i + f_{(i, \vec{m}, r)} g_{r1} B_r^i, \quad (3.8)$$

with $i < \vec{m} < l < r \leq N$ and

$$f_{(i, \vec{\ell}, l)} f_{(l, \vec{\rho}, k)} g_{k1} A_l^i + f_{(i, \vec{\ell}, l)} f_{(l, r)} f_{(r, \vec{\rho}, k)} g_{k1} A_l^i A_r^i + f_{(i, \vec{\ell}, r)} f_{(r, \vec{\rho}, k)} g_{k1} A_r^i \quad (3.9)$$

with $i < \vec{\ell} < l < r < \vec{\rho} < k \leq N$. The 2-chains are defined as

$$g_{i1} + f_{(i, r)} g_{r1} B_r^i, \quad f_{(i, \vec{m}, k)} g_{k1} + f_{(i, r)} f_{(r, \vec{m}, k)} g_{k1} A_r^i \quad (3.10)$$

where r is such that $\varepsilon_i - \varepsilon_r \in \Pi_{\mathfrak{g}'}^+$ and $i < r < \vec{m} < k \leq N$. In all cases the empty routes \vec{m} are admissible.

Lemma 3.8. *For all $\alpha \in \Pi_{\mathfrak{g}'}^+$ and all $(l, r) \in P(\alpha)$ the (l, r) -chains are annihilated by ∂_{lr} .*

Proof. Suppose that $i = l$ and apply ∂_{ir} to the left 2-chain in (3.10). The result is

$$g_{r1} + [h_\alpha]_q g_{r1} B_r^i = g_{r1} (1 + [h_\alpha + (\alpha, \varepsilon_1 - \varepsilon_r)]_q B_r^i) = g_{r1} (1 + [\eta_{i1} - \eta_{r1}]_q B_r^i) = 0,$$

by Lemma 3.2. Applying ∂_{ir} to the right 2-chain in (3.10) we get

$$f_{(r, \vec{m}, k)} g_{k1} (-q^{-\eta_{i1} + \eta_{r1}} + [\eta_{i1} - \eta_{r1}]_q A_r^i) = 0.$$

Now consider 3-chains. The action of ∂_{lr} on the (3.9) produces

$$-f_{(i, \vec{\ell}, l)} q^{-h_\alpha} f_{(r, \vec{\rho}, k)} g_{k,1} A_l^i + f_{(i, \vec{\ell}, l)} [h_\alpha]_q f_{(r, \vec{\rho}, k)} g_{k,1} A_l^i A_r^i + f_{(i, \vec{\ell}, l)} q^{h_\alpha} f_{(r, \vec{\rho}, k)} g_{k,1} A_r^i,$$

which turns zero since $-q^{\eta_{r1} - \eta_{i1}} A_l^i + [\eta_{l1} - \eta_{r1}]_q A_l^i A_r^i + q^{\eta_{i1} - \eta_{r1}} A_r^i = 0$. The action of ∂_{lr} on (3.8) yields

$$f_{(i, \vec{m}, l)} g_{r1} B_l^i + f_{(i, \vec{m}, l)} [h_\alpha]_q g_{r1} A_l^i B_r^i + f_{(i, \vec{m}, l)} q^{h_\alpha} g_{r1} B_r^i.$$

This is vanishing since $B_l^i + [\eta_{l1} - \eta_{r1}]_q A_l^i B_r^i + q^{\eta_{i1} - \eta_{r1}} B_r^i = B_l^i + \frac{[\eta_{i1} - \eta_{r1}]_q}{[\eta_{i1} - \eta_{l1}]_q} B_r^i = 0$. \square

Given a route $\vec{m} = (m_1, \dots, m_k)$ such that $i < \vec{m}$ let $A_{\vec{m}}^i$ denote the product $A_{m_1}^i \dots A_{m_k}^i$. Introduce elements $z_i \in \hat{U}_q(\mathfrak{g}_-)\mathfrak{g}_+$ of weight $\varepsilon_1 - \varepsilon_i$ by

$$z_{i-1} = g_{i1} + \sum_{i < \vec{m} < k \leq N} f_{(i, \vec{m}, k)} g_{k1} A_{\vec{m}}^i B_k^i, \quad i = 2, \dots, N.$$

Again, the summation includes empty \vec{m} .

Proposition 3.9. $e_\alpha z_i = 0 \pmod{\hat{U}_q(\mathfrak{g})e_\alpha}$, for all $\alpha \in \Pi_{\mathfrak{g}'}^+$ and $i = 1, \dots, N - 1$.

Proof. By Lemma 3.6, the vectors $g_{2'1}$ and $z_{N-1} = g_{1'1}$ are normalizing the left ideal $\hat{U}_q(\mathfrak{g})\mathfrak{g}'_+$, so is $z_{N-2} = g_{2'1} + f_{1'} g_{1'1} B_{2'}^{1'}$. Once the cases $i = 2', 1'$ are proved, we further assume $i < 2'$. In view of Lemma 3.7, it is sufficient to show that z_{i-1} is killed, modulo $\hat{U}_q(\mathfrak{g})\mathfrak{g}'_+$, by all ∂_{lr} such that $\varepsilon_l - \varepsilon_r \in \Pi_{\mathfrak{g}'}^+$. Observe that z_{i-1} can be arranged into a linear combination of chains, which are killed by ∂_{lr} , as in Lemma 3.8. \square

The elements z_i , $i = 1, \dots, N - 1$, belong to the normalizer \mathcal{N} . They form the set of positive generators of $Z_q(\mathfrak{g}, \mathfrak{g}')$ for symplectic \mathfrak{g} . In the orthogonal case, the positive part of $Z_q(\mathfrak{g}, \mathfrak{g}')$ is generated by z_i , $i = 1, \dots, N - 2$.

Acknowledgements. This research is supported in part by the RFBR grant 12-01-00207-a. We are grateful to the Max-Planck Institute for Mathematics in Bonn for hospitality and excellent research atmosphere.

References

- [1] Mickelsson, J.: Step algebras of semisimple Lie algebras, Rev. Mod. Phys. **4** (1973), 307–18.
- [2] Zhelobenko, D. P.: Extremal projectors and generalized Mickelsson algebras over reductive Lie algebras Izv. Akad. Nauk. SSSR Ser. Mat. **52** (1988), 758–73 (in Russian) (Engl. transl. Math. USSR Izv. **33** (1989))
- [3] Molev, A. I.: Gelfand-Tsetlin bases for classical Lie algebras, in Handbook of algebra. **4** 109–170, Elsevier/North-Holland, Amsterdam, 2006.
- [4] Zhelobenko D. P.: S-algebras and Verma modules over reductive Lie algebras Dokl. Akad. Nauk. SSSR **273** (1983), 785–788 (in Russian) (Engl. transl. Sov. Math. Dokl. **28** (1983), 696–700)

- [5] Kekäläinen, P.: Step algebras of quantum algebras of type A , B and D , J. Phys. A: Math. Gen. **29** (1996), 1045–1053.
- [6] Asherova, R. M., Smirnov, Yu. F. and Tolstoy, V. N.: Projection operators for simple Lie groups, Theor. Math. Phys. **8** (1971), 813–825.
- [7] Asherova, R. M., Smirnov, Yu. F. and Tolstoy, V. N.: Projection operators for simple Lie groups. II. General scheme for constructing lowering operators. The groups $SU(n)$, Theor. Math. Phys. **15** (1973), 392–401.
- [8] Tolstoy, V. N.: Projection operator method for quantum groups, NATO Sci. Ser. II Math. Phys. Chem., **30**, 457–488, Kluwer Acad. Publ., Dordrecht, 2001.
- [9] Nagel, J. G., Moshinsky, M.: Operators that lower or raise the irreducible vector spaces of U_{n-1} contained in an irreducible vector space of U_n , J. Math. Phys. **6** (1965), 682–694.
- [10] Hou Pei-yu: Orthonormal bases and infinitesimal operators of the irreducible representations of group U_n , Scientia Sinica **15** (1966), 763–772.
- [11] Pang, S. C., Hecht, K. T.: Lowering and raising operators for the orthogonal group in the chain $O(n) \supset O(n-1) \supset \dots$, and their graphs, J. Math. Phys. **8** (1967), 1233–1251.
- [12] Wong, M. K. F.: Representations of the orthogonal group. I. Lowering and raising operators of the orthogonal group and matrix elements of the generators, J. Math. Phys. **8** (1967), 1899–1911.
- [13] Asherova, R. M., Burdík, Č. Havlíček, M., Smirnov, Yu. F. and Tolstoy, V. N.: q-Analog of Gelfand-Graev Basis for the Noncompact Quantum Algebra $U_q(u(n, 1))$, SIGMA **6** (2010), 010, 13 pages.
- [14] Chari, V., Pressley, A.: A guide to quantum groups. Cambridge University Press, Cambridge, 1995.
- [15] Drinfeld, V.: Quantum Groups. In Proc. Int. Congress of Mathematicians, Berkeley 1986, Gleason, A. V. (eds), AMS, Providence (1987) 798–820.