# DENSE BANACH SUBALGEBRAS OF THE NULL SEQUENCE ALGEBRA WHICH DO NOT SATISFY A DIFFERENTIAL SEMINORM CONDITION

LARRY B. SCHWEITZER<sup>1</sup>

ABSTRACT. We construct dense Banach subalgebras A of the null sequence algebra  $c_0$  which are spectral invariant, but do not satisfy the  $D_1$ -condition  $\|ab\|_A \leq K(\|a\|_{\infty} \|b\|_A + \|a\|_A \|b\|_{\infty})$ , for all  $a, b \in A$ . The sequences in A vanish in a skewed manner with respect to an unbounded function  $\sigma \colon \mathbb{N} \to [1, \infty)$ .

### 1. INTRODUCTION

We say that A is a dense Banach subalgebra of a  $C^*$ -algebra B if A is a dense subalgebra of B, and A is a Banach algebra in some norm  $\|\cdot\|_A$ , which is stronger than the restriction to A of the norm on B. We say that A is spectral invariant in B if the quasi-invertible elements of A are precisely those elements of A which are quasi-invertible in B. Recall that for  $a, b \in A$ ,  $a \circ b = a + b - ab$ , and b is a quasi-inverse for a if and only if  $a \circ b = b \circ a = 0$  (see [4, Definition 2.1.1]). Dense subalgebras of a  $C^*$ -algebra which are spectral invariant, or which satisfy a differential seminorm condition such as  $D_1$ , are used to give differential structure to the  $C^*$ -algebra, and they also have applications in noncommutative differential geometry (see [2][1]).

Let  $c_0$  be the  $C^*$ -algebra of complex-valued vanishing sequences, or null sequences, on the natural numbers  $\mathbb{N} = \{0, 1, 2, ...\}$ , with pointwise multiplication and involution. For  $n \in \mathbb{N}$ , let  $e_n \colon \mathbb{N} \to \{0, 1\}$  be the unit step function at n, where  $e_n(k) = 1$  if k = n and  $e_n(k) = 0$  for  $k \in \mathbb{N} \setminus \{n\}$ . Let  $c_f$  denote the linear span of  $\{e_n\}_{n=0}^{\infty}$ , which is a dense ideal in  $c_0$ . The following result shows that it is relatively easy for a dense subalgebra of  $c_0$  to be spectral invariant.

**Theorem 1.1.** Let A be a dense Banach subalgebra of  $c_0$ . Assume that the subalgebra of finite support functions  $c_f$  is a dense subset of A. Then A is spectral invariant in  $c_0$ .

Proof. Let  $A_{qG}$  denote the group of quasi-invertible elements of A, with group operation  $\circ$  and identity 0. Let  $a \in A \setminus A_{qG}$ . Then  $A \circ a$  cannot intersect  $A_{qG}$ , or else a would have a quasi-inverse. So  $(a + A(1 - a)) \cap A_{qG} = \emptyset$ . Note that J = A(1 - a) is an ideal in A, and is proper since  $J \cap (A_{qG} - a) = \emptyset$ . Since Ais a Banach algebra,  $A_{qG}$  and its translate  $A_{qG} - a$  are open sets in A. Hence J

Date: Received: May 12, 2016; Accepted: July 20, 2016.

<sup>2010</sup> Mathematics Subject Classification. Primary 46L87; Secondary 46H10, 46J10, 46B45, 46K99.

Key words and phrases.  $D_1$ -subalgebra, spectral invariance, null sequence algebra, differential structure in  $C^*$ -algebras.

#### L.B. SCHWEITZER

cannot be dense in A, and  $c_f \not\subseteq J$ , using the hypothesis. Since J is a linear space, there is some  $n_0 \in \mathbb{N}$  for which  $e_{n_0} \notin J$ . Since J is a  $c_f$ -module not containing  $e_{n_0}$ , every element of J must vanish at  $n_0$ . This can only happen if  $a(n_0) = 1$ . Hence a is not quasi-invertible in  $c_0$ .

A dense Banach subalgebra A is a  $D_1$ -subalgebra of B if for some constant  $K_A > 0$ , the  $D_1$ -condition

$$||ab||_{A} \le K_{A} \{ ||a||_{A} ||b||_{B} + ||a||_{B} ||b||_{A} \}$$
(1.1)

is satisfied for all  $a, b \in A$ , where  $\|\cdot\|_A$  is the norm on A and  $\|\cdot\|_B$  is the norm on B (see [3]). Being a  $D_1$ -subalgebra implies spectral invariance (see [3, Theorem 5 and Lemma 4]), which raises the question: Is every dense Banach subalgebra of the null sequence algebra, which satisfies the hypotheses of Theorem 1.1, also  $D_1$ ? The purpose of the present paper is to provide a counterexample. To this end, we think of  $c_0$  as  $\mathbb{C}^2$ -valued sequences vanishing at infinity,  $c_0(\mathbb{N}, \mathbb{C}^2)$ , where  $\mathbb{C}^2$  denotes the 2-dimensional commutative  $C^*$ -algebra, with coordinatewise multiplication and involution. We identify the two  $C^*$ -algebras using the isomorphism  $\theta: c_0 \cong c_0(\mathbb{N}, \mathbb{C}^2)$ ,

$$\theta(f)(n) = \left(f(2n), f(2n+1)\right),\tag{1.2}$$

for  $f \in c_0$ ,  $n \in \mathbb{N}$ . If A is a dense subalgebra,  $D_1$  is satisfied along each  $\mathbb{C}^2$ summand, because it is finite dimensional. In Section 2, we construct submultiplicative norms  $\|\cdot\|_n$  on the *n*th copy of  $\mathbb{C}^2$ , which make the  $D_1$ -constants  $K_n$ become unbounded as *n* increases. In Section 3, we define the Banach algebra norm  $\|\cdot\|_A$  as the sup of these  $\mathbb{C}^2$ -norms to construct the counterexample.

## 2. Some norms on $\mathbb{C}^2$

Let  $r \in \mathbb{R}$  and let  $\sigma$  be a constant in  $[1, \infty)$ . Let  $\|\vec{v}\|_{\max} = \max\{|x|, |y|\}, \vec{v} = (x, y) \in \mathbb{C}^2$ , denote the  $C^*$ -norm on  $\mathbb{C}^2$ . Define a seminorm on  $\mathbb{C}^2$  by

$$\|\vec{v}\|_{r,\sigma} = \|T_{r,\sigma}\vec{v}\|_{\max} = \max\{|x+ry|,\sigma|y-rx|\},$$
(2.1)

where  $T_{r,\sigma}$  is the  $(2 \times 2)$ -matrix

$$\begin{pmatrix} 1 & r \\ -\sigma r & \sigma \end{pmatrix}.$$
 (2.2)

Note that  $\Delta = \det(T_{r,\sigma}) = \sigma(1+r^2) > 0$ , so  $T_{r,\sigma}$  is invertible, and  $\|\cdot\|_{r,\sigma}$  is a norm on  $\mathbb{C}^2$ .

We want to find the smallest constant D > 0 which satisfies  $\|\vec{v}\|_{\max} \leq D \|\vec{v}\|_{r,\sigma}$ for all  $\vec{v} \in \mathbb{C}^2$ . Then D is the norm of

$$T_{r,\sigma}^{-1} = \frac{1}{\Delta} \begin{pmatrix} \sigma & -r \\ \sigma r & 1 \end{pmatrix}$$

as an operator on  $\mathbb{C}^2$  with max-norm,

$$D = ||T_{r,\sigma}^{-1}||_{\text{op}} = \max\{||(\sigma, -r)||_{1}, ||(\sigma r, 1)||_{1}\}/\Delta$$
  

$$= \frac{1}{\sigma(1+r^{2})} \max\{\sigma + |r|, \sigma|r| + 1\}$$
  

$$= \frac{1}{1+r^{2}} \max\{1 + |r|/\sigma, |r| + 1/\sigma\}$$
  

$$\leq \frac{1}{1+r^{2}} \max\{1 + |r|, |r| + 1\} \quad \text{since } \sigma \geq 1$$
  

$$= \frac{1+|r|}{1+r^{2}},$$
(2.3)

since the dual of the max-norm is the  $\ell^1$ -norm. As r ranges over all real numbers, the last expression in (2.3) is bounded by 1.21.

Next we want to find a constant C > 0 satisfying  $\|\vec{v}_1 * \vec{v}_2\|_{r,\sigma} \leq C \|\vec{v}_1\|_{r,\sigma} \|\vec{v}_2\|_{r,\sigma}$ , for all  $\vec{v}_1, \vec{v}_2 \in \mathbb{C}^2$ , where  $\vec{v}_1 * \vec{v}_2$  denotes pointwise multiplication. This is equivalent to finding the norm of the operator  $\vec{u}_1 \otimes \vec{u}_2 \mapsto T_{r,\sigma}(T_{r,\sigma}^{-1}\vec{u}_1 * T_{r,\sigma}^{-1}\vec{u}_2)$  from  $\mathbb{C}^2 \otimes \mathbb{C}^2$  to  $\mathbb{C}^2$ . The operator is  $\frac{1}{\Delta^2}$  times

$$\begin{pmatrix} 1 & r \\ -\sigma r & \sigma \end{pmatrix} \left( \begin{pmatrix} \sigma & -r \\ \sigma r & 1 \end{pmatrix} \begin{pmatrix} u_{11} \\ u_{12} \end{pmatrix} * \begin{pmatrix} \sigma & -r \\ \sigma r & 1 \end{pmatrix} \begin{pmatrix} u_{21} \\ u_{22} \end{pmatrix} \right)$$

$$= \begin{pmatrix} 1 & r \\ -\sigma r & \sigma \end{pmatrix} \begin{pmatrix} (\sigma u_{11} - r u_{12})(\sigma u_{21} - r u_{22}) \\ (\sigma r u_{11} + u_{12})(\sigma r u_{21} + u_{22}) \end{pmatrix}$$

$$= \begin{pmatrix} 1 & r \\ -\sigma r & \sigma \end{pmatrix} \begin{pmatrix} \sigma^2 & -r\sigma & -r\sigma & r^2 \\ \sigma^2 r^2 & \sigma r & \sigma r & 1 \end{pmatrix} \begin{pmatrix} u_{11} u_{21} \\ u_{11} u_{22} \\ u_{12} u_{21} \\ u_{12} u_{22} \end{pmatrix}$$

$$= \begin{pmatrix} \sigma^2 (1 + r^3) & \sigma (r^2 - r) & \sigma (r^2 - r) & r^2 + r \\ \sigma^3 (r^2 - r) & \sigma^2 (r^2 + r) & \sigma^2 (r^2 + r) & \sigma (1 - r^3) \end{pmatrix} \begin{pmatrix} u_{11} u_{21} \\ u_{12} u_{22} \\ u_{12} u_{21} \\ u_{12} u_{22} \end{pmatrix}$$

So the constant C is

$$C = \max\{\|(\sigma^{2}(1+r^{3}), \sigma(r^{2}-r), \sigma(r^{2}-r), r^{2}+r)\|_{1}, \\\|(\sigma^{3}(r^{2}-r), \sigma^{2}(r^{2}+r), \sigma^{2}(r^{2}+r), \sigma(1-r^{3}))\|_{1}\}/\Delta^{2} \\ = \frac{\sigma \max\{\frac{|1+r^{3}|}{\sigma} + \frac{2|r^{2}-r|}{\sigma^{2}} + \frac{|r^{2}+r|}{\sigma^{3}}, |r^{2}-r| + \frac{2|r^{2}+r|}{\sigma} + \frac{|1-r^{3}|}{\sigma^{2}}\}}{(1+r^{2})^{2}} \\ \leq \frac{\sigma \max\{|1+r^{3}| + 2|r^{2}-r| + |r^{2}+r|, |r^{2}-r| + 2|r^{2}+r| + |1-r^{3}|\}}{(1+r^{2})^{2}},$$

$$(2.4)$$

where the first step used that the dual of the max-norm is the  $\ell^1$ -norm, and the last step used that  $\sigma \geq 1$ . As r ranges over all real numbers, the last expression in (2.4) is bounded by  $2\sigma$ .

### L.B. SCHWEITZER

It follows that  $2\sigma \|\vec{v}_1 * \vec{v}_2\|_{r,\sigma} \leq (2\sigma \|\vec{v}_1\|_{r,\sigma}) (2\sigma \|\vec{v}_2\|_{r,\sigma})$ , for  $\vec{v}_1, \vec{v}_2 \in \mathbb{C}^2$ . By the paragraph before,  $\|\vec{v}\|_{\max} \leq D \|\vec{v}\|_{r,\sigma} < 2 \|\vec{v}\|_{r,\sigma} = \frac{1}{\sigma} (2\sigma \|\vec{v}\|_{r,\sigma})$ , for  $\vec{v} \in \mathbb{C}^2$ .

# 3. The Banach algebras $A_{r,\sigma}$

In this section, we pass from the case of a single copy of  $\mathbb{C}^2$  (see Section 2) to infinitely many copies of  $\mathbb{C}^2$ . Let  $r \in \mathbb{R}$  and let  $\sigma$  be any unbounded function from  $\mathbb{N}$  to  $[1, \infty)$ . Define an infinite matrix  $S_{r,\sigma}$  as the direct sum of  $(2 \times 2)$ -matrices

$$S_{r,\sigma} = \bigoplus_{n=0}^{\infty} 2\sigma(n) T_{r,\sigma(n)},$$

where each  $T_{r,\sigma(n)}$  is defined as in (2.2). Let  $A_{r,\sigma} = \{ f \in c_0 \mid S_{r,\sigma}\theta f \in c_0(\mathbb{N}, \mathbb{C}^2) \}$ , where  $\theta$  was defined in the Introduction (see (1.2)). By [5, Theorem 4.3.1],  $A_{r,\sigma}$ is complete in the norm

$$||f||_{r,\sigma} = ||S_{r,\sigma}\theta f||_{c_0(\mathbb{N},\mathbb{C}^2)} = \sup_{n=0}^{\infty} \{2\sigma(n) || (f(2n), f(2n+1)) ||_{r,\sigma(n)} \}, \quad (3.1)$$

where each  $\|\cdot\|_{r,\sigma(n)}$  is defined as in (2.1), and where by the final remarks of Section 2,  $\|fg\|_{r,\sigma} \leq \|f\|_{r,\sigma} \|g\|_{r,\sigma}$  and  $\|f\|_{\infty} \leq \|f\|_{r,\sigma}$ , for  $f, g \in A_{r,\sigma}$ . Since  $S_{r,\sigma}\theta f \in c_0(\mathbb{N}, \mathbb{C}^2)$  for  $f \in A_{r,\sigma}$ , then

$$0 = \lim_{n \to \infty} \left\| \left( S_{r,\sigma} \theta f \right)(n) \right\|_{\mathbb{C}^2} = \lim_{n \to \infty} 2\sigma(n) \left\| \left( f(2n), f(2n+1) \right) \right\|_{r,\sigma(n)}$$

For  $\epsilon > 0$ , let  $N_{\epsilon}$  be large enough so that the argument of the limit is smaller than  $\epsilon$  if  $n > N_{\epsilon}$ . Then for  $n \ge N_{\epsilon}$ ,

$$\begin{split} \left\| f - \left( f(0), \dots, f(2n+1), 0, 0, \dots \right) \right\|_{r,\sigma} &= \left\| \left( \underbrace{0, \dots, 0}_{2n+2 \text{ zeros}}, f(2n+2), \dots \right) \right\|_{r,\sigma} \\ &= \sup_{k>n} 2\sigma(k) \left\| \left( f(2k), f(2k+1) \right) \right\|_{r,\sigma(k)} \\ &< \epsilon, \end{split}$$

using the definition of  $\|\cdot\|_{r,\sigma}$  (3.1). It follows that  $c_f$  is dense in  $A_{r,\sigma}$ , and we can apply Theorem 1.1 to see that  $A_{r,\sigma}$  is spectral invariant in  $c_0$ .

**Theorem 3.1.** The dense Banach subalgebra  $A_{r,\sigma}$  of  $c_0$  is not a  $D_1$ -subalgebra of  $c_0$  for  $r \in \mathbb{R} \setminus \{0, 1\}$ .

*Proof.* For  $n \in \mathbb{N}$ , define  $a_n \in A_{r,\sigma}$  by

$$a_n = \left(\underbrace{0, 0, \dots, 0, 0}_{2n \text{ zeros}}, 1, r, 0, 0, \dots\right),$$

where  $a_n(2n) = 1$  and  $a_n(2n+1) = r$  are the only nonzero components. Then

$$\begin{aligned} \|a_n\|_{r,\sigma} &= 2\sigma(n)(1+r^2), \\ \|a_n^2\|_{r,\sigma} &= 2\sigma(n)\max(|1+r^3|,\sigma(n)|r^2-r|), \\ \|a_n\|_{\infty} &= \max(1,|r|). \end{aligned}$$

If K > 0 were a constant satisfying the  $D_1$ -condition (1.1) for  $A_{r,\sigma}$  in  $c_0$ , then

$$K \geq \frac{\|a_n^2\|_{r,\sigma}}{2\|a_n\|_{\infty}\|a_n\|_{r,\sigma}} = \frac{\max(|1+r^3|,\sigma(n)|r^2-r|)}{2(1+r^2)\max(1,|r|)}$$
  
$$\geq \frac{\sigma(n)|r^2-r|}{2(1+r^2)\max(1,|r|)}$$

must hold for each n for which  $\sigma(n)|r^2 - r| \ge |1 + r^3|$ . No such constant K can exist if  $r \ne 1$  or 0, since  $\sigma$  is unbounded.

Remark 3.2. Note that  $A_{r,\sigma}$  is a Banach  $\star$ -algebra. The norms defined on  $\mathbb{C}^2$  (2.1) and the norm on  $A_{r,\sigma}$  (3.1) are both left unchanged by the  $\star$ -operation of pointwise complex-conjugation.

Remark 3.3. In the cases r = 0 and r = 1, it can be shown that  $A_{r,\sigma}$  is a  $D_1$ -subalgebra of  $c_0$ . Further,  $A_{r,\sigma}$  is a dense Banach ideal in  $c_0$  if r = 0, but not if r = 1.

## References

- S. J. Bhatt, Smooth Fréchet subalgebras of C<sup>\*</sup>-algebras defined by first order differential seminorms, Proc. Indian Acad. Sci. Math. Sci. 126 (2016), no. 1, 125-141.
- 2. A. Connes, Noncommutative Geometry, Academic Press, San Diego, 1994.
- E. Kissin and V. S. Shulman, Differential properties of some dense subalgebras of C<sup>\*</sup>algebras, Proc. Edinb. Math. Soc. (2) 37 (1994), no. 3, 399-422.
- T. W. Palmer, Banach Algebras and the General Theory of \*-algebras, Volume I: Algebras and Banach Algebras, Encyclopedia Math. Appl. 49, Cambridge Univ. Press, Cambridge, 1994.
- A. Wilansky, Summability through Functional Analysis, North-Holland Math. Stud. 85, North-Holland, Amsterdam, 1984.

<sup>1</sup>DEPARTMENT OF STATISTICS AND BIOSTATISTICS, CALIFORNIA STATE UNIVERSITY EAST BAY, 25800 CARLOS BEE BOULEVARD, HAYWARD, CA 94542.

*E-mail address*: lschweitzer@horizon.csueastbay.edu; lsch@svpal.org.