

# On $\mathcal{H}C$ -subgroups of a finite group<sup>\*†</sup>

Lijun Huo, Xiaoyu Chen<sup>‡</sup>, Wenbin Guo

Department of Mathematics, University of Science and Technology of China,

Hefei, Anhui 230026, P. R. China

e-mails: ljhuo@mail.ustc.edu.cn, jelly@mail.ustc.edu.cn, wbguo@ustc.edu.cn

## Abstract

A subgroup  $H$  of a finite group  $G$  is said to be an  $\mathcal{H}C$ -subgroup of  $G$  if there exists a normal subgroup  $T$  of  $G$  such that  $G = HT$  and  $H^g \cap N_T(H) \leq H$  for all  $g \in G$ . In this paper, we investigate the structure of a finite group  $G$  under the assumption that certain subgroups of  $G$  of arbitrary prime power order are  $\mathcal{H}C$ -subgroups of  $G$ .

**Key words:**  $\mathcal{H}$ -subgroups;  $\mathcal{H}C$ -subgroups;  $p$ -nilpotent group; nilpotent group; supersolvable group.

## 1 Introduction

Throughout this paper, all groups considered are finite.  $G$  always denotes a group,  $p$  denotes a prime, and  $|G|_p$  denotes the order of Sylow  $p$ -subgroups of  $G$ . A class of groups  $\mathfrak{F}$  is called a formation if  $\mathfrak{F}$  is closed under taking homomorphic images and subdirect products. A formation  $\mathfrak{F}$  is said to be saturated if  $G \in \mathfrak{F}$  whenever  $G/\Phi(G) \in \mathfrak{F}$ . All unexplained notation and terminology are standard, as in [7, 10, 12].

Recall that a subgroup  $H$  of  $G$  is said to be an  $\mathcal{H}$ -subgroup of  $G$  if  $H^g \cap N_G(H) \leq H$  for all  $g \in G$ . This concept was introduced by Goldschmidt in [9] and Bianchi et al. in [5]. It is easy to see that normal subgroups, Sylow subgroups and self-normalizing subgroups of  $G$  are all  $\mathcal{H}$ -subgroups of  $G$ . Csörgö, Herzog [6] and Asaad [1] further investigated the influence of  $\mathcal{H}$ -subgroups on the structure of a finite group.

---

<sup>\*</sup>This manuscript was finished in August 2013.

<sup>†</sup>Research is supported by a NNSF grant of China (Grant 11371335) and Research Fund for the Doctoral Program of Higher Education of China(Grant 20113402110036).

<sup>‡</sup>Corresponding author.

2000 AMS Mathematics Subject Classification: 20D10, 20D20.

Besides, Y. Wang [20] introduced the concept of  $c$ -normal subgroups. A subgroup  $H$  of  $G$  is said to be  $c$ -normal in  $G$  if there exists a normal subgroup  $K$  of  $G$  such that  $G = HK$  and  $H \cap K \leq H_G$ , where  $H_G$  is the largest normal subgroup of  $G$  contained in  $H$ . The properties of  $c$ -normal subgroups have been studied by many authors, see for example, [3, 4, 14, 15].

Recently, some attempts were made to give a generalization of both  $c$ -normal subgroups and  $\mathcal{H}$ -subgroups. In [2], M. Asaad et al. introduced the concept of weakly  $\mathcal{H}$ -subgroups: a subgroup  $H$  of  $G$  is called an  $\mathcal{HC}$ -subgroup of  $G$  if there exists a normal subgroup  $T$  of  $G$  such that  $G = HT$  and  $H \cap T$  is an  $\mathcal{H}$ -subgroup of  $G$ . Meanwhile, X. Wei and X. Guo [21] introduced the concept of  $\mathcal{HC}$ -subgroups: a subgroup  $H$  of  $G$  is said to be an  $\mathcal{HC}$ -subgroup of  $G$  if there exists a normal subgroup  $T$  of  $G$  such that  $G = HT$  and  $H^g \cap N_T(H) \leq H$  for all  $g \in G$ . It is easy to see that every weakly  $\mathcal{H}$ -subgroup of  $G$  is an  $\mathcal{HC}$ -subgroup of  $G$ .

In [21], the authors gave some conditions on maximal subgroups or minimal subgroups of Sylow subgroups, which are sufficient to guarantee a group to be  $p$ -nilpotent or supersolvable. In this paper, we continue to investigate the structure of a group  $G$  under the assumption that certain subgroups of  $G$  of arbitrary prime power order are  $\mathcal{HC}$ -subgroups of  $G$ . New characterizations of some classes of finite groups are obtained.

## 2 Preliminaries

**Lemma 2.1.** *Suppose that  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ .*

- (1) ([5, Theorem 6(2)]) *If  $H$  is subnormal in  $G$ , then  $H$  is normal in  $G$ .*
- (2) ([5, Lemma 7(2)]) *If  $H \leq K \leq G$ , then  $H$  is an  $\mathcal{H}$ -subgroup of  $K$ .*
- (3) ([5, Lemma 2(1)]) *If  $N \leq H$  and  $N \trianglelefteq G$ , then  $H$  is an  $\mathcal{H}$ -subgroup of  $G$  if and only if  $H/N$  is an  $\mathcal{H}$ -subgroup of  $G/N$ .*
- (4) ([5, Theorem 6(3)]) *If  $N \trianglelefteq G$  and  $N \leq N_G(H)$ , then  $N_G(HN) = N_G(H)$  and  $HN$  is an  $\mathcal{H}$ -subgroup of  $G$ .*

**Lemma 2.2.** *Let  $H$  and  $K$  be subgroups of  $G$ , and  $N \trianglelefteq G$ .*

- (1) ([21, Lemma 2.3(1)]) *If  $H \leq K$  and  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$ , then  $H$  is an  $\mathcal{HC}$ -subgroup of  $K$ .*
- (2) ([21, Lemma 2.3(2)]) *If  $N \leq H$ , then  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$  if and only if  $H/N$  is an  $\mathcal{HC}$ -subgroup of  $G/N$ .*
- (3) ([21, Lemma 2.4]) *If  $H$  is a  $p$ -group with  $(p, |N|) = 1$  and  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$ , then  $HN$  is an  $\mathcal{HC}$ -subgroup of  $G$  and  $HN/N$  is an  $\mathcal{HC}$ -subgroup of  $G/N$ .*

**Lemma 2.3.** [21, Theorem 3.7] *Let  $p$  be the smallest prime dividing  $|G|$  and let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Then  $G$  is  $p$ -nilpotent if every maximal subgroup of  $P$  is an  $\mathcal{HC}$ -subgroup of  $G$ .*

**Lemma 2.4.** [9, Corollary B3] *Suppose that  $S$  is a 2-subgroup of  $G$  such that  $S$  is an  $\mathcal{H}$ -subgroup of  $G$  and  $N_G(S)/C_G(S)$  is a 2-group. Then  $S$  is a Sylow 2-subgroup of  $S^G$ .*

**Lemma 2.5.** [16, Theorem 1] *Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Then the following two statements are true:*

- (1) *If  $p$  is odd and every minimal subgroup of  $P$  lies in  $Z(N_G(P))$ , then  $G$  is  $p$ -nilpotent.*
- (2) *If  $p = 2$  and every cyclic subgroup of  $P$  of order 2 or 4 is quasinormal in  $N_G(P)$ , then  $G$  is 2-nilpotent.*

Let  $F^*(G)$  denote the generalized Fitting subgroup of  $G$ , that is, the largest normal quasinilpotent subgroup of  $G$ . The following basic facts can be found in [13, Chapter X].

- Lemma 2.6.** (1) *If  $N$  is a normal subgroup of  $G$ , then  $F^*(N) = N \cap F^*(G)$ .*  
(2)  *$F(G) \leq F^*(G) = F^*(F^*(G))$ . If  $F^*(G)$  is solvable, then  $F^*(G) = F(G)$ .*  
(3)  *$C_G(F^*(G)) \leq F(G)$ .*  
(4) *If  $G > 1$ , then  $F^*(G) > 1$ . In fact,  $F^*(G)/F(G) = \text{soc}(F(G)C_G(F(G))/F(G))$ .*

**Lemma 2.7.** [21, Lemma 2.5] *Let  $K$  be a normal subgroup of  $G$  and let  $H$  be a normal subgroup of  $K$ . If  $H$  is an  $\mathcal{H}C$ -subgroup of  $G$ , then  $H$  is  $c$ -normal in  $G$ .*

**Lemma 2.8.** *Let  $H$  be a  $p$ -subgroup of  $G$ . If  $H$  is an  $\mathcal{H}C$ -subgroup of  $G$  and  $H$  is not an  $\mathcal{H}$ -subgroup of  $G$ , then  $G$  has a normal subgroup  $M$  such that  $|G : M| = p$  and  $G = HM$ .*

*Proof.* By hypothesis,  $G$  has a normal subgroup  $T$  such that  $G = HT$  and  $H^g \cap N_T(H) \leq H$  for all  $g \in G$ . Since  $H$  is not an  $\mathcal{H}$ -subgroup of  $G$ , we have that  $T < G$ . Hence  $G/T$  is a  $p$ -group, and so  $G$  has a normal subgroup  $M$  containing  $T$  such that  $|G : M| = p$  and  $G = HM$ .  $\square$

**Lemma 2.9.** [11, Lemma 2.9] *Let  $\mathfrak{F}$  be a saturated formation containing all supersolvable groups and let  $G$  be a group with a normal subgroup  $E$  such that  $G/E \in \mathfrak{F}$ . If  $E$  is cyclic, then  $G \in \mathfrak{F}$ .*

**Lemma 2.10.** *Let  $\mathfrak{F}$  be a saturated formation containing all supersolvable groups. Suppose that  $M$  is a subgroup of  $G$  such that  $|G : M| = p$ ,  $F(G) \not\leq M$  and  $M \in \mathfrak{F}$ . Then  $G \in \mathfrak{F}$ .*

*Proof.* If  $\Phi(G) > 1$ , then it is easy to see that  $G/\Phi(G)$  satisfies the hypothesis of the lemma, and so  $G/\Phi(G) \in \mathfrak{F}$  by induction. This implies that  $G \in \mathfrak{F}$ . We may, therefore, assume that  $\Phi(G) = 1$ . Then  $F(G) = N_1 \times N_2 \cdots \times N_t$ , where  $N_i$  ( $i = 1, \dots, t$ ) is a solvable minimal normal subgroup of  $G$ . Since  $F(G) \not\leq M$ , there exists a solvable minimal normal subgroup  $N_i$  of  $G$  such that  $N_i \not\leq M$ . Then clearly,  $G = N_i M$ , and so  $N_i \cap M = 1$ . Therefore,  $|N_i| = |G : M| = p$  and  $G/N_i \cong M \in \mathfrak{F}$ . It follows from Lemma 2.9 that  $G \in \mathfrak{F}$ .  $\square$

**Lemma 2.11.** *Let  $H$  be an  $\mathcal{HC}$ -subgroup of  $G$ . If  $L/\Phi(L)$  is a chief factor of  $G$  and  $H \leq L$ , then  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ .*

*Proof.* By hypothesis, there exists a normal subgroup  $T$  of  $G$  such that  $G = HT$  and  $H^g \cap N_T(H) \leq H$  for all  $g \in G$ . Since  $L/\Phi(L)$  is a chief factor of  $G$ , either  $(L \cap T)\Phi(L)/\Phi(L) = 1$  or  $(L \cap T)\Phi(L)/\Phi(L) = L/\Phi(L)$ . In the former case,  $L = H(L \cap T) = H$ . This implies that  $H \trianglelefteq G$ , and so  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ . In the latter case,  $(L \cap T)\Phi(L) = L$ , and so  $T = G$ . This also implies that  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ .  $\square$

**Lemma 2.12.** [18, Lemma 2.8] *Let  $P$  be a normal  $p$ -subgroup of  $G$  contained in  $Z_\infty(G)$ . Then  $O^p(G) \leq C_G(P)$ .*

**Lemma 2.13.** [8, Lemma 2.4] *Let  $P$  be a  $p$ -group. If  $\alpha$  is a  $p'$ -automorphism of  $P$  which centralizes  $\Omega_1(P)$ , then  $\alpha = 1$  unless  $P$  is a non-abelian 2-group. If  $[\alpha, \Omega_2(P)] = 1$ , then  $\alpha = 1$  without restriction.*

### 3 Main results

**Theorem 3.1.** *Let  $p$  be the smallest prime divisor of  $|G|$  and let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Suppose that  $P$  is cyclic or  $P$  has a subgroup  $D$  with  $1 < |D| < |P|$  such that every subgroup  $H$  of  $P$  of order  $|D|$  is an  $\mathcal{HC}$ -subgroup of  $G$ . When  $p = 2$  and  $|P : D| > 2$ , suppose further that  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$  if there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4. Then  $G$  is  $p$ -nilpotent.*

*Proof.* Suppose that the result is false and let  $G$  be a counterexample of minimal order. Then we proceed via the following steps.

(1)  $P$  is not cyclic and  $|P : D| > p$ .

If  $P$  is cyclic, then by [19, (10.1.9)],  $G$  is  $p$ -nilpotent, a contradiction. Suppose that  $|P : D| = p$ . Then every maximal subgroup of  $P$  is an  $\mathcal{HC}$ -subgroup of  $G$ . Hence by Lemma 2.3,  $G$  is  $p$ -nilpotent, also a contradiction.

(2) Every proper subgroup of  $G$  containing  $P$  is  $p$ -nilpotent.

Let  $V$  be any proper subgroup of  $G$  containing  $P$ . Then by Lemma 2.2(1),  $V$  satisfies the hypothesis of the theorem. By the choice of  $G$ ,  $V$  is  $p$ -nilpotent. Thus (2) follows.

(3)  $O_{p'}(G) = 1$ .

If not, then by Lemma 2.2(3),  $G/O_{p'}(G)$  satisfies the hypothesis of the theorem. By the choice of  $G$ ,  $G/O_{p'}(G)$  is  $p$ -nilpotent, and so  $G$  is  $p$ -nilpotent, which is impossible.

(4)  $G$  is not a non-abelian simple group.

Assume that  $G$  is a non-abelian simple group. Then by Feit-Thompson's Theorem, we have that  $p = 2$ . Let  $H$  be a subgroup of  $P$  of order  $|D|$ . Then clearly,  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ . Hence by Lemma 2.1(2),  $H$  is an  $\mathcal{H}$ -subgroup of  $P$ , and thus  $H \trianglelefteq P$  by Lemma 2.1(1). It follows from (2) that  $N_G(H)$  is 2-nilpotent for  $H \not\trianglelefteq G$ , and so  $N_G(H)/C_G(H)$  is a 2-group. By Lemma 2.4,  $H$  is a Sylow 2-subgroup of  $G$ , a contradiction. Therefore,  $G$  is not a non-abelian simple group.

(5)  $O_p(G) > 1$ , and every proper normal subgroup of  $G$  is contained in  $O_p(G)$ .

Let  $L$  be a proper normal subgroup of  $G$ . Then we only need to prove that  $L$  is a  $p$ -group. By (3),  $p \mid |L|$ . If  $|L|_p > |D|$ , then  $L$  satisfies the hypothesis of the theorem by Lemma 2.2(1). Hence  $L$  is  $p$ -nilpotent due to the choice of  $G$ . It follows from (3) that  $L$  is a  $p$ -group. Now consider that  $|L|_p \leq |D|$ . Then there exists a normal subgroup  $K$  of  $P$  such that  $P \cap L \leq K$  and  $|K| = p|D|$ . This induces that  $|LK|_p = |K| = p|D|$ , and so  $K$  is a Sylow  $p$ -subgroup of  $LK$ . If  $LK = G$ , then  $|P| = |K| = p|D|$ , which contradicts (1). Thus  $LK < G$ . By Lemma 2.2(1),  $LK$  satisfies the hypothesis of the theorem. Then by the choice of  $G$ ,  $LK$  is  $p$ -nilpotent, and so is  $L$ . Hence  $L$  is a  $p$ -group by (3), and consequently (5) holds.

(6) Every  $\mathcal{HC}$ -subgroup of  $G$  contained in  $P$  is an  $\mathcal{H}$ -subgroup of  $G$ .

Let  $V$  be any  $\mathcal{HC}$ -subgroup of  $G$  contained in  $P$ . Then there exists a normal subgroup  $T$  of  $G$  such that  $G = VT$  and  $V^g \cap N_T(V) \leq V$  for all  $g \in G$ . By (5), since  $G$  is not a  $p$ -group, we have that  $T = G$ . Therefore,  $V$  is an  $\mathcal{H}$ -subgroup of  $G$ .

(7) Let  $N$  be a minimal normal subgroup of  $G$  contained in  $O_p(G)$ . Then  $|N| \leq |D|$  and  $G/N$  is  $p$ -nilpotent.

If  $|N| > |D|$ , then there exists a subgroup  $H$  of  $N$  of order  $|D|$  such that  $H$  is an  $\mathcal{H}$ -subgroup of  $G$  by (6). It follows from Lemma 2.1(1) that  $H$  is normal in  $G$ , which is impossible. Hence  $|N| \leq |D|$ . First suppose that  $|N| < |D|$ . Then by (6) and Lemma 2.1(3),  $G/N$  satisfies the hypothesis of the theorem. By the choice of  $G$ ,  $G/N$  is  $p$ -nilpotent.

Now consider that  $|N| = |D|$ . We claim that every cyclic subgroup of  $P/N$  of order prime or 4 (when  $p = 2$ ) is normal in  $N_G(P)/N$ . Let  $X/N$  be a subgroup of  $P/N$  of order  $p$ . If  $N \leq \Phi(X)$ , then  $X$  is cyclic, and so is  $N$ . This implies that  $|N| = |D| = p$ . Then by (6), every cyclic subgroup of  $P$  of order  $p$  or 4 (when  $p = 2$ ) is an  $\mathcal{H}$ -subgroup of  $G$ . By Lemmas 2.1(1) and 2.1(2), every cyclic subgroup of  $P$  of order  $p$  or 4 (when  $p = 2$ ) is normal in  $N_G(P)$ . Since  $p$  is the smallest prime divisor of  $|G|$ , every minimal subgroup of  $P$  lies in  $Z(N_G(P))$ . Hence by Lemma 2.5,  $G$  is  $p$ -nilpotent, a contradiction. Thus  $N \not\leq \Phi(X)$ , and so  $X$  has a maximal subgroup  $S$  such that  $X = SN$ . Since  $|S| = |N| = |D|$ ,  $S$  is an  $\mathcal{H}$ -subgroup of  $G$  by (6). By Lemmas 2.1(1) and 2.1(2),  $S \trianglelefteq N_G(P)$ , and thus  $X/N = SN/N \trianglelefteq N_G(P)/N$ . This shows that the claim holds when  $p$  is odd. Consider that  $p = 2$ . Then by (1),  $|P : D| > 2$ . Let  $Y/N$  be a cyclic subgroup of  $P/N$  of order 4. If  $N \leq \Phi(Y)$ , then  $Y$  is cyclic. This implies that  $|N| = |D| = 2$ , a contradiction. Thus  $N \not\leq \Phi(Y)$ , and

so  $Y$  has a maximal subgroup  $U$  such that  $Y = UN$ . Clearly,  $|U| = 2|D|$ . Since  $U/U \cap N \cong Y/N$  is a cyclic group of order 4, by hypothesis and (6),  $U$  is an  $\mathcal{H}$ -subgroup of  $G$ . A similar discussion as above shows that  $Y/N = UN/N \trianglelefteq N_G(P)/N$ . Hence the claim holds when  $p = 2$ . Since  $p$  is the smallest prime divisor of  $|G|$ , every minimal subgroup of  $P/N$  lies in  $Z(N_G(P)/N)$ . Therefore,  $G/N$  is  $p$ -nilpotent by Lemma 2.5.

(8) Final contradiction.

By (7),  $G/N$  is  $p$ -nilpotent. Then  $G$  has a normal subgroup  $M$  of  $G$  such that  $|G : M| = p$ . By (1),  $|M|_p > |D|$ . Then by (6) and Lemma 2.1(2),  $M$  satisfies the hypothesis of the theorem. Hence  $M$  is  $p$ -nilpotent due to the choice of  $G$ , and so  $G$  is  $p$ -nilpotent. The final contradiction completes the proof.  $\square$

The following corollary can be deduced immediately from Lemma 2.2(3) and Theorem 3.1.

**Corollary 3.2.** *Suppose that every noncyclic Sylow subgroup  $P$  (if exists) of  $G$  has a subgroup  $D$  such that  $1 < |D| < |P|$  and every subgroup  $H$  of  $P$  of order  $|D|$  is an  $\mathcal{HC}$ -subgroup of  $G$ . When  $P$  is a Sylow 2-subgroup of  $G$  and  $|P : D| > 2$ , suppose further that  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$  if there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4. Then  $G$  has a Sylow tower of supersolvable type.*

**Theorem 3.3.** *Let  $\mathfrak{F}$  be a saturated formation containing all supersolvable groups and let  $G$  be a group with a normal subgroup  $E$  such that  $G/E \in \mathfrak{F}$ . Suppose that every noncyclic Sylow subgroup  $P$  (if exists) of  $F^*(E)$  has a subgroup  $D$  such that  $1 < |D| < |P|$  and every subgroup  $H$  of  $P$  of order  $|D|$  is an  $\mathcal{HC}$ -subgroup of  $G$ . When  $P$  is a Sylow 2-subgroup of  $F^*(E)$  and  $|P : D| > 2$ , suppose further that  $H$  is an  $\mathcal{HC}$ -subgroup of  $G$  if there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4. Then  $G \in \mathfrak{F}$ .*

*Proof.* Suppose that the result is false and let  $(G, E)$  be a counterexample such that  $|G| + |E|$  is minimal. Then we proceed via the following steps.

(1)  $F^*(E) = F(E)$ .

By Lemma 2.2(1) and Corollary 3.2,  $F^*(E)$  has a Sylow tower of supersolvable type, and so  $F^*(E)$  is solvable. It follows from Lemma 2.6(2) that  $F^*(E) = F(E)$ .

(2) There exist a noncyclic Sylow  $p$ -subgroup  $P$  of  $F(E)$  and a subgroup  $H$  of  $P$  of order  $|D|$  or  $2|D|$  (when  $p = 2$ , and there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4) such that  $|P : D| > p$  and  $H$  is not an  $\mathcal{H}$ -subgroup of  $G$ .

Suppose that for every prime divisor  $p$  of  $|F(E)|$  and every noncyclic Sylow  $p$ -subgroup  $P$  of  $F(E)$ , either  $|P : D| = p$  or all subgroups  $H$  of  $P$  of order  $|D|$  or  $2|D|$  (when  $p = 2$ ,  $|P : D| > 2$  and there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$ )

and  $H/D_1$  is a cyclic group of order 4) are  $\mathcal{H}$ -subgroups of  $G$ . In the former case, by Lemma 2.7, all subgroups  $H$  of  $P$  of order  $|D|$  are c-normal in  $G$ . In the latter case, by Lemma 2.1(1), all subgroups  $H$  of  $P$  of order  $|D|$  or  $2|D|$  (when  $p = 2$ ,  $|P : D| > 2$  and there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4) are normal in  $G$ . Hence by [17, Theorem 1.4],  $G \in \mathfrak{F}$ , a contradiction. Thus (2) holds.

(3) Final contradiction.

By hypothesis and (2),  $H$  is an  $\mathcal{H}C$ -subgroup of  $G$  and  $H$  is not an  $\mathcal{H}$ -subgroup of  $G$ . Hence  $G$  has a normal subgroup  $M$  such that  $|G : M| = p$  and  $G = HM$  by Lemma 2.8. Since  $G = HM = EM$ , we have that  $M/E \cap M \cong G/E \in \mathfrak{F}$ . By Lemma 2.6(1),  $F^*(E \cap M) = F^*(E) \cap M = F(E) \cap M$ . Note that  $|F(E) : F(E) \cap M| = |G : M| = p$  and  $|F(E) \cap M|_p > |D|$  by (2). Then clearly,  $(M, E \cap M)$  satisfies the hypothesis of the theorem by Lemma 2.2(1). By the choice of  $(G, E)$ ,  $M \in \mathfrak{F}$ . It follows from Lemma 2.10 that  $G \in \mathfrak{F}$ . The final contradiction completes the proof.  $\square$

**Theorem 3.4.** *Let  $\mathfrak{F}$  be a saturated formation containing all supersolvable groups and let  $G$  be a group with a normal subgroup  $E$  such that  $G/E \in \mathfrak{F}$ . Suppose that every noncyclic Sylow subgroup  $P$  (if exists) of  $E$  has a subgroup  $D$  such that  $1 < |D| < |P|$  and every subgroup  $H$  of  $P$  of order  $|D|$  is an  $\mathcal{H}C$ -subgroup of  $G$ . When  $P$  is a Sylow 2-subgroup of  $E$  and  $|P : D| > 2$ , suppose further that  $H$  is an  $\mathcal{H}C$ -subgroup of  $G$  if there exists  $D_1 \trianglelefteq H \leq P$  such that  $2|D_1| = |D|$  and  $H/D_1$  is a cyclic group of order 4. Then  $G \in \mathfrak{F}$ .*

*Proof.* Suppose that the result is false and let  $(G, E)$  be a counterexample such that  $|G| + |E|$  is minimal. By Lemma 2.2(1) and Corollary 3.2, we see that  $E$  has a Sylow tower of supersolvable type. Without loss of generality, let  $p$  be the largest prime divisor of  $|E|$ . Then  $P \trianglelefteq G$ . By Lemma 2.2(3),  $(G/P, E/P)$  satisfies the hypothesis of the theorem. Then the choice of  $(G, E)$  implies that  $G/P \in \mathfrak{F}$ . Hence  $(G, P)$  satisfies the hypothesis of Theorem 3.3, and so  $G \in \mathfrak{F}$ .  $\square$

**Theorem 3.5.** *Let  $E$  be a normal subgroup of  $G$  such that  $G/E$  is nilpotent. Suppose that every minimal subgroup of  $E$  is contained in  $Z_\infty(G)$ , and every cyclic subgroup of  $E$  of order 4 is an  $\mathcal{H}C$ -subgroup of  $G$ . Then  $G$  is nilpotent.*

*Proof.* Assume that the result is false and let  $(G, E)$  be a counterexample such that  $|G| + |E|$  is minimal. Then we prove the theorem via the following steps.

(1)  $G$  is a minimal nonnilpotent group, that is,  $G = P \rtimes Q$ , where  $P$  is a normal Sylow  $p$ -subgroup of  $G$  and  $Q$  is a nonnormal cyclic Sylow  $q$ -subgroup of  $G$  for some prime  $q \neq p$ ;  $P/\Phi(P)$  is a chief factor of  $G$ ;  $\exp(P) = p$  when  $p > 2$  and  $\exp(P)$  is at most 4 when  $p = 2$ .

Let  $K$  be any proper subgroup of  $G$ . Then  $K/E \cap K \cong EK/E \leq G/E$  is nilpotent, and every minimal subgroup of  $E \cap K$  is contained in  $Z_\infty(G) \cap K \leq Z_\infty(K)$ .

By hypothesis, every cyclic subgroup of  $E \cap K$  of order 4 is an  $\mathcal{HC}$ -subgroup of  $G$ . Thus by Lemma 2.2(1), every cyclic subgroup of  $E \cap K$  of order 4 is an  $\mathcal{HC}$ -subgroup of  $K$ . Hence  $(K, E \cap K)$  satisfies the hypothesis of the theorem. Then the choice of  $(G, E)$  implies that  $K$  is nilpotent. Hence  $G$  is a minimal nonnilpotent group, and so (1) holds by [12, Chapter III, Satz 5.2].

(2)  $P \leq E$ .

If not, then  $P \cap E < P$ , and so  $(P \cap E)Q < G$ . By (1),  $(P \cap E)Q$  is nilpotent. This implies that  $Q \trianglelefteq (P \cap E)Q$ . Since  $G/P \cap E \lesssim G/P \times G/E$  is nilpotent,  $(P \cap E)Q \trianglelefteq G$ , and thus  $Q \trianglelefteq G$ , a contradiction.

(3) Final contradiction.

If  $\exp(P) = p$ , then  $P \leq Z_\infty(G)$ , and so  $G$  is nilpotent, which is impossible. Hence we may assume that  $p = 2$  and  $\exp(P) = 4$ . Then by Lemma 2.11, every cyclic subgroup of  $P$  of order 4 is an  $\mathcal{H}$ -subgroup of  $G$ , and so every cyclic subgroup of  $P$  of order 4 is normal in  $G$  by Lemma 2.1(1). Take an element  $x \in P \setminus \Phi(P)$ . Since  $P/\Phi(P)$  is a chief factor of  $G$ ,  $P = \langle x \rangle^G \Phi(P) = \langle x \rangle^G$ . If  $x$  is of order 2, then  $P = \langle x \rangle^G \leq Z_\infty(G)$ , a contradiction. Now assume that  $x$  is of order 4. Then  $\langle x \rangle \trianglelefteq G$ , and so  $P = \langle x \rangle$  is cyclic. By [19, (10.1.9)],  $G$  is 2-nilpotent, and so  $Q \trianglelefteq G$ . This is the final contradiction.  $\square$

**Theorem 3.6.** *Let  $E$  be a normal subgroup of  $G$  such that  $G/E$  is nilpotent. Suppose that every minimal subgroup of  $F^*(E)$  is contained in  $Z_\infty(G)$ , and every cyclic subgroup of  $F^*(E)$  of order 4 is an  $\mathcal{HC}$ -subgroup of  $G$ . Then  $G$  is nilpotent.*

*Proof.* Assume that the result is false and let  $(G, E)$  be a counterexample such that  $|G| + |E|$  is minimal. Then we prove the theorem via the following steps.

(1) Every proper normal subgroup of  $G$  is nilpotent.

Let  $K$  be any proper normal subgroup of  $G$ . Then  $K/E \cap K \cong EK/E \leq G/E$  is nilpotent. By Lemma 2.6(1),  $F^*(E \cap K) = F^*(E) \cap K$ . Hence by Lemma 2.2(1),  $(K, E \cap K)$  satisfies the hypothesis of the theorem. The the choice of  $(G, E)$  implies that  $K$  is nilpotent.

(2)  $E = G = \gamma_\infty(G)$  and  $F^*(G) = F(G) < G$ , where  $\gamma_\infty(G)$  is the nilpotent residual of  $G$ .

If  $E < G$ , then  $E$  is nilpotent by (1), and so  $F^*(E) = F(E) = E$ . By Theorem 3.5,  $G$  is nilpotent, a contradiction. Thus  $E = G$ . Now suppose that  $F^*(G) = G$ . Then by Theorem 3.5 again,  $G$  is nilpotent, which is impossible. Hence  $F^*(G) < G$ , and  $F^*(G) = F(G)$  by (1). If  $\gamma_\infty(G) < G$ , then by (1),  $\gamma_\infty(G) \leq F(G)$ , and so  $G/F(G)$  is nilpotent. It follows from Theorem 3.5 that  $G$  is nilpotent, a contradiction. Thus  $\gamma_\infty(G) = G$ .

(3) every cyclic subgroup of  $F(G)$  of order 4 is contained in  $Z(G)$ .

By hypothesis and (2), every cyclic subgroup  $H$  of  $F(G)$  of order 4 is an  $\mathcal{HC}$ -subgroup of  $G$ . Then there exists a normal subgroup  $T$  of  $G$  such that  $G = HT$



and  $H^g \cap N_T(H) \leq H$  for all  $g \in G$ . If  $T < G$ , then  $T \leq F(G)$  by (1), and thereby  $F(G) = G$ , a contradiction. Hence  $T = G$ , and so  $H$  is an  $\mathcal{H}$ -subgroup of  $G$ . By Lemma 2.1(1),  $H \trianglelefteq G$ . This implies that  $G/C_G(H)$  is abelian. Then by (2),  $C_G(H) = \gamma_\infty(G) = G$ , and so  $H \leq Z(G)$ . Thus (3) holds.

(4) Final contradiction.

Let  $p$  be any prime divisor of  $|F(G)|$  and let  $P$  be the Sylow  $p$ -subgroup of  $F(G)$ . Then  $P \trianglelefteq G$ . If  $p$  is odd, then by hypothesis,  $\Omega_1(P) \leq Z_\infty(G)$ . It follows from Lemma 2.12 that  $O^p(G) \leq C_G(\Omega_1(P))$ , and so  $O^p(G) \leq C_G(P)$  by Lemma 2.13. Then by (2),  $C_G(P) = \gamma_\infty(G) = G$ . Now consider that  $p = 2$ . Then by hypothesis and (3),  $\Omega_2(P) \leq Z_\infty(G)$ . A similar discussion as above also shows that  $C_G(P) = G$ . Therefore, we have that  $C_G(F(G)) = G$ , which contradicts the fact that  $C_G(F(G)) \leq F(G)$  by (2) and Lemma 2.6(3). The proof is thus completed.  $\square$

## References

- [1] M. Asaad, On  $p$ -nilpotence and supersolvability of finite groups, *Comm. Algebra*, **34** (2006), 189–195.
- [2] M. Asaad, A. A. Heliel, M. M. Al-Mosa Al-Shomrani, On weakly  $\mathcal{H}$ -subgroups of finite groups, *Comm. Algebra*, **40** (2012), 3540–3550.
- [3] M. Asaad, M. E. Mohamed, On  $c$ -normality of finite groups, *J. Aust. Math. Soc.*, **78** (2005), 297–304.
- [4] A. Ballester-Bolinches, Y. Wang, Finite groups with some  $C$ -normal minimal subgroups, *J. Pure Appl. Algebra*, **153** (2000), 121–127.
- [5] M. Bianchi, A. Gillio Berta Mauri, M. Herzog, L. Verardi, On finite solvable groups in which normality is a transitive relation, *J. Group Theory*, **3** (2000), 147–156.
- [6] P. Csörgö, M. Herzog, On supersolvable groups and the nilpotator, *Comm. Algebra*, **32** (2004), 609–620.
- [7] K. Doerk, T. Hawkes, *Finite Soluble Groups*, Berlin, Walter de Gruyter, 1992.
- [8] T. M. Gagen, *Topics in Finite Groups*, Cambridge, Cambridge University Press, 1976.
- [9] D. M. Goldschmidt, Strongly closed 2-subgroups of finite groups, *Ann. of Math.*, **102** (1975), 475–489.
- [10] W. Guo, *The Theory of Classes of Groups*, Dordrecht, Kluwer Academic Publishers, 2000.

- [11] W. Guo, A. N. Skiba, Finite groups with given  $s$ -embedded and  $n$ -embedded subgroups, *J. Algebra*, **321** (2009), 2843–2860.
- [12] B. Huppert, *Endliche Gruppen I*, Berlin, Springer-Verlag, 1967.
- [13] B. Huppert, Blackburn N, *Finite Groups III*, Berlin–New York, Springer-Verlag, 1982.
- [14] J. J. Jaraden, A. N. Skiba, On  $c$ -normal subgroups of finite groups, *Comm. Algebra*, **35** (2007), 3776–3788.
- [15] D. Li, X. Guo, The influence of  $c$ -normality of subgroups on the structure of finite groups, *J. Pure Appl. Algebra*, **150** (2000), 53–60.
- [16] S. Li, On minimal subgroups of finite groups, *Comm. Algebra*, **22** (1994), 1913–1918.
- [17] Y. Li, S. Qiao, Y. Wang, On weakly  $s$ -permutably embedded subgroups of finite groups, *Comm. Algebra*, **37** (2009), 1086–1097.
- [18] Y. Li, Y. Wang, On  $\pi$ -quasinormally embedded subgroups of finite group, *J. Algebra*, **281** (2004), 109–123.
- [19] D. J. S. Robinson, *A Course in the Theory of Groups*, New York, Springer-Verlag, 1996.
- [20] Y. Wang,  $C$ -normality of groups and its properties, *J. Algebra*, **180** (1996), 954–965.
- [21] X. Wei, X. Guo, On  $\mathcal{H}C$ -subgroups and the structure of finite groups, *Comm. Algebra*, **40** (2012), 3245–3256.