
A provable SVD-based algorithm for learning topics in dominant admixture corpus

Trapit Bansal[†], C. Bhattacharyya[‡]
 Department of Computer Science and Automation
 Indian Institute of Science
 Bangalore -560012, India
[†]trapitbansal@gmail.com
[‡]chiru@csa.iisc.ernet.in

Ravindran Kannan
 Microsoft Research
 India
 kannan@microsoft.com

Abstract

Topic models, such as Latent Dirichlet Allocation (LDA), posit that documents are drawn from ad-mixtures of distributions over words, known as topics. The inference problem of recovering topics from such a collection of documents drawn from admixtures, is NP-hard. Making a strong assumption called *separability*, [4] gave the first provable algorithm for inference. For the widely used LDA model, [6] give a provable algorithm using clever tensor-methods. But [4] and [6] do not learn topic vectors with bounded l_1 error (which is the natural measure for probability vectors) and they need assumptions on numerical parameters like the condition number.

Our aim here is to develop a model which makes intuitively relevant and empirically supported assumptions and to design an algorithm with natural, simple components (one of them being SVD), which provably solves the inference problem for the model with bounded l_1 error. A topic in LDA and other models is essentially characterized by a group of co-occurring words. Motivated by this, we introduce topic specific *Catchwords*, a group of words which occur with strictly greater frequency in a topic than any other topic individually and are required to have high frequency together rather than individually. A major contribution of the paper is to show that under this more realistic assumption (as we show empirically for real corpora), a Singular value decomposition (SVD) based algorithm with a crucial pre-processing step of thresholding, can provably recover the topics from a collection of documents drawn from *Dominant admixtures*. Dominant admixtures are convex combination of distributions in which one distribution has a significantly high contribution than the other distributions. It is folklore that SVD can solve inference only if we have documents with pure topics. We overcome this by doing a thresholding procedure first before SVD and a k -means procedure after SVD. Using recent results on k -means, we show that our algorithm correctly identifies the dominant topic in most documents. Apart from the simplicity of the algorithm, the sample complexity (the number of documents needed) has near optimal dependence on w_0 , the lowest probability that a topic is dominant, and is better than [4]. Empirical evidence shows that on several real world corpora, both *Catchwords* and *Dominant admixture* assumptions hold and the proposed algorithm substantially outperforms the state of the art [5].

1 Introduction

Topic models [1] assume that each document in a text corpus is generated from an *ad-mixture* of topics, where, each topic is a distribution over words in a Vocabulary. An admixture is a convex

combination of distributions. Words in the document are then picked in i.i.d. trials, each trial has a multinomial distribution over words given by the weighted combination of topic distributions. The problem of inference, recovering the topic distributions from such a collection of documents, is provably NP-hard. Existing literature pursues techniques such as variational methods [2] or MCMC procedures [3] for approximating the maximum likelihood estimates.

Given the intractability of the problem one needs further assumptions on topics to derive polynomial time algorithms which can provably recover topics. A possible (strong) assumption is that each document has only one topic but the collection can have many topics. A document with only one topic is sometimes referred as a *pure topic* document. [7] proved that a natural algorithm, based on SVD, recovers topics when each document is pure and in addition, for each topic, there is a set of words, called *primary words*, whose total frequency in that topic is close to 1. More recently, [6] show using tensor methods that if the topic weights have Dirichlet distribution, we can learn the topic matrix. Note that while this allows non-pure documents, the Dirichlet distribution gives essentially uncorrelated topic weights.

In an interesting recent development [4, 5] gave the first provable algorithm which can recover topics from a corpus of documents drawn from admixtures, assuming *separability*. Topics are said to be separable if in every topic there exists at least one *Anchor* word. A word in a topic is said to be an *Anchor* word for that topic if it has a high probability in that topic and *zero* probability in remaining topics. The requirement of high probability in a topic for a single word is unrealistic.

Our Contributions: Topic distributions, such as those learnt in LDA, try to model the occurrence of a group of words which describes a theme. Keeping this in mind we introduce the notion of *Catchwords*. A group of words are called *Catchwords* of a topic, if each word occurs strictly more frequently in the topic than other topics and together they have high frequency. This is a much weaker assumption than separability. Furthermore we observe, empirically, that posterior topic weights assigned by LDA to a document often have the property that one of the weights is significantly higher than the rest. Motivated by this observation, which has not been exploited by topic modelling literature, we suggest a new assumption. It is natural to assume that in a text corpus, a document, even if it has multiple themes, will have an overarching dominant theme. In this paper we focus on document collections drawn from *dominant admixtures*. A document collection is said to be drawn from a dominant admixture if for every document, there is one topic whose weight is significantly higher than the other topics and in addition, for every topic, there is a small fraction of documents which are nearly purely on that topic. The main contribution of the paper is to show that under these assumptions, our algorithm, which we call TSVD, indeed provably finds a good approximation in total l_1 error to the topic matrix. We prove a bound on the error of our approximation which does not grow with the dictionary size d , unlike [5], where the error grows linearly with d .

Empirical evidence shows that on semi-synthetic corpora constructed from several real world datasets, as suggested by [5], TSVD substantially outperforms the state of the art [5]. In particular it is seen that compared to [5] TSVD gives 27% lower error in terms of L_1 recovery on 90% of the topics.

Problem Definition: d, k, s will denote respectively, the number of words in the dictionary, number of topics and number of documents. d, s are large, whereas, k is to be thought of as much smaller. Let $\mathcal{S}_k = \{x = (x_1, x_2, \dots, x_k) : x_l \geq 0; \sum_l x_l = 1\}$. For each topic, there is a fixed vector in \mathcal{S}_k giving the probability of each word in that topic. Let \mathbf{M} be the $d \times k$ matrix with these vectors as its columns.

Documents are picked in i.i.d. trials. To pick document j , one first picks a k - vector $W_{1j}, W_{2j}, \dots, W_{kj}$ of topic weights according to a fixed distribution on \mathcal{S}_k . Let $P_{\cdot,j} = \mathbf{M}W_{\cdot,j}$ be the weighted combination of the topic vectors. Then the m words of the document are picked in i.i.d. trials; each trial picks a word according to the multinomial distribution with $P_{\cdot,j}$ as the probabilities. All that is given as data is the frequency of words in each document, namely, we are given the $d \times s$ matrix \mathbf{A} , where $A_{ij} = \frac{\text{Number of occurrences of word } i \text{ in Document } j}{m}$. Note that $E(\mathbf{A}|\mathbf{W}) = \mathbf{P}$, where, the expectation is taken entry-wise.

In this paper we consider the problem of finding \mathbf{M} given \mathbf{A} .

2 Previous Results

In this section we review literature related to designing provable algorithms for topic models. For an overview of topic models we refer the reader to the excellent survey[1]. Provable algorithms for recovering topic models was started by [7]. Latent Semantic Indexing(LSI) [8] remains a successful method for retrieving similar documents by using SVD. [7] showed that one can recover \mathbf{M} from a collection of documents, with *pure topics*, by using SVD based procedure under the additional Primary Words Assumption. [6] showed that in the admixture case, if one assumes Dirichlet distribution for the topic weights, then, indeed, using tensor methods, one can learn \mathbf{M} to l_2 error provided some added assumptions on numerical parameters like condition number are satisfied.

The first provably polynomial time algorithm for admixture corpus was given in [4, 5]. For a topic l , a word i is an **anchor word** if $M_{i,l} \geq p_0$; $M_{i,l'} = 0 \quad \forall l' \neq l$.

Theorem 2.1 [4] *If every topic has an anchor word, there is a polynomial time algorithm that returns an \hat{M} such that with high probability,*

$$\sum_{l=1}^k \sum_{i=1}^d |\hat{M}_{il} - M_{il}| \leq d\varepsilon \quad \text{provided } s \geq \text{Max} \left\{ O \left(\frac{k^6 \log d}{a^4 \varepsilon^2 p_0^6 \gamma^2 m} \right), O \left(\frac{k^4}{\gamma^2 a^2} \right) \right\},$$

where, γ is the condition number of $E(WW^T)$, a is the minimum expected weight of a topic and m is the number of words in each document.

Note that the error grows linearly in the dictionary size d , which is often large. Note also the dependence of s on parameters p_0 , which is, $1/p_0^6$ and on a , which is $1/a^4$. If, say, the word “run” is an anchor word for the topic “baseball” and $p_0 = 0.1$, then the requirement is that every 10 th word in a document on this topic is “run”. This seems too strong to be realistic. It would be more realistic to ask that a set of words like - “run”, “hit”, “score”, etc. together have frequency at least 0.1 which is what our catchwords assumption does.

3 Learning Topics from Dominant Admixtures

Informally, a document is said to be drawn from a Dominant Admixture if the document has one *dominant topic*. Besides its simplicity, we show empirical evidence from real corpora to demonstrate that topic dominance is a reasonable assumption. The Dominant Topic assumption is weaker than the Pure Topic assumption. More importantly SVD based procedures proposed by [7] will not apply. Inspired by the *Primary words* assumption we introduce the assumption that each topic has a set of *Catchwords* which individually occur more frequently in that topic than others. This is again a much weaker assumption than both *Primary Words* and *Anchor Words* assumptions and can be verified experimentally. In this section we establish that by applying SVD on a matrix, obtained by thresholding the word-document matrix, and subsequent k means clustering can learn topics having Catchwords from a Dominant Admixture corpus.

3.1 Assumptions: Catchwords and Dominant admixtures

Let $\alpha, \beta, \rho, \delta, \varepsilon_0$ be non-negative reals satisfying:

$$\beta + \rho \leq (1 - \delta)\alpha. \tag{1}$$

$$\alpha + 2\delta \leq 0.5 \quad ; \quad \delta \leq 0.08. \tag{2}$$

Dominant topic Assumption (a) For $j = 1, 2, \dots, s$, document j has a dominant topic $l(j)$ such that $W_{l(j),j} \geq \alpha$ and $W_{l',j} \leq \beta, \quad \forall l' \neq l(j)$.

(b)For each topic l , there are at least $\varepsilon_0 w_0 s$ documents in each of which topic l has weight at least $1 - \delta$.

Catchwords Assumption: There are k disjoint sets of words - S_1, S_2, \dots, S_k such that with ε defined in (9)

$$\forall i \in S_l, \forall l' \neq l, M_{il'} \leq \rho M_{il} \quad (3)$$

$$\sum_{i \in S_l} M_{il} \geq p_0 \quad (4)$$

$$\forall i \in S_l, m\delta^2\alpha M_{il} \geq 8 \ln \left(\frac{20}{\varepsilon w_0} \right). \quad (5)$$

Part (b.) of the Dominant Topic Assumption is in a sense necessary for ‘‘identifiability’’ - namely for the model to have a set of k document vectors so that every document vector is in the convex hull of these vectors. The Catchwords assumption is natural to describe a theme as it tries to model a unique group of words which is likely to co-occur when a theme is expressed. This assumption is close to topics discovered by LDA like models, which try to model of co-occurrence of words. If $\alpha, \delta \in \Omega(1)$, then, the assumption (5) says $M_{il} \in \Omega^*(1/m)$. In fact if $M_{il} \in o(1/m)$, we do not expect to see word i (in topic l), so it cannot be called a catchword at all.

A slightly different (but equivalent) description of the model will be useful to keep in mind. What is fixed (not stochastic) are the matrices \mathbf{M} and the distribution of the weight matrix \mathbf{W} . To pick document j , we can first pick the dominant topic l in document j and condition the distribution of $W_{\cdot,j}$ on this being the dominant topic. One could instead also think of $W_{\cdot,j}$ being picked from a mixture of k distributions. Then, we let $P_{ij} = \sum_{l=1}^k M_{il} W_{lj}$ and pick the m words of the document in i.i.d multinomial trials as before. We will assume that

$$T_l = \{j : l \text{ is the dominant topic in document } j\} \text{ satisfies } |T_l| = w_l s,$$

where, w_l is the probability of topic l being dominant. This is only approximately valid, but the error is small enough that we can disregard it.

For $\zeta \in \{0, 1, 2, \dots, m\}$, let $p_i(\zeta, l)$ be the probability that $j \in T_l$ and $A_{ij} = \zeta/m$ and $q_i(\zeta, l)$ the corresponding ‘‘empirical probability’’:

$$p_i(\zeta, l) = \int_{W_{\cdot,j}} \binom{m}{\zeta} P_{ij}^\zeta (1 - P_{ij})^{m-\zeta} \text{Prob}(W_{\cdot,j} | j \in T_l) \text{Prob}(j \in T_l), \text{ where, } P_{\cdot,j} = \mathbf{M}W_{\cdot,j}. \quad (6)$$

$$q_i(\zeta, l) = \frac{1}{s} |\{j \in T_l : A_{ij} = \zeta/m\}|. \quad (7)$$

Note that $p_i(\zeta, l)$ is a real number, whereas, $q_i(\zeta, l)$ is a random variable with $E(q_i(\zeta, l)) = p_i(\zeta, l)$. We need a technical assumption on the $p_i(\zeta, l)$ (which is weaker than unimodality).

No-Local-Min Assumption: We assume that $p_i(\zeta, l)$ does not have a local minimum, in the sense:

$$p_i(\zeta, l) > \text{Min}(p_i(\zeta - 1, l), p_i(\zeta + 1, l)) \quad \forall \zeta \in \{1, 2, \dots, m - 1\}. \quad (8)$$

The justification for the this assumption is two-fold. First, generally, Zipf’s law kind of behaviour where the number of words plotted against relative frequencies declines as a power function has often been observed. Such a plot is monotonically decreasing and indeed satisfies our assumption. But for Catchwords, we do not expect this behaviour - indeed, we expect the curve to go up initially as the relative frequency increases, then reach a maximum and then decline. This is a unimodal function and also satisfies our assumption. Indeed, we have empirically observed, see EXPTS, that these are essentially the only two behaviours.

Relative sizes of parameters Before we close the section we discuss the values of the parameters are in order. Here, s is large. For asymptotic analysis, we can think of it as going to infinity. $1/w_0$ is also large and can be thought of as going to infinity. [In fact, if $1/w_0 \in O(1)$, then, intuitively, we see that there is no use of a corpus of more than constant size - since our model has i.i.d. documents, intuitively, the number of samples we need should depend mainly on $1/w_0$]. m is (much) smaller, but need not be constant.

c refers to a generic constant independent of $m, s, 1/w_0, \varepsilon, \delta$; its value may be different in different contexts.

3.2 The TSVD Algorithm

Existing SVD based procedures for clustering on raw word-document matrices fail because the spread of frequencies of a word within a topic is often more (at least not significantly less) than the gap between the word's frequencies in two different topics. Hypothetically the frequency for the word *run*, in the topic *Sports*, may range from say 0.01 on up. But in other topics, it may range from 0 to 0.005 say. The success of the algorithm will lie on correctly identifying the dominant topics such as sports by identifying that the word *run* has occurred with high frequency. In this example, the gap (0.01-0.005) between Sports and other topics is less than the spread within Sports (1.0-0.01), so a 2-clustering approach (based on SVD) will split the topic Sports into two. While this is a toy example, note that if we threshold the frequencies at say 0.01, ideally, sports will be all above and the rest all below the threshold, making the succeeding job of clustering easy.

There are several issues in extending beyond the toy case. Data is not one-dimensional. We will use different thresholds for each word; word i will have a threshold ζ_i/m . Also, we have to compute ζ_i/m . Ideally we would not like to split any T_l , namely, we would like that for each l and each i , either most $j \in T_l$ have $A_{ij} > \zeta_i/m$ or most $j \in T_l$ have $A_{ij} \leq \zeta_i/m$. We will show that our threshold procedure indeed achieves this. One other nuance: to avoid conditioning, we split the data \mathbf{A} into two parts $\mathbf{A}^{(1)}$ and $\mathbf{A}^{(2)}$, compute the thresholds using $\mathbf{A}^{(1)}$ and actually do the thresholding on $\mathbf{A}^{(2)}$. We will assume that the initial \mathbf{A} had $2s$ columns, so each part now has s columns. Also, T_1, T_2, \dots, T_k partitions the columns of $\mathbf{A}^{(1)}$ as well as those of $\mathbf{A}^{(2)}$. The columns of thresholded matrix \mathbf{B} are then clustered, by a technique we call Project and Cluster, namely, we project the columns of \mathbf{B} to its k -dimensional SVD subspace and cluster in the projection. The projection before clustering has recently been proven [9] (see also [10]) to yield good starting cluster centers. The clustering so found is not yet satisfactory. We use the classic Lloyd's k -means algorithm proposed by [12]. As we will show, the partition produced after clustering, $\{R_1, \dots, R_k\}$ of $\mathbf{A}^{(2)}$ is close to the partition induced by the Dominant Topics, $\{T_1, \dots, T_k\}$. Catchwords of topic l are now (approximately) identified as the most frequently occurring words in documents in R_l . Finally, we identify nearly pure documents in T_l (approximately) as the documents in which the catchwords occur the most. Then we get an approximation to $M_{.l}$ by averaging these nearly pure documents. We now describe the precise algorithm.

3.3 Topic recovery using Thresholded SVD

Threshold SVD based K-means (TSVD)

$$\varepsilon = \text{Min} \left(\frac{1}{900c_0^2} \frac{\alpha p_0}{k^3 m}, \frac{\varepsilon_0 \sqrt{\alpha p_0} \delta}{640m\sqrt{k}} \right). \quad (9)$$

1. Randomly partition the columns of \mathbf{A} into two matrices $\mathbf{A}^{(1)}$ and $\mathbf{A}^{(2)}$ of s columns each.

2. Thresholding

(a) **Compute Thresholds on $\mathbf{A}^{(1)}$** For each i , let ζ_i be the highest value of $\zeta \in \{0, 1, 2, \dots, m\}$ such that $|\{j : A_{ij}^{(1)} > \frac{\zeta}{m}\}| \geq \frac{w_0}{2}s$; $|\{j : A_{ij}^{(1)} = \frac{\zeta}{m}\}| \leq 3\varepsilon w_0 s$.

(b) **Do the thresholding on $\mathbf{A}^{(2)}$** : $B_{ij} = \begin{cases} \sqrt{\zeta_i} & \text{if } A_{ij}^{(2)} > \zeta_i/m \text{ and } \zeta_i \geq 8 \ln(20/\varepsilon w_0) \\ 0 & \text{otherwise} \end{cases}$.

3. **SVD** Find the best rank k approximation $\mathbf{B}^{(k)}$ to \mathbf{B} .

4. Identify Dominant Topics

(a) **Project and Cluster** Find (approximately) optimal k -means clustering of the columns of $\mathbf{B}^{(k)}$.

(b) **Lloyd's Algorithm** Using the clustering found in Step 4(a) as the starting clustering, apply Lloyd's algorithm k means algorithm to the columns of \mathbf{B} (\mathbf{B} , not $\mathbf{B}^{(k)}$).

(c) Let R_1, R_2, \dots, R_k be the k -partition of $[s]$ corresponding to the clustering after Lloyd's. // *Will prove that $R_l \approx T_l$ * //

5. Identify Catchwords

(a) For each i, l , compute $g(i, l) =$ the $\lfloor \varepsilon_0 w_0 s / 2 \rfloor$ th highest element of $\{A_{ij}^{(2)} : j \in R_l\}$.

(b) Let $J_l = \{i : g(i, l) > \text{Max} \left(\frac{4}{m\delta^2} \ln(20/\varepsilon w_0), \text{Max}_{l' \neq l} \gamma g(i, l') \right)\}$, where, $\gamma = \frac{1-2\delta}{(1+\delta)(\beta+\rho)}$.

6. **Find Topic Vectors** Find the $\lfloor \varepsilon_0 w_0 s / 2 \rfloor$ highest $\sum_{i \in J_l} A_{ij}^{(2)}$ among all $j \in [s]$ and return the average of these $A_{.,j}$ as our approximation $\hat{M}_{.,l}$ to $M_{.,l}$.

Theorem 3.1 Main Theorem *Under the Dominant Topic, Catchwords and No-Local-Min assumptions, the algorithm succeeds with high probability in finding an \hat{M} so that*

$$\sum_{i,l} |M_{il} - \hat{M}_{il}| \in O(k\delta), \text{ provided }^1 s \in \Omega^* \left(\frac{1}{w_0} \left(\frac{k^6 m^2}{\alpha^2 p_0^2} + \frac{m^2 k}{\varepsilon_0^2 \delta^2 \alpha p_0} + \frac{d}{\varepsilon_0 \delta^2} \right) \right).$$

A note on the sample complexity is in order. Notably, the dependence of s on w_0 is best possible (namely $s \in \Omega^*(1/w_0)$) within logarithmic factors, since, if we had fewer than $1/w_0$ documents, a topic which is dominant with probability only w_0 may have none of the documents in the collection. The dependence of s on d needs to be at least $d/\varepsilon_0 w_0 \delta^2$: to see this, note that we only assume that there are $r = O(\varepsilon_0 w_0 s)$ nearly pure documents on each topic. Assuming we can find this set (the algorithm approximately does), their average has standard deviation of about $\sqrt{M_{il}}/\sqrt{r}$ in coordinate i . If topic vector $M_{.,l}$ has $O(d)$ entries, each of size $O(1/d)$, to get an approximation of $M_{.,l}$ to l_1 error δ , we need the per coordinate error $1/\sqrt{dr}$ to be at most δ/d which implies $s \geq d/\varepsilon_0 w_0 \delta^2$. Note that to get comparable error in [4], we need a quadratic dependence on d .

There is a long sequence of Lemmas to prove the theorem. The Lemmas and the proofs are given in Appendix. The essence of the proof lies in proving that the clustering step correctly identifies the partition induced by the dominant topics. For this, we take advantage of a recent development on the k -means algorithm from [9] [see also [10]], where, it is shown that under a condition called the *Proximity Condition*, Lloyd's k means algorithm starting with the centers provided by the SVD-based algorithm, correctly identifies almost all the documents' dominant topics. We prove that indeed the Proximity Condition holds. This calls for machinery from Random Matrix theory (in particular bounds on singular values). We prove that the singular values of the thresholded word-document matrix are nicely bounded. Once the dominant topic of each document is identified, we are able to find the Catchwords for each topic. Now, we rely upon part (b.) of the Dominant Topic assumption : that is there is a small fraction of nearly Pure Topic-documents for each topic. The Catchwords help isolate the nearly pure-topic documents and hence find the topic vectors. The proofs are complicated by the fact that each step of the algorithm induces conditioning on the data- for example, after clustering, the document vectors in one cluster are not anymore independent.

4 Experimental Results

We compare the thresholded SVD based k -means (**TSVD**) algorithm of Section 3.2 with the algorithms of [5], *Recover-KL* and *Recover-L2*, using the code made available by the authors². We observed the results of Recover-KL to be better than Recover-L2, and report here the results of Recover-KL (R-KL) (full set of results can be found in supplementary section 2). We first provide empirical support for the algorithm assumptions in Section 3.1, namely the dominant topic and the catchwords assumption. Then we show on 4 different semi-synthetic data that TSVD provides as good or better recovery of topics than the Recover algorithms. Finally on real-life datasets, we show that the algorithm performs as well as [5] in terms of perplexity and topic coherence.

Implementation Details: For TSVD, thresholding parameters used were: $w_0 = \frac{1}{k}$, $\varepsilon = \frac{1}{6}$. Using a different value of $\varepsilon = 0.5$ for finding the catchwords (step 5 in 3.2) and taking the 50% fractile for finding the topic vectors (step 6) gave empirically best results. The new algorithm is sensitive to the initialization of the first k -means step in the projected SVD space. To remedy this, we run 10 independent random initialization of the algorithm with K-Means++ [13] and report the best result.

¹The superscript * hides a logarithmic factor in dk/δ_{fail} , where, $\delta_{\text{fail}} > 0$ is the desired upper bound on the probability of failure.

²<http://www.cs.nyu.edu/~halpern/files/anchor-word-recovery.zip>

| Corpus | Documents (s) | Topics (k) | % s with Dominant Topics | | | % Topics with CW | CW Mean Frequency |
|--------|----------------------|-------------------|--------------------------|----------------|----------------|---------------------|----------------------|
| | | | $\alpha = 0.4$ | $\alpha = 0.8$ | $\alpha = 0.9$ | | |
| NIPS | 1,500 | 50 | 56.6% | 10.7% | 4.8% | 95% | 0.05 |
| NYT | 30,000 | 50 | 63.7% | 20.9% | 12.7% | 100% | 0.11 |
| Pubmed | 30,000 | 50 | 62.2% | 20.3% | 10.7% | 90% | 0.05 |
| 20NG | 13,389 | 20 | 74.1% | 54.4% | 44.3% | 100% | 0.06 |

Table 1: Algorithm Assumptions. For dominant topic assumption, fraction of documents with satisfy the assumption for (α, β) are shown. $\beta = 0.3$ for $\alpha = 0.4$ and $\alpha = 0.8, 0.9$ are “almost pure topic” cases (i.e. $\delta = 0.2, 0.1$). Last two columns show results for catchwords (CW) assumption.

Datasets: We use four real word datasets in the experiments. As pre-processing steps we removed standard stop-words, selected the vocabulary size by term-frequency and removed documents with less than 20 words. Datasets are: (1) **NIPS**³: Consists of 1,500 NIPS full papers, vocabulary of 2,000 words and mean document length 1023. (2) **NYT**³: Consists of a random subset of 30,000 documents from the New York Times dataset, vocabulary of 5,000 words and mean document length 238. (3) **Pubmed**³: Consists of a random subset of 30,000 documents from the Pubmed abstracts dataset, vocabulary of 5,030 words and mean document length 58. (4) **20NewsGroup**⁴ (20NG): Consist of 13,389 documents, vocabulary of 7,118 words and mean document length 160.

4.1 Algorithm Assumptions

To check the *dominant topic* and *catchwords* assumptions, we first run 1000 iterations of Gibbs sampling on the real corpus and learn the posterior document-topic distribution ($\{W_{\cdot,j}\}$) for each document in the corpus (by averaging over 10 saved-states separated by 50 iterations after the 500 burn-in iterations). We will use this posterior document-topic distribution as the document generating distribution to check the two assumptions.

Dominant topic assumption: Table 1 shows the fraction of the documents in each corpus which satisfy this assumption with $\alpha = 0.4$ (minimum probability of dominant topic) and $\beta = 0.3$ (maximum probability of non-dominant topics). The fraction of documents which have dominant topic weight at least 0.8 and 0.9 ($\delta = 0.2, 0.1$) are also shown. The results show that the assumption is well justified and there is also a good fraction of documents satisfying almost pure topic assumption.

Catchwords assumption: We first find a k -clustering of the documents $\{T_1, \dots, T_k\}$ by assigning all documents which have highest posterior probability for the same topic into one cluster. Then we use step 5 of TSVD (Section 3.2) to find the set of catchwords for each topic-cluster, i.e. $\{S_1, \dots, S_k\}$, with the parameter $\varepsilon = 0.5$. Table 1 (last two columns) reports the fraction of topics with non-empty set of catchwords and the mean per topic frequency of the catchwords⁵. Results indicate that most topics on real data contain catchwords and the mean per topic frequency of the catchwords is also quite high.

4.2 Empirical Results

Semi-synthetic Data: Following [5], we generate semi-synthetic corpora from LDA model trained by MCMC, to ensure that the synthetic corpora retain the characteristics of real data. Gibbs sampling is run for 1000 iterations on all the four datasets and the final word-topic distribution is used to generate varying number of synthetic documents with document-topic distribution drawn from a symmetric Dirichlet with hyper-parameter 0.01. For NIPS, NYT and Pubmed we use $k = 50$ topics, for 20NewsGroup $k = 20$, and mean document lengths of 1000, 300, 100 and 200 respectively. Note that the synthetic data is *not* guaranteed to satisfy dominant topic assumption for every document (on average about 80% documents satisfy the assumption for value of (α, β) tested in Section 4.1)

Topic Recovery: We learn the word-topic distribution (\hat{M}) for the semi-synthetic corpora using TSVD and the Recover algorithms of [5]. Given these learned topic distributions and the original

³<http://archive.ics.uci.edu/ml/datasets/Bag+of+Words>

⁴<http://qwone.com/~jason/20Newsgroups>

⁵ $\left(\frac{1}{k} \sum_{l=1}^k \frac{1}{|T_l|} \sum_{i \in S_l} \sum_{j \in T_l} A_{ij}\right)$




Figure 1: Histogram of L_1 error across topics (40,000 documents). *TSVD*(blue, solid border) gets smaller error on most topics than *R-KL*(green, dashed border).

| Corpus | Documents | R-KL | TSVD |
|--------|-----------|--------------|----------------------|
| NIPS | 40,000 | 0.308 | 0.094 (69.6%) |
| | 50,000 | 0.308 | 0.135 (56.3%) |
| | 60,000 | 0.311 | 0.124 (60.2%) |
| Pubmed | 40,000 | 0.332 | 0.291 (12.5%) |
| | 50,000 | 0.326 | 0.290 (11.2%) |
| | 60,000 | 0.328 | 0.297 (9.6%) |
| 20NG | 40,000 | 0.120 | 0.125 (-4.1%) |
| | 50,000 | 0.114 | 0.115 (-0.7%) |
| | 60,000 | 0.110 | 0.108 (2.5%) |
| NYT | 40,000 | 0.208 | 0.198 (4.8%) |
| | 50,000 | 0.206 | 0.186 (9.4%) |
| | 60,000 | 0.200 | 0.192 (4.1%) |

Table 2: L_1 reconstruction error on various semi-synthetic datasets. Brackets in the last column give percent improvement over R-KL (best performing Recover algorithm). Full results in supplementary.

data-generating distribution (M), we align the topics of M and \hat{M} by bipartite matching and evaluate the L_1 distance between each pair of topics. We report the average of L_1 error across topics (reconstruction-error) in Table 2 for TSVD and Recover-KL (R-KL). TSVD has smaller error on most datasets than the Recover-KL algorithm. We observed performance of TSVD to be always better than Recover-L2 (see supplement Table 1 for full results). Best performance is observed on NIPS which has largest mean document length, indicating that larger m leads to better recovery. Results on 20NG are slightly worse than R-KL but the difference is small. While the values in Table 2 are averaged values, Figure 1 shows that TSVD algorithm achieves much better topic recovery for majority of the topics (>90%) on most datasets (full results in supplement Figure 1).

Real Data: To evaluate perplexity[2] on real data, the held-out sets consist of 350 documents for NIPS, 10000 documents for NYT and Pubmed, and 6780 documents for 20NewsGroup. In addition to TSVD, we tested for TSVD-100 which uses all the documents (100% as opposed to 50%) in step 6 (Algorithm 3.2). *TSVD-100* achieved perplexity measure of **830 (NIPS)**, **1271 (Pubmed)**, **1521 (NYT)**, **2300 (20NG)** while *Recover-KL* achieved **754 (NIPS)**, **1188 (Pubmed)**, **1579 (NYT)**, **2431 (20NG)** (refer to supplement Table 3 for complete results). TSVD gives comparable perplexity with Recover-KL, results being slightly better on NYT and 20NewsGroup which are larger datasets with moderately high mean document lengths. We also find comparable results on Topic Coherence [11] (see Table 2 in supplementary for the complete results).

4.3 Additional Empirical Results

Topic Recovery Results of L_1 reconstruction error for Recover-L2, Recover-KL and TSVD are given in Table 3. Note that TSVD always performs better than Recover-L2. Recover-KL is slightly better than TSVD on 20NewsGroup dataset, but for larger sample size result of TSVD improves. Figure 2 shows that on most datasets, the performance of TSVD on majority of the topics (> 90%) is much better than Recover-KL (results with Recover-L2 are similar). Averaged across all these datasets and the document sizes, TSVD gives average improvement of 27% over Recover-KL.

Perplexity and Topic Coherence Table 5 gives the perplexity results and Table 4 gives the Topic Coherence results. TSVD-100 refers to the TSVD algorithm using all 100% documents of a topic (in step 6 of TSVD). The proposed TSVD algorithm is comparable with Recover-KL on NIPS and Pubmed, and gives better perplexity on NYT and 20NewsGroup. Topic coherence is slightly better on NIPS, NYT and Pubmed, and slightly worse on 20NG, but note that the difference is within range of variability across topics.

| Corpus | Documents | Recover-L2 | Recover-KL | TSVD | % Improvement |
|--------|-----------|------------|--------------|--------------|---------------|
| NIPS | 40,000 | 0.342 | 0.308 | 0.094 | 69.6% |
| | 50,000 | 0.346 | 0.308 | 0.135 | 56.3% |
| | 60,000 | 0.346 | 0.311 | 0.124 | 60.2% |
| Pubmed | 40,000 | 0.388 | 0.332 | 0.291 | 12.5% |
| | 50,000 | 0.378 | 0.326 | 0.290 | 11.2% |
| | 60,000 | 0.372 | 0.328 | 0.297 | 9.6% |
| 20NG | 40,000 | 0.126 | 0.120 | 0.125 | -4.1% |
| | 50,000 | 0.118 | 0.114 | 0.115 | -0.7% |
| | 60,000 | 0.114 | 0.110 | 0.108 | 2.5% |
| NYT | 40,000 | 0.214 | 0.208 | 0.198 | 4.8% |
| | 50,000 | 0.211 | 0.206 | 0.186 | 9.4% |
| | 60,000 | 0.205 | 0.200 | 0.192 | 4.1% |

Table 3: L1 reconstruction error on various semi-synthetic datasets. Last column is percent improvement over Recover-KL (best performing Recover algorithm).




Figure 2: Histogram of L1 error across topics for 40,000 synthetic documents. *TSVD* (blue, solid border) gets better recovery on most topics (> 90%) for most datasets (leaving small number of outliers) than *Recover-KL* (green, dashed border).

| Corpus | Recover-KL | Recover-L2 | TSVD | TSVD-100 |
|--------|---------------|---------------------|---------------|---------------------|
| NIPS | -86.4 ± 24.5 | -88.6 ± 22.7 | -63.2 ± 36.0 | -54.6 ± 36.4 |
| NYT | -105.2 ± 25.0 | -102.1 ± 28.2 | -106.8 ± 23.0 | -98.1 ± 34.5 |
| Pubmed | -94.0 ± 22.5 | -94.4 ± 22.5 | -86.4 ± 28.1 | -78.5 ± 31.4 |
| 20NG | -93.7 ± 13.6 | -89.4 ± 20.7 | -95.0 ± 15.2 | -92.1 ± 24.7 |

Table 4: Average Topic Coherence (with standard deviation across topics)

Topics on Real Data Table 6 shows the top 5 words of matched pair of topics for *TSVD* and *Recover-KL*, matched with Gibbs sampling, on the real NYT dataset³. The catchwords for *TSVD* and the anchor-words for *Recover* are highlighted (if present in top 5). Note that generally most high-frequency words of topics are identified as catchwords, while for *Recover* there is a significant number of topics (38%) which do not have anchor-word in the top 5 words of the topic. In general the topics given by *TSVD* are closer to those given by Gibbs sampling (total average L_1 error with topics from Gibbs sampling for topics from *TSVD* is 0.037, whereas for *Recover-KL* it is 0.047)

³the “zzz” token is an identifier placed with named entities in the NYT dataset

| Corpus | Recover-KL | Recover-L2 | TSVD | TSVD-100 |
|--------|-------------|------------|------|-------------|
| NIPS | 754 | 749 | 841 | 830 |
| NYT | 1579 | 1685 | 1526 | 1521 |
| Pubmed | 1188 | 1203 | 1295 | 1271 |
| 20NG | 2431 | 2565 | 2400 | 2300 |

Table 5: Perplexity on test data

Table 6: Top 5 words of matched topic pairs for TSVD, Recover-KL and Gibbs sampling. Catch-words and anchor words in top 5 words are highlighted for TSVD and Recover-KL

| TSVD | Recover-KL | Gibbs |
|---|--|--|
| election political zzz_vladimir_putin government minister | zzz_clinton government zzz_pakistan zzz_india zzz_united_states | government political election zzz_vladimir_putin zzz_russia |
| film movie character play show | film show movie music book | film movie character play director |
| zzz_elian zzz_miami boy father relatives | zzz_elian boy zzz_miami father family | zzz_elian zzz_miami boy father zzz_cuba |
| patient doctor drug cancer study | patient drug doctor percent found | patient doctor drug medical cancer |
| music song album band record | black reporter zzz_new_york zzz_black show | music song album band musical |
| show network series television zzz_abc | con zzz_mexico son federal mayor | show television network series zzz_abc |
| film movie actor director movies | goal play team season game | film movie award actor zzz_oscar |
| book writer writing read reader | zzz_john_rocker player team right braves | book word writer author wrote |
| stock percent market zzz_nasdaq fund | million percent tax bond fund | stock market percent fund investor |
| officer police case prosecutor lawyer | zzz_ray lewis police case officer death | police officer official case investigation |
| zzz_russian war rebel troop military | zzz_russian zzz_russia war zzz_vladimir_putin rebel | war military zzz_russian soldier troop |
| friend family home son father | look gun game point shot | family home father son friend |
| company million percent stock analyst | million company stock percent shares | company million companies business market |
| zzz_microsoft window company government software | zzz_microsoft company computer system software | zzz_microsoft company window antitrust government |
| cup minutes tablespoon add oil | cup minutes tablespoon add oil | cup minutes add tablespoon oil |
| tax cut plan billion taxes | zzz_governor_bush tax campaign taxes plan | tax plan billion million cut |
| room home friend mother feet | room show look home house | room look water house hand |
| com question information zzz_eastern sport | com information question zzz_eastern sport | com information daily question zzz_eastern |
| zzz_internet web site online customer | web site zzz_internet company com | web site zzz_internet online sites |
| school student teacher public children | school student program million children | school student teacher program children |
| prices percent economy market zzz_fed | percent stock market economy prices | percent prices economy market oil |

Continued on next page

Table 6: Top 5 words of matched topic pairs for TSVD, Recover-KL and Gibbs sampling. Catch-words and anchor words in top 5 words are highlighted for TSVD and Recover-KL

| TSVD | Recover-KL | Gibbs |
|--|--|--|
| building room wall restaurant house | zzz_kosovo police zzz_serb war official | building town area resident million |
| computer mail web internet user | web www site cookie cookies | computer system software technology mail |
| car wheel seat model zzz_suv | car driver truck system model | car driver truck vehicle wheel |
| right hand give mean need | mate women bird film idea | women look com need telegram |
| com www information syndicate article | chocolate food wine flavor buy | www com hotel room tour |
| book find school woman sales | book find school woman women | book find woman british school |
| zzz_al_gore zzz_bill_bradley campaign president vice | zzz_al_gore zzz_bill_bradley campaign president percent | zzz_al_gore campaign zzz_bill_bradley president democratic |
| race zzz_tiger_wood win won tour | zzz_tiger_wood shot tournament tour player | zzz_tiger_wood tour tournament shot player |
| million percent company newspaper com | dog quick jump altered food | plant human food product scientist |
| team season game player play | team season player coach zzz_cowboy | team player season coach league |
| died family home book zzz_new_york | show film country right women | art artist painting museum show |
| zzz_clinton zzz_israel talk peace zzz_china | zzz_israel zzz_lebanon peace zzz_syria israeli | zzz_israel peace palestinian talk israeli |
| zzz_mccain zzz_bush voter primary campaign | zzz_john_mccain zzz_george_bush campaign republican voter | zzz_john_mccain zzz_george_bush campaign zzz_bush zzz_mccain |
| worker percent million job zzz_u_s | official zzz_iraq government zzz_united_states oil | zzz_mexico drug government zzz_united_states mexican |
| point game shot lead half | point game team season zzz_laker | game point team play season |
| article heart art zzz_discover magazine | air wind snow shower weather | water snow weather air scientist |
| run inning hit game season | run season game inning hit | run season game hit inning |
| yard zzz_ram play game titan | game team season play zzz_ram | team season game coach zzz_nfl |
| zzz_american zzz_china policy zzz_u_s democracy | zzz_china zzz_taiwan government trade zzz_party | zzz_china zzz_united_states zzz_u_s zzz_clinton zzz_american |
| campaign candidates zzz_george_bush presidential zzz_al_gore | zzz_bradley zzz_al_gore campaign zzz_gore zzz_clinton | zzz_clinton president gay mayor zzz_rudolph_giuliani |
| group official law million plan | zzz_bush zzz_mccain campaign republican voter | gun bill law zzz_congress legislation |
| goal game games team season | team game point season player | race won win fight team |
| player team patriot season zzz_bill_belichick | newspaper zzz_chronicle zzz_examiner zzz_hearst million | million money worker company pay |
| coach team player season jet | test women study student found | plane flight passenger pilot zzz_boeing |

Continued on next page

Table 6: Top 5 words of matched topic pairs for TSVD, Recover-KL and Gibbs sampling. Catch-words and anchor words in top 5 words are highlighted for TSVD and Recover-KL

| TSVD | Recover-KL | Gibbs |
|--|--|---|
| season spring team manager player | pope church book jewish religious | religious church jewish jew zzz_god |
| daily american statesman austin kid | company companies deal zzz_internet zzz_time_warner | media zzz_time_warner television newspaper cable |
| zzz_earl_caldwell zzz_black reporter black zzz_harlem | corp group list oil meeting | black white zzz_black hispanic reporter |
| zzz_john_mccain zzz_george_bush campaign zzz_south_carolina republican | flag black zzz_confederate right group | flag zzz_confederate zzz_south_carolina black zzz_south |
| minutes hour starring movie violence | official government case officer security | court law case lawyer right |

Summary: We evaluated the proposed algorithm, **TSVD**, rigorously on multiple datasets with respect to the state of the art (Recover), following the evaluation methodology of [5]. In Table 2 we show that the L1 reconstruction error for the new algorithm is small and on average **20%** better than the best results of the Recover algorithms [5]. We also demonstrate that on real datasets the algorithm achieves comparable perplexity and topic coherence to Recover. Moreover, we show on multiple real datasets that the algorithm assumptions are well justified in practice.

Conclusion

Real world corpora often exhibits the property that in every document there is one topic dominantly present. A standard SVD based procedure will not be able to detect these topics, however TSVD, a thresholded SVD based procedure, as suggested in this paper, discovers these topics. While SVD is time-consuming, there have been a host of recent sampling-based approaches which make SVD easier to apply to massive corpora which may be distributed among many servers. We believe that apart from topic recovery, thresholded SVD can be applied even more broadly to similar problems, such as matrix factorization, and will be the basis for future research.

References

- [1] Blei, D. Introduction to probabilistic topic models. *Communications of the ACM*, pp. 77–84, 2012.
- [2] Blei, D., Ng, A., and Jordan, M. Latent Dirichlet allocation. *Journal of Machine Learning Research*, pp. 3:993–1022, 2003. Preliminary version in *Neural Information Processing Systems* 2001.
- [3] Griffiths, T. L. and Steyvers, M. Finding scientific topics. *Proceedings of the National Academy of Sciences*, 101:5228–5235, 2004.
- [4] Arora, S., Ge, R., and Moitra, A. Learning topic models – going beyond SVD. In *Foundations of Computer Science*, 2012b.
- [5] Arora, S., Ge, R., Halpern, Y., Mimno, D., Moitra, A., Sontag, D., Wu, Y., and Zhu M. A practical algorithm for topic modeling with provable guarantees. In *International Conference on Machine Learning*, 2013
- [6] Anandkumar, A., Foster, D., Hsu, D., Kakade, S., and Liu, Y. A Spectral Algorithm for Latent Dirichlet Allocation In *Neural Information Processing Systems*, 2012.
- [7] Papadimitriou, C., Raghavan, P., Tamaki H., and Vempala S. Latent semantic indexing: a probabilistic analysis. *Journal of Computer and System Sciences*, pp. 217–235, 2000. Preliminary version in PODS 1998.

- [8] Deerwester, S., Dumais, S., Landauer, T., Furnas, G., and Harshman, R. Indexing by latent semantic analysis. *Journal of the American Society for Information Science*, pp. 391–407, 1990.
- [9] Kumar, A., and Kannan, R. Clustering with spectral norm and the k-means algorithm. In *Foundations of Computer Science*, 2010
- [10] Awashti, P., and Sheffet, O. Improved spectral-norm bounds for clustering. In *Proceedings of Approx/Random*, 2012.
- [11] Mimno, D., Wallach, H., Talley, E., Leenders, M. and McCallum, A. Optimizing semantic coherence in topic models. In *Empirical Methods in Natural Language Processing*, pp. 262–272, 2011.
- [12] Lloyd, Stuart P. Least squares quantization in PCM, *IEEE Transactions on Information Theory* 28 (2): 129137,1982.
- [13] Arthur, D., and Vassilvitskii, S. K-means++: The advantages of careful seeding. In *Proceedings of ACM-SIAM symposium on Discrete algorithms*, pp. 1027–1035, 2007
- [14] McDiarmid, C. On the method of Bounded Differences. *Surveys in Combinatorics: London Math. Soc. Lecture Note Series 141*. Cambridge University Press., 1989.
- [15] Vershynin R. Introduction to non-asymptotic analysis of random matrices. In *ArXiv:1011.3027v6 [math.PR]* 4 Oct 2011

5 Appendix

6 Line of Proof

We describe the Lemmas we prove to establish the result. The detailed proofs are in the Section 7.

6.1 General Facts

We start with a consequence of the no-local-minimum assumption. We use that assumption solely through this Lemma.

Lemma 6.1 *Let $p_i(\zeta, l)$ be as in (6). If for some $\zeta_0 \in \{0, 1, \dots, m\}$ and $\nu \geq 0$, $\sum_{\zeta \geq \zeta_0} p_i(\zeta, l) \geq \nu$ and also $\sum_{\zeta \leq \zeta_0} p_i(\zeta, l) \geq \nu$ then, $p_i(\zeta_0, l) \geq \frac{\nu}{m}$.*

Next, we state a technical Lemma which is used repeatedly. It states that for every i, ζ, l , the empirical probability that $A_{ij} = \zeta/m$ is close to the true probability. Unsurprisingly, we prove it using H-C. But we will state a consequence in the form we need in the sequel.

Lemma 6.2 *Let $p_i(\zeta, l)$ and $q_i(\zeta, l)$ be as in (6) and (7). We have*

$$\forall i, l, \zeta : \text{Prob} \left(|p_i(\zeta, l) - q_i(\zeta, l)| \geq \frac{\varepsilon}{2} \sqrt{w_0} \sqrt{p_i(\zeta, l)} + \frac{\varepsilon^2 w_0}{2} \right) \leq 2 \exp(-\varepsilon^2 s w_0 / 8).$$

From this it follows that with probability at least $1 - 2 \exp(-\varepsilon^2 w_0 s / 8)$,

$$\frac{1}{2} q_i(\zeta, l) - \varepsilon^2 w_0 \leq p_i(\zeta, l) \leq 2 q_i(\zeta, l) + 2 \varepsilon^2 w_0.$$

6.1.1 Properties of Thresholding

Say that a threshold ζ_i “splits” $T_l^{(2)}$ if $T_l^{(2)}$ has a significant number of j with $A_{ij} > \zeta_i/m$ and also a significant number of j with $A_{ij} \leq \zeta_i/m$. Intuitively, it would be desirable if no threshold splits any T_l , so that, in \mathbf{B} , for each i, l , either most $j \in T_l^{(2)}$ have $B_{ij} = 0$ or most $j \in T_l^{(2)}$ have $B_{ij} = \sqrt{\zeta_i}$. We now prove that this is indeed the case with proper bounds. We henceforth refer to the conclusion of the Lemma below by the mnemonic “no threshold splits any T_l ”.

Lemma 6.3 (No Threshold Splits any T_l) For a fixed i, l , with probability at least $1 - 2 \exp(-\varepsilon^2 w_0 s / 8)$, the following holds:

$$\text{Min} \left(\text{Prob}(A_{ij}^{(2)} \leq \frac{\zeta_i}{m}; j \in T_l^{(2)}), \text{Prob}(A_{ij}^{(2)} > \frac{\zeta_i}{m}; j \in T_l^{(2)}) \right) \leq 4m\varepsilon w_0.$$

Let μ be a $d \times s$ matrix whose columns are given by

$$\forall j \in T_l^{(2)}, \mu_{\cdot, j} = E(B_{\cdot, j} | j \in T_l).$$

μ 's columns corresponding to all $j \in T_l$ are the same. The entries of the matrix μ are fixed (real numbers) once we have $\mathbf{A}^{(1)}$ (and the thresholds ζ_i are determined). Note: We have “integrated out W ”, i.e.,

$$\mu_{ij} = \int_{W_{\cdot, j}} \text{Prob}(W_{\cdot, j} | j \in T_l) E(B_{ij} | W_{\cdot, j}).$$

(So, think of $W_{\cdot, j}$ for $\mathbf{A}^{(1)}$'s columns being picked first from which ζ_i is calculated. $W_{\cdot, j}$ for columns of $\mathbf{A}^{(2)}$ are not yet picked until the ζ_i are determined.) But μ_{ij} are random variables before we fix $\mathbf{A}^{(1)}$. The following Lemma is a direct consequence of “no threshold splits any T_l ”.

Lemma 6.4 Let $\zeta'_i = \text{Max}(\zeta_i, 8 \ln(20/\varepsilon w_0))$. With probability at least $1 - 4kd \exp(-\varepsilon^2 s w_0 / 8)$ (over the choice of $\mathbf{A}^{(1)}$):

$$\begin{aligned} \forall l, \forall j \in T_l, \forall i : \mu_{ij} \leq \varepsilon_l \sqrt{\zeta'_i} \text{ OR } \mu_{ij} \geq \sqrt{\zeta'_i} (1 - \varepsilon_l) \\ \forall l, \forall i, \text{Var}(B_{ij}) \leq 2\varepsilon_l \zeta'_i, \end{aligned} \quad (10)$$

where, $\varepsilon_l = 4m\varepsilon w_0 / w_l$.

So far, we have proved that for every i , the threshold does not split any T_l . But this is not sufficient in itself to be able to cluster (and hence identify the T_l), since, for example, this alone does not rule out the extreme cases that for most j in every T_l , $A_{ij}^{(2)}$ is above the threshold (whence $\mu_{ij} \geq (1 - \varepsilon_l) \sqrt{\zeta'_i}$ for almost all j) or for most j in no T_l is $A_{ij}^{(2)}$ above the threshold, whence, $\mu_{ij} \leq \varepsilon_l \sqrt{\zeta'_i}$ for almost all j . Both these extreme cases would make us loose all the information about T_l due to thresholding; this scenario and milder versions of it have to be proven not to occur. We do this by considering how thresholds handle catchwords. Indeed we will show that for a catchword $i \in S_l$, a $j \in T_l$ has $A_{ij}^{(2)}$ above the threshold and a $j \notin T_l$ has $A_{ij}^{(2)}$ below the threshold. Both statements will only hold with high probability, of course and using this, we prove that $\mu_{\cdot, j}$ and $\mu_{\cdot, j'}$ are not too close for j, j' in different T_l 's. For this, we need the following Lemmas.

Lemma 6.5 For $i \in S_l$, and $l' \neq l$, we have with $\eta_i = \lfloor M_{il}(\alpha + \beta + \rho)m/2 \rfloor$,

$$\text{Prob}(A_{ij} \leq \eta_i/m | j \in T_l) \leq \varepsilon w_0 / 20, \quad \text{Prob}(A_{ij} \geq \eta_i/m | j \in T_{l'}) \leq \varepsilon w_0 / 20.$$

Lemma 6.6 With probability at least $1 - 8mdk \exp(-\varepsilon^2 w_0 s / 8)$, we have

$$\text{for } j \in T_l, j' \notin T_l, |\mu_{\cdot, j} - \mu_{\cdot, j'}|^2 \geq \frac{2}{9} \alpha p_0 m.$$

6.1.2 sec:Proximity

Next, we wish to show that clustering as in TSVD identifies the dominant topics correctly for most documents, i.e., that $R_l \approx T_l$ for all l . For this, we will use a theorem from [9] [see also [10]] which in this context says:

Theorem 6.7 If all but a f fraction of the the $B_{\cdot, j}$ satisfy the “proximity condition”, then TSVD identifies the dominant topic in all but $c_1 f$ fraction of the documents correctly after polynomial number of iterations.

To describe the proximity condition, first let σ be the maximum over all directions v of the square root of the mean-squared distance of $B_{\cdot, j}$ to $\mu_{\cdot, j}$, i.e.,

$$\sigma^2 = \text{Max}_{\|v\|=1} \frac{1}{s} |v^T (\mathbf{B} - \mu)|^2 = \frac{1}{s} \|\mathbf{B} - \mu\|^2.$$

The parameter σ should remind the reader of standard deviation, which is indeed what this is, since $E(\mathbf{B}|T_1, T_2, \dots, T_l) = \mu$. Our random variables $B_{\cdot, j}$ being d -dimensional vectors, we take the maximum standard deviation in any direction.

Definition: \mathbf{B} is said to satisfy the proximity condition with respect to μ , if for each l and each $j \in T_l$ and and each $l' \neq l$ and $j' \in T_{l'}$, the projection of $B_{\cdot, j}$ onto the line joining $\mu_{\cdot, j}$ and $\mu_{\cdot, j'}$ is closer to $\mu_{\cdot, j}$ by at least

$$\Delta = \frac{c_0 k}{\sqrt{w_0}} \sigma,$$

than it is to $\mu_{\cdot, j'}$. [Here, c_0 is a constant.]

To prove proximity, we need to bound σ . This will be the task of the subsection 7.1 which relies heavily on Random Matrix Theory.

7 Proofs of Correctness

We start by recalling the Höfdding-Chernoff (H-C) inequality in the form we use it.

Lemma 7.1 Höfdding-Chernoff *If X is the average of r independent random variables with values in $[0, 1]$ and $E(X) = \mu$, then, for an $t > 0$,*

$$\text{Prob}(X \geq \mu + t) \leq \exp\left(-\frac{t^2 r}{2(\mu + t)}\right); \text{Prob}(X \leq \mu - t) \leq \exp\left(-\frac{t^2 r}{2\mu}\right).$$

Proof: (of Lemma 6.1) Abbreviate $p_i(\cdot, l)$ by $f(\cdot)$. We claim that either (i) $f(\zeta) \geq f(\zeta - 1) \forall 1 \leq \zeta \leq \zeta_0$ or (ii) $f(\zeta + 1) \leq f(\zeta) \forall m - 1 \geq \zeta \geq \zeta_0$. To see this, note that if both (i) and (ii) fail, we have $\zeta_1 \leq \zeta_0$ and $\zeta_2 \geq \zeta_0$ with $f(\zeta_1) - f(\zeta_1 - 1) < 0 < f(\zeta_2 + 1) - f(\zeta_2)$. But then there has to be a local minimum of f between ζ_1 and ζ_2 . If (i) holds, clearly, $f(\zeta_0) \geq f(\zeta) \forall \zeta < \zeta_0$ and so the lemma follows. So, also if (ii) holds.

Proof: (of Lemma 6.2) Note that $q_i(\zeta, l) = \frac{1}{s} |\{j \in T_l : A_{ij} = \zeta/m\}| = \frac{1}{s} \sum_{j=1}^s X_j$, where, X_j is the indicator variable of $A_{ij} = \zeta/m \wedge j \in T_l$. $\frac{1}{s} \sum_j E(X_j) = p_i(\zeta, l)$ and we apply H-C with $t = \frac{1}{2}\varepsilon\sqrt{w_0}\sqrt{p_i(\zeta, l)} + \frac{1}{2}\varepsilon^2 w_0$ and $\mu = p_i(\zeta, l)$. We have $\frac{t^2}{\mu+t} \geq \varepsilon^2 w_0/4$, as is easily seen by calculating the roots of the quadratic $t^2 - \frac{1}{4}t\varepsilon^2 w_0 - \frac{1}{4}\varepsilon^2 w_0 \mu = 0$. Thus we get the claimed for T_l . Note that the same proof applies for $T_l^{(1)}$ as well as $T_l^{(2)}$.

To prove the second assertion, let $a = q_i(\zeta, l)$ and $b = \sqrt{p_i(\zeta, l)}$, then, b satisfies the quadratic inequalities:

$$b^2 - \frac{1}{2}\varepsilon\sqrt{w_0}b - (a + \frac{1}{2}\varepsilon^2 w_0) \leq 0; b^2 + \frac{1}{2}\varepsilon\sqrt{w_0}b - (a - \frac{1}{2}\varepsilon^2 w_0) \geq 0.$$

By bounding the roots of these quadratics, it is easy to see the second assertion after some calculation.

Proof: (of Lemma 6.3) Note that ζ_i is a random variable which depends only on $A^{(1)}$. So, for $j \in T_l^{(2)}$, A_{ij} are independent of ζ_i . Now, if

$$\text{Prob}(A_{ij} \leq \frac{\zeta_i}{m}; j \in T_l^{(2)}) > 4m\varepsilon w_0 \text{ and } \text{Prob}(A_{ij} > \frac{\zeta_i}{m}; j \in T_l^{(2)}) > 4m\varepsilon w_0,$$

by Lemma (6.1), we have

$$\text{Prob}(A_{ij} = \frac{\zeta_i}{m}; j \in T_l^{(2)}) > 4\varepsilon w_0.$$

Since $\text{Prob}(A_{ij} = \zeta/m; j \in T_l^{(1)}) = \text{Prob}(A_{ij} = \zeta/m; j \in T_l^{(2)})$ for all ζ , we also have

$$\text{Prob}(A_{ij} = \frac{\zeta_i}{m}; j \in T_l^{(1)}) = p_i(\zeta_i, l) > 4\varepsilon w_0. \quad (11)$$

Pay a failure probability of $2 \exp(-\varepsilon^2 s w_0 / 8)$ and assume the conclusion of Lemma (6.2) and we have:

$$\frac{1}{s} \left| \{j \in T_l^{(1)} : A_{ij} = \frac{\zeta_i}{m}\} \right| = q_i(\zeta_i, l) \geq p_i(\zeta_i, l) - \frac{\varepsilon}{2} \sqrt{w_0 p_i(\zeta_i, l)} - \frac{\varepsilon^2}{2} w_0.$$

Now, it is easy to see that $p_i(\zeta, l) - \frac{\varepsilon}{2} \sqrt{w_0 p_i(\zeta, l)}$ increases as $p_i(\zeta, l)$ increases subject to (11). So,

$$p_i(\zeta, l) - \frac{\varepsilon}{2} \sqrt{w_0 p_i(\zeta, l)} - \frac{\varepsilon^2}{2} w_0 > (4\varepsilon - \varepsilon^{3/2} - \frac{1}{2}\varepsilon^2)w_0 \geq 3\varepsilon w_0,$$

contradicting the definition of ζ_i in the algorithm. This completes the proof of the Lemma.

Proof: (of Lemma 6.4): After paying a failure probability of $4kd \exp(-\varepsilon^2 s w_0 / 8)$, assume no threshold splits any T_l . [The factors of k and d come in because we are taking the union bound over all words and all topics.] Then,

$$\text{Prob}(A_{ij}^{(2)} \leq \frac{\zeta_i}{m} \mid j \in T_l^{(2)}) = \sum_{\zeta=0}^{\zeta_i} p_i(\zeta, l) / \text{Prob}(j \in T_l) \leq 4m\varepsilon \frac{w_0}{w_l}$$

$$\text{or Prob}(A_{ij}^{(2)} > \frac{\zeta_i}{m} \mid j \in T_l^{(2)}) = \sum_{\zeta=\zeta_i+1}^m p_i(\zeta, l) / \text{Prob}(j \in T_l) \leq 4m\varepsilon \frac{w_0}{w_l}.$$

Wlg, assume that $\text{Prob}(A_{ij} \leq \zeta_i/m \mid j \in T_l) \leq \varepsilon_l$. Then, with probability, at least $1 - \varepsilon_l$, $A_{ij}^{(2)} > \zeta_i/m$. Now, either $\zeta_i < 8 \ln(20/\varepsilon w_0)$ and all $B_{ij}, j \in T_l$ are zero and then $\mu_{ij} = 0$, or $\zeta_i \geq 8 \ln(20/\varepsilon w_0)$, whence, $E(B_{ij} \mid j \in T_l) \in [(1 - \varepsilon_l)\sqrt{\zeta_i'}, \sqrt{\zeta_i'}]$. So, $\mu_{ij} \geq (1 - \varepsilon_l)\sqrt{\zeta_i'}$ and $\text{Prob}(B_{ij} = 0) \leq \varepsilon_l$. So,

$$\text{Var}(B_{ij}^2 \mid j \in T_l) \leq (\sqrt{\zeta_i'} - (1 - \varepsilon_l)\sqrt{\zeta_i'})^2 \text{Prob}(B_{ij} = \sqrt{\zeta_i'} \mid j \in T_l) + (\sqrt{\zeta_i'} - 0)^2 \text{Prob}(B_{ij} = 0 \mid j \in T_l) \leq 2\varepsilon_l \zeta_i'.$$

This proves the lemma in this case. The other case is symmetric.

Proof: (of Lemma 6.5) Recall that $P_{ij} = \sum_l M_{il} W_{lj}$ is the probability of word i in document j conditioned on \mathbf{W} . Fix an $i \in S_l$. From the dominant topic assumption,

$$\forall j \in T_l, P_{ij} = \sum_{l_1} M_{il_1} W_{l_1, j} \geq M_{il} W_{lj} \geq M_{il} \alpha. \quad (12)$$

The P_{ij} are themselves random variables. Note that (12) holds with probability 1. From Catchword assumption and (1), we get that

$$M_{il} \alpha - (\eta_i/m) \geq M_{il} \alpha - M_{il}((\alpha + \beta + \rho)/2) \geq M_{il} \alpha \delta / 2.$$

Now, we will apply H-C with $\mu - t = \eta_i/m$ and $\mu \geq M_{il} \alpha$ for the m independent words in a document. By Calculus, the probability bound from H-C of $\exp(-t^2 w_l s / 2\mu) = \exp(-(\mu - (\eta_i/m))/2\mu)$ is highest subject to the constraints $\mu \geq M_{il} \alpha; \eta_i \leq m M_{il}(\alpha + \beta + \rho)/2$, when $\mu = M_{il} \alpha$ and $\eta_i = m M_{il}(\alpha + \beta + \rho)/2$, whence, we get

$$\text{Prob}(A_{ij} \leq \eta_i/m \mid j \in T_l) \leq \exp(-M_{il} \alpha \delta^2 m / 8) \leq \varepsilon w_0 / 20,$$

using (5). Now, we prove the second assertion of the Lemma.

$$\begin{aligned} \forall j \in T_{l'}, l' \neq l, \sum_{l_1} M_{il_1} W_{l_1, j} &= M_{il} W_{lj} + \sum_{l_1 \neq l} M_{il_1} W_{l_1, j} \\ &\leq M_{il} W_{lj} + (\text{Max}_{l_1 \neq l} M_{il_1}) (1 - W_{lj}) \\ &\leq M_{il}(\beta + \rho). \end{aligned} \quad (13)$$

$$\frac{\eta_i}{m} - M_{il}(\beta + \rho) \geq \frac{M_{il}(\alpha + \beta + \rho)}{2} - M_{il}(\beta + \rho) - \frac{1}{m} \geq \frac{3M_{il} \alpha \delta}{8},$$

using (5) and (1). Applying the first inequality of Lemma (7.1) with $\mu + t = \eta_i/m$ and $\mu \leq M_{il}(\beta + \rho)$ and again using (5),

$$\text{Prob}(A_{ij} \geq \eta_i/m \mid j \in T_{l'}) \leq \exp(-9M_{il} \alpha \delta^2 m / 64) \leq \varepsilon w_0 / 20.$$

Lemma 7.2 For $i \in S_l$, $\text{Prob}(\zeta_i < \eta_i) \leq 3mke^{-\varepsilon^2 sw_0/8}$, with η_i as defined in Lemma 6.5.

Proof: Fix attention on $i \in S_l$. After paying the failure probability of $3mke^{-\varepsilon^2 sw_0/8}$, assume the conclusions of Lemma (6.2) hold for all ζ, l . It suffices to show that

$$\left| \{j : A_{ij}^{(1)} > \eta_i/m\} \right| \geq \frac{w_0 s}{2}, \quad \left| \{j : A_{ij}^{(1)} = \frac{\eta_i}{m}\} \right| < 3w_0 \varepsilon s,$$

since, η_i is an integer and ζ_i is the largest integer satisfying the inequalities. The first statement follows from first assertion of Lemma 6.5. The second statement is slightly more complicated. Using both the first and second assertions of Lemma 6.5, we get that for all l' (including $l' = l$), we have

$$\text{Prob}(A_{ij} = \eta_i/m | j \in T_{l'}^{(1)}) \leq \varepsilon w_0/20.$$

$$\begin{aligned} \left| \{j \in T_{l'}^{(1)} : A_{ij} = \eta_i/m\} \right| &\leq \varepsilon w_0 w_{l'} s/20 + \frac{\varepsilon}{2} \sqrt{w_0/w_{l'}} \sqrt{\varepsilon w_0/20} w_{l'} s + \frac{\varepsilon^2 w_0 w_{l'}}{2} \\ &\leq \frac{\varepsilon w_0 s}{8} (w_{l'} + \sqrt{\varepsilon w_{l'}}) + \frac{\varepsilon^2 w_0 s}{2}. \end{aligned}$$

Now, adding over all l' and using $\sum_{l'} \sqrt{w_{l'}} \leq \sqrt{k} \sqrt{\sum_{l'} w_{l'}} = \sqrt{k}$, we get

$$\left| \{j : A_{ij}^{(1)} = \eta_i/m\} \right| \leq \varepsilon w_0 s,$$

since $\varepsilon \leq 1/k$.

Lemma 7.3 Define $I_l = \{i \in S_l : \zeta_i \geq \eta_i\}$. With probability at least $1 - 8mdk \exp(-\varepsilon^2 w_0 s/8)$, we have for all l ,

$$\sum_{i \in I_l} \zeta'_i \geq m \alpha p_0/2.$$

Proof: After paying the failure probability, we assume the conclusion of Lemma 6.2 holds for all i, ζ, l . Now, by Lemma 7.2, we have (with $\mathbf{1}$ denoting the indicator function)

$$E \left(\sum_{i \in S_l} M_{il} \mathbf{1}(\zeta_i < \eta_i) \right) \leq 3mk \exp(-\varepsilon^2 sw_0/8) \sum_{i \in S_l} M_{il},$$

which using Markov inequality implies that with probability at least $1 - 6mk \exp(-\varepsilon^2 sw_0/8)$,

$$\sum_{i \in I_l} M_{il} \geq \frac{1}{2} \sum_{i \in S_l} M_{il} \geq p_0/2, \quad (14)$$

using (4). Note that by (5), no catchword has ζ'_i set to zero. So,

$$\sum_{i \in I_l} \zeta'_i = \sum_{i \in I_l} \zeta_i \geq \sum_{i \in I_l} \eta_i \geq \sum_{I_l} m M_{il} \alpha/2 \geq \alpha p_0 m/2.$$

Proof: (of Lemma 6.6) For this proof, i will denote an element of I_l . By Lemma 6.5,

$$\forall i \in I_l, l' \neq l, \text{Prob}(A_{ij} > \frac{\zeta_i}{m} | j \in T_{l'}^{(1)}) \leq \text{Prob}(A_{ij} > \eta_i/m | j \in T_{l'}^{(1)}) \leq \frac{\varepsilon w_0}{20}. \quad (15)$$

This implies by Lemma 6.2, for $l' \neq l$,

$$\left| \{j \in T_{l'}^{(1)} : A_{ij} > \frac{\zeta_i}{m}\} \right| \leq w_{l'} s \left(\frac{\varepsilon w_0}{20} + \varepsilon \sqrt{w_0/w_{l'}} \sqrt{\varepsilon w_0/4} \right) + w_0 \varepsilon^2 s/2. \quad (16)$$

Summing over all $l' \neq l$, we get (using $\sum_{l'} \sqrt{w_{l'}} \leq \sqrt{\sum_{l'} w_{l'}} \sqrt{k} \leq 1/\sqrt{\varepsilon}$ by (9))

$$\sum_{l' \neq l} \left| \{j \in T_{l'}^{(1)} : A_{ij} > \frac{\zeta_i}{m}\} \right| \leq \varepsilon w_0 s.$$

Now the definition of ζ_i in the algorithm implies that:

$$\sum_{\zeta > \zeta_i} q_i(\zeta, l) = \left| \{j \in T_l^{(1)} : A_{ij} > \frac{\zeta_i}{m}\} \right| \geq \left(\frac{w_0}{2} - \varepsilon w_0 \right) s \geq w_0 s / 4.$$

So, by Lemma 6.2,

$$\begin{aligned} \text{Prob}(j \in T_l; A_{ij} > \zeta_i / m) &= \sum_{\zeta > \zeta_i} p_i(\zeta, l) \geq \frac{1}{2} \sum_{\zeta > \zeta_i} q_i(\zeta, l) - \varepsilon^2 w_0 m \\ &\geq \frac{w_0}{8} - \varepsilon^2 w_0 m \geq w_0 / 9, \end{aligned}$$

using (9). Next let $\tilde{p} = \text{Prob}(A_{ij} = \zeta_i / m; j \in T_l)$. Since $|\{j \in T_l^{(1)} : A_{ij} = \zeta_i / m\}| \leq 3\varepsilon w_0 s$, by the definition of ζ_i in the algorithm, we get by a similar argument

$$\tilde{p} \leq 2q_i(\zeta_i, l) + 2\varepsilon^2 w_0 \leq 7\varepsilon w_0. \quad (17)$$

Now, by Lemma 6.1, we have

$$\tilde{p} \geq \text{Min} \left(\frac{w_0}{9m}, \frac{1}{m} \text{Prob}(A_{ij} \leq \zeta_i / m; j \in T_l^{(2)}) \right).$$

By (9), $7\varepsilon w_0 < w_0 / 9m$ and so we get:

$$\text{Prob}(A_{ij} \leq \zeta_i / m; j \in T_l^{(2)}) \leq 7\varepsilon m w_0.$$

Noting that by (5), no catchword has ζ'_i set to zero, $\text{Prob}(B_{ij} = 0 | j \in T_l^{(2)}) \leq 7\varepsilon m w_0 / w_l \leq 1/6$, by (9). This implies

$$\mu_{ij} \geq \frac{5}{6} \sqrt{\zeta'_i}.$$

Now, by (15), we have for $j' \notin T_l$,

$$\mu_{ij'} \leq \sqrt{\zeta'_i} / 6.$$

So, we have

$$|\mu_{.,j} - \mu_{.,j'}|^2 \geq \sum_{i \in I_l} (\mu_{ij} - \mu_{ij'})^2 \geq (4/9) \sum_{i \in I_l} \zeta'_i.$$

Now Lemma (7.3) implies the current Lemma.

7.1 Bounding the Spectral norm

Theorem 7.4 Fix an l . For $j \in T_l$, let $R_{.,j} = B_{.,j} - \mu_{.,j}$. [The $R_{.,j}, j \in T_l$ are vector-valued random variables which are independent, even conditioned on the partition T_1, T_2, \dots, T_k .] With probability at least $1 - 10mdk \exp(-\varepsilon^2 w_0 s / 8)$, we have $\|R\|^2 \leq ckw_0 \varepsilon sm^2$. Thus,

$$\|B - \mu\|^2 \leq c\varepsilon w_0 sm^2 k^2.$$

We will apply Random Matrix Theory, in particular the following theorem, to prove Theorem 7.4.

Theorem 7.5 [15, Theorem 5.44] Suppose R is a $d \times r$ matrix with columns $R_{.,j}$ which are independent identical vector-valued random variables. Let $U = E(R_{.,j} R_{.,j}^T)$ be the inertial matrix of $R_{.,j}$. Suppose $|R_{.,j}| \leq \nu$ always. Then, for any $t > 0$, with probability at least $1 - de^{-ct^2}$, we have⁴

$$\|R\| \leq \|U\|^{1/2} \sqrt{r} + t\nu.$$

We need the following Lemma first.

Lemma 7.6 With probability at least $1 - \exp(-s\varepsilon w_0 / 3)$, we have

$$\zeta_0 \leq 4m\lambda \quad ; \quad \sum_i \zeta'_i \leq 4km \quad (18)$$

⁴ $\|R\|$ denotes the spectral norm of R .

Proof: The probability of word i in document j , is given by: $P_{ij} = \sum_l M_{il} W_{lj} \leq \lambda_i$ (where, $\lambda_i = \max_l M_{il}$). If $\lambda_i < \frac{1}{m} \ln(20/\varepsilon w_0)$, then, $\text{Prob}(A_{ij} > (8/m) \ln(20/\varepsilon w_0)) \leq \varepsilon w_0$ by H-C (since A_{ij} is the average of m i.i.d. trials). Let X_j be the indicator function of $A_{ij} > (8/m) \ln(20/\varepsilon w_0)$. X_j are independent and so using H-C, we see that with probability at least $1 - \exp(-\varepsilon w_0 s/3)$, less than $w_0 s/2$ of the A_{ij} are greater $(8/m) \ln(20/\varepsilon w_0)$, whence, $\zeta'_i = 0$. So we have (using the union bound over all words):

$$\text{Prob} \left(\sum_{i: \lambda_i < (1/m) \ln(20/\varepsilon w_0)} \zeta'_i > 0 \right) \leq d \exp(-\varepsilon w_0 s/3).$$

If $\lambda_i \geq (1/m) \ln(20/\varepsilon w_0)$, then

$$\text{Prob}(A_{ij} > 4\lambda_i) \leq e^{-\lambda_i m} \leq \varepsilon w_0/2,$$

which implies by the same X_j kind of argument that with probability at least $1 - \exp(-\varepsilon w_0 s/4)$, for a fixed i , $\zeta_i \leq 4\lambda_i m$. Using the union bound over all words and adding all i , we get that with probability at least $1 - 2d \exp(-\varepsilon w_0 s/4)$,

$$\sum_i \zeta'_i \leq 4m \sum_i \lambda_i \leq 4m \sum_{i,l} M_{il} \leq 4km.$$

Now we prove the bound on ζ_0 . For each fixed i, j , we have $\text{Prob}(A_{ij} \geq 4\lambda) \leq e^{-m\lambda} \leq \varepsilon w_0$. Now, let Y_j be the indicator variable of $A_{ij} \geq 4\lambda$. The $Y_j, j = 1, 2, \dots, s$ are independent (for each fixed i). So, $\text{Prob}(\zeta_i \geq 4m\lambda) \leq \text{Prob}(\sum_j Y_j \geq w_0 s/2) \leq e^{-\varepsilon w_0 s/3}$. Using an union bound over all words, we get that $\text{Prob}(\zeta_0 > 4m\lambda) \leq d e^{-\varepsilon w_0/3}$ by H-C.

Proof: (of Theorem 7.4) First,

$$\|U\| = \text{Max}_{|v|=1} E(v^T R_{\cdot,j})^2 \leq E(|R_{\cdot,j}|^2) \leq 2\varepsilon_l \sum_i \zeta'_i \leq 8\varepsilon_l km,$$

by Lemma (7.6) and Lemma (6.4). We can also take $\nu = 2\sqrt{km}$ in Theorem 7.5 and with $t = \sqrt{\varepsilon m w_0 s}$, the first statement of the current theorem follows (noting $r = w_l s$). The second statement follows by just paying a factor of k for the k topics.

7.2 Proving Proximity

From Theorem (7.4), the σ in definition 6.1.2 is $\sqrt{c\varepsilon w_0 m^2 k^2}$. So, the Δ in definition 6.1.2 is $cc_0 \sqrt{\varepsilon} k^2 m$. So it suffices to prove:

Lemma 7.7 For $j \in T_l$ and $j' \in T_{l'}, l' \neq l$, let $\hat{B}_{\cdot,j}$ be the projection of $B_{\cdot,j}$ onto the line joining $\mu_{\cdot,j}$ and $\mu_{\cdot,j'}$. The probability that $|\hat{B}_{\cdot,j} - \mu_{\cdot,j'}| \leq |\hat{B}_{\cdot,j} - \mu_{\cdot,j}| + cc_0 k^2 \sqrt{\varepsilon} m$ is at most $c\varepsilon m w_0 \sqrt{k} / \sqrt{\alpha p_0}$. Hence, with probability at least $1 - cmdk \exp(-cw_0 \varepsilon^2 s)$, the number of j for which $B_{\cdot,j}$ does not satisfy the proximity condition is at most $c\varepsilon_0 w_0 \delta s / 10c_1$.

Proof: After paying the failure probability of $cmdk \exp(-w_0 \varepsilon^2/8)$, of Lemmas (7.6) and (6.6), assume that $\zeta_0 \leq 4m\lambda$, $|\mu_{\cdot,j} - \mu_{\cdot,j'}|^2 \geq 2\alpha m p_0/9$ and $\sum_i \zeta'_i \leq 4km$.

Let $X = (B_{\cdot,j} - \mu_{\cdot,j}) \cdot (\mu_{\cdot,j'} - \mu_{\cdot,j})$. X is a random variable, whose expectation is 0 conditioned on $j \in T_l^{(2)}$.

Since $\text{Prob}(B_{ij} = \sqrt{\zeta'_i} | j \in T_l) = \mu_{ij} / \sqrt{\zeta'_i}$, we have:

$$\begin{aligned}
E|X| &\leq E \sum_i |B_{ij} - \mu_{ij}| |\mu_{ij'} - \mu_{ij}| \\
&= \sum_i \left[(\sqrt{\zeta'_i} - \mu_{ij}) \frac{\mu_{ij}}{\sqrt{\zeta'_i}} + (1 - \frac{\mu_{ij}}{\sqrt{\zeta'_i}}) \mu_{ij} \right] |\mu_{ij} - \mu_{ij'}| \\
&\leq 2\varepsilon_l \sum_i \sqrt{\zeta'_i} |\mu_{ij} - \mu_{ij'}| \quad \text{by Lemma 6.4} \\
&\leq 2\varepsilon_l \left(\sum_i \zeta'_i \right)^{1/2} |\mu_{.,j} - \mu_{.,j'}| \leq 4\varepsilon_l \sqrt{km} |\mu_{.,j} - \mu_{.,j'}|.
\end{aligned}$$

Now apply Markov inequality to get

$$\text{Prob}(|X| \geq \frac{1}{8} |\mu_{.,j} - \mu_{.,j'}|^2) \leq 32\varepsilon_l \sqrt{km} / |\mu_{.,j} - \mu_{.,j'}| \leq 80\varepsilon_l \sqrt{k/\alpha p_0}.$$

If $|X| \leq |\mu_{.,j} - \mu_{.,j'}|^2/8$, then, $|\hat{B}_{.,j} - \mu_{.,j'}| \geq |\hat{B}_{.,j} - \mu_{.,j}| + 3|\mu_{.,j} - \mu_{.,j'}|/4 \geq |\hat{B}_{.,j} - \mu_{.,j}| + c_{c_0} k^2 \sqrt{\varepsilon} m$, by (9). This proves the first assertion of the Lemma.

The second statement of the Lemma follows by applying H-C to the random variable $\sum_j Z_j/s$, where, Z_j is the indicator random variable of $B_{.,j}$ not satisfying the proximity condition (and using (9).)

The last Lemma implies that the algorithm TSVD correctly identifies the dominant topic in all but at most $\varepsilon_0 w_0/10$ fraction of the documents by Theorem (6.7).

Lemma 7.8 *With probability at least $1 - \exp(-w_0 s \varepsilon^2/8)$, TSVD correctly identifies the dominant topic in all but at most $\varepsilon_0 w_0 \delta/10$ fraction of documents in each T_l .*

7.3 Identifying Catchwords

Recall the definition of J_l from Step 5a of the algorithm. The two lemmas below are roughly converses of each other which prove roughly that J_l consists of those i for which M_{il} is strictly higher than $M_{il'}$. Using them, Lemma 7.11 says that almost all the $\varepsilon_0 w_0 s/2$ documents found in Step 6 of the algorithm are $1 - \delta$ pure for topic l .

Lemma 7.9 *Let $\nu = \gamma(1 - 2\delta)/(1 + \delta)$. If $i \in J_l$, then for all $l' \neq l$, $M_{il} \geq \nu M_{il'}$ and $M_{il} \geq \frac{3}{m\delta^2} \ln(20/\varepsilon w_0)$.*

Proof: It is easy to check that the assumptions (2) and (1) imply $\nu \geq 2$. Let $i \in J_l$. By the definition of J_l in the algorithm, $g(i, l) \geq (4/m\delta^2) \ln(20/\varepsilon w_0)$. Note that $P_{ij} \leq \text{Max}_{l_1} M_{il_1}$ for all j . So,

$$\max_{l_1} M_{il_1} \geq \frac{3}{m\delta^2} \ln(20/\varepsilon w_0). \quad (19)$$

If the Lemma is false, then, for l' attaining $\text{Max}_{l_1 \neq l} M_{il_1}$, we have $M_{il} < \nu M_{il'}$. Recall $R_{l'}$ defined in Step 4c of the algorithm. Let

$$\hat{T}_{l'} = R_{l'} \cap (\text{the set of } 1 - \delta \text{ pure documents in } T_{l'}).$$

Since all but $\varepsilon_0 w_0 s/10$ documents in $T_{l'}$ belong to $R_{l'}$, we have $|\hat{T}_{l'}| \geq 0.9\varepsilon_0 w_0 s$. For $j \in \hat{T}_{l'}$, $P_{ij} \geq M_{il'} W_{l'j} \geq (1 - \delta) M_{il'}$. So, $\text{Prob}(A_{ij} < M_{il'}(1 - 2\delta)) \leq \exp(-m\delta^2 M_{il'}/3) \leq \varepsilon w_0/4$ using (19). Thus the number of documents in $R_{l'}$ for which $A_{ij} \geq M_{il'}(1 - 2\delta)$ is at least $0.9\varepsilon_0 w_0 s - 3\varepsilon w_0 s \geq .6\varepsilon_0 w_0 s$. This implies that with probability at least $1 - \exp(-c\varepsilon^2 s w_0)$, $g(i, l') \geq M_{il'}(1 - 2\delta)$.

Now, for $j \in T_l$, $P_{ij} \leq \text{Max}(M_{il}, M_{il'}) \leq \nu M_{il'}$. So, $\text{Prob}(A_{ij} > M_{il'}\nu(1 + \delta)) \leq \varepsilon w_0/4$, again using (19). At most $\varepsilon_0 w_0 s/10$ documents of other T_{l_1} , $l_1 \neq l$ are in R_l (by Lemma 7.8). So, whp, $g(i, l) \leq M_{il'}\nu(1 + \delta)$ and so we have

$$g(i, l) \leq \frac{\nu(1 + \delta)}{1 - 2\delta} g(i, l'),$$

contradicting the definition of J_l . So, we must have that $M_{il} \geq \nu M_{il'}$ for all $l' \neq l$. The second assertion of the Lemma now follows from (19).

Lemma 7.10 *If $M_{il} \geq \text{Max} \left(\frac{5}{m\delta^2} \ln(20/\varepsilon w_0), \text{Max}_{l' \neq l} \frac{1}{\rho} M_{il'} \right)$, then, with probability at least $1 - \exp(-c\varepsilon^2 w_0 s)$, we have that $i \in J_l$. So, $S_l \subseteq J_l$.*

Proof: Let $\hat{T}_l = R_l \cap$ (set of $1 - \delta$ pure documents in T_l). For $j \in \hat{T}_l$, $P_{ij} \geq M_{il}(1 - \delta)$ which implies that whp, (since $|\hat{T}_l| \geq 0.9\varepsilon_0 s$, again by Lemma 7.8)

$$g(i, l) \geq M_{il}(1 - 2\delta) \quad (20)$$

On the other hand, for $j \in T_{l'}$ and for $l' \neq l$, $i : M_{il'} \leq \rho M_{il}$ (hypothesis of the Lemma), $P_{ij} \leq M_{il} W_{lj} + \rho M_{il}(1 - W_{lj}) \leq M_{il}(\beta + \rho)$. So whp,

$$g(i, l') \leq M_{il}(\beta + \rho)(1 + \delta). \quad (21)$$

From (20) and (21) and hypothesis of the Lemma, it follows that

$$g(i, l) \geq \text{Max} \left(\frac{4}{m\delta^2} \ln(1/\varepsilon w_0), \frac{(1 - 2\delta)}{(1 + \delta)(\beta + \rho)} g(i, l') \right).$$

So, $i \in J_l$ as claimed. It only remains to check that i in S_l satisfies the hypothesis of the Lemma which is obvious.

Lemma 7.11 *Let $\nu_l = \sum_{i \in J_l} M_{il}$ and let L be the set of $\lfloor (s\varepsilon_0 w_0 / 2) \rfloor$ $A_{.,j}$'s whose average is returned in Step 6 of the TSVD Algorithm as $\hat{M}_{.,l}$. With probability at least $1 - c \exp(-c\varepsilon^2 w_0 s)$, we have:*

$$\left| \frac{1}{|L|} \sum_{j \in L} (A_{.,j} - M_{.,l}) \right|_1 \leq O(\delta). \quad (22)$$

Proof: The proof needs care since J_l is itself a random set dependent on $A^{(2)}$. To understand the proof intuitively, if we pretend that there is no conditioning of J_l on $A^{(2)}$, then, basically, our arguments in Lemma 7.9 would yield this Lemma. However, we have to work harder to avoid conditioning effects. Define

$$K_l = \{i : M_{il} \geq \nu M_{il'} \forall l' \neq l; M_{il} \geq (3/m\delta^2) \ln(20/\varepsilon w_0)\}.$$

Note that K_l is not a random set; it does not depend on A , just on M which is fixed. Lemma 7.9 proved that $J_l \subseteq K_l$. Since $\sum_i M_{il} = 1$, we have $|K_l| \leq m\delta^2/3$. The probability bounds given here will be after conditioning on \mathbf{W} . [In other words, we prove statements of the form $\text{Prob}(\mathcal{E} | \mathbf{W}) \leq a$ which is (the usual) shorthand for: for each possible value w of the matrix W , $\text{Prob}(\mathcal{E} | \mathbf{W} = w) \leq a$.] This will be possible, since, even after fixing W , the $A_{.,j}$ are independent, though certainly not identically distributed now, since the $W_{.,j}$ may differ.

For $i \in K_l$, we have for all j , $P_{ij} = \sum_{l'} M_{il'} W_{l'j} \leq M_{il}$, since, $M_{il'} \leq M_{il}/\nu \leq M_{il}/2$ for $l' \neq l$. For any $x \leq M_{il}$,

$$\text{Prob}(|A_{ij}^{(2)} - P_{ij}| \geq \delta M_{il} | W, P_{ij} = x) \leq 2 \exp \left(-\frac{\delta^2 M_{il}^2 m}{2(1 + \delta)x} \right) \leq 2 \exp \left(-\frac{m\delta^2 M_{il}}{3} \right).$$

Noting that $m\delta^2 M_{il} \geq 3 \ln(20/\varepsilon w_0)$ for $i \in K_l$, we get

$$\text{Prob}(|A_{ij}^{(2)} - P_{ij}| \geq \delta M_{il} | W) \leq \varepsilon w_0 / 20.$$

Using the union bound over all $i \in K_l$ yields (for each $j \in [s]$),

$$\text{Prob}(\exists i \in K_l : |A_{ij}^{(2)} - P_{ij}| \geq \delta M_{il} | W) \leq \frac{m\delta^2 \varepsilon w_0}{20} \leq \frac{\varepsilon_0 w_0 \delta^2}{20},$$

by (9). Let

$$BAD = \{j : \exists i \in K_l : |A_{ij}^{(2)} - P_{ij}| \geq \delta M_{il}\}.$$

Using the independence of $A_{.,j}$, (even conditioned on W), apply H-C to get that for the event

$$\mathcal{E} : |BAD| \geq \frac{s\varepsilon_0 w_0 \delta}{10}$$

$$\text{Prob}(\mathcal{E} \mid W) \leq 2 \exp(-c\varepsilon w_0 s). \quad (23)$$

After paying the failure probability, for the rest of the proof, assume that $\neg \mathcal{E}$ holds. Let $U_l = \{j : W_{lj} \geq 1 - \delta\}$. By the dominant topic assumption, we know that $|U_l| \geq \varepsilon_0 w_0 s$. So, $|U_l \setminus BAD| \geq 4\varepsilon_0 w_0 s/5$ and we get (using (9)):

$$\forall N_l \subseteq K_l, \left| \left\{ j : W_{lj} \geq 1 - \delta ; \sum_{i \in N_l} A_{ij}^{(2)} \geq (1 - 2\delta) \sum_{i \in N_l} M_{il} \right\} \right| \geq 4\varepsilon_0 w_0 s/5. \quad (24)$$

Now consider $j : W_{lj} \leq (1 - 6\delta)$ and $i \in K_l$.

$$P_{ij} \leq M_{il} W_{lj} + \sum_{l' \neq l} M_{il'} W_{l'j} \leq M_{il}(1 - 6\delta) + \frac{M_{il}}{\nu} 6\delta \leq M_{il}(1 - 3\delta),$$

since by (2) and (1), we have that $\nu \geq 2$. So, for a j with $W_{lj} \leq 1 - 6\delta$ to have $\sum_{i \in J_l} A_{ij}^{(2)} \geq (1 - 2\delta)\nu_l$, j must be in BAD . This gives us

$$\forall N_l \subseteq K_l, \left| \left\{ j : W_{lj} \leq (1 - 6\delta) ; \sum_{i \in N_l} A_{ij}^{(2)} \geq (1 - 2\delta) \sum_{i \in N_l} M_{il} \right\} \right| \leq \varepsilon_0 w_0 \delta s/10. \quad (25)$$

Let L be the set of $\lfloor \varepsilon_0 w_0 s/2 \rfloor$ j achieving the highest $\sum_{i \in J_l} A_{ij}^{(2)}$. By the above, L contains at most $\varepsilon_0 \delta s/5$ j 's with $W_{lj} < 1 - 6\delta$, the rest being j with $W_{lj} \geq 1 - 6\delta$. So are we finished with the proof - i.e., does this prove (22)? The answer is unfortunately, no. We can show from the above that $\sum_{i \in J_l} |A_{ij} - M_{il}| \leq O(\delta)$ for most $j \in L$ and so the average of $A_{.,j}$, $j \in L$ is close to $M_{.,l}$ when we restrict only to $i \in J_l$. But, on words not in J_l , we have not proved that the average of $A_{ij}^{(2)}$, $j \in L$ is close to $M_{.,l}$. We will do so presently, but first note that this is not a trivial task. For example, if say, $M_{il} = \Omega(1/d)$ for all $i \notin K_l$ (or for a fraction of them) so that $\sum_{i \notin K_l} M_{il} \in \Omega(1)$, then an individual $A_{.,j}$ could have $O(m)$ of the A_{ij} , $i \notin K_l$ set to $1/m$. [One copy of each of $O(m)$ words picked to be in the document.] But then we would have $|A_{.,j} - M_{.,l}|_1 \in \Omega(1)$ which is too much error. We will show that since we are taking the average over L and not just a single document, this will not happen. But the proof is again tricky because of conditioning: both J_l and L depend on the data. So, to argue that the average over L behaves well, we have to prove it for each possible L . There are at most $\binom{s}{\lfloor \varepsilon_0 w_0 s/2 \rfloor} \leq (2/\varepsilon_0 w_0 s)^{\varepsilon_0 w_0 s/2}$ possible L 's and we will be able to take the union bound over all of them.

Claim 7.1 *With probability at least $1 - c m d k \exp(-c\varepsilon^2 w_0 s)$, we have for each $L \subseteq [s]$ with $|L| = \lfloor \varepsilon_0 w_0 s/2 \rfloor$:*

$$\left| \frac{1}{|L|} \sum_{j \in L} (A_{.,j} - P_{.,j}) \right|_1 \leq O(\delta).$$

Proof: Let $X = \left| \frac{1}{|L|} \sum_{j \in L} (A_{.,j} - P_{.,j}) \right|_1$. Each $A_{.,j}$ is itself the average of m independent choices of words. So

$$X = \left| \frac{1}{m|L|} \sum_{j \in L} \sum_{r=1}^m (A_{.,j}^{(r)} - P_{.,j}) \right|_1.$$

So, X is a function of $m|L|$ independent random variables. Changing any one of these arbitrarily changes X by at most $1/m|L|$.

Recall the Bounded Difference inequality [14]:

Lemma 7.12 Let z_1, \dots, z_n, z'_i are $(n+1)$ independent random variables each taking values in \mathcal{Z} and h be a measurable function from \mathcal{Z}^n to \mathbb{R} with constants $r_i \geq 0, i \in [n]$ such that

$$\max_{z_1, \dots, z_n, z'_i \in \mathcal{Z}} |h(z_1, \dots, z_n) - h(z_1, \dots, z'_i, \dots, z_n)| \leq r_i$$

If $E(h)$ is the expectation of h then $\text{Prob}(|h - E(h)| \geq t) \leq 2 \exp\left(-\frac{t^2}{\sum_{i=1}^n r_i^2}\right)$.

Using this we get

$$\text{Prob}(|X - EX| \geq c\delta) \leq \exp(-c\delta^2 \varepsilon_0 w_0 s m).$$

The “extra” m in the exponent helps kill the upper bound of $(2/\varepsilon_0 w_0 s)^{\varepsilon_0 w_0 s/2}$ on the number of L ’s and gives us

$$|X - EX| \leq O(\delta) \forall L.$$

We still have to bound EX . By Jenson’s inequality,

$$EX \leq \frac{1}{|L|} \sum_i \left(E \left(\left(\sum_{j \in L} (A_{ij} - P_{ij}) \right)^2 \right) \right)^{1/2} \leq \frac{1}{|L|} \sum_i \sqrt{\sum_{j \in L} P_{ij}} \leq \sqrt{d}/\sqrt{|L|},$$

where, we have used the independence of $A_{.,j}$ and the fact that $E(A_{ij} - P_{ij})^2 = \text{Var}(A_{ij})$. This proves the claim.

We now bound $\left| \frac{1}{|L|} \sum_{j \in L} (P_{.,j} - M_{.,l}) \right|_1$. Note that by (24) and (25), all but at most $\varepsilon_0 w_0 \delta s / 10$ of the j ’s in L have $W_{lj} \geq 1 - 6\delta$, whence, we get $|P_{.,j} - M_{.,l}|_1 \leq 6\delta$ for these j . For the j with $W_{lj} < 1 - 6\delta$, we just use $|P_{.,j} - M_{.,l}|_1 \leq 2$. So

$$\left| \frac{1}{|L|} \sum_{j \in L} (P_{.,j} - M_{.,l}) \right|_1 \leq 6\delta + \frac{0.2\varepsilon_0 w_0 \delta s}{10|L|} \in O(\delta).$$

This finishes the proof of (22).