

Unbiased estimates for products of moments and cumulants for finite and infinite populations

by

Christopher S. Withers, Industrial Research Limited, Lower Hutt, New Zealand
email: kit.withers@gmail.com

Saralees Nadarajah, University of Manchester, Manchester M13 9PL, UK
email: mbbssn2@manchester.ac.uk

Abstract: Let $F = F_N$ be the distribution of a finite real population of size N . Let $\hat{F} = F_n$ be the empirical distribution of a sample of size n drawn from the population without replacement. We prove the following remarkable *inversion principle* for obtaining unbiased estimates. Let $T(F_N)$ be any product of the moments or cumulants of F_N . Let $T_{n,N}(F_N) = ET(F_n)$. Then $ET_{n,N}(F_n) = T(F_N)$. We also obtain an explicit expression for $T_{n,N}(F_N)$ for all $T(F_N)$ of order up to 6.

We also prove the following related result. If F_n and F_N are the sample and population distributions, the only functionals for which $ET(F_n) = \lambda_{n,N}T(F_N)$ are noncentral moments, and generalized second and third order central moments. For these three cases the eigenvalues are $\lambda_{n,N} = 1$, $(1 - n^{-1})(1 - N^{-1})^{-1}$, and $(1 - n^{-1})(1 - 2n^{-1})(1 - N^{-1})^{-1}(1 - 2N^{-1})^{-1}$ respectively.

AMS subject classifications: Primary 62G05; Secondary 62G20

Keywords and phrases: Cumulants; Finite population; Moments; Unbiased estimates.

1 Introduction

Given a random sample of size n without replacement say X_1, \dots, X_n from a finite real population x_1, \dots, x_N with distribution $F(x) = N^{-1} \sum_{i=1}^N 1(x_i \leq x)$, mean $\mu = m_1 = N^{-1} \sum_{i=1}^N x_i$,

r th moment $m_r = N^{-1} \sum_{i=1}^N x_i^r$, r th central moment $\mu_r = \mu_r(F) = N^{-1} \sum_{i=1}^N (x_i - \mu)^r$ and r th cumulant κ_r , we obtain unbiased estimates (UEs) for products of them

$$\begin{aligned} m(1^{r_1} 2^{r_2} \dots) &= m_1^{r_1} m_2^{r_2} \dots, \\ \mu(1^{r_1} 2^{r_2} \dots) &= \mu^{r_1} \mu_2^{r_2} \mu_3^{r_3} \dots, \\ \kappa(1^{r_1} 2^{r_2} \dots) &= \kappa_1^{r_1} \kappa_2^{r_2} \dots \end{aligned}$$

and for products of the joint central moments

$$\mu(1^{r_1} 2^{r_2} \dots) = E(\hat{\mu} - \mu)^{r_1} (\hat{\mu}_2 - E \hat{\mu}_2)^{r_2} \dots$$

and for products of the corresponding joint cumulants $\kappa(1^{r_1}2^{r_2}\dots)$, for *weight* ≤ 6 , where the weight is

$$r(1^{r_1}2^{r_2}\dots) = 1 \cdot r_1 + 2 \cdot r_2 + \dots \text{ that is } r(\boldsymbol{\pi}) = \pi_1 + \pi_2 + \dots$$

for $\boldsymbol{\pi} = (\pi_1, \pi_2, \dots)$ a partition of r . We assume all partitions $\boldsymbol{\pi}$ are put into ascending order $(1^{r_1}2^{r_2}\dots)$. For example, we write (1^22) or (112) rather than (21^2) or (121) .

Our UEs are given in terms of

$$\widehat{m}(1^{r_1}2^{r_2}\dots) = \widehat{m}_1^{r_1}\widehat{m}_2^{r_2}\dots, \text{ and } \widehat{\mu}(1^{r_1}2^{r_2}\dots) = \widehat{\mu}^{r_1}\widehat{\mu}_2^{r_2}\dots,$$

respectively, where

$$\widehat{\mu} = \widehat{m}_1 = \bar{X} = n^{-1} \sum_{i=1}^n X_i, \quad \widehat{m}_r = n^{-1} \sum_{i=1}^n X_i^r, \quad \widehat{\mu}_r = \mu_r(\widehat{F}) = n^{-1} \sum_{i=1}^n (X_i - \bar{X})^r$$

and $\widehat{F}(x) = n^{-1} \sum_{i=1}^n 1(X_i \leq x)$, the sample distribution.

For $\boldsymbol{\pi}$ a partition of r , Section 2 gives $E\widehat{m}(\boldsymbol{\pi})$ and an UE of $m(\boldsymbol{\pi})$ for $r \leq 6$. Section 3 gives $E\widehat{\mu}(\boldsymbol{\pi})$ and an UE of $\mu(\boldsymbol{\pi})$ for $r \leq 6$.

We discover a remarkable *inversion principle*. A result of this is that these UEs do not require having to invert any matrices or solve any sets of linear equations. Set

$$\begin{aligned} \mathbf{m}_{(r)} &= \{m(\boldsymbol{\pi}) : \boldsymbol{\pi} \text{ a partition of } r\}, \\ \boldsymbol{\mu}_{(r)} &= \{\mu(\boldsymbol{\pi}) : \boldsymbol{\pi} \text{ a partition of } r\}, \\ \mathbf{K}_{(r)} &= \{K(\boldsymbol{\pi}) : \boldsymbol{\pi} \text{ a partition of } r\}. \end{aligned}$$

In Section 2, we derive a matrix $\mathbf{B}_r = \mathbf{B}_r(N, n)$ from Skellam (1949) such that $E\widehat{\mathbf{m}}_{(r)} = \mathbf{B}_r\mathbf{m}_{(r)}$, so $\mathbf{B}_r(N, n)^{-1}\widehat{\mathbf{m}}_{(r)}$ is an UE of $\mathbf{m}_{(r)}$.

In Section 3, we derive a matrix $\mathbf{C}_r = \mathbf{C}_r(N, n)$ From Sukhatme (1944) such that $E\widehat{\boldsymbol{\mu}}_{(r)} = \mathbf{C}_r\boldsymbol{\mu}_{(r)}$, so $\mathbf{C}_r(N, n)^{-1}\widehat{\boldsymbol{\mu}}_{(r)}$ is an UE of $\boldsymbol{\mu}_{(r)}$. So, expressing cumulants in terms of moments as $\mathbf{K}_{(r)} = \mathbf{G}_r\mathbf{m}_{(r)}$, where \mathbf{G}_r is a matrix of constants, it follows that

$$E\widehat{\mathbf{K}}_{(r)} = \mathbf{D}_r(N, n)\mathbf{K}_{(r)} \text{ where } \mathbf{D}_r(N, n) = \mathbf{G}_r\mathbf{C}_r(N, n)\mathbf{G}_r^{-1}. \quad (1)$$

The inversion principle proved in Section 7 states the following amazing result.

$$\mathbf{B}_r(N, n)^{-1} = \mathbf{B}_r(n, N), \quad \mathbf{C}_r(N, n)^{-1} = \mathbf{C}_r(n, N), \quad \mathbf{D}_r(N, n)^{-1} = \mathbf{D}_r(n, N). \quad (2)$$

This implies that for $T(F)$ a product of moments or cumulants, if $T_{n,N}(F) = ET(\widehat{F})$ then $ET_{N,n}(\widehat{F}) = T(F)$. In a later paper we shall extend this result to more general functionals.

Section 4 gives equivalent multivariate results to those of Section 3. For partitions π_1, π_2, \dots , set

$$\mu(\pi_1, \pi_2, \dots) = \mu(\pi_1)\mu(\pi_2)\dots$$

and

$$\kappa(\pi_1, \pi_2, \dots) = \kappa(\pi_1) \kappa(\pi_2) \cdots$$

Section 5 gives $E\widehat{\mu}(\pi_1, \pi_2, \dots)$ and an UE of $\mu(\pi_1, \pi_2, \dots)$ up to total order 6, in particular UEs for $\mu(1^r)$. MAPLE was used to simplify the UE of $\mu(1^r)$ given by Dwyer and Tracy (1980) for $r \leq 5$ and to confirm they agreed with our results. Earlier, Nath (1968,1969) gave $\mu(1^r)$ for $r = 3, 4$ and UEs for them.

Section 6 gives $E\widehat{\kappa}(\pi_1, \pi_2, \dots)$ and an UE of $\kappa(\pi_1, \pi_2, \dots)$ up to total order 6.

Section 7 also proves a multivariate inversion principle: in this case the dimension $n_r \times n_r$ in (1.7) jumps to $N_r \times N_r$, where

$$n_r \text{ is the number of partitions of } r, \quad N_r = \sum_{\pi} P(\pi)$$

and $P(\pi)$ is the partition function

$$P(1^{r_1} 2^{r_2} \dots) = r! / \prod_{i=1} (i!^{r_i} r_i!) \quad (3)$$

for $r = \sum_{i=1} i r_i$.

Section 8 proves the following related result. The only functionals $T(F)$ for which

$$\lambda_{n,N} = ET(F_n) / T(F) \text{ does not depend on } F$$

are of the form

$$T(F) = ES(X), \quad \mu_{1,2}^t(F), \quad \mu_{1,2,3}^u(F),$$

where

$$\mu_{1,2}^t(F) = E \{t(X, X) - t(X, Y)\}, \quad \mu_{1,2,3}^u(F) = E \{u(X, X, X) - u(X, Y, Y) - u(Y, X, Y) - u(Y, Y, X) + 2u(X, Y, Z)\}$$

where X, Y, Z are independent with distribution F . For these three cases $\lambda_{n,N} = 1, (1 - n^{-1}) / (1 - N^{-1})$ and $\prod_{i=1}^2 \{(1 - i n^{-1}) / (1 - i N^{-1})\}$.

Pierce (1940) gives transformations that allow one to obtain noncentral moments up to order six for finite populations from those for infinite populations. He also covers the case where the sample values X_1, \dots, X_n may have different moments.

We shall use π for a partition of r , π_- for a partition of r excluding one's say $\pi_- = (2^{r_2} 3^{r_3} \dots)$ with $2r_2 + 3r_3 + \dots = r$, and π_+ for a partition of r including at least one 1. Let n_r be the number of partitions of r , and N_r the sum of the partitions of r . We also partition vectors and matrices using subscript $+$ and $-$. For example,

$$\boldsymbol{\mu}_{-(r)} = \{\mu(\pi_-)\}, \quad \boldsymbol{\mu}_{+(r)} = \{\mu(\pi_+)\},$$

and

$$\mathbf{B}_r = \mathbf{B}_r(N, n) = \begin{pmatrix} \mathbf{B}_{--r} & \mathbf{0} \\ \mathbf{B}_{+-r} & \mathbf{B}_{++r} \end{pmatrix}$$

since $\mathbf{B}_{-+r} = \mathbf{0}$.

Since also $\mathbf{C}_{-+r} = \mathbf{D}_{-+r} = \mathbf{0}$, the inversion principle (1) implies

$$\mathbf{B}_{--r}(N, n)^{-1} = \mathbf{B}_{--r}(n, N), \quad \mathbf{C}_{--r}(N, n)^{-1} = \mathbf{C}_{--r}(n, N), \quad \mathbf{D}_{--r}(N, n)^{-1} = \mathbf{D}_{--r}(n, N), \quad (4)$$

so that

$$\mathbf{B}_{--r}(n, N)\widehat{\mathbf{m}}_{-(r)} \text{ is an UE of } \mathbf{m}_{-(r)},$$

$$\mathbf{C}_{--r}(n, N)\widehat{\boldsymbol{\mu}}_{-(r)} \text{ is an UE of } \boldsymbol{\mu}_{-(r)},$$

and

$$\mathbf{D}_{--r}(n, N)\widehat{\mathbf{K}}_{-(r)} \text{ is an UE of } \mathbf{K}_{-(r)}.$$

The number of parts in $\boldsymbol{\pi}$ is denoted by $q(\boldsymbol{\pi})$:

$$q(1^{r_1}2^{r_2}\dots) = r_1 + r_2 + \dots$$

Set $(n)_i = n(n-1)\dots(n-i+1) = n!/(n-i)!$.

2 Products of non central moments

Here, we derive the result

$$E\widehat{\mathbf{m}}(r) = \mathbf{B}_r\mathbf{m}(r), \quad (5)$$

where $\mathbf{m}(r) = \{m(\boldsymbol{\pi})\}$ and $m(\boldsymbol{\pi}) = m_{\pi_1}m_{\pi_2}\dots$ for $\boldsymbol{\pi} = (\pi_1, \pi_2, \dots)$. Skellam (1949) showed that

$$ES_{a_1}\dots S_{a_s} = \sum_{\boldsymbol{\pi}} \lambda(\boldsymbol{\pi}) \sum_{P(\boldsymbol{\pi})} s_{R_{\pi_1}} s_{R_{\pi_2}} \dots$$

for

$$S_r = \sum_{i=1}^n X_i^r = n\widehat{m}_r, \quad s_r = \sum_{i=1}^N x_i^r = Nm_r,$$

where $\sum_{\boldsymbol{\pi}}^s$ sums over partitions $\boldsymbol{\pi}$ of s , the partition function $P(\boldsymbol{\pi})$ is given by (3), $R_{\pi_1} =$

$a_1 + \dots + a_{\pi_1}$, $R_{\pi_2} = a_{\pi_1+1} + \dots + a_{\pi_1+\pi_2}$, etc and $\sum_{P(\boldsymbol{\pi})}$ sums over all $P(\boldsymbol{\pi})$ such terms and $\lambda(\boldsymbol{\pi}) = \lambda_{\boldsymbol{\pi}}$ is the Carver function. For example,

$$E S_{a_1}S_{a_2}S_{a_3} = \lambda(3)s_{a_1+a_2+a_3} + \lambda(12)\sum^3 s_{a_1+a_2}s_{a_3} + \lambda(1^3)s_{a_1}s_{a_2}s_{a_3}$$

and

$$\begin{aligned}
ES_{a_1} \cdots S_{a_5} &= \lambda(5)s_{a_1+\dots+a_5} + \lambda(14) \sum^3 s_{a_1+\dots+a_4} s_{a_5} \\
&+ \lambda(23) \sum^{10} s_{a_1+a_2+a_3} s_{a_4+a_5} + \lambda(1^2 3) \sum^{10} s_{a_1+a_2+a_3} s_{a_4} s_{a_5} \\
&+ \lambda(12^2) \sum^{15} s_{a_1+a_2} s_{a_3+a_4} s_{a_5} + \lambda(1^3 2) \sum^{10} s_{a_1+a_2} s_{a_3} s_{a_4} s_{a_5} \\
&+ \lambda(1^5) s_{a_1} \cdots s_{a_5},
\end{aligned} \tag{6}$$

where

$$\begin{aligned}
\lambda(3) &= e_1 - 3e_2 + 2e_3, \quad \lambda(12) = e_2 - e_3, \quad \lambda(1^3) = e_3, \\
\lambda(5) &= e_1 - 15e_2 + 50e_3 - 60e_4 + 24e_5, \\
\lambda(14) &= e_2 - 7e_3 + 12e_4 - 6e_5, \\
\lambda(23) &= e_2 - 4e_3 + 5e_4 - 2e_5, \\
\lambda(1^2 3) &= e_3 - 3e_4 + 2e_5, \\
\lambda(12^2) &= e_3 - 2e_4 + e_5, \\
\lambda(1^3 2) &= e_4 - e_5, \\
\lambda(1^5) &= e_5,
\end{aligned}$$

where $e_j = (n)_j / (N)_j$. He wrote (5) out in full for $s \leq 4$. We extend this to $s \leq 6$ in Appendix E. Set

$$\mathbf{S}_{(r)} = \{S_{\pi_1} S_{\pi_2} \cdots : (\pi_1, \pi_2, \dots) \text{ a partition of } r\},$$

and similarly for $\mathbf{s}_{(r)}$. It follows from (5) that

$$E\mathbf{S}_{(r)} = \mathbf{A}_r \mathbf{s}_{(r)}, \tag{7}$$

where

$$\mathbf{A}_r = \left\{ A_{\pi, \pi'} : \pi, \pi' \text{ partitions of } r \right\}$$

and $\mathbf{A}_1, \dots, \mathbf{A}_6$ are as follows:

$$\begin{aligned} \mathbf{A}_1 &= \lambda_1, \quad \mathbf{A}_2 = \begin{pmatrix} \lambda_1 & & \\ \lambda_2 & \lambda_{11} & \\ & & \end{pmatrix}, \quad \mathbf{A}_3 = \begin{pmatrix} \lambda_1 & & \\ \lambda_2 & \lambda_{11} & \\ \lambda_3 & 3\lambda_{12} & \lambda_{13} \end{pmatrix}, \\ \mathbf{A}_4 &= \begin{bmatrix} \lambda_1 & & & & \\ \lambda_2 & \lambda_{11} & & & \\ \lambda_2 & 0 & \lambda_{11} & & \\ \lambda_3 & 2\lambda_{12} & \lambda_{12} & \lambda_{13} & \\ \lambda_4 & 4\lambda_{13} & 3\lambda_{22} & 6\lambda_{122} & \lambda_{14} \end{bmatrix}, \\ \mathbf{A}_5 &= \begin{bmatrix} \lambda_1 & & & & & & & \\ \lambda_2 & \lambda_{11} & & & & & & \\ \lambda_2 & 0 & \lambda_{11} & & & & & \\ \lambda_3 & 2\lambda_{12} & \lambda_{12} & \lambda_{13} & & & & \\ \lambda_3 & \lambda_{12} & 2\lambda_{12} & 0 & \lambda_{13} & & & \\ \lambda_4 & 3\lambda_{13} & \lambda_{13} + 3\lambda_{22} & 3\lambda_{122} & 3\lambda_{122} & \lambda_{14} & & \\ \lambda_5 & 5\lambda_{14} & 10\lambda_{23} & 10\lambda_{113} & 15\lambda_{22} & 10\lambda_{12} & 3\lambda_{15} & \end{bmatrix}, \\ \mathbf{A}_6 &= \begin{bmatrix} \lambda_1 & & & & & & & & & & & & & & & & & & & \\ \lambda_2 & \lambda_{11} & & & & & & & & & & & & & & & & & & & \\ \lambda_2 & 0 & \lambda_{11} & & & & & & & & & & & & & & & & & & \\ \lambda_2 & 0 & 0 & \lambda_{11} & & & & & & & & & & & & & & & & & \\ \lambda_3 & 2\lambda_{12} & \lambda_{12} & 0 & \lambda_{11} & & & & & & & & & & & & & & & & \\ \lambda_3 & \lambda_{12} & \lambda_{12} & \lambda_{12} & 0 & \lambda_{13} & & & & & & & & & & & & & & & \\ \lambda_3 & 0 & 3\lambda_{12} & 0 & 0 & 0 & & & & & & & & & & & & & & & \\ \lambda_4 & 3\lambda_{13} & \lambda_{13} + 3\lambda_{22} & 0 & 3\lambda_{122} & 3\lambda_{122} & & & & & & & & & & & & & & & \\ \lambda_4 & 2\lambda_{13} & 2\lambda_{13} + \lambda_{22} & 2\lambda_{22} & \lambda_{122} & 4\lambda_{122} & & & & & & & & & & & & & & & \\ \lambda_5 & 4\lambda_{14} & \lambda_{14} + 6\lambda_{23} & 4\lambda_{23} & 6\lambda_{113} & 4\lambda_{113} + 12\lambda_{122} & 3\lambda_{122} & & & & & & & & & & & & & & \\ \lambda_6 & 6\lambda_{15} & 15\lambda_{24} & 10\lambda_{33} & 15\lambda_{124} & 60\lambda_{123} & 15\lambda_{23} & \lambda_{13} & \lambda_{14} & & & & & & & & & & & & \end{bmatrix}. \end{aligned}$$

Since $S_r = n\widehat{m}_r$ and $s_r = Nm_r$, (7) implies (5) with

$$\mathbf{B}_r = \mathbf{D}_{r,n}^{-1} \mathbf{A}_r \mathbf{D}_{r,N},$$

where $\mathbf{D}_{r,N} = \text{diag} \{N^{q(\pi)} : r(\pi) = r\}$. For example, $\mathbf{D}_{4,N} = \text{diag} (N, N^2, N^2, N^3, N^4)$. Writing

$$\begin{aligned} \mathbf{A}_r(N, n) &= \mathbf{A}_r = \left\{ A_{\pi, \pi'}(N, n) : \pi, \pi' \text{ partition of } r \right\}, \\ \mathbf{B}_r(N, n) &= \mathbf{B}_r = \left\{ B_{\pi, \pi'}(N, n) : \pi, \pi' \text{ partitions of } r \right\}, \end{aligned}$$

it follows from (5) that

$$E\widehat{m}_{\pi_1} \widehat{m}_{\pi_2} \cdots = \sum_{\pi'}^r B_{\pi, \pi'}(N, n) m_{\pi_1} m_{\pi_2} \cdots = b_{\pi}(N, n, \mathbf{m})$$

say. By the inversion principle

$$\mathbf{B}_r(N, n)^{-1} = \mathbf{B}_r(n, N),$$

so $b_{\pi}(n, N, \widehat{\mathbf{m}})$ is an UE of $m_{\pi_1} m_{\pi_2} \cdots$. For example,

$$E\widehat{m}_1^3 = n^{-3} (\lambda_3 N m_3 + 3\lambda_{21} N^2 m_2 m_1 + \lambda_{13} N^3 m_1^3),$$

so $N^{-3} (\bar{\lambda}_3 n \widehat{m}_3 + 3\bar{\lambda}_{21} n^2 \widehat{m}_2 \widehat{m}_1 + \bar{\lambda}_{13} n^3 \widehat{m}_1^3)$ is an UE of m_1^3 , where $\bar{\lambda}_{\pi}$ is λ_{π} with n and N reversed.

Example 2.1 Suppose that the population consists of Np ones and $N(1-p)$ zeros. Then $m_r = p$ for $r > 0$, so

$$E(n\widehat{p})^r = \sum_{\boldsymbol{\pi}} \lambda_{\boldsymbol{\pi}} P(\boldsymbol{\pi}) (Np)^{q(\boldsymbol{\pi})}.$$

So,

$$E\widehat{p}^r = n^{-r} \sum_{i=1}^r a_{r,i}(N, n) (Np)^i = b_r(N, n, p)$$

say, where

$$a_{r,i}(N, n) = \sum_{\boldsymbol{\pi}} \{\lambda_{\boldsymbol{\pi}} P(\boldsymbol{\pi}) : q(\boldsymbol{\pi}) = i\},$$

and an UE of p^r is $b_r(n, N, \widehat{p})$. This implies another inversion principle: $\mathbf{a}_r = \mathbf{a}_r(N, n) = \{a_{i,j}(N, n)\}$, $r \times r$ with $a_{i,j} = 0$ for $i < j$ has inverse $\mathbf{a}_r(n, N)$. The first six are given by

$$\mathbf{a}_r = \begin{pmatrix} \mathbf{a}_{r-1} & 0 \\ \mathbf{c}'_r & \lambda_{1^r} \end{pmatrix},$$

where $\mathbf{c}'_r = (a_{r,1}, \dots, a_{r,r-1})$ are

$$\begin{aligned} c_2 &= \lambda_2, \quad \mathbf{c}'_3 = (\lambda_3, 3\lambda_{12}), \quad \mathbf{c}'_4 = (\lambda_4, 4\lambda_{13} + 3\lambda_{22}, 6\lambda_{122}), \\ \mathbf{c}'_5 &= (\lambda_5, 5\lambda_{14} + 10\lambda_{23}, 10\lambda_{113} + 15\lambda_{122}, 10\lambda_{132}), \\ \mathbf{c}'_6 &= (\lambda_6, 6\lambda_{15} + 15\lambda_{24} + 10\lambda_{33}, 15\lambda_{124} + 60\lambda_{123} + 15\lambda_{23}, 20\lambda_{133} + 45\lambda_{1222}, 15\lambda_{142}). \end{aligned}$$

The block matrix $\mathbf{A} = \begin{pmatrix} \mathbf{A}_{1,1} & \mathbf{0} \\ \mathbf{A}_{2,1} & \mathbf{A}_{2,2} \end{pmatrix}$ has $\mathbf{A}^{-1} = \begin{pmatrix} \mathbf{A}_{1,1}^{-1} & \mathbf{0} \\ \mathbf{A}_{2,1}^{-1} & \mathbf{A}_{2,2}^{-1} \end{pmatrix}$, where $\mathbf{A}_{2,1}^{-1} = -\mathbf{A}_{2,2}^{-1} \mathbf{A}_{2,1} \mathbf{A}_{1,1}^{-1}$. So, if $\mathbf{A} = \mathbf{A}(N, n)$ then $\mathbf{A}(N, n)^{-1} = \mathbf{A}(n, N)$ if and only if $\mathbf{A}_{i,i}(N, n)^{-1} = \mathbf{A}_{i,i}(n, N)$, $i = 1, 2$ and $-\mathbf{A}_{2,2}(n, N) \mathbf{A}_{2,1}(N, n) \mathbf{A}_{1,1}(n, N) = \mathbf{A}_{2,1}(n, N)$.

Example 2.2 Suppose F is Poisson (λ), that is for $i \geq 0$ a proportion $p_i = e^{-\lambda} \lambda^i / i!$ of the population equal i . This implies that $N = \infty$, but the algebra can be carried out as if $N < \infty$, then $N = \infty$ substituted at the end. For $i \geq 1$ the Stirling numbers of the second kind $S_{i,1}, \dots, S_{i,i}$ are defined by

$$j^i = \sum_{k=1}^i S_{i,k}(j)_k.$$

Then $m_i = EX^i = \sum_{k=1}^i S_{i,k} \lambda^k$. So, $(m_1, \dots, m_r) = (\lambda, \dots, \lambda^r) \boldsymbol{\alpha}'_r$, where $\boldsymbol{\alpha}_r = (S_{i,j})$ is $r \times r$ and $S_{i,j} = 0$ for $j > i$. So, $\boldsymbol{\alpha}'_r^{-1}(\widehat{m}_1, \dots, \widehat{m}_r)$ is an UE of $(\lambda, \dots, \lambda^r)$.

But other UEs of λ^r can be obtained from products of $\{\widehat{m}_i\}$. For example,

$$E(n\widehat{m}_1)^r = \sum_{i=1}^r d_{r,i} N^i,$$

where

$$d_{r,i} = \sum_{\boldsymbol{\pi}}^r \{ \lambda_{\boldsymbol{\pi}} P(\boldsymbol{\pi}) m_{\pi_1} m_{\pi_2} \cdots : q(\boldsymbol{\pi}) = i \},$$

Thus,

$$\begin{aligned} E(n\widehat{m}_1)^2 &= (\lambda^2 + \lambda) \lambda_2 N + \lambda^2 \lambda_{11} N^2 \\ &= (\lambda^2 + \lambda) n + \lambda^2 (n)_2 \text{ at } N = \infty. \end{aligned}$$

So, at $N = \infty$, an UE of $\theta = \lambda^2$ is $\widehat{\theta}_1 = \widehat{m}_1^2 - n^{-1}\widehat{m}_1$.

But by above $\widehat{\theta}_2 = \widehat{m}_2 - \widehat{m}_1$ is also an UE.

One can now apply (6) to calculate $\text{var } \widehat{\theta}_1$, $\text{var } \widehat{\theta}_2$ to see when $\widehat{\theta}_1$ is more efficient than $\widehat{\theta}_2$.

3 Products of powers of the mean and central moments and the Inversion Principle

Here, we derive $E\widehat{\mu}(\boldsymbol{\pi})$ and the UE of $\mu(\boldsymbol{\pi}) = \mu_{\pi_1}\mu_{\pi_2}\cdots$, where $\mu_1 = \mu$. We also elaborate on the inversion principle (2).

By Sukhatme (1944),

$$E\widehat{\mu}(\boldsymbol{\pi}_-) = \sum_{\boldsymbol{\pi}'_-}^r C_{\boldsymbol{\pi}_-, \boldsymbol{\pi}'_-} \mu(\boldsymbol{\pi}'_-), \quad (8)$$

where $C_{\boldsymbol{\pi}_-, \boldsymbol{\pi}'_-}$ is given for $r \leq 6$ in terms of $e_j = (n)_j / (N)_j$.

Set

$$\begin{aligned} \boldsymbol{\mu}_{-(r)} &= \{ \mu(\boldsymbol{\pi}_-) : r(\boldsymbol{\pi}_-) = r \}, \quad \boldsymbol{\mu}_{+(r)} = \{ \mu(\boldsymbol{\pi}_+) : r(\boldsymbol{\pi}_+) = r \}, \\ \boldsymbol{\mu}'_{(r)} &= \left(\boldsymbol{\mu}'_{-(r)}, \boldsymbol{\mu}'_{+(r)} \right) = \{ \mu(\boldsymbol{\pi}) : r(\boldsymbol{\pi}) = r \}', \end{aligned}$$

where $r(\boldsymbol{\pi}) = \pi_1 + \pi_2 + \cdots$, the order of $\boldsymbol{\pi}$. In particular,

$$\begin{aligned} \boldsymbol{\mu}_{-(2)} &= \mu_2, \quad \boldsymbol{\mu}_{-(3)} = \mu_3, \quad \boldsymbol{\mu}_{-(4)} = (\mu_4, \mu_2^2)', \quad \boldsymbol{\mu}_{-(5)} = (\mu_5, \mu_2\mu_3)', \\ \boldsymbol{\mu}_{-(6)} &= (\mu_6, \mu_2\mu_4, \mu_3^2, \mu_2^3)', \\ \boldsymbol{\mu}_{+(2)} &= \mu^2, \quad \boldsymbol{\mu}'_{+(3)} = (\mu\mu_2, \mu^3), \\ \boldsymbol{\mu}'_{+(4)} &= (\mu\mu_3, \mu^2\mu_2, \mu^4), \\ \boldsymbol{\mu}'_{+(5)} &= (\mu\mu_4, \mu\mu_2^2, \mu^2\mu_3, \mu^3\mu_2, \mu^5), \\ \boldsymbol{\mu}'_{+(6)} &= (\mu\mu_5, \mu\mu_2\mu_3, \mu^2\mu_4, \mu^2\mu_2^2, \mu^3\mu_3, \mu^4\mu_2, \mu^6). \end{aligned}$$

Let $\widehat{\boldsymbol{\mu}}_{-(r)}, \widehat{\boldsymbol{\mu}}_{+(r)}, \widehat{\boldsymbol{\mu}}_{(r)}$ denote their sample versions, that is with F replaced by \widehat{F} .

Then the result (8) can be written as

$$E\widehat{\boldsymbol{\mu}}_{-(r)} = \mathbf{C}_{--r} \boldsymbol{\mu}_{-(r)},$$

where $\mathbf{C}_{--r} = (C_{\pi_-, \pi'_-} : \pi_-, \pi'_-$ partitions of r excluding ones). So, $\tilde{\boldsymbol{\mu}}_{-(r)} = \mathbf{C}_{--r}^{-1} \hat{\boldsymbol{\mu}}_{-(r)}$ is an UE of $\boldsymbol{\mu}_{-(r)}$. The coefficients of C_{π_-, π'_-} are given in Appendix A for $r \leq 6$.

Sukhatme (1944) gives $\{C_{\pi, \pi'}\}$ such that

$$E(\hat{\boldsymbol{\mu}} - \boldsymbol{\mu})^{r_1} \hat{\boldsymbol{\mu}}_2^{r_2} \hat{\boldsymbol{\mu}}_3^{r_3} \cdots = \sum_{\pi'_-}^r C_{\pi_-, \pi'_-} \boldsymbol{\mu}(\pi'_-)$$

for $\boldsymbol{\pi} = (1^{r_1} 2^{r_2} \cdots)$. To obtain $E \hat{\boldsymbol{\mu}}^{r_1} \hat{\boldsymbol{\mu}}_2^{r_2} \hat{\boldsymbol{\mu}}_3^{r_3} \cdots$, one just expands $\hat{\boldsymbol{\mu}}^{r_1} = (\delta + \boldsymbol{\mu})^{r_1}$, where $\delta = \hat{\boldsymbol{\mu}} - \boldsymbol{\mu}$. So, one obtains

$$E \hat{\boldsymbol{\mu}}(\boldsymbol{\pi}) = \sum_{\pi'}^r C_{\pi, \pi'} \boldsymbol{\mu}(\pi'),$$

that is

$$E \hat{\boldsymbol{\mu}}_{(r)} = \mathbf{C}_r \boldsymbol{\mu}_{(r)},$$

where $\mathbf{C}_r = (C_{\pi, \pi'} : \pi, \pi'$ partitions of r) and

$$C_{1^i \pi_-, 1^j \pi'_-} = \binom{i}{j} C_{1^{i-j} \pi_-, \pi'_-} \quad (9)$$

for π_-, π'_- partitions excluding ones with $i + r(\boldsymbol{\pi}) = j + r(\boldsymbol{\pi}')$, and the left hand side of (9) is zero if $j > i$. For completeness $\{C_{\pi, \pi'}\}$ are given in Appendix A for $r \leq 6$. An UE of $\boldsymbol{\mu}_{(r)}$ is $\tilde{\boldsymbol{\mu}}_{(r)} = \mathbf{C}_r^{-1} \hat{\boldsymbol{\mu}}_{(r)}$. So, by (10) below

$$\tilde{\boldsymbol{\mu}}_{+(r)} = C_r^{2,1} \hat{\boldsymbol{\mu}}_{-(r)} + C_r^{2,2} \hat{\boldsymbol{\mu}}_{+(r)}$$

is an UE of $\boldsymbol{\mu}_{+(r)}$.

Note that for $\boldsymbol{\pi}_1, \boldsymbol{\pi}_2, \dots$ partitions of r_1, r_2, \dots

$$E \hat{\boldsymbol{\mu}}(\boldsymbol{\pi}_1) E \hat{\boldsymbol{\mu}}(\boldsymbol{\pi}_2) \cdots = \sum_{\pi'_1}^{r_1} \sum_{\pi'_2}^{r_2} \cdots C_{\pi_1, \pi'_1} C_{\pi_2, \pi'_2} \cdots \boldsymbol{\mu}(\boldsymbol{\pi}'_1 + \boldsymbol{\pi}'_2 + \cdots),$$

where

$$\boldsymbol{\mu}(1^{r_1,1} 2^{r_1,2} + \cdots + 1^{r_2,1} 2^{r_2,2} + \cdots + \cdots) = \boldsymbol{\mu}(1^{r_1} 2^{r_2} \cdots)$$

for $r_1 = r_{1,1} + r_{2,1} + \cdots$, $r_2 = r_{1,2} + r_{2,2} + \cdots$, etc. So, $\prod_i E \hat{\boldsymbol{\mu}}(\boldsymbol{\pi}_i)$ has the form $\sum_{\boldsymbol{\pi}}^r D_{\boldsymbol{\pi}} \boldsymbol{\mu}(\boldsymbol{\pi})$,

where $r = r_1 + r_2 + \cdots$ and has UE $\sum_{\pi'}^r D_{\pi'}^* \hat{\boldsymbol{\mu}}(\boldsymbol{\pi})$, where $D_{\pi'}^* = \sum_{\boldsymbol{\pi}}^r D_{\boldsymbol{\pi}} C^{\boldsymbol{\pi}, \boldsymbol{\pi}'}$. For example, in this way we can write down UEs for $\boldsymbol{\mu}^{r_1} (E \hat{\boldsymbol{\mu}}_2)^{r_2} (E \hat{\boldsymbol{\mu}}_3)^{r_3} \cdots$ for $r_1 + 2r_2 + 3r_3 + \cdots \leq 6$.

Since \mathbf{C}_r has the form $\begin{pmatrix} \mathbf{C}_{1,1} & \mathbf{0} \\ \mathbf{C}_{2,1} & \mathbf{C}_{2,2} \end{pmatrix}$ with $\mathbf{C}_{1,1} = \mathbf{C}_{--r}$, its inverse is

$$\mathbf{C}_r^{-1} = \begin{pmatrix} \mathbf{C}_r^{1,1} & \mathbf{0} \\ \mathbf{C}_r^{2,1} & \mathbf{C}_r^{2,2} \end{pmatrix} \quad (10)$$

with $\mathbf{C}_r^{1,1} = \mathbf{C}_{1,1}^{-1}$, $\mathbf{C}_r^{2,1} = -\mathbf{C}_{2,2}^{-1}\mathbf{C}_{2,1}\mathbf{C}_{1,1}^{-1}$ and $\mathbf{C}_r^{2,2} = \mathbf{C}_{2,2}^{-1}$.

However, this dimension-reduction method to obtain \mathbf{C}_r^{-1} is not necessary due to the inversion principle, $\mathbf{C}_r(N, n)^{-1} = \mathbf{C}_r(n, N)$. This implies $\mathbf{C}_{-r}(N, n)^{-1} = \mathbf{C}_{-r}(n, N)$ as noted in (4). This result was discovered using MAPLE and has been verified for the \mathbf{C}_{-r} , \mathbf{C}_r of Appendix A, that is for $\{\mathbf{C}_{-r} : r \leq 6\}$, $\{\mathbf{C}_r : r \leq 5\}$. Its proof is given in Section 7. Setting $(C^{\pi, \pi'}) = \mathbf{C}_r^{-1}$ and $\mathbf{C}^{\pi, \pi'}(N, n) = \mathbf{C}^{\pi, \pi'}$, we can write these as $C^{\pi, \pi'}(N, n) = C_{\pi, \pi'}(n, N)$ and $C^{\pi-, \pi'-}(N, n) = C_{\pi-, \pi'-}(n, N)$.

We now show how to obtain UEs for products of cumulants. The r th cumulant κ_r may be written as

$$\kappa_r = \mathbf{a}'_r \boldsymbol{\mu}_{-(r)}$$

for $r \geq 2$, where $a_2 = a_3 = 1$, $\mathbf{a}_4 = (1, -3)$, $\mathbf{a}_5 = (1, -10)$, $\mathbf{a}_6 = (1, -15, -10, 30)$. So, $\tilde{\kappa}_r = \mathbf{a}'_r \tilde{\boldsymbol{\mu}}_{-(r)}$ is an UE for κ_r . For the case $N = \infty$, Fisher (1929) gave these k -statistics, $\boldsymbol{\mu}_{-(r)}$ for $r \leq 6$ and their joint cumulants. Sukhatme obtained $C_{\pi, \pi'}$ using Fisher's method.

Wishart (1952) gave tables for the poly-kays, $k_{r_1, r_2, \dots}$, the UE of $\kappa_{r_1} \kappa_{r_2} \dots$, in terms of the symmetric functions. This is reproduced in Appendix Table 11 of Stuart and Ord (1987) who use $l_{r_1, r_2, \dots}$ for $\kappa_{r_1, r_2, \dots}$. The symmetric functions are given by their Appendix Table 10 in terms of the power sums $s_r = \sum_i x_i^r$. However, no explicit formulae for the polykays in terms of the sample moments or cumulants appear to be available. We rectify this in Appendix D, not by using Tables 11 and 10, but by writing

$$\kappa_{\pi_1} \kappa_{\pi_2} \dots = \begin{cases} \mathbf{a}(\boldsymbol{\pi})' \boldsymbol{\mu}_{-(r)}, & \text{if } \boldsymbol{\pi} = (\pi_1, \pi_2, \dots) \text{ does not contain 1,} \\ \mathbf{b}(\boldsymbol{\pi})' \boldsymbol{\mu}_{(r)}, & \text{if } \boldsymbol{\pi} \text{ contains 1,} \end{cases}$$

where $r = \pi_1 + \pi_2 + \dots$. So the UE of $\kappa_{\pi_1} \kappa_{\pi_2} \dots$ is $\mathbf{a}(\boldsymbol{\pi})' \tilde{\boldsymbol{\mu}}_{-(r)}$ or $\mathbf{b}(\boldsymbol{\pi})' \tilde{\boldsymbol{\mu}}_{-(r)}$.

In Section 12.22, Stuart and Ord (1987) give a number of references on this subject, and say that Dwyer and Tracy (1980) give UEs for products of central moments. This is not so: they give the multivariate equivalent of the UE for $E(\bar{X} - \mu)^r$ for $r \leq 5$. They also give $E\hat{\mu}_r$ and $E(\bar{X} - \mu)^r$ for $r \leq 6$ by a different method, or rather the equivalent multivariate results.

An alternative method of getting UEs for $E(\bar{X} - \mu)^r$, and indeed for $E(\bar{X} - \mu)^i \hat{\mu}(\boldsymbol{\pi}_-)$ is to note that

$$E(\bar{X} - \mu)^i \hat{\mu}(\boldsymbol{\pi}_-) = \sum_{\boldsymbol{\pi}'_-}^{r+i} C_{1^i \boldsymbol{\pi}_-, \boldsymbol{\pi}'_-} \mu(\boldsymbol{\pi}'_-)$$

and so by (1) has UE

$$\sum_{\boldsymbol{\pi}''_-}^{r+i} D_{1^i \boldsymbol{\pi}_-, \boldsymbol{\pi}''_-} \hat{\mu}(\boldsymbol{\pi}''_-),$$

where

$$D_{1^i \boldsymbol{\pi}_-, \boldsymbol{\pi}''_-} = \sum_{\boldsymbol{\pi}'_-}^{r+i} C_{1^i \boldsymbol{\pi}_-, \boldsymbol{\pi}'_-} C^{\boldsymbol{\pi}'_-, \boldsymbol{\pi}''_-},$$

and $C^{\pi', \pi''} = C_{\pi', \pi''}(n, N)$.

Set $\mu_a^* = n^{-1} \sum_{i=1}^n (X_i - \mu)^a$. Given a partition π of r and a set of numbers $\{a_1, \dots, a_r\}$,

the set can be divided into groups of sizes π_1, π_2, \dots in $P(\pi)$ ways. Let $R_j = \sum a_i$ summed over the j th group. By (8),

$$E \prod_{i=1}^r \mu_{a_i}^* = n^{-r} \sum_{\pi} \lambda(\pi) N^{q(\pi)} \sum^{P(\pi)} \mu_{R_1} \mu_{R_2} \cdots. \quad (11)$$

Skellam (1949) checked Sukhatme's first seventeen formulas using (8) and also checked all his $\lambda(\pi)$. We now use (8) to obtain the formula for $E(\hat{\mu} - \mu)\hat{\mu}_5$ overlooked by Sukhatme. Note that

$$(\hat{\mu} - \mu)\hat{\mu}_5 = \mu_1^* \{ \mu_5^* - 5\mu_1^* \mu_4^* + 10\mu_1^{*2} \mu_3^* - 10\mu_1^{*3} \mu_2^* + 4\mu_1^{*5} \}.$$

By (11),

$$\begin{aligned} n^2 E \mu_1^* \mu_5^* &= \lambda(2) N \mu_6, \\ n^3 E \mu_1^{*2} \mu_4^* &= \lambda(3) N \mu_6 + \lambda(21) N^2 \mu_2 \mu_4, \end{aligned}$$

and so on. Collecting terms, $E(\hat{\mu} - \mu)\hat{\mu}_5 = \sum_{\pi_-}^6 C_{15, \pi_-} m(\pi_-)$, where $\{C_{15, \pi_-}\}$ are given in Appendix A.

By (9), $\{C_{15, \pi_+}\}$ are given by $C_{15, 1^j \pi} = C_{5, \pi}$ if $j = 1$ and $C_{15, 1^j \pi} = 0$ if $j > 1$. Note that $\lambda(\pi)$ has the form

$$\lambda(\pi) = \sum_{j=q(\pi)}^{r(\pi)} c_j e_j$$

with

$$c_{q(\pi)} = 1, \quad c_{r(\pi)} = \prod_{i=1}^{r(\pi)} (-1)^{\pi_i - 1} (\pi_i - 1)! = (-1)^{r(\pi) - q(\pi)} \prod_i (\pi_i - 1)!.$$

Now suppose X_1, \dots, X_n is a random sample from an *infinite* population. Then Sukhatme noted that the same results (1.6), (1.17) hold with $N^i e_j = 0$ for $i < j$ and $N^j e_j = (n)_j$. Thus only the first term in the expansion

$$\lambda(\pi) = \sum_{j=q(\pi)}^{r(\pi)} c_j e_j$$

makes a contribution: $N^{q(\pi)} \lambda(\pi) = c_j (n)_j$ at $j = q(\pi)$.

Since the infinite population is that of most interest, we give these simplified formulas for $C_{\pi, \pi'}$ in Appendix B. Because of the inversion principle, for obtaining UEs for $N = \infty$

we need $C_{\boldsymbol{\pi}, \boldsymbol{\pi}'}$ at $n = \infty$. These are given in Appendix C. Putting $n = \infty$ is the same as putting $n^{-i}e_j = 0$ for $i > j$ and $n^{-j}e_j = 1/(N)_j$. Thus only the last term in the expansion

$$\lambda(\boldsymbol{\pi}) = \sum_{j=q(\boldsymbol{\pi})}^{r(\boldsymbol{\pi})} c_j e_j$$

makes a contribution: $n^{-r(\boldsymbol{\pi})}\lambda(\boldsymbol{\pi}) = c_j/(N)_j$ at $j = r(\boldsymbol{\pi})$. For example,

$$\begin{aligned} C_{12,3} &= N \{ \lambda(1)n^{-1} - \lambda(2)n^{-2} \} \\ &= N \{ e_1 n^{-1} - (e_1 - e_2) n^{-2} \} \\ &= \begin{cases} (n)_1 (n^{-1} - n^{-2}), & \text{at } N = \infty, \\ N \{ 1/(N)_1 + 1/(N)_2 \}, & \text{at } n = \infty. \end{cases} \end{aligned}$$

These results agree with James (1958) who gave UEs of μ_r for $r \leq 6$, $N = \infty$.

4 Multivariate extensions

For the multivariate case one considers the population x_1, \dots, x_N to be in R^r not R . For $X = (X_1, \dots, X_r)' \sim F$, define its cross-moments as

$$m_{r.i_1, \dots, i_r} = E X_{i_1} \cdots X_{i_r}, \quad \mu_{r.i_1, \dots, i_r} = E(X_{i_1} - m_{1.i_1}) \cdots E(X_{i_r} - m_{1.i_r})$$

and cross-cumulants as $\kappa_{r.i_1, \dots, i_r}$. (The subscript r . serves to avoid confusion with the univariate notation.)

Fisher noted that the UE of $\kappa_{r.1, \dots, r}$ follows from that of κ_r for the univariate problem. In the same way for t, X in R^r , $\mu_{r.1, \dots, r}$ is just the coefficient of $t_1 \cdots t_r / r!$ in $\mu_r(Y)$ for $Y = t'X$. But for $a(x) : R^r \rightarrow R$ any function, and $Y = a(X)$, the UE of $\mu_r(Y)$ is as given previously with X_j replaced by $a(X_j)$.

For example $\tilde{\mu}_2/(1 - n^{-1})$ implies $\tilde{\mu}_{2.11} = \hat{\mu}_{2.11}/(1 - n^{-1})$ is an UE of $\mu_{2.11}$ where $\hat{\mu}_{2.11} = \mu_{2.11}(F)$.

The same method gives expectations for products of sample cross-moments and UEs for products of cross-moments for the multivariate versions of

$\mu_2, \mu^2, \mu_3, \mu^3, \mu_4, \mu_2^2, \mu^4, \mu_5, \dots$ but fails for the multivariate versions of

$$\mu_2\mu, \mu_3\mu, \mu_2\mu^2, \mu_3\mu_2, \dots \tag{12}$$

For example the coefficient of $2t_1t_2$ in

$$E\hat{\mu}^2 = C_{1^2.2}\mu_2 + C_{1^2.1^2}\mu^2 \text{ is } E\hat{m}_{1.1}\hat{m}_{1.2} = C_{1^2.2}\mu_{2.12} + C_{1^2.1^2}m_{1.1}m_{1.2}$$

and in

$$E(C^{1^2.2}\hat{\mu}_2 + C^{1^2.1^2}\hat{\mu}^2) = \mu^2 \text{ is } E(C^{1^2.2}\hat{\mu}_{2.12} + C^{1^2.1^2}\hat{m}_{1.1}\hat{m}_{1.2}) = m_{1.1}m_{1.2},$$

but the coefficient of $3!t_1t_2t_3$ in $E\widehat{\mu}_2\widehat{\mu} = C_{21.3}\mu_3 + C_{21.21}\mu^2\mu$ is

$$E\left(\sum^3\widehat{\mu}_{2.12}\widehat{m}_{1.3}\right)/3 = C_{21.3}\mu_{3.123} + C_{21.21}\left(\sum^3\mu_{2.12}m_{1.3}\right)/3 \quad (13)$$

where $\sum^3\mu_{2.12}m_{1.3} = \mu_{2.12}m_{1.3} + \mu_{2.23}m_{1.1} + \mu_{2.13}m_{1.2}$.

However from (2.2) one might surmise that

$$E\widehat{\mu}_{2.12}\widehat{m}_{1.3} = C_{21.3}\mu_{3.123} + C_{21.21}\mu_{2.12}m_{1.3}$$

which is correct, as can be shown from the multivariate symmetric function

$$\langle 21 \rangle_{123}^n = \sum^i X_{i1}X_{i2}X_{i3}/(n)_2 \text{ has mean } \langle 21 \rangle_{123}^N = \sum^i x_{i1}x_{i2}x_{i3}/(N)_2.$$

Similarly, the analogous results for (2.1) and (3.1) also hold.

These can be written for fixed i_1, \dots, i_r in $\{1, \dots, r\}$ as

$$E\widehat{m}(\pi)_{r.i_1, \dots, i_r} = \sum_{\pi'}^r \mathbf{B}_{\pi.\pi'} m(\pi')_{r.i_1, \dots, i_r} \quad (14)$$

$$\text{and } E\widehat{m\bar{u}}(\pi)_{r.i_1, \dots, i_r} = \sum_{\pi'}^r \mathbf{C}_{\pi.\pi'} m\bar{u}(\pi')_{r.i_1, \dots, i_r} \quad (15)$$

where for $\pi_1 \leq \pi_2 \leq \dots$, $m(\pi)_{r.i_1, \dots, i_r} = m_{\pi_1.i_1 \dots i_{\pi_1}} m_{\pi_2.j_1 \dots j_{\pi_2}} \dots$ where $j_k = i_{k+\pi_1}$, and $\mu(\pi)_{r.i_1, \dots, i_r}$ is similarly defined,

$$\mathbf{B}_{\pi.\pi'} = P(\pi')^{-1} B_{\pi.\pi'} \sum^{P(\pi)}$$

and

$$\mathbf{C}_{\pi.\pi'} = P(\pi')^{-1} C_{\pi.\pi'} \sum^{P(\pi')}$$

are now operators. For example

$$m(12)_{3.ijk} = m_{1.i}m_{2.jk} \quad (16)$$

$$\begin{aligned} \text{so } E\widehat{m}_{1.i}\widehat{m}_{2.jk} &= \mathbf{B}_{12.3}m_{3.ijk} + \mathbf{B}_{12.12}m_{1.i}m_{2.jk} + \mathbf{B}_{12.13}m_{1.i}m_{2.j}m_{3.k}, \\ &= n^{-2}(\lambda_2 N m_{3.ijk} + \lambda_{11} N^2 m_{1.i}m_{2.jk}) \end{aligned} \quad (17)$$

Similarly $m(1^3)_{3.ijk} = m_{1.i}m_{2.j}m_{3.k}$ so

$$\begin{aligned} E\widehat{m}_{1.i}\widehat{m}_{2.j}\widehat{m}_{3.k} &= \mathbf{B}_{1^3.3}m_{3.ijk} + \mathbf{B}_{1^3.12}m_{1.i}m_{2.jk} + \mathbf{B}_{1^3.13}m_{1.i}m_{1.j}m_{1.k} \\ &= n^{-3}(\lambda_3 N m_{3.ijk} + \lambda_{21} N^2 \sum^3 m_{1.i}m_{2.jk} + \lambda_{13} N^3 m_{1.i}m_{1.j}m_{1.k}). \end{aligned} \quad (18)$$

Note that (3.7) involves $\sum^3 m_{1.i}m_{2.jk}$ but (3.6) only $m_{1.i}m_{2.jk}$.

Set $m_{(i_1 \dots i_r)}$

$$= \{m(\pi)_{r.j_1 \dots j_r} : \pi \text{ partition of } r, (j_1 \dots j_r) \text{ a permutation of } (i_1 \dots i_r)\} \quad (19)$$

For example

$$m_{(ijk)} = (m_{3.ijk}, m_{1.i}m_{2.jk}, m_{1.j}m_{2.ki}, m_{1.k}m_{2.ij}, m_{1.i}m_{1.j}m_{1.k})' \quad (20)$$

So $m_{(i_1 \dots i_r)}$ has dimension $N_r = \sum_{\pi}^r P(\pi)$. Then (4.3), (4.4) can be written as

$$E\widehat{m}_{(i_1 \dots i_r)} = \mathbf{B}_r m_{(i_1 \dots i_r)}, \quad E\widehat{mu}_{(i_1 \dots i_r)} = \mathbf{C}_r mu_{(i_1 \dots i_r)}, \quad (21)$$

where $\mathbf{B}_r, \mathbf{C}_r$ are $N_r \times N_r$.

This form is most useful for $(i_1 \dots i_r) = (1 \dots r)$: other cases where 2 or more of $i_1 \dots i_r$ are equal could have their dimension reduced. But we shall achieve this shortly anyway from (4.10).

Because N_r is also the number of multivariate symmetric functions, the proof of the univariate invariance principle given in Section 7 extends immediately to prove **the multivariate invariance principle** :

$$\mathbf{B}_r(N, n) = \mathbf{B}_r \text{ and } \mathbf{C}_r(N, n) = \mathbf{C}_r \text{ satisfy} \quad (22)$$

$$\mathbf{B}_r(N, n)^{-1} = \mathbf{B}_r(n, N) \text{ and } \mathbf{C}_r(N, n)^{-1} = \mathbf{C}_r(n, N). \quad (23)$$

So $\mathbf{B}_r(n, N)\widehat{m}_{(i_1 \dots i_r)}$ is an UE of $m_{(i_1 \dots i_r)}$,
and $\mathbf{C}_r(n, N)\widehat{mu}_{(i_1 \dots i_r)}$ is an UE of $mu_{(i_1 \dots i_r)}$.

That is,

$$\sum_{\pi'}^r P(\pi')^{-1} B_{\pi.\pi'} \sum_{\pi'}^{P(\pi')} \widehat{m}(\pi')_{r.i_1 \dots i_r} \text{ is an UE of } m(\pi)_{r.i_1 \dots i_r} \quad (24)$$

and

$$\sum_{\pi'}^r P(\pi')^{-1} C_{\pi.\pi'} \sum_{\pi'}^{P(\pi')} \widehat{mu}(\pi')_{r.i_1 \dots i_r} \text{ is an UE of } mu(\pi)_{r.i_1 \dots i_r}. \quad (25)$$

For example letting $\bar{\lambda}_{\pi}$ denote λ_{π} with (N, n) reversed, by (4.6),

$$N^{-2}(\bar{\lambda}_2 n \widehat{m}_{3.ijk} + \bar{\lambda}_{11} N^2 \widehat{m}_{1.i} \widehat{m}_{2.jk}) \text{ is an UE of } m_{1.i} m_{2.jk},$$

$$\text{and } N^{-3}(\bar{\lambda}_3 n \widehat{m}_{3.ijk} + \bar{\lambda}_{21} n^2 \sum^3 \widehat{m}_{1.i} \widehat{m}_{2.jk} + \bar{\lambda}_{13} n^3 \widehat{m}_{1.i} \widehat{m}_{1.j} \widehat{m}_{1.k})$$

is an UE of $m_{1.i} m_{2.j} m_{3.k}$.

Note 4.1 As in (2.4), we can also write $\mathbf{B}_r = \mathbf{B}_r(n, N)$ in the form

$$\mathbf{B}_r = \mathbf{D}_{rn}^{-1} \mathbf{A}_r \mathbf{D}_{rN} \text{ where } \mathbf{D}_{rN} = \text{diag}(N^{q_1}, \dots, N^{q_{N_r}}) \quad (26)$$

and $q_k = q(\pi_k)$ if the k th element of $m_{(i_1 \dots i_r)}$ is $m(\pi_k)_{r.i_1 \dots i_r}$.

Taking $m_{(i)} = m_{1,i}$, $m_{(ij)} = (m_{2,ij}, m_{1,i} m_{1,j})'$ and $m_{(ijk)}$ as in (4.9) this gives $\mathbf{D}_{1N} = N$, $\mathbf{D}_{2N} = \text{diag}(N, N^2)$, $\mathbf{D}_{3N} = \text{diag}(N, N^2, N^2, N^2, N^3)$,

$$\mathbf{A}_1 = \lambda_1, \mathbf{A}_2 = \begin{pmatrix} \lambda_1 & \\ \lambda_2 & \lambda_{11} \end{pmatrix}, \mathbf{A}_3 = \begin{bmatrix} \lambda_1 & & & & \\ \lambda_2 & \lambda_{11} & & & \\ \lambda_2 & 0 & \lambda_{11} & & \\ \lambda_2 & 0 & 0 & \lambda_{11} & \\ \lambda_3 & \lambda_{12} & \lambda_{12} & \lambda_{12} & \lambda_{1^3} \end{bmatrix}$$

As $N_4 = 1 + 4 + 3 + 6 + 1 = 15$, it is not practical (or necessary) to write out any other \mathbf{A}_r in full.

□

If $i_1 = \dots = i_r$ then (4.3), (4.4), (4.13), (4.14) reduce to the univariate formulas. However to apply them at the intermediate level when between 2 and $r - 1$ of i_1, \dots, i_r are equal, one needs to be clear what $\sum^{P(\pi)} m(\pi)_{r.i_1 \dots i_r}$ means, and so what $\sum^{P(\pi)} mu(\pi)_{r.i_1, \dots, i_r}$ means. This can be done by reinterpreting the expressions for $ES_{i_1} \dots S_{i_r}$ in Appendix E.

For example the contribution from $\pi = (13)$ to $ES_{i_1} \dots S_{i_4}$ is $\lambda(13) \sum^4 s_{i_1+i_2+i_3} s_{i_4}$, and to $ES_{i_1}^2 S_{i_2} S_{i_3}$ is $\lambda(13) (\sum^2 s_{2i_1+i_2} s_{i_3} + 2s_{i_1+i_2+i_3} s_{i_1})$,

$$\text{so } \sum^{P(13)} m(13)_{4.i_1 i_1 i_2 i_3} = \sum^2 m_{3.2i_1+i_2} m_{1.i_3} + 2m_{3.i_1+i_2+i_3} m_{1.i_1}.$$

In this way the expressions in Appendix E can be interpreted to give $E\widehat{m}(\pi)_{r.i_1 \dots i_r}$ or an UE for $m(\pi)_{r.i_1 \dots i_r}$:

Set $\mathbf{i}_a = (i_1, \dots, i_a)$, $\mathbf{j}_b = (j_1, \dots, j_b), \dots$. Then

$$\begin{aligned} En\widehat{m}_{a.i_a} &= \lambda_1 N m_{a.i_a}, \\ En^2\widehat{m}_{a.i_a}\widehat{m}_{b.j_b} &= \lambda_2 N m_{a+b.i_a j_b} + \lambda_{11} N^2 m_{a.i_a} m_{b.j_b}, \\ En^3\widehat{m}_{a.i_a}\widehat{m}_{b.j_b}\widehat{m}_{c.k_c} &= \lambda_3 N m_{a+b+c.i_a j_b k_c} \\ &\quad + \lambda_{21} N^2 \sum^2 m_{a+b.i_a j_b} m_{c.k_c} + \lambda_{1^3} N^3 m_{a.i_a} m_{b.j_b} m_{c.k_c}, \\ En^3\widehat{m}_{a.i_a}^2\widehat{m}_{b.j_b} &= \lambda_3 N m_{2m+b.i_a i_a j_b} \\ &\quad + \lambda_{21} N^2 (m_{2a.i_a i_a} m_{b.j_b} + 2m_{a+b.i_a j_b} m_{a.i_a}) + \lambda_{1^3} N^3 m_{a.i_a}^2 m_{b.j_b}, \end{aligned}$$

and so on.

We end this section with UEs for multinomial parameters.

Example 4.1 Suppose F is multinomial $_k(N, \mathbf{p})$, that is for $1 \leq i \leq k$, Np_i of $\{x_1, \dots, x_N\}$ equal e_{ik} , the i th unit vector in R^k .

Take $2 \leq r \leq k$. Then $m_{r,i_1 \dots i_r} = \begin{cases} 0 & \text{if any two of } i_1, \dots, i_r \text{ differ.} \\ p_{i_1} & \text{if } i_1 = \dots = i_r. \end{cases}$

Set $\widehat{p}_i = \widehat{m}_{1,i}$.

Taking $a = 1, b = 2, \dots, e = 5$ in Appendix E, discarding all terms with $+$ as a subscript, and setting $S_a = n\widehat{p}_a, s_{/a} = Np_a$, we have

$$E\widehat{p}_1 \cdots \widehat{p}_r = n^{-r} \lambda_{1^r} N^r p_1 \cdots p_r \text{ for } r \geq 1,$$

$$\begin{aligned} E\widehat{p}_1^2 \widehat{p}_2 \cdots \widehat{p}_{r-1} &= n^{-r} (\lambda_{1^{r-2}2} + \lambda_{1^r} N p_1) N^{r-1} p_1 \cdots p_{r-1} \text{ for } r \geq 3, \\ E\widehat{p}_1^3 \widehat{p}_2 \cdots \widehat{p}_{r-2} &= n^{-r} (\lambda_{1^{r-3}3} + 3\lambda_{1^{r-2}2} N p_1 + \lambda_{1^r} N^2 p_1^2) N^{r-2} p_1 \cdots p_{r-2}, \\ E\widehat{p}_1^2 \widehat{p}_2^2 &= n^{-4} (\lambda_{22} + \lambda_{1^2 2} N (p_1 + p_2) + \lambda_{1^4} N^2 p_1 p_2) N^2 p_1 p_2, \\ E\widehat{p}_1^4 \widehat{p}_2 &= n^{-5} (\lambda_{14} + 4\lambda_{1^2 3} N p_1 + 3\lambda_{1^2 2} N p_1 + 6\lambda_{1^3 2} N^2 p_1^2 + \lambda_{1^5} N^3 p_1^3) N^2 p_1 p_2, \\ E\widehat{p}_1^3 \widehat{p}_2^2 &= n^{-5} (\lambda_{23} + \lambda_{1^2 3} N p_2 + 3\lambda_{1^2 2} N p_1 + \lambda_{1^3 2} N^2 (p_1^2 + 3p_1 p_2) \\ &\quad + \lambda_{1^5} N^3 p_1^2 p_2) N^2 p_1 p_2, \\ E\widehat{p}_1^5 \widehat{p}_2 &= n^{-6} (\lambda_6 + \lambda_{15} + 5\lambda_{1^2 4} N p_1 + 10\lambda_{1^2 3} N p_1 + 10\lambda_{1^3 3} N^2 p_1^2, \\ &\quad + 15\lambda_{1^2 2 2} N^2 p_1^2 + 10\lambda_{1^4 2} N^3 p_1^3 + \lambda_{1^5} N^4 p_1^4) N^2 p_1 p_2, \\ E\widehat{p}_1^4 \widehat{p}_2^2 &= n^{-6} (\lambda_{24} + \lambda_{1^2 4} N p_2 + 4\lambda_{1^2 3} N p_1 + 3\lambda_{23} N p_1 + 4\lambda_{1^3 3} N^2 p_1 p_2 \\ &\quad + 3\lambda_{1^2 2 2} N^2 p_1 (2p_1 + p_2) + \lambda_{1^4 2} N^3 p_1^2 (p_1 + 6p_2) + \lambda_{1^6} N^4 p_1^3 p_2) N^2 p_1 p_2, \\ E\widehat{p}_1^3 \widehat{p}_2^3 &= n^{-6} (\lambda_{32} + 3\lambda_{1^2 3} N (p_1 + p_2) + \lambda_{1^3 3} N^2 (p_1^2 + p_2^2) + 9\lambda_{1^2 2 2} N^2 p_1 p_2 \\ &\quad + 3\lambda_{1^4 2} N^3 p_1 p_2 (p_1 + p_2) + \lambda_{1^6} N^4 p_1^2 p_2^2) N^2 p_1 p_2, \\ E\widehat{p}_1^4 \widehat{p}_2 \widehat{p}_3 &= n^{-6} (\lambda_{1^2 4} + 4\lambda_{1^3 3} N p_1 + 3\lambda_{1^2 2 2} N p_1 + 6\lambda_{1^4 2} N^2 p_1^2 \\ &\quad + \lambda_{1^6} N^3 p_1^3) N^3 p_1 p_2 p_3, \\ E\widehat{p}_1^3 \widehat{p}_2^2 \widehat{p}_3 &= n^{-6} (\lambda_{1^2 3} + \lambda_{1^3 3} N p_2 + 3\lambda_{1^2 2 2} N p_1 + \lambda_{1^4 2} N^2 p_1 (p_1 + p_2) \\ &\quad + \lambda_{1^6} N^3 p_1^2 p_2) N^3 p_1 p_2 p_3, \\ E\widehat{p}_1^2 \widehat{p}_2^2 \widehat{p}_3^2 &= n^{-6} (\lambda_{23} + \lambda_{1^2 2 2} N (p_1 + p_2 + p_3) + \lambda_{1^4 2} N^2 (p_1 p_2 + p_1 p_3 + p_2 p_3), \\ &\quad + \lambda_{1^6} N^3 p_1 p_2 p_3) N^3 p_1 p_2 p_3, \\ E\widehat{p}_1^2 \widehat{p}_2^2 \widehat{p}_3 \widehat{p}_4 &= n^{-6} (\lambda_{1^2 2 2} + \lambda_{1^4 2} N (p_1 + p_2) + \lambda_{1^6} N^2 p_1 p_2) N^4 p_1 p_2 p_3 p_4. \end{aligned}$$

If the expression for $E\widehat{p}_{\pi_1} \widehat{p}_{\pi_2} \cdots$ is $a_{Nn}(\pi, p)$, then $a_{nN}(\pi, \widehat{p})$ is an UE for $p_{\pi_1} p_{\pi_2} \cdots$.

This generates a new set of matrices $A = A(N, n)$ satisfying the invariance principle

$$A(N, n)^{-1} = A(n, N) :$$

from $(1 \dots r)$, $A = \lambda_1^r$;

$$\text{from } (1 \dots r-1), (1^2 2 \dots r-1), A = \begin{pmatrix} \lambda_{1^{r-1}} & & \\ & \lambda_{1^{r-2}} & \\ & & \lambda_{1^r} \end{pmatrix};$$

$$\text{from } (1 \dots r-2), (1^2 2 \dots r-2), (1^3 2 \dots r-2), A = \begin{pmatrix} \lambda_{1^{r-2}} & & & \\ \lambda_{1^{r-3}} & \lambda_{1^{r-1}} & & \\ & & \lambda_{1^{r-3}} & \lambda_{1^r} \\ & & & 3\lambda_{1^{r-2}2} \end{pmatrix};$$

$$\text{from } (12, 1^2 2, 1^3 2), A = \begin{pmatrix} \lambda_2 & & & \\ \lambda_{12} & \lambda_{1^3} & & \\ & & \lambda_{1^3} & \\ \lambda_{13} & 3\lambda_{1^2 2} & \lambda_{1^4} & \end{pmatrix};$$

$$\text{from } (12, 1^2 2, 12^2, 1^2 2^2), A = \begin{pmatrix} \lambda_2 & & & & \\ \lambda_{12} & \lambda_{1^3} & & & \\ \lambda_{12} & 0 & \lambda_{1^3} & & \\ \lambda_{22} & \lambda_{1^2 2} & \lambda_{1^2 2} & \lambda_{1^4} & \end{pmatrix};$$

$$\text{from } (12, 1^2 2, 1^3 2, 1^4 2), A = \begin{pmatrix} \lambda_2 & & & & \\ \lambda_{12} & \lambda_{13} & & & \\ \lambda_{13} & 3\lambda_{1^2 2} & & \lambda_{1^4} & \\ \lambda_{14} & 4\lambda_{1^2 3} + \lambda_{12^2} & 6\lambda_{1^3 2} & \lambda_{1^5} & \\ & & & & \end{pmatrix};$$

from $(12, 12^2, 1^2 2, 1^3 2, 1^2 2^2, 1^3 2^2)$,

$$A = \begin{pmatrix} \lambda_2 & & & & & & \\ \lambda_{12} & \lambda_{13} & & & & & \\ \lambda_{12} & 0 & \lambda_{13} & & & & \\ \lambda_{13} & 0 & 3\lambda_{1^2 2} & \lambda_{1^4} & & & \\ \lambda_{22} & \lambda_{1^2 2} & \lambda_{1^2 2} & 0 & \lambda_{1^4} & & \\ \lambda_{23} & \lambda_{1^2 3} & 3\lambda_{12^2} & \lambda_{1^3 2} & 3\lambda_{1^3 2} & \lambda_{1^5} & \end{pmatrix};$$

$$\text{from } (123, 12^2 3, 1^2 23, 1^2 2^2 3), A = \begin{pmatrix} \lambda_{13} & & & & \\ \lambda_{1^2 2} & \lambda_{1^4} & & & \\ \lambda_{1^2 2} & 0 & \lambda_{1^4} & & \\ \lambda_{12^2} & \lambda_{1^3 2} & \lambda_{1^3 2} & \lambda_{1^5} & \end{pmatrix};$$

and so on. \square

Raghunandan and Srinivasan (1973) give multivariate analogs of $E\widehat{\mu}_r$ for $5 \leq r \leq 8$ in terms of symmetric functions with tables to express these in terms of noncentral moments, and $E(\overline{X} - \mu)^r$ for $r = 5, 6$. The latter agree with Sukhatme except have $-5e_3$ where Sukhatme has $e_2 - 6e_3$ in the coefficient of μ_3^2 in $E(\overline{X} - \mu)^6$. Sukhatme's version is the correct one since $g_9 n^6$ is $C_{33} = C_3 \otimes C_3$ in the notation of Dwyer and Tracy. They also give the multivariate analogs of the UE of $E(\overline{X} - \mu)^4$ in terms of $\widehat{\mu}_4$ and $\sum_{i \neq j} (X_i - \overline{X})^2 (X_j - \overline{X})^2$, and the UE of $E(\overline{X} - \mu)^5$ in terms of $\widehat{\mu}_5$ and $\sum_{i \neq j} (X_i - \overline{X})^3 (X_j - \overline{X})^2$.

It would be useful if Pierce's transformations (to obtain results for finite N from results for $N = \infty$) were extended to cover multivariate problems like (3.2).

5 UEs of products of central moments

Here we express the joint central moments $\mu(\pi)$ of (1.3) as a linear combination of $m(\pi)$ and so obtain

$$E\widehat{\mu}(\pi_1)\widehat{\mu}(\pi_2)\dots$$

and UEs of $\mu(\pi_1)\mu(\pi_2)\dots$ for total order

$$r = r(\pi_1) + r(\pi_2) \dots \leq 6.$$

We derive coefficients $M_{\pi.\pi_-}$ such that

$$\mu(\pi) = \sum_{\pi_-}^r M_{\pi.\pi_-} m u(\pi_-). \quad (27)$$

This enables us to express $\mu(\pi_1)\mu(\pi_2)\dots$ in the form

$$D = \sum_{\pi_-}^r D_{\pi_-} m u(\pi_-). \quad (28)$$

By Section 2, an UE of D is then

$$\tilde{D} = \sum_{\pi'_-}^r D_{\pi'_-}^* \widehat{mu}(\pi'_-) \quad (29)$$

$$\text{where } D_{\pi'_-}^* = \sum_{\pi_-}^r D_{\pi_-} C^{\pi-\pi'_-} \text{ and } (C^{\pi-\pi_-}) = C_{--r}^{-1} \quad (30)$$

We begin with $E\widehat{\mu}(\pi)$ and give the UE of $\mu(\pi)$ in the order

$$\pi = 1^r, 2^r, 3^2, 1^r 2, 1^r 3, 1^r 4, 1^r 2^2, 23, 123.$$

We omit $\pi = 15$ as Sukhatme omitted $\mu_{(51)}$.

As noted in Section 2,

$$\mu(1^r) = E(\bar{X} - \mu)^r = \sum_{\pi_-}^r C_{1^r, \pi_-} mu(\pi_-),$$

$$\text{so in (5.1), } M_{1^r, \pi_-} = C_{1^r, \pi_-} \quad (31)$$

The D^* needed in (5.3) for the UE of $D = \mu(1^r)$ are as follows.

$$\begin{aligned} \mu(1^2) : D_2^* &= (N-n)N^{-1}(n-1)^{-1}, \\ \mu(1^3) : D_3^* &= (N-2n)(N-n)N^{-2}(n-1)_2^{-1}, \\ \mu(1^4) : D_4^* &= -(N-n)n^{-1}(2N^2n - 4N^2 + 6Nn - 3n^3 \\ &\quad - 3n^2)(n-1)_3^{-1}N^{-3}, \end{aligned} \quad (32)$$

$$\begin{aligned} D_{22}^* &= 3(N-n)n^{-1}(N^2n^2 - 4N^2n + 4N^2 - Nn^3 - 6Nn \\ &\quad + 6Nn^2 + 3n^2 - 3n^3)(n-1)_3^{-1}N^{-3}, \\ \mu(1^5) : D_5^* &= -(N-n)(N-2n)n^{-1}(9N^2n - 15N^2 + 20Nn - 8Nn^2 \\ &\quad - 10n^2 - 2n^3)(n-1)_4^{-1}N^{-4}, \\ D_{23}^* &= 10(N-n)(N-2n)n^{-1}(N^2n^2 - 3N^2n + 3N^2 - Nn^3 \\ &\quad - 4Nn - 4Nn + 4Nn^2 - 2n^3 + 2n^2)(n-1)_4^{-1}N^{-4} \end{aligned} \quad (33)$$

$$\begin{aligned}
\mu(1^6) : D_6^* &= (N-n)n^{-3}(-24N^4 + 6N^4n^3 - 84N^4n^2 + 126N^4n \\
&\quad + 120N^3n^3 - 345N^3n^2 + 60N^3n + 45N^3n^4 - 80N^2n^2 \\
&\quad + 50N^2n^4 + 400N^2n^3 - 130N^2n^5 + 75Nn^6 - 225Nn^4 \\
&\quad + 60Nn^3 - 150Nn^5 - 20n^4 + 80n^6 + 5n^7 + 55n^5)(n-1)_5^{-1}N^{-5}, \quad (34) \\
D_{24}^* &= -15(N-n)n^{-3}(24N^4 + 2N^4n^4 - 24N^4n^3 + 82N^4n^2 \\
&\quad - 102N^4n - 2N^3n^5 - 220N^3n^3 + 285N^3n^2 - 60N^3n \\
&\quad + 57N^3n^4 - 3N^2n^6 + 80N^2n^2 + 220N^2n^4 - 320N^2n^3 \\
&\quad - 43N^2n^5 + 3Nn^7 + 6Nn^6 - 90Nn^5 - 60Nn^3 + 165Nn^4 \\
&\quad + 5n^7 + 10n^6 + 20n^4 - 35n^5)(n-1)_5^{-1}N^{-5} \\
D_{32}^* &= 10(N-n)n^{-3}(24N^4 + N^4n^4 - 12N^4n^3 \\
&\quad + 57N^4n^2 - 90N^4n - 5N^3n^5 - 165N^3n^3 + 255N^3n^2 \\
&\quad - 60N^3n + 39N^3n^4 + 8N^2n^6 + 80N^2n^2 + 175N^2n^4 \\
&\quad - 280N^2n^3 - 47N^2n^5 - 4Nn^7 + 24Nn^6 - 75Nn^5 \\
&\quad - 60Nn^3 + 135Nn^4 - 5n^7 + 10n^6 + 20n^4 - 25n^5)(n-1)_5^{-1}N^{-5} \\
D_{23}^* &= 15(24N^4 + N^4n^4 - 12N^4n^3 + 46N^4n^2 - 66N^4n \\
&\quad - 2N^3n^5 + 30N^3n^4 + 180N^3n^2 - 60N^3n - 122N^3n^3 \\
&\quad - 210N^2n^3 + N^2n^6 - 27N^2n^5 + 127N^2n^4 \\
&\quad + 80N^2n^2 + 120Nn^4 - 60Nn^3 - 59Nn^5 + 9Nn^6 \\
&\quad + 10n^6 + 20n^4 - 30n^5)n^{-2}(N-n)(n-1)_5^{-1}N^{-5}.
\end{aligned} \tag{35}$$

Now in the expansion for μ_r in terms of $\{\mu'_i\}$, the coefficient of μ'_r is 1. So in the expansion of $\mu(\pi)$, the coefficient of $m(\pi)$ is 1. So

$$M_{\pi,r} = C_{\pi,r} \tag{36}$$

For example $\mu(2^2) = E\hat{\mu}_2^2 - (E\hat{\mu}_2)^2 = RHS$ (5.1) with

$$\begin{aligned}
M_{2^2,4} &= C_{2^2,4}, \\
M_{2^2,2^2} &= C_{2^2,2^2} - C_{2,2}^2 \\
&= -N(n-1)(N^2n - 3N^2 + 6N - 3n - 3)(N-n)n^{-3}(N-1)^{-2}(N-2)_2^{-1},
\end{aligned}$$

So by (5.4) the UE of $\mu(2^2)$ is (5.3) with

$$D_4^* = 0,$$

$$D_{2^2}^* = 1.$$

$$\text{Since } \mu(2^3) = E\hat{\mu}_2^3 - 3E\hat{\mu}_2^2 E\hat{\mu}_2 + 2(E\hat{\mu}_2)^3,$$

$$\begin{aligned} M_{2^3,24} &= C_{2^3,24} - 3C_{2^2,4}C_{2,2} \\ &= -3N(n-1)(N-n)(N^4n^2 + 5N^4 - 6N^4n \\ &\quad - 2N^3n^3 + 5N^3 + 13N^3n^2 - 53N^2n^2 + 20N^2n \\ &\quad - 45N^2 + 2N^2n^3 + 55N + 30Nn + 35Nn^2 \\ &\quad + 20Nn^3 - 20n - 20n^2 - 20n^3 - 20)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \end{aligned}$$

$$M_{2^3,3^2} = C_{2^3,3^2},$$

$$\begin{aligned} M_{2^3,2^3} &= C_{2^3,2^3} - 3C_{2^2,2^2}C_{2,2} + 2C_{2,2}^3 \\ &= 2N^2(n-1)(N-n)(N^4n^2 + 15N^4 - 12N^4n \\ &\quad - 75N^3 - 2N^3n^3 + 25N^3n^2 + 30N^3n \\ &\quad - 3N^2n^3 + 135N^2 - 139N^2n^2 + 51N^2n \\ &\quad + 167Nn^2 - 144Nn - 105N + 56Nn^3 + 75n - 75n^3 \\ &\quad - 30n^2 + 30)n^{-5}(N-1)^{-3}(N-2)_4^{-1}, \end{aligned}$$

So

$$D_6^* = 0,$$

$$D_{24}^* = 0,$$

$$D_{3^2}^* = 0,$$

$$D_{2^3}^* = 1.$$

$$\begin{aligned} \mu(3^2) : M_{3^2,3^2} &= C_{3^2,3^2} - C_{3,3}^2 \\ &= -N(n-1)_2(N-n)(20N^5 + N^5n^2 - 12N^5n - 100N^4 \\ &\quad + 4N^4n^2 + 42N^4n + 260N^3 - 41N^3n^2 + 24N^3n \\ &\quad + 40N^2n^2 - 460N^2 - 282N^2n + 420Nn + 100Nn^2 \\ &\quad + 440N - 240n - 160 - 80n^2)n^{-5}(N-1)^{-2}(N-2)^{-2}(N-3)_3^{-1}, \end{aligned}$$

$$M_{3^2,2^3} = C_{3^2,2^3},$$

$$D_6^* = 0, D_{24}^* = 0, D_{3^2}^* = 1, D_{2^3}^* = 0$$

$$\begin{aligned} \text{Also } \mu(1^r k) &= E(\bar{x} - \mu)^r (\hat{\mu}_k - E\hat{\mu}_k) \\ &= \sum_{\pi_-}^{r+k} C_{1^r k, \pi_-} m(\pi_-) - \sum_{\pi_-}^r C_{1^r, \pi_-} m(\pi_-) \sum_{\pi_-}^k C_{k, \pi'_-} m(\pi'_-). \end{aligned}$$

So $\mu(12) = M_{12,3}\mu_3$ for $M_{12,3} = C_{12,3}$, as given by (4.9), and

$$D_3^* = (N-n)N^{-1}(n-2)^{-1}.$$

$$\begin{aligned} \mu(1^2 2) : M_{1^2 2, 2^2} &= C_{1^2 2, 2^2} - C_{1^2, 2}C_{2, 2}, \\ &= -N(n-1)(N-n)n^{-3}(N-1)^{-2}(N-2)_2^{-1}(3N^2 - 4Nn \\ &\quad - 6N + 3 + 6n), \end{aligned}$$

$$D_4^* = (N-n)n^{-1}(2N-n-n^2)(n-2)_2^{-1}N^{-2},$$

$$D_{2^2}^* = (N-n)n^{-1}(6N+Nn^2-6Nn-3n+3n^2)(n-2)_2^{-1}N^{-2}.$$

$$\begin{aligned} \mu(1^3 2) : M_{1^3 2, 2^3} &= C_{1^3 2, 2^3} - C_{1^3, 3}C_{2, 2} \\ &= N(n-1)(N-n)(-10N^3 + 3N^3n - 3N^2n^2 \\ &\quad + 15N^2n + 20N^2 - 920n^2 - 36Nn - 10N + 24n^2 \\ &\quad + 12n)n^{-4}(N-1)^{-2}(N-2)_2^{-1}, \end{aligned}$$

$$\begin{aligned}
\mu(1^3 3) : M_{1^3 3, 24} &= C_{1^3 3, 24}, \\
M_{1^3 3, 3^2} &= C_{1^3 3, 3^2} - C_{1^3, 3} C_{3, 3} \\
&= -(N-n)N(n-1)_2(+10N^5 - 33N^4 n - 50N^4 \\
&\quad + 24N^3 n^2 + 130N^3 + 162N^3 n - 303N^2 n - 108N^2 n^2 \\
&\quad - 230N^2 + 132N n^2 + 220N + 270N n - 120n \\
&\quad - 80)n^{-5}(N-1)^{-2}(N-2)^{-2}(N-3)_3^{-1}, \\
M_{1^3 3, 2^3} &= C_{1^3 3, 2^3}, \\
D_6^* &= -(N-n)N^{-3}n^{-3}(12N^2 - 12N^2 n^2 - 27N^2 n \\
&\quad + 3N^2 n^3 - 3N n^4 + 30N n^3 - 12N n + 33N n^2 \\
&\quad - 11n^3 - n^5 + 4n^2 - 16n^4)(n-3)_3^{-1}, \\
D_{24}^* &= 3(N-n)N^{-3}n^{-3}(N^2 n^4 - 3N^2 n^3 - 25N^2 n^2 \\
&\quad + 75N^2 n - 60N^2 + 60N n - N n^5 + 6N n^4 + 40N n^3 \\
&\quad - 105N n^2 + 35n^3 - 20n^2 - 10n^4 - 5n^5)(n-3)_3^{-1}, \\
D_{3^2}^* &= (N-n)N^{-3}n^{-3}(120N^2 + N^2 n^4 - 6N^2 n^3 + 15N^2 n^2 \\
&\quad - 90N^2 n - 120N n - 2N n^5 + 24N n^4 - 60N n^3 + 150N n^2 \\
&\quad - 50n^3 + 40n^2 + 20n^4 - 10n^5)(n-3)_3^{-1}, \\
D_{2^3}^* &= -3(-60N^2 + 3N^2 n^3 - 28N^2 n^2 + 75N^2 n - 3N n^4 + 35N n^3 \\
&\quad - 90N n^2 + 60N n - 10n^4 + 30n^3 - 20n^2)n^{-2}N^{-3}(N-n)(n-3)_3^{-1}.
\end{aligned}$$

$$\begin{aligned}
\mu(14) : M_{14, 23} &= C_{14, 23}, \\
D_5^* &= (n^2 - n - 3)(N-n)n^{-1}N^{-1}(n-4)^{-1}(n-2)^{-1}, \\
D_{23}^* &= -2(n-1)(2n-3)(N-n)n^{-1}N^{-1}(n-4)^{-1}(n-2)^{-1}.
\end{aligned}$$

$$\begin{aligned}
\mu(1^2 4) : M_{1^2 4, 24} &= C_{1^2 4, 24} - C_{1^2, 2} C_{4, 4}, \\
&= -(n-1)N(N-n)(9N^4 n^2 - 39N^4 n + 45N^4 + 39N^3 n^2 \\
&\quad - 81N^3 n - 10N^3 n^3 + 45N^3 + 45N^2 n^2 - 405N^2 \\
&\quad - 6N^2 n^3 + 237N^2 n - 435N n^2 + 495N \\
&\quad + 135N n - 180n + 270n^2 - 180 - 30n^3 \\
&\quad + 70N n^3)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \\
M_{1^2 4, 3^2} &= C_{1^2 4, 3^2}, \\
M_{1^2 4, 2^3} &= C_{1^2 4, 2^3} - C_{1^2, 2} C_{4, 2^2} \\
&= 6(n-1)N(N-n)(2N^4 n^2 - 12N^4 n + 9N^3 n^2 + 15N^4 \\
&\quad - 2N^3 n^3 - 60N^3 + 27N^3 n + 75N^2 - 44N^2 n^2 + 15N^2 n \\
&\quad - 30N n + 9N n^2 - 30N + 26N n^3 + 30n^2 \\
&\quad 30n^3)n^{-5}(N-1)^{-2}(N-2)_4^{-1},
\end{aligned}$$

$$\begin{aligned}
D_6^* &= (-36N^4 n^3 + 8N^4 n^4 + 22N^4 n^2 + 54N^4 n - 24N^4 - N^3 n^6 - 155N^3 n^2 \\
&\quad - 118N^3 n + 3N^3 n^5 + 121N^3 n^3 - 18N^3 n^4 + 72N^3 - N^2 n^6 + N^2 7n^5 \\
&\quad + 60N^2 n + 233N^2 n^3 - 126N^2 n^4 - 5N^2 n^2 + 64N n - 96N n^3 + 4N n^4 \\
&\quad + 32N n^5 + 12N n^6 - 304N n^2 - 112n^5 - 64n^2 + 208n^3 + 168n^4 \\
&\quad - 8n^6)n^{-3}(N-n)(n-2)_4^{-1}(N-3)^{-1}N^{-4}, \tag{37}
\end{aligned}$$

$$\begin{aligned}
D_{24}^* = & (110N^4n^4 - 204N^4n^3 + 450N^4n - 21N^4n^5 + N^4n^6 - 360N^4 \\
& + 1080N^3 + 6N^3n^6 - 6N^3n^5 - 186N^3n^4 - 675N^3n^2 - 690N^3n \\
& + 687N^3n^3 - 408N^2n^4 + 423N^2n^3 - 3N^2n^6 + 135N^2n^5 - 615N^2n^2 \\
& + 900N^2n + 240Nn^5 + 2400Nn^3 - 468Nn^4 + 960Nn - 108Nn^6 \\
& - 3600Nn^2 - 960n^2 + 2160n^3 - 1320n^4 \\
& + 120n^6)n^{-3}(N - n) \\
& (n - 2)_4^{-1}(N - 3)^{-1}N^{-4},
\end{aligned} \tag{38}$$

$$\begin{aligned}
D_{32}^* = & -2(10N^4n^4 - 57N^4n^3 + 65N^4n^2 + 90N^4n - 120N^4 - 310N^3n^2 \\
& + 39N^3n^4 + 360N^3 - 50N^3n - 10N^3n^5 + 71N^3n^3 - 2N^2n^6 \\
& + 2N^2n^5 + 300N^2n + 184N^2n^3 - 45N^2n^4 - 295N^2n^2 + 720Nn^3 \\
& - 124Nn^4 + 28Nn^6 + 320Nn - 64Nn^5 - 1040Nn^2 + 160n^5 \\
& - 320n^2 + 560n^3 - 360n^4)n^{-3}(N - n)(n - 2)_4^{-1}(N - 3)^{-1}N^{-4},
\end{aligned} \tag{39}$$

$$\begin{aligned}
D_{23}^* = & 6(N - n)n^{-2}(2N^4n^4 - 18N^4n^3 + 64N^4n^2 - 105N^4n + 60N^4 \\
& - 180N^3 + 205N^3n - 2n^5N^3 + 15N^3n^4 - 31N^3n^3 \\
& - 32N^3n^2 + 141N^2n^3 + 90N^2n - 228N^2n^2 - 27N^2n^4 \\
& + 24Nn^5 - 160Nn + 320Nn^2 - 72Nn^4 - 72Nn^3 + 200n^4 \\
& - 40n^5 + 160n^2 - 320n^3)(n - 2)_4^{-1}(N - 3)^{-1}N^{-4},
\end{aligned} \tag{40}$$

$$\tag{41}$$

$$\text{Also } \mu(1^r k^2) = E(\overline{X} - \mu)^r (\hat{\mu}_k^2 - 2\hat{\mu}_k E\hat{\mu}_k + (E\hat{\mu}_k)^2)$$

$$\begin{aligned}
\mu(12^2) : M_{12^2, 23} & = C_{12^2, 23} - 2C_{12, 3}C_{2, 2} \\
& = -2(n - 1)N(N - n)(3N^3n - 5N^3 - 4N^2n^2 \\
& \quad + 10N^2 + 8Nn^2 - 11Nn - 5N + 2n^2 \\
& \quad + 2n)n^{-4}(N - 1)^{-2}(N - 2)_3^{-1}, \\
D_5^* & = -(n - 1)n^{-1}N^{-1}(N - n)(n - 4)^{-1}(n - 2)^{-1}, \\
D_{23}^* & = 2(n^2 - 3n + 1)n^{-1}N^{-1}(N - n)(n - 4)^{-1}(n - 2)^{-1}.
\end{aligned}$$

$$\begin{aligned}
\mu(1^2 2^2) : M_{1^2 2^2, 24} &= C_{1^2 2^2, 24} - 2C_{1^2 2, 4} C_{2, 2} \\
&= (n-1)N(N-n)(-12N^4 n + N^4 n^2 + 15N^4 + 31N^3 n^2 \\
&\quad - 35N^3 n - N^3 n^3 + 15N^3 - 22N^2 n^3 - 135N^2 \\
&\quad - 29N^2 n^2 + 90N^2 n + 65Nn + 61Nn^3 + 165N - 109Nn^2 \\
&\quad + 10n^2 - 60n - 60 + 10n^3)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \\
M_{1^2 2^2, 3^2} &= C_{1^2 2^2, 3^2}, \\
M_{1^2 2^2, 2^3} &= C_{1^2 2^2, 2^3} - 2C_{1^2 2, 2^2} C_{2, 2} + C_{1^2, 2} C_{2, 2}^2 \\
&= -N^2(n-1)(N-n)(-21N^4 n + 30N^4 + N^4 n^2 + 44N^3 n^2 \\
&\quad - N^3 n^3 - 150N^3 + 45N^3 n + 270N^2 - 25N^2 n^3 + 93N^2 n \\
&\quad - 194N^2 n^2 + 127Nn^3 - 210N - 237Nn + 178Nn^2 + 60 \\
&\quad + 120n - 5n^2 - 125n^3)n^{-5}(N-1)^{-3}(N-2)_4^{-1}, \\
D_6^* &= -(N-n)(n-1)(+2Nn^3 - 4Nn^2 - 10Nn + 8N - 4n \\
&\quad + 7n^2 + 8n^3 - 3n^4)N^{-2}n^{-3}(n-2)_4^{-1}, \\
D_{24}^* &= (N-n)(120N + 16Nn^4 - 102Nn^3 + 200Nn^2 \\
&\quad - 150Nn - 145n^3 + 105n^2 - 60n + n^5 - 2n^6 \\
&\quad + 53n^4)N^{-2}n^{-3}(n-2)_4^{-1}, \\
D_{3^2}^* &= 2(N-n)(-40N + Nn^5 - 9Nn^4 + 31Nn^3 - 45Nn^2 \\
&\quad + 30Nn + 40n^3 - 25n^2 + 20n + 8n^5 - n^6 \\
&\quad - 26n^4)N^{-2}n^{-3}(n-2)_4^{-1}, \\
D_{2^3}^* &= (-120N - 17Nn^4 + 92Nn^3 - 208Nn^2 + 210Nn + Nn^5 \\
&\quad + 60n - 48n^4 - 120n^2 + 6n^5 + 126n^3)n^{-2}N^{-2}(N-n)(n-2)_4^{-1}. \\
\text{Also, } \mu(23) : M_{23, 23} &= C_{23, 23} - C_{2, 2} C_{3, 3}, \\
&= -2(N-n)N^2(n-1)_2(2N^2 n - 5N^2 + 10N - \\
&\quad - n)n^{-4}(N-1)^{-2}(N-2)_3^{-1}, \\
D_5^* &= 0, \quad D_{23}^* = 1.
\end{aligned}$$

$$\text{and } \mu(123) = E(\overline{X} - \mu)(\hat{\mu}_2 \hat{\mu}_3 - \hat{\mu}_2 E \hat{\mu}_3 - \hat{\mu}_3 E \hat{\mu}_2 + E \hat{\mu}_2 E \hat{\mu}_3)$$

$$\begin{aligned}
\text{so } M_{123.24} &= C_{123.24} - C_{13.4}C_{2.2} \\
&= -(N-n)N(n-1)_2(-15N^4 + 9N^4n - 15N^3 - 2N^3n - 13N^3n^2 \\
&\quad + 135N^2 + 42N^2n^2 + 9N^2n - 165N - 130Nn - 35Nn^2 + 30n^2 \\
&\quad + 90n + 60)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \\
M_{123.32} &= C_{123.32} - C_{12.3}C_{3.3} \\
&= -(N-n)N(n-1)_2(-10N^5 + 4N^5n - 3N^4n + 50N^4 - 5N^4n^2 \\
&\quad - 130N^3 + 18N^3n^2 - 62N^3n + 230N^2 + 13N^2n^2 \\
&\quad - 90Nn^2 + 195N^2n - 230Nn - 220N + 80 + 120n \\
&\quad + 40n^2)n^{-5}(N-1)^{-2}(N-2)^{-2}(N-3)_3^{-1}, \\
M_{123.23} &= C_{123.23} - C_{13.22}C_{2.2} \\
&= 3N^2(n-1)_2(N-n)(-10N^2 + 4N^2n - 5Nn^2 - Nn \\
&\quad + 30N + 15n^2 - 25n - 20)n^{-5}(N-1)_5^{-1}, \\
D_6^* &= -(1+n)(n-1)^2(N-n)n^{-3}N^{-1}(n-3)^{-1}(n-5)^{-1}, \\
D_{24}^* &= (n^4 - 10n^2 - 15)n^{-3}N^{-1}(N-n)(n-3)^{-1}(n-5)^{-1}, \\
D_{32}^* &= (n^4 - 5n^3 + 5n^2 + 5n + 10)n^{-3}N^{-1}(N-n)(n-3)^{-1}(n-5)^{-1}, \\
D_{23}^* &= -3(N-n)N^{-1}n^{-2}(n^3 - 5n^2 + 5n - 5)(n-3)^{-1}(n-5)^{-1}.
\end{aligned} \tag{42}$$

Now we consider products of **two** central moments $\mu(\pi_1)\mu(\pi_2)$, in the order

$$(\pi_1.\pi_2) = (1^2.1^2), (1^2.1^3), (1^2.1^4), (1^3.1^3), (1^2.12), (1^2.1^22), (1^2.2^2), (1^3.12), (12.12),$$

$$\begin{aligned}
\mu(1^2)^2 &= C_{1^2.2}^2 \mu_2^2 \text{ so} \\
D_4^* &= 0, \\
D_{2^2}^* &= (N-n)^2(N^2n^2 - 3N^2n + 3N^2 - 2Nn^2 + 3N + 3n^2 \\
&\quad - 3n)n^{-1}(N-1)^{-1}N^{-3}(n-1)_3^{-1} \\
\mu(1^2)\mu(1^3) &= C_{1^2.2}C_{1^3.3}\mu_2\mu_3 \text{ so} \\
D_5^* &= 0, \\
D_{2^3}^* &= (N-n)^2(N-2n)(N^2n^2 - 5Nn^2 - 2N^2n + 2N^2 + 10N + 10n^2 \\
&\quad - 10n)n^{-1}(N-1)^{-1}N^{-4}(n-1)_4^{-1}, \\
\mu(1^2)\mu(1^4) &= C_{1^2.2}C_{1^4.4}\mu_2\mu_4 + C_{1^2.2}C_{1^4.2^2}\mu_2^3 \\
\text{so } D_6^* &= D_{3^2}^* = 0, \\
D_{2^4}^* &= -(N-n)^2(-63N^4n^3 + 190N^4n^2 - 225N^4n + 8N^4n^4 + 60N^4 \\
&\quad + 60N^3 + 640N^3n^2 - 525N^3n + 98N^3n^4 - 12N^3n^5 - 345N^3n^3 \\
&\quad + 198N^2n^4 + 3N^2n^6 - 45N^2n^5 - 750N^2n^3 + 1140N^2n^2 - 240N^2n \\
&\quad - 9Nn^5 - 990Nn^3 + 405Nn^4 + 18Nn^6 + 360Nn^2 - 90n^5 - 180n^3 \\
&\quad + 315n^4 - 45n^6)N^{-5}(n-1)_5^{-1}(N-1)^{-1}n^{-3}, \\
D_{2^3}^* &= (-60N^2 + n^5N^2 - 11N^2n^4 + 44N^2n^3 - 82N^2n^2 + 90N^2n - 60N \\
&\quad + 21Nn^4 - 96Nn^2 + 180Nn - 24Nn^3 - 3Nn^5 + 60n - 48n^4 \\
&\quad - 120n^2 + 6n^5 + 126n^3)N^{-2}(N-n)(n-2)_4^{-1}(N-1)^{-1}n^{-2}, \\
\mu(1^3)^2 &= C_{1^3.3}^2 \mu_3^2 \text{ so } D_6^* = D_{2^4}^* = D_{2^3}^* = 0, \\
D_{3^2}^* &= (N-2n)^2(N-n)^2(N^3n^4 + 25N^3n^2 - 40N^3 - 10N^3n - 8N^3n^3 \\
&\quad + 8N^2n^3 - 26N^2n^2 + 70N^2n - 4N^2n^4 - 120N^2 + 14Nn^4 - 4Nn^3 \\
&\quad - 70Nn^2 - 80N + 220Nn + 80n - 100n^2 + 40n^3 \\
&\quad - 20n^4)n^{-3}(N-1)_2^{-1}(n-1)_5^{-1}N^{-5}, \\
\mu(1^2)\mu(1^2) &= C_{1^2.2}C_{1^2.3}\mu_2\mu_3 \text{ so } D_5^* = 0, \\
D_{2^3}^* &= (N-n)^2(N^2n^2 - 2N^2n + 2N^2 + 10N - 5Nn^2 + 10n^2 \\
&\quad - 10n)n^{-1}(N-1)^{-1}N^{-3}(n-2)_3^{-1}, \\
\mu(1^2)\mu(1^2) &= C_{1^2.2}C_{1^2.4}\mu_2\mu_4 + C_{1^2.2}C_{1^2.2^2}\mu_2^3 \\
\text{so } D_6^* &= D_{3^2}^4 = 0, \\
D_{2^4}^* &= -(N-n)^2(60N^3 + 2N^3n^4 - 21N^3n^3 + 70N^3n^2 - 105N^3n \\
&\quad - 59N^2n^3 + 60N^2 - 285N^2n + 17N^2n^4 + 190N^2n^2 - n^5N^2 \\
&\quad - 135Nn^3 - 120Nn + 3Nn^4 - 6n^5N + 330Nn^2 + 60n^2 + 15n^5 \\
&\quad - 105n^3 + 30n^4)(n-2)_4^{-1}N^{-4}(N-1)^{-1}n^{-3}, \\
D_{2^3}^* &= (60N^3 + N^3n^4 - 12N^3n^3 + 50N^3n^2 - 90N^3n + 60N^2 + 30N^2n^2 \\
&\quad - 120N^2n + 3N^2n^3 - 3Nn^4 - 57Nn^3 - 120Nn + 210Nn^2 \\
&\quad - 90n^3 + 60n^2 + 30n^4)(N-n)^2(n-2)_4^{-1}N^{-4}(N-1)^{-1}n^{-2}, \\
\mu(1^2)\mu(2^2) &= C_{1^2.2}C_{2^2.4}\mu_2\mu_4 + C_{1^2.2}C_{2^2.2^2}\mu_2^3, \\
\text{so } D_6^* &= D_{3^2}^* = 0, \\
D_{2^4}^* &= -(N-n)^2(-60N + 2Nn^5 - 17Nn^4 + 49Nn^3 - 55Nn^2 \\
&\quad + 45Nn - 60 + n^4 - 145n^2 + 105n + 53n^3 \\
&\quad - 2n^5)N^{-2}(N-1)^{-1}(n-2)_4^{-1}n^{-3}, \\
D_{2^3}^* &= (-60N^2 + N^2n^5 - 11N^2n^4 + 44N^2n^3 - 82N^2n^2 + 90N^2n - 60N \\
&\quad + 21Nn^4 - 96Nn^2 + 180Nn - 24Nn^3 - 3n^5N + 60n - 48n^4
\end{aligned}$$

$$\begin{aligned}
\mu(1^3)\mu(12) &= C_{1^3,3}C_{12,3}\mu_3^2 \\
\text{so } D_6^* &= D_{24}^* = D_{23}^* = 0, \\
D_{32}^* &= (N-2n)(N-n)^2(N^3n^4 - 40N^3 + 25N^3n^2 - 8N^3n^3 \\
&\quad - 10N^3n - 26N^2n^2 + 70N^2n + 8N^2n^3 - 120N^2 - 4N^2n^4 - 70Nn^2 \\
&\quad - 80N + 14Nn^4 + 220Nn - 4Nn^3 - 20n^4 + 40n^3 \\
&\quad - 100n^2 + 80n)n^{-3}(N-1)_2^{-1}N^{-4}(n-2)_4^{-1},
\end{aligned}$$

$$\begin{aligned}
\mu(12)^2 &= C_{12,3}^2\mu_3^2 \\
\text{so } D_6^* &= D_{24}^* = D_{23}^* = 0, \\
D_{32}^* &= (n-1)(N-n)^2(+N^3n^4 - 40N^3 + 25N^3n^2 - 8N^3n^3 - 10N^3n \\
&\quad + 8N^2n^3 - 26N^2n^2 + 70N^2n - 120N^2 - 4N^2n^4 - 4Nn^3 - 70Nn^2 \\
&\quad - 80N + 220Nn + 14Nn^4 + 40n^3 - 100n^2 + 80n \\
&\quad - 20n^4)N^{-3}n^{-3}(N-1)_2^{-1}(n-2)_4^{-1}.
\end{aligned}$$

Finally, we have one product of 3 central moments of order 6:

$$\begin{aligned}
\mu(1^2)^3 &= C_{1^2,2}^3\mu_2^3 \text{ has } D_6^* = D_{24}^* = D_{32}^* = 0, \\
D_{23}^* &= (N-n)^3(N-2)(N^3n^3 + 9Nn^3 - 15n^3 - 3N^2n^3 - 9N^3n^2 \\
&\quad - 9Nn^2 + 9N^2n^2 \\
&\quad + 45n^2 - 30n + 29N^3n - 6N^2n - 45Nn \\
&\quad - 30N^3 + 30N)n^{-1}(N-1)^{-2}(n-1)_5^{-1}N^{-5}.
\end{aligned}$$

When $N = \infty$ the nonzero coefficients $D_{\pi_-}^*$ for the UEs of $\mu(\pi_1)\mu(\pi_2)\cdots$ are as follows.

$$\begin{aligned}
\mu(1^2)^2 &= C_{1^2,2}^2\mu_2^2 \text{ so } D_4^* = 0, \\
D_{2^2}^* &= n^{-1}(n^2 - 3n + 3)(n - 1)_3^{-1}, \\
\mu(1^2)\mu(1^3) &= C_{1^2,2}C_{1^3,3}\mu_2\mu_3 \text{ so } D_5^* = 0, \\
D_{2^3}^* &= n^{-1}(n^2 - 2n + 2)(n - 1)_4^{-1}, \\
\mu(1^2)\mu(1^4) &= C_{1^2,2}C_{1^4,4}\mu_2\mu_4 + C_{1^2,2}C_{1^4,2^2}\mu_2^3 \\
\text{so } D_6^* &= D_{3^2}^* = 0, \\
D_{2^4}^* &= -(8n^4 - 63n^3 + 190n^2 - 225n + 60)n^{-3}(n - 1)_5^{-1}, \\
D_{2^3}^* &= (n^4 - 9n^3 + 26n^2 - 30n + 30)n^{-2}(n - 3)_3^{-1}, \\
\mu(1^3)^2 &= C_{1^3,3}^2\mu_3^2 \text{ so } D_6^* = D_{2^4}^* = D_{2^3}^* = 0, \\
D_{3^2}^* &= n^{-3}(n^4 - 8n^3 + 25n^2 - 10n - 40)(n - 1)_5^{-1}, \\
\mu(1^2)\mu(12) &= C_{1^2,2}C_{12,3}\mu_2\mu_3 \text{ so } D_5^* = 0, \\
D_{2^3}^* &= n^{-1}(n^2 - 2n + 2)(n - 2)_3^{-1}, \\
\mu(1^2)\mu(1^2 2) &= C_{1^2,2}C_{1^2 2,4}\mu_2\mu_4 + C_{1^2,2}C_{1^2 2,2^2}\mu_2^3 \\
\text{so } D_6^* &= D_{3^2}^* = 0, \\
D_{2^4}^* &= -(2n^4 - 21n^3 + 70n^2 - 105n + 60)(n - 2)_4^{-1}n^{-3}, \\
D_{2^3}^* &= (n^3 - 10n^2 + 30n - 30)n^{-2}(n - 3)_3^{-1}, \\
\mu(1^2)\mu(2^2) &= C_{1^2,2}C_{2^2,4}\mu_2\mu_4 + C_{1^2,2}C_{2^2,2^2}\mu_2^3, \\
\text{so } D_6^* &= D_{3^2}^* = 0, \\
D_{2^4}^* &= -(2n^5 - 17n^4 + 49n^3 - 55n^2 + 45n - 60)(n - 2)_4^{-1}n^{-3}, \\
D_{2^3}^* &= (n^4 - 9n^3 + 26n^2 - 30n + 30)n^{-2}(n - 3)_3^{-1}, \\
\mu(1^3)\mu(12) &= C_{1^3,3}C_{12,3}\mu_3^2 \\
\text{so } D_6^* &= D_{2^4}^* = D_{2^3}^* = 0, \\
D_{3^2}^* &= n^{-3}(n^4 - 8n^3 + 25n^2 - 10n - 40)(n - 2)_4^{-1}, \\
\mu(12)^2 &= C_{12,3}^2\mu_3^2 \\
\text{so } D_6^* &= D_{2^4}^* = D_{2^3}^* = 0, \\
D_{3^2}^* &= (n - 1)n^{-3}(n^4 - 8n^3 + 25n^2 - 10n - 40)(n - 2)_4^{-1}.
\end{aligned}$$

Finally, we have one product of 3 central moments of order 6:

$$\begin{aligned}
\mu(1^2)^3 &= C_{1^2,2}^3\mu_2^3 \text{ has } D_6^* = D_{2^4}^* = D_{3^2}^* = 0, \\
D_{2^3}^* &= n^{-1}(n^2 - 7n + 15)(n - 3)_3^{-1}(n - 1)^{-1}.
\end{aligned}$$

6 UEs of products of cumulants

Here we give $E\widehat{\kappa}(\pi_1)\widehat{\kappa}(\pi_2)\dots$ and UEs of $\kappa(\pi_1)\kappa(\pi_2)\dots$ for joint cumulants $\{\kappa(\pi_i)\}$ up to total order 6 not covered by Section 2,3,5.

We first give coefficients K_{π,π_-} such that

$$\kappa(\pi) = \sum_{\pi_-}^r K_{\pi,\pi_-} m u(\pi_-) \text{ for } \pi \text{ of order } r > 1. \quad (43)$$

Its UE is then (5.3) for $D_{\pi_-}^*$ of (5.4) with $D_{\pi_-} = K_{\pi_- \pi_-}$.

$$\begin{aligned} \kappa(1^4) : \text{ Since } \kappa_{1^4} &= \mu_{1^4} - 3\mu_{1^2}^2, \\ K_{1^4,4} &= M_{1^4,4} = C_{1^4,4}, \\ K_{1^4,2^2} &= M_{1^4,2^2} - 3M_{1^2,2}^2 = C_{1^4,2^2} - 3C_{1^2,2}^2 \\ &= -3(N-n)(N^3 + 4Nn^2 - 4N^2n - 2N^2 + N \\ &\quad + 6Nn - 6n^2)n^{-3}(N-1)^{-2}(N-2)_2^{-1}, \end{aligned}$$

so

$$\begin{aligned} D_4^* &= K_{1^4,4} * C^{4,4} + K_{1^4,2^2} C^{22,4} \\ &= (N-n)(nN^3 + N^3 - 6n^2N^2 - nN^2 - 7N^2 + 6n^2N + 6Nn^3 \\ &\quad + 12nN - 6n^3 - 6n^2)(n-1)_3^{-1}N^{-3}(N-1)^{-1}n^{-1} \\ D_{22}^* &= -3(N-n)(nN^3 - N^3 - 4n^2N^2 - nN^2 + 7N^2 + 6n^2N \\ &\quad + 4Nn^3 - 12nN - 6n^3 + 6n^2)(n-1)_3^{-1}N^{-3}(N-1)^{-1}n^{-1}. \end{aligned}$$

$\kappa(1^5)$: similarly

$$K_{1^5,5} = C_{1^5,5}, \tag{44}$$

$$\begin{aligned} K_{1^5,23} &= C_{1^5,23} - 10C_{1^2,2}C_{1^3,3} \\ &= -10(N-n)(N-2n)(N^3 - 6nN^2 + 6n^2N - 2N^2 \\ &\quad + 12nN - 12n^2N)n^{-4}(N-1)^{-2}(N-2)_3, \end{aligned} \tag{45}$$

$$\begin{aligned} D_5^* &= (-N+2n)(-N+n)(5N^3 + nN^3 - 12n^2N^2 - nN^2 - 65N^2 + 12n^3N \\ &\quad + 12n^2N + 120nN - 12n^3 - 60n^2)n^{-1}(n-1)_4^{-1}N^{-4}(N-1)^{-1}, \end{aligned}$$

$$\begin{aligned} D_{23}^* &= -10(N-2n)(N-n)(nN^3 - N^3 - 6n^2N^2 - nN^2 + 13N^2 + 6n^3N \\ &\quad + 12n^2N - 24nN - 12n^3 + 12n^2)n^{-1}(n-1)_4^{-1}N^{-4}(N-1)^{-1}, \end{aligned} \tag{46}$$

$$\kappa(1^6) : K_{1^6,6} = C_{1^6,6},$$

$$\begin{aligned} K_{1^6,24} &= C_{1^6,24} - 15C_{1^4,4}C_{1^2,2} \\ &= -15(N-n)(+N^5 - 16N^4n + N^4 + 30N^3n + 64N^3n^2 \\ &\quad - 9N^3 - 96N^2n^3 + 10N^2n + 11N^2 - 150N^2n^2 + 48Nn^4 \\ &\quad + 240Nn^3 - 10Nn^2 - 4N - 120n^4)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \end{aligned}$$

$$\begin{aligned} K_{1^6,3^2} &= C_{1^6,3^2} - 10C_{1^3,3}^2 \\ &= -10(N-n)(N^6 - 5N^5 - 12N^5n + 48N^4n^2 + 13N^4 \\ &\quad + 54N^4n - 72N^3n^3 - 234N^3n^2 - 23N^3 - 66N^3n + 36N^2n^4 \\ &\quad + 360N^2n^3 + 306N^2n^2 + 22N^2 - 180Nn^4 - 8N - 480Nn^3 \\ &\quad + 240n^4)n^{-5}(N-1)^{-2}(N-2)^{-2}(N-3)_3^{-1}, \end{aligned}$$

$$\begin{aligned} K_{1^6,2^3} &= C_{1^6,2^3} - 15C_{1^4,2^2}C_{1^2,2} + 30C_{1^2,2}^3 \\ &= 30(N-n)(N^6 - 5N^5 - 11N^5n + 9N^4 + 52N^4n + 39N^4 \\ &\quad - 7N^3 - 56N^3n^3 - 176N^3n^2 - 71N^3n + 30N^2n + 28N^2n^4 \\ &\quad + 248N^2n^3 + 2N^2 + 191N^2n^2 - 240Nn^3 - 30Nn^2 - 124Nn^4 \\ &\quad + 120n^4n^2)n^{-5}(N-1)^{-3}(N-2)_4^{-1}, \end{aligned}$$

$$\begin{aligned}
D_6^* = & (N-n)(N^7n^3 - 2880Nn^3 + 3120n^2N^2 - 1440nN^3 + 960n^4 + 248N^4 \\
& + 600n^7N - 3000n^6N^2 + 6150n^5N^3 + 13800n^5N + 6000n^6N \\
& - 16860n^5N^2 - 3840n^6 - 240n^7 - 2640n^5 - 6300N^4n^4 \\
& + 18240N^3n^4 - 20280N^2n^4 + 8400Nn^4 + 16050n^3N^3 \\
& - 10020n^3N^2 - 8402n^3N^4 + 5280n^2N^3 - 7592n^2N^4 \\
& - 922nN^4 + 2945N^5n^3 - 140N^5 - 104N^6 + 1430N^5n^2 \\
& + 2005N^5n - 454N^6n^2 - 134N^6n16N^7n^2 - 94N^6n^3 \\
& + 150N^5n^5 - 30N^6n^4 - 4N^7 - 480n^7N^2 + 120N^3n^7 \\
& - 240N^4n^6 + 960n^6N^3 - 600N^4n^5 + 11N^7n \\
& + 210n^4N^5)N^{-5}(n-1)_5^{-1}(N-1)_2^{-1}(N-1)^{-1}n^{-3}
\end{aligned}$$

$$\begin{aligned}
D_{24}^* = & -15(N-n)(N^7n^3 + 2880Nn^3 - 3120n^2N^2 + 1440nN^3 - 960n^4 - 248N^4 \\
& + 456n^7N - 960n^6N^2 + 722n^5N^3 + 120n^5N + 912n^6N - 1596n^5N^2 \\
& - 480n^6 - 240n^7 + 1680n^5 - 302N^4n^4 + 1512N^3n^4 + 3360N^2n^4 \\
& - 5520Nn^4 - 6650n^3N^3 + 6900n^3N^2 - 626n^3N^4 - 3840n^2N^3 \\
& + 5236n^2N^4 + 674nN^4 + 115N^5n^3 + 140N^5 + 104N^6 - 44N^5n^2 \\
& - 1745N^5n - 34N^6n^2 + 118N^6n + 2N^7n^2 - 4N^6n^3 + 64N^5n^5 - 16N^6n^4 \\
& + 4N^7 - 264n^7N^2 + 48N^3n^7 - 96N^4n^6 + 528n^6N^3 - 342N^4n^5 - 7N^7n \\
& + 78n^4N^5)n^{-3}(n-1)_5^{-1}(N-2)_2^{-1}(N-1)^{-1}N^{-5},
\end{aligned}$$

$$\begin{aligned}
D_{32}^* = & -10(N-n)(N^7n^3 - 2880Nn^3 + 3120n^2N^2 - 1440nN^3 + 960n^4 \\
& + 248N^4 + 408n^7N - 672n^6N^2 + 102n^5N^3 - 600n^5N - 48n^6N \\
& + 1284n^5N^2 + 480n^6 - 240n^7 - 1200n^5 + 354N^4n^4 - 1704N^3n^4 \\
& - 1920N^2n^4 + 4080Nn^4 + 5070n^3N^3 - 5340n^3N^2 + 790n^3N^4 \\
& + 3120n^2N^3 - 4496n^2N^4 - 550nN^4 - 241N^5n^3 - 140N^5 - 104N^6 - 4N^5n^2 \\
& + 1615N^5n + 62N^6n^2 - 110N^6n - 2N^7n^2 - 4N^6n^3 + 48N^5n^5 - 12N^6n^4 \\
& - 4N^7 - 216n^7N^2 + 36N^3n^7 - 72N^4n^6 + 432n^6N^3 - 282N^4n^5 + 5N^7n \\
& + 66n^4N^5)n^{-3}N^{-5}(n-1)_5^{-1}(N-1)_2^{-1}(N-1)^{-1},
\end{aligned}$$

$$\begin{aligned}
D_{23}^* = & 30(N-n)(N^7n^2 + 1440Nn^3 - 1560n^2N^2 + 720nN^3 - 480n^4 - 124N^4 \\
& - 180n^6N^2 + 360n^5N^3 - 384n^5N + 384n^6N - 588n^5N^2 - 240n^6 + 720n^5 \\
& - 239N^4n^4 - 64N^3n^4 + 2292N^2n^4 - 1680Nn^4 - 2796n^3N^3 + 900n^3N^2 \\
& + 563n^3N^4 + 540n^2N^3 + 1278n^2N^4 - 594nN^4 + 44N^5n^3 + 130N^5 - 8N^6 \\
& - 322N^5n^2 - 167N^5n + 11N^6n^2 + 44N^6n - 11N^6n^3 + 2N^7 + 28n^6N^3 \\
& - 56N^4n^5 - 3N^7n + 39n^4N^5)n^{-2}N^{-5}(n-1)_5^{-1}(N-1)_2^{-1}(N-1)^{-1}.
\end{aligned}$$

$\kappa(1^4 2)$: since $\kappa_{1^4 2} = \mu_{1^4 2} - 6\mu_{1^2} \mu_{1^2 2} - 4\mu_{1^3} \mu_{12}$,

$$\begin{aligned}
K_{1^4 2.6} &= M_{1^4 2.6}, \\
K_{1^4 2.24} &= M_{1^4 2.24} - 6C_{1^2.2} C_{1^2 2.4} \\
&= -(N-n)N(n-1)(15N^4 - 118N^3 n + 15N^3 + 240N^2 n^2 \\
&\quad - 135N^2 + 186N^2 n + 160Nn - 144Nn^3 - 540Nn^2 + 165N - 60n \\
&\quad - 60n^2 + 360n^3 - 60)n^{-5}(N-1)^{-2}(N-2)_4^{-1}, \\
K_{1^4 2.3^2} &= M_{1^4 2.3^2} - 4C_{1^3.3} C_{12.3} \\
&= -2(N-n)N(n-1)(5N^5 - 31N^4 n - 25N^4 + 60N^3 n^2 + 65N^3 + 144N^3 n \\
&\quad - 36N^2 n^3 - 211N^2 n - 288N^2 n^2 - 115N^2 + 90Nn + 180Nn^3 + 372Nn^2 \\
&\quad + 110N - 40n - 240n^3 - 40)n^{-5}(N-1)^{-2}(N-2)^{-2}(N-3)_3^{-1}, \\
K_{1^4 2.2^3} &= M_{1^4 2.2^3} - 6C_{1^2.2} M_{1^2 2.2^2} \\
&= 6(N-n)N(n-1)(+5N^5 - 28N^4 n - 25N^4 + 45N^3 + 131N^3 n + 50N^3 n^2 \\
&\quad - 228N^2 n^2 - 28N^2 n^3 - 35N^2 - 178N^2 n + 75Nn + 262Nn^2 + 124Nn^3 \\
&\quad + 10N - 60n^2 - 120n^3)n^{-5}(N-1)^{-3}(N-2)_4^{-1},
\end{aligned}$$

$$\begin{aligned}
D_6^* &= (N-n)(N^6 n^3 - 1416Nn^3 + 480n^2 N + 1244n^2 N^2 - 400nN^2 - 298nN^3 \\
&\quad + 528n^4 - 192n^3 - 92N^4 + 120N^3 + 96n^6 N^2 - 144n^5 N^3 - 1176n^5 N - 120n^6 N \\
&\quad + 540n^5 N^2 - 24N^5 + 768n^5 + 48n^6 + 92N^4 n^4 - 1090N^3 n^4 + 2884N^2 n^4 \\
&\quad - 2376Nn^4 - 2390n^3 N^3 + 2932n^3 N^2 + 907n^3 N^4 - 1934n^2 N^3 + 696n^2 N^4 \\
&\quad + 689nN^4 - 58N^5 n^3 + 16N^6 n^2 + 11N^6 n - 14N^5 n^4 - 24n^6 N^3 + 36N^4 n^5 \\
&\quad - 98N^5 n - 238N^5 n^2 - 4N^6)n^{-3}(n-2)_4^{-1}(N-1)_2^{-1}(N-1)^{-1}N^{-4}, \\
D_{24}^* &= -(N-n)(15N^6 n^3 + 14040Nn^3 - 7200n^2 N - 12660n^2 N^2 + 6000nN^2 \\
&\quad + 2670nN^3 - 5040n^4 + 2880n^3 + 1380N^4 - 1800N^3 + 792n^6 N^2 \\
&\quad - 1260n^5 N^3 - 2376n^5 N - 1368n^6 N + 2244n^5 N^2 + 360N^5 + 1440n^5 \\
&\quad + 720n^6 + 564N^4 n^4 - 1370N^3 n^4 + 3948N^2 n^4 + 360Nn^4 - 2706n^3 N^3 \\
&\quad - 10980n^3 N^2 + 723n^3 N^4 + 16610n^2 N^3 - 504n^2 N^4 - 8235nN^4 \\
&\quad - 84N^5 n^3 + 30N^6 n^2 - 105N^6 n - 118N^5 n^4 - 144n^6 N^3 + 240N^4 n^5 \\
&\quad + 1110N^5 n - 236N^5 n^2 + 60N^6)n^{-3}(n-2)_4^{-1}(N-1)_2^{-1}(N-1)^{-1}N^{-4}
\end{aligned}$$

$$\begin{aligned}
D_{32}^* &= -2(N-n)(+5N^6n^3 - 3480Nn^3 + 2400n^2N + 3220n^2N^2 - 2000nN^2 \\
&\quad - 590nN^3 + 1200n^4 - 960n^3 - 460N^4 + 600N^3 + 216n^6N^2 - 348n^5N^3 \\
&\quad + 168n^5N - 408n^6N + 492n^5N^2 - 120N^5 - 480n^5 + 240n^6 + 163N^4n^4 \\
&\quad + 124N^3n^4 - 1276N^2n^4 + 360Nn^4 + 1130n^3N^3 + 2420n^3N^2 - 391n^3N^4 \\
&\quad - 4480n^2N^3 + 57n^2N^4 + 2395nN^4 - 8N^5n^3 - 10N^6n^2 + 25N^6n - 31N^5n^4 \\
&\quad - 36n^6N^3 + 60N^4n^5 - 310N^5n + 133N^5n^2 \\
&\quad - 20N^6)n^{-3}(n-2)_4^{-1}(N-1)_2^{-1}(N-1)^{-1}N^{-4} \\
D_{23}^* &= 6(N-n)(5N^6n^2 + 1320Nn^3 - 1200n^2N - 180n^2N^2 + 1000nN^2 \\
&\quad - 720nN^3 - 720n^4 + 480n^3 + 350N^4 - 300N^3 - 28n^5N^3 - 384n^5N + 180n^5N^2 \\
&\quad - 60N^5 + 240n^5 + 50N^4n^4 - 298N^3n^4 + 420N^2n^4 + 504Nn^4 + 316n^3N^3 \\
&\quad - 2116n^3N^2 + 116n^3N^4 + 1798n^2N^3 - 496n^2N^4 - 404nN^4 - 28N^5n^3 - 15N^6n \\
&\quad + 139N^5n + 21N^5n^2 + 10N^6)n^{-2}(n-2)_4^{-1}(N-1)_2^{-1}(N-1)^{-1}N^{-4}.
\end{aligned}$$

Similarly we can write down UEs for products of cumulants up to order 6 not covered by Section 2, Section 4. These are

$$\begin{aligned}
\text{of order } 4 &: \mu^2\mu(1^2), \mu\mu(12), m(2)\mu(1^2), \\
\text{of order } 5 &: \mu^3\mu(1^2), \mu^2\mu(12), \mu^2\kappa(1^3), \mu m(2)\mu(1^2), \\
&\quad \mu\mu(13), \mu\mu(2^2), m(2)\mu(12), m(2)\mu(1^3), m(3)\mu(1^2), \\
\text{of order } 6 &: \mu^4\mu(1^2), \mu^3\mu(12), \mu^3\mu(1^3), \mu^2m(2)\mu(1^2), \\
&\quad \mu^2\mu(13), \mu^2\mu(2^2), \mu^2\kappa(1^22), \mu^2\kappa(1^4), \\
&\quad \mu m(2)\mu(12), \mu m(2)\mu(1^3), \mu\mu(14), \mu\mu(23), \\
&\quad m(2)\kappa(1^4), m(2)\mu(1^22), m(2)\mu(2^2), m(2)\mu(13), \\
&\quad m(2)\mu(1^2)^2, m(2)^2\mu(1^2), m(3)\mu(12).
\end{aligned}$$

Similarly we can write down an UE for $m(2)\mu(1^4)$.

7 Proof of the inversion principles

We begin by following a false trail that leads in Section 8 to the solution of the eigenfunction problem stated at the end of the abstract.

A proof for B_r would follow if

$$B_r = H\Lambda H^{-1} \text{ where } \Lambda = \text{diag}(\lambda_1, \lambda_2, \dots), \text{ the evalues of } B, \quad (47)$$

and Λ satisfies the inversion principle, that is

$$\lambda_i(N, n) = \lambda_i \text{ satisfies } \lambda_i^{-1} = \lambda_i(n, N). \quad (48)$$

Writing $H = (u_1, u_2, \dots)$ and $H^{-1} = \begin{pmatrix} v_1' \\ v_2' \\ \vdots \end{pmatrix}$, (6.1) is equivalent to

$$Bu_i = \lambda_i u_i, v_i' B = \lambda_i v_i', v_i' u_j = \delta_{ij}, B = \sum_i \lambda_i u_i v_i'. \quad (49)$$

This implies that $a_i(F) = v_i' \mu_{(r)}$ satisfies

$$Ea_i(\widehat{F}) = \lambda_i a_i(F). \quad (50)$$

If v_i does not depend on (n, N) we call $a_i(F)$ an **r-e function** or r-eigen function). Since $m_r = EX^r = \sum_{j=0}^{\infty} \binom{r}{j} \mu^{r-j} \mu_j = v_1' m_{(r)}$ say and $E\overline{X}^r = EX^r$, for all r there is always an **r-e function** with $\lambda = 1$.

Other examples we have seen are μ_r (with $\lambda = C_{r,r}$) for $r = 2$ or 3 . In each case λ_i satisfies (7.2).

One can show that for $r \leq 3$, one can choose H in (7.1) to be **constant**, so that the number of efunctions equals the number of partitions of r say n_r that is, the dimension of $m_{(r)}$. Unfortunately this is false for $r \geq 4$. In fact in Section 8 we show that the **only** r -efunctions apart from m_r (up to a constant multiple) are $E \prod_{i=1}^s (X^{a_i} - EX^{a_i})$, with $\lambda = C_{s,s}$ for $s = 2$ or 3 where $\sum_1^s a_i = r$.

For $r = 1$, $H = \Lambda = 1$ so $a_1 = \mu$.

For $r = 2$, we can take $\lambda_1 = 1, \lambda_2 = C_{2,2}$,
 $H = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}, H^{-1} = \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$, so $a_1 = \mu_2 + \mu^2 = m_2$ and $a_2 = -\mu_2$.

For $r = 3$, we can take $\lambda_1 = C_{3,3}, \lambda_2 = C_{2,2}, \lambda_3 = 1$,

$$H = \begin{pmatrix} 1 & 0 & 0 \\ -1/2 & -1 & 0 \\ 1/2 & 3 & 1 \end{pmatrix} \text{ and } H^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ -1/2 & -1 & 0 \\ 1 & 3 & 1 \end{pmatrix},$$

so $a_1 = \mu_3, a_2 = -(\mu_3 + 2\mu_2\mu)/2 = -(m_3 - \mu m_2)/2, a_3 = \mu_3 + 3\mu_2\mu + \mu^3 = m_3$.

This was derived using the following result.

Theorem 7.1 Suppose for $i = 1, 2$, that $C_i = H_i \Lambda_i H_i^{-1}$ is $p_i \times p_i$ with $\Lambda_i = \text{diag}(\lambda_{i1}, \lambda_{i2}, \dots)$ and C_1, C_2 have no values in common. Then for D any $p_2 \times p_1$ matrix

$$\begin{pmatrix} C_1 & 0 \\ D & C_2 \end{pmatrix} = H \begin{pmatrix} \Lambda_1 & 0 \\ 0 & \Lambda_2 \end{pmatrix} H^{-1} \text{ where} \quad (51)$$

$$H = \begin{pmatrix} H_1 & 0 \\ H_2 X_0 & H_2 \end{pmatrix}, H^{-1} = \begin{pmatrix} H_1^{-1} & 0 \\ -X_0 H_1^{-1} & H_2^{-1} \end{pmatrix},$$

$$(X_0)_{ij} = (H_2^{-1} D H_1)_{ij} / (\lambda_{1j} - \lambda_{2i}). \quad (52)$$

The proof uses the following easily proved lemma.

Lemma 7.1 *Suppose for $i = 1, 2$ that $C_i = H_i \Lambda_i H_i^{-1}$ is $p_i \times p_i$ and X is $p_2 \times p_1$ satisfying*

$$XC_1 - C_2X = D. \quad (53)$$

Then (6.5) holds with $H = \begin{pmatrix} H_1 & 0 \\ XH_1 & H_2 \end{pmatrix}$, $H^{-1} = \begin{pmatrix} H_1^{-1} & 0 \\ -H_2^{-1}X & H_2^{-1} \end{pmatrix}$.

PROOF OF THEOREM 7.1. Put $X_0 = H_2^{-1}XH_1$ so (6.7) becomes

$$X_0\Lambda_1 - \Lambda_2X_0 = H_2^{-1}DH_1, \text{ which is just (6.6).}\square$$

The theorem can be extended to the case where an evaluate of $\begin{pmatrix} C_1 & 0 \\ 0 & C_2 \end{pmatrix}$ has multiplicity greater than 1. If B_r has repeated evaluate, then in place of (6.1), one has

$$B_r = H\Lambda H^{-1} \text{ where } \Lambda_{ij} = \begin{cases} \lambda_i & \text{for } j = i \\ 1 \text{ or } 0 & \text{for } j = i + 1 \\ 0 & \text{otherwise .} \end{cases}$$

But $\begin{pmatrix} \lambda & 0 \\ 1 & \lambda \end{pmatrix}^{-1} = \begin{pmatrix} \lambda^{-1} & 0 \\ -\lambda^{-2} & \lambda^{-1} \end{pmatrix}$ so Λ does not satisfy the inversion principle, so a different proof is needed.

Recall now the symmetric functions

$$[\pi_1\pi_2\cdots]_n = \sum_n' X_{i_1}^{\pi_1} X_{i_2}^{\pi_2} \cdots, \quad [\pi_1\pi_2\cdots]_N = \sum_N' x_{i_1}^{\pi_1} x_{i_2}^{\pi_2} \cdots$$

where \sum_n' sums over i_1, i_2, \dots distinct in $1, \dots, n$.

The standardised function $\langle \pi_1 \cdots \pi_j \rangle_n = [\pi_1 \cdots \pi_j]_n / (n)_j$ satisfies the invariance principle

$$\begin{aligned} E\langle \pi_1 \cdots \pi_j \rangle_n &= \langle \pi_1 \cdots \pi_j \rangle_N. \\ \text{So for every partition } \pi \text{ of } r, & \quad E\langle \pi \rangle_n = \langle \pi \rangle_N. \end{aligned} \quad (54)$$

$$\text{Now } [1]_N^2 = \sum x_i x_j = \sum_{i \neq j} x_i x_j + \sum x_i^2 = [1^2]_N + [2]_N$$

$$\text{so } [1^2]_N = [1]_N^2 - [2]_N.$$

Similarly we can write

$$S(\pi) = \sum_{\pi'}^r V^{\pi, \pi'} [\pi]_N \text{ and } [\pi]_N = V(\pi)' s_{(r)} = \sum_{\pi'}^r V_{\pi, \pi'} s(\pi') \quad (55)$$

$$\text{where } s_{(r)} = \{s_1^{r_1} s_2^{r_2} \cdots : 1.r_1 + 2.r_2 + \cdots = r\} = \{s(\pi) : r(\pi) = r\},$$

$$s(\pi) = s_{\pi_1} s_{\pi_2} \cdots, s_i = [i]_N = \sum_{j=1}^N x_j^i = N m_i \text{ and } m_i = EX^i.$$

The constant column vector $V(\pi) = \{V_{\pi,\pi'} : r(\pi') = r\}$ is given by the columns of Appendix Table 10 of Stuart and Ord (1987) for $r \leq 6$. The rows of this table give $\{V^{\pi,\pi'}\}$. We can write (7.9) as

$$([\pi]_N : r(\pi) = r) = V s_{(r), n_r} \times 1, \text{ where } V = \{V(\pi)'\} \text{ is } n_r \times n_r.$$

Now $s_1^{r_1} s_2^{r_2} \dots = N^q m_1^{r_1} m_2^{r_2} \dots$ where $q = r_1 + r_2 + \dots = q(1^{r_1} 2^{r_2} \dots)$. That is, as before $q(\pi)$ is the number of partitions in π .

$$\begin{aligned} \text{Also } \langle \pi \rangle_N &= [\pi]_N / (N)_{q(\pi)}, \\ \text{and } s_{(r)} &= D_N m_{(r)} \text{ where } D_N = \text{diag}(N^{q(\pi)} : r(\pi) = r), \\ m_{(r)} &= \{m_1^{r_1} m_2^{r_2} \dots : 1.r_1 + 2.r_2 + \dots = r\}. \\ \text{Also } m u_{(r)} &= G m_{(r)} \text{ where } G \text{ is a constant } n_r \times n_r \text{ matrix.} \\ \text{Set } \overline{D}_N &= \text{diag}\{(N)_{q(\pi)} : r(\pi) = r\}, n_r \times n_r. \end{aligned}$$

(56)

So by (7.9), $\{\langle \pi \rangle_N : r(\pi) = r\} = E_N m_{(r)} = F_N m u_{(r)}$

$$\text{where } E_N = \overline{D}_N^{-1} V D_N \text{ and } F_N = E_N G^{-1}$$

and (7.8) can be written

$$E F_n \widehat{m} u_{(r)} = F_N \widehat{m} u_{(r)}, \quad E E_n \widehat{m}_{(r)} = E_N m_{(r)}, \text{ so that}$$

$$B_r = E_n^{-1} E_N, \quad B_r^{-1} = E_N^{-1} E_n, \quad C_r = F_n^{-1} F_N \text{ and } C_r^{-1} = F_N^{-1} F_n.$$

This proves the univariate inversion principles. We now prove the multivariate inversion principle.

For π an ordered partition of r , $[\pi]_{1\dots r}$ now depends on the order of $\pi_1 \pi_2 \dots$. For example

$$[21]_{123}^N = \sum' x_{i1} x_{i2} x_{j3} \neq [12]_{123}^N = \sum' x_{i1} x_{j2} x_{j3} = [21]_{231} = [21]_{321}$$

Put another way, for π an unordered partition, we need to consider not just $[\pi]_{1\dots r}$ but all distinct $[\pi]_a$ for a any permutation of $1 \dots r$.

For $r = 3$, this gives $[3]_{123}, [21]_{123}, [21]_{231}, [21]_{312}, [1^3]_{123}$, so the dimension of $\mu_{(3)123} = \{\mu(\pi)_{123}\}$ is 5 whereas that for $\mu_{(3)}$ was only 3.

As in (7.9) there exists a constant vector $V(\pi)$ such that $[\pi]_{1\dots r}^N = V_{1\dots r}(\pi)' S_{1\dots r}(\pi)$, that is $([\pi]_{1\dots r}^N) = V s_{(r)1\dots r}$ say. Also $s_{(r)1\dots r} = D_N m_{(r)1\dots r}$ where D_N as before but with dimensions d_r say.

Redefining \overline{D}_N similarly.

$$\begin{aligned} \text{So } E \overline{D}_n^{-1} V D_n \widehat{m}_{(r)1\dots r} &= E \overline{D}_n^{-1} V s_{(r)1\dots r} = E \overline{D}_n^{-1} ([\pi]_{1\dots r}^n) \\ &= \overline{D}_N^{-1} ([\pi]_{1\dots r}^N) = \overline{D}_N^{-1} V s_{(r)1\dots r} = \overline{D}_N^{-1} V D_N m_{(r)1\dots r}. \\ \text{So } E \widehat{m}_{(r)1\dots r} &= E_n^{-1} E_N m_{(r)1\dots r} \text{ where } E_n = \overline{D}_n^{-1} V D_n \end{aligned}$$

and $E_N^{-1}E_n\widehat{m}_{(r)1\dots r}$ is an **UE** of $m_{(r)1\dots r}$. This proves the multivariate inversion principle for $m_{(r)1\dots r}$. That for $\mu_{(r)1\dots r}$ follows since $\mu_{(r)1\dots r} = Gm_{(r)1\dots r}$ where G is a constant non singular matrix.

Note 7.1 The evalues of B_r satisfy

$$|\lambda V\Lambda_{nN} - \overline{\Lambda}_{nN}V| = 0 \text{ where } \Lambda_{nN} = D_n D_N^{-1}, \overline{\Lambda}_{nN} = \overline{D}_n \overline{D}_N^{-1}.$$

Since Λ_{nN} and $\overline{\Lambda}_{nN}$ are diagonal and V is lower-triangular, it follows that the evalues of B_r are $\{\nu_i = e_i e_1^{-i}, 1 \leq i \leq r\}$ where $e_i = (n)_i / (N)_i$, with ν_i having multiplicity equal to the number of partitions π of r with $q(\pi) = i$.

Example 7.1 $\mu'_{(4)} = (\mu_4, \mu_2^2, \mu\mu_3, \mu^2\mu_2, \mu^4)$ so

$$B_4 = \begin{pmatrix} A_4 & 0 \\ B_{21} & B_{22} \end{pmatrix} \text{ has evalues } \nu_1 \nu_2^2 \nu_3 \nu_4, \text{ that is } \{\nu_1 = 1, \nu_2, \nu_2, \nu_3, \nu_4\}.$$

$$\text{But } B_{22} = \begin{pmatrix} C_{3.3} & 0 & 0 \\ 3C_{12.3} & C_{2.2} & 0 \\ 10C_{1^3.3} & 10C_{1^2.2} & 1 \end{pmatrix} \text{ has evalues } \nu_1 = 1, \nu_2 = C_{2.2}, \nu_3 = C_{3.3}.$$

So A_4 has evalues ν_2, ν_4 , as was confirmed using MAPLE.

To see that H for B_4 is not constant, it suffices to show that H for A_4 is not constant.

Set $\begin{pmatrix} a & b \\ d & c \end{pmatrix} = A_4$. Then $Au = \lambda u \iff (a - \lambda)u_1 + bu_2 = 0, du_1 + (c - \lambda)u_2 = 0,$

$$\implies \text{one can take } u \text{ proportional to } \begin{pmatrix} b \\ \lambda - a \end{pmatrix} \text{ or } \begin{pmatrix} \lambda - c \\ d \end{pmatrix}.$$

So if H is constant for A_4 then $(\lambda - a)/b$ and $(\lambda - c)/d$ are constant.

For $\lambda = \nu_4, (\lambda - a)/b = -1/3$ and $(\lambda - c)/d = -3,$

but for $\lambda = \nu_2, (\lambda - a)/b = (Nn - n - 1 - N)/(2Nn - 3n + 3 - 3N)$

and $(\lambda - c)/d = (2Nn - 3n + 3 - 3N)/(Nn - n - 1 - N)$ depend on n and N .

□

Example 7.2 $\mu'_{(5)} = (\mu_5, \mu_2\mu_3, \mu\mu_4, \mu_2^2\mu, \mu^2\mu_3, \mu^3\mu_2, \mu^5)$

so B_6 has evalues $\nu_1 \nu_2^2 \nu_3^2 \nu_4 \nu_5$, that is $\{\nu_1 = 1, \nu_2, \nu_2, \nu_3, \nu_3, \nu_4, \nu_5\}$.

$$\text{But by (2.7)} B_{22} = \begin{bmatrix} A_4 & 0 \\ C & D \end{bmatrix}$$

where D is lower triangular with diagonal $(\nu_3 = C_{3.3}, \nu_2 = C_{2.2}, \nu_1 = 1)$.

So A_5 has evalues ν_3, ν_5 . □

Example 7.3 B_6 has evalues $\nu_1 \nu_2^3 \nu_3^3 \nu_4^2 \nu_5 \nu_6$.

$$\text{But for } r = 6, B_{22} = \begin{bmatrix} A_5 & 0 & 0 & 0 & 0 \\ \cdot & A_4 & 0 & 0 & 0 \\ \cdot & \cdot & C_{3.3} & 0 & 0 \\ \cdot & \cdot & \cdot & C_{2.2} & 0 \\ \cdot & \cdot & \cdot & \cdot & 1 \end{bmatrix}$$

so has evalues $\nu_1 \nu_2^2 \nu_3^2 \nu_4 \nu_5$. So A_6 has evalues $\nu_2 \nu_3 \nu_4 \nu_6$. □

The relation between the values of A_r and the number of π_- with $q(\pi_-) = i$ is not known.
 \square

8 Symmetric functions and efunctions

Here we extend the usual expressions for symmetric functions in terms of products of power sums, and use these relations to show that the only efunctions are the noncentral moments and the 2nd and 3rd order generalised central moments.

The symmetric function $[\pi_1 \pi_2 \dots] = \sum'_N x_{i_1}^{\pi_1} x_{i_2}^{\pi_2} \dots$ is well defined for π_1, π_2, \dots any real or complex numbers.

The constants $V(\pi)$ in (7.9) tabled in Stuart and Ord can be derived as follows. We use an obvious notation.

$$\begin{aligned}
[a][b] &= [ab] + [a+b] \text{ so } [ab] = [a][b] - [a+b], \\
[a][b][c] &= [abc] + \sum^3 [a, b+c] + [a+b+c] \\
\text{so } [abc] &= [a][b][c] - \sum^3 [a][b+c] + 2[a+b+c], \\
[a] \cdot [d] &= [a \dots d] + \sum^6 [a, b, c+d] + \sum^3 [a+b, c+d] \\
&\quad + \sum^4 [a, b+c+d] + [a+b+c+d], \\
\text{so } [a \dots d] &= [a] \dots [d] - \sum^6 [a][b][c+d] + \sum^3 [a+b][c+d] \\
&\quad + 2 \sum^4 [a][b+c+d] - 6[a+b+c+d], \\
[a] \dots [e] &= [a \dots e] + \sum^{10} [abc, d+e] + \sum^{15} [a, b+c, d+e] \\
&\quad + \sum^{10} [ab, c+d+e] + \sum^{10} [a+b, c+d+e] \\
&\quad + \sum^5 [a, b+c+d+e] + [a+..+e], \\
\text{so } [a \dots e] &= [a] \dots [e] - \sum^{10} [a][b][c][d+e] + \sum^{15} [a][b+c][d+e] \\
&\quad + 2 \sum^{10} [a][b][c+d+e] - 2 \sum^{10} [a+b][c+d+e] \\
&\quad - 6 \sum^5 [a][b+c+d+e] + 24[a+..+e].
\end{aligned}$$

$$\begin{aligned}
\text{Setting } S_m(1^{r_1} 2^{r_2} \dots) &= \sum^m [a_1 \dots a_{r_1}, a_{r_1+1} + a_{r_1+2}, \dots, a_{r_1+2r_2-1} + a_{r_1+2r_2}, \\
&\quad a_{r_1+2r_2+1} + a_{r_1+2r_2+2} + a_{r_1+2r_2+3}, \dots] \tag{57}
\end{aligned}$$

$$\begin{aligned}
\text{and } T_m(1^{r_1} 2^{r_2} \dots) &= \sum^m [a_1] \dots [a_{r_1}] [a_{r_1+1} + a_{r_1+2}] \dots [a_{r_1+2r_2-1} + a_{r_1+2r_2}] \\
&\quad [a_{r_1+2r_2+1} + a_{r_1+2r_2+2} + a_{r_1+2r_2+3}] \dots, \tag{58}
\end{aligned}$$

where $m = P(1^{r_1}2^{r_2}\dots)$, we can write these pairs of equations more compactly as

$$T_1(1^2) = S_1(1^2) + S_1(2), \quad (59)$$

$$S_1(1^2) = T_1(1^2) - T_1(2), \quad (60)$$

$$T_1(1^3) = S_1(1^3) + S_3(12) + S_1(3), \quad (61)$$

$$S_1(1^3) = T_1(1^3) - T_3(12) + 2T_1(3), \quad (62)$$

$$T_1(1^4) = S_1(1^4) + S_6(1^22) + S_3(2^2) + S_4(13) + S_1(4), \quad (63)$$

$$S_1(1^4) = T_1(1^4) - T_6(1^22) + T_3(2^2) + 2T_4(13) - 6T_1(4), \quad (64)$$

$$T_1(1^5) = S_1(1^5) + S_{10}(1^32) + S_{15}(12^2) + S_{10}(1^23) \\ + S_{10}(23) + S_5(14) + S_1(5), \quad (65)$$

$$S_1(1^5) = T_1(1^5) - T_{10}(1^32) + T_{15}(12^2) + 2T_{10}(1^23) \\ - 2T_{10}(23) - 6T_5(14) + 24T_1(5). \quad (66)$$

In this way we replace and extend the last row and column of the r^{th} table in Appendix Table 10 of Stuart and Ord by a pair of equations.

The general expression for $T_1(1^r)$ is

$$T_1(1^r) = \sum_{i=1}^r B_{ri}$$

where B_{ri} can be written down from the expression for the partial exponential Bell polynomial $B_{ri}(x)$ tabled on p307 of Comtet (1974). For example

$$B_{64}(x) = 20x_1^3x_3 + 45x_1^2x_2^2 \text{ gives } B_{64} = S_{20}(1^33) + S_{45}(1^22^2).$$

In this way one obtains

$$T_1(1^6) = S_1(1^6) + S_{15}(1^42) + S_{20}(1^33) + S_{45}(1^22^2) + S_{15}(1^24) \\ + S_{60}(123) + S_{15}(2^3) + S_{15}(24) + S_6(15) + S_{10}(3^2) + S_1(6). \quad (67)$$

The coefficients of the reverse series are most easily obtained from the coefficients of the terms in the expansion of (1^r) in Appendix Table 10. In this way we obtain

$$S_1(1^6) = T_1(1^6) - T_{15}(1^42) + 2T_{20}(1^33) + T_{45}(1^22^2) - 6T_{15}(1^24) \\ - 2T_{60}(123) - T_{15}(2^3) + 6T_{15}(24) + 24T_6(15) + 4T_{10}(3^2) - 120T_1(6).$$

(More generally reading their r th table horizontally gives $\{S_m(\pi)/m\}$ in terms of $\{T_{m'}(\pi')/m'\}$, while reading vertically gives the reverse. For example reading line 2 of their 3rd table gives $S_3(12)/3 = 1.T_1(3)/1 + 1.T_3(12)/3$, that is $\sum^3[a, b+c] = 3[a+b+c] + \sum^3[a][b+c]$.) These expressions for $\{S_1(1^i)\}$ can be used to prove our assertion in Section 6 that the only $r-e$ functions are m_r , (with $\lambda = 1$), $\mu_{ab} = m_{a+b} - m_a m_b$ where $a+b=r$, with $\lambda = C_{2,2}$, and $\mu_{abc} = m_{a+b+c} - \sum^3 m_a m_{b+c} + 2m_a m_b m_c$ where $a+b+c=r$, (with $\lambda = C_{3,3}$).

$$\text{We have } \langle a+b \rangle_N = [a+b]_N/N = m_{a+b}, \\ \text{and } \langle ab \rangle_N = [ab]_N/(N)_2 = ([a]_N[b]_N - [a+b]_N)/(N)_2 \\ = (Nm_a m_b - Nm_{a+b})/(N-1).$$

So for $\langle ab \rangle_N + \nu \langle a + b \rangle_N$ to be proportional to an efunction for some constant ν , it must equal $N(m_a m_b + \theta m_{a+b}) / (N - 1)$ for some constant θ , in which case since $E\langle \pi \rangle_n = \langle \pi \rangle_N$,

$$E\alpha(\widehat{F}) = \lambda \alpha(F) \text{ where } \alpha(F) = m_a m_b + \theta m_{a+b} \text{ and } \lambda = (1 - n^{-1}) / (1 - N^{-1}) = C_{2.2}.$$

So we need to solve $(Nm_a m_b - Nm_{a+b}) / (N - 1) = N(m_a m_b + \theta m_{a+b}) / (N - 1)$.

This gives $\theta = \nu = -1$, which proves that (to within a multiplicative constant), $\mu_{ab} = m_{a+b} - m_a m_b$ is the only efunction of this type.

Taking $a + b = r$ makes it an $r - e$ function.

A similar argument shows that for $\langle abc \rangle_N + \nu_1 \sum^3 \langle a, b + c \rangle_N + \nu_2 \langle a + b + c \rangle_N$ to be proportional to an efunction, the efunction must be μ_{abc} and that $\lambda = C_{3.3}$.

However when one looks for a linear combination of say $\langle abcd \rangle_N, \dots, \langle a + b + c + d \rangle_N$ proportional to an efunction, we find that the only solutions have efunctions of the form m'_a or $\mu_{a'b'}$ or $\mu_{a'b'c'}$.

We now give a much more general meaning to the symmetric function relations (8.3) - (8.11). The functions $S(\pi) = S_{m(\pi)}(\pi) = S_m(\pi)$ and $T(\pi) = T_{m(\pi)}(\pi) = T_m(\pi)$ are defined in terms of the functions $[a_1 \dots a_r]$ and $[a_1] \dots [a_r]$ respectively, where $r = r(\pi)$.

Let $t(x_1 \dots x_r) = t(x_1, \dots, x_r)$ be an arbitrary function. Replace $[a_1 \dots a_r]$ in the definition of $S(\pi)$ by $\sum' t(x_{i_1}, \dots, x_{i_r})$ and $[a_1] \dots [a_r]$ in the definition of $T(\pi)$ by $\sum t(x_{i_1}, \dots, x_{i_r})$, where \sum sums over i_1, \dots, i_r in $1, \dots, N$ and \sum' sums over distinct i_1, \dots, i_r in $1, \dots, N$.

Then **the relations (8.3) - (8.11) remain true.**

For example (8.5) gives

$$\begin{aligned} \sum t(x_i x_j x_k) &= T(1^3) = S(1^3) + S(12) + S(3) \\ \text{where } S(1^3) &= \sum' t(x_i x_j x_k), \\ S(12) &= \sum^3 \sum' t(x_i x_j x_j) = \sum' \{t(x_i x_j x_j) + t(x_j x_i x_j) + t(x_j x_j x_i)\} \\ \text{and } S(3) &= \sum t(x_i x_i x_i). \end{aligned}$$

Similarly (8.6) gives

$$\begin{aligned} \sum' t(x_i x_j x_k) &= S(1^3) = T(1^3) - T(12) + 2T(3) \\ \text{where } T(1^3) &= \sum t(x_i x_j x_k), \\ T(12) &= \sum^3 \sum t(x_i x_j x_j), \\ \text{and } T(3) &= \sum t(x_i x_i x_i) = S(3). \end{aligned}$$

If we choose $t(x_1 x_2 \dots) = x_1^{a_1} x_2^{a_2} \dots$, we obtain our original (8.3) - (8.11). However the population need no longer be real numbers : x_1, \dots, x_N can now belong to any space Ω , and $t(x_1, \dots, x_r)$ is any real (or complex) function on Ω^r . The preceding results on r-efunctions may be similarly extended. Let us say that $a(F)$ is an **efunction** with **value**

λ_{nN} if

$$Ea(\widehat{F}) = \lambda_{nN}a(F).$$

Then for any functions $t_r(x_1, \dots, x_r)$ set

$$a_1(F) = Et_1(X'_1), \quad (68)$$

$$a_2(F) = E\{t_2(X'_1, X'_1) - t_2(X'_1, X'_2)\}, \quad (69)$$

$$\text{and } a_3(F) = E\{t_3(X'_1 X'_1 X'_1) - \sum_{i=1}^3 t_3(X'_1 X'_i X'_i) + 2t_3(X'_1 X'_2 X'_3)\} \quad (70)$$

where X'_1, X'_2, X'_3, \dots are **independent** with distribution F .

Then for $1 \leq i \leq 3$, $a_i(F)$ is an efunction with $\lambda_{nN} = C_{i,i}$.

We call a_2 and a_3 **generalised 2nd and 3rd order central moments**.

For if $t_r(x_1 \dots x_r) = f_1(x_1) \dots f_r(x_r)$ and $Y_i = f_i(X)$

then $a_2 = \mu(Y_1, Y_2) = \text{covar}(Y_1, Y_2)$ and $a_3 = \mu(Y_1, Y_2, Y_3)$.

By the same argument as before there are no other efunctions of the form

$$\sum_{r=1}^s \sum_{i_1 \dots i_r=1} \nu_{i_1 \dots i_r} Et_r(X'_{i_1}, \dots, X'_{i_r}).$$

But any smooth functional $T(F)$ can be expanded about a fixed distribution F_0 as

$$T(F) = T(F_0) + \sum_{r=1}^{\infty} E T_{F_0}(X'_1, \dots, X'_r)/r!$$

where $T_{F_0}(x_1, \dots, x_r)$ is the r th functional (von Mises) derivative of $T(F_0)$.

So the only efunctions that exist are those of form (8.12) - (8.14).

Appendix A

Here, we give the coefficients $C_{\pi, \pi'}$ obtainable from Sukhatme (1944) needed for \mathbf{C}_{--r} of (3.4) and \mathbf{C}_r of (1.7). The factorisations were obtained using MAPLE. Recall that $C_{\pi-, \pi'_+} = 0$ so expressions like $C_{2,1^2} = 0$ are not included here.

$$\begin{aligned} \lambda(\pi) & \text{ for } \pi \text{ a partition of } r \leq 6: \lambda(1^i) = e_i, \lambda(1^i 2) = e_{i+1} - 2e_{i+2}, \\ \lambda(1^i 3) & = e_{i+1} - 3e_{i+2} + 2e_{i+3}, \lambda(1^i 4) = e_{i+1} - 7e_{i+2} + 12e_{i+3} - 6e_{i+4}, \\ \lambda(1^i 5) & = e_{i+1} - 15e_{i+2} + 50e_{i+3} - 60e_{i+4} + 24e_{i+5}, \\ \lambda(6) & = e_1 - 31e_2 + 180e_3 - 390e_4 + 360e_5 - 120e_6, \\ \lambda(2^2) & = e_2 - 2e_3 + e_4, \\ \lambda(23) & = e_2 - 4e_3 + 5e_4 - 2e_5, \\ \lambda(24) & = e_2 - 8e_3 + 19e_4 - 18e_5 + 6e_6, \\ \lambda(3^2) & = e_2 - 6e_3 + 13e_4 - 12e_5 + 4e_6, \\ \lambda(2^3) & = e_3 - 3e_4 + 3e_5 - e_6. \end{aligned}$$

Set $f_i = \lambda(i)n^{-i}$, $F_i = Nf_i$, so $F_1 = 1$,

$$F_2 = (N-n)n^{-1}(N-1)^{-1},$$

$$F_3 = (N-2n)(N-n)n^{-2}(N-1)_2^{-1},$$

$$F_4 = (N-n)(N^2-6Nn+N+6n^2)n^{-3}(N-1)_3^{-1},$$

$$F_5 = (N-2n)(N-n)(N^2-12Nn+5N+12n^2)n^{-4}(N-1)_4^{-1},$$

$$F_6 = (N-n)(N^4-30N^3n+16N^3-240Nn^3+150N^2n^2+90Nn^2-90N^2n-4N+11N^2+120n^4)n^{-5}(N-1)_5^{-1},$$

$$g_1 = \lambda(12)n^{-3} = n^{-2}(n-1)(N-n)N^{-1}(N-1)_2^{-1},$$

$$g_2 = \lambda(2^2)n^{-4} = (n-1)(N-n)_2n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_3 = (\lambda(13)+3\lambda(2^2))n^{-4} = (n-1)(4N-5n-2)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_4 = \lambda(23)n^{-5} = (n-1)(N-n)_2(N-2n)n^{-4}N^{-1}(N-1)_4^{-1},$$

$$g_5 = \lambda(1^2)n^{-2} = n^{-1}(n-1)N^{-1}(N-1)^{-1},$$

$$g_6 = (\lambda(14)+6\lambda(23))n^{-5},$$

$$= (n-1)(N-n)(7N^2-24Nn+6n-N+18n^2)n^{-4}N^{-1}(N-1)_4^{-1},$$

$$g_7 = \lambda(24)n^{-6},$$

$$= (n-1)(N-n)_2(N^2-6Nn+3N-4+6n^2)n^{-5}N^{-1}(N-1)_5^{-1},$$

$$g_8 = \lambda(13)n^{-4} = (n-1)(N-2n+1)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_9 = \lambda(3^2)n^{-6},$$

$$= (n-1)(N-n)_2(N^2-4Nn-N+4+4n^2)n^{-5}N^{-1}(N-1)_5^{-1},$$

$$g_{10} = (2\lambda(13)+3\lambda(2^2)) = (n-1)(5N-7n-1)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{11} = (6\lambda(13)+7\lambda(2^2))n^{-4}$$

$$= (n-1)(13N-19n-1)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{12} = (\lambda(14)+6\lambda(23))n^{-5},$$

$$= (n-1)(N-n)(7N^2+18n^2-24Nn+6n-N)n^{-4}N^{-1}(N-1)_4^{-1},$$

$$g_{13} = (2\lambda(13)+9\lambda(2^2))n^{-4}$$

$$= (n-1)(11N-13n-7)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{14} = (2\lambda(13)+\lambda(2^2))n^{-4},$$

$$= (n-1)(3N-5n+1)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{15} = (\lambda(13)+\lambda(2^2))n^{-4} = (n-1)(2N-3n)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{16} = (\lambda(13)+6\lambda(2^2))n^{-4} = (n-1)(7N-8n-5)(N-n)n^{-3}N^{-1}(N-1)_3^{-1},$$

$$g_{17} = (\lambda(14)+6\lambda(23))n^{-5}$$

$$= (n-1)(N-n)(7N^2+18n^2-24nN+6n-N)n^{-4}N^{-1}(N-1)_4^{-1},$$

$$g_{18} = \lambda(3^2)n^{-6}$$

$$= (n-1)(N-n)_2(N^2+4n^2-4nN-N+4)n^{-5}N^{-1}(N-1)_5^{-1},$$

$$h_1 = \lambda(12^2)n^{-5} = (n-1)_2(N-n)_2n^{-4}N^{-1}(N-1)_4^{-1},$$

$$h_2 = \lambda(2^3)n^{-6} = (n-1)_2(N-n)_3n^{-5}N^{-1}(N-1)_5^{-1},$$

$$h_3 = \lambda(1^22)n^{-4} = n^{-3}(n-1)_2(N-n)N^{-1}(N-1)_3^{-1},$$

$$h_4 = \lambda(1^3)n^{-3} = n^{-2}(n-1)_2N^{-1}(N-1)_2^{-1},$$

and $F_r = Nf_r$, $G_r = N^2g_r$, $H_r = N^3h_r$. Then,

$$C_{2,2} = F_1 - F_2 = N(n-1)n^{-1}(N-1)^{-1},$$

$$C_{1^2,2} = F_2 = (N-n)n^{-1}(N-1)^{-1},$$

$$\begin{aligned}
C_{1^2,1^2} &= 1, \\
C_{3,3} &= F_1 - 3F_2 + 2F_3 = N^2(n-1)_2 n^{-2} (N-1)_2^{-1}, \\
C_{12,3} &= F_2 - F_3 = N(n-1)(N-n)n^{-2}(N-1)_2^{-1}, \\
C_{12,12} &= C_{2,2}, \\
C_{12,1^3} &= 0, \\
C_{1^3,3} &= F_3 = (N-2n)(N-n)n^{-2}(N-1)_2^{-1}, \\
C_{1^3,12} &= 3C_{1^2,2}, \\
C_{1^3,1^3} &= 1, \\
C_{4,4} &= F_1 - 4F_2 + 6F_3 - 3F_4 \\
&= N(n-1)(n^2N^2 - 2n^2N + 3n^2 - 3n - 3nN^2 + 3N^2 + 3N)n^{-3}(N-1)_3^{-1}, \\
C_{4,2^2} &= 6G_1 - 9G_2 = 3N(n-1)(2nN - 3n + 3 - 3N)(N-n)n^{-3}(N-1)_3^{-1}, \\
C_{2^2,4} &= F_2 - 2F_3 + F_4 = N(n-1)(nN - n - 1 - N)(N-n)n^{-3}(N-1)_3^{-1}, \\
C_{2^2,2^2} &= G_5 - 2G_1 + 3G_2 \\
&= N(n-1)(n^2N^2 - 3n^2N + 3n^2 + 3n - 2nN^2 + 3N^2 - 3N)n^{-3}(N-1)_3^{-1}, \\
C_{13,4} &= F_2 - 3F_3 + 2F_4 = (n-1)_2(N+1)_2(N-n)n^{-3}(N-1)_3^{-1}, \\
C_{13,2^2} &= -3G_1 + 6G_2 = -3(n-1)_2(N-n)Nn^{-3}(N-2)_2^{-1}, \\
C_{13,13} &= C_{3,3}, \\
C_{13,1^22} &= C_{13,1^4} = 0, \\
C_{1^22,4} &= F_3 - F_4 = -(n-1)N(N-2n+1)(N-n)n^{-3}(N-1)_3^{-1}, \\
C_{1^22,2^2} &= G_1 - 3G_2 = (n-1)N(nN - 3N + 3)(N-n)n^{-3}(N-1)_3^{-1}, \\
C_{1^22,13} &= 2C_{12,3}, \\
C_{1^22,1^22} &= C_{2,2}, \\
C_{1^22,1^4} &= 0, \\
C_{1^4,4} &= F_4 = (N-n)(N^2 + 6n^2 - 6nN + N)n^{-3}(N-1)_3^{-1}, \\
C_{1^4,2^2} &= 3G_2 = 3N(n-1)(N-n)_2 n^{-3}(N-1)_3^{-1}, \\
C_{1^4,13} &= 4C_{1^3,3}, \\
C_{1^4,1^22} &= 6C_{1^2,2}, \\
C_{1^4,1^4} &= 1, \\
C_{5,5} &= F_1 - 5F_2 + 10F_3 - 10F_4 + 4F_5 \\
&= N^2(n-1)_2(n^2N^2 - 5n^2N + 10n^2 - 10n - 2nN^2 + 2N^2 \\
&\quad + 10N)n^{-4}(N-1)_4^{-1}, \\
C_{5,23} &= 10G_1 - 10G_3 + 40G_4 \\
&= 10N^2(n-1)_2(nN - 2n + 2 - 2N)(N-n)n^{-4}(N-1)_4^{-1}, \\
C_{23,5} &= F_2 - 4F_3 + 5F_4 - 2F_5 \\
&= N^2(n-1)_2(nN - n - 5 - N)(N-n)n^{-4}(N-1)_4^{-1}, \\
C_{23,23} &= G_5 - 7G_1 + 5G_3 - 20G_4 \\
&= N^2(n-1)_2(n^2N^2 - 2n^2N + 2n^2 + 10n - 5nN^2 - 10N \\
&\quad + 10N^2)n^{-4}(N-1)_4^{-1}, \\
C_{14,5} &= F_2 - 4F_3 + 6F_4 - 3F_5 \\
&= N(n-1)(N-n)(n^2N^2 - n^2N + 6n^2 - 3nN^2 - 9nN - 6n + 15N \\
&\quad + 3N^2)n^{-4}(N-1)_4^{-1}, \\
C_{14,23} &= -4G_1 + 6G_3 - 30G_4 \\
&= -2N(n-1)(N-n)(2n^2N^2 + n^2N - 6n^2 + 6n - 12nN^2 + 9nN \\
&\quad + 15N^2 - 15N)n^{-4}(N-1)_4^{-1},
\end{aligned}$$

$$\begin{aligned}
C_{14,14} &= C_{4,4}, \\
C_{14,12^2} &= C_{4,2^2}, \\
C_{14,1^23} &= C_{14,1^32} = C_{14,1^5} = 0, \\
C_{12^2,5} &= F_3 - 2F_4 + F_5 \\
&= -N(n-1)(N-n)(-N^2n + 2n^2N - 2n^2 - 2n - 3nN + 5N \\
&\quad + N^2)n^{-4}(N-1)_4^{-1}, \\
C_{12^2,23} &= 2G_1 - 2G_3 + 10G_4 \\
&= 2N(n-1)(N-n)(N^2n^2 - 2Nn^2 + 2n^2 + 3Nn - 4N^2n + 2n - 5N \\
&\quad + 5N^2)n^{-4}(N-1)_4^{-1}, \\
C_{12^2,14} &= C_{2^2,4}, \\
C_{12^2,12^2} &= C_{2^2,2^2}, \\
C_{12^2,1^23} &= C_{12^2,1^32} = C_{12^2,1^5} = 0, \\
C_{1^23,5} &= F_3 - 3F_4 + 2F_5 = -N(n-1)_2(-N^2 + 2Nn + 4n - 5N)(N-n)n^{-4}(N-1)_4^{-1}, \\
C_{1^23,23} &= G_1 - 3G_3 + 20G_4 \\
&= N(n-1)_2(N^2n - 10N^2 + 8Nn - 8n + 10N)(N-n)n^{-4}(N-1)_4^{-1}, \\
C_{1^23,14} &= 2C_{13,4}, \\
C_{1^23,12^2} &= 2C_{13,2^2}, \\
C_{1^23,1^23} &= C_{3,3}, \\
C_{1^23,1^32} &= C_{1^23,1^5} = 0, \\
C_{1^32,5} &= F_4 - F_5 = N(n-1)(N-n)(N^2 + 6n^2 - 6Nn - 6n + 5N)n^{-4}(N-1)_4^{-1}, \\
C_{1^32,23} &= G_3 - 10G_4 \\
&= -N(n-1)(N-n)(-4N^2n + 5Nn^2 + 12n - 12Nn - 10N \\
&\quad + 10N^2)n^{-4}(N-1)_4^{-1}, \\
C_{1^32,14} &= 3C_{1^22,4}, \\
C_{1^32,12^2} &= 3C_{1^22,2^2}, \\
C_{1^32,1^23} &= 3C_{12,3}, \\
C_{1^32,1^32} &= C_{2,2}, \\
C_{1^32,1^5} &= 0, \\
C_{1^5,5} &= F_5 = (N-2n)(N-n)(N^2 + 12n^2 - 12nN + 5N)n^{-4}(N-1)_4^{-1}, \\
C_{1^5,23} &= 10G_4 = 10N(n-1)(N-n)_2(N-2n)n^{-4}(N-1)_4^{-1}, \\
C_{1^5,14} &= 5C_{1^4,4}, \\
C_{1^5,12^2} &= 5C_{1^4,2^2}, \\
C_{1^5,1^23} &= 10C_{1^3,3}, \\
C_{1^5,1^32} &= 10C_{1^2,2}, \\
C_{1^5,1^5} &= 1, \\
C_{6,6} &= F_1 - 6F_2 + 15F_3 - 20F_4 + 15F_5 - 5F_6 \\
&= N(n-1)(n^4N^4 - 9n^4N^3 + 31n^4N^2 - 39Nn^4 + 40n^4 - 5n^3N^4 + 30n^3N^3 \\
&\quad - 80n^3 - 95N^2n^3 + 30n^3N + 120n^2N^2 + 20n^2 + 100n^2N + 10n^2N^4 \\
&\quad - 10N^3n^2 - 75nN - 10nN^4 - 75nN^3 - 100nN^2 + 20n - 20N + 80N^3 \\
&\quad + 55N^2 + 5N^4)n^{-5}(N-1)_5^{-1}, \\
C_{6,24} &= 15(G_1 - 4G_2 + G_6 - 5G_7) \\
&= 15N(n-1)(N-n)(n^3N^3 + 29n^3N - 40n^3 - 8N^2n^3 - 50n^2N + 80n^2 \\
&\quad + 16n^2N^2 - 4N^3n^2 - 10nN - 20n - nN^2 + 7nN^3 - 5N^3 + 35N - 20 \\
&\quad - 10N^2)n^{-5}(N-1)_5^{-1}, \\
C_{6,3^2} &= 10(-2G_8 + 6G_4 - 5G_9) \\
&= 10N(n-1)(N-n)(-2N^3n^2 - 24n^3N + 40n^3 + 4N^2n^3 - 80n^2 - 2n^2N^2
\end{aligned}$$

$$\begin{aligned}
& + 40n^2N + 20n - 11nN^2 + 6nN^3 + 5nN - 25N + 10N^2 - 5N^3 \\
& + 20)n^{-5}(N-1)_5^{-1}, \\
C_{6,2^3} & = 15(3H_1 - 5H_2) \\
& = 15N^2(n-1)_2(3nN - 5N + 10 - 10n)(N-n)_2n^{-5}(N-1)_5^{-1}, \\
C_{24,6} & = F_2 - 5F_3 + 10F_4 - 9F_5 + 3F_6 \\
& = N(n-1)(N-n)(n^3N^3 + 11n^3N - 8n^3 - 4N^2n^3 - 4n^2N^2 - 8n^2 - 8n^2N \\
& - 4N^3n^2 + 12n + 42nN^2 + 12nN + 6nN^3 - 33N - 48N^2 - 3N^3 \\
& + 12)n^{-5}(N-1)_5^{-1}, \\
C_{24,24} & = G_5 - 5G_1 + 6G_{10} - 9G_6 + 45G_7 \\
& = N(n-1)(n^4N^4 - 9n^4N^3 + 53n^4N^2 - 135Nn^4 + 120n^4 - 12n^3N^3 \\
& + 41N^2n^3 - 5n^3N^4 - 60n^3N + 120n^3 + 3N^3n^2 + 51n^2N^2 + 30n^2N^4 \\
& - 210n^2N - 180n^2 - 36nN^3 + 495nN - 63nN^4 - 180n + 45N^4 + 180N \\
& - 315N^2 + 90N^3)n^{-5}(N-1)_5^{-1}, \\
C_{24,3^2} & = -4G_1 + 4G_3 - 36G_4 + 30G_9 \\
& = -2N(n-1)(N-n)(2n^3N^3 - 40n^3 - 14N^2n^3 + 40n^3N - 8N^3n^2 \\
& + 22n^2N^2 - 40n^2 - 10n^2N + 18nN^3 + 60n - 33nN^2 + 15nN - 15N^3 \\
& - 75N + 30N^2 + 60)n^{-5}(N-1)_5^{-1}, \\
C_{24,2^3} & = 3(2H_3 - 9H_1 + 15H_2) \\
& = 3N^2(n-1)_2(N-n)(2n^2N^2 + -9n^2N + 10n^2 + 24nN - 9nN^2 - 45N \\
& + 15N^2 + 30)n^{-5}(N-1)_5^{-1}, \\
C_{3^2,6} & = F_2 - 6F_3 + 13F_4 - 12F_5 + 4F_6 \\
& = N(n-1)_2(N-n)(n^2N^3 - 4n^2 - 2n^2N^2 + 5n^2N \\
& - 12nN^2 + 3Nn - 12n - 3nN^3 + 2N^3 - 8 + 22N + 32N^2)n^{-5}(N-1)_5^{-1}, \\
C_{3^2,24} & = -6G_1 + 3G_{11} - 12G_{12} + 60G_7 \\
& = -3N(n-1)_2(N-n)(2n^2N^3 + 20n^2 - 5n^2N^2 - 5n^2N - 9nN^3 + 12nN^2 - 15Nn + \\
& 60n + 40 - 70N + 20N^2 + 10N^3)n^{-5}(N-1)_5^{-1}, \\
C_{3^2,3^2} & = G_5 - 6G_1 + 2G_{13} - 48G_4 + 40G_9 \\
& = N(n-1)_2(N^4n^3 - 10Nn^3 - 40n^3 + 25N^2n^3 - 8N^3n^3 - 26N^2n^2 \\
& - 4N^4n^2 + 8N^3n^2 + 70Nn^2 - 120n^2 - 4N^3n + 220Nn - 70N^2n - 80n \\
& + 14N^4n + 80N - 100N^2 - 20N^4 + 40N^3)n^{-5}(N-1)_5^{-1}, \\
C_{3^2,2^3} & = 3(3H_3 - 12H_1 + 20H_2) \\
& = 3N^2(n-1)_2(N-n)(3n^2N^2 + 20n^2 - 15n^2N - 12nN^2 + 32Nn \\
& + 40 + 20N^2 - 60N)n^{-5}(N-1)_5^{-1}, \\
C_{2^3,6} & = F_3 - 3F_4 + 3F_5 - F_6 \\
& = -N(n-1)(N-n)(-n^2N^3 + 2n^3N^2 - 6n^3N + 4n^3 - 4n^2N^2 + 4n^2 \\
& + n^2N + 14nN^2 + 4n + 2nN^3 + 4Nn - N^3 + 4 - 16N^2 - 11N)n^{-5}(N-1)_5^{-1}, \\
C_{2^3,24} & = 3(G_1 - G_{14} + G_6 - 5G_7) \\
& = 3N(n-1)(N-n)(n^3N^3 - 20n^3 - 7n^3N^2 + 20n^3N - 3n^2N^3 + 2n^2N^2 \\
& + 15n^2N - 20n^2 - 10Nn - nN^2 + 7nN^3 - 20n + 35N - 5N^3 - 20 \\
& - 10N^2)n^{-5}(N-1)_5^{-1}, \\
C_{2^3,3^2} & = 2(-3G_2 + 6G_4 - 5G_9) \\
& = -2N(n-1)(N-n)_2(3n^2N^2 + 20n^2 - 15n^2N - 6nN^2 + 10Nn - 5N \\
& + 5N^2 + 20)n^{-5}(N-1)_5^{-1}, \\
C_{2^3,2^3} & = H_4 - 3H_3 + 9H_1 - 15H_2 \\
& = N^2(n-1)_2(n^3N^3 - 30n^3 - 9n^3N^2 + 29n^3N + 9n^2N^2 - 6n^2N - 3n^2N^3
\end{aligned}$$

$$+ 30n - 9nN^2 - 45Nn + 9nN^3 - 30N + 45N^2 - 15N^3)n^{-5}(N-1)_5^{-1}.$$

This completes $\{C_{\pi_-, \pi_-}\}$ for \mathbf{C}_6 . We now give $\{C_{\pi_+, \pi_-}\}$ for \mathbf{C}_6 :

$$\begin{aligned} C_{15,6} &= F_2 - 5F_3 + 10F_4 - 10F_5 + 4F_6 \\ &= (N-n)(N+1)_2(n-1)_2(n^2N^2 - 5n^2N + 16n^2 - 2nN^2 - 10nN \\ &\quad - 12n + 2N^2 + 30N - 8)n^{-5}(N-1)_5^{-1}, \\ C_{15,24} &= 5(-G_1 + 6G_2 - 2G_{17} + 12G_7) \\ &= -5N(n-1)_2(N-n)(n^2N^3 - 6n^2N^2 + 29n^2N - 48n^2 - 4nN^3 + 36n \\ &\quad - 8nN + 12N^2 + 6N^3 + 24 - 42N)n^{-5}(N-1)_5^{-1}, \\ C_{15,3^2} &= 10(G_8 - 4G_4 + 4G_{18}) \\ &= -10N(n-1)_2(N-n)(-nN^3 + 2n^2N^2 - 10n^2N + 16n^2 - 12n + 5nN \\ &\quad + 2N^3 - 4N^2 - 8 + 10N)n^{-5}(N-1)_5^{-1}, \\ C_{15,2^3} &= 30(-H_1 + 2H_2) \\ &= -30N^2(n-1)_2(N-n)_2(nN - 3n + 4 - 2N)n^{-5}(N-1)_5^{-1}, \\ C_{1^{24},6} &= F_3 - 4F_4 + 6F_5 - 3F_6 \\ &= -N(n-1)(N-n)(-N^3n^2 + 22n^3 + 2N^2n^3 - 10N^2n^2 - 47Nn^2 \\ &\quad - 38n^2 + 3N^3n + 87Nn + 12n + 42N^2n - 33N - 3N^3 + 12 \\ &\quad - 48N^2)n^{-5}(N-1)_5^{-1}, \\ C_{1^{24},24} &= G_1 - 12G_2 + 6G_6 - 45G_7 \\ &= N(n-1)(N-n)(-12N^3n^2 + 47Nn^3 - 90n^3 + N^3n^3 + 330n^2 - 132Nn^2 \\ &\quad - 24N^2n^2 - 105Nn - 180n + 42N^3n + 99N^2n + 315N - 90N^2 - 45N^3 \\ &\quad - 180)n^{-5}(N-1)_5^{-1}, \\ C_{1^{24},3^2} &= -4G_8 + 24G_4 - 30G_9 \\ &= 2N(n-1)(N-n)(-2N^3n^2 - 12Nn^3 + 20n^3 + 4N^2n^3 + 62Nn^2 - 20N^2n^2 \\ &\quad - 100n^2 + 60n - 15Nn + 3N^2n + 12N^3n - 75N + 30N^2 - 15N^3 \\ &\quad + 60)n^{-5}(N-1)_5^{-1}, \\ C_{1^{24},2^3} &= 18H_1 - 45H_2 \\ &= 9N^2(n-1)_2(2Nn - 5n + 10 - 5N)(N-n-1)(N-n)n^{-5}(N-1)_5^{-1}, \\ C_{123,6} &= F_3 - 4F_4 + 5F_5 - 2F_6 \\ &= -(N+1)_2(n-1)_2(N-n)(-N^2n - 2n^2 + 2Nn^2 - 5Nn - 6n + N^2 \\ &\quad + 15N - 4)n^{-5}(N-1)_5^{-1}, \\ C_{123,24} &= G_1 - 6G_{15} + 5G_6 - 30G_7 \\ &= N(n-1)_2(N-n)(N^3n^2 - 25Nn^2 + 30n^2 + 6N^2n^2 - 20Nn \\ &\quad - 10N^3n + 90n - 105N + 30N^2 + 15N^3 + 60)n^{-5}(N-1)_5^{-1}, \\ C_{123,3^2} &= G_1 - G_{16} + 20G_4 - 20G_9 \\ &= N(n-1)_2(N-n)(N^3n^2 + 15Nn^2 - 20n^2 - 4N^2n^2 - 5N^3n + 25Nn \\ &\quad - 60n - 40 + 50N + 10N^3 - 20N^2)n^{-5}(N-1)_5^{-1}, \\ C_{123,2^3} &= -3H_3 + 15H_1 - 30H_2 \\ &= -3N^2(n-1)_2(N-n)(N^2n^2 - 4Nn^2 + 5n^2 - 5N^2n + 5n + 10Nn \\ &\quad + 20 - 30N + 10N^2)n^{-5}(N-1)_5^{-1}, \\ C_{1^{33},6} &= F_4 - 3F_5 + 2F_6 \\ &= N(n-1)_2(N-n)(N^3 + 18n^2 + 6Nn^2 - 6n - 36Nn - 6N^2n - 4 \\ &\quad + 11N + 16N^2)n^{-5}(N-1)_5^{-1}, \\ C_{1^{33},24} &= 3G_2 - 3G_6 + 30G_7 \\ &= -3N(n-1)_2(N-n)(-N^3n + 9Nn^2 - 10n^2 + N^2n^2 - 5Nn - 12N^2n \end{aligned}$$

$$\begin{aligned}
& + 30n + 20 - 35N + 10N^2 + 5N^3)n^{-5}(N-1)_5^{-1}, \\
C_{1^3 3, 3^2} &= G_8 - 12G_4 + 20G_9 \\
&= -N(n-1)_2(N-n)(-N^3n + 6Nn^2 + 2N^2n^2 + 45Nn - 60n - 24N^2n \\
&\quad - 40 - 20N^2 + 50N + 10N^3)n^{-5}(N-1)_5^{-1}, \\
C_{1^3 3, 2^3} &= -9H_1 + 30H_2 \\
&= -3N^2(n-1)_2(N-n)_2(3Nn - 5n + 20 - 10N)n^{-5}(N-1)_5^{-1}, \\
C_{1^2 2^2, 6} &= F_4 - 2F_5 + F_6 \\
&= N(n-1)(N-n)(N^3n - 6n^3 + 6Nn^3 - 6N^2n^2 - 12Nn^2 - 6n^2 \\
&\quad + 29Nn + 4n + 14N^2n - 16N^2 - 11N - N^3 + 4)n^{-5}(N-1)_5^{-1}, \\
C_{1^2 2^2, 2^4} &= G_{14} - 2G_6 + 15G_7 \\
&= -N(n-1)(N-n)(-3N^3n^2 - 9Nn^3 + 10n^3 + 5N^2n^3 + 21Nn^2 - 22N^2n^2 \\
&\quad + 10n^2 - 35Nn + 33N^2n - 60n + 14N^3n + 105N - 30N^2 - 15N^3 \\
&\quad - 60)n^{-5}(N-1)_5^{-1}, \\
C_{1^2 2^2, 3^2} &= 2G_2 - 8G_4 + 10G_9 \\
&= 2N(n-1)(N-n)_2(N^2n^2 - Nn^2 - 4N^2n - 5N + 20 \\
&\quad + 5N^2)n^{-5}(N-1)_5^{-1}, \\
C_{1^2 2^2, 2^3} &= H_3 - 6H_1 + 15H_2 \\
&= N^2(n-1)_2(N-n)(N^2n^2 - 3Nn^2 + 5n^2 - 6N^2n + 6Nn + 15n + 15N^2 \\
&\quad + 30 - 45N)n^{-5}(N-1)_5^{-1}, \\
C_{1^4 2, 6} &= F_5 - F_6 \\
&= N(n-1)(N-n)(N-2n+1)(N^2 + 12n^2 - 12Nn - 12n - 4 + 15N)n^{-5}(N-1)_5^{-1}, \\
C_{1^4 2, 2^4} &= G_6 - 15G_7 \\
&= N(n-1)(N-n)(7N^3n + 18Nn^3 + 60n^2 - 24N^2n^2 - 54Nn^2 - 60n \\
&\quad + 69N^2n - 40Nn + 105N - 30N^2 - 15N^3 - 60)n^{-5}(N-1)_5^{-1}, \\
C_{1^4 2, 3^2} &= 2(2G_4 - 5G_9) \\
&= 2N(n-1)(N-n)_2(2N^2n - 4Nn^2 + 10Nn - 20 - 5N^2 \\
&\quad + 5N)n^{-5}(N-1)_5^{-1}, \\
C_{1^4 2, 2^3} &= 3(H_1 - 5H_2) \\
&= 3N^2(n-1)_2(N-n)_2(Nn + 10 - 5N)n^{-5}(N-1)_5^{-1}, \\
C_{1^6, 6} &= F_6 \\
&= (N-n)(-30N^3n + 120n^4 - 240Nn^3 + 150N^2n^2 + 90Nn^2 - 90N^2n \\
&\quad - 4N + 11N^2 + 16N^3 + N^4)n^{-5}(N-1)_5^{-1}, \\
C_{1^6, 2^4} &= 15G_7 \\
&= 15N(n-1)(N-n)_2(N^2 + 6n^2 - 6Nn + 3N - 4)n^{-5}(N-1)_5^{-1}, \\
C_{1^6, 3^2} &= 10G_9 \\
&= 10N(n-1)(N-n)_2(N^2 + 4n^2 - 4Nn - N + 4)n^{-5}(N-1)_5^{-1}, \\
C_{1^6, 2^3} &= 15H_2 \\
&= 15N^2(n-1)_2(N-n)_3n^{-5}(N-1)_5^{-1}.
\end{aligned}$$

Finally, $\{C_{\pi_+, \pi_+}\}$ for \mathbf{C}_6 are obtainable using (3.7).

Appendix B

This gives $C_{\pi, \pi'}$ for \mathbf{C}_{-r} of (3.4) and \mathbf{C}_r of (1.7) when $N = \infty$.

The results of Appendix A hold with

$$\begin{aligned}
F_1 &= 1, F_2 = n^{-1}, F_3 = n^{-2}, F_4 = n^{-3}, F_5 = n^{-4}, F_6 = n^{-5}, G_1 = (n-1)n^{-2}, G_2 = \\
&(n-1)n^{-3}, G_3 = 4G_2, G_4 = (n-1)n^{-4}, \\
G_5 &= (n-1)n^{-1}, G_6 = 7G_4, G_7 = (n-1)n^{-5}, \\
G_8 &= G_2, G_9 = G_7, G_{10} = 5G_2, \\
G_{11} &= 13G_2, G_{12} = 7G_4, G_{13} = 11G_2, G_{14} = 3G_2, G_{15} = 2G_2, G_{16} = 7G_2, \\
H_1 &= (n-1)_2n^{-4}, H_2 = (n-1)_2n^{-5}, H_3 = (n-1)_2n^{-3}, H_4 = (n-1)_2n^{-2}, \\
C_{2,2} &= (n-1)n^{-1}, C_{1^2,2} = n^{-1}, C_{1^2,1^2} = 1, \\
C_{3,3} &= (n-1)_2n^{-2}, C_{12,3} = (n-1)n^{-2}, C_{12,12} = (n-1)n^{-1}, C_{12,1^3} = 0, \\
C_{1^3,3} &= n^{-2}, C_{1^3,1^2} = 3n^{-1}, C_{1^3,1^3} = 1, \\
C_{4,4} &= (n-1)(n^2 - 3n + 3)n^{-3}, C_{4,2^2} = 3(n-1)(2n-3)n^{-3}, \\
C_{2^2,4} &= (n-1)^2n^{-3}, C_{2^2,2^2} = (n-1)(n^2 - 2n + 3)n^{-3}, \\
C_{13,4} &= (n-1)_2n^{-3}, C_{13,2^2} = -3(n-1)_2n^{-3}, C_{13,1^3} = C_{3,3}, \\
C_{13,1^2} &= C_{13,1^4} = 0, \\
C_{1^2,2,4} &= (n-1)n^{-3}, C_{1^2,2,2^2} = (n-1)(n-3)n^{-3}, C_{1^2,2,1^3} = 2(n-1)n^{-2}, \\
C_{1^2,1^2,2} &= (n-1)n^{-1}, C_{1^2,2,1^4} = 0, \\
C_{1^4,4} &= n^{-3}, C_{1^4,2^2} = 3(n-1)n^{-3}, C_{1^4,1^3} = 4n^{-2}, C_{1^4,1^2} = 6n^{-1}, C_{1^4,1^4} = 1, \\
C_{5,5} &= (n-1)_2(n^2 - 2n + 2)n^{-4}, C_{5,2^3} = 10(n-1)(n-2)^2n^{-4}, \\
C_{2^3,5} &= (n-2)(n-1)^2n^{-4}, C_{2^3,2^3} = (n-1)_2(n^2 - 5n + 10)n^{-4}, \\
C_{14,5} &= (n-1)(n^2 - 3n + 3)n^{-4}, C_{14,2^3} = -2(n-1)(2n^2 - 12n + 15)n^{-4}, \\
C_{14,1^4} &= C_{4,4}, C_{14,1^2} = C_{4,2^2}, C_{14,1^3} = C_{14,1^3} = C_{14,1^5} = 0, \\
C_{1^2,2,5} &= (n-1)^2n^{-4}, C_{1^2,2,2^3} = 2(n-1)(n^2 - 4n + 5)n^{-4}, C_{1^2,2,1^4} = C_{2^2,4}, \\
C_{1^2,2,1^2} &= C_{2^2,2^2}, C_{1^2,2,1^3} = C_{1^2,2,1^3} = C_{1^2,2,1^5} = 0, \\
C_{1^2,3,5} &= (n-1)_2n^{-4}, C_{1^2,3,2^3} = (n-10)(n-1)_2n^{-4}, C_{1^2,3,1^4} = 2C_{31,4}, \\
C_{1^2,3,1^2} &= 2C_{13,2^2}, C_{1^2,3,1^3} = C_{3,3}, C_{1^2,3,1^3} = C_{1^2,3,1^5} = 0, \\
C_{1^3,2,5} &= (n-1)n^{-4}, C_{1^3,2,2^3} = 2(n-1)(2n-5)n^{-4}, C_{1^3,2,1^4} = 3C_{1^2,2,4}, \\
C_{1^3,2,1^2} &= 3C_{1^2,2,2^2}, C_{1^3,2,1^3} = 3C_{12,3}, C_{1^3,2,1^3} = C_{2,2}, C_{1^3,2,1^5} = 0, \\
C_{1^5,5} &= n^{-4}, C_{1^5,2^3} = 10(n-1)n^{-4}, C_{1^5,1^4} = 5C_{1^4,4}, C_{1^5,1^2} = 15C_{1^4,2^2}, \\
C_{1^5,1^3} &= 10C_{1^3,3}, C_{1^5,1^3} = 10C_{1^2,2}, C_{1^5,1^5} = 1, \\
C_{6,6} &= (n-1)(n^4 - 5n^3 + 10n^2 - 10n + 5)n^{-5}, \\
C_{6,2^4} &= 15(n-1)(n^3 - 4n^2 + 7n - 5)n^{-5}, \\
C_{6,3^2} &= -10(n-1)(2n^2 - 6n + 5)n^{-5}, \\
C_{6,2^3} &= 15(3n-5)(n-1)_2n^{-5}, \\
C_{2^4,6} &= (n^2 - 3n + 3)(n-1)^2n^{-5}, \\
C_{2^4,2^4} &= (n-1)(n^4 - 5n^3 + 30n^2 - 63n + 45)n^{-5}, \\
C_{2^4,3^2} &= -2(n-1)(2n^3 - 8n^2 + 18n - 15)n^{-5}, \\
C_{2^4,2^3} &= 3(n-1)_2(2n^2 - 9n + 15)n^{-5}, \\
C_{3^2,6} &= (n-1)^2(n-2)^2n^{-5}, \\
C_{3^2,2^4} &= -3(n-1)(2n-5)(n-2)^2n^{-5}, \\
C_{3^2,3^2} &= (n-1)(n^2 - 2n + 10)(n-2)^2n^{-5}, \\
C_{3^2,2^3} &= 3(n-1)_2(3n^2 - 12n + 20)n^{-5}, \\
C_{2^3,6} &= (n-1)^3n^{-5}, \\
C_{2^3,2^4} &= 3(n^2 - 2n + 5)(n-1)^2n^{-5}, \\
C_{2^3,3^2} &= -2(n-1)(3n^2 - 6n + 5)n^{-5}, \\
C_{2^3,2^3} &= (n-1)_2(n^3 - 3n^2 + 9n - 15)n^{-5},
\end{aligned}$$

$$\begin{aligned}
C_{1^{24},6} &= (n-1)(n^2-3n+3)n^{-5}, \\
C_{1^{24},24} &= (n-1)(n-3)(n^2-9n+15)n^{-5}, \\
C_{1^{24},3^2} &= -2(n-1)(2n^2-12n+15)n^{-5}, \\
C_{1^{24},3} &= 9(2n-5)(n-1)_2 n^{-5}, \\
C_{1^{23},6} &= (n-2)(n-1)^2 n^{-5}, \\
C_{1^{23},24} &= (n-1)_2 (n^2-10n+15)n^{-5}, \\
C_{1^{23},3^2} &= (n-1)_2 (n^2-5n+10)n^{-5}, \\
C_{1^{23},2^3} &= -3(n-1)_2 (n^2-5n+10)n^{-5}, \\
C_{1^{33},6} &= (n-1)_2 n^{-5}, \\
C_{1^{33},24} &= 3(n-5)(n-1)_2 n^{-5}, \\
C_{1^{33},3^2} &= (n-10)(n-1)_2 n^{-5}, \\
C_{1^{33},2^3} &= -3(3n-10)(n-1)_2 n^{-5}, \\
C_{1^{22}2,6} &= (n-1)^2 n^{-5}, \\
C_{1^{22}2,24} &= (3n-5)(n-1)(n-3)n^{-5}, \\
C_{1^{22}2,3^2} &= 2(n-1)(n^2-4n+5)n^{-5}, \\
C_{1^{22}2,2^3} &= (n-1)_2 (n^2-6n+15)n^{-5}, \\
C_{1^{42},6} &= (n-1)n^{-5}, \\
C_{1^{42},24} &= (n-1)(7n-15)n^{-5}, \\
C_{1^{42},3^2} &= 2(n-1)(2n-5)n^{-5}, \\
C_{1^{42},2^3} &= 3(n-5)(n-1)_2 n^{-5}, \\
C_{1^6,6} &= n^{-5}, \\
C_{1^6,24} &= 15(n-1)n^{-5}, \\
C_{1^6,3^2} &= 10(n-1)n^{-5}, \\
C_{1^6,2^3} &= 15(n-1)_2 n^{-5}.
\end{aligned}$$

Appendix C

Here, we give for $r \leq 6$, $\mathbf{C}_{--r}^{-1} \mathbf{C}_{1,1}^{-1} = \mathbf{C}_r^{1,1} = (C^{\pi^-, \pi'})$ for the case $N = \infty$. By (3.4) this is needed for the UE of $\boldsymbol{\mu}_{-(r)}$. By the inversion principle, these elements are $C^{\pi, \pi'}(\infty, n) = C_{\pi, \pi'}(n, \infty)$. We also give the other elements of \mathbf{C}_r^{-1} , $C_r^{2,1}$ and $C_r^{2,2}$ for $r \leq 5$ defined by (2.9). These are needed for the UE of $\boldsymbol{\mu}_{+(r)}$ in (3.8). Set $\mathbf{B}_r(N, n) = \mathbf{B}_r$, $\mathbf{C}_r(N, n) = \mathbf{C}_r$. Then $\mathbf{B}_r^{-1} = \mathbf{B}_r(n, N)$, $\mathbf{C}_r^{-1} = \mathbf{C}_r(n, N)$,

$$\begin{aligned}
C_2^{1,1} &= C_{2,2}^{-1} = C^{2,2} = n(n-1)^{-1}, \\
C_2^{2,1} &= C^{1^2,2} = -(n-1)^{-1}, \\
C_2^{2,2} &= C^{1^2,1^2} = 1.
\end{aligned}$$

$$C_3^{1,1} = C_{3,3}^{-1} = C^{3,3} = (n-1)_2^{-1} n^2, \mathbf{C}_3^{2,1} = \begin{pmatrix} C^{12,3} \\ C^{1^3,3} \end{pmatrix},$$

$$C^{12,3} = -n(n-1)_2^{-1},$$

$$C^{1^3,3} = 2(n-1)_2^{-1},$$

$$\mathbf{C}_3^{2,2} = \begin{pmatrix} C^{12,12} & C^{12,1^3} \\ C^{1^3,12} & C^{1^3,1^3} \end{pmatrix},$$

$$C^{12,12} = n(n-1)^{-1},$$

$$\begin{aligned}
C^{12,1^3} &= 0, \\
C^{1^3,12} &= -3(n-1)^{-1}, \\
C^{1^3,1^3} &= 1. \\
\mathbf{C}_4^{11} &= \begin{pmatrix} C^{4,4} & C^{4,2^2} \\ C^{2^2,4} & C^{2^2,2^2} \end{pmatrix}, \\
C^{4,4} &= n(n^2 - 2n + 3)(n-1)_3^{-1}, \\
C^{4,2^2} &= -3n(2n-3)(n-1)_3^{-1}, \\
C^{2^2,4} &= -n(n-2)_2^{-1}, \\
C^{2^2,2^2} &= n(n^2 - 3n + 3)(n-1)_3^{-1}, \\
\mathbf{C}_4^{2,1} &= \begin{pmatrix} C^{13,4} & C^{13,2^2} \\ C^{1^2 2,4} & C^{1^2 2,2^2} \\ C^{1^4,4} & C^{1^4,2^2} \end{pmatrix}, \\
C^{13,4} &= -n(1+n)(n-1)_3^{-1}, \\
C^{13,2^2} &= 3n(n-2)_2^{-1}, \\
C^{1^2 2,4} &= 2n(n-1)_3^{-1}, \\
C^{1^2 2,2^2} &= -n^2(n-1)_3^{-1}, \\
C^{1^4,4} &= -6(n-1)_3^{-1}, \\
C^{1^4,2^2} &= 3n(n-1)_3^{-1}, \\
\mathbf{C}_4^{2,2} &= \begin{pmatrix} C^{13,13} & C^{13,1^2 2} & C^{13,1^4} \\ C^{1^2 2,13} & C^{1^2 2,1^2 2} & C^{1^2 2,1^4} \\ C^{1^4,13} & C^{1^4,1^2 2} & C^{1^4,1^4} \end{pmatrix}, \\
C^{13,13} &= n^2(n-1)_2^{-1}, \\
C^{13,1^2 2} &= 0, \\
C^{13,1^4} &= 0, \\
C^{1^2 2,13} &= -2n(n-1)_2^{-1}, \\
C^{1^2 2,1^2 2} &= n(n-1)^{-1}, \\
C^{1^2 2,1^4} &= 0, \\
C^{1^4,13} &= 8(n-1)_2^{-1}, \\
C^{1^4,1^2 2} &= -6(n-1)^{-1}, \\
C^{1^4,1^4} &= 1. \\
\mathbf{C}_5^{1,1} &= \begin{pmatrix} C^{5,5} & C^{5,2^3} \\ C^{2^3,5} & C^{2^3,2^3} \end{pmatrix}, \\
C^{5,5} &= n^2(n^2 - 5n + 10)(n-1)_4^{-1}, \\
C^{5,2^3} &= -10n^2(n-1)^{-1}(n-3)_2^{-1}, \\
C^{2^3,5} &= -n^2(n-2)_3^{-1}, \\
C^{2^3,2^3} &= n^2(n^2 - 2n + 2)(n-1)_4^{-1}, \\
\mathbf{C}_5^{2,1} &= \begin{pmatrix} C^{14,5} & C^{14,2^3} \\ C^{1^2 2,5} & C^{1^2 2,2^3} \\ C^{1^3 2,5} & C^{1^3 2,2^3} \\ C^{1^5,5} & C^{1^5,2^3} \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
C^{14,5} &= -n(n^2 - n + 6)(n - 1)_4^{-1}, \\
C^{14,23} &= 2n(n + 2)(2n - 3)(n - 1)_4^{-1}, \\
C^{12^2,5} &= 2n(n - 2)_3^{-1}, \\
C^{12^2,23} &= -2n(n^2 - 2n + 2)(n - 1)_4, \\
C^{1^23,5} &= 2n(n + 2)(n - 1)_4^{-1}, \\
C^{1^23,23} &= -n(n^2 + 8n - 8)(n - 1)_4^{-1}, \\
C^{1^32,5} &= -6n(n - 1)_4^{-1}, \\
C^{1^32,23} &= 5n^2(n - 1)_4^{-1}, \\
C^{1^5,5} &= 24(n - 1)_4^{-1}, \\
C^{1^5,23} &= -20n(n - 1)_4^{-1}, \\
C_5^{2,2} &= \begin{pmatrix} C^{14,14} & C^{14,12^2} & C^{14,1^23} & C^{14,1^32} & C^{14,1^5} \\ C^{12^2,14} & C^{12^2,12^2} & C^{12^2,1^23} & C^{12^2,1^32} & C^{12^2,1^5} \\ C^{1^23,14} & C^{1^23,12^2} & C^{1^23,1^23} & C^{1^23,1^32} & C^{1^23,1^5} \\ C^{1^32,14} & C^{1^32,12^2} & C^{1^32,1^23} & C^{1^32,1^32} & C^{1^32,1^5} \\ C^{1^5,14} & C^{1^5,12^2} & C^{1^5,1^23} & C^{1^5,1^32} & C^{1^5,1^5} \end{pmatrix}, \\
C^{14,14} &= n(n^2 - 2n + 3)(n - 1)_3^{-1}, \\
C^{14,12^2} &= -3n(2n - 3)(n - 1)_3^{-1}, \\
C^{14,1^23} &= 0, \\
C^{14,1^32} &= 0, \\
C^{14,1^5} &= 0, \\
C^{12^2,14} &= -n(n - 2)_2^{-1}, \\
C^{12^2,12^2} &= n(n^2 - 3n + 3)(n - 1)_3^{-1}, \\
C^{12^2,1^23} &= 0, \\
C^{12^2,1^32} &= 0, \\
C^{12^2,1^5} &= 0, \\
C^{1^23,14} &= -2n(n + 1)(n - 1)_3^{-1}, \\
C^{1^23,12^2} &= 6n(n - 2)_2^{-1}, \\
C^{1^23,1^23} &= n^2(n - 1)_2^{-1}, \\
C^{1^23,1^32} &= 0, \\
C^{1^23,1^5} &= 0, \\
C^{1^32,14} &= 6n(n - 1)_3^{-1}, \\
C^{1^32,12^2} &= -3n^2(n - 1)_3^{-1}, \\
C^{1^32,1^23} &= -3n(n - 1)_2^{-1}, \\
C^{1^32,1^32} &= n(n - 1)^{-1}, \\
C^{1^32,1^5} &= 0, \\
C^{1^5,14} &= -30(n - 1)_3^{-1}, \\
C^{1^5,12^2} &= 15n(n - 1)_3^{-1}, \\
C^{1^5,1^23} &= 20(n - 1)_2^{-1}, \\
C^{1^5,1^32} &= -10(n - 1)^{-1}, \\
C^{1^5,1^5} &= 1.
\end{aligned}$$

$$\mathbf{C}_6^{1,1} = \begin{pmatrix} C^{6,6} & C^{6,24} & C^{6,3^2} & C^{6,2^3} \\ C^{24,6} & C^{24,24} & C^{24,3^2} & C^{24,2^3} \\ C^{3^2,6} & C^{3^2,24} & C^{3^2,3^2} & C^{3^2,2^3} \\ C^{2^3,6} & C^{2^3,24} & C^{2^3,3^2} & C^{2^3,2^3} \end{pmatrix},$$

$$C^{6,6} = n(n^4 - 9n^3 + 31n^2 - 39n + 40)(n-1)_5^{-1},$$

$$C^{6,24} = -15n(n^3 - 8n^2 + 29n - 40)(n-1)_5^{-1},$$

$$C^{6,3^2} = -40n(n^2 - 6n + 10)(n-1)_5^{-1},$$

$$C^{6,2^3} = 15n^2(3n - 10)(n-1)_5^{-1},$$

$$C^{24,6} = -n(n^2 - 3n + 8)(n-2)_4^{-1},$$

$$C^{24,24} = n(n^4 - 9n^3 + 53n^2 - 135n + 120)(n-1)_5^{-1},$$

$$C^{24,3^2} = 4n(n^2 - 5n + 10)(n-1)^{-1}(n-3)_3^{-1},$$

$$C^{24,2^3} = -3n^2(2n - 5)(n-1)^{-1}(n-3)_3^{-1},$$

$$C^{3^2,6} = -n(n^2 - n + 4)(n-2)_4^{-1},$$

$$C^{3^2,24} = 3n(2n^3 - 5n^2 - 5n + 20)(n-1)_5^{-1},$$

$$C^{3^2,3^2} = n(n^4 - 8n^3 + 25n^2 - 10n - 40)(n-1)_5^{-1},$$

$$C^{3^2,2^3} = -3n^2(3n^2 - 15n + 20)(n-1)_5^{-1},$$

$$C^{2^3,6} = 2n(n-3)_3^{-1},$$

$$C^{2^3,24} = -3n(n^2 - 5n + 10)(n-1)^{-1}(n-3)_3^{-1},$$

$$C^{2^3,3^2} = -2n(3n^2 - 15n + 20)(n-1)_5^{-1},$$

$$C^{2^3,2^3} = n^2(n^2 - 7n + 15)(n-1)^{-1}(n-3)_3^{-1},$$

$$C^{1^2 4,6} = 2n(n^2 + 11)(n-1)_5^{-1},$$

$$C^{1^2 4,24} = -n(n^3 + 47n - 90)(n-1)_5^{-1},$$

$$C^{1^2 4,3^2} = -4(n+5)n(n-3)_3^{-1}(n-1)^{-1},$$

$$C^{1^2 4,2^3} = 9n^2(2n - 5)(n-1)_5^{-1},$$

$$C^{1^{23},6} = 2(n+1)n(n-2)_4^{-1},$$

$$C^{1^{23},24} = -n(n^3 + 6n^2 - 25n + 30)(n-1)_5^{-1},$$

$$C^{1^{23},3^2} = -n(n^3 + 15n - 4n^2 - 20)(n-1)_5^{-1},$$

$$C^{1^{23},2^3} = 3n^2(n^2 - 4n + 5)(n-1)_5^{-1},$$

$$C^{1^{33},6} = -6n(3+n)(n-1)_5^{-1},$$

$$C^{1^{33},24} = 3(n+10)n(n-2)_4^{-1},$$

$$C^{1^{33},3^2} = 2n^2(3+n)(n-1)_5^{-1},$$

$$C^{1^{33},2^3} = -3n^2(3n - 5)(n-1)_5^{-1},$$

$$C^{1^{22} 2,6} = -6n(n-2)_4^{-1},$$

$$C^{1^{22} 2,24} = n(5n^2 - 9n + 10)(n-1)_5^{-1},$$

$$C^{1^{22} 2,3^2} = 2n^2(n-2)_4^{-1},$$

$$C^{1^{22} 2,2^3} = -n^2(n^2 - 3n + 5)(n-1)_5^{-1},$$

$$C^{1^{42},6} = 24n(n-1)_5^{-1},$$

$$C^{1^{42},24} = -18n^2(n-1)_5^{-1},$$

$$C^{1^{42},3^2} = -8n^2(n-1)_5^{-1},$$

$$\begin{aligned}C^{1^4 2, 2^3} &= 3n^3(n-1)_5^{-1}, \\C^{1^6, 6} &= -120(n-1)_5^{-1}, \\C^{1^5, 2^4} &= 90n(n-1)_5^{-1}, \\C^{1^5, 3^2} &= 40n(n-1)_5^{-1}, \\C^{1^5, 2^3} &= -15n^2(n-1)_5^{-1}.\end{aligned}$$

Appendix D

Here, we give $\mathbf{a} = \mathbf{a}(\boldsymbol{\pi})$, $\mathbf{b} = \mathbf{b}(\boldsymbol{\pi})$ needed in (1.11), the explicit formula for the UEs of products of cumulants up to order six:

$$\begin{aligned}
\kappa_4 : \mathbf{a} &= (1, -3) \\
\mathbf{aC}_{--4}^{-1} &= n(N-1)N^{-3}(n-1)_3^{-1}(J_1, J_2), \\
J_1 &= N^2n^2 - 6Nn^2 + 6n^2 + 6n + N^2n - 6N, \\
J_2 &= -3(N^2n^2 - N^2n - 4Nn^2 + 6n^2 + 6N - 6n). \\
\kappa_5 : \mathbf{a} &= (1, -10), \\
\mathbf{aC}_{--5}^{-1} &= n^2(N-1)_2N^{-4}(n-1)_4^{-1}(J_1, J_2), \\
J_1 &= N^2n^2 - 12Nn^2 + 12n^2 + 5N^2n + 60n - 60N, \\
J_2 &= -10(N^2n^2 - N^2n - 6Nn^2 + 12n^2 + 12N - 12n). \\
\kappa_{4\mu} : \mathbf{b} &= (0, 0, 1, -3, 0, 0, 0), \\
\mathbf{bC}_5^{-1} &= n(N-1)N^{-4}(n-1)_4^{-1}(J_1, J_2, J_3, J_4, 0, 0, 0), \\
J_1 &= -(N-n)(N^2n^2 + 5N^2n - 18Nn - 6Nn^2 - 12N + 6n^2 + 30n), \\
J_2 &= 2(N-n)(5N^2n^2 - 5N^2n + 18Nn - 24Nn^2 + 12N + 30n^2 \\
&\quad - 30n), \\
J_3 &= N(n-4)(N^2n^2 - 6Nn^2 + 6n^2 + 6n + N^2n - 6N), \\
J_4 &= -3N(n-4)(N^2n^2 - N^2n - 4Nn^2 + 6n^2 + 6N - 6n). \\
\kappa_6 : \mathbf{a} &= (1, -15, -10, 30), \\
\mathbf{aC}_{--6}^{-1} &= n(N-1)(n-1)_5^{-1}N^{-5}(J_1, J_2, J_3, J_4), \\
J_1 &= 1890N^2n^2 - 1800Nn - 2400Nn^2 + 600N^2n - 90N^3n^3 + 480N \\
&\quad - 480n + 480N^2 - 120N^3 + 1320n^2 + 1920n^3 - 30N^3n^4 \\
&\quad + N^4n^4 + 480N^2n^3 - 1800Nn^3 - 390N^3n^2 - 90N^3n + 150N^2n^4 \\
&\quad - 240Nn^4 + 16N^4n^3 + 11N^4n^2 - 4N^4n + 120n^4, \\
J_2 &= -15(6N^2n^2 + 1320Nn - 360N^2n - 480N + 480n - 480N^2 \\
&\quad + 120N^3 - 840n^2 + 240n^3 - 16N^3n^4 + N^4n^4 + 36N^2n^3 \\
&\quad - 96Nn^3 - 26N^3n^2 + 90N^3n + 78N^2n^4 - 168Nn^4 + 2N^4n^3 \\
&\quad - 7N^4n^2 + 4N^4n + 120n^4), \\
J_3 &= -10(-54N^2n^2 - 1080Nn + 120Nn^2 + 240N^2n + 480N - 480n \\
&\quad + 480N^2 - 120N^3 + 600n^2 - 240n^3 - 12N^3n^4 + N^4n^4 \\
&\quad - 60N^2n^3 + 144Nn^3 + 54N^3n^2 - 90N^3n + 66N^2n^4 - 144Nn^4 \\
&\quad - 2N^4n^3 + 5N^4n^2 - 4N^4n + 120n^4), \\
J_4 &= 30n(N-2)(9N^2n^2 - 180Nn - 36Nn^2 + 36N^2n + N^3n^3 \\
&\quad + 120N - 120n + 180n^2 - 60n^3 - 9N^2n^3 + 36Nn^3 - 3N^3n^2 + 2N^3n).
\end{aligned}$$

$$\begin{aligned}
\kappa_5\mu : \mathbf{b} &= (0^4, 1, -10, 0^5), \\
\mathbf{bC}_6^{-1} &= n(N-1)_2 N^{-4} (n-1)_4^{-1} (J_1, J_2, J_3, J_4, J_5, J_6, 0, 0, 0, 0), \\
J_1 &= -(n+1)(N-n)(n^2 N^2 - 12n^2 N + 12n^2 + 15nN^2 - 60nN \\
&\quad + 180n - 4N^2 - 72N - 48)N^{-1}(n-5)^{-1}, \\
J_2 &= 15(N-n)(n^3 N^2 - 7nN^2 + 4N^2 + 2n^2 N^2 - 8n^3 N + 72N \\
&\quad - 16nN + 24n^2 + 12n^3 + 48 - 84n)N^{-1}(n-5)^{-1}, \\
J_3 &= 10(N-n)(n^3 N^2 + 5nN^2 - 4N^2 - 2n^2 N^2 + 30nN - 6n^3 N - 72N \\
&\quad - 48 - 24n^2 + 12n^3 + 60n)N^{-1}(n-5)^{-1}, \\
J_4 &= -30(N-n)n(n^2 N^2 - 3nN^2 + 2N^2 + 12nN + 6N - 6n^2 N \\
&\quad + 12n^2 - 36n + 24)N^{-1}(n-5)^{-1}, \\
J_5 &= n(n^2 N^2 - 12n^2 N + 12n^2 + 60n + 5nN^2 - 60N), \\
J_6 &= -10n(n^2 N^2 - nN^2 - 6n^2 N + 12N - 12n + 12n^2). \\
\kappa_4\mu^2 : \mathbf{b} &= (0^6, 1, -3, 0^3), \\
\mathbf{bC}_6^{-1} &= n(N-1)N^{-5}(n-1)_5^{-1}(J_1, J_2, J_3, J_4, J_5, J_6, J_7, J_8, 0, 0, 0), \\
J_1 &= (N-n)(24 + 174Nn - 28N^2 n^2 + 84Nn^2 - 83N^2 n + 24N - 66n - 56N^2 \\
&\quad + 4N^3 - 96n^2 - 6n^3 + 6Nn^3 - N^2 n^3 + 2N^3 n^2 + 18N^3 n), \\
J_2 &= -(N-n)(-210Nn - 90N^2 n^2 + 198Nn^2 - 69N^2 n + N^3 n^3 - 360N + 630n \\
&\quad + 360N^2 - 60N^3 - 180n^2 - 90n^3 + 84Nn^3 - 21N^2 n^3 + 15N^3 n^2 + 20N^3 n \\
&\quad - 360), \\
J_3 &= -2(N-n)(-26N^2 n^2 - 30Nn + 6Nn^2 + 59N^2 n + 120N - 150n - 100N^2 \\
&\quad + 20N^3 + 60n^2 - 30n^3 - 5N^2 n^3 + 24Nn^3 + 7N^3 n^2 - 15N^3 n + 120), \\
J_4 &= 3n(N-2)(N-n)(3N^2 n - 12Nn^2 + 12Nn + 30n^2 - 90n + 30N + 60 \\
&\quad - 10N^2 + N^2 n^2), \\
J_5 &= -2N(n-5)(N-n)(N^2 n^2 + 5N^2 n - 18Nn - 6Nn^2 - 12N + 6n^2 + 30n), \\
J_6 &= 4N(n-5)(N-n)(5N^2 n^2 - 5N^2 n + 18Nn - 24Nn^2 + 12N + 30n^2 \\
&\quad - 30n), \\
J_7 &= N^2(n-4)_2(N^2 n^2 - 6Nn^2 + 6n^2 + N^2 n + 6n - 6N), \\
J_8 &= -3N^2(n-4)_2(-N^2 n - 4Nn^2 + 6n^2 + 6N - 6n + N^2 n^2).
\end{aligned}$$

$$\begin{aligned}
\kappa_4 \kappa_2 : \mathbf{a} &= (0, 1, 0, -3), \\
\mathbf{aC}_{--6}^{-1} &= n(N-1)(n-1)_5^{-1} N^{-5} (J_1, J_2, J_3, J_4), \\
J_1 &= -(N-n)(24 - 16N^2 n^2 + 24Nn + 84Nn^2 - 5N^2 n + N^3 n^3 + 24N - 66n \\
&\quad + 4N^2 + 4N^3 - 96n^2 - 6n^3 - 7N^2 n^3 + 12Nn^3 + 2N^3 n^2 - 7N^3 n), \\
J_2 &= -93N^2 n^2 + 990Nn - 120N^2 n + 24N^3 n^3 - 360N + 360n - 360N^2 - 60N^3 \\
&\quad - 630n^2 - 60N^4 + 180n^3 - 14N^3 n^4 + N^4 n^4 + 48N^2 n^3 - 72Nn^3 - 121N^3 n^2 \\
&\quad + 255N^3 n + 57N^2 n^4 - 126Nn^4 - 10N^4 n^2 + 45N^4 n + 90n^4, \\
J_3 &= 2(N-n)(58N^2 n^2 + 30Nn - 66Nn^2 - 85N^2 n + 2N^3 n^3 + 120N - 150n \\
&\quad + 20N^2 + 20N^3 + 60n^2 - 30n^3 - 17N^2 n^3 + 36Nn^3 - 5N^3 n^2 - 5N^3 n + 120), \\
J_4 &= -3n(N-2)(9N^2 n^2 - 90Nn - 18Nn^2 + 8N^2 n + N^3 n^3 + 60N - 60n - 20N^3 \\
&\quad + 90n^2 - 30n^3 - 5N^2 n^3 + 18Nn^3 - 7N^3 n^2 + 20N^3 n).
\end{aligned}$$

For $N = \infty$,

$$\begin{aligned}
\kappa_4 : \mathbf{a} &= (1, -3), \\
\mathbf{aC}_{--4}^{-1} &= n^2(n-1)_3^{-1} (n+1, -3(n-1)). \\
\kappa_5 : \mathbf{a} &= (1, -10), \\
\mathbf{aC}_{--5}^{-1} &= n^3(n-1)_4^{-1} (n+5, -10(n+1)). \\
\kappa_4 \mu : \mathbf{b} &= (0, 0, 1, -3, 0, 0, 0), \\
\mathbf{bC}_5^{-1} &= n^2(n-1)_4^{-1} (J_1, J_2, J_3, J_4, 0, 0, 0), \\
J_1 &= -(n+5), \\
J_2 &= 10(n-1), \\
J_3 &= (n+1)(n-4), \\
J_4 &= -3(n-1)(n-4). \\
\kappa_6 : \mathbf{a} &= (1, -15, -10, 30), \\
\mathbf{aC}_{--6}^{-1} &= n^2(n-1)_5^{-1} (J_1, J_2, J_3, J_4), \\
J_1 &= (n+1)(n^2 + 15n - 4), \\
J_2 &= -15(n+4)(n-1)^2, \\
J_3 &= -10(n^2 - n + 4)(n-1), \\
J_4 &= 30n(n-1)_2. \\
\kappa_5 \mu : \mathbf{b} &= (0^4, 1, -10, 0^5), \\
\mathbf{bC}_6^{-1} &= (J_1, J_2, J_3, J_4, J_5, J_6, 0, 0, 0, 0, 0), \\
J_1 &= -n(n+1)(n^2 - 4 + 15n)(n-1)_5^{-1}, \\
J_2 &= 15n(n+4)(n-1)(n-2)_4^{-1}, \\
J_3 &= 10(n^2 - n + 4)n(n-2)_4^{-1}, \\
J_4 &= -30n^2(n-3)_3^{-1}, \\
J_5 &= n^3(n+5)(n-1)_4^{-1}, \\
J_6 &= -10n^3(n-2)_3^{-1}.
\end{aligned}$$

These expressions for $\kappa_4, \kappa_5, \kappa_6$ agree with Fisher's: see equation (12.28) of Stuart and Ord (1987). The UE for $\kappa_4\mu$ above can be derived from $\kappa_{41} = \kappa_4\kappa_1 - \kappa_5/n$ at the top of page 6 of Wishart (1952).

Appendix E

Set $S_a = \sum_{i=1}^n X_i^a, s_a = \sum_{i=1}^N x_i^a$. Here, we extend the table of Skellam (1949) for $ES_{a_1} \cdots S_{a_r}$ in terms of $\{s_a\}$ up to $r = 6$. We exclude the cases where a_1, \dots, a_r are all the same or all different.

$$\begin{aligned}
ES_a^2 S_b &= \lambda_3 s_{a+b+c} + \lambda_{12} (s_{2a} s_b + 2s_{a+b} s_a) + \lambda_{13} s_a^2 s_b, \\
ES_a^2 S_b S_c &= \lambda_4 s_{2a+b+c} + \lambda_{13} \left(\sum_{i=1}^2 s_{2a+b} s_c + 2s_{a+b+c} s_a \right) \\
&\quad + \lambda_{22} (s_{2a} s_{b+c} + 2s_{a+b} s_{a+c}) \\
&\quad + \lambda_{122} (s_{2a} s_b s_c + s_{b+c} s_a^2 + 2 \sum_{i=1}^2 s_{a+b} s_a s_c) + \lambda_{14} s_a^2 s_b s_c, \\
ES_a^2 S_b^2 &= \lambda_4 s_{2a+2b} + 2\lambda_{13} (s_{2a+b} s_b + s_{a+2b} s_a) + \lambda_{22} (s_{2a} s_{2b} + 2s_{a+b}^2) \\
&\quad + \lambda_{122} (s_{2a} s_b^2 + s_{2b} s_a^2 + 4s_{a+b} s_a s_b) + \lambda_{14} s_a^2 s_b^2, \\
ES_a^3 S_b &= \lambda_4 s_{3a+b} + \lambda_{13} (s_{3a} s_b + 3s_{2a+b} s_a) + 3\lambda_{22} s_{2a} s_{a+b} \\
&\quad + \lambda_{122} (3s_{2a} s_a s_b + 3s_{a+b} s_a^2) + \lambda_{14} s_a^3 s_b, \\
ES_a^4 S_b &= \lambda_5 s_{4a+b} + \lambda_{14} (4s_a s_{3a+b} + s_b s_{4a}) + \lambda_{23} (4s_{a+b} s_{3a} + 6s_{2a} s_{2a+b}) \\
&\quad + \lambda_{123} (4s_{3a} s_a s_b + 6s_{2a+b} s_a^2) + \lambda_{122} (3s_{2a}^2 s_b + 12s_{a+b} s_{2a} s_a) \\
&\quad + \lambda_{132} (6s_{2a} s_a^2 s_b + 4s_{a+b} s_a^3) + \lambda_{13} s_a^4 s_b, \\
ES_a^3 S_b^2 &= \lambda_5 s_{3a+2b} + \lambda_{14} (3s_a s_{2a+2b} + 2s_b s_{b+3a}) \\
&\quad + \lambda_{23} (3s_{2a} s_{a+2b} + s_{2b} s_{3b} + 6s_{a+b} s_{2a+b}) \\
&\quad + \lambda_{123} (s_{3a} s_b^2 + 6s_{2a+b} s_a s_b + 3s_{a+2b} s_a^2) \\
&\quad + \lambda_{122} (3s_{2b} s_{2a} s_a + 6s_{a+b}^2 s_a + 6s_{2b} s_{a+b} s_b) \\
&\quad + \lambda_{132} (s_{2b} s_a^3 + 6s_{a+b} s_a^2 s_b + 3s_{2a} s_a s_b^2) + \lambda_{15} s_a^3 s_b^2, \\
ES_a^3 S_b S_c &= \lambda_5 s_{3a+b+c} + \lambda_{14} (3s_{2a+b+c} s_a + \sum_{i=1}^2 s_{3a+c} s_b) \\
&\quad + \lambda_{23} (s_{3a} s_{b+c} + 3 \sum_{i=1}^2 s_{2a+b} s_{a+c} + 3s_{a+b+c} s_{2a}) \\
&\quad + \lambda_{123} (s_{3a} s_b s_c + 3 \sum_{i=1}^2 s_{2a+b} s_a s_c + 3s_{a+b+c} s_a^2) \\
&\quad + \lambda_{122} (3 \sum_{i=1}^2 s_{2a} s_{a+b} s_c + 3s_{2a} s_{b+c} s_a + 6s_{a+b} s_{a+c} s_a) \\
&\quad + \lambda_{132} (s_{b+c} s_a^3 + 3 \sum_{i=1}^2 s_{a+b} s_c s_a^2 + 3s_{2a} s_b s_c) + \lambda_{15} s_a^3 s_b s_c,
\end{aligned}$$

$$\begin{aligned}
ES_a^2 S_b^2 S_c &= \lambda_5 s_{2a+2b+c} + \lambda_{14} (s_{2a+2b} s_c + 2 \sum^2 s_{a+c+2b} s_a) \\
&+ \lambda_{23} (\sum^2 s_{2a+c} s_{2b} + 2 \sum^2 s_{2a+b} s_{b+c} + 4 s_{a+b+c} s_{a+b}) \\
&+ \lambda_{1^2 3} (\sum^2 s_{2b+c} s_a^2 + 2 \sum^2 s_{a+2b} s_a s_c + 4 s_{a+b+c} s_a s_b) \\
&+ \lambda_{12^2} (s_{2a} s_{2b} s_c + 2 \sum^2 s_{2a} s_{b+c} s_b + 2 s_{a+b}^2 s_c + 4 \sum^2 s_{a+b} s_{a+c} s_b) \\
&+ \lambda_{1^3 2} (\sum^2 s_{2a} s_b^2 s_c + 4 s_{a+b} s_a s_b s_c + 2 \sum^2 s_{a+c} s_a s_b^2) + \lambda_{1^5} s_a^2 s_b^2 s_c, \\
ES_a^2 S_b S_c S_d &= \lambda_5 s_{2a+b+c+d} + \lambda_{14} (\sum^3 s_{2a+b+c} s_d + s_{a+b+c+d} s_a) \\
&+ \lambda_{23} (s_{2a} s_{b+c+d} + 2 \sum^3 s_{a+b} s_{a+c+d} + \sum^3 s_{b+c} s_{2a+d}) \\
&+ \lambda_{1^2 3} (s_a^2 s_{b+c+d} + 2 \sum^3 s_a s_b s_{a+c+d} + \sum^3 s_b s_c s_{2a+d}) \\
&+ \lambda_{12^2} (3 \sum^3 s_a s_{a+b} s_{c+d} + 2 \sum^3 s_b s_{a+c} s_{a+d}) \\
&+ \lambda_{1^3 2} (s_{2a} s_b s_c s_d + 2 \sum^3 s_{a+b} s_a s_c s_d + \sum^3 s_a^2 s_{b+c} s_d) + \lambda_{1^5} s_a^2 s_b s_c s_d, \\
ES_a^5 S_b &= \lambda_6 s_5 a s_b + \lambda_{15} (5 s_a s_{4a+b} + s_b s_5 a) + 5 \lambda_{24} (s_{a+b} s_{4a} + 2 s_{2a} s_{3a+b}) \\
&+ 10 \lambda_{33} s_{2a+b} s_{3a} + 5 \lambda_{1^2 4} (s_{4a} s_a s_b + 2 s_{3a+b} s_{2a}) \\
&+ 10 \lambda_{123} (2 s_{3a} s_{a+b} s_a + 3 s_{2a+b} s_{2a} s_a + s_{3a} s_{2a} s_b) + 15 \lambda_{2^3} s_{2a}^2 s_{a+b} \\
&+ 10 \lambda_{1^3 3} (s_{3a} s_a^2 s_b + s_{2a+b} s_a^3) + 15 \lambda_{1^2 2^2} (2 s_{2a} s_{a+b} s_a^2 + s_{2a}^2 s_b s_a) \\
&+ 5 \lambda_{1^4 2} (2 s_{2a} s_a^3 s_b + s_{a+b} s_a^4) + \lambda_{1^6} s_a^5 s_b, \\
ES_a^4 S_b^2 &= \lambda_6 s_{4a+2b} + 2 \lambda_{15} (s_a s_{3a+2b} + s_b s_{4a+b}) + \lambda_{24} (8 s_{a+b} s_{3a+b} \\
&+ 6 s_{2a} s_{2a+2b} + s_{2b} s_{4a}) + 2 \lambda_{33} (3 s_{2a+b}^2 + 2 s_{a+2b} s_{3a}) \\
&+ \lambda_{1^2 4} (8 s_{3a+b} s_a s_b + 6 s_{2a+2b} s_a^2 + s_{4a} s_b^2) + 4 \lambda_{123} (6 s_{2a+b} s_{a+b} s_a \\
&+ 3 s_{a+2b} s_{2a} s_a + 3 s_{2a+b} s_{2a} s_b + s_{3a} s_{2b} s_a + 2 s_{3a} s_{a+b} s_b) \\
&+ 3 \lambda_{2^3} (4 s_{a+b}^2 s_{2a} + s_{2a}^2 s_{2b}) + 4 \lambda_{1^3 3} (3 s_{2a+b} s_a^2 s_b + s_{3a} s_a s_b^2 + s_{a+2b} s_a^3) \\
&+ 3 \lambda_{1^2 2^2} (s_{2a}^2 s_b^2 + 2 s_{2a} s_{2b} s_a^2 + 4 s_{a+b}^2 s_a^2 + 8 s_{a+b} s_{2a} s_a s_b) \\
&+ \lambda_{1^4 2} (8 s_{a+b} s_a^3 s_b + 6 s_{2a} s_a^2 s_b^2 + s_{2b} s_a^4) + \lambda_{1^6} s_a^4 s_b^2, \\
ES_a^3 S_b^3 &= \lambda_6 s_{3a+3b} + 3 \lambda_{15} \sum^2 s_a s_{2a+3b} + 3 \lambda_{24} (s_{a+b} s_{2a+2b} + \sum^2 s_{2a} s_{a+3b}) \\
&+ \lambda_{33} (s_{a+2b} s_{2a+b} + s_{3a} s_{3b}) + 3 \lambda_{1^2 4} (3 s_{2a+2b} s_a s_b + \sum^2 s_{a+3b} s_a^2) \\
&+ 3 \lambda_{123} (\sum^2 s_{3b} s_{2a} s_a + 3 \sum^2 s_{a+2b} s_{2a} s_b + 6 \sum^2 s_{a+2b} s_{a+b} s_a) \\
&+ 3 \lambda_{2^3} (2 s_{a+b}^3 + 3 s_{a+b} s_{2a} s_{2b}) + \lambda_{1^3 3} (9 \sum^2 s_{2a+b} s_a s_b^2 + \sum^2 s_{3b} s_a^3) \\
&+ 9 \lambda_{1^2 2^2} (s_{2b} s_{2a} s_a s_b + 2 s_{a+b}^2 s_a s_b + s_{a+b} \sum^2 s_{2a} s_b^2) \\
&+ 3 \lambda_{1^4 2} (3 s_{a+b} s_a^2 s_b^2 + \sum^2 s_{2a} s_a s_b^3) + \lambda_{1^6} s_a^3 s_b^3,
\end{aligned}$$

$$\begin{aligned}
ES_a^4 S_b S_c &= \lambda_6 s_{4a+b+c} + \lambda_{15} (4s_a s_{3a+b+c} + \sum^2 s_{4a+b} s_c) + \lambda_{24} (4 \sum^2 s_{a+b} s_{3a+c} \\
&+ 6s_{2a} s_{2a+b+c} + s_{b+c} s_{4a}) + 2\lambda_{33} (2s_{a+b+c} s_{3a} + 3s_{2a+b} s_{2a+c}) \\
&+ \lambda_{124} (4s_a \sum^2 s_{3a+b} s_c + 6s_{2a+b+c} s_a^2 + s_{4a} s_b s_c) \\
&+ 2\lambda_{123} (2s_{3a} \sum^3 s_{a+b} s_c + 6s_{a+b+c} s_{2a} s_a + 3s_{2a} \sum^2 s_{2a+b} s_c \\
&+ 6s_a \sum^2 s_{2a+b} s_{a+c}) + 3\lambda_{23} (4s_{2a} s_{a+b} s_{a+c} + s_{2a}^2 s_{b+c}) \\
&+ 2\lambda_{133} (2s_{3a} s_a s_b s_c + 3s_a^2 \sum^2 s_{2a+b} s_c + 2s_{a+b+c} s_a^3) \\
&+ 3\lambda_{122^2} (4s_{2a} s_a \sum^2 s_{a+b} s_c + s_{2a}^2 s_b s_c + 2s_{2a} s_{b+c} s_a^2 + 4s_{a+c} s_{a+b} s_a^2) \\
&+ \lambda_{142} (6s_{2a} s_a^2 s_b s_c + 4s_a^3 \sum^2 s_{a+b} s_c + s_{b+c} s_a^4) + \lambda_{16} s_a^4 s_b s_c,
\end{aligned}$$

$$\begin{aligned}
ES_a^3 S_b^2 S_c &= \lambda_6 s_{3a+2b+c} + \lambda_{15} 3s_a s_{2a+2b+c} + 2s_b s_{3a+b+c} + s_c s_{3a+2b} \\
&+ \lambda_{24} (6s_{a+b} s_{2a+b+c} + 3s_{a+c} s_{2a+2b} + 3s_{2a} s_{a+2b+c} + 2s_{b+c} s_{3a+b} \\
&+ s_{2b} s_{3a+c}) + \lambda_{33} (6s_{a+b+c} s_{2a+b} + 3s_{a+2b} s_{2a+c} + s_{3a} s_{2b+c}) \\
&+ \lambda_{124} (6s_{2a+b+c} s_a s_b + 3s_{2a+2b} s_a s_c + 3s_{a+2b+c} s_a^2 + 2s_{3a+b} s_b s_c \\
&+ s_{3a+c} s_b^2) + \lambda_{123} (3s_{2b+c} s_{2a} s_a + 3s_{a+2b} s_{2a} s_c + 3s_{2a+c} s_{2b} s_a \\
&+ 12s_{a+b+c} s_{a+b} s_a + 6s_{2a+b} s_{a+c} s_b + 6s_{2a+b} s_{a+b} s_c + 6s_{2a+b} s_{b+c} s_a \\
&+ 6s_{a+2b} s_{a+c} s_a + 6s_{2a+c} s_{a+b} s_b + 6s_{a+b+c} s_{2a} s_b + s_{3a} s_{2b} s_c \\
&+ 2s_{3a} s_{b+c} s_b) + 3\lambda_{23} (2s_{a+b}^2 s_{a+c} + 2s_{a+b} s_{2a} s_{b+c} + s_{2a} s_{a+c} s_{2b}) \\
&+ \lambda_{133} (6s_{2a+b} s_a s_b s_c + 6s_{a+b+c} s_a^2 s_b + 3s_{2a+c} s_a s_b^2 + 3s_{a+2b} s_a^2 s_c \\
&+ s_{2b+c} s_a^3 + s_{3a} s_b^2 s_c) + \lambda_{122^2} (3s_{a+c} s_{2a} s_b^2 + 3s_{a+c} s_{2b} s_a^2 \\
&+ 3s_{2a} s_{2b} s_a s_c + 6s_{a+b}^2 s_c s_a + 6s_{b+c} s_{a+b} s_a^2 + 6s_{a+b} s_{2a} s_c s_b \\
&+ 6s_{b+c} s_{2a} s_a s_b + 12s_{a+c} s_{a+b} s_{a+b}) + \lambda_{142} (6s_{a+b} s_a^2 s_b s_c \\
&+ 3s_{2a} s_a s_b^2 s_c + 3s_{a+c} s_a^2 s_b^2 + 2s_{b+c} s_a^3 s_b + s_{2b} s_a^3 s_c) + \lambda_{16} s_a^3 s_b^2 s_c,
\end{aligned}$$

$$\begin{aligned}
ES_a^2 S_b^2 S_c^2 &= \lambda_6 s_{2a+2b+2c} + 2\lambda_{15} \sum^3 s_a s_{a+2b+2c} + \lambda_{24} (4 \sum^3 s_{a+b} s_{a+b+2c} \\
&+ \sum^3 s_{2a} s_{2b+2c}) + 2\lambda_{33} (2s_{a+b+c}^2 + \sum^3 s_{a+2b} s_{a+2c}) \\
&+ \lambda_{124} (4 \sum^3 s_{a+b+2c} s_a s_b + \sum^3 s_{2a+2b} s_c^2) + 2\lambda_{123} (4s_{a+b+c} \sum^3 s_{a+b} s_c \\
&+ 2 \sum^6 s_{2a+b} s_{b+c} s_c + \sum^6 s_{a+2b} s_{2c} s_a) + \lambda_{23} (2 \sum^3 s_{a+b}^2 s_{2c} \\
&+ 8s_{a+b} s_{a+c} s_{b+c} + s_{2c} s_{2a} s_{2b}) + 2\lambda_{133} (4s_{a+b+c} s_a s_b s_c \\
&+ \sum^6 s_{b+2c} s_a^2 s_b) + \lambda_{122^2} (8 \sum^3 s_{b+c} s_{a+b} s_c s_a + 4 \sum^3 s_{2c} s_{a+b} s_b s_a \\
&+ \sum^3 s_{2b} s_{2a} s_c^2 + 2 \sum^3 s_{a+b}^2 s_c^2) + \lambda_{142} (4 \sum^3 s_{a+b} s_a s_b s_c^2 \\
&+ \sum^3 s_{2a} s_b^2 s_c^2) + \lambda_{16} s_a^2 s_b^2 s_c^2,
\end{aligned}$$

$$\begin{aligned}
ES_a^2 S_b^2 S_c S_d = & \lambda_6 s_{2a+2b+c+d} + \lambda_{15} \left(2 \sum^2 s_a s_{a+2b+c+d} + \sum^2 s_c s_{2a+2b+d} \right) \\
& + \lambda_{24} \left(4s_{a+b} s_{a+b+c+d} + 2 \sum^4 s_{a+c} s_{a+2b+d} + \sum^2 s_{2a} s_{2b+c+d} \right. \\
& + s_{c+d} s_{2a+2b} \left. \right) + \lambda_{33} \left(4s_{a+b+c} s_{a+b+d} + 2 \sum^2 s_{a+2b} s_{a+c+d} \right. \\
& + \sum^2 s_{2a+c} s_{2b+d} \left. \right) + \lambda_{124} \left(4s_{a+b+c+d} s_a s_b + 2 \sum^4 s_a s_c s_{a+2b+d} \right. \\
& + \sum^2 s_a^2 s_{2b+c+d} s_a^2 + s_{2a+2b} s_c s_d \left. \right) + \lambda_{123} \left(\sum^4 s_{2a} s_c s_{2b+d} \right. \\
& + 2 \sum^4 s_a s_{2b+c} s_{a+d} + 4s_{a+b} \sum^2 s_c s_{a+b+d} + 4 \sum^4 s_a s_{b+c} s_{a+b+d} \\
& + 2 \sum^4 s_c s_{a+2b} s_{a+d} + 2s_{c+d} \sum^2 s_a s_{a+2b} + 4s_{a+b} \sum^2 s_a s_{b+c+d} \\
& + 2 \sum^2 s_a s_{2b} s_{a+c+d} \left. \right) + \lambda_{23} \left(2s_{a+b}^2 s_{c+d} + 4 \sum^2 s_{a+b} s_{a+c} s_{b+d} \right. \\
& + 2 \sum^2 s_{2a} s_{b+c} s_{b+d} + s_{2a} s_{2b} s_{c+d} \left. \right) + \lambda_{133} \left(4 \sum^2 s_a s_b s_c s_{a+b+d} \right. \\
& + 2 \sum^2 s_a^2 s_b s_{b+c+d} + 2 \sum^2 s_a s_c s_d s_{a+2b} + \sum^4 s_a^2 s_c s_{2b+d} \left. \right) \\
& + \lambda_{122^2} \left(2 \sum^4 s_{2a} s_c s_b s_{b+d} + 2 \sum^2 s_a^2 s_{b+c} s_{b+d} \right. \\
& + s_{c+d} \sum^2 s_a^2 s_{2b} + 4 \sum^4 s_{a+b} s_{b+d} s_a s_c + 4 \sum^2 s_{a+d} s_{b+c} s_a s_b \\
& + 4s_a s_b s_{c+d} s_{a+b} + 2s_c s_d s_{a+b}^2 + s_{2a} s_{2b} s_c s_d \left. \right) \\
& + \lambda_{142} \left(4s_{a+b} s_a s_b s_c s_d + 2 \sum^4 s_a^2 s_b s_c s_{b+d} + s_{c+d} s_a^2 s_b^2 \right. \\
& + s_c s_d \sum^2 s_{2a} s_b^2 \left. \right) + \lambda_{16} s_a^2 s_b^2 s_c s_d,
\end{aligned}$$

$$\begin{aligned}
ES_a^3 S_b S_c S_d &= \lambda_6 s_{3a+b+c+d} + \lambda_{15} (3s_a s_{2a+b+c+d} + \sum^3 s_b s_{3a+c+d}) \\
&+ \lambda_{24} (3 \sum^3 s_{a+b} s_{2a+c+d} + 3s_{2a} s_{a+b+c+d} + \sum^3 s_{b+c} s_{3a+d}) \\
&+ \lambda_{33} (3 \sum^3 s_{2a+b} s_{a+c+d} + 3s_a s_{b+c+d}) + \lambda_{124} (3s_a \sum^3 s_b s_{2a+c+d} \\
&+ 3s_a^2 s_{a+b+c+d} + \sum^3 s_b s_c s_{3a+d}) + \lambda_{123} (s_{3a} \sum^3 s_b s_{c+d} + 3s_a s_{2a} s_{b+c+d} \\
&+ 3s_{2a} \sum^3 s_b s_{a+c+d} + 6s_a \sum^3 s_{a+b} s_{a+c+d} + 3s_a \sum^3 s_{b+c} s_{2a+d} \\
&+ 3 \sum^6 s_b s_{a+c} s_{2a+d}) + 3\lambda_{23} (s_{2a} \sum^3 s_{a+b} s_{c+d} + 2s_{a+d} s_{a+b} s_{a+c}) \\
&+ \lambda_{133} (3s_a \sum^3 s_{2a+b} s_c s_d + 3s_a^2 \sum^3 s_{a+b+c} s_d + s_{b+c+d} s_a^3 \\
&+ s_{3a} s_b s_c s_d) + 3\lambda_{122} (s_{2a} \sum^3 s_b s_c s_{a+d} + 2s_a \sum^3 s_b s_{a+c} s_{a+d} \\
&+ s_a^2 \sum^3 s_{a+b} s_{c+d} + s_a s_{2a} \sum^3 s_b s_{c+d}) + \lambda_{142} (3s_{2a} s_a s_b s_c s_d \\
&+ 3s_a^2 \sum^3 s_b s_c s_{a+d} + s_a^3 \sum^3 s_b s_{c+d}) + \lambda_{16} s_a^3 s_b s_c s_d, \\
ES_a^2 S_b S_c S_d S_e &= \lambda_6 s_{2a+b+c+d+e} + \lambda_{15} (2s_a s_{a+b+c+d+e} + \sum^4 s_b s_{2a+c+d+e}) \\
&+ \lambda_{24} (2 \sum^4 s_{a+b} s_{a+c+d+e} + s_{2a} s_{b+c+d+e} + \sum^6 s_{b+c} s_{2a+d+e}) \\
&+ \lambda_{33} (2 \sum^3 s_{a+b+c} s_{a+d+e} + \sum^4 s_{2a+b} s_{c+d+e}) \\
&+ \lambda_{124} (2s_a \sum^4 s_b s_{a+c+d+e} + s_{b+c+d+e} s_a^2 + \sum^6 s_{2a+d+e} s_b s_c) \\
&+ \lambda_{123} (2 \sum^{12} s_b s_{a+c} s_{a+d+e} + s_{2a} \sum^4 s_b s_{c+d+e} + 2s_a \sum^6 s_{b+c} s_{a+d+e} \\
&+ 2s_a \sum^4 s_{a+b} s_{c+d+e} + \sum^{12} s_b s_{c+d} s_{2a+e}) + \lambda_{23} (2 \sum^6 s_{a+b} s_{a+c} s_{d+e} \\
&+ s_{2a} \sum^3 s_{b+c} s_{d+e}) + \lambda_{133} (2s_a \sum^6 s_b s_c s_{a+b+c} + s_a^2 \sum^4 s_b s_{c+d+e} \\
&+ \sum^4 s_{2a+b} s_c s_d s_e) + \lambda_{122} (2 \sum^6 s_b s_c s_{a+d} s_{a+e} + s_{2a} \sum^6 s_b s_c s_{d+e} \\
&+ s_a^2 \sum^3 s_{b+c} s_{d+e} + 2s_a \sum^{12} s_b s_{a+c} s_{d+e}) + \lambda_{142} (2s_a \sum^4 s_{a+b} s_c s_d s_e \\
&+ s_a^2 \sum^6 s_{b+c} s_d s_e + s_{2a} s_b s_c s_d s_e) + \lambda_{16} s_{2a} s_b s_c s_d s_e.
\end{aligned}$$

References

- [1] Carver, H. C. (1930). Fundamentals of the theory of sampling. *Annals of Mathematical Statistics*, 1, 101-121.
- [2] Carver, H. C. (1930). Fundamentals of the theory of sampling. *Annals of Mathematical Statistics*, 1, 260-274.
- [3] Dwyer, P. S. and Tracy, D. S. (1980). Expectation and estimation of product moments in sampling from a finite population. *Journal of the American Statistical Association*, 75, 431-437.
- [4] Dwyer, P. S. (1938). On combined expansions of products of symmetric power sums and of sums of symmetric power products with applications to sampling (continued). *Annals of Mathematical Statistics*, 9, 97-132.
- [5] Fisher, R. A. (1929). Moments and product moments of sampling distributions. *Proceedings of the London Mathematical Society Series 2*, 30, 199-238.
- [6] James, G. S. (1958). On moments and cumulants of systems of statistics. *Sankhyā*, 20, 1-30.
- [7] Nath, S. N. (1968). On product moments from a finite universe. *Journal of the American Statistical Association*, 63, 535-541.
- [8] Nath, S. N. (1969). More results on product moments from a finite universe. *Journal of the American Statistical Association*, 64, 864-869.
- [9] Pierce, J. A. (1940). A study of a universe of n finite populations with application to moment-function adjustments for grouped data. *Annals of Mathematical Statistics*, 11, 311-334.
- [10] Raghunandan, K. and Srinivasan, R. (1973). Some product moments useful in sampling theory. *Journal of the American Statistical Association*, 68, 409-413.
- [11] Skellam, J. G. (1949). The distribution of moment statistics of samples drawn without replacement from a finite population. *Journal of the Royal Statistical Society, B*, 11, 291-296.
- [12] Stuart, A. and Ord, J. K. (1987). *Kendall's Advanced Theory of Statistics*, fifth edition, volume 1. Griffin, London.
- [13] Sukhatme, P. V. (1944). Moments and product moments of moment-statistics for samples of the finite and infinite populations. *Sankhyā*, 6, 363-382.
- [14] Tiit, E. (1988). Unbiased and n^{-k} -biased estimations of entire rational functions of moments. *Statistical and Probabilistic Models*, Report 798, Tartu State University, Tartu, Estonia, pp. 3-17.
- [15] Wishart, J. (1952). Moment coefficients of the k -statistics in samples from a finite population. *Biometrika*, 39, 1-13.