

Distributed Stochastic Approximation Algorithm With Expanding Truncations

Jinlong Lei, and Han-Fu Chen

Abstract—In this work, a novel distributed stochastic approximation algorithm (DSAA) is proposed to seek roots of the sum of local functions, each of which is associated with an agent from the multiple agents connected over a network. At each iteration, each agent updates its estimate for the root utilizing the noisy observations of its local function and the information derived from the neighboring agents. The key difference of the proposed algorithm from the existing ones consists in the expanding truncations (so it is called as DSAAWET), by which the boundedness of the estimates can be guaranteed without imposing the growth rate constraints on the local functions. The estimates generated by DSAAWET are shown to converge almost surely (a.s.) to a consensus set, which belongs to a connected subset of the root set of the sum function. In comparison with the existing results, we impose weaker conditions on the local functions and on the observation noise. We then apply the proposed algorithm to two applications, one from signal processing and the other one from distributed optimization, and establish the almost sure convergence. Numerical simulation results are also included.

Index Terms—Distributed stochastic approximation, expanding truncation, multi-agent network, distributed optimization.

I. INTRODUCTION

Distributed algorithms have been extensively investigated in connection with the problems arising from sensor networks and networked systems in recent years, such as consensus problem [1], [2], distributed estimation [3], [4], sensor localization [5], and distributed optimization [6]–[8]. The distributed algorithms work in the situation, where the goal is cooperatively accomplished by a multi-agent network with computation and communication abilities allocated in a distributed environment. Their advantages over the centralized approaches for networked problems lie in enhancing the robustness of the networks, preserving privacy, and reducing the communication and computation costs.

Stochastic approximation (SA) was firstly considered by Robbins and Monro in 1950s [9] for finding roots of a function with noisy observations, now it is known as the RM algorithm. Then SA was used by Kiefer and Wolfowitz [11] to estimate the maximum of an expectation-valued function only with noisy function observations. The SA-based schemes have found wide applications in signal processing, communications

and adaptive control, see, e.g., [35]–[37]. Recently, many distributed problems are also solved by SA-based distributed algorithms, e.g., distributed parameter estimation [3], [10], distributed convex optimization over random networks [17], and distributed non-convex stochastic optimization [18]. As such, the distributed variants of SA recently have drawn much attention from researchers.

Distributed stochastic approximation algorithms (DSAA) were proposed in [12], [13] to cooperatively find roots of a function, being a sum of local functions associated with agents connected over a network. Each agent updates its estimate for the root based on its local information composed of the observations of its local function possibly corrupted by noises, and the information obtained from its neighbors. The weak convergence for DSAA with a constant step-size is investigated in [12], while the a.s. convergence for DSAA with a decreasing step-size is studied in [13]. Besides, the performance gap between the distributed and the centralized stochastic approximation algorithms is investigated in [14]. The asynchronous and distributed stochastic approximation algorithms are also addressed in [15], [16] with the components separately estimated at different processors. However, it is noticed that almost all aforementioned SA-based distributed algorithms require rather restrictive conditions to guarantee convergence. For example, in [13] it is required that each local function is globally Lipschitz continuous and the observation noise is a martingale difference sequence (MDS)¹. However, these conditions may not hold for some problems, e.g., the distributed principal component analysis and the distributed gradient-free optimization to be discussed in Section II.B. This work aims at resolving the distributed root-seeking problem under weaker conditions compared with those used in [13].

Contributions of this work are as follows. 1) We propose a novel DSAA with expanding truncations (DSAAWET) over networks with deterministic switching topologies. The key difference from the existing algorithms (cf. [12], [13]) is the network expanding truncation mechanism that adaptively defines enclosing bounds as follows: the initial value lies within the initial bound; when the estimate crosses the enclosing bound, it is reinitialized to the initial point and the bound is enlarged. This together with the decreasing step-sizes makes as if the algorithm returns with the same initial value but with smaller step-sizes and larger bounds. Then it is shown that the reinitialization ceases in a finite number of steps and the estimates are bounded. Further, estimates

Jinlong Lei is with the Department of Industrial and Manufacturing Engineering, Pennsylvania State University, University Park, PA 16802, USA (e-mail: jxl800@psu.edu, lejijinlong11@mails.ucas.ac.cn).

Han-Fu Chen is with the Key Laboratory of Systems and Control, Institute of Systems Science, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, P. R. China. (e-mail: hfchen@iss.ac.cn).

This work was supported by the National Center for Mathematics and Interdisciplinary Sciences, Chinese Academy of Sciences.

¹Consider an adapted sequence $\{X_t, \mathcal{F}_t\}_{t=0}^{\infty}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. X_t is an MDS if it satisfies $\mathbb{E}[X_t] < \infty$ and $\mathbb{E}[X_t | \mathcal{F}_{t-1}] = 0$, a.s., for all t .

generated by DSAAWET at all agents converge a.s. to a consensus set, which is contained in the root set of the sum function. Compared with [13], we neither assume the observation noise to be an MDS nor impose any growth rate constraint and Lipschitz continuity on the local functions. 2) The proposed algorithm is then applied to the distributed principal component analysis and to the distributed gradient-free optimization. Their a.s. convergence is established as well, while the algorithms in [12], [13] might not be applicable to these problems. The proposed method has also been applied in [19] and [20] to the distributed blind channel identification and the output consensus of networked nonlinear systems.

The notations used in this work are listed in Table I and the rest of this work is arranged as follows. The distributed root-seeking problem is formulated in Section II along with two motivation examples. DSAAWET is defined in Section III and the corresponding convergent results are presented as well. The proof of the main results is given in Section IV with some details placed in Appendices. The proposed algorithm is applied to solve the two application problems in Section V with numerical examples included. Some concluding remarks are given in Section VI.

Symbol	Definition
$\ v\ , \ A\ $	Euclidean (l^2) norm of vector v , matrix A
$A \geq 0$	Each entry of matrix A is nonnegative, and A is called the nonnegative matrix.
\mathbf{I}_m	$m \times m$ identity matrix
$\mathbf{1}, \mathbf{0}$	Vector or matrix with all entries equal to 1, 0
X^T, X^{-1}	Transpose of matrix X , inverse of matrix X
$col\{x_1, \dots, x_m\}$	$col\{x_1, \dots, x_m\} \triangleq (x_1^T, \dots, x_m^T)^T$ stacking the vectors or matrices x_1, \dots, x_m
$I_{[\text{Inequality}]}$	Indicator function meaning it equals 1 if the inequality holds and 0 if the inequality does not hold
\otimes	Kronecker product
$a \wedge b$	$\min\{a, b\}$
J	$J \triangleq \{x \in \mathbb{R}^l : f(x) = 0\}$ denotes the root set of $f(\cdot)$
$d(x, J)$	$\min_y \{\ x - y\ : y \in J\}$
$E[\cdot]$	Expectation operator
D_{\perp}	$D_{\perp} \triangleq (\mathbf{I}_N - \frac{\mathbf{1}\mathbf{1}^T}{N}) \otimes \mathbf{I}_l$, where N denotes the number of agents in the network and l denotes the dimension of the root
$\delta(t)$	$\delta(t) \rightarrow 0$ as $t \rightarrow 0$.
$m(k, T)$	$m(k, T) \triangleq \max\{m : \sum_{i=k}^m \gamma_i \leq T\}$ is an integer-valued function for $T > 0$ and integer k
$\sigma_{i,k}$	The truncation number of agent i at time k
$\hat{\sigma}_{i,k}$	$\hat{\sigma}_{i,k} \triangleq \max_{j \in N_i(k)} \sigma_{j,k}$, where $N_i(k)$ is the set of neighboring agents of agent i at time k
σ_k	The largest truncation number among all agents at time k , i.e., $\sigma_k = \max_{i \in \mathcal{V}} \sigma_{i,k} = \max_{i \in \mathcal{V}} \hat{\sigma}_{i,k}$.
$\tau_{i,m}$	$\tau_{i,m} \triangleq \inf\{k : \sigma_{i,k} = m\}$
τ_m	$\tau_m \triangleq \min_{i \in \mathcal{V}} \tau_{i,m}$, the smallest time when at least one of agents has its truncation number reached m
$\tilde{\tau}_{j,m}$	$\tilde{\tau}_{j,m} \triangleq \tau_{j,m} \wedge \tau_{m+1}$

TABLE I: Notations

II. PROBLEM FORMULATION AND MOTIVATIONS

In this section, we first formulate the distributed root-seeking problem with the related communication model. Then we give two motivation problems that cannot be solved by the existing algorithms, but can be solved by DSAAWET to be proposed in this work.

A. Problem Statement

Consider the case where all agents in a network collectively search the root of the sum function given by

$$f(\cdot) = \frac{1}{N} \sum_{i=1}^N f_i(\cdot), \quad (1)$$

where $f_i(\cdot) : \mathbb{R}^l \rightarrow \mathbb{R}^l$ is the local function assigned to agent i and can only be observed by agent i . Let $J \triangleq \{x \in \mathbb{R}^l : f(x) = 0\}$ denote the root set of $f(\cdot)$.

Time is slotted at $k = 0, 1, 2, \dots$. For any $i \in \mathcal{V}$, denote by $x_{i,k} \in \mathbb{R}^l$ the estimate for the root of $f(\cdot)$ given by agent i at time k . Agent i at time $k+1$ has its local noisy observation

$$O_{i,k+1} = f_i(x_{i,k}) + \varepsilon_{i,k+1}, \quad (2)$$

where $\varepsilon_{i,k+1}$ is the observation noise. Agent i is required to update its estimate $x_{i,k}$ on the basis of its local observation and the information obtained from its neighbors.

The information exchange among the N agents at time k is described by a digraph $\mathcal{G}(k) = \{\mathcal{V}, \mathcal{E}(k)\}$, where $\mathcal{V} = \{1, \dots, N\}$ is the node set with node i representing agent i ; $\mathcal{E}(k) \subset \mathcal{V} \times \mathcal{V}$ is the edge set with $(j, i) \in \mathcal{E}(k)$ if agent i can obtain information from agent j at time k . Let the associated adjacency matrix be denoted by $W(k) = [\omega_{ij}(k)]_{i,j=1}^N$, where $\omega_{ij}(k) > 0$ if and only if $(j, i) \in \mathcal{E}(k)$, and $\omega_{ij}(k) = 0$, otherwise. Denote by $N_i(k) \triangleq \{j \in \mathcal{V} : (j, i) \in \mathcal{E}(k)\}$ the set of neighboring agents of agent i at time k .

A time-independent digraph $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ is called strongly connected if for any $i, j \in \mathcal{V}$ there exists a directed path from i to j . By this we mean a sequence of edges $(i, i_1), (i_1, i_2), \dots, (i_{p-1}, j)$ in the digraph with distinct nodes $i_m \in \mathcal{V} \quad \forall m : 0 \leq m \leq p-1$, where p is called the length of the directed path. A nonnegative square matrix A is called doubly stochastic if $A\mathbf{1} = \mathbf{1}$ and $\mathbf{1}^T A = \mathbf{1}^T$.

B. Motivation Examples

We now give two motivation examples that cannot be solved by the existing distributed stochastic approximation algorithm [13]. We will return to these examples in Section V to show that DSAAWET to be proposed in this work can solve them.

1) *Distributed Principal Component Analysis*: In signal processing and pattern recognition, effective clustering of large data sets is an important objective. Principal component analysis (PCA) is a powerful technique to process the multivariate data by constructing a concise data representation through computing the dominant eigenvalues and the corresponding eigenvectors of the data covariance matrix [21]. For recent years, the distributed computation of PCA has attracted much attention from researchers (cf. [22]–[27]). Roughly speaking, there are two different types of distributed PCA: (i) Samples composed of all features collected from the wireless sensor networks (WSNs) [22], [23]. (ii) Agents in the network collect observations for different features of the whole features [24]–[27]. The global PCA is computed in [22] for distributed updating data sets without convergence analysis, while the distributed PCA of a covariance matrix of the fixed data sets is addressed in [23]. The authors of [24] concentrate on computing principal components of a covariance matrix observed with noise, propose an algorithm with each sensor estimating the

corresponding entry of the principal eigenvector, and show that the associated ODE is close to that of Oja's [21]. A distributed PCA algorithm is proposed on the directed Gaussian graphical models in [25]. The goal of [26], [27] is to distributively compute the principal components of a symmetric weighted adjacency matrix associated with the connected network.

We consider the distributed setting similar to that considered in [22], [23] aiming at proposing a distributed algorithm for the global PCA and establishing its a.s. convergence. The global data matrix u_k at time k is distributed among N sensors that are spatially distributed in the network:

$$u_k = \begin{pmatrix} u_{1,k} \\ \cdots \\ u_{N,k} \end{pmatrix},$$

where $u_{i,k} \in \mathbb{R}^{p_i \times l}$ is collected by sensor i . The rows of u_k denote the observations while the columns denote the features. Assume that for each $i \in \mathcal{V}$, $\{u_{i,k}\}$ is an i.i.d. sequence with zero mean. Denote by $A \triangleq E[u_k^T u_k] \in \mathbb{R}^{l \times l}$ the covariance matrix of u_k . The primary objective is to estimate the unit eigenvector corresponding to the largest eigenvalue of A . Because the unit eigenvector corresponding to the largest eigenvalue of A belongs to the set of the nonzero roots of the function $f(x) = Ax - (x^T Ax)x$ on the unit sphere, then the distributed PCA might be solved by virtue of finding roots of the sum function defined as follows:

$$f(x) = \sum_{i=1}^N f_i(x),$$

where $f_i(x) = A_i x - (x^T A_i x)x$ with $A_i \triangleq E[u_{i,k}^T u_{i,k}]$.

Denote by $x_{i,k}$ the estimate given by agent i at time k for the unit eigenvector corresponding to the largest eigenvalue of matrix A . Since A_i cannot be directly derived, by replacing A_i with its sample data $A_{i,k} = u_{i,k}^T u_{i,k}$ at time k , the local observation of agent i at time $k+1$ is defined as follows:

$$O_{i,k+1} \triangleq A_{i,k} x_{i,k} - (x_{i,k}^T A_{i,k} x_{i,k}) x_{i,k}. \quad (3)$$

Then the distributed principal component analysis is in the distributed root-seeking form. Noting that $f_i(\cdot)$ contains a cubic term, the local function is not globally Lipschitz continuous. Though the existing distributed stochastic approximation algorithms might be still applicable since eigenvectors can be chosen to have unit-norm by projecting the estimates onto the unit hypersphere, the exact convergence to the true eigenvector for the distributed data set with updating has not been fully addressed yet. In Section V, we will show that DSAAWET can solve this problem and ensure the a.s. convergence.

2) *Distributed Gradient-free Optimization*: Consider a multi-agent network of N agents, for which the objective is to cooperatively solve the following optimization problem:

$$\min_x c(x) = \sum_{i=1}^N c_i(x), \quad (4)$$

where $c_i(\cdot) : \mathbb{R}^l \rightarrow \mathbb{R}$ is the local objective function of agent i , and $c_i(\cdot)$ is only known by i itself. Assume the optimization problem (4) has solutions. Consider the case where gradients of the cost functions are unavailable but the cost functions can be observed with noises. Then the finite time difference of the cost functions can be adopted to estimate the gradient, see [11], [28], [29]. This problem is referred to as the gradient-free optimization [30]. The gradient-free methods for nonsmooth distributed optimization with convex set constraint are considered in [31] [32], where it is assumed that each local objective function is convex and Lipschitz continuous over the convex set. The underlying network topology is modeled as time varying in [31] while as fixed in [32]. In both [31] and [32], the Gaussian smoothing technique is used, and the convergence to an approximate solution within the error level depending on the smoothing parameter and the Lipschitz constant of local objective function is established. In contrast, we consider the distributed smooth optimization with the aim of designing a distributed gradient-free algorithm that can achieve a.s. convergence while without requiring each objective function be convex and Lipschitz continuous.

Assume that the cost functions $c_i(\cdot)$, $i = 1, \dots, N$ are differentiable and the global function $c(\cdot)$ is convex. Then solving the problem (4) is equivalent to finding roots of the function $f(x) = \sum_{i=1}^N f_i(x)/N$ with $f_i(\cdot) \triangleq -\nabla c_i(\cdot)$, where $\nabla c_i(\cdot)$ denotes the gradient function of $c_i(\cdot)$. Here the randomized KW method proposed in [28] is adopted to estimate gradients of the cost functions. Denote by $x_{i,k}$ the estimate of the solution to the problem (4) given by agent i at time k . Let $\Delta_{i,k} \in \mathbb{R}^l$, $k = 1, 2, \dots$ be a sequence of mutually independent random vectors with each component independently taking values ± 1 with probability $\frac{1}{2}$. The observation of function $-\nabla c_i(\cdot)$ at point $x_{i,k}$ is constructed as

$$O_{i,k+1} = -\frac{c_i(x_{i,k}^+) + \xi_{i,k+1}^+ - c_i(x_{i,k}^-) - \xi_{i,k+1}^-}{2\alpha_k} \Delta_{i,k}, \quad (5)$$

where $x_{i,k}^+ = x_{i,k} + \alpha_k \Delta_{i,k}$, $x_{i,k}^- = x_{i,k} - \alpha_k \Delta_{i,k}$, $\alpha_k > 0$ for any $k \geq 0$, $\xi_{i,k+1}^+$ and $\xi_{i,k+1}^-$ are the observation noises of the cost function $c_i(\cdot)$ at points $x_{i,k}^+$ and $x_{i,k}^-$, respectively.

Thus, we have transformed the distributed gradient-free optimization to the distributed root-seeking form. It is shown

$$\sigma_{i,0} = 0, \quad \hat{\sigma}_{i,k} \triangleq \max_{j \in N_i(k)} \sigma_{j,k}, \quad (6)$$

$$x'_{i,k+1} = \left(\sum_{j \in N_i(k)} \omega_{ij}(k) (x_{j,k} I_{[\sigma_{j,k} = \hat{\sigma}_{i,k}]} + x^* I_{[\sigma_{j,k} < \hat{\sigma}_{i,k}]} + \gamma_k O_{i,k+1}) I_{[\sigma_{i,k} = \hat{\sigma}_{i,k}]} + x^* I_{[\sigma_{i,k} < \hat{\sigma}_{i,k}]} \right), \quad (7)$$

$$x_{i,k+1} = x^* I_{[\|x'_{i,k+1}\| > M\hat{\sigma}_{i,k}]} + x'_{i,k+1} I_{[\|x'_{i,k+1}\| \leq M\hat{\sigma}_{i,k}]}, \quad (8)$$

$$\sigma_{i,k+1} = \hat{\sigma}_{i,k} + I_{[\|x'_{i,k+1}\| > M\hat{\sigma}_{i,k}]}, \quad (9)$$

in [28] that for any $i \in \mathcal{V}$, the observation noise $\varepsilon_{i,k+1} = O_{i,k+1} + \nabla c_i(x_{i,k})$ is not an MDS, and hence DSAA proposed in [13] cannot be directly applied to this problem.

The problems arising from this kind of examples motivate us to propose DSAAWET, by which the two motivation examples can be resolved as to be shown in Section V.

III. DSAAWET AND ITS CONVERGENCE

In this section, we define DSAAWET and formulate the main results of this work.

A. DSAAWET

Let us first explain the idea of expanding truncations. In many important problems, the sequence of estimates generated by the RM algorithm may not be bounded, and it is hard to define in advance a region the sought-for parameter belongs to. This motivates us to adaptively define truncation bounds as follows. When the estimate crosses the current truncation bound, the estimate is reinitialized to some pre-specified point x^* and at the same time the truncation bound is enlarged. Due to the decreasing step-sizes, this operation makes as if we rerun the algorithm with initial value x^* , with smaller step-sizes and larger truncation bounds. The expanding truncation mechanism incorporating with some verifiable conditions makes the reinitialization cease in a finite number of steps, and hence makes the estimates bounded. As results, after a finite number of steps the algorithm runs as the RM algorithm. This is well explained in [36] and the related references therein.

We apply the idea of expanding truncation to the distributed estimation, i.e., for the case $N > 1$. Let $x_{i,k}$ denote agent i ' estimate for the root of $f(\cdot)$ at time k , and $\sigma_{i,k}$ denote the number of truncations for agent i up-to-time k . We now define DSAAWET (Eqns. (6)-(9)) with any initial values $x_{i,0}$, where $O_{i,k+1}$ is defined by (2), $\{\gamma_k\}$ is the steplength, $x^* \in \mathbb{R}^l$ is a fixed vector known to all agents, $\{M_k\}_{k \geq 0}$ is a sequence of positive numbers increasingly diverging to infinity with $M_0 \geq \|x^*\|$, and $M_{\hat{\sigma}_{i,k}}$ serves as the the truncation bound when the $(k+1)$ th estimate for agent i is generated. Note that during an iteration, each agent uses its neighbors' information including estimates for the root and the truncation number.

The algorithm (6)-(9) is performed according to the following three steps.

(i) **Max consensus of truncation numbers** (Eqn. (6)). At time k , the update of agent i may or may not be truncated. This yields that at time k , $\sigma_{i,k}$ may be different for $i = 1, \dots, N$. We then set the truncation number $\hat{\sigma}_{i,k}$ to be the largest one among $\{\sigma_{j,k}, j \in \mathcal{N}_i(k)\}$ of its neighboring agents as indicated by (6). It is worth noting that we do not require the truncation number of each agent to reach a max consensus in finite time. Nevertheless, we will prove in Lemmas 4.9 and 4.10 that all agents' truncation numbers will asymptotically reach a consensus to some positive integer.

(ii) **Average consensus + innovation update** (Eqn. (7)). At time $k+1$, agent i produces an intermediate value $x'_{i,k+1}$ by (7). If $\sigma_{i,k} < \hat{\sigma}_{i,k}$, then $x'_{i,k+1} = x^*$. Otherwise, $x'_{i,k+1}$ is a combination of the consensus part and the innovation part, where the consensus part is a weighted average of estimates

derived at its neighbors, and the innovation part processes its local current observation.

(iii) **Local truncation judgement** (Eqns. (8)-(9)). If $x'_{i,k+1}$ remains inside its local truncation bound $M_{\hat{\sigma}_{i,k}}$, i.e., $\|x'_{i,k+1}\| \leq M_{\hat{\sigma}_{i,k}}$, then $x_{i,k+1} = x'_{i,k+1}$ and $\sigma_{i,k+1} = \hat{\sigma}_{i,k}$. If $x'_{i,k+1}$ exits from the sphere with radius $M_{\hat{\sigma}_{i,k}}$, i.e., $\|x'_{i,k+1}\| > M_{\hat{\sigma}_{i,k}}$, then $x_{i,k+1}$ is pulled back to the pre-specified point x^* , meanwhile, the truncation number $\sigma_{i,k+1}$ is increased to $\hat{\sigma}_{i,k} + 1$.

Denote by

$$\sigma_k \triangleq \max_{i \in \mathcal{V}} \sigma_{i,k}, \quad (10)$$

the largest truncation number among all agents at time k . If $N = 1$, then by denoting $\hat{\sigma}_{i,k} = \sigma_{i,k} \triangleq \sigma_k$, $x_{i,k} \triangleq x_k$, $x'_{i,k} \triangleq x'_k$, $O_{i,k} \triangleq O_k$, it is seen that (6)-(9) becomes

$$\begin{aligned} x'_{k+1} &= x_k + \gamma_k O_{k+1}, \\ x_{k+1} &= x^* I_{[\|x'_{k+1}\| > M_{\sigma_k}]} + x'_{k+1} I_{[\|x'_{k+1}\| \leq M_{\sigma_k}]}, \\ \sigma_{k+1} &= \sigma_k + I_{[\|x'_{k+1}\| > M_{\sigma_k}]}. \end{aligned}$$

The above algorithm is called the stochastic approximation algorithm with expanding truncations (SAAWET), which requires possibly the weakest conditions for its convergence among various modifications of the RM algorithm [33]–[36]. Thus, the advantages of SAAWET over the RM algorithm might remain for its distributed variant in the case $N > 1$.

Remark 3.1: It is noticed that $\sigma_{i,k+1} \geq \hat{\sigma}_{i,k} \geq \sigma_{i,k} \quad \forall k \geq 0$ by (6) and (9). Further, it is concluded that

$$x_{i,k+1} = x^* \text{ if } \sigma_{i,k+1} > \sigma_{i,k}. \quad (11)$$

This can be seen from the following consideration: i) If $\sigma_{j,k} \leq \sigma_{i,k} \quad \forall j \in \mathcal{N}_i(k)$, then from (6) we derive $\hat{\sigma}_{i,k} = \sigma_{i,k}$. Since $\sigma_{i,k+1} > \sigma_{i,k}$, by (9) it follows that $\|x'_{i,k+1}\| > M_{\hat{\sigma}_{i,k}}$, and hence from (8) we derive $x_{i,k+1} = x^*$. ii) If there exists $j \in \mathcal{N}_i(k)$ such that $\sigma_{j,k} > \sigma_{i,k}$, then from (6) we derive $\hat{\sigma}_{i,k} = \max_{j \in \mathcal{N}_i(k)} \sigma_{j,k} > \sigma_{i,k}$, and from (7) we have $x'_{i,k+1} = x^*$. Consequently, by (8) we have $x_{i,k+1} = x^*$.

B. Assumptions

We list the assumptions to be used.

A1 $\gamma_k > 0, \gamma_k \xrightarrow[k \rightarrow \infty]{} 0$, and $\sum_{k=1}^{\infty} \gamma_k = \infty$.

A2 There exists a continuously differentiable function $v(\cdot) : \mathbb{R}^l \rightarrow \mathbb{R}$ such that

$$\text{a) } \sup_{\delta \leq d(x,J) \leq \Delta} f^T(x) v_x(x) < 0 \quad (12)$$

for any $\Delta > \delta > 0$, where $v_x(\cdot)$ denotes the gradient of $v(\cdot)$ and $d(x, J) = \min_y \{\|x - y\| : y \in J\}$,

b) $v(J) \triangleq \{v(x) : x \in J\}$ is nowhere dense,

c) $\|x^*\| < c_0$ and $v(x^*) < \inf_{\|x\|=c_0} v(x)$ for some positive constant c_0 , where x^* is used in (7) (8).

A3 The local functions $f_i(\cdot) \quad \forall i \in \mathcal{V}$ are continuous.

Let us explain the conditions. A1 is a standard assumption for stochastic approximation, see [34], [36]. A2 a) means that $v(\cdot)$ serves as a Lyapunov function for the differential equation $\dot{x} = f(x)$. It is noticed that A2 b) holds if J is finite, and A2 c) takes place if $v(\cdot)$ is radially unbounded.

The following condition A4 imposed on the communication graphs is taken from [6], which the readers are referred to for detailed explanations.

A4 a) For any $k \geq 0$, $W(k)$ is a doubly stochastic matrix;

b) There exists a constant $0 < \eta < 1$ such that

$$\omega_{ij}(k) \geq \eta \quad \forall j \in \mathcal{N}_i(k) \quad \forall i \in \mathcal{V} \quad \forall k \geq 0;$$

c) The digraph $\mathcal{G}_\infty = \{\mathcal{V}, \mathcal{E}_\infty\}$ is strongly connected, where \mathcal{E}_∞ is the set of edges (j, i) such that j is a neighbor of i which communicates with i infinitely often, i.e.,

$$\mathcal{E}_\infty = \{(j, i) : (j, i) \in \mathcal{E}(k) \text{ for infinitely many indices } k\};$$

d) There exists a positive integer B such that for every $(j, i) \in \mathcal{E}_\infty$, agent j sends the information to the neighbor i at least once every B consecutive time slots, i.e.,

$$(j, i) \in \mathcal{E}(k) \cup \mathcal{E}(k+1) \cup \dots \cup \mathcal{E}(k+B-1)$$

for all $(j, i) \in \mathcal{E}_\infty$ and any $k \geq 0$.

Remark 3.2: A4 a) requires the adjacency matrices of the digraphs to be doubly stochastic, which does not hold for all digraphs. Nevertheless, a necessary and sufficient condition is provided in [43] for a digraph to be doubly stochasticable. A4 b) states that each agent gives significant weights to its own and the neighbors' states. It is explained in [6] that A4 c) is equivalent to the assumption that the composite directed graph $\{\mathcal{V}, \cup_{l \geq k} \mathcal{E}_k\}$ is strongly connected for all k . A4 d) implies that for any $i, j \in \mathcal{V}$, the information of j can be propagated to i in less than $B(N-1)$ steps. Define $\Phi(k, k+1) = \mathbf{I}_N$ and $\Phi(k, s) = W(k) \cdots W(s)$ for all $k \geq s$. Then by [6, Proposition 1], there exist $c > 0$ and $0 < \rho < 1$ such that

$$\|\Phi(k, s) - \frac{1}{N} \mathbf{1}\mathbf{1}^T\| \leq c\rho^{k-s+1} \quad \forall k \geq s. \quad (13)$$

Before introducing conditions on noise, let us denote by $(\Omega, \mathcal{F}, \mathbb{P})$ the probability space. For any $i \in \mathcal{V}$, let $\varepsilon_{i,k+1} : (\mathbb{R}^l \times \Omega, \mathbb{B}^l \times \mathcal{F}) \rightarrow (\mathbb{R}^l \times \mathbb{B}^l)$ be a measurable function, where \mathbb{B} is a Borel σ -algebra. Let the noise be given by $\varepsilon_{i,k+1} = \varepsilon_{i,k+1}(x_{i,k}(\omega), \omega)$, $\omega \in \Omega$. Thus, the state-dependent noise is considered. For notational simplicity, $\varepsilon_{i,k+1}$ is used to denote $\varepsilon_{i,k+1}(x_{i,k}(\omega), \omega)$ for the sample path $\omega \in \Omega$ under consideration throughout this work.

A5 For the sample path ω under consideration, the following assertions hold for any $i \in \mathcal{V}$:

a) $\gamma_k \varepsilon_{i,k+1} \xrightarrow[k \rightarrow \infty]{} 0$, and

b) $\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{m=n_k}^{m(n_k, t_k)} \gamma_m \varepsilon_{i,m+1} I_{[\|x_{i,m}\| \leq K]} \right\| = 0$
 $\forall t_k \in [0, T]$ for any sufficiently large K

along indices $\{n_k\}$ whenever $\{x_{i,n_k}\}$ converges, where $m(k, T) \triangleq \max\{m : \sum_{i=k}^m \gamma_i \leq T\}$.

Remark 3.3: It is noticed that A5 b) is convenient for dealing with state-dependent noise. The indicator function $I_{[\|x_{i,m}\| \leq K]}$ in the condition will be casted away if the observation noise does not depend on the estimates. However, for the state-dependent noise, before establishing the boundedness of $\{x_{i,k}\}$, the condition with an indicator function included is

easier to be verified. It is worth noting that in A5 b) we do not assume existence of a convergent subsequence of $\{x_{i,k}\}$ for any i , we only require A5 b) hold along indices of any convergent subsequence if exists. Verification of A5 b) along convergent subsequences is much easier than that along the whole sequence. If $\{\varepsilon_{i,k}\}$ can be decomposed into two parts $\varepsilon_{i,k} = \varepsilon_{i,k}^{(1)} + \varepsilon_{i,k}^{(2)}$ such that $\sum_{k=0}^{\infty} \gamma_k \varepsilon_{i,k+1} I_{[\|x_{i,k}\| \leq K]} < \infty$ and $\varepsilon_{i,k}^{(2)} I_{[\|x_{i,k}\| \leq K]} \xrightarrow[k \rightarrow \infty]{} 0$, then A5 b) holds. So, A5 holds when the observation noise is an i.i.d. sequence or an MDS with bounded second moments if $\sum_{k=1}^{\infty} \gamma_k^2 < \infty$.

C. Main Results

Define the vectors $X_k \triangleq \text{col}\{x_{1,k}, \dots, x_{N,k}\}$, $\varepsilon_k \triangleq \text{col}\{\varepsilon_{1,k}, \dots, \varepsilon_{N,k}\}$, $F(X_k) \triangleq \text{col}\{f_1(x_{1,k}), \dots, f_N(x_{N,k})\}$. Denote by $X_{\perp,k} \triangleq D_{\perp} X_k$ the disagreement vector of X_k with $D_{\perp} \triangleq (\mathbf{I}_N - \frac{11^T}{N}) \otimes \mathbf{I}_l$, and by $x_k = \frac{1}{N} \sum_{i=1}^N x_{i,k}$ the average of the estimates derived at all agents at time k .

Theorem 3.4: Let $\{x_{i,k}\}$ be produced by (6)-(9) with an arbitrary initial value $x_{i,0}$. Assume A1-A4 hold. Then for any sample path ω where A5 holds, we have the following:

i) $\{x_{i,k}\}$ is bounded and there exists a positive integer k_0 possibly depending on ω such that

$$x_{i,k+1} = \sum_{j \in \mathcal{N}_i(k)} \omega_{ij}(k) x_{j,k} + \gamma_k O_{i,k+1} \quad \forall k \geq k_0, \quad (14)$$

or in the following compact form:

$$X_{k+1} = (W(k) \otimes \mathbf{I}_l) X_k + \gamma_k (F(X_k) + \varepsilon_{k+1}) \quad \forall k \geq k_0; \quad (15)$$

ii) $X_{\perp,k} \xrightarrow[k \rightarrow \infty]{} \mathbf{0}$ and $d(x_k, J) \xrightarrow[k \rightarrow \infty]{} 0$; (16)

iii) there exists a connected subset $J^* \subset J$ such that

$$d(x_k, J^*) \xrightarrow[k \rightarrow \infty]{} 0. \quad (17)$$

The proof of Theorem 3.4 is presented in Section IV. It is noticed from the proof that for deriving i), condition A5 a) is not required. Theorem 3.4 establishes that the sequence $\{X_k\}$ is bounded; the algorithm (6)-(9) finally turns to a standard DSAA without truncations; and the estimates given by all agents converge a.s. to a consensus set, contained in a connected subset of the root set J , when A5 holds for almost all sample paths for all agents. As a consequence, if J is not dense in any connected set, then x_k converges to a point in J . However, it is unclear how x_k behaves when J is dense in some connected set. This problem was investigated for the centralized algorithm in [42]. Based on Eqn. (14), we might be able to investigate the rate of convergence of DSAAWET similar to the standard DSAA, while we leave it for future research due to the page limitation.

Remark 3.5: Compared with [13], we impose weaker conditions on the local functions and on the observation noise. In fact, we only require the local functions be continuous, while conditions ST1 and ST2 in [13, Theorem 3] do not allow the functions to grow faster than linearly. In addition, we do not require the observation noise to be an MDS as in [13]. As shown in [36], [37], A5 is probably the weakest requirement

for the noise since it is also necessary for convergence whenever the root x^0 of $f(\cdot)$ is a singleton and $f(\cdot)$ is continuous at x^0 . Compared with the random communication graphs used in [13], here we use the deterministic switching graphs to describe the communication relationships among agents.

IV. PROOF OF MAIN RESULTS

Prior to analyzing $\{x_{i,k}\}$, let us recall the convergence analysis for SAAWET, i.e., DSAWET with $N = 1$. The key step in the analysis is to establish the boundedness of the estimates, or to show that truncations cease in a finite number of steps. If the number of truncations increases unboundedly, then SAAWET is pulled back to a fixed vector x^* infinitely many times. This produces convergent subsequences from the estimation sequence. Then the condition A5 b) is applicable and it incorporating with A2 yields a contradiction. This proves the boundedness of the estimates.

Let us try to use this approach to prove the boundedness of $x_k = \frac{1}{N} \sum_{i=1}^N x_{i,k}$ with $\{x_{i,k}\}$ generated by (6)-(9). In the case $\sigma_k \xrightarrow[k \rightarrow \infty]{} \infty$, we have $\lim_{k \rightarrow \infty} \sigma_{i,k} = \infty \quad \forall i \in \mathcal{V}$ by Corollary 4.3 given below. Then from Remark 3.1 it is known that the estimate $x_{i,k+1}$ given by agent i is pulled back to x^* when the truncation occurs at time $k+1$. This means that $\{x_{i,k}\} \forall i \in \mathcal{V}$ contains convergent subsequences. However, $\{x_k\}$ may still not contain any convergent subsequence to make A5 b) applicable. This is because truncations may occur at different times for different $i \in \mathcal{V}$. Therefore, the conventional approach used for convergence analysis of SAAWET cannot directly be applied to the algorithm (6)-(9).

To overcome the difficulty, we first introduce auxiliary sequences $\{\tilde{x}_{i,k}\}$ and $\{\tilde{\varepsilon}_{i,k+1}\}$ for any $i \in \mathcal{V}$. It will be shown in Lemma 4.1 that $\{\tilde{x}_{i,k}\}$ satisfies the recursions (20)-(22), for which the truncation bound at time k is the same M_{σ_k} for all agents and the estimates $\tilde{x}_{i,k+1} \forall i \in \mathcal{V}$ are pulled back to x^* when $\sigma_{k+1} > \sigma_k$. As a result, the auxiliary sequence $\{\tilde{X}_k\}$ has convergent subsequences, where $\tilde{X}_k \triangleq \text{col}\{\tilde{x}_{1,k}, \dots, \tilde{x}_{N,k}\}$. Besides, it will be shown in Lemma 4.5 that the noise $\{\tilde{\varepsilon}_k\}$ satisfies a condition similar to A5 b) along any convergent subsequence of $\{\tilde{X}_k\}$, where $\tilde{\varepsilon}_k \triangleq \text{col}\{\tilde{\varepsilon}_{1,k}, \dots, \tilde{\varepsilon}_{N,k}\}$. To borrow the analytical method from the centralized stochastic approximation algorithm, we rewrite the algorithm (20)-(22) in the centralized form (34) with observation noise $\{\zeta_{m+1}\}$. By the results given in Lemma 4.1 and Lemma 4.5, it is shown in Lemma 4.7 that the noise sequence $\{\zeta_{m+1}\}$ satisfies (35) along convergent subsequences, which is similar to A5 b) when $N = 1$. Then by algorithm (34) and the noise property (35), we show that the numbers of truncations for all agents converge to the same finite value, and that $\{\tilde{x}_{i,k}\}_{i \in \mathcal{V}}$ reach a consensus located in the root set. Thus, $\{x_{i,k}\}$ and $\{\tilde{x}_{i,k}\}$ coincide in a finite number of steps, and their convergence is equivalent. In the rest of this section, we demonstrate in detail the aforementioned ideas.

A. Auxiliary Sequences

Denote by $\tau_{i,m} \triangleq \inf\{k : \sigma_{i,k} = m\}$ the smallest time when the truncation number of agent i has reached m , and

by $\tau_m \triangleq \min_{i \in \mathcal{V}} \tau_{i,m}$ the smallest time when at least one agent has its truncation number reached m . Set $\tilde{\tau}_{j,m} \triangleq \tau_{j,m} \wedge \tau_{m+1}$, where $a \wedge b = \min\{a, b\}$.

For any $i \in \mathcal{V}$, we construct two auxiliary sequences $\{\tilde{x}_{i,k}\}_{k \geq 0}$ and $\{\tilde{\varepsilon}_{i,k+1}\}_{k \geq 0}$. Note that for the considered ω , for any $k \geq 0$ there exists a unique integer $m \geq 0$ such that $k \in [\tau_m, \tau_{m+1})$. We then define

$$\tilde{x}_{i,k} \triangleq x^*, \quad \tilde{\varepsilon}_{i,k+1} \triangleq -f_i(x^*) \quad \text{if } k \in [\tau_m, \tilde{\tau}_{i,m}), \quad (18)$$

$$\tilde{x}_{i,k} \triangleq x_{i,k}, \quad \tilde{\varepsilon}_{i,k+1} \triangleq \varepsilon_{i,k+1} \quad \text{if } k \in [\tilde{\tau}_{i,m}, \tau_{m+1}). \quad (19)$$

Lemma 4.1: The sequences $\{\tilde{x}_{i,k}\}, \{\tilde{\varepsilon}_{i,k+1}\}$ defined by (18) (19) satisfy the following recursions

$$\hat{x}_{i,k+1} \triangleq \sum_{j \in N_i(k)} \omega_{ij}(k) \tilde{x}_{j,k} + \gamma_k (f_i(\tilde{x}_{i,k}) + \tilde{\varepsilon}_{i,k+1}), \quad (20)$$

$$\begin{aligned} \tilde{x}_{i,k+1} &= \hat{x}_{i,k+1} I_{[\|\hat{x}_{j,k+1}\| \leq M_{\sigma_k} \quad \forall j \in \mathcal{V}]} \\ &+ x^* I_{[\exists j \in \mathcal{V} \quad \|\hat{x}_{j,k+1}\| > M_{\sigma_k}]}, \end{aligned} \quad (21)$$

$$\sigma_{k+1} = \sigma_k + I_{[\exists j \in \mathcal{V} \quad \|\hat{x}_{j,k+1}\| > M_{\sigma_k}]}, \quad \sigma_0 = 0. \quad (22)$$

The proof is given in Appendix A.

Before clarifying the property of the noise sequence $\{\tilde{\varepsilon}_{i,k+1}\}_{k \geq 0}$, we need the following lemma.

Lemma 4.2: Assume A4 holds. Then

$$\text{i) } \sigma_{j,k+Bd_{i,j}} \geq \sigma_{i,k} \quad \forall j \in \mathcal{V} \quad \forall k \geq 0, \quad (23)$$

where $d_{i,j}$ is the length of the shortest directed path from i to j in \mathcal{G}_∞ , and B is the positive integer given in A4 d),

$$\text{ii) } \tilde{\tau}_{j,m} \leq \tau_m + BD \quad \forall j \in \mathcal{V} \quad \text{for } m \geq 1, \quad (24)$$

where $D \triangleq \max_{i,j \in \mathcal{V}} d_{i,j}$.

Proof: i) Since \mathcal{G}_∞ is strongly connected by A4 c), for any $j \in \mathcal{V}$ there exists a sequence of nodes $i_1, i_2, \dots, i_{d_{i,j}-1}$ such that $(i, i_1) \in \mathcal{E}_\infty, (i_1, i_2) \in \mathcal{E}_\infty, \dots, (i_{d_{i,j}-1}, j) \in \mathcal{E}_\infty$.

Noticing that $(i, i_1) \in \mathcal{E}_\infty$, by A4 d) we have

$$(i, i_1) \in \mathcal{E}(k) \cup \mathcal{E}(k+1) \cup \dots \cup \mathcal{E}(k+B-1).$$

Therefore, there exists a positive integer $k' \in [k, k+B-1]$ such that $(i, i_1) \in \mathcal{E}(k')$. So, $i \in N_{i_1}(k')$, and hence by (6) and (9) we derive

$$\sigma_{i_1, k+B} \geq \sigma_{i_1, k'+1} \geq \hat{\sigma}_{i_1, k'} \geq \sigma_{i, k'} \geq \sigma_{i, k}.$$

Similarly, we have $\sigma_{i_2, k+2B} \geq \sigma_{i_1, k+B} \geq \sigma_{i, k}$. Continuing this procedure, we finally reach the inequality (23).

ii) Let $\tau_m = k_1$ for some $m \geq 1$. Then there is an i such that $\tau_{i,m} = k_1$. By (23) we have $\sigma_{j, k_1+Bd_{i,j}} \geq \sigma_{i, k_1} = m \quad \forall j \in \mathcal{V}$.

For the case where $\sigma_{j, k_1+Bd_{i,j}} = m \quad \forall j \in \mathcal{V}$, we have $\tau_{j,m} \leq k_1 + Bd_{i,j} \quad \forall j \in \mathcal{V}$. By noticing $\tau_m = k_1$, from here by the definition of $\tilde{\tau}_{j,m}$ we obtain (24):

$$\tilde{\tau}_{j,m} \leq \tau_{j,m} \leq \tau_m + Bd_{i,j} \leq \tau_m + BD \quad \forall j \in \mathcal{V}.$$

For the case where $\sigma_{j, k_1+Bd_{i,j}} > m$ for some $j \in \mathcal{V}$, we have $\tau_{m+1} \leq k_1 + Bd_{i,j}$ for some $j \in \mathcal{V}$, and hence $\tau_{m+1} \leq \tau_m + BD$. Again, by noticing $\tau_m = k_1$ we obtain (24):

$$\tilde{\tau}_{j,m} \leq \tau_{m+1} \leq \tau_m + BD \quad \forall j \in \mathcal{V}.$$

Proposition 4.3: If $\sigma_k \xrightarrow[k \rightarrow \infty]{} \infty$, then for any $i \in \mathcal{V}$, $\lim_{k \rightarrow \infty} \sigma_{i,k} = \infty$. This is because there exists an $i_0 \in \mathcal{V}$ such that $\sigma_{i_0,k} \xrightarrow[k \rightarrow \infty]{} \infty$. Then from (23) it follows that for any $j \in \mathcal{V}$, $\sigma_{j,k} \xrightarrow[k \rightarrow \infty]{} \infty$.

Proposition 4.4: If $\{\sigma_k\}$ is bounded, then $\{\tilde{x}_{i,k}\}$ and $\{x_{i,k}\}$, $\{\tilde{\varepsilon}_{i,k}\}$ and $\{\varepsilon_{i,k}\}$ coincide in a finite number of steps.

Proof: Since σ_k is defined as the largest truncation number among all agents at time k , from $\lim_{k \rightarrow \infty} \sigma_k = \sigma$ it follows that for any $i \in \mathcal{V}$ and $k \geq 0$, $\sigma_{i,k} \leq \sigma$. Then by the definition of $\tau_{i,m}$ we have that for any $i \in \mathcal{V}$, $\tau_{i,\sigma+1} = \inf\{k : \sigma_{i,k} = \sigma + 1\} = \infty$, and hence

$$\tau_{\sigma+1} = \infty \text{ when } \lim_{k \rightarrow \infty} \sigma_k = \sigma. \quad (25)$$

Then the result is derived by (19). \blacksquare

Lemma 4.5: Assume A5 b) holds at the sample path ω under consideration for all agents. Then for this ω

$$\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, t_k) \wedge (\tau_{\sigma_{n_k}} + 1 - 1)} \gamma_s \tilde{\varepsilon}_{s+1} I_{\|\tilde{X}_s\| \leq K} \right\| = 0 \quad \forall t_k \in [0, T] \text{ for sufficiently large } K > 0 \quad (26)$$

along indices $\{n_k\}$ whenever $\{\tilde{X}_{n_k}\}$ converges at ω .

Proof: The proof is shown in Appendix B. \blacksquare

B. Local Properties Along Convergent Subsequences

Set $\Psi(k, s) \triangleq [D_{\perp}(W(k) \otimes \mathbf{I}_l)][D_{\perp}(W(k-1) \otimes \mathbf{I}_l)] \dots [D_{\perp}(W(s) \otimes \mathbf{I}_l)] \quad \forall k \geq s$, $\Psi(k-1, k) \triangleq \mathbf{I}_{Nl}$.

Beacuse for any $k \geq 1$, the matrix $W(k)$ is doubly stochastic, then by using the rule of Kronecker product

$$(A \otimes B)(C \otimes D) = (AC) \otimes (BD), \quad (27)$$

we conclude that for any $k \geq s-1$,

$$\Psi(k, s) = \left(\Phi(k, s) - \frac{1}{N} \mathbf{1}\mathbf{1}^T \right) \otimes \mathbf{I}_l, \quad (28)$$

$$\Psi(k, s) D_{\perp} = \left(\Phi(k, s) - \frac{1}{N} \mathbf{1}\mathbf{1}^T \right) \otimes \mathbf{I}_l. \quad (29)$$

The following lemma measures the closeness of the auxiliary sequence $\{\tilde{X}_k\}$ along its convergent subsequence $\{\tilde{X}_{n_k}\}$.

Lemma 4.6: Assume A1, A3, A4 hold and that A5 b) holds at the sample path ω under consideration. Let $\{\tilde{X}_{n_k}\}$ be a convergent subsequence of $\{\tilde{X}_k\} : \tilde{X}_{n_k} \xrightarrow[k \rightarrow \infty]{} \bar{X}$ at the considered ω . Then for this ω , there is a $T > 0$ such that for sufficiently large k and any $T_k \in [0, T]$,

$$\tilde{X}_{m+1} = (W(m) \otimes \mathbf{I}_l) \tilde{X}_m + \gamma_m \left(F(\tilde{X}_m) + \tilde{\varepsilon}_{m+1} \right) \quad (30)$$

for any $m = n_k, \dots, m(n_k, T_k)$, and

$$\|\tilde{X}_{m+1} - \tilde{X}_{n_k}\| \leq c_1 T_k + M'_0, \quad (31)$$

$$\|\tilde{x}_{m+1} - \tilde{x}_{n_k}\| \leq c_2 T_k \quad \forall m : n_k \leq m \leq m(n_k, T_k), \quad (32)$$

where $\tilde{x}_k \triangleq \frac{1}{N} (\mathbf{1}^T \otimes \mathbf{I}_l) \tilde{X}_k = \frac{1}{N} \sum_{i=1}^N \tilde{x}_{i,k}$ and c_0, c_1, M'_0 are positive constants which may depend on ω .

The proof is given in Appendix C.

Multiplying both sides of (30) with $\frac{1}{N} (\mathbf{1}^T \otimes \mathbf{I}_l)$ from left, by $\mathbf{1}^T W(m) = \mathbf{1}^T$ and (27) it follows that

$$\begin{aligned} \bar{x}_{m+1} &= \bar{x}_m + \gamma_m f(\bar{x}_m) + (\mathbf{1}^T \otimes \mathbf{I}_l) \gamma_m \tilde{\varepsilon}_{m+1} / N \\ &\quad + \gamma_m \sum_{i=1}^N (f_i(\tilde{x}_{i,m}) - f_i(\bar{x}_m)) / N. \end{aligned} \quad (33)$$

By setting $e_{i,m+1} \triangleq (f_i(\tilde{x}_{i,m}) - f_i(\bar{x}_m)) / N$, $e_{m+1} \triangleq \sum_{i=1}^N e_{i,m+1}$, and $\zeta_{m+1} \triangleq (\mathbf{1}^T \otimes \mathbf{I}_l) \tilde{\varepsilon}_{m+1} / N + e_{m+1}$, we can rewrite (33) in the centralized form as follows:

$$\bar{x}_{m+1} = \bar{x}_m + \gamma_m (f(\bar{x}_m) + \zeta_{m+1}). \quad (34)$$

The following lemma gives the property of the noise sequence $\{\zeta_{k+1}\}$. For the proof we refer to Appendix D.

Lemma 4.7: Assume that all conditions used in Lemma 4.6 are satisfied. Let $\{\tilde{X}_{n_k}\}$ be a convergent subsequence with limit \bar{X} at the considered ω . Then for this ω

$$\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s \zeta_{s+1} \right\| = 0 \quad \forall T_k \in [0, T]. \quad (35)$$

The following lemma gives the crossing behavior of $v(\cdot)$ at the trajectory \bar{x}_k with respect to a non-empty interval that has no intersection with $v(J)$.

Lemma 4.8: Assume A1-A4 hold and that A5 b) holds for the sample path ω under consideration. Then any nonempty interval $[\delta_1, \delta_2]$ with $d([\delta_1, \delta_2], v(J)) > 0$ cannot be crossed by $\{v(\bar{x}_{n_k}), \dots, v(\bar{x}_{m_k})\}$ infinitely many times with $\{\|\tilde{X}_{n_k}\|\}$ bounded, where by ‘‘crossing $[\delta_1, \delta_2]$ by $\{v(\bar{x}_{n_k}), \dots, v(\bar{x}_{m_k})\}$ ’’ it is meant that $v(\bar{x}_{n_k}) \leq \delta_1$, $v(\bar{x}_{m_k}) \geq \delta_2$, and $\delta_1 < v(\bar{x}_s) < \delta_2 \quad \forall s : n_k < s < m_k$.

Proof: Assume the converse: for some nonempty interval $[\delta_1, \delta_2]$ with $d([\delta_1, \delta_2], v(J)) > 0$, there exist infinitely many crossings $\{v(\bar{x}_{n_k}), \dots, v(\bar{x}_{m_k})\}$ with $\{\|\tilde{X}_{n_k}\|\}$ bounded.

By the boundedness of $\{\|\tilde{X}_{n_k}\|\}$, we can extract a convergent subsequence still denoted by $\{\tilde{X}_{n_k}\}$ with $\lim_{k \rightarrow \infty} \tilde{X}_{n_k} = \bar{X}$.

So, $\lim_{k \rightarrow \infty} \bar{x}_{n_k} = \bar{x} = \frac{\mathbf{1}^T \otimes \mathbf{I}_l}{N} \bar{X}$. Setting $T_k = \gamma_{n_k}$ in (32), we derive $\|\bar{x}_{n_k+1} - \bar{x}_{n_k}\| \leq c_2 \gamma_{n_k} \xrightarrow[k \rightarrow \infty]{} 0$. By the definition of crossings, we obtain $v(\bar{x}_{n_k}) \leq \delta_1 < v(\bar{x}_{n_k+1})$, and hence

$$v(\bar{x}_{n_k}) \xrightarrow[k \rightarrow \infty]{} \delta_1 = v(\bar{x}), \quad d(\bar{x}, J) \triangleq \vartheta > 0. \quad (36)$$

Then by (32) it is seen that

$$d(\bar{x}_s, J) > \frac{\vartheta}{2} \quad \forall s : n_k \leq s \leq m(n_k, t) + 1 \quad (37)$$

for sufficiently small $t > 0$ and large k . From (34) we obtain

$$\begin{aligned} v(\bar{x}_{m(n_k, t)+1}) &= v(\bar{x}_{n_k} + \sum_{s=n_k}^{m(n_k, t)} \gamma_s (f(\bar{x}_s) + \zeta_{s+1})) \\ &= v(\bar{x}_{n_k}) + v_x(\xi_k)^T \sum_{s=n_k}^{m(n_k, t)} \gamma_s (f(\bar{x}_s) + \zeta_{s+1}), \end{aligned}$$

where ξ_k is in-between \bar{x}_{n_k} and $\bar{x}_{m(n_k, t)+1}$. Then

$$\begin{aligned} v(\bar{x}_{m(n_k, t)+1}) - v(\bar{x}_{n_k}) &= \sum_{s=n_k}^{m(n_k, t)} \gamma_s v_x(\bar{x}_s)^T f(\bar{x}_s) \\ &\quad + \sum_{s=n_k}^{m(n_k, t)} \gamma_s (v_x(\xi_k) - v_x(\bar{x}_s))^T f(\bar{x}_s) \\ &\quad + v_x(\xi_k)^T \sum_{s=n_k}^{m(n_k, t)} \gamma_s \zeta_{s+1}. \end{aligned} \quad (38)$$

By (12) and (37), there exists a constant $\alpha_1 > 0$ such that

$$v_x(\bar{x}_s)^T f(\bar{x}_s) \leq -\alpha_1 \quad \forall s : n_k \leq s \leq m(n_k, t)$$

for sufficiently small $t > 0$ and large k , and hence

$$\sum_{s=n_k}^{m(n_k,t)} \gamma_s v_x(\bar{x}_s)^T f(\bar{x}_s) \leq -\alpha_1 t. \quad (39)$$

Since $\{\bar{x}_s : n_k \leq s \leq m(n_k, t)\}$ are bounded, by continuity of $f(\cdot)$ there exists a constant $c_6 > 0$ such that

$$\sum_{s=n_k}^{m(n_k,t)} \gamma_s \|f(\bar{x}_s)\| \leq c_6 t. \quad (40)$$

By invoking that ξ_k is in-between \bar{x}_{n_k} and $\bar{x}_{m(n_k,t)+1}$, by continuity of $v_x(\cdot)$ and (32), we know that

$$v_x(\xi_k) - v_x(\bar{x}_s) = \delta(t) \quad \forall s : n_k \leq s \leq m(n_k, t), \quad (41)$$

where $\delta(t) \rightarrow 0$ as $t \rightarrow 0$. Then by (40) we derive

$$\begin{aligned} & \sum_{s=n_k}^{m(n_k,t)} \gamma_s (v_x(\xi_k) - v_x(\bar{x}_s))^T f(\bar{x}_s) \\ & \leq \delta(t) \sum_{s=n_k}^{m(n_k,t)} \gamma_s \|f(\bar{x}_s)\| \leq \delta(t)t. \end{aligned} \quad (42)$$

Since $\bar{x}_{n_k} \xrightarrow{k \rightarrow \infty} \bar{x}$, by continuity of $v_x(\cdot)$ and (32) it follows that for sufficiently small $t > 0$ and large k

$$v_x(\bar{x}_s) - v_x(\bar{x}) = o(1) + \delta(t) \quad \forall s : n_k \leq s \leq m(n_k, t),$$

where $o(1) \rightarrow 0$ as $k \rightarrow \infty$. Then by (41) we derive

$$v_x(\xi_k) - v_x(\bar{x}) = o(1) + \delta(t) \quad \forall s : n_k \leq s \leq m(n_k, t).$$

Consequently, for sufficiently small $t > 0$ and large k

$$\begin{aligned} & v_x(\xi_k)^T \sum_{s=n_k}^{m(n_k,t)} \gamma_s \zeta_{s+1} \\ & = ((v_x(\xi_k) - v_x(\bar{x})) + v_x(\bar{x}))^T \sum_{s=n_k}^{m(n_k,t)} \gamma_s \zeta_{s+1} \\ & \leq (o(1) + \delta(t) + \|v_x(\bar{x})\|) \sum_{s=n_k}^{m(n_k,t)} \gamma_s \zeta_{s+1}. \end{aligned} \quad (43)$$

Substituting (39), (42), and (43) into (38), we obtain

$$\begin{aligned} & v(\bar{x}_{m(n_k,t)+1}) - v(\bar{x}_{n_k}) \leq -\alpha_1 t + \delta(t)t \\ & + (o(1) + \delta(t) + \|v_x(\bar{x})\|) \sum_{s=n_k}^{m(n_k,t)} \gamma_s \zeta_{s+1}. \end{aligned}$$

Then by (36) it follows that

$$\begin{aligned} & \limsup_{k \rightarrow \infty} v(\bar{x}_{m(n_k,t)+1}) \leq \delta_1 - \alpha_1 t + \delta(t)t \\ & + (\delta(t) + \|v_x(\bar{x})\|) \limsup_{k \rightarrow \infty} \sum_{s=n_k}^{m(n_k,t)} \gamma_s \zeta_{s+1}. \end{aligned}$$

Hence from (35), for sufficiently small t we have that

$$\limsup_{k \rightarrow \infty} v(\bar{x}_{m(n_k,t)+1}) \leq \delta_1 - \frac{\alpha_1}{2} t. \quad (44)$$

However, by continuity of $v_x(\cdot)$ and (32) we know that

$$\lim_{t \rightarrow 0} \max_{n_k \leq m \leq m(n_k,t)} \|v(\bar{x}_{m+1}) - v(\bar{x}_{n_k})\| = 0,$$

which implies that $m(n_k, t) + 1 < m_k$ for sufficiently small t . Therefore, $v(\bar{x}_{m(n_k,t)+1}) \in (\delta_1, \delta_2)$, which contradicts with (44). Consequently, the converse assumption is not true. The proof is completed. \blacksquare

C. Finiteness of Number of Truncations

Lemma 4.9: Let $\{x_{i,k}\}$ be produced by (6)-(9) with an arbitrary initial value $x_{i,0}$. Suppose A1, A3, A4, and assume A5 b) holds for the sample path ω under consideration.

i) If $\lim_{k \rightarrow \infty} \sigma_k = \infty$, then there exists an integer sequence $\{n_k\}$ such that $\bar{x}_{n_k} = x^*$. Further, $\{\bar{x}_{n_k}\}$ starting from x^* crosses the sphere with $\|x\| = c_0$ infinitely many times, where $\{\bar{x}_k\}$ is defined in Lemma 4.6 and c_0 is given in A2 c).

ii) If, in addition, A2 also holds, then there exists a positive integer σ possibly depending on ω such that

$$\lim_{k \rightarrow \infty} \sigma_k = \sigma. \quad (45)$$

The proof of the lemma is given in Appendix E. Lemma 4.9 says that the largest truncation number among all agents converges, while the following lemma indicates that the truncation numbers at all agents converge to the same limit.

Lemma 4.10: Assume all conditions required by Lemma 4.9 are satisfied. Then there exists a positive integer σ such that

$$\lim_{k \rightarrow \infty} \sigma_{i,k} = \sigma \quad \forall i \in \mathcal{V}. \quad (46)$$

Proof: Since all conditions required by Lemma 4.9 hold, (45) holds for some positive integer σ . Thus,

$$\sigma_{i,k} \leq \sigma \quad \forall k \geq 0 \quad \forall i \in \mathcal{V}. \quad (47)$$

By (45) and (B.2), we have $\tau_{\sigma+1} = \infty$, and hence $\tilde{\tau}_{i,\sigma} = \tau_{i,\sigma} \leq BD + \tau_\sigma \quad \forall i \in \mathcal{V}$ by (24). This means that the smallest time when the truncation number of agent i reaches σ is not larger than $BD + \tau_\sigma$. So, the truncation number of agent i after time $BD + \tau_\sigma$ is not smaller than σ , i.e., $\sigma_{i,k} \geq \sigma \quad \forall k \geq BD + \tau_\sigma \quad \forall i \in \mathcal{V}$. It incorporating with (47) yields

$$\sigma_{i,k} = \sigma \quad \forall k \geq BD + \tau_\sigma \quad \forall i \in \mathcal{V}. \quad (48)$$

Consequently, (46) holds. \blacksquare

D. Proof of Theorem 3.4

Proof. i) By (45) and (48), there is a positive integer σ possibly depending on ω such that

$$\hat{\sigma}_{i,k} = \sigma_{i,k} = \sigma \quad \forall k \geq k_0 \triangleq BD + \tau_\sigma \quad \forall i \in \mathcal{V}. \quad (49)$$

Then by (7), we have the following for all $k \geq k_0$:

$$x'_{i,k+1} = \sum_{j \in N_i(k)} \omega_{ij}(k) x_{j,k} + \gamma_k O_{i,k+1} \quad \forall i \in \mathcal{V}. \quad (50)$$

By (48) and (49), we obtain $\sigma_{i,k+1} = \hat{\sigma}_{i,k} = \sigma \quad \forall k \geq k_0 \quad \forall i \in \mathcal{V}$. Thus, for any $k \geq k_0$ and any $i \in \mathcal{V}$, we have that $\|x'_{i,k+1}\| \leq M_\sigma$ by (9), and $x_{i,k+1} = x'_{i,k+1}$ by (8). Then for any $i \in \mathcal{V}$, $\{x_{i,k}\}$ is bounded and (14) follows from (50).

ii) By multiplying both sides of (15) with D_\perp from left, we have that

$$X_{\perp,k+1} = D_\perp(W(k) \otimes I_l)X_k + \gamma_k D_\perp(F(X_k) + \varepsilon_{k+1}),$$

and inductively

$$X_{\perp,k+1} = \Psi(k, k_0)X_{k_0} + \sum_{m=k_0}^k \gamma_m \Psi(k-1, m)D_\perp(F(X_m) + \varepsilon_{m+1}) \quad \forall k \geq k_0.$$

Then by (28) and (29), we derive

$$\begin{aligned} X_{\perp, k+1} &= [(\Phi(k, k_0) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l] X_{k_0} \\ &+ \sum_{m=k_0}^k \gamma_m [(\Phi(k-1, m) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l] F(X_m) \\ &+ \sum_{m=k_0}^k \gamma_m [(\Phi(k-1, m) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l] \varepsilon_{m+1}. \end{aligned}$$

Therefore, from (13) by continuity of $F(\cdot)$ and the boundedness of $\{X_s\}$, we conclude that there exist positive constants c_1, c_2, c_3 possibly depending on ω such that

$$\begin{aligned} \|X_{\perp, k+1}\| &\leq c_1 \rho^{k+1-k_0} + c_2 \sum_{m=k_0}^k \gamma_m \rho^{k-m} \\ &+ c_3 \sum_{m=k_0}^k \gamma_m \rho^{k-m} \|\varepsilon_{m+1}\| \quad \forall k \geq k_0. \end{aligned} \quad (51)$$

Note that for any given $\epsilon > 0$, there exists a positive integer k_1 such that $\gamma_k \leq \epsilon \quad \forall k \geq k_1$. We then have

$$\begin{aligned} \sum_{m=0}^k \gamma_m \rho^{k-m} &= \sum_{m=0}^{k_1} \gamma_m \rho^{k-m} + \sum_{m=k_1+1}^k \gamma_m \rho^{k-m} \\ &\leq \rho^{k-k_1} \sum_{m=0}^{k_1} \gamma_m + \epsilon \frac{1}{1-\rho} \xrightarrow[k \rightarrow \infty]{\epsilon \rightarrow 0} 0. \end{aligned}$$

Therefore, the second term at the right-hand side of (51) tends to zero as $k \rightarrow \infty$. Similarly, the last term at the right-hand side of (51) also tends to zero since $\lim_{k \rightarrow \infty} \gamma_k \varepsilon_{k+1} = 0$. Therefore, by $0 < \rho < 1$, from (51) we conclude that

$$X_{\perp, k} \xrightarrow[k \rightarrow \infty]{} \mathbf{0}.$$

By i) and Proposition 4.4, we see that $\{\tilde{x}_{i,k} \quad \forall i \in \mathcal{V}\}$ are bounded for this ω , and hence $\{\bar{x}_k\}$ is bounded. The rest of the proof is similar to that given in [36].

We first show the convergence of $v(\bar{x}_k)$. Since

$$v_1 \triangleq \liminf_{k \rightarrow \infty} v(\bar{x}_k) \leq \limsup_{k \rightarrow \infty} v(\bar{x}_k) \triangleq v_2,$$

we want to prove $v_1 = v_2$. Assume the converse: $v_1 < v_2$. Since $v(J)$ is nowhere dense, there exists a nonnegative interval $[\delta_1, \delta_2] \in (v_1, v_2)$ such that $d([\delta_1, \delta_2], v(J)) > 0$. Then $v(\bar{x}_k)$ crosses the interval $[\delta_1, \delta_2]$ infinitely many times. This contradicts Lemma 4.8. Therefore, $v_1 = v_2$, which implies the convergence of $v(\bar{x}_k)$.

We then prove $d(\bar{x}_k, J) \xrightarrow[k \rightarrow \infty]{} 0$. Assume the converse. Then by the boundedness of $\{\bar{x}_k\}$, there exists a convergent subsequence $\bar{x}_{n_k} \xrightarrow[k \rightarrow \infty]{} \bar{x}$ with $d(\bar{x}, J) \triangleq \vartheta > 0$. From (32) it follows that for sufficiently small $t > 0$ and large k

$$d(\bar{x}_s, J) > \frac{\vartheta}{2} \quad \forall s : n_k \leq s \leq m(n_k, t),$$

and hence by (12) there exists a constant $b > 0$ such that

$$v_x(\bar{x}_s)^T f(\bar{x}_s) < -b \quad \forall s : n_k \leq s \leq m(n_k, t).$$

Thus, similar to the proof for obtaining (44) it is seen that for sufficiently small $t > 0$

$$\limsup_{k \rightarrow \infty} v(\bar{x}_{m(n_k, t)+1}) \leq v(\bar{x}) - \frac{b}{2} t.$$

This contradicts with the convergence of $v(\bar{x}_k)$. Therefore, $d(\bar{x}_k, J) \xrightarrow[k \rightarrow \infty]{} 0$, and hence $d(x_k, J) \xrightarrow[k \rightarrow \infty]{} 0$.

iii) Assume the converse: i.e., J^* is disconnected. Then there exist closed sets J_1^* and J_2^* such that $J^* = J_1^* \cup J_2^*$ and $d(J_1^*, J_2^*) > 0$. Define $\rho = \frac{1}{3} d(J_1^*, J_2^*)$. By $d(\bar{x}_k, J^*) \xrightarrow[k \rightarrow \infty]{} 0$ there exists k_0 such that

$$\bar{x}_k \in B(J_1^*, \rho) \cup B(J_2^*, \rho) \quad \forall k \geq k_0, \quad (52)$$

where $B(A, \rho)$ denotes the ρ -neighborhood of the set A .

Define

$$\begin{aligned} n_0 &= \inf\{k > k_0, d(\bar{x}_k, J_1^*) < \rho\}, \\ m_p &= \inf\{k > n_p, d(\bar{x}_k, J_2^*) < \rho\}, \\ n_p &= \inf\{k > m_p, d(\bar{x}_k, J_1^*) < \rho\}, \quad p \geq 0. \end{aligned}$$

Then by (52), we have that $\bar{x}_{n_p} \in B(J_1^*, \rho)$ and $\bar{x}_{n_{p+1}} \in B(J_2^*, \rho)$ for any $p \geq 0$. Then by $d(J_1^*, J_2^*) = 3\rho$, we derive

$$\|\bar{x}_{n_p} - \bar{x}_{n_{p+1}}\| > \rho. \quad (53)$$

Since $\{\bar{x}_{n_k}\}$ is bounded, we can extract a convergent subsequence, denoted still by $\{\bar{x}_{n_k}\}$. By setting $T_k = \gamma_{n_k}$ in (32) we derive $\|\bar{x}_{n_{k+1}} - \bar{x}_{n_k}\| \leq c_2 \gamma_{n_k} \xrightarrow[k \rightarrow \infty]{} 0$, which contradicts with (53). So, the converse assumption is not true. The proof is completed. ■

V. CONVERGENCE FOR APPLICATION PROBLEMS

In this section, we establish convergence results for the two problems stated in Section II, and present the corresponding numerical simulations.

A. Distributed PCA

We now apply DSAAWET to the problem of distributed principal component analysis, and establish its convergence.

Corollary 5.1: Let $\{x_{i,k}\}$ be produced by (6)–(9), where $O_{i,k+1}$ is defined by (3), $x_{i,0} = x^* = \mathbf{1}/\sqrt{N}$, and $\gamma_k = \frac{1}{k}$. Suppose A4 holds, and, in addition, that

B1) for any $i \in \mathcal{V}$, $\{A_{i,k}\}$ is almost surely bounded;

B2) A is positive definite, and the largest eigenvalue of A has unit multiplicity with the corresponding unit eigenvectors denoted by $u^{(1)}$;

B3) for any $i \in \mathcal{V}$, $\liminf_{k \rightarrow \infty} \mathbb{E}[\|y_k^T N_{i,k} z_k\| | \mathcal{F}_k] > 0$ for any \mathcal{F}_k -measurable y_k and z_k bounded from above and from zero, where $N_{i,k} \triangleq A_{i,k} - A_i$ and $\mathcal{F}_k \triangleq \sigma\{x_{i,p}, i \in \mathcal{V}, 0 \leq p \leq k\}$. Then (15) takes place and $X_{\perp, k} \xrightarrow[k \rightarrow \infty]{} \mathbf{0}$ a.s. Assume further that with probability one X_k does not converge to $\mathbf{0}$. Then for any $i, j \in \mathcal{V}$ and for almost all $\omega \in \Omega$, the following holds:

$$\lim_{k \rightarrow \infty} x_{i,k}(\omega) = \lim_{k \rightarrow \infty} x_{j,k}(\omega) = u^{(1)}. \quad (54)$$

Proof: We prove this result by adopting the similar procedures as that used in the proof of Theorem 3.4. Firstly, we use the auxiliary sequences defined in (18)(19) to prove the finiteness of truncation numbers. Then we show that the estimates are bounded and asymptotically reach consensus. Finally, we show that the estimates converge to the unit eigenvector corresponding to the largest eigenvalue.

It is clear that the step size $\gamma_k = \frac{1}{k}$ satisfies A1. By $f_i(x) = A_i x - (x^T A_i x)x$, we see A3 holds. Then by (2)(3) we have

$$\begin{aligned} \varepsilon_{i,k+1} &= O_{i,k+1} - f_i(x_{i,k}) \\ &= (A_{i,k} - A_i)x_{i,k} - (x_{i,k}^T (A_{i,k} - A_i)x_{i,k})x_{i,k}. \end{aligned}$$

Then from B1 we conclude that $\{\varepsilon_{i,k+1} I_{[\|x_{i,k}\| \leq K]}\}$ is an MDS with bounded second moments for any $K > 0$. Then by the convergence theorem for MDS [40] we have that

$$\sum_{k=0}^{\infty} \gamma_k \varepsilon_{i,k} I_{[\|x_{i,k}\| \leq K]} < \infty \text{ a.s.} \quad (55)$$

So, A5 b) holds for almost all sample path ω for any agent i .

Denote by J^0 the set of nonzero roots of $f(x) = Ax - (x^T Ax)x$ in \mathbb{R}^l . Take some $u_0 \in J^0$. Then we get $Au_0 = (u_0^T Au_0)u_0$, and hence by multiplying both sides with u_0^T from the left we derive $u_0^T Au_0 = (u_0^T Au_0)u_0^T u_0$. Since $u_0^T Au_0 > 0$ by the positive definiteness of A , we have that $u_0^T u_0 = 1$. Thus, J^0 is on the unit sphere. By noticing that the nonzero roots of $f(\cdot)$ on the unit sphere are the unit eigenvectors of A , and hence J^0 coincides with the set of all unit eigenvectors of A . Following [38], we define the Lyapunov function as follows:

$$v(x) = \frac{e^{\|x\|^2}}{x^T Ax} \quad \forall x \neq \mathbf{0}.$$

Then we have

$$v_x(x) = 2xv(x) - \frac{e^{\|x\|^2} 2Ax}{(x^T Ax)^2} = \frac{2v(x)}{x^T Ax} ((x^T Ax)x - Ax),$$

and hence

$$f^T(x)v_x(x) = -\frac{2v(x)}{x^T Ax} \|f(x)\|^2 < 0 \quad \forall x \notin J^0 \cup \{\mathbf{0}\}. \quad (56)$$

We now prove that $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$. Assume the converse: $\lim_{k \rightarrow \infty} \sigma_k = \infty$.

By Lemma 4.9 i), $\{\tilde{X}_k\}$ has a convergent subsequence $\{\tilde{X}_{n_k}\}$ with $\tilde{X}_{n_k} = (\mathbf{1} \otimes \mathbf{I}_m)x^*$. Since A1, A3, A4 and A5 b) hold, Lemmas 4.6 and 4.7 take place. By noticing $\bar{x}_{n_k} = \mathbf{1}/\sqrt{N}$, from (32) we see that for sufficiently small T and large k , all components of $\bar{x}_{m+1} \quad \forall m : n_k \leq m \leq m(n_k, T)$ are uniformly above zero. This together with (56) yields (39) when we follow the proof procedure for Lemma 4.8 with $v(J)$ replaced by $v(J^0)$. It is worth noting that $v(J)$ is not defined since $v(x)$ is not defined at $x = \mathbf{0}$, while $v(J^0)$ is well defined. Then, any nonempty interval $[\delta_1, \delta_2]$ with $d([\delta_1, \delta_2], v(J^0)) > 0$ cannot be crossed by $\{v(\bar{x}_{n_k}), \dots, v(\bar{x}_{m_k})\}$ infinitely many times.

Since $v(x)$ is radially unbounded, there exists $c_0 > 1$ such that $v(x^*) < \inf_{\|x\|=c_0} v(x)$. By Lemma 4.9 i), $\{\bar{x}_{n_k}\}$ starting from x^* crosses the sphere with $\|x\| = c_0$ infinitely many times. Since $v(J^0)$ is nowhere dense, there exists a nonempty interval $[\delta_1, \delta_2] \in (v(x^*), \inf_{\|x\|=c_0} v(x))$ with $d([\delta_1, \delta_2], v(J^0)) > 0$. Therefore, $[\delta_1, \delta_2]$ with $d([\delta_1, \delta_2], v(J^0)) > 0$ is crossed by $\{v(\bar{x}_{n_k}), \dots, v(\bar{x}_{m_k})\}$ infinitely many times. This yields a contradiction, hence the converse assumption is not true. Thus,

$$\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty. \quad (57)$$

Since (57) holds, the proof for Theorem 3.4 i) is still applicable. As a result, $\{X_k\}$ is a.s. bounded and there exists a positive integer k_0 such that the compact form (15) holds. Then by (55) we obtain that $\sum_{k=0}^{\infty} \gamma_k \varepsilon_{i,k} < \infty$ a.s., and hence A5 holds for almost all ω . Since the proof for the first result in Theorem 3.4 ii) still holds, we obtain $X_{\perp,k} \xrightarrow[k \rightarrow \infty]{} \mathbf{0}$ a.s. We further show the rate of convergence of $X_{\perp,k}$. By the boundedness of $\{X_k\}$ and by B1, we see that $\{\|\varepsilon_k\|\}_{k \geq k_0}$ is bounded a.s. Then similar to the proof of (51), for almost all

ω , there exist positive constants c_1, c_2 possibly depending on ω such that for any $k \geq k_0$:

$$\|X_{\perp,k+1}\| \leq c_1 \rho^{k+1-k_0} + c_2 \sum_{m=k_0}^k \gamma_m \rho^{k-m}. \quad (58)$$

Note that

$$\begin{aligned} \frac{1}{\gamma_k} \sum_{m=k_0}^k \gamma_m \rho^{k-m} &= \sum_{m=k_0}^k \frac{k}{m} \rho^{k-m} \\ &= \sum_{m=k_0}^k \left(1 + \frac{k-m}{m}\right) \rho^{k-m} \\ &\leq \sum_{m=k_0}^k \rho^{k-m} + \sum_{m=k_0}^k (k-m+1) \rho^{k-m} \\ &\leq \frac{1}{1-\rho} + \frac{d}{d\rho} \sum_{m=k_0}^k \rho^{k-m+1} \leq \frac{1}{1-\rho} + \frac{1}{(1-\rho)^2}, \end{aligned}$$

which incorporating with (58) yields that for almost all ω :

$$\limsup_{k \rightarrow \infty} \gamma_k^{-1} X_{\perp,k} \leq \frac{c_2(\omega)}{1-\rho} + \frac{c_2(\omega)}{(1-\rho)^2}. \quad (59)$$

By recalling that with probability one X_k does not converge to zero, for sufficiently large $k_1 > k_0$ we have that

$$X_k \neq \mathbf{0} \quad \forall k \geq k_1 \quad a.s. \quad (60)$$

This is because the converse assumption $X_{k_2} = \mathbf{0}$ for some $k_2 > k_1$ implies that $X_k = \mathbf{0} \quad \forall k \geq k_2$ by (3) and (14). This contradicts with the assumption that X_k does not converge to zero. So, (60) holds. Hence by $X_{\perp,k} \xrightarrow[k \rightarrow \infty]{} \mathbf{0}$ it follows that x_k does not converge to zero. Since A1, A3, A4, A5 and (56) hold, similar to the proof of Theorem 3.4 iii) we have that for almost all $\omega \in \Omega$:

$$\lim_{k \rightarrow \infty} d(x_k(\omega), J^*) = 0, \quad (61)$$

where J^* is a connected subset J^0 .

Similar to the procedure for deriving (34), we can rewrite (15) in the following centralized form:

$$x_{k+1} = x_k + \frac{\gamma_k}{N} (f(x_k) + \zeta_{k+1} + \beta_{k+1}), \quad (62)$$

where $\beta_{k+1} \triangleq \sum_{i=1}^N (f_i(x_{i,k}) - f_i(x_k))$ and $\zeta_{k+1} \triangleq (\mathbf{1}^T \otimes \mathbf{I}_l) \varepsilon_{k+1}$. Note that $\{\zeta_k\}$ is an MDS with bounded second moments by B1, $\beta_k = O(\gamma_k)$ a.s. by (64), and $\{x_k\}$ is bounded. Denote by $\lambda^{(1)}, \dots, \lambda^{(l)}$ the eigenvalues of A in a nonincreasing order. Then $\lambda^{(1)} > \lambda^{(p)} \quad \forall p > 1$ by B2. Multiplying both sides of (62) from the left with $(u^{(1)})^T$, by noticing that $(u^{(1)})^T A = \lambda^{(1)} (u^{(1)})^T$ and $f(x) = Ax - (x^T Ax)x$, we obtain the following:

$$\begin{aligned} (u^{(1)})^T x_{k+1} &= (u^{(1)})^T x_k + \frac{\gamma_k}{N} \left(\lambda^{(1)} - x_k^T A x_k \right) (u^{(1)})^T x_k \\ &\quad + \frac{\gamma_k}{N} (u^{(1)})^T (\zeta_{k+1} + \beta_{k+1}). \end{aligned} \quad (63)$$

Notice that $u^{(1)}$ is orthogonal to other eigenvectors of A and that x_k converges to some unit eigenvector of A by (61). So, to prove (54) it suffices to show that $\mathbb{P}(\Omega_1) = 0$, where $\Omega_1 = \{\omega : (u^{(1)})^T x_k(\omega) \rightarrow 0\}$. By definition of $\varepsilon_{i,k+1}$, we get

$$\begin{aligned} &\mathbb{E} \left[\|(u^{(1)})^T \varepsilon_{i,k+1}\| \middle| \mathcal{F}_k \right] \\ &= \mathbb{E} \left[\|(u^{(1)})^T N_{i,k} x_{i,k} - (x_{i,k}^T N_{i,k} x_{i,k}) (u^{(1)})^T x_{i,k}\| \middle| \mathcal{F}_k \right] \\ &\geq \mathbb{E} \left[\|(u^{(1)})^T N_{i,k} x_{i,k}\| \middle| \mathcal{F}_k \right] \\ &\quad - \left\| (u^{(1)})^T x_{i,k} \right\| \cdot \mathbb{E} \left[\|x_{i,k}^T N_{i,k} x_{i,k}\| \middle| \mathcal{F}_k \right]. \end{aligned}$$

Then by B3, $\liminf_k \mathbb{E} [\|(u^{(1)})^T \varepsilon_{i,k+1}\| | \mathcal{F}_k] > 0$ on Ω_1 , and hence $\liminf_k \mathbb{E} [\|(u^{(1)})^T \zeta_{k+1}\| | \mathcal{F}_k] > 0$ on Ω_1 . Note that on Ω_1 , by (61) we have that $\lambda^{(1)} - x_k^T A x_k \rightarrow \lambda^{(1)} - \lambda^{(p)} > 0$ for some $p > 1$. Then by (63) and [41, Lemma 1], we obtain that $\mathbb{P}(\Omega_1) = 0$. The proof is completed. ■

Remark 5.2: [41, Proposition] For any $i \in \mathcal{V}$, assume that $(N_{i,k}, \mathcal{F}_k)$ with $N_{i,k} = \{N_{i,k}(p,q)\}_{p,q=1}^l$ is an MDS, $\mathbb{E} [N_{i,k+1}(p,q)N_{i,k+1}(s,t) | \mathcal{F}_k] = 0$ whenever $(p,q) \neq (s,t)$, and $\liminf_k \mathbb{E} [\|N_{i,k+1}(p,q)\|^2 | \mathcal{F}_k] > 0 \forall p,q = 1, \dots, l$. Then B3 holds.

In Corollary 5.1 “non-convergence of X_k to zero” is assumed rather than proved, but the numerical simulations substantiate this assertion. This is not surprising because if we compare (62) with $u_k^{(i)}$ given in [37], the estimates for unit eigenvectors corresponding to eigenvalues arranged in the decreasing order of a noisy observed matrix A (pp. 289–316 in [37]), we find that (62) and the recursive expression ((5.1.9) of [37]) of $u_k^{(1)}$ are in a complete similarity while the latter converges to the unit eigenvector corresponding to the largest eigenvalue of A as proved in [37]. It is worth noting that normalization technique used in [37] automatically excludes estimates from tending to zero, but this can also be achieved by truncations as done in [36, Section 6.2] when solving the adaptive stabilization problem, where not only the upper side truncations are used to prevent estimates from diverging to infinity but also the lower side truncations for estimates not tending to zero. We might apply this kind of idea to the formulated distributed root-seeking problem, then the modified distributed stochastic approximation algorithm would converge to nonzero roots of sum functions.

Example 5.3: Distributed Principal Component Analysis

Let $N = 1000$. The matrix $W(k)$ is given by the following:

$$W(3k-2) = \begin{pmatrix} W_1 & \mathbf{0} \\ \mathbf{0} & I_{N/2} \end{pmatrix}, \quad W(3k-1) = \begin{pmatrix} I_{N/2} & \mathbf{0} \\ \mathbf{0} & W_2 \end{pmatrix},$$

$$W(3k) = \begin{pmatrix} \frac{1}{2} I_{N/2} & \frac{1}{2} I_{N/2} \\ \frac{1}{2} I_{N/2} & \frac{1}{2} I_{N/2} \end{pmatrix},$$

where matrices $W_1 \in \mathbb{R}^{N/2 \times N/2}$ and $W_2 \in \mathbb{R}^{N/2 \times N/2}$ are doubly stochastic. Further, let W_1 and W_2 be the adjacency matrices of some strongly connected digraphs. Thus, A4 holds.

Each sensor $i = 1, 2, \dots, N$ has access to a 9-dimensional i.i.d. Gaussian sequence $\{u_{i,k}\}$ with zero mean. Set $\gamma_k = \frac{1}{k}$, $M_k = 2^k$. Let the sequence $\{x_{i,k}\}$ be produced by (6)–(9) with $O_{i,k+1}$ defined by (3) and with $x_{i,0} = x^* = \mathbf{1}/\sqrt{N}$. Denote by x_k^i the i th component of $x_k = \frac{1}{N} \sum_{i=1}^N x_{i,k}$, and by $e(k) = \sum_{i=1}^N \|x_{i,k} - u^{(1)}\|_2 / N$ the average of 2-norm errors for all agents at time k . The estimates x_k are demonstrated in Fig. 1 with the solid lines denoting the true values and the dotted, dashed, and dash-dot lines representing the estimates. Meanwhile, the estimation errors $e(k)$ are demonstrated in Fig. 1. From the figure it is seen that the estimates converge to the unit eigenvector corresponding to the largest eigenvalue of A .

For a comparison, we also present the simulation results computed by DSAA proposed in [13] for the same example. The initial value for each agent, α_k and γ_k are set to be the same as those used in DSAAWET. The simulation results are given in Table II with $e + 11$ denoting the exponential 10^{11} ,




Fig. 1: The estimation sequences and estimation errors

from which it is seen that the estimates are unbounded. When a smaller step size, say $\gamma_k = \frac{1}{k+20}$, is taken, then the estimates are bounded for some sample paths and are unbounded for others. It is hard to indicate how small the step-sizes should be for a given sample path in advance. As a matter of fact, DSAAWET adaptively chooses step-sizes for any sample path.

k	0	1	2	3	4
x_k^1	1/3	1.9	-587	$5.46e + 11$	$-1.49e + 41$
x_k^2	1/3	3.2	-593	$3.64e + 11$	$-9.89e + 40$
x_k^3	1/3	0.277	-45.4	$-9.9e + 10$	$2.93e + 40$
x_k^4	1/3	-1.26	402	$-6e + 11$	$1.72e + 41$
x_k^5	1/3	-3.26	736	$-2.23e + 11$	$4.76e + 40$
x_k^6	1/3	0.25	-17.8	$7.86e + 10$	$-2.23e + 40$
x_k^7	1/3	3.3	-711	$5.02e + 11$	$-1.34e + 41$
x_k^8	1/3	0.19	2.48	$-1.76e + 11$	$5.54e + 40$
x_k^9	1/3	-1.63	397	$-2.37e + 11$	$6.38e + 40$

TABLE II: Estimates produced by DSAA in [13]

B. Distributed Gradient-free Optimization

We now consider the convergence of DSAAWET applied to the distributed gradient-free optimization problem.

Corollary 5.4: Let $\{x_{i,k}\}$ be produced by (6)–(9) with $O_{i,k+1}$ given by (5) for any initial value $x_{i,0}$. Assume that A4 holds, and, in addition,

C1 i) $\gamma_k > 0$, $\sum_{k=1}^{\infty} \gamma_k = \infty$, and $\sum_{k=1}^{\infty} \gamma_k^p < \infty$ for some $p \in (1, 2]$;

ii) $\alpha_k > 0$, and $\alpha_k \rightarrow 0$ as $k \rightarrow \infty$;

iii) $\sum_{k=1}^{\infty} \gamma_k^2 / \alpha_k^2 < \infty$;

C2 i) $c_i(\cdot)$, $i = 1, 2, \dots, N$ are continuously differentiable, and $\nabla c(\cdot)$ is locally Lipschitz continuous;

ii) $c(\cdot)$ is convex with a unique global minimum x_{min} ;

iii) $c(x^*) < \sup_{\|x\|=c_0} c(x)$ and $\|x^*\| < c_0$ for some positive constant c_0 , where x^* is used in (7) (8);

C3 $\xi_{i,k+1} = \xi_{i,k+1}^+ - \xi_{i,k+1}^-$ is independent of $\{\Delta_{i,s}, s = 1, 2, \dots, k\}$ for any $k \geq 0$, and $\xi_{i,k+1}$ satisfies one of the following two conditions:

i) $\sup_k |\xi_{i,k}| \leq \xi_i$ a.s., where ξ_i may be random;

ii) $\sup_k E[\xi_{i,k+1}^2] < \infty$
for any $i = 1, 2, \dots, N$. Then

$$x_{i,k} \xrightarrow[k \rightarrow \infty]{} x_{min} \quad a.s. \quad (64)$$

for any $i = 1, 2, \dots, N$.

Proof: To apply Theorem 3.4 to this problem, we have to verify conditions A1-A5.

By C1 the step-size $\{\gamma_k\}$ satisfies A1. Since $c(\cdot)$ is convex and differentiable, x_{min} is the global minimum of $c(\cdot)$ if and only if $\nabla c(x_{min}) = 0$. So, the original problem (4) is equivalent to finding the root $J = \{x_{min}\}$ of $f(\cdot) = \sum_{i=1}^N f_i(\cdot)/N$ with $f_i(\cdot) = -\nabla c_i(\cdot)$. By setting $v(\cdot) = c(\cdot)$, we obtain that $f^T(x)v_x(x) = -\|\nabla c(x)\|^2/N$ and $v(J) = \{c(x_{min})\}$. So, A2 a) and A2 b) hold. By C2 iii) we have A2 c). It is clear that C2 i) and iii) imply A3.

By (2) we have that

$$\varepsilon_{i,k+1} = O_{i,k+1} + \nabla c_i(x_{i,k}), \quad (65)$$

where $O_{i,k+1}$ is given by (5). The analysis for the observation noise $\{\varepsilon_{i,k+1}\}$ is the same as that given in [28], and hence is omitted here. It is shown in [28] that $\{\varepsilon_{i,k+1}\} \forall i \in \mathcal{V}$ is not an MDS, while using the local Lipschitz continuity of $\nabla c(\cdot)$, C1 and C3, by [28, Lemma 2], the following limit takes place:

$$\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{n=k}^{m(k,t_k)} \gamma_m \varepsilon_{i,n+1} I_{[\|x_{i,n}\| \leq K]} \right\| = 0 \quad (66)$$

$\forall t_k \in [0, T]$ for any positive integer K , *a.s.*

Therefore, A5 b) holds with probability one for any $i \in \mathcal{V}$. By Theorem 3.4 i), we see that $\{x_{i,k}\}$ is bounded almost surely. So, A5 a) holds by taking $t_k = \gamma_k$ in (66).

In summary, we have shown that A1-A5 hold. Since $J = \{x_{min}\}$, by Theorem 3.4 ii) we derive (64). ■

Remark 5.5: If the convex function $c(\cdot)$ is allowed to have non-unique minima while other conditions in Corollary 5.4 remain unchanged, then by Theorem 3.4 we know that $x_{i,k} \forall i \in \mathcal{V}$ converge to a connected subset $J^* \subset J \triangleq \{x \in \mathbb{R}^l : \nabla c(x) = 0\}$.

Remark 5.6: Compared with the biased mean error-bounds for the local weighted average vector given in [31] [32], the almost sure convergence of the iterates to the global optimal solution set is established here. Regarding the problem settings: i) each nonsmooth objective is assumed to be convex and Lipschitz continuous in [31] [32], while in this work, each objective function is not necessarily convex but required to be continuously differentiable with the gradient function being locally Lipschitz continuous; ii) a convex set constraint optimization is considered in [31] [32] while the unconstrained optimization is considered here; iii) the noisy observations of the objective functions are considered here in contrast to the exact observations discussed in [31] [32].

Example 5.7: Distributed Gradient-free Optimization

Consider the network of three agents with local cost functions given by

$$\begin{aligned} L_1(x, y) &= x^2 + y^2 + 10 \sin(x); \\ L_2(x, y) &= (x - 4)^2 + (y - 1)^2 - 10 \sin(x); \\ L_3(x, y) &= 0.01(x - 2)^4 + (y - 2)^2. \end{aligned}$$

Let the communication relationship among the agents be described by a strongly connected digraph with the adjacency matrix being doubly stochastic. The task of the network is to find the minimum $(x^0, y^0) = (2, 1)$ of the cost function $L(x, y) = \sum_{i=1}^3 L_i(x, y)$. Though each local cost function is non-convex, the global cost function $L(x, y)$ is convex.

Let the observation noise for the cost function of each agent be a sequence of i.i.d. random vectors $\in \mathcal{N}(0, I)$. The first and second component of the initial values for all agents are set to be mutually independent and uniformly distributed over the intervals $[-2, 6]$ and $[-2, 4]$, respectively. Set $x^* = (-1, 4)^T$, $\gamma_k = \frac{2}{k}$, $\alpha_k = \frac{1}{k^{0.2}}$, and $M_k = 2^k$. Let the estimates for the minimum be produced by (6)–(9) with $O_{i,k+1}$ defined by (5). The estimates of x^0 and y^0 produced by the three agents are demonstrated in Fig. 2, where $x_i(k)$ and $y_i(k)$ denote the estimates for x^0 and y^0 , respectively, given at agent i and at time k . From the figure it is seen that the estimates given by all agents converge to the minimum, which is consistent with the theoretic result.




Fig. 2: Trajectories of the estimates

VI. CONCLUDING REMARKS

In this paper, DSAAWET is defined to solve the formulated distributed root-seeking problem. The estimates are shown to converge to a consensus set belonging to a connected subset of the root set. Two problems as examples of those which can be solved by DSAAWET are demonstrated with numerical simulations provided.

For further research it is of interest to analyze the convergence rate of the proposed algorithm, and to consider the convergence properties of DSAAWET over random networks taking into account the possible packet loss in communication.

APPENDIX A PROOF LEMMA 4.1

Before proving the lemma, we show that for any $k \in [\tau_m, \tau_{m+1})$, the following assertions hold:

$$i) \quad \tilde{x}_{i,k} = x^*, \quad \tilde{\varepsilon}_{i,k+1} = -f_i(x^*) \text{ if } \sigma_{i,k} < m; \quad (A.1)$$

$$\text{ii) } \tilde{x}_{i,k} = x_{i,k}, \quad \tilde{\varepsilon}_{i,k+1} = \varepsilon_{i,k+1} \quad \text{if } \sigma_{i,k} = m; \quad (\text{A.2})$$

$$\text{iii) } \tilde{x}_{j,k} = x^* \quad \text{if } \sigma_{j,k-1} < m; \quad (\text{A.3})$$

$$\text{iv) } \tilde{x}_{j,k+1} = x^* \quad \forall j \in \mathcal{V} \quad \text{if } \sigma_{k+1} = m + 1. \quad (\text{A.4})$$

Proof: i) From $\sigma_{i,k} < m$ we see that the truncation number of agent i is smaller than m at time k , then by the definition of $\tau_{i,m}$ we derive $\tau_{i,m} > k$. Thus, $\tilde{\tau}_{i,m} = \tau_{i,m} \wedge \tau_{m+1} > k$, and hence from (18) we conclude (A.1).

ii) From $\sigma_{i,k} = m$ by definition we have $\tau_{i,m} \leq k$, and hence $\tilde{\tau}_{i,m} = \tau_{m+1} \wedge \tau_{i,m} = \tau_{i,m} \leq k$. Then by (19) it is clear that (A.2) holds.

iii) By $\tau_m \leq k < \tau_{m+1}$ we see $\sigma_{j,k} \leq m \quad \forall j \in \mathcal{V}$. We show (A.3) separately for the cases $\sigma_{j,k} = m$ and $\sigma_{j,k} < m$. 1) Let us first consider the case $\sigma_{j,k} = m$. Since $\sigma_{j,k-1} < m$ and $\sigma_{j,k} = m$, by (11) we obtain $x_{j,k} = x^*$. Hence from (A.2) we see $\tilde{x}_{j,p} = x_{j,p} = x^*$. 2) We now consider the case $\sigma_{j,k} < m$. By (A.1) we see $\tilde{x}_{j,k} = x^*$, which is the assertion of (A.3).

iv) From $k \in [\tau_m, \tau_{m+1})$ we see $\sigma_k = m$. Thus from $\sigma_{k+1} = m + 1$ by definition we derive $\tau_{m+1} = k + 1$, and hence $k + 1 \in [\tau_{m+1}, \tau_{m+2})$. By $\sigma_k = m$ we see $\sigma_{j,k} < m + 1 \quad \forall j \in \mathcal{V}$, then we derive (A.4) by (A.3). ■

Proof of Lemma 4.1: We prove the lemma by induction.

We first prove (20)-(22) for $k = 0$. Since $0 \in [\tau_0, \tau_1)$ and $\sigma_{i,0} = 0 \quad \forall i \in \mathcal{V}$, by (A.2) we derive $\tilde{x}_{i,0} = x_{i,0}$, $\tilde{\varepsilon}_{i,1} = \varepsilon_{i,1} \quad \forall i \in \mathcal{V}$. Then by noticing $\hat{\sigma}_{i,0} = \sigma_{i,0} = 0 \quad \forall i \in \mathcal{V}$, from (7) (20) we see

$$\hat{x}_{i,1} = x'_{i,1} \quad \forall i \in \mathcal{V}. \quad (\text{A.5})$$

We now show that $\tilde{x}_{i,1}$ and σ_1 generated by (20)-(22) are consistent with their definitions (18) (19) (10) by considering the following two cases:

i) There is no truncation at $k = 1$, i.e., $\sigma_{i,1} = 0 \quad \forall i \in \mathcal{V}$. In this case, from $\hat{\sigma}_{i,0} = 0 \quad \forall i \in \mathcal{V}$ by (9) we have $\|x'_{i,1}\| \leq M_0 \quad \forall i \in \mathcal{V}$, and hence $\|\hat{x}_{i,1}\| \leq M_0 \quad \forall i \in \mathcal{V}$ by (A.5). Then $x_{i,1} = x'_{i,1}$ by (8), $\tilde{x}_{i,1} = \hat{x}_{i,1}$ and $\sigma_1 = 0$ by (21) and (22), respectively. These together with (A.5) imply that $\tilde{x}_{i,1} = x_{i,1} \quad \forall i \in \mathcal{V}$, which is consistent with the definition of $\tilde{x}_{i,1}$ given by (19) since $\tilde{\tau}_{i,0} \leq 1 < \tau_1$. By (10) we see $\sigma_1 = \max_{i \in \mathcal{V}} \sigma_{i,1} = 0$, which is consistent with that derived from (20)-(22).

ii) There is a truncation at $k = 1$ for some agent i_0 , i.e., $\sigma_{i_0,1} = 1$. In this case, by (8) (9) we derive $x_{i_0,1} = x^*$, $\|x'_{i_0,1}\| > M_0$, and hence $\|\hat{x}_{i_0,1}\| > M_0$ by (A.5). Therefore, $\tilde{x}_{i,1} = x^* \quad \forall i \in \mathcal{V}$ and $\sigma_1 = 1$ by (21) (22). By (10) from $\sigma_{i_0,1} = 1$ we derive $\sigma_1 = \max_{i \in \mathcal{V}} \sigma_{i,1} = 1$. Since $0 \in [\tau_0, \tau_1)$ and $\sigma_1 = 1$, by (A.4) we see $\tilde{x}_{i,1} = x^* \quad \forall i \in \mathcal{V}$. Thus, $\tilde{x}_{i,1}$ and σ_1 defined by (18) (19) and (10) are consistent with those produced by (20)-(22).

In summary, we have proved the lemma for $k = 0$.

Inductively, we assume (20)-(22) hold for $k = 0, 1, \dots, p$. At the fixed sample path ω for a given integer p there exists a unique integer m such that $\tau_m \leq p < \tau_{m+1}$. We intend to show (20)-(22) also hold for $k = p + 1$.

Before doing this, we first express $\hat{x}_{i,p+1} \quad \forall i \in \mathcal{V}$ produced by (20) for the following two cases:

Case 1: $\sigma_{i,p} < m$. Since $p \in [\tau_m, \tau_{m+1})$, by (A.1) we see

$$\tilde{x}_{i,p} = x^*, \quad \tilde{\varepsilon}_{i,p+1} = -f_i(x^*). \quad (\text{A.6})$$

From $\sigma_{i,p} < m$ it follows that $\sigma_{j,p-1} < m \quad \forall j \in N_i(p)$, because otherwise, there would exist a $j_1 \in N_i(p)$ such that $\sigma_{j_1,p-1} \geq m$. Hence by (6) (9) we would derive $\sigma_{i,p} \geq \hat{\sigma}_{i,p-1} \geq \sigma_{j_1,p-1} \geq m$, yielding a contradiction.

From $\sigma_{j,p-1} < m \quad \forall j \in N_i(p)$ by (A.3) we have $\tilde{x}_{j,p} = x^* \quad \forall j \in N_i(p)$, which incorporating with (20) (A.6) yields

$$\hat{x}_{i,p+1} = x^* \quad \forall i : \sigma_{i,p} < m. \quad (\text{A.7})$$

Case 2: $\sigma_{i,p} = m$. By $\tau_m \leq p < \tau_{m+1}$ we see $\sigma_{j,p} \leq m \quad \forall j \in \mathcal{V}$, and hence by (6) we obtain

$$\hat{\sigma}_{i,p} = m \quad \forall i : \sigma_{i,p} = m. \quad (\text{A.8})$$

Then by (7)

$$\begin{aligned} x'_{i,p+1} &= \sum_{j \in N_i(p)} \omega_{ij}(p) (x_{j,p} I_{[\sigma_{j,p}=m]} + x^* I_{[\sigma_{j,p}<m]}) \\ &+ \gamma_p (f_i(x_{i,p}) + \varepsilon_{i,p+1}). \end{aligned} \quad (\text{A.9})$$

From $\sigma_{i,p} = m$ and $p \in [\tau_m, \tau_{m+1})$, by (A.2) it is clear that

$$\tilde{x}_{i,p} = x_{i,p}, \quad \tilde{\varepsilon}_{i,p+1} = \varepsilon_{i,p+1}. \quad (\text{A.10})$$

By the first term in (A.1) and (A.2), for any $j \in N_i(p)$ we have

$$\tilde{x}_{j,p} = x_{j,p} \quad \text{if } \sigma_{j,p} = m, \quad \tilde{x}_{j,p} = x^* \quad \text{if } \sigma_{j,p} < m. \quad (\text{A.11})$$

Substituting (A.10) (A.11) into (20), from (A.9) we derive

$$\hat{x}_{i,p+1} = x'_{i,p+1} \quad \forall i : \sigma_{i,p} = m. \quad (\text{A.12})$$

Since $\tau_m \leq p < \tau_{m+1}$, from $p < \tau_{m+1}$ it follows that $\sigma_p < m + 1$, and hence $\sigma_{p+1} \leq m + 1$, while from $\tau_m \leq p$ it follows that $\sigma_p = m$, and hence $\sigma_{p+1} \geq m$. Thus, we have $m \leq \sigma_{p+1} \leq m + 1$.

We now show that $\tilde{x}_{i,p+1}$ and σ_{p+1} generated by (20)-(22) are consistent with their definitions (18) (19) (10). We prove this separately for the cases $\sigma_{p+1} = m + 1$ and $\sigma_{p+1} = m$.

Case 1: $\sigma_{p+1} = m + 1$. We first show

$$\sigma_{i,p+1} \leq m \quad \text{if } \sigma_{i,p} < m \quad (\text{A.13})$$

separately for the following two cases 1) and 2): 1) $\sigma_{i,p} < m$ and $\sigma_{j,p} < m \quad \forall j \in N_i(p)$. For this case by (6) we derive $\hat{\sigma}_{i,p} < m$, and hence $\sigma_{i,p+1} \leq \hat{\sigma}_{i,p} + 1 \leq m$ by (9). 2) $\sigma_{i,p} < m$ and $\sigma_{j,p} = m$ for some $j \in N_i(p)$. For this case we obtain $\hat{\sigma}_{i,p} = m$, $x'_{i,p+1} = x^*$ by (6) (7). Since $\|x^*\| \leq M_0 \leq M_m$, then by (9) $\sigma_{i,p+1} = \hat{\sigma}_{i,p} = m$. Thus, $\sigma_{i,p+1} \leq m$ when $\sigma_{i,p} < m$. Thereby (A.13) holds. This means that

$$\sigma_{i,p+1} = m + 1 \quad \text{only if } \sigma_{i,p} = m. \quad (\text{A.14})$$

By definition from $\sigma_{p+1} = m + 1$ we know that there exists some $i_0 \in \mathcal{V}$ such that $\sigma_{i_0,p+1} = m + 1$. Then $\sigma_{i_0,p} = m$ by (A.14), and hence $\hat{\sigma}_{i_0,p} = m$ from (A.8). Then from $\sigma_{i_0,p+1} = m + 1$ by (9) we derive $\|x'_{i_0,p+1}\| > M_m$, and hence $\|\hat{x}_{i_0,p+1}\| = \|x'_{i_0,p+1}\| > M_m$ by (A.12). Then from (21) (22) we derive $\tilde{x}_{i,p+1} = x^* \quad \forall i \in \mathcal{V}$ and $\sigma_{p+1} = m + 1$, which is consistent with σ_{p+1} defined by (10). Since $\sigma_{p+1} = m + 1$ and $p \in [\tau_m, \tau_{m+1})$, by (A.4) or from (18) (19), we see $\tilde{x}_{i,p+1} = x^* \quad \forall i \in \mathcal{V}$. This is consistent with that produced by (20)-(22).

Case 2: We now consider the case $\sigma_{p+1} = m$. In this case, $\sigma_{i,p+1} \leq m \forall i \in \mathcal{V}$. By (A.8), from (8) (9) we see

$$\|x'_{i,p+1}\| \leq M_m, \quad x_{i,p+1} = x'_{i,p+1} \quad \forall i : \sigma_{i,p} = m. \quad (\text{A.15})$$

So, by (A.12) we have

$$\|\hat{x}_{i,p+1}\| = \|x'_{i,p+1}\| \leq M_m \quad \forall i : \sigma_{i,p} = m. \quad (\text{A.16})$$

From $\|x^*\| \leq M_0 \leq M_m$ and (A.7) we derive $\|\hat{x}_{i,p+1}\| \leq M_m \forall i : \sigma_{i,p} < m$, which incorporating with (A.16) yields $\|\hat{x}_{i,p+1}\| \leq M_m \forall i \in \mathcal{V}$. Then from (21) we have

$$\tilde{x}_{i,p+1} = \hat{x}_{i,p+1} \quad \forall i \in \mathcal{V}, \quad \sigma_{p+1} = m. \quad (\text{A.17})$$

Thus σ_{p+1} is consistent with that defined by (10).

It remains to show that $\tilde{x}_{i,p+1}$ generated by (20)-(22) is consistent with that defined by (18) (19). We consider two cases: 1) $\sigma_{i,p} = m$. For this case, by (A.12) (A.15) (A.17) we see $\tilde{x}_{i,p+1} = x_{i,p+1} \forall i : \sigma_{i,p} = m$. By $\sigma_{p+1} = m$ we see $p+1 \in [\tau_m, \tau_{m+1})$, and hence $\tilde{x}_{i,p+1} = x_{i,p+1}$ by (A.2). So, the assertion holds for any i with $\sigma_{i,p} = m$. 2) $\sigma_{i,p} < m$. From $\sigma_{p+1} = m$ we see $p+1 \in [\tau_m, \tau_{m+1})$, and hence by $\sigma_{i,p} < m$ from (A.3) we see that $\tilde{x}_{i,p+1}$ defined by (18) (19) equals x^* . By (A.7) (A.17) we derive $\tilde{x}_{i,p+1} = x^*$, and hence the assertion holds for any i with $\sigma_{i,p} < m$.

In summary, we have shown that $\tilde{x}_{i,p+1}$ and σ_{p+1} generated by (20)-(22) are consistent with their definitions (18) (19) (10). This completes the proof. ■

APPENDIX B PROOF OF LEMMA 4.5

For the lemma it suffices to prove

$$\begin{aligned} & \lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, t_k) \wedge (\tau_{\sigma_{n_k}+1-1})} \gamma_s \tilde{\varepsilon}_{i,s+1} I_{[\|\tilde{x}_{i,s}\| \leq K]} \right\| \\ & = 0 \quad \forall t_k \in [0, T] \text{ for sufficiently large } K > 0 \end{aligned} \quad (\text{B.1})$$

along indices $\{n_k\}$ whenever $\{\tilde{x}_{i,n_k}\}$ converges for the sample path ω where A5 b) holds for agent i .

We consider the following two cases:

Case 1: $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$. We now show

$$\tau_{\sigma+1} = \infty \text{ when } \lim_{k \rightarrow \infty} \sigma_k = \sigma. \quad (\text{B.2})$$

Recall that σ_k is defined as the largest truncation number among all agents at time k , from $\lim_{k \rightarrow \infty} \sigma_k = \sigma$ we have $\sigma_{i,k} \leq \sigma \quad \forall k \geq 0 \quad \forall i \in \mathcal{V}$. From here by the definition of $\tau_{i,m}$ it follows that $\tau_{i,\sigma+1} = \inf\{k : \sigma_{i,k} = \sigma + 1\} = \infty \quad \forall i \in \mathcal{V}$, and hence $\tau_{\sigma+1} = \infty$. Thus, (B.2) holds.

From (24) we have $\tilde{\tau}_{i,\sigma} \leq \tau_{\sigma} + BD$, and hence by (19) (B.2)

$$\tilde{x}_{i,k} = x_{i,k}, \quad \tilde{\varepsilon}_{i,k+1} = \varepsilon_{i,k+1} \quad \forall k \geq \tau_{\sigma} + BD. \quad (\text{B.3})$$

So, $\left\| \sum_{s=k}^{m(k,t) \wedge (\tau_{\sigma_k+1-1})} \gamma_s \tilde{\varepsilon}_{i,s+1} I_{[\|\tilde{x}_{i,s}\| \leq K]} \right\| = \left\| \sum_{s=k}^{m(k,t)} \gamma_s \varepsilon_{i,s+1} I_{[\|x_{i,s}\| \leq K]} \right\|$ for any $t > 0$ and any sufficiently large k . Then we conclude (B.1) by A5 b).

Case 2: $\lim_{k \rightarrow \infty} \sigma_k = \infty$. We prove (B.1) separately for the following three cases:

i) $\tilde{\tau}_{i,\sigma_{n_p}} \leq n_p$. For this case, $[n_p, \tau_{\sigma_{n_p}+1}) \subset [\tilde{\tau}_{i,\sigma_{n_p}}, \tau_{\sigma_{n_p}+1})$, and hence from (19) we derive

$$\tilde{x}_{i,s} = x_{i,s}, \quad \tilde{\varepsilon}_{i,s+1} = \varepsilon_{i,s+1} \quad \forall s : n_p \leq s < \tau_{\sigma_{n_p}+1}. \quad (\text{B.4})$$

Thus, for sufficiently large K and any $t_p \in [0, T]$

$$\begin{aligned} & \left\| \sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \tilde{\varepsilon}_{i,s+1} I_{[\|\tilde{x}_{i,s}\| \leq K]} \right\| \\ & = \left\| \sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \varepsilon_{i,s+1} I_{[\|x_{i,s}\| \leq K]} \right\|. \end{aligned} \quad (\text{B.5})$$

By (B.4) we conclude that $\{x_{i,n_p}\}$ is a convergent subsequence. Noticing $\sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \leq \sum_{s=n_p}^{m(n_p, t_p)} \gamma_s \leq t_p \leq T$, we then conclude (B.1) by (B.5) and A5 b).

ii) $\tilde{\tau}_{i,\sigma_{n_p}} > n_p$ and $\tilde{\tau}_{i,\sigma_{n_p}} = \tau_{\sigma_{n_p}+1}$. By the definitions of τ_k and σ_k we derive $\tau_{\sigma_k} \leq k$, and hence $\tau_{\sigma_{n_p}} \leq n_p$. Then $[n_p, \tau_{\sigma_{n_p}+1}) \subset [\tau_{\sigma_{n_p}}, \tilde{\tau}_{i,\sigma_{n_p}})$, and hence by (18) we have

$$\tilde{x}_{i,s} = x^*, \quad \tilde{\varepsilon}_{i,s+1} = -f_i(x^*) \quad \forall s : n_p \leq s < \tau_{\sigma_{n_p}+1}.$$

From $\tilde{\tau}_{i,\sigma_{n_p}} = \tau_{\sigma_{n_p}+1}$ by (24) we see $\tau_{\sigma_{n_p}+1} \leq \tau_{\sigma_{n_p}} + BD \leq n_p + BD$. Then for sufficiently large K and any $t_p \in [0, T]$

$$\begin{aligned} & \left\| \sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \tilde{\varepsilon}_{i,s+1} I_{[\|\tilde{x}_{i,s}\| \leq K]} \right\| \\ & \leq \sum_{s=n_p}^{n_p + BD} \gamma_s \|f_i(x^*)\| \xrightarrow{p \rightarrow \infty} 0, \end{aligned}$$

and hence (B.1) holds.

iii) $\tilde{\tau}_{i,\sigma_{n_p}} > n_p$ and $\tilde{\tau}_{i,\sigma_{n_p}} < \tau_{\sigma_{n_p}+1}$. By the definition of $\tilde{\tau}_{i,\sigma_{n_p}}$ from $\tilde{\tau}_{i,\sigma_{n_p}} < \tau_{\sigma_{n_p}+1}$ it follows that $\tilde{\tau}_{i,\sigma_{n_p}} = \tau_{i,\sigma_{n_p}}$, and hence by (24) $\tau_{i,\sigma_{n_p}} \leq \tau_{\sigma_{n_p}} + BD$. By noticing $\tau_{\sigma_{n_p}} \leq n_p$ we conclude that

$$\tau_{\sigma_{n_p}} \leq n_p < \tilde{\tau}_{i,\sigma_{n_p}} = \tau_{i,\sigma_{n_p}} \leq n_p + BD. \quad (\text{B.6})$$

So, $[n_p, \tau_{i,\sigma_{n_p}}) \subset [\tau_{\sigma_{n_p}}, \tilde{\tau}_{i,\sigma_{n_p}})$. Then from here and $\tilde{\tau}_{i,\sigma_{n_p}} = \tau_{i,\sigma_{n_p}}$ by (18) (19) we derive

$$\begin{aligned} \tilde{x}_{i,s} &= x^*, \quad \tilde{\varepsilon}_{i,s+1} = -f_i(x^*) \quad \forall s : n_p \leq s < \tau_{i,\sigma_{n_p}}, \\ \tilde{x}_{i,s} &= x_{i,s}, \quad \tilde{\varepsilon}_{i,s+1} = \varepsilon_{i,s+1} \quad \forall s : \tau_{i,\sigma_{n_p}} \leq s < \tau_{\sigma_{n_p}+1}. \end{aligned}$$

Consequently, for sufficiently large K and any $t_p \in [0, T]$

$$\begin{aligned} & \left\| \sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \tilde{\varepsilon}_{i,s+1} I_{[\|\tilde{x}_{i,s}\| \leq K]} \right\| \\ & \leq \left\| \sum_{s=n_p}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s f_i(x^*) I_{[n_p \leq s < \tau_{i,\sigma_{n_p}}]} \right\| \\ & + \left\| \sum_{s=\tau_{i,\sigma_{n_p}}}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \varepsilon_{i,s+1} I_{[\|x_{i,s}\| \leq K]} \right\|. \end{aligned} \quad (\text{B.7})$$

Note that the first term at the right hand of (B.7) is smaller than $\sum_{s=n_p}^{\tau_{i,\sigma_{n_p}}} \gamma_s \|f_i(x^*)\|$, which tends to zero as $k \rightarrow \infty$ by the last inequality in (B.6) and $\lim_{k \rightarrow \infty} \gamma_k = 0$. The truncation number for agent i at time $\tau_{i,\sigma_{n_p}}$ is σ_{n_p} , while it is smaller than σ_{n_p} at time $\tau_{i,\sigma_{n_p}} - 1$ since $\tau_{i,\sigma_{n_p}}$ is the smallest time when the truncation number of i has reached σ_{n_p} . Consequently, by Remark 3.1 we have $x_{i,\tau_{i,\sigma_{n_p}}} = x^*$, and hence $\{x_{i,\tau_{i,\sigma_{n_p}}}\}_{p \geq 1}$ is a convergent subsequence. Noticing

$$\sum_{s=\tau_{i,\sigma_{n_p}}}^{m(n_p, t_p) \wedge (\tau_{\sigma_{n_p}+1-1})} \gamma_s \leq \sum_{s=n_p}^{m(n_p, t_p)} \gamma_s \leq t_p,$$

we conclude (B.1) by (B.7) and A5 b).

Since one of i), ii), iii) must take place for the case $\lim_{k \rightarrow \infty} \sigma_k = \infty$, we thus have proved (B.1) in Case 2.

Combining Case 1 and Case 2 we derive (B.1). ■

APPENDIX C
PROOF OF LEMMA 4.6

Let us consider a fixed ω where A5 b) holds.

Let $C > \|\tilde{X}\|$. There exists an integer $k_C > 0$ such that

$$\|\tilde{X}_{n_k}\| \leq C \quad \forall k \geq k_C. \quad (\text{C.1})$$

By Lemma 4.5, there exist a constant $T_1 > 0$ and a positive integer $k_0 \geq k_C$ such that

$$\left\| \sum_{s=n_k}^{m(n_k, t_k) \wedge (\tau_{\sigma_{n_k}+1}-1)} \gamma_s \tilde{\varepsilon}_{s+1} I_{\|\tilde{X}_s\| \leq K} \right\| \leq T_0 \quad (\text{C.2})$$

$$\forall t_k \in [0, T_0] \quad \forall T_0 \in [0, T_1] \quad \forall k \geq k_0$$

for sufficiently large $K > 0$. Define

$$M'_0 \triangleq 1 + C(c\rho + 2), \quad (\text{C.3})$$

$$H_1 \triangleq \max_X \{ \|F(X)\| : \|X\| \leq M'_0 + 1 + C \}, \quad (\text{C.4})$$

$$c_1 \triangleq H_1 + 3 + \frac{c(\rho + 1)}{1 - \rho}, \quad \text{and} \quad c_2 \triangleq \frac{H_1 + 1}{\sqrt{N}}, \quad (\text{C.5})$$

where c and ρ are given by (13). Select $T > 0$ such that

$$0 < T \leq T_1 \quad \text{and} \quad c_1 T < 1. \quad (\text{C.6})$$

For any $k \geq k_0$ and any $T_k \in [0, T]$, define

$$s_k \triangleq \sup \{ s \geq n_k : \|\tilde{X}_j - \tilde{X}_{n_k}\| \leq c_1 T_k + M'_0 \quad \forall j : n_k \leq j \leq s \}. \quad (\text{C.7})$$

Then from (C.1) and (C.6) it follows that

$$\|\tilde{X}_s\| \leq M'_0 + 1 + C \quad \forall s : n_k \leq s \leq s_k. \quad (\text{C.8})$$

We intend to prove $s_k > m(n_k, T_k)$. Assume the converse that for sufficiently large $k \geq k_0$ and any $T_k \in [0, T]$

$$s_k \leq m(n_k, T_k). \quad (\text{C.9})$$

We first show that there exists a positive integer $k_1 > k_0$ such that for sufficiently large $k \geq k_1$

$$s_k < \tau_{\sigma_{n_k}+1} \quad \forall k \geq k_1 \quad \forall T_k \in [0, T]. \quad (\text{C.10})$$

We prove (C.10) for the two alternative cases: $\lim_{k \rightarrow \infty} \sigma_k = \infty$ and $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$.

i) $\lim_{k \rightarrow \infty} \sigma_k = \infty$. Since $\{M_k\}$ is a sequence of positive numbers increasingly diverging to infinity, there exists a positive integer $k_1 > k_0$ such that $M_{\sigma_{n_k}} > M'_0 + 1 + C$ for all $k \geq k_1$. Hence, from (C.8) we know $s_k < \tau_{\sigma_{n_k}+1}$.

ii) $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$. For this case there exists a positive integer $k_1 > k_0$ such that $\sigma_{n_k} = \sigma$ for all $k \geq k_1$, and hence $\tau_{\sigma_{n_k}+1} = \tau_{\sigma+1} = \infty$ by (B.2). Then $m(n_k, T_k) < \tau_{\sigma_{n_k}+1}$, and hence by (C.9) we derive (C.10).

By (C.6), we see that $T_k \in [0, T] \subset [0, T_1]$. Then from (C.2) it follows that for sufficiently large $k \geq k_1$ and $K > 0$:

$$\left\| \sum_{s=n_k}^{m(n_k, t_k) \wedge (\tau_{\sigma_{n_k}+1}-1)} \gamma_s \tilde{\varepsilon}_{s+1} I_{\|\tilde{X}_s\| \leq K} \right\| \leq T_k \quad (\text{C.11})$$

$$\forall t_k \in [0, T_k] \quad \forall T_k \in [0, T].$$

Set $t_k = \sum_{m=n_k}^s \gamma_m$ for some $s \in [n_k, s_k]$. Then $\sum_{m=n_k}^s \gamma_m \leq \sum_{m=n_k}^{s_k} \gamma_m \leq T_k$ by (C.9). Since $m(n_k, t_k) = s$, from (C.10) we have that $m(n_k, t_k) \wedge (\tau_{\sigma_{n_k}+1} - 1) = s$. Then by setting $K \triangleq M'_0 + 1 + C$, from (C.8) (C.11) it follows that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$:

$$\left\| \sum_{m=n_k}^s \gamma_m \tilde{\varepsilon}_{m+1} \right\| \leq T_k \quad \forall s : n_k \leq s \leq s_k. \quad (\text{C.12})$$

Let us consider the following algorithm starting from n_k without truncation

$$Z_{m+1} = (W(m) \otimes \mathbf{I}_l) Z_m + \gamma_m (F(Z_m) + \tilde{\varepsilon}_{m+1}), \quad Z_{n_k} = \tilde{X}_{n_k}. \quad (\text{C.13})$$

By (C.10) we know that (30) holds for $m = n_k, \dots, s_k - 1$ for $\forall k \geq k_1 \quad \forall T_k \in [0, T]$. Then we have the following:

$$Z_m = \tilde{X}_m \quad \forall m : n_k \leq m \leq s_k. \quad (\text{C.14})$$

Hence by (C.4) (C.8), we know that for $\forall k \geq k_1 \quad \forall T_k \in [0, T]$

$$\|F(Z_m)\| \leq H_1 \quad \forall m : n_k \leq m \leq s_k. \quad (\text{C.15})$$

Set $z_k = \frac{\mathbf{1}^T \otimes \mathbf{I}_l}{N} Z_k$. Multiplying both sides of (C.13) from left with $\frac{1}{N}(\mathbf{1}^T \otimes \mathbf{I}_l)$, by $\mathbf{1}^T W(m) = \mathbf{1}^T$ and (27), we derive

$$z_{s+1} = z_s + \frac{\mathbf{1}^T \otimes \mathbf{I}_l}{N} \gamma_s (F(Z_s) + \tilde{\varepsilon}_{s+1}),$$

and hence

$$\|z_{s+1} - z_{n_k}\| = \left\| \frac{\mathbf{1}^T \otimes \mathbf{I}_l}{N} \sum_{m=n_k}^s \gamma_m (F(Z_m) + \tilde{\varepsilon}_{m+1}) \right\| \leq \frac{1}{\sqrt{N}} \left(\sum_{m=n_k}^s \gamma_m \|F(Z_m)\| + \sum_{m=n_k}^s \gamma_m \|\tilde{\varepsilon}_{m+1}\| \right).$$

Then by (C.9), (C.12), and (C.15), we conclude that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$

$$\|z_{s+1} - z_{n_k}\| \leq \frac{H_1 + 1}{\sqrt{N}} \sum_{m=n_k}^s \gamma_m = c_2 T_k \quad \forall s : n_k \leq s \leq s_k. \quad (\text{C.16})$$

Denote by $Z_{\perp, s} \triangleq D_{\perp} Z_s$ the disagreement vector of Z_s . By multiplying both sides of (C.13) from left with D_{\perp} , we derive

$$Z_{\perp, s+1} = D_{\perp} (W(s) \otimes \mathbf{I}_l) Z_s + \gamma_s D_{\perp} (F(Z_s) + \tilde{\varepsilon}_{s+1}),$$

and inductively

$$Z_{\perp, s+1} = \Psi(s, n_k) Z_{n_k} + \sum_{m=n_k}^s \gamma_m \Psi(s-1, m) D_{\perp} (F(Z_m) + \tilde{\varepsilon}_{m+1}) \quad \forall s \geq n_k.$$

Thus, by (28) and (29) we have that

$$Z_{\perp, s+1} = \left[(\Phi(s, n_k) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l \right] Z_{n_k} + \sum_{m=n_k}^s \gamma_m \left[(\Phi(s-1, m) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l \right] F(Z_m) + \sum_{m=n_k}^s \gamma_m \left[(\Phi(s-1, m) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l \right] \tilde{\varepsilon}_{m+1}. \quad (\text{C.17})$$

By (C.1) and (C.14), we have that $\|Z_{n_k}\| \leq C$. Thus, from (13) and (C.15) it follows that

$$\|Z_{\perp, s+1}\| \leq C c \rho^{s+1-n_k} + \sum_{m=n_k}^s \gamma_m H_1 c \rho^{s-m} + \left\| \sum_{m=n_k}^s \gamma_m \left[(\Phi(s-1, m) - \frac{1}{N} \mathbf{1} \mathbf{1}^T) \otimes \mathbf{I}_l \right] \tilde{\varepsilon}_{m+1} \right\|. \quad (\text{C.18})$$

Define $\Gamma_n \triangleq \sum_{m=1}^n \gamma_m \tilde{\varepsilon}_{m+1}$. Then by (C.12), we obtain that $\|\Gamma_s - \Gamma_{n_k-1}\| \leq T_k \quad \forall s : n_k \leq s \leq s_k$. Note that

$$\begin{aligned} & \sum_{m=n_k}^s \gamma_m (\Phi(s-1, m) \otimes I_l) \tilde{\varepsilon}_{m+1} \\ &= \sum_{m=n_k}^s (\Phi(s-1, m) \otimes I_l) (\Gamma_m - \Gamma_{m-1}) \\ &= \sum_{m=n_k}^s (\Phi(s-1, m) \otimes I_l) (\Gamma_m - \Gamma_{n_k-1}) \\ &\quad - \sum_{m=n_k}^s (\Phi(s-1, m) \otimes I_l) (\Gamma_{m-1} - \Gamma_{n_k-1}). \end{aligned}$$

Summing by parts, by (13) we have

$$\begin{aligned} & \left\| \sum_{m=n_k}^s \gamma_m (\Phi(s-1, m) \otimes I_l) \tilde{\varepsilon}_{m+1} \right\| \leq \|\Gamma_s - \Gamma_{n_k-1}\| + \\ & \sum_{m=n_k}^{s-1} \|\Phi(s-1, m) - \Phi(s-1, m+1)\| \|\Gamma_m - \Gamma_{n_k-1}\| \\ & \leq T_k + \sum_{m=n_k}^{s-1} (c\rho^{s-m-1} + c\rho^{s-m}) T_k \\ & \leq T_k + \frac{c(\rho+1)}{1-\rho} T_k \quad \forall s : n_k \leq s \leq s_k. \end{aligned}$$

This incorporating with (C.12) yields that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$,

$$\begin{aligned} & \left\| \sum_{m=n_k}^s \gamma_m [(\Phi(s-1, m) - \frac{1}{N} \mathbf{1}\mathbf{1}^T) \otimes I_l] \tilde{\varepsilon}_{m+1} \right\| \\ & \leq (2 + \frac{c(\rho+1)}{1-\rho}) T_k \quad \forall s : n_k \leq s \leq s_k. \end{aligned} \quad (\text{C.19})$$

By noticing $\sum_{m=n_k}^s \gamma_m \rho^{s-m} \leq \frac{1}{1-\rho} \sup_{m \geq n_k} \gamma_m$, from $\lim_{k \rightarrow \infty} \gamma_k = 0$, (C.18), and (C.19) it follows that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$:

$$\begin{aligned} & \|Z_{\perp, s+1}\| \leq Cc\rho + \frac{cH_1}{1-\rho} \sup_{m \geq n_k} \gamma_m + (2 + \frac{c(\rho+1)}{1-\rho}) T_k \\ & \leq Cc\rho + 1 + (2 + \frac{c(\rho+1)}{1-\rho}) T_k \quad \forall s : n_k \leq s \leq s_k. \end{aligned} \quad (\text{C.20})$$

By noticing $Z_s = Z_{\perp, s} + (\mathbf{1} \otimes I_l) z_s$, we derive

$$\begin{aligned} & \|Z_{s+1} - Z_{n_k}\| \\ &= \|(\mathbf{1} \otimes I_l) z_{s+1} + Z_{\perp, s+1} - Z_{\perp, n_k} - (\mathbf{1} \otimes I_l) z_{n_k}\| \\ &\leq \|Z_{\perp, s+1}\| + \|Z_{\perp, n_k}\| + \sqrt{N} \|z_{s+1} - z_{n_k}\|. \end{aligned}$$

Since $\|Z_{\perp, n_k}\| \leq 2\|Z_{n_k}\| = 2C$, from (C.16) and (C.20) it follows that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$

$$\begin{aligned} & \|Z_{s+1} - Z_{n_k}\| \leq Cc\rho + 1 + (2 + \frac{c(\rho+1)}{1-\rho}) T_k + 2C + \\ & (H_1 + 1) T_k \leq C(c\rho + 2) + 1 + (3 + H_1 + \frac{c(\rho+1)}{1-\rho}) T_k \\ & = M'_0 + c_1 T_k \quad \forall s : n_k \leq s \leq s_k. \end{aligned} \quad (\text{C.21})$$

Therefore, from (C.6) and $\|Z_{n_k}\| \leq C$ we know that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$

$$\|Z_{s_k+1}\| \leq \|Z_{n_k}\| + M'_0 + c_1 T_k \leq M'_0 + 1 + C.$$

Rewrite (20) in the compact form as follows

$$\hat{X}_{s_k+1} = [W(s_k) \otimes I_l] \tilde{X}_{s_k} + \gamma_{s_k} (F(\tilde{X}_{s_k}) + \tilde{\varepsilon}_{s_k+1}),$$

where $\hat{X}_k \triangleq \text{col}\{\hat{x}_{1,k}, \dots, \hat{x}_{N,k}\}$. Then $\hat{X}_{s_k+1} = Z_{s_k+1}$ by (C.13) (C.14), and hence from (C.22) it follows that

$$\|\hat{X}_{s_k+1}\| \leq M'_0 + 1 + C. \quad (\text{C.22})$$

We now show

$$\tilde{X}_{s_k+1} = \hat{X}_{s_k+1} \text{ and } s_k + 1 < \tau_{\sigma_{n_k+1}} \quad (\text{C.23})$$

for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$. We consider the following two cases: $\lim_{k \rightarrow \infty} \sigma_k = \infty$ and $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$.

i) $\lim_{k \rightarrow \infty} \sigma_k = \infty$. By noting $M_{\sigma_{n_k}} > M'_0 + 1 + C \quad \forall k \geq k_1$, from (C.22), (21) and (22) we get $\tilde{X}_{s_k+1} = \hat{X}_{s_k+1}$ and $\sigma_{s_k+1} = \sigma_{s_k}$. Hence $s_k + 1 < \tau_{\sigma_{n_k+1}}$ by (C.10).

ii) $\lim_{k \rightarrow \infty} \sigma_k = \sigma < \infty$. Since $\tau_{\sigma_{n_k+1}} = \infty \quad \forall k \geq k_1$, by (C.9) we see $s_k + 1 < \tau_{\sigma_{n_k+1}}$. Then by $\sigma_{n_k} = \sigma \quad \forall k \geq k_1$ we conclude $\sigma_{s_k+1} = \sigma_{s_k} = \sigma$. Hence by (21) we derive $\tilde{X}_{s_k+1} = \hat{X}_{s_k+1}$. Thus, (C.23) holds.

From (C.23), we know that (30) holds for $m = s_k$ for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$. By $\hat{X}_{s_k+1} = Z_{s_k+1}$ and (C.23), we obtain $\tilde{X}_{s_k+1} = Z_{s_k+1}$. Hence from (C.21) and $Z_{n_k} = \tilde{X}_{n_k}$ it follows that for sufficiently large $k \geq k_1$ and any $T_k \in [0, T]$

$$\|\tilde{X}_{s_k+1} - \tilde{X}_{n_k}\| \leq M'_0 + c_1 T_k.$$

This contradicts with the definition of s_k given by (C.7). Thus, (C.9) does not hold. Consequently, $s_k > m(n_k, T_k)$, and hence by the definition of s_k given in (C.7) we derive (31).

Since $\{\tilde{X}_s : n_k \leq s \leq m(n_k, T_k)\}$ are bounded, similar to proving (C.10) it can be shown that $m(n_k, T_k) + 1 < \tau_{\sigma_{n_k+1}}$. As a result, (30) holds for $m = n_k, \dots, m(n_k, T_k)$. Similar to (C.16), we have the following assertion for sufficiently large k and any $T_k \in [0, T]$:

$$\|\bar{x}_{m+1} - \bar{x}_{n_k}\| \leq c_2 T_k \quad \forall m : n_k \leq m \leq m(n_k, T_k).$$

Hence, (32) holds. The proof of Lemma 4.6 is completed. ■

APPENDIX D

PROOF OF LEMMA 4.7

Since $\lim_{k \rightarrow \infty} \tilde{X}_{n_k} = \bar{X}$, we then have $\lim_{k \rightarrow \infty} \bar{x}_{n_k} = \bar{x} = \frac{\mathbf{1}^T \otimes I_l \bar{X}}{N}$. Lemma 4.6 ensures that there exists a $T \in (0, 1)$ such that for sufficiently large k , $m(n_k, T) < \tau_{\sigma_{n_k+1}}$ and $\{\tilde{X}_s : n_k \leq s \leq m(n_k, T) + 1\}$ are bounded. So, for any $T_k \in [0, T]$ and any sufficiently large K

$$\left\| \sum_{s=n_k}^{m(n_k, T_k) \wedge (\tau_{\sigma_{n_k+1}} - 1)} \gamma_s \tilde{\varepsilon}_{s+1} I_{\|\tilde{X}_s\| \leq K} \right\| = \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s \tilde{\varepsilon}_{s+1} \right\|.$$

Therefore, by Lemma 4.5 we derive

$$\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s \tilde{\varepsilon}_{s+1} \right\| = 0 \quad \forall T_k \in [0, T].$$

So, for (35) it suffices to show

$$\lim_{T \rightarrow 0} \limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s e_{s+1} \right\| = 0 \quad \forall T_k \in [0, T]. \quad (\text{D.1})$$

Similar to (C.20), we can show that there exist positive constants c_3, c_4, c_5 such that for sufficiently large k

$$\|\tilde{X}_{\perp, s+1}\| \leq c_3 \rho^{s+1-n_k} + c_4 \sup_{m \geq n_k} \gamma_m + c_5 T \quad \forall s : n_k \leq s \leq m(n_k, T). \quad (\text{D.2})$$

Since $0 < \rho < 1$, there exists a positive integer m' such that $\rho^{m'} < T$. Then $\sum_{m=n_k}^{n_k+m'} \gamma_m \xrightarrow[k \rightarrow \infty]{} 0$ by $\lim_{k \rightarrow \infty} \gamma_k = 0$. Thus,

$n_k + m' < m(n_k, T)$ for sufficiently large k . Then by (D.2), we know that for sufficiently large k

$$\|\tilde{X}_{\perp, s+1}\| \leq o(1) + (c_3 + c_5)T \quad \forall s : n_k + m' \leq s \leq m(n_k, T), \quad (\text{D.3})$$

where $o(1) \rightarrow 0$ as $k \rightarrow \infty$.

Since $\tilde{x}_{n_k} \xrightarrow[k \rightarrow \infty]{} \bar{x}$, from (32) and (D.3) it follows that for sufficiently large k and $\forall s : n_k + m' \leq s \leq m(n_k, T)$

$$\begin{aligned} \|\bar{x}_s - \bar{x}\| &\leq \|\bar{x}_s - \tilde{x}_{n_k}\| + \|\tilde{x}_{n_k} - \bar{x}\| = o(1) + \delta(T), \\ \|\tilde{x}_{i,s} - \bar{x}\| &\leq \|\tilde{x}_{i,s} - \bar{x}_s\| + \|\bar{x}_s - \bar{x}\| = o(1) + \delta(T), \end{aligned}$$

where $\delta(T) \rightarrow 0$ as $T \rightarrow 0$. By continuity of $f_i(\cdot)$ we derive $\|f_i(\tilde{x}_{i,s}) - f_i(\bar{x}_s)\| \leq \|f_i(\tilde{x}_{i,s}) - f_i(\bar{x})\| + \|f_i(\bar{x}_s) - f_i(\bar{x})\| = o(1) + \delta(T)$. Consequently,

$$\|e_{i,s+1}\| = \|f_i(\tilde{x}_{i,s}) - f_i(\bar{x}_s)\| / N = o(1) + \delta(T).$$

Then for sufficiently large k

$$\|e_{s+1}\| = o(1) + \delta(T) \quad \forall s : n_k + m' \leq s \leq m(n_k, T). \quad (\text{D.4})$$

By the boundedness of $\{\tilde{X}_s : n_k \leq s \leq m(n_k, T)\}$ and the continuity of $f_i(\cdot)$, we know that there exists a constant $c_e > 0$ such that $\|e_{s+1}\| \leq c_e$. Then from (D.4) we derive

$$\begin{aligned} \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s e_{s+1} \right\| &\leq \sum_{s=n_k}^{n_k+m'} \gamma_s c_e + (o(1) + \delta(T)) \sum_{s=n_k+m'+1}^{m(n_k, T_k)} \gamma_s \\ &\leq c_e m' \sup_{s \geq n_k} \gamma_s + T(\delta(T) + o(1)) \quad \forall T_k \in [0, T] \end{aligned}$$

for sufficiently large k . Then by $\lim_{k \rightarrow \infty} \gamma_k = 0$, we derive

$$\limsup_{k \rightarrow \infty} \frac{1}{T} \left\| \sum_{s=n_k}^{m(n_k, T_k)} \gamma_s e_{s+1} \right\| = \delta(T) \quad \forall T_k \in [0, T].$$

Letting $T \rightarrow 0$, we derive (D.1), and hence (35) holds.

The proof of Lemma 4.7 is completed. \blacksquare

APPENDIX E PROOF OF LEMMA 4.9

i) Assume $\lim_{k \rightarrow \infty} \sigma_k = \infty$. Then there exists a positive integer n_k such that for any $k \geq 1$, $\sigma_{n_k} = k$ and $\sigma_{n_k-1} = k-1$. Hence $n_k - 1 \in [\tau_{k-1}, \tau_k)$ by $\sigma_{n_k} = k$. Thus, from (A.4) it follows that $\tilde{x}_{i, n_k} = x^* \forall i \in \mathcal{V}$. Consequently, $\tilde{X}_{n_k} = (\mathbf{1} \otimes I_l) x^*$ and hence $\{\tilde{X}_{n_k}\}$ is a convergent subsequence with $\bar{x}_{n_k} = x^*$.

Since $\{M_k\}$ is a sequence of positive numbers increasingly diverging to infinity, there exists a positive integer k_0 such that

$$M_k \geq 2\sqrt{N}c_0 + 2 + M'_1 \quad \forall k \geq k_0, \quad (\text{E.1})$$

where c_0 is given in A2 c) and

$$M'_1 = 2 + (2\sqrt{N}c_0 + 2)(c\rho + 2). \quad (\text{E.2})$$

In what follows, we show that under the converse assumption $\{\bar{x}_{n_k}\}$ starting from x^* crosses the sphere with $\|x\| = c_0$ infinitely many times. Define

$$m_k \triangleq \inf\{s > n_k : \|\tilde{X}_s\| \geq 2\sqrt{N}c_0 + 2 + M'_1\}, \quad (\text{E.3})$$

$$l_k \triangleq \sup\{s < m_k : \|\tilde{X}_s\| \leq 2\sqrt{N}c_0 + 2\}. \quad (\text{E.4})$$

Noticing $\|\tilde{X}_{n_k}\| = \sqrt{N}\|x^*\|$ and $\|x^*\| < c_0$, we derive $\|\tilde{X}_{n_k}\| < \sqrt{N}c_0$. Hence from (E.3) (E.4) it is seen that $n_k < l_k < m_k$. By the definition of l_k , we know that $\{\tilde{X}_{l_k}\}$ is bounded. Then there exists a convergent subsequence, denoted still by $\{\tilde{X}_{l_k}\}$. By \bar{X} denoting the limiting point of \tilde{X}_{l_k} , from $\|\tilde{X}_{l_k}\| \leq 2\sqrt{N}c_0 + 2$ it follows that $\|\bar{X}\| \leq 2\sqrt{N}c_0 + 2$.

By Lemma 4.6, there exist constants $M'_0 > 0$ defined by (C.3) with $C = 2\sqrt{N}c_0 + 2$, $c_1 > 0$ and $c_2 > 0$ defined by (C.5), $0 < T < 1$ with $c_1 T \leq 1$ such that

$$\|\tilde{X}_{m+1} - \tilde{X}_{l_k}\| \leq c_1 T + M'_0 \quad \forall m : l_k \leq m \leq m(l_k, T)$$

for sufficiently large $k \geq k_0$. Then for sufficiently large $k \geq k_0$

$$\begin{aligned} \|\tilde{X}_{m+1}\| &\leq \|\tilde{X}_{l_k}\| + c_1 T + M'_0 \\ &\leq 2\sqrt{N}c_0 + 2 + 1 + 1 + (2\sqrt{N}c_0 + 2)(c\rho + 2) \\ &= 2\sqrt{N}c_0 + 2 + M'_1 \quad \forall m : l_k \leq m \leq m(l_k, T). \end{aligned} \quad (\text{E.5})$$

Then $m(l_k, T) \leq n_{k+1}$ for sufficiently large $k \geq k_0$ by (E.1).

From (E.5) by the definition of m_k defined in (E.3), we conclude $m(l_k, T) + 1 \leq m_k$ for sufficiently large $k \geq k_0$. Then by (E.3) (E.4), we know that for sufficiently large $k \geq k_0$

$$2\sqrt{N}c_0 + 2 < \|\tilde{X}_{m+1}\| \leq 2\sqrt{N}c_0 + 2 + M'_1 \quad \forall m : l_k \leq m \leq m(l_k, T). \quad (\text{E.6})$$

Since $0 < \rho < 1$, there exists a positive integer m_0 such that $4c\rho^{m_0} < 1$. Then $\sum_{m=l_k}^{l_k+m_0} \gamma_m \rightarrow 0$ by $\gamma_k \rightarrow 0$, and hence $l_k + m_0 < m(l_k, T) < n_{k+1}$ for sufficiently large $k \geq k_0$. So, from (E.6) it is seen that for sufficiently large $k \geq k_0$

$$\|\tilde{X}_{l_k+m_0}\| > 2\sqrt{N}c_0 + 2. \quad (\text{E.7})$$

Noticing that $\{\tilde{X}_{m+1} : l_k \leq m \leq m(l_k, T)\}$ are bounded, similarly to (C.20) we know that for sufficiently large $k \geq k_0$

$$\begin{aligned} \|\tilde{X}_{\perp, m+1}\| &\leq (2\sqrt{N}c_0 + 2)c\rho^{m+1-l_k} + \frac{cH_1}{1-\rho} \sup_{m \geq l_k} \gamma_m \\ &+ 2T + \frac{c(\rho+1)}{1-\rho} T \quad \forall m : l_k \leq m \leq m(l_k, T). \end{aligned}$$

From here, by $c_1 T < 1$ and $\gamma_k \xrightarrow[k \rightarrow \infty]{} 0$ it follows that

$$\begin{aligned} \|\tilde{X}_{\perp, l_k+m_0}\| &\leq 2c\rho^{m_0}(\sqrt{N}c_0 + 1) + \frac{1}{2} + c_1 T \\ &\leq \frac{1}{2}(\sqrt{N}c_0 + 1) + \frac{1}{2} + 1 = \frac{\sqrt{N}c_0}{2} + 2 \end{aligned} \quad (\text{E.8})$$

for sufficiently large $k \geq k_0$. By noticing $(\mathbf{1} \otimes I_l)\bar{x}_{l_k+m_0} = \tilde{X}_{l_k+m_0} - \tilde{X}_{\perp, l_k+m_0}$, from (E.7) and (E.8) we conclude that

$$\begin{aligned} \sqrt{N} \|\bar{x}_{l_k+m_0}\| &= \|\tilde{X}_{l_k+m_0} - \tilde{X}_{\perp, l_k+m_0}\| \\ &\geq \|\tilde{X}_{l_k+m_0}\| - \|\tilde{X}_{\perp, l_k+m_0}\| > \frac{3}{2}\sqrt{N}c_0. \end{aligned} \quad (\text{E.9})$$

Therefore, $\|\bar{x}_{l_k+m_0}\| > c_0$.

Thus, we have shown that for sufficiently large $k \geq k_0$, starting from x^* , $\{\bar{x}_{n_k}\}$ crosses the sphere with $\|x\| = c_0$ before the time n_{k+1} . So, $\{\bar{x}_{n_k}\}$ starting from x^* crosses the sphere with $\|x\| = c_0$ infinitely many times.

ii) Since $v(J)$ is nowhere dense, there exists a nonempty interval $[\delta_1, \delta_2] \in (v(x^*), \inf_{\|x\|=c_0} v(x))$ with $d([\delta_1, \delta_2], v(J)) > 0$. Assume the converse $\lim_{k \rightarrow \infty} \sigma_k = \infty$.

By i) $\{v(\bar{x}_k)\}$ crosses the interval $[\delta_1, \delta_2]$ infinitely many times while $\{\tilde{X}_{n_k}\}$ converges. This contradicts Lemma 4.8. Therefore, the inverse assumption $\lim_{k \rightarrow \infty} \sigma_k = \infty$ is impossible, and hence (45) holds. The proof is completed. ■

REFERENCES

- [1] W. Ren, and R. W. Beard, "Consensus seeking in multiagent systems under dynamically changing interaction topologies," *IEEE Trans. Autom. Control*, vol. 50, no. 5, pp. 655–661, 2005.
- [2] R. Olfati-Saber, and R. M. Murray, "Consensus problems in networks of agents with switching topology and time-delays," *IEEE Trans. Autom. Control*, vol. 49, no. 9, pp. 1520–1533, 2004.
- [3] Q. Zhang and J. F. Zhang, "Distributed parameter estimation over unreliable networks with Markovian switching topologies," *IEEE Trans. Autom. Control*, vol. 57, no. 10, pp. 2545–2560, 2012.
- [4] C. G. Lopes, and A. H. Sayed, "Diffusion least-mean square over adaptive networks: formulation and performance analysis," *IEEE Trans. Signal Processing*, vol. 56, no. 7, pp. 3122–3136, 2008.
- [5] U. A. Khan, S. Kar, and José M. F. Moura, "Distributed Sensor Localization in Random Environments Using Minimal Number of Anchor Nodes," *IEEE Trans. Signal Processing*, vol. 57, no. 5, pp. 2000–2016, 2009.
- [6] A. Nedić, and A. Ozdaglar, "Distributed subgradient methods for multi-agent optimization," *IEEE Trans. Autom. Control*, vol. 54, no. 1, pp. 48–61, 2009.
- [7] A. Nedić, A. Ozdaglar, and P. A. Parrilo, "Constrained consensus and optimization in multi-agent networks," *IEEE Trans. Autom. Control*, vol. 55, no. 4, pp. 922–938, 2010.
- [8] J. Lei, H. T. Fang, and H. F. Chen, "Primal–dual algorithm for distributed constrained optimization", *Systems & Control Letters*, vol. 96, pp. 110–117, 2016.
- [9] H. Robbins, and S. Monro, "A stochastic approximation method," *Annu. Math. Statist.*, vol. 22, no. 3, pp. 400–405, 1951.
- [10] J. lei and H. F. Chen, "Distributed randomized PageRank algorithm based on stochastic approximation", *IEEE Transactions on Automatic Control*, vol. 60, no. 6, pp. 1641–1646, 2015.
- [11] J. Kiefer, and J. Wolfowitz, "Stochastic estimation of the maximum of a regression function," *Ann. Math. Statist.*, vol. 23, no.3, pp. 462–466, 1952.
- [12] H. J. Kushner, and G. Yin, "Asymptotic properties of distributed and communicating stochastic approximation algorithms," *SIAM Journal on Control and Optimization*, vol. 25, no. 5, pp. 1266–1290, 1987.
- [13] P. Bianchi, G. Fort, and W. Hachem, "Performance of a distributed stochastic approximation algorithm," *IEEE Trans. Inf. Theory*, vol. 59, no. 11, pp. 7405–7418, 2013.
- [14] G. Morral, P. Bianchi, G. Fort, J. Jakubowicz, "Distributed stochastic approximation: The price of non-double stochasticity," *Proc. Asilomar* 2012.
- [15] V. S. Bokar, and V. V. Phansalkar, "Managing interprocessor delays in distributed recursive algorithms," *Sadhana*, vol. 19, pp. 995–1003, 1994.
- [16] V. S. Bokar, "Asynchronous stochastic approximation," *SIAM Journal on Control and Optimization*, vol. 36, no. 3, pp. 840–851, 1998.
- [17] J. Lei, H. T. Fang, and H. F. Chen, "Asymptotic Properties of Primal-Dual Algorithm for Distributed Stochastic Optimization over Random Networks with Imperfect Communications", *SIAM Journal on Control and Optimization*, vol. 56, no. 3, pp. 2159–2188, 2018.
- [18] P. Bianchi, J. Jakubowicz, "Convergence of a multi-agent projected stochastic gradient algorithm for non-convex optimization," *IEEE Trans. Autom. Control*, vol. 58, no. 2, pp. 391–405, 2013.
- [19] R. Liu and H. F. Chen, "Distributed and recursive blind channel identification to sensor networks", *Control Theory and Technology*, vol. 15, no. 4, pp. 274–287, 2017.
- [20] W. Feng and H.F. Chen, "Output Consensus of Networked Hammerstein and Wiener Systems", *SIAM Journal on Control and Optimization*, vol. 57, no. 2, pp. 1230–1254, 2019.
- [21] E. Oja, and J. Karhunen, "On stochastic approximation of the eigenvectors and eigenvalues of the expectation of a random matrix," *Journal of Mathematical Analysis and Applications*, vol. 106, pp. 69–84, 1985.
- [22] Z. J. Bai, R. H. Chan, and F. T. Luk, "Principal component analysis for distributed data sets with updating", *International Workshop on Advanced Parallel Processing Technologies*, pp. 471–483, Springer Berlin Heidelberg, 2005.
- [23] S. V. Macua, P. Belanovic, S. Zazo, "Consensus-based distributed principal component analysis in wireless sensor networks," *International Workshop on SPAWC*, pp. 1–5, 2010.
- [24] Y. A. Borgne, S. Raybaud, and G. Bontempi, "Distributed principal component analysis for wireless sensor networks," *Sensors*, vol. 8, no. 8, pp. 4821–4850, 2008.
- [25] Z. Meng, A. Wiesel, and A.O. Hero, "Distributed principal component analysis on networks via directed graphical models," *IEEE International Conference on ICASSP*, pp. 2877–2880, 2012.
- [26] S. B. Korada, A. Montanari, and S. Oh, "Gossip PCA", *In Proceedings of the ACM SIGMETRICS joint international conference on Measurement and modeling of computer systems*, pp. 209–220, 2011.
- [27] G. Morral, P. Bianchi, and J. Jakubowicz, "Asynchronous distributed principal component analysis using stochastic approximation", *IEEE 51st Annual Conference on Decision and Control*, pp. 1398–1403.
- [28] H. F. Chen, T. Ducan, and B. P. Ducan, "A Kiefer-Wolfowitz algorithm with randomized differences," *IEEE Trans. Autom. Control*, vol. 44, no. 3, pp. 442–453, 1999.
- [29] J. C. Spall, "Multivariate stochastic approximation using a simultaneous perturbation gradient optimization," *IEEE Trans. Automatic Control*, vol. 45, pp. 1839–1853, 1992.
- [30] L. M. Rios, and N. V. Sahinidis, "Derivative-free optimization: A review of algorithms and comparison of software implementations," *Journal of Global Optimization*, vol. 56, no.3, pp. 1247–1293, 2013.
- [31] D. Yuan and D. W. C. Ho, "Randomized gradient-free method for multi-agent optimization over time-varying networks," *IEEE Trans. Neural Netw. Learn. Syst.*, vol. 26, no. 6, pp. 1342–1347, 2015.
- [32] J. Li, C. Wu, Z. Wu, and Q. Long, "Gradient-free method for nonsmooth distributed optimization," *Journal of Global Optimization*, vol. 61, no. 2, pp. 325–340, 2015.
- [33] A. Benveniste, M. Metivier, and P. Priouret, *Adaptive Algorithms and Stochastic Approximation*, Springer-Verlag, New York, 1990.
- [34] V. S. Bokar, *Stochastic approximation: a dynamical systems viewpoint*. Tata Institute of Fundamental Research, Mumbai, 2008.
- [35] H. J. Kushner, and G. Yin, *Stochastic Approximation Algorithms and Applications*. Springer-Verlag, New York, 2003.
- [36] H. F. Chen, *Stochastic Approximation and Its Applications*. Dordrecht, The Netherlands: Kluwer, 2002.
- [37] H. F. Chen and W. X. Zhao, *Recursive Identification and Parameter Estimation*. CRC Press, 2014.
- [38] B. Delyon, "General results on the convergence of stochastic algorithms," *IEEE Transactions on Automatic Control*, vol. 41, no. 9, pp. 1245–1255, 1996.
- [39] J. Lei, and H. F. Chen, "Distributed Stochastic Approximation Algorithm With Expanding Truncations: Algorithm and Applications", *arXiv:1410.7180v4*.
- [40] S. C. Yuan and H. Teicher, *Probability Theory*. Springer, 1997.
- [41] H.F. Chen, H.T. Fang, and L. L. Zhang, "Recursive estimation for ordebblue eigenvectors of symmetric matrix with observation noise," *Journal of Mathematical Analysis and Applications*, no. 382, vol. 2, pp.822–842, 2011.
- [42] H. T. Fang, and H. F. Chen, "Stability and instability of limit point of the stochastic approximation algorithms", *IEEE Trans. Automa. Control*, vol. 45, no. 3, pp. 413–420, 2000.
- [43] B. Gharesifard and J. Cortés, "When does a digraph admit a doubly stochastic adjacency matrix?," *in Proc. Amer. Control Conf.*, Baltimore, MD, 2010, pp. 2440–2445.