THE ZAK TRANSFORM AND THE STRUCTURE OF SPACES INVARIANT BY THE ACTION OF AN LCA GROUP

D. BARBIERI, E. HERNÁNDEZ, V. PATERNOSTRO

ABSTRACT. We study closed subspaces of $L^2(\mathcal{X})$, where (\mathcal{X}, μ) is a σ -finite measure space, that are invariant under the unitary representation associated to a measurable action of a discrete countable LCA group Γ on \mathcal{X} . We provide a complete description for these spaces in terms of range functions and a suitable generalized Zak transform. As an application of our main result, we prove a characterization of frames and Riesz sequences in $L^2(\mathcal{X})$ generated by the action of the unitary representation under consideration on a countable set of functions in $L^2(\mathcal{X})$. Finally, closed subspaces of $L^2(G)$, for G being an LCA group, that are invariant under translations by elements on a closed subgroup Γ of G are studied and characterized. The results we obtain for this case are applicable to cases where those already proven in [5, 7] are not.

1. INTRODUCTION

Closed spaces that are invariant under a certain kind of unitary operators are of particular importance in several areas of Harmonic Analysis such as wavelets, spline systems, Gabor systems and approximation theory [1, 19, 24, 30, 33]. As a particular case of these spaces we have the so-called *shift-invariant spaces* which are closed subspaces of $L^2(\mathbb{R}^n)$ invariant under all integer translations. The structure of shift-invariant spaces has been intensively studied in [4, 11, 12, 36, 37]. Recently, shift-invariant spaces have been considered in the context of locally compact abelian (LCA) groups. That is, when G is an LCA group, a shift-invariant space is a closed subspace of $L^2(G)$ that is invariant under translations by elements of a closed subspace of G. The case when Γ is a discrete countable subgroup of G with G/Γ compact has been treated in [7, 31], and extended to the case of Γ closed and G/Γ compact in [5].

The aim of this paper is twofold. First of all, we will give a comprehensive description of the structure of closed spaces that are invariant by a class of operators that generalize translations, namely unitary representations of discrete groups arising from measurable group actions on measure spaces. Secondly, we will see how a similar approach allows us to treat translations by elements of a closed subgroup without any other assumption on it. Both cases are treated with a similar approach by using a generalized Zak transform, properly adapted to each case.

More precisely, in the first part of the paper we will consider, when (\mathcal{X}, μ) is a σ -finite measure space, closed subspaces of $L^2(\mathcal{X})$ invariant under the unitary representation associated to a measurable action of a discrete countable LCA group Γ on \mathcal{X} . A measurable action of Γ on \mathcal{X} is a measurable map $\sigma : \Gamma \times \mathcal{X} \to \mathcal{X}$ that satisfies $\sigma_{\gamma}(\sigma_{\gamma'}(x)) = \sigma_{\gamma+\gamma'}(x)$ for all $\gamma, \gamma' \in \Gamma$, all $x \in \mathcal{X}$ and $\sigma_e(x) = x$ for all $x \in \mathcal{X}$, where $\sigma_{\gamma} : \mathcal{X} \to \mathcal{X}$ is given by $\sigma_{\gamma}(x) = \sigma(\gamma, x)$. The action is said to be quasi- Γ -invariant if there exists a measurable function $J_{\sigma} : \Gamma \times \mathcal{X} \to \mathbb{R}^+$ such that

²⁰¹⁰ Mathematics Subject Classification. Primary 43A65, 43A70; Secondary 42C15, 42C40.

Key words and phrases. shift-invariant spaces, groups actions, range functions, Riesz bases, frames, Zak transform, LCA groups.

 $d\mu(\sigma_{\gamma}(x)) = J_{\sigma}(\gamma, x)d\mu(x)$. To each quasi- Γ -invariant action σ we can associate a unitary representation Π_{σ} of Γ on $L^{2}(\mathcal{X})$ given by

(1)
$$\Pi_{\sigma}(\gamma)f(x) = J_{\sigma}(-\gamma, x)^{\frac{1}{2}}f(\sigma_{-\gamma}(x)).$$

It is with respect to this unitary representation that we consider invariant subspaces of $L^2(\mathcal{X})$. We say that a closed subspace V of $L^2(\mathcal{X})$ is (Γ, σ) -invariant if whenever $f \in V$ we have that $\Pi_{\sigma}(\gamma)f \in V, \forall \gamma \in \Gamma$.

This setting includes the case of shift-invariant subspaces of $L^2(\mathbb{R}^n)$ treated in [4, 11, 12, 36, 37] as well as those considered in the context of LCA groups [7, 31]. But moreover, it includes other classical operators from Analysis such us the dilation operator used in wavelet theory, which arises from the dilation action $\sigma : \mathbb{Z} \times \mathbb{R}^n \to \mathbb{R}^n$ given by $\sigma(j, x) = 2^j x, j \in \mathbb{Z}, x \in \mathbb{R}^n$. More examples are given in Section 2 (see Example 2.1).

The main issue that arises when trying to address this problem can be understood by noting that the previous papers where the structure of shift-invariant spaces is established, such as [4, 7], use heavily the Fourier transform either in \mathbb{R}^n or in the LCA group G, as well as in the subgroup of translations Γ . But in our situation, $L^2(\mathcal{X})$ with (\mathcal{X}, μ) a measure space, such a tool is not available. The novelty of our approach is that we work with a version of the Zak transform adapted to our setting.

The classical Zak transform is an isometric isomorphism from $L^2(\mathbb{R}^n)$ onto $L^2(\mathbb{T}^{2n})$ given, for $f \in L^2(\mathbb{R}^n)$, by

$$Zf(x,\xi) = \sum_{k \in \mathbb{Z}^n} f(x+k)e^{-2\pi i k\xi}, \qquad (x,\xi) \in \mathbb{T}^{2n}.$$

The history of the Zak transform is well described in Chapter 8 of [19]: The Zak transform was first introduced and used by Gelfand [16] for a problem in differential equations. Weil [41] defined this transform on arbitrary locally compact abelian groups with respect to arbitrary closed subgroups. ... Subsequently, the Zak transform was rediscovered several times, notably by Zak [44, 45] for a problem in solid state physics and Brezin [6] for differential equations. In representation theory and in abstract harmonic analysis Z is often called the Weil-Brezin map, but in applied mathematics and signal analysis it has become customary to refer to Z as the Zak transform. ... The popularity of the Zak transform in engineering seems largely due to Janssen's influential survey article [29].

Properties of the Zak transform, as well as applications in pure and applied mathematics can be found in [28, 29], [19, Chapter 8], [17, 18, 22, 23, 42].

In our setting, the generalized Zak transform is defined in the following way: for $\psi \in L^2(\mathcal{X}), \alpha \in \widehat{\Gamma}$ and $x \in \mathcal{X}$,

$$Z_{\sigma}[\psi](\alpha)(x) := \sum_{\gamma \in \Gamma} \left[(\Pi_{\sigma}(\gamma)\psi)(x) \right] X_{-\gamma}(\alpha),$$

where X_{γ} denotes the character on $\widehat{\Gamma}$ associated to $\gamma \in \Gamma$.

The importance of such generalized Zak transform in the study of invariant subspaces of $L^2(\mathcal{X})$ has been pointed out in [22] where it is used to prove results concerning properties of sequences generated by a single function $\psi \in L^2(\mathcal{X})$ under the representation Π_{σ} given by (1).

Our main result characterizes (Γ, σ) -invariant subspaces of $L^2(\mathcal{X})$, for σ -finite measure space \mathcal{X} , in terms of range functions and the generalized Zak transform. As a consequence, we are able to characterize frames and Riesz sequences in $L^2(\mathcal{X})$ generated by the action of the unitary representation given by (1) on a countable (meaning countable or finite) family of elements in $L^2(\mathcal{X})$. As easy corollaries of these characterizations we deduce the results obtained in [22] concerning characterizations of cyclic sequences generated by a single function in terms of the so-called bracket map. In addition, we also show that every (Γ, σ) -invariant subspace of $L^2(\mathcal{X})$ can be decomposed into an orthogonal sum of (Γ, σ) -invariant subspaces, each of them generated by a single function whose Π_{σ} -orbit is a Parseval frame for the space it spans.

In the second part of the paper, we will then focus on subspaces of $L^2(G)$, with G an LCA group, that are invariant under translations by elements of a closed subgroup Γ of G, and still address this problem in terms of a suitable Zak transform. The subgroup Γ is allowed to be non-discrete and hence the situation does not fit directly in the setting previously considered in this paper. For this reason, we need to work with a non-discrete version of the Zak transform and deal with some technical issues. As a result, also in this case we can characterize the structure of these spaces in terms of range functions but now using the non-discrete version of the Zak transform. Since we ask Γ to be just closed – not discrete nor co-compact – our results include more cases than those covered by [5, 7, 31].

The paper is organized as follows. Section 2 is devoted to give precise definitions of the basic objects involved in the study, while Section 3 contains the main properties of the Zak transform for discrete LCA groups. The main result, where we characterize (Γ, σ) -invariant subspaces of $L^2(\mathcal{X})$ in terms of range functions and the generalized Zak transform is given in Section 4. The proof of it uses the characterization of multiplicative invariant subspaces developed in [5], which is summarized, for the readers convenience, in Subsection 4.1. Applications of this result are given in Section 5 where we characterize frames and Riesz sequences in $L^2(\mathcal{X})$ generated by the action of Π_{σ} on a countable set of $L^2(\mathcal{X})$. Finally, in Section 6 we consider the case of an LCA second countable group G and provide a characterization of subspaces of $L^2(G)$ that are invariant under translations by elements of a closed subgroup Γ of G.

Note: Once this manuscript was submitted for publication we learned of the paper by J. Iverson [27] where similar results are proved independently. In [27], the results we prove in Section 6 for closed subspaces of $L^2(G)$ invariant under translations of by a closed subgroup Γ of an LCA group G, are extended using similar techniques, to the case of closed abelian subgroups Γ of a locally compact group G. On the other hand, in Sections 4 and 5 we obtain results for closed subspaces of a general $L^2(\mathcal{X})$ space invariant under the action of a discrete LCA group Γ , which are not included in [27]. Also we learned of the paper by S. Saliani [39] on linear independence in spaces invariant under the action of LCA groups, which appeared almost in the same days and contains ideas on similar topics.

Acknowledgement: D. Barbieri was supported by a Marie Curie Intra European Fellowship (Prop. N. 626055) within the 7th European Community Framework Programme. D. Barbieri and E. Hernández were supported by Grant MTM2013-40945-P (Ministerio de Economía y Competitividad, Spain). E. Hernández was supported by Grant MTM2010-16518 (Ministerio de Economía y Competitividad, Spain) and V. Paternostro was supported by a fellowship for post-doctoral researchers from the Alexander von Humboldt Foundation.

The authors want to thank the referee for her/his comments which helped us to improve the manuscript.

2. Preliminaries and notation

2.1. **Basic on LCA groups.** Let Γ be a locally compact abelian (LCA) group with operation written additively. By $\widehat{\Gamma}$ we denote the dual group of Γ , that is, the set of continuous characters on Γ . For $\gamma \in \Gamma$ and $\alpha \in \widehat{\Gamma}$ we use the notation (γ, α) for the complex value that α takes at γ . For each $\gamma \in \Gamma$, the corresponding character on $\widehat{\Gamma}$ is denoted by X_{γ} , where $X_{\gamma} : \widehat{\Gamma} \to \mathbb{C}$ and $X_{\gamma}(\alpha) = (\gamma, \alpha)$.

We denote the Haar measures on Γ and $\widehat{\Gamma}$ by m_{Γ} and $m_{\widehat{\Gamma}}$ respectively. The Fourier transform of a Haar integrable function f on Γ , is the function \widehat{f} on $\widehat{\Gamma}$ defined by $\widehat{f}(\alpha) = \int_{\Gamma} f(\gamma)(-\gamma, \alpha) dm_{\Gamma}(\gamma)$. We fix the Haar measures m_{Γ} and $m_{\widehat{\Gamma}}$ such that the inversion formula holds (for details see [38, Section 1.5]). Therefore, the Fourier transform on $L^1(\Gamma) \cap L^2(\Gamma)$ extends to a unitary operator from $L^2(\Gamma)$ to $L^2(\widehat{\Gamma})$ that we denote equally either by \wedge or by \mathcal{F}_{Γ} .

The Fourier transform satisfies the following well known property: for any $\gamma \in \Gamma$,

(2)
$$\widehat{f(\cdot - \gamma)} = (-\gamma, \cdot)\widehat{f(\cdot)}$$

As usual, when Γ is discrete, we choose m_{Γ} to be counting measure. Then, the inversion formula holds with $m_{\widehat{\Gamma}}$ normalized such that $m_{\widehat{\Gamma}}(\widehat{\Gamma}) = 1$. With this normalization of the Haar measure and supposing that Γ is countable, $\{X_{\gamma}\}_{\gamma \in \Gamma}$ turns out to be an orthonormal basis for $L^2(\widehat{\Gamma})$.

2.2. LCA groups acting on σ -finite measure spaces. We now state the precise definition of the actions we shall work with. This type of actions, called *quasi*- Γ -*invariant* actions, were previously considered in [22] within the abelian setting and in [2] for countable discrete groups that can be non-abelian. Fix Γ a countable discrete LCA group. Let (\mathcal{X}, μ) be a σ -finite measure space and $\sigma : \Gamma \times \mathcal{X} \to \mathcal{X}$ an action satisfying the following conditions:

- (i) for each $\gamma \in \Gamma$ the map $\sigma_{\gamma} : \mathcal{X} \to \mathcal{X}$ given by $\sigma_{\gamma}(x) := \sigma(\gamma, x)$ is measurable;
- (ii) $\sigma_{\gamma}(\sigma_{\gamma'}(x)) = \sigma_{\gamma+\gamma'}(x)$, for all $\gamma, \gamma' \in \Gamma$ and for all $x \in \mathcal{X}$;
- (iii) $\sigma_e(x) = x$ for all $x \in \mathcal{X}$, where e is the identity of Γ .

As Γ is discrete and countable, $\sigma : \Gamma \times \mathcal{X} \to \mathcal{X}$ turns out to be a measurable action, since if $E \subseteq \mathcal{X}$ is measurable set, then $\sigma^{-1}(E) = \bigcup_{\gamma \in \Gamma} [\{\gamma\} \times \sigma_{\gamma}^{-1}(E)]$ is measurable in $\Gamma \times \mathcal{X}$.

The action σ is said to be *quasi*- Γ -*invariant* if there exists a measurable positive function $J_{\sigma} : \Gamma \times \mathcal{X} \to \mathbb{R}^+$, called *Jacobian of* σ , such that $d\mu(\sigma_{\gamma}(x)) = J_{\sigma}(\gamma, x)d\mu(x)$. As a consequence of the properties of σ and the definition of the Jacobian we have:

(3)
$$J_{\sigma}(\gamma_1 + \gamma_2, x) = J_{\sigma}(\gamma_1, \sigma_{\gamma_2}(x)) J_{\sigma}(\gamma_2, x), \quad \forall \gamma_1, \gamma_2 \in \Gamma, \ x \in \mathcal{X}.$$

To each quasi- Γ -invariant action σ we can associate a unitary representation Π_{σ} of Γ on $L^2(\mathcal{X})$ given by

$$\Pi_{\sigma}(\gamma)\psi(x) = J_{\sigma}(-\gamma, x)^{\frac{1}{2}}\psi(\sigma_{-\gamma}(x)).$$

From (3) it follows that Π_{σ} is a unitary operator. To prove that Π_{σ} is a representation see [2, Section 4.2].

The action σ has the *tiling property* if there exists a measurable set $C \subseteq \mathcal{X}$ such that $\mu(\sigma_{\gamma}(C) \cap \sigma_{\gamma'}(C)) = 0$ if $\gamma \neq \gamma'$ and $\mu(\mathcal{X} \setminus \bigcup_{\gamma \in \Gamma} \sigma_{\gamma}(C)) = 0$.

We provide here relevant examples of concrete quasi- Γ -invariant actions satisfying the tiling property. In particular, they include the usual operators used in Analysis such us translations, dilations and those arising from the action of the shear matrix. Note that, the tiling set C necessarily has positive measure but, as the examples show, it can have either finite or infinite measure.

Example 2.1.

• Let us consider the group \mathbb{Z} acting on \mathbb{R}^2 by

$$\sigma_1(k,(x,y)) = \begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = (x,kx+y).$$

The action σ_1 satisfies conditions i-iii and $J_1(k, (x, y)) = 1$. Moreover, σ_1 has the tiling property with tiling set $C = \{(x, y) \in \mathbb{R}^2 : 0 \leq y \leq x\} \cup \{(x, y) \in \mathbb{R}^2 : x \leq y \leq 0\}$. The associated unitary representation on $L^2(\mathbb{R}^2)$ is $\Pi_{\sigma_1}(k)\psi(x, y) = \psi(x, -kx + y)$.

• A different action of \mathbb{Z} on \mathbb{R}^2 is given by

$$\sigma_2(j,(x,y)) = \begin{pmatrix} 2^j & 0\\ 0 & 2^{j/2} \end{pmatrix} \begin{pmatrix} x\\ y \end{pmatrix} = (2^j x, 2^{j/2} y).$$

Also in this case σ_2 satisfies conditions i-iii but we have that $J_2(j, (x, y)) = 2^{3j/2}$. A tiling set for σ_2 is $C = C_1 \setminus C_2$ where $C_1 = [-1, 1]^2$ and $C_2 = [-1/2, 1/2] \times [-1/\sqrt{2}, 1/\sqrt{2}]$. The associated unitary representation on $L^2(\mathbb{R}^2)$ is $\Pi_{\sigma_2}(j)\psi(x, y) = 2^{-3j/4}\psi(2^{-j}x, 2^{-j/2}y)$.

• The group \mathbb{Z}^n acts on \mathbb{R}^n by translations $\sigma_3(m, x) = m + x$. For this action $J_3(m, x) = 1$, a tiling set is $C = [-1/2, 1/2]^n$ and the unitary representation associated on $L^2(\mathbb{R}^n)$ to σ_3 is $\prod_{\sigma_3}(m)\psi(x) = \psi(x-m)$.

2.3. (Γ, σ) -invariant spaces on $L^2(\mathcal{X})$. Given σ a quasi- Γ -invariant action of Γ on \mathcal{X} we can define (Γ, σ) -invariant spaces in $L^2(\mathcal{X})$ as spaces that are invariant by the operators $\Pi_{\sigma}(\gamma), \gamma \in \Gamma$. More precisely, we have:

Definition 2.2. A closed subspace V of $L^2(\mathcal{X})$ is said to be (Γ, σ) -invariant if

 $f \in V \Longrightarrow \Pi_{\sigma}(\gamma) f \in V$, for any $\gamma \in \Gamma$.

For any subset $\mathcal{A} \subseteq L^2(\mathcal{X})$ we define

 $S_{\sigma}^{\Gamma}(\mathcal{A}) = \overline{\operatorname{span}} \{ \Pi_{\sigma}(\gamma)\phi \colon \phi \in \mathcal{A}, \gamma \in \Gamma \} \text{ and } E_{\sigma}^{\Gamma}(\mathcal{A}) = \{ \Pi_{\sigma}(\gamma)\phi \colon \phi \in \mathcal{A}, \gamma \in \Gamma \}.$

We call $S_{\sigma}^{\Gamma}(\mathcal{A})$ the (Γ, σ) -invariant space generated by \mathcal{A} . If $V = S_{\sigma}^{\Gamma}(\mathcal{A})$ for some finite set \mathcal{A} we say that V is a *finitely generated* (Γ, σ) -invariant space, and a *principal or cyclic* (Γ, σ) -invariant space if \mathcal{A} has only a single function.

When $L^2(\mathcal{X})$ is separable each (Γ, σ) -invariant space V is of the form $V = S^{\Gamma}_{\sigma}(\mathcal{A})$ for some countable (meaning finite or countable) set $\mathcal{A} \subseteq L^2(\mathcal{X})$.

Remark 2.3. By [9, Proposition 3.4.5], the space $L^2(\mathcal{X})$ is separable if and only if μ is a σ -finite measure and the σ -algebra on \mathcal{X} is countably generated, meaning that it is generated by countable family of subsets of \mathcal{X} . From now on, we will always assume that $L^2(\mathcal{X})$ is separable, even without mentioning it.

2.4. Vector valued spaces. Let (Ω, ν) be a σ -finite measure space and let \mathcal{H} be a separable Hilbert space. The vector valued space $L^2(\Omega, \mathcal{H})$ is the space of measurable functions $\Phi : \Omega \to \mathcal{H}$ such that $\|\Phi\|^2 = \int_{\Omega} \|\Phi(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) < +\infty$. The inner product in $L^2(\Omega, \mathcal{H})$ is given by $\langle \Phi, \Psi \rangle = \int_{\Omega} \langle \Phi(\omega), \Psi(\omega) \rangle_{\mathcal{H}} d\nu(\omega)$.

A function $\Phi: \Omega \to \mathcal{H}$ is called *simple* if there exist $x_1, \ldots, x_n \in \mathcal{H}$ and measurable subsets E_1, \ldots, E_n of Ω such that $\Phi = \sum_{j=1}^n x_j \chi_{E_j}$, where χ_{E_j} denotes the characteristic function of E_j . When Ω is of finite measure, the set of simple functions is dense in $L^2(\Omega, \mathcal{H})$. Since we could not find a reference for this fact, we provide a proof in the Appendix. More details about the subject can be found in [13].

2.5. Frames and Riesz bases. We briefly recall the definitions of frame and Riesz basis. For a detailed exposition on this subject we refer to [8].

Let \mathcal{H} be a separable Hilbert space, I be a finite or countable index set and $\{f_i\}_{i\in I}$ be a sequence in \mathcal{H} . The sequence $\{f_i\}_{i\in I}$ is said to be a *frame* for \mathcal{H} if there exist $0 < A \leq B < +\infty$ such that

$$A||f||^{2} \leq \sum_{i \in I} |\langle f, f_{i} \rangle|^{2} \leq B||f||^{2}$$

for all $f \in \mathcal{H}$. The constants A and B are called *frame bounds*. When A = B = 1, $\{f_i\}_{i \in I}$ is called *Parseval frame*.

The sequence $\{f_i\}_{i \in I}$ is said to be a *Riesz basis* for \mathcal{H} if it is a complete system in \mathcal{H} and if there exist $0 < A \leq B < +\infty$ such that

$$A\sum_{i\in I} |a_i|^2 \le \|\sum_{i\in I} a_i f_i\|^2 \le B\sum_{i\in I} |a_i|^2$$

for all sequences $\{a_i\}_{i \in I}$ of finite support.

The sequence $\{f_i\}_{i \in I}$ is a *frame (or Riesz) sequence*, if it is a frame (or Riesz basis) for the Hilbert space it spans, namely $\overline{\text{span}}\{f_i : i \in I\}$.

3. A Zak transform associated to quasi- Γ -invariant actions

In [22] the authors introduced a definition of the Zak transform associated to quasi- Γ -invariant action σ of a discrete and countable LCA group Γ . Recently, in [2], an analogous noncommutative Zak transform was introduced for non-abelian groups. In this section we will make use of the definition given in [22] and see how the Zak transform allows us to think of $L^2(\mathcal{X})$ as a vector valued space. The Zak transform combined with range functions will allow us to prove a characterization of (Γ, σ) -invariant spaces similar to those previously given in [4, 5, 7, 10].

Assume σ is a quasi- Γ -invariant action of Γ on \mathcal{X} satisfying the tiling property with tiling set C. Let Π_{σ} be its associated unitary representation of Γ on $L^{2}(\mathcal{X})$. Given $\psi \in L^{2}(\mathcal{X})$ we can define the function $\psi_{\sigma} : \Gamma \times \mathcal{X} \to \mathbb{C}$ as

(4)
$$\psi_{\sigma}(\gamma, x) := \Pi_{\sigma}(\gamma)\psi(x)$$

For a fixed $x \in \mathcal{X}$ the function $\psi_{\sigma}(\cdot, x)$ can be understood as the evolution of ψ along the orbit of $x, \{\sigma_{\gamma}(x) : \gamma \in \Gamma\}$.

Lemma 3.1. For every $\psi \in L^2(\mathcal{X})$, the sequence $\psi_{\sigma}(\cdot, x) = \{\psi_{\sigma}(\gamma, x)\}_{\gamma \in \Gamma}$ belongs to $\ell^2(\Gamma)$ for μ -a.e. $x \in \mathcal{X}$.

Proof. Using the tiling property of the action σ , we obtain

$$\begin{split} \|\psi\|_{L^{2}(\mathcal{X})}^{2} &= \int_{\mathcal{X}} |\psi(x)|^{2} d\mu(x) = \sum_{\gamma \in \Gamma} \int_{\sigma_{\gamma}(C)} |\psi(x)|^{2} d\mu(x) \\ &= \sum_{\gamma \in \Gamma} \int_{C} |\psi(\sigma_{\gamma}(y))|^{2} J_{\sigma}(\gamma, y) d\mu(y) \\ &= \int_{C} \sum_{\gamma \in \Gamma} |\psi_{\sigma}(-\gamma, y)|^{2} d\mu(y) = \int_{C} \|\psi_{\sigma}(\cdot, y)\|_{\ell^{2}(\Gamma)}^{2} d\mu(y). \end{split}$$

Since $\psi \in L^2(\mathcal{X})$, it follows that $\|\psi_{\sigma}(\cdot, y)\|_{\ell^2(\Gamma)}^2 < +\infty$ for μ -a.e. $y \in C$.

Finally, by (3) it follows that for $x \in \mathcal{X}$ written as $x = \sigma_{\gamma_0}(y)$ with $\gamma_0 \in \Gamma$ and $y \in C$

$$\psi_{\sigma}(\gamma, x) = \psi_{\sigma}(\gamma, \sigma_{\gamma_0}(y)) = J_{\sigma}(\gamma_0, y)^{-\frac{1}{2}} \psi_{\sigma}(\gamma - \gamma_0, y).$$

Thus, $\|\psi_{\sigma}(\cdot, x)\|_{\ell^2(\Gamma)}^2 = J_{\sigma}(\gamma_0, y)^{-1} \|\psi_{\sigma}(\cdot, y)\|_{\ell^2(\Gamma)}^2$ and the lemma follows. \Box

Definition 3.2 ([22]). Given any $\psi \in L^2(\mathcal{X})$ define for $\alpha \in \widehat{\Gamma}$ and $x \in \mathcal{X}$ such that $\psi_{\sigma}(\cdot, x) \in \ell^2(\Gamma)$

$$Z_{\sigma}[\psi](\alpha)(x) := \sum_{\gamma \in \Gamma} \left[(\Pi_{\sigma}(\gamma)\psi)(x) \right] (-\gamma, \alpha).$$

We call $Z_{\sigma}[\psi]$ the *(generalized)* Zak transform of ψ .

Note that $Z_{\sigma}[\psi](\alpha)(x)$ is the Fourier transform of the sequence $\psi_{\sigma}(\cdot, x)$ evaluated in α ; i.e. $Z_{\sigma}[\psi](\alpha)(x) := \widehat{\psi_{\sigma}(\cdot, x)}(\alpha)$. By Lemma 3.1, $Z_{\sigma}[\psi](\alpha)(x)$ is defined for μ a.e. $x \in \mathcal{X}$ and every $\alpha \in \widehat{\Gamma}$.

In the next proposition we show that $L^2(\mathcal{X})$ is isometrically isomorphic to the vector valued Hilbert space $L^2(\widehat{\Gamma}, L^2(C))$. For this vector valued space, the norm and inner product are

$$\|\Phi\|^2 = \int_{\widehat{\Gamma}} \|\Phi(\alpha)\|^2_{L^2(C)} dm_{\widehat{\Gamma}}(\alpha) \quad \text{and} \quad \langle \Phi, \Psi \rangle = \int_{\widehat{\Gamma}} \langle \Phi(\alpha), \Psi(\alpha) \rangle_{L^2(C)} dm_{\widehat{\Gamma}}(\alpha),$$

respectively.

Proposition 3.3. The mapping $Z_{\sigma}: L^2(\mathcal{X}) \longrightarrow L^2(\widehat{\Gamma}, L^2(C))$ defined by

$$Z_{\sigma}[\psi](\alpha)(x) = \sum_{\gamma \in \Gamma} \left[(\Pi_{\sigma}(\gamma)\psi)(x) \right] (-\gamma, \alpha)$$

is an isometric isomorphism and satisfies

(5)
$$Z_{\sigma}[\Pi_{\sigma}(\gamma)\psi] = X_{\gamma}Z_{\sigma}[\psi], \quad \forall \ \gamma \in \Gamma, \ \psi \in L^{2}(\mathcal{X}).$$

Proof. For each $\psi \in L^2(\mathcal{X})$, $Z_{\sigma}[\psi]$ is measurable as a function defined on $\widehat{\Gamma} \times C$. Now, proceeding as in Lemma 3.1 and by Plancherel and Fubini's Theorems we have

(6)

$$\begin{aligned} \|\psi\|_{L^{2}(\mathcal{X})}^{2} &= \int_{C} \sum_{\gamma \in \Gamma} |(\Pi_{\sigma}(\gamma)\psi)(x)|^{2} d\mu(x) \\ &= \int_{C} \|\psi_{\sigma}(\cdot,x)\|_{\ell^{2}(\Gamma)}^{2} d\mu(x) = \int_{C} \|\widehat{\psi_{\sigma}(\cdot,x)}\|_{L^{2}(\widehat{\Gamma})}^{2} d\mu(x) \\ &= \int_{C} \int_{\widehat{\Gamma}} |\sum_{\gamma \in \Gamma} \left[(\Pi_{\sigma}(\gamma)\psi)(x) \right] (-\gamma,\alpha)|^{2} dm_{\widehat{\Gamma}}(\alpha) d\mu(x) \\ &= \int_{\widehat{\Gamma}} \int_{C} |Z_{\sigma}[\psi](\alpha)(x)|^{2} d\mu(x) dm_{\widehat{\Gamma}}(\alpha). \end{aligned}$$

Therefore, $Z_{\sigma}[\psi](\alpha) \in L^2(C)$ for $m_{\widehat{\Gamma}}$ -a.e $\alpha \in \widehat{\Gamma}$. Moreover, using again Fubini's Theorem, one can see that $Z_{\sigma}[\psi]$ is measurable as a vector valued function from $\widehat{\Gamma}$ to $L^2(C)$. Continuing with (6), we obtain

$$\|\psi\|_{L^{2}(\mathcal{X})}^{2} = \int_{\widehat{\Gamma}} \|Z_{\sigma}[\psi](\alpha)\|_{L^{2}(C)}^{2} dm_{\widehat{\Gamma}}(\alpha) = \|Z_{\sigma}[\psi]\|^{2}.$$

Thus, $Z_{\sigma}[\psi] \in L^2(\widehat{\Gamma}, L^2(C))$ and Z_{σ} is an isometry.

Let us see now that Z_{σ} is onto. Let $\Phi \in L^2(\widehat{\Gamma}, L^2(C))$ be a simple function. Then, $\Phi = \sum_{j=1}^n f_j \chi_{E_j}$, where $f_j \in L^2(C)$ and $E_j \subseteq \widehat{\Gamma}$ is $m_{\widehat{\Gamma}}$ -measurable, for $j = 1, \ldots, n$. Since $\{X_\gamma\}_{\gamma \in \Gamma}$ is an orthonormal basis for $L^2(\widehat{\Gamma})$, we can write $\chi_{E_j} = \sum_{\gamma \in \Gamma} a_{\gamma}^j X_{\gamma}$ where $a^j := \{a_{\gamma}^j\}_{\gamma \in \Gamma} \in \ell^2(\Gamma)$ for every $j = 1, \ldots, n$, and then $\Phi = \sum_{\gamma \in \Gamma} \sum_{j=1}^n f_j a_{\gamma}^j X_{\gamma}$. For $\gamma \in \Gamma$ we define ψ on $\sigma_{\gamma}(C)$ as $\psi(\sigma_{\gamma}(x)) := J(\gamma, x)^{-1/2} \sum_{j=1}^{n} f_j(x) a_{\gamma}^j$ for μ -a.e. $x \in C$. Since C is a tiling set, we then have ψ defined on \mathcal{X} . Furthermore,

$$\begin{split} \|\psi\|_{L^{2}(\mathcal{X})}^{2} &= \sum_{\gamma \in \Gamma} \int_{C} |\psi(\sigma_{\gamma}(x))|^{2} J_{\sigma}(\gamma, x) \, d\mu(x) \\ &= \sum_{\gamma \in \Gamma} \int_{C} |\sum_{j=1}^{n} f_{j}(x) a_{\gamma}^{j}|^{2} \, d\mu(x) \leq \sum_{\gamma \in \Gamma} \int_{C} \sum_{j=1}^{n} |f_{j}(x)|^{2} \sum_{l=1}^{n} |a_{\gamma}^{l}|^{2} \, d\mu(x) \\ &= \sum_{l=1}^{n} \|a^{l}\|_{\ell^{2}(\Gamma)}^{2} \sum_{j=1}^{n} \|f_{j}\|_{L^{2}(C)}^{2} < +\infty, \end{split}$$

and thus $\psi \in L^2(\mathcal{X})$. Now for $\alpha \in \widehat{\Gamma}$ and $x \in C$ we have

$$Z_{\sigma}[\psi](\alpha)(x) = \sum_{\gamma \in \Gamma} \psi(\sigma_{-\gamma}(x)) J(\gamma, x)^{-1/2}(-\gamma, \alpha)$$
$$= \sum_{\gamma \in \Gamma} \sum_{j=1}^{n} f_j(x) a_{-\gamma}^j(-\gamma, \alpha) = \Phi(\alpha)(x).$$

Therefore, the set of simple functions is contained in the range of Z_{σ} . Since Γ is discrete and then $\widehat{\Gamma}$ compact, $m_{\widehat{\Gamma}}(\widehat{\Gamma})$ is finite and the set of simple functions is dense in $L^2(\widehat{\Gamma}, L^2(C))$ (See Proposition A.2 in the Appendix). Hence, due to the fact that the range of Z_{σ} is closed, we conclude that Z_{σ} is onto.

To verify that Z_{σ} satisfies (5) observe that, since Π_{σ} is a unitary representation, and each $\alpha \in \widehat{\Gamma}$ is a homomorphism, we have

$$Z_{\sigma}[\Pi(\gamma_{0})\psi](\alpha)(x) = \sum_{\gamma\in\Gamma}[\Pi_{\sigma}(\gamma)\Pi_{\sigma}(\gamma_{0})\psi(x)](-\gamma,\alpha) = \sum_{\gamma\in\Gamma}[\Pi_{\sigma}(\gamma+\gamma_{0})\psi(x)](-\gamma,\alpha)$$
$$= \sum_{\gamma\in\Gamma}[\Pi_{\sigma}(\gamma)\psi(x)](-\gamma+\gamma_{0},\alpha) = (\gamma_{0},\alpha)Z_{\sigma}[\psi](\alpha)(x).$$

4. Characterization of (Γ, σ) -invariant spaces of $L^2(\mathcal{X})$

In this section we want to characterize (Γ, σ) -invariant spaces in terms of the generalized Zak transform and range functions. Towards this end, we will evoke the general machinery developed by Bownik and Ross in [5] where they defined and characterized multiplicatively invariant spaces on $L^2(\Omega, \mathcal{H})$.

4.1. Multiplicatively invariant spaces in $L^2(\Omega, \mathcal{H})$. In order to make our exposition self contained, we hereby summarize the material we need concerning multiplicatively invariant spaces. See [5, Section 2] for details and proofs.

Fix (Ω, ν) a σ -finite measure space and \mathcal{H} a separable Hilbert space.

A set $D \subseteq L^{\infty}(\Omega)$ is said to be a *determining set* for $L^{1}(\Omega)$ if for every $f \in L^{1}(\Omega)$ such that $\int_{\Omega} f(\omega)g(\omega) d\nu(\omega) = 0 \ \forall g \in D$, one has f = 0. In the setting of Helson [20, 21], a determining set is the set of exponentials with integer parameter, $D = \{e^{2\pi i k \cdot}\}_{k \in \mathbb{Z}} \subseteq L^{\infty}(\mathbb{T}).$

Definition 4.1. A closed subspace $M \subseteq L^2(\Omega, \mathcal{H})$ is *multiplicatively invariant* with respect to the determining set D for $L^1(\Omega)$ (MI space for short) if

$$\Phi \in M \Longrightarrow g\Phi \in M$$
, for any $g \in D$.

For an at most countable subset $\boldsymbol{\Phi} \subseteq L^2(\Omega, \mathcal{H})$ define $M_D(\boldsymbol{\Phi}) = \overline{\operatorname{span}}\{g\Phi \colon \Phi \in \boldsymbol{\Phi}, g \in D\}$. The subspace $M_D(\boldsymbol{\Phi})$ is called the multiplicatively invariant space generated by $\boldsymbol{\Phi}$, and we say that $\boldsymbol{\Phi}$ is a set of generators for $M_D(\boldsymbol{\Phi})$.

MI spaces have been recently characterized in terms of measurable range functions by Bownik and Ross in [5]. We recall that a range function is a mapping $J: \Omega \to \{\text{closed subspaces of } \mathcal{H}\}$ equipped with the orthogonal projections $P_J(\omega)$ of \mathcal{H} onto $J(\omega)$. A range function is said to be measurable if for every $a \in \mathcal{H}$, $\omega \mapsto P_J(\omega)a$ is measurable as a vector valued function.

Theorem 4.2. [5, Theorem 2.4] Suppose that $L^2(\Omega)$ is separable, so that $L^2(\Omega, \mathcal{H})$ is also separable. Let M be a closed subspace of $L^2(\Omega, \mathcal{H})$ and D a determining set for $L^1(\Omega)$. Then, M is an MI space with respect to D if and only if there exists a measurable range function J such that

$$M = \{ \Phi \in L^2(\Omega, \mathcal{H}) : \Phi(\omega) \in J(\omega) \text{ a.e. } \omega \in \Omega \}.$$

Identifying range functions that are equal almost everywhere, the correspondence between MI spaces and measurable range functions is one-to-one and onto.

Moreover, when $M = M_D(\mathbf{\Phi})$ for some at most countable set $\mathbf{\Phi} \subseteq L^2(\Omega, \mathcal{H})$ the range function associated to M is

$$J(\omega) = \overline{\operatorname{span}} \{ \Phi(\omega) : \Phi \in \boldsymbol{\Phi} \}, \quad a.e. \ \omega \in \Omega.$$

4.2. Characterization of (Γ, σ) -invariant spaces. To our purpose of characterizing (Γ, σ) -invariant spaces we will connect them with MI spaces. We start by choosing $\Omega = \widehat{\Gamma}$ and $\mathcal{H} = L^2(C)$. We point out that since we assume that $L^2(\mathcal{X})$ is separable, so is $L^2(C)$. Furthermore, the space $L^2(\widehat{\Gamma})$ is also separable and then $L^2(\widehat{\Gamma}, L^2(C))$ is separable as well. In this setting a *range function* is a mapping

(7)
$$J: \Gamma \longrightarrow \{ \text{closed subspaces of } L^2(C) \}$$

equipped with the orthogonal projections $P_J(\alpha) : L^2(C) \to J(\alpha)$.

Our main result is:

Theorem 4.3. Let $V \subseteq L^2(\mathcal{X})$ be a closed subspace and let Z_{σ} be the Zak transform of Proposition 3.3. Then, the following conditions are equivalent:

- (1) V is a (Γ, σ) -invariant space;
- (2) there exists a measurable range function J such that

$$V = \left\{ f \in L^2(\mathcal{X}) : \ Z_{\sigma}[f](\alpha) \in J(\alpha) \ a.e. \ \alpha \in \widehat{\Gamma} \right\}$$

Identifying range functions which are equal almost everywhere, the correspondence between (Γ, σ) -invariant spaces and measurable range functions is one to one and onto.

Moreover, if $V = S^{\Gamma}_{\sigma}(\mathcal{A})$ for some countable subset \mathcal{A} of $L^{2}(\mathcal{X})$, the measurable range function J associated to V is given by

$$J(\alpha) = \overline{\operatorname{span}}\{Z_{\sigma}[\phi](\alpha) : \phi \in \mathcal{A}\}, \quad a.e. \ \alpha \in \widehat{\Gamma}.$$

The proof of Theorem 4.3 is based in the next lemma.

Lemma 4.4. Let $V \subseteq L^2(\mathcal{X})$ be a closed subspace. Then, V is a (Γ, σ) -invariant space if and only if $Z_{\sigma}[V] \subseteq L^2(\widehat{\Gamma}, L^2(C))$ is a MI space with respect to the determining set $D = \{X_{\gamma}\}_{\gamma \in \Gamma}$. In particular, $\mathcal{A} \subseteq L^2(\mathcal{X})$ is a set of generators for the (Γ, σ) -invariant space V if and only if the set $Z_{\sigma}[\mathcal{A}]$ generates $Z_{\sigma}[V]$ as a D-MI space.

Proof. First note that, as a consequence of the uniqueness of the Fourier transform, the set $D = \{X_{\gamma}\}_{\gamma \in \Gamma}$ is a determining set for $L^1(\widehat{\Gamma})$. Then, the result follows from the fact that the Zak transform is an isomorphism and by property (5).

Proof of Theorem 4.3. Let us first show $(1) \Rightarrow (2)$. By Lemma 4.4, $Z_{\sigma}[V]$ is a D-MI space in $L^{2}(\widehat{\Gamma}, L^{2}(C))$. Then by Theorem 4.2 there exists a measurable range function J such that $Z_{\sigma}[V] = \{\Phi \in L^{2}(\widehat{\Gamma}, L^{2}(C)) : \Phi(\alpha) \in J(\alpha) \text{ a.e. } \alpha \in \widehat{\Gamma}\}$. This, together with Proposition 3.3 implies that $V = \{f \in L^{2}(\mathcal{X}) : Z_{\sigma}[f](\alpha) \in J(\alpha) \text{ a.e. } \alpha \in \widehat{\Gamma}\}$, proving (2).

To prove $(2) \Rightarrow (1)$ observe that if $V = \{f \in L^2(\mathcal{X}) : Z_{\sigma}[f](\alpha) \in J(\alpha) \text{ a.e. } \alpha \in \widehat{\Gamma}\}$, for some measurable range function J, then $Z_{\sigma}[V] = \{\Phi \in L^2(\widehat{\Gamma}, L^2(C)) : \Phi(\alpha) \in J(\alpha) \text{ a.e. } \alpha \in \widehat{\Gamma}\}$. This implies that $Z_{\sigma}[V]$ is a *D*-MI space. Indeed, if $\Phi \in Z_{\sigma}[V]$, $\Phi(\alpha) \in J(\alpha)$ a.e. $\alpha \in \widehat{\Gamma}$ and thus, since $J(\alpha)$ is a subspace, $X_{\gamma}(\alpha)\Phi(\alpha) \in J(\alpha)$ a.e. $\alpha \in \widehat{\Gamma}$. Therefore, $X_{\gamma}\Phi \in Z_{\sigma}[V]$. Then, Lemma 4.4 guarantees that V is a (Γ, σ) -invariant space.

The bijective correspondence between measurable range functions and (Γ, σ) -invariant spaces is due to Theorem 4.2.

Finally, when $V = S_{\sigma}^{\Gamma}(\mathcal{A})$ for some $\mathcal{A} \subseteq L^{2}(\mathcal{X})$, we know by Lemma 4.4 that $Z_{\sigma}[V]$ is generated by $Z_{\sigma}[\mathcal{A}]$ as a *D*-MI space. Thus, Theorem 4.2 guarantees that the range function J is $J(\alpha) = \overline{\operatorname{span}}\{Z_{\sigma}[\phi](\alpha) : \phi \in \mathcal{A}\}$, a.e. $\alpha \in \widehat{\Gamma}$.

When \mathbb{Z}^n acts by translations on \mathbb{R}^n , it is known that there is a relationship between the minimum number of functions that one needs to generate a shiftinvariant space and the dimension of the spaces defined by the range function associated to it (see [40, Proposition 4.1]). This connection is also true for (Γ, σ) invariant spaces. In fact, we shall show that it is true at the general level of MI spaces.

Keeping the notation of Section 4.1, suppose that $M \subseteq L^2(\Omega, \mathcal{H})$ is a MI space with respect to D generated by a finite number of functions in $L^2(\Omega, \mathcal{H})$. Define the *length* of M as

 $\ell(M) := \min\{n \in \mathbb{N} \colon \exists \Phi_1, \cdots, \Phi_n \in M \text{ with } M = M_D(\Phi_1, \dots, \Phi_n)\}.$

The length of M can be expressed in terms of the range function associated to M as follows:

Proposition 4.5. Let $M \subseteq L^2(\Omega, \mathcal{H})$ be a MI space finitely generated and let J be the range function associated to M through Theorem 4.2. Then,

$$\ell(M) = \operatorname{ess\,sup\,dim}_{\omega \in \Omega} (J(\omega)).$$

Proof. First put $\ell(M) = \ell$. Since M can be generated by a set of ℓ functions, Theorem 4.2 gives that $\dim(J(\omega)) \leq \ell$ for a.e. $\omega \in \Omega$ and then $\operatorname{ess\,sup}_{\omega \in \Omega} \dim(J(\omega)) \leq \ell$. To see that indeed the equality must hold, it is enough to prove that $\{\omega \in \Omega : \dim(J(\omega)) = \ell\}$ has positive ν -measure.

Suppose towards a contradiction that $\nu(\{\omega \in \Omega : \dim(J(\omega)) = \ell\}) = 0$. Then, $\dim(J(\omega)) \leq \ell - 1$ for a.e. $\omega \in \Omega$. Following the ideas of the proof of [40, Proposition 4.1], we will construct a set of $\ell - 1$ generators for M.

Let $\{\Phi_1, \ldots, \Phi_\ell\} \in L^2(\Omega, \mathcal{H})$ be a set of generators for M. Then, $J(\omega) = \operatorname{span}\{\Phi_i(\omega) : 1 \le i \le \ell\}$ for a.e. $\omega \in \Omega$. Since $\dim(J(\omega)) \le \ell - 1$ for a.e. $\omega \in \Omega$, there must be a vector of the set $\{\Phi_i(\omega) : 1 \le i \le \ell\}$ belonging to the space spanned by the others $\ell - 1$ vectors. With this in mind, define the $(\nu$ -measurable) function $f: \Omega \to \{1, \ldots, \ell\}$ as $f(\omega) = \min\{1 \le i \le \ell : J(\omega) = \operatorname{span}\{\Phi_j(\omega) : 1 \le j \le \ell, j \ne i\}\}$. Then, if $\Omega_i := \{\omega \in \Omega : f(\omega) = i\}, \nu(\Omega \setminus \bigcup_{i=1}^{\ell} \Omega_i) = 0$ and the union $\bigcup_{i=1}^{\ell} \Omega_i$ is disjoint. Note that if $\omega \in \Omega_i$, then $J(\omega) = \operatorname{span}\{\Phi_j(\omega) : 1 \le j \le \ell, j \ne i\}$.

We define functions $\Psi_1, \ldots, \Psi_{\ell-1} \in L^2(\Omega, \mathcal{H})$ such that: on Ω_1

$$\Psi_j = \Phi_{j+1}$$
, for $1 \le j \le \ell - 1$

and on $\Omega_i, 2 \leq i \leq \ell$,

$$\Psi_j = \begin{cases} \Phi_j & \text{if } 1 \le j \le i-1\\ \Phi_{j+1} & \text{if } i \le j \le \ell-1. \end{cases}$$

Therefore, $J(\omega) = \operatorname{span}\{\Psi_j(\omega) : 1 \le j \le \ell - 1\}$ for a.e. $\omega \in \Omega$ and by Theorem 4.2, $M = M_D(\Psi_1, \ldots, \Psi_{\ell-1})$ which is a contraction and the proposition is proven. \Box

For a (Γ, σ) -invariant space V which is finitely generated, we can define the *length* of V as $\ell(V) := \min\{n \in \mathbb{N} : \exists \phi_1, \cdots, \phi_n \in V \text{ with } V = S_{\sigma}^{\Gamma}(\phi_1, \ldots, \phi_n)\}$. As an immediate consequence of the above result and Lemma 4.4 we get:

Corollary 4.6. Let $V \subseteq L^2(\mathcal{X})$ be (Γ, σ) -invariant space finitely generated and let J be be the range function associated to V through Theorem 4.3. Then,

$$\ell(V) = \operatorname{ess\,sup\,dim}_{\alpha \in \widehat{\Gamma}} (J(\alpha))$$

5. Frame and Riesz systems arising from quasi- Γ -invariant actions

In this section we investigate necessary and sufficient conditions for the system $E_{\sigma}^{\Gamma}(\mathcal{A})$ to be a frame or a Riesz basis for $S_{\sigma}^{\Gamma}(\mathcal{A})$.

Theorem 5.1. Let $\mathcal{A} \subseteq L^2(\mathcal{X})$ be a countable set and let J be the measurable range function associated to $V = S_{\sigma}^{\Gamma}(\mathcal{A})$. Then, the following two conditions are equivalent:

- (i) $E_{\sigma}^{\Gamma}(\mathcal{A})$ is a frame for V with frame bound $0 < A \leq B < +\infty$.
- (ii) For almost every $\alpha \in \widehat{\Gamma}$, the set $\{Z_{\sigma}[\phi](\alpha) : \phi \in \mathcal{A}\}$ is a frame for $J(\alpha)$ with uniform frame bounds A, B.

Proof. By the isometry property of Z_{σ} and by (5) we have

$$\sum_{\phi \in \mathcal{A}} \sum_{\gamma \in \Gamma} |\langle f, \Pi_{\sigma}(\gamma)\phi \rangle_{L^{2}(\mathcal{X})}|^{2} = \sum_{\phi \in \mathcal{A}} \sum_{\gamma \in \Gamma} |\langle Z_{\sigma}[f], Z_{\sigma}[\Pi_{\sigma}(\gamma)\phi] \rangle_{L^{2}(\widehat{\Gamma}, L^{2}(C))}|^{2}$$

$$(8) = \sum_{\phi \in \mathcal{A}} \sum_{\gamma \in \Gamma} |\int_{\widehat{\Gamma}} \langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)} \overline{(\gamma, \alpha)} \, dm_{\widehat{\Gamma}}(\alpha)|^{2}.$$

Assume (i) holds. For a fixed $\phi \in \mathcal{A}$, call $F(\alpha) := \langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^2(C)}$ and note that $F \in L^1(\widehat{\Gamma})$. Then, by the Pontrjagin Duality [38, Theorem 1.7.2],

$$\widehat{F}(\gamma) = \int_{\widehat{\Gamma}} \langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)} \overline{(\gamma, \alpha)} \, dm_{\widehat{\Gamma}}(\alpha)$$

and thus, (8) implies

c

$$\sum_{\gamma \in \Gamma} |\widehat{F}(\gamma)|^2 < +\infty.$$

As a consequence we have that $\widehat{F} \in \ell^2(\Gamma)$ and then $F \in L^2(\widehat{\Gamma})$. By Plancherel's Theorem we then get,

$$\sum_{\gamma \in \Gamma} |\int_{\widehat{\Gamma}} \langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)} \overline{(\gamma, \alpha)} \, dm_{\widehat{\Gamma}}(\alpha)|^{2} \, dm_{\Gamma}(\gamma)$$

$$= \|\widehat{F}\|_{\ell^{2}(\Gamma)}^{2} = \|F\|_{L^{2}(\widehat{\Gamma})}^{2}$$

$$= \int_{\widehat{\Gamma}} |\langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)}|^{2} \, dm_{\widehat{\Gamma}}(\alpha).$$

From here the arguments leading to (ii) are an effortless adaptation of those in [4] and [7, Theorem 4.1], hence we leave them for the reader.

Conversely, from (ii) and since Z_{σ} is an isometry, it follows that for every $f \in$ $S^{\Gamma}_{\sigma}(\mathcal{A})$ and for a.e. $\alpha \in \widehat{\Gamma}$,

(9)
$$A \| Z_{\sigma}[f](\alpha) \|_{L^{2}(C)}^{2} \leq \sum_{\phi \in \mathcal{A}} |\langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)}|^{2} \leq B \| Z_{\sigma}[f](\alpha) \|_{L^{2}(C)}^{2}.$$

Therefore, integrating over $\widehat{\Gamma}$, (9) becomes

$$A\|Z_{\sigma}[f]\|_{L^{2}(\widehat{\Gamma},L^{2}(C))}^{2} \leq \sum_{\phi \in \mathcal{A}} \int_{\widehat{\Gamma}} |\langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)}|^{2} \leq B\|Z_{\sigma}[f]\|_{L^{2}(\widehat{\Gamma},L^{2}(C))}^{2}$$

and in particular, we have that $\alpha \mapsto \langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle$ belongs to $L^{2}(\widehat{\Gamma})$ for each $\phi \in \mathcal{A}$. Thus, with similar arguments to those we used in the other implication, we conclude

$$\sum_{\phi \in \mathcal{A}} \sum_{\gamma \in \Gamma} |\langle f, \Pi_{\sigma}(\gamma)\phi \rangle_{L^{2}(\mathcal{X})}|^{2} = \sum_{\phi \in \mathcal{A}} \int_{\widehat{\Gamma}} |\langle Z_{\sigma}[f](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)}|^{2},$$

and hence, the frame property of $E_{\sigma}^{\Gamma}(\mathcal{A})$ follows immediately.

In [22] it was proven that the unitary representation Π_{σ} is dual integrable with bracket map given by $[\psi, \phi](\alpha) = \langle Z_{\sigma}[\psi](\alpha), Z_{\sigma}[\phi](\alpha) \rangle_{L^{2}(C)}$. Then, if the set of generators \mathcal{A} has only one single function, Theorem 5.1 reduces to [22, Theorem 5.7] when the unitary representation involved is Π_{σ} :

Corollary 5.2. Let $\psi \in L^2(\mathcal{X})$ and define $\Omega_{\psi} = \{\alpha \in \widehat{\Gamma} : \|Z_{\sigma}[\psi](\alpha)\|_{L^2(C)}^2 > 0\}.$ Then, the following are equivalent:

- (i) $E_{\sigma}^{\Gamma}(\psi)$ is a frame for $S_{\sigma}^{\Gamma}(\psi)$ with frame bound $0 < A \leq B < +\infty$. (ii) For almost every $\alpha \in \Omega_{\psi}$, $A \leq \|Z_{\sigma}[\psi](\alpha)\|_{L^{2}(C)}^{2} \leq B$.

In the next theorem we state the analogous result to Theorem 5.1 for Riesz bases.

Theorem 5.3. Let $\mathcal{A} \subseteq L^2(\mathcal{X})$ be a countable set and let J be the measurable range function associated to $V = S_{\sigma}^{\Gamma}(\mathcal{A})$. Then, the following two conditions are equivalent:

- (i) $E_{\sigma}^{\Gamma}(\mathcal{A})$ is a Riesz basis for V with bound $0 < A \leq B < +\infty$.
- (ii) For almost every $\alpha \in \widehat{\Gamma}$, the set $\{Z_{\sigma}[\phi](\alpha) : \phi \in \mathcal{A}\}$ is a Riesz basis for $J(\alpha)$ with uniform bounds A, B.

Proof. We proceed as in [7, Theorem 4.3] and [4]. Let $\{b_{(\gamma,\phi)}\}_{\gamma\in\Gamma,\phi\in\mathcal{A}}$ be a scalar sequence with finite support. Then we have,

(10)
$$\|\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}b_{(\gamma,\phi)}\Pi_{\sigma}(\gamma)\phi\|_{L^{2}(\mathcal{X})}^{2} = \|\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}b_{(\gamma,\phi)}Z_{\sigma}[\Pi_{\sigma}(\gamma)\phi]\|^{2}$$
$$= \int_{\widehat{\Gamma}}\|\sum_{\phi\in\mathcal{A}}[\sum_{\gamma\in\Gamma}b_{(\gamma,\phi)}X_{\gamma}(\alpha)]Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2}dm_{\widehat{\Gamma}}(\alpha).$$

For each $\phi \in \mathcal{A}$, let P_{ϕ} be the trigonometric polynomial $P_{\phi} = \sum_{\gamma \in \Gamma} b_{(\gamma,\phi)} X_{\gamma}$. Note that since $\{b_{(\gamma,\phi)}\}_{\gamma\in\Gamma,\phi\in\mathcal{A}}$ has finite support, only finitely many polynomials P_{ϕ} are non-zero. In particular, for every $\alpha \in \widehat{\Gamma}$, the scalar sequence $\{P_{\phi}(\alpha)\}_{\phi \in \mathcal{A}}$ has finite support and moreover

(11)
$$\int_{\widehat{\Gamma}} \|P_{\phi}(\alpha)\|_{\ell^{2}(\mathcal{A})}^{2} dm_{\widehat{\Gamma}}(\alpha) = \sum_{\phi \in \mathcal{A}} \|P_{\phi}\|_{L^{2}(\widehat{\Gamma})}^{2} = \sum_{\phi \in \mathcal{A}} \sum_{\gamma \in \Gamma} |b_{(\gamma,\phi)}|^{2},$$

for a.e. $\alpha \in \widehat{\Gamma}$.

Suppose (*ii*) holds. Then, for $\{a_{\phi}\}_{\phi \in \mathcal{A}} = \{P_{\phi}(\alpha)\}_{\phi \in \mathcal{A}}$ we have

(12)
$$A\sum_{\phi\in\mathcal{A}}|P_{\phi}(\alpha)|^{2} \leq \|\sum_{\phi\in\mathcal{A}}P_{\phi}(\alpha)Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2} \leq B\sum_{\phi\in\mathcal{A}}|P_{\phi}(\alpha)|^{2}$$

for a.e. $\alpha \in \widehat{\Gamma}$. Integrating over $\widehat{\Gamma}$ in (12) and using (10) and (11) we get

$$A\sum_{\phi\in\mathcal{A}}\sum_{\gamma\in\Gamma}|b_{(\gamma,\phi)}|^2 \le \|\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}b_{(\gamma,\phi)}\Pi(\gamma)\phi\|^2 \le B\sum_{\phi\in\mathcal{A}}\sum_{\gamma\in\Gamma}|b_{(\gamma,\phi)}|^2.$$

Hence, (i) holds.

Conversely, suppose, towards a contradiction, that (ii) fails. Then, by a similar argument as the one given in the proof of [7, Theorem 4.1], there must exist a measurable set $W \subseteq \widehat{\Gamma}$ with $m_{\widehat{\Gamma}}(W) > 0$, a sequence with finite support $a = \{a_{\phi}\}_{\phi \in \mathcal{A}}$ and $\varepsilon > 0$ such that

(13)
$$\|\sum_{\phi\in\mathcal{A}}a_{\phi}Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2} > (B+\varepsilon)\|a\|_{\ell^{2}(\mathcal{A})}^{2}, \quad \forall \alpha \in W$$

or

and

(14)
$$\|\sum_{\phi\in\mathcal{A}}a_{\phi}Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2} < (A-\varepsilon)\|a\|_{\ell^{2}(\mathcal{A})}^{2}, \quad \forall \alpha \in W.$$

For each $\phi \in \mathcal{A}$ such that $a_{\phi} \neq 0$, let $m_{\phi} := a_{\phi}\chi_W \in L^{\infty}(\widehat{\Gamma})$. By [7, Lemma 4.4] there exists a sequence of trigonometric polynomials $\{P_n^{\phi}\}_{n\in\mathbb{N}}$ such that $P_n^{\phi}(\alpha) \rightarrow m_{\phi}(\alpha)$ when $n \to +\infty$ for a.e. $\alpha \in \widehat{\Gamma}$ and $\|P_n^{\phi}\|_{L^{\infty}(\widehat{\Gamma})} \leq C$ for all $n \in \mathbb{N}$ and some C > 0. For every $n \in \mathbb{N}$, let $\{b_{(\gamma,\phi)}^n\}_{\gamma\in\Gamma,\phi\in\mathcal{A}}$ be the sequence of coefficient of P_n^{ϕ} . Since, $E_{\sigma}(\mathcal{A})$ is a Riesz sequence we have

(15)
$$A\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}|b^{n}_{(\gamma,\phi)}|^{2} \leq \|\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}b^{n}_{(\gamma,\phi)}\Pi(\gamma)\phi\|^{2}_{L^{2}(\mathcal{X})} \leq B\sum_{\gamma\in\Gamma,\phi\in\mathcal{A}}|b^{n}_{(\gamma,\phi)}|^{2}.$$

Using (11) and (10) we can rewrite (15) as

(16)
$$A\|P_n^{\phi}\|_{L^2(\widehat{\Gamma})}^2 \leq \int_{\widehat{\Gamma}} \|\sum_{\phi \in \mathcal{A}} P_n^{\phi}(\alpha) Z_{\sigma}[\phi](\alpha)\|_{L^2(C)}^2 dm_{\widehat{\Gamma}}(\alpha) \leq B\|P_n^{\phi}\|_{L^2(\widehat{\Gamma})}^2.$$

Note that (16) extends to m_{ϕ} leading to

$$A\|m_{\phi}\|_{L^{2}(\widehat{\Gamma})}^{2} \leq \int_{\widehat{\Gamma}}\|\sum_{\phi\in\mathcal{A}}m_{\phi}(\alpha)Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2}dm_{\widehat{\Gamma}}(\alpha) \leq B\|m_{\phi}\|_{L^{2}(\widehat{\Gamma})}^{2}.$$

This last inequality contradicts (13), (14) because by integrating over $\widehat{\Gamma}$ in (13) and (14) we get

$$\int_{\widehat{\Gamma}} \|\sum_{\phi \in \mathcal{A}} m_{\phi}(\alpha) Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2} dm_{\widehat{\Gamma}}(\alpha) \geq (B+\varepsilon) \|m_{\phi}\|_{L^{2}(\widehat{\Gamma})}^{2}$$
$$\int_{\widehat{\Gamma}} \|\sum_{\phi \in \mathcal{A}} m_{\phi}(\alpha) Z_{\sigma}[\phi](\alpha)\|_{L^{2}(C)}^{2} dm_{\widehat{\Gamma}}(\alpha) \leq (A-\varepsilon) \|m_{\phi}\|_{L^{2}(\widehat{\Gamma})}^{2}.$$

As for the frame case, when \mathcal{A} has only one element, Theorem 5.3 recovers [22, Proposition 5.3].

Corollary 5.4. Let $\psi \in L^2(\mathcal{X})$. Then, they are equivalent

- (i) $E_{\sigma}^{\Gamma}(\psi)$ is a Riesz basis for $S_{\sigma}^{\Gamma}(\psi)$ with bounds $0 < A \leq B < +\infty$.
- (ii) For almost every $\alpha \in \widehat{\Gamma}$, $A \leq \|Z_{\sigma}[\psi](\alpha)\|_{L^{2}(C)}^{2} \leq B$.

In the next theorem, we prove that each (Γ, σ) -invariant space can be decomposed into an orthogonal sum of principal (Γ, σ) -invariant spaces. This is not new or even surprising so far, since it follows from the well-known fact that every unitary representation is the orthogonal sum of cyclic representations (cf. [15, Proposition 3.3]). But we shall actually prove more, namely that each principal (Γ, σ) -invariant space involved in the decomposition can be chosen to be generated by a function ψ such that $E_{\sigma}^{\Gamma}(\psi)$ is a Parseval frame for $S_{\sigma}^{\Gamma}(\psi)$. This result is based in the Decomposition Theorem for MI spaces [5, Theorem 2.6] (see also [21]).

Theorem 5.5. Let $V \subseteq L^2(\mathcal{X})$ be a (Γ, σ) -invariant space. Then, there exist a sequence of functions $\{\psi_n\}_{n\in\mathbb{N}} \subseteq L^2(\mathcal{X})$ such that V can be decomposed into an orthogonal sum

$$V = \bigoplus_{n=1}^{\infty} S_{\sigma}^{\Gamma}(\psi_n)$$

and, for each $n \in \mathbb{N}$, $E_{\sigma}^{\Gamma}(\psi_n)$ is a Parseval frame for $S_{\sigma}^{\Gamma}(\psi_n)$.

Proof. Since V is a (Γ, σ) -invariant space, by Lemma 4.4, $M := Z_{\sigma}[V]$ is a MI space with respect to $D = \{X_{\gamma}\}_{\gamma \in \Gamma}$. Thus, by [5, Theorem 2.6] M can be decomposed into an orthogonal sum $M = \bigoplus_{n=1}^{\infty} M_n$ where each M_n is a MI space and the range function associated to M_n is $J_n(\alpha) = \operatorname{span}\{\Phi_n(\alpha)\}$ with $\Phi_n \in L^{\infty}(\widehat{\Gamma}, L^2(C))$ such that $\|\Phi_n(\alpha)\|_{L^2(C)} \in \{0, 1\}$ for a.e. $\alpha \in \widehat{\Gamma}$.

Since, $\widehat{\Gamma}$ has finite measure, it follows that $\Phi_n \in L^2(\widehat{\Gamma}, L^2(C))$ and $M_n = M_D(\Phi_n)$. Now, define $\psi_n \in L^2(\mathcal{X})$ such that $Z_{\sigma}[\psi_n] = \Phi_n$. Then, since Z_{σ} is an isometric isomorphism and by Lemma 4.4 we have that $V = \bigoplus_{n=1}^{\infty} S_{\sigma}^{\Gamma}(\psi_n)$. Moreover, since $\|\Phi_n(\alpha)\|_{L^2(C)} \in \{0,1\}$ for a.e. $\alpha \in \widehat{\Gamma}$, Corollary 5.2 implies that $E_{\sigma}^{\Gamma}(\psi_n)$ is a Parseval frame for $S_{\sigma}^{\Gamma}(\psi_n)$ and the theorem is proven.

Remark 5.6. If $\{\psi_n\}_{n\in\mathbb{N}}$ is a sequence as in Theorem 5.5 for the (Γ, σ) -invariant space V, it follows that $E_{\sigma}^{\Gamma}(\mathcal{A})$ is a Parseval frame for V, where $\mathcal{A} = \{\psi_n : n \in \mathbb{N}\}$. In particular this guarantees that every (Γ, σ) -invariant space V has a frame of the form $E_{\sigma}^{\Gamma}(\mathcal{A})$ for some set \mathcal{A} . This was already known when the group involved acts by translations (see [4, 5, 7]).

So far, we know that every (Γ, σ) -invariant space has a frame of the form $E_{\sigma}^{\Gamma}(\mathcal{A})$ for some set \mathcal{A} . But the statement is not true anymore if we replace frame for Riesz basis. That is, not every (Γ, σ) -invariant space has a Riesz basis of the form $E_{\sigma}^{\Gamma}(\mathcal{A})$ for some set \mathcal{A} . We now provide an example of this situation.

Example 5.7. Let $W_1 \subseteq \widehat{\Gamma}$ be a measurable set with $0 < m_{\widehat{\Gamma}}(W_1) < 1/2$. In particular, we have $m_{\widehat{\Gamma}}(\widehat{\Gamma} \setminus W_1) > 0$. Further, let $W_2 \subseteq C$ be a measurable set with $0 < \mu(W_2) < +\infty$ and consider the function $\Phi(\alpha)(x) := \chi_{W_1}(\alpha)\chi_{W_2}(x)$. It is immediately seen that $\Phi \in L^2(\widehat{\Gamma}, L^2(C))$ and that for all $\alpha \in \widehat{\Gamma}$, $\|\Phi(\alpha)\|_{L^2(C)}^2 = \chi_{W_1}(\alpha)\mu(W_2)$. Define $\psi \in L^2(\mathcal{X})$ by $Z_{\sigma}[\psi] = \Phi$ and put $V = S_{\sigma}^{\Gamma}(\psi)$. Then, the following hold:

(i) $E_{\sigma}^{\Gamma}(\psi)$ is a frame for V but not a Riesz basis.

(*ii*) If $\mathcal{A} \subseteq L^2(\mathcal{X})$ and $V = S^{\Gamma}_{\sigma}(\mathcal{A})$ then, $E^{\Gamma}_{\sigma}(\mathcal{A})$ cannot be a Riesz basis for V.

Item (i) simply follows from Corollaries 5.2 and 5.4. To see (ii), suppose that \mathcal{A} is a set of generator for V and that $E_{\sigma}^{\Gamma}(\mathcal{A})$ is a Riesz basis for V. Then, by Theorem 4.3, for a.e. $\alpha \in \widehat{\Gamma}$, $\{Z_{\sigma}[\phi](\alpha) : \phi \in \mathcal{A}\}$ is a Riesz basis for $J(\alpha)$ where J is the range function associated to V. Thus, denoting by $|\mathcal{A}|$ the number of elements of \mathcal{A} , we have that $|\mathcal{A}| = \dim(J(\alpha))$ for a.e. $\alpha \in \widehat{\Gamma}$. But, on the other hand, V has length 1 and then $|\mathcal{A}| = 1$ (cf. Corollary 4.6). Now, take $\phi \in V$ such that $V = S^{\Gamma}_{\sigma}(\phi)$. Then, since V is generated also by ψ and $E^{\Gamma}_{\sigma}(\psi)$ is a frame for V we have that $\phi = \sum_{\gamma \in \Gamma} c_{\gamma} \Pi_{\sigma}(\gamma) \psi$ for some sequence of coefficients $\{c_{\gamma}\}_{\gamma \in \Gamma} \in \ell^{2}(\Gamma)$. By (5), $Z_{\sigma}[\phi] = \left(\sum_{\gamma \in \Gamma} c_{\gamma} X_{\gamma}\right) Z_{\sigma}[\psi]$ and then, $\{\alpha \in \widehat{\Gamma} : Z_{\sigma}[\phi](\alpha) \neq 0\} \subseteq \{\alpha \in \widehat{\Gamma} : Z_{\sigma}[\psi](\alpha) \neq 0\} = W_{1}$. This means that $Z_{\sigma}[\phi]$ vanishes on a set of positive measure of $\widehat{\Gamma}$ (namely $\widehat{\Gamma} \setminus W_{1}$) and then condition (*ii*) in Corollary 5.4 is not satisfied for ϕ , implying that $E^{\Gamma}_{\sigma}(\phi)$ cannot be a Riesz basis for V.

6. CLOSED SUBGROUPS ACTING BY TRANSLATION

Let G be a second countable LCA group and let Γ be a closed subgroup of G which does not need to be discrete. We devote this section to study closed subspaces of $L^2(G)$ that are invariant under translations in Γ . When G/Γ is compact, the structure of theses spaces has been analyzed first in [7] for the case when Γ is discrete, and recently in [5] when Γ is not required to be discrete. The main results of [5, 7] are a characterization of spaces invariant under translations in terms of range function using fiberization techniques.

Our aim is to show how these spaces can be characterized in terms of range functions without assuming that G/Γ is compact. For this purpose, we shall work with an appropriate non-discrete generalized Zak transform Z which will reduce to the mapping Z_{σ} introduced in Section 3 when the action $\sigma : \Gamma \times G \to G$ is $\sigma(\gamma, x) = \gamma + x$ and Γ is discrete. The mapping Z will be the one that will replace the fiberization techniques of [5, 7]. Since we only assume that Γ is closed, our results will be applicable to cases where those in [5, 7] are not.

6.1. Setting. Let G be a second countable LCA group. As it was pointed out in [5] this is equivalent to say that G (or \widehat{G}) is σ -compact and metrizable. The translation operator by an element $y \in G$ is denoted by $T_y : L^2(G) \to L^2(G)$ and is given by $T_y f(x) = f(x - y)$.

Let $\Gamma \subseteq G$ be a closed subgroup of G. We fix C a m_G -measurable section of the quotient G/Γ . Its existence is guaranteed by [32, Lemma 1.1] or [14, Theorem 1].

The classes in G/Γ are denoted by $[x] = x + \Gamma$, for $x \in G$. We define the *cross-section* mapping $\tau : G/\Gamma \to C$ by $\tau([x]) = [x] \cap C$. Through τ we can carry over the topological and algebraic structure of G/Γ onto C. Therefore, in what follows we will consider C as a topological space with the topology it inherits from G/Γ through τ . Thus, τ is bijective continuous mapping with continuous inverse. Furthermore, we can define a measure μ on C by

(17)
$$\mu(E) := m_{G/\Gamma}(\tau^{-1}(E))$$

for every set E in the Borel σ -algebra $\Sigma = \{E \subseteq C : \tau^{-1}(E) \text{ is } m_{G/\Gamma}\text{-measurable}\}$. Since $m_{G/\Gamma}$ is Borel regular and τ and τ^{-1} are continuous, μ is also a Borel regular measure on C. Moreover, since C is Borel measurable with respect to m_G , by [14, Theorem 1], we have that $\{A \subseteq C : A \text{ is } m_G\text{-measurable}\} \subseteq \Sigma$. Measures constructed as in (17) are usually called *pushforward or image measures*. For details on the subject we refer to [3, Chapter 3, Section 3.6].

We need the following version of the so-called Weil's formula, [26, Theorem (28.54)].

Theorem 6.1. Let G be an LCA group and Γ be a closed subgroup of G. Consider $C \subseteq G$ a m_G -measurable section for the quotient G/Γ and let μ be the measure on C given by (17). Then, for every $f \in L^1(G)$ one has that for μ -almost $x \in C$, the function $\gamma \mapsto f(x + \gamma)$ is m_{Γ} -measurable and belongs to $L^1(\Gamma)$. Furthermore, the

function $x \mapsto \int_{\Gamma} f(x+\gamma) dm_{\Gamma}(\gamma)$ is μ -measurable, belongs to $L^1(C) := L^1(C,\mu)$ and, if m_{Γ} and μ are fixed, the Haar measure on G, m_G can be chosen so that

(18)
$$\int_{G} f(y) \, dm_G(y) = \int_{C} \int_{\Gamma} f(x+\gamma) \, dm_{\Gamma}(\gamma) \, d\mu(x).$$

Proof. First note that for every $F \in L^1(G/\Gamma)$, $F \circ \tau^{-1} \in L^1(C)$ and

(19)
$$\int_{G/\Gamma} F([x]) \, dm_{G/\Gamma}([x]) = \int_C F(\tau^{-1}(x)) \, d\mu(x).$$

This is in fact [3, Theorem 3.6.1], but it is easily seen since, by definition of μ , (19) holds for $F = \chi_A$ with $A \subseteq G/\Gamma m_{G/\Gamma}$ -measurable. Then, by linearity, (19) holds for simple functions in $L^1(G/\Gamma)$. Finally, using standard arguments of measure theory (cf. [43, Theorems 10.3 and 10.20]), one can see that (19) extends to all $F \in L^1(G/\Gamma)$. It is also true that every function in $L^1(C)$ is of the form $F \circ \tau^{-1}$ for some $F \in L^1(G/\Gamma)$.

Now, if $f \in L^1(G)$, [26, Theorem (28.54)] implies that $F([x]) = \int_{\Gamma} f(x + \gamma) dm_{\Gamma}(\gamma)$ belongs to $L^1(G/\Gamma)$ and the Haar measures can be chosen in order to have $\int_G f(y) dm_G(y) = \int_{G/\Gamma} F([x]) dm_{G/\Gamma}([x])$. Then, the result follows by applying (19).

Remark 6.2.

- (1) When Γ is assumed to be discrete and countable and m_{Γ} is chosen to be the counting measure, the measure μ in (17) is the restriction of m_G to C. This follows from an easily adaptation of the proof of [7, Lemma 2.10]. In particular, we have that $0 < m_G(C)$.
- (2) When Γ is not discrete, it could be that $0 = m_G(C)$. Then, $\mu \neq m_G|_C$. Indeed, if μ were $m_G|_C$, the right hand side of (18) would be 0 for every $f \in L^1(G)$ which is a contraction. As an example, consider $\Gamma = \mathbb{R} \times \{0\} \subseteq G = \mathbb{R}^2$. Then, $C = \{0\} \times \mathbb{R}$ and it has zero Lebesgue measure. In this case, formula (18) becomes Fubini's formula, $\int_{\mathbb{R}^2} f(x, y) d(x, y) = \int_{\mathbb{R}} \left(\int_{\mathbb{R}} f(x, y) dx \right) dy$.

The annihilator of Γ is the (closed) subgroup $\Gamma^* = \{\delta \in \widehat{G} : (\gamma, \delta) = 1 \text{ for all } \gamma \in \Gamma\}$ of the dual group \widehat{G} of G. Let Ω be a $m_{\widehat{G}}$ -measurable section of \widehat{G}/Γ^* . The section Ω of \widehat{G}/Γ^* can be identified with $\widehat{\Gamma}$, through the bijective mapping $\pi : \Omega \to \widehat{\Gamma}$ given by

(20)
$$\pi(\omega): \Gamma \longrightarrow \mathbb{C}, \qquad \pi(\omega)(\gamma) = (\gamma, \omega).$$

If $\tilde{\tau}$ is the cross-section mapping for the quotient \widehat{G}/Γ^* and the section Ω , then, $\pi \circ \tilde{\tau} : \widehat{G}/\Gamma^* \to \widehat{\Gamma}$ is an isomorphism which is in fact an homeomorphism (see [38, Theorem 2.1.2]). Furthermore, it can be proven that $m_{\widehat{G}/\Gamma^*}((\pi \circ \tilde{\tau})^{-1}(\cdot))$ defines a Haar measure on $\widehat{\Gamma}$, that we call $m_{\widehat{\Gamma}}$. In the same way we saw (19), one can show that for every $F \in L^1(\widehat{G}/\Gamma^*)$,

$$\int_{\widehat{G}/\Gamma^*} F([\xi]) \, dm_{\widehat{G}/\Gamma^*}([\xi]) = \int_{\widehat{\Gamma}} F((\pi \circ \widetilde{\tau})^{-1}(\xi)) \, dm_{\widehat{\Gamma}}(\xi).$$

Now, we consider on Ω the measure ν defined by (17) but with $m_{\widehat{G}/\Gamma^*}$ and the cross-section mapping $\tilde{\tau}$ associated to \widehat{G}/Γ^* and Ω . Then, Theorem 6.1 and the above discussion prove that for every $G \in L^1(\Omega) := L^1(\Omega, \nu)$,

(21)
$$\int_{\Omega} G(\omega) \, d\nu(\omega) = \int_{\widehat{\Gamma}} G(\pi^{-1}((\cdot, \omega)) \, dm_{\widehat{\Gamma}}((\cdot, \omega)),$$

where (\cdot, ω) are all the elements of $\widehat{\Gamma}$.

As we specified in Section 2.1, for each $\gamma \in \Gamma$, the corresponding character on $\widehat{\Gamma}$ is denoted by X_{γ} . We then define for each $\gamma \in \Gamma$, the functions $\tilde{X}_{\gamma} : \Omega \to \mathbb{C}$ by $\tilde{X}_{\gamma}(\omega) := X_{\gamma} \circ \pi^{-1}((\cdot, \omega)) = (\gamma, \omega)$. In particular, by the uniqueness of the Fourier transform and (21), we conclude that $D = {\tilde{X}_{\gamma}}_{\gamma \in \Gamma}$ is a determining set for $L^{1}(\Omega)$. This shall be relevant in Subsection 6.3.

6.2. A non-discrete generalized Zak transform. Let us keep the notation and the hypotheses on G and Γ of Subsection 6.1. Throughout this section Ω is a $m_{\widehat{G}}$ measurable section for the quotient \widehat{G}/Γ^* and C is a m_G -measurable section for the quotient G/Γ . We regard C and Ω as measure spaces with measures μ and ν respectively, where both are given by (17). Furthermore, we consider the following normalization of the Haar measures involved: first, we fix m_{Γ} such that the Inversion formula [38, Section 1.5] holds for m_{Γ} and $m_{\widehat{\Gamma}} = m_{\widehat{G}/\Gamma^*}((\pi \circ \tilde{\tau})^{-1}(\cdot))$. In particular, we have that m_{Γ} and $m_{\widehat{\Gamma}}$ are dual measures, meaning that Plancherel's Theorem is true. Then, for μ and the already fixed m_{Γ} , we choose m_G such that Theorem 6.1 holds.

Definition 6.3. Let $\psi \in L^1(G)$. For $\omega \in \Omega$ and $x \in C$ such that $\gamma \mapsto \psi(x - \gamma) \in L^1(\Gamma)$ we define the *Zak transform* of ψ as

$$Z[\psi](\omega)(x) := \int_{\Gamma} \psi(x-\gamma)(-\gamma,\omega) \, dm_{\Gamma}(\gamma).$$

Note that, due to Theorem 6.1, the above definition makes complete sense for all $\omega \in \Omega$ and for μ -a.e. $x \in C$. Moreover, $Z[\psi](\omega)(x) = \mathcal{F}_{\Gamma}(\psi(x-\cdot))(\omega)$ where \mathcal{F}_{Γ} denotes the Fourier transform on Γ .

Our aim is to show that Z defines an isometric isomorphism between $L^2(G)$ and $L^2(\Omega, L^2(C))$. We need the following intermediate lemma. From now on, $C_c(G)$ will stand for the set of continuous functions on G with compact support.

Lemma 6.4. For $\psi \in C_c(G)$, we have $Z[\psi] \in L^2(\Omega, L^2(C))$ and $\|\psi\|_{L^2(G)} = \|Z[\psi]\|$. Moreover, $Z[T_{\gamma}\psi] = \tilde{X}_{\gamma}Z[\psi]$ for all $\gamma \in \Gamma$.

Proof. First note that since $\psi \in C_c(G)$, $Z[\psi]$ is a continuous function on both variables. Now, by Theorem 6.1, we know that

(22)
$$\int_{G} |\psi(y)|^2 dm_G(y) = \int_{C} \int_{\Gamma} |\psi(x-\gamma)|^2 dm_{\Gamma}(\gamma) d\mu(x).$$

This implies that for μ -a.e. $x \in C$, $\psi(x - \cdot) \in L^2(\Gamma)$. By Plancherel's Theorem and (21), we have

(23)
$$\int_{\Gamma} |\psi(x-\gamma)|^2 dm_{\Gamma}(\gamma) = \int_{\Omega} |\mathcal{F}_{\Gamma}(\psi(x-\cdot))(\omega)|^2 d\nu(\omega).$$

Thus, we can replace (23) in (22) and then, since $Z[\psi]$ is continuous, we can apply Fubini's Theorem to obtain

$$\int_{G} |\psi(y)|^2 dm_G(y) = \int_{\Omega} \int_{C} |\mathcal{F}_{\Gamma}(\psi(x-\cdot))(\omega)|^2 d\mu(x) d\nu(\omega)$$

From here, we conclude that $Z[\psi] \in L^2(\Omega, L^2(C))$ and $\|\psi\|_{L^2(G)} = \|Z[\psi]\|$.

To see that $Z[T_{\gamma}\psi] = \tilde{X}_{\gamma}Z[\psi]$, use (2) to obtain,

$$Z[T_{\gamma}\psi](\omega)(x) = \mathcal{F}_{\Gamma}[T_{\gamma}\psi(x-\cdot)](\omega)$$

= $\mathcal{F}_{\Gamma}[\psi(x-(\gamma+\cdot))](\omega) = (\gamma,\omega)\mathcal{F}_{\Gamma}[\psi(x-\cdot)](\omega)$
= $(\gamma,\omega)Z[\psi](\omega)(x).$

As a consequence of Lemma 6.4, $Z : C_c(G) \to L^2(\Omega, L^2(C))$ defines an isometry that can be extended by density to the whole $L^2(G)$. We keep on denoting by Zthe extension of $Z : C_c(G) \to L^2(\Omega, L^2(C))$ to $L^2(G)$. As we shall show in the upcoming result, Z turns out to be an isomorphism.

Proposition 6.5. Let $Z : L^2(G) \to L^2(\Omega, L^2(C))$ be the extension to $L^2(G)$ of the mapping of Lemma 6.4. Then, Z is an isometric isomorphism satisfying $Z[T_\gamma \psi] = \tilde{X}_\gamma Z[\psi]$ for all $\gamma \in \Gamma$ and all $\psi \in L^2(G)$.

Proof. We only need to prove that Z is onto. This will require to deal with some technical issues but roughly speaking, the idea is to show that there is a dense set of $L^2(\Omega, L^2(C))$ contained in the image of Z. We structure the proof on steps.

Step 1: Consider the set $\mathcal{D} := \{\Phi(\omega; x) = \mathcal{F}_{\Gamma}(\Psi(\cdot; x))(\omega) : \Psi \in C_{c}(\Gamma \times C)\}$. Then, by Plancherel's Theorem, $\mathcal{D} \subseteq C(\Omega \times C) \cap L^{2}(\Omega \times C)$. We shall show now that \mathcal{D} is dense in $L^{2}(\Omega \times C)$. Towards this end, take $\Phi \in L^{2}(\Omega \times C)$ and $\varepsilon > 0$. Now, choose $\Phi_{1}(\omega; x) = \sum_{j=1}^{n} \alpha_{j} f_{j}(\omega) g_{j}(x)$ with $f_{j} \in L^{2}(\Omega), g_{j} \in L^{2}(C)$ and $\alpha_{j} \in \mathbb{C}$ for each $1 \leq j \leq n$ such that $\|\Phi - \Phi_{1}\|_{L^{2}(\Omega \times C)} < \varepsilon/2$. This can be done since the set span $\{f.g: f \in L^{2}(\Omega), g \in L^{2}(C)\}$ is dense in $L^{2}(\Omega \times C)$ (e.g. [34, pp. 51]).

Put $\Psi_1(\gamma; x) = \sum_{j=1}^n \alpha_j \mathcal{F}_{\Gamma}^{-1}(f_j)(\gamma) g_j(x)$. Then, $\Psi_1 \in L^2(\Gamma \times C)$ and, due to the density of $C_c(\Gamma \times C)$ in $L^2(\Gamma \times C)$, we can choose $\Psi \in C_c(\Gamma \times C)$ with $\|\Psi_1 - \Psi\|_{L^2(\Gamma \times C)} < \varepsilon/2$. If we call $\Phi_2(\omega; x) = \mathcal{F}_{\Gamma}(\Psi(\cdot; x))(\omega) \in \mathcal{D}$, then, by Plancherel's and Fubini's Theorems we have that $\|\Phi_1 - \Phi_2\|_{L^2(\Omega \times C)} = \|\Psi_1 - \Psi\|_{L^2(\Gamma \times C)} < \varepsilon/2$. Therefore, $\|\Phi - \Phi_2\|_{L^2(\Omega \times C)} < \varepsilon$ and consequently, \mathcal{D} is dense in $L^2(\Omega \times C)$. Finally note that, since \widehat{G}/Γ^* is σ -compact, so is Ω . Then, by Proposition A.3, $L^2(\Omega, L^2(C)) = L^2(\Omega \times C)$ and we have that \mathcal{D} is dense in $L^2(\Omega, L^2(C))$.

Step 2: We now prove that \mathcal{D} is contained in the range of Z. Let $\Phi \in \mathcal{D}$, $\Phi(\omega; x) = \mathcal{F}_{\Gamma}(\Psi(\cdot; x))(\omega)$ with $\Psi \in C_c(\Gamma \times C)$. Consider the mapping $y \mapsto (\tau([y]) - \tau([y]))$ $y; \tau([y])$ from G to $\Gamma \times C$, where τ is the cross-section mapping for G/Γ and C, and define the function ψ on G as $\psi(y) := \Psi(\tau([y]) - y; \tau([y]))$. Since $\Psi \in C_c(\Gamma \times C)$ and $\tau([\cdot]) : (G, m_G) \to (C, \mu)$ is measurable, to see that ψ is m_G -measurable it is enough to prove that $\xi(y) := \tau([y]) - y$ is a measurable function from (G, m_G) to (Γ, m_{Γ}) . Observe first that by the discussion after (17) in Subsection 6.1, we can conclude that $\tau([\cdot]): G \to C \subseteq G$ is a measurable function when regarding C with the Haar measure of G, m_G . Thus, $\xi : G \to \Gamma \subseteq G$ is also measurable when we consider in Γ the measure m_G . Note now, that since the topology in Γ is the relative topology as a subgroup of G, any open set of Γ , U is the intersection of an open set V of G with Γ , i.e. $U = V \cap \Gamma$. It follows then, that open sets of Γ are m_G -measurable and therefore, $\beta(\Gamma) \subseteq \beta(G)$ where $\beta(\Gamma)$ (resp. $\beta(G)$) denotes the Borel σ -algebra in Γ (resp. in G). In particular, this implies that ξ is a measurable function from (G, m_G) to (Γ, m_{Γ}) as we wanted to prove. Moreover, ψ is bounded because so is Ψ and hence, ψ is locally integrable on G. Now, for every compact set $F \subseteq G$, we can apply (18) to $\chi_F |\psi| \in L^1(G)$ to obtain,

(24)

$$\int_{G} \chi_{F}(y) |\psi(y)| \, dm_{G}(y) = \int_{C} \int_{\Gamma} \chi_{F}(x+\gamma) |\psi(x+\gamma)| \, dm_{\Gamma}(\gamma) \, d\mu(x)$$

$$= \int_{C} \int_{\Gamma} \chi_{F}(x+\gamma) |\Psi(-\gamma;x)| \, dm_{\Gamma}(\gamma) \, d\mu(x)$$

$$\leq \int_{C} \int_{\Gamma} |\Psi(\gamma;x)| \, dm_{\Gamma}(\gamma) \, d\mu(x) < +\infty.$$

Since G is σ -compact, $G = \bigcup_{n \in \mathbb{N}} F_n$ where $F_n \subseteq G$ is compact for every $n \in \mathbb{N}$ and $F_n \subseteq F_{n+1}$. Thus, since $\chi_{F_n} |\psi| \to |\psi| m_G$ -a.e. on G, by (24) and [43, Corollary 10.30] we have that $\psi \in L^1(G)$. With the same argument applied to $|\psi|^2$, we also see that $\psi \in L^2(G) \cap L^1(G)$.

Finally,

$$Z[\psi](\omega)(x) = \int_{\Gamma} \psi(x-\gamma)(-\gamma,\omega) \, dm_{\Gamma}(\gamma)$$
$$= \int_{\Gamma} \Psi(\gamma;x)(-\gamma,\omega) \, dm_{\Gamma}(\gamma)$$
$$= \mathcal{F}_{\Gamma}(\Psi(\cdot;x))(\omega) = \Phi(\omega;x).$$

Therefore, \mathcal{D} is contained in the range of Z as we wanted to prove.

Step 3: Since the range of Z contains a dense set and it is closed, Z must be onto. \Box

Remark 6.6.

- (1) Suppose that Γ is discrete and countable. Then, since the Haar measure m_G on G is invariant under translations, the action σ : Γ × G → G, σ(γ, x) = γ + g is a quasi-Γ-invariant action with J_σ(γ, x) ≡ 1. Note that the unitary representation associated to σ is Π_σ(γ) = T_γ. Moreover, the section C of G/Γ is a tiling set for σ and then we are in a setting where the results of Section 3 can be applied. Then, if Z_σ is as in Definition 3.2 and Z as in Definition 6.3, we have Z_σ = Z.
- (2) The construction of the mapping Z was proposed by Weil in [41]. In that work, the author works with the quotients instead of sections and describes the range of Z in a different way than here. To our purpose, it is very important to exactly know that the range of Z is a vector valued space since this is what allows us to connect closed subspaces of $L^2(G)$ that are invariant under translations with MI spaces.

6.3. Subspaces invariant under translations. As we already said, we want to give a characterization in terms of range functions of closed subspaces of $L^2(G)$ that are invariant under translations in Γ , where G is a second countable LCA group and Γ is a closed subgroup of G. To be precise, we say that a closed subspace V of $L^2(G)$ is invariant under translation in Γ , or Γ -invariant for short, if $T_{\gamma}V \subseteq V$ for all $\gamma \in \Gamma$. As usual, we say that V is generated by an at most countable set $\mathcal{A} \in L^2(G)$ when $V = S^{\Gamma}(\mathcal{A}) := \overline{\operatorname{span}}\{T_{\gamma}\phi : \gamma \in \Gamma, \phi \in \mathcal{A}\}.$

If $Z : L^2(G) \to L^2(\Omega, L^2(C))$ is the isomorphism of Proposition 6.5, then Z turns Γ -invariant spaces in $L^2(G)$ into MI spaces in $L^2(\Omega, L^2(C))$ with respect to the determining set $D = \{\tilde{X}_{\gamma}\}_{\gamma \in \Gamma}$ and vice versa. This fact can be proven in an analogous way to that used to show Lemma 4.4. As a consequence, the following characterization for Γ -invariant spaces holds. Its proof is a straightforward adaptation of that of Theorem 4.3 and therefore we leave it for the reader.

Theorem 6.7. Let $V \subseteq L^2(G)$ be a closed subspace and Z be the mapping of Proposition 6.5. Then, the following conditions are equivalent:

- (1) V is a Γ -invariant space;
- (2) there exists $J: \Omega \to \{ closed \ subspaces \ of \ L^2(C) \}$, a measurable range function, such that

 $V = \{ f \in L^2(G) : Z[f](\omega) \in J(\omega), \nu - a.e. \ \omega \in \Omega \}.$

Identifying range functions which are equal almost everywhere, the correspondence between Γ -invariant spaces and measurable range functions is one to one and onto.

Moreover, if $V = S^{\Gamma}(\mathcal{A})$ for some countable subset \mathcal{A} of $L^{2}(G)$, the measurable range function J associated to V is given by

$$J(\omega) = \overline{\operatorname{span}}\{Z[\phi](\omega) : \phi \in \mathcal{A}\}, \qquad \nu - a.e. \ \omega \in \Omega.$$

Remark 6.8. Theorem 6.7 characterizes Γ -invariant spaces in terms of range function in the same spirit as in [5, Theorem 3.8]. The main difference with the results in [5] is that we do not require Γ to be co-compact (i.e. G/Γ compact). The co-compact assumption on Γ guaranties that Γ^* is discrete and this is what is crucial to define the mapping \mathcal{T} of [5, Proposition 3.7] (see also [7, Proposition 3.3]). Working with the isomorphism Z instead of \mathcal{T} allows as to withdraw the co-compactness on Γ and as a consequence, our results are applicable to cases where those in [5] are not. To give a simple example, consider $G = \mathbb{R}^3$ and $\Gamma = \mathbb{Z} \times \mathbb{R} \times \{0\}$. Then Γ is closed but $G/\Gamma \approx \mathbb{T} \times \{0\} \times \mathbb{R}$ is not compact. Furthermore, when Γ is discrete (but not necessarily co-compact) Theorem 6.7 is still more general than [7, Theorem 3.10]. For instance, it covers the (simple) case when $\Gamma = \mathbb{Z} \times \{0\} \subseteq G = \mathbb{R}^2$ which does not fit in the setting of [7, Theorem 3.10] because $G/\Gamma \approx \mathbb{T} \times \mathbb{R}$ is not compact.

Remark 6.9. As a consequence of Theorem 6.7 we recover Corollary 3.9 in [5]. That is, for a second countable LCA group G, a closed subspace V of $L^2(G)$ is invariant under all translations by G if and only if there exists a measurable set $E \subset \widehat{G}$ such that $V = \{f \in L^2(G) : \operatorname{supp} \widehat{f} \subset E\}$. Indeed, take $\Gamma = G$ in Theorem 6.7; then $L^2(C) = L^2(\{e\}) = \mathbb{C}, \ \Omega = \widehat{G}, \ \text{each } J(\omega) \text{ is a closed subspace of } \mathbb{C}$ and hence $J(\omega) = \mathbb{C}$ or $\{0\}, \ \text{and } Z[f](\omega)(e) = \mathcal{F}f(-\omega)$. Thus, it is enough to take $E = \{\omega \in \widehat{G} : J(-\omega) = \mathbb{C}\}.$

As one may expect, it is also possible to characterize frame and Riesz conditions of the system $E^{\Gamma}(\mathcal{A}) := \{T_{\gamma}\phi : \gamma \in \Gamma, \phi \in \mathcal{A}\}$ in terms of Z. Since $E^{\Gamma}(\mathcal{A})$ is indexed by the - in general - non-discrete set $\Gamma \times \mathcal{A}$, the frame and Riesz properties have to be regarded in the continuous sense as in Definitions 5.2 and 5.3 in [5]. Then, we have:

Theorem 6.10. Let $\mathcal{A} \subseteq L^2(G)$ be a countable set and let J the a measurable range function associated to $V = S^{\Gamma}(\mathcal{A})$ through Theorem 6.7. Then, the following two conditions are equivalent:

- (i) $E^{\Gamma}(\mathcal{A})$ is a continuous frame (resp., continuous Riesz basis) for V with bound $0 < A \leq B < +\infty$.
- (ii) For ν -almost every $\omega \in \Omega$, the set $\{Z[\phi](\omega) : \phi \in A\}$ is a frame (resp., Riesz basis) for $J(\omega)$ with uniform bounds A, B.

The proof of this result is, once again, a straightforward adaptation of the proof of [5, Theorem 5.1] and therefore we omit it. However, we do want to emphasize what was noticed in [5]: first, despite item (i) in Theorem 6.10 involves the concept of continuous frames (resp., Riesz bases), the frame and Riesz basis notions in (ii) are the usual ones. Second, with the same strategies than those used in the proof of Theorem 5.1 for Riesz bases in [5], it can be seen that both item (i) and item (ii) in Theorem 6.10 for Riesz bases imply that Γ is discrete. Then, the result reduces to Theorem 5.3 where the action involved is $\sigma : \Gamma \times G \to G$, $\sigma(\gamma, x) = \gamma + x$.

6.4. Connection with the so-called Fiberization mapping. As we already pointed out, when Γ is assumed to be co-compact, Γ -invariant spaces are characterized in terms of range functions either by Theorem 6.7 or [5, Theorem 3.8]. What we shall investigate now, is the relationship of the main tools used in those theorems, mainly Z, the non-discrete generalized Zak transform, and \mathcal{T} the fiberization mapping of [5, Proposition 3.7].

Let G, Γ , \widehat{G} , Γ^* , Ω and C be as in Section 6.1 with their respective measures and suppose, additionally, that G/Γ is compact. Then, Γ^* is discrete and countable (cf. [25, Theorem 24.15]). By Remark 6.2, we have that ν , the measure on Ω given by (17), is the restriction of $m_{\widehat{G}}$ to Ω . As we identified Ω with $\widehat{\Gamma}$, we can identify C with the dual group of Γ^* .

The isometry \mathcal{T} proposed in [5, 7] is the mapping $\mathcal{T}: L^2(G) \to L^2(\Omega, \ell^2(\Gamma^*))$ defined by

$$\mathcal{T}f(\omega) = \{f(\omega+\delta)\}_{\delta\in\Gamma^*}.$$

Proposition 6.11. For $f \in \mathcal{C}_c(G)$ such that $\widehat{f} \in L^1(\widehat{G})$ it holds that

$$\mathcal{F}_{\Gamma^*}(\mathcal{T}f(\omega))(x) = (x,\omega)Z_{\sigma}[f](-w)(-x),$$

for $x \in C$ and $\omega \in \Omega$, where \mathcal{F}_{Γ^*} denotes the Fourier transform with respect to the group Γ^* .

Proof. Fix $x \in C$ and $\omega \in \Omega$. Denoting by M_{ω} the modulation operator by ω , i.e $M_{\omega}\phi(\cdot) = (\cdot, \omega)\phi(\cdot)$, we can rewrite $Z_{\sigma}[f](-w)(-x)$ as

$$Z_{\sigma}[f](-\omega)(-x) = \int_{\Gamma} f(-x+\gamma)(\gamma,-\omega) \, dm_{\Gamma}(\gamma)$$
$$= (x,-\omega) \int_{\Gamma} T_x M_{-\omega} f(\gamma) \, dm_{\Gamma}(\gamma)$$

Since $f \in \mathcal{C}_c(G)$ and $\widehat{f} \in L^1(\widehat{G})$, the same holds for $T_x M_{-\omega} f$. This is, $T_x M_{-\omega} f \in \mathcal{C}_c(G)$ and $\widehat{T_x M_{-\omega}} f \in L^1(\widehat{G})$. Therefore, by the Poisson Formula [35, Theorem 5.5.2],

(25)
$$\int_{\Gamma} T_x M_{-\omega} f(\gamma) \, dm_{\Gamma}(\gamma) = \sum_{\delta \in \Gamma^*} \widehat{T_x M_{-\omega}} f(\delta) = \sum_{\delta \in \Gamma^*} \widehat{f}(\omega + \delta)(-x, \delta).$$

Since the right hand side of (25) is the Fourier transform of $\mathcal{T}f(\omega)$ at x with respect to the group Γ^* , the result follows.

As a consequence of Plancherel's Theorem, we immediately obtain:

Corollary 6.12. If f and g are as in Proposition 6.11 then,

 $\langle \mathcal{T}f(\omega), \mathcal{T}g(\omega) \rangle_{\ell^2(\Gamma^*)} = \langle Z_{\sigma}[f](-\omega), Z_{\sigma}[g](-\omega) \rangle_{L^2(C)}.$

Appendix A.

In this section we shall prove two results concerning vector valued spaces. The first one says that, when Ω has finite measure and \mathcal{H} is any separable Hilbert space, the set of simple functions of $L^2(\Omega, \mathcal{H})$ is dense. The second one shows that, when Ω is σ -finite and the Hilbert space \mathcal{H} is an L^2 -space, this is $\mathcal{H} = L^2(C)$ for some measure space (C, μ) , then $L^2(\Omega, L^2(C)) = L^2(\Omega \times C)$. We believe that both results are known, but since we could not find an explicit reference for them, we decided to provide a proof. We will make use of the following intermediate result. Its proof can be found in [13, Proof of Theorem 2.1, Chapter II].

Lemma A.1. Let (Ω, ν) be a measure space such that $\nu(\Omega) < +\infty$ and let \mathcal{H} be a separable Hilbert space. If $\Phi : \Omega \to \mathcal{H}$ is a measurable function, then, for every $\varepsilon > 0$ there exist a partition of Ω into measurable sets, $\{B_n\}_{n\in\mathbb{N}}$ and a sequence $\{x_n\}_{n\in\mathbb{N}} \subseteq \mathcal{H}$ such that $\Psi := \sum_{n=1}^{\infty} x_n \chi_{B_n}$ satisfy $\|\Phi(\omega) - \Psi(\omega)\|_{\mathcal{H}} < \varepsilon$ for a.e. $\omega \in \Omega$.

Proposition A.2. Suppose that $\nu(\Omega) < +\infty$. Then for each $\Phi \in L^2(\Omega, \mathcal{H})$ there exists a sequence $\{\Phi_n\}_{n\in\mathbb{N}} \subseteq L^2(\Omega, \mathcal{H})$ of simple functions such that $\lim_{n\to\infty} \|\Phi - \Phi_n\| = 0$.

Proof. Fix $\Phi \in L^2(\Omega, \mathcal{H})$. For each $n \in \mathbb{N}$, by Lemma A.1 we can choose $\Psi_n = \sum_{j=1}^{\infty} x_j^n \chi_{B_j^n}$ with $\{B_j^n\}_{j \in \mathbb{N}}$ being a partition of Ω in measurable sets and $\{x_j^n\}_{j \in \mathbb{N}} \subseteq \mathcal{H}$ such that $\|\Phi(\omega) - \Psi_n(\omega)\|_{\mathcal{H}} < \frac{1}{n}$ for a.e. $\omega \in \Omega$. Since $\Phi \in L^2(\Omega, \mathcal{H})$, we have that $\Psi_n \in L^2(\Omega, \mathcal{H})$ for all $n \in \mathbb{N}$. Indeed, for a.e. $\omega \in \Omega$, $\|\Psi_n(\omega)\|_{\mathcal{H}} \leq \frac{1}{n} + \|\Phi(\omega)\|_{\mathcal{H}} \leq 2 \max\{\frac{1}{n}, \|\Phi(\omega)\|_{\mathcal{H}}\}$ and then

(26)
$$\|\Psi_n(\omega)\|_{\mathcal{H}}^2 \le 4 \max\{\frac{1}{n^2}, \|\Phi(\omega)\|_{\mathcal{H}}^2\} \le 4(\frac{1}{n^2} + \|\Phi(\omega)\|_{\mathcal{H}}^2).$$

Now, integrating over Ω in (26) we get $\|\Psi_n\|^2 = \int_{\Omega} \|\Psi_n(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) \le 4(\frac{1}{n^2}\nu(\Omega) + \|\Phi\|^2) < +\infty.$

Note that, for every $n \in \mathbb{N}$, $\|\Psi_n\|^2 = \sum_{j=1}^{\infty} \|x_j^n\|_{\mathcal{H}}^2 \nu(B_j^n)$ and then we can choose N(n) large enough so that

(27)
$$\sum_{j=N(n)+1}^{\infty} \|x_j^n\|_{\mathcal{H}}^2 \nu(B_j^n) = \int_{\bigcup_{j=N(n)+1}^{\infty} B_j^n} \|\Psi_n(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) < \frac{1}{n}.$$

Define $\Phi_n = \sum_{j=1}^{N(n)} x_j^n \chi_{B_j^n}$ and let us see that $\Phi_n \to \Phi$, as $n \to +\infty$ in $L^2(\Omega, \mathcal{H})$. Proceeding as before and using (27), we have

$$\begin{split} \|\Phi_n - \Phi_n\|^2 &= \int_{\Omega} \|\Phi_n(\omega) - \Phi(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) \\ &\leq 4 \left(\int_{\Omega} \|\Phi_n(\omega) - \Psi_n(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) + \int_{\Omega} \|\Psi_n(\omega) - \Phi(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) \right) \\ &\leq 4 \left(\int_{\bigcup_{j=N(n)+1}^{\infty} B_j^n} \|\Psi_n(\omega)\|_{\mathcal{H}}^2 d\nu(\omega) + \int_{\Omega} \frac{1}{n^2} d\nu(\omega) \right) \\ &\leq 4 \left(\frac{1}{n} + \frac{\nu(\Omega)}{n^2} \right), \end{split}$$

and the result follows.

As it was pointed out to us by the referee, the next proposition is a generalization of [34, Theorem II.10 (c)].

Proposition A.3. Let (Ω, ν) be a σ -finite measure space and (C, μ) be a measure space such that $L^2(C)$ is separable. Then, $L^2(\Omega, L^2(C)) = L^2(\Omega \times C)$.

Proof. Clearly, by Fubini's Theorem, $L^2(\Omega \times C) \subseteq L^2(\Omega, L^2(C))$. The claim $L^2(\Omega, L^2(C)) = L^2(\Omega \times C)$ will follow from the fact that Ω is σ -finite and Proposition A.2. To see this, first write $\Omega = \bigcup_{n \in \mathbb{N}} \Omega_n$ where, for every $n \in \mathbb{N}$, $\Omega_n \subseteq \Omega$ is of finite ν -measure and $\Omega_n \subseteq \Omega_{n+1}$.

For every $n \in \mathbb{N}$, each simple function in $L^2(\Omega_n, L^2(C))$ (simple in the sense of vector valued functions) is a $(\nu \times \mu)$ -measurable function from $\Omega_n \times C$ to \mathbb{C} . Therefore, if $\mathcal{D}_n = \{$ simple functions of $L^2(\Omega_n, L^2(C)) \}$ we then have that $\mathcal{D}_n \subseteq$ $L^2(\Omega_n \times C) \subseteq L^2(\Omega_n, L^2(C))$. Now, since the norm in $L^2(\Omega_n, L^2(C))$ restricted to $L^2(\Omega_n \times C)$ is the norm on $L^2(\Omega_n \times C)$ and $L^2(\Omega_n \times C)$ is closed, we conclude from Proposition A.2 that $L^2(\Omega_n \times C) = L^2(\Omega_n, L^2(C))$.

Finally, take $\Phi \in L^2(\Omega, L^2(C))$ and put $\Phi_n := \chi_{\Omega_n} \Phi$. Then, for all $n \in \mathbb{N}$, $\Phi_n \in L^2(\Omega_n, L^2(C)) = L^2(\Omega_n \times C)$ and in particular, Φ_n is a $(\nu \times \mu)$ -measurable function on the product space $\Omega_n \times C$. Thus, since $\lim_{n\to\infty} \Phi_n(\omega)(x) = \Phi(\omega)(x)$ for $(\nu \times \mu)$ -a.e. $(\omega; x) \in \Omega \times C$, Φ is a $(\nu \times \mu)$ -measurable function on the product space $\Omega \times C$ and therefore, $\Phi \in L^2(\Omega \times C)$.

References

- A. Aldroubi and K. Gröchenig. Nonuniform sampling and reconstruction in shift-invariant spaces. SIAM Rev., 43(4):585–620 (electronic), 2001.
- [2] D. Barbieri, E. Hernández, and J. Parcet. Riesz and frame systems generated by unitary actions of discrete groups. *To appear in Appl. Comput. Harmon. Anal.*, DOI: 10.1016/j.acha.2014.09.007, 2014.
- [3] V. I. Bogachev. Measure theory. Vol. I, II. Springer-Verlag, Berlin, 2007.
- M. Bownik. The structure of shift-invariant subspaces of L²(Rⁿ). J. Funct. Anal., 177(2):282–309, 2000.
- [5] M. Bownik and K. A. Ross. The structure of translation-invariant spaces on locally compact abelian groups. To appear in J. Fourier Anal. Appl., DOI: 10.1007/s00041-015-9390-5, 2015.
- [6] J. Brezin. Harmonic analysis on manifolds. Trans. on Amer. Math. Soc., 150:611–618, 1970.
 [7] C. Cabrelli and V. Paternostro. Shift-invariant spaces on LCA groups. J. Funct. Anal., 258(6):2034–2059, 2010.
- [8] O. Christensen. An introduction to frames and Riesz bases. Birkhäuser, Boston, Basel, Berlin 2003.
- [9] D. L. Cohn. Measure theory. Birkhäuser Boston, Inc., Boston, MA, 1993. Reprint of the 1980 original.
- [10] B. Currey, A. Mayeli, and V. Oussa. Characterization of shift-invariant spaces on a class of nilpotent Lie groups with applications. J. Fourier Anal. Appl., 20(2):384–400, 2014.
- [11] C. de Boor, R. DeVore and A. Ron. The structure of finitely generated shift-invariant spaces in L²(R^d). J. Funct. Anal., 119(1):37 –78, 1994.
- [12] C. de Boor, R. DeVore and A. Ron. Approximation from shift-invariant subspaces of L²(R^d). Trans. Amer. Math. Soc., 341:787 – 806, 1994.
- [13] J. Diestel and Jr.,J.J.Uhl. Vector measures. With a foreword by B. J. Pettis, Mathematical Surveys, No. 15, American Mathematical Society, Providence, R.I., xiii+322, 1977.
- [14] J. Feldman and F. P. Greenleaf. Existence of Borel transversals in groups. Pacific J. Math., 25:455–461, 1968.
- [15] G. B. Folland. A course in abstract harmonic analysis. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995.
- [16] I.M. Gelfand. Eigenfunction expansions for equations with periodic coefficients. Dokl. Akad. Nauk. SSR., 73:1117–1120, 1950.
- [17] I. Gladkova. The Zak transform and a new approach to waveform design. IEEE Transactions on Aerospace and Electronic Systems., 37(4):1458–1464, 2001.
- [18] I. Gladkova. Design of frequency modulation waveforms via the Zak transform. IEEE Transactions on Aerospace and Electronic Systems., 40(1):355–359, 2004.
- [19] K. Gröchenig. Foundations of time-frequency analysis. Birkhäuser Boston, Inc., Boston, MA, 2001.
- [20] H. Helson. Lectures on invariant subspaces. Academic Press, New York, 1964.
- [21] H. Helson. The Spectral Theorem. Springer-Verlag New York, Inc. New York, USA, 1986.
- [22] E. Hernández, H. Šikić, G. Weiss, and E. Wilson. Cyclic subspaces for unitary representations of LCA groups; generalized Zak transform. *Collog. Math.*, 118(1):313–332, 2010.
- [23] E. Hernández, H. Šikić, G. Weiss, and E. Wilson. The Zak transform(s). In Wavelets and Mutiscale Analysis., 151–157, Birhäuser/Springer 2011.
- [24] E. Hernández and G. Weiss. A first course on wavelets. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1996. With a foreword by Yves Meyer.
- [25] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. I: Structure of topological groups, integration theory, group representations. Second edition. Springer-Verlag, New York-Berlin, 1979.
- [26] E. Hewitt and K. A. Ross. Abstract harmonic analysis. Vol. II: Structure and analysis for compact groups. Analysis on locally compact Abelian groups. Springer-Verlag, New York-Berlin, 1970.
- [27] J. W. Iverson. Subspaces of $L^2(G)$ invariant under translations by an abelian subgroup. J. Funct. Anal., 269:865 913, 2015.
- [28] A. J. E. M. Janssen. Bargmann transform, Zak transform, and coherent states. J. Math. Phys., 23:720–731, 1982.
- [29] A. J. E. M. Janssen. The Zak transform: A signal transform for sampled time-continuous signals. *Philips Journal of Research*, 34(1):23–69, 1988.
- [30] A. J. E. M. Janssen. The duality condition for Weyl-Heisenberg frames. In *Gabor analysis and algorithms*, Appl. Numer. Harmon. Anal., pages 33–84, Birkhäuser Boston, Boston, MA, 1998.

- [31] R. A. Kamyabi Gol and R. Raisi Tousi. A range function approach to shift-invariant spaces on locally compact abelian groups. Int. J. Wavelets Multiresolut. Inf. Process., 8(1):49–59, 2010.
- [32] G. W. Mackey. Induced representations of locally compact groups. I. Ann. of Math. (2), 55:101–139, 1952.
- [33] S. G. Mallat. Multiresolution approximations and wavelet orthonormal bases of L²(R). Trans. Amer. Math. Soc., 315(1):69–87, 1989.
- [34] M. Reed and B. Simon. Methods of modern mathematical physics. I. Second edition. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1980.
- [35] H. Reiter. Classical harmonic analysis and locally compact groups. Clarendon Press, Oxford, 1968.
- [36] A. Ron and Z. Shen. Frames and stable bases for shift-invariant subspaces of L²(ℝ^d). Canad. J. Math., 47:1051 − 1094, 1995.
- [37] A. Ron and Z. Shen. Affine systems in $L^2(\mathbb{R}^d)$: the analysis of the analysis operator. J. Funct. Anal., 148:408 – 447, 1997.
- [38] W. Rudin. Fourier analysis on groups. Interscience Tracts in Pure and Applied Mathematics, No. 12. Interscience Publishers (a division of John Wiley and Sons), New York-London, 1962.
- [39] S. Saliani. Linear independence of translates implies linear independence of affine Parseval frames on LCA groups. Preprint, 2014. Available at http://arxiv.org/abs/1411.1252
- [40] T. Tessera and H. Wang, Uncertainty principles in finitely generated shift-invariant spaces with additional invariance. J. Math. Anal. Appl., 410(1):134–143, 2014.
- [41] A. Weil, Sur certains groupes d'opérateurs unitaires. Acta Math., 111:143–211, 1964.
- [42] E. N. Wilson, The importance of Zak transforms for harmonic analysis. Revista de la Unión Matemática Argentina, 52(2):105–113, 2011.
- [43] R. L. Wheeden and A. Zygmund. Measure and integral. An introduction to real analysis. Pure and Applied Mathematics, Vol. 43. Marcel Dekker, Inc., New York-Basel, 1977.
- [44] J. Zak, Finite translations in solid state physics. Phys. Rev. Letter., 19:1385–1387, 1967.
- [45] J. Zak, The k_q -representation in the dynamics in electrons in solids. *Phys. Rev.*, 168:686–747, 1968.

(D. BARBIERI) UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN E-mail address: davide.barbieri@uam.es

(E. HERNÁNDEZ) UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN

E-mail address: eugenio.hernandez@uam.es

(V. PATERNOSTRO) DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NAT-URALES, UNIVERSIDAD DE BUENOS AIRES, ARGENTINA AND IMAS-CONICET, CONSEJO NACIONAL DE INVESTIGACIONES CIENTÍFICAS Y TÉCNICAS, ARGENTINA

E-mail address: vpater@dm.uba.ar