A new multivariate dependence measure based on comonotonicity

Ying Zhang

School of Mathematical Sciences, Qufu Normal University Shandong 273165, China e-mail: zhangying0513@gmail.com

Chuancun Yin *

School of Statistics, Qufu Normal University Shandong 273165, China e-mail: ccyin@mail.qfnu.edu.cn

August 15, 2018

Abstract In this paper we introduce a new multivariate dependence measure based on comonotonicity by means of product moment which motivated by the recent papers of Koch and Schepper (ASTIN Bulletin 41 (2011) 191-213) and Dhaene et al. (Journal of Computational and Applied Mathematics 263 (2014) 78-87). Some differences and relations between the new dependence measure and other multivariate measures are analyzed. We also give several characteristics of this measure and estimations based on the definitions and its property are presented.

Keywords: Comonotonicity; Product moment; Tail dependence measure; Concordance order; Estimation.

^{*}Corresponding author.

1 Introduction

Early in 1966, a pioneering paper by E. L. Lehmann [1] which gave numerous useful results in both statistical theory and application started the study of concepts of dependence for random vectors. Applications of these concepts in actuarial science have recently received increased interest. Scholars have proposed many concepts to formalize the notation of dependence existing between risks. Early sources include Lehmann [1], Esary, Proschan and Walkup [2], Esary and Proschan [3] and Kimeldorf and Sampson [4]. We are grateful that Mari and Kotz [5] pointed out the concept of correlation introduced by Galton in 1885 dominated statistics until the 1970s serving as practically the only measure of dependence. This often resulted in somewhat misleading conclusions, for reasons becomes clear in the later literatures.

The following problem which can't be avoided is how to measure a dependent random vector in practice. A risk measure is defined as a mapping from a set of random variables or vectors representing the risks at hand to the real numbers. Scarsini [6] defined certain desirable properties for a measure of association between two random variables (see also Definition 5.1.7 and Theorem 5.1.8 in Nelsen [7]) and coined the name concordance measures for those satisfying these conditions. There are a lots of ways to discuss and to measure dependence. First and foremost is Pearson's correlation coefficient [8] which captures the linear dependence between couples of random variables, but which is not invariant under monotone transformations of coordinate axes. Then, Kendall's tau and Spearman's rho are proposed to measure a form of dependence known as concordance, which are scale-invariant. The rank correlation coefficients fulfill all the desirable properties provided X_1 and X_2 are continuous. After that, Kaas et al. [9] and Dhaene et al. [10] defined bounds in convex order sense for the multivariate dependence measure, calculated by means of comonotonic vectors. Recently Koch and Schepper [11] proposed a comonotonicity coefficient for arbitrary *m*-dimensional vectors to measure the degree of comonotonicity, they also explained why in the case of cash flows the convex bounds reveal such effective approximations in Kaas et al. [9] and Dhaene et al. [10].

The impetus for writing this paper came while reading Dhaene et al. [12], which defined a multivariate dependence measure for aggregating risks as follows.

$$\rho_C(\mathbf{X}) = \frac{\operatorname{Var}(S) - \operatorname{Var}(S^{\perp})}{\operatorname{Var}(S^C) - \operatorname{Var}(S^{\perp})} = \frac{\sum_{i=1}^m \sum_{j < i} \operatorname{Cov}(X_i, X_j)}{\sum_{i=1}^m \sum_{j < i} \operatorname{Cov}(X_i^C, X_j^C)}$$

provided the covariances exist. In this definition, without any doubt, the calculation is largely simplified. However, the dependence between each pair of components is considered while ignoring the dependence among all the components. Especially in the case that all pairs of components are uncorrelated, the dependence measures are always 0. So in this paper we introduce a new multivariate dependence measure ρ that takes all aspects into account. Then, we also find the new dependence measure is surprisingly similar to Koch and Schepper's comonotonicity coefficient κ in [11] when it comes to non-negative random vectors. However, the new one focus on the tail distribution function rather than on the distribution function. Of course, the new measure is also different from other multivariate dependence measures in e.g. [11]-[17] when the dimension $m \geq 3$. When m = 2we can relate our new dependence measure to the classical ones.

The paper is organised as follows. In Section 2 we recall some definitions and notions on distributions and copulas, and we repeat the most common dependence measures. Section 3 is the most important part, we introduce our new dependence measure and discuss its major properties. Special attentions are given to the relationships between the new dependence measure and Koch and Schepper's comonotonicity coefficient [11] and Dhaene et al.'s dependence measure [12] in Section 4. Afterwards in Section 5 we give some comments on the estimation.

2 Preliminaries

First of all, we briefly review the Fréchet space in this section, which is an essential concept when investigating the dependence structure in multivariate vectors.

Definition 2.1. (Fréchet space) Let $F_1, F_2, ..., F_m$ be univariate df s. The Fréchet space $\Re_m(F_1, F_2, ..., F_m)$ consists of all the m-dimensional (df s F_X of) random vectors X possessing $F_1, F_2, ..., F_m$ as marginal dfs, that is,

$$F_i(x) = P(X_i \le x), \ x \in \mathbb{R}, \ i = 1, 2, ..., m.$$

Define the Fréchet upper bound as

$$W_m(\mathbf{x}) = \min \{F_1(x_1), F_2(x_2), ..., F_m(x_m)\}, \ \mathbf{x} \in \mathbb{R}^m,$$

and the Fréchet lower bound as

$$M_m(\mathbf{x}) = \max \{ \sum_{i=1}^m F_i(x_i) - n + 1, 0 \}, \ \mathbf{x} \in \mathbb{R}^m.$$

Then the inequality

$$M_m(\mathbf{x}) \le F_{\mathbf{X}}(\mathbf{x}) \le W_m(\mathbf{x})$$

holds for all $\mathbf{x} \in \mathbb{R}^m$ and $\mathbf{X} \in \Re_m(F_1, F_2, ..., F_m)$.

An important concept of positive dependence in a multivariate context is positive orthant dependence.

Definition 2.2. (POD) The random vector $\mathbf{X} = (X_1, X_2, ..., X_m)$ is said to be positively lower orthant dependence (PLOD) if

$$P(X_1 \le x_1, X_2 \le x_2, ..., X_m \le x_m) \ge \prod_{i=1}^m P(X_i \le x_i), \ \forall (x_1, x_2, ..., x_m) \in \mathbb{R}^m,$$
(2.1)

and it is said to be positively upper orthant dependence (PUOD) if

$$P(X_1 > x_1, X_2 > x_2, ..., X_m > x_m) \ge \prod_{i=1}^m P(X_i > x_i), \ \forall (x_1, x_2, ..., x_m) \in \mathbb{R}^m.$$
(2.2)

The random vector $\mathbf{X} = (X_1, X_2, ..., X_m)$ is said to be positively orthant dependence (POD) when it's both PLOD and PUOD.

Comonotonic vector and independent vector are two important and at the same time extreme elements of a Fréchet space.

Definition 2.3. (Comonotonic vector and independent vector) For every Fréchet space $\Re_m(F_1, F_2, ..., F_m)$, we define the independent vector $\mathbf{X}^I = (X_1^I, X_2^I, ..., X_m^I)$ as the vector with distribution $F_{\mathbf{X}}^I(x_1, x_2, ..., x_m) = \prod_{i=1}^m F_i(x_i)$, and the comonotonic vector $\mathbf{X}^C = (X_1^C, X_2^C, ..., X_m^C)$ as the vector with distribution $F_{\mathbf{X}}^C(x_1, x_2, ..., x_m) = \min_{i=1}^m F_i(x_i)$.

In Kaas et al. [18] (Section 10.6), an alternative characterization for comonotonic random vector is discussed. A random vector \mathbf{X} is comontonic if and only if there exist an rv Z and non-decreasing functions $t_1, t_2, ..., t_m$ such that

$$\mathbf{X} =_d (t_1(Z), t_2(Z), ..., t_m(Z))'.$$

Note that with the inverse function defined as $F_i^{-1}(p) = \inf\{x \in \mathbb{R} | F_i(x) \ge p\}, p \in [0, 1],$ the comonotonic vector can be easily constructed as

$$\mathbf{X}^{C} =_{d} \left(F_{1}^{-1}(U), F_{2}^{-1}(U), ..., F_{m}^{-1}(U) \right), \ U \sim U[0, 1].$$
(2.3)

In order to construct the equivalent expression of our new dependence measure, we also need to introduce the concept of copula. A copula is a joint distribution function with uniform margins. One of the most fundamental results about copulas is summarized in the following lemma.

Lemma 2.1. (Sklar's Theorem) Every df F_X of a random vector $X \in \Re_m(F_1, F_2, ..., F_m)$ can be represented as

$$F_{\mathbf{X}}(\mathbf{x}) = C(F_1(x_1), F_2(x_2), ..., F_m(x_m)), \ \mathbf{x} \in \mathbb{R}^m,$$
(2.4)

in terms of a copula C. If the marginals $F_1, F_2, ..., F_m$ are continuous then the copula involved in (2.4) is unique and explicitly given by

$$C(\boldsymbol{u}) = F_{\boldsymbol{X}}(F_1^{-1}(u_1), F_2^{-1}(u_2), ..., F_m^{-1}(u_m)), \ \boldsymbol{u} \in [0, 1]^m$$

In order to compare our new dependence measure with the common ones, we recall them here; see e.g. [11, 12, 16, 17, 19].

(i) Peareson correlation coefficient

$$\gamma_p(X,Y) = \frac{\operatorname{Cov}(X,Y)}{\sqrt{\operatorname{Var}(X)\operatorname{Var}(Y)}};$$

(ii) Kendall's rank correlation coefficient

$$\tau(X,Y) = 4 \int_0^1 \int_0^1 C(u_1, u_2) dC(u_1, u_2) - 1;$$

(iii) Spearman's rank correlation coefficient

$$\rho_s(X,Y) = 12 \int_0^1 \int_0^1 (C(u_1, u_2) - u_1 u_2) du_1 du_2$$

(iv) Gini's correlation coefficient

$$G(X,Y) = 2\int_0^1 \int_0^1 (|u_1 + u_2 - 1| - |u_1 - u_2|) dC(u_1, u_2);$$

(v) Blomqvist's correlation coefficient

$$\beta(X,Y) = 4F_{\mathbf{X}}(F_1^{-1}(1/2), F_2^{-1}(1/2)) - 1;$$

(vi) Koch and Schepper's comonotonicity coefficient

$$\kappa(\mathbf{X}) = \frac{\int \cdots \int (F_{\mathbf{X}}(\mathbf{x}) - F_{\mathbf{X}}^{I}(\mathbf{x})) d\mathbf{x}}{\int \cdots \int (F_{\mathbf{X}}^{C}(\mathbf{x}) - F_{\mathbf{X}}^{I}(\mathbf{x})) d\mathbf{x}};$$

(vii) Dhaene et al.'s multivariate dependence measure for aggregating risks

$$\rho_C(\mathbf{X}) = \frac{\operatorname{Var}(S) - \operatorname{Var}(S^{\perp})}{\operatorname{Var}(S^C) - \operatorname{Var}(S^{\perp})} = \frac{\sum_{i=1}^m \sum_{j < i} \operatorname{Cov}(X_i, X_j)}{\sum_{i=1}^m \sum_{j < i} \operatorname{Cov}(X_i^C, X_j^C)}.$$

3 A new dependence measure

Definition 3.1. The dependence measure $\rho(\mathbf{X})$ of a random vector \mathbf{X} with non-degenerate margins is defined as

$$\rho(\mathbf{X}) = \frac{E[\prod_{i=1}^{m} X_i] - \prod_{i=1}^{m} E[X_i]}{E[\prod_{i=1}^{m} X_i^C] - \prod_{i=1}^{m} E[X_i]}$$
(3.1)

provided the expectations exist.

When it comes to non-negative random vectors, the new dependence measure has a equivalent form of the ratio of two hypervolumes.

Proposition 3.1. Let $F_{\mathbf{X}}^C$ and $F_{\mathbf{X}}^I$ be the joint distributions of the comonotonic and the independent vector of the Fréchet space $\Re_m(F_1, F_2, ..., F_m)$ and $\bar{F}_{\mathbf{X}}^C$ and $\bar{F}_{\mathbf{X}}^I$ be the joint tail distributions, respectively. For any non-negative random vector \mathbf{X} of $\Re_m(F_1, F_2, ..., F_m)$ with joint tail distribution $\bar{F}_{\mathbf{X}}$, the dependence measure $\rho(\mathbf{X})$ can be defined as:

$$\rho(\mathbf{X}) = \frac{\int \cdots \int \left(\bar{F}_{\mathbf{X}}(\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}}{\int \cdots \int \left(\bar{F}_{\mathbf{X}}^{C}(\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}}$$
(3.2)

where the integrations are performed over the whole domain of X.

Proof. As we all know, the product moment of the components of an m-dimensional non-negative random vector \mathbf{X} can be written as

$$E[\prod_{i=1}^{m} X_i] = \int_{x_1=0}^{+\infty} \cdots \int_{x_m=0}^{+\infty} \bar{F}_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$

which concludes the proof.

Remark 3.1. In the case of continuous distribution functions, there exists a unique survival copula $\bar{C}_{\mathbf{X}}$ for which

$$\bar{F}_{\mathbf{X}}(x_1,...,x_m) = \bar{C}_{\mathbf{X}}(\bar{F}_1(x_1),...,\bar{F}_m(x_m)).$$

Inserting this into the formula for $\rho(\mathbf{X})$ in (3.2) results in the following equivalent expression for the new dependence measure:

$$\rho(\mathbf{X}) = \frac{\int_0^1 \cdots \int_0^1 (\bar{C}_{\mathbf{X}}(1-u_1, \dots, 1-u_m) - \prod_{i=1}^m (1-u_i)) dF_1^{-1}(u_1) \cdots dF_m^{-1}(u_m)}{\int_0^1 \cdots \int_0^1 (\bar{C}_{\mathbf{X}^C}(1-u_1, \dots, 1-u_m) - \prod_{i=1}^m (1-u_i)) dF_1^{-1}(u_1) \cdots dF_m^{-1}(u_m)}.$$
 (3.3)

Actually, the multivariate distributions $\bar{C}_{\mathbf{X}}(1-u_1,...,1-u_m)$ and $C_{\mathbf{X}}(u_1,...,u_m)$ are different in general.

Our new dependence measure has several interesting properties. For instance, it satisfies the axioms of normalization, monotonicity, permutation invariance and duality in Taylor ([15]) just as most dependence measures.

Definition 3.2. A random vector $\mathbf{X} = (X_1, X_2, ..., X_m)$ is said to be smaller than the random vector $\mathbf{Y} = (Y_1, Y_2, ..., Y_m)$ in the concordance order, written as $\mathbf{X} \leq_c \mathbf{Y}$, if both

$$P(X_1 \le t_1, X_2 \le t_2, ..., X_m \le t_m) \le P(Y_1 \le t_1, Y_2 \le t_2, ..., Y_m \le t_m)$$

and

$$P(X_1 > t_1, X_2 > t_2, ..., X_m > t_m) \le P(Y_1 > t_1, Y_2 > t_2, ..., Y_m > t_m)$$

hold for all $(t_1, t_2, ..., t_m) \in \mathbb{R}^m$.

Proposition 3.2. For any two random vectors $\mathbf{X} = (X_1, X_2, ..., X_m)$ and $\mathbf{Y} = (Y_1, Y_2, ..., Y_m)$, ρ has the following properities.

(i) (normalization) If **X** has comonotonic components, then $\rho(\mathbf{X}) = 1$; if **X** has independent components, then $\rho(\mathbf{X}) = 0$.

(ii) (monotonicity) If **X** is smaller than **Y** in the concordance order, then $\rho(\mathbf{X}) \leq \rho(\mathbf{Y})$.

(iii) (permutation invariance) For any permutation $(i_1, i_2, ..., i_m)$ of (1, 2, ..., m), we have that $\rho(X_{i_1}, X_{i_2}, ..., X_{i_m}) = \rho(X_1, X_2, ..., X_m)$.

(iv) (duality) $\rho(-X_1, -X_2, ..., -X_m) = \rho(X_1, X_2, ..., X_m).$

Proof. The proofs of (i) and (iii) are straightforward.

(ii) Any random vectors that are ordered in concordance order obviously have the same marginal distributions. Consequently, $X_i =_d Y_i$ and $\prod_{i=1}^m E[X_i] = \prod_{i=1}^m E[Y_i]$, i = 1, 2, ..., m. Therefore, for any comonotonic vectors $\mathbf{X}^C = (X_1^C, X_2^C, ..., X_m^C)$ and $\mathbf{Y}^C = (Y_1^C, Y_2^C, ..., Y_m^C)$, we have $F_{\mathbf{X}^C}(x_1, x_2, ..., x_m) = F_{\mathbf{Y}^C}(x_1, x_2, ..., x_m)$. So $\rho(\mathbf{X})$ and $\rho(\mathbf{Y})$ have the same denominators. On the other hand, $\mathbf{X} \leq_c \mathbf{Y}$ implies that $F_{\mathbf{X}}(x_1, x_2, ..., x_m) \leq F_{\mathbf{Y}}(x_1, x_2, ..., x_m)$, which concludes the proof.

(iv) Obviously, $E[\prod_{i=1}^{m}(-X_i)] = (-1)^m E[\prod_{i=1}^{m} X_i]$. For the comonotic vector we find

$$\begin{split} E[\prod_{i=1}^{m} (-X_i)^C] &= E[\prod_{i=1}^{m} F_{-X_i}^{-1}(U)], \ U \sim U(0,1) \\ &= E[\prod_{i=1}^{m} (-F_{X_i}^{-1}(1-U))] \\ &= E[\prod_{i=1}^{m} -F_{X_i}^{-1}(V)], \ V \sim U(0,1) \\ &= (-1)^m E[\prod_{i=1}^{m} X_i^C]. \end{split}$$

Hence,

$$\rho(-X_1, -X_2, ..., -X_m) = \frac{E[\prod_{i=1}^m (-X_i)] - \prod_{i=1}^m E[-X_i]}{E[\prod_{i=1}^m (-X_i)^C] - \prod_{i=1}^m E[X_i]} \\
= \frac{(-1)^m E[\prod_{i=1}^m X_i] - (-1)^m \prod_{i=1}^m E[X_i]}{(-1)^m E[\prod_{i=1}^m X_i^C] - (-1)^m \prod_{i=1}^m E[X_i]} \\
= \rho(X_1, X_2, ..., X_m)$$

which concludes the proof.

Proposition 3.3. For any random vector \mathbf{X} we have $\rho(\mathbf{X}) \leq 1$. If $\rho(\mathbf{X}) = 1$, then $\mathbf{X} =_d \mathbf{X}^C$.

Proof. From the fact that

$$\begin{aligned} F_{\mathbf{X}}(y_1, y_2, ..., y_m) &\leq \min\{F_{X_1}(y_1), F_{X_2}(y_2), ..., F_{X_m}(y_m)\} \\ &= F_{\mathbf{X}}^C(y_1, y_2, ..., y_m), \; \forall \; (y_1, y_2, ..., y_m) \in \mathbb{R}^m, \end{aligned}$$

it follows that $E[\prod_{i=1}^{m} X_i] \leq E[\prod_{i=1}^{m} X_i^C]$, which implies that $\rho(\mathbf{X}) \leq 1$. If $\rho(\mathbf{X}) = 1$, then $E[\prod_{i=1}^{m} X_i] = E[\prod_{i=1}^{m} X_i^C]$ and $F_{\mathbf{X}}(x_1, ..., x_m) = F_{\mathbf{X}}^C(x_1, ..., x_m)$, so $\mathbf{X} =_d \mathbf{X}^C$.

The reverse implication for $\rho(\mathbf{X}) = 0$ does not hold in general. Indeed, one can easily construct several non-independent random vectors for which $\rho(\mathbf{X}) = 0$.

Example 3.1. Let be Y an rv taking the values 0, $\pi/2$ and π with probability 1/3 each. Then, it's easy to see that $X_1 = \sin Y$ and $X_2 = \cos Y$ are uncorrelated (i.e. $\rho(X, Y) = 0$). However, they are not independent since X_1 and X_2 are functionally connected (by the relation $X_1^2 + X_2^2 = 1$).

For a continuous counterexample, take $Z \sim U(0,1)$ and $X_1 = \sin Z$, $X_2 = \cos Z$. Then,

$$E[X_1] = E[X_2] = E[X_1X_2] = 0$$

so that X_1 and X_2 are uncorrelated but not independent since the relation $X_1^2 + X_2^2 = 1$ holds.

For the comonotonic random vector the joint tail distribution function has a similar form as the joint distribution function.

Proposition 3.4. For any comonotonic random vector $\mathbf{X}^{C} = (X_{1}^{C}, X_{2}^{C}, ..., X_{m}^{C})$, the joint tail distribution function $\bar{F}_{\mathbf{X}^{C}}(\mathbf{x}), \ \mathbf{x} \in \mathbb{R}^{m}$ has the following expression:

$$\bar{F}_{\mathbf{X}^{C}}(\mathbf{x}) = \min\{\bar{F}_{1}(x_{1}), \bar{F}_{2}(x_{2}), ..., \bar{F}_{m}(x_{m})\}, \ \mathbf{x} \in \mathbb{R}^{m}.$$

Proof. From (2.3), for any comonotonic random vector $\mathbf{X}^{C} = (X_{1}^{C}, ..., X_{m}^{C})$, we have

$$\begin{split} \bar{F}_{\mathbf{X}^{C}}(\mathbf{x}) &= P(X_{1}^{C} > x_{1}, X_{2}^{C} > x_{2}, ..., X_{m}^{C} > x_{m}) \\ &= P(F_{1}^{-1}(U) > x_{1}, F_{2}^{-1}(U) > x_{2}, ..., F_{m}^{-1}(U) > x_{m}) \\ &= P(U > F_{1}(x_{1}), U > F_{2}(x_{2}), ..., U > F_{m}(x_{m})) \\ &= P(U > \max\{F_{1}(x_{1}), F_{2}(x_{2}), ..., F_{m}(x_{m})\}) \\ &= 1 - \max\{F_{1}(x_{1}), F_{2}(x_{2}), ..., F_{m}(x_{m})\} \\ &= \min\{\bar{F}_{1}(x_{1}), \bar{F}_{2}(x_{2}), ..., \bar{F}_{m}(x_{m})\}, \ \mathbf{x} \in \mathbb{R}^{m} \end{split}$$

which concludes the proof.

Proposition 3.5. For any non-negative PUOD random vector \mathbf{X} we have $\rho(\mathbf{X}) \geq 0$.

Proof. From the definition of PUOD we know that $\bar{F}_{\mathbf{X}}^{C}(x_{1},...,x_{m}) \geq \bar{F}_{\mathbf{X}}(x_{1},...,x_{m}) \geq \prod_{i=1}^{m} \bar{F}_{X_{i}}(x_{i})$, so the integrands in Proposition 3.1 are non-negative, which concludes the proof.

4 The relationships between the new dependence measure and the classical ones

For m = 2 we can relate ρ to Pearson correlation coefficient γ , the comonotonicity coefficient κ in Koch and Schepper [11] and ρ_c defined in Dhaene et al. [12].

Proposition 4.1. For any random couple (X, Y) we have $\rho(X, Y) = \gamma(X, Y)$ if and only if the marginal distributions differ only in location and/or scale parameters.

Proof. The couples (X, Y) and (X^C, Y^C) have the same marginals, so

$$\rho(X,Y) = \frac{E[XY] - E[X]E[Y]}{E[X^{C}Y^{C}] - E[X]E[Y]} = \frac{Cov(X,Y)}{Cov(X^{C},Y^{C})} = \frac{\gamma(X,Y)}{\gamma(X^{C},Y^{C})}$$

which equals $\gamma(X, Y)$ if and only if $\gamma(X^C, Y^C) = 1$. Hence, $Y^C = aX^C + b$ with probability 1 for some constants a > 0 and $b \in \mathbb{R}$, and thus $F_Y^{-1}(p) = aF_X^{-1}(p) + b$ from which we can conclude that the marginal distributions differ only in location and/or scale parameters.

Proposition 4.2. For any random couple (X, Y), we have $\rho(X, Y) = \rho_c(X, Y)$.

Proof. The proof is obvious.

Proposition 4.3. For any random non-negative random vector $\mathbf{X} = (X_1, X_2, ..., X_m)$, we have $\rho(\mathbf{X}) = \kappa(-\mathbf{X})$. Especially, when m = 2, $\rho(\mathbf{X}) = \kappa(\mathbf{X})$.

Proof. For any non-negative random vector **X**,

$$F_{-\mathbf{X}}(\mathbf{x}) = P(-X_1 \le x_1, -X_2 \le x_2, ..., -X_m \le x_m)$$

= $P(X_1 \ge -x_1, X_2 \ge -x_2, ..., X_m \ge -x_m)$
= $\bar{F}_{\mathbf{X}}(-\mathbf{x}),$

 \mathbf{SO}

$$\begin{split} \rho(\mathbf{X}) &= \frac{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(\bar{F}_{\mathbf{X}}(\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}}{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(\bar{F}_{\mathbf{X}}^{C}(\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}} \\ &= \frac{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(\bar{F}_{\mathbf{X}}^{C}(-\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(-\mathbf{x})\right) d\mathbf{x}}{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(\bar{F}_{\mathbf{X}}^{C}(-\mathbf{x}) - \bar{F}_{\mathbf{X}}^{I}(-\mathbf{x})\right) d\mathbf{x}} \\ &= \frac{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(F_{-\mathbf{X}}(\mathbf{x}) - F_{-\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}}{\int_{0}^{+\infty} \cdots \int_{0}^{+\infty} \left(F_{-\mathbf{X}}^{C}(\mathbf{x}) - F_{-\mathbf{X}}^{I}(\mathbf{x})\right) d\mathbf{x}} \\ &= \kappa(-\mathbf{X}). \end{split}$$

When m = 2, for any non-negative random couple (X_1, X_2) ,

$$\bar{F}_{X_1,X_2}(x_1,x_2) = 1 - F_{X_1}(x_1) - F_{X_2}(x_2) + F_{X_1,X_2}(x_1,x_2).$$

Therefore,

$$\rho(X_1, X_2) = \frac{\int_0^{+\infty} \int_0^{+\infty} \left(\bar{F}_{X_1, X_2}(x_1, x_2) - \bar{F}_{X_1}(x_1)\bar{F}_{X_2}(x_2)\right) dx_1 dx_2}{\int_0^{+\infty} \int_0^{+\infty} \left(\bar{F}_{X_1, X_2}^C(x_1, x_2) - \bar{F}_{X_1}(x_1)\bar{F}_{X_2}(x_2)\right) dx_1 dx_2} \\
= \frac{\int_0^{+\infty} \int_0^{+\infty} \left(F_{X_1, X_2}(x_1, x_2) - F_{X_1}(x_1)F_{X_2}(x_2)\right) dx_1 dx_2}{\int_0^{+\infty} \int_0^{+\infty} \left(F_{X_1, X_2}^C(x_1, x_2) - F_{X_1}(x_1)F_{X_2}(x_2)\right) dx_1 dx_2} \\
= \kappa(X_1, X_2)$$

which concludes the proof.

Remark From last formula we can write

$$\rho(X_1, X_2) = \frac{\int_0^1 \int_0^1 \left(C(u_1, u_2) - u_1 u_2 \right) dF_{X_1}^{-1}(u_1) dF_{X_2}^{-1}(u_2)}{\int_0^1 \int_0^1 \left(\min\{u_1, u_2\} - u_1 u_2 \right) dF_{X_1}^{-1}(u_1) dF_{X_2}^{-1}(u_2)}.$$

So ρ depends not only on the copula, but also on the marginals.

Example 4.1. In order to illustrate the phenomenon mentioned above, let us consider the *Farlie-Gumbel-Morgenstern* copula which is given by

$$C_{\alpha}(\mathbf{u}) = u_1 u_2 [1 + \alpha (1 - u_1)(1 - u_2)], \ \alpha \in [-1, 1].$$

It can be shown that $\rho = \frac{1}{3}\alpha$ when both marginals are U(0, 1). So the range of ρ for this family is [-1/3, 1/3] which obtains the minimum and the maximum. All other marginals will result in a measure smaller than 1/3. For example, if we insert Exp(1) marginals in the *Farlie-Gumbel-Morgenstern* copula, we have $\rho = \frac{1}{4}\alpha$.

However, the new dependence measure is different from κ in [11] and ρ_c in [12] for $m \geq 3$. The following examples can clarify it.

Example 4.2. Let's consider the multivariate *Eyraud-Gumbel-Morgenstern* function [20] for *n*-dimension

$$H(x_1, x_2, ..., x_n) = F_1(x_1) F_2(x_2) \cdots F_n(x_n) [1 + \sum_{1 \le j_1 < j_2 \le n} \alpha_{j_1 j_2} (1 - F_{j_1}(x_{j_1})) (1 - F_{j_2}(x_{j_2})) + \cdots + \alpha_{12...n} (1 - F_1(x_1)) (1 - F_2(x_2)) \cdots (1 - F_n(x_n))]$$

where the coefficients $\alpha's$ are real constants. For the marginals are U(0, 1), after complicated calculation we can get the following results for n = 3

$$\rho(U_1, U_2, U_3) = \frac{E[U_1 U_2 U_3] - E[U_1]E[U_2]E[U_3]}{E[U_1^C U_2^C U_3^C] - E[U_1]E[U_2]E[U_3]} \\
= \frac{\frac{1}{8} + \frac{1}{72}(\alpha_{12} + \alpha_{23} + \alpha_{13}) - \frac{1}{216}\alpha_{123} - \frac{1}{8}}{\frac{1}{4} - \frac{1}{8}} \\
= \frac{1}{9}(\alpha_{12} + \alpha_{23} + \alpha_{13}) - \frac{1}{27}\alpha_{123}.$$

However,

$$\kappa(U_1, U_2, U_3) = \frac{\int_0^1 \int_0^1 \int_0^1 \left(F_{U}(x_1, x_2, x_3) - F_{U_1}(x_1) F_{U_2}(x_2) F_{U_3}(x_3) \right) dx_1 dx_2 dx_3}{\int_0^1 \int_0^1 \int_0^1 \left(\min\{F_{U_1}(x_1), F_{U_2}(x_2), F_{U_3}(x_3) \right) - F_{U_1}(x_1) F_{U_2}(x_2) F_{U_3}(x_3) \right) dx_1 dx_2 dx_3} \\
= \frac{\frac{1}{72} (\alpha_{12} + \alpha_{23} + \alpha_{13}) + \frac{1}{216} \alpha_{123}}{\frac{1}{8}} \\
= \frac{1}{9} (\alpha_{12} + \alpha_{23} + \alpha_{13}) + \frac{1}{27} \alpha_{123}.$$

Example 4.3. Suppose non-negative random vector **X** obeys a multivariate Pareto distribution of the second kind introduced by Alexandru et al. [21]. If $\mathbf{X} \sim Par_m(II)(\boldsymbol{\mu}, \boldsymbol{\sigma}, \boldsymbol{\alpha}, \alpha_0)$, then the decumulative distribution function is

$$\bar{F}_{\mathbf{X}}(\mathbf{x}) = (1 + \max_{j=1}^{m} \frac{x_j - \mu_j}{\sigma_j})^{-\alpha_0} \prod_{j=1}^{m} (1 + \frac{x_j - \mu_j}{\sigma_j})^{-\alpha_j}, \ x_j > \mu_j, \ j = 1, 2, ..., m,$$

where μ_j are real and σ_j , α_j , α_0 are positive constants. The marginal distribution is

$$\bar{F}_{X_j}(x) = (1 + \frac{x_j - \mu_j}{\sigma_j})^{-\alpha_j}, \ x_j > \mu_j, \ \sigma_j > 0, \ \alpha_j > 0,$$

which is a heavy tail distribution. In addition, setting $\alpha_0 = 0$ yields a probabilistic model having independent Pareto distributed marginals. In order to simplify our calculation, we set $\mathbf{X} \sim Par_m(II)(\mathbf{0}, \mathbf{1}, \boldsymbol{\alpha}, \alpha_0)$, where $\boldsymbol{\alpha} = (\alpha, ..., \alpha)$, i.e. the marginal distributions are same.

After complicated calculation, we can get the results when m = 3, $\alpha > 3$ as follows.

$$E[X_j] = \frac{1}{\alpha_0 + \alpha - 1}, \ j = 1, 2, 3,$$
$$E[X_1 X_2 X_3] = \frac{6}{(\alpha_0 + \alpha - 1)(\alpha_0 + 2\alpha - 2)(\alpha_0 + 3\alpha - 3)},$$
$$E[X_1^C X_2^C X_3^C] = \frac{1}{(\alpha - 1)(\alpha - 2)(\alpha - 3)}.$$

Therefore, we have

$$\rho(X_1, X_2, X_3) = \frac{E[X_1 X_2 X_3] - E[X_1]E[X_2]E[X_3]}{E[X_1^C X_2^C X_3^C] - E[X_1]E[X_2]E[X_3]} \\
= \frac{\frac{6}{(\alpha_0 + \alpha - 1)(\alpha_0 + 2\alpha - 2)(\alpha_0 + 3\alpha - 3)} - \frac{1}{(\alpha_0 + \alpha - 1)^3}}{\frac{1}{(\alpha - 1)(\alpha - 2)(\alpha - 3)} - \frac{1}{(\alpha_0 + \alpha - 1)^3}}.$$

For the covariances, we get the following results.

$$Cov(X_i, X_j) = \frac{\alpha_0}{(\alpha_0 + \alpha - 1)^2(\alpha_0 + 2\alpha - 2)}, \ i, j = 1, 2, 3,$$

$$Cov(X_i^C, X_j^C) = \frac{\alpha_0 + \alpha}{(\alpha_0 + \alpha - 1)^2(\alpha_0 + \alpha - 2)}, \ i, j = 1, 2, 3.$$

 So

$$\rho_C(X_1, X_2, X_3) = \frac{\sum_{i=1}^3 \sum_{j < i} \operatorname{Cov}(X_i, X_j)}{\sum_{i=1}^3 \sum_{j < i} \operatorname{Cov}(X_i^C, X_j^C)}$$
$$= \frac{\alpha_0(\alpha_0 + \alpha - 2)}{(\alpha_0 + \alpha)(\alpha_0 + 2\alpha - 2)}.$$



Figure 1: $\rho(X_1, X_2, X_3)$ (the red) and $\rho_C(X_1, X_2, X_3)$ (the green) for $(X_1, X_2, X_3) \sim Par_3(II)(\mathbf{0}, \mathbf{1}, (\alpha, \alpha, \alpha), \alpha_0), \alpha > 3.$

We can give the figure of ρ and ρ_C in Figure 1. From the figure, it's not difficult to know that this multivariate Pareto distribution can't be used to describe the comonotonic case no matter what α_0 and α are.

Example 4.4. Suppose that risks

$$\mathbf{X} = (X_1, X_2, \dots, X_m)' \sim Nor((\mu_1, \mu_2, \dots, \mu_m)', \Sigma),$$
$$\Sigma = \begin{pmatrix} \sigma_1^2 & \sigma_{12} & \cdots & \sigma_{13} \\ \sigma_{12} & \sigma_2^2 & \cdots & \sigma_{23} \\ \vdots & \vdots & \vdots \\ \sigma_{1m} & \sigma_{2m} & \cdots & \sigma_m^2 \end{pmatrix},$$

then

$$\mathbf{X}^{C} = (X_{1}^{C}, X_{2}^{C}, ..., X_{m}^{C})' \sim Nor((\mu_{1}, \mu_{2}, ..., \mu_{m})', \Sigma^{C}),$$
$$\Sigma^{C} = \begin{pmatrix} \sigma_{1}^{2} & \sigma_{1}\sigma_{2} & \cdots & \sigma_{1}\sigma_{m} \\ \sigma_{1}\sigma_{2} & \sigma_{2}^{2} & \cdots & \sigma_{2}\sigma_{m} \\ \vdots & \vdots & \vdots \\ \sigma_{1}\sigma_{m} & \sigma_{2}\sigma_{m} & \cdots & \sigma_{m}^{2} \end{pmatrix},$$

where $\sigma_{ij} = \rho_{ij}\sigma_i\sigma_j$, i, j = 1, 2, ..., m.

In order to calculate the product moments easily, we can refer to the characteristic function

$$\varphi_{\mathbf{X}}(\mathbf{t}) = e^{i\mathbf{t}'\boldsymbol{\mu} - \frac{1}{2}\mathbf{t}'\boldsymbol{\Sigma}\mathbf{t}}$$

for $\mathbf{X} \sim Nor(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $\mathbf{t} \in \mathbb{R}^m$.

As is well known, for any random vector \mathbf{X} , the characteristic function $\varphi_{\mathbf{X}}(\mathbf{t})$ always exists. So suppose that for a random vector \mathbf{X} the expectation $E[\prod_{i=1}^{m} X_i^{r_k}]$ exists for some set of non-negative integers $r_1, r_2, ..., r_m$. Then this expectation can be found from the relation

$$E[\prod_{i=1}^{m} X_{i}^{r_{k}}] = \frac{1}{i^{r_{1}+r_{2}+\dots+r_{m}}} \left(\frac{\partial^{r_{1}+r_{2}+\dots+r_{m}}}{\partial^{r_{1}} t_{1} \partial^{r_{2}} t_{2} \cdots \partial^{r_{m}} t_{m}} \varphi_{\mathbf{X}}(\mathbf{t}) \right) |_{\mathbf{t}=\mathbf{0}}$$

where $\mathbf{0} = (0, 0, ..., 0)'$.

In this example we can get the following characteristic functions for ${\bf X}$ and ${\bf X}^C$

$$\begin{split} \varphi_{\mathbf{X}}(\mathbf{t}) &= e^{i\sum_{i=1}^{m}\mu_{i}t_{i}-\frac{1}{2}\sum_{i=1}^{m}\sum_{j=1}^{m}\rho_{ij}\sigma_{i}\sigma_{j}t_{i}t_{j}}, \\ \varphi_{\mathbf{X}}c(\mathbf{t}) &= e^{i\sum_{i=1}^{m}\mu_{i}t_{i}-\frac{1}{2}\sum_{i=1}^{m}\sum_{j=1}^{m}\sigma_{i}\sigma_{j}t_{i}t_{j}}. \end{split}$$

Differentiate the characteristic function with respect to \mathbf{t} and let $\mathbf{t} = \mathbf{0}$, we can get the following product moments:

$$E[\prod_{i=1}^{m} X_{i}] = (-i)^{m} \frac{\partial^{m} \varphi_{\mathbf{X}}(\mathbf{t})}{\partial t_{1} \partial t_{2} \dots \partial t_{m}} |_{\mathbf{t}=\mathbf{0}}$$

$$= \prod_{k=1}^{m} \mu_{j_{k}} + \sum_{S(m,1)} \sigma_{j_{1}j_{2}} \prod_{k=3}^{m} \mu_{j_{k}} + \sum_{S(m,2)} \sigma_{j_{1}j_{2}} \sigma_{j_{3}j_{4}} \prod_{k=5}^{m} \mu_{j_{k}} + \cdots$$

$$+ \sum_{S(m,\frac{2m-5+(-1)^{m}}{4})} \sigma_{j_{1}j_{2}} \cdots \sigma_{j_{A}j_{B}} \prod_{k=C}^{m} \mu_{j_{k}} + \sum_{S(m,\frac{2m-3+(-1)^{m}}{4})} \sigma_{j_{1}j_{2}} \cdots \sigma_{j_{D}j_{E}} \prod_{k=F}^{m} \mu_{j_{k}}$$

where $A = m - 4 + \frac{1 + (-1)^m}{2}$, $B = m - 3 + \frac{1 + (-1)^m}{2}$, $C = D = m - 2 + \frac{1 + (-1)^m}{2}$, $E = m - 1 + \frac{1 + (-1)^m}{2}$, $F = \frac{[m - (-1)^m m]}{2}$, $S(m, k) = \frac{m!}{2^k k! (m-k)!}$ and $\sum_{S(m,k)}$ means the sum of S(m,k) cases; see [22].

Therefore,

$$\rho_{C}(\mathbf{X}) = \frac{\sum_{i=1}^{m} \sum_{j < i} \operatorname{Cov}(X_{i}, X_{j})}{\sum_{i=1}^{m} \sum_{j < i} \operatorname{Cov}(X_{i}^{C}, X_{j}^{C})} = \frac{\sum_{i=1}^{m} \sum_{j < i} \rho_{ij} \sigma_{i} \sigma_{j}}{\sum_{i=1}^{m} \sum_{j < i} \sigma_{i} \sigma_{j}}$$
$$\rho(\mathbf{X}) = \frac{E[\prod_{i=1}^{m} X_{i}] - \prod_{i=1}^{m} E[X_{i}]}{E[\prod_{i=1}^{m} X_{i}^{C}] - \prod_{i=1}^{m} E[X_{i}]} = \frac{I}{J}$$

where

$$I = \sum_{S(m,1)} \rho_{j_1 j_2} \sigma_{j_1} \sigma_{j_2} \prod_{k=3}^m \mu_{j_k} + \sum_{S(m,2)} \rho_{j_1 j_2} \sigma_{j_1} \sigma_{j_2} \rho_{j_3 j_4} \sigma_{j_3} \sigma_{j_4} \prod_{k=5}^m \mu_{j_k} + \cdots$$

+
$$\sum_{S(m, \frac{2m-5+(-1)^m}{4})} \rho_{j_1 j_2} \sigma_{j_1} \sigma_{j_2} \cdots \rho_{j_A j_B} \sigma_{j_A} \sigma_{j_B} \prod_{k=C}^m \mu_{j_k}$$

+
$$\sum_{S(m, \frac{2m-3+(-1)^m}{4})} \rho_{j_1 j_2} \sigma_{j_1} \sigma_{j_2} \cdots \rho_{j_D j_E} \sigma_{j_D} \sigma_{j_E} \prod_{k=F}^m \mu_{j_k},$$

$$J = \sum_{S(m,1)} \sigma_{j_1} \sigma_{j_2} \prod_{k=3}^{m} \mu_{j_k} + \sum_{S(m,2)} \sigma_{j_1} \sigma_{j_2} \sigma_{j_3} \sigma_{j_4} \prod_{k=5}^{m} \mu_{j_k} + \cdots$$
$$+ \sum_{S(m,\frac{2m-5+(-1)^m}{4})} \sigma_{j_1} \sigma_{j_2} \cdots \sigma_{j_A j_B} \prod_{k=C}^{m} \mu_{j_k} + \sum_{S(m,\frac{2m-3+(-1)^m}{4})} \sigma_{j_1} \sigma_{j_2} \cdots \sigma_{j_D j_E} \prod_{k=F}^{m} \mu_{j_k},$$

and $A = m - 4 + \frac{1 + (-1)^m}{2}, B = m - 3 + \frac{1 + (-1)^m}{2}, C = D = m - 2 + \frac{1 + (-1)^m}{2}, E = m - 1 + \frac{1 + (-1)^m}{2}, F = \frac{[m - (-1)^m m]}{2}, S(m, k) = \frac{m!}{2^{kk!(m-k)!}}.$

Remark 4.1. It's obvious that $\rho(\mathbf{X})$ not only depends on σ_{ij} but also on μ_j while $\rho_C(\mathbf{X})$ only relys on σ_{ij} , i, j = 1, 2, ..., m. Especially, when m = 3 and $\mu_1 = \mu_2 = \mu_3$,

$$\rho(\mathbf{X}) = \frac{\rho_{12}\sigma_1\sigma_2 + \rho_{13}\sigma_1\sigma_3 + \rho_{23}\sigma_2\sigma_3}{\sigma_1\sigma_2 + \sigma_1\sigma_3 + \sigma_2\sigma_3} = \rho_C(\mathbf{X}).$$

5 Estimation

In this section we give some comments on the estimation of ρ from a sample of **X**. Inference on $\rho(\mathbf{X})$ in Definition 3.1 boils down to inference on the expectations $E[\prod_{i=1}^{m} X_i]$, $E[\prod_{i=1}^{m} X_i^C]$ and $\prod_{i=1}^{m} E[X_i]$. These can be estimated by moments estimation. This is of great significance in a real-life data set when the product moment is difficult to find.

Suppose we want to estimate the new dependence measure for m variables, for which we have n coupled observations, not necessarily independent, summarized in a data matrix Y (dimension $n \times m$).

So we can get the the estimations

$$\widehat{E}[\prod_{j=1}^{m} X_j] = \frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} Y_{ij}$$

$$\widehat{E}[X_j] = \frac{1}{n} \sum_{i=1}^n Y_{ij}.$$

For the estimation of $E[\prod_{j=1}^{m} X_{j}^{C}]$, we need a sample of \mathbf{X}^{C} . Dhaene et al.[8] show that for any \mathbf{x} and \mathbf{y} in the range of a comonotonic vector either $\mathbf{x} \leq \mathbf{y}$ or $\mathbf{x} \geq \mathbf{y}$ holds. In other words, all possible outcomes of \mathbf{X}^{C} are ordered componentwise. As \mathbf{X} and \mathbf{X}^{C} also have the same marginal distributions, we can easily turn the sample of \mathbf{X} into a sample of \mathbf{X}^{C} . Denoting the *i*th order statistic of X_{j} by $Y_{j}^{(i)}$, we find the following sample of \mathbf{X}^{C} : $(Y_{1}^{(i)}, ..., Y_{m}^{(i)}), i = 1, 2, ..., n$. Accordingly, we can get the product moment estimation

$$\widehat{E}[\prod_{j=1}^{m} X_{j}^{C}] = \frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} Y_{j}^{(i)}.$$

Summarizing, we find the following estimator for ρ :

$$\widehat{\rho}(\mathbf{X}) = \frac{\frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} Y_{ij} - \prod_{j=1}^{m} \frac{1}{n} \sum_{i=1}^{n} Y_{ij}}{\frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} Y_{j}^{(i)} - \prod_{j=1}^{m} \frac{1}{n} \sum_{i=1}^{n} Y_{ij}}.$$

When the random vectors are non-negative, the estimation has another form. Actually, the empirical tail distribution can be written as

$$\hat{F}_{\mathbf{X}}(x_1, x_2, \dots, x_m) = P(X_1 > x_1, X_2 > x_2, \dots, X_m > x_m) = \frac{1}{n} \sum_{i=1}^n \prod_{j=1}^m I(Y_{ij} > x_j);$$

for the empirical versions of the distribution for independent and comonotonic vectors of the same $Fr\acute{e}$ chet space, we have

$$\hat{\bar{F}}_{\mathbf{X}}^{I}(x_{1}, x_{2}, ..., x_{m}) = \prod_{j=1}^{m} \hat{\bar{F}}_{j}(x_{j}) = \prod_{j=1}^{m} (\frac{1}{n} \sum_{i=1}^{n} I(Y_{ij} > x_{j})),$$
$$\hat{\bar{F}}_{\mathbf{X}}^{C}(x_{1}, x_{2}, ..., x_{m}) = P(X_{1}^{C} > x_{1}, X_{2}^{C} > x_{2}, ..., X_{m}^{C} > x_{m}) = \frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} I(Y_{j}^{(i)} > x_{j})$$

Now, define $m_j = \min_{i=1}^n Y_{ij}$ and $M_j = \max_{i=1}^n Y_{ij}$. As a consequence, $Y_j^{(1)} = m_j$ and $Y_j^{(n)} = M_j$ for any j = 1, 2, ..., m, and $\int_{m_j}^{M_j} I(Y_{ij} > x_j) dx_j = Y_{ij} - m_j$ for any $i = 1, 2, ..., n, \ j = 1, 2, ..., m$.

If we replace the tail distribution functions in Proposition 3.1 by their empirical versions, we get the estimator for ρ :

$$\widehat{\rho}(\mathbf{X}) = \frac{\frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} (Y_{ij} - m_j) - \frac{1}{n^m} \prod_{j=1}^{m} (\sum_{i=1}^{n} Y_{ij} - nm_j)}{\frac{1}{n} \sum_{i=1}^{n} \prod_{j=1}^{m} (Y_j^{(i)} - m_j) - \frac{1}{n^m} \prod_{j=1}^{m} (\sum_{i=1}^{n} Y_{ij} - nm_j)}.$$

Acknowledgements. The research was supported by the National Natural Science Foundation of China (No. 11171179), the Research Fund for the Doctoral Program of Higher Education of China (No. 20133705110002) and the Program for Scientific Research Innovation Team in Colleges and Universities of Shandong Province.

References

- E.L. Lehmann, Some concepts of dependence, The Annals of Mathematical Statistics, 37 (5) (1966) 1137-1153.
- [2] J.D. Esary, F. Proschan, D.W. Walkup, Association of random variables, with applications, The Annals of Mathematical Statistics, 38 (5) (1967) 1466-1474.
- J.D. Esary, F. Proschan, Relationships among some concepts of bivariate dependence, The Annals of Mathematical Statistics, 43 (2) (1972) 651-655.
- [4] G. Kimeldorf, A.R. Sampson, Monotone dependence, The Annals of Statistics, 6 (4) (1978) 895-903.
- [5] D.D. Mari, S. Kotz, Correlation and Dependence, Imperial College Press, London, 2001.
- [6] M. Scarsini, On measures of concordance, Stochastica, 8 (3) (1984) 201-218.
- [7] R. B. Nelsen, An Introduction to Copulas, Springer, New York, 1999.
- [8] M. Denuit, P. Lambert, Constraints on concordance measures in bivariate discrete data, Journal of Multivariate Analysis, 93 (1) (2005) 40-57.
- [9] R. Kaas, J. Dhaene, M.J. Goovaerts, Upper and lower bounds for sums of random variables, Insurance: Mathematics and Economics, 27 (2) (2000) 151-168.
- [10] J. Dhaene, M. Denuit, M.J. Goovaerts, R. Kaas, D. Vyncke, The concept of comonotonicity in actuarial science and finance: theory, Insurance: Mathematics and Economics, 31 (1) (2002) 3-33.
- [11] I. Koch, D.A. Schepper, Measuring comonotonicity in *m*-dimensional vectors, ASTIN Bulletin, 41 (1) (2011) 191-213.
- [12] J. Dhaene, D. Linders, W. Schoutens, D. Vyncke, A multivariate dependence measure for aggregating risks, Journal of Computational and Applied Mathematics, 263 (2014) 78-87.
- [13] E.F. Wolff, N-dimensional measures of dependence, Stochastica 4 (3) (1980) 175-188.
- [14] B.F. Fernndez, J.M. Gonzlez-Barrios, Multidimensional dependency measures, Journal of Multivariate Analysis, 89 (2) (2004) 351-370.

- [15] M.D. Taylor, Multivariate measures of concordance, Annals of the Institute of Statistical Mathematics, 59 (4) (2007) 789-806.
- [16] J. Behboodian, A. Dolati, M. Úbeda-Flores, A multivariate version of Gini's rank association coefficient, Statistical Papers, 48 (2) (2007) 295-304.
- [17] F. Schmid, R. Schmidt, Multivariate extensions of Spearman's rho and related statistics, Statistics & Probability Letters, 77 (4) (2007) 407-416.
- [18] R. Kaas, M. Goovaerts, J. Dhaene, M. Denuit, Modern Actuarial Risk Theory, Kluwer Academic Publishers, Boston, 2001.
- [19] M. Denuit, J. Dhaene, M. Goovaerts, R. Kaas, Actuarial Theory for Dependent Risks: Measures, Orders and Models, John Wiley & Sons, 2006.
- [20] S. Cambanis, Some properties and generalizations of multivariate Eyraud-Gumbel-Morgenstern distributions, Journal of Multivariate Analysis, 7 (4) (1977) 551-559.
- [21] A.V. Asimit, E. Furman, R. Vernic, On a multivariate Pareto distribution, Insurance: Mathematics and Economics, 46 (2) (2010) 308-316.
- [22] Y. Tang, Q. Meng, High order moment of multivariate normal random vector, Journal of North China Institute of Technology, 20 (1) (1999) 45-48.