

Recurrence Relations of the Multi-Indexed Orthogonal Polynomials : II

Satoru Odake

Department of Physics, Shinshu University,
Matsumoto 390-8621, Japan

Abstract

In a previous paper we presented $3+2M$ term recurrence relations with variable dependent coefficients for M -indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types. In this paper we present (conjectures of) the recurrence relations with constant coefficients for these multi-indexed orthogonal polynomials. The simplest recurrence relations have $3+2\ell$ terms, where $\ell (\geq M)$ is the degree of the lowest member of the multi-indexed orthogonal polynomials.

1 Introduction

Exactly solvable quantum mechanical systems in one dimension have seen remarkable developments in recent years and the central role is played by exceptional orthogonal polynomials [1]–[30] (and the references therein). A set of polynomials $\{\mathcal{P}_n(\eta) | n \in \mathbb{Z}_{\geq 0}\}$ is called exceptional orthogonal polynomials, when the following conditions (i)–(iii) plus (iv) are satisfied; (i) they are orthogonal with respect to some inner product, (ii) there are missing degrees, i.e., $\{\deg \mathcal{P}_n | n \in \mathbb{Z}_{\geq 0}\} \subsetneq \mathbb{Z}_{\geq 0}$, (iii) but they form a complete set, and (iv) they satisfy second order differential or difference equations. The constraints of Bochner's theorem and its generalizations [31, 32] are avoided by the condition (ii). We want to distinguish the following two cases; the set of missing degrees $\mathcal{I} = \mathbb{Z}_{\geq 0} \setminus \{\deg \mathcal{P}_n | n \in \mathbb{Z}_{\geq 0}\}$ is case (1): $\mathcal{I} = \{0, 1, \dots, \ell - 1\}$, case (2) $\mathcal{I} \neq \{0, 1, \dots, \ell - 1\}$, where ℓ is a positive integer. The situation of case (1) is called stable in [12]. The first example of the case (1) exceptional orthogonal polynomials, X_1 Laguerre and Jacobi, was found by Gómez-Ullate, Kamran and Milson [1], and its quantum mechanical formulation was given by Quesne [2]. Based on the quantum mechanical

formulation (ordinary quantum mechanics (oQM), discrete quantum mechanics with pure imaginary shifts (idQM)[16]), Sasaki and the present author constructed X_ℓ polynomials and their generalizations, multi-indexed orthogonal polynomials [4, 5, 17, 23]. The multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types, which are obtained by multi-step Darboux transformations [33, 34, 35, 3, 9] with virtual state wavefunctions as seed solutions [17, 23], correspond to the case (1). The exceptional orthogonal polynomials, which are obtained by multi-step Darboux transformations with eigenstate or pseudo virtual state wavefunctions as seed solutions [25, 26, 9, 29], correspond to the case (2). For those having purely discrete orthogonality weight functions, the number of orthogonal polynomials may be finite. The multi-indexed (q -)Racah polynomials [13, 21], which are constructed based on discrete quantum mechanics with real shifts (rdQM) [16], correspond to the case (1).

The ordinary orthogonal polynomials $\{P_n(\eta)|n \in \mathbb{Z}_{\geq 0}, \deg P_n = n\}$ satisfy the three term recurrence relations, $\eta P_n(\eta) = A_n P_{n+1}(\eta) + B_n P_n(\eta) + C_n P_{n-1}(\eta)$ (A_n, B_n, C_n : constants), and conversely the polynomials satisfying the three term recurrence relations are orthogonal polynomials (Favard's theorem [32]). Since the exceptional orthogonal polynomials are not ordinary orthogonal polynomials, they do not satisfy the three term recurrence relations. In a previous paper [27], we showed that M -indexed orthogonal polynomials $P_{\mathcal{D},n}(\eta)$ of Laguerre, Jacobi, Wilson and Askey-Wilson types satisfy $3 + 2M$ term recurrence relations ($\mathcal{D} = \{d_1, \dots, d_M\}$),

$$R_{n,0}^{[M]}(\eta)P_{\mathcal{D},n}(\eta) = - \sum_{\substack{k=-M-1 \\ k \neq 0}}^{M+1} R_{n,k}^{[M]}(\eta)P_{\mathcal{D},n+k}(\eta), \quad (1.1)$$

where $R_{n,k}^{[M]}(\eta)$'s are polynomials of degree $M + 1 - |k|$ in η . In contrast to the three term recurrence relations, the coefficients of (1.1) are not constants. The three term recurrence relations are used to study bispectral properties or dual polynomials [32, 36], in which the constant coefficients of the recurrence relations are important. To study bispectral properties, recurrence relations with constant coefficients are desired,

$$X(\eta)P_{\mathcal{D},n}(\eta) = \sum_{k=-L}^L r_{n,k}^{X,\mathcal{D}} P_{\mathcal{D},n+k}(\eta) \quad (\forall n \in \mathbb{Z}_{\geq 0}), \quad (1.2)$$

where $r_{n,k}^{X,\mathcal{D}}$'s are constants and $X(\eta)$ is some polynomial of degree L in η . Such recurrence relations for $M = 1$ case were first given by Sasaki, Tsujimoto and Zhedanov [11]. They found

$1 + 4\ell$ term recurrence relations. Recently Miki and Tsujimoto found different recurrence relations with $3 + 2\ell$ terms [37]. Their choices of $X(\eta)$ are $\Xi_\ell(\eta)^2$ and $\int_0^\eta \Xi_\ell(y)dy$, respectively. Durán studied recurrence relations with constant coefficients for several exceptional orthogonal polynomials including multi-indexed Laguerre polynomials by using duality [38].

In this paper we present infinitely many (conjectures of) recurrence relations with constant coefficients for M -indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types, namely we present infinitely many polynomials $X(\eta)$ leading to (1.2). The minimal degree of $X(\eta)$ is (conjectured as) $\ell_{\mathcal{D}} + 1$, where $\ell_{\mathcal{D}}$ is the degree of the lowest member multi-indexed orthogonal polynomial $P_{\mathcal{D},0}(\eta)$, and this gives $3 + 2\ell_{\mathcal{D}}$ term recurrence relations with constant coefficients.

This paper is organized as follows. In section 2 we recapitulate some fundamental formulas of the multi-indexed orthogonal polynomials and present a method deriving recurrence relations with constant coefficients. Section 3 is the main part of the paper. After discussing a necessary condition for $X(\eta)$, we present (conjectures of) recurrence relations with constant coefficients for the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types, Conjecture 1 and Conjecture 2. The final section is for a summary and comments. Some useful formulas of the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types are listed in Appendix.

2 Method

In this section we explain an idea for deriving the recurrence relations with constant coefficients. We follow the notation of [27]. The virtual state wavefunction $\tilde{\phi}(x)$ is characterized by the degree v and the type t (I or II), like $\tilde{\phi}_v^t(x)$. For simplicity, we suppress type t in many places.

The fundamental formulas of the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types are found in [27]. Among them we recall that

$$\hat{\mathcal{A}}_{d_1 \dots d_s} \phi_{d_1 \dots d_{s-1} n}(x) = \phi_{d_1 \dots d_s n}(x), \quad \hat{\mathcal{A}}_{d_1 \dots d_s}^\dagger \phi_{d_1 \dots d_s n}(x) = (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_s}) \phi_{d_1 \dots d_{s-1} n}(x), \quad (2.1)$$

$$\phi_{d_1 \dots d_s n}(x) = \Psi_{d_1 \dots d_s}(x) P_{d_1 \dots d_s, n}(\eta(x)) \quad (n \in \mathbb{Z}_{\geq 0}), \quad P_{d_1 \dots d_s, n}(\eta) \stackrel{\text{def}}{=} 0 \quad (n < 0), \quad (2.2)$$

$$\deg P_{d_1 \dots d_s, n}(\eta) = \ell_{d_1 \dots d_s} + n, \quad \deg \Xi_{d_1 \dots d_s}(\eta) = \ell_{d_1 \dots d_s}, \quad d_j > 0,$$

$$\ell_{d_1 \dots d_s} = \sum_{j=1}^s d_j - \frac{1}{2}s(s-1) + 2s_{\text{I}}s_{\text{II}}, \quad s_t = \#\{d_j | d_j: \text{type } t, j = 1, \dots, s\} \quad (t = \text{I, II}), \quad (2.3)$$

$$\begin{aligned}
(\phi_{d_1 \dots d_s n}, \phi_{d_1 \dots d_s m}) &= h_{d_1 \dots d_s n} \delta_{nm}, \quad h_{d_1 \dots d_s n} = \prod_{j=1}^s (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_j}) \cdot h_n, \\
&= (\Psi_{d_1 \dots d_s}^2 P_{d_1 \dots d_s n}, P_{d_1 \dots d_s m}).
\end{aligned} \tag{2.4}$$

The relations (2.1) are rewritten by using the forward ($\hat{\mathcal{F}}$) and backward ($\hat{\mathcal{B}}$) shift operators as

$$\hat{\mathcal{F}}_{d_1 \dots d_s} P_{d_1 \dots d_{s-1}, n}(\eta) = P_{d_1 \dots d_s, n}(\eta), \quad \hat{\mathcal{B}}_{d_1 \dots d_s} P_{d_1 \dots d_s, n}(\eta) = (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_s}) P_{d_1 \dots d_{s-1}, n}(\eta). \tag{2.5}$$

Here $\hat{\mathcal{F}}_{d_1 \dots d_s}$ and $\hat{\mathcal{B}}_{d_1 \dots d_s}$ are defined by

$$\hat{\mathcal{F}}_{d_1 \dots d_s} \stackrel{\text{def}}{=} \Psi_{d_1 \dots d_s}(x)^{-1} \circ \hat{\mathcal{A}}_{d_1 \dots d_s} \circ \Psi_{d_1 \dots d_{s-1}}(x), \quad \hat{\mathcal{B}}_{d_1 \dots d_s} \stackrel{\text{def}}{=} \Psi_{d_1 \dots d_{s-1}}(x)^{-1} \circ \hat{\mathcal{A}}_{d_1 \dots d_s}^\dagger \circ \Psi_{d_1 \dots d_s}(x), \tag{2.6}$$

and their explicit forms are given in (A.1)–(A.2) and (A.8)–(A.9). This gives Rodrigues type formula, $P_{d_1 \dots d_s, n}(\eta) = \hat{\mathcal{F}}_{d_1 \dots d_s} \cdots \hat{\mathcal{F}}_{d_1 d_2} \hat{\mathcal{F}}_{d_1} P_n(\eta)$. These formulas were not presented explicitly in our previous papers. We remark that, from their explicit forms, $\hat{\mathcal{F}}_{d_1 \dots d_s}$ and $\hat{\mathcal{B}}_{d_1 \dots d_s}$ map rational functions of η to rational functions of η . For an appropriate parameter range, the Hamiltonians $\mathcal{H}_{d_1 \dots d_s}$ are non-singular and their eigenfunctions $\{\phi_{d_1 \dots d_s n}(x)\}_{n=0}^\infty$ form a complete set of the Hilbert space. For any polynomial $X(\eta)$ in η , the function $X(\eta(x))\phi_{d_1 \dots d_s n}(x)$ belongs to the Hilbert space.

The Hamiltonian $\mathcal{H}_{d_1 \dots d_s}$ does not depend on the order of d_j 's. On the other hand, the multi-indexed orthogonal polynomial $P_{d_1 \dots d_s, n}(\eta)$ changes the sign under a permutation of d_j 's, $P_{d_{\sigma_1} \dots d_{\sigma_s}, n}(\eta) = \text{sgn}(\begin{smallmatrix} 1 & \dots & s \\ \sigma_1 & \dots & \sigma_s \end{smallmatrix}) P_{d_1 \dots d_s, n}(\eta)$. The denominator polynomial $\Xi_{d_1 \dots d_s}(\eta)$ also changes the sign, $\Xi_{d_{\sigma_1} \dots d_{\sigma_s}}(\eta) = \text{sgn}(\begin{smallmatrix} 1 & \dots & s \\ \sigma_1 & \dots & \sigma_s \end{smallmatrix}) \Xi_{d_1 \dots d_s}(\eta)$. We write $\mathcal{H}_{d_1 \dots d_M}$, $\phi_{d_1 \dots d_M n}(x)$, $P_{d_1 \dots d_M, n}(\eta)$, $\Xi_{d_1 \dots d_M}(\eta)$, $h_{d_1 \dots d_M, n}$, etc. as $\mathcal{H}_{\mathcal{D}}$, $\phi_{\mathcal{D} n}(x)$, $P_{\mathcal{D}, n}(\eta)$, $\Xi_{\mathcal{D}}(\eta)$, $h_{\mathcal{D}, n}$, etc., respectively ($\mathcal{D} = \{d_1, \dots, d_M\}$).¹

First we note the following property of orthogonal polynomials.

Lemma 1 *Let us assume for a certain polynomial $X(\eta)$ of degree L in η that*

$$X(\eta) P_{\mathcal{D}, n}(\eta) = \sum_{k=-n}^L r_{n,k}^{X, \mathcal{D}} P_{\mathcal{D}, n+k}(\eta) \quad (\forall n \in \mathbb{Z}_{\geq 0}). \tag{2.7}$$

Here $r_{n,k}^{X, \mathcal{D}}$'s are constants. The sum $\sum_{k=-n}^L$ can be replaced by $\sum_{k=-L}^L$.

¹ In the Appendix of [27], we assumed the ‘standard order’ $\mathcal{D} = \{d_1^{\text{I}}, \dots, d_{M_1}^{\text{I}}, d_1^{\text{II}}, \dots, d_{M_1}^{\text{II}}\}$ for simplicity. We do not assume it in this paper.

Proof Multiplying by $\Psi_{\mathcal{D}}(x)$ to (2.7), we have

$$X(\eta)\phi_{\mathcal{D},n}(x) = \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} \phi_{\mathcal{D},n+k}(x).$$

By using (2.4) we have

$$\begin{aligned} (\phi_{\mathcal{D}m}, X\phi_{\mathcal{D}n}) &= \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} h_{\mathcal{D},m} \delta_{m,n+k} = \theta(m \leq n+L) r_{n,m-n}^{X,\mathcal{D}} h_{\mathcal{D},m} \\ &= (X\phi_{\mathcal{D}m}, \phi_{\mathcal{D}n}) = \sum_{k=-m}^L r_{m,k}^{X,\mathcal{D}} h_{\mathcal{D},n} \delta_{n,m+k} = \theta(m \geq n-L) r_{m,n-m}^{X,\mathcal{D}} h_{\mathcal{D},n}, \end{aligned} \quad (2.8)$$

where $\theta(P)$ is a step function for a proposition P , $\theta(P) = 1$ (P : true), 0 (P : false). This means $(\phi_{\mathcal{D}m}, X\phi_{\mathcal{D}n}) = 0$ unless $n-L \leq m \leq n+L$. Namely $r_{n,k}^{X,\mathcal{D}} = 0$ unless $-L \leq k \leq L$.

□

Remark Although the inner product formulas used in the proof are valid only for ‘real’ $X(\eta)$ ($X^* = X$) and an appropriate parameter range such that the Hamiltonian is non-singular, the final result, which represents the polynomial equations, is valid for any parameter values and complex $X(\eta)$.

Next we explain a method to obtain recurrence relations with constant coefficients. Let $X(\eta)$ be a polynomial of degree L in η . Since $X(\eta)\phi_{\mathcal{D}n}(x)$ belongs to the Hilbert space and $\{\phi_{\mathcal{D}n}(x)\}_{n=0}^{\infty}$ is a complete set, we have the expansion

$$X(\eta)\phi_{\mathcal{D}n}(x) = \sum_{k=-n}^{\infty} r_{n,k}^{X,\mathcal{D}} \phi_{\mathcal{D},n+k}(x), \quad (2.9)$$

where $r_{n,k}^{X,\mathcal{D}}$ ’s are constants. By using this and (2.4), we obtain

$$(\phi_{\mathcal{D}m}, X\phi_{\mathcal{D}n}) = \sum_{k=-n}^{\infty} r_{n,k}^{X,\mathcal{D}} h_{\mathcal{D},m} \delta_{m,n+k} = r_{n,m-n}^{X,\mathcal{D}} h_{\mathcal{D},m}. \quad (2.10)$$

On the other hand, by using (2.1)–(2.2) and (2.6), we obtain

$$\begin{aligned} &(\phi_{\mathcal{D}m}, X\phi_{\mathcal{D}n}) \\ &= (\hat{\mathcal{A}}_{d_1 \dots d_M} \cdots \hat{\mathcal{A}}_{d_1 d_2} \hat{\mathcal{A}}_{d_1} \phi_m, X\phi_{\mathcal{D}n}) \\ &= (\phi_m, \hat{\mathcal{A}}_{d_1}^\dagger \hat{\mathcal{A}}_{d_1 d_2}^\dagger \cdots \hat{\mathcal{A}}_{d_1 \dots d_M}^\dagger (X\phi_{\mathcal{D}n})) \\ &= (\phi_0 P_m, (\phi_0 \hat{\mathcal{B}}_{d_1} \Psi_{d_1}^{-1}) (\Psi_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \Psi_{d_1 d_2}^{-1}) \cdots (\Psi_{d_1 \dots d_{M-1}} \hat{\mathcal{B}}_{d_1 \dots d_M} \Psi_{d_1 \dots d_M}^{-1}) (\Psi_{d_1 \dots d_M} X P_{\mathcal{D},n})) \end{aligned}$$

$$\begin{aligned}
&= (\phi_0 P_m, \phi_0 \hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n})) \\
&= (\phi_0^2 P_m, \hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n})). \tag{2.11}
\end{aligned}$$

From the property of $\hat{\mathcal{B}}_{d_1 \dots d_s}$, the function $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n})$ is a rational function of η . When it is not a polynomial in η , we have infinitely many m such that $(\phi_{\mathcal{D}m}, X \phi_{\mathcal{D}n}) \neq 0$, namely r.h.s. of (2.9) is an infinite sum. Let us consider the case that $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n})$ is a polynomial of degree $n + L'$ in η . Since any polynomial in η can be expanded in $P_n(\eta)$, we have

$$\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n}) = \sum_{k=-n}^{L'} r_{n,k}^{(0)X,\mathcal{D}} P_{n+k}(\eta), \tag{2.12}$$

where $r_{n,k}^{(0)X,\mathcal{D}}$'s are constants. Substituting this to (2.11), we obtain

$$(\phi_{\mathcal{D}m}, X \phi_{\mathcal{D}n}) = \sum_{k=-n}^{L'} r_{n,k}^{(0)X,\mathcal{D}} h_m \delta_{m,n+k}, = \theta(m \leq L' + n) r_{n,m-n}^{(0)X,\mathcal{D}} h_m. \tag{2.13}$$

Eqs.(2.10) and (2.13) imply

$$r_{n,k}^{X,\mathcal{D}} = 0 \quad (k > L'), \quad r_{n,k}^{X,\mathcal{D}} h_{\mathcal{D},n+k} = r_{n,k}^{(0)X,\mathcal{D}} h_{n+k} \quad (-n \leq k \leq L'). \tag{2.14}$$

Thus we obtain

$$X(\eta) \phi_{\mathcal{D}n}(x) = \sum_{k=-n}^{L'} r_{n,k}^{X,\mathcal{D}} \phi_{\mathcal{D}n+k}(x), \tag{2.15}$$

namely,

$$X(\eta) P_{\mathcal{D},n}(\eta) = \sum_{k=-n}^{L'} r_{n,k}^{X,\mathcal{D}} P_{\mathcal{D},n+k}(\eta). \tag{2.16}$$

By comparing the degree of both sides, we have $L' = L$. By Lemma 1, the sum $\sum_{k=-n}^L$ can be replaced by $\sum_{k=-L}^L$.

We summarize this argument as the following proposition.

Proposition 1 *Let us assume for a certain polynomial $X(\eta)$ of degree L in η that the function $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} (X P_{\mathcal{D},n})$ is a polynomial in η . Expand it as (2.12). We have $1 + 2L$ term recurrence relations with constant coefficients for $P_{\mathcal{D},n}(\eta)$:*

$$X(\eta) P_{\mathcal{D},n}(\eta) = \sum_{k=-L}^L r_{n,k}^{X,\mathcal{D}} P_{\mathcal{D},n+k}(\eta) \quad (\forall n \in \mathbb{Z}_{\geq 0}), \quad r_{n,k}^{X,\mathcal{D}} = \frac{r_{n,k}^{(0)X,\mathcal{D}}}{\prod_{j=1}^M (\mathcal{E}_{n+k} - \tilde{\mathcal{E}}_{d_j})}. \tag{2.17}$$

Remark 1 See Remark below Lemma 1.

Remark 2 Under the assumption of this proposition, the functions $\hat{\mathcal{B}}_{d_1 \dots d_j} \hat{\mathcal{B}}_{d_1 \dots d_j d_{j+1}} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M}$ ($XP_{\mathcal{D},n}$) ($j = 2, \dots, M$) are also polynomials in η .

Remark 3 The function $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M}(XP_{\mathcal{D},n})$ is rewritten as

$$\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M}(XP_{\mathcal{D},n}) = (\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} X \hat{\mathcal{F}}_{d_1 \dots d_M} \cdots \hat{\mathcal{F}}_{d_1 d_2} \hat{\mathcal{F}}_{d_1}) P_n. \quad (2.18)$$

This operator $\hat{\mathcal{B}}_{d_1} \cdots X \cdots \hat{\mathcal{F}}_{d_1}$ maps polynomials in η to rational functions of η . To find a proper polynomial $X(\eta)$ giving recurrence relations with constant coefficients is rephrased as follows; Find a polynomial $X(\eta)$ such that the operator $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \cdots \hat{\mathcal{B}}_{d_1 \dots d_M} X \hat{\mathcal{F}}_{d_1 \dots d_M} \cdots \hat{\mathcal{F}}_{d_1 d_2} \hat{\mathcal{F}}_{d_1}$ maps polynomials in η to polynomials in η .

3 Recurrence Relations with Constant Coefficients

In this section we present recurrence relations with constant coefficients (1.2) for the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types.

3.1 Multi-indexed Laguerre and Jacobi polynomials

In this subsection we discuss the recurrence relations with constant coefficients for the multi-indexed Laguerre and Jacobi polynomials. We note that the first order differential operator of the form $a(x) \frac{d}{dx} + b(x)$ ($a(x), b(x)$: functions of x) acts on the product of two functions $f(x)$ and $g(x)$ as

$$\left(a(x) \frac{d}{dx} + b(x) \right) (f(x)g(x)) = f(x) \left(a(x) \frac{d}{dx} + b(x) \right) g(x) + a(x) \frac{df(x)}{dx} g(x). \quad (3.1)$$

First we consider a necessary condition for $X(\eta)$ giving recurrence relations with constant coefficients. Let us assume (2.7) for a polynomial $X(\eta)$ of degree L in η . Applying $\hat{\mathcal{B}}_{\mathcal{D}} = \hat{\mathcal{B}}_{d_1 \dots d_M}$ (A.2) to (2.7), we have

$$\begin{aligned} \hat{\mathcal{B}}_{\mathcal{D}}(X(\eta)P_{\mathcal{D},n}(\eta)) &= \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} \hat{\mathcal{B}}_{\mathcal{D}} P_{\mathcal{D},n+k}(\eta) = \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} (\mathcal{E}_{n+k} - \tilde{\mathcal{E}}_{d_M}) P_{d_1 \dots d_{M-1},n}(\eta) \\ &= X(\eta) \hat{\mathcal{B}}_{\mathcal{D}} P_{\mathcal{D},n}(\eta) - c_{\mathcal{F}}^2 e_{\mathcal{D}}^{\hat{\mathcal{B}}}(\eta) \frac{\Xi_{d_1 \dots d_{M-1}}(\eta)}{\Xi_{\mathcal{D}}(\eta)} \frac{dX(\eta)}{d\eta} P_{\mathcal{D},n}(\eta) \\ &= (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_M}) X(\eta) P_{d_1 \dots d_{M-1},n}(\eta) - c_{\mathcal{F}}^2 e_{\mathcal{D}}^{\hat{\mathcal{B}}}(\eta) \frac{\Xi_{d_1 \dots d_{M-1}}(\eta)}{\Xi_{\mathcal{D}}(\eta)} \frac{dX(\eta)}{d\eta} P_{\mathcal{D},n}(\eta), \end{aligned} \quad (3.2)$$

where (3.1) and (2.5) are used. Since the expression in the first line is a polynomial in η , the expression in the last line should be so. The denominator polynomial $\Xi_{\mathcal{D}}(\eta)$ does not have common roots with $e_{\mathcal{D}}^{\hat{\mathcal{B}}}(\eta)$ and $P_{\mathcal{D},n}(\eta)$ for some n . Therefore, if $\Xi_{\mathcal{D}}(\eta) = \Xi_{d_1 \dots d_M}(\eta)$ does not have common roots with $\Xi_{d_1 \dots d_{M-1}}(\eta)$, the polynomial $\frac{dX(\eta)}{d\eta}$ should be divisible by $\Xi_{\mathcal{D}}(\eta)$.

We summarize this argument as follows.

Proposition 2 *Let $X(\eta)$ be a polynomial of degree L in η . Assume (2.7) and*

$$\frac{dX(\eta)}{d\eta} = \Xi_{\mathcal{D}}(\eta)Y(\eta), \quad Y(\eta) : \text{a polynomial in } \eta. \quad (3.3)$$

Then one action of $\hat{\mathcal{B}}_{\mathcal{D}}$ to the both sides of (2.7) keeps the polynomiality intact.

Remark If two polynomials in η , $\Xi_{\mathcal{D}}(\eta) = \Xi_{d_1 \dots d_M}(\eta)$ and $\Xi_{d_1 \dots d_{M-1}}(\eta)$, do not have common roots, the polynomial of degree L in η , $X(\eta)$, satisfying (2.7) should satisfy (3.3) for some polynomial $Y(\eta)$.

The overall normalization and the constant term of $X(\eta)$ are not important, because the change of the former induces that of the overall normalization of $r_{n,k}^{X,\mathcal{D}}$ and the shift of the latter induces that of $r_{n,0}^{X,\mathcal{D}}$. By taking the constant term of $X(\eta)$ as $X(0) = 0$, the condition for the candidate of $X(\eta)$ (3.3) gives

$$X(\eta) = \int_0^\eta \Xi_{\mathcal{D}}(y)Y(y)dy, \quad \deg X(\eta) = L = \ell_{\mathcal{D}} + \deg Y(\eta) + 1. \quad (3.4)$$

The minimal degree candidate of $X(\eta)$, which corresponds to $Y(\eta) = 1$, is

$$X_{\min}(\eta) = \int_0^\eta \Xi_{\mathcal{D}}(y)dy, \quad \deg X_{\min}(\eta) = \ell_{\mathcal{D}} + 1. \quad (3.5)$$

Based on these properties we present our main result. After one action of $\hat{\mathcal{B}}_{\mathcal{D}}$ to (2.7), further actions of $\hat{\mathcal{B}}_{d_1 \dots d_{M-1}}, \hat{\mathcal{B}}_{d_1 \dots d_{M-2}}, \dots$ give more conditions for $X(\eta)$. However, it seems that these additional conditions are satisfied automatically by the original condition (3.3) and by the properties of $P_{d_1 \dots d_s, n}(\eta)$ and $\Xi_{d_1 \dots d_s}(\eta)$, for example, see the proof of $M = 2$ case in Remark 5 below. We conjecture that this candidate $X(\eta)$ (3.4) actually gives recurrence relations with constant coefficients.

Conjecture 1 *For any polynomial $Y(\eta)$, we take $X(\eta)$ as (3.4). Then the multi-indexed Laguerre and Jacobi polynomials $P_{\mathcal{D},n}(\eta)$ satisfy $1 + 2L$ term recurrence relations with constant coefficients (1.2).*

Remark 1 If two polynomials in η , $\Xi_{\mathcal{D}}(\eta) = \Xi_{d_1 \dots d_M}(\eta)$ and $\Xi_{d_1 \dots d_{M-1}}(\eta)$, do not have common roots, this conjecture exhausts all possible $X(\eta)$ giving recurrence relations with constant coefficients, and the minimal degree choice $X(\eta) = X_{\min}(\eta)$ (3.5) gives $3 + 2\ell_{\mathcal{D}}$ term recurrence relations.

Remark 2 The minimal degree polynomial $X_{\min}(\eta)$ can be divisible by η . The degree of $\frac{X_{\min}(\eta)}{\eta}$ is $\ell_{\mathcal{D}}$, which is the lowest degree of $P_{\mathcal{D},n}(\eta)$. The recurrence relations (1.2), $\frac{X_{\min}(\eta)}{\eta} \times \eta P_{\mathcal{D},n}(\eta) = \sum \dots$, can be regarded as a natural generalization of the three term recurrence relations of the ordinary orthogonal polynomial $P_n(\eta)$, $1 \times \eta P_n(\eta) = A_n P_{n+1}(\eta) + B_n P_n(\eta) + C_n P_{n-1}(\eta)$.

Remark 3 Since $Y(\eta)$ is arbitrary, we obtain infinitely many recurrence relations. However not all of them are independent. For ‘ $M = 0$ case’ (namely, ordinary orthogonal polynomials), it is trivial that recurrence relations obtained from arbitrary $Y(\eta)$ ($\deg Y \geq 1$) are derived by the three term recurrence relations.

Remark 4 For $M = 1$ case ($\mathcal{D} = \{\ell\}$), the minimal degree choice $X(\eta) = X_{\min}(\eta)$, which gives $3 + 2\ell$ term recurrence relations, was given by Miki and Tsujimoto [37], and the choice $X(\eta) = \Xi_{\ell}(\eta)^2$, which corresponds to $Y(\eta) = 2\Xi_{\ell}(\eta)$ and gives $1 + 4\ell$ term recurrence relations, was given by Sasaki, Tsujimoto and Zhedanov [11]. For general M , $Y(\eta) = 2\partial_{\eta}\Xi_{\mathcal{D}}(\eta)p(\eta) + \Xi_{\mathcal{D}}(\eta)\partial_{\eta}p(\eta)$, where $p(\eta)$ is any polynomial in η , gives $X(\eta) = \Xi_{\mathcal{D}}(\eta)^2 p(\eta)$.

Remark 5 Direct verification of this conjecture is rather straightforward for lower M and smaller d_j , n and $\deg Y$, by a computer algebra system, e.g. Mathematica. For $M = 1, 2$ we can prove this conjecture. Since we have Proposition 1, it is sufficient to show that $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \dots \hat{\mathcal{B}}_{d_1 \dots d_M}(X P_{\mathcal{D},n})$ is a polynomial in η .

$M = 1$ From (3.2) we have

$$\hat{\mathcal{B}}_{d_1}(X(\eta)P_{d_1,n}(\eta)) = (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_1})X(\eta)P_n(\eta) - c_{\mathcal{F}}^2 e_{d_1}^{\hat{\mathcal{B}}}(\eta)Y(\eta)P_{d_1,n}(\eta).$$

This is a polynomial in η . Thus $M = 1$ case is proved. □

$M = 2$ From (3.2) and (A.2) we have

$$\begin{aligned} & \hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2}(X(\eta)P_{d_1 d_2, n}(\eta)) \\ &= \hat{\mathcal{B}}_{d_1}((\mathcal{E}_n - \tilde{\mathcal{E}}_{d_2})X(\eta)P_{d_1, n}(\eta) - c_{\mathcal{F}}^2 e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta)\Xi_{d_1}(\eta)Y(\eta)P_{d_1 d_2, n}(\eta)) \\ &= (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_2})\left((\mathcal{E}_n - \tilde{\mathcal{E}}_{d_1})X(\eta)P_n(\eta) - c_{\mathcal{F}}^2 \frac{e_{d_1}^{\hat{\mathcal{B}}}(\eta)}{\Xi_{d_1}(\eta)}\Xi_{d_1 d_2}(\eta)Y(\eta)P_{d_1, n}(\eta)\right) \\ & \quad + c_{\mathcal{F}}^4 e_{d_1}^{\hat{\mathcal{B}}}(\eta)\partial_{\eta}(e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta)Y(\eta)P_{d_1 d_2, n}(\eta)) + c_{\mathcal{F}}^4 \frac{e_{d_1}^{\hat{\mathcal{B}}}(\eta)}{\Xi_{d_1}(\eta)}e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta)\partial_{\eta}\Xi_{d_1}(\eta)Y(\eta)P_{d_1 d_2, n}(\eta) \end{aligned}$$

$$- c_{\mathcal{F}}^3 \tilde{e}_{d_1}^{\hat{\mathcal{B}}} e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta) Y(\eta) P_{d_1 d_2, n}(\eta).$$

By using (A.6) with $s = 2$, this becomes

$$\begin{aligned} &= (\mathcal{E}_n - \tilde{\mathcal{E}}_{d_2})(\mathcal{E}_n - \tilde{\mathcal{E}}_{d_1})X(\eta)P_n(\eta) \\ &\quad + c_{\mathcal{F}}^4 e_{d_1}^{\hat{\mathcal{B}}}(\eta) \partial_{\eta}(e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta)Y(\eta)P_{d_1 d_2, n}(\eta)) + c_{\mathcal{F}}^4 e_{d_1}^{\hat{\mathcal{B}}}(\eta) e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta) Y(\eta) \partial_{\eta} P_{d_1 d_2, n}(\eta) \\ &\quad - c_{\mathcal{F}}^3 (\tilde{e}_{d_1}^{\hat{\mathcal{B}}} e_{d_1 d_2}^{\hat{\mathcal{B}}}(\eta) + e_{d_1}^{\hat{\mathcal{B}}}(\eta) \tilde{e}_{d_1 d_2}^{\hat{\mathcal{B}}}) Y(\eta) P_{d_1 d_2, n}(\eta). \end{aligned}$$

This is a polynomial in η . Thus $M = 2$ case is proved. \square

3.2 Multi-indexed Wilson and Askey-Wilson polynomials

In this subsection we discuss the recurrence relations with constant coefficients for the multi-indexed Wilson and Askey-Wilson polynomials. We restrict the parameters: $\{a_1^*, a_2^*\} = \{a_1, a_2\}$ (as a set) and $\{a_3^*, a_4^*\} = \{a_3, a_4\}$ (as a set).

The sinusoidal coordinate $\eta(x)$ is $\eta(x) = x^2$ for Wilson case and $\eta(x) = \cos x$ for Askey-Wilson case. They satisfy [39]

$$\frac{\eta(x - i\frac{\gamma}{2})^{n+1} - \eta(x + i\frac{\gamma}{2})^{n+1}}{\eta(x - i\frac{\gamma}{2}) - \eta(x + i\frac{\gamma}{2})} = \sum_{k=0}^n g_n'^{(k)} \eta(x)^{n-k} \quad (n \in \mathbb{Z}_{\geq 0}), \quad (3.6)$$

where $g_n'^{(k)}$ is given by [40]

$$\begin{aligned} \eta(x) = x^2 : \quad g_n'^{(k)} &= \frac{(-1)^k}{2^{2k+1}} \binom{2n+2}{2k+1}, \\ \eta(x) = \cos x : \quad g_n'^{(k)} &= \theta(k : \text{even}) \frac{(n+1)!}{2^k} \sum_{r=0}^{\frac{k}{2}} \binom{n-k+r}{r} \frac{(-1)^r \llbracket n-k+1+2r \rrbracket'}{(\frac{k}{2}-r)! (n-\frac{k}{2}+1+r)!}, \\ \llbracket n \rrbracket' &\stackrel{\text{def}}{=} \frac{e^{-\frac{\gamma}{2}n} - e^{\frac{\gamma}{2}n}}{e^{-\frac{\gamma}{2}} - e^{\frac{\gamma}{2}}}. \end{aligned} \quad (3.7)$$

For a polynomial $p(\eta)$ in η , when it is regarded as a function of x , we denote it by adding a check,

$$\check{p}(x) \stackrel{\text{def}}{=} p(\eta(x)). \quad (3.8)$$

Since $\eta(x - i\frac{m}{2}\gamma) + \eta(x + i\frac{m}{2}\gamma)$ and $\eta(x - i\frac{m}{2}\gamma)\eta(x + i\frac{m}{2}\gamma)$ ($m \in \mathbb{Z}$) are expressed as polynomials in $\eta(x)$, any symmetric polynomial in $\eta(x - i\frac{m}{2}\gamma)$ and $\eta(x + i\frac{m}{2}\gamma)$ is expressed as a polynomial in $\eta(x)$ [39, 27]. For example, the followings are polynomials in $\eta(x)$ (p, p_1, p_2 : polynomials in η):

$$\check{p}(x - i\frac{\gamma}{2}) + \check{p}(x + i\frac{\gamma}{2}), \quad \frac{\check{p}(x - i\frac{\gamma}{2}) - \check{p}(x + i\frac{\gamma}{2})}{\eta(x - i\frac{\gamma}{2}) - \eta(x + i\frac{\gamma}{2})},$$

$$\begin{aligned}
& \check{p}(x - i\frac{\gamma}{2})\check{p}(x + i\frac{\gamma}{2}), \quad \frac{\check{p}_1(x - i\gamma)\check{p}_2(x - i\frac{\gamma}{2}) - \check{p}_1(x + i\gamma)\check{p}_2(x + i\frac{\gamma}{2})}{\eta(x - i\frac{\gamma}{2}) - \eta(x + i\frac{\gamma}{2})}, \\
\eta(x) = x^2 : & \quad x\check{p}_1(x + i\frac{\gamma}{2})\check{p}_2(x - i\frac{\gamma}{2}) + x\check{p}_1(x - i\frac{\gamma}{2})\check{p}_2(x + i\frac{\gamma}{2}), \\
\eta(x) = \cos x : & \quad e^{\pm i x}\check{p}_1(x + i\frac{\gamma}{2})\check{p}_2(x - i\frac{\gamma}{2}) + e^{\mp i x}\check{p}_1(x - i\frac{\gamma}{2})\check{p}_2(x + i\frac{\gamma}{2}). \tag{3.9}
\end{aligned}$$

For a polynomial $p(\eta)$ in η , let us define a polynomial in η , $I[p](\eta)$, as follows:

$$p(\eta) = \sum_{k=0}^n a_k \eta^k \mapsto I[p](\eta) = \sum_{k=0}^{n+1} b_k \eta^k, \tag{3.10}$$

where b_k 's are defined by

$$b_{k+1} = \frac{1}{g_k'} \left(a_k - \sum_{j=k+1}^n g_j'^{(j-k)} b_{j+1} \right) \quad (k = n, n-1, \dots, 1, 0), \quad b_0 = 0. \tag{3.11}$$

The constant term of $I[p](\eta)$ is chosen to be zero. It is easy to show that this polynomial $I[p](\eta) = P(\eta)$ satisfies

$$\frac{\check{P}(x - i\frac{\gamma}{2}) - \check{P}(x + i\frac{\gamma}{2})}{\eta(x - i\frac{\gamma}{2}) - \eta(x + i\frac{\gamma}{2})} = \check{p}(x). \tag{3.12}$$

The operator of the form $a(x)e^{\frac{\gamma}{2}p} - b(x)e^{-\frac{\gamma}{2}p}$ ($a(x), b(x)$: functions of x) acts on the product of two functions $f(x)$ and $g(z)$ as

$$\begin{aligned}
& (a(x)e^{\frac{\gamma}{2}p} - b(x)e^{-\frac{\gamma}{2}p})(f(x)g(x)) \\
& = a(x)f(x - i\frac{\gamma}{2})g(x - i\frac{\gamma}{2}) - b(x)f(x + i\frac{\gamma}{2})g(x + i\frac{\gamma}{2}) \\
& = f^{(+)}(x)(a(x)g(x - i\frac{\gamma}{2}) - b(x)g(x + i\frac{\gamma}{2})) - if^{(-)}(x)(a(x)g(x - i\frac{\gamma}{2}) + b(x)g(x + i\frac{\gamma}{2})) \\
& = f^{(+)}(x)(a(x)e^{\frac{\gamma}{2}p} - b(x)e^{-\frac{\gamma}{2}p})g(x) - if^{(-)}(x)(a(x)g(x - i\frac{\gamma}{2}) + b(x)g(x + i\frac{\gamma}{2})), \tag{3.13}
\end{aligned}$$

where $f^{(\pm)}(x)$ are defined by [27]

$$f^{(+)}(x) \stackrel{\text{def}}{=} \frac{1}{2}(f(x - i\frac{\gamma}{2}) + f(x + i\frac{\gamma}{2})), \quad f^{(-)}(x) \stackrel{\text{def}}{=} \frac{i}{2}(f(x - i\frac{\gamma}{2}) - f(x + i\frac{\gamma}{2})). \tag{3.14}$$

The auxiliary function $\varphi(x)$ is rewritten as

$$\varphi(x) = ic_\varphi(\eta(x - i\frac{\gamma}{2}) - \eta(x + i\frac{\gamma}{2})), \quad c_\varphi = \begin{cases} 1 & : \text{W} \\ (\sinh \frac{-\gamma}{2})^{-1} & : \text{AW} \end{cases}. \tag{3.15}$$

First we consider a necessary condition for $X(\eta)$ giving recurrence relations with constant coefficients. Let us assume (2.7) for a polynomial $X(\eta)$ of degree L in η . Applying $\hat{\mathcal{B}}_{\mathcal{D}} = \hat{\mathcal{B}}_{d_1 \dots d_M}$ (A.9) to (2.7), we have

$$\hat{\mathcal{B}}_{\mathcal{D}}(\check{X}(x)\check{P}_{\mathcal{D},n}(x)) = \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} \hat{\mathcal{B}}_{\mathcal{D}}\check{P}_{\mathcal{D},n+k}(x) = \sum_{k=-n}^L r_{n,k}^{X,\mathcal{D}} (\mathcal{E}_{n+k} - \tilde{\mathcal{E}}_{d_M}) \check{P}_{d_1 \dots d_{M-1},n}(\eta)$$

$$\begin{aligned}
&= \check{X}^{(+)}(x)\hat{\mathcal{B}}_{\mathcal{D}}\check{P}_{\mathcal{D},n}(x) - i\check{X}^{(-)}(x)\left(\frac{ic_{\mathcal{D}}^{\hat{\mathcal{B}}}}{\varphi(x)\check{\Xi}_{\mathcal{D}}(x)}e_{\mathcal{D}}^{\hat{\mathcal{B}}}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x+i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x-i\frac{\gamma}{2})\right. \\
&\quad \left. + \frac{ic_{\mathcal{D}}^{\hat{\mathcal{B}}}}{\varphi(x)\check{\Xi}_{\mathcal{D}}(x)}e_{\mathcal{D}}^{\hat{\mathcal{B}}^*}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x-i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x+i\frac{\gamma}{2})\right) \\
&= (\mathcal{E}_n - \check{\mathcal{E}}_{d_M})\check{X}^{(+)}(x)\check{P}_{d_1\dots d_{M-1},n}(x) \\
&\quad + \frac{c_{\mathcal{D}}^{\hat{\mathcal{B}}}}{\check{\Xi}_{\mathcal{D}}(x)}\frac{\check{X}^{(-)}(x)}{\varphi(x)}\left(e_{\mathcal{D}}^{\hat{\mathcal{B}}}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x+i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x-i\frac{\gamma}{2})\right. \\
&\quad \left. + e_{\mathcal{D}}^{\hat{\mathcal{B}}^*}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x-i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x+i\frac{\gamma}{2})\right), \tag{3.16}
\end{aligned}$$

where (3.13) and (2.5) are used. Since the expression in the first line is a polynomial in η , the expression in the last line should be so. By (3.9) and (A.12), $\frac{\check{X}^{(-)}(x)}{\varphi(x)} = \frac{\check{X}(x-i\frac{\gamma}{2})-\check{X}(x+i\frac{\gamma}{2})}{2c_{\varphi}(\eta(x-i\frac{\gamma}{2})-\eta(x+i\frac{\gamma}{2}))}$ and $e_{\mathcal{D}}^{\hat{\mathcal{B}}}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x+i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x-i\frac{\gamma}{2}) + e_{\mathcal{D}}^{\hat{\mathcal{B}}^*}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x-i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x+i\frac{\gamma}{2})$ are polynomials in $\eta(x)$. If the latter is not divisible by $\check{\Xi}_{\mathcal{D}}(x)$, the former should be divisible by $\check{\Xi}_{\mathcal{D}}(x)$.

We summarize this argument as follows.

Proposition 3 *Let $X(\eta)$ be a polynomial of degree L in η . Assume (2.7) and*

$$\frac{\check{X}(x-i\frac{\gamma}{2})-\check{X}(x+i\frac{\gamma}{2})}{\eta(x-i\frac{\gamma}{2})-\eta(x+i\frac{\gamma}{2})} = \check{\Xi}_{\mathcal{D}}(x)\check{Y}(x), \quad Y(\eta) : \text{a polynomial in } \eta. \tag{3.17}$$

Then one action of $\hat{\mathcal{B}}_{\mathcal{D}}$ to the both sides of (2.7) keeps the polynomiality intact.

Remark If two polynomials in η , $\check{\Xi}_{\mathcal{D}}(x) = \check{\Xi}_{d_1\dots d_M}(x)$ and $e_{\mathcal{D}}^{\hat{\mathcal{B}}}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x+i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x-i\frac{\gamma}{2}) + e_{\mathcal{D}}^{\hat{\mathcal{B}}^*}(x)\check{\Xi}_{d_1\dots d_{M-1}}(x-i\frac{\gamma}{2})\check{P}_{\mathcal{D},n}(x+i\frac{\gamma}{2})$, have no common roots for some n , the polynomial of degree L in η , $X(\eta)$, satisfying (2.7) should satisfy (3.17) for some polynomial $Y(\eta)$.

By taking the constant term of $X(\eta)$ as $X(0) = 0$, the condition for the candidate of $X(\eta)$ (3.17) gives

$$X(\eta) = I[\Xi_{\mathcal{D}}Y](\eta), \quad \deg X(\eta) = L = \ell_{\mathcal{D}} + \deg Y(\eta) + 1. \tag{3.18}$$

The minimal degree candidate of $X(\eta)$, which corresponds to $Y(\eta) = 1$, is

$$X_{\min}(\eta) = I[\Xi_{\mathcal{D}}](\eta), \quad \deg X_{\min}(\eta) = \ell_{\mathcal{D}} + 1. \tag{3.19}$$

Based on these properties we present our another main result. Like as § 3.1, we conjecture that this candidate $X(\eta)$ (3.18) actually gives recurrence relations with constant coefficients.

Conjecture 2 For any polynomial $Y(\eta)$, we take $X(\eta)$ as (3.18). Then the multi-indexed Wilson and Askey-Wilson polynomials $P_{\mathcal{D},n}(\eta)$ satisfy $1 + 2L$ term recurrence relations with constant coefficients (1.2).

Remark 1 If two polynomials in η , $\check{\Xi}_{\mathcal{D}}(x) = \check{\Xi}_{d_1 \dots d_M}(x)$ and $e_{\mathcal{D}}^{\hat{\mathcal{B}}}(x) \check{\Xi}_{d_1 \dots d_{M-1}}(x + i\frac{\gamma}{2}) \check{P}_{\mathcal{D},n}(x - i\frac{\gamma}{2}) + e_{\mathcal{D}}^{\hat{\mathcal{B}}^*}(x) \check{\Xi}_{d_1 \dots d_{M-1}}(x - i\frac{\gamma}{2}) \check{P}_{\mathcal{D},n}(x + i\frac{\gamma}{2})$, have no common roots for some n , this conjecture exhausts all possible $X(\eta)$ giving recurrence relations with constant coefficients, and the minimal degree choice $X(\eta) = X_{\min}(\eta)$ (3.19) gives $3 + 2\ell_{\mathcal{D}}$ term recurrence relations.

Remark 2 See Remark 2 and 3 below Conjecture 1.

Remark 3 By (3.9), for any polynomial $p(\eta)$, $\check{Y}(x) = \frac{\check{\Xi}_{\mathcal{D}}(x-i\gamma)\check{p}(x-i\frac{\gamma}{2}) - \check{\Xi}_{\mathcal{D}}(x+i\gamma)\check{p}(x+i\frac{\gamma}{2})}{\eta(x-i\frac{\gamma}{2}) - \eta(x+i\frac{\gamma}{2})}$ is also a polynomial in η . This $Y(\eta)$ gives $\check{X}(x) = \check{\Xi}_{\mathcal{D}}(x - i\frac{\gamma}{2}) \check{\Xi}_{\mathcal{D}}(x + i\frac{\gamma}{2}) \check{p}(x)$, which corresponds to Remark 4 below Conjecture 1.

Remark 4 Direct verification of this conjecture is rather straightforward for lower M and smaller d_j , n and $\deg Y$, by a computer algebra system, e.g. Mathematica. For $M = 1$ we can prove this conjecture. Since we have Proposition 1, it is sufficient to show that $\hat{\mathcal{B}}_{d_1} \hat{\mathcal{B}}_{d_1 d_2} \dots \hat{\mathcal{B}}_{d_1 \dots d_M}(X P_{\mathcal{D},n})$ is a polynomial in η .

$M = 1$ From (3.16) we have

$$\begin{aligned} & \hat{\mathcal{B}}_{d_1}(\check{X}(x) \check{P}_{d_1,n}(x)) \\ &= (\mathcal{E}_n - \check{\mathcal{E}}_{d_1}) \check{X}^{(+)}(x) \check{P}_n(x) + \frac{c_{d_1}^{\hat{\mathcal{B}}}}{2c_{\varphi}} \check{Y}(x) (e_{d_1}^{\hat{\mathcal{B}}}(x) \check{P}_{d_1,n}(x - i\frac{\gamma}{2}) + e_{d_1}^{\hat{\mathcal{B}}^*}(x) \check{P}_{d_1,n}(x + i\frac{\gamma}{2})). \end{aligned}$$

This is a polynomial in η . Thus $M = 1$ case is proved. \square

4 Summary and Comments

In addition to $3+2M$ term recurrence relations with variable dependent coefficients presented in [27], we have presented (conjectures of) the recurrence relations with constant coefficients for the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types, Conjecture 1 and Conjecture 2. Since $Y(\eta)$ is arbitrary, we obtain infinitely many recurrence relations. However not all of them are independent. The most important one is the minimal degree one $X_{\min}(\eta)$ (3.5) or (3.19), which gives $3+2\ell_{\mathcal{D}}$ term recurrence relations. Here $\ell_{\mathcal{D}} (\geq M)$ is the degree of the lowest member polynomial $P_{\mathcal{D},0}(\eta)$. For this case, the coefficients $r_{n,k}^{X_{\min}, \mathcal{D}}$ may have nice forms. Both derivations given in [27] and present paper are based on multi-step Darboux transformations. Although we have discussed Laguerre, Jacobi,

Wilson and Askey-Wilson cases, the method is applicable to the multi-indexed (q -)Racah polynomials (which correspond to the case (1) $\mathcal{I} = \{0, 1, \dots, \ell - 1\}$) and various case (2) polynomials. The results in [11, 37] correspond to special cases of our results. Concerning the results in [38], we hope to clarify its relation to our results. We hope also that Conjecture 1 and 2 will be proved in the near future.

Multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types are labeled by an index set \mathcal{D} but different index sets may give the same multi-indexed orthogonal polynomials, $P_{\mathcal{D},n}(\eta; \boldsymbol{\lambda}) \propto P_{\mathcal{D}',n}(\eta; \boldsymbol{\lambda}')$ [30]. For example, $\mathcal{D}_1 = \{1^{\text{II}}, 3^{\text{II}}, 4^{\text{II}}, 5^{\text{II}}, 8^{\text{II}}\}$ with $\boldsymbol{\lambda}$, $\mathcal{D}_2 = \{1^{\text{I}}, 2^{\text{I}}, 6^{\text{I}}, 8^{\text{I}}\}$ with $\boldsymbol{\lambda} - 9\tilde{\boldsymbol{\delta}}_1$ and $\mathcal{D}_3 = \{3^{\text{I}}, 5^{\text{I}}, 2^{\text{II}}\}$ with $\boldsymbol{\lambda} - 6\tilde{\boldsymbol{\delta}}_1$ give the same multi-indexed orthogonal polynomials, $P_{\mathcal{D}_1,n}(\eta; \boldsymbol{\lambda}) \propto P_{\mathcal{D}_2,n}(\eta; \boldsymbol{\lambda} - 9\tilde{\boldsymbol{\delta}}_1) \propto P_{\mathcal{D}_3,n}(\eta; \boldsymbol{\lambda} - 6\tilde{\boldsymbol{\delta}}_1)$. The $3 + 2M$ term recurrence relations given in [27] states that these polynomials $P_{\mathcal{D}_1,n}(\eta; \boldsymbol{\lambda})$, $P_{\mathcal{D}_2,n}(\eta; \boldsymbol{\lambda} - 9\tilde{\boldsymbol{\delta}}_1)$ and $P_{\mathcal{D}_3,n}(\eta; \boldsymbol{\lambda} - 6\tilde{\boldsymbol{\delta}}_1)$ satisfy 13, 11 and 9 term recurrence relations with variable dependent coefficients, respectively. But the above equivalence implies that all of them satisfy 9 term recurrence relations with variable dependent coefficients. On the other hand, the degrees of lowest members $P_{\mathcal{D},0}(\eta)$ are $\ell_{\mathcal{D}_1} = \ell_{\mathcal{D}_2} = \ell_{\mathcal{D}_3} = 11$, and for each case the minimal degree polynomial $X_{\min}(\eta)$ gives 25 term recurrence relations with constant coefficients. The above equivalence, which gives $\Xi_{\mathcal{D}_1}(\eta; \boldsymbol{\lambda}) \propto \Xi_{\mathcal{D}_2}(\eta; \boldsymbol{\lambda} - 9\tilde{\boldsymbol{\delta}}_1) \propto \Xi_{\mathcal{D}_3}(\eta; \boldsymbol{\lambda} - 6\tilde{\boldsymbol{\delta}}_1)$, implies that these three 25 term recurrence relations are essentially same.

The $3 + 2M$ term recurrence relations with variable dependent coefficients can be used to calculate the multi-indexed orthogonal polynomials effectively and it needs $M + 1$ initial data $P_{\mathcal{D},n}(\eta)$ ($n = 0, 1, \dots, M$) [27]. The simplest recurrence relations with constant coefficients corresponding to $X_{\min}(\eta)$ has $3 + 2\ell_{\mathcal{D}}$ terms and it needs $\ell_{\mathcal{D}} + 1$ initial data. The difference of M and $\ell_{\mathcal{D}}$, $\ell_{\mathcal{D}} - M = \sum_{j=1}^M (d_j - 1) + 2M_{\text{I}}M_{\text{II}}$, becomes large for large d_j . In order to calculate the multi-indexed orthogonal polynomials by using recurrence relations, the $3 + 2M$ term recurrence relations with variable dependent coefficients are useful. On the other hand, in order to study bispectral properties etc., recurrence relations with constant coefficients are needed.

We hope that the recurrence relations with constants coefficients obtained in this paper will be used as a starting point for theoretical developments of various problems involving bispectrality, generalizations of the Jacobi matrix, spectral theory, etc.

Acknowledgments

I thank the organizers and participants of the workshop “Exceptional orthogonal polynomials and exact solutions in Mathematical Physics” at Segovia (Spain), 7-12 September 2014. This workshop was exciting and I learned importance of the recurrence relations with constant coefficients. I thank R.Sasaki for discussion and reading of the manuscript, and S.Tsujimoto and H.Miki for discussion. I am supported in part by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), No.25400395.

A Some formulas

The notation and fundamental formulas of the multi-indexed orthogonal polynomials of Laguerre, Jacobi, Wilson and Askey-Wilson types are found in [27]. See also [23]. In this appendix we present other basic formulas. See footnote in §2. We write parameter $(\boldsymbol{\lambda})$ dependence explicitly.

A.1 Multi-indexed Laguerre and Jacobi polynomials

Explicit forms of the operators (2.6) are

$$\hat{\mathcal{F}}_{d_1 \dots d_s}(\boldsymbol{\lambda}) = c_{\mathcal{F}}^{-1} \frac{\Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda})}{\Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda})} \left(c_{\mathcal{F}} e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\eta) \left(\frac{d}{d\eta} - \frac{\partial_{\eta} \Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda})}{\Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda})} \right) + \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda}) \right), \quad (\text{A.1})$$

$$\hat{\mathcal{B}}_{d_1 \dots d_s}(\boldsymbol{\lambda}) = c_{\mathcal{F}} \frac{\Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda})}{\Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda})} \left(c_{\mathcal{F}} e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\eta) \left(-\frac{d}{d\eta} + \frac{\partial_{\eta} \Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda})}{\Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda})} \right) + \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda}) \right), \quad (\text{A.2})$$

where $e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\eta)$, $e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\eta)$, $\tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda})$ and $\tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda})$ are given by

$$\begin{aligned} \text{L: } \quad e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\eta) &= \begin{cases} 1 & : d_s = d_s^{\text{I}} \\ \eta & : d_s = d_s^{\text{II}} \end{cases}, \quad e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\eta) = \begin{cases} \eta & : d_s = d_s^{\text{I}} \\ 1 & : d_s = d_s^{\text{II}} \end{cases}, \\ \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda}) &= \begin{cases} -2 & : d_s = d_s^{\text{I}} \\ 2(g + s_{\text{I}} - s_{\text{II}}) + 1 & : d_s = d_s^{\text{II}} \end{cases}, \\ \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda}) &= \begin{cases} -2(g + s_{\text{I}} - s_{\text{II}}) + 1 & : d_s = d_s^{\text{I}} \\ 2 & : d_s = d_s^{\text{II}} \end{cases}, \\ \text{J: } \quad e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\eta) &= \begin{cases} \frac{1+\eta}{2} & : d_s = d_s^{\text{I}} \\ \frac{1-\eta}{2} & : d_s = d_s^{\text{II}} \end{cases}, \quad e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\eta) = \begin{cases} \frac{1-\eta}{2} & : d_s = d_s^{\text{I}} \\ \frac{1+\eta}{2} & : d_s = d_s^{\text{II}} \end{cases}, \\ \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda}) &= \begin{cases} -2(h + s_{\text{II}} - s_{\text{I}}) - 1 & : d_s = d_s^{\text{I}} \\ 2(g + s_{\text{I}} - s_{\text{II}}) + 1 & : d_s = d_s^{\text{II}} \end{cases}, \end{aligned} \quad (\text{A.3})$$

$$\tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda}) = \begin{cases} -2(g + s_{\text{I}} - s_{\text{II}}) + 1 & : d_s = d_s^{\text{I}} \\ 2(h + s_{\text{II}} - s_{\text{I}}) - 1 & : d_s = d_s^{\text{II}} \end{cases}. \quad (\text{A.4})$$

By (A.1)–(A.2), eqs. (2.5) are

$$\begin{aligned} & P_{d_1 \dots d_s, n}(\eta; \boldsymbol{\lambda}) \\ &= \frac{1}{\Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\eta) (\Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda}) \partial_{\eta} P_{d_1 \dots d_{s-1}, n}(\eta; \boldsymbol{\lambda}) - \partial_{\eta} \Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda}) P_{d_1 \dots d_{s-1}, n}(\eta; \boldsymbol{\lambda})) \right. \\ & \quad \left. + c_{\mathcal{F}}^{-1} \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda}) \Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda}) P_{d_1 \dots d_{s-1}, n}(\eta; \boldsymbol{\lambda}) \right), \end{aligned} \quad (\text{A.5})$$

$$\begin{aligned} & (\mathcal{E}_n(\boldsymbol{\lambda}) - \tilde{\mathcal{E}}_d(\boldsymbol{\lambda})) P_{d_1 \dots d_{s-1}, n}(\eta; \boldsymbol{\lambda}) \\ &= \frac{c_{\mathcal{F}}^2}{\Xi_{d_1 \dots d_s}(\eta; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\eta) (-\Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda}) \partial_{\eta} P_{d_1 \dots d_s, n}(\eta; \boldsymbol{\lambda}) + \partial_{\eta} \Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda}) P_{d_1 \dots d_s, n}(\eta; \boldsymbol{\lambda})) \right. \\ & \quad \left. + c_{\mathcal{F}}^{-1} \tilde{e}_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda}) \Xi_{d_1 \dots d_{s-1}}(\eta; \boldsymbol{\lambda}) P_{d_1 \dots d_s, n}(\eta; \boldsymbol{\lambda}) \right). \end{aligned} \quad (\text{A.6})$$

A.2 Multi-indexed Wilson and Askey-Wilson polynomials

We restrict the parameters: $\{a_1^*, a_2^*\} = \{a_1, a_2\}$ (as a set) and $\{a_3^*, a_4^*\} = \{a_3, a_4\}$ (as a set).

Explicit form of the potential function $\hat{V}_{d_1 \dots d_s}(x; \boldsymbol{\lambda})$ is

$$\begin{aligned} \hat{V}_{d_1 \dots d_s}(x; \boldsymbol{\lambda}) &= \frac{\check{\Xi}_{d_1 \dots d_{s-1}}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x - i\gamma; \boldsymbol{\lambda})}{\check{\Xi}_{d_1 \dots d_{s-1}}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x; \boldsymbol{\lambda})} \\ & \quad \times \begin{cases} \kappa^{s_{\text{I}} - s_{\text{II}} - 1} \alpha^{\text{I}}(\boldsymbol{\lambda}) V(x; \mathbf{t}^{\text{I}}(\boldsymbol{\lambda}^{[s_{\text{I}} - 1, s_{\text{II}}]})) & : d_s = d_s^{\text{I}} \\ \kappa^{s_{\text{II}} - s_{\text{I}} - 1} \alpha^{\text{II}}(\boldsymbol{\lambda}) V(x; \mathbf{t}^{\text{II}}(\boldsymbol{\lambda}^{[s_{\text{I}}, s_{\text{II}} - 1]})) & : d_s = d_s^{\text{II}} \end{cases}. \end{aligned} \quad (\text{A.7})$$

Explicit forms of the operators (2.6) are

$$\begin{aligned} \hat{\mathcal{F}}_{d_1 \dots d_s}(\boldsymbol{\lambda}) &= \frac{ic_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda})}{\varphi(x) \check{\Xi}_{d_1 \dots d_{s-1}}(x; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) e^{\frac{\gamma}{2}p} \right. \\ & \quad \left. - e_{d_1 \dots d_s}^{\hat{\mathcal{F}}^*}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) e^{-\frac{\gamma}{2}p} \right), \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} \hat{\mathcal{B}}_{d_1 \dots d_s}(\boldsymbol{\lambda}) &= \frac{ic_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda})}{\varphi(x) \check{\Xi}_{d_1 \dots d_s}(x; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_{s-1}}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) e^{\frac{\gamma}{2}p} \right. \\ & \quad \left. - e_{d_1 \dots d_s}^{\hat{\mathcal{B}}^*}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_{s-1}}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) e^{-\frac{\gamma}{2}p} \right), \end{aligned} \quad (\text{A.9})$$

where $c_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda})$, $c_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda})$, $e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(x; \boldsymbol{\lambda})$ and $e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(x; \boldsymbol{\lambda})$ are given by

$$c_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda})^{-1} = c_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda}) = \begin{cases} \kappa^{\frac{s-1}{2} + s_{\text{II}}} \alpha^{\text{I}}(\boldsymbol{\lambda})^{\frac{1}{2}} & : d_s = d_s^{\text{I}} \\ \kappa^{\frac{s-1}{2} + s_{\text{I}}} \alpha^{\text{II}}(\boldsymbol{\lambda})^{\frac{1}{2}} & : d_s = d_s^{\text{II}} \end{cases}, \quad (\text{A.10})$$

$$e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(x; \boldsymbol{\lambda}) = \begin{cases} v_1(x; \boldsymbol{\lambda}^{[s_{\text{I}}, s_{\text{II}}]}) & : d_s = d_s^{\text{I}} \\ v_2(x; \boldsymbol{\lambda}^{[s_{\text{I}}, s_{\text{II}}]}) & : d_s = d_s^{\text{II}} \end{cases}, \quad (\text{A.11})$$

$$e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(x; \boldsymbol{\lambda}) = \begin{cases} v_2(x; \boldsymbol{\lambda}^{[s_{\text{I}}-1, s_{\text{II}}]}) & : d_s = d_s^{\text{I}} \\ v_1(x; \boldsymbol{\lambda}^{[s_{\text{I}}, s_{\text{II}}-1]}) & : d_s = d_s^{\text{II}} \end{cases}. \quad (\text{A.12})$$

Here $v_1(x; \boldsymbol{\lambda})$ and $v_2(x; \boldsymbol{\lambda})$ are given in [23]:

$$v_1(x; \boldsymbol{\lambda}) = \begin{cases} \prod_{j=1}^2 (a_j + ix) & : \text{W} \\ e^{-ix} \prod_{j=1}^2 (1 - a_j e^{ix}) & : \text{AW} \end{cases}, \quad v_2(x; \boldsymbol{\lambda}) = \begin{cases} \prod_{j=3}^4 (a_j + ix) & : \text{W} \\ e^{-ix} \prod_{j=3}^4 (1 - a_j e^{ix}) & : \text{AW} \end{cases}. \quad (\text{A.13})$$

Note that $v_1^*(x; \boldsymbol{\lambda}) = v_1(-x; \boldsymbol{\lambda})$ and $v_2^*(x; \boldsymbol{\lambda}) = v_2(-x; \boldsymbol{\lambda})$. By (A.8)–(A.9), eqs. (2.5) are

$$\begin{aligned} & \check{P}_{d_1 \dots d_s, n}(x; \boldsymbol{\lambda}) \\ &= \frac{i c_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(\boldsymbol{\lambda})}{\varphi(x) \check{\Xi}_{d_1 \dots d_{s-1}}(x; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{F}}}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) P_{d_1 \dots d_{s-1}, n}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) \right. \\ & \quad \left. - e_{d_1 \dots d_s}^{\hat{\mathcal{F}}^*}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_s}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) P_{d_1 \dots d_{s-1}, n}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) \right), \quad (\text{A.14}) \end{aligned}$$

$$\begin{aligned} & (\mathcal{E}_n(\boldsymbol{\lambda}) - \tilde{\mathcal{E}}_{d_s}(\boldsymbol{\lambda})) P_{d_1 \dots d_{s-1}, n}(x; \boldsymbol{\lambda}) \\ &= \frac{i c_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(\boldsymbol{\lambda})}{\varphi(x) \check{\Xi}_{d_1 \dots d_s}(x; \boldsymbol{\lambda})} \left(e_{d_1 \dots d_s}^{\hat{\mathcal{B}}}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_{s-1}}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) P_{d_1 \dots d_s, n}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) \right. \\ & \quad \left. - e_{d_1 \dots d_s}^{\hat{\mathcal{B}}^*}(x; \boldsymbol{\lambda}) \check{\Xi}_{d_1 \dots d_{s-1}}(x - i\frac{\gamma}{2}; \boldsymbol{\lambda}) P_{d_1 \dots d_s, n}(x + i\frac{\gamma}{2}; \boldsymbol{\lambda}) \right). \quad (\text{A.15}) \end{aligned}$$

References

- [1] D. Gómez-Ullate, N. Kamran and R. Milson, “An extension of Bochner’s problem: exceptional invariant subspaces,” J. Approx Theory **162** (2010) 987-1006, [arXiv:0805.3376\[math-ph\]](#); “An extended class of orthogonal polynomials defined by a Sturm-Liouville problem,” J. Math. Anal. Appl. **359** (2009) 352-367, [arXiv:0807.3939\[math-ph\]](#).
- [2] C. Quesne, “Exceptional orthogonal polynomials, exactly solvable potentials and supersymmetry,” J. Phys. **A41** (2008) 392001 (6pp), [arXiv:0807.4087\[quant-ph\]](#).
- [3] S. Odake and R. Sasaki, “Crum’s theorem for ‘discrete’ quantum mechanics,” Prog. Theor. Phys. **122** (2009) 1067-1079, [arXiv:0902.2593\[math-ph\]](#).
- [4] S. Odake and R. Sasaki, “Infinitely many shape invariant potentials and new orthogonal polynomials,” Phys. Lett. **B679** (2009) 414-417, [arXiv:0906.0142\[math-ph\]](#); “Another set of infinitely many exceptional (X_ℓ) Laguerre polynomials,” Phys. Lett. **B684** (2010) 173-176, [arXiv:0911.3442\[math-ph\]](#).

- [5] S. Odake and R. Sasaki, “Infinitely many shape invariant discrete quantum mechanical systems and new exceptional orthogonal polynomials related to the Wilson and Askey-Wilson polynomials,” *Phys. Lett.* **B682** (2009) 130-136, [arXiv:0909.3668 \[math-ph\]](#).
- [6] S. Odake and R. Sasaki, “Infinitely many shape invariant potentials and cubic identities of the Laguerre and Jacobi polynomials,” *J. Math. Phys.* **51** (2010) 053513 (9pp), [arXiv:0911.1585 \[math-ph\]](#).
- [7] C.-L. Ho, S. Odake and R. Sasaki, “Properties of the exceptional (X_ℓ) Laguerre and Jacobi polynomials,” *SIGMA* **7** (2011) 107 (24pp), [arXiv:0912.5447 \[math-ph\]](#).
- [8] D. Gómez-Ullate, N. Kamran and R. Milson, “Exceptional orthogonal polynomials and the Darboux transformation,” *J. Phys.* **A43** (2010) 434016, [arXiv:1002.2666 \[math-ph\]](#).
- [9] L. García-Gutiérrez, S. Odake and R. Sasaki, “Modification of Crum’s theorem for ‘discrete’ quantum mechanics,” *Prog. Theor. Phys.* **124** (2010) 1-26, [arXiv:1004.0289 \[math-ph\]](#).
- [10] S. Odake and R. Sasaki, “Exceptional Askey-Wilson type polynomials through Darboux-Crum transformations,” *J. Phys.* **A43** (2010) 335201 (18pp), [arXiv:1004.0544 \[math-ph\]](#).
- [11] R. Sasaki, S. Tsujimoto and A. Zhedanov, “Exceptional Laguerre and Jacobi polynomials and the corresponding potentials through Darboux-Crum transformations,” *J. Phys.* **A43** (2010) 315204, [arXiv:1004.4711 \[math-ph\]](#).
- [12] D. Gómez-Ullate, N. Kamran and R. Milson, “On orthogonal polynomials spanning a non-standard flag,” *Contemp. Math.* **563** (2011) 51-72, [arXiv:1101.5584 \[math-ph\]](#).
- [13] S. Odake and R. Sasaki, “Exceptional (X_ℓ) (q)-Racah polynomials,” *Prog. Theor. Phys.* **125** (2011) 851-870, [arXiv:1102.0812 \[math-ph\]](#).
- [14] Y. Grandati, “Solvable rational extensions of the Morse and Kepler-Coulomb potentials,” *J. Math. Phys.* **52** (2011) 103505 (12pp), [arXiv:1103.5023 \[math-ph\]](#).

- [15] D. Gómez-Ullate, N. Kamran and R. Milson, “Two-step Darboux transformations and exceptional Laguerre polynomials,” *J. Math. Anal. Appl.* **387** (2012) 410-418, [arXiv:1103.5724\[math-ph\]](#).
- [16] S. Odake and R. Sasaki, “Discrete quantum mechanics,” (Topical Review) *J. Phys.* **A44** (2011) 353001 (47pp), [arXiv:1104.0473\[math-ph\]](#).
- [17] S. Odake and R. Sasaki, “Exactly solvable quantum mechanics and infinite families of multi-indexed orthogonal polynomials,” *Phys. Lett.* **B702** (2011) 164-170, [arXiv:1105.0508\[math-ph\]](#).
- [18] C.-L. Ho, “Prepotential approach to solvable rational extensions of harmonic oscillator and Morse potentials,” *J. Math. Phys.* **52** (2011) 122107 (8pp), [arXiv:1105.3670\[math-ph\]](#).
- [19] C. Quesne, “Revisiting (quasi-)exactly solvable rational extensions of the Morse potential,” *Int. J. Mod. Phys. A* **27** (2012) 1250073 (18pp), [arXiv:1203.1812\[math-ph\]](#).
- [20] Y. Grandati, “New rational extensions of solvable potentials with finite bound state spectrum,” *Phys. Lett.* **A376** (2012) 2866-2872, [arXiv:1203.4149\[math-ph\]](#).
- [21] S. Odake and R. Sasaki, “Multi-indexed (q -)Racah polynomials,” *J. Phys.* **A 45** (2012) 385201 (21pp), [arXiv:1203.5868\[math-ph\]](#).
- [22] D. Gómez-Ullate, N. Kamran and R. Milson, “A conjecture on exceptional orthogonal polynomials,” *Found. Comput. Math.* In press, [arXiv:1203.6857\[math-ph\]](#).
- [23] S. Odake and R. Sasaki, “Multi-indexed Wilson and Askey-Wilson polynomials,” *J. Phys.* **A46** (2013) 045204 (22pp), [arXiv:1207.5584\[math-ph\]](#).
- [24] C. Quesne, “Novel enlarged shape invariance property and exactly solvable rational extensions of the Rosen-Morse II and Eckart potentials,” *SIGMA* **8** (2012) 080 (19pp), [arXiv:1208.6165\[math-ph\]](#).
- [25] S. Odake and R. Sasaki, “Krein-Adler transformations for shape-invariant potentials and pseudo virtual states,” *J. Phys.* **A46** (2013) 245201 (24pp), [arXiv:1212.6595\[math-ph\]](#).

- [26] S. Odake and R. Sasaki, “Extensions of solvable potentials with finitely many discrete eigenstates,” J. Phys. **A46** (2013) 235205 (15pp), [arXiv:1301.3980 \[math-ph\]](#).
- [27] S. Odake, “Recurrence Relations of the Multi-Indexed Orthogonal Polynomials,” J. Math. Phys. **54** (2013) 083506 (18pp), [arXiv:1303.5820 \[math-ph\]](#).
- [28] D. Gómez-Ullate, Y. Grandati and R. Milson, “Rational extensions of the quantum harmonic oscillator and exceptional Hermite polynomials,” [arXiv:1306.5143 \[math-ph\]](#).
- [29] S. Odake and R. Sasaki, “Casoratian Identities for the Wilson and Askey-Wilson Polynomials,” To appear in Journal of Approximation Theory, [arXiv:1308.4240 \[math-ph\]](#).
- [30] S. Odake, “Equivalences of the Multi-Indexed Orthogonal Polynomials,” J. Math. Phys. **55** (2014) 013502 (17pp), [arXiv:1309.2346 \[math-ph\]](#).
- [31] E. Routh, “On some properties of certain solutions of a differential equation of the second order,” Proc. London Math. Soc. **16** (1884) 245-261; S. Bochner, “Über Sturm-Liouvillesche Polynomsysteme,” Math. Zeit. **29** (1929) 730-736.
- [32] G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc., Providence, RI (1939); T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York (1978); M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, vol. 98 of Encyclopedia of mathematics and its applications, Cambridge Univ. Press, Cambridge (2005).
- [33] G. Darboux, *Théorie générale des surfaces* vol 2 (1888) Gauthier-Villars, Paris.
- [34] M. M. Crum, “Associated Sturm-Liouville systems,” Quart. J. Math. Oxford Ser. (2) **6** (1955) 121-127, [arXiv:physics/9908019](#).
- [35] M. G. Krein, “On continuous analogue of a formula of Christoffel from the theory of orthogonal polynomials,” (Russian) Doklady Acad. Nauk. CCCP, **113** (1957) 970-973; V. É. Adler, “A modification of Crum’s method,” Theor. Math. Phys. **101** (1994) 1381-1386.
- [36] F. A. Grünbaum and L. Haine, “Bispectral Darboux transformations: an extension of the Krall polynomials,” IMRN (1997) No. 8, 359–392.

- [37] H. Miki and S. Tsujimoto, “A new recurrence formula for generic exceptional orthogonal polynomials,” [arXiv:1410.0183 \[math.CA\]](#)
- [38] A. J. Durán, “Higher order recurrence relation for exceptional Charlier, Meixner, Hermite and Laguerre orthogonal polynomials,” [arXiv:1409.4697 \[math.CA\]](#).
- [39] S. Odake and R. Sasaki, “Unified theory of exactly and quasi-exactly solvable ‘discrete’ quantum mechanics: I. Formalism,” *J. Math. Phys* **51** (2010) 083502 (24pp), [arXiv:0903.2604 \[math-ph\]](#).
- [40] S. Odake and R. Sasaki, “Solvable Discrete Quantum Mechanics: q -Orthogonal Polynomials with $|q| = 1$ and Quantum Dilogarithm,” [arXiv:1406.2768 \[math-ph\]](#).