

ORBITAL CIRCULARIZATION OF A PLANET ACCRETING DISK GAS: FORMATION OF DISTANT JUPITERS IN CIRCULAR ORBITS BASED ON CORE ACCRETION MODEL

AKIHIRO KIKUCHI¹, ARIKA HIGUCHI¹, AND SHIGERU IDA²

¹*Department of Earth and Planetary Sciences, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan*

²*Earth-Life Science Institute, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8550, Japan*

kikuchi.a@geo.titech.ac.jp, higuchia@geo.titech.ac.jp, ida@elsi.jp

ABSTRACT

Recently, gas giant planets in nearly circular orbits with large semimajor axes ($a \sim 30\text{--}1000\text{AU}$) have been detected by direct imaging. We have investigated orbital evolution in a formation scenario for such planets, based on core accretion model: i) Icy cores accrete from planetesimals at $\lesssim 30\text{AU}$, ii) they are scattered outward by an emerging nearby gas giant to acquire highly eccentric orbits, and iii) their orbits are circularized through accretion of disk gas in outer regions, where they spend most of time. We analytically derived equations to describe the orbital circularization through the gas accretion. Numerical integrations of these equations show that the eccentricity decreases by a factor of more than 5 during the planetary mass increases by a factor of 10. Because runaway gas accretion increases planetary mass by $\sim 10\text{--}300$, the orbits are sufficiently circularized. On the other hand, a is reduced at most only by a factor of 2, leaving the planets in outer regions. If the relative velocity damping by shock is considered, the circularization is slowed down, but still efficient enough. Therefore, this scenario potentially accounts for the formation of observed distant jupiters in nearly circular orbits. If the apocenter distances of the scattered cores are larger than the disk sizes, their a shrink to a quarter of the disk sizes; the a -distribution of distant giants could reflect outer edges of the disks in a similar way that those of hot jupiters may reflect inner edges.

Subject headings: planetary systems — planets and satellites: formation — accretion, accretion disks

1. Introduction

Distant extrasolar gaseous giant planets in nearly circular orbits have been detected by direct imaging observations in several systems (e.g., Kalas et al. 2008; Marois et al. 2008; Kuzuhara et al.

2013). In the conventional core accretion model, it is difficult to form cores that are massive enough to undergo runaway gas accretion at $\gtrsim 30\text{AU}$ within disk lifetime (\sim a few million years)¹, because the core growth timescale is roughly proportional to a cube of the distance from the central star (e.g., Ida & Lin 2004a). Although the cores cannot be formed in such distant regions, gas giant planets formed interior to 30AU can be scattered by other giant planets to attain semimajor axes $a \gtrsim 100\text{AU}$ (e.g., Marzari & Weidenschilling 2002; Nagasawa et al. 2008). Because the dynamical energy is lower in outer regions, a of scattered planets are broadly distributed up to $\sim 1000\text{AU}$. However, due to the conservation of the total angular momentum, the eccentricities e of the scattered orbits must be excited to be close to unity. While disk-planet interactions tend to damp such high values of e , they may not be efficient enough to account for the observed low eccentricities of distant, gas giant planets, because local protoplanetary disk mass may not be massive enough in the distant regions (Ida et al. 2013).

This difficulty has raised the possibility of formation of gas giants by disk gravitational instability (e.g., Boss 2001; Helled et al. 2014 and references therein). Kratter et al. (2010) showed that if disk instability forms planetary mass clumps, it would form more abundant brown-dwarfs and M-star companions. A population synthesis simulation based on the disk instability model (Forgan & Rice 2013) showed that most of such brown-dwarfs and M-star companions may be retained in outer regions. However, it is not consistent with direct imaging surveys so far done. Furthermore, the observationally clear correlation between fraction of stars with gas giants and stellar metallicity (Fischer & Valenti 2005) is not easy for the disk instability model to explain, while the correlation is consistent with the core accretion model (e.g., Ida & Lin 2004b).

Based on the core accretion model, Crida et al. (2009) proposed outward type II migration for the origin of the distant gas giants in nearly circular orbits. However, the outward type II migration requires a pair of giant planets in a common gap with the inner one more massive than the outer one and appropriate disk conditions.

Ida et al. (2013) found out another path to form the distant gas giants in nearly circular orbits, based on the core accretion model: outward scattering of cores by a nearby gas giant followed by accretion of gas in outer regions. Because the orbital circularization through gas accretion was also shown by a hybrid N-body and 2D hydrodynamical simulation (E. Thommes 2010, private communication), this path is one of promising mechanisms. Details of the path they found are as follows. Oligarchic growth produces similar-sized multiple cores (Kokubo & Ida 1998). Once some core starts runaway gas accretion, the planet’s mass rapidly increases (e.g., Bodenheimer & Pollack 1986; Ikoma et al. 2000). When the mass increase is fast enough, the planet undergoes close encounters with nearby cores to strongly scatter them rather than shepherd them (Zhou & Lin

¹Recently, “pebble accretion” has been proposed, which is a rapid growth process accreting small bodies suffering strong gas drag (e.g., Lambrechts & Johansen 2012; Youdin & Kenyon 2013, and references therein). If pebble accretion works well in outer disk regions, cores could be formed even outside of 30 AU. This possibility should also be pursued, although it is not discussed here.

2007; Shiraishi & Ida 2008). Some cores are scattered to large distance where the surface densities of both residual planetesimals and gas are relatively low. If the scattered planet is a gas giant, its mass could be comparable to or larger than the local disk mass, because the scattering may occur in a late disk evolution stage after the formation of the gas giant. Then, eccentricity damping due to dynamical friction from local disk gas is inefficient.

On the other hand, in the core scattering model, cores' masses are usually well below the local disk mass. However, since the scattered cores have highly eccentric orbits and the relative velocity between the cores and the disk gas should be highly supersonic, dynamical friction from the local disk gas is less efficient (Ostriker 1999; Papaloizou & Larwood 2000; Muto et al. 2011). Muto et al. (2011) showed that the dynamical friction timescale in supersonic regime for a planet with mass M_p , orbital eccentricity e and semimajor axis a in a gas disk with surface density Σ is

$$\tau_{\text{DF}} \sim \frac{1}{8\pi} \left(\frac{M_*}{\Sigma a^2} \right) \left(\frac{M_*}{M_p} \right) \left(\frac{c_s}{v_K} \right) \left(\frac{v}{v_K} \right)^3 T_K \sim 10^4 \left(\frac{\Sigma a^2}{0.01 M_*} \right)^{-1} \left(\frac{M_p}{10 M_\oplus} \right)^{-1} \left(\frac{c_s/v_K}{0.1} \right) e^3 T_K, \quad (1)$$

where M_* is the host star's mass, c_s is local sound velocity, and v_K and T_K are Kepler velocity and its orbital period at a . For a planet with mass $10M_\oplus$ in a highly eccentric orbit ($e \sim 1$) with $a \sim 100\text{AU}$, τ_{DF} is as long as $\sim 10^7$ years, which is longer than an observationally inferred disk lifetime \sim a few $\times 10^6$ years. Thereby, the dynamical friction from local disk gas may be less effective than the orbital circularization via planetary gas accretion, which we discuss in this paper (see section 5). Furthermore, since the dynamical friction also damps semimajor axis efficiently in the course of eccentricity damping from values close to unity, we may not be able to retain the cores in outer regions.

Reduction in the planetesimal accretion rate decreases the critical core mass for the onset of gas accretion (e.g., Ikoma et al. 2000). The scattered cores in highly eccentric orbits spend most of time at large distances, where planetesimal accretion rate is significantly low, so that it is possible for the scattering to trigger gas accretion onto the cores. In the case of highly eccentric orbits, since the cores spend most of time near apocenters at large distance, their orbits are circularized there through accretion of local gas with higher specific angular momentum. As a result, it is expected that the cores' orbits are circularized keeping their apocenters almost fixed, in the course of gas accretion. We will show that the apocenter shrinks because the semimajor axis slightly shrinks due to energy dissipation by collision between the disk gas and the planet and that the energy dissipation accelerates the orbital circularization (in the case of moderate eccentricity, the energy dissipation is more important for the orbital circularization than accretion of high angular momentum gas).

Assuming that the orbits of the scattered cores are quickly circularized to the degree that depends on the ratio between the planet mass and local disk mass with their semimajor axes kept fixed, Ida et al. (2013) performed a population synthesis simulation to predict statistical distributions of distant gas giants formed by this mechanism. They showed that the fraction of systems with the distant gas giants is a ~ 0.1 – 1% and most of them have low eccentricities ($e \lesssim 0.1$).

Although the fraction is further lower, systems with multiple distant gas giants are also formed, because a single gas giant can scatter multiple cores in inner regions. HR8799 system has four distant gas giants and the outer three planets could be in 4:2:1 resonance. In the scattering core model, formation of four distant gas giants is extremely rare and probability for capture into the resonances during inward migration associated with eccentricity damping is not clear. However, the inner two planets have semimajor axes ~ 15 and 27 AU, which could be formed *in situ* without the scattering process. Formation of HR8799 system by core accretion scenario is a very interesting problem that should be addressed in the future. In this paper, we focus on a fundamental process of orbital circularization of an isolated planet through gas accretion.

In the core scattering model, positive correlation between the semimajor axis and the mass of the distant gas giants is predicted. The critical planet mass for gap opening is higher in larger orbital radius (e.g., Ida & Lin 2004a). If the gap opening halts growth of gas giants, the correlation is established, as shown in the population synthesis calculation in Ida et al. (2013).

Note that Ida et al. (2013) assumed that the eccentricities of the scattered cores are efficiently damped without any decrease in semimajor axes, implicitly assuming very efficient damping due to gas accretion, although they did not incorporate detailed orbital evolution by the gas accretion. As we will show, the eccentricity damping due to gas accretion is indeed efficient, while the degree of damping depends on how much the planets grow by accreting gas and semimajor axes are also damped by a factor of ~ 2 . The population synthesis simulation must be improved by incorporating the damping formulas due to gas accretion derived in this paper, in order to discuss the distribution of distant gas giants in comparison with observation when the number of discovered planets becomes large enough for statistical arguments.

Here, we investigate the orbital circularization of the scattered planets during gas accretion through detailed analytical calculations. Section 2 describes the assumptions of gas accretion onto the cores. In section 3, we analytically derive the formulas for the orbital evolution in the course of gas accretion. In section 4, we describe the orbital evolution by numerically solving the formulas. In section 5, we show some results of the population synthesis calculations, by incorporating the prescriptions of orbital evolution through the mass growth due to accretion of disk gas. Section 6 is devoted for summary.

2. Model

We start our calculation from the stage at which a core has already been scattered outward by a gas giant to attain eccentricity close to unity (e_{ini}) and semimajor axis (a_{ini}) that is much larger than the original one (a_{ori}). Note that the pericenter distance of the scattered planet's orbit must be close to a_{ori} : $a_{\text{ori}} \simeq q_{\text{ini}} = a_{\text{ini}}(1 - e_{\text{ini}})$. Since a_{ori} should be close to the gas giant's orbital radius, we can regard that $q_{\text{ini}} \sim 1\text{--}10\text{AU}$, based on core accretion model (e.g., Ida & Lin 2004a). As we will show later, the pericenter distance of the scattered planet quickly increases.

Accordingly, the planet immediately becomes isolated from the perturbing gas giant, so that we neglect its further perturbations. We do not calculate the initial scattering process by the gas giant, but study evolution of e and a of the scattered planet due to accretion of gas, for given e_{ini} , a_{ini} and disk radius r_{d} , to derive general formulas for the orbital evolution. In the following, we explain our prescriptions for accretion of gas onto the scattered planet.

After a core mass exceeds a critical core mass, pressure gradient no longer supports gas envelope of the planet against the planetary gravitational force and quasi-static contraction of the gas envelope starts (e.g., Mizuno 1980; Bodenheimer & Pollack 1986). The critical core mass is given by (Ikoma et al. 2000)

$$M_{\text{c,crit}} \simeq 10 \left(\frac{\dot{M}_{\text{c}}}{M_{\oplus}/10^6 \text{yr}} \right)^{(0.2-0.3)} \left(\frac{\kappa}{\kappa_0} \right)^{(0.2-0.3)} M_{\oplus}, \quad (2)$$

where κ is opacity of the gas envelope and κ_0 is that of the minimum-mass solar nebula model (Hayashi 1981). Since the planetesimal accretion rate, \dot{M}_{c} , determines heat energy source to support the envelope, a lower value of \dot{M}_{c} leads to a smaller value of $M_{\text{c,crit}}$. In general, planetesimal accretion rate rapidly decreases with orbital radius (e.g., Ida & Lin 2004a). After a core is scattered outward to acquire high orbital eccentricity, the core spends most of time at much larger orbital radii than near the original location. Thereby, an orbit-averaged value of \dot{M}_{c} is significantly lowered and it is likely that quasi-static contraction of gas envelope is initiated by the outward scattering.

According to the quasi-static contraction, disk gas can be supplied to Hill radius or Bondi radius of the planet. This means that gas accretion rate onto the planet is regulated by heat transfer through the envelope rather than environmental disk gas conditions except in the final stage in which the contraction is very fast. The (Kelvin-Helmholtz) timescale of the envelope contraction is given by (Ikoma et al. 2000; Ikoma & Genda 2006)

$$\tau_{\text{KH}} \simeq 10^{10} \left(\frac{M}{M_{\oplus}} \right)^{-(3-4)} \left(\frac{\kappa}{\kappa_{\text{ini}}} \right) \text{years}. \quad (3)$$

From these arguments, we assume that gas accretion rate does not depend on the position of the eccentric orbit during an orbital period, although environmental disk conditions considerably change during one orbital cycle for highly eccentric orbits.

In general, the dependence of τ_{KH} on orbital radius r is weak for radiation-dominated envelope (e.g. Ikoma et al. 2000). If convective envelope develops, envelope contraction rate can be affected by disk temperature and density (Ikoma et al. 2001; Piso & Youdin 2014). However, even if τ_{KH} has the r -dependence, the dependence may be smoothed out when τ_{KH} is longer than the orbital period, that is, when $M \lesssim 100M_{\oplus}$, because the response time of the envelope structure is given by τ_{KH} . As we show in the following, gas accretion rate onto the planet may be regulated by disk gas supply rather than by envelope contraction for $M \gtrsim 100M_{\oplus}$.

Note that in the highly eccentric orbit, Bondi and Hill radii significantly change during one orbital circulation. The change might also induce oscillation of gas envelope that could affect gas

accretion and heat generation/cooling. Investigation of this effect is left for a future work. We will only assume the constant accretion rate during one orbit inside the disk, but not adopt any particular form of τ_{KH} .

When the envelope contraction is faster than the supply of gas and the supply has the r -dependence, the assumption of the constant accretion rate is violated. The supply can be limited by global disk accretion and Bondi accretion. The limit by global disk accretion becomes important for $M \gtrsim 100M_{\oplus}$, because the quasi-static contraction rate is given by $\dot{M}_{\text{KH}} \sim M/\tau_{\text{KH}} \sim 10^{-10}(M/M_{\oplus})^{(4-5)}(\kappa/\kappa_{\text{ini}})^{-1}M_{\oplus}/\text{yr}$ (Eq. (3)) and the observationally inferred typical value of disk accretion rates onto T Tauri stars is $\dot{M}_{\text{disk}} \sim 10^{-8}M_{\odot}/\text{yr} \sim 3 \times 10^{-3}M_{\oplus}/\text{yr}$. However, the r -dependence of disk accretion rate is very weak in the regions of $r \ll r_{\text{d}}$, where a steady-accretion-disk approximation is valid.

The Bondi accretion rate is given by $\dot{M}_{\text{Bondi}} \sim \pi\rho_g(GM/v^2)^2v$ where ρ_g is disk gas density and v is the relative velocity between the planet and disk gas. Both ρ_g and v sensitively depend on r . In general, $\dot{M}_{\text{Bondi}} < \dot{M}_{\text{KH}}$ for high e and large M_{p} . But, our calculations start from small M_{p} and e is already damped when M_{p} becomes large. We found that in most of orbital evolution we consider, $\dot{M}_{\text{Bondi}} \gtrsim \dot{M}_{\text{KH}}$ and the supply limit by Bondi accretion does not occur.

Thus, our assumption of constant gas accretion may be justified. As a result of the time-independent gas accretion rate, the planet accretes gas preferentially in outer regions where the planet spends most of time. The specific angular momentum of the planetary orbit is given by $\ell_{\text{p}} = \sqrt{GM_*a(1-e^2)} \sim \sqrt{2GM_*q}$, where $q = a(1-e)$ is the pericenter distance and $e \sim 1$ is assumed. That of the local gas near the apocenter ($Q = a(1+e) \simeq 2a$) is given by $\ell_{\text{g}} \sim \sqrt{2GM_*a}$ (we assume circular Keplerian motion for the disk gas). Since $q = a(1-e) \ll a$ for $e \sim 1$, the planet's specific angular momentum is increased by the accretion of local gas. Then, the planetary orbit tends to be circularized with the apocenter distance (Q) fixed.

If the orbit deviates from the disk with a finite size r_{d} , beyond which gas density is significantly declined, we halt gas accretion at $r > r_{\text{d}}$. In the following derivations, we consider two cases: i) the apocenter is inside the disk ($Q < r_{\text{d}}$) and ii) it is outside the disk ($Q > r_{\text{d}}$). We will refer to cases i) and ii) as "embedded case" and "deviated case," respectively. In the "deviated case," the planet mostly accretes gas at $r \sim r_{\text{d}}$ and planetary orbits tend to be fitted to circular orbits at r_{d} rather than those at Q .

The specific orbital energies of the planet and local gas near the apocenter are $\epsilon_{\text{p}} = -GM_*/2a$ and $\epsilon_{\text{g}} = -GM_*/2a(1+e) \sim -GM_*/4a$, respectively. Near the apocenter, the planet's specific orbital energy is increased by the accretion of local gas near the apocenter. However, the accretion of lower specific energy near the pericenter is significant due to a deep potential near the pericenter, in spite of fast passage of the pericenter. As we show in section 4.1, in the embedded case, the orbit-averaged specific orbital energy of accreting gas is exactly the same as that of the planet, irrespective of orbital eccentricity. The planet's specific orbital energy actually decreases if collisional dissipation between the planet and disk gas is taken into account. It also contributes to eccentricity damping,

with slight decay of the semimajor axis.

In embedded case, gas accretion rate onto the planet is independent of phase of the orbits. In the deviated case, we assume a constant gas accretion rate at $r < r_d$ and zero accretion rate at $r > r_d$. We do not assume even the value of the constant accretion rate, because we will derive orbital evolution as a function of planetary mass M_p but not as a function of time.

The relative velocity between the planet and the disk gas would be supersonic almost everywhere for highly eccentric orbits with $e \gtrsim h/r \sim 0.1$ where h is the disk scale height. For incident supersonic gas flow to stay in Hill radius or Bondi radius, we need some energy dissipation. Bow shock in front of the planet may provide the energy dissipation. We will leave full hydrodynamic simulations on the bow shock for future work and assume that the planet accretes disk gas in unperturbed flow and the accretion rate is independent of orbital phase in most of calculations. Note, however, that the relative velocity between the gas flow and the planet is smaller in the post-shock flow than in the unperturbed flow, which may make the eccentricity damping less efficient. In section 4.3, we perform calculations taking into account the effect of the shock with a simple 1D model and show that the eccentricity damping is indeed slowed down but does not significantly change our conclusion.

When cores are scattered by a gas giant, in early stage, eccentricities are preferentially pumped up compared with inclinations. But, if the core undergoes repeated close encounters before its orbit is circularized, orbital inclinations are also excited. We also calculated e and a evolution with non-zero inclinations. We have found that the final values of e and a change by less than 5% if the inclination is smaller than 30 degrees. So, we here show the results with zero inclinations.

In summary, the assumptions we use in most of runs are:

1. The gas disk is in Keplerian rotation. Because the relative velocity between the gas and the planet is generally supersonic, the gas is hardly perturbed by the planetary gravitational perturbations.
2. The motions of the planet and the gas disk are coplanar.
3. The gas accretion rate onto the planet is constant of during one orbit, so that orbit-averaging can be done. The planet captures the local unperturbed disk gas, conserving mass and angular momentum (energy is not conserved).
4. If the gas disk has the finite size, we truncate gas accretion during the period in which the planet goes out of the disk.

In section 3 through section 4.2, we adopt the above assumptions and derive analytical formulas to describe the orbital circularization process through the gas accretion. Even if we adopt assumption 4., analytical formulas are derived, since the constant gas accretion rate is still applied at $r < r_d$ and analytical orbit averaging can be done. If we include the effect of shock dissipation, analytical

integration is not possible, so that we show the orbital evolution obtained by numerical integration (section 4.3).

3. Derivation of formulas for orbital changes

With the assumptions 1 to 4 described in the above, we analytically derive formulas to calculate the orbital evolution in the form of differential equations. Numerically integrating the differential equations, we will show the evolution paths of e and a that are uniquely determined by their initial values and r_d .

According to discussions in section 2, we first calculate changes in the angular momentum and energy of the planet, ΔL and ΔE , during one orbital period, assuming that the mass accretion rate is constant with time during one orbital period. We also assume that the changes in orbital elements are small enough over one orbital period, in other words, the mass of the captured gas during one orbit (ΔM) is much smaller than the instantaneous planetary mass (M).

The changes ΔL and ΔE are then given by

$$\Delta L \simeq \int l_{\text{gas}} dM = \Delta M \frac{1}{t_d} \int_{-t_d/2}^{t_d/2} l_{\text{gas}} dt, \quad (4)$$

$$\Delta E \simeq \int (\epsilon_{\text{gas}} - \epsilon_{\text{coll}}) dM = \Delta M \frac{1}{t_d} \int_{-t_d/2}^{t_d/2} (\epsilon_{\text{gas}} - \epsilon_{\text{coll}}) dt, \quad (5)$$

where the integral is during one orbit, t_d is a duration at $r < r_d$ ($t_d \leq T_K$), $t = 0$ is a pericenter passage, l_{gas} and ϵ_{gas} are specific angular momentum and orbital energy of accreting gas and ϵ_{coll} is energy dissipation by collision between the planet and accreting gas.

Through ΔL and ΔE during mass growth of ΔM , specific angular momentum and orbital energy of the planet, ℓ_p and ϵ_p , are changed. Since $\Delta L = (\Delta \ell_p + \ell_p)(M + \Delta M) - M \ell_p \simeq \Delta \ell_p \cdot M + \ell_p \Delta M$ and $\Delta E \simeq \Delta \epsilon_p \cdot M + \epsilon_p \Delta M$, the change rate of ℓ_p and ϵ_p of the planet in one orbital period are expressed as

$$\frac{\Delta \ell_p}{\ell_p} \simeq \frac{\Delta L}{M \ell_p} - \frac{\Delta M}{M} = \frac{\Delta M}{M} f_\ell, \quad (6)$$

$$\frac{\Delta \epsilon_p}{\epsilon_p} \simeq \frac{\Delta E}{M \epsilon_p} - \frac{\Delta M}{M} = \frac{\Delta M}{M} f_\epsilon, \quad (7)$$

where

$$f_\ell = \frac{1}{t_d} \int_{-t_d/2}^{t_d/2} \left(\frac{l_{\text{gas}}}{\ell_p} - 1 \right) dt, \quad (8)$$

$$f_\epsilon = \frac{1}{t_d} \int_{-t_d/2}^{t_d/2} \left(\frac{\epsilon_{\text{gas}}}{\epsilon_p} - \frac{\epsilon_{\text{coll}}}{\epsilon_p} - 1 \right) dt. \quad (9)$$

Since $\ell_p = \sqrt{GM_*a(1-e^2)}$ and $\epsilon_p = -GM_*/2a$, where M_* is the host star's mass, G is the gravitational constant, the changes $\Delta\ell_p$ and $\Delta\epsilon_p$ are related with the changes of the eccentricity and semimajor axis (Δe and Δa) as

$$\frac{\Delta\ell_p}{\ell_p} = \frac{\Delta(\sqrt{a})}{\sqrt{a}} + \frac{\Delta(\sqrt{1-e^2})}{\sqrt{1-e^2}} \simeq \frac{\Delta a}{2a} - \frac{e}{1-e^2}\Delta e, \quad (10)$$

$$\frac{\Delta\epsilon_p}{\epsilon_p} = \frac{\Delta(a^{-1})}{a^{-1}} \simeq -\frac{\Delta a}{a}. \quad (11)$$

So, Δe and Δa are given by

$$\Delta e \simeq -\frac{1-e^2}{e} \left(\frac{\Delta\ell_p}{\ell_p} + \frac{1}{2} \frac{\Delta\epsilon_p}{\epsilon_p} \right) = -\frac{\Delta M}{M} f_e, \quad (12)$$

$$\frac{\Delta a}{a} \simeq -\frac{\Delta\epsilon_p}{\epsilon_p} = -\frac{\Delta M}{M} f_a, \quad (13)$$

where

$$f_e = \frac{1-e^2}{e} \left(f_\ell + \frac{1}{2} f_\epsilon \right), \quad (14)$$

$$f_a = f_\epsilon. \quad (15)$$

We finally derive the differential equations for the orbit-averaged evolution of e and a in terms of planetary mass M :

$$\frac{de}{d \log M} \simeq \frac{\Delta e}{\Delta M/M} = -f_e, \quad (16)$$

$$\frac{d \log a}{d \log M} \simeq \frac{\Delta a/a}{\Delta M/M} = -f_a. \quad (17)$$

From these equations, we also obtain

$$\frac{d \log a}{de} \simeq \frac{f_a}{f_e}. \quad (18)$$

So far, we have not assumed any specific forms for ℓ_{gas} and ϵ_{gas} . Here we assume that the planet captures gas in circular Keplerian motion to analytically derive formulas f_e and f_a . (In section 4.3, we calculate f_e and f_a for post-shocked gas flow using a simple 1D model.) Note that analytical formulas of f_e and f_a are derived even for deviated case where $Q > r_d$.

For unperturbed gas flow (circular Keplerian flow),

$$\ell_{\text{gas}} = \sqrt{GM_*r}, \quad (19)$$

$$\epsilon_{\text{gas}} = -\frac{GM_*}{2r}, \quad (20)$$

$$\epsilon_{\text{coll}} = \frac{v_{\text{rel}}(r)^2}{2}, \quad (21)$$

where $v_{\text{rel}}(r)$ is the relative velocity between the planet and the local gas, and r is the instantaneous distance of the planet from the central star. The radial and tangential components of instantaneous velocity of an eccentric Keplerian orbit of the planet at r are given by

$$v_r = v_K \sqrt{2 - \frac{r}{a} - \frac{a}{r}(1 - e^2)}, \quad (22)$$

$$v_\phi = v_K \sqrt{\frac{a}{r}(1 - e^2)}, \quad (23)$$

where a and e are the planet's semimajor axis and eccentricity. Because the local Keplerian velocity is given by $v_K = \sqrt{GM_*/r}$, the square of relative velocity is

$$v_{\text{rel}}(r)^2 = v_r^2 + (v_\phi - v_K)^2 \quad (24)$$

$$= \frac{GM_*}{r} \left(3 - \frac{r}{a} - 2\sqrt{\frac{a}{r}(1 - e^2)} \right). \quad (25)$$

For integrating Eqs. (8) and (9), we convert time to eccentric anomaly using the Kepler equation. The time average of powers of r^α ($\alpha = 1/2, -1, -3/2$) are analytically integrated, using the conversion:

$$\frac{1}{t_d} \int_{-t_d/2}^{t_d/2} \left(\frac{r}{a}\right)^{1/2} dt = \frac{2\sqrt{1+e}}{3} f_{1/2}(e, u_d);$$

$$f_{1/2}(e, u_d) = \frac{4(E(k) - E(y, k)) - (1 - e)(K(k) - F(y, k)) - e \sin 2y \sqrt{1 - k^2 \sin^2 y}}{u_d - e \sin u_d}, \quad (26)$$

$$\frac{1}{t_d} \int_{-t_d/2}^{t_d/2} \left(\frac{r}{a}\right)^{-1} dt = f_{-1}(e, u_d); \quad f_{-1}(e, u_d) = \frac{u_d}{u_d - e \sin u_d}, \quad (27)$$

$$\frac{1}{t_d} \int_{-t_d/2}^{t_d/2} \left(\frac{r}{a}\right)^{-3/2} dt = \frac{2}{\sqrt{1+e}} f_{-3/2}(e, u_d); \quad f_{-3/2}(e, u_d) = \frac{K(k) - F(y, k)}{u_d - e \sin u_d}, \quad (28)$$

where $k \equiv \sqrt{2e/(1+e)}$, $y \equiv (\pi - u_d)/2$, $K(k)$ is the complete elliptic integral of the first kind, $E(k)$ is the complete elliptic integral of the second kind, $F(y, k)$ is the elliptic integral of the first kind, $E(y, k)$ is the elliptic integral of the second kind, and u_d is the maximum eccentric anomaly ($0 < u_d < \pi$) within the disk ($r < r_d$), which is given by

$$u_d \equiv \begin{cases} \cos^{-1} \left[\frac{1}{e} \left(1 - \frac{r_d}{a} \right) \right] & [\text{for } Q > r_d], \\ \pi & [\text{for } Q < r_d]. \end{cases} \quad (29)$$

In embedded case, $u_d = \pi$ and $t_d = T_K$, while $u_d < \pi$ and $t_d < T_K$, depending on r_d , in deviated cse.

With Eqs. (26) to (28), Eqs. (8) and (9) are written as

$$f_\ell(e, u_d) = \frac{2}{3\sqrt{1-e}} f_{1/2}(e, u_d) - 1, \quad (30)$$

$$f_\epsilon(e, u_d) = 4f_{-1}(e, u_d) - 4\sqrt{1-e} f_{-3/2}(e, u_d) - 2. \quad (31)$$

Note that f_ℓ and f_ϵ are functions of only e , independent of a , in embedded case. Even in deviated case, the a -dependence enters f_ℓ and f_ϵ only through the scaled quantity a/r_d in u_d .

From Eqs. (14) and (15),

$$\frac{de}{d \log M} = -f_e(e, u_d) = -\frac{1-e^2}{e} \left(f_\ell(e, u_d) + \frac{1}{2} f_\epsilon(e, u_d) \right), \quad (32)$$

$$\frac{d \log a}{d \log M} = -f_a(e, u_d) = -f_\epsilon(e, u_d). \quad (33)$$

These equations show that while the semimajor axis is damped only by $\Delta\epsilon_p$, the orbital eccentricity is damped by both $\Delta\ell_p$ and $\Delta\epsilon_p$.

By numerically integrating Eqs. (32) and (33), we obtain the evolution of the orbital elements according to the planetary mass growth. In Fig. 1, we plot $f_\ell(e, u_d)$, $f_a(e, u_d)[= f_\epsilon(e, u_d)]$ and $f_e(e, u_d)$ as a function of e for $u_d = \pi, \pi/2$ and $\pi/4$. Because $f_e(e, u_d), f_a(e, u_d) > 0$ for any values of e and u_d , e and a monotonically decrease as the planet grows through accretion of disk gas. For $u_d = \pi$ (embedded case), f_ℓ dominates the eccentricity damping. That is, accretion of disk gas with high specific angular momentum near the apocenter is responsible for the eccentricity damping. On the other hand, for $u_d = \pi/2$ and $\pi/4$ (deviated cases), f_ℓ is small or negative except for high e . In these cases, the energy dissipation by collision between incident gas and the planet is responsible for the eccentricity damping (see section 4.2).

When e becomes small enough, $d \log a / d \log M$ quickly approaches zero ($f_a \rightarrow 0$). Thus, the asymptotic values of a are uniquely determined by the initial values of e , a , and r_d . In the next section, we show the numerically obtained evolution paths.

4. Evolution paths of e and a

4.1. Embedded case

First, we consider embedded case, that is, whole parts of a planetary orbit is embedded in the disk. Since equation (16) is independent of multiplication of M by a constant factor, we can adopt a scaled quantity M/M_{ini} as a variable, where M_{ini} is the initial value of M . Then, Eq. (16) includes only e and M/M_{ini} , so that the evolution of e is uniquely given as a function of M/M_{ini} for any initial values of eccentricity (e_{ini}). The evolutionary paths for representative values of e_{ini} that obtained by numerical integration of Eq. (16) are shown in Figure 2a. This figure shows that e decreases to values below $0.2e_{\text{ini}}$ when M attains $10M_{\text{ini}}$. Since a typical core mass to start runaway gas accretion is $\sim 10M_\oplus$, it means that e is reduced to values smaller than 0.2 when the planet acquires Saturnian mass ($\sim 100M_\oplus$), even if its initial orbit was close to a parabolic orbit ($e_{\text{ini}} \sim 1$). If the mass is scaled by that at $e = 0.1$, denoted by M_{01} , e is uniquely determined by M/M_{01} . Figure 2b shows the self-similar solution of e as a function of M/M_{01} (the solid curve). The fitting formula given by Eq. (41), which is presented in section 5 is also plotted with the dashed curve.




Fig. 1.— The functions $f_l(e, u_d)$, $f_a(e, u_d)[= f_\epsilon(e, u_d)]$ and $f_e(e, u_d)$. They are plotted as a function of e for $u_d = \pi$ (solid lines), $\pi/2$ (dashed lines) and $\pi/4$ (dotted lines).




Fig. 2.— Evolution of e as a function of M . (a) M is scaled by M_{ini} and $e_{\text{ini}} = 0.8$ (the solid line), 0.9 (the dashed line) and 0.99 (the dotted line) are plotted. (b) M is scaled by M_{01} , where M_{01} is M at $e = 0.1$. In this case, e is uniquely determined by M/M_{01} (the solid line). The fitting formula, Eq. (41), which is presented in section 5 is also plotted with the dashed line.




Fig. 3.— Evolution of e as a function of a . (a) a is scaled by a_{ini} and $e_{\text{ini}} = 0.8$ (the solid line), 0.9 (the dashed line) and 0.99 (the dotted line) are plotted. (b) a is scaled by a_0 , where a_0 is the asymptotic values of a at $e \rightarrow 0$. In this case, e is uniquely determined by a/a_0 (the solid line). The fitting formula, Eq. (43), which is presented in section 5 is also plotted with the dashed line.




Fig. 4.— The ratios of specific orbital angular momentum (energy) of local gas (l_{gas} , ϵ_{gas}) to those of the planet (l_{p} , ϵ_{p}) as a function of t/T_{K} . For (a) $e = 0.9$ and (b) 0.2 , $l_{\text{gas}}/l_{\text{p}}$ (the solid lines), $\epsilon_{\text{gas}}/\epsilon_{\text{p}}$ (the dashed lines) and $(\epsilon_{\text{gas}} - \epsilon_{\text{coll}})/\epsilon_{\text{p}}$ (the dotted lines) are plotted. The pericenter and apocenter correspond to $t/T_{\text{K}} = 0$ and $t/T_{\text{K}} = 0.5$, respectively.




Fig. 5.— The ratio $a_{\text{final}}/q_{\text{ini}}$ as a function of e_{ini} , where a_{final} is asymptotic semimajor axis after the orbital circularization and q_{ini} is initial pericenter distance before the circularization.

Since Eq. (18) has a similar structure to Eq. (16), the evolution of e is uniquely given as a function of a/a_{ini} . Figure 3a shows the evolutionary paths on the a - e plane for representative values of e_{ini} . Because both e and a keep decreasing, the evolution starts at the right end and moves leftward. Damping of e is dominated over that of a except for $e \simeq 1$. Even if $e_{\text{ini}} = 0.99$, the asymptotic value of a for $e \rightarrow 0$, which we denote as a_{final} , is as much as $\sim 0.48a_{\text{ini}}$. As in the case of e - M relation, if the semimajor axis is scaled by that at a specific value of e , the evolution of e is expressed by a single curve, irrespective of e_{ini} and a_{ini} . Figure 3b shows the self-similar relation, a/a_0 as a function of e (the solid curve), where a_0 is a at $e = 0$. The fitting formula given by Eq. (43), which is presented in section 5 is also plotted with the dashed curve. This plot clearly shows that damping of a is much smaller than that of e : for the damping of e from ~ 1 to 0, a is decreased by $\sim 50\%$, and for that from 0.8 to 0, the decrease in a is only $\sim 30\%$.

Figure 4a shows specific orbital angular momentum and energy of local gas (ℓ_{gas} and ϵ_{gas}) scaled by those of the planet (ℓ_{p} and ϵ_{p}) for $e = 0.9$ as functions of t/T_{K} . The pericenter passage is at $t/T_{\text{K}} = 0$ and $t/T_{\text{K}} = 1$, and the apocenter passage is at $t/T_{\text{K}} = 0.5$, respectively. Because the orbit is highly eccentric, in most of time of an orbital period, $\ell_{\text{gas}} > \ell_{\text{p}}$ and $\epsilon_{\text{gas}} > \epsilon_{\text{p}}$ (since the energy is negative, $|\epsilon_{\text{gas}}| < |\epsilon_{\text{p}}|$) except in the regions close to pericenter ($t/T_{\text{K}} = 0$ and $t/T_{\text{K}} = 1$). As a result, an orbit-averaged value of $\ell_{\text{gas}}/\ell_{\text{p}}$ is considerably larger than unity (in this case, it is $\langle \ell_{\text{gas}}/\ell_{\text{p}} \rangle = f_{\ell}(0.9, \pi) + 1 = 2.66$) and the specific angular momentum of the planet increases through accretion of disk gas.

On the other hand, it is analytically shown that an orbit-averaged value of $\epsilon_{\text{gas}}/\epsilon_{\text{p}}$ is unity, since

$$\left\langle \frac{\epsilon_{\text{gas}}}{\epsilon_{\text{p}}} \right\rangle = \frac{1}{T_{\text{K}}} \int_{-T_{\text{K}}/2}^{T_{\text{K}}/2} \left(\frac{\epsilon_{\text{gas}}}{\epsilon_{\text{p}}} \right) dt = \frac{1}{T_{\text{K}}} \int_{-T_{\text{K}}/2}^{T_{\text{K}}/2} \left(\frac{r}{a} \right)^{-1} dt = f_{-1}(e, \pi) = \frac{\pi}{\pi - e \sin \pi} = 1. \quad (34)$$

Although $|\epsilon_{\text{gas}}| < |\epsilon_{\text{p}}|$ in most of time, $|\epsilon_{\text{gas}}|$ is much larger than $|\epsilon_{\text{p}}|$ near the pericenter passage (Fig. 4a). The contribution of large $|\epsilon_{\text{gas}}|$ compensates for the excess energy accretion in outer regions. But, the orbital energy of the planet decreases because it is also contributed by the collisional energy dissipation $\epsilon_{\text{coll}} (= v_{\text{rel}}^2/2)$. Equation (12) shows that the collisional energy dissipation also damps e . If the collisional energy dissipation is neglected, a is conserved and e is damped by the accretion of higher specific angular momentum gas. With the effect of the collisional energy dissipation, damping of e is faster and a is also damped while the a -damping is slower than the e -damping.

From these relations, together with $\epsilon_{\text{p}} = -GM_*/2a$ and $\ell_{\text{p}} = \sqrt{GM_*a(1-e^2)}$, it is readily found that both a and e always decrease in the case of constant gas accretion rate. Figure 4b shows $\ell_{\text{gas}}/\ell_{\text{p}}$ and $\epsilon_{\text{gas}}/\epsilon_{\text{p}}$ at $e = 0.2$. In this case, the integrals are more symmetric about $\ell_{\text{gas}}/\ell_{\text{p}} = 1$ and $\epsilon_{\text{gas}}/\epsilon_{\text{p}} = 1$. Thereby, when e is reduced to $\lesssim 0.2$, the orbit-averaged values of $\ell_{\text{gas}}/\ell_{\text{p}}$ and $\epsilon_{\text{gas}}/\epsilon_{\text{p}}$ are nearly unity and the decrease in e and a due to planetary mass growth slows down.

Initial pericenter distance q_{ini} of the core's orbit before the e -damping process would correspond to the original semimajor axis of the core (a_{ori}) before the core was scattered by a gas giant, which

may be $\sim 1\text{--}10\text{AU}$. Figure 5 shows asymptotic semimajor axis a_{final} scaled by q_{ini} . Because a_{final} is the final semimajor axis of a gas giant formed from a scattered core after e is damped, $a_{\text{final}}/q_{\text{ini}}$ indicates an efficiency to send a planet to outer regions. In this figure, we find that a core originally at inner region ($a_{\text{ori}} \sim q_{\text{ini}} \sim 10\text{AU}$) become a gas giant with large radius ($a_{\text{final}} \gtrsim 30\text{AU}$ when $e_{\text{ini}} \gtrsim 0.73$, and $a_{\text{final}} \gtrsim 100\text{AU}$ when $e_{\text{ini}} \gtrsim 0.94$).

Figure 6 shows the evolution of pericenter distance scaled by initial one, q/q_{ini} , due to planetary mass growth, for representative values of e_{ini} . It is shown that q/q_{ini} quickly increases, which justifies our assumption that the scattered planet becomes quickly isolated and the further perturbations from the gas giant in inner region are neglected.

4.2. Deviated case

Next, we consider deviated case in which $Q > r_{\text{d}}$. Since in this case, gas accretion is halted at $r > r_{\text{d}}$, the planet cannot accrete gas with higher specific angular momentum and energy, resulting in smaller f_{ℓ} and larger $f_{\epsilon}(= f_a)$ (see Fig. 1). The increase of $f_{\epsilon}(= f_a)$ is more effective than the decrease of f_{ℓ} , so f_e is larger. Thus, both e and a dampings in the deviated case are more efficient than in the embedded case.

Since there is a characteristic length r_{d} , a self-similar solution like Fig. 3b does not exist. However, it is clear that evolutions of Q/r_{d} and q/r_{d} should be the same for the same initial values. Figure 7 shows the evolutions of Q/r_{d} and q/r_{d} . The evolutions are to the right-down direction. The evolutions in deviated case correspond to those in the region of $Q > r_{\text{d}}$. We also added following embedded evolutions in the region of $Q < r_{\text{d}}$. Because the evolution paths do not cross each other, we can parameterize the evolution paths with one parameter. In Fig. 7, we used the value of q/r_{d} at the time when Q is reduced to be r_{d} , as the parameter.

The evolutions of e and a corresponding to the solutions in Figure 7 are plotted in Figure 8. The orbital evolution is toward the left-down direction. This figure shows that in the early phase of $Q > r_{\text{d}}$, the semimajor axis is predominantly damped. Note that even in this phase where a is rapidly reduced, Figure 7 shows that q is increased so quickly that the planet becomes isolated from the perturbing gas giant. For $e_{\text{ini}} \sim 1$, a is damped by order of magnitude until $Q \sim a(1 + e) \sim 2a$ is reduced to $\sim r_{\text{d}}$. In the following embedded phase of $Q < r_{\text{d}}$, however, a is reduced at most by a factor of 2, as we showed. Thus, in this case, $a_{\text{final}} \sim r_{\text{d}}/4$, independent of the values of a_{ini} , as long as $e_{\text{ini}} \sim 1$. In other words, we can infer the values of r_{d} from a_{final} .

Figure 9 shows the evolution of e as a function of the planetary mass M/M_Q , where M_Q is the planetary mass at $Q = r_{\text{d}}$. In the deviated phase, the e -damping is more efficient than in the embedded phase, although it is slightly slower because of high eccentricity.




Fig. 6.— The evolution of q/q_{ini} as a function of M/M_{ini} . $e_{\text{ini}} = 0.8$ (the solid line), 0.9 (the dashed line) and 0.99 (the dotted line) are plotted.




Fig. 7.— The evolution of apocenter distance Q and pericenter distance q . Both are scaled by the disk size r_d . The evolution paths are parameterized by the values of q/r_d at $Q = r_d$ at which the deviated-case evolution is switched to embedded-case evolution; the paths with $q/r_d|_{Q=r_d} = 0.01$ (the solid line), 0.03 (the dashed line), 0.1 (the dotted line) and 0.3 (the dot-dashed line) are plotted.




Fig. 8.— The evolutions of e and a/r_d corresponding to the solutions in Fig. 7.




Fig. 9.— The evolution of e as a function of M/M_Q corresponding to the solutions in Fig. 7, where M_Q is the planetary mass at $Q = r_d$.

4.3. Effect of shock

We have considered the energy dissipation by collision between incident gas flow and the planet. The dissipation is needed to bind the gas around the planet and it accelerates eccentricity damping as we showed. The dissipation should occur through bow shock in front of the planet. The shock not only causes the energy dissipation but also makes the relative velocity lower. So far, we have neglected the relative velocity damping by shock, which should weaken the eccentricity and semimajor axis damping. Here we evaluate the effect of the shock using a simple 1D model. Full 2D or 3D hydrodynamical simulations will be done in a separate paper.

The simple 1D model we use is as follows. The ratio of post-shock velocity (sv_{rel}) to pre-shock one (v_{rel}) is

$$s = \frac{(\gamma - 1)\mathcal{M}^2 + 2}{(\gamma + 1)\mathcal{M}^2} \simeq \frac{1}{4} \left(1 + \frac{3}{\mathcal{M}^2} \right), \quad (35)$$

where γ is the specific heat ratio ($\gamma = 5/3$ in the monatomic molecule) and \mathcal{M} is Mach number for pre-shock gas flow, which is given by

$$\mathcal{M} = \frac{v_{\text{rel}}}{c_s} = 30 \left(\frac{r}{1\text{AU}} \right)^{-1/4} \left(3 - \frac{r}{a} - 2\sqrt{\frac{a}{r}(1 - e^2)} \right)^{1/2}, \quad (36)$$

where we used an optically thin disk temperature, $T = 280(r/1\text{AU})^{-1/2}\text{K}$, for evaluation of sound velocity c_s . For subsonic case ($\mathcal{M} < 1$), $s = 1$. The radial and tangential components of velocity of post-shock gas (u_r, u_ϕ) and those of pre-shock gas ($0, v_{\text{K}}$) are related to those of planet (v_r, v_ϕ), which are given by Eqs.(22) and (23), as

$$u_r - v_r = s(0 - v_r), \quad (37)$$

$$u_\phi - v_\phi = s(v_{\text{K}} - v_\phi). \quad (38)$$

Then, the integrands of Eqs. (8) and (9) for f_ℓ and f_ϵ are

$$\begin{aligned} \frac{l_{\text{gas}}}{l_{\text{p}}} - 1 &= \frac{ru_\phi}{rv_\phi} - 1 = s \left(\frac{rv_{\text{K}}}{rv_\phi} - 1 \right) = s \left(\frac{l_{\text{gas},0}}{l_{\text{p}}} - 1 \right), \quad (39) \\ \frac{\epsilon_{\text{gas}}}{\epsilon_{\text{p}}} - \frac{\epsilon_{\text{coll}}}{\epsilon_{\text{p}}} - 1 &= \frac{(u_r^2 + u_\phi^2)/2 - GM_*/r - s^2v_{\text{rel}}^2/2}{(v_r^2 + v_\phi^2)/2 - GM_*/r} - 1 \\ &= s \left(\frac{v_{\text{K}}^2/2 - GM_*/r - v_{\text{rel}}^2/2}{(v_r^2 + v_\phi^2)/2 - GM_*/r} - 1 \right) = s \left(\frac{\epsilon_{\text{gas},0}}{\epsilon_{\text{p}}} - \frac{\epsilon_{\text{coll},0}}{\epsilon_{\text{p}}} - 1 \right), \quad (40) \end{aligned}$$

where quantities with subscript ',0' means those for unperturbed gas flow neglecting shock ($l_{\text{gas},0} = \sqrt{GM_*r}$, $\epsilon_{\text{gas},0} = -GM_*/2r$, $\epsilon_{\text{coll},0} = v_{\text{rel}}^2/2$).

As we showed in the previous subsections, the integrations for f_ℓ and f_ϵ (Eqs. (8) and (9)) can be analytically done in the case neglecting the damping of the relative velocity (equivalently, $s = 1$). However, since in the present case, s varies along the orbit (Fig. 10), we integrate Eqs. (8)

and (9) numerically. Here we consider embedded case. Since f_ℓ and f_ϵ depend on a through s in the present case, we assume $a = 100\text{AU}$ (a is variable when we consider the evolution paths). Note that evolution paths in the a - e plane is the same as those in the case without the relative velocity damping. In Eq. (18), $d \log a/de$, is given approximately by orbit-averaged f_a and f_e . But, more exactly, $d \log a/de$ must be integrated every time. From Eqs. (39) and (40), it is apparent that s completely cancels and $d \log a/de$ is the same.

The functions f_ℓ , $f_a(= f_\epsilon)$ and f_e in the case with shock are compared with those without shock in Fig. 11. The effect of shock lowers all the functions. Accordingly, both e and a dampings are slowed down, while the evolution paths on the a - e plane do not change. The evolution paths on the e - M , a - M , and q - M planes are shown in Fig. 12. The initial conditions are $q_{\text{ini}} = 10\text{AU}$, $e_{\text{ini}} = 0.9$ and $M_{\text{ini}} = 10M_\oplus$, respectively. While e declines to values < 0.2 at $M_p \sim 50M_\oplus$ in non-shock case, it does not becomes < 0.2 until $M_p \sim 3000M_\oplus$ in shock case. However, since the masses of direct-imaged planets are relatively large ($\sim 10M_J$), the orbital circularization is still effective. The a damping is also slowed down, but the semimajor axis at $M_p \sim 10M_J$ is not significantly larger than that in non-shock case. The pericenter distance q is still quickly increased, so that the assumption that the scattered planet becomes quickly isolated and the further perturbations from the gas giant in inner region are neglected is justified. Thus, although the eccentricity damping is less efficient, formation of distant gas giants in nearly circular orbits is not significantly inhibited by the effect of shock.




Fig. 10.— The time dependence of s in one orbit where s is the ratio of velocity of after and before shock. $e = 0.9$ (the solid line), 0.5 (the dashed line) and 0.2 (the dotted line) are plotted. The pericenter and apocenter correspond to $t/T_K = 0$ and $t/T_K = 0.5$, respectively. We assume an optically thin disk temperature, $T = 280(r/1\text{AU})^{-1/2}\text{K}$, and $a = 100\text{AU}$.




Fig. 11.— The functions $f_l, f_a(=f_\epsilon)$ and f_e . The solid lines and dotted lines represent the functions without shock and with shock, respectively. We assume $a = 100\text{AU}$ in shock case.




Fig. 12.— The evolution paths on the e - M , a - M , and q - M planes. The initial conditions are $q_{ini} = 10\text{AU}$, $e_{ini} = 0.9$ and $M_{ini} = 10M_{\oplus}$, respectively. The solid lines and dotted lines represent the functions without shock and with shock, respectively.

5. Fitting formulas and population synthesis simulation

The self-similar solution in Fig. 2b can be approximately fitted by

$$e \simeq 0.1 \left[1 - \log_{10} \left(\frac{M}{M_{01}} \right) \right]^3. \quad (41)$$

where M_{01} is M at $e = 0.1$. Then, for any given e_{ini} and M_{ini} , e for $M (> M_{\text{ini}})$ are evaluated by

$$\begin{aligned} e &\simeq 0.1 \left[1 - \log_{10} \left(\frac{M_{\text{ini}}}{M_{01}} \right) - \log_{10} \left(\frac{M}{M_{\text{ini}}} \right) \right]^3 \\ &= 0.1 \left[\left(\frac{e_{\text{ini}}}{0.1} \right)^{1/3} - \log_{10} \left(\frac{M}{M_{\text{ini}}} \right) \right]^3. \end{aligned} \quad (42)$$

Because this is an approximate formula, Eq. (42) can be negative for large values of M/M_{ini} . In such cases, we set $e = 0$, because e has very small values in the exact solution. On the other hand, the self-similar solution in Fig. 3b can be fitted by

$$\frac{a}{a_0} \simeq 1 + 0.6e^3 + \frac{0.007}{1 - e}. \quad (43)$$

where a_0 is the asymptotic values of a at $e \rightarrow 0$. Since a similar relation holds for a_{ini} and e_{ini} ,

$$\frac{a}{a_{\text{ini}}} \simeq \frac{1 + 0.6e^3 + 0.007/(1 - e)}{1 + 0.6e_{\text{ini}}^3 + 0.007/(1 - e_{\text{ini}})}. \quad (44)$$

In the population synthesis simulation, when a core with mass M_c closely encounters with a gas giant, eccentricity and semimajor axis that are excited by the scattering are evaluated with a Monte-Carlo procedure (see, e.g., Ida et al. 2013). We set these eccentricity, semimajor axis, and M_c as e_{ini} , a_{ini} , and M_{ini} in the above equations, respectively. The mass growth of the planet due to gas accretion after the scattering is also calculated in the population synthesis simulation. The mass growth is truncated when disk gas is severely depleted or a clear gap along the planetary orbit is opened (see, e.g., Ida & Lin 2004a; Ida et al. 2013). From Eq. (42), the value of e when the planet mass increases to M can be derived from e_{ini} and M_{ini} . The semimajor axis a at M is derived from e_{ini} , e and a_{ini} from Eq. (44). Note that Ida et al. (2013) simply assumed $e = 0$ and $a = a_{\text{ini}}$.

Figures 13 show the e - a distributions of gas giant planets around solar-type stars, obtained by a population synthesis calculation with similar parameters to those of the results in Fig. 7 of Ida et al. (2013). Note that rocky and icy planets with smaller masses are not plotted here. In panel a, neither the dynamical friction to cores nor the damping via gas accretion is included. Most of giant planets at $\gtrsim 30\text{AU}$ have large eccentricities, because they suffered strong gravitational scattering by other giants. In panel b, only the dynamical friction is included. Eccentricities of small number of planets are damped, but the effects are not significant. On the other hand, both effects are included in panel c. Eccentricities are damped to values below 0.2 for $\sim 30\%$ of giant




Fig. 13.— The distributions of e and a of gas giant planets around solar-type stars, obtained by population synthesis calculations with similar parameters to those of the results in Fig. 7 of Ida et al. (2013). For the details of calculations, see Ida et al. (2013). In panel a, neither the dynamical friction to cores nor the damping via gas accretion is included. In panel b, only the dynamical friction is included. Both effects are included in panel c.

planets. However, since the damping is not as efficient as the simple treatment in Ida et al. (2013), the fraction of systems that have gas giants with $a > 30\text{AU}$ and $e < 0.2$ is $\sim 0.1\%$, which is smaller than the probability ($\sim 0.4\%$) in Fig. 7 of Ida et al. (2013). It is a future problem to check if such low fraction is consistent with direct imaging surveys. As already pointed out in Ida et al. (2013), the formation rate of high eccentricity gas giants at $\sim O(1)\text{AU}$ is lower in the theoretical prediction than that found by radial velocity surveys. If the theoretical prediction is improved so that more frequent formation of high eccentricity gas giants is reproduced, the theoretically predicted fraction of systems with distant gas giants in nearly circular orbits may also be increased.

6. Summary

We have investigated orbital circularization due to planet growth through accreting disk gas. We have analytically derived the differential equations for evolutions of orbital eccentricity e and semimajor axis a and numerically integrated them to discuss the solutions.

The motivation of these calculations is to examine our scenario for the formation of the distant gas giants in nearly circular orbits, which are recently being discovered by direct imaging surveys. Our scenario is based on the conventional core accretion model as follows: i) Icy cores accrete from planetesimals in inner regions at $a \lesssim 30\text{AU}$, ii) they are scattered outward by a nearby gas giant to acquire highly eccentric orbits, iii) their orbits are circularized through accretion of local protoplanetary disk gas, and iv) through the local gas accretion, the planets become gas giants. We started our calculations after step ii). For given initial e and a , we followed the process in step iii).

For highly eccentric orbits, the planet spends most of time in the outer regions where disk gas has higher specific orbital angular momentum than the planet. Since the gas accretion rate from the disk is regulated by envelope contraction except for final gas accretion phase, we assume that disk gas accretion rate is constant within one orbit. Even in the final phase when the accretion rate is limited by the supply of gas due to global disk accretion, the assumption is valid if steady disk accretion is established. Thus, specific angular momentum of the planet increases with planet accretion, resulting in circularization of the planetary orbit. Energy dissipation by collision between disk gas and the planet also induces the eccentricity damping.

Just after step ii), core's pericenter distance must be close to its original location. We found that pericenter distance is quickly raised by the orbital circularization, so that perturbations of the gas giant in the inner region can be neglected in the orbital circularization process. Thereby, we investigated orbital evolution of isolated planets accreting disk gas.

The orbital evolutions that we found are:

- 1) The eccentricity is reduced to < 0.2 before the planetary mass is increased by a factor of 10 (for example, if an icy core with $\sim 10M_{\oplus}$ starts gas accretion, its orbit is circularized with $e < 0.2$

before it acquires a Saturnian-mass.)

2) The eccentricity damping is dominated over the semimajor axis damping. During e is reduced from ~ 1 to zero, a is decreased only by a factor of 2.

These show that planetary growth and orbital circularization concurrently proceed and the orbital circularization is very efficient. If we take into account the effect of bow shock for supersonic incident gas flow, the orbital circularization becomes slower, but it is still efficient enough to account for the observed orbital properties of distant gas giants. The orbit is left in large orbital radii, which are about half of the semimajor axes that the scattered icy cores initially acquire.

We performed the population synthesis calculation by incorporating the fitting formulas for the eccentricity and semimajor axis damping by planet mass growth to indeed show that the damping is efficient and giants with $e \lesssim 0.2$ are left in distant regions at $a \sim 30\text{--}300\text{AU}$. However, with more detailed prescription using the formulas derived here, the fraction of systems that have such distant jupiters is as small as $\sim 0.1\%$, which is lower by a factor 4 than that predicted in Ida et al. (2013) using simpler prescription.

We also consider the effect of the finite disk size. If the eccentric orbits of the scattered cores are deviated from the protoplanetary disk near their apocenters, their semimajor axes shrink to a quarter of the disk sizes. In other words, if observations show a concentration of distant gas giants at some orbital radius, it could reflect typical sizes of the protoplanetary disks, in a similar way that the pile-up location of hot jupiters could reflect the size of magnetospheric cavity (the size of the disk inner edge) where type II migration could be stalled.

We thank Prof. Andrew Youdin for valuable and useful comments as a referee. We also thank Takayuki Muto and Taku Takeuchi for discussions. Our study was supported by JSPS KAKENHI Grant Number 23103005.

REFERENCES

- Bodenheimer, P., & Pollack, J. B. 1986, *Icar*, 67, 391
- Boss, A. 2001, *ApJ*, 551, L167
- Crida, A. et al. 2009, *ApJ*, 705, L148
- Fischer D. A. & Valenti J. 2005, *ApJ*, 622, 1102
- Forgan, D. & Rice, K. 2013, *MNRAS*, 432, 3168
- Hayashi, C. 1981, *Prog. Theor. Phys. Suppl.*, 70, 35
- Helled et al. 2014, a chapter in *Protostars and Planets VI* (Univ. Arizona Press), in press, 2013arXiv1311.1142H

- Ida, S., & Lin, D. N. C. 2004a, *ApJ*, 604, 388
- Ida, S., & Lin, D. N. C. 2004b, *ApJ*, 616, 567
- Ida, S., Lin, D. N. C. & Nagasawa, M. 2013, *ApJ*, 775, 42
- Ikoma, M., Nakazawa, K., & Emori, H. 2000, *ApJ*, 537, 1013
- Ikoma, M., Emori, H. & Nakazawa, K. 2001, *ApJ*, 553, 999
- Ikoma, M., & Genda, H., 2006. *ApJ*, 648, 696
- Kalas, P., Graham, J. R., Chiang, E., et al. 2008, *Science*, 322, 1345
- Kokubo, E., & Ida, S. 1998, *Icar*, 131, 171
- Kratter, K. M., Murray-Clay, R. A. & Youdin, A. N. 2010, *ApJ*, 710, 1375
- Kuzuhara et al. 2013, *ApJ*, 763, L32
- Lambrechts, M. & Johansen, A. 2012, *A&A*, 544, A32
- Marois et al. 2008, *Science*, 322, 1348
- Marzari, F., & Weidenschilling, S. J. 2002, *Icarus*, 156, 570
- Mizuno, H. 1980, *Prog. Theor. Phys.*, 64, 544
- Muto, T., Takeuchi, T., & Ida, S. 2011, *ApJ*, 737, 37
- Nagasawa, M., Ida, S., & Bessho, T. 2008, *ApJ*, 678, 498
- Ostriker, E. C. 1999, *ApJ*, 513, 252
- Papaloizou, J. C. B., & Larwood, J. D. 2000, *MNRAS*, 315, 823
- Piso, Ana-Maria A. & Youdin, Andrew N. 2014, *ApJ*, 786, 21
- Shiraishi, M., & Ida, S. 2008, *ApJ*, 684, 1416
- Youdin, A. & Kenyon, S. 2012, a chapter in *Planets, Stars and Stellar Systems* (Springer Science+Business Media Dordrecht), 1
- Zhou, J., Lin, D. N. C., & Sun, Y. 2007, *ApJ*, 666, 423