

# Multispecies quantum Hurwitz numbers\*

J. Harnad<sup>1,2</sup>

<sup>1</sup> *Centre de recherches mathématiques, Université de Montréal  
C. P. 6128, succ. centre ville, Montréal, QC, Canada H3C 3J7  
e-mail: harnad@crm.umontreal.ca*

<sup>2</sup> *Department of Mathematics and Statistics, Concordia University  
7141 Sherbrooke W., Montréal, QC Canada H4B 1R6*

## Abstract

The construction of hypergeometric 2D Toda  $\tau$ -functions as generating functions for quantum Hurwitz numbers is extended here to multispecies families. Both the enumerative geometrical significance of these multispecies quantum Hurwitz numbers as weighted enumerations of branched coverings of the Riemann sphere and their combinatorial significance in terms of weighted paths in the Cayley graph of  $S_n$  are derived.

## 1 Introduction

In [4, 5, 8] a general method was developed for constructing parametric families of 2D Toda  $\tau$ -functions of hypergeometric type [17, 15, 16, 14] that are generating functions for weighted enumeration of branched coverings of the Riemann sphere. These were shown to be interpretable equivalently as weighted sums over paths in the Cayley graph of the symmetric group  $S_n$  generated by transpositions.

The notion of 2D Toda  $\tau$ -functions of generalized hypergeometric type, as introduced in [14], included families which, when the flow variables are restricted to the trace invariants of a pair of  $N \times N$  matrices, may be understood as hypergeometric functions of matrix arguments [3]. The combinatorial significance of the coefficient in the double power sum symmetric function expansions was explained in [4, 5] in terms of path counting in the Cayley graph consisting of strictly and weakly monotonically increasing subsequences of transpositions having given lengths. This was shown in [8] to be equivalent in geometrical terms to the (signed) enumeration of branched covers of the Riemann sphere with multispecies branch points constrained to have fixed total ramification index within each species.

Let  $H(\mu^{(1)}, \dots, \mu^{(k)})$ ,  $|\mu^{(i)}| = n$ , denote the number of  $n$ -sheeted branched coverings of the Riemann sphere (not necessarily connected), with  $k$  branch points, whose ramification profiles are given by the  $k$  partitions  $(\mu^{(1)}, \dots, \mu^{(k)})$  of  $n$ ; the so-called Hurwitz numbers. The

---

\*Work of JH supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds de recherche du Québec - Nature et technologies (FRQNT); work of AO supported by RFBR grant 14-01-00860.

Frobenius-Schur formula [10, 6] expresses these in terms of the irreducible group characters for  $S_n$

$$H(\mu^{(1)}, \dots, \mu^{(k)}) = \sum_{\lambda, |\lambda|=n} h_\lambda^{k-2} \prod_{i=1}^k \frac{\chi_\lambda(\mu^{(i)})}{Z_{\mu^{(i)}}}, \quad (1.1)$$

where  $\chi_\lambda(\mu)$  is the character of the irreducible representation of symmetry type  $\lambda$  evaluated on the conjugacy class  $\text{cyc}(\mu)$  consisting of elements with cycle lengths equal to the parts of the partition  $\mu$ ,

$$h_\lambda = \det \left( \frac{1}{(\lambda_i - i + j)!} \right)^{-1} \quad (1.2)$$

is the product of the hook lengths in the Young diagram of partition  $\lambda$  and

$$Z_\mu = \prod_{i \in Nb} i^{j_i} (j_i)!, \quad (1.3)$$

where  $j_i$  the number of parts of the partition  $\mu$  equal to  $i$ , is the order of the stabilizer under conjugation of any element of the conjugacy class  $\text{cyc}(\mu)$ .

As defined in [5], weighted double Hurwitz numbers  $H_G^d(\mu, \nu)$  for  $n$  sheeted branched coverings of the sphere, corresponding to a weight generating function  $G$ , that include a pair of fixed branch points with profile types given by the pair of partitions  $(\mu, \nu)$  at, say,  $(0, \infty)$ , and genus  $g$  given by the Riemann Hurwitz formula

$$2 - 2g = \ell(\mu) + \ell(\nu) - d, \quad (1.4)$$

$$d = \sum_{i=1}^k \ell^*(\mu^{(i)}), \quad \ell^*(\mu) := |\mu| - \ell(\mu) \quad (1.5)$$

are given by the weighted sum

$$H_G^d(\mu, \nu) := \sum_{k=0}^{\infty} \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)})=d}} W_G(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu), \quad (1.6)$$

over all  $k$ -tuples of nontrivial ramification profiles with weight given by  $W_G(\mu^{(1)}, \dots, \mu^{(k)})$ , satisfying the condition (1.5).

A special class of weighted Hurwitz numbers was introduced in [5], depending on an auxiliary quantum deformation parameter  $q$ , and the weighted counting of branched covers was related to the energy distributions of a Bosonic gas. In the present work, we extend these results to multispecies generalizations of such quantum Hurwitz numbers and construct multiparametric families of 2D Toda  $\tau$ -functions, depending on an arbitrary number of auxiliary parameters, that may be viewed as generating functions for multispecies quantum Hurwitz numbers, with independent quantum deformation parameters  $\{q_i, p_j\}$  for each species. The weightings may be also be interpreted in terms of the energy distributions of noninteracting multispecies mixtures of Bose gases.

## 1.1 Quantum Hurwitz numbers

Quantum Hurwitz numbers were introduced in [5], in four variants, and a 1-parameter family of  $2D$  Toda  $\tau$ -function generating functions was constructed for each. In the first case, the weight generating function is:

$$E(q, z) := \prod_{i=0}^{\infty} (1 + q^i z) = 1 + \sum_{i=1}^{\infty} E_i(q) z^i, \quad (1.7)$$

$$E_i(q) := \left( \prod_{j=1}^i \frac{q^{j-1}}{1 - q^j} \right). \quad (1.8)$$

The second is a slight modification of this, with weight generating function

$$E'(q, z) := \prod_{i=1}^{\infty} (1 + q^i z) = 1 + \sum_{i=1}^{\infty} E'_i(q) z^i, \quad (1.9)$$

$$E'_i(q) := \left( \prod_{j=1}^i \frac{q^j}{1 - q^j} \right). \quad (1.10)$$

The third case is based on the weight generating function

$$H(q, z) := \prod_{i=1}^{\infty} (1 - q^i z)^{-1} = 1 + \sum_{i=1}^{\infty} H_i(q) z^i, \quad (1.11)$$

$$H_i(q) := \left( \prod_{j=1}^i \frac{1}{1 - q^j} \right), \quad (1.12)$$

and the final case is a hybrid, formed from the product of the first and third for two distinct quantum deformation parameters  $q$  and  $p$ , with weight generating function

$$Q(q, p, z) := \prod_{k=0}^{\infty} (1 + q^k z) (1 - p^k z)^{-1} = \sum_{i=0}^{\infty} Q_i(q, p) z^i, \quad (1.13)$$

$$Q_i(q, p) := \sum_{m=0}^i q^{\frac{1}{2}m(m-1)} \left( \prod_{j=1}^m (1 - q^j) \prod_{j=1}^{i-m} (1 - p^j) \right)^{-1}, \quad Q_{\lambda}(q, p) = \prod_{i=1}^{\ell(\lambda)} Q_{\lambda_i}(q, p) \quad (1.14)$$

These are all expressible as exponentials of the quantum dilogarithm function

$$\text{Li}_2(q, z) := \sum_{k=1}^{\infty} \frac{z^k}{k(1 - q^k)}. \quad (1.15)$$

$$E(q, z) = e^{-\text{Li}_2(q, -z)}, \quad E'(q, z) = (1 + z)^{-1} e^{-\text{Li}_2(q, -z)} \quad (1.16)$$

$$H(q, z) = e^{\text{Li}_2(q, z)}, \quad Q(q, p, z) = e^{\text{Li}_2(p, z) - \text{Li}_2(q, -z)}, \quad . \quad (1.17)$$

The content products for the first and third of these are defined to be

$$r_\lambda^{E(q,z)}(N) := \prod_{(ij) \in \lambda} E(q, (N + j - i)z) \quad (1.18)$$

$$r_\lambda^{H(q,z)}(N) := \prod_{(ij) \in \lambda} H(q, (N + j - i)z). \quad (1.19)$$

The associated hypergeometric 2D Toda  $\tau$ -functions have diagonal double Schur function expansions with these as coefficients:

$$\tau^{E(q,z)}(N, \mathbf{t}, \mathbf{s}) = \sum_\lambda r_\lambda^{E(q,z)}(N) S_\lambda(\mathbf{t}) S_\lambda(\mathbf{s}) \quad (1.20)$$

$$\tau^{H(q,z)}(N, \mathbf{t}, \mathbf{s}) = \sum_\lambda r_\lambda^{H(q,z)}(N) S_\lambda(\mathbf{t}) S_\lambda(\mathbf{s}). \quad (1.21)$$

Using the Frobenius character formula [11],

$$S_\lambda = \sum_{\mu, |\mu|=|\lambda|} \frac{\chi_\lambda P_\mu}{Z_\mu} \quad (1.22)$$

and setting  $N = 0$ , these may be rewritten as double expansions in the power sum symmetric functions [5]:

$$\tau^{E(q,z)}(\mathbf{t}, \mathbf{s}) := \tau^{E(q,z)}(0, \mathbf{t}, \mathbf{s}) = \sum_{d=0}^{\infty} z^d \sum_{\mu, \nu, |\mu|=|\nu|} H_{E(q)}^d(\mu, \nu) P_\mu(\mathbf{t}) P_\nu(\mathbf{s}), \quad (1.23)$$

$$\tau^{H(q,z)}(\mathbf{t}, \mathbf{s}) := \tau^{H(q,z)}(0, \mathbf{t}, \mathbf{s}) = \sum_{d=0}^{\infty} z^d \sum_{\mu, \nu, |\mu|=|\nu|} H_{H(q)}^d(\mu, \nu) P_\mu(\mathbf{t}) P_\nu(\mathbf{s}). \quad (1.24)$$

The coefficients are the corresponding quantum Hurwitz numbers  $H_{E(q)}^d(\mu, \nu)$ ,  $H_{H(q)}^d(\mu, \nu)$ , which count weighted  $n$ -sheeted branched coverings of the Riemann sphere, defined by

$$H_{E(q)}^d(\mu, \nu) := \sum_{k=0}^{\infty} \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} W_{E(q)}(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu), \quad (1.25)$$

$$H_{H(q)}^d(\mu, \nu) := \sum_{k=0}^{\infty} (-1)^{k+d} \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} W_{H(q)}(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu). \quad (1.26)$$

where the weightings for such covers with  $k$  additional branch points are:

$$W_{E(q)}(\mu^{(1)}, \dots, \mu^{(k)}) = \frac{1}{k!} \sum_{\sigma \in S_k} \frac{q^{(k-1)\ell^*(\mu^{(1)})} \dots q^{\ell^*(\mu^{(k-1)})}}{(1 - q^{\ell^*(\mu^{(\sigma(1))})}) \dots (1 - q^{\ell^*(\mu^{(\sigma(k))})}) \dots q^{\ell^*(\mu^{(\sigma(k))})}}, \quad (1.27)$$

$$W_{H(q)}(\mu^{(1)}, \dots, \mu^{(k)}) = \frac{1}{k!} \sum_{\sigma \in S_k} \frac{1}{(1 - q^{\ell^*(\mu^{(\sigma(1))})}) \dots (1 - q^{\ell^*(\mu^{(\sigma(1))})}) \dots q^{\ell^*(\mu^{(\sigma(k))})}}. \quad (1.28)$$

Here

$$\ell^*(\mu) := |\mu| - \ell(\mu) \quad (1.29)$$

is the *colength* of the partition  $\mu$ , which is the index of coalescence of the sheets of the branched cover over the branch points with ramification profiles  $\mu^{(i)}$ , the sum  $\sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}}$  is over all  $k$ -tuples of partitions having nontrivial ramification profiles that satisfy the constraint  $\sum_{i=1}^k \ell^*(\mu^{(i)}) = d$ , and  $H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu)$  is the number of branched  $n$ -sheeted coverings, up to isomorphism, having  $k+2$  branch points with ramification profiles  $(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu)$ .

These thus count weighted covers with a pair of branch points, say  $(0, \infty)$ , having ramification profiles of type  $(\mu, \nu)$  and an arbitrary number of further branch points, whose profiles  $(\mu^{(1)}, \dots, \mu^{(k)})$  are constrained only by the requirement that the sum of the colengths, which is related to the genus by the Riemann-Hurwitz formula

$$\sum_{i=1}^k \ell^*(\mu^{(i)}) = 2g - 2 + \ell(\mu) + \ell(\nu) = d, \quad (1.30)$$

be fixed to equal  $d$ .

Another interpretation that is purely combinatorial can be given to the quantum Hurwitz numbers  $H_{E(q)}^d(\mu, \nu)$  and  $H_{H(q)}^d(\mu, \nu)$  appearing in (1.24), as follows. Let  $(a_1 b_1) \cdots (a_d b_d)$  be a product of  $d$  transpositions  $(a_i b_i) \in S_n$  in the symmetric group  $S_n$  with  $a_i < b_i$ ,  $i = 1, \dots, d$ . If  $h \in S_n$  is in the conjugacy class  $\text{cyc}(\mu) \subset S_n$  consisting of elements with cycle lengths equal to the parts  $(\mu_1, \dots, \mu_{\ell(\mu)})$  of the partition  $\mu$  of weight  $|\mu| = n$ , and length  $\ell(\mu)$ , we may view the successive steps in the product

$$(a_1 b_1) \cdots (a_d b_d) h \quad (1.31)$$

as a path in the Cayley graph generated by all transpositions. To each such path, we attach a *signature* consisting of a partition  $\lambda$  of  $d$ ,  $|\lambda| = d$ , whose parts  $\lambda_i$  consist of the number of transpositions  $(a_i b_i)$  sharing the same second element. If we further require that the ones with equal second elements be grouped together into consecutive subsequences in which these second elements are constant, with the consecutive subsequences strictly increasing in their second elements, then the number  $\tilde{N}_\lambda$  of elements with signature  $\lambda$  is related to the number  $N_\lambda$  of such ordered sequences by

$$\tilde{N}_\lambda = \frac{|\lambda|!}{\prod_{i=1}^{\ell(\lambda)} \lambda_i!} N_\lambda \quad (1.32)$$

Denote the number of such paths from the conjugacy class of cycle type  $\text{cyc}(\mu)$  to the one of type  $\text{cyc}(\nu)$  having signature  $\lambda$  as  $\tilde{m}_{\mu\nu}^\lambda$ , and the number of ordered sequences of type  $\lambda$  as  $m_{\mu\nu}^\lambda$ . Thus

$$\tilde{m}_{\mu\nu}^\lambda = \frac{|\lambda|!}{\prod_{i=1}^{\ell(\lambda)} \lambda_i!} m_{\mu\nu}^\lambda. \quad (1.33)$$

For all paths of signature  $\lambda$  we assign weights

$$\tilde{E}_\lambda(q) = \prod_{i=1}^{\ell(\lambda)} \lambda_i! E_{\lambda_i}(q) = \prod_{i=1}^{\ell(\lambda)} \frac{\lambda_i! q^{\frac{1}{2}\lambda_i(\lambda_i-1)}}{\prod_{j=1}^{\lambda_i} (1-q^j)}, \quad (1.34)$$

$$\tilde{H}_\lambda(q) = \prod_{i=1}^{\ell(\lambda)} \lambda_i! H_{\lambda_i}(q) = \prod_{i=1}^{\ell(\lambda)} \frac{\lambda_i!}{\prod_{j=1}^{\lambda_i} (1-q^j)} \quad (1.35)$$

to paths in the Cayley graph, and a pair of corresponding combinatorial weighted Hurwitz numbers

$$F_{E(q)}^d(\mu, \nu) := \frac{1}{d!} \sum_{\lambda, |\lambda|=d} \tilde{E}_\lambda \tilde{m}_{\mu\nu}^\lambda, \quad (1.36)$$

$$F_{H(q)}^d(\mu, \nu) := \frac{1}{d!} \sum_{\lambda, |\lambda|=d} \tilde{H}_\lambda \tilde{m}_{\mu\nu}^\lambda, \quad (1.37)$$

that give the weighted enumeration of paths, using the weighting factors  $\tilde{E}_\lambda(q)$  and  $\tilde{H}_\lambda(q)$  respectively for all paths of signature  $\lambda$ .

It was shown in [5] that the enumerative geometrical and combinatorial definitions of these quantum weighted Hurwitz numbers coincide:

$$H_{E(q)}^d(\mu, \nu) = F_{E(q)}^d(\mu, \nu), \quad H_{H(q)}^d(\mu, \nu) = F_{H(q)}^d(\mu, \nu). \quad (1.38)$$

A similar result holds for weights generated by the function  $E'(q, z)$ , with corresponding quantum Hurwitz numbers defined by

$$H_{E'(q)}^d(\mu, \nu) := \sum_{k=0}^{\infty} \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)})=d}} W_{E'(q)}(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu), \quad (1.39)$$

where the weights  $W_{E'(q)}(\mu^{(1)}, \dots, \mu^{(k)})$  are given by

$$W_{E'(q)}(\mu^{(1)}, \dots, \mu^{(k)}) := \frac{1}{k!} \sum_{\sigma \in S_k} \frac{q^{(k)\ell^*(\mu^{(1)})} \dots q^{\ell^*(\mu^{(k)})}}{(1 - q^{\ell^*(\mu^{(\sigma(1))})}) \dots (1 - q^{\ell^*(\mu^{(\sigma(1))})} \dots q^{\ell^*(\mu^{(\sigma(k))})})}, \quad (1.40)$$

$$:= \frac{1}{k!} \sum_{\sigma \in S_k} \frac{1}{(q^{-\ell^*(\mu^{(\sigma(1))})} - 1) \dots (q^{-\ell^*(\mu^{(\sigma(1))})} \dots q^{-\ell^*(\mu^{(\sigma(k))})} - 1)}. \quad (1.41)$$

Choosing  $q$  as a positive real number, parametrizing it as

$$q = e^{-\beta h \omega}, \quad \beta = \frac{1}{kT} \quad (1.42)$$

and identifying the energy levels  $\epsilon_k$  as those for a Bose gas with linear spectrum in the integers, as for a 1-D harmonic oscillator

$$\epsilon_k := k\hbar\omega, \quad k \in \mathbf{N}, \quad (1.43)$$

we see that if we assign the energy

$$\epsilon(\mu) := \epsilon_{\ell^*(\mu)} = \ell^*(\mu)\hbar\omega \quad (1.44)$$

to each branch point with ramification profile of type  $\mu$ , it contributes a factor

$$n(\mu) = \frac{1}{e^{\beta\epsilon(\mu)} - 1} \quad (1.45)$$

to the weighting distributions, the same as that for a bosonic gas.

## 2 2D Toda $\tau$ -functions as generating functions for multispecies quantum Hurwitz numbers

### 2.1 The multiparameter family of $\tau$ -functions $\tau^{Q(\mathbf{q};\mathbf{p},\mathbf{z},\mathbf{w})}(N, \mathbf{t}, \mathbf{s})$

We now consider a multiparameter family of weight generating functions  $Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})$  obtained by taking the product of any number of the generating functions  $E(q_i, z_i)$  and  $H(p_j, w_j)$  for distinct sets of generating function parameters  $\mathbf{z} = (z_1, \dots, z_l)$ ,  $\mathbf{w} = (w_1, \dots, w_m)$  and quantum parameters  $\mathbf{q} = (q_1, \dots, q_l)$ ,  $\mathbf{p} = (p_1, \dots, p_m)$

$$Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w}) := \prod_{i=1}^l E(q_i, z_i) \prod_{j=1}^m H(p_j, w_j). \quad (2.1)$$

Following the approach developed in [5], we define an associated element of the center  $\mathbf{Z}(\mathbf{C}[S_n])$  of the group algebra  $\mathbf{C}[S_n]$  by

$$\hat{Q}(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w}, \mathcal{J}) := \prod_{a=1}^n Q(\mathbf{q}, \mathcal{J}_a \mathbf{z}; \mathbf{p}, \mathcal{J}_a \mathbf{w}), \quad (2.2)$$

where  $\mathcal{J} := (\mathcal{J}_1, \dots, \mathcal{J}_n)$  are the Jucys-Murphy elements [9, 12, 1]

$$\mathcal{J}_1 := 0, \quad \mathcal{J}_b := \sum_{a=1}^{b-1} (ab), \quad b = 1, \dots, n, \quad (2.3)$$

which generate an abelian subalgebra of  $\mathbf{Z}(\mathbf{C}[S_n])$ . The element  $\hat{Q}(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w}, \mathcal{J})$  defines an endomorphism of  $\mathbf{Z}(\mathbf{C}[S_n])$  under multiplication, which is diagonal in the basis  $\{F_\lambda\}$  of

$\mathbf{Z}(\mathbf{C}[S_n])$  consisting of the orthogonal idempotents, corresponding to irreducible representations, labelled by partitions  $\lambda$  of  $n$ :

$$\hat{Q}(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w}, \mathcal{J})F_\lambda = r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})} F_\lambda \quad (2.4)$$

where

$$r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})} = \prod_{i=1}^l r_\lambda^{E(q_i)}(z_i) \prod_{j=1}^m r_\lambda^{H(p_j)}(w_j). \quad (2.5)$$

More generally, defining

$$r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(N) = \prod_{i=1}^l r_\lambda^{E(q_i)}(N, z_i) \prod_{j=1}^m r_\lambda^{H(p_j)}(N, w_j), \quad (2.6)$$

where

$$r_\lambda^{E(q)}(N, z) := \prod_{(ij) \in \lambda} E(q, (N + j - i)z) \quad (2.7)$$

$$r_\lambda^{H(q)}(N, z) := \prod_{(ij) \in \lambda} H(q, (N + j - i)z), \quad (2.8)$$

we have

$$r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})} = r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(0). \quad (2.9)$$

From general considerations [14, 8], the double Schur function series

$$\tau^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(N, \mathbf{t}, \mathbf{s}) := \sum_{\lambda} r_\lambda^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(N) S_\lambda(\mathbf{t}) S_\lambda(\mathbf{s}) \quad (2.10)$$

is known to define a family of  $2D$  Toda  $\tau$ -functions of hypergeometric type.

## 2.2 Multispecies geometric quantum Hurwitz numbers

We now consider coverings in which the branch points are divided into two different classes,  $\{\mu_+^{i, u_i}\}_{i=1, \dots, l; u_i=1, \dots, k_i^-}$  and  $\{\mu_-^{j, v_j}\}_{j=1, \dots, m; v_j=1, \dots, k_j^-}$  corresponding to the weight generating functions of type  $E(q_i)$  and  $H(p_j)$  respectively, each of which is further divided into  $l$  and  $m$  distinct species (or ‘‘colours’’), of which there are  $\{k_i^+\}$  and  $\{k_j^-\}$  branch points of types  $E$  and  $H$  and colours  $i$  and  $j$  respectively. The weighted number of such coverings with specified total colengths  $\mathbf{c} = (c_1, \dots, c_l)$ ,  $\mathbf{d} = (d_1, \dots, d_m)$ ,  $c_i, d_j \in \mathbf{N}$  for each class and colour is the multispecies quantum Hurwitz number

$$H_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu) := \sum_{\substack{\{k_i^+=1; k_j^-=1\} \\ i=1, \dots, l; j=1, \dots, m}}^{\infty} \sum_{\substack{\{\mu_+^{(i, u_i)}\}; \mu_-^{(j, v_j)}\} \\ \sum_{u_i=1}^{k_i^+} \ell^*(\mu_+^{(i, u_i)})=c_i, \sum_{v_j=1}^{k_j^-} \ell^*(\mu_-^{(j, v_j)})=d_j}}$$



$$\times \prod_{i=1}^l W_{E(q_i)}(\mu_+^{(i,1)}, \dots, \mu_+^{(i,k_i^+)}) \prod_{j=1}^m W_{H(p_j)}(\mu_-^{(j,1)}, \dots, \mu_-^{(j,k_j^-)}) H(\{\mu_+^{(i,u_i)}; \mu_-^{(j,v_j)}\}, \mu, \nu). \quad (2.11)$$

Substituting the Frobenius-Schur formula (1.1) and the Frobenius character formula (1.22) into (2.10), it follows that  $\tau_{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(N, \mathbf{t}, \mathbf{s})$  is the generating function for  $H_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu)$ . Using multi-index notation to denote

$$\prod_{i=1}^l z_i^{c_i} \prod_{j=1}^m w_j^{d_j} =: \mathbf{z}^{\mathbf{c}} \mathbf{w}^{\mathbf{d}}, \quad (2.12)$$

we have:

**Theorem 2.1.**

$$\tau_{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(0, \mathbf{t}, \mathbf{s}) := \sum_{\mathbf{c}=(0, \dots, 0); \mathbf{d}=(0, \dots, 0)}^{(\infty, \dots, \infty)} \mathbf{z}^{\mathbf{c}} \mathbf{w}^{\mathbf{d}} \sum_{\mu, \nu} H_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu) P_{\mu}(\mathbf{t}) P_{\nu}(\mathbf{s}). \quad (2.13)$$

### 2.3 Multispecies combinatorial quantum Hurwitz numbers

Another basis for  $\mathbf{Z}(\mathbf{C}[S_n])$  consists of the cycle sums

$$C_{\mu} := \sum_{h \in \text{cyc}(\mu)} h, \quad (2.14)$$

where  $\text{cyc}(\mu)$  denotes the conjugacy class of elements  $h \in \text{cyc}(\mu)$  with cycle lengths equal to the parts of  $\mu$ . The two are related by

$$F_{\lambda} = h_{\lambda}^{-1} \sum_{\mu, |\mu|=|\lambda|} \chi_{\lambda}(\mu) C_{\mu}, \quad (2.15)$$

where  $\chi_{\lambda}(\mu)$  denotes the irreducible character of the irreducible representation of type  $\lambda$  evaluated on the conjugacy class  $\text{cyc}(\mu)$ . Under the characteristic map, this is equivalent to the Frobenius character formula (1.22). The Macmahon generating function

$$\prod_{i=1}^{\infty} (1 - q^i)^{-1} = \sum_{n=0}^{\infty} D_n q^n, \quad (2.16)$$

gives the number  $D_n$  of partitions of  $n$ . We denote by  $\mathbf{F}_{E(q_i)}^{(n, c_i)}$  and  $\mathbf{F}_{H(q_j)}^{(n, d_j)}$  the  $D_n \times D_n$  matrices, whose elements are  $\left( F_{E(q_i)}^{c_i}(\mu, \nu) \right)_{|\mu|=|\nu|=n}$  and  $\left( F_{H(q_j)}^{d_j}(\mu, \nu) \right)_{|\mu|=|\nu|=n}$ , respectively,

for  $i = 1, \dots, l$ ,  $j = 1, \dots, m$ . Since these represent central elements of the group algebra  $\mathbf{C}[S_n]$ , they commute amongst themselves. Denoting the matrix product:

$$\mathbf{F}_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})} = \prod_{i=1}^l \mathbf{F}_{E(q_i)}^{(n, c_i)} \prod_{j=1}^m \mathbf{F}_{Hq_j}^{(n, d_j)}, \quad (2.17)$$

its matrix elements, denoted  $F_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu)$ , may be interpreted as the weighted number of

$$d := \sum_{i=1}^l c_i + \sum_{j=1}^m d_j \quad (2.18)$$

step paths in the Cayley graph from the conjugacy class of cycle type  $\mu$  to the one of type  $\nu$ , where all paths are divided into equivalence classes, according to their multisignature. This consists of partitions of  $\lambda_+^{(i)}$  of  $c_i$  and  $\lambda_-^{(j)}$  of  $d_j$  with

$$\lambda_+^{(i)} = \ell(\lambda_+^{(i)}), \quad \lambda_-^{(j)} = \ell(\lambda_-^{(j)}) \quad (2.19)$$

parts  $\lambda_+^{(i)}$  and  $\lambda_-^{(j)}$ , each of which is itself subdivided into parts  $(\lambda_{+, u_i}^{(i)}, \lambda_{-, v_j}^{(j)})$ , in which the second elements of the transpositions are constant, and distinct for each  $\lambda_{+, u_i}^{(i)}$ , or  $\lambda_{-, v_j}^{(j)}$ . We define the hypersignature of such a path as the set of numbers  $\{\lambda_{+, u_i}^{(i)}, \lambda_{-, v_j}^{(j)}\}_{u_i=1, \dots, k_i^+, v_j=1, \dots, k_j^-}$ . The weight for each path within such an equivalence class is then the product of the weights for each subsegment:

$$\prod_{i=1}^l \prod_{u_i=1}^{k_i^+} E(q_i)_{\lambda_{+, u_i}^{(i)}} \prod_{j=1}^m \prod_{v_j=1}^{k_j^-} H(p_j)_{\lambda_{-, v_j}^{(j)}} \quad (2.20)$$

and  $F_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu)$  is the sum of these, each multiplied by the number of elements of the equivalence class of paths. with the given hypersignature. The multispecies generalization of (1.38) is equality of the geometric and combinatorial Hurwitz numbers:

**Theorem 2.2.**

$$F_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu) = H_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu). \quad (2.21)$$

*Proof.* Applying the central element (2.4) to the cycle sum  $C_\mu$  gives

$$\hat{Q}(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w}, \mathcal{J}) C_\mu = \sum_{\nu, |\nu|=|\mu|} F_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu) C_\nu \quad (2.22)$$

and the orthogonality of group characters that

$$\tau^{Q(\mathbf{q}, \mathbf{z}; \mathbf{p}, \mathbf{w})}(\mathbf{t}, \mathbf{s}) := \sum_{\mathbf{c}=(0, \dots, 0); \mathbf{d}=(0, \dots, 0)}^{(\infty, \dots, \infty)} \mathbf{z}^{\mathbf{c}} \mathbf{w}^{\mathbf{d}} \sum_{\mu, \nu} F_{(\mathbf{q}, \mathbf{p})}^{(\mathbf{c}, \mathbf{d})}(\mu, \nu) P_\mu(\mathbf{t}) P_\nu(\mathbf{s}). \quad (2.23)$$

Comparing this with eq. (2.13) proves the result. □

**Remark 2.1. Multispecies Bose gases**

Returning to the interpretation of the quantum Hurwitz weighting distributions in terms of Bose gases, if we choose each  $q_i$  to be a positive real number with  $q_i < 1$ , and parametrize it, as before,

$$q_i = e^{-\beta \hbar \omega_i} \tag{2.24}$$

for some ground state energy  $\hbar \omega_i$ , and again choose a linear energy spectrum, with energy assigned to the branchpoint  $\mu$  of type  $i$  with profile type  $\mu$  to be

$$\epsilon^{(i)}(\mu) := \ell^*(\mu) \hbar \omega_i \tag{2.25}$$

we see that the resulting contributions to the weighting distributions from each species of branch points of ramification type  $\mu^{(j)}$  are given by

$$n_B^{(i)}(\mu) = \frac{1}{e^{\beta \epsilon^{(i)}(\mu)} - 1}, \tag{2.26}$$

which are those of a multispecies mixture of Bose gases.

## 2.4 General multispecies Hurwitz numbers

The concept of multispecies weighted Hurwitz numbers is of course not restricted to the particular quantum weightings considered here. For any choice of weight generating functions  $G_1(z_1), G_2(z_2), \dots$ , we may form composites by similarly using the product  $\prod_i G_i(z_i)$  as generating function for multiple weighting types. The resulting content product coefficients  $r_\lambda^{\prod_i G_i(z_i)}$  are again just the product of the individual ones

$$r_\lambda^{\prod_i G_i(z_i)} = \prod_i r_\lambda^{G_i(z_i)}. \tag{2.27}$$

The weighted enumerative geometrical and combinatorial interpretations remain the same as in the cases studied here, but with the individual weight generators  $E(q, z_i)$  or  $H(p_j, w_j)$  replaced by  $G_1(z_1), G_2(z_2), \dots$ . In particular, we may include factors of the type  $E'(q, z)$  or  $Q(q, p, z)$  to arrive at similar formulae for the weighted Hurwitz numbers, both in the weighted enumerative geometrical and combinatorial sense, differing only in the detailed form of the various component weights.

We may also include weight factors in which some or all of the parameters  $z_1, z_2 \dots$  are repeated in the product. This only affects the constraints on the sums of the colengths in the weighted multispecies Hurwitz numbers. (See, e.g. example 3.3 in [5], in which the weights are uniform, but the linear generating function that generates the Hurwitz numbers for Belyi curves, and strictly monotonic paths is replaced by a power of the latter, resulting in multiple branch points, with the total colength fixed, and multimonic paths. )

General multispecies Hurwitz numbers, with arbitrary multiplicative weight generating functions, including alternative approaches to the quantum multispecies case are the subject

of [7]. An interesting open question is: does there exist a choice of weight generating functions that has the effect of replacing the energy distributions (2.28) for the individual components by

$$n_F^{(i)}(\mu) = \frac{1}{e^{\beta\epsilon^{(i)}(\mu)} + 1}, \quad (2.28)$$

i.e., with those for a fermionic gas?

*Acknowledgements.* This work began as a joint project with A. Yu. Orlov, with whom the notion of (signed) multispecies Hurwitz numbers was first considered in [8]. It grew into two separate approaches to multispecies quantum Hurwitz numbers, one of which is presented here; the other may be found in [13]. The author would also like to thank M. Guay-Paquet for helpful discussions.

## References

- [1] P. Diaconis and C. Greene, “Applications of Murphy’s elements”, *Stanford Technical Report* 335 (1989).
- [2] G. Frobenius, “Über die Charaktere der symmetrischen Gruppe”, *Sitzber. Preuss. Akad. Berlin*, 516-534 (1900).
- [3] K.I. Gross and D.S. Richards, “Special functions of matrix arguments. I: Algebraic induction, zonal polynomials, and hypergeometric functions”, *Transactions Amer Math Soc.*, **301**, 781-811 (1987).
- [4] M. Guay-Paquet and J. Harnad, “2D Toda  $\tau$ -functions as combinatorial generating functions”, [arXiv:1405.6303](https://arxiv.org/abs/1405.6303).
- [5] M. Guay-Paquet and J. Harnad, “Generating functions for weighted Hurwitz numbers”, [arXiv:1408.6766](https://arxiv.org/abs/1408.6766).
- [6] G. Frobenius, “Über die Charaktere der symmetrischen Gruppe”, *Sitzber. Preuss. Akad. Berlin*, 516-534 (1900).
- [7] J. Harnad “Generating functions for multispecies weighted Hurwitz numbers”, (in preparation, 2014)
- [8] J. Harnad and A. Yu. Orlov, “Hypergeometric  $\tau$ -functions, Hurwitz numbers and enumeration of paths”, [arXiv:1407.7800](https://arxiv.org/abs/1407.7800).
- [9] A. A. Jucys, “Symmetric polynomials and the center of the symmetric group ring”, *Reports on Mathematical Physics* **5**(1) (1974), 107–112.

- [10] S.K. Lando and A.K. Zvonkin *Graphs on Surfaces and their Applications*, Encyclopaedia of Mathematical Sciences, Volume **141**, with appendix by D. Zagier, Springer, N.Y. (2004).
- [11] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, Clarendon Press, Oxford, (1995).
- [12] G. E. Murphy, “A new construction of Young’s seminormal representation of the symmetric groups”, *Journal of Algebra* **69**, 287–297 (1981).
- [13] S. M. Natanzon and A. Yu. Orlov, “Hurwitz numbers and BKP hierarchy”, . arXiv:1407.8323.
- [14] A. Yu. Orlov and D. M. Scherbin, “Hypergeometric solutions of soliton equations”, *Theoretical and Mathematical Physics* **128**, 906-926 (2001).
- [15] K. Takasaki, “Initial value problem for the Toda lattice hierarchy”, in: *Group Representation and Systems of Differential Equations*, 139–163, *Adv. Stud. in Pure Math.* **4**, (1984).
- [16] T. Takebe, “Representation theoretical meaning of the initial value problem for the Toda lattice hierarchy I”, *Lett. Math. Phys.* **21** 77–84, (1991).
- [17] K. Ueno and K. Takasaki, “Toda Lattice Hierarchy”, in: *Group Representation and Systems of Differential Equations*, *Adv. Stud. in Pure Math.* **4**, , 1–95 (1984).