PAINLEVÉ 2 EQUATION WITH ARBITRARY MONODROMY PARAMETER, TOPOLOGICAL RECURSION AND DETERMINANTAL FORMULAS

KOHEI IWAKI AND OLIVIER MARCHAL

ABSTRACT. The goal of this article is to prove that the determinantal formulas of the Painlevé 2 system identify with the correlation functions computed from the topological recursion on their spectral curve for an arbitrary non-zero monodromy parameter. The result is established for a WKB expansion of two different Lax pairs associated to the Painlevé 2 system, namely the Jimbo-Miwa Lax pair and the Harnad-Tracy-Widom Lax pair, where a small expansion parameter \hbar is introduced by a proper rescaling. The proof is based on showing that these systems satisfy the topological type property introduced in [2, 3]. In the process, we explain why the insertion operator method traditionally used to prove the topological type property is currently incomplete and we propose new algebraic methods to bypass the issue. Our work generalizes similar results obtained from random matrix theory in the special case of vanishing monodromies [7, 8]. Explicit computations up to q = 3 are provided along the paper as an illustration of the results. Eventually, taking the time parameter t to infinity we observe that the symplectic invariants $F^{(g)}$ of the Jimbo-Miwa and Harnad-Tracy-Widom spectral curves converge to the Euler characteristic of moduli space of genus q Riemann surfaces.

Contents

1. Introduction	2
2. Summary of the main results	4
3. Jimbo-Miwa Lax pair for Painlevé 2	5
3.1. Introduction of \hbar in the Jimbo-Miwa Lax pair	5
3.2. Hamiltonian system and tau-function of the JM Lax pair	7
3.3. First orders of the JM tau-function	8
3.4. Spectral curve and topological recursion for the JM pair	9
3.5. Tau-function and symplectic invariants for the JM pair	12
3.6. Limit at ∞_B and ∞_C : The Hermite-Weber curve	12
4. The Harnad-Tracy-Widom Lax Pair	16
4.1. Introduction of \hbar in the Harnad-Tracy-Widom Lax pair	16

Date: September 28, 2018.

¹⁹⁹¹ Mathematics Subject Classification. Primary 34M55, 81T45; Secondary 34M56, 34M60. Key words and phrases. Painlevé 2 equation, tau-functions, topological recursion, determinantal formulas.

4.2. Spectral curve and topological recursion for the HTW pair	17
4.3. Tau-function and symplectic invariants for the HTW pair	19
4.4. Limit at $t = \infty_A$: the Bessel curve	19
5. Determinantal formulas and topological type property	21
5.1. Determinantal formulas	21
5.2. Topological type property	22
5.3. Proof of the leading order property of W_n using loop equations	24
6. Outlooks	29
Appendix A. Proof of the parity symmetry	29
Appendix B. Proof of the pole property	31
B.1. Jimbo-Miwa Lax pair	31
B.2. Harnad-Tracy-Widom Lax pair	33
Appendix C. Incomplete proof using an insertion operator	34
Appendix D. Tau-function and symplectic invariants	38
Appendix E. Constant term in (1.2) and the symplectic invariance	38
Acknowledgment	39
References	39

1. INTRODUCTION

In the past decade, the connection between random matrix theory, topological recursion and integrable systems has been developed intensively. Indeed, it was first proved that the partition function describing Hermitian random matrix models (first one matrix models and later two matrix models) are isomonodromic tau-functions [5, 6], a central element of integrable systems. Additionally, the local statistics of eigenvalues in Hermitian matrix models have been proved to be universal and related to Fredholm determinants whose kernels are determined by the nature of the point in the global distribution (edge, bulk point, critical points, etc.) [30, 33]. Lately, these Fredholm determinants were expressed with some Painlevé transcendents [33]. Recently, Evnard and Orantin provided a recursive algorithm, known as "the topological recursion" [17], to compute the (possibly formal) 1/Nexpansion of the correlation functions and partition function of any Hermitian matrix model. This recursion was generalized almost immediately to any "spectral curve" that may or may not come from a matrix model. This topological recursion has been proved very useful in enumerative geometry where many combinatorial results were recently obtained or rediscovered with this formalism [1, 10, 14, 15]. In particular the main interest of the topological recursion is the definition of a series of numbers $F^{(g)}$ known as "symplectic invariants" that are invariant under a certain class of symplectic transformations of the initial spectral curve and that reconstruct the logarithm of the partition function when the spectral curve arises from a matrix model. In a more recent article, Bergère and Eynard [3] were able

 $\mathbf{2}$

to associate a natural spectral curve to any 2×2 Lax pair and provided some determinantal formulas attached to the Lax pair. These determinantal formulas match the correlation functions and symplectic invariants obtained from the computation of the topological recursion on the spectral curve when some additional conditions, known generically as the "topological type" (TT) property, are satisfied. More recently, these notions were extended successfully to $n \times n$ Lax pairs by Bergère, Borot and Eynard [2]. These results are important since they can be used to prove that the determinantal formulas and the tau-function of the Lax pair can be computed perturbatively to any order with the topological recursion associated to the spectral curve, which in general is relatively easy. So far, the TT property has been discussed in three different cases:

- First in [4], in relation with the local statistics of eigenvalues near the edge of the distribution for a Hermitian matrix model, the authors proved the TT property for the Painlevé 2 system (with the Jimbo-Miwa Lax pair) with vanishing monodromy. The approach was generalized in the case of a critical edge with the (2m, 1) hierarchy in [27]. These results where recently recovered and precised in [7, 8].
- In [28], in relation with local statistics of eigenvalues in the bulk of the distribution for a Hermitian matrix model, the authors proved that the result holds for the Painlevé 5 system with vanishing monodromy parameters.
- Eventually in [2], the authors were able to prove the TT property for the q-th reduction of the KP hierarchy, that is to say all (p,q) models. In particular this includes the Painlevé 1 equation (for which there is no monodromy parameter).

However it is worth mentioning that all these articles use at some point the method of the insertion operator. In this article we show that the current proof regarding the insertion operator is incomplete and we present another way based on loop equations to bypass this issue (see Section 5.3 and Appendix C).

In this article, our main goal is to prove that the TT property holds for Lax pairs of the Painlevé 2 equation:

$$\hbar^2 \ddot{q} = 2q^3 + tq - \theta + \frac{\hbar}{2} \tag{1.1}$$

(where $\dot{=} \frac{d}{dt}$ and \hbar is a small parameter) with arbitrary non-zero monodromy parameter θ . More precisely, we will prove the result for two different (\hbar -deformed) Lax pairs frequently used to describe the Painlevé 2 system: the Jimbo-Miwa (JM) Lax pair and the Harnad-Tracy-Widom (HTW) Lax pair. (See [20, 24].) For these two Lax pairs we will review how to produce the spectral curve and the tau-function. Although these Lax pairs describe the same integrable system (Painlevé 2), their spectral curves are totally different. Then, after presenting the topological recursion and the determinantal formulas, we will prove the TT property by proving the three conditions proposed in [3]. This result proves that the generating functions for both sets of symplectic invariants $F_{\rm JM}^{(g)}(t)$ and $F_{\rm HTW}^{(g)}(t)$ defined from the spectral curves of JM pair and HTW pair give the corresponding tau-functions of Painlevé 2 (see Theorem 3.4 and 4.2). Note that, although $F_{\rm JM}^{(g)}(t)$ and $F_{\rm HTW}^{(g)}(t)$ are different as a computational result, both of them gives tau-functions of Painlevé 2 equation. As presented in Theorem 3.7 and 4.3 the connection between the two sets of symplectic invariants appears from constant terms (a tau-function is always defined up to a constant) that are fixed by the topological recursion. More specifically one of our main results is that:

$$F_{\rm JM}^{(g)}(t) = F_{\rm HTW}^{(g)}(t) - \frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
(1.2)

Along the article B_{2g} stands for the Bernoulli numbers defined by

$$\frac{t}{e^t - 1} - 1 + \frac{t}{2} = \sum_{m=1}^{\infty} B_{2m} \frac{t^{2m}}{(2m)!}.$$

We could find the reason why the specific constant terms appear and we connect them to two simple spectral curves: the Hermite-Weber curve (semi-circle curve) for which the symplectic invariants have been known and the Bessel curve for which we could not find any reference in the literature. Actually, the JM curve and HTW curve are related by a symplectic transformation (see Remark 4.4; we thank Professor Yasuhiko Yamada who suggests this fact to the authors). We also show that the above discrepancy between $F^{(g)}$'s is consistent with the integration constant computed in [18] (see Appendix E).

2. Summary of the main results

This article aims at a better understanding between the integrable structure of the Painlevé equations and the topological recursion. Our main results are:

- We prove that the determinantal formulas and tau-functions associated to two different Painlevé 2 Lax pairs (Jimbo-Miwa and Harnad-Tracy-Widom) are identical to the correlation functions and symplectic invariants computed by the topological recursion applied to the corresponding spectral curves (Theorem 3.4, 4.2 and 5.4). Explicit results for the expansion of the tau-function are also presented for both Lax pairs.
- We also show that some limit $t \to \infty$ of the symplectic invariants of both Lax pairs coincide with the Euler characteristic of the moduli space \mathcal{M}_g of Riemann surfaces of genus g computed in [19, 32], up to sign (Theorem 3.7 and 4.3). This also prove the constant term in the relation (1.2) between two symplectic invariants for JM and HTW spectral curves (Appendix E).
- New methods of proof of the TT property are introduced in this article (Section 5.3). In particular the presence of a compatible time differential system is shown to be of critical importance to the TT property (Appendix

B). Moreover these new methods can be easily applied to more general situations and should provide a way to perform the same analysis for the other Painlevé equations.

• We show that the method of the insertion operator used in several papers is incomplete since there is a subtle gap in the proof (Appendix C). This issue was the main reason for the introduction of new methods to prove the TT property.

3. JIMBO-MIWA LAX PAIR FOR PAINLEVÉ 2

In this section we present the (\hbar -depending) Jimbo-Miwa Lax pair as well as the tau-function and its expansion in \hbar . Finally we compute the spectral curve and we illustrate our results with the computation of the first few symplectic invariants $F^{(g)}$ defined by performing the topological recursion, and compare them to the tau-function.

3.1. Introduction of \hbar in the Jimbo-Miwa Lax pair. The following 2×2 Lax pair is equivalent (up to a certain gauge transformation) to the one given in Appendix C of [24]:

$$\begin{pmatrix}
\frac{\partial\Psi}{\partial x}(x,t) = \begin{pmatrix}
x^2 + p + \frac{t}{2} & x - q \\
-2(xp + qp + \theta) & -\left(x^2 + p + \frac{t}{2}\right)
\end{pmatrix}\Psi(x,t), \\
\frac{\partial\Psi}{\partial t}(x,t) = \begin{pmatrix}
\frac{x + q}{2} & \frac{1}{2} \\
-p & -\frac{x + q}{2}
\end{pmatrix}\Psi(x,t).$$
(3.1)

The Lax pair we will use in this article is a \hbar -deformed version of (3.1). The introduction of \hbar is essential in the relation to topological recursion, and it is done by the following rescaling of all quantities involved in the former Lax pair:

$$x \to \hbar^{-\frac{1}{3}}\tilde{x} \ , \ p \to \hbar^{-\frac{2}{3}}\tilde{p} \ , \ t \to \hbar^{-\frac{2}{3}}\tilde{t} \ , \ q \to \hbar^{-\frac{1}{3}}\tilde{q} \ , \ \theta \to \hbar^{-1}\tilde{\theta}.$$
(3.2)

The scaling degrees are chosen so that the resulting Lax pair can be treated in terms of WKB method. The above rescaling provides the \hbar -deformed Lax pair

(we omit \sim for clarity):

$$\begin{cases}
\hbar \frac{\partial \Psi}{\partial x}(x,t) = \begin{pmatrix} x^2 + p + \frac{t}{2} & x - q \\ -2(xp + qp + \theta) & -\left(x^2 + p + \frac{t}{2}\right) \end{pmatrix} \Psi(x,t) \\
\stackrel{\text{def}}{=} \mathcal{D}(x,t)\Psi(x,t), \\
\hbar \frac{\partial \Psi}{\partial t}(x,t) = \begin{pmatrix} \frac{x+q}{2} & \frac{1}{2} \\ -p & -\frac{x+q}{2} \end{pmatrix} \Psi(x,t) \\
\stackrel{\text{def}}{=} \mathcal{R}(x,t)\Psi(x,t).
\end{cases}$$
(3.3)

We call the Lax pair (3.3) the Jimbo-Miwa pair (JM pair, for short).

We remind the reader that q and p are implicitly assumed to depend on the time variable t (and also on \hbar) but not on x. Moreover θ , called the monodromy parameter, is independent of x, t and \hbar . Throughout of the paper, we assume that

$$\theta \neq 0. \tag{3.4}$$

The compatibility equations of the differential system (also known as zerocurvature equations) are given by:

$$\hbar \left(\frac{\partial \mathcal{D}}{\partial t} - \frac{\partial \mathcal{R}}{\partial x} \right) + [\mathcal{D}, \mathcal{R}] = 0.$$
(3.5)

From (3.3) they are equivalent to:

$$\hbar \dot{p} = -2qp - \theta , \ \hbar \dot{q} = p + q^2 + \frac{t}{2}.$$
 (3.6)

Here and in what follows a dot is used to denote the derivative relatively to t when no ambiguity appears. Differentiating the last equation and eliminating p with the first equation gives that q satisfies the Painlevé 2 equation (with the small parameter \hbar):

$$\hbar^2 \ddot{q} = 2q^3 + tq - \theta + \frac{\hbar}{2}.$$
 (3.7)

This type of Painlevé equations, and associated Lax pairs with a small parameter \hbar were studied in [26].

In this paper we are interested in the \hbar -perturbative expansion of a solution of Painlevé 2 equation:

$$q(t) = \sum_{k=0}^{\infty} q_k(t)\hbar^k = q_0(t) + q_1(t)\hbar + q_2(t)\hbar^2 + \cdots .$$
 (3.8)

The top term $q_0(t)$ satisfies

$$2q_0(t)^3 + tq_0(t) - \theta = 0 \text{ and } \dot{q}_0(t) = -\frac{q_0(t)^2}{4q_0(t)^3 + \theta}.$$
(3.9)

As is clear from (3.9), $\dot{q}_0(t)$ has singularity where $4q_0(t)^3 + \theta = 0$ holds. Such a point on t-plane is called a turning point of Painlevé 2 in [26]. In what follows, we assume that t lies on a domain on which $4q_0(t)^3 + \theta \neq 0$ holds. Note also that $q_0(t) \neq 0$ holds for any t under the assumption (3.4).

Since $q_0(t)$ is a solution of a cubic equation, there are 3 possible choices of branches for $q_0(t)$. In particular when $t \to \infty$ there are three possible behaviors depending on the chosen branch:

$$q_0(t) \sim_{t \to \infty_A} \frac{\theta}{t}$$
, $q_0(t) \sim_{t \to \infty_B} \sqrt{\frac{-t}{2}}$, $q_0(t) \sim_{t \to \infty_C} -\sqrt{\frac{-t}{2}}$. (3.10)

It is easy to see that, once we fix the branch of the algebraic function $q_0(t)$, the coefficients $\{q_i(t)\}_{i>1}$ appearing in (3.8) are determined recursively. Thanks to the relation (3.6), we also get a similar \hbar -expansion of p(t).

3.2. Hamiltonian system and tau-function of the JM Lax pair. The taufunction are classically defined since the works of Jimbo-Miwa-Ueno [23]. For the JM pair, these quantities are easy to derive (the leading order of the matrices $\mathcal{D}(x,t)$ and $\mathcal{R}(x,t)$ when $x \to \infty$ are both diagonal) and can be directly adapted from the known $\hbar = 1$ case. The Hamiltonian system attached to the JM pair (3.3) is:

$$\begin{cases} \hbar \dot{q} = \frac{\partial H_{\rm JM}}{\partial p} = p + q^2 + \frac{t}{2}, \\ \hbar \dot{p} = -\frac{\partial H_{\rm JM}}{\partial q} = -2qp - \theta, \end{cases}$$
(3.11)

where $H_{\rm JM}$ is the Hamiltonian for Painlevé 2:

$$H_{\rm JM} = \frac{1}{2}p^2 + \left(q^2 + \frac{t}{2}\right)p + \theta q.$$
 (3.12)

Let $\sigma(t)$ be the corresponding Hamiltonian function, that is, the function obtained by substituting a solution (q, p) of (3.11) into $H_{\rm JM}$. It satisfies:

$$\dot{\sigma} = \frac{p}{2} \text{ and } \hbar \ddot{\sigma} = -qp - \frac{\theta}{2}.$$
 (3.13)

as well as the σ -form of the Painlevé 2 equation:

$$(\hbar\ddot{\sigma})^2 + 4(\dot{\sigma})^3 + 2t(\dot{\sigma})^2 - 2\sigma\dot{\sigma} - \frac{\theta^2}{4} = 0.$$
(3.14)

Then, the tau-function for JM Lax pair is defined by:

$$-\hbar^2 \frac{d}{dt} \ln \tau_{\rm JM} = \sigma(t). \tag{3.15}$$

7

Since (3.14) only involves even power of \hbar then the \hbar -expansion of $\sigma(t)$ and $\ln \tau(t)$ only involve even powers of \hbar :

$$\sigma(t) = \sum_{k=0}^{\infty} \sigma_{2k}(t)\hbar^{2k}, \qquad (3.16)$$

$$\ln \tau_{\rm JM} \stackrel{\text{def}}{=} \sum_{k=0}^{\infty} \tau_{2k}(t) \hbar^{2k-2}, \quad \tau_{2k}(t) = -\int^t \sigma_{2k}(s) ds. \tag{3.17}$$

Moreover, (3.13) implies that p(t) also only contains even order terms:

$$p(t) = \sum_{k=0}^{\infty} p_{2k}(t)\hbar^{2k}.$$
(3.18)

3.3. First orders of the JM tau-function. In this section we present the computation of the first orders of the tau-function for the JM Lax pair. In what follows, we choose to express all quantities as function of $q_0(t)$ which is a solution of (3.9). Straightforward computations give:

$$\begin{aligned}
\sigma_0(t) &= \frac{\theta(8q_0^3 - \theta)}{8q_0^2}, \\
\sigma_2(t) &= -\frac{\theta q_0}{8(4q_0^3 + \theta)^2}, \\
\sigma_4(t) &= -\frac{3\theta q_0^4(560q_0^6 - 184\,\theta q_0^3 + 3\theta^2)}{32(4q_0^3 + \theta)^7}, \\
\sigma_6(t) &= -\frac{\theta q_0^7}{32(4q_0^3 + \theta)^{12}} (3203200\,q_0^{12} - 3668064\,\theta q_0^9) \\
&+ 838632\,\theta^2 q_0^6 - 39482\,\theta^3 q_0^3 + 189\,\theta^4).
\end{aligned}$$
(3.19)

In particular one can also verify directly that the coefficients presented here satisfy the differential equation (3.14). Integrating over t leads to:

$$\tau_{0}(t) = \frac{4}{3}\theta q_{0}^{3} + \frac{\theta^{3}}{24q_{0}^{3}} + \frac{\theta^{2}}{2}\ln q_{0} + \text{Cste},$$

$$\tau_{2}(t) = \frac{1}{24}\ln(1 + \frac{\theta}{4q_{0}^{3}}) + \frac{1}{12}\ln 2 + \text{Cste},$$

$$\tau_{4}(t) = \frac{\theta(700 q_{0}^{6} - 85 \theta q_{0}^{3} - 2\theta^{2})}{480(4q_{0}^{3} + \theta)^{5}} + \text{Cste},$$

$$\tau_{6}(t) = \frac{\theta}{4032 (4q_{0}^{3} + \theta)^{10}} (6726720 q_{0}^{15} - 5017712 \theta q_{0}^{12} + 541132 \theta^{2} q_{0}^{9} - 1089 \theta^{3} q_{0}^{6} + 160 \theta^{4} q_{0}^{3} + 4 \theta^{5}) + \text{Cste}$$
(3.20)

Here the constant terms are to be understood as not depending on t. We will see that these integration constants are specified by the topological recursion, and correspond to lower end-points for the integral (3.17) defining τ_{2k} taken at $t = \infty$ for $k \geq 2$. Actually, we can choose ∞ (for any ∞_A, ∞_B and ∞_C) as the lower end-point since $\sigma_{2k} \underset{t \to \infty}{=} O(t^{-2})$ holds if $k \geq 2$. This property can easily be proved by a recursion relation satisfied by $\{q_i(t)\}_{i\geq 0}$ appearing in (3.8).

3.4. Spectral curve and topological recursion for the JM pair. From Bergère and Eynard [3] we know that the spectral curve of a Lax pair is given by the leading order in \hbar of the characteristic polynomial of $\mathcal{D}(x, t)$. Thus we find:

$$y^{2} = (x - q_{0})^{2} (x^{2} + 2q_{0}x + 3q_{0}^{2} + t)$$

= $(x - q_{0})^{2} \left(x + q_{0} - \sqrt{-\frac{\theta}{q_{0}}} \right) \left(x + q_{0} + \sqrt{-\frac{\theta}{q_{0}}} \right)$ (3.21)

where $q_0(t)$ is the solution of (3.9). We call (3.21) the Jimbo-Miwa spectral curve (JM curve, for short). JM curve is of genus 0 with two branch points. It can be parametrized with a global Zhukovsky variable:

$$\begin{cases} x(z) = -q_0 + \frac{1}{2}\sqrt{-\frac{\theta}{q_0}\left(z + \frac{1}{z}\right)}, \\ y(z) = \frac{1}{2}\sqrt{-\frac{\theta}{q_0}\left(z - \frac{1}{z}\right)\left(-2q_0 + \frac{1}{2}\sqrt{-\frac{\theta}{q_0}\left(z + \frac{1}{z}\right)}\right)}. \end{cases}$$
(3.22)

With this parametrization, the branch points are located at $z = \pm 1$ and the differential ydx has two poles at z = 0 and $z = \infty$.

Definition 3.1 (Definition 4.2 of [17]). For $g \ge 0$ and $n \ge 1$, define the Eynard-Orantin differential $\omega_n^{(g)}(z_1, \ldots, z_n)$ of the type (g, n) for the spectral curve (3.22) by the following topological recursion relation ([17]):

$$\begin{aligned}
\omega_{1}^{(0)}(z_{1}) &= y(z_{1})dx(z_{1}), \\
\omega_{2}^{(0)}(z_{1}, z_{2}) &= \frac{dz_{1}dz_{2}}{(z_{1} - z_{2})^{2}}, \\
\omega_{n+1}^{(g)}(z_{0}, z_{1}, \dots, z_{n}) &= \sum_{r \in R} \operatorname{Res}_{z=r} K(z_{0}, z) \Big[\omega_{n+1}^{(g-1)}(z, \bar{z}, z_{1}, \dots, z_{n}) \\
&+ \sum_{\substack{r \in R \\ I \cup J = \{1, \dots, n\}}}' \omega_{1+|I|}^{(g_{1})}(z, z_{I}) \omega_{1+|J|}^{(g_{2})}(\bar{z}, z_{J}) \Big].
\end{aligned}$$
(3.23)

Here R is the set of branch points, \bar{z} is the local conjugate of z near a branch point r,

$$K(z_0, z) = \frac{1}{2} \frac{\int_z^z \omega_2^{(0)}(\cdot, z_0)}{(y(z) - y(\bar{z}))dx(z)}$$
(3.24)

is called the recursion kernel, and the summation in the last line of (3.23) means "except for the cases $(g_1, I) = (0, \emptyset)$ and $(g_2, J) = (0, \emptyset)$ ".

For JM curve (3.22), $R = \{\pm 1, -1\}$ and $\bar{z} = z^{-1}$ for both branch points $r = \pm 1$. The Eynard-Orantin differentials $\omega_n^{(g)}$ are meromorphic multi-differentials on the *n*-times product of spectral curves, and known to be holomorphic except for the branch points if $(g, n) \neq (0, 1), (0, 2)$. Since our spectral curve has genus 0, the topological recursion becomes easier (see [17] for general case).

We also introduce symplectic invariants $F^{(g)}$ for a genus 0 spectral curve, following [17]:

Definition 3.2 (Definition 4.3 of [17]). The *g*-th symplectic invariant (or genus g free energy) of the spectral curve is defined as follows:

• For g = 0, set

$$F^{(0)} = \frac{1}{2} \sum_{\alpha} \operatorname{Res}_{z=\alpha} V_{\alpha}(z) y(z) dx(z) + \frac{1}{2} \sum_{\alpha} t_{\alpha} \mu_{\alpha}, \qquad (3.25)$$

where the sum is taken over all poles of y(z)dx(z), and for a pole α of y(z)dx(z), we define

$$t_{\alpha} = \operatorname{Res}_{z=\alpha} y(z) dx(z) \tag{3.26}$$

$$V_{\alpha}(z) = \operatorname{Res}_{p=\alpha} \ln\left(1 - \frac{\xi_{\alpha}(z)}{\xi_{\alpha}(p)}\right) y(p) dx(p), \qquad (3.27)$$

$$\mu_{\alpha} = \int_{\alpha}^{z_o} \left(y(z) dx(z) - dV_{\alpha}(z) - t_{\alpha} \frac{d\xi_{\alpha}(z)}{\xi_{\alpha}(z)} \right) + V_{\alpha}(z_o) + t_{\alpha} \ln \xi_{\alpha}(z_o).$$
(3.28)

Here z_o is a fixed arbitrary generic point on the spectral curve, and $\xi_{\alpha}(z)$ is a certain function of z which is chosen depending on the behavior of y(z)dx(z) near $z = \alpha$ (see §3.4.2 of [17]).

• For g = 1, set

$$F^{(1)} = -\frac{1}{24} \ln \left(\tau_B(\{x(a_i)\})^{12} \prod_{r \in R} y'(r) \right), \qquad (3.29)$$

where $\tau_B(\{x_i\})$ is a function of $x_i = x(r_i)$ $(r_i \in R)$ which is defined (up to constant) by the following property:

$$\frac{\partial}{\partial x_i} \ln \tau_B(\{x_i\}) = \operatorname{Res}_{z=a_i} \frac{B(z,\bar{z})}{dx(z)}.$$

 $(\tau_B({x_i})$ is called the Bergman tau-function of the spectra curve.) We also set

$$y'(r) = \lim_{z \to r} \frac{y(z) - y(r)}{\sqrt{x(z) - x(r)}}$$
(3.30)

for $r \in R$.

• For $g \ge 2$, set

$$F^{(g)} = \frac{1}{2 - 2g} \sum_{r \in R} \operatorname{Res}_{z=r} \Phi(z) \omega_1^{(g)}(z), \qquad (3.31)$$

where

$$\Phi(z) = \int_{z_o}^{z} y(z) dx(z)$$

with an arbitrary chosen generic point z_o .

Denote by $F_{\rm JM}^{(g)}$ the symplectic invariants for JM curve (3.22). In the above definition of $F_{\rm JM}^{(0)}$, y(z)dx(z) has poles at 0 and ∞ , and we chose $\xi_0(z) = \xi_{\infty}(z) = x(z)$. Also, for $F^{(1)}$, we fix the ambiguity of the normalization constant of the Bergman tau-function as

$$F_{\rm JM}^{(1)} = -\frac{1}{24} \ln \left(\gamma^3 y'(1) y'(-1) \right), \quad \gamma = \frac{1}{2} \sqrt{-\frac{\theta}{q_0}}.$$

(See Chapter 7 of [16] for the above formula.) Then, we find:

$$F_{\rm JM}^{(0)} = \frac{4\theta}{3}q_0^3 - \frac{\theta^2}{4} + \frac{\theta^3}{24q_0^3} - \frac{\theta^2}{2}\ln\left(-\frac{\theta}{4q_0}\right)$$

$$F_{\rm JM}^{(1)} = -\frac{1}{24}\ln\left(\theta^2\left(1 + \frac{\theta}{4q_0^3}\right)\right),$$

$$F_{\rm JM}^{(2)} = \frac{(2048\,q_0^{12} + 2560\,\theta\,q_0^9 + 1280\,\theta^2q_0^6 + 1020\,\theta^3q_0^3 - 45\,\theta^4)\,q_0^3}{480\,\theta^2\,(4\,q_0^3 + \theta)^5},$$

$$F_{\rm JM}^{(3)} = -\frac{q_0^6}{4032\theta^4\,(\theta + 4\,q_0^3)^{10}}\left(4194304\,q_0^{24} + 10485760\,\theta\,q_0^{21}\right) + 11796480\,\theta^2q_0^{18} + 7864320\,\theta^3q_0^{15} + 3440640\,\theta^4q_0^{12}}{-5694528\,\theta^5q_0^9 + 5232752\,\theta^6q_0^6 - 510412\,\theta^7q_0^3 + 3969\,\theta^8}\right).$$
(3.32)

Moreover, it is easy to prove that when $q_0 \to 0$ (i.e., $t \to \infty_A$) the correlation functions $\omega_n^{(g)}$ and the symplectic invariants $F_{\rm JM}^{(g)}$ (identified with $\omega_0^{(g)}$ in the next formula) behave like:

$$\omega_n^{(g)}(z_1,\ldots,z_n) \underset{q_0\to 0}{\sim} \operatorname{Cste} q_0^{\frac{3}{2}(2g-2+n)} dz_1\cdots dz_n.$$
(3.33)

Indeed, the kernel $K(z_0, z)$ used in the recursion behaves like:

$$K(z_0, z) = -\frac{4z^4}{(z^2 - 1)(z_0 z - 1)(z_0 - z)\left((1 + z)\left(\frac{-\theta}{q_0}\right)^{\frac{3}{2}} + 4\theta z\right)}\frac{dz_0}{dz} = O\left(q_0^{\frac{3}{2}}\right).$$
(3.34)

Thus adding a power $q_0^{3/2}$ at each step of the recursion. In particular, we get:

$$\lim_{t \to \infty_A} F_{\rm JM}^{(g)}(t) = 0 \quad \text{for } g \ge 2.$$
(3.35)

3.5. Tau-function and symplectic invariants for the JM pair. In this subsection we state one of our main results regarding the relationship between symplectic invariants $F_{\rm JM}^{(g)}$ and the tau-function of Painlevé 2.

Theorem 3.3. The JM pair is of topological type (in the sense of Section 5) and we have:

$$\frac{dF_{\rm JM}^{(g)}(t)}{dt} = -\sigma_{2g}(t) \quad for \ g \ge 2.$$
(3.36)

Theorem 3.3 will be proved in Section 5 and in appendix. From former results (3.19) and (3.32) we can verify that $\frac{dF_{\rm JM}^{(g)}}{dt} = -\sigma_{2g}$ also hold for g = 0 and g = 1. This implies the following:

Theorem 3.4. The generating function of symplectic invariants of JM curve (3.22) gives a tau-function of Painlevé 2. In other words,

$$\ln \tau_{\rm JM} = \sum_{g=0}^{\infty} \hbar^{2g-2} F_{\rm JM}^{(g)}(t)$$
(3.37)

satisfies (3.15). Furthermore, we have

$$F_{\rm JM}^{(g)}(t) = -\int_{\infty_A}^t \sigma_{2g}(s)ds \quad for \ g \ge 2.$$
 (3.38)

Theorem 3.4 follows from Theorem 3.3 and (3.35).

3.6. Limit at ∞_B and ∞_C : The Hermite-Weber curve. We already know that the functions $F_{\rm JM}^{(g)}(t)$ vanish for $g \ge 2$ when $t \to \infty_A$ (i.e., $q_0 \to 0$). We find that some interesting numbers appear when taking the limit of $F_{\rm JM}^{(g)}(t)$ to both $t \to \infty_B$ and $t \to \infty_C$. Unfortunately taking these limits $t \to \infty_{B,C}$ in the spectral curve (3.22) is not directly possible since the coefficients diverge in the limit. To avoid this difficulty, we perform the following affine symplectic transformation

$$x = \frac{1}{2\sqrt{-q_0}}X - q_0 \ , \ y = 2\sqrt{-q_0}Y$$
(3.39)

giving the curve:

$$Y^{2} = \left(\frac{X^{2}}{4} - \theta\right) \left(1 + \frac{X}{4(-q_{0})^{\frac{3}{2}}}\right)^{2}.$$
 (3.40)

It is clear that the spectral curve (3.40) has the same symplectic invariants $F_{\rm JM}^{(g)}$ since the coordinate transformation (3.39) preserves the 1-form $\omega_1^{(0)}$ and the set of

12

branch points. In the limit $t \to \infty_{B,C}$ (i.e., $q_0 \to \infty$), the curve (3.40) is reduced to the Hermite-Weber curve:

$$Y^{2} = \frac{X^{2}}{4} - \theta \tag{3.41}$$

which can be parametrized into:

$$\begin{cases} X(z) = \sqrt{\theta} \left(z + \frac{1}{z} \right), \\ Y(z) = \frac{\sqrt{\theta}}{2} \left(z - \frac{1}{z} \right). \end{cases}$$
(3.42)

Proposition 3.5 (Cf. [19, 32]). The symplectic invariants $F_{\text{Weber}}^{(g)}$ for the spectral curve (3.42) are given by

$$F_{\text{Weber}}^{(0)} = \frac{3\theta^2}{4} - \frac{\theta^2}{2}\ln\theta,$$
 (3.43)

$$F_{\text{Weber}}^{(1)} = -\frac{1}{12}\ln\theta, \qquad (3.44)$$

$$F_{\text{Weber}}^{(g)} = -\frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
(3.45)

Proof. Although the statement is well-known in the physics literature (see Remark 3.6 below), we could not find a specific proof in the literature. Therefore we propose here an alternative proof based on the results obtained in [13, 31]. In order to use these results, let us consider the spectral curve:

$$\begin{cases} \tilde{X}(t) = \sqrt{\theta} \left(2 + \frac{4}{t^2 - 1} \right), \\ \tilde{Y}(t) = \sqrt{\theta} \frac{t + 1}{t - 1}. \end{cases}$$
(3.46)

It is obtained from the spectral curve (3.42) by the affine symplectic transformation $(X, Y) \mapsto (\tilde{X}, \tilde{Y}) = (X, Y + \frac{X}{2})$ which preserves the set $(F^{(g)})_{g\geq 2}$ (Theorem 7.1 of [17]), combined with the change of parametrization $z \mapsto z(t) = \frac{t+1}{t-1}$. In [31] the authors introduced the functions (called the Poincaré polynomials):

$$F_n^{(g)}(t_1,\ldots,t_n) = \left(\sum_{\substack{\Gamma: \text{ ribbon graphs}\\\text{ of type } (g,n)}} \frac{(-1)^{e(\Gamma)}}{|\operatorname{Aut}(\Gamma)|} \prod_{\substack{\eta: \text{ edges}\\\text{ in } \Gamma}} \frac{(t_{i_\eta}+1)(t_{j_\eta}+1)}{2(t_{i_\eta}+t_{j_\eta})}\right) \theta^{2-2g-n}$$

for $g \ge 0$ and $n \ge 1$ (see [31] for the precise definitions of the involved quantities). In [13, 31], the authors proved that these functions satisfy the following properties:

• For $g \ge 0$ and $n \ge 1$, the multi-differentials

$$\tilde{\omega}_n^{(g)}(t_1,\ldots,t_n) = d_{t_1}\cdots d_{t_n} F_n^{(g)}(t_1,\ldots,t_n)$$

satisfy the topological recursion applied to the spectral curve (3.46) (See Theorem 2.13 of [13]).

- For $g \ge 0$ and $n \ge 1$, we have $F_n^{(g)}(t_1, \ldots, t_n)|_{t_i=-1} = 0$ for each $1 \le i \le n$. For $g \ge 0$ and $n \ge 1$, we have $F_n^{(g)}(1, \ldots, 1) = (-1)^n \chi(\mathcal{M}_{g,n}) \theta^{2-2g-n}$, where $\mathcal{M}_{g,n}$ is the moduli space of genus g Riemann surfaces with n marked points. In particular, it is proved in [19, 32] that:

$$\chi(\mathcal{M}_{g,n}) = \frac{(2g-3+n)!}{(2g)!}(2g-1)B_{2g}.$$
(3.47)

Precisely speaking, [13, 31] considered the case $\theta = 1$. But these results can be generalized as above since we can introduce the parameter θ by a trivial scaling of variables. In particular, we have $\int_{-1}^{t} \tilde{\omega}_{1}^{(g)}(t) = \tilde{F}_{1}^{(g)}(t)$ and thus

$$\int_{-1}^{1} \tilde{\omega}_{1}^{(g)}(t) = F_{1}^{(g)}(1) = -\chi(\mathcal{M}_{g,1})\theta^{1-2g} = \frac{B_{2g}}{2g\,\theta^{2g-1}}$$

On the other hand, since the parameter θ is the filling fraction (or equivalently $\pm \theta$ are the temperatures at $t = \pm 1$, respectively), the variation formula (see Section 5.3 of [17]) implies that:

$$\frac{dF_{\text{Weber}}^{(g)}}{d\theta} = \int_{-1}^{1} \tilde{\omega}_1^{(g)}(t) = \frac{B_{2g}}{2g \,\theta^{2g-1}} \quad \text{for } g \ge 2.$$

Thus we have the desired equality (3.45) modulo an additive constant which is independent of θ . However, this additive constant must vanish due to the homogeneity relation

$$F_{\text{Weber}}^{(g)}\Big|_{\theta\mapsto\lambda\theta} = \lambda^{2-2g} F_{\text{Weber}}^{(g)} \text{ for } g \ge 2.$$

(See Theorem 5.3 of [17]). Thus, we have finally proved (3.45) for $g \ge 2$. Eventually, formulas (3.43) and (3.44) are obtained from a direct explicit computation.

Remark 3.6. The curve (3.42) also appears as the spectral curve arising in the following Gaussian Hermitian matrix model (with an appropriate normalization):

$$Z_{\rm G} = \frac{1}{(2\pi)^N N!} \int_{\mathbb{R}^N} d\lambda_1 \dots d\lambda_N \,\Delta(\lambda_1, \dots, \lambda_N)^2 \, e^{-\frac{1}{\hbar} \sum_{i=1}^N \frac{\lambda_i^2}{2}}.$$
 (3.48)

Here $\Delta(\lambda_1, \ldots, \lambda_N) = \prod_{1 \le i < j \le N} (\lambda_j - \lambda_i)$ is the standard Vandermonde determinant. The value of the partition function $Z_{\rm G}$ is explicitly known from Mehta's integral (a case of Selberg-like integrals, see [9] for details):

$$Z_{\rm G} = \frac{\hbar^{\frac{N^2}{2}}}{(2\pi)^{\frac{N}{2}}N!} \prod_{i=1}^{N} \Gamma(1+i) = \frac{\hbar^{\frac{N^2}{2}}}{(2\pi)^{\frac{N}{2}}} \prod_{i=1}^{N} i!.$$
(3.49)

The parameter θ in (3.42) is the so-called 't Hooft parameter in the analysis of the large N limit: $N \to \infty$, $\hbar \to 0$ with $\theta = N\hbar$ fixed. The above explicit formula enables us to find the explicit large N expansion (see [29] for example):

$$-\ln Z_G \Big|_{N=\hbar\theta} \sim \sum_{g\geq 0} \hbar^{2g-2} F_{\rm G}^{(g)}(\theta),$$

$$F_{\rm G}^{(0)}(\theta) = \frac{3}{4} \theta^2 - \frac{1}{2} \theta^2 \ln \theta, \quad F_{\rm G}^{(1)}(\theta) = -\frac{1}{12} \ln \theta + \zeta'(-1),$$

$$F_{\rm G}^{(g)}(\theta) = -\frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \geq 2.$$
(3.50)

Here $\zeta'(-1)$ is the derivative of Riemann's ζ -function $\zeta(s)$ evaluated at s = -1. The right hand side of (3.50) coincides with the Euler characteristic of the moduli space \mathcal{M}_q of genus g Riemann surfaces (see [19, 32]).

For $\theta = 1$, general results regarding the connection of the topological recursion with standard large N limit asymptotic expansions of Hermitian matrix integrals (with polynomial potentials) claim that the generating function of the symplectic invariants $F_{\text{Weber}}^{(g)}|_{\theta=1}$ matches with the asymptotic expansion of $-\ln Z_G|_{\theta=1}$ (which is the standard large N limit of the Gaussian Hermitian matrix integral) up to some additive constants: (See Corollary 5.1 in [12] or main theorems of [2, 9]). More precisely, we have:

$$F_{\text{Weber}}^{(g)}\Big|_{\theta=1} = F_G^{(g)}\Big|_{\theta=1} + C^{(g)} \text{ for } g \ge 0.$$

Since θ may be introduced by a trivial rescaling of the parameter $N \mapsto N\theta$ in the Gaussian matrix integral, the last equality may be extended to:

$$F_{\text{Weber}}^{(g)}(\theta) = F_G^{(g)}(\theta) + C^{(g)}\theta^{2-2g} \quad \text{for } g \ge 0.$$
(3.51)

Using the exact expression (3.50) and the values of $F_{\text{Weber}}^{(g)}(\theta)$ proved in Proposition 3.5, we may obtain the constants and we find:

$$C^{(0)} = 0, C^{(1)} = \zeta'(-1)$$
 and $C^{(g)} = 0$ for $g \ge 2$.

Note in particular that, as claimed in the physics literature, the constants $(C^{(g)})_{g\geq 2}$ are vanishing. However, we stress here that current theorems relating the topological recursion with Hermitian matrix integrals are not sufficient to determine the constants and thus prove Proposition 3.5.

Eventually, we note that the symplectic invariants (3.45) are closely related to the Voros coefficients in the theory of the exact WKB analysis. This issue will be discussed in the forthcoming paper [22].

Since it is known from [17] that the symplectic invariants (and correlation functions) obtained for a limiting curve are equal to the limit of symplectic invariants, we have from proposition 3.5 the following theorem: **Theorem 3.7.** Both of the limits $t \to \infty_B$ and $t \to \infty_C$ of the symplectic invariant $F_{\text{JM}}^{(g)}(t)$ of JM curve are given by

$$\lim_{t \to \infty_{B,C}} F_{\rm JM}^{(g)}(t) = -\frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad for \ g \ge 2.$$
(3.52)

In particular taking $q_0 \rightarrow 0$ in (3.32) we can verify that this result holds for g = 2 and g = 3. Theorem 3.7 and equation (3.38) also imply the following:

$$\int_{\infty_A}^{\infty_{B,C}} \sigma_{2g}(t) dt = \frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
(3.53)

Note that a path connecting ∞_A and $\infty_{B,C}$ never exists when $\theta = 0$ since the equation (3.9) defining $q_0(t)$ splits in that case. This is consistent with the fact that the r.h.s. of (3.53) blows up when $\theta \to 0$.

4. The Harnad-Tracy-Widom Lax Pair

In this section, we develop the same approach for the Painlevé 2 system but with another Lax pair, called Harnad-Tracy-Widom Lax pair, as a starting point. This Lax pair is connected to the previous one (3.3) by a Laplace-type integral transformation ([25]), but their relation is non-trivial in terms of the topological recursion. To our knowledge the two Lax pairs (up to trivial transformations) studied in this article represent the two usual pairs used to describe the Painlevé 2 system.

4.1. Introduction of \hbar in the Harnad-Tracy-Widom Lax pair. The following Lax pair is introduced in [20]:

$$\begin{cases} \frac{\partial}{\partial x}\Psi(x,t) = \begin{pmatrix} -q + \frac{\theta}{2x} & x - p - 2q^2 - t \\ \frac{1}{2} + \frac{p}{2x} & q - \frac{\theta}{2x} \end{pmatrix} \Psi(x,t), \\ \frac{\partial}{\partial t}\Psi(x,t) = \begin{pmatrix} q & -x \\ -\frac{1}{2} & -q \end{pmatrix} \Psi(x,t). \end{cases}$$
(4.1)

Like in the JM case we can introduce a small expansion parameter \hbar with a suitable rescaling of the variables:

$$x \to \hbar^{-\frac{2}{3}}\tilde{x} , \ q \to \hbar^{-\frac{1}{3}}\tilde{q} , \ p \to \hbar^{-\frac{2}{3}}\tilde{p} , \ t \to \hbar^{-\frac{2}{3}}\tilde{t} , \ \theta \to \hbar^{-1}\tilde{\theta}$$
 (4.2)

as well as a suitable gauge transformation $\Psi \to \text{diag}(\hbar^{-\frac{1}{6}}, \hbar^{\frac{1}{6}})\Psi$. Omitting ~ for clarity, we get:

$$\begin{cases}
\hbar \frac{\partial}{\partial x} \Psi(x,t) = \begin{pmatrix} -q + \frac{\theta}{2x} & x - p - 2q^2 - t \\ \frac{1}{2} + \frac{p}{2x} & q - \frac{\theta}{2x} \end{pmatrix} \Psi(x,t) \\
\stackrel{\text{def}}{=} \mathcal{D}(x,t) \Psi(x,t), \qquad (4.3) \\
\hbar \frac{\partial}{\partial t} \Psi(x,t) = \begin{pmatrix} q & -x \\ -\frac{1}{2} & -q \end{pmatrix} \Psi(x,t) \\
\stackrel{\text{def}}{=} \mathcal{R}(x,t) \Psi(x,t).
\end{cases}$$

We call the Lax pair (4.3) Harnad-Tracy-Widom pair (HTW pair, for short).

The compatibility equations for this Lax pair are given by:

$$\hbar \dot{q} = q^2 + p + \frac{t}{2}$$
 and $\hbar \dot{p} = -2qp - \theta.$ (4.4)

As in the Jimbo-Miwa case, we recover that q is a solution of the Painlevé 2 equation:

$$\hbar^2 \ddot{q} = 2q^3 + tq - \theta + \frac{\hbar}{2}.$$
 (4.5)

Although the compatibility equations are the same as for the JM pair, the definition of Jimbo-Miwa-Ueno tau-function is a little different ([23, 24]; see also Section 5.2). The Hamiltonian for the HTW pair is:

$$H_{\rm HTW} = \frac{1}{2}p^2 + \left(q^2 + \frac{t}{2}\right)p + \theta q + \frac{t^2}{8}$$
(4.6)

and the tau-function for HTW pair is defined as

$$-h^2 \frac{d}{dt} \ln \tau_{\rm HTW} = H_{\rm HTW} = \sigma(t) + \frac{t^2}{8}$$

$$(4.7)$$

where $\sigma(t)$ is the solution of the σ -form of Painlevé 2 in (3.14).

4.2. Spectral curve and topological recursion for the HTW pair. As usual the spectral curve is given by the leading order in \hbar of the characteristic polynomial of $\mathcal{D}(x,t)$. We find:

$$y^{2} = \frac{1}{2x^{2}} \left(x - \frac{\theta}{2q_{0}} \right)^{2} \left(x + 2q_{0}^{2} \right)$$
(4.8)

where $q_0(t)$ is a solution of (3.9). It is a genus 0 curve with a single branch point arising at $x = -2q_0^2$ but with a pole singularity at x = 0. It can be parametrized

globally with:

$$\begin{cases} x(z) = 2q_0^2(z^2 - 1), \\ y(z) = \frac{z\left(\theta - 4q_0^3(z^2 - 1)\right)}{4q_0^2\left(z^2 - 1\right)} = \frac{z\left(q_0x(z) - \frac{\theta}{2}\right)}{x(z)}. \end{cases}$$
(4.9)

We call this spectral curve the HTW spectral curve. Note that the branch point is located at z = 0 and the local conjugate point around the branch point is given by $\bar{z} = -z$. We define the Eynard-Orantin differentials and the symplectic invariants $F_{\text{HTW}}^{(g)}$ for the HTW spectral curve (4.9) in the same way as (3.23), (3.25), (3.29) and (3.31). To be more precise, in the definition of $F_{\text{HTW}}^{(0)}$, the sum in (3.25) runs over $\alpha \in \{+1, -1, \infty\}$ and $\xi_1(z) = \xi_{-1}(z) = x(z)^{-1}$ and $\xi_{\infty}(z) = x(z)^{1/2}$. On the other hand, following Chapter 7 of [16], we define $F_{\text{HTW}}^{(1)}$ by

$$F_{\rm HTW}^{(1)} = -\frac{1}{24} \ln \left(y'(0) \right).$$

Here y'(0) is defined in (3.30). Then we find:

$$F_{\rm HTW}^{(0)}(t) = \frac{q_0^6}{3} + \frac{5\theta}{6}q_0^3 - \frac{3\theta^2}{4} + \frac{\theta^2}{4}\ln\left(-8q_0^2\right)$$

$$F_{\rm HTW}^{(1)}(t) = -\frac{1}{24}\ln\left(\frac{1}{\sqrt{2}}\left(1 + \frac{\theta}{4q_0^3}\right)\right),$$

$$F_{\rm HTW}^{(2)}(t) = \frac{\theta\left(700\,q_0^6 - 85\,\theta\,q_0^3 - 2\theta^2\right)}{480\,(4q_0^3 + \theta)^5},$$

$$F_{\rm HTW}^{(3)}(t) = \frac{\theta}{4032\,(4q_0^3 + \theta)^{10}}\left(6726720q_0^{15} - 5017712\theta q_0^{12} + 541132\theta^2 q_0^9 - 1089\theta^3 q_0^6 + 160\theta^4 q_0^3 + 4\theta^5\right). \quad (4.10)$$

Additionally, as $q_0 \to \infty$ (i.e., $t \to \infty_{B,C}$), it is easy to prove by recursion that the correlation functions and symplectic invariants $F_{\text{HTW}}^{(g)}$ (identified with $\omega_0^{(g)}$ in the next formula) generated by the topological recursion on (4.9) behave like:

$$\omega_n^{(g)}(z_1,\ldots,z_n) \underset{q_0 \to \infty}{\sim} \operatorname{Cste} q_0^{-3(2g-2+n)} dz_1 \cdots dz_n.$$
(4.11)

Indeed, the recursion kernel behaves like

$$K(z_0, z) = \frac{(z^2 - 1)}{2z(z^2 - z_0^2)(\theta - 4q_0^3(z^2 - 1))} \frac{dz_0}{dz} = O\left(q_0^{-3}\right).$$
(4.12)

Thus adding q_0^{-3} at each step of the recursion. In particular we find that:

$$\lim_{t \to \infty_{B,C}} F_{\text{HTW}}^{(g)}(t) = 0 \quad \text{for } g \ge 2.$$
(4.13)

18

4.3. Tau-function and symplectic invariants for the HTW pair. In this section we state the second main result of this paper.

Theorem 4.1. The HTW pair is of topological type (in the sense of Section 5) and we have:

$$\frac{dF_{\rm HTW}^{(g)}(t)}{dt} = -\sigma_{2g}(t) \quad for \ g \ge 2.$$
(4.14)

In particular we can verify from (4.10) that (4.14) is correct for g = 2 and g = 3. Proof of Theorem 4.1 will be given in Section 5 and in appendix. We can also verify that

$$\frac{dF_{\rm HTW}^{(0)}(t)}{dt} = -\left(\sigma_0(t) + \frac{t^2}{8}\right), \quad \frac{dF_{\rm HTW}^{(1)}(t)}{dt} = -\sigma_2(t) \tag{4.15}$$

holds in accordance with (4.7). This leads to the following theorem:

Theorem 4.2. The generating function of symplectic invariants of the HTW curve (4.9) gives a τ -function of Painlevé 2. In other words,

$$\ln \tau_{\rm HTW} = \sum_{g=0}^{\infty} \hbar^{2g-2} F_{\rm HTW}^{(g)}(t)$$
 (4.16)

satisfies (4.7). Furthermore, in the both of limits $t \to \infty_B$ and $t \to \infty_C$ we have:

$$F_{\rm HTW}^{(g)}(t) = -\int_{\infty_{B,C}}^{t} \sigma_{2g}(s)ds \quad \text{for } g \ge 2.$$
 (4.17)

4.4. Limit at $t = \infty_A$: the Bessel curve. Let us realize the following affine symplectic transformation on the spectral curve (4.8):

$$x = \frac{2q_0^2}{\theta^2}X$$
, $y = \frac{\theta^2}{2q_0^2}Y$ (4.18)

we get the new spectral curve:

$$Y^{2} = \frac{(X + \theta^{2}) \left(1 - \frac{4q_{0}^{3}}{\theta^{3}}X\right)^{2}}{4X^{2}}.$$
(4.19)

When $t \to \infty_A$ (i.e., $q_0 \to 0$) we get that the limiting curve becomes

$$Y^2 = \frac{X + \theta^2}{4X^2},$$
 (4.20)

what we call the Bessel curve. It can be parametrized into:

$$\begin{cases} X(z) = \theta^2 (z^2 - 1), \\ Y(z) = \frac{z}{2\theta(z^2 - 1)}. \end{cases}$$
(4.21)

In particular straightforward computations of the topological recursion gives:

$$F_{\text{Bessel}}^{(0)} = -\frac{3\theta^2}{4} - \frac{\theta^2}{4} \ln\left(-\frac{1}{4\theta^2}\right)$$

$$F_{\text{Bessel}}^{(1)} = -\frac{1}{24} \ln\left(-\frac{1}{4\theta^2}\right),$$

$$F_{\text{Bessel}}^{(2)} = -\frac{1}{240\theta^2},$$

$$F_{\text{Bessel}}^{(3)} = \frac{1}{1008\theta^4}.$$
(4.22)

General properties regarding limits and symplectic transformations of the curve in the topological recursion tell us that:

$$\lim_{t \to \infty_A} F_{\text{HTW}}^{(g)}(t) = F_{\text{Bessel}}^{(g)} \quad \text{for } g \ge 2.$$
(4.23)

On the other hand, it follows from (3.53) and (4.17) that we have

$$\lim_{t \to \infty_A} F_{\text{HTW}}^{(g)} = -\int_{\infty_{B,C}}^{\infty_A} \sigma_{2g}(t) = \frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
(4.24)

Therefore, as a corollary of our main theorems, we have computed $F_{\text{Bessel}}^{(g)}$ explicitly:

Theorem 4.3. The symplectic invariants $F_{\text{Bessel}}^{(g)}$ of the Bessel curve (4.21) are given by

$$F_{\text{Bessel}}^{(g)} = \frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
(4.25)

To our knowledge, (4.25) has not been mentioned in the literature.

Remark 4.4. Y. Yamada pointed out to the authors that the JM curve and HTW curve are related by the following symplectic transformation:

$$x_{\rm HTW} = x_{\rm JM}^2 + y_{\rm JM} + \frac{t}{2}, \quad y_{\rm HTW} = -x_{\rm JM} + \frac{\theta}{2x_{\rm JM}^2 + 2y_{\rm JM} + t},$$
 (4.26)

where $(x_{\text{JM/HTW}}, y_{\text{JM/HTW}})$ are coordinates in the expressions (3.21) and (4.8) of JM/HTW curve. Note that the parametrizations (3.22) and (4.9) of JM/HTW curve are not compatible with this symplectic transformation. As is shown in [18], a class of symplectic transformations (which includes the transformation $x \leftrightarrow y$) do not preserve $F^{(g)}$ in general; this explains why there is a discrepancy between $F_{\text{JM}}^{(g)}$ and $F_{\text{HTW}}^{(g)}$. Our computation shows that the difference is explicitly described by the Bernoulli numbers:

$$F_{\rm JM}^{(g)} - F_{\rm HTW}^{(g)} = -\frac{B_{2g}}{2g(2g-2)\theta^{2g-2}} \quad \text{for } g \ge 2.$$
 (4.27)

In Appendix E we show that the r.h.s. of (4.27) coincides with the integration constant computed in [18].

20

5. Determinantal formulas and topological type property

In this section we review the determinantal formulas formalism and the issue of the topological type property. Then we mention our main results and discuss about the consequences. The proof of the topological type properties are postponed in Subsection 5.3, Appendix A and B.

5.1. **Determinantal formulas.** Here we remind the reader about determinantal formulas developed in [3].

Determinantal formulas are built from a solution

$$\Psi(x,t) = \begin{pmatrix} \psi(x,t) & \phi(x,t) \\ \tilde{\psi}(x,t) & \tilde{\phi}(x,t) \end{pmatrix}$$
(5.1)

of the JM or the HTW Lax pair:

$$\hbar \frac{\partial}{\partial x} \Psi = \mathcal{D}\Psi , \ \hbar \frac{\partial}{\partial t} \Psi = \mathcal{R}\Psi$$
 (5.2)

where the matrices \mathcal{D} and \mathcal{R} are traceless and have formal series expansions

$$\mathcal{D}(x,t) = \sum_{k=0}^{\infty} D^{(k)}(x,t)\hbar^k , \ \mathcal{R}(x,t) = \sum_{k=0}^{\infty} R^{(k)}(x,t)\hbar^k$$
(5.3)

We take a matrix-type WKB formal solution

$$\Psi(x,t) = \left(\sum_{k=0}^{\infty} \Psi^{(k)}(x,t)\hbar^k\right) \exp\left(\frac{T(x,t)}{\hbar}\right), \quad T(x,t) = \operatorname{diag}(s(x,t), -s(x,t))$$
(5.4)

which is normalized by det $\Psi = 1$. Here the phase function s(x, t) satisfies

$$\frac{\partial}{\partial x}s(x,t) = \sqrt{E_{\infty}(x,t)},\tag{5.5}$$

where

$$E_{\infty}(x,t) = -\det \mathcal{D}^{(0)}(x,t)$$

=
$$\begin{cases} (x-q_0)^2(x^2+2q_0x+3q_0^2+t) & \text{for JM case,} \\ \frac{1}{2x^2}\left(x-\frac{\theta}{2q_0}\right)^2(x+2q_0^2) & \text{for HTW case.} \end{cases}$$
(5.6)

Determinantal formulas are obtained from the Christoffel-Darboux kernel:

$$K(x_1, x_2) = \frac{\psi(x_1)\tilde{\phi}(x_2) - \tilde{\psi}(x_1)\phi(x_2)}{x_1 - x_2}$$
(5.7)

(here we are omitting the *t*-dependence for simplicity) with the following definition:

Definition 5.1 (Definition 2.3 of [3]). The (connected) correlation functions are defined by:

$$W_{1}(x) = \frac{\partial \psi}{\partial x}(x)\tilde{\phi}(x) - \frac{\partial \psi}{\partial x}(x)\phi(x),$$

$$W_{n}(x_{1},...,x_{n}) = -\frac{\delta_{n,2}}{(x_{1}-x_{2})^{2}} + (-1)^{n+1}\sum_{\sigma:n-\text{cycles}}\prod_{i=1}^{n}K(x_{i},x_{\sigma(i)})$$

for $n \geq 2.$
(5.8)

The correlation functions are formal power series of \hbar whose coefficients are symmetric functions of x_1, \ldots, x_n . Note that there exists an alternative expression for the correlation functions [3]. Define the rank 1 projector by

$$M(x,t) = \Psi(x,t) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \Psi^{-1}(x,t) = \begin{pmatrix} \psi \tilde{\phi} & -\psi \phi \\ \tilde{\psi} \tilde{\phi} & -\phi \tilde{\psi} \end{pmatrix}.$$
 (5.9)

It is in fact the canonical projector on the first coordinate taken into the basis defined by $\Psi(x,t)$. The rank 1 projector satisfies:

$$M^2 = M$$
, $\text{Tr}M = 1$, $\det M = 0.$ (5.10)

Theorem 2.1 of [3] gives an alternative expression for $W_n(x_1, \ldots, x_n)$:

$$W_{1}(x) = -\frac{1}{\hbar} \operatorname{Tr}(\mathcal{D}(x)M(x)),$$

$$W_{2}(x_{1}, x_{2}) = \frac{\operatorname{Tr}(M(x_{1})M(x_{2})) - 1}{(x_{1} - x_{2})^{2}},$$

$$W_{n}(x_{1}, \dots, x_{n}) = (-1)^{n+1} \operatorname{Tr} \sum_{\sigma: n- \text{cycles}} \prod_{i=1}^{n} \frac{M(x_{\sigma(i)})}{x_{\sigma(i)} - x_{\sigma(i+1)}}$$

$$= \frac{(-1)^{n+1}}{n} \sum_{\sigma \in S_{n}} \frac{\operatorname{Tr}M(x_{\sigma(1)}) \dots M(x_{\sigma(n)})}{(x_{\sigma(1)} - x_{\sigma(2)}) \dots (x_{\sigma(n-1)} - x_{\sigma(n)})}$$

for $n \geq 3.$ (5.11)

5.2. Topological type property. Now we give the definition of the topological type property for the differential equation (5.2) having the spectral curve of genus 0.

Definition 5.2 (Definition 3.3 of [2], Section 2.5 of [3]). The differential system (5.2) is said to be of topological type if the correlation functions W_n given in Definition 5.1 satisfy the following three conditions:

- (1) Parity property: $W_n|_{\hbar \mapsto -\hbar} = (-1)^n W_n$ hold for $n \ge 1$.
- (2) Leading order property: The leading order of the series expansion of the correlation function W_n is at least of order \hbar^{n-2} . When these two conditions are satisfied, W_n has the following expansion (called a topological

expansion):

$$W_n(x_1, \dots, x_n) = \sum_{g=0}^{\infty} \hbar^{2g-2+n} W_n^{(g)}(x_1, \dots, x_n) \quad \text{for } n \ge 1.$$
 (5.12)

(3) Pole property: The coefficients $W_n^{(g)}(x_1, \ldots, x_n)$ of correlation functions $\overline{W_n}$ have no poles at even zeros of $E_{\infty}(x)$ given in (5.6).

The authors of [2] and [3] proved that:

Proposition 5.3 (Theorem 2.1 of [3], Theorem 3.1 and Corollary 4.2 of [2]).

(i) If the differential system (5.2) is of topological type, then the coefficients $W_n^{(g)}(x_1, \ldots, x_n)$ appearing in the expansion (5.12) of the correlation function $W_n(x_1, \ldots, x_n)$ are identical to Eynard-Orantin differentials $\omega_n^{(g)}(z_1, \ldots, z_n)$ obtained from the topological recursion applied on the spectral curve $y^2 = E_{\infty}(x)$ in the following way:

$$W_n^{(g)}(x(z_1), \dots, x(z_n))dx(z_1) \cdots dx(z_n) = \omega_n^{(g)}(z_1, \dots, z_n)$$

for $g \ge 0$ and $n \ge 1$, (5.13)

where x(z) is the rational function of z appearing in the parametrization (3.22) or (4.9) of the spectral curve.

(ii) Suppose that the differential system (5.2) is of topological type. Then, the generating function

$$\sum_{g=0}^{\infty} \hbar^{2g-2} F_g(t)$$

of the symplectic invariants F_g obtained from the topological recursion applied to $y^2 = E_{\infty}(x)$ gives the isomonodromic tau-function associated with (5.2) in the sense of Jimbo-Miwa-Ueno ([23]).

Here we omit the general definition of the Jimbo-Miwa-Ueno's isomonodromic tau-function (see Section 5 of [23]; see also Section 4.2 in [2] and Section 1.5 in [7]). In the case of JM/HTW Lax pair, the isomonodromic tau-function $\tau_{\rm JM/HTW}$ is defined (up to constant) by

$$2 \operatorname{Res}_{x=\infty} \left(\frac{1}{\hbar} \frac{\partial s_{\infty}}{\partial t}(x) W_1(x) dx \right) = \begin{cases} \frac{d}{dt} \ln \tau_{\mathrm{JM}}(t,\hbar) & \text{for JM case,} \\ \frac{d}{dt} \ln \tau_{\mathrm{HTW}}(t,\hbar) & \text{for HTW case.} \end{cases}$$
(5.14)

where $W_1(x)$ is given in (5.8), and $s_{\infty}(x)$ is the divergent part of s(x) given in (5.4) when $x \to \infty$:

$$s_{\infty}(x) = \begin{cases} \frac{x^3}{3} + \frac{tx}{2} & \text{for JM case,} \\ \frac{\sqrt{2}x^{\frac{3}{2}}}{3} - \frac{tx^{\frac{1}{2}}}{\sqrt{2}} & \text{for HTW case.} \end{cases}$$
(5.15)

We can verify that the above definition of the tau-function is consistent with (3.15) and (4.7), respectively, since $W_1(x)$ behaves as

$$W_{1}(x) = \begin{cases} \frac{x^{2}}{\hbar} + \frac{t}{2\hbar} - \frac{\theta}{\hbar x} + \frac{\sigma(t)}{\hbar x^{2}} + O(x^{-2}) & \text{for JM case,} \\ \frac{x^{\frac{1}{2}}}{\sqrt{2\hbar}} - \frac{tx^{-\frac{1}{2}}}{2\sqrt{2\hbar}} - \frac{\sigma(t) + \frac{t^{2}}{8}}{\sqrt{2\hbar}} x^{-\frac{3}{2}} + O(x^{-2}) & \text{for HTW case.} \end{cases}$$
(5.16)

when $x \to \infty$. As is explained in [2, 7], the claim (ii) in Proposition 5.3 is a consequence of the claim (i) and the "variation formula" established in [17]. To make our paper self-contained, we will explain this point in Appendix D.

Our context fits perfectly with Proposition 5.3 and thus only the proof of the topological type property remains. Our main theorem (including the statement of Theorem 3.4 and 4.2) claims that the conditions $(1) \sim (3)$ in Definition 5.2 hold for both JM and HTW Lax pairs:

Theorem 5.4. For any choice of the monodromy parameter $\theta \neq 0$, both of the Jimbo-Miwa Lax pair (3.3) and the Harnad-Tracy-Widom Lax pair (4.3) are of topological type. Therefore, the \hbar -expansion of the tau-function τ_{2g} and correlation functions $W_n^{(g)}$ respectively identify with the symplectic invariants $F^{(g)}$ and the Eynard-Orantin differentials $\omega_n^{(g)}$ computed from the topological recursion applied to the corresponding spectral curves (3.22) and (4.9).

We will prove that the three conditions $(1) \sim (3)$ in Definition 5.2 hold for both Lax pairs in the rest of Section 5 and Appendix A, B. We note that, in the previous works [2, 3, 28], the leading order property (2) of correlation function W_n was derived by using so-called "insertion operator". However, we find that there is an incompleteness in the proof using the insertion operators. In next subsection we give another proof of the leading order property without using the insertion operator. Our new method shows that the leading order property (2) follows from the pole property (3). The parity property (1) and the no pole property (3) will be proved in Appendix A and B, respectively. We will also explain why the proof based on the insertion operator is incomplete in Appendix C.

5.3. Proof of the leading order property of W_n using loop equations. This subsection is dedicated to prove that the both JM and HTW Lax pair enjoy the

property (2) in Definition 5.2; that is, the leading order of the series expansion in \hbar of W_n is at least of order \hbar^{n-2} :

$$W_n(x_1, \dots, x_n) = O(\hbar^{n-2}) \text{ for } n \ge 1.$$
 (5.17)

Determinantal formulas in Definition 5.1 have been introduced so that they satisfy a set of equations known as the loop equations. These loop equations (also known as Schwinger-Dyson equations) originate in random matrix theory where they are crucial. We recall here the main result of [3]:

Proposition 5.5 (Theorem 2.9 of [3]). Let us define the following functions (we denote by L_n the set of variables $\{x_1, \ldots, x_n\}$):

$$P_{1}(x) = \frac{1}{\hbar^{2}} \det \mathcal{D}(x,t),$$

$$P_{2}(x;x_{2}) = \frac{1}{\hbar} \operatorname{Tr} \left(\frac{\mathcal{D}(x,t) - \mathcal{D}(x_{2},t) - (x-x_{2})\mathcal{D}'(x_{2},t)}{(x-x_{2})^{2}} M(x_{2}) \right),$$

$$P_{n+1}(x;L_{n}) = (-1)^{n} \left[Q_{n+1}(x;L_{n}) - \sum_{j=1}^{n} \frac{1}{x-x_{j}} \operatorname{Res}_{x' \to x_{j}} Q_{n+1}(x',L_{n}) \right],$$

$$Q_{n+1}(x;L_{n}) = \frac{1}{\hbar} \sum_{\sigma \in S_{n}} \frac{\operatorname{Tr} \left(\mathcal{D}(x)M(x_{\sigma(1)}) \dots M(x_{\sigma(n)}) \right)}{(x-x_{\sigma(2)}) \dots (x_{\sigma(n-1)} - x_{\sigma(n)})(x_{\sigma(n)} - x)}.$$
(5.18)

Then, the correlation functions satisfy

$$P_1(x) = W_2(x, x) + W_1(x)^2, (5.19)$$

and

$$0 = P_{n+1}(x; L_n) + W_{n+2}(x, x, L_n) + 2W_1(x)W_{n+1}(x, L_n) + \sum_{J \subset L_n, J \notin \{\emptyset, L_n\}} W_{1+|J|}(x, J)W_{1+n-|J|}(x, L_n \setminus J) + \sum_{j=1}^n \frac{d}{dx_j} \frac{W_n(x, L_n \setminus x_j) - W_n(L_n)}{x - x_j} \quad \text{for } n \ge 1.$$
(5.20)

Moreover $P_{n+1}(x; L_n)$ is a rational function of x whose poles are at the poles of $\mathcal{D}(x, t)$.

The equations (5.19) and (5.20) are called the loop equations. As we will see this proposition and a subtle induction are sufficient to prove that W_n is at least of order \hbar^{n-2} . Let us now make the following crucial observation:

Theorem 5.6. In the JM Lax pair case, $P_{n+1}(x; L_n)$ does not depend on x for $n \ge 1$. In the HTW Lax pair case, the functions $P_{n+1}(x; L_n)$ are of the form $P_{n+1}(x; L_n) = \frac{1}{x} \tilde{P}_{n+1}(L_n)$.

Proof. In the JM case, since the entries of $\mathcal{D}(x,t)$ is polynomial of x, $P_{n+1}(x;L_n)$ may only have singularity at $x = \infty$. However looking at large x the definition shows that $P_{n+1}(x;L_n)$ can only be a polynomial of degree 0 and hence does not depend on x. In addition we get an explicit formula:

$$P_{n+1}(x; L_n) = P_{n+1}(L_n)$$

= $\frac{(-1)^{n+1}}{\hbar} \sum_{\sigma \in S_n} \frac{\text{Tr}\left(\sigma_3 M(x_{\sigma(1)}) \dots M(x_{\sigma(n)})\right)}{(x_{\sigma(1)} - x_{\sigma(2)}) \dots (x_{\sigma(n-1)} - x_{\sigma(n)})} \text{ for } n \ge 1.$ (5.21)

For example, using directly the definition we have $P_2(x; x_2) = \frac{1}{\hbar} \operatorname{Tr}(\sigma_3 M(x_2))$ which is indeed independent of x. In the HTW case, the form of $\mathcal{D}(x, t)$ implies that $P_{n+1}(x; L_n)$ may only have simple poles at x = 0 and a simple zero at infinity (degree of numerator-denominator shows that it behaves as $O(x^{-1})$ at infinity). Hence we conclude that it is proportional to x^{-1} (one could even get a complete expression by taking the residue at x = 0).

We now have all the ingredients to prove the following theorem:

Theorem 5.7. The correlation functions $W_n(x_1, \ldots, x_n)$ admit a series expansion in \hbar starting at least at order \hbar^{n-2} .

The rest of Section 5.3 is devoted to the proof of Theorem 5.7. Our proof is done by induction. We will denote $L_i = \{x_1, \ldots, x_i\}$. Let us define the following statement:

$$\mathcal{P}_k : W_j(x_1, \dots, x_j) \text{ is at least of order } \hbar^{k-2} \text{ for } j \ge k.$$
 (5.22)

We can easily verify that the correlation functions W_n for $n \ge 2$ has the formal series expansion of the form

$$W_n(x_1, \dots, x_n) = \sum_{k=0}^{\infty} w_n^{(k)}(x_1, \dots, x_n)\hbar^k \text{ for } n \ge 2.$$
 (5.23)

It also follows from the first loop equation (5.19) that $W_1(x)$ has the formal series expansion of the form

$$W_1(x) = \sum_{k=-1}^{\infty} w_1^{(k)}(x)\hbar^k, \qquad (5.24)$$

where the leading term $w_1^{(-1)}(x)$ coincides with $\sqrt{E_{\infty}(x)}$. Indeed, let us look the first loop equation (5.19). Since $W_2(x_1, x_2)$ only starts at \hbar^0 while $P_1(x) = \frac{1}{\hbar^2} \det \mathcal{D}$ looking at order \hbar^{-2} in the last equation provides the result. Thus we have checked that the statement \mathcal{P}_1 is true. The statement \mathcal{P}_2 is obviously true in view of (5.23).

Let us assume that the statement \mathcal{P}_i is true for all $i \leq n$ for some integer $n \geq 2$. Now we look at the loop equation (5.20). By the induction hypothesis, we have that the last two terms are at least of order \hbar^{n-2} . Indeed in the sum we have terms of order $\hbar^{1+|J|-2+1+n-|J|-2} = \hbar^{n-2}$. Moreover we also have from the same assumption that $W_{n+2}(x, x, L_n)$ is also of order at least \hbar^{n-2} (since $n+2 \ge n$). Eventually $W_{n+1}(x, L_n)$ is at least of order \hbar^{n-2} so we get when looking at order \hbar^{n-3} in (5.20):

$$0 = P_{n+1}^{(n-3)}(x; L_n) + 2w_1^{(-1)}(x)w_{n+1}^{(n-2)}(x, L_n).$$
(5.25)

Here $P_{n+1}^{(\ell)}(x; L_n)$ is the coefficient of \hbar^{ℓ} in the \hbar -expansion of $P_{n+1}(x; L_n)$. If we assume that $w_{n+1}^{(n-2)}(x, L_n) \neq 0$ then we have:

$$w_{n+1}^{(n-2)}(x,L_n) = \frac{P_{n+1}^{(n-3)}(x;L_n)}{2w_1^{(-1)}(x)}.$$
(5.26)

In our cases we get:

$$w_{n+1}^{(n-2)}(x, L_n) = \begin{cases} \frac{P_{n+1}^{(n-3)}(L_n)}{2(x-q_0)\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} & \text{for JM case,} \\ \frac{\tilde{P}_{n+1}^{(n-3)}(L_n)}{\left(x - \frac{\theta}{2q_0}\right)\sqrt{x+2q_0^2}} & \text{for HTW case.} \end{cases}$$
(5.27)

In both cases we obtain that $w_{n+1}^{(n-2)}(x, L_n)$ must have a simple pole at the even zero of $E_{\infty}(x)$, and this contradicts the pole property which will be proved in Appendix B. Consequently we must have $w_{n+1}^{(n-2)}(x, L_{i_0}) = 0$. This proves that $w_{n+1}(x, L_n)$ is at least of order \hbar^{n-1} .

To complete the proof of \mathcal{P}_{n+1} , we now need to prove the same statement for higher correlation functions. Let us prove it by a second induction by defining:

$$\tilde{\mathcal{P}}_i : W_i(x_1, \dots, x_i) \text{ is of order at least } \hbar^{n-1}.$$
 (5.28)

We want to prove $\tilde{\mathcal{P}}_i$ for all $i \geq n+1$ by induction. We just proved it for i = n+1 so initialization is done. Let us assume that $\tilde{\mathcal{P}}_j$ is true for all j satisfying $n+1 \leq j \leq i_0$. We look at the loop equation:

$$0 = P_{i_0+1}(x; L_{i_0}) + W_{i_0+2}(x, x, L_{i_0}) + 2W_1(x)W_{i_0+1}(x, L_{i_0}) + \sum_{J \subset L_{i_0}, J \notin \{\emptyset, L_{i_0}\}} W_{1+|J|}(x, J)W_{1+i_0-|J|}(x, L_{i_0} \setminus J) + \sum_{j=1}^{i_0} \frac{d}{dx_j} \frac{W_{i_0}(x, L_{i_0} \setminus x_j) - W_{i_0}(L_{i_0})}{x - x_j}.$$
(5.29)

By assumption on $\tilde{\mathcal{P}}_{i_0}$, the last sum with the derivatives contains terms of order at least \hbar^{n-1} . In the sum involving the subsets of L_{i_0} it is straightforward to see that the terms are all of order at least \hbar^{n-1} . Indeed, as soon as one of the index is greater then n + 1, the assumption of $\tilde{\mathcal{P}}_i$ for $n + 1 \leq i \leq i_0$ tells us that this term is already at order at least \hbar^{n-1} . Since the second factor of the product is at least of order \hbar^0 then it does not decrease the order. Now if both factors have indexes strictly lower than n + 1, then the assumption of \mathcal{P}_j for all $j \leq n$ tell us that the order of the product is at least of $\hbar^{|J|+1-2+1+i_0-|J|-2} = \hbar^{i_0-2}$ which is greater than n-1 since $i_0 \ge n+1$. Additionally by induction on \mathcal{P}_n we know that $W_{i_0+1}(x, L_{i_0})$ is at least of order \hbar^{n-2} as well as $W_{i_0+2}(x, x, L_{i_0})$. Consequently looking at order \hbar^{n-3} in (5.29) gives:

$$0 = P_{i_0+1}^{(n-3)}(x; L_{i_0}) + 2w_1^{(-1)}(x)w_{i_0+1}^{(n-2)}(x, L_{i_0}).$$
(5.30)

We can apply a similar reasoning as for (5.25). If we assume that $w_{i_0+1}^{(n-2)}(x, L_{i_0}) \neq 0$, then we have:

$$w_{i_0+1}^{(n-2)}(x, L_{i_0}) = \frac{P_{i_0+1}^{(n-3)}(x; L_{i_0})}{2w_1^{(-1)}(x)}.$$
(5.31)

In our two cases we get:

$$w_{i_0+1}^{(n-2)}(x, L_{i_0}) = \begin{cases} \frac{P_{i_0+1}^{(n-3)}(L_{i_0})}{2(x-q_0)\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} & \text{for JM case} \\ \frac{\tilde{P}_{i_0+1}^{(n-3)}(L_{i_0})}{\left(x - \frac{\theta}{2q_0}\right)\sqrt{x+2q_0^2}} & \text{for HTW case.} \end{cases}$$
(5.32)

In both cases we obtain that $w_{i_0+1}^{(n-2)}(x, L_{i_0})$ must have a simple pole at the even zero of $E_{\infty}(x)$ which contradicts the pole property that will be proved in Appendix B. Consequently we must have $w_{i_0+1}^{(n-2)}(x, L_{i_0}) = 0$. In particular it means that $w_{i_0+1}(x, L_{i_0})$ (which by assumption of \mathcal{P}_n was already known to be of order \hbar^{n-2}) is at least of order \hbar^{n-1} thus making the induction on $\tilde{\mathcal{P}}_{i_0}$. Hence by induction we have proved that $\forall i \geq n+1$, $\tilde{\mathcal{P}}_i$ holds which exactly proves that \mathcal{P}_{n+1} is true. Eventually by induction we have just proved that \mathcal{P}_n holds for $n \geq 1$, which implies the desired property (5.17).

Remark 5.8. Several important observations can be made about this proof:

- The proof heavily relies on the pole property of the correlation functions $W_n^{(g)}$. In particular it is central to know that the correlation functions are regular at the even zeros of $E_{\infty}(x)$ since it provides the contradiction in (5.25) and (5.30).
- The possible poles of $\mathcal{D}(x,t)$ are irrelevant in the proof. Indeed, they specify the form of $P_{n+1}(x;L_n)$ but do not play an important role in the contradiction of (5.25) and (5.30).
- The presence of at least one even zero in the spectral curve is necessary in our proof because $\mathcal{D}(x,t)$ is a polynomial of degree 2 in the JM case or has a simple pole at x = 0 in the HTW case. In the case when $\mathcal{D}(x,t)$ is a polynomial of degree 1, then $P_{n+1}(x; L_n)$ would be identically zero and thus our proof would also work in a simpler way. Hence the central element is the balance between the order of the singularity of $\mathcal{D}(x,t)$ and the fact that the spectral curve is of genus 0.
- This proof can be applied to more general cases: as soon as the spectral curve has a double zero and the pole structure is proved then the method

28

can be applied. In the case of Lax pairs, the pole property is usually easy to obtain from the t-differential equation like we did in Appendix B and the spectral curve is even simpler to obtain.

6. Outlooks

We believe that the topological type property (TT property) should hold more general Lax pairs including those for all six Painlevé equations with arbitrary (generic) monodromy parameters. The previous works in this direction treats Painlevé equation with a specific monodromy parameters (Cf. [7, 28]), but our work shows that the absence of monodromy parameter is not a necessary condition to obtain the TT property. Moreover, in this paper we have introduced new general methods to prove the TT property (see Section 5.3 and Appendix B) that should generalize easily for other Lax pairs. In fact, while this article was under peer review, the authors together with A. Saenz succeeded in proving the TT property for all six Painlevé equations ([21]). Several natural questions arise from this work:

- At the conceptual level a better understanding would definitely be an interesting development. Indeed it is even unclear so far to what extent the TT property is connected with Lax pairs. Does it work for any Lax pairs? Only specific integrable systems ? Our new methods seem general enough to work for many Lax pairs but a better understanding of the scope and limits of the methods is required.
- On a totally different perspective, we have shown here that studying Lax pairs and their symplectic invariants may lead to explicit formulas for symplectic invariants of new spectral curves. Cases where general formulas for the symplectic invariants are known explicitly are extremely rare and this could provide a way to improve the classification of symplectic invariants for simple spectral curves. This knowledge might be of some use for enumerative geometry where a list of spectral curves and associated symplectic invariants would be helpful.
- In this paper we also drew attention on the incompleteness of the insertion operator method. This calls for a better definition of the insertion operator to fix the current problem.

APPENDIX A. PROOF OF THE PARITY SYMMETRY

We want to prove that the \hbar series expansion of the determinantal formulas $W_n(x_1, \ldots, x_n)$ only involves powers of \hbar of the same parity. (Cf., (5.12)). In order to do this, we use Proposition 3.3 of [2] that gives a sufficient criteria to obtain the $\hbar \leftrightarrow -\hbar$ symmetry. We recall their proposition here:

Proposition A.1 (Proposition 3.3 of [2]). Let us denote \dagger the operator that change \hbar into $-\hbar$. If there exists an invertible matrix $\Gamma(t)$ independent of x such that:

$$\Gamma(t)\mathcal{D}^{t}(x,t)\Gamma^{-1}(t) = \mathcal{D}^{\dagger}(x,t), \qquad (A.1)$$

then the correlators W_n satisfy

$$W_n^{\dagger} = (-1)^n W_n \quad \text{for } n \ge 1. \tag{A.2}$$

In particular if this proposition is satisfied then it automatically follows that the \hbar expansion a given function $W_n(x_1, \ldots, x_n)$ may only involve powers of \hbar with the same parity (given by the parity of n). Therefore all we have to do is prove the existence of a suitable matrix $\Gamma(t)$ for our two Lax pairs.

Recall that $\sigma^{\dagger} = \sigma$ and $p^{\dagger} = p$ hold (see (3.16) and (3.18)). Then, it follows from (3.6) that q satisfies

$$q^{\dagger} = -q - \frac{\theta}{p}.\tag{A.3}$$

Using this relation we can obtain $\mathcal{D}^{\dagger}(x,t)$ and we can find an invertible matrix Γ satisfying (A.1) as follows:

Theorem A.2. We can find suitable matrices $\Gamma(t)$ for our Lax pairs:

• For the Jimbo-Miwa case, the matrix

$$\Gamma(t) = \begin{pmatrix} 1 & 0\\ 0 & -2p(t) \end{pmatrix} \tag{A.4}$$

satisfies (A.1).

• For the Harnad-Tracy-Widom case, the matrix

$$\Gamma(t) = \begin{pmatrix} 1 & \frac{p(t)}{2\theta} \\ \frac{p(t)}{2\theta} & 0 \end{pmatrix}$$
(A.5)

satisfies (A.1).

Consequently, the series expansion in \hbar for W_n only involves even (resp. odd) powers of \hbar when n is even (resp. odd).

Theorem A.2 follows from (A.3) immediately. In Jimbo-Miwa case we have

$$\mathcal{D}^{\dagger}(x,t) = \begin{pmatrix} x^2 + p + \frac{t}{2} & x + q + \frac{\theta}{p} \\ -2p(x-q) & -\left(x^2 + p + \frac{t}{2}\right) \end{pmatrix}$$

and

$$\mathcal{D}^{t}(x,t) = \begin{pmatrix} x^{2} + p + \frac{t}{2} & -2p\left(x + q + \frac{\theta}{p}\right) \\ x - q & -\left(x^{2} + p + \frac{t}{2}\right) \end{pmatrix},$$

while in the Harnad-Tracy-Widom case we have:

$$\mathcal{D}^{\dagger}(x,t) = \begin{pmatrix} q + \frac{\theta}{p} + \frac{\theta}{2x} & x - p - 2\left(q + \frac{\theta}{p}\right)^2 - t\\ \frac{1}{2} + \frac{p}{2x} & -\left(q + \frac{\theta}{p} + \frac{\theta}{2x}\right) \end{pmatrix}$$

and

$$\mathcal{D}^{t}(x,t) = \begin{pmatrix} -q + \frac{\theta}{2x} & \frac{1}{2} + \frac{p}{2x} \\ x - p - 2q^{2} - t & q - \frac{\theta}{2x} \end{pmatrix}.$$

Then (A.1) can be checked easily by matrix multiplication in both cases.

30

APPENDIX B. PROOF OF THE POLE PROPERTY

As mentioned earlier we are interested in this article about solutions q(t) of Painlevé 2 admitting a formal expansion in the \hbar parameter. Consequently for both Lax pairs, this implies a series expansion in \hbar for M(x,t), W_n , $\mathcal{D}(x,t)$ and $\mathcal{R}(x,t)$ of the form:

$$\mathcal{D}(x,t) = \sum_{k=0}^{\infty} \mathcal{D}^{(k)}(x,t)\hbar^{k}$$
$$\mathcal{R}(x,t) = \sum_{k=0}^{\infty} \mathcal{R}^{(k)}(x,t)\hbar^{k},$$
$$M(x,t) = \sum_{k=0}^{\infty} M^{(k)}(x,t)\hbar^{k}.$$
(B.1)

In this appendix an exponent ^(k) denotes the coefficient of \hbar^k in the \hbar -expansion. Moreover, it is also obvious from the definitions of both Lax pairs that $\mathcal{D}^{(k)}(x,t)$ and $\mathcal{R}^{(k)}(x,t)$ do not depend on x for $k \geq 1$.

B.1. Jimbo-Miwa Lax pair. We want to prove that the matrices $M^{(k)}(x,t)$ only have singularities (as a function of x) at the branch points of the JM spectral curve and possibly a pole at infinity. The plan is first to compute explicitly $M^{(0)}(x,t)$ and then find a recursive relation between the matrices.

B.1.1. Computation of $M^{(0)}(x,t)$. Inserting the series expansion of M(x,t) into the differential system for M(x,t):

$$\hbar \partial_x M(x,t) = [\mathcal{D}(x,t), M(x,t)] \tag{B.2}$$

$$\hbar \partial_t M(x,t) = [\mathcal{R}(x,t), M(x,t)] \tag{B.3}$$

gives that:

$$0 = \left[\mathcal{D}^{(0)}(x,t), M^{(0)}(x,t)\right] \text{ and } 0 = \left[\mathcal{R}^{(0)}(x,t), M^{(0)}(x,t)\right].$$
(B.4)

Additionally, since M(x,t) is a rank 1 projector we know that TrM(x,t) = 1and det M(x,t) = 0. At order \hbar^0 this is equivalent to $\text{Tr}M^{(0)}(x,t) = 1$ and det $M^{(0)}(x,t) = 0$. Since $M^{(0)}(x,t)_{2,2} = 1 - M^{(0)}(x,t)_{1,1}$, the second equation of (B.4) only gives 3 different equations:

$$0 = \frac{1}{2} \left(M^{(0)}(x,t) \right)_{2,1} - \frac{\theta}{2q_0} \left(M^{(0)}(x,t) \right)_{1,2},
0 = \frac{1}{2} \left(1 - 2 \left(M^{(0)}(x,t) \right)_{1,1} \right) + (x+q_0) \left(M^{(0)}(x,t) \right)_{1,2},
0 = (x+q_0) \left(M^{(0)}(x,t) \right)_{2,1} + \frac{\theta}{2q_0} \left(1 - 2 \left(M^{(0)}(x,t) \right)_{1,1} \right).$$
(B.5)

We have used here the fact that $p_0 = -\frac{\theta}{2q_0}$ and $t = -2q_0^2 + \frac{\theta}{q_0}$. It is easy to observe that only two of the previous equations are independent. Therefore we have so far

a system of 2 independent equations with 3 unknowns. In order to complete it, we must use the fact that det $M^{(0)}(x,t) = 0$. In the end we find the following system:

$$0 = \frac{1}{2} \left(M^{(0)}(x,t) \right)_{2,1} - \frac{\theta}{2q_0} \left(M^{(0)}(x,t) \right)_{1,2},
0 = \frac{1}{2} \left(1 - 2 \left(M^{(0)}(x,t) \right)_{1,1} \right) + (x+q_0) \left(M^{(0)}(x,t) \right)_{1,2},
0 = \left(M^{(0)}(x,t) \right)_{1,1} \left(1 - \left(M^{(0)}(x,t) \right)_{1,1} \right) \\ - \left(M^{(0)}(x,t) \right)_{1,2} \left(M^{(0)}(x,t) \right)_{2,1}.$$
(B.6)

It is also important to note that the first equation of (B.4) would have lead exactly to the same system of equations. This system can be solved explicitly and we find:

$$M^{(0)}(x,t) = \begin{pmatrix} \frac{1}{2} + \frac{x+q_0}{2\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} & \frac{1}{2\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} \\ \frac{\theta}{2q_0\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} & \frac{1}{2} - \frac{x+q_0}{2\sqrt{(x+q_0)^2 + \frac{\theta}{q_0}}} \end{pmatrix}.$$
 (B.7)

It is obvious that $M^{(0)}(x,t)$ only have singularities at the branch points of the spectral curve. In particular, it is holomorphic at the even zero $x = q_0$ of $E_{\infty}(x)$ given by (5.6).

B.1.2. Recursive system for higher orders. Since $\text{Tr}M^{(k)}(x,t) = 0$ for $k \geq 1$, it suffices to consider the pole structure of $M^{(k)}(x,t)_{1,1}$, $M^{(k)}(x,t)_{1,2}$ and $M^{(k)}(x,t)_{2,1}$. Looking at order \hbar^k with $k \geq 1$ in (B.3), we have

$$\begin{bmatrix} \mathcal{R}^{(0)}(x,t), M^{(k)}(x,t) \end{bmatrix} = \partial_t M^{(k-1)}(x,t) -\sum_{i=0}^{k-1} \left[\mathcal{R}^{(k-i)}(x,t), M^{(i)}(x,t) \right] - \left[\mathcal{R}^{(k)}(x,t), M^{(0)}(x,t) \right].$$

Thus we get the following linear system

$$\begin{pmatrix} 0 & -\frac{\theta}{2q_0} & \frac{1}{2} \\ -1 & x+q_0 & 0 \\ x+q_0 & \frac{\theta}{2q_0} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} M^{(k)}(x,t)_{1,1} \\ M^{(k)}(x,t)_{1,2} \\ M^{(k)}(x,t)_{2,1} \end{pmatrix} = \begin{pmatrix} J^{(k)}(x,t)_1 \\ J^{(k)}(x,t)_2 \\ J^{(k)}(x,t)_3 \end{pmatrix},$$

where $J^{(k)}(x,t)_{\ell}$ are polynomials which are written in terms of $M^{(i)}(x,t)$ for $0 \leq i \leq k-1$ and their t-derivatives, and $\mathcal{R}^{(i)}(x,t) \ 0 \leq i \leq k$. Since

$$\det \begin{pmatrix} 0 & -\frac{\theta}{2q_0} & \frac{1}{2} \\ -1 & x+q_0 & 0 \\ x+q_0 & \frac{\theta}{2q_0} & \frac{1}{2} \end{pmatrix} = -\frac{1}{2} \left((x+q_0)^2 + \frac{\theta}{q_0} \right),$$
(B.8)

we get

$$\begin{pmatrix} M^{(k)}(x,t)_{1,1} \\ M^{(k)}(x,t)_{1,2} \\ M^{(k)}(x,t)_{2,1} \end{pmatrix} = \frac{1}{(x+q_0)^2 + \frac{\theta}{q_0}}$$

$$\times \begin{pmatrix} -(x+q_0) & -\frac{\theta}{q_0} & x+q_0\\ -1 & x+q_0 & 1\\ 2(x+q_0)^2 + \frac{\theta}{q_0} & \frac{(x+q_0)\theta}{q_0} & \frac{\theta}{q_0} \end{pmatrix} \begin{pmatrix} J^{(k)}(x,t)_1\\ J^{(k)}(x,t)_2\\ J^{(k)}(x,t)_3 \end{pmatrix}.$$
 (B.9)

Then, since we know that $\mathcal{R}^{(i)}(x,t)$'s are holomorphic at $x = q_0$ for all $i \ge 0$, a straightforward induction using (B.9) shows that the only singularities of $M^{(k)}(x,t)$ are at the branch points and possibly a pole at infinity:

Theorem B.1. For $k \ge 0$, the matrices $M^{(k)}(x,t)$ only have singularities at the branch points of the spectral curve $x = -q_0 \pm \sqrt{\theta/q_0}$ and a possible pole singularity at infinity. In particular they are holomorphic at the even zero $x = q_0$ of $E_{\infty}(x)$ given by (5.6). Consequently the same singularity structure holds for the functions $W_n^{(g)}(x_1,\ldots,x_n)$ thanks to the relation (5.11).

Remark B.2. It is also interesting to observe that using the differential equation in x rather than the one in t provides a similar linear equation instead of (B.9):

$$\begin{pmatrix} M^{(k)}(x,t)_{1,1} \\ M^{(k)}(x,t)_{1,2} \\ M^{(k)}(x,t)_{2,1} \end{pmatrix} = \frac{1}{(x-q_0)\left((x+q_0)^2 + \frac{\theta}{q_0}\right)} \\ \times \begin{pmatrix} -(x+q_0) & -\frac{\theta}{q_0} & x+q_0 \\ -1 & x+q_0 & 1 \\ 2(x+q_0)^2 + \frac{\theta}{q_0} & \frac{(x+q_0)\theta}{q_0} & \frac{\theta}{q_0} \end{pmatrix} \begin{pmatrix} \tilde{J}^{(k)}(x,t)_1 \\ \tilde{J}^{(k)}(x,t)_2 \\ \tilde{J}^{(k)}(x,t)_3 \end{pmatrix}.$$

However in this case, it is not easy to exclude the pole at the double zero $x = q_0$ of the spectral curve. Thus we understand here the importance of the differential equation with respect to t in the context of the determinantal formulas.

B.2. Harnad-Tracy-Widom Lax pair. Most of the arguments of the previous section also apply to the Harnad-Tracy-Widom Lax pair.

B.2.1. Computation of $M^{(0)}(x,t)$. Looking at $\left[\mathcal{R}^{(0)}(x,t), M^{(0)}(x,t)\right] = 0$ and det $M^{(0)}(x,t) = 0$ leads to the following system:

$$0 = -xM^{(0)}(x,t)_{2,1} + \frac{1}{2}M^{(0)}(x,t)_{1,2},$$

$$0 = -x\left(1 - 2M^{(0)}(x,t)_{1,1}\right) + 2q_0M^{(0)}(x,t)_{1,2},$$

$$0 = -2q_0M^{(0)}(x,t)_{2,1} + \frac{1}{2}\left(1 - 2M^{(0)}(x,t)_{1,1}\right),$$

$$0 = M^{(0)}(x,t)_{1,1}\left(1 - M^{(0)}(x,t)_{1,1}\right) - M^{(0)}(x,t)_{1,2}M^{(0)}(x,t)_{2,1}.$$

(B.10)

Note that only two of the first three equations are independent. This system of equations admits a unique solution given by:

$$M^{(0)}(x,t) = \begin{pmatrix} \frac{1}{2} - \frac{q_0}{\sqrt{2}\sqrt{x+2q_0^2}} & \frac{x}{\sqrt{2}\sqrt{x+2q_0^2}} \\ \frac{1}{2\sqrt{2}\sqrt{x+2q_0^2}} & \frac{1}{2} + \frac{q_0}{\sqrt{2}\sqrt{x+2q_0^2}} \end{pmatrix}.$$
 (B.11)

It is then obvious that $M^{(0)}(x,t)$ only have singularities at $x = -2q_0^2$ the unique branch point of the spectral curve and at infinity. Note also that $\left[\mathcal{D}^{(0)}(x,t), M^{(0)}(x,t)\right] = 0$ would have provided an equivalent system of equations.

B.2.2. Recursive system for higher orders. By the same method presented in Subsection B.1.2, we can derive a linear equation satisfied by the entries of $M^{(k)}(x,t)$:

$$\begin{pmatrix} 0 & \frac{1}{2} & -x \\ 2x & 2q_0 & 0 \\ 2q_0 & -\frac{1}{2} & -x \end{pmatrix} \begin{pmatrix} M^{(k)}(x,t)_{1,1} \\ M^{(k)}(x,t)_{1,2} \\ M^{(k)}(x,t)_{2,1} \end{pmatrix} = \begin{pmatrix} J^{(k)}(x,t)_1 \\ J^{(k)}(x,t)_2 \\ J^{(k)}(x,t)_3 \end{pmatrix},$$

where, as well as in the JM case, $J^{(k)}(x,t)_{\ell}$ are polynomials written in terms of $M^{(i)}(x,t)$ for $0 \leq i \leq k-1$ and $\mathcal{R}^{(i)}(x,t)$ $0 \leq i \leq k$. A straightforward computation shows that:

$$\det \begin{pmatrix} 0 & \frac{1}{2} & -x\\ 2x & 2q_0 & 0\\ 2q_0 & -\frac{1}{2} & -x \end{pmatrix} = 2x(x+2q_0^2)$$
(B.12)

which is non-vanishing at $x = \theta/(2q_0)$. Therefore, the same discussion given in Subsection B.1.2 shows the following.

Theorem B.3. For $k \ge 0$, the matrices $M^{(k)}(x,t)$ only have singularities at the branch point of the spectral curve $x = -2q_0^2$ and poles at x = 0 and $x = \infty$. In particular they are holomorphic at the even zero $x = \theta/(2q_0)$ of $E_{\infty}(x)$ given by (5.6). Consequently the same singularity structure holds for the functions $W_n^{(g)}(x_1,\ldots,x_n)$ thanks to the relation (5.11).

Here we again note that the differential equation in x is not helpful to show Theorem B.3 as well as in the JM case (see Remark B.2).

APPENDIX C. INCOMPLETE PROOF USING AN INSERTION OPERATOR

In [2, 3, 28] the various authors presented the construction of an insertion operator δ_{η} to prove the leading order of the \hbar expansion of W_n . Unfortunately this proof is incomplete and requires an important modification to be correct that is currently being investigated. We present here the main reason for the incompleteness of this method.

The method of the insertion operators naturally applies to the Picard-Vessiot (PV) ring \mathbb{B}_1 attached to the Lax pair, that is to say to the differential ring generated by the entries of $\Psi(x)$ and the scalar function $(\det \Psi(x))^{-1}$ over the

differential ring \mathbb{K}_1 of rational functions of x. Taking an arbitrary large number of spatial variables, we end up with the projective limit: $\mathbb{B}_{\infty} = \bigcup_{i=1}^{\infty} \mathbb{B}_i$ over the field $\mathbb{K}_{\infty} = \bigcup_{i=1}^{\infty} \mathbb{K}_i$ of rational functions in any arbitrary large number of variables x_i . Most quantities defined in this paper belongs to the PV ring \mathbb{B}_{∞} since they can be expressed with the entries of $\Psi(x)$ and $(\det \Psi(x))^{-1}$. For example, matrix elements of $\Psi^{-1}(x)$, $\mathcal{D}(x)$, $\mathcal{R}(x)$, $K(x_1, x_2)$, M(x) and all $W_n(x_1, \ldots, x_n)$ belong to \mathbb{B}_{∞} . The idea of the insertion operator is to create an operator δ_η acting on the PV ring that satisfies the following properties (i) ~ (v) (Cf. Definition 2.5, Definition 4.2 and Section 5.7.2 of [2]):

- (i) $\delta_{\eta}(\mathbb{K}_{\infty}) = 0$, and $\delta_{\eta}(\mathbb{B}_n) \subset \mathbb{B}_{n+1}$.
- (ii) δ_{η} is a derivation operator: $\delta_{\eta}(fg) = (\delta_{\eta}f)g + f(\delta_{\eta}g)$.
- (iii) δ_{η} inserts a variable into the correlation functions:

$$\delta_{\eta} W_n(x_1, \dots, x_n) = W_{n+1}(x_1, \dots, x_n, \eta).$$
(C.1)

This property is equivalent to impose that

$$\delta_{\eta} K(x_1, x_2) = -K(x_1, \eta) K(\eta, x_2).$$
(C.2)

- (iv) $\delta_{\eta}M(x)$ is of order \hbar , and is expressed in terms of M(x), $M(\eta)$ and their *t*-derivatives.
- (v) δ_{η} commutes with ∂_t .

With these properties it is then possible to show that the \hbar expansion of W_n must start at least at \hbar^{n-2} : Firstly, Property (C.1) implies

$$W_n(x_1, \dots, x_n) = \delta_{x_n} \cdots \delta_{x_3} W_2(x_1, x_2).$$
 (C.3)

On the other hand, Properties (iv) and (v) imply

$$\delta_{\eta_1} \cdots \delta_{\eta_n} M(x) = O(\hbar^n). \tag{C.4}$$

Then, since $W_2(x_1, x_2) = O(\hbar^0)$ is expressed by M (see (5.11)), we get the desired property (5.17).

In [2] and [28] explicit formulas are proposed for the definition of a suitable insertion operator through its action on the solution of the isomonodromy system:

$$\delta_{\eta}\Psi(x) = \left(\frac{M(\eta)}{x-\eta} + Q(\eta)\right)\Psi(x),\tag{C.5}$$

where $Q(\eta)$ is a matrix that depends on the Lax pair and is determined by imposing the above properties. Note that, the insertion operator (C.5) satisfies (C.1) for any choice of $Q(\eta)$. Property (iv) requires

$$\delta_{\eta} M(x) = -\frac{[M(x), M(\eta)]}{x - \eta} + [Q(\eta), M(x)] = O(\hbar).$$
(C.6)

Moreover, the condition (v) implies $[\delta_{\eta}, \partial_t]\Psi(x) = 0$; namely,

$$\delta_{\eta} \mathcal{R}(x) = \hbar \partial_t Q(\eta) + [Q(\eta), \mathcal{R}(x)] + \left[M(\eta), \frac{\mathcal{R}(x) - \mathcal{R}(\eta)}{x - \eta} \right].$$
(C.7)

KOHEI IWAKI AND OLIVIER MARCHAL

Condition (C.7) almost determines the action of δ_n , as we explain below.

Now let us consider the JM Lax pair to explain how the insertion operator is specified. First, look at the equality (C.7). The l.h.s. given by

$$\delta_{\eta} \mathcal{R}(x) = \begin{pmatrix} \frac{1}{2} \delta_{\eta} q & 0\\ -\delta_{\eta} p & -\frac{1}{2} \delta_{\eta} q \end{pmatrix}$$

is independent of x. (The action of δ_{η} on q and p will be defined later so that (C.7) is satisfied; see (C.10) and (C.11) below.) Since $\mathcal{R}(x)$ is linear in x, the only x-depending term in the r.h.s. of (C.7) is

$$\begin{bmatrix} Q(\eta), \mathcal{R}(x) \end{bmatrix} = \begin{pmatrix} -p Q(\eta)_{1,2} - \frac{1}{2}Q(\eta)_{2,1} & \frac{1}{2}(Q(\eta)_{1,1} - Q(\eta)_{2,2}) - (x+q)Q(\eta)_{1,2} \\ p (Q(\eta)_{1,1} - Q(\eta)_{2,2}) - (x+q)Q(\eta)_{2,1} & p Q(\eta)_{1,2} + \frac{1}{2}Q(\eta)_{2,1} \end{pmatrix}$$

Therefore, the *x*-independence requires

$$Q(\eta)_{1,2} = Q(\eta)_{2,1} = 0, \tag{C.8}$$

and (C.7) is reduced to the following set of equalities:

$$2M(\eta)_{1,2} = Q(\eta)_{1,1} - Q(\eta)_{1,2}, \qquad (C.9)$$

$$\delta_{\eta}q = 2\hbar Q(\eta)_{1,1},$$
 (C.10)

$$\delta_{\eta} p = -2\hbar \partial_t M(\eta)_{1,1}. \tag{C.11}$$

Here we have used the following equalities (Cf. (B.3)):

$$\hbar \partial_t M(x)_{1,1} = \frac{1}{2} M(x)_{2,1} + p M(x)_{1,2}, \hbar \partial_t M(x)_{1,2} = (x+q) M(x)_{1,2} - M(x)_{1,1} + \frac{1}{2}, \hbar \partial_t M(x)_{2,1} = -2p M(x)_{1,1} - (x+q) M(x)_{2,1} + p, \hbar \partial_t M(x)_{2,2} = -\hbar \partial_t M(x)_{1,1}.$$
(C.12)

Thanks to (C.12), if we take $Q(\eta)$ satisfying (C.8) and (C.9), straightforward computations shows that:

$$\delta_{\eta} M(x)_{1,1} = -\frac{2\hbar}{x-\eta} \left(M(x)_{1,2} \partial_{t} M(\eta)_{1,1} - M(\eta)_{1,2} \partial_{t} M(x)_{1,1} \right), \\\delta_{\eta} M(x)_{1,2} = -\frac{2\hbar}{x-\eta} \left(M(x)_{1,2} \partial_{t} M(\eta)_{1,2} - M(\eta)_{1,2} \partial_{t} M(x)_{1,2} \right), \\\delta_{\eta} M(x)_{2,1} = \frac{\hbar}{p(x-\eta)} \left(M(x)_{2,1} \partial_{t} M(\eta)_{2,1} - M(\eta)_{2,1} \partial_{t} M(x)_{2,1} \right) \\-\frac{2\hbar}{p} M(x)_{2,1} \partial_{t} M(\eta)_{1,1}, \\\delta_{\eta} M(x)_{2,2} = -\delta_{\eta} M(x)_{1,1}.$$
(C.13)

36

which are obviously of order $O(\hbar)$. As a conclusion, in our JM case we can take:

$$Q(\eta) = M(\eta)_{1,2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$
 (C.14)

$$\delta_{\eta}q = 2\hbar\partial_t M(\eta)_{1,2}, \quad \delta_{\eta}p = -2\hbar\partial_t M(\eta)_{1,1}. \tag{C.15}$$

Then, the important conditions (C.6) and (C.7) are satisfied.

From (C.13) it is tempting to conclude that the insertion operator δ_{η} satisfies (C.4). However (C.13) is not sufficient to prove (C.4) due to the following reason. Since the r.h.s. of (C.13) involves time derivatives of M(x), iterative application of the insertion operators creates terms of the form $\delta_{\eta}\partial_t M(x)$, $\delta_{\eta}\partial_t^2 M(x)$ and so on. The problem is that the condition (C.7) is not enough to prove $[\delta_{\eta}, \partial_t] = 0$ as operators acting on PV ring. For example, the condition (C.7) is not enough to prove $[\delta_{\eta}, \partial_t^2]\Psi(x) = 0$. To have this identity, we also need to require

$$[\delta_{\eta}, \partial_t] \mathcal{R}(x) = 0, \tag{C.16}$$

and previous works never checked this identity. In the JM case, this condition is equivalent to $[\delta_{\eta}, \partial_t]q = [\delta_{\eta}, \partial_t]p = 0$. Since $\hbar \partial_t q = p + q^2 + \frac{t}{2}$, straightforward computation shows

$$\begin{split} \hbar[\delta_{\eta},\partial_{t}]q &= \delta_{\eta}p + 2q\,\delta_{\eta}q + \frac{1}{2}\delta_{\eta}t - \hbar\partial_{t}\delta_{\eta}q \\ &= -2\hbar\partial_{t}M(\eta)_{1,1} + 4q\,\hbar\partial_{t}M(\eta)_{1,2} + \frac{1}{2}\delta_{\eta}t - 2\hbar^{2}\partial_{t}^{2}M(\eta)_{1,2}. \end{split}$$

Thus we need to further require

$$\delta_{\eta}t = 4\hbar\partial_t M(\eta)_{1,1} - 8q\,\hbar\partial_t M(\eta)_{1,2} + 4\hbar^2\partial_t^2 M(\eta)_{1,2} \tag{C.17}$$

to have $[\delta_{\eta}, \partial_t]q = 0$. The equality (C.17) contradicts to Condition (v) since the r.h.s. of (C.17) depends on t. Indeed, it follows from (B.7) that the first term of its \hbar -expansion is given by

$$\delta_{\eta}t = \hbar \Big(4\partial_t M^{(0)}(\eta)_{1,2} - 8q_0 \partial_t M^{(0)}(\eta)_{1,1} \Big) + O(\hbar^2)$$

= $-\hbar \frac{\eta + q_0}{\left((\eta + q_0)^2 + \frac{\theta}{q_0}\right)^{3/2}} + O(\hbar^2).$

This implies that $[\delta_{\eta}, \partial_t]t \neq 0$ which proves that the insertion operator δ_{η} does not satisfy Condition (v). Consequently this method does not prove (C.4) and the leading order property of W_n .

Unfortunately this problem is not specific to the JM Lax pair and is really intrinsic to the current method of the insertion operator. For example it also arises in the HTW Lax pair as well. We could not find any simple way to fix the problem and it is likely that substantial modifications of the insertion operator are required. However since the insertion operator exists in the context of random matrix models, we believe that it should exist in the context of determinantal formulas too.

Appendix D. Tau-function and symplectic invariants

Here we describe a relation between the tau-function (5.14) and the (generating function of) symplectic invariants, following the idea of [2, 7].

Theorem D.1 (Theorem 5.1 of [17]). For $g \ge 1$, both $F^{(g)} = F_{\rm JM}^{(g)}$ and $F_{\rm HTW}^{(g)}$ satisfy

$$\frac{dF^{(g)}}{dt} = 2 \operatorname{Res}_{x=\infty} \left(\frac{\partial s_{\infty}}{\partial t}(x) W_1^{(g)}(x) dx \right).$$
(D.1)

Here $W_1(x)$ and $s_{\infty}(x)$ are given in (5.8) and (5.15), respectively.

Proof. The fact that the JM and HTW Lax pair are of topological type and the result of Proposition 5.3 (i) imply

$$W_1^{(g)}(x(z))dx(z) = \omega_1^{(g)}(z),$$
 (D.2)

where x(z) appears in the parametrization (3.22) or (4.9) of the spectral curve, and $\omega_1^{(g)}(z)$ is the Eynard-Orantin differential of type (g, 1). On the other hand, the function

$$\Lambda(z) = \frac{\partial s_{\infty}}{\partial t}(x(z))$$

satisfies

$$\frac{\partial x}{\partial t}(z)dy(z) - \frac{\partial y}{\partial t}(z)dx(z)
= \begin{cases}
\operatorname{Res}_{w=\infty} \Lambda(w)\omega_2^{(0)}(w,z) - \operatorname{Res}_{w=0} \Lambda(w)\omega_2^{(0)}(w,z) & \text{for JM case,} \\
\operatorname{Res}_{w=\infty} \Lambda(w)\omega_2^{(0)}(w,z) & \text{for HTW case.}
\end{cases}$$
(D.3)

This is the required condition for $\Lambda(z)$ to apply Theorem 5.1 of [17] which proves (D.1). This completes the proof of Theorem 3.3 and Theorem 4.1.

Appendix E. Constant term in (1.2) and the symplectic invariance

The symplectic transformation (4.26) from JM curve to HTW curve is obtained from the composition of following three symplectic transformations:

$$(x_1, y_1) = (x_{\text{JM}}, y_{\text{JM}}) \mapsto (x_2, y_2) \text{ by } x_2 = x_1, \ y_2 = y_1 + x_1^2 + \frac{\iota}{2}$$
 (E.1)

$$(x_2, y_2) \mapsto (x_3, y_3)$$
 by $x_3 = y_2, y_3 = -x_2$ (E.2)

$$(x_3, y_3) \mapsto (x_4, y_4) = (x_{\text{HTW}}, y_{\text{HTW}}) \text{ by } x_4 = x_3, \ y_4 = y_3 + \frac{\theta}{2x_3}.$$
 (E.3)

Since the symplectic transformations (E.1) and (E.3) preserve the branch points and the recursion kernel $K(z_0, z)$, we can conclude that the free energy $F^{(g)}$ is invariant (Cf. Theorem 7.1 in [17]). On the other hand, the second symplectic transformation (E.2) does not preserve $F^{(g)}$ in general as shown in [11, 18], and Theorem 3.1 in [18] proves that an integral of $\omega_1^{(g)}$ appears as their difference. In our case Theorem 3.1 in [18] shows

$$F_{\rm JM}^{(g)} - \frac{\theta}{2 - 2g} \int_0^\infty \omega_{\rm JM,1}^{(g)} = F_{\rm HTW}^{(g)} + \frac{\theta}{2(2 - 2g)} \int_{-1}^1 \omega_{\rm HTW,1}^{(g)} \quad \text{for } g \ge 2.$$
(E.4)

This is equivalent to say that:

$$F_{\rm JM}^{(g)} - F_{\rm HTW}^{(g)} = \frac{\theta}{2 - 2g} \left(\int_0^\infty \omega_{\rm JM,1}^{(g)} + \frac{1}{2} \int_{-1}^1 \omega_{\rm HTW,1}^{(g)} \right) \quad \text{for } g \ge 2.$$
 (E.5)

This relation for $g \in \{2,3\}$ can be directly tested from our computations. In fact from our work in Section 3.6 and Section 4.4, we obtained the exact value of the r.h.s. of (E.5) (Cf. (4.27)). Moreover, it is straightforward to observe that $\omega_{\text{JM},1}^{(g)} \xrightarrow{\to} 0$ and $\omega_{\text{HTW},1}^{(g)} \xrightarrow{\to} 0$ for all $g \geq 2$. Therefore, we obtain:

$$\int_{0}^{\infty} \omega_{\text{Weber},1}^{(g)} = \frac{B_{2g}}{2g \,\theta^{2g-1}} \text{ and } \int_{-1}^{1} \omega_{\text{Bessel},1}^{(g)} = \frac{B_{2g}}{g \,\theta^{2g-1}} \quad \text{for } g \ge 2.$$
(E.6)

In both cases, the choice of integration contour does not matter since the differential forms do not have any residue. The only requirement is that the contours avoid $\{\pm 1\}$ in the Hermite-Weber case and $\{0\}$ in the Bessel case.

Acknowledgment

The authors would like to thank Gaëtan Borot, Bertrand Eynard, Alba Grassi, Masahide Manabe, Motohico Mulase, Yasuhiko Yamada for fruitful discussion. O. Marchal would like to thank Université Lyon 1 and particularly Université Jean Monnet and Institut Camille Jordan for the opportunity to make this research possible. O. Marchal would also like to thank his family and friends for moral support during the preparation of this article. K. Iwaki is supported by JSPS KAKENHI Grant Number 13J02831, 16K17613 and 16H06337. O. Marchal is supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). The authors also grateful to Springer International Publishing. The final publication is available at Springer via http://dx.doi.org/10.1007/s00023-017-0576-z.

References

- J. Ambjorn and L. Chekhov, "The matrix model for dessins d'enfants", arxiv:1404.4240, 2014.
- [2] M. Bergère, G. Borot and B. Eynard, "Rational differential systems, loop equations, and application to the qth reductions of KP", Annales de l'institut Henri Poincaré, DOI: 10.1007/s00023-014-0391-8, 2015.
- [3] M. Bergère and B. Eynard, "Determinantal formulae and loop equations", arxiv:0901.3273, 2009.

- [4] M. Bergère and B. Eynard, "Universal scaling limits of matrix models, and (p,q) Liouville gravity", arxiv:0909.0854, 2009.
- [5] M. Bertola, B. Eynard and J. Harnad, "Partition functions for matrix models and isomonodromic tau functions", *Journal of Physics A*, Vol. 36, 3067-3083, 2003.
- [6] M. Bertola and O. Marchal, "The partition function of the two-matrix model as an isomonodromic tau-function", Journal of Mathematical Physics, Vol. 50, 013529, 2009.
- [7] G. Borot and B. Eynard, "Tracy-Widom GUE law and symplectic invariants", arXiv:1011.1418, 2010.
- [8] G. Borot and B. Eynard, "The asymptotic expansion of Tracy-Widom GUE law and symplectic invariants", arXiv:1012.2752, 2010.
- [9] G. Borot, A. Guionnet, "Asymptotic expansion of β matrix models in the one-cut regime", Communications in Mathematical Physics, Vol. **317**, Issue 2, 447-483, 2013.
- [10] V. Bouchard, A. Catuneanu, O. Marchal and P. Sułkowski, "The Remodeling Conjecture and the Faber-Pandharipande Formula", *Letters in Mathematical Physics*, Vol. 103, Issue 1, 59-77, 2013.
- [11] V. Bouchard and P. Sułkowski, "Topological recursion and mirror curves", Advances in Theoretical and Mathematical Physics 16 (2012), 1443–1483.
- [12] L. Chekhov, B. Eynard and N. Orantin, "Free energy topological expansion for the 2-matrix model", Journal of High Energy Physics, JHEP12(2006)053.
- [13] O. Dumitrescu and M. Mulase, "Lectures on the topological recursion for Higgs bundles and quantum curves", arXiv:1509.09007, 2015.
- [14] O. Dumitrescu, M. Mulase, B. Safnuk and A. Sorkin, "The spectral curve of the Eynard-Orantin recursion via the Laplace transform", *Contemporary Mathematics*, Vol. 593, 2013.
- [15] B. Eynard, "Intersection numbers of spectral curves", arxiv:1104.0176, 2011.
- [16] B. Eynard, "Counting Surfaces", CRM Aisenstadt Chair lectures, Progress in Mathematical Physics, Vol. 70, Springer, 2016.
- [17] B. Eynard and N. Orantin, "Invariants of algebraic curves and topological expansion", Communications in Number Theory and Physics, Vol. 1, 347-452, 2007.
- [18] B. Eynard and N. Orantin, "About the x-y symmetry of the F_g algebraic invariants", arXiv:1311.4993, 2013.
- [19] J. Harer and D. Zagier, "The Euler characteristic of the moduli space of curves", *Inventiones mathematicae*, Vol. 85, 457-485, 1986.
- [20] J. Harnad, C. A. Tracy and H. Widom, "Hamiltonian structure of equations appearing in random matrices", in Low-Dimensional Topology and Quantum Field Theory, ed. H. Osborn, NATO ASI Series B, Vol. **314**, Plenum Press, New York, pp. 231-245, 1993.
- [21] K. Iwaki, O. Marchal and A. Saenz, "Painlevé equations, topological type property and reconstruction by the topological recursion", arXiv:1601.02517.
- [22] K. Iwaki, T. Koike and Y. Takei, in preparation.
- [23] M. Jimbo, T. Miwa and K. Ueno, "Monodromy preserving deformation of linear ordinary differential equations with rational coefficients I. General theory and τ -function", *Physica 2D*, Vol. **2**, 306-352, 1981.
- [24] M. Jimbo and T. Miwa, "Monodromy preserving deformation of linear ordinary differential equations with rational coefficients II", *Physica 2D*, Vol. 2, 407-448, 1981.
- [25] N. Joshi, A. V. Kitaev and P. A. Treharne, "On the linearization of the first and second Painlevé equations", *Journal of Physics A: Mathematical and Theoretical*, Vol. 42, 055208, 2009.
- [26] T. Kawai and Y. Takei, "Algebraic Analysis of Singular Perturbation", Iwanami Series in Modern Mathematics, Translations of Mathematical Monographs, Vol. 227, 2005.

- [27] O. Marchal and M. Cafasso, "Double-scaling limits of random matrices and minimal (2m, 1) models: the merging of two cuts in a degenerate case", *Journal of Statistical Mechanics*, P04013, 2011.
- [28] O. Marchal, B. Eynard and M. Bergère, "The sine-law gap probability, Painlevé 5, and asymptotic expansion by the topological recursion", *Random Matrices: Theory and Applications*, Vol. 3, 1450013 (2014)
- [29] M. Mariño, "Les Houches lectures on matrix models and topological strings", arxiv:hep-th/0410165.
- [30] M. L. Mehta, "Random matrices", Pure and applied mathematics, Vol. 142, Elsevier academic press, Amsterdam, third edition, 2004.
- [31] M. Mulase and M. Penkava, "Topological recursion for the Poincaré polynomial of the combinatorial moduli space of curves", Advances in Mathematics, Vol. 230, 1322–1339, 2012.
- [32] R. C. Penner, "Perturbative series and the moduli space of Riemann surfaces". Journal of Differential Geometry, Vol. 27, 35–53, 1988.
- [33] C. Tracy and H. Widom, "Introduction to Random Matrices", Springer Lecture Notes in Physics, Vol. 424, 103-130, 1993.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, JAPAN *E-mail address*: iwaki@math.nagoya-u.ac.jp

UNIVERSITÉ DE LYON, CNRS UMR 5208, UNIVERSITÉ JEAN MONNET, INSTITUT CAMILLE JORDAN, FRANCE

E-mail address: olivier.marchal@univ-st-etienne.fr