

Saha Equation in Rindler Space

Sanchari De^a and Somenath Chakrabarty^{a†}^aDepartment of Physics, Visva-Bharati, Santiniketan-731235, India[†]E-mail: somenath.chakrabarty@visva-bharati.ac.in

Abstract

The Saha equations for photo-ionization process of hydrogen atoms and the creation of electron-positron pairs at high temperature are investigated in a reference frame undergoing a uniform accelerated motion in an otherwise flat Minkowski space-time geometry or equivalently, in a rest frame in presence of a strong uniform gravitational field. It is known as the Rindler space.

Keywords: Saha equation, Uniformly accelerated frame, Rindler coordinates, Photo ionization, Pair production

PACS:03.65.Ge,03.65.Pm,03.30.+p,04.20.-q

1 Introduction

The well known Lorentz transformations are the space-time coordinate transformation between two inertial frame of references [1]. However, with the help of the principle of equivalence, one may obtain the space-time transformations between a uniformly accelerated frame and an inertial frame and vice-versa in the same manner as it is done in special theory of relativity [2, 3, 4, 5]. In the present scenario the flat geometry is called the Rindler space. For an illustration, let us consider two reference frames- the frame S is an inertial one, which is at rest but there is a uniform gravitational field and the frame S' , which is undergoing a uniform accelerated motion with respect to some inertial frame, but in absence of gravitational field. Then in accordance with the principle of equivalence these two frames are physically equivalent. Alternatively, one may state, that a reference frame in uniform acceleration in absence of gravitational field is equivalent to a frame at rest in presence of a uniform gravitational field. We may assume that the strong gravitational field is produced by a black hole like gravitating object. Therefore the gravitational field may be approximated by a constant value within

a small region at a distance x from the centre of the gravitating object. This is also called the local acceleration of the frame.

To investigate Saha equation in a uniformly accelerated frame of reference or in Rindler space, we first develop a formalism with the physical concepts of principle of equivalence to obtain the elements of the metric tensor $g^{\mu\nu}$. We shall show that analogous to the Minkowski space-time metric tensor, the off-diagonal elements of $g^{\mu\nu}$ are also zero in Rindler space. In the next step, with the conventional form of action as has been defined in special theory of relativity [1], which is invariant here also, the Lagrangian of a particle (in our study, which may be a hydrogen atom or a hydrogen ion or an electron) is derived from Hamilton's principle. Which further gives the momentum and energy or Hamiltonian of the particle from the standard relations of classical mechanics. Then considering a partially ionized hydrogen plasma which is a reactive mixture of neutral hydrogen atoms, hydrogen ions, electrons and photons or considering an electron-positron plasma composed of an interacting mixture of electrons, positrons and photons, we shall obtain the modified form of Saha equations for both the cases when observed from a uniformly accelerated frame of reference or in Rindler space. To the best of our knowledge, the study of Saha equation in Rindler space has not been reported earlier. We shall also compare our findings with the conventional results.

The article is organized in the following manner: In the next section we shall establish some of the useful relations of special theory of relativity in uniformly accelerated frame. In this section we shall also obtain the relativistic version of Hamiltonian in Rindler coordinate system. In section-3 we shall investigate photo-ionization of hydrogen atoms in partially ionized hydrogen plasma in Rindler space. In section-4 we shall study the electron-positron pair production from intense electromagnetic radiation at high temperature in an interacting electron-positron plasma, observed from the same kind of uniformly accelerated frame. In the last section we give the conclusion of our findings.

2 Basic Formalism

For the sake of completeness, in this section, following [6, 7, 8] we shall establish some of the useful formulas of special theory of relativity for a uniformly accelerated frame of reference. Let us first assume that the frame S'

has rectilinear motion with uniform velocity v along x -direction with respect to some inertial frame S . Further the coordinates of an event occurred at the point P (say) is indicated by (x, y, z, t) in S -frame and with (x', y', z', t') in the frame S' . The primed and the un-primed coordinates are related by the conventional form of Lorentz transformations and are given by

$$\begin{aligned} x' &= \gamma(x - vt), \quad y' = y, \quad z' = z \text{ and} \\ t' &= \gamma \left(t - \frac{vx}{c^2} \right) \text{ with } \gamma = \left(1 - \frac{v^2}{c^2} \right)^{-1/2} \end{aligned} \quad (1)$$

is the Lorentz factor. Next we consider a uniformly accelerated frame S' . Now for the sake of simplicity, we assume that the acceleration of the test particle is also along x -direction in S -frame and is α_x . Then in the primed frame it is given by

$$\alpha'_x = \alpha_x \gamma^{-3} \left(1 - \frac{vu_x}{c^2} \right)^{-3}, \quad \text{where } \alpha_x = \frac{du_x}{dt} \quad (2)$$

Therefore for this particle having constant acceleration in S' -frame, in which it was instantaneously in coincidence with its origin at some earlier time, we have

$$\alpha'_x = g \text{ (say)} = \gamma^3 \alpha_x \quad (3)$$

Here it has further been assumed that the motion of the particle is also rectilinear and is along x -direction with $u_x = v$. Then with the initial condition that $u_x = 0$ at $t = 0$, we have

$$u_x = gt \left[1 + \left(\frac{gt}{c} \right)^2 \right]^{-1/2} \quad (4)$$

Therefore the proper time of the particle is given by

$$\tau = \frac{c}{g} \sinh^{-1} \left(\frac{gt}{c} \right) \quad (5)$$

Hence

$$ct = \frac{c^2}{g} \sinh \left(\frac{g\tau}{c} \right) \quad (6)$$

Then from the expression $u_x = dx/dt$, the velocity of the particle along x -direction and using eqn.(4), we have

$$x = \frac{c^2}{g} \cosh \left(\frac{g\tau}{c} \right) \quad (7)$$

Eqns.(6) and (7) are the parametric form of the equation for the world line of the particle undergoing constant accelerated motion. Hence the hyperbolic motion of the particle can be represented by the equation

$$x^2 - (ct)^2 = \left(\frac{c^2}{g}\right)^2 \quad (8)$$

To obtain the Rindler coordinates, which are used for a system / observer undergoing constant proper acceleration in a Minkowski flat space-time, we have followed the scheme described below. Following [6, 7] (see also [9]), we have constructed a vector which is tangential with the world line and is proportional to

$$\left(\cosh\left(\frac{g\tau}{c}\right), \sinh\left(\frac{g\tau}{c}\right)\right) \quad (9)$$

These are again the parametric form of coordinates of the points moving along the proper time axis of the uniformly accelerated observer. Next consider a vector which is orthogonal to above vector. The parametric form of which is proportional to

$$\left(\sinh\left(\frac{g\tau}{c}\right), \cosh\left(\frac{g\tau}{c}\right)\right) \quad (10)$$

Then the parametric form of the world line of a second observer, which is h distance ahead of the first one as measured by the first observer is given by

$$\begin{aligned} ct &= \left(\frac{c^2}{g} + h\right) \sinh\left(\frac{g\tau}{c}\right) \quad \text{and} \\ x &= \left(\frac{c^2}{g} + h\right) \cosh\left(\frac{g\tau}{c}\right) \end{aligned} \quad (11)$$

For the second accelerated observer, the hyperbolic motion is described by

$$x^2 - (ct)^2 = \left(\frac{c^2}{g} + h\right)^2 = \left(\frac{c^2}{g}\right)^2 \left(1 + \frac{gh}{c^2}\right)^2 \quad (12)$$

The magnitude of constant acceleration for the second observer is therefore

$$g' = g \left(1 + \frac{gh}{c^2}\right)^{-1} \quad (13)$$

i.e., the two observers undergoing accelerated motions with different magnitudes of acceleration.

To obtain the Rindler coordinates, we consider an event with space time coordinate ct' and x' as measured by the first observer. Further it has been assumed that x' is the distance from the origin of the first observer and t' is the corresponding proper-time. Then with respect to an inertial frame, we have from eqns.(9)-(11) and considering simultaneity of these events

$$\begin{aligned} ct &= \left(\frac{c^2}{g} + x'\right) \sinh\left(\frac{gt'}{c}\right) \quad \text{and} \\ x &= \left(\frac{c^2}{g} + x'\right) \cosh\left(\frac{gt'}{c}\right) \end{aligned} \quad (14)$$

Hence one can also express the inverse relations

$$ct' = \frac{c^2}{2g} \ln\left(\frac{x+ct}{x-ct}\right) \quad \text{and} \quad x' = (x^2 - (ct)^2)^{1/2} - \frac{c^2}{g} \quad (15)$$

The Rindler space-time coordinates, given by eqns.(14) and (15) are then just an accelerated frame transformation of the Minkowski metric of special relativity. The Rindler coordinate transformations change the Minkowski line element from

$$\begin{aligned} ds^2 &= d(ct)^2 - dx^2 - dy^2 - dz^2 \quad \text{to} \\ ds^2 &= \left(1 + \frac{gx'}{c^2}\right)^2 d(ct')^2 - dx'^2 - dy'^2 - dz'^2 \end{aligned} \quad (16)$$

Since the motion is assumed to be rectilinear and along x -direction, $dy' = dy$ and $dz' = dz$. The form of the metric tensor can then be written as

$$g^{\mu\nu} = \text{diag}\left(\left(1 + \frac{gx}{c^2}\right)^2, -1, -1, -1\right) \quad (17)$$

Now following the concept of dynamics of special theory of relativity [1], the action integral may be written as (see also [10])

$$S = -\alpha_0 \int_a^b ds \equiv \int_a^b L dt \quad (18)$$

Then using eqns.(16) and (18) and putting $\alpha_0 = -m_0c$, where m_0 is the rest mass of the particle, the Lagrangian of the particle is given by [11]

$$L = -m_0c^2 \left[\left(1 + \frac{gx}{c^2}\right)^2 - \frac{v^2}{c^2} \right] \quad (19)$$

where \vec{v} is the velocity of the particle. The momentum of the particle is then given by

$$\vec{p} = m_0 \vec{v} \left[\left(1 + \frac{gx}{c^2} \right)^2 - \frac{v^2}{c^2} \right]^{-1/2} \quad (20)$$

Hence the Hamiltonian of the particle or the single particle energy is given by

$$H = \varepsilon_p = m_0 c^2 \left(1 + \frac{gx}{c^2} \right) \left(1 + \frac{p^2}{m_0^2 c^2} \right)^{1/2} \quad (21)$$

In the new sections, using this simple form of single particle energy, we shall obtain the Saha equations for photo-ionization process of hydrogen atoms and also for electron-positron pair creation. We shall follow the standard formalism as discussed in the text books on astrophysics and statistical mechanics [12, 13].

3 Photo-Ionization of Hydrogen Atoms

To study the Saha ionization process for hydrogen atoms, we consider a partially ionized hydrogen plasma, consisting of neutral hydrogen atoms, hydrogen ions, electrons and electromagnetic radiation or photons. For the sake of mathematical simplicity, the system is assumed to have cylindrical geometry and the plasma is expanding with a constant acceleration g along the positive x -direction, which is also the symmetry axis of the cylinder. Then according to the principle of equivalence, any ionization or de-ionization of the accelerated particles taking place at some point is equivalent to their occurrence at rest frame in presence of a gravitational field g . Now it is well known that depending on the amount of energy transferred, the interaction with photon will either excite the hydrogen atoms or ionize them. The electrons can also be captured by the hydrogen ions, which is the de-ionization process. Ultimately a dynamic dead lock situation will be reached, when the rate of ionization process and the rate of de-ionization process become just equal, which may be represented by the reaction equation



where H_n indicates the n th excited state of hydrogen atom with $n = 1$ the ground state. Under such physical condition the partially ionized hydrogen plasma is assumed to be in thermodynamic equilibrium with all of its

constituents. In the equilibrium condition the chemical potentials of the components are related by the equation

$$\mu(H_n) = \mu(H^+) + \mu(e) \quad (23)$$

Since the number of photons is not conserved, $\mu(\gamma) = 0$.

Now in the non-relativistic situation, the single particle energy as shown in eqn.(21) may be approximated as

$$\varepsilon_p = m_0 c^2 \left(1 + \frac{gx}{c^2}\right) + \left(1 + \frac{gx}{c^2}\right) \frac{p^2}{2m_0} \quad (24)$$

Then defining

$$m' = m_0 \left(1 + \frac{gx}{c^2}\right)^{-1} \quad \text{and} \quad m'' = m_0 \left(1 + \frac{gx}{c^2}\right) \quad (25)$$

we can write

$$\varepsilon_p = m'' c^2 + \frac{p^2}{2m'}$$

as the effective single particle energy or Hamiltonian of the particle. Now it can very easily be shown from the standard results of text books on statistical mechanics [14, 13] that the number density of a particular component (except for photon) is given by

$$\begin{aligned} n &= \frac{N}{\Delta V} \\ &= \frac{4\pi g_d}{h^3} \int_0^\infty p^2 dp \exp \left[-\frac{1}{kT} \left(\frac{p^2}{2m'} + m'' c^2 - \mu \right) \right] \end{aligned} \quad (26)$$

where g_d is the degeneracy of the species, $\Delta V = A\Delta x$ is a small volume element, Δx is a small length element in the x -direction at a distance x from the origin and A is the cross-sectional area of the cylinder. The length Δx is such that g is constant within ΔV . On evaluating the above integral, we have

$$n = n_Q g_d \exp \left[\frac{1}{kT} (\mu - m'' c^2) \right] \quad (27)$$

Hence

$$\mu = m'' c^2 - kT \ln \left(\frac{g_d n_Q}{n} \right) \quad (28)$$

This is the general expression for chemical potential for a particular species in terms of its concentration. Further,

$$n_Q = \left(\frac{2\pi m' kT}{h^2} \right)^{3/2} \quad (29)$$

is called the quantum concentration for the particular species in the mixture. Then using eqn.(23) for chemical equilibrium condition and eqn.(28), the expression for the chemical potential, we have

$$\frac{n(H^+)n(e)}{n(H_n)} = \frac{n_{Q_e}}{g_n} \exp\left(-\frac{\Delta E_n}{kT}\right) = R_{g>0} \text{ (say)} \quad (30)$$

where $n(i)$ is the number density for the species i and $g_n = n^2$ is the degeneracy of the neutral hydrogen atom in the n th excited state (since the degeneracy of electrons has been considered separately in the derivation, the factor 2 does not appear in the expression for g_n) and

$$\begin{aligned} \Delta E_n &= \Delta m'' c^2 = \left(1 + \frac{gx}{c^2}\right) \Delta m c^2, \text{ with} \\ \Delta m &= m(H_n) - m(H^+) - m(e) \end{aligned} \quad (31)$$

is a measure of excitation energy in the present scenario.

Eqn.(30) gives the ratio of equilibrium concentration at a given temperature as measured in an accelerated frame or in other words in a local rest frame in presence of a constant gravitational field g . Whereas the conventional form of Saha equation, i.e., in an inertial frame with $g = 0$ is given by

$$\frac{n(H^+)n(e)}{n(H_n)} = \frac{n_{Q_e g=0}}{g_n} \exp\left(-\frac{\Delta m c^2}{kT}\right) = R_{g=0} \text{ (say)} \quad (32)$$

Hence the ratio with $g > 0$ and $g = 0$ can be written as

$$\frac{R_{g>0}}{R_{g=0}} = \left(1 + \frac{gx}{c^2}\right)^{-3/2} \exp\left(-\frac{gx\Delta m}{kT}\right) \quad (33)$$

4 Pair Production at High Temperature

We next consider the electron-positron pair creation at high temperature in a local rest frame in presence of a constant gravitational field g . The

temperature is assumed to be such that kT is larger than at least two times the electron rest mass. The pair production process can be represented by the reaction equation

$$\gamma + \gamma \leftrightarrow e^- + e^+ \quad (34)$$

Similar to the photo-ionization of hydrogen atoms, we assume that the interacting electron-positron plasma along with the photons is in thermodynamic equilibrium. In this case the chemical equilibrium condition is given by

$$\mu(e^+) + \mu(e^-) = 0 \quad (35)$$

where $\mu(\gamma) = 0$ and

$$\begin{aligned} \mu(e^-) &= m''(e)c^2 - kT \ln \left(\frac{g_e n_Q}{n_{e^-}} \right) \text{ and} \\ \mu(e^+) &= m''(e)c^2 - kT \ln \left(\frac{g_e n_Q}{n_{e^+}} \right) \end{aligned} \quad (36)$$

Then from eqn.(35), we have

$$n_{e^-} n_{e^+} = g_e^2 n_Q^2 \exp \left(-\frac{2m''(e)c^2}{kT} \right) \quad (37)$$

where

$$n_Q = \left(\frac{2\pi m'(e)kT}{h^2} \right)^{3/2} \quad (38)$$

is the quantum concentration for electron and positron. Here $m'(e)$ and $m''(e)$ are given by eqn.(25) with m_0 replaced by $m(e)$, the electron rest mass.

If the system is assumed to be charge neutral then $n_{e^-} = n_{e^+}$. Further eqn.(37) can also be expressed as a ratio of the product of electron-positron concentration with $g > 0$ and $g = 0$, which is given by

$$\frac{n_{e^-} n_{e^+} g > 0}{n_{e^-} n_{e^+} g = 0} = \left(1 + \frac{gx}{c^2} \right)^{-3/2} \exp \left(-\frac{2gxm(e)}{kT} \right) \quad (39)$$

5 Conclusion

To analyze the findings of this article, let us consider a set of uniformly accelerated frames in the space outside the event horizon of the black hole of

mass M . Then in accordance with the principle of equivalence, each of the frames are characterized by a local acceleration g . The position of the frames in Rindler space are indicated by the spatial coordinates x_l , measured from the centre of the black hole. The values for g and x_l are obviously different for different uniformly accelerated frames. The reference frames are distributed uniformly in the space, so that some of them are at a very close proximity with the event horizon, while some others are far away from the black hole surface.

Now consider a particular frame which is far away from the event horizon, i.e., $x_l \gg R_s$, where $R_s = 2GM/c^2$, the Schwarzschild radius of the black hole. Then the acceleration g of the frame, or equivalently the gravitational field of the frame at rest may be expressed in Newtonian form $g = GM/x_l^2$. Then it can very easily be shown that the factor as shown below may be expressed as

$$1 + \frac{gx_l}{c^2} = 1 + \frac{R_s}{2x_l}$$

Therefore for $x_l \gg R_s$, one can very easily verify from eqn.(25) that $m' = m'' \approx$, the rest mass of the particle. Further, in such asymptotic condition, it can very easily be shown from eqns.(33) and (39) that $R_{g>0}/R_{g=0} \approx 1$ and $(n_{e^-} - n_{e^+})_{g>0}/(n_{e^-} - n_{e^+})_{g=0} \approx 1$ respectively.

On the other hand, if we consider a frame very close to the surface of the black hole, or the spatial coordinate x_l for this frame is very close to R_s , then both the gravitational field and the temperature can be expressed in blue shifted forms [15] and are given by

$$g(x_l) = \frac{\frac{GM}{x_l^2}}{\left(1 - \frac{R_s}{x_l}\right)^{1/2}} \quad \text{and} \quad T(x_l) = \frac{T(\infty)}{\left(1 - \frac{R_s}{x_l}\right)^{1/2}} \quad (40)$$

Hence we have

$$1 + \frac{gx_l}{c^2} = 1 + \frac{\frac{R_s}{2x_l}}{\left(1 - \frac{R_s}{x_l}\right)^{1/2}}$$

Obviously the gravitational field, temperature and the above factor diverges for $x_l \rightarrow R_s$. Further, in this limit $m' \rightarrow 0$ and $m'' \rightarrow \infty$. It is now a matter of little algebra to show that the quantum concentration for a particular species in this frame is given by

$$n_Q = \left(\frac{2\pi kT(\infty)}{h^2}\right)^{3/2} \left[\left(1 - \frac{R_s}{x_l}\right)^{1/2} + \frac{R_s}{2x_l}\right]^{-3/2} \quad (41)$$

Hence for $x_l \rightarrow R_s$, we have

$$n_Q = \left(\frac{4\pi kT(\infty)}{h^2} \right)^{3/2} \quad (42)$$

i.e., the blue shifted nature of temperature and gravitational field counter balance each other.

In this case the arguments of the exponential terms for photo-ionization of hydrogen and for $e^- - e^+$ pair creation as shown by the eqns.(33) and (39) are given by

$$A1 = \frac{R_s}{x_l} \frac{\Delta mc^2}{2kT(\infty)} \quad \text{and} \quad A2 = \frac{R_s}{x_l} \frac{m(e)c^2}{2kT(\infty)}$$

respectively. In these quantities also the divergence of g and that of T cancel each other in the blue shifted region. Then in the limiting case $x_l \rightarrow R_s$, we have

$$A1 = \frac{\Delta mc^2}{2kT(\infty)} \quad \text{and} \quad A2 = \frac{m(e)c^2}{2kT(\infty)}$$

On the other hand, in such extreme condition the pre-exponential terms in both eqns.(33) and (39) become zero, making both the ratios equal to zero. Hence we may conclude that strong gravitational field suppresses photo-ionization of hydrogen and also $e^- - e^+$ pair creation in chemical equilibrium in a partially ionized hydrogen plasma or in an $e^- - e^+$ plasma respectively. Since the blue shifted nature of g and T are counter balanced, the exponential parts do not take important role. Whereas the blue shifted form of g in the pre-exponential terms in eqns.(33) and (39) makes these two quantities exactly zero in the limiting condition. Therefore photo-ionization of hydrogen is possible at the surface of a main-sequence star or a post main-sequence star or at the surface of a white dwarf. For these stellar objects the strength of gravitational field is moderate, whereas for a neutron stars, a very compact stellar objects with extremely strong surface gravitational field, it is not likely to have hydrogen ions at the vicinity of its surface.

References

- [1] L.D. Landau and E.M. Lifshitz, The Classical Theory of Fields, Butterworth-Heimenann, Oxford, (1975.)
- [2] S. Weinberg, Gravitation and Cosmology, Wiley, New York, (1972).

- [3] C.W. Misner, Kip S. Thorne and J.A. Wheeler, Gravitation, W.H. Freeman and Company, New York, (1972).
- [4] W. Rindler, Essential Relativity, Springer-Verlag, New York, (1977).
- [5] C. Moller, The Theory of Relativity, Calarendon Press, Oxford, (1972).
- [6] M. Socolovsky, Annales de la Foundation Louis de Broglie **39**, 1, (2014).
- [7] G.F. Torres del Castillo Torres and C.L. Perez Sanchez, Revista Mexican De Fisika, **52**, 70 (2006).
- [8] D. Percoco and V.N. Villaba, Class. Quantum Grav., **9**, 307, (1992).
- [9] N.D. Birrell and P.C.W. Davies, Quantum Field Theory in Curved Space, Cambridge University Press, Cambridge, (1982).
- [10] C-G Huang and J-R Sun, arXiv:gr-qc/0701078,(2007).
- [11] Domingo J. Louis-Martinez, Class Quantum Grav., **28**, 035004, (2011).
- [12] A.C. Phillips, The Physics of Stars, John Wiley & Sons, New York, (1996).
- [13] L.D. Landau and E.M. Lifshitz, Statistical Physics, Part-I, Butterworth-Heimenann, Oxford, (1980).
- [14] K. Huang, Statistical Mechanics, Wiley-Eastern Pvt. Ltd., New Delhi, (1975).
- [15] R.C. Tolman, Relativity, Thermodynamics and Cosmology, Claredon Press, Oxford, (1934).