Variance Reduced Multilevel Monte Carlo Path Simulation: going beyond the complexity ε^{-2}

Denis Belomestny and Tigran Nagapetyan

Abstract In this paper a novel modification of the Multilevel Monte Carlo approach (MLMC), allowing for further significant complexity reduction, is proposed. The idea of the modification is to use a specifically designed control variate in the first level of MLMC. We show that under a proper choice of the control variate, one can reduce the complexity order of the modified MLMC algorithm down to $\varepsilon^{-2+\delta}$ for any $\delta \in [0,1)$ with ε being the precision to be achieved. These theoretical results are illustrated by several numerical examples.

1 Introduction

The multilevel path simulation method introduced in Giles [3] has recently gained a lot of popularity as a complexity reduction tool. The main advantage of the MLMC methodology is that it can be straightforwardly applied to various situations and requires almost no prior knowledge on the path generating process. Any multilevel Monte Carlo (MLMC) algorithm uses a number of levels of resolution, l = 0, 1, ..., L, with l = 0 being the coarsest, and l = L being the finest. In the context of SDE path simulation problem on the interval [0, T], level 0 corresponds to one time step $\Delta_0 = T$, whereas the level L is related to 2^L uniform timesteps $\Delta_L = 2^{-L} \cdot T$.

The problem of interest in this paper is to estimate the quantity $a := E(f(X_T)) \in \mathbb{R}$ for a real-valued functional f with a given accuracy ε , where $(X_t)_{t \in [0,T]}$ is a stochastic process, which can't sampled directly. We compare different Monte Carlo algorithms with their cost error relations, where by the cost and the error of the

Tigran Nagapetyan

Denis Belomestny

Duisburg-Essen University, Duisburg, Germany

e-mail: belomestny@uni-due.de

Department of Statistics, University of Oxford 24–29 St Giles', Oxford OX1 3LB, England, e-mail: nagapetyan@stats.ox.ac.uk

Monte Carlo algorithm \mathscr{M} we understand $\operatorname{cost}(\mathscr{M}) = \operatorname{E}(\# \text{ operations and random number calls})$ and $\operatorname{error}(\mathscr{M}) = \left(\operatorname{E}(a - \mathscr{M})^2\right)^{1/2}$, respectively. We say that a sequence of the Monte Carlo algorithms \mathscr{M}_n with $\lim_{n \to \infty} \operatorname{error}(\mathscr{M}_n) = 0$ achieves the order of convergence $\gamma > 0$ if there exists c > 0 and $\eta \in \mathbb{R}$, such that

$$\forall n \in \mathbb{N} : \operatorname{cost}(\mathscr{M}_n) \leq c \cdot (\operatorname{error}(\mathscr{M}_n))^{-\gamma} \cdot (-\log \operatorname{error}(\mathscr{M}_n))^{\eta}$$

Note, that the smaller γ is, the better is the performance of the Monte Carlo algorithm. In our setting we can't sample X_T directly, but we can sample $X_{l,T}$, where we assume that

$$\lim_{\to\infty} \operatorname{cost}(f(X_{l,T})) = \infty, \ \operatorname{E} f(X_{l,T}) \to \operatorname{E} f(X_T).$$

The multilevel path simulation approach consists in first writing the expectation of the finest approximation $E[f(X_{L,T})]$ as a telescopic sum

$$\mathbf{E}[f(X_{L,T})] = \mathbf{E}[f(X_{0,T})] + \sum_{l=1}^{L} \mathbf{E}[f(X_{l,T}) - f(X_{l-1,T})]$$
(1)

and then applying Monte Carlo to estimate each expectation in this sum. One important prerequisite for MLMC to work is that $X_{l,T}$ and $X_{l-1,T}$ are coupled in some way and this can be achieved by using the same discretized trajectories of the underlying diffusion process to construct the consecutive approximations $X_{l,T}$ and $X_{l-1,T}$. The degree of coupling is usually measured in terms of the variance $Var[f(X_{l,T}) - f(X_{l-1,T})]$. It is shown in Giles [3], that under the conditions:

$$\left| \mathbf{E}[f(X_{L,T})] - \mathbf{E}[f(X_T)] \right| \le c_1 \Delta_L^{\alpha}, \quad \operatorname{Var}\left[f(X_{l,T}) - f(X_{l-1,T}) \right] \le c_2 \Delta_l^{\beta}, \qquad (2)$$

with some $\alpha \ge 1/2$, $\beta > 0$, $c_1 > 0$, $c_2 > 0$ and with cost of sampling $f(X_{l,T})$ bounded by $c_3 \cdot \Delta_l^{-1}$, the computational complexity of the resulting multilevel estimate needed to achieve the accuracy ε (in terms of RMSE) is proportional to

$$\mathscr{C} \asymp \begin{cases} \varepsilon^{-2}, & \beta > 1, \\ \varepsilon^{-2} \log^2(\varepsilon), & \beta = 1, \\ \varepsilon^{-2-(1-\beta)/\alpha}, & 0 < \beta < 1. \end{cases}$$
(3)

This is a significant improvement over the classical standard Monte Carlo approach, which has complexity $\varepsilon^{-2-1/\alpha}$. The above asymptotic estimates however show that reduction of complexity beyond the order ε^{-2} is not possible, doesn't matter how large is $\beta > 1$. Moreover, there is an issue about achieving $\beta > 1$ in higher dimensions with an implementable algorithm, which has bees successfully resolved in [2] under certain regularity assumptions. This fact motivates a question on existence of algorithms with complexity order of order $\varepsilon^{-2+\delta}$, where $\delta > 0$. This is our main concern, and here we propose a modification of the original MLMC algorithm which makes further complexity reduction possible. Let us note that existence of such modification does not contradict the general lower bound in [1], as the au-

thors in [1] consider the case of general path dependent functionals of $(X_t)_{t \in [0,T]}$ and we study here functionals of the form $f(X_T)$ under some additional smoothness assumption on f. In this context let us mention the work [6], where a deterministic quadrature rule based on the distribution of a simplified weak Ito-Taylor step is proposed. In fact, the algorithm presented in [6] also provides complexity rates better than ones of the MLMC algorithm, but its application is limited to one-dimensional case.

The plan of the paper is as follows. The main idea of the variance reduced MLMC approach is introduced in (2). Section 3 is devoted to the construction of control variate.

2 Variance Reduced MLMC

Fix some $0 < L_0 < L$ and consider a random variable M_{L_0} with $E[M_{L_0}] = 0$, then

$$\mathbf{E}[f(X_{L,T})] = \mathbf{E}[f(X_{L_0,T}) - M_{L_0}] + \sum_{l=L_0+1}^{L} \mathbf{E}[f(X_{l,T}) - f(X_{l-1,T})].$$

As opposite to the representation (1), we start the telescopic sum not at the roughest approximation $\Delta_0 = T$, but at some intermediate one corresponding to Δ_{L_0} . Moreover, at level zero we subtract a zero mean random variable M_{L_0} , which can be viewed as a control variate. By fixing a vector of natural numbers $\mathbf{n} = (n_{L_0}, \dots, n_L) \in \mathbb{N}_0^{L-L_0+1}$, we can construct a modified multilevel Monte Carlo estimate for $Y = \mathbb{E}[f(X_{L,T})]$ via

$$\widehat{Y} \doteq \frac{1}{n_{L_0}} \sum_{i=1}^{n_{L_0}} \left[f(X_{L_0,T}^{(i)}) - M_{L_0}^{(i)} \right] + \sum_{l=L_0+1}^{L} \frac{1}{n_l} \sum_{i=1}^{n_l} \left[f(X_{l,T}^{(i)}) - f(X_{l-1,T}^{(i)}) \right],$$

where all pairs $(X_{l-1}^{(i)}, X_{l,T}^{(i)})$ are independent. Obviously $E[\widehat{Y}] = E[f(X_{L,T})]$ and

$$\operatorname{Var}\left[\widehat{Y}\right] \lesssim \frac{1}{n_{L_0}} \operatorname{Var}\left[f(X_{L_0,T}) - M_{L_0}\right] + \sum_{l=L_0+1}^{L} n_l^{-1} \Delta_l^{\beta},$$

where \leq stands for inequality up to a some constant not depending on L and **n**, provided the assumption (2) is fulfilled and f is Lipschitz continuous. So we have for the mean square error of \hat{Y} ,

$$\mathbb{E}\left[|\widehat{Y} - \mathbb{E}\left[f(X_T)\right]|^2\right] \lesssim \Delta_L^{2\alpha} + \frac{1}{n_{L_0}} \operatorname{Var}\left[f(X_{L_0,T}) - M_{L_0}\right] + \sum_{l=L_0+1}^L n_l^{-1} \Delta_l^{\beta}.$$
 (4)

Page:3

job:MCQMC_submit mac

macro:svmult.cls

Note that the cost of calculating $\sum_{l=L_0+1}^{L} \frac{1}{n_l} \sum_{i=1}^{n_l} \left[f(X_{l,T}^{(i)}) - f(X_{l-1,T}^{(i)}) \right]$ needs to be proportional to $\Delta_{L_0}^{\beta-1} \cdot \varepsilon^{-2}$ under conditions (2) with $\beta > 1$. So we need to construct the control variate M_{L_0} and calculate $\frac{1}{n_{L_0}} \sum_{i=1}^{n_{L_0}} \left[f(X_{L_0,T}^{(i)}) - M_{L_0}^{(i)} \right]$ with the cost lower than ε^{-2} . We will measure the complexity of building and using the control variate M_{L_0} in terms of Δ_{L_0} . Moreover, we will assume, that the control variate M_{L_0} satisfies

$$\operatorname{Var}\left[f(X_{L_0,T}) - M_{L_0}\right] \lesssim \Delta_{L_0}^{\mu}$$

with some $\mu > 0$. Our assumptions can be formalised as follows for certain M > 1 and $\Delta_{\ell} = M^{-\ell}$.

- Sampling cost: $\operatorname{cost}(f(X_{l,T}), f(X_{l-1,T})) \leq c \cdot M^{l}$ (5)
- Weak convergence: $\left| \mathsf{E}(f(X_T)) \mathsf{E}(f(X_{l,T})) \right| \le c \cdot M^{-l \cdot \alpha}, \ \alpha \ge 1$ (6)
- Degree of coupling: $\mathbb{E}\left[\left(f(X_{l,T}) f(X_{l-1,T})\right)^2\right] \le c \cdot M^{-l \cdot \beta}, \ \beta > 1$ (7)
- CV construction cost: $\operatorname{cost}_{c}(M_{L_{0}}) \leq c \cdot \Delta_{L_{0}}^{-\mu_{1}}, \ \mu_{1} \geq 0$ (8)
- CV usage cost: $\operatorname{cost}_{u}(M_{L_0}) \leq c \cdot \Delta_{L_0}^{-\mu_2}, \ \mu_1 \geq \mu_2 \geq 0$ (9)

CV effectiveness:
$$\operatorname{Var}[f(X_{L_0,T}) - M_{L_0}] \le c \cdot \Delta_{L_0}^{\mu_3}, \mu_3 > \max(1,\mu_2)$$
 (10)

The above assumptions lead to the following complexity theorem.

Theorem 1. Under the assumptions in (5)-(10), we have with $\eta = \min(\beta - 1, \mu_3 - \max(\mu_2, 1))$, the overall cost of the variance reduced MLMC algorithm is proportional to

$$\mathscr{C}^{\varepsilon}_{\mu_{1},\mu_{2},\mu_{3},\beta} = \varepsilon^{-\max\left(2 - \frac{2\eta}{\eta + \max(\mu_{1},1)}, \frac{1}{\alpha}\right)}, \quad \varepsilon \to 0.$$

Proof. It's easy to see, that with $\beta > 1$ we have the overall cost proportional to

$$\max\left(\Delta_{L_0}^{\beta-1}\cdot\varepsilon^{-2},\varepsilon^{-\frac{1}{\alpha}}\right)+\Delta_{L_0}^{-\mu_1}+\max\left(\varepsilon^{-2}\cdot\Delta_{L_0}^{\mu_3},1\right)\cdot\left(\Delta_{L_0}^{-1}+\Delta_{L_0}^{-\mu_2}\right)$$

or equivalently

$$\begin{split} \Delta_{L_0}^{\beta-1} \cdot \varepsilon^{-2} + \varepsilon^{-\frac{1}{\alpha}} + \Delta_{L_0}^{-\mu_1} + \Delta_{L_0}^{-1} + \Delta_{L_0}^{-\mu_2} + \varepsilon^{-2} \cdot \Delta_{L_0}^{\mu_3 - \max(\mu_2, 1)} \asymp \\ \Delta_{L_0}^{\eta} \cdot \varepsilon^{-2} + \varepsilon^{-\frac{1}{\alpha}} + \Delta_{L_0}^{-\max(\mu_1, \mu_2, 1)} \Rightarrow \Delta_{L_0} \asymp \varepsilon^{\frac{2}{\eta + \max(\mu_1, 1)}}. \end{split}$$

Hence the overall cost is bounded by

$$\mathscr{C}^{\varepsilon}_{\mu_1,\mu_2,\mu_3,\beta} = \varepsilon^{-\max\left(2 - \frac{2\eta}{\eta + \max(\mu_1,1)}, \frac{1}{\alpha}\right)}.$$

Remark 1 We see that, provided $\alpha > \frac{1}{2}$, it doesn't matter how difficult it is to construct the control variate M_{L_0} or to use it (i.e. how large is $\max(\mu_1, \mu_2, 1)$), if the goal

is to get the complexity rate less than ε^{-2} (i.e. to get $\eta > 0$); what matters only is that the efficiency of the control variate M_{L_0} is sufficiently large (i.e. $\mu_3 > \max\{1, \mu_2\}$).

3 Construction of control variates for SDE

In this section we are going to present a method of constructing control variates satisfying the assumptions (5)-(10). Let T > 0 be a fixed time horizon. Consider a *d*-dimensional diffusion process $(X_t; t \in [0, T])$ defined by the Itô stochastic differential equation

$$dX_t = \mu(X_t) dt + \sigma(X_t) dW_t, \quad X_0 = x \tag{11}$$

for continuous functions $\mu : \mathbb{R}^d \to \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \to \mathbb{R}^{d \times m}$, where $(W_t \in \mathbb{R}^m; t \in [0, T])$ is a standard *m*-dimensional Brownian motion. The coefficients μ and σ are assumed to be such that there exists unique strong solution for (11).

3.1 Some observations in one-dimensional case

Our construction of the control variate will be connected to the Wiener Chaos decomposition (see [7] for a detailed exposition). Let $(\phi_i)_{i\geq 1}$ be an orthonormal basis in $L^2(0,T)$. The Wiener chaos of order $p \in \mathbb{N}$ is the L^2 -closure of the vector field spanned by

$$\left\{\prod_{i\geq 1}H_{p_i}\left(\int_0^T\phi_i(s)\,dW_s\right):\,\sum_{i\geq 1}p_i=p\right\},\,$$

where H_p is the Hermite polynomial of order p given by the formula

$$H_p(x) \doteq \frac{(-1)^p}{\sqrt{p!}} e^{x^2/2} \frac{d^p}{dx^p} e^{-x^2/2}, \quad p \in \mathbb{N}_0.$$

It is well known that $(H_p)_{p\geq 0}$ is a sequence of orthogonal polynomials in $L^2(\mathbb{R},\mu)$, where μ stands for centered Gaussian measure. Every square integrable random variable *F*, measurable with respect to \mathscr{F}_T , admits the decomposition

$$F = \mathbf{E}[F] + \sum_{k \ge 1} \sum_{|p|=k} c_p \prod_{i \ge 1} H_{p_i} \left(\int_0^T \phi_i(s) \, dW_s \right)$$
(12)

with $p = (p_1, \ldots, p_k, \ldots) \in \mathbb{N}^{\mathbb{N}}$ and $|p| = \sum_{i \ge 1} p_i$. Taking into account the orthogonality of Hermite polynomials, we derive an expression for the coefficients c_p :

Page: 5 job: MCQMC_submit macro: svmult.cls date/time: 3-Mar-2022/2:45

Denis Belomestny and Tigran Nagapetyan

$$c_p = \mathbb{E}\left[F \times \prod_{i \ge 1} H_{p_i}\left(\int_0^T \phi_i(s) dW_s\right)\right].$$

In the situation where $F = f(X_{\Delta,T})$ and $X_{\Delta,T}$ comes from a discretisation of (11) with a time step $\Delta = T/J$ for some $J \in \mathbb{N}$, it is natural to take $\phi_i(t) \doteq \mathbb{I}(t \in](i-1)\Delta, i\Delta])/\sqrt{\Delta}$, i = 1, ..., J. If $X_{\Delta,T}$ is measurable with respect to $\mathscr{G}_J \doteq \sigma(\Delta_1 W, ..., \Delta_J W)$ with $\Delta_i W \doteq W_{i\Delta} - W_{(i-1)\Delta}$ and $f(X_{\Delta,T}) \in L^2(\mathscr{G}_J, \mathbb{P})$, then we obtain the decomposition

$$f(X_{\Delta,T}) = \mathbb{E}[f(X_{\Delta,T})] + \sum_{k \ge 1} \sum_{|p|=k} c_p \prod_{i=1}^J H_{p_i}(\Delta_i W / \sqrt{\Delta})$$
(13)

with $p = (p_1, ..., p_J) \in \mathbb{N}_J^0$. The above measurability assumption means that the approximation $X_{\Delta,T}$ involves only uniformly-spaced discrete Brownian increments. This is, for example, the case for the Euler scheme and the Milstein scheme under the commutativity condition. Furthermore, Giles and Szpruch [4] constructed a coupled Milstein scheme that fulfils both the above measurability assumption and the condition (2) with $\beta > 1$. Let us further analyse the decomposition (13). First note that the coefficients in (13) can be computed via

$$c_p = \mathbf{E}\left[f(X_{\Delta,T}) \times \prod_{i=1}^J H_{p_i}(\Delta_i W/\sqrt{\Delta})\right].$$

So now we can consider a control variate of the form

$$M_{K,\Delta} \doteq \sum_{k=1}^{K} \sum_{|p|=k} c_p \prod_{i=1}^{J} H_{p_i} \left(\Delta_i W / \sqrt{\Delta} \right).$$

Note that in order to compute all coefficients appearing in $M_{K,\Delta}$ we need $O(J^K)$ operations, which is unfeasible. We overcome this issue in the next section, where we suggest another representation.

3.2 Control variate construction in multidimensional case

Let $d, J \in \mathbb{N}$, let $(\Omega, \mathscr{F}, \mathbb{P}, (\mathscr{F}_t)_{t \in [0,T]})$ be a filtered probability space and let $W : [0,T] \times \Omega \to \mathbb{R}^m$ be a standard $(\mathscr{F}_t)_{t \in [0,T]}$ -Brownian motion. For $J \in \mathbb{N}$ and $j \in \{1, \dots, J\}$ we define $\Delta_j W = W_{j\Delta} - W_{(j-1)\Delta}$, where $\Delta = T/J$, and by W^i we denote the *i*-th component of the vector. Let $\Phi_{\delta} : \mathbb{R}^{d+m} \to \mathbb{R}^d$ be measurable and suppose that for all $s, t \in [0,T]$ satisfying s < t and $\delta \in [0,\infty)$, there exists a constant $C \in [0,\infty)$ not depending on δ and t - s, such that for all $X \in L^2((\Omega, \mathscr{F}, \mathbb{P}); \mathbb{R}^d)$ it holds that

Page: 6 job: MCQMC_submit macro: svmult.cls date/time: 3-Mar-2022/2:45

Variance Reduced MLMC Path Simulation: going beyond the complexity ε^{-2}

$$\mathbf{E} \left\| \Phi_{\delta} \left(X, \frac{W_t - W_s}{\sqrt{t - s}} \right) \right\|^2 \le C^2 \mathbf{E} \, \|X\|^2.$$
(14)

7

Theorem 2. Let $J \in \mathbb{N}$. Let $X_{\Delta,0} \in L^2((\Omega, \mathscr{F}_0, \mathbb{P}); \mathbb{R}^d)$, and define $(X_{\Delta,j\Delta})_{j=1}^J \in L^2(\Omega; (\mathbb{R}^d)^J)$ by

$$X_{\Delta,j\Delta} = \Phi_{\Delta} \left(X_{\Delta,(j-1)\Delta}, \frac{\Delta_j W}{\sqrt{\Delta}} \right)$$
(15)

for all $j \in \{1, 2, ..., J\}$. Let $f : \mathbb{R}^d \to \mathbb{R}$ be measurable and satisfy $E |f(X_{\Delta,T})|^2 < \infty$. Then

$$f(X_{\Delta,T}) = \mathbb{E}[f(X_{\Delta,T})|X_0]$$

$$+ \sum_{k=1}^{\infty} \sum_{j=1}^{J} \sum_{i=1}^{m} a_{k,j,i} (X_{\Delta,(j-1)\Delta}, (\Delta_j W^r)_{r=1}^{i-1}) H_k\left(\frac{\Delta_j W^i}{\sqrt{\Delta}}\right),$$
(16)

where the coefficients $a_{k,j,i}$: $\mathbb{R}^{d+i-1} \to \mathbb{R}$ in (16) are given by

$$a_{k,j,i}(x,y) = \mathbb{E}\left[f(X_{\Delta,T})H_k\left(\frac{\Delta_j W^i}{\sqrt{\Delta}}\right) \middle| X_{\Delta,(j-1)\Delta} = x, (\Delta_j W^r)_{r=1}^{i-1} = (y^r)_{r=1}^{i-1}\right]$$
(17)

for all $k \in \mathbb{N}$, $j \in \{1, \ldots, J\}$ and $i \in \{1, \ldots, m\}$.

Remark 2 The analogue of the main representation (16) is of the form

$$f(X_{\Delta,T}) = \mathbb{E}[f(X_{\Delta,T})] + \sum_{j=1}^{J} \sum_{i=1}^{m} \sum_{1 \le q_1 < \ldots < q_i \le m} \sum_{k \in \mathbb{N}^i} a_{k,j,i,q}(X_{\Delta,(j-1)\Delta}) \prod_{r=1}^{i} H_{k_r}\left(\frac{\Delta_j W^{q_r}}{\sqrt{\Delta}}\right),$$

with

$$a_{k,j,i,q}(x) = \mathbb{E}\left[f(X_{\Delta,T})\prod_{r=1}^{i}H_{k_r}(\Delta_j W^{q_r}/\sqrt{\Delta}) \middle| X_{\Delta,(j-1)\Delta} = x\right]$$

and can be proved along the same lines as (16).

Let us compare (in the one-dimensional case for the ease of notation) the representations (16) and (13). First of all, (16) has the form

$$f(X_{\Delta,T}) = \mathbb{E}[f(X_{\Delta,T})|X_0] + \sum_{k=1}^{\infty} \sum_{j=1}^{J} a_{k,j}(X_{\Delta,(j-1)\Delta}) H_k\left(\frac{\Delta_j W}{\sqrt{\Delta}}\right)$$

with

$$a_{k,j}(X_{\Delta,(j-1)\Delta}) = \sum_{p \in I_{j,k}} c_p(f(X_{\Delta,J\Delta})) \left(\prod_{\ell=0}^j H_{p_\ell}(\Delta_\ell W/\sqrt{\Delta})\right).$$

Denote

Page:7

job:MCQMC_submit ma

macro:svmult.cls date/

Denis Belomestny and Tigran Nagapetyan

$$\mathscr{M}_{K,J} := \sum_{k=1}^{K} \sum_{j=1}^{J} a_{k,j} (X_{\Delta,(j-1)\Delta}) H_k\left(\frac{\Delta_j W}{\sqrt{\Delta}}\right)$$
(18)

and

$$M_{K,J} := \sum_{k=1}^{K} \sum_{|p|=k} c_p \prod_{i=1}^{J} H_{p_i}(\Delta_i W/\sqrt{\Delta}).$$

The difference between the control variates $\mathcal{M}_{K,J}$ and M_K can be written as

$$\mathscr{M}_{K,J} - M_{K,J} = \sum_{k \ge K+1} \sum_{j=1}^{J} \sum_{k'=1}^{K} \left[\sum_{|p|=k, p \in I_{j,k'}} c_p \prod_{i=1}^{J} H_{p_i} \left(\frac{\Delta_i W}{\sqrt{\Delta}} \right) \right].$$

This implies that

$$\operatorname{Var}(f(X_{\Delta,T}) - \mathcal{M}_{K,J}) \leq \operatorname{Var}(f(X_{\Delta,T}) - M_{K,J}).$$

This latter inequality turns out to be very useful, as it is easier to analyze the truncation error related to the control variates $M_{K,J}$ than the one connected to $\mathcal{M}_{K,J}$. We can write $X_{\Delta,j\Delta} = F(\sqrt{\Delta} \cdot \xi_1, \dots, \sqrt{\Delta} \cdot \xi_j)$ for some function $F : \mathbb{R}^{j\cdot m} \to \mathbb{R}^d$, $j = 1, \dots, J$, where $\xi_j = \frac{\Delta_j W}{\sqrt{\Delta}}$, $j = 1, \dots, J$, are *m*-dimensional Brownian increments and $X_{\Delta,0} = x$. This notation will be used in the next theorem, which assesses the efficiency of the control variate $M_{K,\Delta}$.

Theorem 3. Consider the equation (11) and its discretization given by

$$X_{\Delta,j\Delta} = \Phi_{\Delta}\left(X_{\Delta,(j-1)\Delta}, \frac{\Delta_j W}{\sqrt{\Delta}}\right).$$

Assume that function $f(F(x, y_1, ..., y_J))$ is p times differentiable in $y \in \mathbb{R}^J$ such that

$$\mathbf{E}\left[\frac{\partial^{p_1+\ldots+p_J}f(F(\sqrt{\Delta}\cdot\xi_1,\ldots,\sqrt{\Delta}\cdot\xi_J))}{\partial\xi_1^{p_1}\ldots\partial\xi_J^{p_J}}\right]$$

is uniformly bounded in $J(\Delta = T/J)$ and $p \in \mathbb{N}^J$ with $|p| \leq K$. Then

$$\operatorname{Var}(f(X_{\Delta,J\Delta}) - M_{K,\Delta}) \lesssim \Delta^{K}.$$

Proof. Due to the independence of Brownian increments and orthogonality of Hermite polynomials, we get

$$\operatorname{Var}\left[\sum_{\substack{k\geq K+1}}^{\infty}\sum_{\substack{p\in\mathbb{N}_{0}^{(J+1)\times m}\\|p|=k}}c_{p}(f(X_{\Delta,J\Delta}))\prod_{j=0}^{J}\prod_{i=1}^{m}H_{p_{j,i}}\left(\frac{\Delta_{j}W^{i}}{\sqrt{\Delta}}\right)\right]=\sum_{\substack{k\geq K+1}}^{\infty}\sum_{\substack{p\in\mathbb{N}_{0}^{(J+1)\times m}\\|p|=k}}c_{p}^{2}.$$

Page:8

job:MCQMC_submit mac

macro:svmult.cls

Variance Reduced MLMC Path Simulation: going beyond the complexity ε^{-2}

Using the integration by parts in the case of one-dimensional diffusion (the multidimensional case is absolutely the same), we get for any c_p :

$$\mathbf{E}\left[f(X_{\Delta,T}) \times \prod_{i=1}^{J} H_{p_i}\left(\frac{\Delta_i W}{\sqrt{\Delta}}\right)\right] = \mathbf{E}\left[F(x,\sqrt{\Delta}\cdot\xi_1,\dots,\sqrt{\Delta}\cdot\xi_J) \times \prod_{i=1}^{J} H_{p_i}(\xi_i)\right]$$
$$= \frac{\Delta^{(K+1)/2}}{\prod_{i=1}^{J} p_i!} \mathbf{E}\left[\frac{\partial^{p_1+\dots+p_J} f(F(x,\sqrt{\Delta}\cdot\xi_1,\dots,\sqrt{\Delta}\cdot\xi_J)))}{\partial\xi_1^{p_1}\dots\partial\xi_J^{p_J}}\right].$$

Remark 1. Suppose that $f(x) \equiv x$. Then using the chain rule, we get for the Euler scheme:

$$\frac{\partial f(F(\sqrt{\Delta} \cdot \xi_1, \dots, \sqrt{\Delta} \cdot \xi_J))}{\partial \xi_j} = \prod_{i=j+1}^J \left(1 + \frac{\partial \mu}{\partial x}(X_i) \Delta + \frac{\partial \sigma}{\partial x}(X_i) \sqrt{\Delta} \cdot \xi_i \right) \sigma(X_j).$$

Taking expectation and using conditioning, we get

$$\mathbf{E}\left[\frac{\partial f(F(\sqrt{\Delta}\cdot\xi_1,\ldots,\sqrt{\Delta}\cdot\xi_J))}{\partial\xi_j}\right] = \prod_{i=j+1}^J \left(1 + \mathbf{E}\left[\frac{\partial\mu}{\partial x}(X_i)\right]\Delta\right) \mathbf{E}[\sigma(X_j)].$$
(19)

Hence the left hand side of (19) is uniformly bounded in *J*, provided the expectations $E\left[\frac{\partial \mu}{\partial x}(X_i)\right]$, i = j + 1, ..., J, and $E[\sigma(X_j)]$ are bounded.

4 Regression approach to control variate construction

In order to use the control variate (18) we need to compute the coefficients . Since

$$a_{k,j,i}(x,y) = \mathbb{E}\left[f(X_{\Delta,T})H_k\left(\frac{\Delta_j W^i}{\sqrt{\Delta}}\right) \middle| X_{\Delta,(j-1)\Delta} = x, (\Delta_j W^r)_{r=1}^{i-1} = (y^r)_{r=1}^{i-1}\right]$$
(20)

we can use nonparametric regression to estimate them, and therefore define coefficients μ_1 , μ_2 and μ_3 .

4.1 General nonparametric approach

To ease the explanation we now consider on D + 1-dimensional random vector (X, Y) where X is \mathbb{R}^D -valued and Y is \mathbb{R} -valued. Suppose that we want to find an approximation which is "close to" the \mathbb{R} -valued function

$$a(x) := \mathbb{E}[Y|X = x].$$
 (21)

Page: 9 job: MCQMC_submit macro: svmult.cls date/time: 3-Mar-2022/2:45

Let us choose Q real-valued functions ψ_1, \ldots, ψ_Q on \mathbb{R}^D and simulate a big number N of samples from the distributions of X and Y. In what follows these N samples are denoted by \mathcal{D}_N :

$$\mathscr{D}_N \doteq \left\{ (X^{(n)}, Y^{(n)}) : n = 1, \dots, N \right\}.$$

Let $\beta = (\beta_1, \dots, \beta_Q)$ be a solution of the following least squares optimisation problem:

$$\operatorname{argmin}_{\beta \in \mathbb{R}^{Q}} \sum_{n=1}^{N} \left[Y^{(n)} - \sum_{k=1}^{Q} \beta_{k} \psi_{k}(X^{(n)}) \right]^{2}.$$

Define an estimate for the function *a* via

$$\hat{a}(x) \doteq \hat{a}(x, \mathscr{D}_N) \doteq \sum_{k=1}^{Q} \beta_k \psi_k(x), \quad x \in \mathbb{R}^d.$$

The intermediate expression $\hat{a}(x, \mathcal{D}_N)$ in the above formula emphasises that the estimates \hat{a} of the functions a are random in that they depend on the simulated samples. The cost of computing β is of order NQ^2 , since β is of the form $\beta = B^{-1}b$ with

$$B_{k,l} \doteq \frac{1}{N} \sum_{n=1}^{N} \psi_k(X^{(n)}) \psi_l(X^{(n)})$$
(22)

and

$$b_k \doteq rac{1}{N} \sum_{n=1}^N \psi_k ig(X^{(n)} ig) Y^{(n)},$$

where $k, l \in \{1, ..., Q\}$. In what follows, we use the notation \mathbb{P}_X for the distribution of *X*. In particular, we will work with the corresponding L^2 -norm:

$$\|g\|_{L^2(\mathbb{P}_X)}^2 \doteq \int_{\mathbb{R}^D} g^2(x) \mathbb{P}_X(dx) = \mathbb{E}\left[g^2(X)\right].$$

We assume that, for some positive constants Σ and A, it holds

(A1) $\sup_{x \in \mathbb{R}^D} \operatorname{Var}[Y|X = x] \le \Sigma < \infty,$ (A2) $\sup_{x \in \mathbb{R}^D} |a(x)| \le A < \infty.$

Next we denote by \tilde{a} the truncated regression estimate, which is defined as follows:

$$\tilde{a}(x) \doteq T_A \hat{a}(x) \doteq \begin{cases} \hat{a}(x) & \text{if } |\hat{a}(x)| \le A, \\ A \operatorname{sgn}(\hat{a}(x)) & \text{otherwise.} \end{cases}$$
(23)

Page:10 job:MCQMC_submit macro:svmult.cls date/time:3-Mar-2022/2:45

Variance Reduced MLMC Path Simulation: going beyond the complexity $arepsilon^{-2}$

We again emphasise that, in fact, $\tilde{a}(x) = \tilde{a}(x, \mathcal{D}_N)$, that is, the estimates \tilde{a} of the functions *a* depend on the simulated samples. Under (A1)–(A2) we obtain the following L^2 -upper bound:

$$\mathbb{E} \|\tilde{a} - a\|_{L^{2}(\mathbb{P}_{X})}^{2} \leq \tilde{c} \left(\Sigma + A^{2}(\log N + 1)\right) \frac{Q}{N} + 8 \inf_{g \in \Psi_{Q}} \|a - g\|_{L^{2}(\mathbb{P}_{X})}^{2}, \quad (24)$$

where $\Psi_Q \doteq \text{span}(\{\psi_1, \dots, \psi_Q\})$ and $\tilde{c} > 0$ is a universal constant (cf. Theorem 11.3 in [5]). Let us introduce the assumption that the function *a* can be well approximated by the functions from Ψ_Q in the sense that there are constants $\kappa > 0$ and $D_{\kappa} > 0$ such that

$$\inf_{g \in \Psi_Q} \|a - g\|_{L^2(\mathbb{P}_X)}^2 \le \frac{D_{\kappa}}{Q^{\kappa}}.$$
(25)

Note that this is a natural condition to be satisfied for good choices of Ψ_Q . So under assumptions (A1), (A2) and (25), we get

$$\mathbb{E} \|\tilde{a} - a\|_{L^2(\mathbb{P}_X)}^2 \lesssim \frac{Q}{N} + \frac{1}{Q^{\kappa}}.$$

Let us now consider the control variate

$$\tilde{\mathscr{M}}_{K,J} = \sum_{k=1}^{K} \sum_{j=1}^{J} \sum_{i=1}^{m} \tilde{a}_{k,j,i} (X_{\Delta,(j-1)\Delta}, (\Delta_j W^r)_{r=1}^{i-1}) H_k \left(\frac{\Delta_j W^i}{\sqrt{\Delta}}\right),$$
(26)

where $\tilde{a}_{k,j,i}$ are estimated using nonparametric regression with Q basis functions. It's easy to see, that

$$\operatorname{Var}(f(X_{\Delta,J\Delta}) - \tilde{\mathscr{M}}_{K,J}) \preceq \Delta^{K} + \Delta^{-1} \cdot \left(\frac{Q}{N} + \frac{1}{Q^{\kappa}}\right)$$

under the corresponding assumptions on the functions $a_{k,j,i}$. At this point it is very important to emphasize, that we still have

$$\mathbf{E}[\tilde{\mathscr{M}}_{K,J}|\mathscr{D}_N]=0,$$

which means, that at no point we introduce additional bias due to the finite number of basis functions or due to the numerical discretization. Now taking into account that the number of coefficients to compute is of order JK^dNQ^2 , we can take for any fixed K > 1, $\mu_3 = K$, $\mu_2 = (K+1)/\kappa + 1$, $\mu_1 = (K+1)(1+3/\kappa) + 1$ to get for the overall cost of the variance reduced MLMC algorithm

$$\mathscr{C}^{\varepsilon}_{\mu_{1},\mu_{2},\mu_{3},\beta} = \varepsilon^{-\max\left(2-\frac{2\eta}{\eta+\mu_{1}},\frac{1}{\alpha}\right)}, \quad \varepsilon \to 0.$$

Page:11

job:MCQMC_submit

macro:svmult.cls

with $\eta = \min(\beta - 1, K(1 - 1/\kappa) - 1/\kappa - 1)$. So the complexity order is alway below ε^{-2} , provided $\beta > 1$, $K > (\kappa + 1)/(\kappa - 1)$, $\kappa > 1$ and $\alpha > 1/2$.

5 Numerical experiments

We consider the SDE

$$dX_{t}^{i} = -\sin\left(X_{t}^{i}\right)\cos^{3}\left(X_{t}^{i}\right)dt + \cos^{2}\left(X_{t}^{i}\right)dW_{t}^{i}, \quad X_{0}^{i} = 0, \quad i \in \{1, 2, 3, 4\},$$

$$dX_{t}^{5} = \sum_{i=1}^{4} \left[-\frac{1}{2}\sin\left(X_{t}^{i}\right)\cos^{2}\left(X_{t}^{i}\right)dt + \cos\left(X_{t}^{i}\right)dW_{t}^{i}\right] + dW_{t}^{5}, \quad X_{0}^{5} = 0.$$
(27)

The solution of (27) is given by

$$X_t^i = \arctan(W_t^i), \ i \in \{1, 2, 3, 4\}, X_t^5 = \sum_{i=1}^4 \operatorname{arsinh}(W_t^i) + W_t^5.$$

for $t \in [0, 1]$. Further, we consider the functional

$$f(x) = \cos\left(\sum_{i=1}^{5} x^{i}\right) - 20\sum_{i=1}^{4} \sin(x^{i}),$$

that is, we have

$$\mathbb{E}\left[f\left(X_{1}\right)\right] = \left(\mathbb{E}\left[\cos\left(\arctan\left(W_{1}^{1}\right) + \operatorname{arsinh}\left(W_{1}^{1}\right)\right)\right]\right)^{4} \mathbb{E}\left[\cos\left(W_{1}^{5}\right)\right]$$

We use the an antithetic MLMC approach from [2], where the following Milstein alike discretization scheme has been utilized:

$$X_{n+1}^{i} = X_{n}^{i} - \sin(X_{n}^{i})\cos^{3}(X_{n}^{i})\Delta + \cos^{2}(X_{n}^{i})\Delta W_{n}^{i} - \cos^{3}(X_{n}^{i}) \cdot \sin(X_{n}^{i}) \cdot (\Delta^{2}W_{n}^{i} - \Delta)$$

$$X_{n+1}^{5} = \sum_{i=1}^{4} \left[-\frac{1}{2}\sin(X_{n}^{i})\cos^{2}(X_{n}^{i})\Delta + \cos(X_{t}^{i})\Delta W_{n}^{i} + \frac{1}{2}\cos(X_{t}^{i})\sin(X_{t}^{i}) \cdot (\Delta^{2}W_{n}^{i} - \Delta) \right] + \Delta W_{n}^{5}$$

for i = 1, 2, 3, 4 and $X_0^i = X_0^5 = 0$. We use the updated Antithetic MLMC estimator

$$\begin{split} \widehat{Y} &\doteq \frac{1}{n_{L_0}} \sum_{i=1}^{n_{L_0}} \left[f(X_{L_0,T}^{(i)}) - M_{2,2^{L_0}}^{(i)} \right] + \sum_{l=L_0+1}^{L} Y_l = \\ \frac{1}{n_{L_0}} \sum_{i=1}^{n_{L_0}} \left[f(X_{L_0,T}^{(i)}) - M_{2,2^{L_0}}^{(i)} \right] + \sum_{l=L_0+1}^{L} \frac{1}{n_l} \sum_{i=1}^{n_l} \left[\frac{1}{2} \left(f(X_{f,l,T}^{(i)}) + f(X_{a,l,T}^{(i)}) \right) - f(X_{l-1,T}^{(i)}) \right], \end{split}$$

where subindices stand for discretized paths with antithetic approach, discussed in details in [2], which we refer to due to the length constraints. This scheme with

coupling recovers variance decay rate $\beta = 2$. We consider accuracies $\log_2(\varepsilon) = -4, -7, -10, -13$, and we set $L_0 = 1, 2, 3, 4$ respectively, which corresponds to our approach to use polynomial partitioning with p = 3.

5.1 Regression for control variate

We consider the control variate (26), with K = 2 and we want to find an estimator for it $\tilde{\mathcal{M}}_{2,2L_0}$. It's easy to see, that

$$\operatorname{Var}(f(X_{\Delta,J\Delta}) - \tilde{\mathscr{M}}_{2,2^{L_0}}) \leq \Delta_{L_0}^2 + \Delta_{L_0}^{-1} \cdot \left(\frac{Q}{N} + \frac{1}{Q^{\kappa}}\right)$$

that all the drift, diffusion and the functional are sufficiently regular, so in the case of polynomial regression of order p, we will have $\kappa = 2 \cdot p$. Taking into account that the overall regression cost is of order $\Delta^{-1}NQ^2$, we set

$$Q \simeq \Delta^{-\frac{3}{\kappa}}, N \simeq \Delta^{-3-\frac{3}{\kappa}} \Rightarrow \mu_1 = 4 + \frac{9}{\kappa}, \ \mu_2 = 1 + \frac{3}{\kappa}, \ \mu_3 = 2$$

In our numerical experiments we will focus on piecewise polynomial approximation of order p = 3, hence $\kappa = 6$, which leads to

$$\eta = \min(\beta - 1, K(1 - 1/\kappa) - 1/\kappa - 1) = \min(1, 2(1 - 1/6) - 1/6 - 1) = 0.5.$$
$$\mathscr{C}^{\varepsilon}_{\mu_1, \mu_2, \mu_3, \beta} = \varepsilon^{-1\frac{5}{6}}, \quad \Delta_{L_0} \asymp \varepsilon^{\frac{1}{3}}.$$

The results, describing the effectiveness of the control variate construction and the variance decay of the antithetic MLMC can be seen on Figure 5.1. Due to sufficient regularity of drift, diffusion and functional (moreover, we work here with the bounded functionals), our numerical results are able to reproduce perfectly our expectations from the theory. The variance for MLMC and Single level MC methods are estimated based on $2.5 \cdot 10^6$ paths, while the variance of Single level MC is estimated on 10^6 paths, with the control variates constructed from $N = \max(\Delta_l^{-3.5}, 100)$ paths. The complexity of $\hat{\mathcal{M}}_{2,2^{L_0}}$ estimation increases with L_0 , so we need so set it accurately. According to our parameters choice we set $\Delta_{L_0} \simeq \varepsilon^{\frac{1}{3}}$, so for accuracies $\log_2(\varepsilon) = -4, -7, -10, -13$ we have $L_0 = 1, 2, 3, 4$ respectively. The cost of constructing the control variates in our simulation has the form

$$\Delta_{L_0}^{-1} \cdot N \cdot Q^2 = \Delta_{L_0}^{-1} \cdot \max(\Delta_l^{-3.5}, 100) \cdot \Delta_{L_0}^{-1},$$

which is presented on Figure 5.1. There one can see reference dotted straight lines, which correspond to the expected Antithetic MLMC complexity of order ε^{-2} for different values of ε . As an illustration of accuracy of our chosen parameters, we

Page:13 job:MCQMC_submit macro:svmult.cls date/time:3-Mar-2022/2:45



Fig. 1 Variance decay of Antithetic MLMC (square markers), Single level MC (circle markers), and Single level with control variate $\tilde{\mathcal{M}}_{2,2^{L_0}}$ (diamond markers).



Fig. 2 Dotted straight lines correspond to the expected MLMC complexity of order ε^{-2} for different values of ε . Decaying lines correspond to MLMC complexities from level of the form $\varepsilon^{-2} \cdot 2^{-(\beta-1)} = \varepsilon^{-2} \cdot 2^{-l}$. The cost of $\tilde{\mathcal{M}}_{2,2^{L_0}}$ construction is presented on a dashed line with starshaped markers.

also plot the cost of estimating $\sum_{l=L_0+1}^{L} Y_l$, which in the case of $\beta = 2$ is proportional to $\varepsilon^{-2} \cdot 2^{-L_0}$. According to the complexity Theorem 1, L_0 corresponds to a level,

Page:14 job:MCQMC_submit macro:svmult.cls date/time:3-Mar-2022/2:45

when the overall cost of construction of control variate $\tilde{\mathcal{M}}_{2,2^{L_0}}$ (and the cost of simulating level L_0 along with the cost of sampling $\tilde{\mathcal{M}}_{2,2^{L_0}}$) is proportional to the cost of simulating all forthcoming Antithetic MLMC levels. As one can see on Figure 5.1, our constraint on number of training paths N doesn't satisfy this rule of thumb for $\varepsilon = 2^{-4}$, while for other accuracies it matches almost perfectly. Moreover, the gain of Antithetic MLMC approach with control variate is clearly represented by the difference between the dotted straight lines for each accuracy and the level, where line of constructing the control variate cost intersects the decaying line of leftover MLMC levels calculation cost.

6 Conclusion

We presented a MLMC method with control variate on its starting level, which allows us to have a randomised Monte Carlo algorithm with complexity of order less than ε^{-2} . This approach is implementable in an arbitrary dimension, which is its strong advantage. Our analysis also suggests that other possible improvements for MLMC can be made, if one can improve the cost of the initial level in MLMC approach, as its cost without variance reduction (determined by central limit theorem) prevent us from improving ε^{-2} complexity bound, regardless of how good is the coupling. The disadvantage of the proposed nonparametric regression algorithm, but not the general control variate approach, is that it requires regularity of the problem and a sophisticated choice of basis functions. Both of these issues should be treated for considered problem specifically, but our numerical results suggest, that even very simple regression approach can give substantial saving, provided that the problem of interest is regular enough.

7 Proofs

7.1 Proof of Theorem 2

Without loss of generality we may assume there exists a Gaussian vector, suggestively denoted by $\Delta_0 W \colon \Omega \to \mathbb{R}^m$ with a covariance operator $\Delta I_{\mathbb{R}^d}$, such that $\mathscr{F}_0 = \sigma(\Delta_0 W)$. For $p \in \mathbb{N}_0^{m \times (J+1)}$ and $X \in L^2((\Omega, \mathscr{F}, \mathbb{P}), \mathbb{R}^d)$ define

$$c_p(X) = \mathbb{E}\left[X\prod_{j=0}^J \prod_{i=1}^m H_{p_{i,j}}(\Delta_j W^i/\sqrt{\Delta})\right].$$
(28)

For $j \in \{0, 1, ..., J\}$, $i \in \{1, ..., m\}$, and $k \in \mathbb{N}$, define

Page:15 job:MCQMC_submit macro:svmult.cls date/time:3-Mar-2022/2:45

Denis Belomestny and Tigran Nagapetyan

$$I_{j,i,k} = \left\{ p \in \mathbb{N}_0^{(J+1) \times m} \colon \begin{array}{l} p_{j,i} = k \text{ and } (\forall r \in \{i+1,\dots,m\} \colon p_{j,r} = 0) \\ \text{and } (\forall \ell \in \{j+1,\dots,J\}, \forall r \in \{1,\dots,m\} \colon p_{\ell,r} = 0) \end{array} \right\}.$$
(29)

The Wiener chaos expansion of $f(X_{\Delta,J\Delta})$ with respect to $(\Delta_j W)_{j=0}^J$ is given by

$$f(X_{\Delta,J\Delta}) = \mathbb{E}[f(X_{\Delta,J\Delta})] + \sum_{k=1}^{\infty} \sum_{\substack{p \in \mathbb{N}_{0}^{(J+1) \times m} \\ |p| = k}} c_{p}(f(X_{\Delta,J\Delta})) \prod_{j=0}^{J} \prod_{i=1}^{m} H_{p_{j,i}}(\Delta_{j}W^{i}/\sqrt{\Delta})$$
$$= \mathbb{E}[f(X_{\Delta,J\Delta})] + \sum_{j=0}^{J} \sum_{i=1}^{m} \sum_{\substack{p \in I_{j,i,k}}}^{\infty} c_{p}(f(X_{\Delta,J\Delta}))H_{k}(\frac{\Delta_{j}W^{i}}{\sqrt{\Delta}})$$
$$\times \left(\prod_{\ell=0}^{j-1} \prod_{r=1}^{m} H_{p_{\ell,r}}(\Delta_{\ell}W^{r}/\sqrt{\Delta})\right) \left(\prod_{r=1}^{i-1} H_{p_{j,r}}(\Delta_{j}W^{r}/\sqrt{\Delta})\right)$$
(30)

Then it follows that

$$E[f(X_{\Delta,J\Delta})|X_0] = E[f(X_{\Delta,J\Delta})|\Delta_0 W] =$$

$$E[f(X_{\Delta,J\Delta})] + \sum_{i=1}^m \sum_{k=1}^\infty \sum_{p \in I_{0,i,k}} c_p(f(X_{\Delta,J\Delta})) H_k\left(\frac{\Delta_0 W^i}{\sqrt{\Delta}}\right) \times \left(\prod_{r=1}^m H_{p_{0,r}}\left(\frac{\Delta_0 W^r}{\sqrt{\Delta}}\right)\right),$$
(31)

and for all $j \in \{1, ..., J\}$, all $i \in \{1, ..., m\}$, and all $k \in \mathbb{N}$ it holds that

$$E\left[f(X_{\Delta,J\Delta})H_{k}(\Delta_{j}W^{i}/\sqrt{\Delta})\middle|(\Delta_{\ell}W)^{j-1}_{\ell=1},(\Delta_{j}W^{r})^{i-1}_{r=1}\right]$$

$$= E\left[f(X_{\Delta,J\Delta})H_{k}(\Delta_{j}W^{i}/\sqrt{\Delta})\middle|X_{\Delta,(j-1)\Delta},(\Delta_{j}W^{r})^{i-1}_{r=1}\right]$$

$$= \sum_{p\in I_{j,i,k}}c_{p}(f(X_{\Delta,J\Delta}))\left(\prod_{\ell=0}^{j-1}\prod_{r=1}^{m}H_{p_{\ell,r}}(\Delta_{\ell}W^{r}/\sqrt{\Delta})\right)\left(\prod_{r=1}^{i-1}H_{p_{j,r}}(\Delta_{j}W^{r}/\sqrt{\Delta})\right).$$
(32)

We can use now (32) to rewrite (30) as (16).

References

- Creutzig, J., Dereich, S., Müller-Gronbach, T., Ritter, K.: Infinite-dimensional quadrature and approximation of distributions. Foundations of Computational Mathematics 9(4), 391–429 (2009)
- Giles, M., Szpruch, L.: Antithetic multilevel Monte Carlo estimation for multi-dimensional SDEs without Lévy area simulation. Annals of Applied Probability 24(4), 1585–1620 (2014)
- Giles, M.B.: Multilevel Monte Carlo path simulation. Operations Research 56(3), 607–617 (2008)
- Giles, M.B., Szpruch, L.: Antithetic multilevel Monte Carlo estimation for multi-dimensional SDEs without Lévy area simulation. Annals of Applied Probability 24(4), 1585–1620 (2014)

Page:16 job:MCQMC_submit macro:svmult.cls

Variance Reduced MLMC Path Simulation: going beyond the complexity $arepsilon^{-2}$

- Györfi, L., Kohler, M., Krzyżak, A., Walk, H.: A distribution-free theory of nonparametric regression. Springer Series in Statistics. Springer-Verlag, New York (2002). DOI 10.1007/ b97848. URL http://dx.doi.org/10.1007/b97848
- Müller-Gronbach, T., Yaroslavtseva, L.: Deterministic quadrature formulas for SDEs based on simplified weak Itô-Taylor steps. DFG SPP 1324 (Preprint 167) (2013)
- 7. Nualart, D.: The Malliavin calculus and related topics. Springer (2006)

Page:17

job:MCQMC_submit

macro:svmult.cls