

Entanglement Spectrum of a Random Partition: Connection with the Localization Transition

Sagar Vijay¹ and Liang Fu¹

¹*Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

We study the entanglement spectrum of a translationally-invariant lattice system under a random partition, implemented by choosing each site to be in one subsystem with probability $p \in [0, 1]$. We apply this random partitioning to a translationally-invariant (i.e., clean) topological state, and argue on general grounds that the corresponding entanglement spectrum captures the universal behavior about its *disorder-driven* transition to a trivial localized phase. Specifically, as a function of the partitioning probability p , the entanglement Hamiltonian H_A must go through a topological phase transition driven by the percolation of a random network of edge-states. As an example, we analytically derive the entanglement Hamiltonian for a one-dimensional topological superconductor under a random partition, and demonstrate that its phase diagram includes transitions between Griffiths phases.

PACS numbers: 73.43.Cd, 03.67.Mn

In recent years, systematic studies of quantum entanglement have greatly advanced our understanding of topological states of matter that cannot be adiabatically connected to a trivial product state. For example, topological entanglement entropy is directly related to the total quantum dimension of fractional quasi-particles [1, 2]. More recently, there has been a growing interest in utilizing the full entanglement spectrum to extract other universal properties [3], especially in chiral topological phases (e.g., quantum Hall states) or symmetry-protected topological phases (e.g., topological insulators). As a common feature, these phases have topologically-protected gapless excitations on a physical boundary. When the ground state is spatially cut into left and right halves, the low-lying part of the entanglement spectrum shares the same universal characteristics as the energy spectrum of these boundary excitations [3–9].

A recent work studied the entanglement spectrum obtained from an extensive partition that divides a system into two *extensive* subsystems [10]. For topological states that support gapless edge states, the corresponding entanglement spectrum was found to encode a wealth of information about the universal quantum critical behavior that would arise at its phase transition to a trivial direct-product state, despite the fact that the system under study itself is non-critical. It has been further shown that the entanglement spectra of extensive partitions can be directly computed from the matrix product state or tensor network representation of ground-state wavefunctions [11], which may offer insights into topological phase transitions [12–15].

In this Letter, we study the entanglement spectrum generated from a random partition that spatially bipartitions a system in a *probabilistic* manner: each physical site is chosen to be in subsystem B (or A) with a probability $p \in [0, 1]$ (or $1 - p$). We apply this random partition to a translationally-invariant (i.e., clean) topological state, and argue on general grounds that the cor-

responding entanglement spectrum reproduces the universal behavior about its *disorder-driven* transition to a trivial localized phase. As an example, we analytically derive the form of the entanglement Hamiltonian for a clean one-dimensional topological superconductor under random partition, and establish the entanglement phase diagram as a function of probability p , finding agreement with the physical phase diagram of a disordered superconductor [16].

We begin by considering a translationally-invariant topological state, which can be either a topological insulator/superconductor or a bosonic symmetry-protected topological phase. It has been shown [10] that upon varying the geometry of A and B subsystems in an extensive partition, the corresponding entanglement Hamiltonian undergoes a gap-closing transition that lies in the same universality class as the transition to a topologically trivial state realized by tuning the *physical* Hamiltonian. This intriguing connection is not a coincidence, but follows from the nature of topological phase transitions, which are driven by the percolation of gapless edge-states. For example, the transition from a quantum Hall insulator to a trivial insulator is described by the quantum percolation of chiral edge-states in a Chalker-Coddington network model [17]. Extensive partitioning of a quantum Hall insulator precisely creates, in the low-lying part of the entanglement spectrum, a network of chiral edge states moving along the percolating borders between A and B . This mapping explains why topological phase transitions and entanglement spectra of extensive partitions are intimately related. It further motivates us to study the random partitioning of a topological ground-state, for which the entanglement spectrum is expected to mimic the network model with randomness and thus connect with the localization transition.

We define the probabilistic partitioning of a clean, topological state $|\Psi\rangle$ as follows. We independently choose each physical site in the full system to be in the B subsystem

tem with probability p ; the remaining sites are defined to be in subsystem A . A partial trace of the density matrix over sites in subsystem B yields a reduced density matrix for the A subsystem $\rho_A \equiv \text{Tr}_B |\Psi\rangle\langle\Psi|$, which can be interpreted as the *thermal* density matrix at temperature $T = 1$ for an entanglement Hamiltonian H_A : $\rho_A \equiv e^{-H_A}$.

Our goal is to determine the phase diagram of H_A as a function of the partitioning probability p . First, when $p \rightarrow 0$, the probabilistic partitioning yields a vanishingly small B subsystem, with most sites belonging to the A subsystem. In this limit, the ground-state of the entanglement Hamiltonian, denoted by $|\psi_A\rangle$, must share the same topological index as the original ground state $|\Psi\rangle$. As $p \rightarrow 1$, however, most sites become part of the B subsystem, so that $|\psi_A\rangle$ becomes a trivial product state over the disjoint regions of the A subsystem, and hence must be topologically trivial. Since the topological character of $|\psi_A\rangle$ changes as we tune the partitioning probability p , we conclude that the entanglement Hamiltonian H_A must go through a *phase transition* at some critical partitioning probability $p = p_c$. Physically, the transition is driven by the percolation of a random network of gapless edge-states propagating around traced-out regions of the B subsystem, as in the case of the aforementioned checkerboard-type extensive partition. Even though the original state $|\Psi\rangle$ is translationally-invariant, the probabilistic partitioning procedure introduces randomness into the entanglement Hamiltonian H_A , with the probability p effectively tuning disorder strength.

The phase diagram of the entanglement Hamiltonian as a function of p satisfies additional constraints. For a given bipartition, the eigenvalue spectra of the reduced density matrices ρ_A and ρ_B are identical, though their Hilbert spaces are distinct. By definition, for a fixed partitioning probability p , the B subsystem is, on average, equivalent to the A subsystem obtained with a partitioning probability $1 - p$. Therefore, the ensemble-averaged spectra of the entanglement Hamiltonians $H_A(p)$ and $H_A(1 - p)$ must be identical. As a result, the presence of a phase transition in the entanglement Hamiltonian with partition probability p implies another transition at probability $1 - p$. In the case where the topological index of $|\Psi\rangle$ cannot be evenly divided between two subsystems, as is the case for topological insulators with a \mathbb{Z}_2 index or quantum Hall insulators with an odd Chern number, we further expect that H_A exhibits at least a topological phase transition at partitioning probability $p = 1/2$, when the two subsystems are equivalent on average.

For the remainder of the paper, we apply our random partitioning procedure to Kitaev's model [18] for a clean one-dimensional p -wave superconductor, extract the phase diagram of the entanglement Hamiltonian as a function of partitioning probability, and demonstrate its correspondence with a disordered superconductor. The




FIG. 1: The two ground-states of the Kitaev p -wave superconductor with dimerization of Majorana fermions *across* (top) and *within* (bottom) lattice sites, corresponding to a TSC and a trivial p -wave superconductor, respectively.

Kitaev model is described by the Hamiltonian

$$H = -w \sum_n \left(c_{n+1}^\dagger c_n + \text{h.c.} \right) + \sum_n \left(\Delta c_{n+1}^\dagger c_n^\dagger + \text{h.c.} \right) - \mu \sum_n \left(c_n^\dagger c_n - \frac{1}{2} \right) \quad (1)$$

with fermion operators c_n, c_n^\dagger satisfying canonical anti-commutation relations. To simplify the calculation below, we restrict ourselves to the case where $w = \Delta \in \mathbb{R}$. Introducing two species of Majorana fermions at each lattice site $\gamma_n \equiv c_n + c_n^\dagger$ and $\chi_n \equiv (c_n^\dagger - c_n)/i$, we may write the Hamiltonian as

$$H = \frac{iw}{2} \sum_n [\eta \gamma_n \chi_n + \chi_n \gamma_{n+1}] \quad (2)$$

where the dimensionless parameter $\eta \equiv \mu/2w$ distinguishes the topologically trivial strong pairing phase ($|\eta| > 1$) and topologically non-trivial weak pairing phase ($|\eta| < 1$). As shown in Figure 1, in the topological superconductor (TSC) phase, Majorana fermions couple more strongly across two adjacent lattice sites than within a lattice site, leading to two unpaired Majorana fermions at the ends of the chain. In the extreme limit when $\eta = 0$ in the topological phase, pairs of Majorana fermions within lattice sites completely decouple. We will refer to this special point $\eta = 0$ as the *Kitaev limit* of the Hamiltonian (2), which proves to be a useful starting point for our analysis below.

Let $|\Psi(\eta)\rangle$ be the ground-state of the Kitaev Hamiltonian with parameter $|\eta| < 1$ in the topological regime. Applying the random partitioning procedure to this ground-state, we trace over *physical* lattice sites (each of which contains two Majorana fermions) with probability p , and obtain an entanglement Hamiltonian $H_A(\eta; p)$. Clearly, $H_A(\eta; p)$ contains random couplings between sites in the A subsystem, and mimics the physical Hamiltonian of a disordered superconductor, with the partitioning probability p playing the role of disorder strength.

To derive $H_A(\eta; p)$, we note that the entanglement Hamiltonian of an N -site free fermion system such as the Kitaev model can only contain fermion bilinear terms [19]. The spectrum of the entanglement Hamiltonian

can be determined from the correlation matrices in the original ground-state $C_{nm} \equiv \langle \Psi | c_n^\dagger c_n | \Psi \rangle$ and $F_{nm} \equiv \langle \Psi | c_n^\dagger c_m^\dagger | \Psi \rangle$, by solving the eigenvalue problem [19]:

$$(2\hat{C} - 2\hat{F} - 1)(2\hat{C} + 2\hat{F} - 1)\phi_\ell = \tanh^2\left(\frac{\epsilon_\ell}{2}\right)\phi_\ell \quad (3)$$

where ϵ_ℓ is an eigenvalue of the entanglement Hamiltonian, with eigenvector ϕ_ℓ . Rewriting the complex fermions in terms of Majorana operators, we may define the $2N \times 2N$ skew-symmetric correlation matrix $\hat{\Gamma}$ for the Majorana fermions which has eigenvalues $\pm \tanh(\epsilon_\ell/2)$ [20, 26]. Since we are interested in the low-lying part of the entanglement spectrum $\epsilon_\ell \rightarrow 0$, $\tanh(\epsilon_\ell/2) \rightarrow \epsilon_\ell/2$ and hence $\hat{\Gamma}$ satisfies:

$$2\hat{\Gamma}\phi_\ell \approx \pm \epsilon_\ell \phi_\ell \quad (4)$$

Therefore, the correlation matrix for Majorana fermions in the original ground-state is *equivalent* to the entanglement Hamiltonian acting on low-lying states $\epsilon_\ell \rightarrow 0$ in the entanglement spectrum. By building the correlation matrix for Majorana fermions in the ground-state $|\Psi(\eta)\rangle$, we may now construct the entanglement Hamiltonian for the Kitaev model after a random partition.

For arbitrary η , performing a random partition will generally produce an entanglement Hamiltonian with highly non-local couplings, due to the non-vanishing correlations between distant Majorana fermions in the ground state. However, for a sufficiently small $|\eta|$, i.e., when the system is close to the Kitaev limit, we may derive the form of $H_A(\eta; p)$ *analytically*. Let us first consider the case $\eta = 0$, when pairs of Majorana fermions decouple. A single cut between two adjacent lattice sites then produces, in the entanglement spectrum of the A subsystem, an unpaired Majorana fermion at the end of A . Aside from this, the entanglement spectrum at $\eta = 0$ is identical to the energy spectrum (properly normalized) of decoupled Majorana pairs in the Kitaev Hamiltonian. Therefore, performing a random partition with several cuts will yield an A subsystem that consists of disjoint segments, each of which hosts unpaired Majorana fermions at the two ends.

We now explicitly construct the entanglement Hamiltonian $H_A(\eta; p)$ near the Kitaev limit by perturbing away from $\eta = 0$. As one may expect, a small η induces a small coupling between the unpaired Majorana fermions at ends of disjoint segments with the rest of the A subsystem. By an analytical calculation [26], we find that couplings between two Majorana fermions in $H_A(\eta; p)$ decrease exponentially with their separation in the original lattice. Therefore, it suffices to include *nearest-neighbor* couplings within subsystem A only in $H_A(\eta; p)$.

Two types of nearest-neighbor couplings appear in $H_A(\eta; p)$. First, to leading order in η , couplings belonging to a connected sequence of sites in the A subsystem are identical to those appearing in the original Kitaev




FIG. 2: The entanglement Hamiltonian $H_A(\eta; p)$, derived in the Kitaev limit $|\eta| \ll 1$, consists of several couplings between adjacent Majorana fermions in the subsystem A . Blue and green hoppings between nearest-neighbor Majorana fermions appear with dimensionless coupling 1 and η , respectively. Tracing over N lattice sites induces a coupling $O(\eta^N)$ between the dangling Majorana modes on the adjacent chains in the A subsystem.

Hamiltonian, after a proper normalization. Second, couplings between Majorana fermions belonging to different segments in the A subsystem are computed from their two-point correlation function. If a series of N consecutive lattice sites – between sites $m-1$ and $m+N$ – are determined to be within the B subsystem and traced over in the random partition, a coupling will be induced between the Majorana fermions at the right and left edges of the two lattice sites, which is found to be proportional to $C(N) \equiv \langle \Psi(\eta) | i\chi_{m-1}\gamma_{m+N} | \Psi(\eta) \rangle$. An explicit calculation [26] yields the result that at long distances, i.e. large N ,

$$C(N) \sim \eta^N / \sqrt{N} + O(\eta^{N+1}) \quad (5)$$

We then conclude that the entanglement Hamiltonian takes the form:

$$H_A(\eta; p) = \frac{iw}{2} \sum_{n \in A} \eta \gamma_n \chi_n + \frac{iw}{2} \sum_{m > n \in A} f_{nm} \chi_n \gamma_m \quad (6)$$

where f_{nm} has non-zero elements $f_{n, n+1} = 1$ and $f_{nm} = C(m-n-1)$ if n and m label lattice sites at the right and left edges of two adjacent segments in the A subsystem. The couplings in the entanglement Hamiltonian are illustrated in Fig. 2.

We now demonstrate that the topological character of the entanglement Hamiltonian (6) changes at a critical partitioning probability $p = 1/2$. Specifically, we demonstrate that when $p < 1/2$ the entanglement Hamiltonian in the A subsystem supports an exponentially localized edge-state $|\psi\rangle$ at zero energy and corresponds to a topological superconductor phase, but that such a state does not exist for $p > 1/2$.

To see this, we first note that a random partitioning of a chain with probability p will produce an A subsystem that consists of clusters of lattice sites. Let us now introduce a boundary in the A subsystem and explicitly construct a zero-energy state of the entanglement spectrum under a random partition. Recall that an exact zero-energy left-boundary eigenstate of the translationally-invariant Kitaev Hamiltonian (2) takes the form $|\psi\rangle =$

$(1, 0, \eta, 0, \eta^2, 0, \dots)$, with $|\eta| < 1$, in the basis of Majorana sites on a semi-infinite chain [18]. We find a similar zero-energy state can be obtained for the entanglement Hamiltonian (6) of the A subsystem, which consists of consecutive clusters of sites of lengths $\{\ell_k\}$, each separated by distance $\{d_k\}$. An edge-state of the entanglement Hamiltonian now takes the form:

$$|\psi_A\rangle \propto \left(\underbrace{1, 0, \eta, 0, \dots, \eta^{\ell_1}, 0, \frac{\eta^{\ell_1}}{C(d_1)}, 0, \dots, \frac{\eta^{\ell_1+\ell_2}}{C(d_1)}, 0, \dots}_{\text{1st cluster of sites}}, \underbrace{\prod_{k=1}^{N-1} \frac{\eta^{\ell_k}}{C(d_k)}, 0, \dots, \eta^{\ell_N} \prod_{k=1}^{N-1} \frac{\eta^{\ell_k}}{C(d_k)}, 0, \dots}_{\text{Nth cluster of sites}} \right) \quad (7)$$

in a basis of Majorana sites in A . From the form of the two-point function computed previously we see that amplitude for the edge-state on the first Majorana site in the N th cluster of the A subsystem is given by:

$$\psi_N = \prod_{k=1}^{N-1} \frac{\eta^{\ell_k}}{C(d_k)} \sim \prod_{k=1}^{N-1} \eta^{\ell_k - d_k} = \eta^{\sum_k \ell_k - \sum_k d_k} \quad (8)$$

Now, if we consider the amplitude at the end of the chain, we see that $\psi_N \rightarrow \eta^{L_A - L_B}$ where L_A and L_B are the sizes of the A and B subsystems, respectively. For a system of size L , regardless of the probability distributions for the lengths $\{\ell_k\}$ and $\{d_k\}$, the sizes of the two subsystems are determined from the partitioning probability to be $L_A = (1-p)L$ and $L_B = pL$ in the thermodynamic limit $L \rightarrow +\infty$ so that

$$\psi_N \rightarrow [\eta^{1-2p}]^L \quad (9)$$

When $p < 1/2$ we observe that the state (7) is an exact zero-energy eigenstate of the entanglement Hamiltonian. When $p > 1/2$, however, the amplitude at the end of the chain diverges and the above state becomes non-normalizable for an infinite set of clusters.

The above calculation of an edge-state immediately implies that the entanglement Hamiltonian H_A changes from being topologically non-trivial at partitioning probability $p < 1/2$ to trivial when $p > 1/2$, and hence must be critical at the point $p = 1/2$. This transition can also be understood by integrating out the Majorana fermions in the interior of the clusters in the A subsystem and constructing an *effective* entanglement Hamiltonian H_A^{eff} acting exclusively on the dangling Majorana modes at the ends of each cluster. In this case, H_A^{eff} will describe a dimerized Majorana fermion chain, in which two adjacent Majorana fermions correspond to sites separated by lengths $\{\dots \ell_k, d_k, \ell_{k+1}, d_{k+1} \dots\}$ in the original lattice. The nearest-neighbor hopping in A , which is proportional to the corresponding correlation function in the ground-state, is determined by the lengths $\{d_k\}$ for intra-cluster hoppings or $\{\ell_{k+1}\}$ for inter-cluster

hoppings. At $p = 1/2$, the A and B subsystems are equivalent on average, so that the length distributions $\{\ell_k\}$ and $\{d_k\}$ are identical, and the ensemble of H_A^{eff} is translationally-invariant, instead of dimerized. The corresponding ground state of a one-dimensional Majorana fermion chain is well-known to be critical [21].

We now demonstrate that in the vicinity of $p = 1/2$, the entanglement Hamiltonian is in Griffiths phases, characterized by a singularity in the density of states at zero energy due to the proliferation of segments of the topologically-ordered or trivial phase, respectively. Recall that when $p < 1/2$, near $p = 1/2$, the characteristic size of clusters in the A subsystem is larger than that of the B subsystem. Then, the dangling Majorana modes on adjacent clusters in the A subsystem, separated by distance x will mix to form localized bound-states with finite energy $\epsilon \sim \exp[cx \ln|\eta|]$, with $c > 0$ a constant. Since the probability of such a configuration of sites in the A subsystem is $p^x(1-p)^2$, the contribution of these low-energy modes to the density of states in the entanglement ground-state is [16]:

$$\rho(\epsilon) = \int_0^\infty dx (1-p)^2 p^x \delta(\epsilon - e^{cx \ln|\eta|}) \propto \frac{1}{\epsilon^{1-\beta(p)}} \quad (10)$$

with the non-universal exponent $\beta(p) \equiv \ln(p)/c \ln|\eta|$. The power-law singularity in the density of states signals the presence of a Griffiths region for an entanglement ground-state with p near $1/2$, due to the proliferation of low-energy configurations of Majorana edge-modes dimerizing across lattice sites. $p > 1/2$ also correspond to a Griffiths phase, with exponent $\beta(1-p)$ due to Majorana modes at the ends of the *same* chain forming bound-states with exponentially small energy. The two Griffiths phases at $p < 1/2$ and $p > 1/2$ are both characterized by a power-law singularity in the density of states at zero energy, but are topologically distinct, as shown by the presence and absence of zero-energy Majorana fermion at the boundary.

To summarize, we have introduced a random partitioning scheme to study the disorder-driven quantum critical behavior of a topological phase; applying this procedure to the one-dimensional p -wave superconductor yields an interesting phase diagram, consisting of two topologically distinct Griffiths phases separated by a critical point. The random-partitioning scheme may be naturally used to study spin-chains [22, 23], as well as higher-dimensional systems to numerically extract critical exponents of disorder-driven phase transitions, such as localization transitions in all symmetry classes of non-interacting topological phases [24, 25]. It might also be interesting to study the entanglement spectrum of fractional topological phases under a random partition, although the connection with topological phase transitions appears to be less direct.

Acknowledgment: We thank Tim Hsieh for helpful discussions. This work is supported by DOE Office of Basic

Energy Sciences, Division of Materials Sciences and Engineering under award de-sc0010526.

-
- [1] M. Levin and X.G. Wen, Phys. Rev. Lett. **96**, 110405 (2006).
 - [2] A. Kitaev and J. Preskill, Phys. Rev. Lett. **96** 110404 (2006).
 - [3] Li, H. and Haldane, F. D. M., Phys. Rev. Lett. **101**, 010504 (2008).
 - [4] A. Chandran, M. Hermanns, N. Regnault, and B.A. Bernevig, Phys. Rev. B **84**, 205136 (2011).
 - [5] X-L Qi, H. Katsura and A. W. W. Ludwig, Phys. Rev. Lett. **108**, 196402 (2012).
 - [6] B. Swingle and T. Senthil, Phys. Rev. B **86**, 045117 (2012).
 - [7] F. Pollmann, A. M. Turner, E. Berg and M. Oshikawa, Phys. Rev. B **81**, 064439 (2010).
 - [8] L. Fidkowski, Phys. Rev. Lett. **104**, 130502 (2010).
 - [9] J. Dubail and N. Read, Phys. Rev. Lett. **107**, 157001 (2011).
 - [10] T. Hsieh and L. Fu, Phys. Rev. Lett. **113**, 106801 (2014).
 - [11] T. Hsieh, L. Fu and Xiao-Liang Qi, Phys. Rev. B **90**, 085137 (2014).
 - [12] W. J. Rao, X. Wan and G. Zhang, Phys. Rev. B **90**, 075151 (2014).
 - [13] R. Santos, arXiv:1408.1716
 - [14] T. Fukui and Y. Hatsugai, Journal of the Physical Society of Japan, **83**, 113705 (2014).
 - [15] J. Borchmann, A. Farrell, S. Matsuura, and T. Peregr-Barnea, arXiv:1407.5980v1 [cond-mat.str-el] (2014).
 - [16] O. Motrunich, K. Damle, and D. A. Huse, Phys. Rev. B **63**, 224204 (2001).
 - [17] J. T. Chalker and P. D. Coddington, J. Phys. C **21**, 2665 (1988).
 - [18] A. Kitaev, arXiv:0010440v2 [cond-mat] (2000).
 - [19] I. Peschel, J. Phys. A **36**, L205 (2003).
 - [20] I. Peschel and V. Eisler, J. Phys. A **42**, 504003 (2009).
 - [21] I. C. Fulga, B. van Heck, J. M. Edge, and A. R. Akhmerov, Phys. Rev. B **89** 155424 (2014).
 - [22] D. S. Fisher, Phys. Rev. B **51**, 6411 (1995).
 - [23] R. A. Hyman and K. Yang, Phys. Rev. Letters **78**, 1783 (1997).
 - [24] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
 - [25] A. Kitaev, arXiv:0901.2686 (2009).
 - [26] Supplementary Material

SUPPLEMENTARY MATERIAL

In this section, we briefly review the calculation of the entanglement Hamiltonian for the ground-state of a free-fermion system as presented in [1, 2] and compute the two-point correlation function between Majorana fermions on even and odd Majorana sites in the ground-state of the Kitaev chain [3], in the limit that the coupling $\eta \equiv \mu/2w$ is small. In the main text, this calculation is used to construct the associated entanglement Hamiltonian, which is directly proportional to the corre-

lation matrix for the Majorana fermions in the ground-state, when acting on the low-lying states in the entanglement spectrum.

The Entanglement Hamiltonian from a Bipartition of a Free-Fermion Ground State

For a free-fermion Hamiltonian, the entanglement Hamiltonian obtained after a bipartition of the ground-state $|\Psi\rangle$ may be determined entirely from the two-point correlation functions in the ground-state. For a free-fermion system, any n -point correlation function can be written as a sum over two-point functions by Wick's Theorem. Furthermore, the reduced density matrix ρ_A is defined to reproduce all observables in the A subsystem. Therefore, for a fermion number-conserving, free-fermion Hamiltonian, the reduced density matrix must take the form:

$$\rho_A = \frac{e^{-\mathcal{H}_A}}{Z} \quad \mathcal{H}_A \equiv \sum_{i,j \in A} h_{ij} c_i^\dagger c_j \quad (11)$$

with $Z \equiv \text{Tr} [e^{-\mathcal{H}_A}]$ and fermion operators c_i, c_j^\dagger satisfying canonical anti-commutation relations. In this case, we may explicitly diagonalize the entanglement Hamiltonian \mathcal{H}_A , and express the correlation matrix $C_{nm} \equiv \langle c_n^\dagger c_m \rangle$ (with $n, m \in A$) evaluated in the ground-state in terms of the eigenvalues $\{\epsilon_j\}$ and eigenvectors $\{\phi_j\}$ of \mathcal{H}_A as [1]:

$$C_{nm} = \text{Tr} [\rho_A c_n^\dagger c_m] = \sum_j \phi_j^*(n) \phi_j(m) (e^{\epsilon_j} + 1)^{-1} \quad (12)$$

In matrix form, this is equivalent to the statement that the entanglement Hamiltonian may be written as $(\mathcal{H}_E)^T = \ln[(1 - \hat{C})/\hat{C}]$, and that \hat{C} solves the eigenvalue problem [2]:

$$(1 - 2\hat{C})\phi_j = \tanh\left(\frac{\epsilon_j}{2}\right)\phi_j \quad (13)$$

A similar calculation may be performed for a free-fermion Hamiltonian defined on N lattice sites that only conserves fermion number parity. In this case, the expansion of an n -point function in terms of two-point correlators will include contributions from both the regular $C_{nm} \equiv \langle c_n^\dagger c_m \rangle$ and ‘anomalous’ correlation functions $F_{nm} \equiv \langle c_n^\dagger c_m^\dagger \rangle$. In order for the reduced density matrix to reproduce all observables in A , the entanglement Hamiltonian must also be a free-fermion Hamiltonian with pairing terms [1], i.e. of the form:

$$\mathcal{H}_E = \sum_{n,m} [f_{nm} c_n^\dagger c_m + \Delta_{nm} c_n^\dagger c_m^\dagger + \text{h.c.}] \quad (14)$$

As explained in [2], we may diagonalize the Hamiltonian by Bogoliubov transformation, so that the \hat{C} and \hat{F} ma-

trices now satisfy the eigenvalue problem:

$$(2\hat{C} - 2\hat{F} - 1)(2\hat{C} + 2\hat{F} - 1)\phi_j = \tanh^2\left(\frac{\epsilon_j}{2}\right)\phi_j \quad (15)$$

Equivalently, the entanglement Hamiltonian may be expressed in terms of Majorana operators $\gamma_{2n-1} \equiv c_n + c_n^\dagger$ and $\gamma_{2n} \equiv -i(c_n - c_n^\dagger)$. In this case, \mathcal{H}_E will only contain terms bilinear in the Majorana operators, and may be written in the form:

$$\mathcal{H}_E = \frac{i}{4} \sum_{n,m \in A} B_{nm} \gamma_n \gamma_m \quad (16)$$

with \hat{B} a $2N \times 2N$ real, skew-symmetric matrix. We may diagonalize the entanglement Hamiltonian by block-diagonalizing \hat{B} by an appropriate orthogonal transformation and bringing it into canonical form:

$$\mathcal{O}^T \hat{B} \mathcal{O} = \begin{pmatrix} 0 & \epsilon_1 & & & & \\ -\epsilon_1 & 0 & & & & \\ & & 0 & \epsilon_2 & & \\ & & -\epsilon_2 & 0 & & \\ & & & & \ddots & \end{pmatrix} \quad (17)$$

with $\mathcal{O} \in O(2N)$ so that the entanglement eigenvalues are $\pm\epsilon_\ell$. Taking the appropriate real linear combination of the Majorana operators $\{\gamma_n\}$ given by the eigenvectors of \hat{B} that diagonalize the entanglement Hamiltonian, it is possible to compute the two-point correlation function $\Gamma_{nm} \equiv \langle \gamma_n \gamma_m \rangle$ ($n \neq m$) in the original ground-state. Similar to the correlation function for a free-fermion system conserving fermion number, this yields the result [2] that Γ solves the eigenvalue problem:

$$\hat{\Gamma} \phi_\ell = \pm \tanh\left(\frac{\epsilon_\ell}{2}\right) \phi_\ell \quad (18)$$

Therefore, in the limit $\epsilon_\ell \rightarrow 0$, we see that $2\hat{\Gamma} \phi_\ell \approx \pm \epsilon_\ell \phi_\ell$, so that the two-point function for Majorana fermions in the original ground-state is equivalent to the entanglement Hamiltonian when acting on the low-lying ($\epsilon_\ell \rightarrow 0$) states in the entanglement spectrum.

The Two-Point Correlation Function for the Kitaev Majorana Chain

We now compute the two-point function for Majorana fermions in the ground-state of the Kitaev Hamiltonian [3]

$$H = -w \sum_n (c_{n+1}^\dagger c_n + \text{h.c.}) + \sum_n (\Delta c_{n+1}^\dagger c_n^\dagger + \text{h.c.}) - \mu \sum_n \left(c_n^\dagger c_n - \frac{1}{2} \right) \quad (19)$$

with $w = \Delta \in \mathbb{R}$. The complex fermion operators satisfy canonical anti-commutation relations $\{c_n, c_m\} =$

$\{c_n^\dagger, c_m^\dagger\} = 0$ and $\{c_n, c_m^\dagger\} = \delta_{nm}$. In the topologically non-trivial phase, the ground-state of the Hamiltonian on an N -site chain with periodic boundary conditions takes the form of a BCS wavefunction with odd fermion parity:

$$|\Psi\rangle = \prod_{k \neq 0} \left[u_k + v_k \cdot a_k^\dagger a_{-k}^\dagger \right] a_0^\dagger |0\rangle \quad (20)$$

where the parameter u_k is given by:

$$u_k^2 = \frac{1}{2} \left(1 + \frac{2w \cos k + \mu}{E(k)} \right) \quad (21)$$

with $u_k^2 + v_k^2 = 1$. Here, the quantity $E(k) = \sqrt{(2w \cos k + \mu)^2 + 4w^2 \sin^2 k}$ is the bulk dispersion [3]. We note that the non-vanishing correlation functions for the ground-state take the form:

$$\langle \Psi | c_p^\dagger c_k^\dagger | \Psi \rangle = \delta_{k,-p} \cdot u_k v_k \quad (22)$$

$$\langle \Psi | c_k^\dagger c_p | \Psi \rangle = \delta_{k,p} \cdot v_k^2 \quad (23)$$

We now explicitly compute the two-point correlation functions for the Majorana fermions in the ground-state of the clean Kitaev chain. The real-space correlation functions for the complex fermions may be written by performing a Fourier transform:

$$\langle c_n^\dagger c_m^\dagger \rangle = \frac{1}{N} \sum_k e^{ik(m-n)} u_k v_k \quad (24)$$

$$\langle c_n^\dagger c_m \rangle = \frac{1}{N} \sum_k e^{ik(m-n)} v_k^2 \quad (25)$$

Now, we define the Majorana operators γ_{2n} and γ_{2n-1} for $n = 1, \dots, N$ as $\gamma_{2n-1} \equiv c_n + c_n^\dagger$ and $\gamma_{2n} \equiv -i(c_n - c_n^\dagger)$ so that the operators satisfy anti-commutation relations $\{\gamma_n, \gamma_m\} = 2\delta_{nm}$. When we perform a random partitioning of our system, we will be tracing over physical lattice sites. For the topologically non-trivial ground-state, this tracing procedure will produce dangling Majorana modes at the edges of connected clusters in the A subsystem. To determine the entanglement Hamiltonian obtained from a random partitioning, we are interested in computing the two-point function between Majorana fermions on odd and even Majorana sites that are separated by physical lattice sites belonging to the B subsystem. From the two-point functions for the complex fermions, we find that this correlation function for the Majorana fermions is given by the expression

$$\langle \gamma_{2n} \gamma_{2m+1} \rangle = -i\delta_{n,m+1} + \frac{i}{N} \sum_k e^{ik(n-m-1)} \cdot v_k^2 \quad (26)$$

We now define the quantity $\eta \equiv \mu/2w$. In the limit $|\eta| \ll 1$, (near the Kitaev limit, as explained in the main text) we expand v_k^2 about $\eta = 0$, and extract the

first non-trivial contribution to the two-point function $\langle \gamma_{2n} \gamma_{2m+1} \rangle$ to lowest order in η . Note that:

$$v_k^2 = \frac{1}{2} - \frac{\eta + \cos k}{2\sqrt{1 + 2\eta \cos k + \eta^2}} \quad (27)$$

We now expand the above expression about $\eta = 0$. At each order in η , we wish to extract the term appearing in the expansion with the largest power of $\cos k$; this is the term that, when substituted into the sum appearing in (26) will provide the first non-zero contribution to the two-point function for a certain pair of well-separated Majorana fermions. At $O(\eta^\ell)$, we find that the term appearing in the expansion of v_k^2 with the highest power of $\cos k$ takes the form

$$\begin{aligned} & \left(\frac{1}{2} \cdot \frac{3}{2} \cdots \frac{2\ell-1}{2} \right) \frac{(-1)^{\ell+1} (2 \cos k + 2\eta)^\ell}{2(1 + 2\eta \cos k + \eta^2)^{2\ell+1}} \Big|_{\eta=0} \frac{\eta^\ell}{\ell!} \cos k \\ &= (-1)^{\ell+1} \frac{2^{\ell-1} \eta^\ell}{\sqrt{\pi}} \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + 1)} (\cos k)^{\ell+1} \end{aligned} \quad (28)$$

where $\Gamma(x)$ is the gamma function. Here we have made use of the following identity:

$$(2\ell - 1)!! = 1 \cdot 3 \cdot 5 \cdots (2\ell - 1) = \frac{2^\ell}{\sqrt{\pi}} \Gamma\left(\ell + \frac{1}{2}\right) \quad (29)$$

Now, substituting this term into the sum appearing in (26) for the two-point function for the Majorana fermions, we see that

$$\begin{aligned} & (-1)^{\ell+1} \frac{2^{\ell-1} \eta^\ell}{\sqrt{\pi}} \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + 1)} \frac{i}{N} \sum_k e^{ik(n-m-1)} (\cos k)^{\ell+1} = (-1)^{\ell+1} \frac{\eta^\ell}{2\sqrt{\pi}} \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + 1)} \frac{i}{N} \sum_k e^{ik(n-m-1)} \left[e^{ik(\ell+1)} + \dots \right] \\ &= i(-1)^{\ell+1} \frac{\eta^\ell}{2\sqrt{\pi}} \frac{\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + 1)} \delta_{m-n, \ell} + \left[\text{shorter-range couplings, i.e. terms proportional to } \delta_{m-n, s} \text{ with } s < \ell \right] \end{aligned} \quad (30)$$

Therefore, we observe that near the Kitaev limit, the above term provides the lowest-order contribution to the two-point function $C(\ell) \equiv \langle \gamma_{2n} \gamma_{2n+2\ell+1} \rangle$, with $\ell \geq 0$ i.e.

$$C(\ell) = (-1)^{\ell+1} \frac{\eta^\ell}{2\sqrt{\pi}} \frac{i\Gamma(\ell + \frac{1}{2})}{\Gamma(\ell + 1)} + O(\eta^{\ell+1}) \quad (31)$$

This is precisely the two-point correlation function that determines the coupling in the entanglement Hamiltonian between two Majorana fermions at the edges of adjacent chains in the A subsystem that are separated by ℓ physical lattice sites belonging to the B subsystem. Our calculation shows that these dangling Majorana modes in the A subsystem, when separated by ℓ physical lattice sites, will have an $O(\eta^\ell)$ coupling. We also see that when $\ell = 0$, we have $|2C(\ell)| = 1$, in agreement with the dimen-

sionless coupling between two nearest-neighbor Majorana fermions belonging to different physical lattice sites close to the Kitaev limit. Finally, when ℓ is sufficiently large, $\Gamma(\ell + 1/2)/\Gamma(\ell + 1) \rightarrow 1/\sqrt{\ell}$, so that:

$$C(\ell) \sim \frac{\eta^\ell}{\sqrt{\ell}} \quad (32)$$

at long distances.

-
- [1] I. Peschel, J. Phys. A **36**, L205 (2003).
 - [2] I. Peschel and V. Eisler, J. Phys. A **42**, 504003 (2009).
 - [3] A. Kitaev, arXiv:0010440v2 [cond-mat] (2000).