

Calculation of the molecular integrals with the range-separated correlation factorMichał Silkowski,¹ Michał Lesiuk,^{1, a)} and Robert Moszynski¹*Faculty of Chemistry, University of Warsaw, Pasteura 1, 02-093 Warsaw, Poland*

(Dated: 20 April 2019)

Explicitly correlated quantum chemical calculations require calculations of five types of molecular integrals beyond the standard electron repulsion integrals. We present a novel scheme, which utilises general ideas of the McMurchie-Davidson technique, to compute these integrals when the so-called “range-separated” correlation factor is used. This correlation factor combines the well-known short range behaviour, resulting from the electronic cusp condition, with the exact long-range asymptotics found for the helium atom [M. Lesiuk, B. Jeziorski, and R. Moszynski, *J. Chem. Phys.* **139**, 134102 (2013)]. Almost all steps of the presented procedure are formulated recursively, so that an efficient implementation and control of the precision are possible. Additionally, the present formulation is very flexible and general, and it allows for use of an arbitrary correlation factor in the electronic structure calculations with minor or no changes.

Keywords: explicitly correlated electronic structure theory, correlated wave function, correlation factor, molecular integrals, Gaussian-type orbitals

^{a)}Electronic mail: lesiuk@tiger.chem.uw.edu.pl.

I. INTRODUCTION

Most of the traditional *ab initio* quantum chemistry methods represent the electronic wavefunction as a linear combination of the Slater determinants. This leads to severe difficulties in describing the Coulomb hole because the electronic cusp condition requires that the electronic wavefunction behaves linearly in the interelectronic distance, r_{12} , near the coalescence points of the electrons¹⁻⁴. To overcome this difficulty explicitly correlated methods have been introduced which overtly include the r_{12} factor into the wavefunction⁵⁻⁷. Typically, the aforementioned r_{12} dependence is implemented through the product of two occupied molecular orbitals with the correlation factor, f_{12} . The correlation factor depends solely on r_{12} and makes it possible to satisfy the cusp conditions, so that the description of the Coulomb hole is dramatically improved. However, an important issue in the explicitly correlated methods is the choice of the correlation factor itself. Originally, the straightforward choice $f_{12} = r_{12}$ was made which gave rise to the class of R12 methods⁸⁻¹⁷. The theory which allows for an arbitrary correlation factor (F12) was formulated somehow later by Manby¹⁹. It was also found that the choice $f_{12} = r_{12}$ is responsible for the bulk of the error in the early R12 methods¹⁸. In 2004, Ten-no proposed the exponential correlation factor²⁰, which is nowadays routinely applied in various explicitly correlated approaches²¹⁻²⁷, and was shown to improve the results significantly as compared with R12. This suggests that the actual form of the correlation factor is important and one should continue searching for a more optimal form of f_{12} . In fact, Tew and Klopper²⁸ investigated numerically the shape of the correlation factor for the helium atom and helium-like ions and compared it with several simple analytic forms. They found that in the short-range r_{12} regime the exponential correlation factor of Ten-no is close to the optimal but they did not have access to moderate and long-range r_{12} distances. In a recent work, the form of correlation factor for the helium atom has been reconsidered by using purely analytic methods²⁹. It has been rigorously found that the asymptotic (long-range) form of the correlation factor is $f_{12} \rightarrow r_{12}^\rho e^{Br_{12}}$, as $r_{12} \rightarrow \infty$, for the helium atom with several plausible models of the electronic wavefunction. It is obvious that none of the correlation factors which are used presently obey this asymptotic behaviour. Therefore, we proposed the so-called *range-separated* correlation factor where the short- and long-range r_{12} regimes are approximated by different formulae and sewed together by using a switching function. A detailed functional form will be specified further in the paper.

One may wonder about the motivation behind the introduction of the range-separated correlation factor. From a brief inspection of the problem it appears that the long-range part of the correlation factor is completely irrelevant and the only thing that matters is the ability to model the correlation hole. However, recent results have proven that standard basis sets used for the conventional (determinants-based) calculations are suboptimal for the F12 methods^{30–32}. This is due to the fact that the cusp region is described reasonably well by the explicitly correlated part of the trial wavefunction and basis set functions are mainly necessary to model the longer-range part of the electronic wavefunction. Therefore, one can expect that a correlation factor which includes a physically motivated long-range ingredient can further reduce the basis set requirements of the current F12 approaches. However, the largest gain in the accuracy can be reached with methods where the correlation factor is responsible for a dominant fraction of the so-called dynamical correlation effects. A prominent method of this type is the geminal-augmented CASSCF method of Martinez *et al.* (G-CASSCF)²⁷. In this method the wavefunction is modelled as a standard CASSCF expansion, multiplied by a proper explicitly correlated part. Therefore, the situation is clear - the better and more physically motivated the correlation factor is, the larger percentage of the (dynamical) correlation energy is retrieved which directly improves the results. Note additionally that “guide” wavefunctions used in the Quantum Monte Carlo (QMC) methods^{33–35} also take the form of the Slater determinants multiplied by the (explicitly correlated) Jastrow factor. Unfortunately, it is not clear whether the present form of $f(r_{12})$, introduced in the context of F12 methods, is advantageous when applied to QMC as well. Nonetheless, this possibility is worth investigating, particularly in the light of recent remarkable improvements in the QMC technology and quality of the results that can be routinely obtained (see Refs. 36–45 as representative examples).

However, before the actual performance of the range-separated correlation factor can be tested, one needs to evaluate the necessary matrix elements. Efficient and stable evaluation of molecular integrals is a constant challenge for the electronic structure theory and a lot of efforts have been devoted to the development of fast and reliable integral codes. For the standard electron repulsion integrals including the Coulomb potential, $1/r_{12}$, several procedures were developed in the works of Dupuis, Rys, and King^{46–48}, McMurchie and Davidson^{49,50}, Pople and Hehre⁵¹, Obara and Saika⁵², and others^{53,55–58}. The review paper of Gill⁵⁹ is very informative in this respect. Introduction of the explicitly correlated methods

into the arsenal of quantum chemistry raised the number of types of molecular integrals which need to be routinely evaluated. Typically, each specific form of the correlation factor was handled separately by modifying the existing algorithms. The works of Klopper and Röhse⁶⁰, Bearpart *et al.*⁶¹, Valeev and Schaefer⁶², Ten-no^{20,23}, Samson *et al.*⁶³, and Weber and Daul⁶⁴ are prominent examples of this approach. In this paper we consider the evaluation of all two-electron integrals that are necessary to perform F12 calculations with the range-separated correlation factor²⁹. They are far from being trivial, mostly due to a complicated form of the correlation factor itself. We show that these integrals can be evaluated by applying the general idea of the McMurchie-Davidson recursive scheme^{49,50}. Noteworthy, our work provides a large level of generality. In fact, with minor or no changes the present scheme can be applied to an *arbitrary* correlation factor. Ahlrichs⁶⁵ presented a method to compute two-electron integrals with any kernel within the Obara-Saika technique, but his general recursive procedure is multidimensional and complicated, especially when the analytical form of the correlation factor is extensive. As far as we know, the present level of generality has never been achieved thus far within the McMurchie-Davidson scheme.

This paper is organised as follows. In Sec. II A we introduce the two-electron molecular integrals necessary for the F12 calculations and perform an initial reduction. Next, in Sec. II A the McMurchie-Davidson technique is applied which allows to express these integrals through the derivatives of the so-called basic integrals with respect to the orbitals locations. In Sec. II C we establish an analytical equation for the basic integrals. Sections II D and II E detail evaluation of the required derivatives of the basic integrals with the help of the Hobson theorem. Section III presents application of the developed scheme to the range-separated correlation factor. The generating integrals $S(\alpha, \beta, \gamma)$ are introduced and their evaluation is discussed. Finally, Sec. IV concludes our paper.

II. MOLECULAR INTEGRALS WITH A GENERAL CORRELATION FACTOR

A. Cartesian two-electron integrals

It is now a well established fact that in the F12 theories the following six types of molecular two-electron integrals appear:

$$I_1 = (ab|r_{12}^{-1}|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) r_{12}^{-1} \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (1)$$

$$I_2 = (ab|r_{12}^{-1}f_{12}|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) r_{12}^{-1} f_{12}(r_{12}) \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (2)$$

$$I_3 = (ab|f_{12}|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) f_{12}(r_{12}) \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (3)$$

$$I_4 = (ab|f_{12}f'_{12}|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) f_{12}(r_{12})f'_{12}(r_{12}) \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (4)$$

$$I_5 = (ab|[f_{12},\hat{T}_1]|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) [f_{12}(r_{12}),\hat{T}_1] \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (5)$$

$$I_6 = (ab|[f_{12},\hat{T}_1],f'_{12}|cd) = \iint d\mathbf{r}_1 d\mathbf{r}_2 \Phi_a^*(\mathbf{r}_1) \Phi_c^*(\mathbf{r}_2) [[f_{12}(r_{12}),\hat{T}_1],f'_{12}(r_{12})] \Phi_b(\mathbf{r}_1) \Phi_d(\mathbf{r}_2), \quad (6)$$

where $\hat{T}_1 = -\frac{1}{2}\nabla_1^2$ stands for the kinetic energy operator of the first electron. We have additionally increased the flexibility of our approach by allowing two different correlation factors, f_{12} and f'_{12} , to appear simultaneously. Functions on the brackets are the (unnormalised) primitive Cartesian Gaussian-type orbitals (GTOs)

$$\Phi(\mathbf{r}, a, \mathbf{A}) = x_A^i y_A^k z_A^m \exp(-ar_A^2), \quad (7)$$

where a denotes the exponent and $\mathbf{A} = (A_x, A_y, A_z)$ is the vector specifying the location of the orbital. In the above expression $x_A = x - A_x$, and similarly for the other components, and $r_A^2 = x_A^2 + y_A^2 + z_A^2$. The same notation is preserved for the remaining orbitals present in Eqs. (1)-(6), which are located at centres B, C, D , and have the respective exponents b, c, d .

In a majority of the explicitly correlated approaches three- and possibly also four-electron integrals appear, as the result of action of the Coulomb, exchange, or strong orthogonality operators on the pair functions^{8-14,17}. However, all these difficult integrals are typically not evaluated explicitly but rather approximated *e.g.* by the insertion of the resolution of

identity (RI) in terms of an auxiliary basis set^{66–68}. This approximation leaves only two-electron integrals in the form (1)-(6). Other techniques for handling many-electron integrals were also suggested^{69–72}.

Let us first simplify the r_{12} -dependent kernels in Eqs. (5) and (6). Firstly, by applying the second Green's identity in Eq. (5), the integral I_5 can be rewritten as

$$I_5 = \frac{1}{2} (\nabla_{\mathbf{A}}^2 - \nabla_{\mathbf{B}}^2) I_3. \quad (8)$$

Secondly, the kernel of Eq. (6) can be simplified by noting that

$$[[f_{12}(r_{12}), \hat{T}_1], f'_{12}(r_{12})] = (\nabla_1 f_{12}) \cdot (\nabla_1 f'_{12}) = \frac{\partial f_{12}}{\partial r_{12}} \frac{\partial f'_{12}}{\partial r_{12}}, \quad (9)$$

so that the difficult double commutator was replaced by the product of the derivatives of the correlation factors. Thus, Eq. (6) takes now a more explicit form

$$I_6 = (ab) \left. \frac{\partial f_{12}}{\partial r_{12}} \frac{\partial f'_{12}}{\partial r_{12}} \right| cd. \quad (10)$$

The above rearrangements are valid for any correlation factors, f_{12} and f'_{12} , for which the initial integrals exist.

B. Application of the McMurchie-Davidson scheme

Once the simplifications introduced in the previous section have been acknowledged, we can attack the integrals (1)-(6) by using the McMurchie-Davidson (MD) scheme. Note that our presentation of this well-established technique is necessarily brief and we consequently do not immerse into the issues which are already solved in the literature. For a more elaborate discussion of the MD method we address the reader to the original papers^{49,50}, or the excellent review of Gill⁵⁹.

The main idea behind the MD method is the following expansion of the product of two Cartesian Gaussian orbitals

$$\begin{aligned} \Phi(\mathbf{r}_1, a, \mathbf{A}) \Phi(\mathbf{r}_1, b, \mathbf{B}) &= \sum_{t=0}^{i+j} E_t^{ij} \sum_{u=0}^{k+l} E_u^{kl} \sum_{v=0}^{m+n} E_v^{mn} \Lambda_{tuv}(\mathbf{r}_1, p, \mathbf{P}) = \\ &= \sum_{tuv} E_{tuv}^{ab} \Lambda_{tuv}(\mathbf{r}_1, p, \mathbf{P}), \end{aligned} \quad (11)$$

where the latter formulae is simply a shorthand notation for the former. In the above expression the so-called Hermite Gaussian functions were introduced

$$\Lambda_{tuv}(\mathbf{r}_1, p, \mathbf{P}) = \frac{d^t}{dP_x^t} \frac{d^u}{dP_y^u} \frac{d^v}{dP_z^v} \exp(-pr_{1P}^2), \quad (12)$$

with the new exponent, $p = a + b$, and the displaced centre, $\mathbf{P} = \frac{a\mathbf{A}+b\mathbf{B}}{p}$. The coefficients E_t^{ij} obey the following recursion relations which can be used to calculate them efficiently

$$E_0^{00} = \exp\left(-\frac{ab}{a+b}(A_x - B_x)^2\right), \quad (13)$$

$$E_0^{i+1,j} = -\frac{b}{p}(A_x - B_x)E_0^{ij} + E_1^{ij}, \quad (14)$$

$$E_0^{i,j+1} = \frac{a}{p}(A_x - B_x)E_0^{ij} + E_1^{ij}E_0, \quad (15)$$

$$E_t^{ij} = \frac{1}{2pt} (iE_{t-1}^{i-1,j} + jE_{t-1}^{i,j-1}), \quad (16)$$

and analogous for E_u^{kl} and E_v^{mn} . Of course, a very similar formulae hold for the product of orbitals of the second electron

$$\begin{aligned} \Phi_c(\mathbf{r}_2, c, \mathbf{C})\Phi_d(\mathbf{r}_2, d, \mathbf{D}) &= \sum_{t'=0}^{i'+j'} E_{t'}^{i'j'} \sum_{u'=0}^{k'+l'} E_{u'}^{k'l'} \sum_{v'=0}^{m'+n'} E_{v'}^{m'n'} \Lambda_{t'u'v'}(\mathbf{r}_2, q, \mathbf{Q}) = \\ &= \sum_{t'u'v'} E_{t'u'v'}^{cd} \Lambda_{t'u'v'}(\mathbf{r}_2, q, \mathbf{Q}), \end{aligned} \quad (17)$$

with $q = c + d$, $\mathbf{Q} = \frac{c\mathbf{C}+d\mathbf{D}}{q}$ and an identical recurrence structure for the primed indices. Let us now insert the expressions (11) and (17) into Eqs. (1)-(4) and (6), taking into account the simplified formula (10). The integral I_5 will be treated separately. Note that the differentiation with respect to the coordinates of \mathbf{P} can be replaced by the differentiation with respect to the components of $-\mathbf{Q}$, simply because of the translational invariance of the resulting expressions. After some rearrangements we arrive at the following formulae:

$$I_i = \sum_{tuv} E_{tuv}^{ab} (-1)^{t+u+v} \sum_{t'u'v'} E_{t'u'v'}^{cd} R_i^{t+t',u+u',v+v'}, \quad i = 1, 2, 3, 4, 6, \quad (18)$$

where

$$R_i^{tuv} = \frac{d^t}{dQ_x^t} \frac{d^u}{dQ_y^u} \frac{d^v}{dQ_z^v} B_i, \quad (19)$$

and B_i are the basic integrals that are given by the following generic expression

$$B_i = \iint d\mathbf{r}_1 d\mathbf{r}_2 e^{-pr_{1P}^2} k_i(r_{12}) e^{-qr_{2Q}^2}. \quad (20)$$

In the above equation $k_i(r_{12})$ are the kernel functions dependent solely on r_{12} and, in general, different for each $i = 1, 2, 3, 4, 6$. In the most general case, the kernel functions take the following form

$$k_1(r_{12}) = r_{12}^{-1}, \quad (21)$$

$$k_2(r_{12}) = r_{12}^{-1} f_{12}(r_{12}), \quad (22)$$

$$k_3(r_{12}) = f_{12}(r_{12}), \quad (23)$$

$$k_4(r_{12}) = f_{12}(r_{12}) f'_{12}(r_{12}), \quad (24)$$

$$k_6(r_{12}) = \frac{\partial f_{12}}{\partial r_{12}} \frac{\partial f'_{12}}{\partial r_{12}}. \quad (25)$$

Note that in some situations the functional forms of the kernels coincide or do not require a separate treatment. For instance, when $f_{12} = f'_{12} = r_{12}$ one has $k_2 = k_6 = 1$ and $k_4 = r_{12}^2$. Note additionally that in this example I_2 , I_4 and I_6 reduce to a combination of one-electron integrals.

As mentioned beforehand, the integral I_5 does not follow the scheme given above and requires a separate method. Let us recall Eq. (8) and follow the approach advocated by Helgaker *et al.*⁵⁰ in the context of the geometrical derivatives. Firstly, the differentiation with respect to the coordinates of \mathbf{A} and \mathbf{B} is rewritten as

$$\frac{d}{dA_x} = -\frac{a}{p} \frac{d}{dQ_x} + \frac{d}{dAB_x}, \quad (26)$$

$$\frac{d}{dB_x} = -\frac{b}{p} \frac{d}{dQ_x} - \frac{d}{dAB_x}, \quad (27)$$

where the vector \mathbf{AB} is simply $\mathbf{A} - \mathbf{B}$. With the above equations at hand, it becomes simple to express the operator $\frac{1}{2}(\nabla_{\mathbf{A}}^2 - \nabla_{\mathbf{B}}^2)$, present in Eq. (8), through the derivatives with respect to the components of \mathbf{Q} and \mathbf{AB} . Next, by applying it to the integral I_3 , given by Eq. (18) with $i = 3$, one obtains

$$\begin{aligned} I_5 = & \frac{a-b}{2p} \sum_{tuv} E_{tuv}^{ab} (-1)^{t+u+v} \sum_{t'u'v'} E_{t'u'v'}^{cd} \nabla_{\mathbf{Q}}^2 R_3^{t+t', u+u', v+v'} \\ & - \frac{a-b}{p} \sum_{tuv} \vec{\nabla}_{AB} E_{tuv}^{ab} (-1)^{t+u+v} \cdot \sum_{t'u'v'} E_{t'u'v'}^{cd} \vec{\nabla}_{\mathbf{Q}} R_3^{t+t', u+u', v+v'}, \end{aligned} \quad (28)$$

where $\vec{\nabla}_{\mathbf{Q}}$ and $\nabla_{\mathbf{Q}}^2$ are the gradient and Laplace operators with respect to \mathbf{Q} , and $\vec{\nabla}_{AB}$ stands for a gradient operator with respect to \mathbf{AB} coordinates. The main difficulty now lies

in the evaluation of the derivatives of R_3^{tuv} and the coefficients E_{tuv} . In the case of R_3^{tuv} one simply recalls Eq. (19) which leads to

$$\nabla_{\mathbf{Q}}^2 R_3^{tuv} = R_3^{t+2,uv} + R_3^{t,u+2,v} + R_3^{tu,v+2}, \quad (29)$$

$$\vec{\nabla}_{\mathbf{Q}} R_3^{tuv} = [R_3^{t+1,uv}, R_3^{t,u+1,v}, R_3^{tu,v+1}]. \quad (30)$$

Additionally, as pointed out by Samson *et al.*⁶³, the action of the $\vec{\nabla}_{\mathbf{AB}}$ on the E_{tuv}^{ab} can be resolved by the following recurrence relations:

$$F_0^{00} = -\frac{2ab}{p} AB_x E_0^{00}, \quad (31)$$

$$F_0^{i+1,j} = -\frac{b}{p} AB_x F_0^{ij} + F_1^{ij} - \frac{b}{p} E_0^{ij}, \quad (32)$$

$$F_0^{i,j+1} = \frac{a}{p} AB_x F_0^{ij} + F_1^{ij} + \frac{a}{p} E_0^{ij}, \quad (33)$$

$$F_t^{ij} = \frac{1}{2pt} (iF_{t-1}^{i-1,j} + jF_{t-1}^{i,j-1}), \quad (34)$$

where $F_t^{ij} = \frac{dE_t^{ij}}{dAB_x}$ and a similar notation (F_u^{kl} and F_v^{mn}) holds for the components y and z . The above expressions are the most straightforwardly obtained by a direct differentiation of Eqs. (13)-(16). One may notice, that in the second term of Eq. (28) both gradient operators produce vector quantities, but a scalar form remains afterwards as the dot product is taken. To sum up, the integral I_5 is also expressed as a linear combination of the quantities R_3^{tuv} .

Let us summarise the progress made in this section by the use of the MD scheme. Firstly, the integrals I_1 - I_4 and I_6 have been reduced to linear combinations of the R_i^{tuv} functions. We have shown that the latter quantities are the derivatives of B_i with respect to the components of \mathbf{Q} . Finally, the integral I_5 , resistant to the standard approach, has also been written down as a linear combination of R_i^{tuv} . All the coefficients appearing in the expressions obey simple recursive schemes and thus can be computed highly efficiently. Therefore, the remaining issues which need to be addressed are the evaluation of the basic integrals, B_i , and their differentiation with respect to the components of \mathbf{Q} . Note, that the approach adopted in this section is similar, but more general, as compared to the work of Samson *et al.*⁶³ However, at this point our derivations separate completely.

C. Basic integrals

In order to evaluate the quantities R_i^{tuv} , $i = 1, 2, 3, 4, 6$, one has to establish an analytical expression for the basic integrals, B_i , given by Eq. (20). The main difficulty lies in the fact that the form of the correlation factors, and thus the form of the kernels k_i , has not been specified yet. In typical situations, an analogue of the basic integral is evaluated with use of the integral transforms of the kernel functions. For instance, in the case when $k_1 = r_{12}^{-1}$ the following transformation can be adopted

$$r_{12}^{-1} = \sqrt{\frac{2}{\pi}} \int_0^\infty \exp(-t^2 r_{12}^2) dt, \quad (35)$$

and a similar one was used by Ten-no for the exponential correlation factor^{20,23}. In the case of an arbitrary correlation factor such an integral transform may not exist or can be overwhelmingly complicated. Thus, another line of attack needs to be devised. As already noted before, all basic integrals are translationally invariant. In other words, these integrals are dependent only on the length of the vector $\mathbf{P} - \mathbf{Q}$ but not on its individual components. Therefore, without any loss of generality, let us make a shift and consequently put $\mathbf{P} = 0$ further in the paper. The basic integrals now take the following form

$$B_i = \iint d\mathbf{r}_1 d\mathbf{r}_2 e^{-pr_1^2} k_i(r_{12}) e^{-qr_2^2}, \quad (36)$$

where the kernel functions, k_i , are defined exactly in the same way as in the previous subsection. Let us now recall the formula for translation of the 1s Gaussian function

$$e^{-qr_2^2} = 4\pi e^{-qQ^2} e^{-qr_2^2} \sum_{l=0}^{\infty} i_l(2qQr_2) \sum_{m=-l}^{+l} Y_{lm}^*(\hat{Q}) Y_{lm}(\hat{r}_2), \quad (37)$$

where $Q = |\mathbf{Q}|$, $i_l(z) = \sqrt{\frac{\pi}{2z}} I_{n+\frac{1}{2}}(z)$, $I_\nu(z)$ are the modified Bessel functions of the first kind⁷³, and $Y_{lm}(\hat{r})$ are the spherical harmonics⁷³. In fact, this formula is very well known to the quantum chemists community, as it is routinely used in the evaluation of the matrix elements of the effective core potentials (see, for example, the work of Kahn *et al.*⁷⁴). Let us now insert Eq. (37) into Eq. (36). An important observation here is that all terms, apart from $l = 0$ and $m = 0$, vanish due to the spherical symmetry of the integrand. Therefore, one is left with

$$B_i = e^{-qQ^2} \iint d\mathbf{r}_1 d\mathbf{r}_2 e^{-pr_1^2 - qr_2^2} \frac{\sinh(2qQr_2)}{2qQr_2} k_i(r_{12}), \quad (38)$$

where we have additionally used the well-known formula $i_0(z) = \sinh(z)/z$. Note that the integrand in Eq. (38) is essentially a one-centre distribution. Therefore, it is convenient to utilise the variables r_1 , r_2 and r_{12} supplemented with three Eulerian angles. After the change of variables and straightforward integration over the angles the following triple integral remains

$$B_i = 8\pi^2 \frac{e^{-qQ^2}}{2qQ} \int_0^\infty dr_{12} k_i(r_{12}) r_{12} \int_0^\infty dr_2 e^{-qr_2^2} \sinh(2qQr_2) \int_{|r_2-r_{12}|}^{r_2+r_{12}} dr_1 r_1 e^{-pr_1^2}. \quad (39)$$

The inner two integrations in the above expression are elementary. After some rearrangements one arrives at

$$B_i = \sqrt{\frac{\pi^5}{p+q}} \frac{1}{qp} \int_0^\infty dr_{12} k_i(r_{12}) r_{12} \left[\frac{e^{-\frac{pq}{p+q}(r_{12}-Q)^2} - e^{-\frac{pq}{p+q}(r_{12}+Q)^2}}{Q} \right]. \quad (40)$$

The above expression is universal *i.e.* it is valid for any kernel for which the initial integrals exist. Additionally, only the terms in the square brackets in Eq. (40) are dependent on Q . This fact is crucial for developments of the further sections.

D. Evaluation of the integrals R_i^{tuv}

The integrals R_i^{tuv} , defined formally as derivatives of the basic integrals, B_i , with respect to the components of \mathbf{Q} can now formally be calculated by a consecutive differentiation of Eq. (40) with respect to Q_x , Q_y and Q_z . However, this straightforward approach is quite cumbersome since the differentiation produces very lengthy expressions. Apart from that, the integrals B_i depend only on the length of \mathbf{Q} , but not on its individual components, which automatically suggests that considerable simplifications are possible.

Let us designate by $g(Q)$ an arbitrary well-behaved and sufficiently many times differentiable function, which is dependent solely on the length of the vector \mathbf{Q} . Now, the spherical tensor gradient operators, $\mathcal{Y}_{lm}(\bar{\nabla})$, are obtained by taking the explicit formulae for the regular solid spherical harmonics, $r^l Y_{lm}(\hat{r})$, and replacing the Cartesian coordinates, x , y , z , by the differentials, $\frac{\partial}{\partial x}$, $\frac{\partial}{\partial y}$, $\frac{\partial}{\partial z}$, respectively. A special case of the Hobson theorem^{75,76} reads

$$\mathcal{Y}_{lm}(\bar{\nabla}_Q)g(Q) = Q^l \left[\left(\frac{1}{Q} \frac{d}{dQ} \right)^l g(Q) \right] Y_{lm}(\hat{Q}). \quad (41)$$

Exactly the same formula is valid for the real versions of the spherical harmonics - one simply takes a combination of the above expression with m and $-m$. This simple observation

removes the need to use complex variables in actual computations. Concerning Eq. (41), see the works of Weniger and Steinborn^{77,78}, for a more elaborate discussion of the Hobson theorem.

The Hobson theorem makes it simple to operate with $\mathcal{Y}_{lm}(\bar{\nabla})$, but the original definition of R_i^{tuv} is given in terms of the derivatives with respect to Cartesian coordinates. Therefore, one has to relate an arbitrary differential with respect to the Cartesian coordinates with the spherical tensor gradient operators. Obviously, the transformation formulae are exactly the same as between Cartesian coordinates and the spherical harmonics, which follows directly from the definition of $\mathcal{Y}_{lm}(\bar{\nabla})$. In our notation, the necessary transformation reads

$$\frac{d^t}{dQ_x^t} \frac{d^u}{dQ_y^u} \frac{d^v}{dQ_z^v} = N(l, t, u, v) \sum_{l \leq t+u+v} c(l, m, t, u, v) \mathcal{Y}_{lm}(\bar{\nabla}). \quad (42)$$

in accordance with the work of Schlegel⁷⁹. For completeness, the analytical expressions for the coefficients $N(l, t, u, v)$ and $c(l, m, t, u, v)$ are given in Supplemental Material⁵⁴. The former are fairly simple and can be computed on the fly. The latter are very sparse *i.e.* a huge number of combinations of the parameters gives zero value. Therefore, they can be calculated in advance and stored in the memory for further use.

Making use of Eqs. (41) and (42) one can write

$$\begin{aligned} R_i^{tuv} &= \frac{d^t}{dQ_x^t} \frac{d^u}{dQ_y^u} \frac{d^v}{dQ_z^v} R_i^{000} = N(l, t, u, v) \sum_{l \leq t+u+v} c(l, m, t, u, v) \mathcal{Y}_{lm}(\bar{\nabla}) R_i^{000} = \\ &= N(l, t, u, v) \sum_{l \leq t+u+v} c(l, m, t, u, v) Q^l \left[\left(\frac{1}{Q} \frac{d}{dQ} \right)^l B_i \right] Y_{lm}(\hat{Q}), \end{aligned} \quad (43)$$

where the identity $R_i^{000} = B_i$ has been used. By inspection of the above expression one immediately recognises that only the derivatives of B_i with respect to Q are necessary, rather than with respect to Q_x , Q_y and Q_z separately. Evaluation of the derivatives with respect to Q is handled in the next subsection. Other quantities appearing in the above expression are the spherical harmonics but they can be efficiently computed with help of the well-known recursive formulae⁷³.

E. Evaluation of the radial derivatives

Due to the advances presented in the previous subsection it becomes clear that a major effort of the whole scheme lies in evaluation of the following terms

$$Q^l \mathcal{D}_Q^l B_i, \quad (44)$$

where the notation $\mathcal{D}_Q = \frac{1}{Q} \frac{d}{dQ}$ has been implied. In the evaluation of the differentials (44) an additional attention must be paid to the numerical issues. In fact, the Q -dependent factor in Eq. (40) consists of two terms which are of opposite signs and nearly equal in magnitude when the value of Q is small. Therefore, a significant digital erosion can be expected when Eq. (40) is evaluated as it stands for small Q . This problem is also largely magnified when a direct differentiation of Eq. (40) is attempted to calculate the radial derivatives, Eq. (44). Therefore, we need to adopt a special approach when the value of Q is small.

1. The case of small Q

Let us rewrite the Q -dependent part of the integrand in Eq. (40) as follows

$$\frac{e^{-\xi(r_{12}-Q)^2} - e^{-\xi(r_{12}+Q)^2}}{Q} = \frac{e^{-\xi(r_{12}^2+Q^2)}}{Q} \sinh(2\xi r_{12}Q) = \frac{e^{-\xi(r_{12}^2+Q^2)}}{Q} \sum_{n=0}^{\infty} \frac{(2\xi r_{12}Q)^{2n+1}}{(2n+1)!}, \quad (45)$$

where $\xi = pq/(p+q)$. The above infinite series is everywhere convergent. Therefore, it can be inserted back into Eq. (40) and the order of summation and integration can be interchanged. This leads to the following series expansion of B_i around $Q = 0$

$$B_i = 2\pi e^{-\xi Q^2} \left(\frac{\pi}{p+q} \right)^{3/2} \sum_{n=0}^{\infty} \frac{(2\xi Q)^{2n}}{(2n+1)!} \int_0^{\infty} dr_{12} k_i(r_{12}) r_{12}^{2n+2} e^{-\xi r_{12}^2}. \quad (46)$$

Passing to the evaluation of the differentials (44), it is now straightforward to apply the operator $Q^l \mathcal{D}_Q^l$ to both sides of Eq. (46) which gives

$$Q^l \mathcal{D}_Q^l B_i = 2\pi Q^l \left(\frac{\pi}{p+q} \right)^{3/2} \sum_{n=0}^{\infty} \frac{(2\xi)^{2n}}{(2n+1)!} \mathcal{D}_Q^l \left(Q^{2n} e^{-\xi Q^2} \right) \int_0^{\infty} dr_{12} k_i(r_{12}) r_{12}^{2n+2} e^{-\xi r_{12}^2}. \quad (47)$$

The action of \mathcal{D}_Q^l on the term $Q^{2n} e^{-\xi Q^2}$ is conveniently resolved by means of the following recursion relation

$$\mathcal{D}_Q^l \left(Q^{2n} e^{-\xi Q^2} \right) = n \mathcal{D}_Q^{l-1} \left(Q^{2n-2} e^{-\xi Q^2} \right) - 2\xi \mathcal{D}_Q^{l-1} \left(Q^{2n} e^{-\xi Q^2} \right) \quad (48)$$

which is sufficiently numerically stable for small Q . Let us conclude by noting that the series expansion (47) converges reasonably well provided that the values of Q and ξ are simultaneously small. In the opposite situation the approach developed in the next subsection is appropriate.

2. The case of moderate or large Q

In this subsection we shall attack the evaluation of the radial derivatives, $Q^l \mathcal{D}_Q^l B_i$, by means of the recursive techniques. Since the problem of small values of Q has already been resolved, we can perform separation of two terms in the square brackets in Eq. (40). To do so productively let us introduce the following quantities

$${}^i \mathcal{H}_{jkl}^\pm = \sqrt{\frac{\pi^5}{p+q}} \frac{1}{qp} \int_0^\infty dr_{12} r_{12}^j k_i(r_{12}) \mathcal{D}_Q^l \left(\frac{e^{-\xi(r_{12} \pm Q)^2}}{Q^k} \right), \quad (49)$$

so that

$$Q^l \mathcal{D}_Q^l B_i = Q^l \left({}^i \mathcal{H}_{11l}^- - {}^i \mathcal{H}_{11l}^+ \right). \quad (50)$$

We assume here that the values of ${}^i \mathcal{H}_{j00}^\pm$ are available at the start and their evaluation will be considered in details further. It is obvious that ${}^i \mathcal{H}_{jk0}^\pm$ are trivial to obtain from the initial ${}^i \mathcal{H}_{j00}^\pm$, so that our task is to increase the value of l at the cost of j and k . Let us first inspect the action of \mathcal{D}_Q on the Q -dependent part of the integrand in Eq. (49) with $l = 0$

$$\mathcal{D}_Q \left(\frac{e^{-\xi(r_{12} \pm Q)^2}}{Q^k} \right) = \mp 2\xi r_{12} \frac{e^{-\xi(r_{12} \pm Q)^2}}{Q^{k+1}} - 2\xi \frac{e^{-\xi(r_{12} \pm Q)^2}}{Q^k} - k \frac{e^{-\xi(r_{12} \pm Q)^2}}{Q^{k+2}}. \quad (51)$$

Let us now multiply both sides of the above expression by $r_{12}^j k_i(r_{12})$ and integrate over r_{12} on the interval $[0, +\infty]$. One obtains the following relation

$${}^i \mathcal{H}_{jk1}^\pm = \mp 2\xi {}^i \mathcal{H}_{j+1,k+1,0}^\pm - 2\xi {}^i \mathcal{H}_{jk0}^\pm - k {}^i \mathcal{H}_{j,k+2,0}^\pm, \quad (52)$$

which can be further generalised by applying \mathcal{D}_Q^{l-1} to both sides

$${}^i \mathcal{H}_{jkl}^\pm = \mp 2\xi {}^i \mathcal{H}_{j+1,k+1,l-1}^\pm - 2\xi {}^i \mathcal{H}_{jk,l-1}^\pm - k {}^i \mathcal{H}_{j,k+2,l-1}^\pm, \quad (53)$$

since the coefficients in this expression are independent of Q . This gives us the desired recursion relation which can be used to calculate ${}^i \mathcal{H}_{11l}^\pm$ starting only with ${}^i \mathcal{H}_{jk0}^\pm$. Note that

the recursive process (53) is rather simple, despite operating in three dimensions, and can be implemented efficiently. Therefore, the presented recursive method offers an attractive way for the evaluation of the radial derivatives.

No progress can be made now unless a concrete functional form of the correlation factor is specified. In the next section we apply the theory developed here to the case of the range-separated correlation factor. However, before passing further let us observe that ${}^i\mathcal{H}_{j00}^\pm$ can be reduced to well-known quantities when the kernel $k_i(r_{12})$ is a familiar function. For instance, when $k_i(r_{12}) = 1/r_{12}$ *i.e.* for the standard electron repulsion integrals, we obtain up to a trivial multiplicative constant, ${}^i\mathcal{H}_{j+1,00}^\pm \sim \int_0^\infty dr_{12} r_{12}^j e^{-\xi(r_{12}\pm Q)^2}$. These integrals are reduced to a combination of the so-called Boys functions⁸⁰, $F_n(z)$, with different arguments. Therefore, we conclude that for a familiar kernel function our scheme requires no new starting quantities, as compared with the dedicated procedures which are in a widespread use.

III. THE RANGE-SEPARATED CORRELATION FACTOR

A. Preliminaries

Let us now consider the so-called range-separated correlation factor²⁹

$$f_{12}(r_{12}; c_0, \rho, B, \mu) = \left(1 + \frac{1}{2}r_{12}\right) e^{-\mu r_{12}^2} + S_n(\mu r_{12}^2) c_0 r_{12}^\rho e^{Br_{12}}, \quad (54)$$

where $c_0, \rho, B, \mu > 0$ are variational parameters and

$$S_n(x) = 1 - e^{-x} \sum_{k=0}^n \frac{x^k}{k!}, \quad (55)$$

is the Tang-Toennies damping function⁸¹. The integer n is assumed to be greater than $-\frac{\rho}{2} - 1$, if $\rho < -2$, or $n = 0$ otherwise [note that $S_n(x) = \mathcal{O}(x^{n+1})$]. It is clear that the term $\left(1 + \frac{1}{2}r_{12}\right) e^{-\mu r_{12}^2}$ is responsible for the short-range correlation and necessary to fulfil the cusp condition. Even greater flexibility can be achieved if the exponential formula of Ten-no is used in the short-range regime²⁹. The long-range part $r_{12}^\rho e^{Br_{12}}$ represents the correct asymptotics, as found for the helium atom²⁹. The role of the Gaussian and Tang-Toennies damping functions is to interpolate smoothly between these two regimes, providing at the same time a large degree of flexibility. It is important to note that the parameters ρ, B can be arbitrary real numbers. In particular, it was found for the helium atom that the value of

B is always positive and the value of ρ depends significantly on the orbital part of the wave function. In the case when ρ is negative the damping function S_n removes the corresponding singularity as $r_{12} \rightarrow 0$. This is the reason why the aforementioned constraint on n strictly holds.

Before the theory developed in the previous section can be applied, one mathematical difficulty needs to be addressed. Let us consider the kernel $k_3(r_{12})$ given by Eq. (23). To calculate the necessary molecular integrals by using the recursive method, one starts with the quantities ${}^3\mathcal{H}_{j00}^\pm$. One can formally insert the explicit form of the correlation factor, Eq. (54), into the expression (49). After several simplifications, this leads to the formula

$${}^3\mathcal{H}_{j00}^\pm = \mathcal{N} \left[\int_0^\infty dr_{12} r_{12}^j e^{-(\mu+\xi)r_{12}^2} e^{\mp 2\xi r_{12}Q} + \frac{1}{2} \int_0^\infty dr_{12} r_{12}^{j+1} e^{-(\mu+\xi)r_{12}^2} e^{\mp 2\xi r_{12}Q} \right. \\ \left. + c_0 \int_0^\infty dr_{12} r_{12}^{j+\rho} S_n(\mu r_{12}^2) e^{(B \mp 2\xi Q)r_{12}} \right], \quad (56)$$

where $\mathcal{N} = \sqrt{\frac{\pi^5}{p+q} \frac{e^{-\xi Q^2}}{qp}}$. An obvious move now is to insert the explicit form of the damping function, Eq. (55), into the above expression. However, the summation present in Eq. (55) cannot be straightforwardly interchanged with the integration in Eq. (56) since some of the resulting integrals can become singular. Therefore, we need to change the integration range from $[0, +\infty]$ to $[\varepsilon, +\infty]$ and take the limit $\varepsilon \rightarrow 0^+$ once the separation of the individual terms is performed. This gives

$${}^3\mathcal{H}_{j00}^\pm = \mathcal{N} \left[\int_0^\infty dr_{12} r_{12}^j e^{-(\mu+\xi)r_{12}^2} e^{\mp 2\xi r_{12}Q} + \frac{1}{2} \int_0^\infty dr_{12} r_{12}^{j+1} e^{-(\mu+\xi)r_{12}^2} e^{\mp 2\xi r_{12}Q} \right. \\ \left. + c_0 \lim_{\varepsilon \rightarrow 0^+} \left(\int_\varepsilon^\infty dr_{12} r_{12}^{j+\rho} e^{(B \mp 2\xi Q)r_{12}} - \sum_{k=0}^n \frac{\mu^k}{k!} \int_\varepsilon^\infty dr_{12} r_{12}^{j+\rho+2k} e^{(B \mp 2\xi Q)r_{12}} e^{-\mu r_{12}^2} \right) \right]. \quad (57)$$

These findings suggest that the all necessary integrals can be expressed through the following family of functions

$$S(\alpha, \beta, \gamma) = \Re \int_0^\infty dx x^\alpha \exp(\beta x - \gamma x^2), \quad (58)$$

where α, β are arbitrary real numbers and $\gamma \geq 0$ and \Re denotes the finite (regular) part of the integral. The precise meaning of the symbol \Re is defined as follows. When $\alpha > -1$ the integrals (58) are regular and can be treated with standard methods. For $\alpha \leq -1$ they are divergent and have to be understood differently. Therefore, when $\alpha \leq -1$ we proceed

as follows. First, the integration range is changed to $[\varepsilon, +\infty]$ and this modified integral is evaluated. Next, the small ε expansion is considered and one obtains an analytical result which consist of a regular, ε -independent part of the integral plus terms which contain ε . The latter terms split into two groups: vanishing as the limit $\varepsilon \rightarrow 0$ is taken and singular terms which are *finite* combinations of inverse powers of ε and possibly logarithms of ε . In the present regularisation scheme all singular terms are simply dropped and the limit $\varepsilon \rightarrow 0$ is taken. The result is a finite, regular part of the otherwise diverging integrals. This approach is justified because in all expressions for the relevant integrals, such as Eq. (57), the integrals S appear in specific combinations, inherited over the form of the damping functions. In these combinations, all singular terms must cancel out since the expressions for the initial molecular matrix elements are regular. In other words, there is no need to keep track of the terms containing ε because we know in advance that they must cancel out exactly when the final expression, such as (57), is evaluated.

As a simple example, let us consider $S(\alpha, \beta, \gamma)$ with $\alpha = -2$, $\gamma = 0$ and $\beta < 0$. Integration over the range $[\varepsilon, +\infty]$, followed by a small ε expansion gives

$$\int_{\varepsilon}^{\infty} dx \frac{e^{\beta x}}{x^2} = -\beta \left[\gamma_E + \log(-\beta) - 1 \right] + \frac{1}{\varepsilon} - \beta \log(\varepsilon) - \frac{\beta^2}{2} \varepsilon + \mathcal{O}(\varepsilon^2), \quad (59)$$

where γ_E is the Euler-Mascheroni constant. One can see that the exemplary integral consists of the regular part (terms in the square bracket), a finite number of singular terms ($\frac{1}{\varepsilon}$ and $\log \varepsilon$) and an infinite number of terms which vanish as $\varepsilon \rightarrow 0$. As the singular terms are dropped and the limit $\varepsilon \rightarrow 0$ is taken one obtains in our notation

$$S(-2, \beta, 0) = \Re \int_0^{\infty} dx \frac{e^{\beta x}}{x^2} = -\beta \left[\gamma_E + \log(-\beta) - 1 \right]. \quad (60)$$

Clearly, this technique is a fairly straightforward way to manage the spurious singularities.

Let us note in passing that in practical situations the form of the correlation factor is fixed *i.e.* one adopts some recommended values of the parameters in Eq. (54). By recalling Eq. (57) as an example, one finds that in actual computations a set of basic integrals in the form $S(\alpha_0 + N, \beta, \gamma)$ is required, where α_0 is a real parameter, $0 < \alpha_0 < 1$, and N is an arbitrary integer. The values of β and γ depend on the nonlinear parameters of the GTOs, whilst α_0 is inherited over the adopted formula for f_{12} . This observation is rather obvious, but important in the context of the next subsections where calculation of $S(\alpha, \beta, \gamma)$ is discussed in details.

In Supplemental Material⁵⁴ we give the analytical forms of the kernel functions, k_i , with the range-separated correlation factor, and express the corresponding basic integrals, B_i , through the $S(\alpha, \beta, \gamma)$. These expressions are long but very regular, and can be turned into an efficient computer code.

B. Calculation of $S(\alpha, \beta, \gamma)$ for $\alpha > -1$

In the case when the value of α is larger than -1 the integrals $S(\alpha, \beta, \gamma)$ can be reduced to the Tricomi's confluent hypergeometric function, $\mathcal{U}(a, b, z)$, by a simple variable transformation $t = \sqrt{2\gamma}x$. In this way one obtains

$$S(\alpha, \beta, \gamma) = (2\sqrt{\gamma})^{-\alpha-1} \Gamma(\alpha + 1) \mathcal{U}\left(\frac{\alpha + 1}{2}, \frac{1}{2}, \frac{\beta^2}{4\gamma}\right). \quad (61)$$

At this point the problem is solved because methods of calculation of $\mathcal{U}(a, b, z)$ for arbitrary real (or even complex) values of the parameters a , b , and z exist^{82,83}. However, here we deal with an exceptionally special case of $\mathcal{U}(a, b, z)$ where $b = \frac{1}{2}$, $a > 0$ and $z > 0$. This signals that a dedicated procedure for the calculation of $\mathcal{U}(a, b, z)$ in this special case should be designed. In fact, the general algorithm is considerably more complicated and connected with a large computational overhead. Our method for the evaluation of $\mathcal{U}(a, 1/2, z)$ with $a > 0$, $z > 0$ is presented in Supplemental Material⁵⁴. To utilise the expression (61) it is also necessary to compute the Euler Gamma function, $\Gamma(z)$, for the positive real values of z but efficient methods exist⁷³.

C. Calculation of $S(\alpha, \beta, \gamma)$ for $\alpha \leq -1$

A slight inconvenience connected with the adopted regularisation scheme for the integrals $S(\alpha, \beta, \gamma)$ is that they are not continuous as a function of α . Strictly speaking, there are discontinuities for each negative integer value of α . As a result, negative integer and noninteger values of α have to be treated separately which is pursued in the next subsections. Let us note, however, that once the summation such as in Eq. (57) is performed, the result is a continuous function of all parameters, as it should be, and as we also checked numerically.

1. *Noninteger values of α*

By integrating Eq. (58) by parts and dropping the singular terms one obtains the following recursion relation

$$S(\alpha - 1, \beta, \gamma) = -\frac{\beta}{\alpha}S(\alpha, \beta, \gamma) + \frac{2\gamma}{\alpha}S(\alpha + 1, \beta, \gamma), \quad (62)$$

when α is a noninteger number, $\alpha < -1$. At this point it becomes advantageous to introduce a slightly simplified family of integrals

$$T_s^\pm(x) = \Re \int_0^\infty \frac{dz}{z^s} e^{\pm z - xz^2}, \quad (63)$$

with $x > 0$ and (noninteger) real $s > -1$. The above integrals are related to the initial $S(\alpha, \beta, \gamma)$ by an elementary exchange of variables, giving

$$S(\alpha, \beta, \gamma) = |\beta|^{-\alpha-1} T_{-\alpha}^{\text{sgn } \beta}(\gamma/\beta^2), \quad (64)$$

where sgn denotes the sign function. Starting with Eq. (62) it becomes simple to show that the integrals $T_s^\pm(x)$ obey the following recursion relation

$${}_s T_{s+1}^\pm(x) = \pm T_s^\pm(x) - 2x T_{s-1}^\pm(x), \quad (65)$$

which is more straightforward to handle than the initial Eq. (62). Let us now consider numerical evaluation of $T_s^\pm(x)$ with the help of the recursive process (65). Firstly, one requires two starting values, $T_{s_0}^\pm(x)$ and $T_{s_0-1}^\pm(x)$, with $s_0 \in (0, 1)$. These integrals are regular and can be evaluated with methods described in the previous section. Namely, one has

$$T_{s_0}^\pm(x) = (2\sqrt{x})^{s_0-1} \Gamma(1 - s_0) \mathcal{U}\left(\frac{1 - s_0}{2}, \frac{1}{2}, \frac{1}{4x}\right), \quad (66)$$

and similarly for the second starting value. The next issue that needs to be addressed is numerical stability of the recursive process (65). We tested the reliability of Eq. (65) for the calculation of $T_s^\pm(x)$ up to $s = 30$ which is sufficient for the practical purposes, and for a broad range of x . The general conclusion, both for T_s^+ and T_s^- , is that Eq. (65) is numerically satisfactory only when $x > 1$ and a significant loss of digits is observed for the smaller values of x . However, it is natural to expect that for $x < 1$ the downward recursion is stable, so let us rewrite Eq. (65) as follows:

$$r_{s-1}^\pm(x) = \frac{2x}{\pm 1 - s r_s^\pm(x)}, \quad (67)$$

where $r_s^\pm(x) = T_{s+1}^\pm(x)/T_s^\pm(x)$ is the Miller-like quotient⁸⁴. Since r_s^\pm vanishes quickly with increasing s one can initiate the downward recursion at some large s with $r_s^\pm = 0$. The recursion is carried out until a value, s_0 , is reached for which $0 < s_0 < 1$. The desired quantities T_s^\pm are then recovered consecutively by using the definition of r_s^\pm *i.e.* $T_{s_0+1}^\pm = r_{s_0}^\pm T_{s_0}^\pm$, $T_{s_0+2}^\pm = r_{s_0+1}^\pm T_{s_0+1}^\pm$ *etc.*, where the starting value, $T_{s_0}^\pm$, is given by Eq. (66).

Let us also note that for β equal to zero, which corresponds to $S(\alpha, 0, \gamma)$, the strategy adopted in this subsection is no longer valid since the relation Eq. (64) becomes ill-defined. However, one can return to Eq. (62) which immediately leads to

$$S(\alpha, 0, \gamma) = \frac{1}{2} \frac{\Gamma(\frac{\alpha+1}{2})}{\gamma^{\frac{\alpha+1}{2}}}, \quad (68)$$

by imposing proper initial conditions.

2. Integer values of α

Similarly as for the noninteger values of α , one starts with the definition, Eq. (58), and integrates it by parts. However, as now the values of α are restricted to the negative integers, boundary terms in the integration by parts give ε -independent, finite contributions. Because of that, the corresponding recursion relation is inhomogeneous

$$S(\alpha - 1, \beta, \gamma) = \sum_{l=0}^{-\alpha/2} \frac{\beta^{-\alpha-2l}}{(-\alpha - 2l)!} \frac{(-\gamma)^l}{l!} - \frac{\beta}{\alpha} S(\alpha, \beta, \gamma) + \frac{2\gamma}{\alpha} S(\alpha + 1, \beta, \gamma). \quad (69)$$

With formally the same definition of $T_s^\pm(x)$, Eq. (63), one can derive the corresponding recursion

$${}_s T_{s+1}^\pm(x) = \pm \sum_{l=0}^{s/2} \frac{(-x)^l}{l!(s-2l)!} \pm T_s^\pm(x) - 2x T_{s-1}^\pm(x), \quad (70)$$

which is more compact and numerically convenient. To initiate the recursive process one needs two starting values, $T_1^\pm(x)$ and $T_0^\pm(x)$. The expression for the latter is rather simple

$$T_0^\pm(x) = \sqrt{\frac{\pi}{4x}} e^{\frac{1}{4x}} \left[1 \pm \text{Erf} \left(\frac{1}{\sqrt{2x}} \right) \right], \quad (71)$$

where $\text{Erf}(\ast)$ is the error function. To bring the former integral into a more convenient form let us introduce the following family of functions

$$\omega_k(x) = \int_0^\infty dz z^k \log(z) e^{-xz-z^2}. \quad (72)$$

With help of these integrals $T_1^\pm(x)$ can be rewritten as

$$T_1^\pm(x) = \mp \omega_0 \left(\pm \frac{1}{\sqrt{x}} \right) + 2\omega_1 \left(\pm \frac{1}{\sqrt{x}} \right). \quad (73)$$

Note, that in contrast with the left-hand-side of the above expression, the integrals present on the right-hand-side are regular and can be understood in the ordinary sense. We developed a dedicated procedure for the calculation of the integrals $\omega_0(x)$ and $\omega_1(x)$, which is based on the series expansions and polynomial interpolation. All necessary expressions are given in Supplemental Material⁵⁴, along with comments and a sketch of the algorithm.

The remaining issue is the numerical stability of the recursive process (70). For $x > 1$ the expression (70) can be used as it stands *i.e.* in the direction of increasing s . However, for $x < 1$ it is necessary to apply the downward recursion. Unfortunately, since the recursion (70) is inhomogeneous, a trick similar as in Eq. (67) cannot be utilised here. Therefore, one has to start at some large s_{max} (significantly larger than the maximal desired s) with $T_{s_{max}+1}^\pm(x) = 0$ and $T_{s_{max}}^\pm(x) = 1$, and carry out the recursion downward (70) until $s = 0$ is reached. A slight inconvenience connected with this approach is that the results of the recursion are not directly $T_s^\pm(x)$, but rather $T_s^\pm(x)$ multiplied by a constant (the same for each s). To determine this constant one simply evaluates $T_0^\pm(x)$ independently by using Eq. (71), compares with the corresponding result of the downward recursion and rescales all values properly.

IV. CONCLUSIONS

We have presented how six types of molecular integrals, necessary for the explicitly correlated electronic structure calculations, can be reduced to simple quantities. We have applied the McMurchie-Davidson scheme first, and reduced all two-electron integrals to combinations of the derivatives of the basic integrals. The differentiations were carried out, without specifying the correlation factor, with the help of the Hobson theorem. Finally, the resulting radial derivatives were evaluated recursively starting with a class of ordinary one-dimensional integrals.

When the developed scheme is applied to the range-separated correlation factor, all basic quantities can be written down as linear combinations of the integrals $S(\alpha, \beta, \gamma)$. Their calculation is not trivial and we have discussed in details the necessary numerical procedures

applicable for specific values of the parameters. We would like to stress that virtually all novel expressions presented in this work were extensively checked with the help of the symbolic environment of MATHEMATICA package⁸⁵. Implementation of the production code is in progress.

The present work opens up a possibility for the practical use of the range-separated factor in explicitly correlated calculations on many-electron atoms and molecules. We believe that the introduction of this new correlation factor will noticeably improve the level of accuracy which can routinely be reached with these methods. Extensive benchmarks will be reported shortly.

ACKNOWLEDGMENTS

This work was supported by the Polish Ministry of Science and Higher Education, grant NN204 182840. RM acknowledges the Foundation for Polish Science for the support through the “*Mistrz*” program. We would like to thank Bogumił Jeziorski for discussions, as well as reading and commenting on the manuscript.

REFERENCES

- ¹T. Kato, Commun. Pure Appl. Math. **10**, 151 (1957).
- ²R. T. Pack and W. Byers Brown, J. Chem. Phys. **45**, 556 (1966).
- ³S. Furnais, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, T. Ø. Sørensen, Commun. Math. Phys. **255**, 183 (2005).
- ⁴M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, H. Stremnitzer, Phys. Rev. Lett. **68**, 3857 (1992).
- ⁵C. Hättig, W. Klopper, A. Köhn, and D. P. Tew, Chem. Rev. **112**, 4 (2012).
- ⁶L. Kong, F. A. Bischoff, and E. F. Valeev, Chem. Rev. **112**, 75 (2012).
- ⁷S. Ten-no, Theor. Chem. Acc. **131**, 1070 (2012).
- ⁸W. Kutzelnigg, Theor. Chim. Acta **68**, 445 (1985).
- ⁹W. Klopper and W. Kutzelnigg, Chem. Phys. Lett. **134**, 17 (1987).
- ¹⁰W. Kutzelnigg and W. Klopper, J. Chem. Phys. **94**, 1985 (1991).
- ¹¹J. Noga, W. Kutzelnigg, and W. Klopper, Chem. Phys. Lett. **199**, 497 (1992).
- ¹²J. Noga and W. Kutzelnigg, J. Chem. Phys. **101**, 7738 (1994).

- ¹³W. Klopper and J. Noga, in *Explicitly correlated functions in chemistry and physics. Theory and applications*, edited by J. Rychlewski (Kluwer, Dordrecht, 2003), p. 149.
- ¹⁴K. Szalewicz and B. Jeziorski, *Mol. Phys.* **108**, 3091 (2010).
- ¹⁵K. Szalewicz, B. Jeziorski, H. J. Monkhorst, and J. G. Zabolitzky, *Chem. Phys. Letters* **91**, 169-172 (1982).
- ¹⁶R. Bukowski, B. Jeziorski, and K. Szalewicz, *J. Chem. Phys.* **110**, 4165-4183 (1999).
- ¹⁷R. Bukowski, B. Jeziorski, and K. Szalewicz, in *Explicitly correlated functions in chemistry and physics. Theory and applications*, edited by J. Rychlewski (Kluwer, Dordrecht, 2003), p. 185.
- ¹⁸A. J. May, E. Valeev, R. Polly, and F. R. Manby, *Phys. Chem. Chem. Phys.* **7**, 2710 (2005).
- ¹⁹A. J. May and F. R. Manby, *J. Chem. Phys.* **121**, 4479 (2004).
- ²⁰S. Ten-no, *Chem. Phys. Lett.* **398**, 56 (2004).
- ²¹T. Shiozaki, G. Knizia, and H.-J. Werner, *J. Chem. Phys.* **134**, 034113 (2011).
- ²²T. Shiozaki and H.-J. Werner, *J. Chem. Phys.* **134**, 184104 (2011).
- ²³S. Ten-no, *Chem. Phys. Lett.* **447**, 185 (2007).
- ²⁴M. Torheyden and E. F. Valeev, *J. Chem. Phys.* **131**, 171103 (2009).
- ²⁵T. Shiozaki and H.-J. Werner, *J. Chem. Phys.* **133**, 141103 (2010).
- ²⁶S. Kedzuch, O. Demel, J. Pittner, S. ten-no, and J. Noga, *Chem. Phys. Lett.* **511**, 418 (2011).
- ²⁷T. J. Martinez and S. Varganov, *J. Chem. Phys.* **132**, 054103 (2010).
- ²⁸D. P. Tew and W. Klopper, *J. Chem. Phys.* **123**, 074101 (2005).
- ²⁹M. Lesiuk, B. Jeziorski, and R. Moszynski, *J. Chem. Phys.* **139**, 134102 (2013).
- ³⁰K. A. Peterson, T. B. Adler, and H. J. Werner, *J. Chem. Phys.* **128**, 084102 (2008).
- ³¹J. G. Hill and K. A. Peterson, *Phys. Chem. Chem. Phys.* **12**, 10460 (2010).
- ³²J. G. Hill, S. Mazumder, K. A. Peterson, *J. Chem. Phys.* **132**, 054108 (2010).
- ³³A. Lüchow and J. B. Anderson, *Annu. Rev. Phys. Chem.* **51**, 501 (2000).
- ³⁴W. M. C Foulkes, L. Mitas, R. J. Needs, and G. Rajagopal, *Rev. Mod. Phys.* **73**, 33 (2001).
- ³⁵B. M. Austin, D. Y. Zubarev and W. A. Lester, *Chem. Rev.* **112**, 263 (2012).
- ³⁶C. Diedrich, A. Lüchow, and S. Grimme, *J. Chem. Phys.* **123**, 184106 (2005).
- ³⁷M. Korth, A. Lüchow and S. Grimme, *J. Phys. Chem. A* **112**, 2104 (2008).
- ³⁸M. Dubecký, P. Jurečka, R. Derian, P. Hobza, M. Otyepka, and L. Mitas, *J. Chem. Theory Comput.* **9**, 4287 (2013).
- ³⁹T. D. Beaudet, M. Casula, J. Kim, S. Sorella, and R. M. Martin, *J. Chem. Phys.* **129**, 164711

- (2008).
- ⁴⁰P. T. A. Galeka, N. C. Handy, and W. A. Lester Jr., *Mol. Phys.* **104**, 3069 (2007).
- ⁴¹N. A. Benedek, I. K. Snook, M. D. Towler, and R. J. Needs, *J. Chem. Phys.* **125**, 104302 (2006).
- ⁴²M. Korth, S. Grimme, and M. D. Towler, *J. Phys. Chem. A* **115**, 11734 (2011).
- ⁴³M. Dubecký, R. Derian, L. Mitas, and I. Štich, *J. Chem. Phys.* **133**, 244301 (2010).
- ⁴⁴M. Dubecký, R. Derian, P. Jurecka, L. Mitas, P. Hobza, and M. Otyepka, *Phys. Chem. Chem. Phys.* **16**, 20915 (2014).
- ⁴⁵L. Horváthová, M. Dubecký, L. Mitas, I. Štich, *J. Chem. Theory Comp.* **9**, 390 (2013).
- ⁴⁶M. Dupuis, J. Rys and H. F. King, *J. Chem. Phys.* **65**, 111 (1976).
- ⁴⁷H. F. King and M. Dupuis, *J. Comput. Phys.* **21**, 144 (1976).
- ⁴⁸J. Rys, M. Dupuis and H. F. King, *J. Comput. Chem.* **4**, 154 (1983).
- ⁴⁹L. E. McMurchie and E. R. Davidson, *J. Comput. Phys.* **26**, 218 (1978).
- ⁵⁰T. Helgaker and P. R. Taylor, *Theor. Chim. Acta* **83**, 177 (1992).
- ⁵¹J. A. Pople and W. J. Hehre, *J. Comput. Phys.* **27**, 161 (1978).
- ⁵²S. Obara and A. Saika, *J. Chem. Phys.* **84**, 3963 (1986).
- ⁵³H. B. Schlegel, *J. Chem. Phys.* **77**, 3676 (1982).
- ⁵⁴See supplemental material at [URL will be inserted by AIP] for additional derivations mentioned in the main text.
- ⁵⁵M. Head-Gordon and J. A. Pople, *J. Chem. Phys.* **89**, 5777 (1988).
- ⁵⁶R. Lindh, U. Ryu and B. Liu, *J. Chem. Phys.* **95**, 5889 (1991).
- ⁵⁷T. P. Hamilton and H. F. Schaefer, *Chem. Phys.* **150**, 163 (1991).
- ⁵⁸B. G. Johnson, P. M. W. Gill and J. A. Pople, *Chem. Phys. Lett.* **206**, 229 (1993).
- ⁵⁹P. M. W. Gill, *Adv. Quantum Chem.* **25**, 141 (1994).
- ⁶⁰W. Klopper and R. Röhse, *Chem. Phys. Lett.* **261**, 35 (1996).
- ⁶¹M. J. Bearpart, N. C. Handy, R. D. Amos, P. E. Maslen, *Theor. Chim. Acta.* **79**, 361 (1991).
- ⁶²E. F. Valeev and H. F. Schaefer III, *J. Chem. Phys.* **113**, 3990 (2000).
- ⁶³C. C. M. Samson, W. Klopper, T. Helgaker, *Comput. Phys. Comm.* **149**, 1 (2002).
- ⁶⁴V. Weber and C. Daul, *Comput. Phys. Comm.* **158**, 1 (2004).
- ⁶⁵R. Ahlrichs, *Phys. Chem. Chem. Phys.* **8**, 3072 (2006).
- ⁶⁶W. Klopper and C. C. M. Samson, *J. Chem. Phys.* **116**, 6297 (2002).
- ⁶⁷S. Ten-no and F. R. Manby, *J. Chem. Phys.* **119**, 5358 (2003).

- ⁶⁸E. F. Valeev, Chem. Phys. Lett. **395**, 190 (2004).
- ⁶⁹B. J. Persson and P. R. Taylor, J. Chem. Phys. **105**, 5915 (1996).
- ⁷⁰B. J. Persson and P. R. Taylor, Theor. Chem. Acc. **97**, 240 (1997).
- ⁷¹P. Dahle and P. R. Taylor, Theor. Chem. Acc. **105**, 401 (2001).
- ⁷²S. Ten-no, J. Chem. Phys. **121**, 117 (2004).
- ⁷³M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, (Dover, New York, 1972).
- ⁷⁴L. R. Kahn, P. Baybutt and D. G. Truhlar, J. Chem. Phys. **65**, 3826 (1976).
- ⁷⁵E. W. Hobson, *The Theory of Spherical and Ellipsoidal Harmonics*, Chelsea, New York, 1965.
- ⁷⁶E. W. Hobson, Proc. London Math. Soc. **24**, 55 (1892).
- ⁷⁷E. J. Weniger and E. O. Steinborn, J. Chem. Phys. **78**, 6121 (1983).
- ⁷⁸E. J. Weniger and E. O. Steinborn, J. Math. Phys. **26**, 664 (1985).
- ⁷⁹H. B. Schlegel, M. J. Frisch, Int. J. Quant. Chem. **54**, 83-87 (1995).
- ⁸⁰S. F. Boys, Proc. Roy. Soc. A **200**, 542 (1950).
- ⁸¹K. T. Tang, J. P. Toennies, J. Chem. Phys. **80**, 3726 (1984).
- ⁸²N. M. Temme, Numer. Math. **41**, 63 (1983).
- ⁸³J. Wimp, *Computation with Recurrence Relations*, Pitman, Boston, MA, 1984.
- ⁸⁴W. Gautschi, SIAM Rev. **9**, 24 (1967).
- ⁸⁵Wolfram Research, Inc., Mathematica, Version 7.0, Champaign, IL (2008).

Calculation of the molecular integrals with the range-separated correlation factor.**Supplemental material.**

Michał Silkowski,¹ Michał Lesiuk,^{1, a)} and Robert Moszynski¹

*Faculty of Chemistry, University of Warsaw, Pasteura 1, 02-093 Warsaw,
Poland*

(Dated: 20 April 2019)

The present document serves as a supplemental material for the publication *Calculation of the molecular integrals with the range-separated correlation factor*. It contains data or derivations which are of some importance but are not necessary for an overall understanding of the manuscript. However, the material presented here will be useful for a reader who wishes to repeat the derivations in details or implement the formulae independently.

arXiv:1412.5070v1 [physics.chem-ph] 16 Dec 2014

^{a)}Electronic mail: lesiuk@tiger.chem.uw.edu.pl.

A. Transformation between spherical and Cartesian differentials

An arbitrary mixed Cartesian differential can be rewritten as an adequate linear combination of the spherical tensor gradient operators by means of the following formula

$$\left(\frac{\partial}{\partial x}\right)^{l_x} \left(\frac{\partial}{\partial y}\right)^{l_y} \left(\frac{\partial}{\partial z}\right)^{l_z} = N(l, l_x, l_y, l_z) \sum_{l \leq l_x + l_y + l_z} c(l, m, l_x, l_y, l_z) \mathcal{Y}_{lm}(\bar{\nabla}), \quad (1)$$

where

$$N(l, l_x, l_y, l_z) = \left(\frac{2^3 \pi (l+1)! (2l_x)! (2l_y)! (2l_z)!}{(2l+2)! l_x! l_y! l_z!} \right)^{\frac{1}{2}}, \quad (2)$$

and

$$\begin{aligned} c(l, m, l_x, l_y, l_z) = & \sum_{l_{x'}+l_{y'}+l_{z'}=l} \frac{(l_x+l_{x'})!(l_y+l_{y'})!(l_z+l_{z'})!}{\left(\frac{l_x+l_{x'}}{2}\right)! \left(\frac{l_y+l_{y'}}{2}\right)! \left(\frac{l_z+l_{z'}}{2}\right)!} \\ & \sqrt{\frac{l! l_x! l_y! l_z! (l-|m|)!}{(2l)!(2l_{x'})!(2l_{y'})!(2l_{z'})!(l+|m|)!}} \\ & \frac{1}{2^l l!} \sum_{i=0}^{(l-|m|)/2} \binom{l}{i} \binom{i}{\frac{l_{x'}+l_{y'}-|m|}{2}} \frac{(-1)^i (2l-2i)!}{(l-|m|-2i)!} \\ & \sum_{k=0}^{\frac{l_{x'}+l_{y'}-|m|}{2}} \binom{\frac{l_{x'}+l_{y'}-|m|}{2}}{k} \binom{|m|}{l_{x'}-2k} (-1)^{-\frac{(|m|-l_{x'}+2k)}{2}}. \end{aligned} \quad (3)$$

Note, that the term under the outer sum is zero unless $\frac{l_x+l_{x'}}{2}$, $\frac{l_y+l_{y'}}{2}$, and $\frac{l_z+l_{z'}}{2}$ are the all integers. The above expressions can be compared with the relevant formulae from the work of Schlegel¹.

B. Computation of $\mathcal{U}(a, 1/2, z)$ with $a > 0$ and $z > 0$

Before presenting the detailed algorithm let us recall several useful formulae which are obeyed by the desired quantities, $\mathcal{U}(a, 1/2, z)$. Details and comments on these expressions can be found in Ref. 2 and references therein. Firstly, the small z expansion of the Kummer's hypergeometric function, $M(a, b, z)$, is

$$M(a, b, z) = \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(b)_n n!}, \quad (4)$$

where $(x)_n$ are the Pochhammer symbols. The corresponding expansion for the Tricomi's hypergeometric function can be written in terms of $M(a, b, z)$ as

$$\mathcal{U}(a, 1/2, z) = \sqrt{\pi} \left[\frac{M(a, 1/2, z)}{\Gamma(a + 1/2)} - 2\sqrt{z} \frac{M(a + 1/2, 3/2, z)}{\Gamma(a)} \right]. \quad (5)$$

Let us also recall the following recursive formulae for increasing the value of a

$$\mathcal{U}(a - 1, 1/2, z) + (1/2 - 2a - z)\mathcal{U}(a, 1/2, z) + a(a + 1/2)\mathcal{U}(a + 1, 1/2, z) = 0, \quad (6)$$

and the corresponding continued fraction (CF)

$$\begin{aligned} a \frac{\mathcal{U}(a + 1, 1/2, z)}{\mathcal{U}(a, 1/2, z)} &= \frac{a_1}{b_{1+}} \frac{a_2}{b_{2+}} \cdots, \\ a_n &= -(a + n - 1/2)(a + n - 1), \\ b_n &= (1/2 - 2a - z - 2n). \end{aligned} \quad (7)$$

Note additionally, that for very large values of z the Poincaré-type asymptotic formula can be used to simplify the computation. This formula is a simple inverse powers series in z

$$\mathcal{U}(a, 1/2, z) = z^{-a} \sum_{s=0}^{\infty} \frac{(a)_s (a + 1/2)_s}{s!} \frac{1}{(-z)^s}, \quad (8)$$

and can be efficiently implemented.

Let us now present the complete algorithm for calculation of $\mathcal{U}(a, 1/2, z)$. First of all, when $z < 1/2$ the small z expansion is supposed to be used, see Eqs. (4) and (5). Typically, these series converge fast in this regime and give accurate results despite a cancellation between two terms in Eq. (5). An exception is the region when the value of a is very large but such case is unlikely to occur in our applications.

The second ingredient of the algorithm is supposed to be used when $z > 1/2$. Let us assume we would like to compute $\mathcal{U}(a, 1/2, z)$ for a fixed z , up to some predefined value a_{max} . Firstly, we compute the ratio $\mathcal{U}(a_{max}, 1/2, z)/\mathcal{U}(a_{max} - 1, 1/2, z)$ by using the continued fraction formula (7). The evaluation can be performed with the Steed's algorithm and convergence of CF is acceptable for virtually every $z > 1/2$ and any reasonable a_{max} . The only inconvenience is the fact that the consecutive numerators and denominators in Eq. (7) grow very fast with n and need to be scaled by a small number from time to time to avoid overflows. Once the ratio $\mathcal{U}(a_{max}, 1/2, z)/\mathcal{U}(a_{max} - 1, 1/2, z)$ has been obtained, we rewrite the recursion relation (6) as follows

$$r_{a-1}(z) = -\frac{1}{(1/2 - 2a - z) + a(a + 1/2)r_a(z)}, \quad (9)$$

where the ratio $r_a(z) = \mathcal{U}(a+1, 1/2, z)/\mathcal{U}(a, 1/2, z)$. One sees that the value obtained from the CF formula is simply $a_{max} r_{a_{max}-1}$. Starting with this value we carry out the recursion (9) downwards, until a minimal value, a_0 , is reached which satisfies $0 < a_0 < 1$. At this point, $\mathcal{U}(a_0, 1/2, z)$ can be computed with expressions Eqs. (4) and (5) for $z < 10$ and Eq. (8) for $z > 10$ since the corresponding series converge acceptably for small a . Finally, using the definition of r_{a_0} one can see that $\mathcal{U}(a_0+1, 1/2, z) = r_{a_0}\mathcal{U}(a_0, 1/2, z)$, then $\mathcal{U}(a_0+2, 1/2, z) = r_{a_0+1}\mathcal{U}(a_0+1, 1/2, z)$ and so forth, until the value of a_{max} is reached. This furnishes the computation of the $\mathcal{U}(a, 1/2, z)$ functions.

C. Basic integrals with the range-separated correlation factor

Before expressing all basic integrals, B_i , through the $S(\alpha, \beta, \gamma)$ integrals, let us introduce a notation that makes the expressions more clear and succinct. Supposing one has two distinct correlation factors in the form given by Eq. (54) in the paper, with two sets of the parameters: c_0, ρ, B, μ and c'_0, ρ', B', μ' , respectively. Firstly, the operator \wp' acting on an expression changes all unprimed quantities into their primed counterparts (and *vice-versa*), for instance:

$$f'_{12}(r) = \wp' f_{12}(r) = \left(1 + \frac{1}{2}r\right) e^{-\mu'r^2} + c'_0 \left(1 - e^{-\mu'r^2} \sum_{\kappa=0}^n \frac{\mu'^{\kappa} r^{2\kappa}}{\kappa!}\right) r^{\rho'} e^{B'r}. \quad (10)$$

Hereafter, the notation $[\dots]_-$ (and $\{\dots\}_-$), should be considered as exactly the same expression as in the bracket before, but with $\xi \rightarrow -\xi$ in the second argument of S *e.g.*

$$S(\rho, B + 2\xi Q, \xi) + \left[\dots\right]_- = S(\rho, B + 2\xi Q, \xi) + S(\rho, B - 2\xi Q, \xi). \quad (11)$$

Finally, $S(\alpha, \dots)$ should be considered as integral S with exactly the same β and γ arguments as in the preceding integral S , for example:

$$S(\alpha, \beta, \gamma) + S(\alpha', \dots) + S(\alpha'', \dots) = S(\alpha, \beta, \gamma) + S(\alpha', \beta, \gamma) + S(\alpha'', \beta, \gamma). \quad (12)$$

With help of this notation one can represent the basic integrals as follows (note that B_5 expresses similarly as B_3 and thus a separate treatment is not required here):

$$B_1 = \sqrt{\frac{\pi^5}{p+q} \frac{1}{qp} \frac{1}{Q}} e^{-\xi Q^2} \left[S(0, 2\xi Q, \xi) - S(0, -2\xi Q, \xi) \right]. \quad (13)$$

$$B_2 = \sqrt{\frac{\pi^5}{p+q} \frac{1}{qp} \frac{1}{Q}} e^{-\xi Q^2} \left[S(0, 2\xi Q, \xi + \mu) + \frac{1}{2} S(1, 2\xi Q, \xi + \mu) + c_0 S(\rho, B + 2\xi Q, \xi) \right. \\ \left. - c_0 \sum_{\kappa=0}^n \frac{\mu^\kappa}{\kappa!} S(\rho + 2\kappa, B + 2\xi Q, \xi + \mu) \right] + \left[\dots \right]_- \quad (14)$$

$$B_3 = \sqrt{\frac{\pi^5}{p+q} \frac{1}{qp} \frac{1}{Q}} e^{-\xi Q^2} \left[S(1, 2\xi Q, \xi + \mu) + \frac{1}{2} S(2, 2\xi Q, \xi + \mu) + c_0 S(\rho + 1, B + 2\xi Q, \xi) \right. \\ \left. - c_0 \sum_{\kappa=0}^n \frac{\mu^\kappa}{\kappa!} S(\rho + 2\kappa + 1, B + 2\xi Q, \xi + \mu) \right] + \left[\dots \right]_- \quad (15)$$

$$B_4 = \sqrt{\frac{\pi^5}{p+q} \frac{1}{qp} \frac{1}{Q}} e^{-\xi Q^2} \left\{ S(1, 2\xi Q, \xi + \mu + \mu') + (1 + \wp') \frac{1}{2} S(2, \dots) + \frac{1}{4} S(3, \dots) + \right. \\ \left. + (1 + \wp') c_0 \left[S(\rho + 1, 2\xi Q + B, \xi + \mu') + \frac{1}{2} S(\rho + 2, \dots) \right] - \right. \\ \left. - (1 + \wp') c_0 \sum_{\kappa=0}^n \frac{\mu^\kappa}{\kappa!} \left[S(\rho + 2\kappa + 1, 2\xi Q + B, \xi + \mu' + \mu) + \frac{1}{2} S(\rho + 2\kappa + 2, \dots) \right] + \right. \\ \left. + c_0 c'_0 \sum_{k=0}^n \sum_{k'=0}^{n'} \frac{\mu^k \mu'^{k'}}{k! k'!} S(\rho + \rho' + 2k + 2k' + 1, 2\xi Q + B + B', \xi + \mu + \mu') + \right. \\ \left. + c_0 c'_0 S(\rho + \rho' + 1, 2\xi Q + B + B', \xi) \right\} + \left\{ \dots \right\}_- \quad (16)$$

$$\begin{aligned}
B_6 = & \sqrt{\frac{\pi^5}{p+q}} \frac{1}{qp} \frac{1}{Q} e^{-\xi Q^2} \left\{ \frac{1}{4} S(1, 2\xi Q, \xi + \mu + \mu') + 4\mu\mu' S(3, \dots) + \mu\mu' S(5, \dots) + 4\mu\mu' S(4, \dots) + \right. \\
& + (1 + \wp') \left(-\mu S(2, \dots) - \frac{\mu}{2} S(3, \dots) \right) + (1 + \wp') \left[c'_0 \left(\frac{1}{2} B' S(\rho' + 1, 2\xi Q + B', \xi + \mu) + \right. \right. \\
& + \left. \left. \frac{1}{2} \rho' S(\rho', \dots) - 2\mu B' S(\rho' + 2, \dots) - \mu B' S(\rho' + 3, \dots) - 2\mu\rho' S(\rho' + 1, \dots) - \mu\rho' S(\rho' + 2, \dots) \right) \right] + \\
& + c_0 c'_0 \left[BB' S(\rho + \rho' + 1, 2\xi Q + B + B', \xi) + \rho\rho' S(\rho + \rho' - 1, \dots) + \right. \\
& + (1 + \wp') \left(B\rho' S(\rho + \rho', \dots) \right) \left. \right] + (1 + \wp') \left[c_0 \sum_{\kappa=0}^n \frac{\mu^\kappa}{\kappa!} \right. \\
& \left(\frac{B}{2} S(\rho + 2\kappa + 1, 2\xi Q + B, \xi + \mu + \mu') - \mu S(\rho + 2\kappa + 2, \dots) + \kappa S(\rho + 2\kappa, \dots) - \right. \\
& - 2\mu' B S(\rho + 2\kappa + 2, \dots) + 2\mu\mu' S(\rho + 2\kappa + 3, \dots) - 4\mu'\kappa S(\rho + 2\kappa + 1, \dots) - \\
& - \mu' B S(\rho + 2\kappa + 3, \dots) + 2\mu\mu' S(\rho + 2\kappa + 4, \dots) - \mu'\kappa S(\rho + 2\kappa + 2, \dots) \left. \right) \left. \right] + \\
& + c_0 c'_0 (1 + \wp') \left[\sum_{\kappa=0}^n \frac{\mu^\kappa}{\kappa!} \left(BB' S(\rho + \rho' + 2\kappa + 1, 2\xi Q + B + B', \xi + \mu) + \right. \right. \\
& + 2\mu B' S(\rho + \rho' + 2\kappa + 2, \dots) + 2\kappa B' S(\rho + \rho' + 2\kappa, \dots) + \rho' B S(\rho + \rho' + 2\kappa, \dots) - \\
& - 2\mu\rho' S(\rho + \rho' + 2\kappa + 1, \dots) + 2\kappa\rho' S(\rho + \rho' + 2\kappa - 1, \dots) \left. \right) \left. \right] + \\
& + c_0 c'_0 \sum_{\kappa=0}^n \sum_{\kappa'=0}^{n'} \frac{\mu^\kappa \mu'^{\kappa'}}{\kappa! \kappa'!} \left[BB' S(\rho + \rho' + 2(\kappa + \kappa') + 1, 2\xi Q + B + B', \xi + \mu + \mu') + \right. \\
& + 4\mu\mu' S(\rho + \rho' + 2(\kappa + \kappa') + 3, \dots) + 4\kappa\kappa' S(\rho + \rho' + 2(\kappa + \kappa') - 1, \dots) + \\
& + (1 + \wp') \left(-2\mu B' S(\rho + \rho' + 2(\kappa + \kappa') + 2, \dots) - 4\mu\kappa' S(\rho + \rho' + 2(\kappa + \kappa') + 1, \dots) + \right. \\
& + \left. \left. 2B'\kappa S(\rho + \rho' + 2(\kappa + \kappa'), \dots) \right) \right] \left. \right\} + \left\{ \dots \right\}_-,
\end{aligned} \tag{17}$$

D. Calculation of the ω_k integrals

Before passing to the actual algorithm let us note that for evaluation of the desired molecular integrals one requires $\omega_0(x)$ and $\omega_1(x)$ with arbitrary real x . Herein, we present the algorithm for $x > 0$ only since the treatment of negative values of x is completely analogous. Considering the evaluation of $\omega_0(x)$ and $\omega_1(x)$ let us first note that the following

small x expansions hold

$$\omega_0(x) = \sum_{k=0}^{\infty} \frac{(-x)^k}{k!} \Gamma\left(\frac{k+1}{2}\right) \psi\left(\frac{k+1}{2}\right), \quad (18)$$

$$\omega_1(x) = \sum_{k=0}^{\infty} \frac{(-x)^k}{k!} \Gamma\left(\frac{k}{2} + 1\right) \psi\left(\frac{k}{2} + 1\right), \quad (19)$$

where $\psi(z)$ is the digamma function. The above expressions can be derived by simply expanding the exponential term in the integrand and integrating term by term. These series converge reasonably well for small x and the coefficients can be tabulated in advance or computed on the fly recursively. Let us also recall that for the half-integer values the digamma function takes the following closed form

$$\psi(n + 1/2) = -\gamma_E - 2 \log 2 + \sum_{k=1}^n \frac{2}{2k - 1}, \quad (20)$$

where γ_E is the Euler-Mascheroni constant.

By performing a linear exchange of variables the integral $\omega_0(x)$ can be rewritten as

$$\omega_0(x) = \int_0^{\infty} dz \log(z) e^{-xz - z^2} = \frac{1}{x} \int_0^{\infty} dt \log t e^{-t - t^2/x^2} - \frac{\log x}{x} \int_0^{\infty} dt e^{-t - t^2/x^2}. \quad (21)$$

By expanding the exponential terms containing $1/x^2$ in both integrals the following large x expansion is obtained

$$\omega_0(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (2k)! \psi(2k + 1)}{x^{2k+1} k!} - \log x \sum_{k=0}^{\infty} \frac{(-1)^k (2k)!}{x^{2k+1} k!}. \quad (22)$$

The above series is clearly divergent *i.e.* it provides asymptotic information about $\omega_0(x)$ when x is large. In practice, the above expression is very useful for calculations with $x > 50$, say, when typically only several terms are necessary to converge to the machine precision. Calculation of the individual terms is fairly straightforward, taking into account that for the integer values of n the digamma function is given by

$$\psi(n) = \sum_{k=1}^{n-1} \frac{1}{k} - \gamma_E. \quad (23)$$

A completely analogous expression to Eq. (22) holds for the second integral, $\omega_1(x)$.

Having resolved the cases of small and large x , let us discuss the calculation of $\omega_0(x)$ and $\omega_1(x)$ for moderate x where neither of the above schemes give a sufficient accuracy. After

numerical experiments we found that the following expansions give reasonably accurate piecewise fits for the desired quantities

$$F_n(x) = \sum_{k=0}^n \frac{c_k}{x^k} + \log x \sum_{k=0}^n \frac{d_k}{x^k}. \quad (24)$$

Our scheme for calculation of $\omega_0(x)$ and $\omega_1(x)$ goes as follows: for $x < 3$ we use the small- x infinite series expansion described above, where it converges to the accuracy of 15 digits in less than 50 terms. Obviously, for $x > 50$ the aforementioned large- x asymptotic expansion is used, which gives the machine precision after about 20 terms. For the remaining values of x we use the following approximations:

- $3 < x < 5$: $F_n(x)$ with $n = 6, 5$ for $\omega_0(x)$ and $\omega_1(x)$, respectively,
- $5 < x < 10$: $F_n(x)$ with $n = 6$ for both $\omega_0(x)$ and $\omega_1(x)$,
- $10 < x < 20$: $F_n(x)$ with $n = 5$ for both $\omega_0(x)$ and $\omega_1(x)$,
- $20 < x < 50$: $F_n(x)$ with $n = 4, 5$ for $\omega_0(x)$ and $\omega_1(x)$, respectively.

The coefficients c_k, d_k are different for each interval and are obtained from the least-squares fit on a equally spaced ($\Delta = 0.001$) grid. These reference data were obtained by integrating numerically $\omega_0(x)$ and $\omega_1(x)$ in an extended arithmetic precision. In the following Tables we report the values of the coefficients c_k, d_k for each interval. Accuracy of these fits is at least 13 significant digits and much better on the average, which is sufficient for the present purposes. Tests were performed on a large sample of randomly generated points.

REFERENCES

- ¹H. B. Schlegel, J. Chem. Phys. **77**, 3676 (1982).
²M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, (Dover, New York, 1972).

TABLE I. Coefficients c_k and d_k in the analytical fits of $\omega_0(x)$ and $\omega_1(x)$ with the functional form given by Eq. (24) for $3 < x < 5$. The symbol $[k]$ denotes the powers of 10, 10^k .

n	c_n	d_n
$\omega_0(x)$		
0	-2.79 418 516 254 156[-01]	+4.35 827 381 193 435[-02]
1	-2.90 713 893 948 490[+01]	+6.03 123 542 136 014[+00]
2	-4.95 348 276 834 514[+02]	+1.90 103 007 336 098[+02]
3	-1.87 541 627 996 346[+03]	+1.37 709 919 986 376[+03]
4	-2.97 088 892 758 365[+02]	+2.88 186 650 129 034[+03]
5	+2.15 515 331 846 835[+03]	+1.58 883 573 230 859[+03]
6	+5.41 315 200 857 445[+02]	+1.45 477 988 235 751[+02]
$\omega_1(x)$		
0	-6.81 608 673 256 649[-03]	-7.81 053 598 547 109[-04]
1	-6.03 136 412 874 311[+00]	+1.26 540 880 331 240[+00]
2	-1.23 965 414 329 573[+02]	+4.59 860 722 344 580[+01]
3	-2.92 393 636 456 307[+02]	+2.90 445 255 218 630[+02]
4	+2.42 606 911 139 842[+02]	+3.44 114 756 369 323[+02]
5	+1.79 651 772 296 869[+02]	+5.59 224 732 011 467[+01]

TABLE II. Coefficients c_k and d_k in the analytical fits of $\omega_0(x)$ and $\omega_1(x)$ with the functional form given by Eq. (24) for $5 < x < 10$. The symbol $[k]$ denotes the powers of 10, 10^k .

n	c_n	d_n
$\omega_0(x)$		
0	+2.67 336 306 337 877[-02]	-3.83 390 302 078 230[-03]
1	+4.47 174 316 838 360[+00]	-2.09 112 260 860 070[+00]
2	+1.69 542 503 617 498[+02]	-5.29 464 781 748 209[+01]
3	+1.39 875 310 206 529[+03]	-7.03 047 033 787 913[+02]
4	+2.02 964 879 213 659[+03]	-2.77 647 534 682 290[+03]
5	-2.17 744 743 464 807[+03]	-2.80 275 812 247 144[+03]
6	-1.42 551 927 395 417[+03]	-4.46 967 011 675 890[+02]
$\omega_1(x)$		
0	+2.89 074 504 802 456[-02]	-4.23 814 942 235 530[-03]
1	+4.44 902 630 985 788[+00]	-1.00 783 068 266 863[+00]
2	+1.08 226 256 016 102[+02]	-3.85 434 024 325 027[+01]
3	+4.27 985 618 179 230[+02]	-3.17 198 102 880 598[+02]
4	-7.33 719 621 295 158[+02]	-3.53 983 894 638 774[+02]
5	-6.13 015 358 740 435[+02]	+7.33 534 471 812 601[+02]
6	+8.04 702 720 006 004[+02]	+3.22 960 340 787 190[+02]

TABLE III. Coefficients c_k and d_k in the analytical fits of $\omega_0(x)$ and $\omega_1(x)$ with the functional form given by Eq. (24) for $10 < x < 20$. The symbol $[k]$ denotes the powers of 10, 10^k .

n	c_n	d_n
$\omega_0(x)$		
0	+9.85 170 564 088 696[-04]	-1.34 352 155 222 340[-04]
1	-3.14 495 326 884 370[-01]	-1.05 330 699 931 180[+00]
2	+1.14 410 655 314 381[+01]	-3.35 296 508 844 390[+00]
3	+1.01 789 460 952 360[+02]	-4.91 298 273 165 823[+01]
4	+5.99 368 814 053 674[+01]	-1.75 836 403 023 178[+02]
5	-1.74 101 744 430 719[+02]	-7.11 820 445 035 004[+01]
$\omega_1(x)$		
0	+1.79 379 586 198 002[-04]	-2.67 961 445 256 187[-05]
1	+4.80 466 657 853 673[-03]	-2.80 747 000 711 979[-03]
2	-2.08 437 134 302 115[+00]	-5.01 274 816 607 827[-01]
3	-6.61 049 670 065 573[+01]	+2.30 089 102 665 028[+01]
4	-2.02 378 315 248 675[+02]	+1.79 404 122 905 036[+02]
5	+2.61 597 803 283 919[+02]	+1.59 307 953 057 555[+02]

TABLE IV. Coefficients c_k and d_k in the analytical fits of $\omega_0(x)$ and $\omega_1(x)$ with the functional form given by Eq. (24) for $20 < x < 50$. The symbol $[k]$ denotes the powers of 10, 10^k .

n	c_n	d_n
$\omega_0(x)$		
0	+1.39 870 024 574 414[-05]	-1.76 030 960 012 356[-06]
1	-5.70 280 882 024 327[-01]	-1.00 126 191 147 610[+00]
2	+5.28 077 445 114 726[-01]	-1.35 267 549 073 835[-01]
3	+5.54 783 065 129 888[+00]	-1.14 510 519 374 110[+00]
4	-2.07 952 101 488 230[+00]	-1.13 489 580 958 509[+01]
$\omega_1(x)$		
0	-8.04 790 438 983 053[-06]	+9.92 668 209 243 167[-07]
1	-4.83 240 835 649 057[-03]	+8.49 304 239 905 638[-03]
2	-4.74 103 840 956 836[-02]	-8.86 737 271 752 571[-01]
3	-9.57 722 093 234 973[+00]	+3.53 498 860 651 889[+00]
4	-2.32 473 977 612 046[+01]	+2.77 433 057 711 969[+01]
5	+4.72 702 979 536 110[+01]	-5.06 583 892 061 286[+00]