

QUANTUM DILOGARITHM IDENTITIES AT ROOT OF UNITY

IVAN CHI-HO IP AND MASAHITO YAMAZAKI

Dedicated to the memory of Kentaro Nagao

ABSTRACT. We study the root of unity degeneration of cluster algebras and quantum dilogarithm identities. We prove identities for the cyclic dilogarithm associated with a mutation sequence of a quiver, and as a consequence new identities for the non-compact quantum dilogarithm at $b = 1$.

CONTENTS

1. Introduction	1
2. Quantum Dilogarithm Identities	2
2.1. Cluster Algebras	2
2.2. Quantum y -variables	3
2.3. Identities for $\Psi_q(x)$ and $g_b(x)$	4
2.4. Modular Double	7
3. Cyclic Dilogarithm Identities (I)	9
3.1. Root of Unity Limit	9
3.2. Identities for $d_N(x)$	9
3.3. Examples	14
4. Cyclic Dilogarithm Identities (II)	14
4.1. Root of Unity Limit of $g_b(x)$	14
4.2. Identities for $\mathfrak{g}_{1,N}(u)$	15
Appendix A. Proof of (80)	17
References	19

1. INTRODUCTION

Quantum dilogarithm identities [31, 24, 23] are remarkable identities satisfied by the quantum dilogarithm function. They are associated with a mutation sequence of a quiver, and fit nicely with the framework of the cluster algebras [13, 14] and their quantization [11, 12]. The simplest case of the quantum dilogarithm identity, associated with the A_2 quiver, is the celebrated pentagon identity [9]. More general quiver mutations can generate identities of increasing complexity. These general quantum dilogarithm identities have applications to a wide-ranging topics in mathematics and physics—the literature is too vast to be exhausted here (see e.g. references in [23]), but to name a few includes Donaldson-Thomas theory [25, 26, 24], integrable lattice models [2, 33, 36], 3d hyperbolic geometry [28, 18], 4d [16, 5, 35] and 3d [34] supersymmetric gauge theories.

2010 *Mathematics Subject Classification.* 13F60.

Quantum cluster algebra has a one deformation parameter q . In this paper we analyze the case where q is a root of unity. In this limit, the compact quantum dilogarithm function $\Psi_q(x)$ degenerates into a cyclic dilogarithm, and we obtain identities for the cyclic dilogarithm, again associated with a mutation sequence of a quiver.

It is known [23] that identities for $\Psi_q(x)$ takes two different forms, the tropical form (Theorem 2.12) and the universal form (Theorem 2.13), where the arguments are given respectively by the tropical quantum y -variables and the (full) quantum y -variables, respectively. Our cyclic dilogarithm identities also can be written in two different ways (Theorem 3.4 and Theorem 3.9), but their arguments are given by quantum y -variables as well as the cyclic y -variables (see Definition 3.2), where the latter can be thought of as the N -th root of the quantum y -variable for the modular double.

We also study the root of unity limit for the non-compact quantum dilogarithm $\Phi_b(x)$, which is discussed in a recent preprint [17]. Using the results from the cyclic dilogarithm identities, in Theorem 4.2 and Theorem 4.6 we found a whole new family of identities for the non-compact quantum dilogarithm at $b = 1$, corresponding to $q = -1$, which plays a very special role in the analysis. These identities are non-trivial since we are taking integral roots of non-commuting functions. The function $\Phi_1(x)$ may also provide new classical limits of the representation theory of the quantum plane $AB = q^2BA$ [19], which has only been previously studied in the $q \rightarrow 1$ limit with the quantum dilogarithm function appearing as the matrix coefficients.

Our identities is a vast generalization of the root-of-unity degeneration of the pentagon identity [1], which has been utilized for example in the study of 3-manifold invariants [22, 17] and solvable 3-dimensional lattice models [3]. We would like to urge the readers of this paper to find concrete applications of our results in their favorite mathematics/physics problems.

This paper is organized as follows. In Section 2 we set up our conventions and summarize the quantum dilogarithm identities. We then consider the root of unity limit of the quantum dilogarithm identities and their consequences, first in Section 3 for the compact dilogarithm $\Psi_q(x)$ and then for the non-compact quantum dilogarithm $\Phi_b(x)$ in Section 4. In the Appendix we present a new proof of the root of unity limit of the non-compact dilogarithm function $\Phi_b(x)$.

Acknowledgments. This work has grown up out of a over-the-tea conversation. We would like to thank the Kavli IPMU tea time for inspiration, and the WPI program (MEXT, Japan) and Kavli foundation for generous support.

2. QUANTUM DILOGARITHM IDENTITIES

In this section we review the basic of classical/quantum cluster algebras and the quantum dilogarithm identities. Our presentation throughout follows closely [23]. We here take q to be of a generic value.

2.1. Cluster Algebras.

Definition 2.1 (Mutation and Classical y -variables). *Let I be a finite set. We consider a pair (initial seed) (B, y) where $B = (b_{ij})_{i,j \in I}$ is a skew symmetric integer matrix, and $y = (y_i)_{i \in I}$ is an I -tuple of commutative variables. Let us define the mutation of (B, y) at k by $\mu_k(B, y) = (B', y')$, where*

$$(1) \quad b'_{ij} = \begin{cases} -b_{ij}, & i = k \text{ or } j = k, \\ b_{ij} + [-b_{ik}]_+ b_{kj} + b_{ik} [b_{kj}]_+ \\ = b_{ij} + [b_{ik}]_+ b_{kj} + b_{ik} [-b_{kj}]_+, & \text{otherwise,} \end{cases}$$

$$(2) \quad y'_i = \begin{cases} y_k^{-1}, & i = k, \\ y_i y_k^{\lfloor b_{ki} \rfloor_+} (1 + y_k)^{-b_{ki}} \\ = y_i y_k^{\lfloor -b_{ki} \rfloor_+} (1 + y_k^{-1})^{-b_{ki}}, & i \neq k, \end{cases}$$

with $\lfloor x \rfloor_+ := \max(x, 0)$. Starting from the initial seed, we can repeat the mutations to obtain a pair (B', y') . We call them seeds of (B, y) . Such y 's take values in the semifield

$$(3) \quad \mathbb{Q}_+(y) := \{\text{rational function of } y \text{ with subtraction-free expression}\},$$

and are called (classical) y -variables.

Remark 2.2. When B correspond to the incident matrix of a quiver Q , this mutation will be a quiver mutation of Q at the vertex $i = k$, and we have $\mu_k^2 = \text{id}$. Such a Q is necessarily acyclic.

Definition 2.3 (Tropicalization). Let us define the tropical semifield by

$$(4) \quad \text{Trop}(y) := \left\{ \prod_{i \in I} y_i^{a_i} \mid a_i \in \mathbb{Z} \right\}$$

with tropical sum

$$(5) \quad \prod_{i \in I} y_i^{a_i} \oplus \prod_{i \in I} y_i^{b_i} := \prod_{i \in I} y_i^{\min(a_i, b_i)}.$$

Then there exists a canonical surjective semifield homomorphism

$$(6) \quad \pi_{\text{trop}} : \mathbb{Q}_+(y) \rightarrow \text{Trop}(y),$$

defined by

$$(7) \quad y_i \mapsto y_i, \quad c \in \mathbb{Q} \mapsto 1.$$

Definition 2.4 (c -vector [14]). The c -vector of a cluster y -variable $y' \in \mathbb{Q}_+(y)$ is the vector $c = (c_i)_{i \in I} \in \mathbb{Z}^{|I|}$ such that

$$(8) \quad \pi_{\text{trop}}(y') = \prod_{j \in I} (y_j)^{c_j}.$$

Definition 2.5 (Tropical Sign). For a cluster variable $y' \in \mathbb{Q}_+(y)$, we set the tropical sign $\epsilon(y') = 1$ (resp. -1) if the components of the c -vector for y' are all non-negative (resp. all non-positive).

Remark 2.6. The tropical sign is always well-defined thanks to the sign coherence proven in [6, 27, 29].

2.2. Quantum y -variables. Let us next discuss the quantum version of cluster y -variables.

Definition 2.7 (Quantum y -variables [11, 12]). Consider a pair (B, Y) where $B = (b_{ij})_{i, j \in I}$ is a skew symmetric integer matrix, and $Y = (Y_i)_{i \in I}$ is an I -tuple of non-commutative variables such that

$$(9) \quad Y_i Y_j = q^{2b_{ji}} Y_j Y_i.$$

We define the quantum mutation of (B, Y) at $k \in I$ by $\mu_k(B, Y) := (B', Y')$ where b_{ij} mutates the same as before, and Y' is given by

$$(10) \quad Y'_i = \begin{cases} Y_k^{-1}, & i = k \\ Y_i \prod_{m=1}^{b_{ik}} (1 + q^{2m-1} Y_k), & i \neq k, b_{ik} \geq 0 \\ Y_i \prod_{m=1}^{|b_{ik}|} (1 + q^{2m-1} Y_k^{-1})^{-1}, & i \neq k, b_{ik} < 0. \end{cases}$$

Starting from the initial seed (B, Y) , we can repeat the mutations to obtain a pair (seed) (B', Y') . The variables Y'_i 's obtained this way are called quantum y -variables.

Remark 2.8. The mutation rule for the quantum y -variable (10) reduces to the mutation rule for the classical y -variables (2) in the limit $q \rightarrow 1$,

For our purposes it is useful to consider more generally non-commutative variables Y_α with $\alpha \in \mathbb{Z}^{|I|}$, satisfying the relation

$$(11) \quad Y_{\alpha+\beta} = q^{\langle \alpha, \beta \rangle} Y_\alpha Y_\beta, \quad \langle \alpha, \beta \rangle = -\langle \beta, \alpha \rangle := \alpha^T B \beta,$$

with the identification that $Y_i^{\pm 1} = Y_{\pm e_i}$ for the unit vector $e_i \in \mathbb{Z}^{|I|}$.

2.3. Identities for $\Psi_q(x)$ and $g_b(x)$.

Definition 2.9 (Period). For any I -sequence $\mathbf{k} = (k_1, \dots, k_L)$, $L > 1$, we consider a sequence of mutations. We fix our initial seed $(B(1), Y(1)) := (B, Y)$, and set

$$(12) \quad (B(t+1), Y(t+1)) := \mu_{k_t}(B(t), Y(t)).$$

Let σ be a permutation of I , we call the sequence \mathbf{k} of mutations a σ -period if

$$(13) \quad b_{\sigma(i)\sigma(j)}(L+1) = b_{ij}(1), \quad Y_{\sigma(i)}(L+1) = Y_i(1), \quad i, j \in I.$$

Remark 2.10. We can give alternative definition of the σ -period by replacing the quantum y -variables $Y_i(t)$ by their classical counterparts $y_i(t)$ in the definition (13), where $(B(1), y(1)) := (B, y)$ and

$$(14) \quad (B(t+1), y(t+1)) := \mu_{k_t}(B(t), y(t)).$$

The two definitions of σ -periods coincide [11].

Definition 2.11 (Compact Quantum Dilogarithm Function [9, 10]). We define the (compact) quantum dilogarithm function $\Psi_q(x)$ by¹

$$(15) \quad \Psi_q(x) := \frac{1}{(-qx; q^2)_\infty}, \quad (x; q)_\infty := \prod_{k=0}^{\infty} (1 - q^k x).$$

Theorem 2.12 (Quantum Dilogarithm Identity in Tropical Form [31, 24]). Consider a sequence of mutation labeled by an I -sequence \mathbf{k} , and suppose that \mathbf{k} is a σ -period for a permutation σ . Let c_t ($t = 1 \dots, L$) be the c -vector of the classical y -variables $y_{k_t}(t)$, and let us denote their tropical signs by ϵ_t ($t = 1 \dots, L$). We then have

$$(16) \quad \Psi_q(Y_{\epsilon_1 c_1})^{\epsilon_1} \dots \Psi_q(Y_{\epsilon_L c_L})^{\epsilon_L} = 1.$$

¹ In some literature q^2 is denoted by q . In our notation we always have integer powers of q .

We can convert this identity to another form, with the help of the following identity [23]:

$$(17) \quad Y_{k_t}(t) = \text{Ad} \left[\Psi_q(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots \Psi_q(Y_{\epsilon_{t-1} c_{t-1}})^{\epsilon_{t-1}} \right] (Y_{c_t}),$$

which holds for an arbitrary sign sequence $(\epsilon_1, \dots, \epsilon_t)$. This identity means we can shuffle the position of Ψ_q as

$$(18) \quad \begin{aligned} & \Psi_q(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots \Psi_q(Y_{\epsilon_{L-1} c_{L-1}})^{\epsilon_{L-1}} \Psi_q(Y_{\epsilon_L})^{\epsilon_L} \\ &= \Psi_q(Y_{k_L}(L)^{\epsilon_L c_L})^{\epsilon_L} \Psi_q(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots \Psi_q(Y_{\epsilon_{L-1} c_{L-1}})^{\epsilon_{L-1}}, \end{aligned}$$

and by repeating this we arrive at the following yet another form of the quantum dilogarithm identity:

Theorem 2.13 (Quantum Dilogarithm Identity in Universal Form). *Under the assumptions of Theorem 2.12, we have*

$$(19) \quad \Psi_q(Y_{k_L}(L)^{\epsilon_L})^{\epsilon_L} \cdots \Psi_q(Y_{k_1}(1)^{\epsilon_1})^{\epsilon_1} = 1.$$

Example 2.14 (A_2 quiver). *Let us consider the A_2 quiver*

$$(20) \quad 1 \longleftarrow 2.$$

We have $I = \{1, 2\}$, $b_{ij} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, $y = (y_1, y_2)$. Quantum y -variables Y_1, Y_2 satisfy $Y_1 Y_2 = q^2 Y_2 Y_1$. Let the I -sequence be $(1, 2, 1, 2, 1)$ with $L = 5$. This is a (12) -period. Indeed, we have

$$(21) \quad \begin{aligned} Y(1) &= (Y_1, Y_2), & b(1) &= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ Y(2) &= (Y_1^{-1}, Y_2(1 + qY_1)), & b(2) &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ Y(3) &= (Y_1^{-1}(1 + qY_2 + Y_1 Y_2), Y_2^{-1}(1 + q^{-1}Y_1)^{-1}), & b(3) &= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ Y(4) &= (Y_1(1 + qY_2 + Y_1 Y_2)^{-1}, q^{-1}Y_1^{-1}Y_2^{-1}(1 + qY_2)), & b(4) &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ Y(5) &= (Y_2^{-1}, q^{-1}Y_1 Y_2(1 + q^{-1}Y_2)^{-1}), & b(5) &= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ Y(6) &= (Y_2, Y_1), & b(6) &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \end{aligned}$$

where the c -vectors $c_t := c(y_{k_t}(t))$ and tropical signs ϵ_t are computed to be

$$(22) \quad \begin{aligned} c_1 &= (1, 0), & \epsilon_1 &= +, \\ c_2 &= (0, 1), & \epsilon_2 &= +, \\ c_3 &= (-1, 0), & \epsilon_3 &= -, \\ c_4 &= (-1, -1), & \epsilon_4 &= -, \\ c_5 &= (0, -1), & \epsilon_5 &= -. \end{aligned}$$

The quantum dilogarithm identity is, in the tropical form

$$(23) \quad \Psi_q(Y_1) \Psi_q(Y_2) \Psi_q(Y_1)^{-1} \Psi_q(q^{-1}Y_1 Y_2)^{-1} \Psi_q(Y_2)^{-1} = 1,$$

where we used $Y_{e_1+e_2} = q^{-1}Y_{e_1}Y_{e_2} = q^{-1}Y_1Y_2$ (recall (11)). This is the famous pentagon identity [9] for $\Psi_q(x)$. In the universal form, we have

$$(24) \quad \begin{aligned} & \Psi_q(Y_2)^{-1} \Psi_q \left(q(1 + qY_2)^{-1} Y_2 Y_1 \right)^{-1} \\ & \Psi_q \left((1 + qY_2 + Y_1 Y_2)^{-1} Y_1 \right)^{-1} \Psi_q(Y_2(1 + qY_1)) \Psi_q(Y_1) = 1. \end{aligned}$$

Example 2.15 (A_3 quiver). Consider A_3 quiver, where we choose the orientation

$$(25) \quad 1 \longleftarrow 2 \longrightarrow 3$$

such that

$$(26) \quad B = (b)_{ij} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}.$$

and $y = (y_1, y_2, y_3)$. Quantum y -variables Y_1, Y_2, Y_3 satisfy

$$(27) \quad Y_1 Y_2 = q^2 Y_2 Y_1, \quad Y_3 Y_2 = q^2 Y_2 Y_3, \quad Y_1 Y_3 = Y_3 Y_1.$$

Consider a mutation sequence $(1, 3, 2, 1, 3, 2, 1, 3, 2)$. The quantum y -variables are then computed to be

$$(28) \quad \begin{aligned} Y(1) &= (Y_1, Y_2, Y_3), \\ Y(2) &= (Y_1^{-1}, Y_2(1 + qY_1), Y_3), \\ Y(3) &= (Y_1^{-1}, Y_2 X_q, Y_3^{-1}), \\ Y(4) &= (Y_1^{-1}(1 + qY_2 X_q), Y_2^{-1} X_q^{-1}, Y_3^{-1}(1 + qY_2 X_q)), \\ Y(5) &= (Y_1(1 + q^{-1} Y_2 X_q)^{-1}, qY_2^{-1} Y_1^{-1} X_q^{-1} (1 + q^{-1} Y_1 + qY_2 X_q), Y_3(1 + q^{-1} Y_2 X_q)^{-1}), \\ Y(6) &= (Y_1(1 + q^{-1} Y_2 X_q)^{-1}, q^2 Y_2^{-1} Y_1^{-1} Y_3^{-1} X_q^{-1} (1 + q^{-1} Y_1 + q^3 Y_2 X_q) (1 + q^{-1} Y_3 + qY_2 X_q), \\ & \quad Y_3(1 + q^{-1} Y_2 X_q)^{-1}) \\ &= (Y_1(1 + q^{-1} Y_2 X_q)^{-1}, Y_1^{-1} Y_2^{-1} Y_3^{-1} (1 + q^3 Y_2 + q^2 Y_2 Y_3) (1 + qY_2 + q^2 Y_2 Y_1), \\ & \quad Y_3(1 + q^{-1} Y_2 X_q)^{-1}), \\ Y(7) &= (Y_1(1 + q^{-1} Y_2 X_q)^{-1} (1 + qY_1^{-1} Y_2^{-1} Y_3^{-1} (1 + q^3 Y_2 + q^2 Y_2 Y_3) (1 + qY_2 + q^2 Y_2 Y_1)), \\ & \quad (1 + qY_2 + q^2 Y_2 Y_3)^{-1} (1 + q^3 Y_2 + q^2 Y_2 Y_1)^{-1} Y_3 Y_2 Y_1, \\ & \quad Y_3(1 + q^{-1} Y_2 X_q)^{-1} (1 + qY_1^{-1} Y_2^{-1} Y_3^{-1} (1 + q^3 Y_2 + q^2 Y_2 Y_3) (1 + qY_2 + q^2 Y_2 Y_1))) \\ &= (q^{-1} Y_3^{-1} Y_2^{-1} (1 + qY_2), (1 + qY_2 + q^2 Y_2 Y_3)^{-1} (1 + q^3 Y_2 + q^2 Y_2 Y_1)^{-1} Y_3 Y_2 Y_1, \\ & \quad q^{-1} Y_1^{-1} Y_2^{-1} (1 + qY_2)), \\ Y(8) &= (Y_3(1 + qY_2^{-1})^{-1}, (1 + qY_2 + q^2 Y_2 Y_1)^{-1} Y_1, qY_2^{-1} Y_1^{-1} (1 + qY_2)), \\ Y(9) &= (Y_3(1 + qY_2^{-1})^{-1}, Y_2^{-1}, Y_1(1 + qY_2^{-1})^{-1}), \\ Y(10) &= (Y_3, Y_2, Y_1), \end{aligned}$$

where

$$(29) \quad X_q := (1 + qY_1)(1 + qY_3) = (1 + qY_1 + qY_3 + q^2 Y_1 Y_3).$$

The mutation sequence is therefore a (13)-period. The c -vectors and tropical signs are given by

$$(30) \quad \begin{array}{ll} c_1 = (1, 0, 0), & \epsilon_1 = +, \\ c_2 = (0, 0, 1), & \epsilon_2 = +, \\ c_3 = (0, 1, 0), & \epsilon_3 = +, \\ c_4 = (-1, 0, 0), & \epsilon_4 = -, \\ c_5 = (0, 0, -1), & \epsilon_5 = -, \\ c_6 = (-1, -1, -1), & \epsilon_6 = -, \\ c_7 = (0, -1, -1), & \epsilon_7 = -, \\ c_8 = (-1, -1, 0), & \epsilon_8 = -, \\ c_9 = (0, -1, 0), & \epsilon_9 = -. \end{array}$$

The quantum dilogarithm identity in the tropical form is²

$$(31) \quad \begin{aligned} & \Psi_q(Y_1)\Psi_q(Y_3)\Psi_q(Y_2)\Psi_q(Y_1)^{-1}\Psi_q(Y_3)^{-1} \\ & \Psi_q(Y_1Y_2Y_3)^{-1}\Psi_q(qY_2Y_3)^{-1}\Psi_q(qY_2Y_1)^{-1}\Psi_q(Y_2)^{-1} = 1, \end{aligned}$$

This can be easily seen as a consequence of the pentagon (23).

The universal form for the quantum dilogarithm identity is given by

$$(32) \quad \begin{aligned} & \Psi_q(Y_2(1+qY_1)(1+qY_3))\Psi_q(Y_3)\Psi_q(Y_1) \\ & = \Psi_q((1+qY_2X_q)^{-1}Y_1)\Psi_q((1+qY_2X_q)^{-1}Y_3) \\ & \Psi_q((1+qY_2+Y_1Y_2)^{-1}(1+q^3Y_2+Y_3Y_2)^{-1}Y_3Y_2Y_1) \\ & \Psi_q(q(1+qY_2)^{-1}Y_2Y_3)\Psi_q(q(1+qY_2)^{-1}Y_1Y_2)\Psi_q(Y_2). \end{aligned}$$

2.4. Modular Double. Let $b \in \mathbb{C}$ and write $q = e^{i\pi b^2}$, and define a new variable \tilde{q} by $\tilde{q} = e^{i\pi b^{-2}}$.

Definition 2.16 (Non-Compact Quantum Dilogarithm [7, 8]). We define the non-compact quantum dilogarithm function $\Phi_b(x)$ by an integral³

$$(33) \quad \Phi_b(x) := \exp\left(\int_{\mathbb{R}+i0} \frac{e^{-2ixt}}{4 \sinh(bt) \sinh(b^{-1}t)} \frac{dt}{t}\right)$$

in the strip $|\operatorname{Im}(z)| < \left|\operatorname{Im}\left(\frac{i(b+b^{-1})}{2}\right)\right|$. This function can then be analytically continued into the whole complex plane. For $\operatorname{Im}(b^2) > 0$, we have $|q|, |\tilde{q}^{-1}| < 1$ and it turns out that $\Phi_b(x)$ can be written as an infinite product

$$(34) \quad \Phi_b(x) = \frac{\prod_{k=0}^{\infty} (1 + q^{2k+1} e^{2\pi b x})}{\prod_{k=0}^{\infty} (1 + \tilde{q}^{-(2k+1)} e^{2\pi b^{-1} x})} = \frac{\Psi_{\tilde{q}^{-1}}(e^{2\pi b^{-1} x})}{\Psi_q(e^{2\pi b x})}.$$

In the following we will also use a function $g_b(u)$, which is related to the function $\Phi_b(x)$ by the relation $g_b(e^{2\pi b x}) = \Phi_b(x)$. Let x, p with $[x, p] = \frac{1}{2\pi i}$ be the standard position and momentum operators acting on $L^2(\mathbb{R})$. Also denote by

$$(35) \quad u = e^{2\pi b x}, \quad v = e^{2\pi b p},$$

² This identity can also be written as

$$(Y_1; q^2)_{\infty} (Y_3; q^2)_{\infty} (Y_2; q^2)_{\infty} = (Y_2; q^2)_{\infty} (-Y_2 Y_1; q^2)_{\infty} (-Y_2 Y_3; q^2)_{\infty} (Y_2 Y_1 Y_3; q^2)_{\infty} (Y_3; q^2)_{\infty} (Y_2; q^2)_{\infty}.$$

³ This is also denoted by $e_b(x)$ in the literature. Note that this is the inverse of Φ_b used in [23].

such that

$$(36) \quad uv = q^2vu$$

on a natural dense core of their domain [32]. Then the functions $\Phi_b(x), g_b(u)$ satisfy the pentagon identity as unitary operators on $L^2(\mathbb{R})$:⁴

$$(37) \quad \begin{aligned} \Phi_b(p)\Phi_b(x) &= \Phi_b(x)\Phi_b(x+p)\Phi_b(p), \\ g_b(v)g_b(u) &= g_b(u)g_b(qvu)g_b(v). \end{aligned}$$

The concept of modular double is introduced in [8] and has been considered in various contexts of mathematics and physics, in particular to representation theory of split real quantum groups [15, 20] and quantum Liouville theory [30, 4] involving the Weyl operators. The idea is that the Weyl pair $\{u, v\}$ has a ‘‘modular double counterpart’’ which mutually commutes with them, and it is natural to consider both pairs of Weyl operators in their functional analysis.

Given a Weyl pair $\{u, v\}$ as above, their modular double counterpart is defined through the use of functional calculus on positive (essentially) self-adjoint operators

$$(38) \quad \widetilde{u} := u^{\frac{1}{b^2}} = e^{2\pi b^{-1}x}, \quad \widetilde{v} := v^{\frac{1}{b^2}} = e^{2\pi b^{-1}p},$$

so that

$$(39) \quad \widetilde{uv} = \widetilde{q^2vu}$$

and consequently the pair $\{\widetilde{u}, \widetilde{v}\}$ mutually commute with $\{u, v\}$ on certain dense domain. With this notation, we see that

$$(40) \quad g_b(u) = \frac{\Psi_{\widetilde{q}^{-1}}(\widetilde{u})}{\Psi_q(u)},$$

and in particular it is manifest that we have the modular duality:

$$(41) \quad g_b(u) = g_{b^{-1}}(\widetilde{u}).$$

Following the same idea, we introduce the modular dual variables formally by

$$(42) \quad \widetilde{Y}_i := Y_i^{\frac{1}{b^2}}$$

such that

$$(43) \quad \widetilde{Y}_i \widetilde{Y}_j = \widetilde{q}^{2b_{ji}} \widetilde{Y}_j \widetilde{Y}_i.$$

and use them as the initial seed of the *dual quantum y-variables*, defined below:

Definition 2.17 (Dual Quantum y-variable). *Let us define the dual quantum cluster y-variables by replacing q by \widetilde{q} in the definition of quantum y-variables (Definition (2.7)). The mutation rule (10) is then replaced by*

$$(44) \quad \widetilde{Y}'_i = \begin{cases} \widetilde{Y}_k^{-1}, & i = k, \\ \widetilde{Y}_i \prod_{m=1}^{b_{ik}} (1 + \widetilde{q}^{2m-1} \widetilde{Y}_k), & i \neq k, b_{ik} \geq 0, \\ \widetilde{Y}_i \prod_{m=1}^{|b_{ik}|} (1 + \widetilde{q}^{2m-1} \widetilde{Y}_k^{-1})^{-1}, & i \neq k, b_{ik} < 0. \end{cases}$$

Moreover, the dual quantum y-variables \widetilde{Y}_i commute with the original quantum y-variables Y_i .

⁴The identity for $\Phi_b(x)$ is called the local form in [23].

Since Y_i 's and their duals \widetilde{Y}_i 's commute, the ratio of compact dilogarithms in (40) can be factorized, and we immediately learn that we can derive the quantum dilogarithm identities (with reverse order) for $g_b(u)$.

Theorem 2.18 (Identities for Non-Compact Quantum Dilogarithms [23]). *Under the assumptions of Theorem 2.12, we have, in tropical form*

$$(45) \quad g_b(Y_{\epsilon_L c_L})^{\epsilon_L} \cdots g_b(Y_{\epsilon_1 c_1})^{\epsilon_1} = 1,$$

and in universal form

$$(46) \quad g_b(Y_{k_1}(1)^{\epsilon_1})^{\epsilon_1} \cdots g_b(Y_{k_L}(L)^{\epsilon_L})^{\epsilon_L} = 1.$$

3. CYCLIC DILOGARITHM IDENTITIES (I)

3.1. Root of Unity Limit. Let us choose a positive integer N . In this section we will choose the value of q to be $q = \zeta := e^{\frac{\pi i}{N}}$, a primitive $2N$ -th root of unity. Then q^2 is the N -th primitive root of unity, and $q^N = -1$. We also have

$$(47) \quad \widetilde{q} = \widetilde{\zeta} := e^{\pi i N} = (-1)^N.$$

With this specialization, the relation (9) reduces to

$$(48) \quad Y_i Y_j = \zeta^{2b_{ji}} Y_j Y_i,$$

and the dual quantum y -variable is simply given by

$$(49) \quad \widetilde{Y}_i = Y_i^N,$$

which by definition commute with all the Y_j .

Proposition 3.1. *When $\widetilde{q} = \widetilde{\zeta} = e^{\pi i N} = (-1)^N$, the transformation property of the dual quantum y -variable (44) reduces to*

$$(50) \quad \widetilde{Y}'_i = \begin{cases} \widetilde{Y}_k^{-1}, & i = k, \\ \widetilde{Y}_i(1 + \widetilde{\zeta} \widetilde{Y}_k)^{b_{ik}}, & i \neq k, b_{ik} \geq 0, \\ \widetilde{Y}_i(1 + \widetilde{\zeta} \widetilde{Y}_k^{-1})^{b_{ik}}, & i \neq k, b_{ik} < 0. \end{cases}$$

Proof. When $\widetilde{q} = \widetilde{\zeta} = (-1)^N$, we have

$$\widetilde{q}^{2m-1} = (-1)^{(2m-1)N} = (-1)^N = \widetilde{\zeta}.$$

Hence the product $\prod_{m=1}^{|b_{ik}|}$ in (44) becomes trivial, leading to the formula above. \square

3.2. Identities for $d_N(x)$.

Definition 3.2 (Cyclic y -variable). *Let us define the cyclic y -variables \overline{Y}_i . They satisfy the commutation relation*

$$(51) \quad \overline{Y}_i \overline{Y}_j = \zeta^{2b_{ji}} \overline{Y}_j \overline{Y}_i,$$

which coincides with the relation (9) for the quantum y -variables. We also take the initial value to be $\overline{Y}_i = Y_i$. However, we require \overline{Y}_i to transform formally as the dual quantum y -variable $\overline{Y}_i = \widetilde{Y}_i^{\frac{1}{N}}$, so that we define

$$(52) \quad \overline{Y}'_i = \begin{cases} \overline{Y}_k^{-1}, & i = k, \\ \overline{Y}_i(1 + \widetilde{\zeta} \overline{Y}_k^N)^{\frac{b_{ik}}{N}}, & i \neq k, b_{ik} \geq 0, \\ \overline{Y}_i(1 + \widetilde{\zeta} \overline{Y}_k^{-N})^{\frac{b_{ik}}{N}}, & i \neq k, b_{ik} < 0. \end{cases}$$

We can repeat this for a mutation sequence, to obtain cyclic y -variables. Similar to the previous cases, we define $(B(1), \bar{Y}(1)) := (B, \bar{Y}) = (B, Y(1))$ and $(B(t+1), \bar{Y}(t+1)) := \mu_{k_i}(B(t), \bar{Y}(t))$. We can also extend the definition of \bar{Y}_i to \bar{Y}_α with $\alpha \in \mathbb{Z}^{|\mathbb{I}|}$, as in (11) in the case of Y_i .

Definition 3.3 (Cyclic Dilogarithm Function). *For a given positive integer N , we define the cyclic dilogarithm function $d_N(x)$ by⁵*

$$(54) \quad d_N(x) := \left(1 + (-x)^N\right)^{\frac{N-1}{2N}} \prod_{k=1}^{N-1} (1 + \zeta^{2k+1}x)^{-\frac{k}{N}}.$$

We are now ready to state one of our main theorems.

Theorem 3.4 (Cyclic Dilogarithm Identity in Dual Universal Form). *Under the assumptions of Theorem (2.12), we have*

$$(55) \quad d_N(\bar{Y}_{k_L}(L)^{\epsilon_L})^{\epsilon_L} \cdots d_N(\bar{Y}_{k_1}(1)^{\epsilon_1})^{\epsilon_1} = 1.$$

Proof. The rest of this subsection is devoted to a proof of this theorem.

Following [1], the basic strategy is to take the limit of the quantum dilogarithm identity for $\Psi_q(x)$. First of all let us consider the classical limit by taking

$$(56) \quad q := e^{-\frac{\tau}{2}}, \quad \tau \in \mathbb{R}_{>0}$$

so that $|q| < 1$, and we take the limit when $\tau \rightarrow 0$. Then it is well-known that

$$(57) \quad \Psi_q(x)^{-1} = (-qx; q^2)_\infty = R_\tau(x) (1 + \mathcal{O}(\tau)),$$

with $R_\tau(x)$ defined by

$$(58) \quad R_\tau(x) := e^{-\frac{\text{Li}_2(-x)}{\tau}}.$$

The quantum dilogarithm identity in the tropical form (2.12) implies the following

Proposition 3.5. *Let the quantum Y -variable be associated to $q = e^{-\tau/2}$. Then we have*

$$(59) \quad \lim_{\tau \rightarrow 0} R_\tau(Y_{\epsilon_L c_L})^{\epsilon_L} \cdots R_\tau(Y_{\epsilon_1 c_1})^{\epsilon_1} = 1.$$

This will be important in our later analysis. The limit here is analyzed for example in [23]. As proven there, the condition that the resulting expression is trivial gives the classical dilogarithm identity.

Now let us consider the case of the root of unity limit. We let

$$(60) \quad q := e^{-\frac{\tau}{2N^2}} \zeta,$$

⁵ Another definition found in the literature is

$$(53) \quad d_N^*(x) := \left(1 - x^N\right)^{\frac{N-1}{2N}} \prod_{k=1}^{N-1} (1 - \zeta^{2k}x)^{-\frac{k}{N}}.$$

This is related to our $d_N(x)$ by $d_N(x) = d_N^*(-\zeta x)$.

and take the limit as $\tau \rightarrow 0$. In this limit, the quantum dilogarithm $\Psi_q(x)$ behaves as [1]⁶

$$(63) \quad \Psi_q(x)^{-1} = (-qx; q^2)_\infty = R_\tau((-x)^N) d_N(x) (1 + \mathcal{O}(\tau)),$$

with $d_N(x)$ defined previously in (54) and $R_\tau(x)$ in (57). To save notations let us denote by

$$(64) \quad R_{\tau,N}(x) := R_\tau((-x)^N).$$

Now, substituting the limiting form (63) into (16), we obtain an identity involving the functions $R_{\tau,N}(x)$ and $d_N(x)$:

$$(65) \quad \lim_{\tau \rightarrow 0} R_{\tau,N}(Y_{\epsilon_L c_L})^{\epsilon_L} d_N(Y_{\epsilon_L c_L})^{\epsilon_L} \cdots R_{\tau,N}(Y_{\epsilon_1 c_1})^{\epsilon_1} d_N(Y_{\epsilon_1 c_1})^{\epsilon_1} = 1.$$

To simplify the resulting expression, we can shuffle the positions of $R_{\tau,N}(x)$, using the following Lemma:

Lemma 3.6 (Conjugation by $R_{\tau,N}$ [1]). *Let $Y_i Y_j = q^{2b_{ji}} Y_j Y_i$. We have*

$$(66) \quad \lim_{\tau \rightarrow 0} R_{\tau,N}(Y_i) \cdot Y_j \cdot R_{\tau,N}(Y_i)^{-1} = Y_j (1 + \tilde{q} Y_i^N)^{-\frac{b_{ji}}{N}}.$$

Proof. We here provide a self-contained proof using functional calculus. We can commute u_j to the left and get

$$\begin{aligned} (\text{LHS}) &= Y_j \cdot \lim_{\tau \rightarrow 0} R_{\tau,N}(q^{2b_{ji}} Y_i) R_{\tau,N}(Y_i)^{-1} \\ &= Y_j \cdot \lim_{\tau \rightarrow 0} \frac{R_\tau(e^{-\tau b_{ji}/N} (-Y_i)^N)}{R_\tau((-Y_i)^N)}. \end{aligned}$$

Now the limit can be computed directly by calculus. Using $\text{Li}_2(x) = -\int_0^x \frac{\log(1-t)}{t} dt$, and let $q(\tau) = e^{-\tau b_{ji}/N}$, we have

$$\begin{aligned} (67) \quad \lim_{\tau \rightarrow 0} \log \frac{R_\tau(q(\tau)(-x)^N)}{R_\tau((-x)^N)} &= \lim_{\tau \rightarrow 0} \frac{1}{\tau} \left(\int_0^{-q(\tau)(-x)^N} \frac{\log(1-t)}{t} dt - \int_0^{(-x)^N} \frac{\log(1-t)}{t} dt \right) \\ &= \lim_{\tau \rightarrow 0} \frac{1}{\tau} \left(\int_{(-x)^N}^{-q(\tau)(-x)^N} \frac{\log(1-t)}{t} dt \right) \\ &= \lim_{\tau \rightarrow 0} \frac{\log(1 - (-q(\tau)(-x)^N))}{-q(\tau)(-x)^N} \cdot \left(-\frac{dq(\tau)}{d\tau} (-x)^N \right) \\ &= -\log(1 + (-x)^N) \frac{b_{ji}}{N} \\ &= -\log(1 + \tilde{q} x^N) \frac{b_{ji}}{N}. \end{aligned}$$

This proves (66). □

Remark 3.7. *In the limit $\tau \rightarrow 0$, Y_i^N , and hence any function of them, commute with all the Y_j . However, in the analysis above the shuffling should first be performed for finite τ , and there remains a non-trivial finite contribution in the limit $\tau \rightarrow 0$ due to the canceling factor $1/\tau$ in the definition of $R_{\tau,N}(x)$.*

⁶ This is equivalent to the formula

$$(61) \quad (x; q^2)_\infty = (1 - x^N)^{\frac{1-N}{2N}} R_\tau^*(x^N) d_N^*(x) (1 + \mathcal{O}(\tau)),$$

with

$$(62) \quad R_\tau^*(x) := (1 - x)^{\frac{1}{2}} e^{-\frac{\text{Li}_2(x)}{\tau}}.$$

Now the crucial observation is that the conjugation by Y_i (66), taking into accounts the tropical signs ϵ_i , has exactly the same effect as the mutation rule for the cyclic y -variables (52). In particular we can pull out all the $R_{\tau,N}$'s to the left or right, as the cost of replacing the arguments of d_N 's by their mutated versions. To be more precise, by mimicking the proof of (17), we obtain

$$(68) \quad \bar{Y}_{k_i}(t) = \lim_{\tau \rightarrow 0} \text{Ad}^{-1} [R_{\tau,N}(Y_{\epsilon_{-1}c_{i-1}})^{\epsilon_{i-1}} \cdots R_{\tau,N}(Y_{\epsilon_i c_i})^{\epsilon_i}] (\bar{Y}_{c_i}).$$

We can now appeal to the shuffling argument which we utilized in the proof of Theorem 2.13. By shuffling the position of $R_{\tau,N}$, we obtain (recall $Y_\alpha = Y_\alpha(1) = \bar{Y}_\alpha$)

$$\begin{aligned} & R_{\tau,N}(Y_{c_L \epsilon_L})^{\epsilon_L} d_N(Y_{c_L \epsilon_L})^{\epsilon_L} \cdots d_N(Y_{c_2 \epsilon_2})^{\epsilon_2} R_{\tau,N}(Y_{c_1 \epsilon_1})^{\epsilon_1} d_N(Y_{c_1 \epsilon_1})^{\epsilon_1} \\ &= R_{\tau,N}(Y_{c_L \epsilon_L})^{\epsilon_L} d_N(Y_{c_L \epsilon_L})^{\epsilon_L} \cdots \\ & \quad R_{\tau,N}(Y_{c_1 \epsilon_1})^{\epsilon_1} d_N(\bar{Y}_{k_2}(2)^{\epsilon_2})^{\epsilon_2} d_N(\bar{Y}_{k_1}(1)^{\epsilon_1})^{\epsilon_1} (1 + \mathcal{O}(\tau)). \end{aligned}$$

By repeating this, we arrive at

$$R_{\tau,N}(Y_{c_L \epsilon_L})^{\epsilon_L} \cdots R_{\tau,N}(Y_{c_1 \epsilon_1})^{\epsilon_1} d_N(\bar{Y}_{k_L}(L)^{\epsilon_L})^{\epsilon_L} \cdots d_N(\bar{Y}_{k_1}(1)^{\epsilon_1})^{\epsilon_1} (1 + \mathcal{O}(\tau)) = 1.$$

The remaining task is to show that the product of R 's vanish in the limit $\tau \rightarrow 0$. Recall $R_{\tau,N}(Y_i) := R_\tau((-Y_i)^N)$. Let $Y'_i := (-Y_i)^N$. Then

$$Y'_i Y'_j = (-Y_i)^N (-Y_j)^N = q^{2N^2} (-Y_j)^N (-Y_i)^N = e^{-\tau} Y'_j Y'_i,$$

hence all the arguments of R satisfy the same commutation relation as the Y_i 's with $q = e^{-\tau/2}$. Hence by Proposition 3.5, we conclude that the product of R is trivial. Since $d_N(x)$ does not depend on τ in the limit, we have proven the identity (55) for the function $d_N(x)$. \square

Remark 3.8. *In this derivation, we shuffled all the R 's to the left of the product. Alternatively we can choose to move $L' = 0, \dots, L$ of the R 's to the left and $L - L'$ to the right, and cancel the product of R 's in the middle. We then obtain variants of the expression above, with some of the variables rescaled according to (66).*

We next rewrite the cyclic dilogarithm identity in another form.

Theorem 3.9 (Cyclic Dilogarithm Identities in Standard Universal Form). *We have the cyclic dilogarithm identities*

$$(69) \quad d_N(Y_{k_1}(1)^{\epsilon_1})^{\epsilon_1} \cdots d_N(Y_{k_L}(L)^{\epsilon_L})^{\epsilon_L} = 1.$$

for the standard quantum y -variables, with the reverse ordering.

Proof. We start from the cyclic dilogarithm identity of (55), we can mimic the arguments of the shuffle formula. First of all, we note that $d(Y_i)$ instead of conjugating tropical form to universal form, actually interchange universal form for \bar{Y}_i and that for the standard Y_i variables.

Lemma 3.10. *We have for $b_{ji} \geq 0$*

$$(70) \quad d_N(Y_i) \cdot Y_j \cdot d_N(Y_i)^{-1} = Y_j \frac{(1 + \bar{q} Y_i^N)^{\frac{b_{ji}}{N}}}{\prod_{k=1}^{b_{ji}} (1 + q^{2k-1} Y_i)},$$

and for $b_{ji} < 0$

$$(71) \quad d_N(Y_i) \cdot Y_j \cdot d_N(Y_i)^{-1} = Y_j \frac{(1 + \bar{q} Y_i^N)^{\frac{b_{ji}}{N}}}{\prod_{k=1}^{|b_{ji}|} (1 + q^{-(2k-1)} Y_i)^{-1}}.$$

Proof. First assume $b_{ji} > 0$. We then have

$$\begin{aligned}
 d_N(Y_i) \cdot Y_j \cdot d_N(Y_i)^{-1} &= Y_j \frac{d_N(q^{2b_{ji}} Y_i)}{d_N(Y_i)} \\
 &= Y_j \frac{\prod_{k=1}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{k}{N}}}{\prod_{k=1}^{N-1} (1 + q^{2k+1+2b_{ji}} Y_i)^{\frac{k}{N}}} \\
 &= Y_j \frac{\prod_{k=1}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{k}{N}}}{\prod_{k=1+b_{ji}}^{N-1+b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{k-b_{ji}}{N}}} \\
 &= Y_j \prod_{k=1}^{b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{k}{N}} \prod_{k=1+b_{ji}}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}} \prod_{k=N}^{N-1+b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}-k}{N}} \\
 &= Y_j \prod_{k=1}^{b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{k}{N}} \prod_{k=1+b_{ji}}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}} \prod_{k=0}^{b_{ji}-1} (1 + q^{2k+1} Y_i)^{-1 + \frac{b_{ji}-k}{N}}.
 \end{aligned}$$

Now note that

$$\begin{aligned}
 \prod_{k=1+b_{ji}}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}} &= \frac{\prod_{k=0}^{N-1} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}}}{\prod_{k=0}^{b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}}} \\
 &= \frac{(1 + \bar{q} Y_i^N)^{\frac{b_{ji}}{N}}}{\prod_{k=0}^{b_{ji}} (1 + q^{2k+1} Y_i)^{\frac{b_{ji}}{N}}}.
 \end{aligned}$$

Hence combining, most of the terms cancel out, and we are left with

$$\begin{aligned}
 d_N(Y_i) \cdot Y_j \cdot d_N(Y_i)^{-1} &= Y_j \frac{(1 + \bar{q} Y_i^N)^{\frac{b_{ji}}{N}}}{\prod_{k=0}^{b_{ji}-1} (1 + q^{2k+1} Y_i)} \\
 &= Y_j \frac{(1 + \bar{q} Y_i^N)^{\frac{b_{ji}}{N}}}{\prod_{k=1}^{b_{ji}} (1 + q^{2k-1} Y_i)},
 \end{aligned}$$

which is precisely the ratio of the “non-tropical terms”. The case for $b_{ji} < 0$ is similar. \square

Therefore conjugation by d_N will replace the non-tropical terms with its dual version. Now note that the non-tropical terms for Y_i commute with \bar{Y}_i and vice versa. Hence the only arguments that matter in the subsequent conjugations are from the tropical terms only. So the adjoint action follows from the shuffling formula as in the case of the R function, but instead, with the tropical term replaced by the universal form of the dual variables.

In particular, we have

$$\text{Ad} [d_N(Y_{k_1}(1)^{\epsilon_1})^{\epsilon_1} \dots d_N(Y_{k_{r-1}}(t-1)^{\epsilon_{r-1}})^{\epsilon_{r-1}}] (Y_{k_r}(t)) = \bar{Y}_{k_r}(t).$$

The rest of the proof is parallel to the proof of Theorem 2.13, see in particular (17) and (18). \square

Remark 3.11. *The commutation relations (48) and (50) has a finite-dimensional matrix representation, and consequently the cyclic dilogarithm identity also reduces to a matrix identity. For example, $Y_1 Y_2 = \zeta^2 Y_2 Y_1$ (see Example 3.12 below) can be represented by a $N \times N$ matrix*

$$(72) \quad (Y_1)_{ab} = \zeta^{2a} \delta_{ab}, \quad (Y_2)_{ab} = \delta_{a,b+1},$$

where the indices a, b are to be understood modulo N .

3.3. Examples. For concreteness let us now discuss two examples.

Example 3.12 (A_2 quiver). *Let us consider the mutation sequence discussed in Example 2.14. The identity in the dual universal form (3.4) reads*

$$(73) \quad \begin{aligned} & d_N \left(Y_2 (1 + \widetilde{\zeta} \widetilde{Y}_1)^{\frac{1}{N}} \right) d_N(Y_1) \\ &= d_N \left(\frac{Y_1}{(1 + \widetilde{\zeta} \widetilde{Y}_2 + \widetilde{Y}_1 \widetilde{Y}_2)^{\frac{1}{N}}} \right) d_N \left(\frac{\zeta Y_2 Y_1}{(1 + \widetilde{\zeta} \widetilde{Y}_1)^{\frac{1}{N}}} \right) d_N(Y_2) \end{aligned}$$

for $Y_1 Y_2 = \zeta^2 Y_2 Y_1$. One can verify that this coincides with the identity derived in [1]. The identity in the standard form (69) reads

$$(74) \quad \begin{aligned} & d_N(Y_1) d_N(Y_2(1 + qY_1)) d_N \left((1 + qY_2 + Y_1 Y_2)^{-1} Y_1 \right)^{-1} \\ & d_N \left(q(1 + qY_2)^{-1} Y_2 Y_1 \right)^{-1} d_N(Y_2)^{-1} = 1. \end{aligned}$$

Example 3.13 (A_3 quiver). *For the mutation sequence for the A_3 quiver considered in Example 2.15, we have, for $Y_1 Y_2 = \zeta^2 Y_2 Y_1$, $Y_3 Y_2 = \zeta^2 Y_3 Y_2$,*

The identity in the dual universal form (3.4) reads

$$(75) \quad \begin{aligned} & d_N(Y_2 \widetilde{X}_{\widetilde{\zeta}}) d_N(Y_3) d_N(Y_1) \\ &= d_N \left(Y_1 (1 + \widetilde{\zeta} \widetilde{Y}_2 \widetilde{X}_{\widetilde{\zeta}})^{-\frac{1}{N}} \right) d_N \left(Y_3 (1 + \widetilde{\zeta} \widetilde{Y}_2 \widetilde{X}_{\widetilde{\zeta}})^{-\frac{1}{N}} \right) \\ & d_N \left(Y_1 Y_2 Y_3 (1 + \widetilde{\zeta} \widetilde{Y}_2 + \widetilde{Y}_1 \widetilde{Y}_2)^{-\frac{1}{N}} (1 + \widetilde{\zeta} \widetilde{Y}_2 + \widetilde{Y}_3 \widetilde{Y}_2)^{-\frac{1}{N}} \right) \\ & d_N \left(\zeta Y_2 Y_3 (1 + \widetilde{\zeta} \widetilde{Y}_2)^{-\frac{1}{N}} \right) d_N \left(\zeta Y_2 Y_1 (1 + \widetilde{\zeta} \widetilde{Y}_2)^{-\frac{1}{N}} \right) d_N(Y_2), \end{aligned}$$

where $\widetilde{X}_{\widetilde{\zeta}} := (1 + \widetilde{\zeta} \widetilde{Y}_1)^{\frac{1}{N}} (1 + \widetilde{\zeta} \widetilde{Y}_3)^{\frac{1}{N}}$, while the identity in the standard form (69) is given by

$$(76) \quad \begin{aligned} & d_N(Y_1) d_N(Y_3) d_N(Y_2(1 + qY_1)(1 + qY_3)) \\ &= d_N(Y_2) d_N(q(1 + qY_2)^{-1} Y_1 Y_2) d_N(q(1 + qY_2)^{-1} Y_2 Y_3) \\ & d_N((1 + qY_2 + Y_1 Y_2)^{-1} (1 + q^3 Y_2 + Y_3 Y_2)^{-1} Y_3 Y_2 Y_1) \\ & d_N((1 + qY_2 X_q)^{-1} Y_3) d_N((1 + qY_2 X_q)^{-1} Y_1). \end{aligned}$$

with X_q as in (29).

4. CYCLIC DILOGARITHM IDENTITIES (II)

In this section we consider the root of unity limit of the quantum dilogarithm identities for $\Phi_b(x)$ and $g_b(x)$.

4.1. Root of Unity Limit of $g_b(x)$. Let $q = e^{\pi i \frac{M}{N}}$ where M, N are coprime, such that $b = \sqrt{\frac{M}{N}}$. It turns out the integral expression for $\Phi_b(x)$ can be evaluated explicitly, which is discussed in a recent preprint [17]. Let $Q = b + b^{-1}$.⁷

Proposition 4.1. *Let $s = \sqrt{MN}$. Then*

$$(77) \quad \Phi_b \left(\frac{z}{2\pi s} - \frac{iQ}{2} \right) = \frac{e^{\frac{i}{2\pi s^2} \text{Li}_2(e^z)} (1 - e^z)^{1 + \frac{iz}{2\pi s^2}}}{D_N(e^{z/N}; q^2) D_M(e^{z/M}; q^{-2})},$$

⁷In [17] they denote by $c_b = \frac{iQ}{2}$.

where the (Kashaev's) cyclic dilogarithm $D_N(x; q)$ is defined as

$$(78) \quad D_N(x; q) := \prod_{k=1}^{N-1} (1 - q^k x)^{k/N}.$$

In particular, for $M = N = 1$, we have⁸

$$(79) \quad \Phi_1(z) = e^{\frac{i}{2\pi} \text{Li}_2(e^{2\pi z})} (1 - e^{2\pi z})^{iz}.$$

For $M = 1$ but N general, we have

$$(80) \quad \Phi_{\frac{1}{\sqrt{N}}}\left(z - \frac{iQ}{2}\right) = \frac{e^{\frac{i}{2\pi s^2} \text{Li}_2(e^{2\pi s z})} (1 - e^{2\pi s z})^{1 + \frac{iz}{s}}}{D_N(e^{2\pi s z/N}; q^2)} \\ = \frac{(1 - e^{2\pi s z}) \Phi_1(s z)^{\frac{1}{s^2}}}{D_N(e^{2\pi s z/N}; q^2)}.$$

The proof of Proposition 4.1 can be found in [17]. We observe from (80) that the quantum dilogarithm function at $b = 1$ plays a special role in the case of root of unity. Using the results from previous analysis of the cyclic dilogarithm, we derive new identities involving this special $\Phi_1(x)$ function. In the appendix we will present a new proof of (80) based on the τ analysis.

4.2. **Identities for $\mathfrak{g}_{1,N}(u)$.** Let $u = e^{2\pi b z}$. Note that in (80),

$$(81) \quad \Phi_1(s z) = g_1(u^N).$$

Hence we can rewrite (80) using $g_b(u)$ as

$$(82) \quad g_{\frac{1}{\sqrt{N}}}(e^{-\pi i} q^{-1} u) = \frac{(1 - u^N) g_1(u^N)^{\frac{1}{N}}}{D_N(u; q^2)}.$$

(Note we need to keep track of $-1 = e^{\pm\pi i}$ since $g_b(u)$ depends on $\log(u)$.) Hence replacing $u \mapsto e^{\pi i} q u$, we have

$$(83) \quad g_{\frac{1}{\sqrt{N}}}(u) = \frac{(1 + \tilde{q} u^N) g_1(e^{\pi i} \tilde{q} u^N)^{\frac{1}{N}}}{D_N(-q u; q^2)}.$$

Let us denote by

$$(84) \quad \mathfrak{g}_{1,N}(u) := \widehat{g}_1(u^N)^{\frac{1}{N}} (1 + \tilde{q} u^N)^{\frac{1+N}{2N}}, \quad \widehat{g}_1(u) := g_1(e^{\pi i} \tilde{q} u),$$

Then

$$(85) \quad g_{\frac{1}{\sqrt{N}}}(u) = d_N(u) \mathfrak{g}_{1,N}(u),$$

where $d_N(u)$ is our cyclic dilogarithm defined in Definition 3.3. We see that the function $\mathfrak{g}_{1,N}(u)$ plays the role of $R_{\tau,N}(x)$ in our previous analysis. In fact we have

Theorem 4.2. $\mathfrak{g}_{1,N}(u)$ satisfies the quantum dilogarithm identities in the tropical form:

$$(86) \quad \mathfrak{g}_{1,N}(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots \mathfrak{g}_{1,N}(Y_{\epsilon_L c_L})^{\epsilon_L} = 1.$$

Proof. Since $g_{\frac{1}{\sqrt{N}}}(u)$ satisfy the quantum dilogarithm identities, we have

$$d_N(Y_{\epsilon_1 c_1})^{\epsilon_1} \mathfrak{g}_{1,N}(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots d_N(Y_{\epsilon_L c_L})^{\epsilon_L} \mathfrak{g}_{1,N}(Y_{\epsilon_L c_L})^{\epsilon_L} = 1.$$

Now the proof follows from the following lemma:

⁸ This identity is well-known in the literature of supersymmetric gauge theories [21].

Lemma 4.3. *The commutation relation for $\mathfrak{g}_{1,N}(Y_i)$ with Y_j is exactly the same as $R_{\tau,N}(Y_i)$ in the $\tau \rightarrow 0$ limit, i.e., it shuffles as the mutation for the cyclic y -variables:*

$$(87) \quad \mathfrak{g}_{1,N}(Y_i) \cdot Y_j \cdot \mathfrak{g}_{1,N}(Y_i)^{-1} = Y_j(1 + \widetilde{q}Y_i^N)^{-\frac{b_{ji}}{N}}.$$

Proof. We only need to consider \widehat{g}_1 since the extra factor commutes with everything. We have

$$\begin{aligned} \widehat{g}_1(Y_i^N)Y_j &= e^{\frac{i}{2\pi}\text{Li}_2(Y_i^N)}(1 + \widetilde{q}Y_i^N)^{-\frac{\log Y_i^N}{2\pi i}}Y_j \\ &= e^{\frac{i}{2\pi}\text{Li}_2(Y_i^N)}Y_j(1 + \widetilde{q}Y_i^N)^{-\frac{\log q^{2b_{ji}N}Y_i^N}{2\pi i}} \\ &= Y_j e^{\frac{i}{2\pi}\text{Li}_2(Y_i^N)}(1 + \widetilde{q}Y_i^N)^{-b_{ji} - \frac{\log Y_i^N}{2\pi i}} \\ &= Y_j(1 + \widetilde{q}Y_i^N)^{-b_{ji}}\widehat{g}_1(Y_i^N), \end{aligned}$$

and hence

$$\widehat{g}_1(Y_i^N)^{\frac{1}{N}} \cdot Y_j \cdot \widehat{g}_1(Y_i^N)^{-\frac{1}{N}} = Y_j(1 + \widetilde{q}Y_i^N)^{-\frac{b_{ji}}{N}}.$$

□

We therefore obtain, after repetition of shuffling,

$$\mathfrak{g}_{1,N}(Y_{\epsilon_1 c_1})^{\epsilon_1} \cdots \mathfrak{g}_{1,N}(Y_{\epsilon_L c_L})^{\epsilon_L} d_N(\overline{Y}_{k_L}(L)^{\epsilon_L})^{\epsilon_L} \cdots d_N(\overline{Y}_{k_1}(1)^{\epsilon_1})^{\epsilon_1} = 1.$$

Since the product of d_N 's cancel thanks to Theorem 3.4, we obtain (86). □

From the Lemma we see that $\mathfrak{g}_{1,N}(u)$ satisfies the shuffle relations for the cyclic y -variables, and consequently we can derive

Corollary 4.4. *$\mathfrak{g}_{1,N}(u)$ also satisfies the quantum dilogarithm identities in the dual universal form:*

$$(88) \quad \mathfrak{g}_{1,N}(\overline{Y}_{k_L}(L)^{\epsilon_L})^{\epsilon_L} \cdots \mathfrak{g}_{1,N}(\overline{Y}_{k_1}(1)^{\epsilon_1})^{\epsilon_1} = 1.$$

Remark 4.5. *Using the functional equations for g_b , one can see that equation (83) is also equivalent to the identity for $\zeta = e^{2\pi i/N}$:*

$$(89) \quad \prod_{k=1}^N g_{\frac{1}{\sqrt{N}}}(\zeta^k z) = g_1(e^{\pi i} \widetilde{q} z^N),$$

which is a generalization of the classical dilogarithm identity

$$(90) \quad N \sum_{k=1}^N \text{Li}_2(\zeta^k x) = \text{Li}_2(x^N).$$

Finally, similarly analysis can be done for general $q = e^{\pi i \frac{M}{N}}$ with M, N coprime, using the fact that the commutation relation for $\mathfrak{g}_{1,N}(Y_i)$ is also true for the modular double variables due to the modular duality of $g_b(x)$. Let $b = \sqrt{\frac{M}{N}}$ and $u = e^{2\pi b z}$, and denote by

$$(91) \quad \mathfrak{g}_{1,N,M}(u) := g_1(e^{\pi i(M+N)} u^N)^{\frac{1}{MN}} (1 + e^{\pi i(M+N)} u^N)^{\frac{1}{2N} + \frac{1}{2M}},$$

note that $u^N = \widetilde{u}^M$. Then using the fact that $D_N(u; q^2)$ commutes with $D_M(\widetilde{u}; \widetilde{q}^2)$, we have

Theorem 4.6. *The function $\mathfrak{g}_{1,N,M}(u)$ satisfies the same quantum dilogarithm identities as $\mathfrak{g}_{1,N}(u)$, namely we have (86) with $\mathfrak{g}_{1,N}(u)$ replaced by $\mathfrak{g}_{1,N,M}(u)$.*

APPENDIX A. PROOF OF (80)

In this subsection we present a new proof of (80), and as a consequence another proof for Theorem 4.2. Similar analysis can be used to prove Proposition 4.1 in its full generality for $q = e^{\pi i \frac{M}{N}}$ but we will omit it for simplicity.

Proof of (80). By definition, we have for $\text{Im}(b^2) > 0$

$$g_b(x) = \frac{(-qx; q^2)_\infty}{(-\bar{q}^{-1}\bar{x}; \bar{q}^{-2})_\infty},$$

or under rescaling

$$g_b(e^{\pi i} q^{-1} x) = \frac{(x; q^2)_\infty}{(\bar{x}; \bar{q}^{-2})_\infty}.$$

Again, note that we need to keep track of $-1 = e^{\pm \pi i}$ since $g_b(x)$ involves $\log(x)$.

Now recall from (62) we have

$$(92) \quad \begin{aligned} (x; q)_\infty &= (1 - x^N)^{\frac{1-N}{2N}} R_\tau^*(x^N) d_N^*(x) (1 + O(\tau)) \\ &= \frac{R_\tau^*(x^N)}{D_N(x; q)} (1 + O(\tau)), \end{aligned}$$

where $R_\tau^*(x) := (1 - x)^{\frac{1}{2}} e^{-\text{Li}_2(x)/\tau}$.

Let $q = e^{-\tau/2N^2} \zeta = e^{-\tau/2N^2} e^{\pi i/N}$. If we write $q = e^{\pi i b^2}$, then $b^2 = \frac{2\pi N + i\tau}{2\pi N^2}$ and

$$\bar{q}^{-1} = e^{-\pi i b^{-2}} = e^{-\frac{2\pi^2 N^2 i}{2\pi N + i\tau}} = e^{-\pi i N - \tau/2 + i\tau^2/4\pi N + O(\tau^3)}.$$

Let us now derive the asymptotics of $(x; \bar{q}^{-2})_\infty$. We apply Euler-Maclaurin formula to its logarithm, following the similar idea from [1]. We obtain

$$\begin{aligned} (x; \bar{q}^{-2})_\infty &= (1 - x)^{\frac{1}{2}} e^{\frac{\text{Li}_2(x)}{\log(\bar{q}^{-2})}} \\ &= (1 - x)^{\frac{1}{2}} e^{-\tau + \frac{i\tau^2}{2\pi N} + O(\tau^3)} \\ &= (1 - x)^{\frac{1}{2}} e^{\frac{\text{Li}_2(x)}{-\tau} (1 + \frac{\tau}{2\pi i N})} (1 + O(\tau)) \\ &= R_\tau^*(x) e^{-\frac{1}{2\pi i N} \text{Li}_2(x)} (1 + O(\tau)), \end{aligned}$$

and hence

$$(93) \quad \begin{aligned} g_{\frac{1}{\sqrt{N}}}(e^{\pi i} q^{-1} x) &= \lim_{\tau \rightarrow 0} \frac{(x; q^2)_\infty}{(\bar{x}; \bar{q}^{-2})_\infty} \\ &= \frac{R_\tau^*(x^N)}{R_\tau^*(\bar{x})} \Big|_{\tau \rightarrow 0} \frac{e^{-\frac{1}{2\pi i N} \text{Li}_2(\bar{x})}}{D_N(x; q^2)} (1 + O(\tau)). \end{aligned}$$

The limit

$$\lim_{\tau \rightarrow 0} \frac{R_\tau^*(x^N)}{R_\tau^*(\bar{x})} = \lim_{\tau \rightarrow 0} e^{-\text{Li}_2(x^N)/\tau + \text{Li}_2(\bar{x})/\tau}$$

can be computed as (see (67) for a similar computation)

$$\begin{aligned} \lim_{\tau \rightarrow 0} -\text{Li}_2(x)/\tau + \text{Li}_2(\tilde{x})/\tau &= \lim_{\tau \rightarrow 0} \frac{1}{\tau} \left(\int_0^{x^N} \frac{\log(1-t)}{t} dt - \int_0^{\tilde{x}} \frac{\log(1-t)}{t} dt \right) \\ &= \lim_{\tau \rightarrow 0} -\frac{\log(1-x(\tau))}{x(\tau)} \frac{d\tilde{x}}{d\tau} \\ &= -\frac{\log(1-x^N)\log(x)}{2\pi i}, \end{aligned}$$

where we used $\tilde{x} = x^{\frac{1}{b^2}} = x^{\frac{2\pi N^2}{2\pi N + i\tau}} = x^{N + \frac{\tau}{2\pi i} + O(\tau^2)}$. We therefore obtain

$$(94) \quad \begin{aligned} g_{\frac{1}{\sqrt{N}}}(e^{\pi i} q^{-1} x) \Big|_{b^2 \rightarrow \frac{1}{N}} &= \frac{e^{\frac{i}{2\pi N} \text{Li}_2(x^N)} (1-x^N)^{-\frac{\log(x)}{2\pi i}}}{D_N(x; q^2)} \\ &= \frac{g_1(x^N)^{\frac{1}{N}}}{D_N(x; q^2)}. \end{aligned}$$

Shifting $x \rightarrow e^{-2\pi i} x$ to the other side of the branch cut, we obtain

$$g_{\frac{1}{\sqrt{N}}}(e^{-\pi i} q^{-1} x) \Big|_{b^2 \rightarrow \frac{1}{N}} = \frac{e^{\frac{i}{2\pi N} \text{Li}_2(x^N)} (1-x^N)^{1-\frac{\log(x)}{2\pi i}}}{D_N(x; q^2)},$$

or in terms of $x = e^{2\pi b z}$:

$$\Phi_{\frac{1}{\sqrt{N}}}\left(z - \frac{iQ}{2}\right) = \frac{e^{\frac{i}{2\pi N} \text{Li}_2(e^{2\pi N b z})} (1 - e^{2\pi N b z})^{1+ibz}}{D_N(e^{2\pi b z}; q^2)},$$

which is identical to (80). \square

As a consequence of this analysis, we provide another proof of Theorem 4.2.

Proof of Theorem 4.2. Again let $q = e^{\pi i b^2} = e^{-\tau/2N^2} e^{\pi i/N}$. From equation (92), (93) and (94), we conclude that

$$\Phi_1(s z)^{\frac{1}{N}} = g_1(e^{2\pi \sqrt{N} z})^{\frac{1}{N}} = \lim_{\tau \rightarrow 0} \frac{R_{\tau}^*(x^N)}{(\tilde{x}; \tilde{q}^{-2})_{\infty}},$$

where $x = e^{2\pi b z}$. Now we substitute $z \mapsto z - \frac{iQ}{2}$ so that $\tilde{x} \mapsto -\tilde{q}^{-1} \tilde{x}$, then by definition we have

$$x^N = e^{2\pi b N z} \mapsto -e^{\tau/2N} (-x)^N (1 + O(\tau)),$$

hence

$$\begin{aligned} R_{\tau}^*(x^N) &\rightarrow R_{\tau}^*(-e^{\tau/2N} (-x)^N) \\ &= R_{\tau}^*(-e^{\tau(1+N)/2N} e^{-\tau/2} (-x)^N) \\ &= (1 + (-x)^N)^{\frac{1+N}{2N}} R_{\tau}^*(-e^{-\tau/2} (-x)^N) (1 + O(\tau)) \\ &= (1 + \tilde{q} x^N)^{\frac{1+N}{2N}} R_{\tau, N}(x) (1 + O(\tau)), \end{aligned}$$

where $R_{\tau, N}(x)$ is defined in (64), and we have used the identity

$$R_{\tau}^*(e^{\alpha\tau} x) = (1-x)^{\alpha} R_{\tau}^*(x) (1 + O(\tau)).$$

Hence we have

$$\Phi_1\left(s\left(z - \frac{iQ}{2}\right)\right)^{\frac{1}{N}} = \lim_{\tau \rightarrow 0} (1 + \tilde{q} x^N)^{\frac{1+N}{2N}} R_{\tau, N}(x) \Psi_{\tilde{q}^{-1}}(\tilde{x}).$$

Now note that \tilde{x} commute with x^N (this does not involve the τ limit), hence we can move the $R_{\tau,N}$ to one side and $\Psi_{\tilde{q}^{-1}}(\tilde{x})$ to the other side. Then since both $R_{\tau,N}(x)$ and $\Psi_{\tilde{q}^{-1}}(\tilde{x})$ satisfy the quantum dilogarithm identities, we conclude that

$$\begin{aligned} (1 + \tilde{q}x^N)^{-\frac{1+N}{2N}} \Phi_1\left(s\left(z - \frac{iQ}{2}\right)\right)^{\frac{1}{N}} &= (1 + \tilde{q}x^N)^{-\frac{1+N}{2N}} g_1(e^{-\pi i \tilde{q}^{-1}} x^N)^{\frac{1}{N}} \\ &= (1 + \tilde{q}x^N)^{\frac{1+N}{2N}} g_1(e^{\pi i \tilde{q}} x^N)^{\frac{1}{N}} \\ &= \mathfrak{g}_{1,N}(x) \end{aligned}$$

also satisfies the quantum dilogarithm identities. \square

REFERENCES

- [1] V. V. Bazhanov and N. Y. Reshetikhin, *Remarks on the quantum dilogarithm*, J. Phys. A **28** (1995), no. 8, 2217–2226. MR 1338071 (96k:81087)
- [2] V. V. Bazhanov, V. V. Mangazeev, and S. M. Sergeev, *Faddeev-Volkov solution of the Yang-Baxter equation and discrete conformal symmetry*, Nucl.Phys. **B784** (2007), 234–258.
- [3] V. Bazhanov and R. Baxter, *Star triangle relation for a three-dimensional model*, J.Statist.Phys. **71** (1993), 839–864.
- [4] A. G. Bytsko and J. Teschner, *R-operator, co-product and Haar-measure for the modular double of $U_q(\mathfrak{sl}(2, \mathbb{R}))$* , Comm. Math. Phys. **240** (2003), no. 1-2, 171–196. MR 2004985 (2004m:17019)
- [5] S. Cecotti, A. Neitzke, and C. Vafa, *R-Twisting and 4d/2d Correspondences*, (2010).
- [6] H. Derksen, J. Weyman, and A. Zelevinsky, *Quivers with potentials and their representations II: applications to cluster algebras*, J. Amer. Math. Soc. **23** (2010), no. 3, 749–790. MR 2629987 (2012c:16044)
- [7] L. Faddeev, *Current - like variables in massive and massless integrable models*, (1994).
- [8] ———, *Discrete Heisenberg-Weyl group and modular group*, Lett.Math.Phys. **34** (1995), 249–254.
- [9] L. Faddeev and R. Kashaev, *Quantum Dilogarithm*, Mod.Phys.Lett. **A9** (1994), 427–434.
- [10] L. Faddeev and A. Y. Volkov, *Abelian current algebra and the Virasoro algebra on the lattice*, Phys.Lett. **B315** (1993), 311–318.
- [11] V. V. Fock and A. B. Goncharov, *The quantum dilogarithm and representations of quantum cluster varieties*, Invent. Math. **175** (2009), no. 2, 223–286. MR 2470108 (2010e:14021)
- [12] V. V. Fock and A. B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 6, 865–930. MR 2567745 (2011f:53202)
- [13] S. Fomin and A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529 (electronic). MR 1887642 (2003f:16050)
- [14] ———, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164. MR 2295199 (2008d:16049)
- [15] I. B. Frenkel and I. C. H. Ip, *Positive representations of split real quantum groups and future perspectives*, Int. Math. Res. Not. IMRN (2014), no. 8, 2126–2164. MR 3194015
- [16] D. Gaiotto, G. W. Moore, and A. Neitzke, *Framed BPS States*, Adv.Theor.Math.Phys. **17** (2013), 241–397.
- [17] S. Garoufalidis and R. Kashaev, *Evaluation of state integrals at rational points*, (2014).
- [18] K. Hikami and R. Inoue, *Braiding Operator via Quantum Cluster Algebra*, J. Phys. A **47** (2014), 474006.
- [19] I. C. H. Ip, *The classical limit of representation theory of the quantum plane*, Internat. J. Math. **24** (2013), no. 4, 1350031, 40. MR 3062971
- [20] I. C.-H. Ip, *Representation of the quantum plane, its quantum double, and harmonic analysis on $GL_q^+(2, \mathbb{R})$* , Selecta Math. (N.S.) **19** (2013), no. 4, 987–1082. MR 3131494
- [21] D. L. Jafferis, *The Exact Superconformal R-Symmetry Extremizes Z*, JHEP **1205** (2012), 159.
- [22] R. M. Kashaev, *Quantum dilogarithm as a 6j-symbol*, Modern Phys. Lett. A **9** (1994), no. 40, 3757–3768. MR 1317945 (96c:17024)
- [23] R. M. Kashaev and T. Nakanishi, *Classical and quantum dilogarithm identities*, SIGMA Symmetry Integrability Geom. Methods Appl. **7** (2011), Paper 102, 29. MR 2861174
- [24] B. Keller, *On cluster theory and quantum dilogarithm identities*, Representations of algebras and related topics, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011, pp. 85–116. MR 2931896
- [25] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, (2008).

- [26] K. Nagao, *Quantum dilogarithm identities*, Infinite analysis 2010—Developments in quantum integrable systems, RIMS Kokyuroku Bessatsu, B28, Res. Inst. Math. Sci. (RIMS), Kyoto, 2011, pp. 165–170. MR 2885304
- [27] ———, *Donaldson-Thomas theory and cluster algebras*, Duke Math. J. **162** (2013), no. 7, 1313–1367. MR 3079250
- [28] K. Nagao, Y. Terashima, and M. Yamazaki, *Hyperbolic 3-manifolds and Cluster Algebras*, (2011).
- [29] P.-G. Plamondon, *Cluster algebras via cluster categories with infinite-dimensional morphism spaces*, Compos. Math. **147** (2011), no. 6, 1921–1954. MR 2862067
- [30] B. Ponsot and J. Teschner, *Liouville bootstrap via harmonic analysis on a noncompact quantum group*, (1999).
- [31] M. Reineke, *Poisson automorphisms and quiver moduli*, J. Inst. Math. Jussieu **9** (2010), no. 3, 653–667. MR 2650811 (2011h:16021)
- [32] K. Schmüdgen, *Integrable operator representations of \mathbb{R}_q^2 , $X_{q,\gamma}$ and $SL_q(2, \mathbb{R})$* , Comm. Math. Phys. **159** (1994), no. 2, 217–237. MR 1256987 (94k:81129)
- [33] Y. Terashima and M. Yamazaki, *Emergent 3-manifolds from 4d Superconformal Indices*, Phys.Rev.Lett. **109** (2012), 091602.
- [34] ———, *3d N=2 Theories from Cluster Algebras*, PTEP **023** (2014), B01.
- [35] D. Xie, *BPS spectrum, wall crossing and quantum dilogarithm identity*, (2012).
- [36] M. Yamazaki, *Quivers, YBE and 3-manifolds*, JHEP **1205** (2012), 147.

KAVLI IPMU (WPI), UNIVERSITY OF TOKYO, CHIBA 277-8583, JAPAN

E-mail address: ivan.ip@ipmu.jp, masahito.yamazaki@ipmu.jp