

Masses of physical scalars in two Higgs doublet models

Ambalika Biswas* and Amitabha Lahiri†

*S. N. Bose National Centre For Basic Sciences,
Block JD, Sector III, Salt Lake, Kolkata 700098, INDIA*

(Dated: May 9, 2022)

Abstract

We find bounds on scalar masses resulting from a criterion of naturalness, in a broad class of two Higgs doublet models (2HDMs). Specifically, we assume the cancellation of quadratic divergences in what are called the type I, type II, lepton-specific and flipped 2HDMs, with an additional U(1) symmetry. This results in a set of relations among masses of the physical scalars and coupling constants, a generalization of the Veltman conditions of the Standard Model. Assuming that the lighter CP -even neutral Higgs particle is the observed scalar particle of mass ~ 125 GeV, and imposing further the constraints from the electroweak T-parameter, stability, and perturbative unitarity, we calculate the range of the mass of each of the remaining physical scalars.

arXiv:1412.6187v1 [hep-ph] 19 Dec 2014

* ambalika12t@boson.bose.res.in

† amitabha@bose.res.in

I. INTRODUCTION

With the discovery of a 125 GeV neutral scalar boson [1, 2], the menagerie of fundamental particles in the Standard Model seems to be complete. Some questions still remain unanswered, including the origins of neutrino mass and dark matter, keeping the door open for physics beyond the Standard Model. Among the simplest extensions of the Standard Model are two Higgs doublet models (2HDMs). Originally motivated by supersymmetry, where a second Higgs doublet is essential, 2HDMs have also been studied independently (for a recent review see [3]).

Axions provide another motivation for 2HDMs. Peccei-Quinn symmetry [4, 5] solves the strong CP problem, but must be spontaneously broken. The corresponding Goldstone boson is the axion, which can be a combination of the phases of two Higgs doublets. Models of baryogenesis often involve 2HDMs [6] because their mass spectrum can be adjusted to produce CP violation, both explicit and spontaneous. Another motivation, one that is important to us, is their use in models of dark matter [7–9]. These models are the inert doublet models, so called because one of the Higgs doublets does not couple to the fermions. Of the 2HDMs we will consider, one model tends to the inert doublet model for large values of the ratio of the vacuum expectation values (VEVs) of the two Higgs fields. The other models also have small couplings to one or more types of fermions in that limit.

In this paper we will consider 2HDMs with a softly broken global U(1) symmetry [4, 10]. We choose the fermion transformations under this symmetry, and impose a naturalness condition of vanishing quadratic divergences on the scalar sector of the models. Using additional restrictions coming from partial wave unitarity, vacuum stability, and the T parameter measuring ‘new physics’, we find bounds on the masses of the additional scalar fields for each of the 2HDMs.

We will work with the scalar potential [11, 12]

$$\begin{aligned}
 V = & \lambda_1 \left(\Phi_1^\dagger \Phi_1 - \frac{v_1^2}{2} \right)^2 + \lambda_2 \left(\Phi_2^\dagger \Phi_2 - \frac{v_2^2}{2} \right)^2 \\
 & + \lambda_3 \left(\Phi_1^\dagger \Phi_1 + \Phi_2^\dagger \Phi_2 - \frac{v_1^2 + v_2^2}{2} \right)^2 + \lambda_4 \left((\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) - (\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1) \right) \\
 & + \lambda_5 \left(\Re(\Phi_1^\dagger \Phi_2) - \frac{1}{2}v_1v_2 \right)^2 + \lambda_6 \left(\Im(\Phi_1^\dagger \Phi_2) \right)^2, \tag{1.1}
 \end{aligned}$$

where all the parameters λ_i are real. This potential is invariant under the symmetry $\Phi_1 \rightarrow -\Phi_1, \Phi_2 \rightarrow \Phi_2$, except for the term $\lambda_5 v_1 v_2 \Re(\Phi_1^\dagger \Phi_2)$, which softly breaks the symmetry. Additional dimension-4 terms are also set to zero by this symmetry.

The scalar doublets take the vacuum expectation values (vevs) v_1, v_2 , which can be taken to be real and positive without any loss of generality. We can then parametrize the scalar doublets as

$$\Phi_i = \left(\begin{array}{c} w_i^+(x) \\ \frac{v_i + h_i(x) + iz_i(x)}{\sqrt{2}} \end{array} \right). \quad i = 1, 2 \tag{1.2}$$

Three of these fields get ‘eaten’ by the W^\pm and Z^0 gauge bosons; the remaining five are physical scalar (Higgs) fields. There is a pair of charged scalars denoted by ξ^\pm , two neutral CP even scalars, the heavy one denoted by H and the light one denoted by h , and one CP

odd pseudoscalar denoted by A . These fields are given by the combinations

$$\begin{pmatrix} \omega^\pm \\ \xi^\pm \end{pmatrix} = \begin{pmatrix} c_\beta & s_\beta \\ -s_\beta & c_\beta \end{pmatrix} \begin{pmatrix} w_1^\pm \\ w_2^\pm \end{pmatrix}, \quad (1.3)$$

$$\begin{pmatrix} \zeta \\ A \end{pmatrix} = \begin{pmatrix} c_\beta & s_\beta \\ -s_\beta & c_\beta \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \quad (1.4)$$

$$\begin{pmatrix} H \\ h \end{pmatrix} = \begin{pmatrix} c_\alpha & s_\alpha \\ -s_\alpha & c_\alpha \end{pmatrix} \begin{pmatrix} h_1 \\ h_2 \end{pmatrix}, \quad (1.5)$$

where we have written $c_\alpha \equiv \cos \alpha$ and $s_\alpha \equiv \sin \alpha$, and likewise for c_β, s_β , and

$$\tan \beta = \frac{v_2}{v_1}. \quad (1.6)$$

We note here that if we rotate the basis $h_1 - h_2$ by the angle β , we obtain the states

$$\begin{pmatrix} H^0 \\ R \end{pmatrix} = \begin{pmatrix} c_\beta & s_\beta \\ -s_\beta & c_\beta \end{pmatrix} \begin{pmatrix} h_1 \\ h_2 \end{pmatrix}, \quad (1.7)$$

which has the property that H^0 has the exact Standard Model couplings with the fermions and gauge bosons [13]. The field R does not have any cubic gauge coupling at the tree level. It can however have flavor changing Yukawa couplings. The lighter CP even mass eigenstate h is related to H^0 and R via the transformation;

$$h = \sin(\beta - \alpha)H^0 + \cos(\beta - \alpha)R \quad (1.8)$$

Thus in order for h to be the Higgs boson of the Standard Model, we require

$$\sin(\beta - \alpha) \approx 1, \quad (1.9)$$

which has been referred to as the *decoupling limit* [14]. Accordingly, we will assume $\beta - \alpha = \frac{\pi}{2}$ in the rest of the paper.

II. VELTMAN CONDITIONS

Naturalness conditions play an important role in constructing models of physics beyond the Standard Model. The scalar masses get quadratically divergent contributions which need to be cancelled if we wish to avoid fine-tuning of parameters. We will thus require a generalization of the Veltman conditions [15], that the masses and couplings in our 2HDM are such that the quadratically divergent contributions to all the scalar masses cancel. Of course, the resulting condition should not then require a fine-tuning.

The Yukawa potential for the 2HDMs is of the form

$$\mathcal{L}_Y = \sum_{i=1,2} \left[-\bar{l}_L \Phi_i G_e^i e_R - \bar{Q}_L \tilde{\Phi}_i G_u^i u_R - \bar{Q}_L \Phi_i G_d^i d_R + h.c. \right], \quad (2.1)$$

where l_L, Q_L are 3-vectors of isodoublets in the space of generations, e_R, u_R, d_R are 3-vectors of singlets, G_e^1 etc. are complex 3×3 matrices in generation space containing the Yukawa coupling constants, and $\tilde{\Phi}_i = i\tau_2 \Phi_i^*$.

Cancellation of quadratic divergences in the scalar masses gives rise to four mass relations, which we may call the Veltman conditions for the 2HDMs being considered [16],

$$2\text{Tr } G_e^1 G_e^{1\dagger} + 6\text{Tr } G_u^{1\dagger} G_u^1 + 6\text{Tr } G_d^1 G_d^{1\dagger} = \frac{9}{4}g^2 + \frac{3}{4}g'^2 + 6\lambda_1 + 10\lambda_3 + \lambda_4 + \frac{1}{2}(\lambda_5 + \lambda_6), \quad (2.2)$$

$$2\text{Tr } G_e^2 G_e^{2\dagger} + 6\text{Tr } G_u^{2\dagger} G_u^2 + 6\text{Tr } G_d^2 G_d^{2\dagger} = \frac{9}{4}g^2 + \frac{3}{4}g'^2 + 6\lambda_2 + 10\lambda_3 + \lambda_4 + \frac{1}{2}(\lambda_5 + \lambda_6), \quad (2.3)$$

$$2\text{Tr } G_e^1 G_e^{2\dagger} + 6\text{Tr } G_u^{1\dagger} G_u^2 + 6\text{Tr } G_d^1 G_d^{2\dagger} = 0, \quad (2.4)$$

where g, g' are the $SU(2)$ and $U(1)_Y$ coupling constants. A fourth equation is the complex conjugate of the third one. Note that these equalities have a slightly different form in [16] because the coefficients of the quartic terms in the scalar potential are also a little different there. As we will see below, the last equation vanishes identically for all the 2HDMs we consider. The mass relations come from the first two equations above.

When the fermions are in the mass eigenstates, the Yukawa interactions are automatically diagonal if there is only one Higgs doublet as in the Standard Model, so there is no FCNC at the tree level. But in the presence of a second scalar doublet, the two Yukawa matrices will not be simultaneously diagonalizable in general, and thus the Yukawa couplings will not be flavor diagonal. Neutral Higgs scalars will mediate FCNCs. It was shown in [17, 18] that the necessary and sufficient condition for the absence of FCNC at tree level is that all fermions of a given charge and helicity transform according to the same irreducible representation of $SU(2)$, corresponding to the same eigenvalue of T_3 , and that a basis exists in which they receive their contributions in the mass matrix from a single source.

For the fermions of the Standard Model, this theorem implies that all right-handed singlets of a given charge must couple to the same Higgs doublet. In this paper we will ensure this by imposing a global $U(1)$ symmetry, a generalization of the Z_2 symmetry more commonly employed for this purpose, which was used in restricting the form of the scalar potential in Eq. (1.1). Under this $U(1)$ symmetry, the scalar doublets transform as $\Phi_1 \rightarrow e^{i\theta}\Phi_1, \Phi_2 \rightarrow \Phi_2$. On the scalar potential of Eq. (1.1) this imposes the condition $\lambda_5 = \lambda_6$, so that the quartic part of the last two terms combine as

$$\lambda_5 \left| \Phi_1^\dagger \Phi_2 - \frac{v_1 v_2}{2} \right|^2, \quad (2.5)$$

and the quadratic part of this is $\lambda_5 v_1 v_2 \Re(\Phi_1^\dagger \Phi_2)$ as before, which again breaks the symmetry softly.

The left handed fermion doublets remain unchanged under this symmetry, $Q_L \rightarrow Q_L, l_L \rightarrow l_L$. The transformations of right handed fermion singlets determine the type of 2HDM. There are four such possibilities, which may be identified by the right-handed fields which transform under the $U(1)$: type I (none), type II ($d_R \rightarrow e^{-i\theta}d_R, e_R \rightarrow e^{-i\theta}e_R$), lepton specific ($e_R \rightarrow e^{-i\theta}e_R$), flipped ($d_R \rightarrow e^{-i\theta}d_R$).

We note in passing that another way of avoiding FCNCs at tree level is by aligning the Yukawa and mass matrices in flavor space [19]. We do not consider that mechanism in this paper. We also note that for large $\tan\beta$ the type I model approaches the inert doublet model.

III. VELTMAN CONDITIONS FOR THE VARIOUS TYPES OF 2HDMS

The fermion mass matrix is diagonalized by the usual procedure of independent unitary transformations on the left and right-handed fermion fields. This produces the physical fermion fields, and the fermion masses are related to the Yukawa matrices by [16]

$$\text{Tr} [G_{1f}^\dagger G_{1f}] = \frac{2}{v^2} \cos^2 \beta \sum m_f^2, \quad (3.1)$$

$$\text{Tr} [G_{2f}^\dagger G_{2f}] = \frac{2}{v^2} \sin^2 \beta \sum m_f^2, \quad (3.2)$$

for each of the cases where f stands for charged leptons, up-type quarks, or down-type quarks, and the sum is taken over generations. Let us now see what these equations and the Veltman conditions imply for the different 2HDMS, with the global $U(1)$ symmetry.

We first multiply Eq. (2.2) by v^2 to write the constants in terms of particle masses. Then the right hand side becomes

$$6M_W^2 + 3M_Z^2 + v^2 (6\lambda_1 + 10\lambda_3 + \lambda_4 + \lambda_5). \quad (3.3)$$

Next we look at the scalar potential of 2HDM given in Eq. (1.1), which depends on the independent parameters v_1, v_2 and the six λ parameters. In the decoupling limit and with the global $U(1)$ symmetry, the independent parameters may be taken to be the masses m_h, m_H, m_ξ , the angle β , the electroweak VEV v , and the constant λ_5 . The masses of the physical scalar particles are related to these parameters by [20]

$$\lambda_1 = \frac{1}{2v^2 c_\beta^2} m_H^2 - \frac{\lambda_5}{4} (\tan^2 \beta - 1), \quad (3.4)$$

$$\lambda_2 = \frac{1}{2v^2 s_\beta^2} m_H^2 - \frac{\lambda_5}{4} \left(\frac{1}{\tan^2 \beta} - 1 \right), \quad (3.5)$$

$$\lambda_3 = -\frac{1}{2v^2} (m_H^2 - m_h^2) - \frac{\lambda_5}{4}, \quad (3.6)$$

$$\lambda_4 = \frac{2}{v^2} m_\xi^2, \quad (3.7)$$

$$\lambda_6 = \frac{2}{v^2} m_A^2 = \lambda_5. \quad (3.8)$$

Inserting these equations for λ into the expression for VC1 (3.3) we can write it in terms of the physical masses,

$$6M_W^2 + 3M_Z^2 + m_H^2 (3 \tan^2 \beta - 2) + 5m_h^2 + 2m_\xi^2 - \frac{3v^2}{2} \lambda_5 \tan^2 \beta. \quad (3.9)$$

Similarly, multiplying Eq. (2.3) by v^2 we get the corresponding expression for what we will call the second Veltman condition (VC2) for each of the 2HDMS.

$$6M_W^2 + 3M_Z^2 + m_H^2 \left(\frac{3}{\tan^2 \beta} - 2 \right) + 5m_h^2 + 2m_\xi^2 - \lambda_5 \frac{3v^2}{2 \tan^2 \beta}. \quad (3.10)$$

For the different types of 2HDMS, these equal different combinations of the fermion masses. For example, in the Type II 2HDM, Φ_2 couples to up-type quarks, while Φ_1 couples to

down-type quarks and charged leptons, so $G_{2e} = G_{2d} = G_{1u} = 0$. The quantity in (3.9) is then equal to $4 [\sum m_e^2 + 3 \sum m_d^2] \cos^2 \beta$.

Table I lists the quantities corresponding to each 2HDM which equals (3.9) and (3.10). The second column lists the Yukawa matrices which vanish for each type of 2HDM. For each 2HDM, the entries in the third and fourth columns of Table I are equated to these expressions to provide the Veltman conditions specific to each type.

Model	vanishing Yukawa	VC1	VC2
		$6M_W^2 + 3M_Z^2 + 5m_h^2 + 2m_\xi^2 + m_H^2(3 \tan^2 \beta - 2) - \frac{3v^2}{2} \lambda_5 \tan^2 \beta$	$6M_W^2 + 3M_Z^2 + 5m_h^2 + 2m_\xi^2 + m_H^2 \left(\frac{3}{\tan^2 \beta} - 2 \right) - \lambda_5 \frac{3v^2}{2 \tan^2 \beta}$
Type I	G_{1e}, G_{1d}, G_{1u}	0	$4 [\sum m_e^2 + 3 \sum m_u^2 + 3 \sum m_d^2] \sin^2 \beta$
Type II	G_{2e}, G_{2d}, G_{1u}	$4 [\sum m_e^2 + 3 \sum m_d^2] \cos^2 \beta$	$12 \sum m_u^2 \sin^2 \beta$
Lepton-specific	G_{2e}, G_{1d}, G_{1u}	$4 \sum m_e^2 \cos^2 \beta$	$12 [\sum m_u^2 + \sum m_d^2] \sin^2 \beta$
Flipped	G_{1e}, G_{2d}, G_{1u}	$12 \sum m_d^2 \cos^2 \beta$	$4 [\sum m_e^2 + 3 \sum m_u^2] \sin^2 \beta$

TABLE I: Veltman conditions for the different 2HDMs

IV. BOUNDS ON THE MASSES OF HEAVY AND CHARGED SCALARS

We now display our main results, the bounds we have obtained for the masses of the heavy and charged Higgs particles using the Veltman conditions found earlier. For our calculations, we will assume that the h particle is the one that has been observed at the LHC, so that $m_h = 125$ GeV, and $v = \sqrt{v_1^2 + v_2^2} = 246$ GeV. We note that the mass of the charged Higgs particle is known to have a rough lower bound $m_\xi > 100$ GeV, obtained from direct searches [21], however we will not need to use that bound, but will obtain bounds on m_ξ from our calculations. Let us consider one example, say the type II model, to explain our derivation of the bounds.

Consider VC1 for the type II 2HDM, as given in Table I. Since we want the bounds on m_H and m_ξ , let us rewrite this in a form that will prove convenient,

$$m_H^2 (3 \tan^2 \beta - 2) + 2m_\xi^2 = 4 \left[\sum m_e^2 + 3 \sum m_d^2 \right] \cos^2 \beta - 6M_W^2 - 3M_Z^2 - 5m_h^2 + \lambda_5 \frac{3v^2}{2} \tan^2 \beta. \quad (4.1)$$

We note that on the right hand side, all but the last term are experimentally known. The other Veltman condition for the type II model gives the equation

$$m_H^2 \left(\frac{3}{\tan^2 \beta} - 2 \right) + 2m_\xi^2 = 12 \sum m_u^2 \sin^2 \beta - 6M_W^2 - 3M_Z^2 - 5m_h^2 + \lambda_5 \frac{3v^2}{2 \tan^2 \beta}. \quad (4.2)$$

The $U(1)$ symmetry we have assumed implies that $\lambda_5 > 0$, while tree-level unitarity requires [14, 20, 22] $|\lambda_i| \leq 4\pi$. For a fixed value of $\tan \beta$, we plot both equations on the $m_H - m_\xi$ plane for various values of λ_5 . The point where the two curves for a given value of λ_5 cross, is an allowed value of the pair (m_H, m_ξ) .

We can restrict the allowed range of the masses even further by imposing constraints coming from stability, perturbative unitarity, and the oblique electroweak T -parameter.

Conditions for stability, i.e. for the scalar potential being bounded from below were examined in [3, 14, 25], and found to provide lower bounds on certain combinations of the quartic couplings λ_i . On the other hand, the requirement of perturbative unitarity translates into upper limits on combinations of the λ_i , which for multi-Higgs models have been derived by many authors [20, 26–28].

The conditions of stability and unitarity imply, among other things, that

$$0 \leq \lambda_1 + \lambda_2 + 2\lambda_3 \leq \frac{16\pi}{3} \quad (4.3)$$

In terms of the scalar masses in Eq.(3.4) - (3.8), this reads

$$0 \leq (m_H^2 - m_A^2)(\tan^2 \beta + \cot^2 \beta) + 2m_h^2 \leq \frac{32\pi v^2}{3}. \quad (4.4)$$

For $\tan \beta$ far away from unity, this inequality implies that m_H and m_A are almost degenerate as we can see in Fig. 1, a result also found recently in [29]. There is also a similar correlation




FIG. 1: Degeneracy of $m_A - m_H$ for increasing $\tan \beta$

between m_H and m_ξ , but without any dependence on $\tan \beta$. This can be seen from the inequality

$$|2m_\xi^2 - m_H^2 - m_A^2 + m_h^2| \leq 16\pi v^2 \quad (4.5)$$

which also follows from the unitarity constraint. The oblique parameter T for the 2HDMs in the decoupling limit has the expression [30, 31]

$$T = \frac{1}{16\pi \sin^2 \theta_W M_W^2} [F(m_\xi^2, m_H^2) + F(m_\xi^2, m_A^2) - F(m_H^2, m_A^2)], \quad (4.6)$$




FIG. 2: The allowed mass range for the charged Higgs and the heavy CP even Higgs in (a) type I (b) type II (c) lepton specific and (d) flipped 2HDM for $|\lambda_5| \leq 4\pi$ and $\tan\beta = 5$

with

$$F(x, y) = \begin{cases} \frac{x+y}{2} - \frac{xy}{x-y} \ln \frac{x}{y}, & x \neq y \\ 0 & x = y \end{cases} \quad (4.7)$$

The T parameter is constrained by the global fit to precision electroweak data to be [32]

$$T = 0.05 \pm 0.12. \quad (4.8)$$

Our results consist of the pairs (m_H, m_ξ) for each type of 2HDM, satisfying the two Veltman conditions, and consistent with the constraints from stability, tree-level unitarity and the T parameter. For $\tan\beta = 5$, we have plotted the $m_H - m_\xi$ curves corresponding to VC1 and VC2 shown in Table I, for several values of λ_5 . These have been superimposed on the bound determined by the T -parameter, perturbative unitarity, and stability conditions, i.e. the inequalities (4.4), (4.5), and (4.8). The resulting plot is shown in Fig. 2. VC1 produces ellipses, and VC2 gives a narrow band of hyperbolae. Their crossings which fall inside the band representing the bound from the inequalities are the allowed masses. From the plot we can estimate the individual bounds – for all four models, we find approximately $500 \text{ GeV} \lesssim m_\xi \lesssim 700 \text{ GeV}$, and about $400 \text{ GeV} \lesssim m_H \lesssim 600 \text{ GeV}$, with a higher m_H implying a higher m_ξ . As mentioned earlier, m_A is close to m_H as a result of (4.4).

V. DISCUSSION

The most important parameter in the 2HDMs is $\tan\beta$. There is no consensus on the value of $\tan\beta$, except that it should be larger than unity, based on constraints coming from $Z \rightarrow b\bar{b}$ and $B_q\bar{B}_q$ mixing [24]. It was argued in [23] using muon $g - 2$ that a large $\tan\beta$ was needed in lepton specific 2HDM, whereas [33] argued for $\tan\beta \geq 4$ in type I and flipped models using $b \rightarrow s\gamma$. This range of $\tan\beta$ also suppresses $B(t \rightarrow bH^+)$ below 4%. This maximal top-quark branching fraction coincides roughly with the numerical values of the 95% CL limits from the light charged Higgs searches at the LHC [34, 35]. A large value of $\tan\beta$ makes the heavy Higgs particle difficult to detect [36]. In particular, type I 2HDM tends to the inert doublet model for large values of $\tan\beta$, which is a model for dark matter. We have used a conservative $\tan\beta = 5$ to estimate the scalar masses m_H and m_{ξ^\pm} — note that m_A is not very far from m_H because of the degeneracy relation (4.4). A larger $\tan\beta$ makes the $m_H - m_A$ degeneracy more pronounced, so the inequality band becomes narrower. This narrows the ranges of m_H and m_{ξ^\pm} , also pushing the region of overlap upwards, making the heavy and charged Higgses more difficult to detect.

-
- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012).
 - [2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).
 - [3] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, Phys. Rept. **516**, 1 (2012)
 - [4] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977).
 - [5] R. D. Peccei and H. R. Quinn, Phys. Rev. D **16**, 1791 (1977).
 - [6] N. Turok and J. Zadrozny, Nucl. Phys. B **358**, 471 (1991).
 - [7] E. Ma, Phys. Rev. D **73**, 077301 (2006).
 - [8] E. Ma, Mod. Phys. Lett. A **23**, 647 (2008).
 - [9] R. Barbieri, L. J. Hall and V. S. Rychkov, Phys. Rev. D **74**, 015007 (2006) [hep-ph/0603188].
 - [10] P. M. Ferreira and D. R. T. Jones, JHEP **0908**, 069 (2009) [arXiv:0903.2856 [hep-ph]].
 - [11] T. D. Lee, Phys. Rev. D **8**, 1226 (1973).
 - [12] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *The Higgs Hunter's Guide*, Front. Phys. **80**, 1 (2000).
 - [13] G. C. Branco, W. Grimus and L. Lavoura, Phys. Lett. B **380**, 119 (1996)
 - [14] J. F. Gunion and H. E. Haber, Phys. Rev. D **67**, 075019 (2003)
 - [15] M. J. G. Veltman, Acta Phys. Polon. B **12**, 437 (1981).
 - [16] C. Newton and T. T. Wu, Z. Phys. C **62**, 253 (1994).
 - [17] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977).
 - [18] E. A. Paschos, Phys. Rev. D **15**, 1966 (1977).
 - [19] A. Pich and P. Tuzon, Phys. Rev. D **80**, 091702 (2009) [arXiv:0908.1554 [hep-ph]].
 - [20] A. G. Akeroyd, A. Arhrib and E. M. Naimi, Phys. Lett. B **490**, 119 (2000)
 - [21] J. Beringer *et al.* [Particle Data Group Collaboration], *Review of Particle Physics (RPP)*, Phys. Rev. D **86**, 010001 (2012) and 2013 partial update for the 2014 edition.
 - [22] B. W. Lee, C. Quigg and H. B. Thacker, Phys. Rev. D **16**, 1519 (1977).
 - [23] J. Cao, P. Wan, L. Wu and J. M. Yang, Phys. Rev. D **80**, 071701 (2009).
 - [24] A. Arhrib, R. Benbrik, C. H. Chen, R. Guedes and R. Santos, JHEP **0908**, 035 (2009).

- [25] M. Sher, Phys. Rept. **179**, 273 (1989).
- [26] J. Maalampi, J. Sirkka and I. Vilja, Phys. Lett. B **265**, 371 (1991).
- [27] S. Kanemura, T. Kubota and E. Takasugi, Phys. Lett. B **313**, 155 (1993).
- [28] J. Horejsi and M. Kladiva, Eur. Phys. J. C **46**, 81 (2006).
- [29] G. Bhattacharyya, D. Das, P. B. Pal and M. N. Rebelo, JHEP **1310**, 081 (2013).
- [30] H. J. He, N. Polonsky and S. f. Su, Phys. Rev. D **64**, 053004 (2001).
- [31] W. Grimus, L. Lavoura, O. M. Ogreid and P. Osland, J. Phys. G **35**, 075001 (2008).
- [32] M. Baak and R. Kogler, “*The global electroweak Standard Model fit after the Higgs discovery*”, arXiv:1306.0571 [hep-ph].
- [33] J. h. Park, JHEP **0610**, 077 (2006).
- [34] G. Aad *et al.* [ATLAS Collaboration], JHEP **1206**, 039 (2012).
- [35] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1207**, 143 (2012).
- [36] L. Randall, JHEP **0802**, 084 (2008).