

The asymmetry and the diurnal variation in axionic dark matter searches

Y. Semertzidis and J.D. Vergados

KAIST University, Daejeon, Republic of Korea

Abstract

In the present work we propose to study the modulation signal in Axion Dark Matter searches. This can be seen in directional experiments employing cylindrical resonant cavities, which exploit the axion photon-conversion in the presence of strong magnetic fields.

1. Introduction

The axion has been proposed a long time ago as a solution to the strong CP problem [1] resulting to a pseudo Goldstone Boson [2, 3], but it has also been recognized as a prominent dark matter candidate [4]. Axions are cold non baryonic particles, which, in principle, can be distinguished from other dark matter candidates [5]. In fact experiments, e.g. ADMX and ADMX-HF collaborations [6, 7], using SQUID and HFET technologies [8], [9], are now planned to search for them. We will take the view that the axion is non relativistic with mass in μeV - meV scale moving with an average velocity which is $\approx 0.8 \times 10^{-3}c^2$. The allowed parameter space has been presented in a nice slide by Raffelt [10] in the recent Multidark-IBS workshop, which is shown in Fig. 1.1.

In the present work we will explore some special signatures, which may aid the analysis axion dark experiments in discriminating against possible backgrounds.

2. Brief summary of the formalism

The photon axion interaction is dictated by the Lagrangian:

$$\mathcal{L}_{a\gamma\gamma} = g_{a\gamma\gamma} a \mathbf{E} \cdot \mathbf{B}, \quad g_{a\gamma\gamma} = \frac{ag_\gamma}{2\pi f_a}, \quad (2.1)$$

where \mathbf{E} and \mathbf{B} are the electric and magnetic fields. Axion dark matter detectors [11] employ an external magnetic field, $\mathbf{B} \rightarrow \mathbf{B}_0$ in the previous equation, in which case one of the photons is replaced by a virtual photon, while the other




Figure 1.1: The parameter space for axion like particles (courtesy of professor Raffelt)

maintains the energy of the axion, which is its mass plus a small fraction of kinetic energy.

The flux of the axions is given by:

$$\frac{dN}{dt} = \frac{\rho_a}{m_a} v dS \quad (2.2)$$

where v is the velocity of the axion, ρ_a its density and m_a its mass. In practice vdt is larger than the length of the detector and so one may write

$$N = \frac{\rho_a}{m_a} V, V = S\ell, \ell = vT$$

with T some suitable time.

So the power produced by the cavity, see e.g. [6], is given by:

$$P_{mnp} = g_{a\gamma\gamma}^2 \frac{\rho_a}{m_a} B_0^2 V C_{mnp} Q_L \quad (2.3)$$

Q_L is the loaded quality factor of the cavity and the subscript mnp indicates that the power produced is mode dependent. For our purposes we will assume a velocity dependence with the understanding $\rho_a \rightarrow \rho_a v(T/\ell)$ in Eq. (2.3).

3. Time dependence (modulation) of the results

One then can fold the above expression with the velocity distribution of axions. The axion velocity distribution is taken to be Bose-Einstein, see e.g. [12]. One, however, would like the axion dark matter density to fit the rotational curves. Thus for temperatures T such that $m_a/T \approx 4 \times 10^6$ the velocity distribution can be taken to be analogous to that assumed for WIMPs, i.e. a M-B distribution with a characteristic velocity which equals the velocity of the sun around the center of the galaxy, i.e. $v_0 \approx 220\text{km/s}$. So we will employ the distribution:

$$f(\vec{v}) = \frac{1}{(\sqrt{\pi}v_0)^3} e^{-\frac{v^2}{v_0^2}} \quad (3.4)$$

The explicit inclusion of the velocity distribution may not affect significantly the experimental results already obtained with the use of some average velocity. The average velocity, however, is affected by the form of the velocity distribution in the local frame. This is true in particular for the annual modulation and the diurnal variation in directional experiments. It also affects the width in the power spectrum (see below section 4).

3.1. The annual modulation in non directional experiments

The flux of axions in the detector due to the motion of the Earth around the sun, which is called the modulation effect [13] (the velocity of the detector due to the rotation of the Earth around its own axis is too small to have a significant effect). This will affect the observed rates as is well known from the standard dark matter searches. Our next task is to transform the velocity distribution from the galactic to the local frame. The needed equation, see e.g. [14], is:

$$\mathbf{y} \rightarrow \mathbf{y} + \hat{v}_s + \delta (\sin \alpha \hat{x} - \cos \alpha \cos \gamma \hat{y} + \cos \alpha \sin \gamma \hat{v}_s), \quad y = \frac{v}{v_0} \quad (3.5)$$

with $\gamma \approx \pi/6$, \hat{v}_s a unit vector in the Sun's direction of motion, \hat{x} a unit vector radially out of the galaxy in our position and $\hat{y} = \hat{v}_s \times \hat{x}$. The last term in the first expression of Eq. (3.5) corresponds to the motion of the Earth around the Sun with δ being the ratio of the modulus of the Earth's velocity around the Sun divided by the Sun's velocity around the center of the Galaxy, i.e. $v_0 \approx 220\text{km/s}$ and $\delta \approx 0.135$. The above formula assumes that the motion of both the Sun around the Galaxy and of the Earth around the Sun are uniformly circular. The exact orbits are, of course, more complicated but such deviations are not expected to significantly modify our results. In Eq. (3.5) α is the phase of the Earth ($\alpha = 0$ around June 3rd)¹.

The variation of the total flux as a function of time (the phase of the Earth α) is given in Fig. 3.2.




Figure 3.2: The variation of the average axion flux, in units of $(\rho_a/m_a)v_0$, due to the motion of the Earth around the sun. As expected, the maximum occurs when $\alpha = 0$ (around June 3rd).

3.2. Asymmetry of the rates in directional experiments

Consideration of the velocity distribution will give an important signature, if directional experiments become feasible. This can be seen as follows:

¹One could, of course, make the time dependence of the rates due to the motion of the Earth more explicit by writing $\alpha \approx (6/5)\pi (2(t/T) - 1)$, where t/T is the fraction of the year.

- The flux is replaced by the flux of axions in the particular direction of observation $v \rightarrow \vec{v} \cdot \hat{n} = \xi v$, $0 \leq \xi \leq 1$.
- The unit vector \hat{n} is specified by two angles Θ and Φ . The angle Θ is the polar angle between the sun's velocity and the direction of observation. The angle Φ is measured in a plane perpendicular to the sun's velocity, starting from the line coming radially out of the galaxy and passing through the sun's location.
- The axion velocity, in units of the solar velocity, is given as

$$\mathbf{y} = y \left(\hat{x} \sqrt{1 - \xi^2} \cos \phi + \hat{y} \sqrt{1 - \xi^2} \sin \phi + \hat{z} \xi \right) \quad (3.6)$$

- The sun's velocity is given by:

$$\mathbf{y}_s = (\hat{x} \sin \Theta \cos \phi + \hat{y} \sin \Theta \sin \phi + \hat{z} \cos \Theta) \quad (3.7)$$

Then the velocity distribution in the local frame is obtained by the substitution:

$$v^2 \rightarrow v_0^2 \left(y^2 + 1 + 2y \left(\xi \cos \Theta + \sqrt{1 - \xi^2} \sin \Theta (\cos \Phi \cos \phi + \sin \Phi \sin \phi) \right) \right) \quad (3.8)$$

One then can integrate over ξ and ϕ . The results become essentially independent of Φ , so long as the motion of the Earth around the sun is ignored². Thus we obtain the angle averaged flux as a function of the axion velocity for various polar angles Θ . The results obtained are shown in Fig. 3.3. We notice the large directional dependence of the flux, "solar wind" of axions, not, of course, created there. The direction opposite to the sun's motion is favored.

Integrating over the magnitude of the velocity we obtain the total flux as a function of the polar angle Θ . The obtained results are shown in Fig. 3.4. We notice again the large asymmetry. The above results apply in the case one can distinguish the sense of direction. If experimentally this is not feasible, one expects both directions along the sun's velocity to be favored, with event rates the average of the two, while directions of observation on a plane perpendicular to the sun's velocity to be suppressed. This shown in Figs. 3.6 and 3.6

²The annual modulation of the results due to the motion of the Earth around the sun appears in the directional experiments as well, but it will not be discussed here.




Figure 3.3: The angular average $\rho_a/m_a \langle \vec{v} \cdot \hat{n} \rangle$, essentially the corresponding axion flux, in units of $(\rho_a/m_a)v_0$, is shown as a function of its velocity in units of v_0 for the values of $\Theta = 0, \pi/4, \pi/2, 5\pi/8, 3\pi/4, 7\pi/8$ and π increasing upwards. We note the large asymmetry. The most favored direction is opposite to the sun's velocity.

4. Implications on the power spectrum

The axion power spectrum, which is of great interest to experiments, is written as a Breit-Wigner shape [11], [15]:

$$|\mathcal{A}(\omega)|^2 = \frac{\rho_D}{m_a^2} \frac{\Gamma}{(\omega - \omega_a)^2 + (\Gamma/2)^2}, \Gamma = \frac{\omega_a}{Q_a} \quad (4.9)$$

with $\omega_a = m_a (1 + (1/6) \langle v^2 \rangle)$ and $Q_a = m_a / (m_a / (m_a \langle v^2 \rangle / 3))$. The width explicitly depends on the average axion velocity squared in the laboratory. Thus the width in the laboratory is affected by the sun's motion. In the non directional experiments $\langle v^2 \rangle = (3/2)v_0^2$ becomes $\langle v^2 \rangle = (5/2)v_0^2$ (v_0 the velocity of the sun around the center of the galaxy). The situation becomes more dramatic as soon the experiment is directional. In this case we will write the width as

$$\Gamma = \Gamma_{st} r(\Theta, \Phi)$$

where Γ_{st} is the width in the standard experiments. Ignoring the motion of the Earth around the sun the factor r depends only on Θ . Furthermore, if for simplicity we ignore the upper velocity bound in the M-B distribution, i.e. the




Figure 3.4: The average $\langle \vec{v} \cdot \hat{n} \rangle / v_0$, the average total axion flux in units $(\rho_a/m_a)v_0$ as a function of the polar angle Θ . We note the large asymmetry. The favored direction is opposite to the velocity of the sun's motion ($\Theta = \pi$).

escape velocity $v_{esc} = 2.84v_0$, we can get the solution in analytic form. We find:

$$r(\Theta) = \frac{2}{5} \frac{e^{-1}}{2} \left(e^{\cos^2 \Theta} (\cos 2\Theta + 4) \operatorname{erfc}(\cos \Theta) - \frac{2 \cos \Theta}{\sqrt{\pi}} \right), \text{ (sense known)} \quad (4.10)$$

where

$$\operatorname{erfc}(z) = 1 - \operatorname{erf}(z), \operatorname{erf}(z) = \int_0^z dt e^{-t^2} \text{ (error function).}$$

$$r(\Theta) = \frac{2}{5} \frac{1}{2} e^{-\sin^2 \Theta} (\cos 2\Theta + 4), \text{ (sense of direction not known)} \quad (4.11)$$

The adoption of an upper cut off has little effect. In Fig. 4.7 we present the exact results.

The modification of the width in the numerator may not be necessary, if the modification of the flux is suitably taken into account by modifying ρ_D . Anyway the width will exhibit diurnal variation! This diurnal variation is going to be similar to that discussed in the next section for the flux.

5. The diurnal variation in directional experiments

The apparatus will be oriented in a direction specified in the local frame, e.g. by a point in the sky specified, in the equatorial system, by right ascension




Figure 3.5: The average $\langle \vec{v} \cdot \hat{n} \rangle / v_0$, essentially the axion flux in units of $(\rho_a/m_a)v_0$, as a function of the polar angle Θ . We note that some of the asymmetry remains. The favored direction is still along the sun's direction of motion (the angles $\Theta = 0$, and $\Theta = \pi$ cannot be distinguished).

$\tilde{\alpha}$ and inclination $\tilde{\delta}^3$. This will lead to a diurnal variation⁴ of the event rate [16]. This situation has already been discussed in the case of standard WIMPs [17]. We will briefly discuss the transformation into the relevant astronomical coordinates here.

The galactic frame, in the so called J2000 system, is defined by the galactic pole with ascension $\tilde{\alpha}_1 = 12^h 51^m 26.282^s$ and inclination $\tilde{\delta}_1 = +27^\circ 7' 42.01''$ and the galactic center at $\tilde{\alpha}_2 = 17^h 45^m 37.224^s$, $\tilde{\delta}_2 = -(28^\circ 56' 10.23'')$. Thus the galactic unit vector \hat{y} , specified by $(\tilde{\alpha}_1, \tilde{\delta}_1)$, and the unit vector \hat{s} , specified by $(\tilde{\alpha}_2, \tilde{\delta}_2)$, can be expressed in terms of the celestial unit vectors \hat{i} (beginning of measuring the right ascension), \hat{k} (the axis of the Earth's rotation) and $\hat{j} = \hat{k} \times \hat{i}$.

³We have chosen to adopt the notation $\tilde{\alpha}$ and $\tilde{\delta}$ instead of the standard notation α and δ employed by the astronomers to avoid possible confusion stemming from the fact that α is usually used to designate the phase of the Earth and δ for the ratio of the rotational velocity of the Earth around the Sun by the velocity of the sun around the center of the galaxy

⁴This should not be confused with the diurnal variation expected even in non directional experiments due to the rotational velocity of the Earth, which is expected to be too small.




Figure 3.6: The average $\langle \vec{v} \cdot \hat{n} \rangle / v_0$, essentially the average total axion flux in units $(\rho_a/m_a)v_0$ as a function of the polar angle Θ , when the sense of direction cannot be distinguished. The angles $\Theta = 0$, and $\Theta = \pi$ cannot be distinguished. The asymmetry, however, with the $\Theta = \pi/2$ remains.

The direction of observation, if it does not coincide with the axis of rotation of the Earth, during the day will point to a different point of the sky. So as Θ will be changing due to the Earth's rotation, the data will exhibit a characteristic diurnal variation.

One finds

$$\begin{aligned}
 \hat{y} &= -0.868\hat{i} - 0.198\hat{j} + 0.456\hat{k} \text{ (galactic axis) ,} \\
 \hat{x} &= -\hat{s} = 0.055\hat{i} + 0.873\hat{j} + 0.483\hat{k} \text{ (radially out to the sun) ,} \\
 \hat{z} &= \hat{x} \times \hat{y} = 0.494\hat{i} - 0.445\hat{j} + 0.747\hat{k} \text{ (sun's velocity).} \quad (5.12)
 \end{aligned}$$

Note in our system the x-axis is opposite to the s-axis used by the astronomers. Thus a vector oriented by $(\tilde{\alpha}, \tilde{\delta})$ in the laboratory is given in the galactic frame by a unit vector with components:

$$\begin{pmatrix} y \\ x \\ z \end{pmatrix} = \begin{pmatrix} -0.868 \cos \tilde{\alpha} \cos \tilde{\delta} - 0.198 \sin \tilde{\alpha} \cos \tilde{\delta} + 0.456 \sin \tilde{\delta} \\ 0.055 \cos \tilde{\alpha} \cos \tilde{\delta} + 0.873 \sin \tilde{\alpha} \cos \tilde{\delta} + 0.4831 \sin \tilde{\delta} \\ 0.494 \cos \tilde{\alpha} \cos \tilde{\delta} - 0.445 \sin \tilde{\alpha} \cos \tilde{\delta} + 0.747 \sin \tilde{\delta} \end{pmatrix}. \quad (5.13)$$




Figure 4.7: The ratio of the width expected in a directional experiment divided by that expected in a standard experiment. The solid line is expected, if the sense of direction is known, while the dotted will show up, if the sense of direction is not known.

This can also be parametrized as:

$$x = \cos \gamma \cos \tilde{\delta} \cos (\tilde{\alpha} - \tilde{\alpha}_0) - \sin \gamma \left(\cos \tilde{\delta} \cos \theta_P \sin (\tilde{\alpha} - \tilde{\alpha}_0) + \sin \tilde{\delta} \sin \theta_P \right), \quad (5.14)$$

$$y = \cos (\theta_P) \sin \tilde{\delta} - \cos \tilde{\delta} \sin (\tilde{\alpha} - \tilde{\alpha}_0) \sin \theta_P, \quad (5.15)$$

$$z = \cos \delta \cos (\tilde{\alpha} - \tilde{\alpha}_0) \sin \gamma + \cos \gamma \left(\cos \tilde{\delta} \cos \theta_P \sin (\tilde{\alpha} - \tilde{\alpha}_0) + \sin \tilde{\delta} \sin \theta_P \right), \quad (5.16)$$

where $\tilde{\alpha}_0 = 282.25^0$ is the right ascension of the equinox, $\gamma \approx 33^0$ was given above and $\theta_P = 62.6^0$ is the angle the Earth's north pole forms with the axis of the galaxy.

Due to the Earth's rotation the unit vector (x, y, z) , with a suitable choice of the initial time, $\tilde{\alpha} - \tilde{\alpha}_0 = 2\pi(t/T)$, is changing as a function of time

$$\begin{aligned}
x &= \cos \gamma \cos \tilde{\delta} \cos \left(\frac{2\pi t}{T} \right) - \sin \gamma \left(\cos \delta \cos \theta_P \sin \left(\frac{2\pi t}{T} \right) \right. \\
&\quad \left. + \sin \tilde{\delta} \sin \theta_P \right), \tag{5.17}
\end{aligned}$$

$$y = \cos(\theta_P) \sin \tilde{\delta} - \cos \tilde{\delta} \sin \left(\frac{2\pi t}{T} \right) \sin \theta_P, \tag{5.18}$$

$$\begin{aligned}
z &= \cos \left(\frac{2\pi t}{T} \right) \cos \tilde{\delta} \sin \gamma \\
&\quad + \cos \gamma \left(\cos \tilde{\delta} \cos \theta_P \sin \left(\frac{2\pi t}{T} \right) + \sin \tilde{\delta} \sin \theta_P \right), \tag{5.19}
\end{aligned}$$

where T is the period of the Earth's rotation.

Some points of interest are:

The celestial pole:

$$(y, x, z) = (0.460, 0.484, 0.745) \Rightarrow (\theta = 62.6^0, \phi = 57^0),$$

The ecliptic pole:

$$(y, x, z) = (0.497, 0.096, 0.863) \Rightarrow (\theta = 62.6^0, \phi = 83.7^0),$$

The equinox:

$$(y, x, z) = (-0.868, 0.055, 0.494) \Rightarrow (\theta = 150.2^0, \phi = 83.7^0). \tag{5.20}$$

where θ is defined with respect to the polar axis (here y) and ϕ is measured from the x axis towards the z axes.

Thus the angles Θ , which is of interest to us in directional experiments, is given by

$$\Theta = \cos^{-1} z, \tag{5.21}$$

An analogous, albeit a bit more complicated expression can be derived for the angle Φ .

The angle Θ scanned by the direction of observation is shown, for various inclinations $\tilde{\delta}$, in Fig. 16. We see that for negative inclinations, the angle Θ can take values near π , i.e. opposite to the direction of the sun's velocity, where the rate attains its maximum (see Fig. 5.8).

The equipment scans different parts of the galactic sky, i.e. observes different angles Θ . So the rate will change with time depending on whether the sense of observation. We assume that the sense of direction can be distinguished in the experiment. The total flux is exhibited in Fig. 5.9.

Acknowledgments: One of the authors (JDV) is indebted to Professor J. E. Kim for useful discussions.





Figure 5.8: Due to the diurnal motion of the Earth different angles Θ in galactic coordinates are sampled as the earth rotates. The angle Θ scanned by the direction of observation is shown for various inclinations δ . We see that, for negative inclinations, the angle Θ can take values near π , i.e. opposite to the direction of the sun's velocity, where the rate attains its maximum. For an explanation of the curves see Fig. 5.9


References

- [1] R. Peccei, H. R. Quinn, Phys. Rev. Lett 38 (1977) 1440.
- [2] S. Weinberg, Phys. Rev. Lett 40 (1978) 223.
- [3] F. Wilczek, Phys. Rev. Lett 40 (1978) 279.
- [4] J. Primack, D. Seckel, B. Sadoulet, Ann. Rev. Nuc. Par. Sci. 38 (1988) 751.
- [5] P. Sikivie, Phys. Lett. B695 (2011) 22.
- [6] I. P. Stern, ArXiv 1403.5332 (2014) physics.ins-det, on behalf of ADMX and ADMX-HF collaborations, Axion Dark Matter Searches.
- [7] Gray Rybka, The Axion Dark Matter Experiment, IBS MultiDark Joint Focus Program WIMPs and Axions, Daejeon, S. Korea October 2014.
- [8] S. Asztalos, et al., Nucl. Instrum. Meth. A 656, (2011) 30.
- [9] J. Hoskins, et al., Phys. Rev. D 84, (2011) 121302.

- [10] G. Raffelt, Astrophysical Axion Bounds , IBS MultiDark Joint Focus Program WIMPs and Axions, Daejeon, S. Korea October 2014.
- [11] J. Hong, J. E. Kim, S. Nam, Y. Semertzidis, ArXiv arXiv:1403.1576 (2014) physics.ins-det, calculations of Resonance enhancement factor in axion-search tube experiments.
- [12] S. Hannestad, A. Mirizzi, G. Raffelt, JCAP 0507 (2005) 002, arXiv: hep-ph/0504059.
- [13] K. F. A. K. Drukier, D. N. Spergel, Phys. Rev. D 33 (1986) 3495.
- [14] J. D. Vergados, Phys. Rev D 85 (2012) 123502, arXiv:1202.3105 (hep-ph).
- [15] L. Krauss, J. Moody, F. Wilczek, D. Morris, Phys. Rev. Lett. 55, (1985) 1797.
- [16] S. Ahlen, IJMPA 25 (2010) 1, the case for a directional dark matter detector and the status of current experimental efforts, cygnus2009Whitepaper, Edited by J. B. R. Battat; arXiv:0911.0323.
- [17] J. D. Vergados, C. C. Moustakidis, Eur. J. Phys. 9(3) (2011) 628, arXiv:0912.3121 [astro-ph.CO].



(a)



(b)

Figure 5.9: The time dependence (in units of the Earth's rotation period) of the average the total flux, in units of $(\rho_a/m_a)v_0$ for various inclinations $\tilde{\delta}$, when the sense can be determined (a) or both senses are included (b). In the curves indicated by intermediate thickness solid, the short dash, thick solid line, long dash, dashed, fine solid line, and the long-short dashed correspond to inclination $\tilde{\delta} = -\pi/2, -3\pi/10, -\pi/10, 0, \pi/10, 3\pi/10$ and $\pi/2$ respectively. We see that, for negative inclinations, the angle Θ can take values near π , i.e. opposite to the direction of the sun's velocity, where the rate attains its maximum if the sense of direction is known. There is no time variation, of course, when $\tilde{\delta} = \pm\pi/2$.