

Interference and non-linear properties of four-wave mixing resonances in thermal vapor: analytical results and experimental verification

Michał Parniak* and Wojciech Wasilewski

Institute of Experimental Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland

We develop a model to calculate non-linear polarization in a non-degenerate four-wave mixing in diamond configuration which includes the effects of hyperfine structure and Doppler broadening. We verify it against the experiment with $5^2S_{1/2}$, $5^2P_{3/2}$, $5^2D_{3/2}$ and $5^2P_{1/2}$ levels of rubidium 85. Uncomplicated algebra enables us to express the non-linear susceptibility of a thermal ensemble in low intensity regime in terms of Voigt-like profiles and conforms precisely with the experiment. The agreement is also satisfactory at high intensity and the analytical model correctly predicts the position and shape of resonances. Our intelligible results elucidate the physics of coherent interaction of light with atoms involving higher excited levels in vapors at room temperature, which is used in an increasing range of applications.

PACS numbers: 32.80.Wr, 32.80.Qk, 42.50.Gy, 42.65.Ky

I. INTRODUCTION

Coherent interactions of light and atomic vapors involving higher excited states have attracted much attention recently. A seminal work by Peyronel *et al.* [1] demonstrated electromagnetically induced transparency resonance with extreme single-photon sensitivity due to utilization of Rydberg levels. A host of other works explore the use of multi-photon transitions involving higher states in the context of nonlinear optics and quantum information.

The so-called diamond configuration of atomic levels, as sketched in Fig. 1, is frequently used. Applications include: coherent interaction of atoms with ultrashort laser pulses [2], useful for interferometric measurements [3], and coherent control [4]. In continuous wave regime atoms in diamond configuration demonstrate high resonant nonlinearities that enable generation of coherent blue and infrared light [5–8] via four-wave mixing (4WM). This up-conversion scheme also enables coherent transfer of light angular momentum [9].

Atoms in diamond configuration have been successfully used as a non-linear frequency conversion medium [10–13], applicable to light at single photon level. Both spectral [14, 15] and temporal [16] properties of the diamond configuration have been studied.

When the diamond configuration involved a Rydberg state, 4WM has also been observed [17] and coherent revival effects have been seen even in a warm ensemble of atoms [18]. A similar configuration is also proposed to be a room-temperature single photon source [19].

Finally, 4WM in diamond configuration has been used as a source of single photon pairs [20, 21] with well-defined temporal properties [22] which, thanks to proper postselection, can be perfectly matched to absorption by single atoms [23].




Figure 1. The diamond scheme of atomic levels and transitions. Rubidium 85 is used to generate light at the wavelength $\lambda_3 = 762$ nm via 4WM of incident light fields at the wavelengths $\lambda_1 = 780$ nm, $\lambda_2 = 776$ nm and $\lambda_4 = 795$ nm, and Rabi frequencies Ω_1 , Ω_2 and Ω_4 . Levels |1> to |4> correspond to $5^2S_{1/2}$, $5^2P_{3/2}$, $5^2D_{3/2}$ and $5^2P_{1/2}$ states of rubidium 85, respectively. The optical frequencies are parametrized by two one-photon detunings Δ_2 and Δ_4 and a two-photon detuning Δ_3 .

In most of the above-mentioned works a maximally five-level atom model was used to model the experiment. However, actual atoms have rich hyperfine structure which inevitably influences any room-temperature experiments where different transitions are driven in classes of atoms depending on their velocity. The susceptibility of a room-temperature ensemble is a convolution of respective cold-atomic function in a multidimensional space of frequencies of the driving fields with a Gaussian corresponding to thermal velocity distribution. This particular convolution reshapes resonances and forms band structures which extend in certain directions while they stay sub-Doppler in others.

In this paper we develop simple theoretical tools for predicting the influence of this effect on a coherent process in which interference between contributions of different atomic levels plays a role. For weak driving fields the final result is a superposition of Voigt-like profiles, and

* michal.parniak@fuw.edu.pl

a relatively simple recipe is provided for higher intensities. The simplicity worked out enables us to explain the intricacies of interference between contributions of various hyperfine sublevels taking into account the Doppler broadening in a precise and exact way.

We verify our results against experiment in which we measure the intensity of 4WM in diamond configuration in warm rubidium 85 vapor as a function of laser frequencies. The comparison covers both low and high intensity regimes.

The effects we describe and explain herein are especially important to studying phase-dependent interactions [3, 4], to optimizing single-photon generation setups [21, 22], and to explaining 4WM spectra observed in many experiments [8, 14, 15] or possible magnetic field effects.

This paper is organized as follows. In section II we introduce a theoretical description of 4WM in an inhomogeneously broadened multi-level atomic medium. In section III we describe the details of our 4WM experiment in rubidium vapors. Section IV presents a comparison of experimental and theoretical data. The last section V concludes the paper.

II. THEORY

In this section we calculate the intensity of the light generated by 4WM in diamond configuration depicted in Fig. 1. We start with calculating the optical coherence in a four-level atom model. Next we take into account the hyperfine structure, and finally the Doppler broadening.

A. Nonlinear polarization of a four-level atom

In the experiment we observe the intensity of light generated at the transition between levels $|3\rangle$ and $|4\rangle$ at $\lambda_3 = 762$ nm via 4WM. The starting point for the theoretical treatment is the calculation of steady-state optical coherence ρ_{43} in a four-level atom in diamond configuration. For the sake of simplicity we assume that the intensity of the incident laser light remains constant through the atomic medium. When the geometry of the laser beams enables perfect wave-vector matching, the amplitude of the emitted light wave E_3 is proportional to the optical polarization P_3 , which, in turn, is proportional to the density of atoms n , the dipole moment of the transition μ_{43}^* and the optical coherence

$$E_3 \propto P_3 = n\mu_{43}^*\rho_{43} + \text{c.c.} \quad (1)$$

The steady-state coherence ρ_{43} can be calculated from the Liouville equation with relaxation, constructed as in Ref. [17]. For low light intensities we may use the lowest non-vanishing order of the perturbative solution and obtain a compact formula for the optical coherence:

$$\rho_{43} = \frac{\Omega_1\Omega_2\Omega_4^*}{8\tilde{\Delta}_2\tilde{\Delta}_3\tilde{\Delta}_4^*}, \quad (2)$$

where $\tilde{\Delta}_j = \Delta_j + i\Gamma_j/2$ are the complex detunings, Γ_j are the decay rates and Ω_j are the Rabi frequencies, accordingly to Fig. 1. This leads to the following expression for the optical polarization:

$$P_3 = n \frac{\mu_{12}\mu_{23}\mu_{43}^*\mu_{14}^*}{8\epsilon_0\hbar^3\tilde{\Delta}_2\tilde{\Delta}_3\tilde{\Delta}_4^*} E_1 E_2 E_4^* + \text{c.c.} \quad (3)$$

For pump light intensities exceeding saturation $\Omega_i \gtrsim \Gamma_i$ we need to solve the Liouville equation exactly. A computer algebra system provides an exact solution in the form of a rational function of complex detunings $\tilde{\Delta}_i$ and Rabi frequencies Ω_i . We state this solution in a general way:

$$\rho_{43} = \rho_{43}(\tilde{\Delta}_2, \tilde{\Delta}_3, \tilde{\Delta}_4^*, \Omega_1, \Omega_2, \Omega_4^*). \quad (4)$$

In Fig. 2 we plot the optical polarization squared $|P_3|^2$ for different Rabi frequencies Ω_i , calculated using the solution (4) in Eq. (1), to illustrate an already complex behavior of the system. For higher Rabi frequencies the system reveals Autler-Townes-like splitting [15, 24]. The above solutions can be readily applied to describe 4WM in a cold ensemble of atoms or a single atom where the hyperfine structure of each of the intermediate levels is fully resolved.

B. Hyperfine structure: interference of paths

In order to fully account for the hyperfine structure of a real atom, we would have to solve the Liouville equation for an excessively large density matrix. In our experimental case it would have to take into account at least 64 atomic states. Solution of the resulting steady-state Liouville equation for a range of parameters, such as detunings would be hard to compute.

Instead, we propose an alternative solution to this problem that is much more robust in terms of computation. For the sake of simplicity we assume that all light fields, both the pumping ones and the signal one, have a fixed polarization parallel to the quantization axis x , as this is our experimental situation.

In our method we treat the full system as a set of many four-level systems. Each of these can be represented as a certain path $\mathcal{P} = \{|1 F m_{F_1}\rangle, |2 F_2 m_{F_2}\rangle, |3 F_3 m_{F_3}\rangle, |4 F m_{F_4}\rangle, |1 F_1 m_{F_1}\rangle\}$, through intermediate levels, where F_i is the total angular momentum of the state $|i\rangle$, and m_{F_i} is its projection. Each path can be characterized by different dipole moments $\mu_{ij}(\mathcal{P})$, and different hyperfine detunings $\Delta_i(\mathcal{P})$. Angular momenta for all of the excited levels $|2\rangle$, $|3\rangle$ and $|4\rangle$ vary through all possible values, but for




Figure 2. Absolute value squared of the optical polarization $|P_3|^2$ in the regime of high Rabi frequencies for four different configurations. The two-photon detuning $\Delta_3 = \Delta_2$, as the frequency of Ω_3 field is kept constant.

the ground state $|1\rangle$ we choose only one ground state angular momentum F_1 and vary only its projection m_{F_1} . This is due to the fact that the ground state hyperfine manifold is fully resolved in our experimental case.

The dipole moments $\mu_{ij}(\mathcal{P}) = \langle J_i \mathcal{I} F_i m_{F_i} | \hat{\mu} | J_j \mathcal{I} F_j m_{F_j} \rangle$, where $\hat{\mu}$ is the dipole moment operator, \mathcal{I} is the nuclear spin and J_i is the total angular momentum of the electron on the level $|i\rangle$, are calculated using known values of reduced dipole moments $\mu_{ij} = \langle J_i || \hat{\mu} || J_j \rangle$ according to the formulas given in Ref. [25]. The complex detunings $\tilde{\Delta}_i(\mathcal{P})$ calculated from a centroid of respectful lines are affected by hyperfine shifts in the following way:

$$\tilde{\Delta}_i(\mathcal{P}) = \tilde{\Delta}_i + \Delta F_i - \Delta F_1, \quad (5)$$

where ΔF_i is the shift of a given F_i from the centroid of the manifold $|i\rangle$. It follows that the detunings $\tilde{\Delta}_i$ are measured to the manifold centroids.

In our experimental case 61 paths through intermediate states contribute to the optical polarization P_3 . For each path \mathcal{P} we calculate its contribution $P_3(\mathcal{P})$ by inserting solution (4) into Eq. (1). In total, we obtain:

$$P_3 = \sum_{\mathcal{P}} n(\mathcal{P}) \mu_{43}^*(\mathcal{P}) \rho_{43}(\mathcal{P}) + \text{c.c.}, \quad (6)$$

where $n(\mathcal{P})$ is the density of the atoms in the ground state sublevel F_1, m_{F_1} of a particular path \mathcal{P} , while $\rho_{43}(\mathcal{P})$ is calculated taking into account hyperfine shifts and dipole

$\chi(\mathcal{P})$	$F_2 = 2$				$F_2 = 3$				$F_2 = 4$			
$ m_{F_1} $	3	2	1	0	3	2	1	0	3	2	1	0
$F_4 = 2$	0	-40	32	72	0	140	56	0	0	180	360	432
$F_4 = 3$	0	80	32	0	360	0	-21	0	216	144	45	0

Table I. Theoretical relative amplitudes of 4WM resonances in rubidium 85 parametrized by spin of $|2\rangle$ and $|4\rangle$ states F_2 and F_4 for each spin projection of the ground state m_{F_1} . One hyperfine component of the ground state with spin $F_1 = 3$ is taken and averaged over the spin of highest excited state $F_3 = 1 \dots 4$ due to negligible hyperfine splitting.

moments corresponding to the given path \mathcal{P} . The above equation expresses an approximation neglecting coherent interplay of different paths in a single atom.

However, such interplays may only matter when two paths share common levels and some coherences oscillate at similar frequencies. In our experimental situation all of the optical fields are polarized linearly in the same direction x , which we take as quantization axis. In this case all m_{F_i} in each path are the same due to selection rules for dipole moments. Consequently if two distinct paths share a level $|i\rangle$, then they must have a different total angular momentum F_j at some other level $|j\rangle$. The natural oscillation frequencies of coherences for these paths differ by the hyperfine splitting. As the Rabi frequencies in our experiment are smaller than the hyperfine splittings, we conjecture such paths will not interplay coherently. This is verified in the experiment.

Strong driving beams significantly redistribute the atoms among the ground state sublevels F_1, m_{F_1} altering $n(\mathcal{P})$. We find the steady state of a set of rate equations to determine $n(\mathcal{P})$ and thus the relative contributions of paths.

To conclude this section, let us note that within the limit of low intensities of driving fields $\Omega_i \ll \Gamma_i$ the total polarization can be calculated exactly using perturbation calculus. In the lowest nonvanishing order Eq. (3) can be inserted into Eq. (6) leading to a simple result:

$$P_3 = \epsilon_0 E_1 E_2 E_4^* \sum_{\mathcal{P}} \chi(\mathcal{P}) + \text{c.c.}, \quad (7)$$

where we identified the susceptibility $\chi(\mathcal{P})$ of the system:

$$\chi(\mathcal{P}) = n(\mathcal{P}) \frac{\mu_{12}(\mathcal{P}) \mu_{23}(\mathcal{P}) \mu_{43}^*(\mathcal{P}) \mu_{14}^*(\mathcal{P})}{8\epsilon_0 \hbar^3 \tilde{\Delta}_2(\mathcal{P}) \tilde{\Delta}_3(\mathcal{P}) \tilde{\Delta}_4^*(\mathcal{P})}. \quad (8)$$

The above expression approximates the rough features of full non-perturbative solution quite well and we find it worthwhile to track the contributions of various hyperfine components to the final result. We list them in Table I for the 4WM process we study in experiment.

C. Doppler broadening

The final step in the construction of our model consists in averaging the contributions of atoms moving with various velocities v along almost parallel laser beams. We take the one-dimensional Maxwell-Boltzmann velocity distribution $g(v) = \sqrt{\frac{m}{2\pi k_B T}} \exp(-\frac{mv^2}{2k_B T})$, where m is the atomic mass, T is the temperature and k_B is the Boltzmann's constant, to calculate the number of atoms with velocity v . To perform the average over the thermal ensemble, we introduce velocity-dependent detunings: $\tilde{\Delta}_2^{(v)} = \tilde{\Delta}_2 + \frac{v}{\lambda_1}$, $\tilde{\Delta}_3^{(v)} = \tilde{\Delta}_3 + \frac{v}{\lambda_1} + \frac{v}{\lambda_2}$, $\tilde{\Delta}_4^{(v)} = \tilde{\Delta}_4 + \frac{v}{\lambda_4}$. Then we integrate the optical polarization P_3 with the velocity distribution $g(v)$ obtaining:

$$\langle P_3 \rangle_T = \int_{-\infty}^{+\infty} \sum_{\mathcal{P}} g(v) n(\mathcal{P}) \mu_{43}^*(\mathcal{P}) \rho_{43}(\mathcal{P}, v) dv + c.c., \quad (9)$$

where we have neglected velocity-dependent pumping effects and assumed the distribution of the atoms among ground levels $n(\mathcal{P})$ to be velocity independent.

In the perturbative case $\Omega_i \ll \Gamma_i$ we can perform the above integration analytically. Since only the susceptibility $\chi(\mathcal{P})$ given in Eq. (8) is velocity-dependent, the velocity-averaged version of Eq. (7) is $\langle P_3 \rangle_T = E_1 E_2 E_4^* \sum_{\mathcal{P}} n(\mathcal{P}) \langle \chi(\mathcal{P}) \rangle_T$. Here the susceptibility $\langle \chi(\mathcal{P}) \rangle_T$ is calculated by integrating $g(v) / \tilde{\Delta}_2^{(v)} \tilde{\Delta}_3^{(v)} \tilde{\Delta}_4^{(v)*}$, which in turn is reduced into a sum of Voigt-like profile integrals. This is achieved by partial fractions decomposition as detailed in the Appendix. The result can be cast into the following form:

$$\langle \chi(\mathcal{P}) \rangle_T = -i \sqrt{\frac{\pi m}{2k_B T}} \sum_i \frac{\lambda_i \mathcal{V} \left(\tilde{\Delta}_i(\mathcal{P}) \lambda_i \sqrt{\frac{k_B T}{2m}} \right)}{Q_i(\{\tilde{\Delta}_j(\mathcal{P})\})}, \quad (10)$$

where $\mathcal{V}(z)$ is the profile function we define in the Appendix, while $Q_i(\{\tilde{\Delta}_j\})$ are second order polynomials of detunings. The polynomials $Q_i(\{\tilde{\Delta}_j\})$, given in the Appendix as well, are responsible for the sub-Doppler features in the 4WM spectrum. Their real parts are zero when real parts of any two out of three factors in the susceptibility denominator $\tilde{\Delta}_2^{(v)} \tilde{\Delta}_3^{(v)} \tilde{\Delta}_4^{(v)*}$ are zero for the same velocity v . This determines position of the resonances, as we will see in section IV.

III. EXPERIMENT

The heart of the experimental setup is a magnetically shielded rubidium vapor cell heated to the temperature of 373 K. The 2.5 cm in diameter and 7.5 cm long cell contains a natural mixture of rubidium isotopes and no buffer gas. We have decided to use only ^{85}Rb (nuclear spin $\mathcal{I} = 5/2$) due to its higher concentration and, in consequence, higher optical depth. For the ground state $|1\rangle$




Figure 3. Schematic of the experimental situation. Generation and detection stage (a) of the experimental setup; beams enter the rubidium vapor cell and the four-wave mixing signal is separated from the pump light using spatial separation, band-pass interference (BIF) filter and a notch filter (NF). The signal is detected using an avalanche photodiode (APD). The 3D configuration of beams (b) enables phase matching without beam overlap. In the focal plane (c) we see four distinct points corresponding to four beams. The signal at 762 nm is marked with an arrow.

we have chosen the $F_1 = 3$ hyperfine component of the $5^2S_{1/2}$ state. For the intermediate states we have chosen $5^2P_{3/2}$ for the $|2\rangle$ state and $5^2P_{1/2}$ for the last intermediate state $|4\rangle$. As the highest excited state $|3\rangle$ we use $5^2D_{3/2}$ level. The respectful wavelengths for the resonant transitions are $\lambda_1=780$ nm (the D2 line), $\lambda_2=776$ nm, $\lambda_3=762$ nm and $\lambda_4=795$ nm (the D1 line). Out of many possible diamond configurations the one we use has the main advantage of high efficiency of detectors for each of the wavelengths, thus enabling future quantum optics applications.

Inside the cell three beams from three different lasers at 780, 795 and 776 nm intersect at a small angle as depicted in Fig. 3. The fourth beam of light at 762 nm is generated in the cell according to the phase matching condition $\mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4$. The values of wave vectors are approximately $\mathbf{k}_1 \approx \frac{2\pi}{\lambda_1} [0, -\theta, 1]$, $\mathbf{k}_2 \approx \frac{2\pi}{\lambda_2} [0, \theta, 1]$, $\mathbf{k}_3 \approx \frac{2\pi}{\lambda_3} [-\theta, 0, 1]$ and $\mathbf{k}_4 \approx \frac{2\pi}{\lambda_4} [\theta, 0, 1]$, where $\theta = 8$ mrad. Inside the cell the beams are collimated with a $1/e^2$ diameter of approximately 3 mm. We found that the width of the phase matching is quite high, at least 2 mrad.

We use three Toptica lasers: two distributed Bragg reflector diodes at 780 nm and 795 nm and an external cavity laser at 776 nm. The beams are combined to intersect at a small angle inside the cell and then we employ a high-extinction Wollaston polarizer to prepare the light of all beams in the x polarization state.

Finally, the generated 762 nm light needs to be separated from the pump beams. We use three distinct filtering methods to obtain high signal-to-background ratio, as shown in Fig. 3(c). A tilted interference bandpass filter comes first, then we use an iris diaphragm in the focal plane to cover the pump beams, and finally we apply a notch filter with a central wavelength of 785 nm (Thorlabs NF785-33). The only residual light from the

pump lasers we detect comes from the amplified spontaneous emission at 762 nm in laser diodes. The 4WM signal is detected using an avalanche photodiode (Thorlabs APD120A).

The measurement scheme is designed to obtain maps of the 4WM signal as a function of two out of three pump laser detunings. We lock the frequency of 776 nm laser using a wavelength meter (Angstrom WS-7) at a fixed detuning Δ_{776} , nearly resonant to the $|2\rangle - |3\rangle$ transition. During the measurement, the detuning Δ_{795} of the 795 nm laser frequency is altered in small steps of $4 \times 2\pi$ MHz, while the detuning Δ_{780} of 780 nm laser is scanned over the relevant range of optical frequencies. The exact scan rate varied for different measurements, but was of the order of $50 \times 2\pi$ MHz/ms. Data from 800 to 1000 scans were collected and averaged to give a dependence of the 4WM signal from Δ_{780} for fixed Δ_{795} and Δ_{776} .

All detunings are measured from the centroid of respectful resonance line. The detunings of lasers resonant to D1 and D2 lines are determined using saturated absorption signal obtained in auxiliary rubidium vapor cells. Note that through comparing the laser detunings with the theoretical model we obtain $\Delta_2 = \Delta_{780}$, $\Delta_3 = \Delta_{780} + \Delta_{776}$ and $\Delta_4 = \Delta_{795}$.

IV. RESULTS

The first measurement we present was aimed to determine the shape of a single 4WM resonance in the regime of a perturbative solution. To verify the linear perturbative theory we performed a measurement of the 4WM signal, where all pump intensities were low and corresponded to the Rabi frequencies of $\Omega_1/2\pi=5$ MHz, $\Omega_2/2\pi=0.3$ MHz and $\Omega_4/2\pi=4$ MHz. Note that these are all lower than corresponding transitions linewidths. We chose the strongest resonance that corresponded to the path leading through $F_2 = 4$, $F_3 = 3$ and then $F_4 = 2$ levels. The choice of such a configuration entails that there should be no splitting of this resonance due to the hyperfine structure of the highest $|3\rangle$ state, as there is only one possible spin F_3 if $F_2 = 4$ and $F_4 = 2$.

Figures 4(a) and 4(b) present the experimental and theoretical 4WM signal intensity, respectively. The presented maps show an excellent conformity of theory [Eq. (10)] and experiment. The only free parameter for the theory is the intensity multiplicative factor. Additionally, we need to take into account the laser linewidth, which is of the order of several MHz for each laser when averaged over duration of a single measurement.

In Figs. 4(c) and 4(d) we present two cross-sections of the maps. The cross-sections, denoted by dashed lines on the maps, are taken far from the resonance, as the structure there is non-trivial. It exhibits two peaks and it is instructive to observe their separation and relative intensities. The cross-sections show a quantitative agreement between theory (solid line) and experiment (dots with




Figure 4. Experimental (a) and theoretical (b) maps of the intensity of the 4WM signal as a function of 780 nm and 795 nm laser detunings in case of low pump intensities around the $F_2 = 2$ and $F_4 = 4$ resonance. The cross-sections from experimental (dots with errorbars) and theoretical (solid lines) data are presented in figures (c) and (d). The detuning of 776 nm laser frequency was $\Delta_{776}/2\pi = -51$ MHz.

errorbars), confirming the correctness of our theoretical approach. On the maps we observe small discrepancy of intensity and shape in the very center of the resonance that are due to small non-linearity, as the Rabi frequencies are only slightly lower than the linewidths.

The results of second measurement we present here demonstrate the influence of the hyperfine structure of intermediate levels on the 4WM signal. For this measurement we increased the power of each of the pumping lasers, and obtained Rabi frequencies of $\Omega_1/2\pi=34$ MHz, $\Omega_2/2\pi=0.7$ MHz and $\Omega_4/2\pi=28$ MHz. As the ground




Figure 5. Experimental (a) and theoretical (b) 4WM signal demonstrating all possible resonances arising due to the hyperfine structure. The detuning of 776 nm laser frequency was $\Delta_{776}/2\pi = 30$ MHz.

state is the $F_1 = 3$ state, we expect six possible resonances, as this ground state is coupled to three hyperfine levels ($F_2 = 2, 3, 4$) of the $|2\rangle$ state and two hyperfine levels ($F_4 = 2, 3$) of the $|4\rangle$ state. We do not expect to see different resonances arising due to the hyperfine structure of the highest state $|3\rangle$, as the hyperfine splitting of this state is smaller than the resolution of our experiment.

Figures 5(a) and 5(b) present the experimental and theoretical results for the 4WM signal, respectively. We identify the positions of the resonances corresponding to the crossings of the dashed lines. Each line corresponds to a certain hyperfine level of the $|2\rangle$ or the $|4\rangle$ state; these lines correspond to zeros of the real part of denominator of Eq. (10), or more precisely the polynomials L_{ij} , defined in the Appendix. This brings about two resonance conditions:

$$\Delta_{780} = -\Delta F_1 + \frac{\lambda_1 + \lambda_2}{\lambda_1} \Delta F_2 + \frac{\lambda_1}{\lambda_2} \Delta_{776}, \quad (11)$$

$$\Delta_{795} = \frac{\lambda_1 \lambda_2}{(\lambda_1 + \lambda_2) \lambda_4} (\Delta_{780} + \Delta_{776}) + \left(\frac{\lambda_1 \lambda_2}{(\lambda_1 + \lambda_2) \lambda_4} - 1 \right) \Delta F_1 + \Delta F_4, \quad (12)$$

where again we neglected hyperfine splitting of the highest excited state $|3\rangle$. The first equation defines the lines corresponding to hyperfine levels of the $|2\rangle$ state. These are vertical on our map, but their separation is twice the hyperfine splitting, as $\frac{\lambda_1 + \lambda_2}{\lambda_1} \approx 2$. The second equation defines the lines corresponding to the $|4\rangle$ state. These lines have a slope of $\frac{\lambda_1 \lambda_2}{(\lambda_1 + \lambda_2) \lambda_4} \approx \frac{1}{2}$, but their separation is simply equal to the splitting of the $|4\rangle$ state.

For the full theoretical calculation (Fig. 5(b)) we use the non-perturbative framework introduced in section II. The results show good qualitative agreement with the experiment. We observe that both in theory and experiment the strongest resonances are the rightmost ones on our map, corresponding to highest F_2 spin. This is due to the relative amplitudes of resonances, that can be calculated even in the perturbative regime from the products of dipole moments. In table I we give these relative amplitudes for each resonance as a function of the ground state spin projection $|m_{F_1}|$. Note, that different ground state populations give rise to different 4WM resonances. Consequently, non-equilibrium populations may significantly change observed relative amplitudes. In our calculation we take this into account by modifying ground state populations according to the numerical solution of Liouville equation with relaxation and repopulation [26].

One of the non linear effects we predict theoretically and observe experimentally is Autler-Townes-like splitting [15, 24] of the $F_2 = 4, F_4 = 3$ resonance, that does not occur in the linear regime. Another effect observed near the $F_2 = 4, F_4 = 2$ resonance is bending of the resonance line towards the vertical direction. Namely, we note relatively higher intensity as compared to the linear case in the top-left and bottom-right corners of the resonance.

In the experiment we witness two more effects that are not present in the theoretical results. The first effect is that the leftmost resonances, corresponding to lowest F_2 spin values, are even less intense than predicted. The second effect is the appearance of narrow diagonal lines in the 4WM signal, which should be less intense according to our theory. We believe that these two effects stem from velocity selective optical pumping, that our model does not take into account at all. Nevertheless, we find the agreement between our theoretical and experimental results satisfactory.

V. CONCLUSIONS

Beginning with the theoretical description of a four-level atom, we have derived a model for the the intensity

of 4WM in a multi-level atomic medium with inhomogeneous broadening. Our model is a reasonable compromise between simplicity and accuracy for predicting intensities, shapes and positions of 4WM resonances.

In the experimental part we have demonstrated the influence of the rich level structure of rubidium on the 4WM process. We have obtained excellent agreement between theory and experiment for low pumping intensities, as well as good qualitative agreement for higher pump intensities, where more non-linear processes contribute.

Combined with optical pumping, our methods may facilitate the design of coherent control in a multi-level atomic medium with light. We have shown that light at 762 nm couples to the population of the ground state. Future investigations could include coupling of 762 nm light to the ground state coherence in order to create Raman-scattering-like light-atom interface [27] or engineering the 4WM signal by optically pumping the ground state. This could be accomplished in the cells with buffer gas that makes the atomic motion diffusive [28], the cell with anti-relaxation coating [29] or in a cold atomic ensemble. Another approach could involve control of 4WM signal through velocity-selective optical pumping.

ACKNOWLEDGMENTS

We acknowledge the generous support from Konrad Banaszek, Czesław Radzewicz and Tadeusz Stacewicz. This work was supported by the National Science Center grant no. DEC-2011/03/D/ST2/01941 and by the Polish Ministry of Science and Higher Education ‘‘Diamantowy Grant’’ project no. DI2013 011943.

APPENDIX: ENSEMBLE-AVERAGED SUSCEPTIBILITY

We decompose the formula (2) for the optical coherence ρ_{43} in the perturbative regime with velocity-

dependent detunings, obtaining a sum of three expressions with first order polynomials in terms of velocity in the denominators:

$$\left(\tilde{\Delta}_2^{(v)}\tilde{\Delta}_3^{(v)}\tilde{\Delta}_4^{(v)*}\right)^{-1} = \sum_i \left(\tilde{\Delta}_i^{(v)}Q_i(\{\tilde{\Delta}_j\})\right)^{-1} \quad (13)$$

The polynomials are: $Q_2 = L_{23}L_{24}$, $Q_3 = \frac{\lambda_4(\lambda_1+\lambda_2)^2}{\lambda_1\lambda_2^2}L_{23}L_{34}$, $Q_4 = \frac{\lambda_1}{\lambda_2}L_{24}L_{34}$, where $L_{23} = \left(1 + \frac{\lambda_1}{\lambda_2}\right)\tilde{\Delta}_2 - \tilde{\Delta}_3$, $L_{24} = \frac{\lambda_1}{\lambda_2}\tilde{\Delta}_2 - \tilde{\Delta}_4$ and $L_{34} = \lambda_4\left(\frac{1}{\lambda_1} + \frac{1}{\lambda_2}\right)\tilde{\Delta}_4 - \tilde{\Delta}_3$. Equating the real part of L_{ij} to zero yields resonance conditions given by Eq. (11) and (12). The expression can now be easily integrated with velocity distribution $g(v)$, as each component can be integrated separately to give the Voigt-like profile used in Eq. (10):

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{e^{-v^2/2\langle v^2 \rangle}}{v/\lambda + \tilde{\Delta}} dv &= \\ &= -i\pi\lambda \exp\left(-\frac{\tilde{\Delta}^2\lambda^2}{2\langle v^2 \rangle}\right) \left(\operatorname{erf}\left(i\frac{\tilde{\Delta}\lambda}{\sqrt{2\langle v^2 \rangle}}\right) \pm 1\right) = \\ &= -i\pi\lambda\mathcal{V}\left(\frac{\tilde{\Delta}\lambda}{\sqrt{2\langle v^2 \rangle}}\right), \quad (14) \end{aligned}$$

where we take +1 for $\Im m\tilde{\Delta} > 0$ and -1 otherwise and $\langle v^2 \rangle = \frac{k_B T}{m}$. The profile function is defined as:

$$\mathcal{V}(z) = \begin{cases} e^{-z^2}(\operatorname{erf}(iz) + 1) & \Im m(z) > 0 \\ e^{-z^2}(\operatorname{erf}(iz) - 1) & \Im m(z) < 0 \end{cases} \quad (15)$$

Note that it is customary to use the Faddeeva function $w(z) = e^{-z^2}(\operatorname{erf}(iz) + 1)$ [26], but neither this function nor the plasma dispersion function can be used here, as they would yield incorrect results in the lower half of the complex plane.

-
- [1] T. Peyronel, O. Firstenberg, Q.-Y. Liang, S. Hofferberth, A. V. Gorshkov, T. Pohl, M. D. Lukin, and V. Vuletić, *Nature* **488**, 57 (2012).
 - [2] G. Morigi, S. Franke-Arnold, and G.-L. Oppo, *Phys. Rev. A* **66**, 053409 (2002).
 - [3] S. Clow and T. Weinacht, *Phys. Rev. A* **82**, 023411 (2010).
 - [4] H.-g. Lee, H. Kim, J. Lim, and J. Ahn, *Phys. Rev. A* **88**, 053427 (2013).
 - [5] J. F. Sell, M. A. Gearba, B. D. DePaola, and R. J. Knize, *Opt. Lett.* **39**, 528 (2014).
 - [6] T. Meijer, J. D. White, B. Smeets, M. Jeppesen, and R. E. Scholten, *Opt. Lett.* **31**, 1002 (2006).
 - [7] A. S. Zibrov, M. D. Lukin, L. Hollberg, and M. O. Scully, *Phys. Rev. A* **65**, 051801 (2002).
 - [8] A. M. Akulshin, R. J. McLean, A. I. Sidorov, and P. Hanaford, *Opt. Express* **17**, 22861 (2009).
 - [9] G. Walker, A. S. Arnold, and S. Franke-Arnold, *Phys. Rev. Lett.* **108**, 243601 (2012).
 - [10] T. Chanelière, D. N. Matsukevich, S. D. Jenkins, T. A. B. Kennedy, M. S. Chapman, and A. Kuzmich, *Phys. Rev. Lett.* **96**, 093604 (2006).
 - [11] P. S. Donvalkar, V. Venkataraman, S. Clemmen, K. Saha, and A. L. Gaeta, *Opt. Lett.* **39**, 1557 (2014).
 - [12] A. G. Radnaev, Y. O. Dudin, R. Zhao, H. H. Jen, S. D. Jenkins, A. Kuzmich, and T. A. B. Kennedy, *Nat. Phys.* **6**, 894 (2010).
 - [13] H. H. Jen and T. A. B. Kennedy, *Phys. Rev. A* **82**, 023815

- (2010).
- [14] R. T. Willis, F. E. Becerra, L. A. Orozco, and S. L. Rolston, *Phys. Rev. A* **79**, 033814 (2009).
- [15] F. E. Becerra, R. T. Willis, S. L. Rolston, and L. A. Orozco, *Phys. Rev. A* **78**, 013834 (2008).
- [16] F. E. Becerra, R. T. Willis, S. L. Rolston, H. J. Carmichael, and L. A. Orozco, *Phys. Rev. A* **82**, 043833 (2010).
- [17] A. Kölle, G. Epple, H. Kübler, R. Löw, and T. Pfau, *Phys. Rev. A* **85**, 063821 (2012).
- [18] B. Huber, A. Kölle, and T. Pfau, *Phys. Rev. A* **90**, 053806 (2014).
- [19] M. M. Müller, A. Kölle, R. Löw, T. Pfau, T. Calarco, and S. Montangero, *Phys. Rev. A* **87**, 053412 (2013).
- [20] R. T. Willis, F. E. Becerra, L. A. Orozco, and S. L. Rolston, *Opt. Express* **19**, 14632 (2011).
- [21] W. Zhang, D.-S. Ding, J.-S. Pan, and B.-S. Shi, *Chinese Phys. Lett.* **31**, 064208 (2014).
- [22] B. Srivathsan, G. K. Gulati, B. Chng, G. Maslennikov, D. Matsukevich, and C. Kurtsiefer, *Phys. Rev. Lett.* **111**, 123602 (2013).
- [23] G. K. Gulati, B. Srivathsan, B. Chng, A. Cerè, D. Matsukevich, and C. Kurtsiefer, *Phys. Rev. A* **90**, 033819 (2014).
- [24] S. H. Autler and C. H. Townes, *Phys. Rev.* **100**, 703 (1955).
- [25] D. A. Steck, “Rubidium 85 D line data,” (2013).
- [26] W. Happer, Y.-Y. Jau, and T. Walker, *Optically Pumped Atoms* (Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2010).
- [27] R. Chrapkiewicz and W. Wasilewski, *Opt. Express* **20**, 29540 (2012).
- [28] M. Parniak and W. Wasilewski, *Appl. Phys. B* **116**, 415 (2013).
- [29] M. V. Balabas, K. Jensen, W. Wasilewski, H. Krauter, L. S. Madsen, J. H. Müller, T. Fernholz, and E. S. Polzik, *Opt. Express* **18**, 5825 (2010).