

# Metric-torsion preheating: cosmic dynamo mechanism?

By L.C. Garcia de Andrade<sup>1</sup>

## Abstract

Earlier Bassett et al [Phys Rev D 63 (2001) 023506] investigated the amplification of large scale magnetic fields during preheating and inflation in several different models. They argued that in the presence of conductivity resonance effect is weakened. From a dynamo equation in spacetimes endowed with torsion recently derived by Garcia de Andrade [Phys Lett B 711: 143 (2012)] it is shown that in a universe with pure torsion in Minkowski spacetime the cosmological magnetic field is enhanced by ohmic or non-conductivity effect, which shows that the metric-torsion effects is worth while of being studied. In this paper we investigated the metric-torsion preheating perturbation, which leads to the seed cosmological magnetic field in the universe with torsion is of the order of  $B_{seed} \sim 10^{-37} Gauss$  which is several orders of magnitude weaker than the decoupling value obtained from pure metric preheating of  $10^{-15} Gauss$ . Despite of the weakness of the magnetic field this seed field may seed the galactic dynamo.

Key-words: Modified gravity theories, primordial magnetic fields.

---

<sup>1</sup>Departamento de Física Teórica - IF - UERJ - Rua São Francisco Xavier 524, Rio de Janeiro, RJ, Maracanã, CEP:20550. e-mail:garcia@dft.if.uerj.br

# 1 Introduction

The excitement of BICEP2 experiment at south pole [1] to observe inflation and primordial gravitational waves, and recent paper by Bonvin et al [2] by considering that BICEP2 could also have primordial magnetic field observed leads us to revisit Bassett et al [3] where they consider that metric preheating may induce amplification of the magnetic fields at cosmological scales as a cosmic dynamo. As is well-known the preheating is a phase after inflation where particles are created. Bassett et al [3] investigated the amplification of large scale magnetic fields during preheating and inflation in different models with different conformal and gauge couplings. In the presence of conductivity resonance effect is weakened. From dynamo equation in spacetimes endowed with torsion recently derived by Garcia de Andrade [4] it is shown that in a universe with pure torsion in Minkowski spacetime the cosmological magnetic field is enhanced by ohmic or low conductivity effects, which shows that the metric-torsion effects is worth while of being studied. In this paper we investigated the metric-torsion preheating perturbation, which leads to the seed cosmological magnetic field in the universe with torsion is of the order of  $B_{seed} \sim 10^{-37} Gauss$  several orders of magnitude weaker than the decoupling value obtained from pure metric preheating of  $10^{-25} Gauss$ . Despite of the weakness of the magnetic field this seed field may seed the galactic dynamo. The paper is organised as follows: In section 2 the torsionic dynamo equation is revisited. In section 3 we present the solution of electromagnetic wave equation for the perturbed metric in Riemann-Cartan geometry with torsion. Conclusions are presented in section 4.

## 2 Dynamo equation endowed with torsion

In this section we shall consider the torsion dynamo equation obtained recently from a parity violating torsion lagrangean [4]. Here  $\mathbf{T}$  is a torsion vector. The self-induction magnetic equation in the spacetime torsion background is [5]

$$\partial_t \mathbf{B} = \nabla \times [\mathbf{v} \times \mathbf{B} + \eta[\Delta \mathbf{B} - (\nabla \cdot \mathbf{T})\mathbf{B}]] \quad (1)$$

Since we are only interested here on torsion effects given by ohmic losses we express the dynamo equation as

$$\partial_t B \sim [\eta(\Delta B - (\nabla \cdot \mathbf{T})B)] \quad (2)$$

Considering that dimensionally the LHS of this equation is equivalent to the magnetic B field divided by the decay time  $\tau_{torsion}$  and that the  $\Delta \sim -k^2$  where k is the inverse of the coherent length L, one obtains

$$\frac{1}{\tau} \sim \eta[-k^2 + ikT] \quad (3)$$

By taking only into account torsion ohmic losses origin, one obtains in modulus

$$\frac{1}{\tau} \sim \eta[kT] \quad (4)$$

Now making use of the astronomical data  $L \sim 10^2 pc \sim 10^{10} cm$ , the ohmic resistivity  $\eta \sim 10^{21} cm^2 s^{-1}$  and torsion given by Lammerzahl [6] estimate of Hughes-Drever experiment  $T \sim 10^{-17} cm^{-1}$  after a trivial computation one obtains  $\tau_{torsion} \sim 10^{11} years$ , which exceeds the age of the universe  $10^{10} years$ . Thus with this data the origin of magnetic fields from torsion ohmic losses is primordial. Nevertheless if one uses stronger ohmic losses such as  $10^{26} cm^2 s^{-1}$  this situation changes and the time decay of magnetic fields is  $10^9 years$  which does not exceed the universe age and therefore an efficient dynamo mechanism is necessary for the remnant magnetic fields present in the actual universe. Now let us compute the cosmological magnetic field obtained from the solution of the above dynamo equation. The solution of this equation is then given by

$$B_G \sim \eta k T \tau_{torsion} B_{seed} \quad (5)$$

By taking into account that the galactic magnetic field  $B_G \sim 10^{-6} Gauss$  one obtains a seed field of  $B_{seed} \sim 10^{-17} Gauss$ . Now let us discuss the mild

galactic dynamos which gives rise to  $10^3 s^{-1}$  amplification of magnetic fields and obtain stringent limits for Lorentz violation than the one obtained so far by Kostelecky et al [7]. Earlier Berazhiani and Dolgov [8] have obtained mild galactic dynamo with amplification factor of  $10^4 s^{-1}$ , which shows that the dynamo considered here is still more mild. To obtain this result we start by solving the dynamo equation above to find a general solution

$$\partial_t \mathbf{B}(t) = \nabla \times \mathbf{B} + \frac{1}{\kappa} \nabla \times \mathbf{J} - \eta(\nabla \cdot \mathbf{T}) \mathbf{B} \quad (6)$$

where  $\eta = \frac{1}{\kappa}$  is the ohmic loss and  $\kappa$  is the conductivity. Here  $\mathbf{J} = \nabla \times \mathbf{B}$  is the electric current. Formally this equation can be written as

$$\partial_t \ln B(t) = \frac{1}{L} v + \frac{\eta}{L^2} [1 - LT] \quad (7)$$

and its solution is given by the integral

$$B(t) = \int B_{seed} e^{\frac{v}{L} + \frac{\eta}{L^2} [1 - LT]} dt \quad (8)$$

the exponential is considered as the protogalactic dynamo. Now let us use the expression

$$B(t) = B_{seed} \eta \frac{T}{L} \quad (9)$$

to obtain the amplification factor  $\gamma$  as

$$\gamma = \eta \frac{T}{L} \quad (10)$$

Thus to an amplification factor of  $10^4 s^{-1}$  one obtains  $T \sim 10^{-23} GeV$ , while for a mild galactic dynamo of  $\gamma \sim 10^3 s^{-1}$  one obtains. Let us now compute the magnetic field seeds  $B_{seed}$  based on the last computation if the torsion field is  $T \sim 10^{-9} cm^{-1}$  and the  $L \sim 10 Mpc \sim 10^{15} cm$ , then the magnetic field seed is  $B_{seed} \sim 10^{-10} Gauss$ . This is well within the limits for galactic dynamo seeds.

### 3 Metric-torsion preheating as cosmic dynamos?

In this section we discuss the magnetic field amplification due to large metric-torsion perturbations. Here we extend the analysis of metric perturbations due to Maroto [9] to spacetimes with torsion. Let us start with the line element for a flat FLRW model with scalar metric perturbations in the conformal Newtonian or longitudinal gauge

$$ds^2 = a^2(\eta)[-(1 + 2\Phi)d\eta^2 + (1 - 2\Phi)\delta_{ij}dx^i dx^j] \quad (11)$$

where  $\eta$  is the conformal time. Considering the minimal coupling between the e.m fields tensor and torsion tensor  $S^\alpha{}_{\mu\nu}$  as given by  $F_{\mu\nu} = 2\partial_{[\mu}A_{\nu]} + 2S^\alpha{}_{\mu\nu}A^\alpha$ . From Lagrangean for the two-field model the e.m field equation yields

$$\nabla_\mu F^{\mu\nu} = 0 \quad (12)$$

where  $\nabla$  is the Riemann-Cartan connection, taking only first-order torsion terms one obtains the following wave equation

$$\frac{\partial}{\partial\eta}[(1-2\Phi)[\partial_i A_0 - \partial_0 A_i + 2S^k{}_{i0}A_k]] + \frac{\partial}{\partial x^j}[(1+2\Phi)[\partial_j A_i - \partial_i A_j + 2S^k{}_{ji}A_k]] = 0 \quad (13)$$

where we use the relations  $\sqrt{-g} = a^4(1 - 2\Phi)$ ,  $g^{00} = -a^{-2}(1 - 2\Phi)$ ,  $g^{ii} = a^{-2}(1 + 2\Phi)$  in the perturbed metric (11). From Coulomb gauge  $A_0 = 0$ ,  $\partial^i A_i = 0$  one obtains

$$\frac{d^2}{d\eta^2}A_i - \nabla^2 A_i + 2(1+2\Phi)(\partial_j S^k{}_{ji})A_k + 2(1+2\Phi)S^k{}_{ji}\partial_j A_k + \frac{\partial\Phi}{\partial x^j}S^k{}_{ji}A_k + \frac{\partial\Phi}{\partial x^j}S^k{}_{ji}\partial_j A_k = 0 \quad (14)$$

Now by Fourier transforming this equation one obtains

$$\frac{d^2}{d\eta^2}A_i + k^2 A_k - 2\frac{d\Phi}{d\eta}\frac{dA_k}{d\eta} + 4k^2\Phi A_k = +2ik_j S^k{}_{ji}A_k + 2(1 - 2\Phi)\frac{dS^l{}_{k0}}{d\eta} \quad (15)$$

splitting the real and imaginary parts one obtains the following constraint on torsion  $S^k{}_{ji}k_j = 0$  and the wave equation

$$\frac{d^2}{d\eta^2}A_i + k^2 A_k - 2\frac{d\Phi}{d\eta}\frac{dA_k}{d\eta} + 4k^2\Phi A_k = +2(1 - 2\Phi)\frac{dS^l{}_{k0}}{d\eta}A_k \quad (16)$$

The term  $\frac{dA_k}{d\eta}$  can be absorbed in a electric current term to take into account the effects of conductivity as in Maroto [9] using torsionless preheating metric perturbations. Taking the torsion tensor as  $S^l_{k0} := \delta^l_k S_0$  one reduces the last equation to

$$\frac{d^2}{d\eta^2} A_k + k^2 A_k - 2 \frac{d\Phi}{d\eta} \frac{dA_k}{d\eta} + 4k^2 \Phi A_k = +2(1 - 2\Phi) \frac{dS_0}{d\eta} A_k \quad (17)$$

Now taking the transformation [3]  $\tilde{A}_k := (1 - 2\Phi)A_k$  one obtains from the last equation the following solution

$$\tilde{A}_k(\eta) = \tilde{A}^{in}_k + \frac{1}{k} \int \left[ \frac{d^2\Phi}{d\eta^2} + 4 \frac{d\Phi}{d\eta} (1 + \Phi) S_0 \right] A_k \times \sin(\eta - \eta^*) d\eta^* \quad (18)$$

This last equation shows that a proper torsion gauge can lead to a differential equation constraint to torsion and metric scalar perturbation  $\Phi$  given by

$$\frac{d^2}{d\eta^2} \Phi + 2 \frac{d\Phi}{d\eta} + \Phi S_0 = 0 \quad (19)$$

The solution of this equation can be simplify with the approximation  $\frac{d^2}{d\eta^2} \Phi \sim 0$  yields

$$\Phi(\eta) = \Phi_0 e^{-S_0 \eta} \quad (20)$$

This equation shows that for a positive torsion the metric perturbation decays while a negative torsion induces an amplification in the metric perturbation. Now let us relax this strong constrain and consider the solution above in general form to obtain the magnetic field. This can be done in analogy to Bassett et al [3]

$$|B_k| \approx \frac{k}{a^2} \int \left( \frac{d\Psi}{d\eta} \right)^2 d\eta \quad (21)$$

where now by analogy  $\Psi$  is given by

$$\frac{d^2\Psi}{d\eta^2} = \left[ \frac{d^2}{d\eta^2} \Phi + 4(1 + \Phi) \frac{d\Phi}{d\eta} + \Phi S_0 \right] \quad (22)$$

Thus from Bogoliubov coefficient  $\beta_k$  one considers the magnetic energy density

$$\rho_B = \left( \frac{k}{a^4} \right)^4 |\beta_k|^2 \quad (23)$$

Therefore from these last expressions one obtains finally the magnetic field expression from solely torsion contributions

$$|B_k| \approx \frac{kS_0}{a^2} \int \left( \frac{d\Phi^2}{d\eta} \right) d\eta \quad (24)$$

which in turn yields

$$|B_k|_{torsion} \approx \frac{kS_0}{a^2} (\Phi)^2 \approx 10^{-17} \frac{k}{a^2} (\Phi)^2 \quad (25)$$

where we have used the Hughes-Drever experiment torsion result of  $S_0 \approx 10^{-17} cm^{-1}$ . For appropriated coherent scales such as the ones used in Bassett et al [3] one obtains  $|B_k|_{torsion} \approx 10^{-37} G$  which is much weaker than the general relativistic preheating metric scalar perturbation obtained by Bassett et al [3] which is  $|B_k| \sim 10^{-25} G$ . In spite of that this value is still able to seed galactic dynamo.

## 4 Conclusions

The main result of this note is that primordial magnetic fields could be of the torsion origin depending of the amount of the ohmic losses obtained from spacetime torsion. The cosmological magnetic field seeds range between  $10^{-17} G \leq B \leq 10^{-10} G$  which is a more stringent limit than the ones obtained by Barrow et al [10]. An important feature of this paper is that galactic dynamo mechanisms allowed us to place limits on Lorentz violation similarly to what has been recently proposed by the author [11]. Actually the limit  $B \sim 10^{-22} Gauss$  which has been recently obtained by us [11], falls very well within the magnetic field to seed galactic dynamos as shown recently by Barrow et al [10]. It falls shortly below the interval for the B-field given by K. Pandey et al [12] for primordial magnetic field lines using  $Ly\alpha$  clouds. Recently R. Banerjee [13] has informed us that values of  $B - field$  as low as  $10^{-32} Gauss$  may also seed galactic dynamos. Another important source of magnetic field amplification can be metric preheating [9] in a spacetime with torsion which we discuss here in full detail. In this paper we also show that torsion Ohmic effects also serve to amplify magnetic fields during preheating, though the effects seems to be minimal due to the fact that the magnetic field at decoupling is  $B_{dec} \sim 10^{-25} G$  while respective torsion effects

are  $B_{torsion} \sim 10^{-37}G$ . However, despite of the weakness of this field this value is still able to seed galactic dynamo.

## 5 Acknowledgements

We would like to express my gratitude to D Sokoloff and A Brandenburg for helpful discussions on the problem of dynamos and torsion. An anonymous referee is also grateful acknowledged for pointing out a mistake on a previous draft of this paper. Special thanks go to my wife Telma Mussel Rabello and Dr Carlos de Souza Lima for technical support. Financial support from CNPq. and University of State of Rio de Janeiro (UERJ) are grateful acknowledged.

## References

- [1] P. A. R. A de et al. [BICEP2 Collaboration], arXiv: 1403.3985.
- [2] C Bonvim, R Durrer and Roy maartens, Phys Rev Lett 112, 191303 (2014).
- [3] B A Bassett, G Polifrone, S Tsujikawa and F Viniegra, Phys Rev D 63 (2001) 023506. C Tsagas, Phys Rev D 72: 123509 (2005).
- [4] L C Garcia de Andrade, Phys Lett B 711: 143 (2012).
- [5] V de Sabbata and C Sivaram, Spin and Torsion and Gravitation, World Scientific (1995).
- [6] C Laemmerzahl, Phys Lett A 228, 223 (1997).
- [7] A Kostelecky, N Russell and J Tasson, Phys Rev Lett **100**: 111102 (2008).
- [8] Z Berezhiani and A D Dolgov, Astroparticle Phys 21; 59 (2004).
- [9] A L Maroto, Phys Rev D 64: 083006, (2001). L C Garcia de Andrade, Class and Quantum Gravity vol 18,18, 3097 (2001).
- [10] J D Barrow, C Tsagas and K. Yamamoto, Phys Rev D 86, 023533 (2012).



- [11] L C Garcia de Andrade, Nuclear Phys **B** (2011). L Garcia de Andrade, Phys Lett **B** 468, 28 (2011). L Garcia de Andrade, Lorentz violation bounds from torsion trace and radio galactic dynamos, Phys Rev D (Brief Reports) (2011). L Garcia de Andrade, Galactic dynamo seeds from Einstein-Cartan gravity and f-meson dominance gravity, Advances studies in theoretical physics (2014) in press.
- [12] K.L. Pandey and S.K. Sethi, Astrophysical J., 762: 15, (2013).
- [13] R. Banerjee, Private communication (2013).