

# Neutralino-Neutralino pair annihilation crosssections with flavour violation

---

**V. Suryanarayana Mummidi**

*Centre for High Energy Physics, Indian Institute of Science, Bangalore 560012*

*E-mail:* [soori9@cts.iisc.ernet.in](mailto:soori9@cts.iisc.ernet.in)

ABSTRACT: We present the first exact and most general neutralino-neutralino annihilation cross sections in the presence of flavor violation. The existing set of calculations in literature do not include flavour violation. These calculations are important for the calculation of relic density and indirect detection of dark matter in various experiments.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Thermal relic density</b>	<b>4</b>
<b>3</b>	<b>Cross sections</b>	<b>6</b>
3.1	$\chi_i^0 \chi_j^0 \rightarrow H_m^0 H_n^0$	6
3.2	$\chi_i^0 \chi_j^0 \rightarrow A_1^0 A_1^0$	8
3.3	$\chi_i^0 \chi_j^0 \rightarrow H_1^0 A_1^0$	10
3.4	$\chi_i^0 \chi_j^0 \rightarrow H_2^0 A_1^0$	12
3.5	$\chi_i^0 \chi_j^0 \rightarrow H^+ H^-$	13
3.6	$\chi_i^0 \chi_j^0 \rightarrow W^\pm H^\mp$	15
3.7	$\chi_i^0 \chi_j^0 \rightarrow Z H_n^0$	17
3.8	$\chi_i^0 \chi_j^0 \rightarrow Z H_n^0$	19
3.9	$\chi_i^0 \chi_j^0 \rightarrow W^\pm W^\mp$	21
3.10	$\chi_i^0 \chi_j^0 \rightarrow Z Z$	23
3.11	$\chi_i^0 \chi_j^0 \rightarrow f_m \bar{f}_n$	24
<b>4</b>	<b>summary</b>	<b>30</b>
<b>A</b>	<b>Mixing matrices and Coupling constants</b>	<b>30</b>
<b>B</b>	<b>Auxiliary functions</b>	<b>34</b>

---

## 1 Introduction

One of the most astounding discoveries of the twentieth century in our understanding of the Universe' content is that ordinary baryonic matter, that is, matter made up of protons and neutrons, is not the dominant form of material in the Universe. Rather, some strange new form of matter, dubbed 'dark matter', fills our Universe, and it is roughly six times more abundant than ordinary matter. Although we are yet to detect this strange material in the laboratory, there is a great deal of evidence which points to the necessity of its existence.

The existence of dark matter is well motivated by several lines of evidence but the exact nature of dark matter remains unknown. One of the most popular dark matter candidates are WIMPs (Weakly Interacting Massive Particles); in other words, they are massive particles that are electrically neutral which do not interact very strongly with other matter. Standard Model (SM) does not contain any particle that could act as the dark matter. The only stable, electrically neutral, and weakly interacting particles in the SM are the neutrinos. Neutrinos being relativistic particles they cannot form cold dark matter. The

lack of a dark matter candidate in the SM suggests that it must be extended. The most promising model is Minimal Supersymmetric Standard Model (MSSM)[1]. An interesting feature of MSSM (most supersymmetric models) is the existence of a multiplicative conserved quantum number called R-parity, in which each superpartner is assigned  $R=-1$ , and each ordinary particle is assigned  $R=+1$ . This quantum number implies that supersymmetric particles must be created or destroyed in pairs, and that the lightest supersymmetric particle (LSP) is absolutely stable; just as the electron is stable since electric charge is conserved and there is no lighter charged particle into which it could decay. This fact is what makes supersymmetric particles dark matter candidates. If supersymmetry exists and R-parity is conserved, then some LSP's must exist from the Early Universe. They annihilate and coannihilate into SM particles and some of them will be left over as thermal relics in the present Universe. More detailed calculation of thermal relic density of dark matter is presented in the next section.

In MSSM, superpartners of neutral gauge bosons and Higgs; Bino, Wino and Higgsinos respectively have same quantum numbers after electroweak symmetry breaking. So they all mix and form four mass eigen states called Neutralinos, denoted as  $\chi_1^0, \chi_2^0, \chi_3^0$  and  $\chi_4^0$ . By convention, these are labeled in ascending order, so that  $m_{\chi_1^0} < m_{\chi_2^0} < m_{\chi_3^0} < m_{\chi_4^0}$ . If R-parity is conserved, lightest neutralino is the LSP in most of the models. It is natural to consider the possibility that the neutralino is the dark matter candidate in MSSM. In gauge eigen state basis,  $\psi^T = (\tilde{B}, \tilde{W}, \tilde{H}_1, \tilde{H}_2)$ , the neutralino mass part of the Lagrangian is

$$\mathcal{L}_{neutralino} = -\frac{1}{2} \psi^T M_\chi \psi \quad (1.1)$$

Neutralino mass matrix  $M_\chi$  is given by

$$\begin{pmatrix} M_1 & 0 & -\frac{e v_1}{2 c_W} & \frac{e v_2}{2 c_W} \\ 0 & M_2 & \frac{e v_1}{2 s_W} & -\frac{e v_2}{2 s_W} \\ -\frac{e v_1}{2 c_W} & \frac{e v_1}{2 s_W} & 0 & -\mu \\ \frac{e v_2}{2 c_W} & -\frac{e v_2}{2 s_W} & -\mu & 0 \end{pmatrix} \quad (1.2)$$

where  $M_1$  and  $M_2$  are the Bino and Wino masses respectively.  $\mu$  is the higgsino mass parameter, 'e' is the gauge coupling,  $v_1$  and  $v_2$  are vacuum expectation values and  $\theta_W$  ( $c_W = \cos \theta_W$  and  $s_W = \sin \theta_W$ ) is Weinberg angle. The mass matrix  $M_\chi$  can be diagonalized by a unitary matrix  $Z_N$  to obtain mass eigenstates:

$$\chi_i^0 = (Z_N)_{ij} \psi_j \quad (1.3)$$

so that

$$Z_N^{-1} M_\chi Z_N = \begin{pmatrix} m_{\chi_1^0} & 0 & 0 & 0 \\ 0 & m_{\chi_2^0} & 0 & 0 \\ 0 & 0 & m_{\chi_3^0} & 0 \\ 0 & 0 & 0 & m_{\chi_4^0} \end{pmatrix} \quad (1.4)$$

So  $\chi_1^0$  is the weakly interacting massive particle which is a dark matter candidate with mass  $m_{\chi_1^0}$ . With high precision measurements of dark matter relic density, it may not be of distant possibility to constrain supersymmetric models. If supersymmetry is discovered at colliders, it may also be possible to test the hypothesis that neutralino is the dark matter candidate. In either case, it is important to know the thermal relic density of neutralino as accurately as possible. Thermal relic density of dark matter is given by

$$\Omega_\chi = \frac{n_\chi m_\chi}{\rho_c} \quad (1.5)$$

where  $\rho_c$  is the critical density given by  $\frac{3H^2}{8\pi G}$ .  $G$  and  $H$  are gravitational constant and present day Hubble parameter respectively. The solution of Boltzmann equation gives the number density of any particle in the Early Universe as it evolves. As we have assumed that dark matter is thermally produced and is in thermal equilibrium with the background in the early phase of the Universe, its number density  $n_\chi$  can be calculated by solving Boltzmann rate equation

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -\langle\sigma|v|\rangle(n_\chi^2 - n_{\chi,eq}^2) \quad (1.6)$$

Where  $t$  is time, and  $\langle\sigma_{X\bar{X}}|v|\rangle$  is the thermally averaged cross section times the relative velocity of the interacting particles. The first term on the right-hand side is the reduction in LSP number density due to the Hubble expansion, the second term is reduction due to self-annihilation and coannihilations. One can solve eq. (1.6) numerically or using the standard "freeze-out" approximation, and obtains the number density at  $t=0$  (today) provided  $\langle\sigma|v|\rangle$  is known. As it was pointed out in [2], coannihilations between neutralino and next to lightest supersymmetric particle (nLSP) play a significant role in the determination of the dark matter number density if their masses are nearly degenerate. So it is necessary to consider coannihilation of neutralino(LSP) with nLSP. In MSSM, nLSP could be neutralino, chargino or a slepton depending on the boundary conditions at the supersymmetry breaking scale<sup>1</sup>. Therefore, the general calculation of neutralino number density involves consideration of cross sections:

- |                                       |   |
|---------------------------------------|---|
| (i) Neutralino (LSP)-Neutralino(LSP)  | (ii) Neutralino (LSP)-Neutralino (nLSP) |
| (iii)Neutralino (LSP)-Chargino (nLSP) | (iv) Neutralino (LSP)- Slepton (nLSP)   |

So the problem of calculation of today's neutralino dark matter relic density is usually the calculation of the cross section of the initial states (i), (ii), (iii) and (iv) into all standard model particles. The relic density of neutralinos in the MSSM has been calculated by several authors over the years [2–12]. Nihei etal. [9, 10] have considered Neutralino cross sections extensively except for the case (ii). They also have assumed neutralino mixing matrix  $Z_N$  to be real for simplicity and not included flavour violation in the supersymmetric soft sector. Authors in[11, 12] have considered effect of flavor violation on coannihilation and annihilations. In most calculations of the cross sections flavor violation is not considered seriously because of the existing low energy constraints from rare decays like  $\mu \rightarrow e\gamma$ ,

---

<sup>1</sup>Stops or gluinos could also be nLSP. One has to do case by case study of cross sections

Process	Exchanged Particle	
	s-channel	t- and u-channel
$\chi_i^0 \chi_j^0 \rightarrow H_m^0 H_n^0$	$H_k^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow A_1^0 A_1^0$	$H_k^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow H_1^0 A_1^0$	$A_1^0, Z$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow H_2^0 A_1^0$	$H_k^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow H_1^+ H_1^-$	$H_k^0, Z$	$\chi_k^\pm$
$\chi_i^0 \chi_j^0 \rightarrow W^\pm H_1^\mp$	$H_k^0, A_1^0$	$\chi_k^\pm$
$\chi_i^0 \chi_j^0 \rightarrow Z H_n^0$	$Z, A_1^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow Z A_1^0$	$H_k^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow W^+ W^-$	$H_k^0, Z$	$\chi_k^\pm$
$\chi_i^0 \chi_j^0 \rightarrow Z Z$	$H_k^0$	$\chi_k^0$
$\chi_i^0 \chi_j^0 \rightarrow f_m f_n$	$H_k^0, A_1^0, Z$	$f_k$

**Table 1.** A complete set of neutralino pair-annihilation channels into tree-level two-body final states in the MSSM. The index k runs as follows: For neutralinos k=1,...,4; for charginos k=1,2; for sfermions k=1,...,6; for Higgs k=1,2 and for fermions m,n=1,...,6

$\tau \rightarrow \mu\gamma$ , etc. in the leptonic sector and  $b \rightarrow s\gamma, B - \bar{B}^0, B_s - \bar{B}_s$ , in the hadronic sector. However, it is known that there are regions in the parameter space where these constraints are highly weakened. In the leptonic sector cancellations of amplitude of  $l_j \rightarrow l_i\gamma$  are widely noted [13]. These parameter regions increase if one considers a more general SUSY breaking framework like the one with the non-universal gauge masses[14, 15]. In the hadronic sector, constraints are weakened typically for a relatively heavy squark spectrum  $\sim 1$  TeV or so. In fact, so much so that, in mini-split models one expects a little bit of flavor violation to be present in the 2-3 sector[16, 17].

The purpose of our paper is to present most general neutralino(LSP)-neutralino (LSP/nLSP) cross sections without assuming minimal flavor violation or real mixing matrix for neutral gauge eigen state matrix  $Z_N$ . And we present case (iii) and (iv) in our future work. We hope to extend this analysis Sommerfeld enhancement phenomena in a future work. It should be noted that these cross sections are also useful to study indirect detection of dark matter. With flavor becoming important in the future experiments, the large presence of flavor violation in the supersymmetric soft sector could possibly be probed by indirect experiments.

The neutralino pair annihilation and coannihilation cross section with neutralino in this paper is organized and presented as follows. In table.1, we present the various possible SM final states. For each set of final states, we present the channels and the corresponding mediators. In section.2, we sketch the calculation of thermal relic density. In section.3, we present cross sections corresponding to final states presented in table.1.

## 2 Thermal relic density

In this section, we outline the calculation of relic density and write it in a convenient form as it is done in [2, 8, 10].

Consider annihilation of  $N$  supersymmetric particles  $\chi_i$  ( $i=1,2,\dots,N-1,N$ ). And their mass and number density are given by  $m_i$  and  $n_i$  respectively. Normally, the decay rate of supersymmetric particles  $\chi_i$ , heavier than the LSP is much faster than the age of the Universe. In the case of R-parity conserved models, like the one we have considered here, these particles decay into the LSP. If LSP were the dark matter, its final abundance is described by the sum of the density of all supersymmetric particles:

$$n = \sum_i^N n_i \quad (2.1)$$

For  $n$ , Boltzmann evolution equation is given by [8]

$$\frac{dn}{dt} + 3Hn = - \sum_{i,j}^N \langle \sigma_{ij} v_{ij} \rangle (n_i n_j - n_i^{eq} n_j^{eq}) \quad (2.2)$$

where

$$\sigma_{ij} = \sigma(\chi_i \chi_j \rightarrow \text{All}) \quad (2.3)$$

SM particles which are lighter than supersymmetric particles act as background and they are relativistic at a given temperature while are non-relativistic. So supersymmetric particles which are non relativistic are suppressed by a Boltzmann factor. In this scenario, thermal distribution of density  $n_i$  remain in equilibrium. So the ratio  $\frac{n_i}{n}$  is equal to the equilibrium value:

$$\frac{n_i}{n} \sim \frac{n_i^{eq}}{n^{eq}} \quad (2.4)$$

then the eq.2.2 is simplified to

$$\frac{dn}{dt} + 3Hn = - \sum_{i,j}^N \langle \sigma_{eff} v \rangle (n^2 - n_{eq}^2) \quad (2.5)$$

where

$$\langle \sigma_{eff} v \rangle = \sum_{i,j}^N \langle \sigma_{ij} v_{ij} \rangle \frac{n_i^{eq}}{n^{eq}} \frac{n_j^{eq}}{n^{eq}} \quad (2.6)$$

The thermal average  $\langle \sigma_{ij} v_{ij} \rangle$  is defined with Maxwell-Boltzmann equilibrium distribution as

$$\langle \sigma_{ij} v_{ij} \rangle = \frac{\int d^3 p_i d^3 p_j \sigma v_{ij} e^{-E_i/T} e^{-E_j/T}}{\int d^3 p_i d^3 p_j e^{-E_i/T} e^{-E_j/T}} \quad (2.7)$$

where  $p_i = (E_i, \mathbf{p}_i)$  are the four momenta of the two annihilating particles and  $T$  is the temperature of the background bath. Eq.2.6 can be rewritten into a convenient one dimensional integral

$$\langle \sigma_{eff} v \rangle = \frac{\int_{4m_1^2}^{\infty} ds s^{3/2} K_1 \left( \frac{\sqrt{s}}{T} \right) \sum_{ij}^N \beta_f^2(s, m_i^2, m_j^2) \frac{g_i g_j}{g_1^2} \sigma_{ij}(s)}{8 m_1^4 T \left[ \sum_i^N \frac{g_i}{g_1} \frac{m_i^2}{m_1^2} K_2 \left( \frac{m_i}{T} \right) \right]^2} \quad (2.8)$$

where  $g_i$  gives number of degrees of freedom,  $K_i$  is modified Bessel function of order ‘i’ and  $\beta_f(s, m_i, m_j)$  is the kinetic function given by

$$\beta_f(s, m_i, m_j) = \frac{\sqrt{s - (m_i^2 + m_j^2)} \sqrt{s - (m_i^2 - m_j^2)}}{s} \quad (2.9)$$

### 3 Cross sections

In this section, general cross section for the cases (i) and (ii) mentioned in the introduction. To ease the process of comparison of our results with the ones already present in the literature[10], we recast  $\sigma_{ij}(s)$  as

$$\sigma_{ij}(s) = \frac{s}{16\pi \beta_f(s, m_i, m_j)} \sum_{\phi_1 \phi_2} [c \theta (s - (m_1^2 + m_2^2)^2) \beta_f(s, m_1, m_2) \tilde{w}_{\phi_1 \phi_2}(s)] \quad (3.1)$$

where  $m_1$  and  $m_2$  are the masses of the final states  $\phi_1$  and  $\phi_2$  respectively. And ‘c’ is the colour factor. If  $\chi_i^0 \chi_j^0 \rightarrow \phi_1 \phi_2$ , then the function  $\tilde{w}(s)_{\phi_1 \phi_2}$ , in CM frame, can be recast as

$$\tilde{w}(s)_{\phi_1 \phi_2} \equiv \frac{1}{2} \int_{-1}^1 d \cos \theta_{CM} |\mathcal{A}(\chi_i^0 \chi_j^0 \rightarrow \phi_1 \phi_2)|^2 \quad (3.2)$$

where  $\theta_{CM}$  denotes the scattering angle in the center of mass frame. We will follow Table.1 in presenting explicit expressions for  $\tilde{w}_{\phi_1 \phi_2}(s)$  for all the two-body final states.

#### 3.1 $\chi_i^0 \chi_j^0 \rightarrow H_m^0 H_n^0$

This process involves the s-channel CP even Higgs boson ( $H_1^0$  and  $H_2^0$ ) exchange and the t- and u-channel neutralino ( $\chi_k^0$ ,  $k = 1, \dots, 4$ ) exchange. The resulting cross section is sum of that of the s, t, u and possible interference processes among themselves.

$$\tilde{w}_{H_m^0, H_n^0} = \sum_{k, l=1, 2} \tilde{w}_{H_m^0, H_n^0}^{(H_k^0, H_l^0)} + \sum_{k=1, 2} \sum_{l=1, 4} \tilde{w}_{H_m^0, H_n^0}^{(H_k^0, \chi_l^0)} + \sum_{k, l=1, 4} \tilde{w}_{H_m^0, H_n^0}^{(\chi_k^0, \chi_l^0)} \quad (3.3)$$

- CP-even Higgs boson ( $H_k^0, H_l^0$ ) exchange:

$$\tilde{w}_{H_m^0, H_n^0}^{(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 H_m^0 H_n^0} C_{H_l^0 H_m^0 H_n^0}^*}{(s - m_{H_k^0}^2)(s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} (s - m_i^2 - m_j^2) - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) \quad (3.4)$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{H_m^0, H_n^0}^{t(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_m^0 H_n^0}^1(\chi_k^0 - \chi_l^0) + \beta_1(H_m^0, H_n^0, \chi_k^0, \chi_l^0) \Gamma_{H_m^0 H_n^0}^2(\chi_k^0 - \chi_l^0) \\
&\quad - 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_m^0 H_n^0}^3(\chi_k^0 - \chi_l^0) - m_i m_j \beta_4(H_m^0, H_n^0, \chi_k^0, \chi_l^0) \Gamma_{H_m^0 H_n^0}^4(\chi_k^0 - \chi_l^0) \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{H_m^0 H_n^0}^5(\chi_k^0 - \chi_l^0) + m_{\chi_l^0} \Gamma_{H_m^0 H_n^0}^6(\chi_k^0 - \chi_l^0)) \beta_3(H_m^0, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad + (4 d m_i - \frac{m_i \omega_{H_m^0 H_n^0}}{s}) (m_{\chi_k^0} \Gamma_{H_m^0 H_n^0}^7(\chi_k^0 - \chi_l^0) + m_{\chi_l^0} \Gamma_{H_m^0 H_n^0}^8(\chi_k^0 - \chi_l^0)) \quad (3.5)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_n^0, H_m^0}^{u(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_n^0 H_m^0}^1(\chi_k^0 - \chi_l^0) + \beta_1(H_n^0, H_m^0, \chi_k^0, \chi_l^0) \Gamma_{H_n^0 H_m^0}^2(\chi_k^0 - \chi_l^0) \\
&\quad - 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_n^0 H_m^0}^3(\chi_k^0 - \chi_l^0) - m_i m_j \beta_2(H_n^0, H_m^0, \chi_k^0, \chi_l^0) \Gamma_{H_n^0 H_m^0}^4(\chi_k^0 - \chi_l^0) \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{H_n^0 H_m^0}^5(\chi_k^0 - \chi_l^0) + m_{\chi_l^0} \Gamma_{H_n^0 H_m^0}^6(\chi_k^0 - \chi_l^0)) \beta_3(H_n^0, H_m^0, \chi_k^0, \chi_l^0) \\
&\quad + (4 d m_i - \frac{m_i \omega_{H_n^0 H_m^0}}{s}) (m_{\chi_k^0} \Gamma_{H_n^0 H_m^0}^7(\chi_k^0 - \chi_l^0) + m_{\chi_l^0} \Gamma_{H_n^0 H_m^0}^8(\chi_k^0 - \chi_l^0)) \quad (3.6)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_m^0, H_n^0}^{t-u(\chi_k^0, \chi_l^0)} &= d m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_m^0, H_n^0, \chi_k^0, \chi_l^0) I_{H_m^0 H_n^0}^1(\chi_k^0 - \chi_l^0) + J_2(H_m^0, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_m^0 H_n^0}^2(\chi_k^0 - \chi_l^0) - m_i m_j m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_m^0, H_n^0, \chi_k^0, \chi_l^0) I_{H_m^0 H_n^0}^3(\chi_k^0 - \chi_l^0) \\
&\quad + m_i m_j s^2 (l_i/2 - d_{H_m^0 H_n^0} - m_i^2) J_1(H_m^0, H_n^0, \chi_k^0, \chi_l^0) I_{H_m^0 H_n^0}^4(\chi_k^0 - \chi_l^0) \\
&\quad + \frac{1}{4} m_j m_{\chi_k^0} s J_3(H_m^0, H_n^0, \chi_k^0, \chi_l^0) I_{H_m^0 H_n^0}^5(\chi_k^0 - \chi_l^0) + \frac{1}{4} m_j m_{\chi_l^0} s \\
&\quad J_3(H_m^0, H_n^0, \chi_l^0, \chi_k^0) I_{H_m^0 H_n^0}^6(\chi_k^0 - \chi_l^0) - \frac{1}{4} m_i m_{\chi_k^0} s J_4(H_m^0, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_m^0 H_n^0}^7(\chi_k^0 - \chi_l^0) + \frac{1}{4} m_i m_{\chi_l^0} s J_5(H_m^0, H_n^0, \chi_k^0, \chi_l^0) I_{H_m^0 H_n^0}^8(\chi_k^0 - \chi_l^0) \quad (3.7)
\end{aligned}$$

$$\tilde{w}_{H_n^0, H_m^0}^{(\chi_k^0, \chi_l^0)} = \tilde{w}_{H_m^0, H_n^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{H_m^0, H_n^0}^{u(\chi_k^0, \chi_l^0)} + \left( \tilde{w}_{H_m^0, H_n^0}^{t-u(\chi_k^0, \chi_l^0)} + c.c. \right) \quad (3.8)$$



- CP-even Higgs boson ( $H_k^0$ ) and Neutralino ( $\chi_l^0$ ) Interference:

$$\begin{aligned}
\tilde{w}_{H_m^0, H_n^0}^{st(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, H_m^0, H_n^0) J_{H_m^0 H_n^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{H_m^0 H_n^0} - (R_{H_m^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, H_m^0, H_n^0) J_{H_m^0 H_n^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, H_m^0, H_n^0) J_{H_m^0 H_n^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{H_m^0 H_n^0} - (R_{H_m^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{H_m^0 H_n^0} - \omega_{H_m^0 H_n^0}) \mathcal{A}(\chi_l^0, H_m^0, H_n^0) \right) J_{H_m^0 H_n^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.9)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_m^0, H_n^0}^{su(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, H_n^0, H_m^0) J_{H_n^0 H_m^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{H_m^0 H_n^0} - (R_{H_n^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, H_n^0, H_m^0) J_{H_n^0 H_m^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, H_n^0, H_m^0) J_{H_n^0 H_m^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{H_m^0 H_n^0} - (R_{H_n^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{H_m^0 H_n^0} - \omega_{H_m^0 H_n^0}) \mathcal{A}(\chi_l^0, H_n^0, H_m^0) \right) J_{A_1^0 H_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.10)
\end{aligned}$$

$$\tilde{w}_{H_m^0, H_n^0}^{(H_k^0, \chi_l^0)} = \frac{1}{b_{H_m^0 H_n^0} (s - m_{H_k^0}^2)} \left( \tilde{w}_{H_m^0, H_n^0}^{st(H_k^0, \chi_l^0)} + \tilde{w}_{H_m^0, H_n^0}^{su(H_k^0, \chi_l^0)} \right) + c.c. \quad (3.11)$$

The functions  $\Gamma$ 's,  $J$ 's,  $I$ 's,  $f$ 's and  $g$ 's, used in the above expressions depend on mixing matrices and the couplings in the MSSM and are presented in the Appendix.B. General expression, for the CP-even final state, presented above can be used to obtain processes with all possible CP-even Higgs final states by taking  $m, n=1, 2$ . Please note that in case of identical final states ( $H_1^0 H_1^0$  and  $H_2^0 H_2^0$ )  $\tilde{w}$  has to be multiplied by a factor  $\frac{1}{2}$

### 3.2 $\chi_i^0 \chi_j^0 \rightarrow A_1^0 A_1^0$

This process is the sum of s-channel diagrams though CP even Higgs boson ( $H_1^0$  and  $H_2^0$ ) and the t- and u-channel neutralino ( $\chi_k^0$ ,  $k = 1, \dots, 4$ ) exchange (Similar to the  $\chi_i^0 \chi_j^0 \rightarrow H_m^0 H_n^0$ )

$$\tilde{w}_{A_1^0, A_1^0} = \sum_{k, l=1, 2} \tilde{w}_{A_1^0, A_1^0}^{(H_k^0, H_l^0)} + \sum_{k=1, 2} \sum_{l=1, 4} \tilde{w}_{A_1^0, A_1^0}^{(H_k^0, \chi_l^0)} + \sum_{k, l=1, 4} \tilde{w}_{A_1^0, A_1^0}^{(\chi_k^0, \chi_l^0)} \quad (3.12)$$

- CP-odd Higgs boson ( $A_1^0 - A_1^0$ ) exchange:

$$\tilde{w}_{A_1^0, A_1^0}^{(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 A_1^0 A_1^0} C_{H_l^0 A_1^0 A_1^0}^*}{(s - m_{H_k^0}^2) (s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} (s - m_i^2 - m_j^2) - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) \quad (3.13)$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{A_1^0, A_1^0}^{t(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} \\
&- 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&- m_j (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&+ (4 d m_i - \frac{m_i \omega_{A_1^0 A_1^0}}{s}) (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)}) \quad (3.14)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{A_1^0, A_1^0}^{u(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} \\
&- 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&- m_j (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&+ (4 d m_i - \frac{m_i \omega_{A_1^0 A_1^0}}{s}) (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)}) \quad (3.15)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{A_1^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} &= d m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(A_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{A_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + J_2(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&I_{A_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} - m_i m_j m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(A_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{A_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} \\
&+ m_i m_j s^2 (l_i/2 - d_{A_1^0 A_1^0} - m_i^2) J_1(A_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{A_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&+ \frac{1}{4} m_j m_{\chi_k^0} s J_3(A_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{A_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_j m_{\chi_l^0} s \\
&J_3(A_1^0, A_1^0, \chi_l^0, \chi_k^0) I_{A_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)} - \frac{1}{4} m_i m_{\chi_k^0} s J_4(A_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&I_{A_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_i m_{\chi_l^0} s J_5(A_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{A_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)} \quad (3.16)
\end{aligned}$$

$$\tilde{w}_{A_1^0, A_1^0}^{(\chi_k^0, \chi_l^0)} = \tilde{w}_{A_1^0, A_1^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{A_1^0, A_1^0}^{u(\chi_k^0, \chi_l^0)} + \left( \tilde{w}_{A_1^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} + c.c. \right) \quad (3.17)$$

- CP-odd Higgs boson ( $A_1^0$ ) and Neutralino ( $\chi_l^0$ ) Interference:

$$\begin{aligned}
\tilde{w}_{A_1^0, A_1^0}^{st(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&- \frac{m_j}{4} \left( b_{A_1^0 A_1^0} - (R_{H_m^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&+ \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{A_1^0 A_1^0} - (R_{A_1^0 \chi_l^0} \right. \\
&\left. + 4 d s - \tau_{A_1^0 A_1^0} - \omega_{A_1^0 A_1^0}) \mathcal{A}(\chi_l^0, A_1^0, A_1^0) \right) J_{A_1^0 A_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.18)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{A_1^0 A_1^0}^{su(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{A_1^0 A_1^0} - (R_{A_1^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, A_1^0, A_1^0) J_{A_1^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{A_1^0 A_1^0} - (R_{A_1^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{A_1^0 A_1^0} - \omega_{A_1^0 A_1^0}) \mathcal{A}(\chi_l^0, A_1^0, A_1^0) \right) J_{A_1^0 A_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.19)
\end{aligned}$$

$$\tilde{w}_{A_1^0 A_1^0}^{(H_k^0, \chi_l^0)} = \frac{1}{b_{A_1^0 A_1^0} (s - m_{H_k^0}^2)} \left( \tilde{w}_{A_1^0 A_1^0}^{st(H_k^0, \chi_l^0)} + \tilde{w}_{A_1^0 A_1^0}^{su(H_k^0, \chi_l^0)} \right) + c.c. \quad (3.20)$$

Note: As there are identical final states total cross section has to be multiplied by a factor  $\frac{1}{2}$

### 3.3 $\chi_i^0 \chi_j^0 \rightarrow H_1^0 A_1^0$

This process is sum of s-channel diagrams through CP-odd Higgs boson  $A_1^0$  and Z boson and t and u channel neutralino exchange. The resulting cross section is given as

$$\tilde{w}_{H_1^0 A_1^0} = \tilde{w}_{H_1^0 A_1^0}^{(A_1^0, A_1^0)} + \tilde{w}_{H_1^0 A_1^0}^{(A_1^0, A_1^0)} + \tilde{w}_{H_1^0 A_1^0}^{(A_1^0, Z)} + \sum_{k,l=1,4} \tilde{w}_{H_1^0 A_1^0}^{(\chi_k^0, \chi_l^0)} + \sum_{l=1,4} \tilde{w}_{H_1^0 A_1^0}^{(A_1^0, \chi_l^0)} + \sum_{l=1,4} \tilde{w}_{H_1^0 A_1^0}^{(Z, \chi_l^0)} \quad (3.21)$$

- CP-odd Higgs boson  $A_1^0$  and Z exchange:

$$\begin{aligned}
\tilde{w}_{H_1^0 A_1^0}^{(A_1^0, A_1^0)} &= \frac{1}{2} \frac{|C_{A_1^0 H_1^0 A_1^0}|^2}{(s - m_{A_1^0}^2)^2} \left( f_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} (s - m_i^2 - m_j^2) - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} \right) \quad (3.22) \\
\tilde{w}_{H_1^0 A_1^0}^{(Z, Z)} &= \frac{1}{2} \frac{|C_{Z H_1^0 A_1^0}|^2}{(s - M_Z^2)^2 M_Z^4} \left( f_{\chi_i^0 \chi_j^0}^{(Z - Z)} \mathcal{B}_1(Z, H_1^0, A_1^0) - g_{\chi_i^0 \chi_j^0}^{(Z - Z)} \mathcal{B}_2(Z, H_1^0, A_1^0) \right) \\
\tilde{w}_{H_1^0 A_1^0}^{(A_1^0, Z)} &= \frac{1}{2} \frac{C_{A_1^0 H_1^0 A_1^0} C_{Z H_1^0 A_1^0}^*}{(s - m_{A_1^0}^2)^2 M_Z^2 s} \left( f_{\chi_i^0 \chi_j^0}^{(A_1^0 - Z)} m_j \tau_{H_1^0 A_1^0} - g_{\chi_i^0 \chi_j^0}^{(Z - Z)} m_i \tau_{A_1^0 H_1^0} \right) \\
&\quad (m_{H_1^0}^2 - m_{A_1^0}^2) + c.c. \quad (3.23)
\end{aligned}$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{H_1^0 A_1^0}^{t(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1(H_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{H_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} \\
&\quad - 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2(H_1^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{H_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{H_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{H_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3(H_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&\quad + \left( 4 d m_i - \frac{m_i \omega_{H_1^0 A_1^0}}{s} \right) \left( m_{\chi_k^0} \Gamma_{H_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{H_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)} \right) \quad (3.24)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{A_1^0, H_1^0}^{u(\chi_k^0, \chi_l^0)} &= 4d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1(A_1^0, H_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} \\
&\quad - 4m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2(A_1^0, H_1^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3(A_1^0, H_1^0, \chi_k^0, \chi_l^0) \\
&\quad + (4d m_i - \frac{m_i \omega_{A_1^0 A_1^0}}{s}) (m_{\chi_k^0} \Gamma_{A_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)}) \quad (3.25)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_1^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} &= d m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_1^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + J_2(H_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_1^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} - m_i m_j m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_1^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} \\
&\quad + m_i m_j s^2 (l_i/2 - d_{H_1^0 A_1^0} - m_i^2) J_1(H_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_1^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{4} m_j m_{\chi_k^0} s J_3(H_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_1^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_j m_{\chi_l^0} s \\
&\quad J_3(H_1^0, A_1^0, \chi_l^0, \chi_k^0) I_{H_1^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)} - \frac{1}{4} m_i m_{\chi_k^0} s J_4(H_1^0, A_1^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_1^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_i m_{\chi_l^0} s J_5(H_1^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_1^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)} \quad (3.26)
\end{aligned}$$

$$\tilde{w}_{H_1^0, A_1^0}^{(\chi_k^0, \chi_l^0)} = \tilde{w}_{H_1^0, A_1^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{H_1^0, A_1^0}^{u(\chi_k^0, \chi_l^0)} + (\tilde{w}_{H_1^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} + c.c.) \quad (3.27)$$

- CP-odd Higgs boson  $A_1^0$ , Z boson and Neutralino ( $\chi_l^0$ ) Interference:

$$\begin{aligned}
\tilde{w}_{H_1^0, A_1^0}^{st(Z, \chi_k^0)} &= \frac{C_{ZH_1^0 A_1^0}}{s - M_Z^2} \left( \frac{m_j m_{\chi_k^0}}{2b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_3(Z, \chi_k^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(5)(Z-\chi_k^0)} \right. \\
&\quad + \frac{m_j m_{\chi_k^0}}{4b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_4(Z, \chi_k^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(6)(Z-\chi_k^0)} + \mathcal{B}_5(Z, \chi_k^0, H_1^0, A_1^0) \\
&\quad \left. J_{H_1^0 A_1^0}^{(7)(Z-\chi_k^0)} + \frac{m_i m_j}{2b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_6(Z, \chi_k^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(8)(Z-\chi_k^0)} \right) \quad (3.28)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_1^0, A_1^0}^{su(Z, \chi_k^0)} &= \frac{C_{ZH_1^0 A_1^0}}{s - M_Z^2} \left( -\frac{m_j m_{\chi_k^0}}{2b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_3(Z, \chi_k^0, A_1^0, H_1^0) J_{A_1^0 H_1^0}^{(5)(Z-\chi_k^0)} \right. \\
&\quad - \frac{m_j m_{\chi_k^0}}{4b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_4(Z, \chi_k^0, A_1^0, H_1^0) J_{A_1^0 H_1^0}^{(6)(Z-\chi_k^0)} - \mathcal{B}_5(Z, \chi_k^0, A_1^0, H_1^0) \\
&\quad \left. J_{A_1^0 H_1^0}^{(7)(Z-\chi_k^0)} + \frac{m_i m_j}{2b_{H_1^0 A_1^0} M_Z^2} \mathcal{B}_6(Z, \chi_k^0, A_1^0, H_1^0) J_{A_1^0 H_1^0}^{(8)(Z-\chi_k^0)} \right) \quad (3.29)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_1^0, A_1^0}^{st(A_1^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} (b_{H_1^0 A_1^0} - (R_{H_1^0 \chi_l^0} - 4m_i^2 s)) \mathcal{A}(\chi_l^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, H_1^0, A_1^0) J_{H_1^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} (b_{H_1^0 A_1^0} - (R_{H_1^0 \chi_l^0} \\
&\quad + 4ds - \tau_{H_1^0 A_1^0} - \omega_{H_1^0 A_1^0}) \mathcal{A}(\chi_l^0, H_1^0, A_1^0)) J_{H_1^0 A_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.30)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_1^0 A_1^0}^{su(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, A_1^0, H_1^0) J_{A_1^0 H_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{H_1^0 A_1^0} - (R_{A_1^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, A_1^0, H_1^0) J_{A_1^0 H_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, A_1^0, H_1^0) J_{H_1^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{H_1^0 A_1^0} - (R_{A_1^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{H_1^0 A_1^0} - \omega_{H_1^0 A_1^0}) \mathcal{A}(\chi_l^0, A_1^0, H_1^0) \right) J_{A_1^0 H_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.31)
\end{aligned}$$

$$\tilde{w}_{H_1^0 A_1^0}^{(A_1^0, \chi_l^0)} = \frac{1}{b_{H_1^0 A_1^0} (s - m_{A_1^0}^2)} \left( \tilde{w}_{H_1^0 A_1^0}^{st(A_1^0, \chi_l^0)} + \tilde{w}_{H_1^0 A_1^0}^{su(A_1^0, \chi_l^0)} \right) + c.c. \quad (3.32)$$

$$\tilde{w}_{H_1^0 A_1^0}^{(Z, \chi_l^0)} = \frac{1}{b_{H_1^0 A_1^0} (s - M_Z^2)} \left( \tilde{w}_{H_1^0 A_1^0}^{st(Z, \chi_l^0)} + \tilde{w}_{H_1^0 A_1^0}^{su(Z, \chi_l^0)} \right) + c.c. \quad (3.33)$$

### 3.4 $\chi_i^0 \chi_j^0 \rightarrow H_2^0 A_1^0$

This process is the sum of CP even Higgs boson  $H_k^0$  in s-channel and neutralino exchange in t and u channel. Resulting crosssection is given by:

$$\tilde{w}_{H_2^0 A_1^0} = \sum_{k,l=1,2} \tilde{w}_{H_2^0 A_1^0}^{(H_k^0, H_l^0)} + \sum_{k,l=1,4} \tilde{w}_{H_2^0 A_1^0}^{(\chi_k^0, \chi_l^0)} + \sum_{k=1,2} \sum_{l=1,4} \tilde{w}_{H_2^0 A_1^0}^{(H_k^0, \chi_l^0)} \quad (3.34)$$

- CP-even Higgs boson ( $H_k^0, H_l^0$ ) exchange:

$$\tilde{w}_{H_2^0 A_1^0}^{(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 H_2^0 A_1^0} C_{H_l^0 H_2^0 A_1^0}^*}{(s - m_{H_2^0}^2)(s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} (s - m_i^2 - m_j^2) - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) \quad (3.35)$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{H_2^0 A_1^0}^{t(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_2^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1 (H_2^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{H_2^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} \\
&\quad - 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{H_2^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2 (H_2^0, A_1^0, \chi_k^0, \chi_l^0) \Gamma_{H_2^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{H_2^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{H_2^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3 (H_2^0 A_1^0, \chi_k^0, \chi_l^0) \\
&\quad + \left( 4 d m_i - \frac{m_i \omega_{H_2^0 A_1^0}}{s} \right) (m_{\chi_k^0} \Gamma_{H_2^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{H_2^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)}) \quad (3.36)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{A_1^0 H_2^0}^{u(\chi_k^0, \chi_l^0)} &= 4 d m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 H_2^0}^{1(\chi_k^0 - \chi_l^0)} + \beta_1 (A_1^0, H_2^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 H_2^0}^{2(\chi_k^0 - \chi_l^0)} \\
&\quad - 4 m_i m_j m_{\chi_k^0} m_{\chi_l^0} \Gamma_{A_1^0 H_2^0}^{3(\chi_k^0 - \chi_l^0)} - m_i m_j \beta_2 (A_1^0, H_2^0, \chi_k^0, \chi_l^0) \Gamma_{A_1^0 H_2^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad - m_j (m_{\chi_k^0} \Gamma_{A_1^0 H_2^0}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 H_2^0}^{6(\chi_k^0 - \chi_l^0)}) \beta_3 (A_1^0, H_2^0, \chi_k^0, \chi_l^0) \\
&\quad + \left( 4 d m_i - \frac{m_i \omega_{A_1^0 H_2^0}}{s} \right) (m_{\chi_k^0} \Gamma_{A_1^0 H_2^0}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_l^0} \Gamma_{A_1^0 H_2^0}^{8(\chi_k^0 - \chi_l^0)}) \quad (3.37)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_2^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} &= d m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_2^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_2^0 A_1^0}^{1(\chi_k^0 - \chi_l^0)} + J_2(H_2^0, A_1^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_2^0 A_1^0}^{2(\chi_k^0 - \chi_l^0)} - m_i m_j m_{\chi_k^0} m_{\chi_l^0} s^2 J_1(H_2^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_2^0 A_1^0}^{3(\chi_k^0 - \chi_l^0)} \\
&\quad + m_i m_j s^2 (l_i/2 - d_{H_2^0 A_1^0} - m_i^2) J_1(H_2^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_2^0 A_1^0}^{4(\chi_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{4} m_j m_{\chi_k^0} s J_3(H_2^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_2^0 A_1^0}^{5(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_j m_{\chi_l^0} s \\
&\quad J_3(H_2^0, A_1^0, \chi_l^0, \chi_k^0) I_{H_2^0 A_1^0}^{6(\chi_k^0 - \chi_l^0)} - \frac{1}{4} m_i m_{\chi_k^0} s J_4(H_2^0, A_1^0, \chi_k^0, \chi_l^0) \\
&\quad I_{H_2^0 A_1^0}^{7(\chi_k^0 - \chi_l^0)} + \frac{1}{4} m_i m_{\chi_l^0} s J_5(H_2^0, A_1^0, \chi_k^0, \chi_l^0) I_{H_2^0 A_1^0}^{8(\chi_k^0 - \chi_l^0)} \quad (3.38)
\end{aligned}$$

$$\tilde{w}_{H_2^0, A_1^0}(\chi_k^0, \chi_l^0) = \tilde{w}_{H_2^0, A_1^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{H_2^0, A_1^0}^{u(\chi_k^0, \chi_l^0)} + \left( \tilde{w}_{H_2^0, A_1^0}^{t-u(\chi_k^0, \chi_l^0)} + c.c. \right) \quad (3.39)$$

- CP-even Higgs boson ( $H_k^0$ ) and Neutralino ( $\chi_l^0$ ) Interference:

$$\begin{aligned}
\tilde{w}_{H_2^0, A_1^0}^{st(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, H_2^0, A_1^0) J_{H_2^0 A_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{H_2^0 A_1^0} - (R_{H_2^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, H_2^0, A_1^0) J_{H_2^0 A_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, H_2^0, A_1^0) J_{H_2^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{H_2^0 A_1^0} - (R_{H_2^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{H_2^0 A_1^0} - \omega_{H_2^0 A_1^0}) \mathcal{A}(\chi_l^0, H_2^0, A_1^0) \right) J_{H_2^0 A_1^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.40)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{H_2^0, A_1^0}^{su(H_k^0, \chi_l^0)} &= -\frac{1}{2} m_i m_j m_{\chi_l^0} s L(\chi_l^0, A_1^0, H_2^0) J_{A_1^0 A_1^0}^{(1)(H_k^0 - \chi_l^0)} \\
&\quad - \frac{m_j}{4} \left( b_{H_2^0 A_1^0} - (R_{H_2^0 \chi_l^0} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^0, A_1^0, H_2^0) J_{A_1^0 A_1^0}^{(2)(H_k^0 - \chi_l^0)} \\
&\quad + \frac{1}{2} d m_{\chi_l^0} s L(\chi_l^0, A_1^0, H_2^0) J_{A_1^0 A_1^0}^{(3)(H_k^0 - \chi_l^0)} - \frac{m_i}{4} \left( b_{H_2^0 A_1^0} - (R_{H_2^0 \chi_l^0} \right. \\
&\quad \left. + 4 d s - \tau_{H_2^0 A_1^0} - \omega_{H_2^0 A_1^0}) \mathcal{A}(\chi_l^0, A_1^0, H_2^0) \right) J_{A_1^0 H_2^0}^{(4)(H_k^0 - \chi_l^0)} \quad (3.41)
\end{aligned}$$

$$\tilde{w}_{H_2^0, A_1^0}^{(H_k^0, \chi_l^0)} = \frac{1}{b_{H_2^0 A_1^0} (s - m_{H_k^0}^2)} \left( \tilde{w}_{H_2^0, A_1^0}^{st(H_k^0, \chi_l^0)} + \tilde{w}_{H_2^0, A_1^0}^{su(H_k^0, \chi_l^0)} \right) + c.c. \quad (3.42)$$

### 3.5 $\chi_i^0 \chi_j^0 \rightarrow H^+ H^-$

This process is the sum of the s-channel Z and CP even Higgs boson ( $H_1^0$ ,  $H_2^0$ ) exchange and t- and u-channel chargino ( $\chi_k^\pm$ ,  $k = 1, 2$ ) exchange. Resulting cross section including considering interference via s, t and u channel diagrams is given by:

$$\tilde{w}_{H_1^+, H_1^-} = \sum_{k,l=1,2} \tilde{w}_{H_1^+, H_1^-}^{(H_k^0, H_l^0)} + \tilde{w}_{H_1^+, H_1^-}^{(Z, Z)} + \sum_{k=1,2} \sum_{l=1,2} \tilde{w}_{H_1^+, H_1^-}^{(H_k^0, \chi_l^\pm)} + \sum_{k,l=1,2} \tilde{w}_{H_1^+, H_1^-}^{(\chi_k^\pm, \chi_l^\pm)} \quad (3.43)$$

- CP even Higgs boson ( $H_k^0, H_l^0$ ) and Z exchange:

$$\tilde{w}_{H_1^+, H_1^-}^{s(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 H_1^+ H_1^-} C_{H_l^0 H_1^+ H_1^-}^*}{(s - m_{H_k^0}^2)(s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} (s - m_i^2 - m_j^2) - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) \quad (3.44)$$

$$\tilde{w}_{H_1^+, H_1^-}^{s(Z, Z)} = \frac{1}{2} \frac{|C_{ZH_1^+ H_1^-}|^2}{(s - M_Z^2)^2 M_Z^4} \left( f_{\chi_i^0 \chi_j^0}^{(Z-Z)} \mathcal{B}_1(Z, H_1^+ H_1^-) - g_{\chi_i^0 \chi_j^0}^{(Z-Z)} \mathcal{B}_2(Z, H_1^+ H_1^-) \right) \quad (3.45)$$

$$\tilde{w}_{H_1^+, H_1^-}^{s(H_k^0, Z)} = 0 \quad (3.46)$$

- Chargino ( $\chi_k^\pm, \chi_l^\pm$ ) exchange:

$$\begin{aligned} \tilde{w}_{H_1^+, H_1^-}^{t(\chi_k^\pm, \chi_l^\pm)} &= 4d m_{\chi_k^\pm} m_{\chi_l^\pm} \Gamma_{H_1^+ H_1^-}^{1(\chi_k^\pm - \chi_l^\pm)} + \beta_1(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \Gamma_{H_1^+ H_1^-}^{2(\chi_k^\pm - \chi_l^\pm)} \\ &\quad - 4m_i m_j m_{\chi_k^\pm} m_{\chi_l^\pm} \Gamma_{H_1^+ H_1^-}^{3(\chi_k^\pm - \chi_l^\pm)} - m_i m_j \beta_2(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \Gamma_{H_1^+ H_1^-}^{4(\chi_k^\pm - \chi_l^\pm)} \\ &\quad - m_j \left( m_{\chi_k^\pm} \Gamma_{H_1^+ H_1^-}^{5(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_l^\pm} \Gamma_{H_1^+ H_1^-}^{6(\chi_k^\pm - \chi_l^\pm)} \right) \beta_3(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\ &\quad + \left( 4d m_i - \frac{m_i \omega_{H_1^+ H_1^-}}{s} \right) \left( m_{\chi_k^\pm} \Gamma_{H_1^+ H_1^-}^{7(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_l^\pm} \Gamma_{H_1^+ H_1^-}^{8(\chi_k^\pm - \chi_l^\pm)} \right) \end{aligned} \quad (3.47)$$

$$\begin{aligned} \tilde{w}_{H_1^+, H_1^-}^{u(\chi_k^\pm, \chi_l^\pm)} &= 4d m_{\chi_k^\pm} m_{\chi_l^\pm} \Gamma_{H_1^- H_1^+}^{1(\chi_k^\pm - \chi_l^\pm)*} + \beta_1(H_1^-, H_1^+, \chi_k^\pm, \chi_l^\pm) \Gamma_{H_1^- H_1^+}^{2(\chi_k^\pm - \chi_l^\pm)*} \\ &\quad - 4m_i m_j m_{\chi_k^\pm} m_{\chi_l^\pm} \Gamma_{H_1^- H_1^+}^{3(\chi_k^\pm - \chi_l^\pm)*} - m_i m_j \beta_2(H_1^-, H_1^+, \chi_k^\pm, \chi_l^\pm) \Gamma_{H_1^- H_1^+}^{4(\chi_k^\pm - \chi_l^\pm)*} \\ &\quad - m_j \left( m_{\chi_k^\pm} \Gamma_{H_1^- H_1^+}^{5(\chi_k^\pm - \chi_l^\pm)*} + m_{\chi_l^\pm} \Gamma_{H_1^- H_1^+}^{6(\chi_k^\pm - \chi_l^\pm)*} \right) \beta_3(H_1^-, H_1^+, \chi_k^\pm, \chi_l^\pm) \\ &\quad + \left( 4d m_i - \frac{m_i \omega_{H_1^- H_1^+}}{s} \right) \left( m_{\chi_k^\pm} \Gamma_{H_1^- H_1^+}^{7(\chi_k^\pm - \chi_l^\pm)*} + m_{\chi_l^\pm} \Gamma_{H_1^- H_1^+}^{8(\chi_k^\pm - \chi_l^\pm)*} \right) \end{aligned} \quad (3.48)$$

$$\begin{aligned} \tilde{w}_{H_1^+, H_1^-}^{t-u(\chi_k^\pm, \chi_l^\pm)} &= d m_{\chi_k^\pm} m_{\chi_l^\pm} s^2 J_1(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) I_{H_1^+, H_1^-}^{1(\chi_k^\pm - \chi_l^\pm)} + J_2(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\ &\quad I_{H_m^0 H_n^0}^{2(\chi_k^\pm - \chi_l^\pm)} - m_i m_j m_{\chi_k^\pm} m_{\chi_l^\pm} s^2 J_1(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) I_{H_1^+, H_1^-}^{3(\chi_k^\pm - \chi_l^\pm)} \\ &\quad + m_i m_j s^2 (l_i/2 - d_{H_1^+, H_1^-} - m_i^2) J_1(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) I_{H_1^+, H_1^-}^{4(\chi_k^\pm - \chi_l^\pm)} \\ &\quad + \frac{1}{4} m_j m_{\chi_k^\pm} s J_3(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) I_{H_1^+ H_1^-}^{5(\chi_k^\pm - \chi_l^\pm)} + \frac{1}{4} m_j m_{\chi_l^\pm} s \\ &\quad J_3(H_1^+, H_1^-, \chi_l^\pm, \chi_k^\pm) I_{H_1^+ H_1^-}^{6(\chi_k^\pm - \chi_l^\pm)} - \frac{1}{4} m_i m_{\chi_k^\pm} s J_4(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\ &\quad I_{H_1^+ H_1^-}^{7(\chi_k^\pm - \chi_l^\pm)} + \frac{1}{4} m_i m_{\chi_l^\pm} s J_5(H_1^+, H_1^-, \chi_k^\pm, \chi_l^\pm) I_{H_1^+ H_1^-}^{8(\chi_k^\pm - \chi_l^\pm)} \end{aligned} \quad (3.49)$$

$$\tilde{w}_{H_1^+, H_1^-}^{(\chi_k^\pm, \chi_l^\pm)} = \tilde{w}_{H_1^+, H_1^-}^{t(\chi_k^\pm, \chi_l^\pm)} + \tilde{w}_{H_1^+, H_1^-}^{u(\chi_k^\pm, \chi_l^\pm)} + \left( \tilde{w}_{H_1^+, H_1^-}^{t-u(\chi_k^\pm, \chi_l^\pm)} + c.c. \right) \quad (3.50)$$

- CP-even Higgs boson ( $H_k^0$ ) and Neutralino ( $\chi_l^\pm$ ) Interference:

$$\begin{aligned} \tilde{w}_{H_1^+, H_1^-}^{st(H_k^0, \chi_l^\pm)} &= -\frac{1}{2} m_i m_j m_{\chi_l^\pm} s L(\chi_l^\pm, H_1^+, H_1^-) J_{H_1^+ H_1^-}^{(1)(H_k^0 - \chi_l^\pm)} \\ &\quad - \frac{m_j}{4} \left( b_{H_1^+ H_1^-} - (R_{H_1^+ \chi_l^\pm} - 4m_i^2 s) \right) \mathcal{A}(\chi_l^\pm, H_1^+, H_1^-) J_{H_1^+ H_1^-}^{(2)(H_k^0 - \chi_l^0)} \\ &\quad + \frac{1}{2} d m_{\chi_l^\pm} s L(\chi_l^\pm, H_1^+, H_1^-) J_{H_1^+ H_1^-}^{(3)(H_k^0 - \chi_l^\pm)} - \frac{m_i}{4} \left( b_{H_1^+ H_1^-} - (R_{H_1^+ \chi_l^\pm} \right. \\ &\quad \left. + 4d s - \tau_{H_1^+ H_1^-} - \omega_{H_1^+ H_1^-}) \mathcal{A}(\chi_l^\pm, H_1^+, H_1^-) \right) J_{H_1^+ H_1^-}^{(4)(H_k^0 - \chi_l^\pm)} \end{aligned} \quad (3.51)$$

$$\begin{aligned}
\tilde{w}_{H_1^+, H_1^-}^{su(H_k^0, \chi_l^\pm)} &= -\frac{1}{2} m_i m_j m_{\chi_l^\pm} s L(\chi_l^\pm, H_1^-, H_1^+) J_{H_1^- H_1^+}^{(1)(H_k^0 - \chi_l^\pm)} \\
&\quad - \frac{m_j}{4} \left( b_{H_1^+ H_1^-} - (R_{H_1^- \chi_l^\pm} - 4 m_i^2 s) \right) \mathcal{A}(\chi_l^\pm, H_1^-, H_1^+) J_{H_1^- H_1^+}^{(2)(H_k^0 - \chi_l^\pm)} \\
&\quad + \frac{1}{2} d m_{\chi_l^\pm} s L(\chi_l^\pm, H_1^-, H_1^+) J_{H_1^+ H_1^-}^{(3)(H_k^0 - \chi_l^\pm)} - \frac{m_i}{4} \left( b_{H_1^+ H_1^-} - (R_{H_1^- \chi_l^\pm} \right. \\
&\quad \left. + 4 d s - \tau_{H_1^+ H_1^-} - \omega_{H_1^+ H_1^-}) \mathcal{A}(\chi_l^\pm, H_1^-, H_1^+) \right) J_{A_1^0 H_1^0}^{(4)(H_k^0 - \chi_l^\pm)} \quad (3.52)
\end{aligned}$$

$$\tilde{w}_{H_1^+, H_1^-}^{(H_k^0, \chi_l^\pm)} = \frac{1}{b_{H_1^+ H_1^-} (s - m_{H_k^0}^2)} \left( \tilde{w}_{H_1^+, H_1^-}^{st(H_k^0, \chi_l^\pm)} + \tilde{w}_{H_1^+, H_1^-}^{su(H_k^0, \chi_l^\pm)} \right) + c.c. \quad (3.53)$$

### 3.6 $\chi_i^0 \chi_j^0 \rightarrow W^\pm H^\mp$

This process is the sum of the s-channel CP even ( $H_1^0, H_2^0$ ), CP odd ( $A_1^0$ ) Higgs boson exchange and the t- and u-channel chargino ( $\chi_k^\pm, k = 1, 2$ ) exchange

$$\tilde{w}_{W^+ H_1^-} = \tilde{w}_{W^+ H_1^-}^{(A_1^0, A_1^0)} + \sum_{k,l=1,2} \tilde{w}_{W^+ H_1^-}^{(H_k^0, H_l^0)} + \sum_{k=1,2} \tilde{w}_{W^+ H_1^-}^{(H_k^0, A_1^0)} + \sum_{k,l=1,2} \left( \tilde{w}_{W^+ H_1^-}^{(\chi_k^\pm, \chi_l^\pm)} + \tilde{w}_{W^+ H_1^-}^{(H_k^0, \chi_l^\pm)} \right) \quad (3.54)$$

- CP-even Higgs boson ( $H_k^0 - H_l^0$ ) and CP-odd Higgs boson ( $A_1^0$ ):

$$\tilde{w}_{W^+ H_1^-}^{(A_1^0, A_1^0)} = \frac{1}{2} \frac{|C_{A_1^0 W^+ H_1^-}|^2}{(s - m_{A_1^0}^2)^2} \left( f_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} d - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} \right) (s - l_{W^+ H^-}) \quad (3.55)$$

$$\tilde{w}_{W^+ H_1^-}^{(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 W^+ H_1^-} C_{H_l^0 W^+ H_1^-}^*}{(s - m_{H_k^0}^2) (s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} d - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) (s - l_{W^+ H^-}) \quad (3.56)$$

$$\tilde{w}_{W^+ H_1^-}^{(H_k^0, A_1^0)} = \frac{1}{2} \frac{C_{H_k^0 W^+ H_1^-} C_{A_1^0 W^+ H_1^-}^*}{(s - m_{H_k^0}^2) (s - m_{A_1^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - A_1^0)} d - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - A_1^0)} \right) (s - l_{W^+ H^-}) \quad (3.57)$$

- Chargino ( $\chi_k^\pm - \chi_l^\pm$ ) exchange:

$$\begin{aligned}
\tilde{w}_{W^+ H_1^-}^{t(\chi_k^\pm, \chi_l^\pm)} &= \Gamma_{W^+ H_1^-}^{1(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_1(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) + \Gamma_{W^+ H_1^-}^{2(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_2(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) \\
&\quad - 3 m_i m_j s^2 \left( M_w^2 \Gamma_{W^+ H_1^-}^{3(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_k^\pm} m_{\chi_l^\pm} \Gamma_{W^+ H_1^-}^{4(\chi_k^\pm - \chi_l^\pm)} \right) \\
&\quad \mathcal{C}_3(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) - \left( \Gamma_{W^+ H_1^-}^{5(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} + \Gamma_{W^+ H_1^-}^{6(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} \right) \\
&\quad \mathcal{C}_4(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) + \left( \Gamma_{W^+ H_1^-}^{7(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} + \Gamma_{W^+ H_1^-}^{8(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} \right) \\
&\quad \mathcal{C}_5(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) + \left( \Gamma_{W^+ H_1^-}^{5(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} + \Gamma_{W^+ H_1^-}^{6(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} \right. \\
&\quad \left. - 2 \Gamma_{W^+ H_1^-}^{3(\chi_k^\pm - \chi_l^\pm)} m_i \right) \mathcal{C}_6(\chi_k^\pm, \chi_l^\pm, W^+, H_1^-) \quad (3.58)
\end{aligned}$$



$$\begin{aligned}
\tilde{w}_{W^+H_1^-}^{u(\chi_k^\pm, \chi_l^\pm)} &= \left( \Gamma_{H_1^-W^+}^{1(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} + \Gamma_{H_1^-W^+}^{2(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} \right) \mathcal{C}_{11}(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) \\
&+ \left( \Gamma_{H_1^-W^+}^{3(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} + \Gamma_{H_1^-W^+}^{4(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} \right) \mathcal{C}_{10}(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) \\
&+ \left( \Gamma_{H_1^-W^+}^{3(\chi_k^\pm - \chi_l^\pm)} m_i m_{\chi_k^\pm} + \Gamma_{H_1^-W^+}^{4(\chi_k^\pm - \chi_l^\pm)} m_i m_{\chi_l^\pm} + \Gamma_{H_1^-W^+}^{5(\chi_k^\pm - \chi_l^\pm)} m_{\chi_l^\pm} m_{\chi_k^\pm} \right) \\
&\mathcal{C}_9(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) + \Gamma_{H_1^-W^+}^{6(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_8(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) + \Gamma_{H_1^-W^+}^{7(\chi_k^\pm - \chi_l^\pm)} \\
&\mathcal{C}_6(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) + \Gamma_{H_1^-W^+}^{8(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_7(\chi_k^\pm, \chi_l^\pm, H_1^-, W^+) \quad (3.59)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{W^+H_1^-}^{tu(\chi_k^\pm, \chi_l^\pm)} &= I_{W^+H_1^-}^{1(\chi_k^\pm - \chi_l^\pm)} J_6(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) + I_{W^+H_1^-}^{2(\chi_k^\pm - \chi_l^\pm)} J_7(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\
&+ I_{W^+H_1^-}^{3(\chi_k^\pm - \chi_l^\pm)} J_8(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) + I_{W^+H_1^-}^{4(\chi_k^\pm - \chi_l^\pm)} J_9(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\
&+ I_{W^+H_1^-}^{5(\chi_k^\pm - \chi_l^\pm)} J_{10}(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) + I_{W^+H_1^-}^{6(\chi_k^\pm - \chi_l^\pm)} J_{11}(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \\
&+ I_{W^+H_1^-}^{7(\chi_k^\pm - \chi_l^\pm)} J_{12}(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) + I_{W^+H_1^-}^{8(\chi_k^\pm - \chi_l^\pm)} J_{13}(W^+, H_1^-, \chi_k^\pm, \chi_l^\pm) \quad (3.60)
\end{aligned}$$

$$\tilde{w}_{W^+H_1^-}^{(\chi_k^\pm, \chi_l^\pm)} = \tilde{w}_{W^+H_1^-}^{t(\chi_k^\pm, \chi_l^\pm)} + \tilde{w}_{W^+H_1^-}^{u(\chi_k^\pm, \chi_l^\pm)} + \left( \tilde{w}_{W^+H_1^-}^{tu(\chi_k^\pm, \chi_l^\pm)} + c.c. \right) \quad (3.61)$$

- Chargino ( $H_k^0 - \chi_l^\pm$ ) exchange:

$$\begin{aligned}
\tilde{w}_{W^+H_1^-}^{st(H_k^0, \chi_l^\pm)} &= \left( (-2dl_i + l_{ia} m_i^2) J_{W^+H_1^-}^{(5)(H_k^0 - \chi_l^\pm)} + J_{W^+H_1^-}^{(6)(H_k^0 - \chi_l^\pm)} l_i m_i m_j + J_{W^+H_1^-}^{(7)(H_k^0 - \chi_l^\pm)} \right. \\
&\left. l_i m_j m_{\chi_l^\pm} - J_{W^+H_1^-}^{(8)(H_k^0 - \chi_l^\pm)} l_{ia} m_i m_{\chi_l^\pm} \right) \mathcal{C}_{12}(H_k^0, \chi_l^\pm, H_1^-, W^+) \\
&+ \left( \left( 2l_{H_1^-W^+} (l_i - l_{ia}) M_W^2 - 2d_{W^+H_1^-} l_i s \right) J_{W^+H_1^-}^{(5)(H_k^0 - \chi_l^\pm)} - 2d_{W^+H_1^-} s \right. \\
&\left. \left( 2J_{W^+H_1^-}^{(6)(H_k^0 - \chi_l^\pm)} m_i m_j - 2J_{W^+H_1^-}^{(8)(H_k^0 - \chi_l^\pm)} m_i m_{\chi_l^\pm} + 2J_{W^+H_1^-}^{(7)(H_k^0 - \chi_l^\pm)} m_j m_{\chi_l^\pm} \right) \right) \\
&\frac{-b_{W^+H_1^-} + (\tau_{W^+H_1^-} - R_{W^+\chi_k^\pm}) \mathcal{A}(\chi_l^\pm, W^+, H^-)}{16b_{W^+H_1^-} M_W^2 s (m_{H_k^0}^2 - s)} \quad (3.62)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{W^+H_1^-}^{su(H_k^0, \chi_l^\pm)} &= \left( 4s \left( J_{H_1^-W^+}^{(6)(H_k^0-\chi_l^\pm)} \mathcal{C}_{13}(H_k^0, \chi_l^\pm, H_1^-, W^+) l_{ia} m_i m_{\chi_l^\pm} + J_{H_1^-W^+}^{(5)(H_k^0-\chi_l^\pm)} \right. \right. \\
&\quad m_j s l_i m_{\chi_l^\pm} \left( l_{W^+H_1^-} - 6M_W^2 \right) + J_{H_1^-W^+}^{(8)(H_k^0-\chi_l^\pm)} m_i m_j s \left( (M_W^2 - m_{H_1^-}^2) l_i \right. \\
&\quad \left. \left. + 6M_W^2 (M_W^2 - m_{H_1^-}^2 - m_i^2 + m_j^2) + 2b_{W^+H_1^-} s - l_{ia} s \right) \right) + 2J_{H_1^-W^+}^{(5)(H_k^0-\chi_l^\pm)} \\
&\quad (2ds \mathcal{C}_{14}(H_k^0, \chi_l^\pm, H_1^-, W^+) + 2\mathcal{C}_{13}(H_k^0, \chi_l^\pm, H_1^-, W^+) l_{ia} m_i^2 s \\
&\quad \left. + l_{H_1^-W^+} l_i l_{ia} (M_W^2 - m_{H_1^-}^2) (M_W^2 - s) \right) \frac{\mathcal{A}(\chi_l^\pm, H^-, W^+)}{16b_{W^+H_1^-} M_W^2 (m_{H_k^0}^2 - s) s} \\
&\quad \left( \left( J_{H_1^-W^+}^{(8)(H_k^0-\chi_l^\pm)} m_i m_j - J_{H_1^-W^+}^{(6)(H_k^0-\chi_l^\pm)} m_i m_{\chi_l^\pm} + J_{H_1^-W^+}^{(5)(H_k^0-\chi_l^\pm)} m_j m_{\chi_l^\pm} \right) \right. \\
&\quad \left. 4s (2M_W^2 - s) - 2J_{H_1^-W^+}^{(7)(H_k^0-\chi_l^\pm)} \left( l_i (M_W^2 - m_{H_1^-}^2) (M_W^2 - s) + 4M_W^2 s \right. \right. \\
&\quad \left. \left. (2m_i^2 - m_j^2) - 2m_i^2 s^2 \right) \right. \\
&\quad \left. - b_{W^+H_1^-} + (\tau_{H_1^-W^+} - R_{H_1^- \chi_k^\pm}) \mathcal{A}(\chi_l^\pm, H^-, W^+) \right) \\
&\quad \frac{1}{16b_{W^+H_1^-} M_W^2 (m_{H_k^0}^2 - s) s} \tag{3.63}
\end{aligned}$$

$$\tilde{w}_{W^+H_1^-}^{(H_k^0, \chi_l^\pm)} = \left( \tilde{w}_{W^+H_1^-}^{st(H_k^0, \chi_l^\pm)} + \tilde{w}_{W^+H_1^-}^{su(H_k^0, \chi_l^\pm)} \right) + c.c. \tag{3.64}$$

Note that  $\tilde{w}_{WH}$  given above does not include the contribution for  $W^- H^+$  final state. The contributions to the  $W^- H^+$  and  $W^+ H^-$  final states are obviously identical and so the total contribution for  $W^\pm H^\mp$  is twice that of  $W^+ H^-$

### 3.7 $\chi_i^0 \chi_j^0 \rightarrow ZH_n^0$

This process is sum of the s-channel CP odd Higgs boson and Z boson besides t and u channel neutralino exchange. Resulting cross section is obtained by taking the sum of the s, t, and u channel cross sections and possible cross sections obtained considering interference among themselves. And it is given by:

$$\tilde{w}_{Z, H_n^0} = \tilde{w}_{Z, H_n^0}^{s(A_1^0, A_1^0)} + \tilde{w}_{Z, H_n^0}^{s(Z, Z)} + \tilde{w}_{Z, H_n^0}^{s(A_1^0, Z)} + \sum_{k, l=1,4} \tilde{w}_{ZH_n^0}^{(\chi_k^0, \chi_l^0)} + \sum_{l=1,4} \tilde{w}_{Z, H_n^0}^{(Z, \chi_l^0)} + \sum_{l=1,4} \tilde{w}_{Z, H_n^0}^{(A_1^0, \chi_l^0)} \tag{3.65}$$

- CP-odd Higgs boson  $A_1^0$  and Z exchange:

$$\tilde{w}_{Z, H_n^0}^{s(A_1^0, A_1^0)} = \frac{1}{2} \frac{|C_{A_1^0 Z H_n^0}|^2}{(s - m_{A_1^0}^2)^2} \left( f_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} d - 2m_i m_j g_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} \right) (s - l_{ZH_n^0}) \tag{3.66}$$

$$\tilde{w}_{Z, H_n^0}^{s(Z, Z)} = \frac{|C_{ZZH_n^0}|^2}{(s - M_Z^2)^2} \left( f_{\chi_i^0 \chi_j^0}^{(Z-Z)} \mathcal{B}_7(Z, Z, H_n^0) + m_i m_j g_{\chi_i^0 \chi_j^0}^{(Z-Z)} \mathcal{B}_8(Z, Z, H_n^0) \right) \tag{3.67}$$

$$\begin{aligned}
\tilde{w}_{Z, H_n^0}^{s(A_1^0, Z)} &= \frac{C_{ZZH_n^0} C_{A_1^0 Z H_n^0}^*}{16M_Z^4 s (s - m_A^2)} \left( -l_i m_j f_{\chi_i^0 \chi_j^0}^{(Z-A_1^0)} + l_{ia} m_i g_{\chi_i^0 \chi_j^0}^{(Z-A_1^0)} \right) \\
&\quad \left( 2d_{ZH_n^0} l_{ZH_n^0} + l_{ZH_n^0}^2 - 2M_Z^2 (l_{H_n^0 Z} + 2s) \right) \tag{3.68}
\end{aligned}$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{ZH_n^0}^{t(\chi_k^0, \chi_l^0)} &= \Gamma_{ZH_n^0}^{1(\chi_k^0 - \chi_l^0)} \mathcal{C}_1(\chi_k^0, \chi_l^0, Z, H_n^0) + \Gamma_{ZH_n^0}^{2(\chi_k^0 - \chi_l^0)} \mathcal{C}_2(\chi_k^0, \chi_l^0, Z, H_n^0) - 3 m_i m_j s^2 \\
&\quad \left( M_Z^2 \Gamma_{ZH_n^0}^{3(\chi_k^0 - \chi_l^0)} + m_{\chi_k^0} m_{\chi_l^0} \Gamma_{ZH_n^0}^{4(\chi_k^0 - \chi_l^0)} \right) \mathcal{C}_3(\chi_k^0, \chi_l^0, Z, H_n^0) - \left( \Gamma_{ZH_n^0}^{5(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} \right. \\
&\quad \left. + \Gamma_{ZH_n^0}^{6(\chi_k^\pm - \chi_l^\pm)} m_{\chi_k^\pm} \right) \mathcal{C}_4(\chi_k^0, \chi_l^0, Z_n, H_n^0) + \left( \Gamma_{ZH_n^0}^{7(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} + \Gamma_{ZH_n^0}^{8(\chi_k^0 - \chi_l^0)} m_{\chi_k^0} \right) \\
&\quad \mathcal{C}_5(\chi_k^0, \chi_l^0, Z_n, H_n^0) + \left( \Gamma_{ZH_n^0}^{5(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} + \Gamma_{ZH_n^0}^{6(\chi_k^0 - \chi_l^0)} m_{\chi_k^0} - 2 \Gamma_{ZH_n^0}^{3(\chi_k^0 - \chi_l^0)} m_i \right) \\
&\quad \mathcal{C}_6(\chi_k^0, \chi_l^0, Z_n, H_n^0) \tag{3.69}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{ZH_n^0}^{u(\chi_k^0, \chi_l^0)} &= \left( \Gamma_{H_n^0 Z}^{1(\chi_k^0 - \chi_l^0)} m_{\chi_k^0} + \Gamma_{H_n^0 Z}^{2(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} \right) \mathcal{C}_{11}(\chi_k^0, \chi_l^0, H_n^0, Z) \\
&\quad + \left( \Gamma_{H_n^0 Z}^{3(\chi_k^0 - \chi_l^0)} m_{\chi_k^0} + \Gamma_{H_n^0 Z}^{4(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} \right) \mathcal{C}_{10}(\chi_k^0, \chi_l^0, H_n^0, Z) + \left( \Gamma_{H_n^0 Z}^{3(\chi_k^0 - \chi_l^0)} \right. \\
&\quad \left. m_i m_{\chi_k^0} + \Gamma_{H_n^0 Z}^{4(\chi_k^0 - \chi_l^0)} m_i m_{\chi_l^0} + \Gamma_{H_n^0 Z}^{5(\chi_k^0 - \chi_l^0)} m_{\chi_l^0} m_{\chi_k^0} \right) \mathcal{C}_9(\chi_k^0, \chi_l^0, H_n^0, Z) \\
&\quad + \Gamma_{H_n^0 Z}^{6(\chi_k^0 - \chi_l^0)} \mathcal{C}_8(\chi_k^0, \chi_l^0, H_n^0, Z) + \Gamma_{H_n^0 Z}^{7(\chi_k^0 - \chi_l^0)} \mathcal{C}_6(\chi_k^0, \chi_l^0, H_n^0, Z) \\
&\quad + \Gamma_{H_n^0 Z}^{8(\chi_k^0 - \chi_l^0)} \mathcal{C}_7(\chi_k^0, \chi_l^0, H_n^0, Z) \tag{3.70}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{ZH_n^0}^{tu(\chi_k^0, \chi_l^0)} &= I_{ZH_n^0}^{1(\chi_k^0 - \chi_l^0)} J_6(Z, H_n^0, \chi_k^0, \chi_l^0) + I_{ZH_n^0}^{2(\chi_k^0 - \chi_l^0)} J_7(Z, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad + I_{ZH_n^0}^{3(\chi_k^0 - \chi_l^0)} J_8(Z, H_n^0, \chi_k^0, \chi_l^0) + I_{ZH_n^0}^{4(\chi_k^0 - \chi_l^0)} J_9(Z, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad + I_{ZH_n^0}^{5(\chi_k^0 - \chi_l^0)} J_{10}(Z, H_n^0, \chi_k^0, \chi_l^0) + I_{ZH_n^0}^{6(\chi_k^0 - \chi_l^0)} J_{11}(Z, H_n^0, \chi_k^0, \chi_l^0) \\
&\quad + I_{ZH_n^0}^{7(\chi_k^0 - \chi_l^0)} J_{12}(Z, H_n^0, \chi_k^0, \chi_l^0) + I_{ZH_n^0}^{8(\chi_k^0 - \chi_l^0)} J_{13}(Z, H_n^0, \chi_k^0, \chi_l^0) \tag{3.71}
\end{aligned}$$

$$\tilde{w}_{ZH_n^0}^{(\chi_k^0, \chi_l^0)} = \tilde{w}_{ZH_n^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{ZH_n^0}^{u(\chi_k^0, \chi_l^0)} + \left( \tilde{w}_{ZH_n^0}^{tu(\chi_k^0, \chi_l^0)} + c.c. \right) \tag{3.72}$$

- Z boson, CP odd Higgs boson ( $A_1^0$ ) and neutralino ( $\chi_l^0$ ) exchange:

$$\begin{aligned}
\tilde{w}_{Z, H_n^0}^{st(Z, \chi_l^0)} &= \frac{C_{ZZH_n^0}}{s - M_Z^2} \left( \mathcal{B}_9(Z, \chi_l^0, Z, H_n^0) J_{ZH_n^0}^{(5)(Z - \chi_l^0)} + \mathcal{B}_{10}(Z, \chi_l^0, Z, H_n^0) J_{ZH_n^0}^{(6)(Z - \chi_l^0)} \right. \\
&\quad \left. + \mathcal{B}_{11}(Z, \chi_l^0, Z, H_n^0) J_{ZH_n^0}^{(7)(Z - \chi_l^0)} + \mathcal{B}_{12}(Z, \chi_l^0, Z, H_n^0) J_{ZH_n^0}^{(8)(Z - \chi_l^0)} \right) \tag{3.73}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{Z, H_n^0}^{su(Z, \chi_l^0)} &= \frac{C_{ZZH_n^0}}{s - M_Z^2} \left( \mathcal{B}_{13}(Z, \chi_l^0, H_n^0, Z) J_{H_n^0 Z}^{(5)(Z - \chi_l^0)} + \mathcal{B}_{14}(Z, \chi_l^0, H_n^0, Z) J_{H_n^0 Z}^{(6)(Z - \chi_l^0)} \right. \\
&\quad \left. + \mathcal{B}_{15}(Z, \chi_l^0, H_n^0, Z) J_{H_n^0 Z}^{(7)(Z - \chi_l^0)} + \mathcal{B}_{16}(Z, \chi_l^0, H_n^0, Z) J_{H_n^0 Z}^{(8)(Z - \chi_l^0)} \right) \tag{3.74}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{ZH_n^0}^{st(A_1^0, \chi_l^0)} &= \left( (-2dl_i + l_{ia} m_i^2) J_{ZH_n^0}^{(5)(A_1^0 - \chi_l^0)} + J_{ZH_n^0}^{(6)(A_1^0 - \chi_l^0)} l_i m_i m_j + J_{ZH_n^0}^{(7)(A_1^0 - \chi_l^0)} \right. \\
&\quad \left. l_i m_j m_{\chi_l^0} - J_{ZH_n^0}^{(8)(A_1^0 - \chi_l^0)} l_{ia} m_i m_{\chi_l^0} \right) \mathcal{C}_{12}(A_1^0, \chi_l^0, H_n^0, Z) \\
&\quad + \left( (2l_{H_n^0 Z} (l_i - l_{ia}) M_Z^2 - 2d_{ZH_n^0} l_i s) J_{ZH_n^0}^{(5)(A_1^0 - \chi_l^0)} - 2d_{ZH_n^0} s \right. \\
&\quad \left. \left( 2J_{ZH_n^0}^{(6)(A_1^0 - \chi_l^0)} m_i m_j - 2J_{ZH_n^0}^{(8)(A_1^0 - \chi_l^0)} m_i m_{\chi_l^0} + 2J_{ZH_n^0}^{(7)(A_1^0 - \chi_l^0)} m_j m_{\chi_l^0} \right) \right) \\
&\quad \frac{-b_{ZH_n^0} + (\tau_{ZH_n^0} - R_{Z\chi_k^0}) \mathcal{A}(\chi_l^0, Z, H_n^0)}{16b_{ZH_n^0} M_Z^2 s (m_{A_1^0}^2 - s)} \tag{3.75}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{ZH_n^0}^{su(A_1^0, \chi_l^0)} &= \left( 4s \left( J_{H_n^0 Z}^{(6)(A_1^0 - \chi_l^0)} \mathcal{C}_{13}(A_1^0, \chi_l^0, H_n^0, Z) l_{ia} m_i m_{\chi_l^0} + J_{H_n^0 Z}^{(5)(A_1^0 - \chi_l^0)} \right. \right. \\
&\quad \left. \left. m_j s l_i m_{\chi_l^0} (l_{ZH_n^0} - 6M_Z^2) + J_{H_n^0 Z}^{(8)(A_1^0 - \chi_l^0)} m_i m_j s \left( (M_Z^2 - m_{H_n^0}^2) l_i \right. \right. \right. \\
&\quad \left. \left. + 6M_Z^2 (M_Z^2 - m_{H_n^0}^2 - m_i^2 + m_j^2) + 2b_{ZH_n^0} s - l_{ia} s \right) \right) + 2J_{H_n^0 Z}^{(5)(A_1^0 - \chi_l^0)} \\
&\quad \left( 2ds \mathcal{C}_{14}(A_1^0, \chi_l^0, H_n^0, Z) + 2\mathcal{C}_{13}(A_1^0, \chi_l^0, H_n^0, Z) l_{ia} m_i^2 s \right. \\
&\quad \left. + l_{H_n^0 Z} l_i l_{ia} (M_Z^2 - m_{H_n^0}^2) (M_Z^2 - s) \right) \frac{\mathcal{A}(\chi_l^0, H_n^0, Z)}{16b_{ZH_n^0} M_Z^2 (m_{A_1^0}^2 - s) s} \\
&\quad \left( \left( J_{H_n^0 Z}^{(8)(A_1^0 - \chi_l^0)} m_i m_j - J_{H_n^0 Z}^{(6)(A_1^0 - \chi_l^0)} m_i m_{\chi_l^0} + J_{H_n^0 Z}^{(5)(A_1^0 - \chi_l^0)} m_j m_{\chi_l^0} \right) \right. \\
&\quad \left. 4s (2M_Z^2 - s) - 2J_{H_n^0 Z}^{(7)(A_1^0 - \chi_l^0)} \left( l_i (M_Z^2 - m_{H_n^0}^2) (M_Z^2 - s) + 4M_Z^2 s \right. \right. \\
&\quad \left. \left. (2m_i^2 - m_j^2) - 2m_i^2 s^2 \right) \right) \\
&\quad \frac{-b_{ZH_n^0} + (\tau_{H_n^0 Z} - R_{H_n^0 \chi_k^0}) \mathcal{A}(\chi_l^0, H_n^0, Z)}{16b_{ZH_n^0} M_Z^2 (m_{A_1^0}^2 - s) s} \tag{3.76}
\end{aligned}$$

$$\tilde{w}_{ZH_n^0}^{(A_1^0, \chi_l^0)} = \left( \tilde{w}_{ZH_n^0}^{st(A_1^0, \chi_l^0)} + \tilde{w}_{ZH_n^0}^{su(A_1^0, \chi_l^0)} \right) + c.c. \tag{3.77}$$

$$\tilde{w}_{Z, H_n^0}^{(Z, \chi_l^0)} = \left( \tilde{w}_{Z, H_n^0}^{st(Z, \chi_l^0)} + \tilde{w}_{Z, H_n^0}^{su(Z, \chi_l^0)} \right) + c.c. \tag{3.78}$$

$$\tilde{w}_{Z, H_n^0}^{(A_1^0, \chi_l^0)} = \left( \tilde{w}_{Z, H_n^0}^{st(A_1^0, \chi_l^0)} + \tilde{w}_{Z, H_n^0}^{su(A_1^0, \chi_l^0)} \right) + c.c. \tag{3.79}$$

### 3.8 $\chi_i^0 \chi_j^0 \rightarrow ZH_n^0$

This process is sum of the s-channel CP even Higgs boson besides t and u channel neutralino exchange. Resulting cross section is obtained by taking the sum of the s, t, and u channel cross sections and possible cross sections obtained considering interference among themselves. And it is given by:

$$\tilde{w}_{Z, A_1^0} = \sum_{k, l=1, 2} \tilde{w}_{Z, A_1^0}^{s(H_k^0, H_l^0)} + \sum_{k, l=1, 4} \tilde{w}_{ZH_n^0}^{(\chi_k^0, \chi_l^0)} + \sum_{k=1, 2} \sum_{l=1, 4} \tilde{w}_{H_k^0, A_1^0}^{(H_k^0, \chi_l^0)} \tag{3.80}$$

- CP-even Higgs boson  $H_k^0 - H_l^0$  exchange:

$$\tilde{w}_{ZA_1^0}^{(H_k^0, H_l^0)} = \frac{1}{2} \frac{C_{H_k^0 ZA_1^0} C_{H_l^0 ZA_1^0}^*}{(s - m_{H_k^0}^2)(s - m_{H_l^0}^2)} \left( f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} d - 2 m_i m_j g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} \right) (s - l_{ZA_1^0}) \quad (3.81)$$

- Neutralino ( $\chi_k^0 - \chi_l^0$ ) exchange:

$$\begin{aligned} \tilde{w}_{ZA_1^0}^{t(\chi_k^0, \chi_l^0)} &= \Gamma_{ZA_1^0}^1(\chi_k^0 - \chi_l^0) \mathcal{C}_1(\chi_k^0, \chi_l^0, Z, A_1^0) + \Gamma_{ZA_1^0}^2(\chi_k^0 - \chi_l^0) \mathcal{C}_2(\chi_k^0, \chi_l^0, Z, A_1^0) - 3 m_i m_j s^2 \\ &\quad \left( M_Z^2 \Gamma_{ZA_1^0}^3(\chi_k^0 - \chi_l^0) + m_{\chi_k^0} m_{\chi_l^0} \Gamma_{ZA_1^0}^4(\chi_k^0 - \chi_l^0) \right) \mathcal{C}_3(\chi_k^0, \chi_l^0, Z, A_1^0) - \left( \Gamma_{ZA_1^0}^5(\chi_k^0 - \chi_l^0) m_{\chi_l^0} \right. \\ &\quad \left. + \Gamma_{ZA_1^0}^6(\chi_k^\pm - \chi_l^\pm) m_{\chi_k^\pm} \right) \mathcal{C}_4(\chi_k^0, \chi_l^0, Z_n, A_1^0) + \left( \Gamma_{ZA_1^0}^7(\chi_k^0 - \chi_l^0) m_{\chi_l^0} + \Gamma_{ZA_1^0}^8(\chi_k^0 - \chi_l^0) m_{\chi_k^0} \right) \\ &\quad \mathcal{C}_5(\chi_k^0, \chi_l^0, Z_n, A_1^0) + \left( \Gamma_{ZA_1^0}^5(\chi_k^0 - \chi_l^0) m_{\chi_l^0} + \Gamma_{ZA_1^0}^6(\chi_k^0 - \chi_l^0) m_{\chi_k^0} - 2 \Gamma_{ZA_1^0}^3(\chi_k^0 - \chi_l^0) m_i \right) \\ &\quad \mathcal{C}_6(\chi_k^0, \chi_l^0, Z_n, A_1^0) \end{aligned} \quad (3.82)$$

$$\begin{aligned} \tilde{w}_{ZA_1^0}^{u(\chi_k^0, \chi_l^0)} &= \left( \Gamma_{A_1^0 Z}^1(\chi_k^0 - \chi_l^0) m_{\chi_k^0} + \Gamma_{A_1^0 Z}^2(\chi_k^0 - \chi_l^0) m_{\chi_l^0} \right) \mathcal{C}_{11}(\chi_k^0, \chi_l^0, A_1^0, Z) \\ &\quad + \left( \Gamma_{A_1^0 Z}^3(\chi_k^0 - \chi_l^0) m_{\chi_k^0} + \Gamma_{A_1^0 Z}^4(\chi_k^0 - \chi_l^0) m_{\chi_l^0} \right) \mathcal{C}_{10}(\chi_k^0, \chi_l^0, A_1^0, Z) + \left( \Gamma_{A_1^0 Z}^3(\chi_k^0 - \chi_l^0) \right. \\ &\quad \left. m_i m_{\chi_k^0} + \Gamma_{A_1^0 Z}^4(\chi_k^0 - \chi_l^0) m_i m_{\chi_l^0} + \Gamma_{A_1^0 Z}^5(\chi_k^0 - \chi_l^0) m_{\chi_l^0} m_{\chi_k^0} \right) \mathcal{C}_9(\chi_k^0, \chi_l^0, A_1^0, Z) \\ &\quad + \Gamma_{A_1^0 Z}^6(\chi_k^0 - \chi_l^0) \mathcal{C}_8(\chi_k^0, \chi_l^0, A_1^0, Z) + \Gamma_{A_1^0 Z}^7(\chi_k^0 - \chi_l^0) \mathcal{C}_6(\chi_k^0, \chi_l^0, A_1^0, Z) \\ &\quad + \Gamma_{A_1^0 Z}^8(\chi_k^0 - \chi_l^0) \mathcal{C}_7(\chi_k^0, \chi_l^0, A_1^0, Z) \end{aligned} \quad (3.83)$$

$$\begin{aligned} \tilde{w}_{ZA_1^0}^{tu(\chi_k^0, \chi_l^0)} &= I_{ZA_1^0}^1(\chi_k^0 - \chi_l^0) J_6(Z, A_1^0, \chi_k^0, \chi_l^0) + I_{ZA_1^0}^2(\chi_k^0 - \chi_l^0) J_7(Z, A_1^0, \chi_k^0, \chi_l^0) \\ &\quad + I_{ZA_1^0}^3(\chi_k^0 - \chi_l^0) J_8(Z, A_1^0, \chi_k^0, \chi_l^0) + I_{ZA_1^0}^4(\chi_k^0 - \chi_l^0) J_9(Z, A_1^0, \chi_k^0, \chi_l^0) \\ &\quad + I_{ZA_1^0}^5(\chi_k^0 - \chi_l^0) J_{10}(Z, A_1^0, \chi_k^0, \chi_l^0) + I_{ZA_1^0}^6(\chi_k^0 - \chi_l^0) J_{11}(Z, A_1^0, \chi_k^0, \chi_l^0) \\ &\quad + I_{ZA_1^0}^7(\chi_k^0 - \chi_l^0) J_{12}(Z, A_1^0, \chi_k^0, \chi_l^0) + I_{ZA_1^0}^8(\chi_k^0 - \chi_l^0) J_{13}(Z, A_1^0, \chi_k^0, \chi_l^0) \end{aligned} \quad (3.84)$$

$$\tilde{w}_{ZA_1^0}^{(\chi_k^0, \chi_l^0)} = \tilde{w}_{ZA_1^0}^{t(\chi_k^0, \chi_l^0)} + \tilde{w}_{ZA_1^0}^{u(\chi_k^0, \chi_l^0)} + \left( \tilde{w}_{ZA_1^0}^{tu(\chi_k^0, \chi_l^0)} + c.c. \right) \quad (3.85)$$

- CP even Higgs boson ( $H_k^0$ ) and neutralino ( $\chi_l^0$ ) exchange:

$$\begin{aligned} \tilde{w}_{ZA_1^0}^{st(H_k^0, \chi_l^0)} &= \left( (-2 d l_i + l_{ia} m_i^2) J_{ZA_1^0}^{(5)(H_k^0 - \chi_l^0)} + J_{ZA_1^0}^{(6)(H_k^0 - \chi_l^0)} l_i m_i m_j + J_{ZA_1^0}^{(7)(H_k^0 - \chi_l^0)} \right. \\ &\quad \left. l_i m_j m_{\chi_l^0} - J_{ZA_1^0}^{(8)(H_k^0 - \chi_l^0)} l_{ia} m_i m_{\chi_l^0} \right) \mathcal{C}_{12}(H_k^0, \chi_l^0, A_1^0, Z) \\ &\quad + \left( \left( 2 l_{A_1^0 Z} (l_i - l_{ia}) M_Z^2 - 2 d_{ZA_1^0} l_i s \right) J_{ZA_1^0}^{(5)(H_k^0 - \chi_l^0)} - 2 d_{ZA_1^0} s \right. \\ &\quad \left. \left( 2 J_{ZA_1^0}^{(6)(H_k^0 - \chi_l^0)} m_i m_j - 2 J_{ZA_1^0}^{(8)(H_k^0 - \chi_l^0)} m_i m_{\chi_l^0} + 2 J_{ZA_1^0}^{(7)(H_k^0 - \chi_l^0)} m_j m_{\chi_l^0} \right) \right) \\ &\quad \frac{-b_{ZA_1^0} + (\tau_{ZA_1^0} - R_{Z\chi_k^0}) \mathcal{A}(\chi_l^0, Z, A_1^0)}{16 b_{ZA_1^0} M_Z^2 s (m_{H_k^0}^2 - s)} \end{aligned} \quad (3.86)$$

$$\begin{aligned}
\tilde{w}_{ZA_1^0}^{su(H_k^0, \chi_l^0)} &= \left( 4s \left( J_{A_1^0 Z}^{(6)(H_k^0 - \chi_l^0)} \mathcal{C}_{13}(H_k^0, \chi_l^0, A_1^0, Z) l_{ia} m_i m_{\chi_l^0} + J_{A_1^0 Z}^{(5)(H_k^0 - \chi_l^0)} \right. \right. \\
&\quad m_j s l_i m_{\chi_l^0} \left( l_{ZA_1^0} - 6M_Z^2 \right) + J_{A_1^0 Z}^{(8)(H_k^0 - \chi_l^0)} m_i m_j s \left( (M_Z^2 - m_{A_1^0}^2) l_i \right. \\
&\quad \left. \left. + 6M_Z^2 (M_Z^2 - m_{A_1^0}^2 - m_i^2 + m_j^2) + 2b_{ZA_1^0} s - l_{ia} s \right) \right) + 2J_{A_1^0 Z}^{(5)(H_k^0 - \chi_l^0)} \\
&\quad \left( 2ds \mathcal{C}_{14}(H_k^0, \chi_l^0, A_1^0, Z) + 2\mathcal{C}_{13}(H_k^0, \chi_l^0, A_1^0, Z) l_{ia} m_i^2 s \right. \\
&\quad \left. \left. + l_{A_1^0 Z} l_i l_{ia} (M_Z^2 - m_{A_1^0}^2) (M_Z^2 - s) \right) \right) \frac{\mathcal{A}(\chi_l^0, A_1^0, Z)}{16 b_{ZA_1^0} M_Z^2 (m_{H_k^0}^2 - s) s} \\
&\quad \left( \left( J_{A_1^0 Z}^{(8)(H_k^0 - \chi_l^0)} m_i m_j - J_{A_1^0 Z}^{(6)(H_k^0 - \chi_l^0)} m_i m_{\chi_l^0} + J_{A_1^0 Z}^{(5)(H_k^0 - \chi_l^0)} m_j m_{\chi_l^0} \right) \right. \\
&\quad \left. 4s (2M_Z^2 - s) - 2J_{A_1^0 Z}^{(7)(H_k^0 - \chi_l^0)} \left( l_i (M_Z^2 - m_{A_1^0}^2) (M_Z^2 - s) + 4M_Z^2 s \right. \right. \\
&\quad \left. \left. (2m_i^2 - m_j^2) - 2m_i^2 s^2 \right) \right. \\
&\quad \left. - b_{ZA_1^0} + (\tau_{A_1^0 Z} - R_{A_1^0 \chi_k^0}) \mathcal{A}(\chi_l^0, A_1^0, Z) \right) \\
&\quad \frac{1}{16 b_{ZA_1^0} M_Z^2 (m_{H_k^0}^2 - s) s} \tag{3.87}
\end{aligned}$$

$$\tilde{w}_{ZA_1^0}^{(H_k^0, \chi_l^0)} = \left( \tilde{w}_{ZA_1^0}^{st(H_k^0, \chi_l^0)} + \tilde{w}_{ZA_1^0}^{su(H_k^0, \chi_l^0)} \right) + c.c. \tag{3.88}$$

### 3.9 $\chi_i^0 \chi_j^0 \rightarrow W^\pm W^\mp$

This process is the sum of s-channel diagram through CP even ( $H_1^0, H_2^0$ ), Z boson and t- and u-channel chargino ( $\chi_k^\pm, k = 1, 2$ ) exchange

$$\tilde{w}_{W^\pm W^\mp} = \sum_{k,l=1,2} \tilde{w}_{W^\pm W^\mp}^{(H_k^0, H_l^0)} + \tilde{w}_{W^\pm W^\mp}^{(Z,Z)} + \sum_{k,l=1,2} \tilde{w}_{W^\pm W^\mp}^{(\chi_k^\pm, \chi_l^\pm)} + \sum_{k=1,2} \sum_{l=1,4} \tilde{w}_{W^\pm W^\mp}^{(H_k^0, \chi_l^\pm)} + \sum_{l=1,4} \tilde{w}_{W^\pm W^\mp}^{(Z, \chi_l^\pm)} \tag{3.89}$$

- CP-even Higgs boson ( $H_k^0 - H_l^0$ ) and CP-odd Higgs boson (Z):

$$\begin{aligned}
\tilde{w}_{WW}^{(H_k^0 - H_l^0)} &= \frac{C_{H_k^0 WW} C_{H_l^0 WW}^*}{(S - m_{H_k^0}^2)(S - m_{H_l^0}^2)} \frac{1}{8M_W^4} (12M_W^4 - 4M_W^2 S + S^2) \\
&\quad (f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} d - g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} m_i m_j) \tag{3.90}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{WW}^{(Z-Z)} &= \frac{C_{ZWW} C_{ZWW}^*}{(S - M_Z^2)^2} \left( -\frac{f_{\chi_i^0 \chi_j^0}^{(Z-Z)}}{48M_W^2 S^2} (24l_i l_{ia} M_W^2 (4M_W^2 - S) S^2 \right. \\
&\quad \left. + b_{WW}^2 (12M_W^4 - 4M_W^2 S + S^2)) + \frac{g_{\chi_i^0 \chi_j^0}^{(Z-Z)} m_i m_j}{M_W^2} S (S - 4M_W^2) \right. \\
&\quad \left. + \frac{f_{\chi_i^0 \chi_j^0}^{(Z-Z)} d + g_{\chi_i^0 \chi_j^0}^{(Z-Z)} m_i m_j}{8M_W^4} (80M_W^6 - 36M_W^4 S - 8M_W^2 S^2 + 3S^3) \right) \tag{3.91}
\end{aligned}$$

$$\tilde{w}_{WW}^{(Z-H_l^0)} = 0 \tag{3.92}$$

- Chargino exchange:

$$\begin{aligned}
\tilde{w}_{WW}^{t(\chi_k^\pm - \chi_l^\pm)} &= \Gamma_{WW}^{1(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{12}(\chi_k^\pm, \chi_l^\pm, W, W) + \Gamma_{WW}^{2(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{13}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \Gamma_{WW}^{3(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{14}(\chi_k^\pm, \chi_l^\pm, W, W) + \Gamma_{WW}^{4(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{15}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \left( m_{\chi_l^\pm} \Gamma_{WW}^{5(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_k^\pm} \Gamma_{WW}^{6(\chi_k^\pm - \chi_l^\pm)} \right) \mathcal{C}_{16}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \left( m_{\chi_l^\pm} \Gamma_{WW}^{7(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_k^\pm} \Gamma_{WW}^{8(\chi_k^\pm - \chi_l^\pm)} \right) \mathcal{C}_{17}(\chi_k^\pm, \chi_l^\pm, W, W) \quad (3.93)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{WW}^{u(\chi_k^\pm - \chi_l^\pm)} &= \Gamma_{WW}^{1(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{12}(\chi_k^\pm, \chi_l^\pm, W, W) + \Gamma_{WW}^{2(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{13}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \Gamma_{WW}^{3(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{14}(\chi_k^\pm, \chi_l^\pm, W, W) + \Gamma_{WW}^{4(\chi_k^\pm - \chi_l^\pm)} \mathcal{C}_{15}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \left( m_{\chi_l^\pm} \Gamma_{WW}^{5(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_k^\pm} \Gamma_{WW}^{6(\chi_k^\pm - \chi_l^\pm)} \right) \mathcal{C}_{16}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ \left( m_{\chi_l^\pm} \Gamma_{WW}^{7(\chi_k^\pm - \chi_l^\pm)} + m_{\chi_k^\pm} \Gamma_{WW}^{8(\chi_k^\pm - \chi_l^\pm)} \right) \mathcal{C}_{17}(\chi_k^\pm, \chi_l^\pm, W, W) \quad (3.94)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{WW}^{tu(\chi_k^\pm - \chi_l^\pm)} &= I_{WW}^{1(\chi_k^\pm - \chi_l^\pm)} J_{14}(\chi_k^\pm, \chi_l^\pm, W, W) + I_{WW}^{2(\chi_k^\pm - \chi_l^\pm)} J_{15}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ I_{WW}^{3(\chi_k^\pm - \chi_l^\pm)} J_{16}(\chi_k^\pm, \chi_l^\pm, W, W) + I_{WW}^{4(\chi_k^\pm - \chi_l^\pm)} J_{17}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ I_{WW}^{5(\chi_k^\pm - \chi_l^\pm)} J_{18}(\chi_k^\pm, \chi_l^\pm, W, W) + I_{WW}^{6(\chi_k^\pm - \chi_l^\pm)} J_{19}(\chi_k^\pm, \chi_l^\pm, W, W) \\
&+ I_{WW}^{7(\chi_k^\pm - \chi_l^\pm)} J_{20}(\chi_k^\pm, \chi_l^\pm, W, W) + I_{WW}^{8(\chi_k^\pm - \chi_l^\pm)} J_{21}(\chi_k^\pm, \chi_l^\pm, W, W) \quad (3.95)
\end{aligned}$$

- CP-even Higgs boson ( $H_k^0$ ), Z boson interference with chargino ( $\chi_l^\pm$ ):

$$\begin{aligned}
\tilde{w}_{WW}^{st(H_k^0 - \chi_l^\pm)} &= \frac{C_{H_k^0 WW}}{s - m_{H_k^0}^2} \left( J_{WW}^{(5)(H_k^0 - \chi_l^\pm)} \mathcal{B}_{20}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(6)(H_k^0 - \chi_l^\pm)} \right. \\
&\mathcal{B}_{21}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(7)(H_k^0 - \chi_l^\pm)} \mathcal{B}_{22}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(8)(H_k^0 - \chi_l^\pm)} \\
&\left. \mathcal{B}_{23}(H_k^0, \chi_l^\pm, W, W) \right) \quad (3.96)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{WW}^{su(H_k^0 - \chi_l^\pm)} &= \frac{C_{H_k^0 WW}}{s - m_{H_k^0}^2} \left( J_{WW}^{(5)(H_k^0 - \chi_l^\pm)*} \mathcal{B}_{20}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(6)(H_k^0 - \chi_l^\pm)*} \right. \\
&\mathcal{B}_{21}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(7)(H_k^0 - \chi_l^\pm)*} \mathcal{B}_{22}(H_k^0, \chi_l^\pm, W, W) + J_{WW}^{(8)(H_k^0 - \chi_l^\pm)*} \\
&\left. \mathcal{B}_{23}(H_k^0, \chi_l^\pm, W, W) \right) \quad (3.97)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{WW}^{st(Z - \chi_l^\pm)} &= \frac{C_{Z WW}}{s - M_Z^2} \left( J_{WW}^{(5)(Z - \chi_l^\pm)} \mathcal{B}_{24}(Z, \chi_l^\pm, W, W) + J_{WW}^{(6)(Z - \chi_l^\pm)} \right. \\
&\mathcal{B}_{25}(Z, \chi_l^\pm, W, W) + J_{WW}^{(7)(Z - \chi_l^\pm)} \mathcal{B}_{26}(Z, \chi_l^\pm, W, W) + J_{WW}^{(8)(Z - \chi_l^\pm)} \\
&\left. \mathcal{B}_{27}(Z, \chi_l^\pm, W, W) \right) \quad (3.98)
\end{aligned}$$

$$\begin{aligned} \tilde{w}_{WW}^{su(Z-\chi_l^\pm)} &= -\frac{C_{ZWW}}{s-M_Z^2} \left( J_{WW}^{(5)(Z-\chi_l^\pm)*} \mathcal{B}_{24}(Z, \chi_l^\pm, W, W) + J_{WW}^{(6)(Z-\chi_l^\pm)*} \right. \\ &\quad \mathcal{B}_{25}(Z, \chi_l^\pm, W, W) + J_{WW}^{(7)(Z-\chi_l^\pm)*} \mathcal{B}_{26}(Z, \chi_l^\pm, W, W) + J_{WW}^{(8)(Z-\chi_l^\pm)*} \\ &\quad \left. \mathcal{B}_{27}(Z, \chi_l^\pm, W, W) \right) \end{aligned} \quad (3.99)$$

$$\tilde{w}_{WW}^{(H_k^0, \chi_l^\pm)} = \left( \tilde{w}_{WW}^{st(H_k^0, \chi_l^\pm)} + \tilde{w}_{WW}^{su(H_k^0, \chi_l^\pm)} \right) + c.c. \quad (3.100)$$

$$\tilde{w}_{WW}^{(Z, \chi_l^\pm)} = \left( \tilde{w}_{WW}^{st(Z, \chi_l^\pm)} + \tilde{w}_{WW}^{su(Z, \chi_l^\pm)} \right) + c.c. \quad (3.101)$$

### 3.10 $\chi_i^0 \chi_j^0 \rightarrow ZZ$

This process is the sum of s-channel CP even ( $H_1^0, H_2^0$ ) and the t- and u-channel neutralino ( $\chi_k^0, k = 1, 4$ ) exchange. The resulting cross section is

$$\tilde{w}_{ZZ} = \sum_{k,l=1,2} \tilde{w}_{ZZ}^{(H_k^0, H_l^0)} + \sum_{k,l=1,2} \tilde{w}_{ZZ}^{(\chi_k^0, \chi_l^0)} + \sum_{l=1,4} \tilde{w}_{ZZ}^{(H_k^0, \chi_l^0)} \quad (3.102)$$

- CP-even Higgs boson ( $H_k^0 - H_l^0$ ) :

$$\begin{aligned} \tilde{w}_{ZZ}^{(H_k^0 - H_l^0)} &= \frac{C_{H_k^0 ZZ} C_{H_l^0 ZZ}^*}{(S - m_{H_k^0}^2)(S - m_{H_l^0}^2)} \frac{1}{8 M_Z^4} (12 M_Z^4 - 4 M_Z^2 S + S^2) \\ &\quad (f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} d - g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} m_i m_j) \end{aligned} \quad (3.103)$$

- Chargino exchange:

$$\begin{aligned} \tilde{w}_{ZZ}^{t(\chi_k^0 - \chi_l^0)} &= \Gamma_{ZZ}^{1(\chi_k^0 - \chi_l^0)} \mathcal{C}_{12}(\chi_k^0, \chi_l^0, Z, Z) + \Gamma_{ZZ}^{2(\chi_k^0 - \chi_l^0)} \mathcal{C}_{13}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \Gamma_{ZZ}^{3(\chi_k^0 - \chi_l^0)} \mathcal{C}_{14}(\chi_k^0, \chi_l^0, Z, Z) + \Gamma_{ZZ}^{4(\chi_k^0 - \chi_l^0)} \mathcal{C}_{15}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \left( m_{\chi_l^0} \Gamma_{ZZ}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_k^0} \Gamma_{ZZ}^{6(\chi_k^0 - \chi_l^0)} \right) \mathcal{C}_{16}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \left( m_{\chi_l^0} \Gamma_{ZZ}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_k^0} \Gamma_{ZZ}^{8(\chi_k^0 - \chi_l^0)} \right) \mathcal{C}_{17}(\chi_k^0, \chi_l^0, Z, Z) \end{aligned} \quad (3.104)$$

$$\begin{aligned} \tilde{w}_{ZZ}^{u(\chi_k^0 - \chi_l^0)} &= \Gamma_{ZZ}^{1(\chi_k^0 - \chi_l^0)} \mathcal{C}_{12}(\chi_k^0, \chi_l^0, Z, Z) + \Gamma_{ZZ}^{2(\chi_k^0 - \chi_l^0)} \mathcal{C}_{13}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \Gamma_{ZZ}^{3(\chi_k^0 - \chi_l^0)} \mathcal{C}_{14}(\chi_k^0, \chi_l^0, Z, Z) + \Gamma_{ZZ}^{4(\chi_k^0 - \chi_l^0)} \mathcal{C}_{15}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \left( m_{\chi_l^0} \Gamma_{ZZ}^{5(\chi_k^0 - \chi_l^0)} + m_{\chi_k^0} \Gamma_{ZZ}^{6(\chi_k^0 - \chi_l^0)} \right) \mathcal{C}_{16}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + \left( m_{\chi_l^0} \Gamma_{ZZ}^{7(\chi_k^0 - \chi_l^0)} + m_{\chi_k^0} \Gamma_{ZZ}^{8(\chi_k^0 - \chi_l^0)} \right) \mathcal{C}_{17}(\chi_k^0, \chi_l^0, Z, Z) \end{aligned} \quad (3.105)$$

$$\begin{aligned} \tilde{w}_{ZZ}^{tu(\chi_k^0 - \chi_l^0)} &= I_{ZZ}^{1(\chi_k^0 - \chi_l^0)} J_{14}(\chi_k^0, \chi_l^0, Z, Z) + I_{ZZ}^{2(\chi_k^0 - \chi_l^0)} J_{15}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + I_{ZZ}^{3(\chi_k^0 - \chi_l^0)} J_{16}(\chi_k^0, \chi_l^0, Z, Z) + I_{ZZ}^{4(\chi_k^0 - \chi_l^0)} J_{17}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + I_{ZZ}^{5(\chi_k^0 - \chi_l^0)} J_{18}(\chi_k^0, \chi_l^0, Z, Z) + I_{ZZ}^{6(\chi_k^0 - \chi_l^0)} J_{19}(\chi_k^0, \chi_l^0, Z, Z) \\ &\quad + I_{ZZ}^{7(\chi_k^0 - \chi_l^0)} J_{20}(\chi_k^0, \chi_l^0, Z, Z) + I_{ZZ}^{8(\chi_k^0 - \chi_l^0)} J_{21}(\chi_k^0, \chi_l^0, Z, Z) \end{aligned} \quad (3.106)$$



- CP-even Higgs boson ( $H_k^0$ ) interference with neutralino ( $\chi_l^0$ ) :

$$\begin{aligned} \tilde{w}_{ZZ}^{st(H_k^0-\chi_l^0)} &= \frac{C_{H_k^0 ZZ}}{s - m_{H_k^0}^2} \left( J_{ZZ}^{(5)(H_k^0-\chi_l^0)} \mathcal{B}_{20}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(6)(H_k^0-\chi_l^0)} \right. \\ &\quad \mathcal{B}_{21}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(7)(H_k^0-\chi_l^0)} \mathcal{B}_{22}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(8)(H_k^0-\chi_l^0)} \\ &\quad \left. \mathcal{B}_{23}(H_k^0, \chi_l^0, Z, Z) \right) \end{aligned} \quad (3.107)$$

$$\begin{aligned} \tilde{w}_{ZZ}^{su(H_k^0-\chi_l^0)} &= \frac{C_{H_k^0 ZZ}}{s - m_{H_k^0}^2} \left( J_{ZZ}^{(5)(H_k^0-\chi_l^0)} \mathcal{B}_{20}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(6)(H_k^0-\chi_l^0)} \right. \\ &\quad \mathcal{B}_{21}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(7)(H_k^0-\chi_l^0)} \mathcal{B}_{22}(H_k^0, \chi_l^0, Z, Z) + J_{ZZ}^{(8)(H_k^0-\chi_l^0)} \\ &\quad \left. \mathcal{B}_{23}(H_k^0, \chi_l^0, Z, W) \right) \end{aligned} \quad (3.108)$$

$$\tilde{w}_{ZZ}^{(H_k^0, \chi_l^0)} = \left( \tilde{w}_{ZZ}^{st(H_k^0, \chi_l^0)} + \tilde{w}_{ZZ}^{su(H_k^0, \chi_l^0)} \right) + c.c. \quad (3.109)$$

### 3.11 $\chi_i^0 \chi_j^0 \rightarrow f_m \bar{f}_n$

This process is sum of s channel diagrams through CP even Higgs boson, CP odd Higgs boson and Z boson and sfermion t and u channel. This is the only process which involves flavour violation effects from SUSY soft sector. The resultant cross section is given by:

$$\begin{aligned} \tilde{w}_{f_m \bar{f}_n} &= \sum_{k,l=1,2} \tilde{w}_{f_m \bar{f}_n}^{(H_k^0, H_l^0)} + \tilde{w}_{f_m \bar{f}_n}^{(A_1^0, A_1^0)} + \tilde{w}_{f_m \bar{f}_n}^{(Z, Z)} + \sum_{k=1,2} \tilde{w}_{f_m \bar{f}_n}^{(H_k^0, Z)} + \tilde{w}_{f_m \bar{f}_n}^{(A_1^0, Z)} + \sum_{k,l=1,p} \tilde{w}_{f_m \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} \\ &+ \sum_{k=1,2} \sum_{l=1,p} \tilde{w}_{f_m \bar{f}_n}^{(H_k^0 - \tilde{f}_l)} + \sum_{l=1,p} \tilde{w}_{f_m \bar{f}_n}^{(A_1^0 - \tilde{f}_l)} + \sum_{l=1,p} \tilde{w}_{f_m \bar{f}_n}^{(Z - \tilde{f}_l)} \end{aligned} \quad (3.110)$$

- $H_k^0 - H_l^0$ ,  $A_1^0$  and Z boson exchange exchange:

$$\begin{aligned} \tilde{w}_{f_m \bar{f}_n}^{(H_k^0, H_l^0)} &= 2 \frac{C_{H_k^0 f_m \bar{f}_n} C_{H_l^0 f_m \bar{f}_n}^*}{(s - m_{H_k^0}^2)(s - m_{H_l^0}^2)} \left( d f_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} - g_{\chi_i^0 \chi_j^0}^{(H_k^0 - H_l^0)} m_i m_j \right) \\ &\quad (d_{f_m \bar{f}_n} - m_{f_m} m_{\bar{f}_n}) \end{aligned} \quad (3.111)$$

$$\begin{aligned} \tilde{w}_{f_m \bar{f}_n}^{(A_1^0, A_1^0)} &= -2 \frac{C_{A_1^0 f_m \bar{f}_n} C_{A_1^0 f_m \bar{f}_n}^*}{(s - m_{A_1^0}^2)(s - m_{A_1^0}^2)} \left( d f_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} - g_{\chi_i^0 \chi_j^0}^{(A_1^0 - A_1^0)} m_i m_j \right) \\ &\quad (d_{f_m \bar{f}_n} + m_{f_m} m_{\bar{f}_n}) \end{aligned} \quad (3.112)$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{(Z,Z)} &= \frac{|C_{Z f_m \bar{f}_n}^V|^2 + |C_{Z f_m \bar{f}_n}^A|^2}{12 M_Z^4 s^2 (s - M_Z^2)^2} (12 s^2 (2 M_Z^2 - s) (l_{f_m \bar{f}_n} l_{\bar{f}_n f_m} - 2 d_{f_m \bar{f}_n} s) \\
&\quad (d f_{\chi_i^0 \chi_j^0}^{(Z-Z)} + g_{\chi_i^0 \chi_j^0}^{(Z-Z)} m_i m_j) + f_{\chi_i^0 \chi_j^0}^{(Z-Z)} (2 b_{f_m \bar{f}_n}^2 M_Z^4 + 6 l_i l_{ia} s^2 (l_{f_m \bar{f}_n} l_{\bar{f}_n f_m} \\
&\quad + 4 d_{f_m \bar{f}_n} M_Z^2 - 2 d_{f_m \bar{f}_n} s) + 3 (\tau_{\bar{f}_n f_m} \omega_{f_m \bar{f}_n} + \omega_{\bar{f}_n f_m} \tau_{f_m \bar{f}_n}) (M_Z^4 - 2 M_Z^2 s)) \\
&\quad + 48 g_{\chi_i^0 \chi_j^0}^{(Z-Z)} s^2 M_Z^4 d_{f_m \bar{f}_n} m_i m_j) + \frac{m_{f_m} m_{\bar{f}_n} (|C_{Z f_m \bar{f}_n}^V|^2 - |C_{Z f_m \bar{f}_n}^A|^2)}{M_Z^4 (s - M_Z^2)^2} \\
&\quad \left( (d f_{\chi_i^0 \chi_j^0}^{(Z-Z)} + g_{\chi_i^0 \chi_j^0}^{(Z-Z)} m_i m_j) (-4 M_Z^2 s + 2 s^2) + 4 M_Z^4 (d f_{\chi_i^0 \chi_j^0}^{(Z-Z)} \right. \\
&\quad \left. + 2 g_{\chi_i^0 \chi_j^0}^{(Z-Z)} m_i m_j) + f_{\chi_i^0 \chi_j^0}^{(Z-Z)} l_i l_{ia} (2 M_Z^2 + s) \right) \\
&\quad - \frac{(\tau_{f_m \bar{f}_n} \omega_{\bar{f}_n f_m} - \tau_{\bar{f}_n f_m} \omega_{f_m \bar{f}_n}) (s + M_Z^2)}{4 s^2 (s - M_Z^2)^2 M_Z^2} \\
&\quad \left( C_{\chi_i^0 \chi_j^0 Z}^L C_{\chi_i^0 \chi_j^0 Z}^{L*} - C_{\chi_i^0 \chi_j^0 Z}^R C_{\chi_i^0 \chi_j^0 Z}^{R*} \right) \left( C_{Z f_m \bar{f}_n}^{A*} C_{Z f_m \bar{f}_n}^V + C_{Z f_m \bar{f}_n}^A C_{Z f_m \bar{f}_n}^{V*} \right)
\end{aligned} \tag{3.113}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(H_k^0, Z)} = \frac{(l_{\bar{f}_n f_m} m_{f_m} - l_{f_m \bar{f}_n} m_{\bar{f}_n})}{2 M_Z^2 s (s - m_{H_k^0}^2)} C_{Z f_m \bar{f}_n}^{V*} \left( f_{\chi_i^0 \chi_j^0}^{(Z-Z)} l_i m_j - g_{\chi_i^0 \chi_j^0}^{(Z-Z)} l_{ia} m_i \right) \tag{3.114}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(A_1^0, Z)} = -\frac{(l_{\bar{f}_n f_m} m_{f_m} + l_{f_m \bar{f}_n} m_{\bar{f}_n})}{2 M_Z^2 s (s - m_{H_k^0}^2)} C_{Z f_m \bar{f}_n}^{A*} \left( f_{\chi_i^0 \chi_j^0}^{(Z-Z)} l_i m_j - g_{\chi_i^0 \chi_j^0}^{(Z-Z)} l_{ia} m_i \right) \tag{3.115}$$

- Sfermion ( $\tilde{f}_k - \tilde{f}_l$ ) exchange:

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{t(\tilde{f}_k - \tilde{f}_l)} &= \frac{1}{8 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} - R_{f_m \tilde{f}_l})} \left( 2 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} - R_{f_m \tilde{f}_l}) \right. \\
&\quad \left. f_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} \right) - \frac{\mathcal{X}_1(\tilde{f}_k, \tilde{f}_l, f_m, \bar{f}_n)}{2 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} - R_{f_m \tilde{f}_l})} \left( g_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} m_i m_{f_m} s \right. \\
&\quad \left. (f_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} (\omega_{\bar{f}_n f_m} - \tau_{f_m \bar{f}_n}) + 4 g_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} m_j m_{\bar{f}_n} s) \right) \\
&\quad + \frac{\mathcal{Y}_1(\tilde{f}_k, \tilde{f}_l, f_m, \bar{f}_n)}{8 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} - R_{f_m \tilde{f}_l})} \left( f_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} \right. \\
&\quad \left. (\tau_{f_m \bar{f}_n} - \omega_{\bar{f}_n f_m}) - 4 f_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} g_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} m_i m_{f_m} s - 4 f_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} g_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} \right. \\
&\quad \left. m_j m_{\bar{f}_n} s) - \frac{\mathcal{Z}_1(\tilde{f}_k, \tilde{f}_l, f_m, \bar{f}_n)}{8 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} - R_{f_m \tilde{f}_l})} \left( f_{\chi_i^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} \right) \right)
\end{aligned} \tag{3.116}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{u(\tilde{f}_k - \tilde{f}_l)} &= \frac{1}{8 b_{f_m \bar{f}_n} (R_{\bar{f}_n \tilde{f}_k} - R_{\bar{f}_n \tilde{f}_l})} \left( 2 b_{f_m \bar{f}_n} (R_{\bar{f}_n \tilde{f}_k} - R_{\bar{f}_n \tilde{f}_l}) f_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} \right) \\
&+ \frac{\mathcal{X}_1(\tilde{f}_k, \tilde{f}_l, \bar{f}_n, f_m)}{2 b_{f_m \bar{f}_n} (R_{\bar{f}_n \tilde{f}_k} - R_{\bar{f}_n \tilde{f}_l})} \left( g_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} m_i m_{\bar{f}_n} s (f_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} (\tau_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n}) \right. \\
&+ 4 g_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} m_j m_{f_m} s) \left. \right) + \frac{\mathcal{Y}_1(\tilde{f}_k, \tilde{f}_l, \bar{f}_n, f_m)}{8 b_{f_m \bar{f}_n} (R_{\bar{f}_n \tilde{f}_k} - R_{\bar{f}_n \tilde{f}_l})} \left( f_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} \right. \\
&(\omega_{f_m \bar{f}_n} - \tau_{\bar{f}_n f_m}) - 4 f_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} g_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} m_i m_{\bar{f}_n} s + 4 f_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} g_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} \\
&m_j m_{f_m} s) - \frac{\mathcal{Z}_1(\tilde{f}_k, \tilde{f}_l, \bar{f}_n, f_m)}{8 b_{f_m \bar{f}_n} (R_{\bar{f}_n \tilde{f}_k} - R_{\bar{f}_n \tilde{f}_l})} \left( f_{\chi_i^0 \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} f_{\chi_j^0 f_m}^{(\tilde{f}_k - \tilde{f}_l)} \right) \quad (3.117)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{tu(\tilde{f}_k - \tilde{f}_l)} &= -\frac{1}{4} J_{f_m \bar{f}_n}^{(1)(\tilde{f}_k - \tilde{f}_l)} + \frac{J_{f_m \bar{f}_n}^{(1)(\tilde{f}_k - \tilde{f}_l)}}{16 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} + R_{\bar{f}_n \tilde{f}_l} - 2 s l_i)} \left( \mathcal{Y}_2(\tilde{f}_k, \tilde{f}_l, f_m, \bar{f}_n) \right. \\
&(\omega_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n} - (3 R_{f_m \tilde{f}_k} L(\tilde{f}_k, f_m, \bar{f}_n) + R_{\bar{f}_n \tilde{f}_l} L(\tilde{f}_l, \bar{f}_n, f_m)) \tau_{f_m \bar{f}_n}) \\
&- (R_{f_m \tilde{f}_k} L(\tilde{f}_k, f_m, \bar{f}_n) + 3 R_{\bar{f}_n \tilde{f}_l} L(\tilde{f}_l, \bar{f}_n, f_m)) \tau_{\bar{f}_n f_m}) + 2 (R_{f_m \tilde{f}_k}^2 \\
&L(\tilde{f}_k, f_m, \bar{f}_n) + R_{\bar{f}_n \tilde{f}_l}^2 L(\tilde{f}_l, \bar{f}_n, f_m)) + 4 l_i s^2 (l_{f_m \bar{f}_n} L(\tilde{f}_k, f_m, \bar{f}_n) \\
&+ l_{\bar{f}_n f_m} L(\tilde{f}_l, \bar{f}_n, f_m)) \left. \right) + \frac{s (L(\tilde{f}_k, f_m, \bar{f}_n) + L(\tilde{f}_l, \bar{f}_n, f_m))}{4 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} + R_{\bar{f}_n \tilde{f}_l} - 2 s l_i)} (4 s (-d d_{f_m \bar{f}_n} \\
&J_{f_m \bar{f}_n}^{(1)(\tilde{f}_k - \tilde{f}_l)} + d_{f_m \bar{f}_n} J_{f_m \bar{f}_n}^{(4)(\tilde{f}_k - \tilde{f}_l)} m_i m_j + d J_{f_m \bar{f}_n}^{(2)(\tilde{f}_k - \tilde{f}_l)} m_{f_m} m_{\bar{f}_n} + J_{f_m \bar{f}_n}^{(3)(\tilde{f}_k - \tilde{f}_l)} \\
&m_i m_j m_{f_m} m_{\bar{f}_n}) - \frac{s}{4 b_{f_m \bar{f}_n} (R_{f_m \tilde{f}_k} + R_{\bar{f}_n \tilde{f}_l} - 2 s l_i)} \left( \mathcal{Y}_2(\tilde{f}_k, \tilde{f}_l, f_m, \bar{f}_n) \right. \\
&- \tau_{f_m \bar{f}_n} L(\tilde{f}_k, f_m, \bar{f}_n) + \tau_{\bar{f}_n f_m} L(\tilde{f}_l, \bar{f}_n, f_m) \left. \right) \left( -J_{f_m \bar{f}_n}^{(8)(\tilde{f}_k - \tilde{f}_l)} m_i m_{f_m} \right. \\
&+ J_{f_m \bar{f}_n}^{(5)(\tilde{f}_k - \tilde{f}_l)} m_j m_{f_m} + J_{f_m \bar{f}_n}^{(7)(\tilde{f}_k - \tilde{f}_l)} m_i m_{\bar{f}_n} - J_{f_m \bar{f}_n}^{(6)(\tilde{f}_k - \tilde{f}_l)} m_j m_{\bar{f}_n} \left. \right) \quad (3.118)
\end{aligned}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(\tilde{f}_k - \tilde{f}_l)} = \tilde{w}_{f_m \bar{f}_n}^{t(\tilde{f}_k - \tilde{f}_l)} + \tilde{w}_{f_m \bar{f}_n}^{u(\tilde{f}_k - \tilde{f}_l)} - (\tilde{w}_{f_m \bar{f}_n}^{tu(\tilde{f}_k - \tilde{f}_l)} + c.c) \quad (3.119)$$

- Higgs boson ( $H_k^0$ ,  $A_1^0$ ), Z boson and sfermion ( $\tilde{f}_l$ ) exchange:

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{st(H_k^0 - \tilde{f}_l)} &= -\frac{C_{H_k^0 f_m \bar{f}_n}}{16 b_{f_m \bar{f}_n} (s - m_{H_k^0}^2)} \left( J_{f_m \bar{f}_n}^{(5)(H_k^0 - \tilde{f}_l)} (4 b_{f_m \bar{f}_n} s + (\tau_{f_m \bar{f}_n} + \omega_{f_m \bar{f}_n}) l_i \right. \\
&L(\tilde{f}_l, f_m, \bar{f}_n) - 2 s (4 d (d_{f_m \bar{f}_n} - m_{f_m} m_{\bar{f}_n}) + R_{f_m \tilde{f}_l}) L(\tilde{f}_l, f_m, \bar{f}_n) \left. \right) \\
&- 8 s m_i m_j L(\tilde{f}_l, f_m, \bar{f}_n) J_{f_m \bar{f}_n}^{(6)(H_k^0 - \tilde{f}_l)} (d_{f_m \bar{f}_n} - m_{f_m} m_{\bar{f}_n}) - 2 m_j \\
&J_{f_m \bar{f}_n}^{(7)(H_k^0 - \tilde{f}_l)} \left( 2 b_{f_m \bar{f}_n} (m_{f_m} + m_{\bar{f}_n}) - ((m_{f_m} + m_{\bar{f}_n}) R_{f_m \tilde{f}_l} - 2 s l_i m_{f_m}) \right. \\
&L(\tilde{f}_l, f_m, \bar{f}_n) \left. \right) - 2 m_i J_{f_m \bar{f}_n}^{(8)(H_k^0 - \tilde{f}_l)} (2 b_{f_m \bar{f}_n} (m_{f_m} + m_{\bar{f}_n}) - ((m_{f_m} + m_{\bar{f}_n}) \\
&+ (m_{f_m} (\tau_{f_m \bar{f}_n} - \omega_{\bar{f}_n f_m}) - (m_{f_m} + m_{\bar{f}_n}) R_{f_m \tilde{f}_l}) + 2 s m_{\bar{f}_n} l_{f_m \bar{f}_n}) \\
&L(\tilde{f}_l, f_m, \bar{f}_n) \left. \right) \quad (3.120)
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{su(H_k^0 - \tilde{f}_l)} &= -\frac{C_{H_k^0 f_m \bar{f}_n}}{16 b_{f_m \bar{f}_n} (s - m_{H_k^0}^2)} \left( J_{\bar{f}_n f_m}^{(5)(H_k^0 - \tilde{f}_l)} (4 b_{f_m \bar{f}_n} s + (\tau_{\bar{f}_n f_m} + \omega_{\bar{f}_n f_m}) l_i \right. \\
&\quad \left. L(\tilde{f}_l, \bar{f}_n, f_m) - 2 s (4 d (d_{f_m \bar{f}_n} - m_{f_m} m_{\bar{f}_n}) + R_{\bar{f}_n \tilde{f}_l}) L(\tilde{f}_l, \bar{f}_n, f_m) \right) \\
&\quad - 8 s m_i m_j L(\tilde{f}_l, \bar{f}_n, f_m) J_{\bar{f}_n, f_m}^{(6)(H_k^0 - \tilde{f}_l)} (d_{f_m \bar{f}_n} - m_{f_m} m_{\bar{f}_n}) - 2 m_j \\
&\quad J_{\bar{f}_n f_m}^{(7)(H_k^0 - \tilde{f}_l)} \left( 2 b_{f_m \bar{f}_n} (m_{f_m} + m_{\bar{f}_n}) - ((m_{f_m} + m_{\bar{f}_n}) R_{\bar{f}_n \tilde{f}_l} - 2 s l_i m_{\bar{f}_n}) \right. \\
&\quad \left. L(\tilde{f}_l, \bar{f}_n, f_m) \right) - 2 m_i J_{\bar{f}_n f_m}^{(8)(H_k^0 - \tilde{f}_l)} \left( 2 b_{f_m \bar{f}_n} (m_{f_m} + m_{\bar{f}_n}) - ((m_{f_m} + m_{\bar{f}_n}) \right. \\
&\quad \left. + (m_{\bar{f}_n} (\tau_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n}) - (m_{f_m} + m_{\bar{f}_n}) R_{\bar{f}_n \tilde{f}_l}) + 2 s m_{f_m} l_{\bar{f}_n f_m}) \right. \\
&\quad \left. L(\tilde{f}_l, \bar{f}_n, f_m) \right) \Big) \tag{3.121}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{st(A_1^0 - \tilde{f}_l)} &= \frac{C_{A_1^0 f_m \bar{f}_n}}{8 b_{f_m \bar{f}_n} (s - m_{A_1^0}^2)} \left( -s J_{f_m \bar{f}_n}^{(9)(A_1^0 - \tilde{f}_l)} (-2 b_{f_m \bar{f}_n} + (-l_{f_m \bar{f}_n} l_i \right. \\
&\quad \left. + 4 d (d_{f_m \bar{f}_n} + m_{f_m} m_{\bar{f}_n}) + R_{f_m \tilde{f}_l}) L(\tilde{f}_l, f_m, \bar{f}_n) \right) + 4 m_i m_j s \\
&\quad J_{f_m \bar{f}_n}^{(10)(A_1^0 - \tilde{f}_l)} L(\tilde{f}_l, f_m, \bar{f}_n) (d_{f_m \bar{f}_n} + m_{f_m} m_{\bar{f}_n}) - m_j J_{f_m \bar{f}_n}^{(11)(A_1^0 - \tilde{f}_l)} \\
&\quad \left( (m_{f_m} - m_{\bar{f}_n}) (2 b_{f_m \bar{f}_n} - R_{f_m \tilde{f}_l}) L(\tilde{f}_l, f_m, \bar{f}_n) + 2 l_i m_{f_m} s L(\tilde{f}_l, f_m, \bar{f}_n) \right) \\
&\quad - m_i J_{f_m \bar{f}_n}^{(12)(A_1^0 - \tilde{f}_l)} \left( 2 b_{f_m \bar{f}_n} (m_{f_m} - m_{\bar{f}_n}) + L(\tilde{f}_l, f_m, \bar{f}_n) (m_{f_m} \right. \\
&\quad \left. (\tau_{f_m \bar{f}_n} - \omega_{\bar{f}_n f_m}) - R_{f_m \tilde{f}_l} (m_{f_m} - m_{\bar{f}_n}) - 2 l_{f_m \bar{f}_n} m_{\bar{f}_n} s) \right) \Big) \tag{3.122}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{su(A_1^0 - \tilde{f}_l)} &= \frac{C_{A_1^0 f_m \bar{f}_n}}{8 b_{f_m \bar{f}_n} (s - m_{A_1^0}^2)} \left( -s J_{\bar{f}_n f_m}^{(9)(A_1^0 - \tilde{f}_l)} (-2 b_{f_m \bar{f}_n} + (-l_{\bar{f}_n f_m} l_i \right. \\
&\quad \left. + 4 d (d_{f_m \bar{f}_n} + m_{f_m} m_{\bar{f}_n}) + R_{\bar{f}_n \tilde{f}_l}) L(\tilde{f}_l, \bar{f}_n, f_m) \right) + 4 m_i m_j s \\
&\quad J_{\bar{f}_n f_m}^{(10)(A_1^0 - \tilde{f}_l)} L(\tilde{f}_l, \bar{f}_n, f_m) (d_{f_m \bar{f}_n} + m_{f_m} m_{\bar{f}_n}) - m_j J_{\bar{f}_n f_m}^{(11)(A_1^0 - \tilde{f}_l)} \\
&\quad \left( (m_{\bar{f}_n} - m_{f_m}) (2 b_{f_m \bar{f}_n} - R_{\bar{f}_n \tilde{f}_l}) L(\tilde{f}_l, \bar{f}_n, f_m) + 2 l_i m_{\bar{f}_n} s L(\tilde{f}_l, \bar{f}_n, f_m) \right) \\
&\quad - m_i J_{\bar{f}_n f_m}^{(12)(A_1^0 - \tilde{f}_l)} \left( 2 b_{f_m \bar{f}_n} (m_{\bar{f}_n} - m_{f_m}) + L(\tilde{f}_l, \bar{f}_n, f_m) (m_{\bar{f}_n} \right. \\
&\quad \left. (\tau_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n}) - R_{\bar{f}_n \tilde{f}_l} (m_{\bar{f}_n} - m_{f_m}) - 2 s l_{\bar{f}_n f_m} m_{f_m}) \right) \Big) \tag{3.123}
\end{aligned}$$

$$\begin{aligned}
\tilde{\omega}_{f_m \bar{f}_n}^{st(Z-\tilde{f}_i)} &= \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(5)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(9)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} m_j m_{\bar{f}_n} (s - 4 M_Z^2) (2 b_{f_m \bar{f}_n} \\
&- R_{f_m \bar{f}_i} L(\tilde{f}_l, f_m, \bar{f}_n)) + \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(5)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(9)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_j m_{f_m} \left( (s - 2 M_Z^2) ((R_{f_m \bar{f}_i} - 2 l_i s) L(\tilde{f}_l, f_m, \bar{f}_n) - 2 b_{f_m \bar{f}_n}) \right. \\
&+ 2 s \tau_{\bar{f}_n f_m} L(\tilde{f}_l, f_m, \bar{f}_n) \left. \right) + \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(6)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(10)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_i m_{\bar{f}_n} \left( (s - 2 M_Z^2) ((R_{f_m \bar{f}_i} - 2 l_{f_m \bar{f}_n} s) L(\tilde{f}_l, f_m, \bar{f}_n) - 2 b_{f_m \bar{f}_n}) \right. \\
&+ 2 s \omega_{f_m \bar{f}_n} L(\tilde{f}_l, f_m, \bar{f}_n) \left. \right) + \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(6)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(10)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_i m_{f_m} (s - 4 M_Z^2) \left( (\tau_{f_m \bar{f}_n} - \omega_{\bar{f}_n f_m} - R_{f_m \bar{f}_i}) L(\tilde{f}_l, f_m, \bar{f}_n) + 2 b_{f_m \bar{f}_n} \right) \\
&- \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(7)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(11)(Z-\tilde{f}_i)}}{32 s b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} (8 b_{f_m \bar{f}_n} M_Z^2 \\
&(\omega_{\bar{f}_n f_m} + R_{f_m \bar{f}_i}) + 2 b_{f_m \bar{f}_n} s (\tau_{\bar{f}_n f_m} - \tau_{f_m \bar{f}_n} + \omega_{f_m \bar{f}_n} - \omega_{\bar{f}_n f_m}) \\
&- (4 M_Z^2 R_{f_m \bar{f}_i} (R_{f_m \bar{f}_i} - \tau_{f_m \bar{f}_n}) + \tau_{f_m \bar{f}_n}^2 s - 3 \tau_{\bar{f}_n f_m} \omega_{f_m \bar{f}_n} s \\
&+ s \tau_{\bar{f}_n f_m} R_{f_m \bar{f}_i} + \tau_{f_m \bar{f}_n} R_{f_m \bar{f}_i} s - \tau_{f_m \bar{f}_n} (\tau_{\bar{f}_n f_m} + \omega_{f_m \bar{f}_n} + R_{f_m \bar{f}_i}) s \\
&- 16 d d_{f_m \bar{f}_n} s^3 - \omega_{\bar{f}_n f_m} R_{f_m \bar{f}_i} (s - 4 M_Z^2) + 8 d_{f_m \bar{f}_n} s^2 l_i l_{ia} \\
&+ 8 d s^2 l_{f_m \bar{f}_n} l_{\bar{f}_n f_m}) L(\tilde{f}_l, f_m, \bar{f}_n) \\
&+ \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(7)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(11)(Z-\tilde{f}_i)}}{4 b_{f_m \bar{f}_n} M_Z^2} m_{f_m} m_{\bar{f}_n} s L(\tilde{f}_l, f_m, \bar{f}_n) \\
&(l_i l_{ia} - 2 d (s - 2 M_Z^2)) + \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(8)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(12)(Z-\tilde{f}_i)}}{4 b_{f_m \bar{f}_n} M_Z^2} \\
&(l_{f_m \bar{f}_n} l_{\bar{f}_n f_m} - 2 d_{f_m \bar{f}_n} (s - 2 M_Z^2)) \\
&+ \frac{C_{Zf_m \bar{f}_n}^V J_{f_m \bar{f}_n}^{(8)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{f_m \bar{f}_n}^{(12)(Z-\tilde{f}_i)}}{2 b_{f_m \bar{f}_n} M_Z^2} m_i m_j m_{f_m} m_{\bar{f}_n} (s - 4 M_Z^2) \\
&L(\tilde{f}_l, \bar{f}_n, f_m) \tag{3.124}
\end{aligned}$$

$$\begin{aligned}
\tilde{w}_{f_m \bar{f}_n}^{su(Z-\tilde{f}_i)} &= \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(5)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(9)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} m_j m_{f_m} (s - 4 M_Z^2) (2 b_{f_m \bar{f}_n} \\
&- R_{\bar{f}_n \tilde{f}_i} L(\tilde{f}_l, \bar{f}_n, f_m)) + \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(5)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(9)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_j m_{\bar{f}_n} \left( (s - 2 M_Z^2) ((R_{\bar{f}_n \tilde{f}_i} - 2 l_i s) L(\tilde{f}_l, \bar{f}_n, f_m) - 2 b_{f_m \bar{f}_n}) \right. \\
&+ 2 s \tau_{f_m \bar{f}_n} L(\tilde{f}_l, \bar{f}_n, f_m) \left. \right) + \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(6)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(10)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_i m_{f_m} \left( (s - 2 M_Z^2) ((R_{\bar{f}_n \tilde{f}_i} - 2 l_{\bar{f}_n f_m} s) L(\tilde{f}_l, \bar{f}_n, f_m) - 2 b_{f_m \bar{f}_n}) \right. \\
&+ 2 s \omega_{\bar{f}_n f_m} L(\tilde{f}_l, \bar{f}_n, f_m) \left. \right) + \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(6)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(10)(Z-\tilde{f}_i)}}{8 b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} \\
&m_i m_{\bar{f}_n} (s - 4 M_Z^2) \left( (\tau_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n} - R_{\bar{f}_n \tilde{f}_i}) L(\tilde{f}_l, \bar{f}_n, f_m) + 2 b_{f_m \bar{f}_n} \right) \\
&- \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(7)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(11)(Z-\tilde{f}_i)}}{32 s b_{f_m \bar{f}_n} M_Z^2 (s - M_Z^2)} (8 b_{f_m \bar{f}_n} M_Z^2 \\
&(\omega_{f_m \bar{f}_n} + R_{\bar{f}_n \tilde{f}_i}) + 2 b_{f_m \bar{f}_n} s (\tau_{f_m \bar{f}_n} - \tau_{\bar{f}_n f_m} + \omega_{\bar{f}_n f_m} - \omega_{f_m \bar{f}_n}) \\
&- (4 M_Z^2 R_{\bar{f}_n \tilde{f}_i} (R_{\bar{f}_n \tilde{f}_i} - \tau_{\bar{f}_n f_m}) + \tau_{\bar{f}_n f_m}^2 s - 3 \tau_{f_m \bar{f}_n} \omega_{\bar{f}_n f_m} s \\
&+ s \tau_{f_m \bar{f}_n} R_{\bar{f}_n \tilde{f}_i} + \tau_{\bar{f}_n f_m} R_{\bar{f}_n \tilde{f}_i} s - \tau_{\bar{f}_n f_m} (\tau_{f_m \bar{f}_n} + \omega_{\bar{f}_n f_m} + R_{\bar{f}_n \tilde{f}_i}) s \\
&- 16 d d_{f_m \bar{f}_n} s^3 - \omega_{f_m \bar{f}_n} R_{\bar{f}_n \tilde{f}_i} (s - 4 M_Z^2) + 8 d_{f_m \bar{f}_n} s^2 l_i l_{ia} \\
&+ 8 d s^2 l_{f_m \bar{f}_n} l_{\bar{f}_n f_m}) L(\tilde{f}_l, \bar{f}_n, f_m) \left. \right) \\
&+ \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(7)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(11)(Z-\tilde{f}_i)}}{4 b_{f_m \bar{f}_n} M_Z^2} m_{f_m} m_{\bar{f}_n} s L(\tilde{f}_l, \bar{f}_n, f_m) \\
&(l_i l_{ia} - 2 d (s - 2 M_Z^2)) + \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(8)(Z-\tilde{f}_i)} + C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(12)(Z-\tilde{f}_i)}}{4 b_{f_m \bar{f}_n} M_Z^2} \\
&(l_{f_m \bar{f}_n} l_{\bar{f}_n f_m} - 2 d_{f_m \bar{f}_n} (s - 2 M_Z^2)) \\
&+ \frac{C_{Zf_m \bar{f}_n}^V J_{\bar{f}_n f_m}^{(8)(Z-\tilde{f}_i)} - C_{Zf_m \bar{f}_n}^A J_{\bar{f}_n f_m}^{(12)(Z-\tilde{f}_i)}}{2 b_{f_m \bar{f}_n} M_Z^2} m_i m_j m_{f_m} m_{\bar{f}_n} (s - 4 M_Z^2) \\
&L(\tilde{f}_l, \bar{f}_n, f_m) \tag{3.125}
\end{aligned}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(H_k^0 - \tilde{f}_i)} = (\tilde{w}_{f_m \bar{f}_n}^{st(H_k^0 - \tilde{f}_i)} - \tilde{w}_{f_m \bar{f}_n}^{su(H_k^0 - \tilde{f}_i)}) + c.c. \tag{3.126}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(A_1^0 - \tilde{f}_i)} = (\tilde{w}_{f_m \bar{f}_n}^{st(A_1^0 - \tilde{f}_i)} - \tilde{w}_{f_m \bar{f}_n}^{su(A_1^0 - \tilde{f}_i)}) + c.c. \tag{3.127}$$

$$\tilde{w}_{f_m \bar{f}_n}^{(Z - \tilde{f}_i)} = (\tilde{w}_{f_m \bar{f}_n}^{s(Z - \tilde{f}_i)} - \tilde{w}_{f_m \bar{f}_n}^{su(Z - \tilde{f}_i)}) + c.c. \tag{3.128}$$

where  $\tilde{f}_k$  is a sfermion mass eigen stat with  $k=1, \dots, p$ ; for squarks and charged leptons  $p=6$  and for  $s$  neutrinos  $p=3$

## 4 summary

We have presented, for the first time the full neutralino neutralino annihilation cross section with flavour violation. And we compared our result with the ones in NIhei et al. We hope to extend this analysis to neutralino-slepton co-annihilation where we expect large changes and also to sommerfeld enhancement.

## A Mixing matrices and Coupling constants

In this appendix , we collect various couplings involving mixing matrices are listed. Feynman diagrams corresponding to all the annihilation channels are given in [18]. We closely follow Ref.[18] for notational consistency. For the sake of ease and completeness we present in this section relevant spectrum which is used in calculations in this paper. And for more details we urge readers to refer to [18]

- Neutral Higgs scalars. If the Lagrangian contains only real parameters the neutral Higgs particles have well defined CP eigenvalues - two of them are scalars, the other two are pseudoscalars. This is no longer true if some parameters are complex. Nevertheless, in both cases it is convenient to divide neutral Higgses into two classes.

i) Scalar particles  $H_i^0$  ,  $i = 1, 2$ , defined as:

$$\sqrt{2}\Re H_i^0 = Z_R^{ij} H_j^0 + v_i \text{ (no sum over } i \text{)}$$

ii) ‘‘Pseudoscalar’’ particles  $A_i^0$  ,  $i = 1, 2$ :

$$\sqrt{2}\Im H_i^0 = Z_H^{ij} A_j^0 \text{ (no sum over } i \text{)}$$

$A_1^0(\equiv A^0)$  has the mass  $M_A^2 = m_{H_1}^2 + m_{H_2}^2 + 2|\mu|^2$ ,  $A_2^0(\equiv G^0)$  is the massless Goldstone boson which disappears in the unitary gauge

$$Z_H = \begin{pmatrix} \sin\beta & -\cos\beta \\ \cos\beta & \sin\beta \end{pmatrix} \quad Z_R = \begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix} \quad (\text{A.1})$$

where:

$$\tan\beta = \frac{v_2}{v_1} \quad (\text{A.2})$$

$$\tan 2\alpha = \tan 2\beta \frac{M_A^2 + M_Z^2}{M_A^2 - M_Z^2} \quad (\text{A.3})$$

$$(\text{A.4})$$

$$A_M^{ij} = Z_R^{1i} Z_H^{1j} - Z_R^{2i} Z_H^{2j} \quad A_R^{ij} = Z_R^{1i} Z_R^{1j} - Z_R^{2i} Z_R^{2j} \quad (\text{A.5})$$

$$A_H^{ij} = Z_H^{1i} Z_H^{1j} - Z_H^{2i} Z_H^{2j} \quad B_R^i = v_1 Z_R^{1i} - v_2 Z_R^{2i} \quad (\text{A.6})$$

$$A_P^{ij} = Z_R^{1i} Z_H^{2j} + Z_R^{2i} Z_H^{1j} \quad (\text{A.7})$$

- Charged Higgs scalars. Four charged Higgs scalars exist, two of them with the mass

$$M_{H_1^\pm}^2 = M_W^2 + m_{H_1}^2 + m_{H_2}^2 + 2|\mu|^2 \quad (\text{A.8})$$

and the other two massless. In the physical (unitary) gauge  $H_2^\pm (\equiv G^\pm)$  are eaten by  $W$  bosons and disappear from the Lagrangian. Fields  $H_1^+$  and  $H_2^+$  are related to the initial Higgs fields by the rotation matrix  $Z_H$

- Charginos. Four 2-component spinors  $(\lambda_A^1, \lambda_A^2, \Psi_{H_2}^1, \Psi_{H_1}^2)$  combine to give two 4-component Dirac fermions  $\chi_1, \chi_2$  corresponding to two physical charginos. The chargino mixing matrices  $Z_+$  and  $Z_-$  are defined by the condition:

$$(Z_-)^T \begin{pmatrix} M_2 & \frac{ev_2}{\sqrt{2}s_W} \\ \frac{ev_1}{\sqrt{2}s_W} & \mu \end{pmatrix} Z_+ = \begin{pmatrix} M_{\chi_1} & 0 \\ 0 & M_{\chi_2} \end{pmatrix} \quad (\text{A.9})$$

The unitary matrices  $Z_-, Z_+$  are not uniquely specified - by changing their relative phases and the ordering of the eigenvalues it is possible to choose  $M_{\chi_i}$  to be positive and  $M_{\chi_2} > M_{\chi_1}$ . The fields  $\chi_i$  are related to the initial spinors as below:

$$\Psi_{H_1}^2 = Z_+^{2i} \kappa_i^+ \quad (\text{A.10})$$

$$\Psi_{H_2}^1 = Z_-^{2i} \kappa_i^- \quad \chi_i = \begin{pmatrix} \kappa_i^+ \\ \bar{\kappa}_i^- \end{pmatrix} \quad (\text{A.11})$$

$$\lambda_A^\pm \equiv \frac{\lambda_A^1 \mp i\lambda_A^2}{\sqrt{2}} = iZ_\pm^{1i} \kappa_i^\pm \quad (\text{A.12})$$

- Four 2-component spinors  $(\lambda_B, \lambda_A^3, \Psi_{H_1}^1, \Psi_{H_2}^2)$  combine into four Majorana fermions  $\chi_i^0, i = 1 \dots 4$ , called neutralinos. The formulas for mixing and mass matrices are the following:

$$Z_N^T \begin{pmatrix} M_1 & 0 & \frac{-ev_1}{2c_W} & \frac{ev_2}{2c_W} \\ 0 & M_2 & \frac{ev_1}{2s_W} & \frac{-ev_2}{2s_W} \\ \frac{-ev_1}{2c_W} & \frac{ev_1}{2s_W} & 0 & -\mu \\ \frac{ev_2}{2c_W} & \frac{-ev_2}{2s_W} & -\mu & 0 \end{pmatrix} Z_N = \begin{pmatrix} M_{\chi_1^0} & & & 0 \\ & \ddots & & \\ 0 & & & M_{\chi_4^0} \end{pmatrix} \quad (\text{A.13})$$

$$\lambda_B = iZ_N^{1i} \kappa_i^0 \quad (\text{A.14})$$

$$\lambda_A^3 = iZ_N^{2i} \kappa_i^0 \quad (\text{A.15})$$

$$\Psi_{H_1}^1 = Z_N^{3i} \kappa_i^0 \quad \chi_i^0 = \begin{pmatrix} \kappa_i^0 \\ \bar{\kappa}_i^0 \end{pmatrix} \quad (\text{A.16})$$

$$\Psi_{H_2}^2 = Z_N^{4i} \kappa_i^0 \quad (\text{A.17})$$



$$C_{A_i^0 A_j^0 H_k^0} = -\frac{ie^2}{4s_W^2 c_W^2} A_R^{ij} B_R^k \quad (\text{A.18})$$

$$C_{H_i^0 H_j^0 H_k^0} = -\frac{ie^2}{4s_W^2 c_W^2} (A_R^{ij} B_R^k + A_R^{jk} B_R^i + A_R^{ki} B_R^j) \quad (\text{A.19})$$

$$C_{H_i^0 H_1^+ H_1^-} = -i \left( \frac{e^2}{4s_W^2 c_W^2} A_H^{11} B_R^i + \frac{e M_W}{s_W} A_P^{i1} \right) \quad (\text{A.20})$$

$$C_{H_i^0 A_j^0 Z} = \frac{e}{2s_W c_W} A_M^{ij} \quad (\text{A.21})$$

$$C_{H_1^+ H_1^- Z} = ie \frac{c_W^2 - s_W^2}{2s_W c_W} \quad (\text{A.22})$$

$$C_{A_1^0 W^+ H^-} = -\frac{e}{2s_W} \quad (\text{A.23})$$

$$C_{H_i^0 W^+ H^-} = -\frac{e}{2s_W} A_M^{i1} \quad (\text{A.24})$$

$$C_{H_i^0 ZZ} = \frac{ie^2}{2s_W^2 c_W^2} g^{\mu\nu} C_R^i \quad (\text{A.25})$$

$$C_{H_k^0 ZZ} = \frac{ie^2}{2s_W^2 c_W^2} g^{\mu\nu} [v_1 Z_R^{1k} + v_2 Z_R^{2k}] \quad (\text{A.26})$$

$$C_{H_k^0 WW} = \frac{ie^2}{2s_W^2} g^{\mu\nu} [v_1 Z_R^{1k} + v_2 Z_R^{2k}] \quad (\text{A.27})$$

$$C_{Z f_m \bar{f}_n}^V = \frac{ie^2}{4s_W^2 c_W^2} (1 - 2s_W^2) \quad (\text{A.28})$$

$$C_{Z f_m \bar{f}_n}^A = \frac{ie^2}{4s_W^2 c_W^2} \quad (\text{A.29})$$

$$C_{\chi_i^0 \chi_l^0 H_m^0}^L = \frac{e}{2s_W c_W} [(Z_R^{1m} Z_N^{3i} - Z_R^{2m} Z_N^{4i})(Z_N^{1l} s_W - Z_N^{2l} c_W)] \quad (\text{A.30})$$

$$C_{\chi_i^0 \chi_l^0 H_m^0}^R = \frac{e}{2s_W c_W} [(Z_R^{1m} Z_N^{3l*} - Z_R^{2m} Z_N^{4l*})(Z_N^{1i*} s_W - Z_N^{2i*} c_W)] \quad (\text{A.31})$$

$$C_{\chi_l^0 \chi_j^0 A_n^0}^L = \frac{ie}{2s_W c_W} [(Z_H^{1n} Z_N^{3l} - Z_H^{2n} Z_N^{4l})(Z_N^{1j} s_W - Z_N^{2j} c_W)] \quad (\text{A.32})$$

$$C_{\chi_l^0 \chi_j^0 A_n^0}^R = \frac{-ie}{2s_W c_W} [(Z_H^{1n} Z_N^{3j*} - Z_H^{2n} Z_N^{4j*})(Z_N^{1l*} s_W - Z_N^{2l*} c_W)] \quad (\text{A.33})$$

$$C_{\chi_i^0 \chi_j^0 Z}^L = \frac{e}{4s_W c_W} (Z_N^{4j*} Z_N^{4i} - Z_N^{3j*} Z_N^{3i}) P_L \quad (\text{A.34})$$

$$C_{\chi_i^0 \chi_j^0 Z}^R = \frac{-e}{4s_W c_W} (Z_N^{4j} Z_N^{4i*} - Z_N^{3j} Z_N^{3i*}) P_R \quad (\text{A.35})$$

$$C_{\chi_i^0 \chi_l^0 Z}^L = \frac{ie}{2s_W c_W} [(Z_N^{4i*} Z_N^{4l} - Z_N^{3i*} Z_N^{3l})] \quad (\text{A.36})$$

$$C_{\chi_i^0 \chi_l^0 Z}^R = \frac{ie}{2s_W c_W} [(Z_N^{3i} Z_N^{3l*} - Z_N^{4i} Z_N^{4l*})] \quad (\text{A.37})$$

$$C_{\chi_i^0 \chi_j^0 H_k^+}^L = \frac{ie}{s_W c_W} \left[ Z_H^{1k} \left( \frac{1}{\sqrt{2}} Z_-^{2j} (Z_N^{1i} s_W + Z_N^{2i} c_W) - Z_-^{1j} Z_N^{3i} c_W \right) \right] \quad (\text{A.38})$$

$$C_{\chi_i^0 \chi_j^0 H_k^+}^R = -\frac{ie}{s_W c_W} \left[ Z_H^{2k} \left( \frac{1}{\sqrt{2}} Z_+^{2j*} (Z_N^{1i*} s_W + Z_N^{2i*} c_W) + Z_+^{1j*} Z_N^{4i*} c_W \right) \right] \quad (\text{A.39})$$

$$C_{\chi_i^0 \chi_l^{\pm} W}^L = \frac{ie}{s_W} \left[ (Z_N^{2l} Z_+^{1i*} - \frac{1}{\sqrt{2}} Z_N^{4l} Z_+^{2i*}) \right] \quad (\text{A.40})$$

$$C_{\chi_i^0 \chi_l^{\pm} W}^R = \frac{ie}{s_W} \left[ (Z_N^{2l*} Z_-^{1i} + \frac{1}{\sqrt{2}} Z_N^{4l*} Z_-^{2i}) \right] \quad (\text{A.41})$$



## B Auxilary functions

In this appendix, we collected various kinetic functions and auxilary functions used in the main text.

$$\begin{aligned}
l_i &= m_i^2 - m_j^2 + s \\
l_{ia} &= m_j^2 - m_i^2 + s \\
l_{\phi_1\phi_2} &= m_1^2 - m_2^2 + s \\
\tau_{\phi_1\phi_2} &= l_i (m_1^2 - m_2^2 + s) \\
\omega_{\phi_1\phi_2} &= l_{ia} (m_1^2 - m_2^2 + s) \\
R_{\phi_1\phi_2} &= 2(m_1^2 - m_2^2 + m_i^2)s \\
b_{\phi_1\phi_2} &= \sqrt{s^2 + m_i^4 + m_j^4 - 2s(m_i^2 + m_j^2) - 2m_i^2 m_j^2} \\
&\quad \sqrt{s^2 + m_1^4 + m_2^4 - 2s(m_1^2 + m_2^2) - 2m_1^2 m_2^2} \\
d &= \frac{1}{2} (s - (m_i^2 + m_j^2)) \\
d_{\phi_1\phi_2} &= \frac{1}{2} (s - (m_1^2 + m_2^2))
\end{aligned}$$

$$L(\chi, \phi_1, \phi_2) = \log \frac{R_{\phi_1\chi} + b_{\phi_1\phi_2} - \tau_{\phi_1\phi_2}}{R_{\phi_1\chi} - b_{\phi_1\phi_2} - \tau_{\phi_1\phi_2}}$$

$$\mathcal{A}(\chi, \phi_1, \phi_2) = \operatorname{arctanh} \frac{b_{\phi_1\phi_2}}{R_{\phi_1\chi} - \tau_{\phi_1\phi_2}}$$

$$\mathcal{B}(\chi, \phi_1, \phi_2) = R_{\phi_1\chi} - \tau_{\phi_1\phi_2} - \omega_{\phi_1\phi_2}$$

$$\mathcal{C}(\chi, \phi_1, \phi_2) = R_{\phi_1\chi} - \tau_{\phi_1\phi_2} + \omega_{\phi_2\phi_1}$$

$$\rho(\chi, \phi_1, \phi_2) = R_{\phi_1\chi} - \omega_{\phi_1\phi_2}$$

$$\begin{aligned}
\mathcal{X}(\phi_1, \phi_2, \chi_1) &= -8s^2 (2d d_{\phi_1\phi_2} - (2d + l_{ia} m_i^2) + 2R_{\phi_1\chi_1}^2 - R_{\phi_1\chi_1} (3\tau_{\phi_1\phi_2} + \tau_{\phi_2\phi_1} \\
&\quad + \omega_{\phi_1\phi_2} - \omega_{\phi_2\phi_1}) + \tau_{\phi_1\phi_2} (\tau_{\phi_1\phi_2} + \tau_{\phi_2\phi_1} + \omega_{\phi_1\phi_2} + \omega_{\phi_2\phi_1})
\end{aligned}$$

$$\mathcal{Q}_1(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^4 L(\chi_1, \phi_1, \phi_2) - R_{\phi_1\chi_2}^4 L(\chi_2, \phi_1, \phi_2)$$

$$\mathcal{R}_1(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^3 L(\chi_1, \phi_1, \phi_2) - R_{\phi_1\chi_2}^3 L(\chi_2, \phi_1, \phi_2)$$

$$\mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^2 L(\chi_1, \phi_1, \phi_2) - R_{\phi_1\chi_2}^2 L(\chi_2, \phi_1, \phi_2)$$

$$\mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1} L(\chi_1, \phi_1, \phi_2) - R_{\phi_1\chi_2} L(\chi_2, \phi_1, \phi_2)$$

$$\mathcal{X}_1(\chi_1, \chi_2, \phi_1, \phi_2) = L(\chi_1, \phi_1, \phi_2) - L(\chi_2, \phi_1, \phi_2)$$

$$\mathcal{Q}_2(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^4 L(\chi_1, \phi_1, \phi_2) - R_{\phi_2\chi_2}^4 L(\chi_2, \phi_2, \phi_1)$$

$$\mathcal{R}_2(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^3 L(\chi_1, \phi_1, \phi_2) - R_{\phi_2\chi_2}^3 L(\chi_2, \phi_2, \phi_1)$$

$$\mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1}^2 L(\chi_1, \phi_1, \phi_2) - R_{\phi_2\chi_2}^2 L(\chi_2, \phi_2, \phi_1)$$

$$\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2) = R_{\phi_1\chi_1} L(\chi_1, \phi_1, \phi_2) - R_{\phi_2\chi_2} L(\chi_2, \phi_2, \phi_1)$$

$$\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2) = L(\chi_1, \phi_1, \phi_2) - L(\chi_2, \phi_2, \phi_1)$$

$$\begin{aligned}
\beta_1(\phi_1, \phi_2, \chi_1, \chi_2) &= -4d(m_1^2 - m_i^2) - \frac{1}{4s}(s + 8m_i^2\omega_{\phi_1\phi_2}) - \frac{1}{4b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_1\chi_2})} \\
&\quad [R_{\phi_1\chi_2}\mathcal{B}(\chi_2, \phi_1, \phi_2)\mathcal{A}(\chi_2, \phi_1, \phi_2) - \{\chi_2 \rightarrow \chi_1\}] \\
\beta_2(\phi_1, \phi_2, \chi_1, \chi_2) &= 4(m_1^2 + m_i^2) + \frac{4s}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_1\chi_2})} \\
&\quad [R_{\phi_1\chi_2}\mathcal{B}(\chi_1, \phi_1, \phi_2)\mathcal{A}(\chi_1, \phi_1, \phi_2) - \{\chi_1 \rightarrow \chi_2\}] \\
\beta_3(\phi_1, \phi_2, \chi_1, \chi_2) &= 4m_i^2 + \frac{s}{2b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_1\chi_2})} \\
&\quad [R_{\phi_1\chi_2}\mathcal{B}(\chi_1, \phi_1, \phi_2)\mathcal{A}(\chi_1, \phi_1, \phi_2) - \{\chi_1 \rightarrow \chi_2\}] \\
J_1(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{L(\chi_1, \phi_1, \phi_2) + L(\chi_2, \phi_2, \phi_1)}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} + R_{\phi_2\chi_2} - 2s)} \\
J_2(\phi_1, \phi_2, \chi_1, \chi_2) &= -\frac{1}{4} + \frac{\mathcal{X}(\phi_1, \phi_2, \chi_1)L(\chi_1, \phi_1, \phi_2) + \mathcal{X}(\phi_2, \phi_1, \chi_2)L(\chi_2, \phi_2, \phi_1)}{16b_{\phi_1\phi_2}(R_{\phi_1\chi_1} + R_{\phi_2\chi_2} - 2s)} \\
J_3(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{1}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} + R_{\phi_2\chi_2} - 2s)}(R_{\phi_2\chi_2} - 4m_i^2s)L(\chi_2, \phi_2, \phi_1) \\
&\quad - (R_{\phi_1\chi_1} + 4m_i^2s - 2sl_i)L(\chi_1, \phi_1, \phi_2) \\
J_4(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{1}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} + R_{\phi_2\chi_2} - 2s)}((\mathcal{C}(\chi_1, \phi_1, \phi_2) - 4ds)L(\chi_1, \phi_1, \phi_2) - \\
&\quad 2(4ds + \mathcal{B}(\chi_2, \phi_2, \phi_1))\mathcal{A}(\chi_2, \phi_2, \phi_1)) \\
J_5(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{1}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} + R_{\phi_2\chi_2} - 2s)}((4ds + \mathcal{B}(\chi_1, \phi_1, \phi_2))L(\chi_1, \phi_1, \phi_2) \\
&\quad - 2(\mathcal{C}(\chi_2, \phi_2, \phi_1) - 4ds)\mathcal{A}(\chi_2, \phi_2, \phi_1)) \\
J_6(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_j m_{\chi_2}}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})}(2b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2}) - 8m_i^2m_1^2s^2 \\
&\quad \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2) + 6m_1^2s\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2) - \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)) \\
J_7(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{3m_j m_{\chi_1} s}{4b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})}(-2l_i s + 4m_i^2s)\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
&\quad + \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
J_8(\phi_1, \phi_2, \chi_1, \chi_2) &= -\frac{3m_i m_{\chi_2} s}{4b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})}((4ds - 2l_{\phi_1\phi_2}s)\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
&\quad + \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)) \\
J_9(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_i m_{\chi_1}}{4m_1^2} + \frac{m_i m_{\chi_1} \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})}(2m_1^2s(\tau_{\phi_1\phi_2} - \tau_{\phi_2\phi_1}) \\
&\quad - 8d_{\phi_1\phi_2}(l_{\phi_1\phi_2} - m_1^2)s^2) + \frac{m_i m_{\chi_1} \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
&\quad (4d_{\phi_1\phi_2}s + 2l_{\phi_1\phi_2}s - 2m_1^2s) - \frac{m_i m_{\chi_1} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})}
\end{aligned}$$

$$\begin{aligned}
J_{10}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{1}{2} + \frac{\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-b_{\phi_1\phi_2} + s^2(4\tau_{\phi_1\phi_2} + 3\omega_{\phi_1\phi_2} \\
&\quad - \omega_{\phi_2\phi_1})m_i^2m_1^2 + (4d_{\phi_1\phi_2}l_{\phi_1\phi_2}m_i^2 + (-3\tau_{\phi_1\phi_2} + 4d(d_{\phi_1\phi_2} + 2l_i \\
&\quad - 3m_i^2))m_1^2)s) - \frac{\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{32b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})s} (2m_1^2s(-7\tau_{\phi_1\phi_2} \\
&\quad - 3(\tau_{\phi_2\phi_1} + \omega_{\phi_1\phi_2}) + \omega_{\phi_2\phi_1}) \\
&\quad + 16(d + m_i^2)(d_{\phi_1\phi_2} + 2m_1^2)s^2) + \frac{\mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)(4s - 8m_1^2s)}{32b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})s} \\
J_{11}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_{\chi_1}m_{\chi_2}}{4m_1^2} + \frac{ds^2m_{\chi_1}m_{\chi_2}\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
&\quad + \frac{l_{\phi_1\phi_2}sm_{\chi_1}m_{\chi_2}\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} - \frac{m_{\chi_1}m_{\chi_2}\mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
J_{12}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_im_j}{4m_1^2} + \frac{s^2m_im_j\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{2b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-6d_{\phi_1\phi_2} + l_i - 2m_i^2) \\
&\quad + \frac{sm_im_j\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{2b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (d_{\phi_1\phi_2} + m_1^2) \\
&\quad - \frac{m_im_j\mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^2(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
J_{13}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{3m_im_jm_{\chi_1}m_{\chi_2}s^2\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1\phi_2}(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
J_{14}(\phi_1, \phi_2, \chi_1, \chi_2) &= -\frac{b_{\phi_1\phi_2}^2}{48m_1^4s^2} - \frac{1}{16m_1^4s^2} (R_{\phi_1\chi_1}^2 + R_{\phi_2\chi_2}^2 + R_{\phi_1\chi_1}R_{\phi_2\chi_2} - 3l_is \\
&\quad (R_{\phi_1\chi_1} + R_{\phi_2\chi_2})) - \frac{1}{16m_1^4} (-8dd_{\phi_1\phi_2} + 2l_i^2 + l_ila + 8d_{\phi_1\phi_2}m_i^2 \\
&\quad - 4m_1^2(l_{ia} + m_1^2 + 2s)) + \frac{s^2\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2}m_1^4(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-2l_i^2m_1^2s \\
&\quad + 4d_{\phi_1\phi_2}m_i^2(-4l_{ia}m_1^2 + l_is) - m_1^4(-12l_{ia}m_i^2 + 2l_is)) \\
&\quad - \frac{ds^2\mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1\phi_2}m_1^4(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-2d_{\phi_1\phi_2}^2m_i^2 + d_{\phi_1\phi_2}l_im_1^2 + (5d_{\phi_1\phi_2} \\
&\quad - 4l_i + 5m_i^2)m_1^4) - \frac{l_is\mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{16b_{\phi_1\phi_2}m_1^4(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \\
&\quad (-8dd_{\phi_1\phi_2} + l_i^2 + l_ila + 8d_{\phi_1\phi_2}m_i^2 - 4m_1^2(l_{ia} + m_1^2 + 2s)) \\
&\quad + \frac{\mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{16b_{\phi_1\phi_2}m_1^4(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-4dd_{\phi_1\phi_2} + 2l_i^2 + 4d_{\phi_1\phi_2}m_i^2 + l_is \\
&\quad - 2m_1^2(l_{ia} + m_1^2 + 2s)) - \frac{4l_i\mathcal{R}_1(\chi_1, \chi_2, \phi_1, \phi_2) - \mathcal{Q}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{32b_{\phi_1\phi_2}m_1^4(R_{\phi_1\chi_1} - R_{\phi_2\chi_2})s^2}
\end{aligned}$$

$$\begin{aligned}
J_{15}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_{\chi_1} m_{\chi_2} (m_1^2 - d_{\phi_1 \phi_2})}{2 m_1^2} - \frac{m_{\chi_1} m_{\chi_2} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (l_i (l_i - l_{ia}) \\
&\quad m_1^2 + 12 d m_1^4 + d_{\phi_1 \phi_2} (8 d d_{\phi_1 \phi_2} - 2 l_i s)) \\
&\quad - \frac{l_i m_{\chi_1} m_{\chi_2} s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{2 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (d_{\phi_1 \phi_2} - m_1^2) \\
&\quad + \frac{m_{\chi_1} m_{\chi_2} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (d_{\phi_1 \phi_2} - m_1^2) \\
J_{16}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_i m_j (-d_{\phi_1 \phi_2} + m_1^2)}{2 m_1^4} + \frac{m_i m_j s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{2 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (4 d_{\phi_1 \phi_2}^2 m_i^2 \\
&\quad - l_i^2 m_1^2 - (6 d_{\phi_1 \phi_2} + l_i - 6 m_i^2) m_1^4 + 4 d_{\phi_1 \phi_2} m_1^2 s) \\
&\quad + \frac{m_i m_j (d_{\phi_1 \phi_2} - m_1^2)}{4 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (-2 l_i s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
&\quad + \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)) \\
J_{17}(\phi_1, \phi_2, \chi_1, \chi_2) &= -\frac{m_i m_j m_{c j i_1} m_{\chi_2} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (-2 d_{\phi_1 \phi_2}^2 + 5 m_1^4) \\
J_{18}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_j m_{\chi_2} (2 d_{\phi_1 \phi_2} + m_1^2)}{4 m_1^4} + \frac{m_j m_{\chi_2} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} \\
&\quad (-2 d_{\phi_1 \phi_2}^2 m_i^2 + (d_{\phi_1 \phi_2} - l_i) l_i m_1^2 + 3 m_i^2 m_1^4) \\
&\quad + \frac{m_j m_{\chi_2} s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (2 d_{\phi_1 \phi_2} l_i - 2 (d_{\phi_1 \phi_2} - 2 l_i) m_1^2 - 5 m_1^4) \\
&\quad - \frac{m_j m_{\chi_2} (2 d_{\phi_1 \phi_2} + m_1^2)}{8 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
J_{19}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_j m_{\chi_1} (2 d_{\phi_1 \phi_2} + m_1^2)}{4 m_1^4} + \frac{m_j m_{\chi_1} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{2 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} \\
&\quad (-4 d_{\phi_1 \phi_2}^2 m_i^2 + l_i^2 m_1^2 + (-5 l_i + 6 m_i^2) m_1^4) \\
&\quad + \frac{m_j m_{\chi_1} s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (-2 l_i m_1^2 + 5 m_1^4 \\
&\quad + 2 d_{\phi_1 \phi_2} (l_i + m_1^2)) - \frac{m_j m_{\chi_1} (2 d_{\phi_1 \phi_2} + m_1^2)}{8 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2) \\
J_{20}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_i m_{\chi_2} (2 d_{\phi_1 \phi_2} + m_1^2)}{4 m_1^4} - \frac{m_i m_{\chi_2} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} ((4 d_{\phi_1 \phi_2} - l_i) \\
&\quad (l_i - l_{ia}) m_1^2 + d (-8 d_{\phi_1 \phi_2}^2 + 12 m_1^4) + 2 (d_{\phi_1 \phi_2} l_i + m_1^4) s) \\
&\quad + \frac{m_i m_{\chi_2} s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} (4 d_{\phi_1 \phi_2} l_i + (8 d_{\phi_1 \phi_2} - l_i + 3 l_{ia}) m_1^2 \\
&\quad + 2 m_1^4) - \frac{m_i m_{\chi_2} (2 d_{\phi_1 \phi_2} + m_1^2)}{8 b_{\phi_1 \phi_2} m_1^4 (R_{\phi_1 \chi_1} - R_{\phi_2 \chi_2})} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)
\end{aligned}$$

$$\begin{aligned}
J_{21}(\phi_1, \phi_2, \chi_1, \chi_2) &= \frac{m_i m_{\chi_1} (2d_{\phi_1\phi_2} + m_1^2)}{4m_1^4} + \frac{m_i m_{\chi_1} s^2 \mathcal{X}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{4b_{\phi_1\phi_2} m_1^4 (R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (-m_1^2 (l_i - l_{ia}) \\
&\quad (2l_i - m_1^2) + 4d(2d_{\phi_1\phi_2}^2 - 3m_1^4) - 2d_{\phi_1\phi_2} (l_i - 4m_1^2) s) \\
&\quad \frac{m_i m_{\chi_1} s \mathcal{Y}_2(\chi_1, \chi_2, \phi_1, \phi_2)}{8b_{\phi_1\phi_2} m_1^4 (R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} (4d_{\phi_1\phi_2} l_i + (-8d_{\phi_1\phi_2} + 5l_i - 3l_{ia}) m_1^2 \\
&\quad - 2m_1^4) - \frac{m_i m_{\chi_1} (2d_{\phi_1\phi_2} + m_1^2)}{8b_{\phi_1\phi_2} m_1^4 (R_{\phi_1\chi_1} - R_{\phi_2\chi_2})} \mathcal{Z}_2(\chi_1, \chi_2, \phi_1, \phi_2)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_1(\chi, \phi_1, \phi_2) &= -\frac{b_{\phi_1\phi_2} m_\chi^4}{6s^2} + \frac{8}{s^2} (s - m_\chi^2)^2 (m_1^2 - m_2^2)^2 \tau_{\phi_1\phi_2} \tau_{\phi_2\phi_1} - \frac{d}{6} ((m_1^2 + m_2^2 \\
&\quad - 2d_{\phi_1\phi_2}) m_\chi^4 + (s - 2m_\chi^2) (m_1^2 - m_2^2)^2)
\end{aligned}$$

$$\mathcal{B}_2(\chi, \phi_1, \phi_2) = m_i m_j ((s - 2m_\chi^2) (m_1^2 - m_2^2)^2 + (m_i^2 + m_j^2 - 2d_{\phi_1\phi_2}) m_\chi^4)$$

$$\begin{aligned}
\mathcal{B}_3(\phi, \chi, \phi_1, \phi_2) &= b_{\phi_1\phi_2} m_\phi^2 + (m_\phi^2 (R_{\phi_1\chi} - 2sl_i) - 2sl_i (m_1^2 - m_2^2)) \\
&\quad \mathcal{A}(\chi, \phi_1, \phi_2)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_4(\phi, \chi, \phi_1, \phi_2) &= 2b_{\phi_1\phi_2} m_\phi^2 + (2(m_1^2 - m_2^2) l_{ia} + m_\phi^2 (\mathcal{C}(\chi, \phi_1, \phi_2) - 2sl_{\phi_1\phi_2})) \\
&\quad \mathcal{A}(\chi, \phi_1, \phi_2)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_5(\phi, \chi, \phi_1, \phi_2) &= \frac{R_{\phi_1\chi}^2}{8b_{\phi_1\phi_2} s} L(\chi, \phi_1, \phi_2) + \frac{R_{\phi_1\chi}}{8b_{\phi_1\phi_2} m_\phi^2 s} (-2b_{\phi_1\phi_2} m_\phi^2 - L(\chi, \phi_1, \phi_2) \\
&\quad (l_{\phi_2\phi_1} m_\phi^2 (m_i^2 - m_j^2) + (2m_i^2 m_\phi^2 - (m_1^2 - m_2^2) (m_i^2 - m_j^2)) s \\
&\quad + l_{\phi_2\phi_1} m_\phi^2 (2\tau_{\phi_1\phi_2} + 3s))) + \frac{1}{4m_\phi^2} ((\tau_{\phi_1\phi_2} + 2m_i^2) m_\phi^2 + (m_1^2 - m_2^2) \\
&\quad (m_i^2 - m_j^2)) - \frac{s^2 L(\chi, \phi_1, \phi_2)}{4b_{\phi_1\phi_2} m_\phi^2 s} ((m_1^2 - m_2^2) m_i^2 ((m_1^2 - m_2^2) + m_\phi^2 \\
&\quad + (m_i^2 - m_j^2)) + m_\phi^2 (2d(-d_{\phi_1\phi_2} + m_1^2) + 2l_{\phi_1\phi_2} (l_i + m_i^2) + m_i^2 \\
&\quad (m_i^2 - m_j^2)) + \frac{l_{ia}}{4s} (2(m_1^2 - m_2^2) + s)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_6(\phi, \chi, \phi_1, \phi_2) &= b_{\phi_1\phi_2} m_\phi^2 + (2(m_1^2 - m_2^2) (l_{\phi_1\phi_2} - l_i) s + m_\phi^2 (R_{\phi_1\chi} \\
&\quad + 2(d_{\phi_1\phi_2} - m_1^2 - l_i) s)) \mathcal{A}(\chi, \phi_1, \phi_2)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_7(\chi, \phi_1, \phi_2) &= -\frac{1}{48m_\chi^4 s^2} (b_{\phi_1\phi_2}^2 m_\chi^2 + 24d(m_\chi^2 - s)^2 s^2 + 12l_i l_{ia} (2m_\chi^2 - s) s^2) \\
&\quad - \frac{l_{\phi_1\phi_2}^2}{48m_\chi^6 s^2} (l_i l_{ia} (m_\chi^2 - s)^2 + 2d(2m_\chi^2 - s) s^2)
\end{aligned}$$

$$\mathcal{B}_8(\chi, \phi_1, \phi_2) = \frac{1}{8m_\chi^6} (l_{\phi_1\phi_2}^2 (-2m_\chi^2 + s) - 4m_\chi^2 (3m_\chi^4 - 2m_\chi^2 s + s^2))$$

$$\begin{aligned}
\mathcal{B}_9(\phi, \chi, \phi_1, \phi_2) &= -\frac{m_j}{8m_\phi^4 s} (l_{\phi_1\phi_2} l_i (m_\phi^2 - s) + m_\phi^2 (R_{\phi_1\chi} + 2s(-3m_\phi^2 + s))) \\
&\quad + \frac{m_j L(\chi, \phi_1, \phi_2)}{16b_{\phi_1\phi_2} m_\phi^4 s} (2m_\phi^4 s (-3R_{\phi_1\chi} + 4m_i^2 s) + l_{\phi_1\phi_2} s (-l_i R_{\phi_1\chi} \\
&\quad + 2l_{\phi_1\phi_2} m_i^2 s) + m_\phi^2 (R_{\phi_1\chi}^2 + 2(-\tau_{\phi_1\phi_2} + 2l_i^2 + R_{\phi_1\chi}) s^2 - 8m_i^2 s^3))
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_{10}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_i}{16 m_\phi^4} (\tau_{\phi_1 \phi_2} - \omega_{\phi_1 \phi_2} + 12 m_\phi^4 - 4 m_\phi^2 s) + \frac{m_i L(\chi, \phi_1, \phi_2)}{32 b_{\phi_1 \phi_2} m_\phi^4 s} \\
&\quad (\tau_{\phi_1 \phi_2} - \omega_{\phi_1 \phi_2} (12 m_\phi^4 - R_{\phi_1 \chi}) - 4 \tau_{\phi_1 \phi_2} m_\phi^2 s + l_{\phi_1 \phi_2}^2 (l_i^2 - l_i l_{ia} \\
&\quad + 4 d s) + 4 m_\phi^2 (3 m_\phi^2 - s) (-R_{\phi_1 \chi} + 4 d s)) \\
\mathcal{B}_{11}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_\chi}{8 m_\phi^4 s} (m_\phi^2 R_{\phi_1 \chi} - l_{\phi_1 \phi_2} (l_i m_\phi^2 + s (-2 m_\phi^2 + s))) + \frac{m_\chi L(\chi, \phi_1, \phi_2)}{16 b_{\phi_1 \phi_2} m_\phi^4 s} \\
&\quad (l_{\phi_1 \phi_2}^2 s (l_i l_{ia} - (2 d + l_i) s) + l_{\phi_1 \phi_2} R_{\phi_1 \chi} (2 l_i m_\phi^2 + s (-2 m_\phi^2 + s)) \\
&\quad - m_\phi^2 (R_{\phi_1 \chi}^2 + 4 s^2 (l_i l_{ia} + 2 d m_\phi^2 - 2 d s))) \\
\mathcal{B}_{12}(\phi, \chi, \phi_1, \phi_2) &= -\frac{m_i m_j m_\chi s L(\chi, \phi_1, \phi_2)}{8 b_{\phi_1 \phi_2} m_\phi^4} (l_{\phi_1 \phi_2}^2 + 12 m_\phi^4 - 4 m_\phi^2 s) \\
\mathcal{B}_{13}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_\chi L(\chi, \phi_1, \phi_2)}{16 b_{\phi_1 \phi_2} m_\phi^4} (l_{\phi_2 \phi_1} \rho(\chi, \phi_1, \phi_2) (-2 m_\phi^2 + s) + l_{\phi_2 \phi_1}^2 (l_i l_{ia} - 2 d s) \\
&\quad - 4 m_\phi^2 s (l_i l_{ia} + 2 d m_\phi^2 - 2 d s)) + \frac{m_\chi l_{\phi_2 \phi_1}}{4 b_{\phi_1 \phi_2} m_\phi^2} \\
\mathcal{B}_{14}(\phi, \chi, \phi_1, \phi_2) &= -\frac{m_i m_j m_\chi s L(\chi, \phi_1, \phi_2)}{8 b_{\phi_1 \phi_2} m_\phi^4} (l_{\phi_2 \phi_1}^2 + 12 m_\phi^4 - 4 m_\phi^2 s) \\
\mathcal{B}_{15}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_j L(\chi, \phi_1, \phi_2)}{32 b_{\phi_1 \phi_2} m_\phi^4} (l_{\phi_2 \phi_1} (-l_{\phi_1 \phi_2} \rho(\chi, \phi_1, \phi_2) + 4 d_{\phi_1 \phi_2} l_i s) \\
&\quad + l_{\phi_2 \phi_1}^2 (\rho(\chi, \phi_1, \phi_2) - 4 m_i^2 s) + 4 m_\phi^2 (3 m_\phi^2 - s) (R_{\phi_1 \chi} - 4 m_i^2 s)) \\
&\quad - \frac{m_j}{2 m_\phi^4} (l_{\phi_2 \phi_1}^2 - l_{\phi_2 \phi_1} l_{\phi_1 \phi_2} + 12 m_\phi^4 - 4 m_\phi^2 s) \\
\mathcal{B}_{16}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_i L(\chi, \phi_1, \phi_2)}{32 b_{\phi_1 \phi_2} m_\phi^4} (2 l_{\phi_2 \phi_1}^2 l_i l_{ia} + 8 d_{\phi_1 \phi_2} l_{\phi_2 \phi_1} l_{ia} m_\phi^2 + (l_{\phi_2 \phi_1}^2 \\
&\quad - l_{\phi_2 \phi_1} l_{\phi_1 \phi_2}) \rho(\chi, \phi_1, \phi_2) - 4 m_\phi^2 (l_{\phi_2 \phi_1} - m_1^2) R_{\phi_1 \chi} - 2 (l_{\phi_2 \phi_1}^3 \\
&\quad + 4 l_i l_{ia} m_\phi^2 + 8 d m_\phi^4 + 12 l_{\phi_2 \phi_1} m_\phi^4 - 4 l_{\phi_1 \phi_2} m_\phi^4 - l_{\phi_2 \phi_1} R_{\phi_1 \chi}) s \\
&\quad + 8 (2 d + 2 s) m_\phi^2 s^2) + \frac{m_i}{32 m_\phi^4} (2 l_{\phi_2 \phi_1} (-l_{\phi_2 \phi_1} + l_{\phi_1 \phi_2}) + 8 m_\phi^2 (m_\phi^2 + s)) \\
\mathcal{B}_{17}(\phi, \chi, \phi_1, \phi_2) &= s (m_\phi^4 + (m_1^2 - s)^2 - 2 m_2^2 (m_1^2 + s)) \frac{\mathcal{A}(\chi, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^2 (m_2^2 - s) s} \\
\mathcal{B}_{18}(\phi, \chi, \phi_1, \phi_2) &= (6 m_\phi^2 - s) s - (m_1^2 - m_2^2) (2 m_\phi^2 + s) \\
\mathcal{B}_{19}(\phi, \chi, \phi_1, \phi_2) &= 4 d_{\phi_1 \phi_2} s^2 - 3 m_\phi^2 (m_2^2 - m_1^2 + 2 s^2) \\
\mathcal{B}_{20}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_j}{8 m_1^4} (d_{\phi_1 \phi_2} l_i + 2 (-d_{\phi_1 \phi_2} + l_i) m_1^2 + 4 m_1^4) + \frac{s m_j L(\chi, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} m_1^2} \\
&\quad (2 d_{\phi_1 \phi_2}^2 m_i^2 + l_i (-d_{\phi_1 \phi_2} + l_i) m_1^2) - \frac{m_j R_{\phi_1 \chi}}{8 b_{\phi_1 \phi_2} m_1^4 s} (b_{\phi_1 \phi_2} (d_{\phi_1 \phi_2} + 2 m_1^2) \\
&\quad + (d_{\phi_1 \phi_2} (l_i - m_1^2) + 2 m_1^2 (l_i + m_1^2)) s L(\chi, \phi_1, \phi_2))
\end{aligned}$$



$$\begin{aligned}
\mathcal{B}_{21}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_i}{8 b_{\phi_1 \phi_2} m_1^4} (d_{\phi_1 \phi_2} l_i + 2(-d_{\phi_1 \phi_2} + l_i) m_1^2 + 4 m_1^4) + \frac{m_i s L(\chi, \phi_1, \phi_2)}{8 b_{\phi_1 \phi_2} m_1^4} \\
&\quad (-4 d d_{\phi_1 \phi_2}^2 - (d_{\phi_1 \phi_2} - l_i)(l_i - l_{ia}) m_1^2 + (d_{\phi_1 \phi_2} l_i + 4 m_1^4) s) \\
&\quad - \frac{m_i R_{\phi_1 \chi}}{8 b_{\phi_1 \phi_2} s m_1^4} (b_{\phi_1 \phi_2} (d_{\phi_1 \phi_2} + 2 m_1^2) + (d_{\phi_1 \phi_2} (l_i - m_1^2) + 2 m_1^2 (l_i + m_1^2)) \\
&\quad s L(\chi, \phi_1, \phi_2)) \\
\mathcal{B}_{22}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_\chi s}{8 b_{\phi_1 \phi_2} m_1^4} (-2 b_{\phi_1 \phi_2} d_{\phi_1 \phi_2} + (4 d (d_{\phi_1 \phi_2}^2 + 2 m_1^4) + d_{\phi_1 \phi_2} (R_{\phi_1 \chi} - l_i s)) \\
&\quad L(\chi, \phi_1, \phi_2)) \\
\mathcal{B}_{23}(\phi, \chi, \phi_1, \phi_2) &= -\frac{m_i m_j m_\chi s L(\chi, \phi_1, \phi_2)}{2 b_{\phi_1 \phi_2} m_1^4} (d_{\phi_1 \phi_2}^2 + 2 m_1^4)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_{24}(\phi, \chi, \phi_1, \phi_2) &= \frac{1}{192 b_{\phi_1 \phi_2} m_1^4 m_\phi^2 s^2} (2 b_{\phi_1 \phi_2} (2 b_{\phi_1 \phi_2}^2 m_\phi^2 (2 m_1^2 - s) + 24 d_{\phi_1 \phi_2} (d - m_i^2) \\
&\quad m_\phi^2 s^3 - 12 m_1^4 s^2 (l_i^2 - 3 l_i l_{ia} + 4 d s + 2 m_\phi^2 (-2 (2 d + l_i + m_i^2) + 5 s)) \\
&\quad + 6 m_1^2 s^2 ((-2 d_{\phi_1 \phi_2} + s) (l_i^2 - 3 l_i l_{ia} + 4 d s) + m_\phi^2 (-l_i^2 + l_i (7 l_{ia} - 2 s) \\
&\quad + 2 s (4 d_{\phi_1 \phi_2} + 2 m_i^2 + s)))) + 3 s^2 (-m_\phi^2 (l_i^3 + 8 d_{\phi_1 \phi_2} l_i m_i^2 - 8 d_{\phi_1 \phi_2} l_{ia} m_i^2 \\
&\quad + l_i^2 (l_{ia} - 2 s)) s^2 + 16 m_1^6 s (l_i l_{ia} + 6 d m_\phi^2 - 2 d s) + 4 m_1^4 s (-l_i^3 + 3 l_i^2 l_{ia} \\
&\quad - 2 l_i (2 d + l_{ia}) s + 4 d s^2 + 2 m_\phi^2 (2 (l_i - 4 l_{ia}) m_i^2 - l_i s + 2 d (-6 d_{\phi_1 \phi_2} + 6 l_i \\
&\quad + s))) + 2 m_1^2 ((-2 d_{\phi_1 \phi_2} + s) s (l_i (l_i^2 + 4 d_{\phi_1 \phi_2} l_{ia} - 3 l_i l_{ia}) + 4 d (-2 d_{\phi_1 \phi_2} \\
&\quad + l_i) s) + m_\phi s (4 d_{\phi_1 \phi_2} (l_i^2 - l_i l_{ia} + 4 l_{ia} m_i^2) - 8 d d_{\phi_1 \phi_2} (2 l_i + s) + l_i (-l_i^2 \\
&\quad + 7 l_i l_{ia} - 2 l_i s + 4 m_i^2 s + 2 s^2)))) L(\chi, \phi_1, \phi_2)) + \frac{R_{\phi_1 \chi}}{64 b_{\phi_1 \phi_2} m_1^2 m_\phi^2 s} \\
&\quad (4 b_{\phi_1 \phi_2} m_\phi^2 (-4 m_1^2 (-d_{\phi_1 \phi_2} + l_i + 2 m_1^2) + 2 (l_i + m_1^2) s - s^2) - s \\
&\quad (2 m_1^2 (-2 l_i (l_i - 3 l_{ia}) (d_{\phi_1 \phi_2} + m_1^2) + (3 l_i^2 + 4 d_{\phi_1 \phi_2} l_{ia} + 7 l_i l_{ia} \\
&\quad + 8 (2 d + 2 l_i + m_i^2) m_1^2) m_\phi^2) + 4 (2 d m_1^2 + (l_i + m_1^2) m_\phi^2) s^2 - s (2 m_1^2 (-l_i \\
&\quad (l_i - 3 l_{ia}) + 8 d (d_{\phi_1 \phi_2} + m_1^2)) + m_\phi^2 (-8 d d_{\phi_1 \phi_2} + 8 d_{\phi_1 \phi_2} m_i^2 + 8 (l_i - m_i^2) \\
&\quad m_1^2 + 40 m_1^4 + l_i (4 l_i + 2 s)))) L(\chi, \phi_1, \phi_2)) + \frac{R_{\phi_1 \chi}^2}{32 b_{\phi_1 \phi_2} m_1^2 s} (b_{\phi_1 \phi_2} \\
&\quad (4 m_1^2 - 2 s) + s (-4 d_{\phi_1 \phi_2} m_1^2 + 6 l_i m_1^2 + 8 m_1^4 - 3 l_i s - 2 m_1^2 s + s^2) \\
&\quad L(\chi, \phi_1, \phi_2)) + \frac{6 R_{\phi_1 \chi}^3 L(\chi, \phi_1, \phi_2)}{192 b_{\phi_1 \phi_2} m_1^4 s^2} (s - 2 m_1^2)
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_{25}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_i m_j}{8 b_{\phi_1 \phi_2} m_1^2 m_\phi^2} (2 d_{\phi_1 \phi_2} (2 m_\phi^2 - s) + 2 m_1^2 (5 m_\phi^2 - s) + s (3 m_\phi^2 + s)) \\
&\quad (2 b_{\phi_1 \phi_2} - (R_{\phi_1 \chi} + (2 d_{\phi_1 \phi_2} - l_i - 2 m_1^2) s) L(\chi, \phi_1, \phi_2))
\end{aligned}$$

$$\begin{aligned}
\mathcal{B}_{26}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_\chi m_j}{8 b_{\phi_1 \phi_2} m_1^4} (2 b_{\phi_1 \phi_2} (-4 m_1^4 + d_{\phi_1 \phi_2} (2 m_1^2 + s)) - (l_i m_1^2 s^2 - 2 m_1^4 (2 R_{\phi_1 \chi} \\
&\quad + 5 l_i s) + d_{\phi_1 \phi_2} (R_{\phi_1 \chi} s + 2 m_1^2 (R_{\phi_1 \chi} + 2 l_i s))) L(\chi, \phi_1, \phi_2))
\end{aligned}$$

$$\begin{aligned}\mathcal{B}_{27}(\phi, \chi, \phi_1, \phi_2) &= \frac{m_\chi m_i}{8 b_{\phi_1 \phi_2} m_1^4} \left( -2 b_{\phi_1 \phi_2} (-4 m_1^4 + d_{\phi_1 \phi_2} (2 m_1^2 + s)) + (l_{ia} m_1^2 s^2 - 2 m_1^4 (2 R_{\phi_1 \chi} \right. \\ &\quad \left. - 2 l_i s + 7 l_{ia} s) + d_{\phi_1 \phi_2} (s (R_{\phi_1 \chi} - l_i s + l_{ia} s) + 2 m_1^2 (R_{\phi_1 \chi} - l_i s + 3 l_{ia} s)) \right) \\ &\quad L(\chi, \phi_1, \phi_2)\end{aligned}$$

$$\begin{aligned}\mathcal{C}_1(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{1}{8 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left( (\omega_{\phi_1 \phi_2} - \tau_{\phi_1 \phi_2}) (m_i^2 - 2 m_1^2) + 4 (2 d + m_i^2) \right. \\ &\quad \left. m_1^2 s \mathcal{X}_1(\chi_1, \chi_2, \phi_1, \phi_2) + \frac{s \mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left( (\tau_{\phi_1 \phi_2} - \omega_{\phi_1 \phi_2}) m_i^2 \right. \right. \\ &\quad \left. \left. + 2 d (m_i^2 - m_1^2) s \right) + \frac{2 d - m_i^2 + 2 m_1^2}{8 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_2} - R_{\phi_1 \chi_1}) \right. \\ &\quad \left. + \mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2) \right)\end{aligned}$$

$$\begin{aligned}\mathcal{C}_2(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{m_{\chi_1} m_{\chi_2} (\tau_{\phi_1 \phi_2} - \omega_{\phi_1 \phi_2})}{8 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \mathcal{X}_1(\chi_1, \chi_2, \phi_1, \phi_2) \\ &\quad + \frac{s^2 d m_{\chi_1} m_{\chi_2} \mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} + \frac{m_{\chi_1} m_{\chi_2}}{8 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \\ &\quad (2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_2} - R_{\phi_1 \chi_1}) + \mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2))\end{aligned}$$

$$\mathcal{C}_3(\chi_1, \chi_2, \phi_1, \phi_2) = \frac{\mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})}$$

$$\begin{aligned}\mathcal{C}_4(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{m_j}{8 b_{\phi_1 \phi_2} m_1^2 (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_2} - R_{\phi_1 \chi_1}) \\ &\quad + \mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2))\end{aligned}$$

$$\begin{aligned}\mathcal{C}_5(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{3 m_i s}{4 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left[ ((\omega_{\phi_1 \phi_2} - \tau_{\phi_1 \phi_2}) - R_{\phi_1 \chi_1} + 4 d s) \right. \\ &\quad \left. L(\chi_1, \phi_1, \phi_2) - ((\omega_{\phi_1 \phi_2} - \tau_{\phi_1 \phi_2}) - R_{\phi_1 \chi_2} + 4 d s) \right. \\ &\quad \left. L(\chi_2, \phi_1, \phi_2) \right]\end{aligned}$$

$$\begin{aligned}\mathcal{C}_6(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{3 m_j s}{4 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left[ (3 R_{\phi_1 \chi_1} - 4 m_i^2 s) L(\chi_1, \phi_1, \phi_2) \right. \\ &\quad \left. - (3 R_{\phi_1 \chi_2} - 4 m_i^2 s) L(\chi_2, \phi_1, \phi_2) \right]\end{aligned}$$

$$\begin{aligned}\mathcal{C}_7(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{3 m_i m_j s}{2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left[ (-R_{\phi_1 \chi_1} + 2 (m_1^2 - m_i^2) s) L(\chi_1, \phi_1, \phi_2) \right. \\ &\quad \left. - (-R_{\phi_1 \chi_2} + 2 (m_1^2 - m_i^2) s) L(\chi_2, \phi_1, \phi_2) \right]\end{aligned}$$

$$\begin{aligned}\mathcal{C}_8(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{\mathcal{X}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{4 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \left( (m_1^2 - m_i^2) (-\tau_{\phi_1 \phi_2} + l_{\phi_1 \phi_2} s) + m_2^2 (\tau_{\phi_1 \phi_2} \right. \\ &\quad \left. - l_{\phi_2 \phi_1} s + 2 m_i^2 s) + d_{\phi_1 \phi_2} (\tau_{\phi_1 \phi_2} - \omega_{\phi_2 \phi_1} + 4 m_i^2 s) \right) \\ &\quad + \frac{\mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{8 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (8 d_{\phi_1 \phi_2} m_i^2 s (\tau_{\phi_1 \phi_2} - \omega_{\phi_2 \phi_1}) \\ &\quad + 4 m_i^2 m_2^2 s (\tau_{\phi_1 \phi_2} - \omega_{\phi_1 \phi_2}) - (m_1^2 - m_i^2) (8 d m_2^2 s^2 + \tau_{\phi_2 \phi_1} \omega_{\phi_2 \phi_1} \\ &\quad - 2 l_{\phi_2 \phi_1} s \tau_{\phi_1 \phi_2} + \tau_{\phi_1 \phi_2}^2)) - \frac{s - 2 m_1^2 + m_i^2}{8 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (2 b_{\phi_1 \phi_2} \\ &\quad (R_{\phi_1 \chi_2} - R_{\phi_1 \chi_1}) + \mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2))\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_9(\chi_1, \chi_2, \phi_1, \phi_2) &= \frac{\mathcal{X}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{4 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (\tau_{\phi_1 \phi_2} - l_{\phi_2 \phi_1} s) \\
&+ \frac{\mathcal{Y}_1(\chi_1, \chi_2, \phi_1, \phi_2)}{8 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (8 d m_2^2 s^2 + \omega_{\phi_2 \phi_1} \tau_{\phi_2 \phi_1} \\
&- 2 l_{\phi_2 \phi_1} s \tau_{\phi_1 \phi_2} + \tau_{\phi_1 \phi_2}^2) + \frac{1}{8 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \\
&(2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_2} - R_{\phi_1 \chi_1}) + \mathcal{Z}_1(\chi_1, \chi_2, \phi_1, \phi_2))
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{10}(\chi_1, \chi_2, \phi_1, \phi_2) &= -\frac{m_i s L(\chi_1, \phi_1, \phi_2)}{4 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (2 d_{\phi_1 \phi_2} (\omega_{\phi_2 \phi_1} + R_{\phi_1 \chi_1} - \tau_{\phi_1 \phi_2})) \\
&+ m_2^2 (\omega_{\phi_1 \phi_2} + R_{\phi_1 \chi_1} - \tau_{\phi_1 \phi_2}) + \frac{m_i s L(\chi_2, \phi_1, \phi_2)}{4 m_2^2 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} \\
&(2 d_{\phi_1 \phi_2} (\omega_{\phi_2 \phi_1} + R_{\phi_1 \chi_2} - \tau_{\phi_1 \phi_2}) + m_2^2 (\omega_{\phi_1 \phi_2} + R_{\phi_1 \chi_2} - \tau_{\phi_1 \phi_2}))
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{11}(\chi_1, \chi_2, \phi_1, \phi_2) &= -\frac{m_j s L(\chi_1, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (3 R_{\phi_1 \chi_1} - 4 m_i^2 s) \\
&+ \frac{m_j s L(\chi_1, \phi_1, \phi_2)}{4 b_{\phi_1 \phi_2} (R_{\phi_1 \chi_1} - R_{\phi_1 \chi_2})} (3 R_{\phi_1 \chi_2} - 4 m_i^2 s)
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{12}(\chi_1, \chi_2, \phi, \phi) &= -\frac{1}{48 m_\phi^4 S^2} (b_{\phi\phi}^2 + 3 (R_{\phi\chi_1}^2 + R_{\phi\chi_1} R_{\phi\chi_2} + R_{\phi\chi_2}^2 - (2 l_i - l_{ia} + 4 m_\phi^2) \\
&(R_{\phi\chi_1} + R_{\phi\chi_2}) S - (l_i l_{ia} - 8 d_{\phi\phi} m_i^2 + 4 (2 d + 4 d_{\phi\phi} - 2 l_i + l_{ia}) \\
&m_\phi^2 - 4 m_\phi^4) S^2)) + \frac{S^2 \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi)}{4 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} (-(4 d_{\phi\phi} - l_i) (l_i - l_{ia}) \\
&m_i^2 + 2 (l_i + 2 m_i^2) m_j^2 m_\phi^2 + 4 d m_\phi^4) - \frac{S \mathcal{Y}_1(\chi_1, \chi_2, \phi, \phi)}{4 b_{\phi\phi} m_\phi^4 (R_{\phi\chi_1} - R_{\phi\chi_2})} (d_{\phi\phi} \\
&(l_i - l_{ia}) m_i^2 + 4 d_{\phi\phi} (-2 m_i^2 + m_j^2) m_\phi^2 + m_\phi^2 ((l_i - l_{ia}) (l_i + m_i^2) \\
&+ m_\phi^2 (-4 d + 2 m_j^2 + S))) + \frac{\mathcal{Z}_1(\chi_1, \chi_2, \phi, \phi)}{16 b_{\phi\phi} m_\phi^4 (R_{\phi\chi_1} - R_{\phi\chi_2})} (l_i^2 - l_i l_{ia} + 4 d_{\phi\phi} m_i^2 \\
&- 2 (2 d + 4 d_{\phi\phi} - 3 l_i + l_{ia}) m_\phi^2 + 2 m_\phi^4) + \frac{\mathcal{R}_1(\chi_1, \chi_2, \phi, \phi)}{32 b_{\phi\phi} m_\phi^4 S (R_{\phi\chi_1} - R_{\phi\chi_2})} \\
&(-3 l_i + l_{ia} - 4 m_\phi^2) + \frac{\mathcal{Q}_1(\chi_1, \chi_2, \phi, \phi)}{32 b_{\phi\phi} m_\phi^4 S^2 (R_{\phi\chi_1} - R_{\phi\chi_2})}
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}_{13}(\chi_1, \chi_2, \phi, \phi) &= \frac{m_{\chi_k^\pm} m_{\chi_l^\pm} (d_{\phi\phi} - m_\phi^2)}{2 m_\phi^4} - \frac{m_{\chi_k^\pm} m_{\chi_l^\pm} S \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi)}{4 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} (l_i (-l_i + l_{ia})) \\
&+ 4 d m_\phi^2 + \frac{m_{\chi_k^\pm} m_{\chi_l^\pm} S \mathcal{Y}_1(\chi_1, \chi_2, \phi, \phi)}{4 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} (d_{\phi\phi} (l_i - l_{ia}) - 2 l_i m_\phi^2) \\
&+ \frac{m_{\chi_k^\pm} m_{\chi_l^\pm} (d_{\phi\phi} - m_\phi^2)}{4 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} \\
\mathcal{C}_{14}(\chi_1, \chi_2, \phi, \phi) &= \frac{9 m_i m_j S}{2 b_{\phi\phi} (R_{\phi\chi_1} - R_{\phi\chi_2})} (2 m_i^2 S \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi) + 2 m_\phi^2 S \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi) \\
&- \mathcal{Y}_1(\chi_1, \chi_2, \phi, \phi)) \\
\mathcal{C}_{15}(\chi_1, \chi_2, \phi, \phi) &= \frac{9 m_i m_j m_{\chi_k^\pm} m_{\chi_l^\pm} S}{b_{\phi\phi} (R_{\phi\chi_1} - R_{\phi\chi_2})} \\
\mathcal{C}_{16}(\chi_1, \chi_2, \phi, \phi) &= \frac{3 m_j}{4 m_\phi^2} + \frac{3 m_j}{8 m_\phi^2 b_{\phi\phi} (R_{\phi\chi_1} - R_{\phi\chi_2})} (-8 m_i^2 m_\phi^2 S^2 \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi) \\
&+ 6 m_\phi^2 S \mathcal{Y}_1(\chi_1, \chi_2, \phi, \phi) - \mathcal{Z}_1(\chi_1, \chi_2, \phi, \phi)) \\
\mathcal{C}_{17}(\chi_1, \chi_2, \phi, \phi) &= \frac{3 m_i}{4 m_\phi^2} + \frac{3 m_i S^2 \mathcal{X}_1(\chi_1, \chi_2, \phi, \phi)}{4 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} ((2 d_{\phi\phi} - l_i) (l_i - l_{ia}) \\
&- m_\phi^2 (-4 d + 2 S)) - \frac{3 m_i S \mathcal{Y}_1(\chi_1, \chi_2, \phi, \phi)}{8 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})} ((4 d_{\phi\phi} - 3 l_i + l_{ia} - 2 m_\phi^2) \\
&- \frac{3 m_i \mathcal{Z}_1(\chi_1, \chi_2, \phi, \phi)}{8 b_{\phi\phi} m_\phi^2 (R_{\phi\chi_1} - R_{\phi\chi_2})}
\end{aligned}$$

Where  $m_i, m_j$  are masses of  $\chi_i^0$  and  $\chi_j^0$  respectively.

## References

- [1] S. P. Martin, ‘‘A Supersymmetry primer,’’ *Adv.Ser.Direct.High Energy Phys.*, vol. 21, pp. 1–153, 2010.
- [2] K. Griest and D. Seckel, ‘‘Three exceptions in the calculation of relic abundances,’’ *Phys.Rev.*, vol. D43, pp. 3191–3203, 1991.
- [3] M. Srednicki, R. Watkins, and K. A. Olive, ‘‘Calculations of Relic Densities in the Early Universe,’’ *Nucl.Phys.*, vol. B310, p. 693, 1988.
- [4] J. McDonald, K. A. Olive, and M. Srednicki, ‘‘Relic densities of neutralinos,’’ *Phys.Lett.*, vol. B283, pp. 80–84, 1992.
- [5] S. Mizuta and M. Yamaguchi, ‘‘Coannihilation effects and relic abundance of Higgsino dominant LSP(s),’’ *Phys.Lett.*, vol. B298, pp. 120–126, 1993.
- [6] M. Drees and M. M. Nojiri, ‘‘The Neutralino relic density in minimal  $N = 1$  supergravity,’’ *Phys.Rev.*, vol. D47, pp. 376–408, 1993.
- [7] M. Drees, M. M. Nojiri, D. Roy, and Y. Yamada, ‘‘Light Higgsino dark matter,’’ *Phys.Rev.*, vol. D56, pp. 276–290, 1997.

- [8] J. Edsjo and P. Gondolo, “Neutralino relic density including coannihilations,” *Phys.Rev.*, vol. D56, pp. 1879–1894, 1997.
- [9] T. Nihei, L. Roszkowski, and R. Ruiz de Austri, “Exact cross-sections for the neutralino WIMP pair annihilation,” *JHEP*, vol. 0203, p. 031, 2002.
- [10] T. Nihei, L. Roszkowski, and R. Ruiz de Austri, “Exact cross-sections for the neutralino slepton coannihilation,” *JHEP*, vol. 0207, p. 024, 2002.
- [11] D. Choudhury, R. Garani, and S. K. Vempati, “Flavored Co-annihilations,” *JHEP*, vol. 1206, p. 014, 2012.
- [12] B. Herrmann, M. Klasen, and Q. Le Boulc’h, “Impact of squark flavour violation on neutralino dark matter,” *Phys.Rev.*, vol. D84, p. 095007, 2011.
- [13] J. Hisano, T. Moroi, K. Tobe, and M. Yamaguchi, “Lepton flavor violation via right-handed neutrino Yukawa couplings in supersymmetric standard model,” *Phys.Rev.*, vol. D53, pp. 2442–2459, 1996.
- [14] S. Profumo, “Neutralino dark matter, b tau Yukawa unification and nonuniversal sfermion masses,” *Phys.Rev.*, vol. D68, p. 015006, 2003.
- [15] H. Baer, A. Mustafayev, S. Profumo, A. Belyaev, and X. Tata, “Direct, indirect and collider detection of neutralino dark matter in SUSY models with non-universal Higgs masses,” *JHEP*, vol. 0507, p. 065, 2005.
- [16] R. Sundrum, “SUSY Splits, But Then Returns,” *JHEP*, vol. 1101, p. 062, 2011.
- [17] N. Arkani-Hamed, S. Dimopoulos, G. Giudice, and A. Romanino, “Aspects of split supersymmetry,” *Nucl.Phys.*, vol. B709, pp. 3–46, 2005.
- [18] J. Rosiek, “Complete Set of Feynman Rules for the Minimal Supersymmetric Extension of the Standard Model,” *Phys.Rev.*, vol. D41, p. 3464, 1990.