# ON NONLOCAL NONLINEAR ELLIPTIC PROBLEM WITH THE FRACTIONAL LAPLACIAN

#### LI MA

ABSTRACT. In this paper, we study a nonlocal elliptic problem with the fractional Laplacian on  $R^n$ . We show that the problem has infinite positive solutions in  $C^{\tau}(R^n) \cap H^{\alpha}_{loc}(R^n)$ . Moreover each of these solutions tends to some positive constant limit at infinity.

**Mathematics Subject Classification** (2000): 35J60, 53C21, 58J05 **Keywords**: fractional Laplacian,

### 1. Introduction

We prove the following result.

**Theorem 1.** Assume  $0 < \alpha < n$  and p > 1. Let  $\omega : R_+ \to R_+$  be the monotone non-increasing function such that

(1) 
$$\int_{1}^{\infty} \frac{\omega(r)}{r} dr = A < \infty \pounds$$

Then there is a positive constant  $\theta$  such that for smooth functions k(x) and K(x) with  $|K(x)| \leq \theta \omega(|x|)(1+|x|)^{-\tau}$  and  $0 \leq k(x) \leq \theta \omega(|x|)(1+|x|)^{-\tau}$  on  $R^n$  for some  $\tau \geq \alpha$ , the problem

$$(2) \qquad (-\Delta)^{\alpha/2}u + k(x)u = K(x)u^p, \quad in \ R^n$$

has infinite positive solutions in  $C^{\tau}(R^n) \cap H^{\alpha}_{loc}(R^n)$ . Moreover, each of these solutions tends to some positive constant limit at infinity.

We prove the above result by using the Perron method and similar argument as in Lin's work [3]. The Perron method is based on the maximum principle developed in [1]. Similar argument had been carried out in [4] for nonlinear subelliptic equations on  $\mathbb{R}^n$ .

We denote by C the uniform constants, which may vary in different inequalities or formulae.

Denote by  $B_1(0)$  the unit ball in  $\mathbb{R}^n$ .

We present the potential analysis as in [2] in section 2 and we prove the main result in section 3.

Date: May 26th, 2014.

<sup>\*</sup> The research is partially supported by the National Natural Science Foundation of China (No. 11271111) and SRFDP 20090002110019.

2 LI MA

#### 2. Preliminary

Let  $\omega: \mathbb{R}^n \to \mathbb{R}$  be a radially symmetric monotone non-increasing function. Let  $f: \mathbb{R}^n \to \mathbb{R}$  be a locally Holder continuous function with the decay growth as below:

$$|f(x)| \le C\omega(|x|)|x|^{-\tau}$$

where C > 0 and  $\tau > 2\alpha$  are uniform constants.

# Lemma 2. Define

$$w(x) = c_n \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n - \alpha}} dy,$$

where  $c_n = [n(n-2)|B_1(0)|]^{-1}$ . Then w(x) is well-defined and near  $\infty$  we have

(4) 
$$|w(x)| \le \begin{cases} C|x|^{\alpha-n}\omega(|x|) & \text{if } \tau > n \\ C\omega(|x|)\log|x| & \text{if } \tau = n \\ C\omega(|x|)|x|^{\alpha-\tau} & \text{if } \alpha \le \tau \end{cases}$$

where  $c_n = 1/n(n-2)|B_1(0)|$ .

Proof. Clearly we have

$$|w(x)| \le C \int_{R^n} \frac{\omega(|x|)}{|x - y|^{n - \alpha} (1 + |y|^{\tau}} dy.$$

We now divide the the whole space  $R^n$  into three parts:

$$D_1 = \{ y \in R^n; |y - x| \le |x|/2 \},$$
  
$$D_2 = \{ y \in R^n; |x|/2 \le |y - x| \le 2|x| \},$$

and

$$D_3 = \{ y \in R^n; |y - x| \ge 2|x| \}.$$

Set, for i = 1, 2, 3,

$$I_j = \int_{D_i} \frac{\omega(|x|)}{|x - y|^{n - \alpha} (1 + |y|^{\tau}} dy.$$

Then

$$|w(x)| \le C(I_1 + I_2 + I_3).$$

One can show as in [2] that w(x) has the upper bounds as in (4).

We now assume that  $\omega(x)$  satisfies (1).

**Lemma 3.** Assume  $f \ge 0$  on  $R^n$  and  $f(x) \ge C\omega(|x|)|x|^{-\tau}$  for some  $\tau \ge \alpha$  and C > 0. The Riesz potential w(x) defined in Lemma 1 has the following lower bounds at  $\infty$ :

(5) 
$$w(x) \ge \begin{cases} C|x|^{\alpha - n}\omega(|x|) & \text{if } \tau > n \\ C\omega(|x|)\log|x| & \text{if } \tau = n \\ C\omega(|x|)|x|^{\alpha - \tau} & \text{if } \alpha \le \tau n \end{cases}$$

*Proof.* As in the proof above, we have

$$w(x) = \left(\int_{D_1} + \int_{D_2} + \int_{D_3} \right) \frac{c_n f(y)}{|x - y|^{n - \alpha}} dy \ge \int_{D_2} \frac{c_n f(y)}{|x - y|^{n - \alpha}} dy := J.$$

One can show as in [2] that J has the lower bounds as in (5).

#### 3. Proof of Theorem 1

The argument presented below is similar to LIn's work [3]. We just do it briefly.

Take some constants  $\theta_1 > 0$  and 0 < a < 1. We first define the function  $U_a(x)$  on  $\mathbb{R}^n$  by solving the nonlocal equation

$$(-\Delta)^{\alpha/2}u = -\frac{C\omega(x)}{(1+|x|)^{\tau}}, \quad on \ R^n$$

for  $C \in (0, \theta_1)$  with  $0 < U_a(x) \le a$  and  $U_a(x) \to a$  at infinity. We can verify that  $U_a$  is the lower solution to (2).

Then we define the function  $U^a(x)$  on  $\mathbb{R}^n$  by solving the nonlocal equation

$$(-\Delta)^{\alpha/2}u = \frac{C\omega(x)}{(1+|x|)^{\tau}}, \text{ on } R^n$$

for the same C > 0 with  $a \le U^a(x) < 1$  and  $U^a(x) \to a$  at infinity. We can verify that  $U^a$  is the upper solution to (2).

Then we can use the Perron method [1][3] to get the desired solution u(x) to (2) such that  $U_a(x) \le u(x) \le U^a(x)$  on  $\mathbb{R}^n$ .

This then completes the proof of Theorem 1.

## References

- [1] L.Caffarelli, L. Silvestre, regularity Theory for fully nonlinear integro-Differential Equations Comm. Pure Appl. Math. Vol. LXXII 0597-0638(2009.
- [2] Y.Li, W.M.Ni, On conformally scalar curvature equations in  $\mathbb{R}^n$ , Duke Math. J., 57(1988)895-924
- [3] F.H.Lin, On the elliptic equation  $D_i[a_{ij}(x)D_jU k(x)U + K(x)U^p] = 0$ , PAMS,95(2)(1985)219-226
- [4] Li MA, Dezhong Chen, Yang Yang, *Some results for subelliptic equations*, Acta. Math.Sinica. English series, 22(2006)1695-1704.

Ma: Department of mathematics, Henan Normal University, Xinxiang, 453007, China *E-mail address*: lma@tsinghua.edu.cn