Extremal functions for Trudinger-Moser inequalities of Adimurthi-Druet type in dimension two

Yunyan Yang

Department of Mathematics, Renmin University of China, Beijing 100872, P. R. China

Abstract

Combining Carleson-Chang's result [9] with blow-up analysis, we prove existence of extremal functions for certain Trudinger-Moser inequalities in dimension two. This kind of inequality was originally proposed by Adimurthi and O. Druet [1], extended by the author to high dimensional case and Riemannian surface case [40, 41], generalized by C. Tintarev to wider cases including singular form [36] and by M. de Souza and J. M. do Ó [14] to the whole Euclidean space \mathbb{R}^2 . In addition to the Euclidean case, we also consider the Riemannian surface case. The results in the current paper complement that of L. Carleson and A. Chang [9], M. Struwe [35], M. Flucher [16], K. Lin [19], and Adimurthi-Druet [1], our previous ones [41, 26], and part of C. Tintarev [36].

Key words: Extremal function, Trudinger-Moser inequality, Blow-up analysis 2010 MSC: 46E35; 58J05

u

1. Introduction

Let Ω be a smooth bounded domain in \mathbb{R}^2 and $W_0^{1,2}(\Omega)$ be the usual Sobolev space. The classical Trudinger-Moser inequality [44, 33, 32, 37, 30] says

$$\sup_{W_0^{1,2}(\Omega), \|\nabla u\|_2 \le 1} \int_{\Omega} e^{4\pi u^2} dx < \infty.$$
⁽¹⁾

Here and throughout this paper we denote the L^p -norm by $\|\cdot\|_p$. This inequality is sharp in the sense that for any $\alpha > 4\pi$, the integrals in (1) are still finite but the supremum is infinite. Let $u_k \in W_0^{1,2}(\Omega)$ be such that $\|\nabla u_k\|_2 = 1$ and $u_k \rightarrow u$ weakly in $W_0^{1,2}(\Omega)$. Then P. L. Lions [20] proved that for any $p < 1/(1 - \|\nabla u\|_2^2)$, there holds

$$\limsup_{k \to \infty} \int_{\Omega} e^{4\pi p u_k^2} dx < \infty.$$
⁽²⁾

This inequality gives more information than the Trudinger-Moser inequality (1) in case $u \neq 0$. While in case $u \equiv 0$, it is weaker than (1). However Adimurthi and O. Druet [1] proved that for

August 3, 2018

Email address: yunyanyang@ruc.edu.cn (Yunyan Yang)

Preprint submitted to ***

any α , $0 \le \alpha < \lambda_1(\Omega)$,

$$\sup_{u \in W_0^{1,2}(\Omega), \|\nabla u\|_2 \le 1} \int_{\Omega} e^{4\pi u^2 (1+\alpha \|u\|_2^2)} dx < \infty,$$
(3)

and that the supremum is infinity when $\alpha \ge \lambda_1(\Omega)$, where $\lambda_1(\Omega)$ is the first eigenvalue of the Laplacian operator with respect to Dirichlet boundary condition. For any sequence of functions $u_k \in W_0^{1,2}(\Omega)$ with $\|\nabla u_k\|_2 = 1$ and $u_k \rightharpoonup u$ weakly in $W_0^{1,2}(\Omega)$, if $u \ne 0$, it then follows from (3) that for any $\alpha, 0 \le \alpha < \lambda_1(\Omega)$,

$$\limsup_{k \to \infty} \int_{\Omega} e^{4\pi u_k^2 (1+\alpha \|u_k\|_2^2)} dx < \infty.$$
(4)

Note that $1 + \alpha ||u_k||_2^2 < 1 + ||\nabla u||_2^2 < 1/(1 - ||\nabla u||_2^2)$ for sufficiently large k. (4) is weaker than (2). If $u \equiv 0$, we already see that (2) is weaker than (1), and obviously (4) is stronger than (1).

A natural question is to find the high dimensional analogue of (3). Let Ω be a smooth bounded domain in \mathbb{R}^n ($n \ge 3$). We proved in [40] that for any $0 \le \alpha < \lambda_1(\Omega)$,

$$\sup_{u \in W_0^{1,n}(\Omega), \|\nabla u\|_n^n \le 1} \int_{\Omega} e^{\alpha_n |u|^{\frac{n}{n-1}} (1+\alpha \|u\|_n^n)^{\frac{1}{n-1}}} dx < \infty,$$
(5)

and that the supremum is infinite when $\alpha \ge \lambda_1(\Omega)$, where $\alpha_n = n\omega_{n-1}^{1/(n-1)}$, ω_{n-1} is the area of the unit sphere in \mathbb{R}^n , and $\lambda_1(\Omega)$ is defined by

$$\lambda_1(\Omega) = \inf_{u \in W_0^{1,n}(\Omega), \, u \neq 0} \frac{\int_{\Omega} |\nabla u|^n dx}{\int_{\Omega} |u|^n dx}.$$

Trudinger-Moser inequalities on Riemannian manifolds were due to T. Aubin [7], J. Moser [30], P. Cherrier [12, 13], and L. Fontana [17]. Also a few results was recently obtained, on complete noncompact Riemannian manifolds, by G. Mancini and K. Sandeep [27, 28] and the author [43]. One may ask whether or not the analogue of (3) holds on compact Riemannian surface. Let (Σ, g) be a compact Riemannian surface without boundary. In [41], we proved the following: For any α , $0 \le \alpha < \lambda_1(\Sigma)$, there holds

$$\sup_{u \in W^{1,2}(\Sigma), \|\nabla_{g}u\|_{2} \le 1, \int_{\Sigma} u dv_{g} = 0} \int_{\Sigma} e^{4\pi u^{2}(1+\alpha\|u\|_{2}^{2})} < \infty,$$
(6)

and the supremum is infinite when $\alpha \ge \lambda_1(\Sigma)$, where $W^{1,2}(\Sigma)$ is the usual Sobolev space and $\lambda_1(\Sigma)$ is defined by

$$\lambda_1(\Sigma) = \inf_{u \in W^{1,2}(\Sigma), \int_{\Sigma} u dv_g = 0, \, u \neq 0} \frac{\int_{\Sigma} |\nabla_g u|^2 dv_g}{\int_{\Sigma} u^2 dv_g}.$$
(7)

If (Σ, g) is a compact Riemannian surface with smooth boundary, the trace Trudinger-Moser inequalities were also established in [22, 42].

Existence of extremal functions for Trudinger-Moser inequality (1) was first obtained by L. Carleson and A. Chang [9] when Ω is a unit ball. This result was extended by M. Struwe [35] to domains close to a disc in a measure sense, and by M. Flucher and K. Lin [16, 19] to general bounded smooth domains. Later these results were extended by B. Ruf [34] and Li-Ruf [24] to the whole Euclidean space. The existence result on compact Riemannian manifold was first

obtained by Y. Li [21], then by Y. Li and P. Liu [22], and by the author [39]. For existence of extremal functions for Trudinger-Moser inequality of Adimurthi-Druet type (Trudinger-Moser inequalities analogous to (3) above or (9) below), we proved in [41, 26] that supremums in (3) and (6) are attained for sufficiently small $\alpha \ge 0$, and that the supremum in (5) ($n \ge 3$) is attained for all α , $0 \le \alpha < \lambda_1(\Omega)$. In this direction, M. de Souza and J. M. do Ó [14] generalized (3) to the whole Euclidean space \mathbb{R}^2 , and the existence of extremal functions was also obtained.

Recently G. Wang and D. Ye [38] proved the existence of extremal functions for a singular Trudinger-Moser inequality. Precisely, let \mathbb{B} be a unit disc in \mathbb{R}^2 , there holds

$$\sup_{\int_{\mathbb{B}} |\nabla u|^2 dx - \int_{\mathbb{B}} \frac{u^2}{(1-|x|^2)^2} dx \le 1} \int_{\Omega} e^{4\pi u^2} dx < \infty, \tag{8}$$

and the supremum is attained. Another Trudinger-Moser inequality with interior singularity had been established by Adimurthi-Sandeep [2] on bounded smooth domain and Adimurthi and the author [4] on the whole Euclidean space. Moreover C. Tintarev [36] modified the classical Trudinger-Moser inequality as follows: Let Ω be a smooth bounded domain in \mathbb{R}^2 . There holds

$$\sup_{\int_{\Omega} |\nabla u|^2 dx - \int_{\Omega} V(x)u^2 dx \le 1} \int_{\Omega} e^{4\pi u^2} dx < \infty$$
⁽⁹⁾

for some class of V(x) > 0 including (3) and (8). For extremal functions for Trudinger-Moser inequalities on the hyperbolic space, we refer the reader to G. Mancini, K. Sandeep and C. Tintarev [29] and the references therein.

One of our goals in the current paper is to prove that the supremum in (9) is attained in case $V(x) \equiv \alpha$ with $0 \le \alpha < \lambda_1(\Omega)$. Also we consider similar problem for $0 \le \alpha < \lambda_{\ell+1}(\Omega)$, the $(\ell + 1)$ th eigenvalue of the Laplacian operator with respect to Dirichlet boundary condition. Moreover the Riemannian surface case are discussed. Our method is combining Carleson-Chang's result [9] with blow-up analysis. For earlier works involving this method, we refer the reader to [23, 40, 24, 26, 38]. Before ending this section, we remark that for results in this paper, there is a possibility of another proof, which is based on the explicit structure of putative weakly vanishing maximizing sequences as concentrating Moser functions. For details about this new method, we refer the reader to Adimurthi and C. Tintarev [3].

2. Main Results

In this paper we concern extremal functions for Trudinger-Moser inequalities of Adimurthi-Druet type. Let us first consider the Euclidean case. Let Ω be a smooth bounded domain in \mathbb{R}^2 and $\lambda_1(\Omega)$ be the first eigenvalue of the Laplacian operator with respect to Dirichlet boundary condition. Denote

$$||u||_{1,\alpha} = \left(\int_{\Omega} |\nabla u|^2 dx - \alpha \int_{\Omega} u^2 dx\right)^{1/2} \tag{10}$$

for any $u \in W_{0,2}^{1,2}(\Omega)$ with $\int_{\Omega} |\nabla u|^2 dx - \alpha \int_{\Omega} u^2 dx \ge 0$. Clearly $\|\cdot\|_{1,\alpha}$ is equivalent to the Sobolev norm $\|\cdot\|_{W_{0,2}^{1,2}(\Omega)}$ when $0 \le \alpha < \lambda_1(\Omega)$. Our first result can be stated as follows:

Theorem 1. Let Ω be a smooth bounded domain in \mathbb{R}^2 , $\lambda_1(\Omega)$ be the first eigenvalue of the Laplacian operator with respect to Dirichlet boundary condition. If $0 \leq \alpha < \lambda_1(\Omega)$, then the supremum

$$\sup_{u \in W_0^{1,2}(\Omega), \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{4\pi u^2} dx \tag{11}$$

can be attained by some function $u_0 \in W_0^{1,2}(\Omega) \cap C^1(\overline{\Omega})$ with $||u_0||_{1,\alpha} = 1$, where $|| \cdot ||_{1,\alpha}$ is defined as in (10).

Theorem 1 obviously implies C. Tintarev's inequality (9) in the case $V(x) \equiv \alpha$, and whence leads to Adimurthi and O. Druet's original inequality (3). It should be remarked that Theorem 1 does not imply that the supremum in (3) is attained for all α , $0 \le \alpha < \lambda_1(\Omega)$. Indeed, ([26], Theorem 1.2, the case p = 2) has not been improved so far. When $\alpha = 0$, Theorem 1 recovers the results of L. Carleson and A. Chang [9], M. Struwe [35], M. Flucher [16] and K. Lin [19] in dimension two.

Obviously the supremum (11) is infinite if $\alpha \ge \lambda_1(\Omega)$. It is natural to ask what we can say when other eigenvalues of the Laplacian operator are involved. Precisely, let $\lambda_1(\Omega) < \lambda_2(\Omega) < \cdots$ be all distinct eigenvalues of the Laplacian operator with respect to Dirichlet boundary condition and $E_{\lambda_j(\Omega)}$'s be associated eigenfunction spaces, namely

$$E_{\lambda_j(\Omega)} = \left\{ u \in W_0^{1,2}(\Omega) : -\Delta u = \lambda_j(\Omega) u \right\}.$$

Note that $W_0^{1,2}(\Omega)$ is a Hilbert space when it is equipped with the inner product

$$\langle u, v \rangle = \int_{\Omega} \nabla u \nabla v dx, \quad \forall u, v \in W_0^{1,2}(\Omega).$$

For any positive integer ℓ , We set

$$E_{\ell} = E_{\lambda_1(\Omega)} \oplus E_{\lambda_2(\Omega)} \oplus \cdots \oplus E_{\lambda_{\ell}(\Omega)}$$

and

$$E_{\ell}^{\perp} = \left\{ u \in W_0^{1,2}(\Omega) : \int_{\Omega} uv dx = 0, \forall v \in E_{\ell} \right\}.$$
 (12)

It is clear that

$$W_0^{1,2}(\Omega) = E_\ell \oplus E_\ell^\perp, \quad \forall \ell = 1, 2, \cdots$$

Similar to Theorem 1, we have the following:

Theorem 2. Let Ω be a smooth bounded domain in \mathbb{R}^2 , ℓ be any positive integer, $\lambda_{\ell+1}(\Omega)$ be the $(\ell + 1)$ th eigenvalue of the Laplacian operator with respect to Dirichlet boundary condition, and E_{ℓ}^{\perp} be a function space defined as in (12). Then for any α , $0 \leq \alpha < \lambda_{\ell+1}(\Omega)$, the supremum

$$\sup_{\boldsymbol{u}\in E_{\ell}^{\perp}, \|\boldsymbol{u}\|_{1,\alpha}\leq 1} \int_{\Omega} e^{4\pi u^2} dx \tag{13}$$

can be attained by some $u_0 \in E_{\ell}^{\perp} \cap C^1(\overline{\Omega})$ with $||u_0||_{1,\alpha} = 1$, where $|| \cdot ||_{1,\alpha}$ is defined as in (10).

A quite interesting case of Theorem 2 is $\alpha = 0$. It follows that for any positive integer ℓ , the supremum

$$\sup_{u\in E^{\perp}_{\ell}, \|\nabla u\|_{2}\leq 1} \int_{\Omega} e^{4\pi u^{2}} dx$$

can be attained by some $u_0 \in E_{\ell}^{\perp}$ with $\|\nabla u_0\|_2 = 1$, which is new so far. If we denote $E_0 = \{0\}$ and $E_0^{\perp} = W_0^{1,2}(\Omega)$, then Theorem 1 is exactly Theorem 2 in case that $\ell = 0$.

Now we consider the manifold case. Let (Σ, g) be a compact Riemannian surface without boundary, ∇_g and Δ_g be its gradient operator and Laplace-Beltrami operator respectively, and $\lambda_1(\Sigma)$ be the first eigenvalue of Δ_g (see (7) above). We denote

$$||u||_{1,\alpha} = \left(\int_{\Sigma} |\nabla_g u|^2 dv_g - \alpha \int_{\Sigma} u^2 dv_g\right)^{1/2} \tag{14}$$

for all $u \in W^{1,2}(\Sigma)$ with $\int_{\Sigma} |\nabla_g u|^2 dv_g - \alpha \int_{\Sigma} u^2 dv_g \ge 0$. Now we state an analogue of Theorem 1 as follows:

Theorem 3. Let (Σ, g) be a compact Riemannian surface without boundary, $\lambda_1(\Sigma)$ be the first eigenvalue of the Laplace-Beltrami operator. If $0 \le \alpha < \lambda_1(\Sigma)$, then the supremum

$$\sup_{u \in W^{1,2}(\Sigma), \int_{\Sigma} u dv_g = 0, ||u||_{1,\alpha} \le 1} \int_{\Sigma} e^{4\pi u^2} dv_g$$

can be attained by some $u_0 \in W^{1,2}(\Sigma) \cap C^1(\Sigma)$ with $\int_{\Sigma} u_0 dv_g = 0$ and $||u_0||_{1,\alpha} = 1$, where $|| \cdot ||_{1,\alpha}$ is defined as in (14).

In case $\alpha = 0$, Theorem 3 reduces to a result of Y. Li [21]. Also it should be remarked that when $0 \le \alpha < \lambda_1(\Sigma)$, the inequality

$$\Lambda_{1,\alpha} = \sup_{u \in W^{1,2}(\Sigma), \int_{\Sigma} u dv_g = 0, \|u\|_{1,\alpha} \le 1} \int_{\Sigma} e^{4\pi u^2} dv_g < +\infty$$
(15)

is stronger than that

$$\Lambda_{\alpha} = \sup_{u \in W^{1,2}(\Sigma), \int_{\Sigma} u dv_g = 0, \, \|\nabla_g u\|_2 \le 1} \int_{\Sigma} e^{4\pi u^2 (1+\alpha \|u\|_2^2)} dv_g < +\infty, \tag{16}$$

which was studied by the author in [41]. In fact, if $u \in W^{1,2}(\Sigma)$ satisfies $\int_{\Sigma} u dv_g = 0$ and $||\nabla_g u||_2 \le 1$, then $||u||_{1,\alpha}^2 \le 1 - \alpha ||u||_2^2$. Since $1 + a \le \frac{1}{1-a}$ for all a < 1, it follows from (15) that

$$\int_{\Sigma} e^{4\pi u^2 (1+\alpha ||u||_2^2)} dv_g \leq \int_{\Sigma} e^{4\pi \frac{u^2}{1-\alpha ||u||_2^2}} dv_g \leq \int_{\Sigma} e^{4\pi \frac{u^2}{||u||_{1,\alpha}^2}} dv_g \leq \Lambda_{1,\alpha}.$$

Hence we have $\Lambda_{\alpha} \leq \Lambda_{1,\alpha}$. This was also observed by C. Tintarev [36] in the Euclidean case. But we caution the reader that Theorem 3 does not imply the existence of extremal functions for (16). So it is still open whether or not extremal functions for (16) exit for all $0 \leq \alpha < \lambda_1(\Sigma)$. Let $\lambda_1(\Sigma) < \lambda_2(\Sigma) < \cdots$ be all distinct eigenvalues of the Laplace-Beltrami operator Δ_g , and $E_{\lambda_i(\Sigma)}$'s be associated eigenfunction spaces, namely

$$E_{\lambda_i(\Sigma)} = \left\{ u \in W^{1,2}(\Sigma) : \Delta_g u = \lambda_i(\Sigma) u \right\}, \quad i = 1, 2, \cdots$$

For any positive integer ℓ we write

$$E_{\ell} = E_{\lambda_1(\Sigma)} \oplus E_{\lambda_2(\Sigma)} \oplus \cdots \oplus E_{\lambda_{\ell}(\Sigma)}$$

and

$$E_{\ell}^{\perp} = \left\{ u \in W^{1,2}(\Sigma) : \int_{\Sigma} uv dv_g = 0, \forall v \in E_{\ell} \right\}.$$
(17)

Similar to Theorem 2, we have the following:

ı

Theorem 4. Let (Σ, g) be a compact Riemannian surface without boundary, ℓ be any positive integer, $\lambda_{\ell+1}(\Sigma)$ be the $(\ell+1)$ th eigenvalue of the Laplace-Beltrami operator, and E_{ℓ}^{\perp} be a function space defined as in (17). Then for any α , $0 \le \alpha < \lambda_{\ell+1}(\Sigma)$, the supremum

$$\sup_{u \in E_{\ell}^{\perp}, \int_{\Sigma} u dv_g = 0, \|u\|_{1,\alpha} \le 1} \int_{\Sigma} e^{4\pi u^2} dv_g$$

can be attained by some $u_0 \in E_{\ell}^{\perp} \cap C^1(\Sigma)$ with $\int_{\Sigma} u_0 dv_g = 0$ and $||u_0||_{1,\alpha} = 1$, where $|| \cdot ||_{1,\alpha}$ is defined as in (14).

It would be also interesting to find extremal functions for improved trace Trudinger-Moser inequality on compact Riemannian surface with smooth boundary by blow-up analysis. We would not treat this issue here, but refer the reader to B. Osgood, R. Phillips and P. Sarnak [31], P. Liu [25], Y. Li and P. Liu [22], and the author [42] for its development.

The proofs of Theorems 1 to 4 are all based on a result of Carleson-Chang [9] and blow-up analysis. Pioneer works related to this procedure can be found in Ding et al [15], Adimurthi and M. Struwe [5], Y. Li [21], Adimurthi and O. Druet [1]. Throughout this paper, $o_j(1)$ denotes the infinitesimal as $j \to \infty$, $o_{\epsilon}(1)$ denotes the infinitesimal as $\epsilon \to 0$, and so on. In addition we do not distinguish sequence and subsequence, the reader can recognize it easily from the context. Before ending this section, we quote Carleson-Chang's result [9] for our use later:

Lemma 5 (Carleson-Chang). Let \mathbb{B} be the unit disc in \mathbb{R}^2 . Assume $\{v_{\epsilon}\}_{\epsilon>0}$ is a sequence of functions in $W_0^{1,2}(\mathbb{B})$ with $\int_{\mathbb{B}} |\nabla v_{\epsilon}|^2 dx = 1$. If $|\nabla v_{\epsilon}|^2 dx \rightarrow \delta_0$ as $\epsilon \rightarrow 0$ weakly in sense of measure. Then $\limsup_{\epsilon \rightarrow 0} \int_{\mathbb{B}} (e^{4\pi v_{\epsilon}^2} - 1) dx \leq \pi e$.

The remaining part of this paper is organized as follows: In Section 3, we deal with the Euclidean case and prove Theorems 1 and 2; In Section 4, we deal with the case of manifold without boundary and prove Theorems 3 and 4.

3. The Euclidean case

In this section, using Carleson-Chang's result (Lemma 5) and blow-up analysis, we prove Theorems 1 and 2. Since the procedure is now standard [26] (for earlier works, see [15, 5, 21, 1]), we give the outline of the proof and emphasize the difference between our case and the previous ones. In particular, the essential difference between the proofs of Theorem 1 and ([26], Theorem 1.2) is the test function computation in the final step. In the proof of Theorem 2, since the maximizers u_{ϵ} 's may change signs, hence Gidas-Ni-Nirenberg's result [18] can not be applied to our case. However we can exclude the possibility of boundary blow-up via Agmon's regularity theorem ([6], page 444) in an indirect way. In the final step (test function computation), we must ensure that those test functions belong to the space E_{ℓ}^{\perp} , which is different from the counterpart of the proof of Theorem 1.

Proof of Theorem 1. Let α be fixed with $0 \le \alpha < \lambda_1(\Omega)$. We divide the proof into several steps as following:

Step 1. Maximizers for subcritical functionals

In this step, we shall prove that for any $0 < \epsilon < 4\pi$, there exists some $u_{\epsilon} \in W_0^{1,2}(\Omega) \cap C^1(\overline{\Omega})$ with $||u_{\epsilon}||_{1,\alpha} = 1$ such that

$$\int_{\Omega} e^{(4\pi-\epsilon)u_{\epsilon}^2} dx = \sup_{u \in W_0^{1,2}(\Omega), \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{(4\pi-\epsilon)u^2} dx, \tag{18}$$

where $\|\cdot\|_{1,\alpha}$ is defined as in (10). Here we do not assume in advance the above supremum is finite.

This is based on a direct method in the calculus of variations. For any $0 < \epsilon < 4\pi$, we take a sequence of functions $u_j \in W_0^{1,2}(\Omega)$ verifying that

$$\int_{\Omega} |\nabla u_j|^2 dx - \alpha \int_{\Omega} u_j^2 dx \le 1$$
(19)

and that as $j \to \infty$,

$$\int_{\Omega} e^{(4\pi-\epsilon)u_j^2} dx \to \sup_{u \in W_0^{1,2}(\Omega), \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{(4\pi-\epsilon)u^2} dx.$$
(20)

It follows from (19) and $0 \le \alpha < \lambda_1(\Omega)$ that u_j is bounded in $W_0^{1,2}(\Omega)$. Thus we can assume up to a subsequence, $u_j \rightharpoonup u_{\epsilon}$ weakly in $W_0^{1,2}(\Omega)$, $u_j \rightarrow u_{\epsilon}$ strongly in $L^2(\Omega)$, and $u_j \rightarrow u_{\epsilon}$ a.e. in Ω . Clearly we have that

$$0 \le \int_{\Omega} |\nabla u_{\epsilon}|^2 dx - \alpha \int_{\Omega} u_{\epsilon}^2 dx \le \liminf_{j \to \infty} \left(\int_{\Omega} |\nabla u_j|^2 dx - \alpha \int_{\Omega} u_j^2 dx \right) \le 1$$
(21)

and that

$$\int_{\Omega} |\nabla u_j - \nabla u_{\epsilon}|^2 dx = \int_{\Omega} |\nabla u_j|^2 dx - \int_{\Omega} |\nabla u_{\epsilon}|^2 dx + o_j(1)$$

$$\leq 1 - \int_{\Omega} |\nabla u_{\epsilon}|^2 dx + \alpha \int_{\Omega} u_{\epsilon}^2 dx + o_j(1).$$
(22)

Combining (21) and (22), we conclude

$$\limsup_{j \to \infty} \int_{\Omega} |\nabla u_j - \nabla u_{\epsilon}|^2 dx \le 1.$$

It follows from Lion's inequality (2) that $e^{(4\pi-\epsilon)u_j^2}$ is bounded in $L^q(\Omega)$ for some q > 1. Hence $e^{(4\pi-\epsilon)u_j^2} \rightarrow e^{(4\pi-\epsilon)u_\epsilon^2}$ strongly in $L^1(\Omega)$. This together with (20) immediately leads to (18). Obviously the supremum in (18) is strictly greater than $|\Omega|$, the volume of Ω . Therefore $u_\epsilon \neq 0$. If $||u_\epsilon||_{1,\alpha} < 1$, we set $\tilde{u}_\epsilon = u_\epsilon/||u_\epsilon||_{1,\alpha}$, then we obtain $||\tilde{u}_\epsilon||_{1,\alpha} = 1$ and

$$\sup_{\epsilon \in W_0^{1,2}(\Omega), \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{(4\pi - \epsilon)u^2} dx \ge \int_{\Omega} e^{(4\pi - \epsilon)\widetilde{u}_{\epsilon}^2} dx > \int_{\Omega} e^{(4\pi - \epsilon)u_{\epsilon}^2} dx.$$

This contradicts (18). Hence $||u_{\epsilon}||_{1,\alpha} = 1$.

It is not difficult to see that u_{ϵ} satisfies the Euler-Lagrange equation

$$\begin{cases} -\Delta u_{\epsilon} - \alpha u_{\epsilon} = \frac{1}{\lambda_{\epsilon}} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} \text{ in } \Omega, \\ u_{\epsilon} > 0 \text{ in } \Omega, \\ \int_{\Omega} |\nabla u_{\epsilon}|^{2} dx - \alpha \int_{\Omega} u_{\epsilon}^{2} dx = 1, \\ \lambda_{\epsilon} = \int_{\Omega} u_{\epsilon}^{2} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dx. \end{cases}$$

$$(23)$$

Applying elliptic estimates to (23), we have $u_{\epsilon} \in C^{1}(\overline{\Omega})$. Let $c_{\epsilon} = u_{\epsilon}(x_{\epsilon}) = \max_{\Omega} u_{\epsilon}$. Since $||u_{\epsilon}||_{1,\alpha} = 1$, without loss of generality, we assume u_{ϵ} converges to u^{*} weakly in $W_{0}^{1,2}(\Omega)$, strongly in $L^{2}(\Omega)$, and almost everywhere in Ω . If c_{ϵ} is bounded, then $e^{(4\pi-\epsilon)u_{\epsilon}^{2}}$ is bounded in $L^{\infty}(\Omega)$, and thus $e^{(4\pi-\epsilon)u_{\epsilon}^{2}}$ converges to $e^{4\pi u^{*2}}$ in $L^{1}(\Omega)$. Hence for any $u \in W_{0}^{1,2}(\Omega)$ with $||u||_{1,\alpha} \leq 1$, we have by (18) that

$$\int_{\Omega} e^{4\pi u^2} dx = \lim_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)u^2} dx \le \lim_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)u_{\epsilon}^2} dx = \int_{\Omega} e^{4\pi u^{*2}} dx.$$

This implies that

$$\sup_{u \in W_0^{1,2}(\Omega), \|u\|_{1,a} \le 1} \int_{\Omega} e^{4\pi u^2} dx = \int_{\Omega} e^{4\pi u^{*2}} dx.$$
(24)

So $u^* \in W_0^{1,2}(\Omega)$ attains the above supremum. Obviously $||u^*||_{1,\alpha} = 1$. Applying elliptic estimates to its Euler-Lagrange equation, we obtain $u^* \in C^1(\overline{\Omega})$. Therefore u^* is the desired extremal function. Hence we assume $c_{\epsilon} \to \infty$ in the sequel. Without loss of generality, we assume $x_{\epsilon} \to x_0 \in \overline{\Omega}$. By a result of Gidas-Ni-Nirenberg ([18], page 223), the distance between x_{ϵ} and $\partial\Omega$ must be greater than $\delta > 0$ depending only on Ω . Therefore $x_0 \notin \partial\Omega$.

Step 2. Energy concentration phenomenon

In this step we shall prove that $u_{\epsilon} \to 0$ weakly in $W_0^{1,2}(\Omega)$, $u_{\epsilon} \to 0$ strongly in $L^q(\Omega)$ for any q > 1, and $|\nabla u_{\epsilon}|^2 dx \to \delta_{x_0}$ weakly in sense of measure as $\epsilon \to 0$, where δ_{x_0} is the usual Dirac measure centered at x_0 .

Noting that $||u_{\epsilon}||_{1,\alpha} = 1$, we can assume $u_{\epsilon} \rightharpoonup u_0$ weakly in $W_0^{1,2}(\Omega)$, and $u_{\epsilon} \rightarrow u_0$ strongly in $L^q(\Omega)$ for any q > 1. It follows that

$$\int_{\Omega} |\nabla u_{\epsilon}|^2 dx = 1 + \alpha \int_{\Omega} u_0^2 dx + o(1),$$
(25)

and that

$$\int_{\Omega} |\nabla (u_{\epsilon} - u_0)|^2 dx = 1 - \int_{\Omega} |\nabla u_0|^2 dx + \alpha \int_{\Omega} u_0^2 dx + o(1).$$
(26)

Suppose $u_0 \neq 0$. In view of (26), Lions' inequality (2) implies that $e^{4\pi u_{\epsilon}^2}$ is bounded in $L^q(\Omega)$ for any fixed q with $1 < q < 1/(1 - ||u_0||_{1,\alpha}^2)$. Then applying elliptic estimates to (23), we have that u_{ϵ} is uniformly bounded in Ω , which contradicts $c_{\epsilon} \to \infty$. Therefore $u_0 \equiv 0$ and (25) becomes

$$\int_{\Omega} |\nabla u_{\epsilon}|^2 dx = 1 + o_{\epsilon}(1).$$
⁽²⁷⁾

Suppose $|\nabla u_{\epsilon}|^2 dx \rightarrow \mu$ in sense of measure. If $\mu \neq \delta_{x_0}$, then in view of (27) and $u_0 \equiv 0$, we can choose some $r_0 > 0$ and a cut-off function $\phi \in C_0^1(\mathbb{B}_{r_0}(x_0))$, which is equal to 1 on $\mathbb{B}_{r_0/2}(x_0)$, such that $\mathbb{B}_{r_0}(x_0) \subset \Omega$ and

$$\limsup_{\epsilon \to 0} \int_{\mathbb{B}_{r_0}(x_0)} |\nabla(\phi u_{\epsilon})|^2 dx < 1.$$

By the classical Trudinger-Moser inequality (1), $e^{(4\pi-\epsilon)(\phi u_{\epsilon})^2}$ is bounded in $L^r(\mathbb{B}_{r_0}(x_0))$ for some r > 1. Applying elliptic estimates to (23), we have that u_{ϵ} is uniformly bounded in $\mathbb{B}_{r_0/2}(x_0)$, which contradicts $c_{\epsilon} \to \infty$ again. Therefore $|\nabla u_{\epsilon}|^2 dx \to \delta_{x_0}$ and Step 2 is finished.

Step 3. Blow-up analysis for u_{ϵ}

We set

$$r_{\epsilon} = \sqrt{\lambda_{\epsilon}} c_{\epsilon}^{-1} e^{-(2\pi - \epsilon/2)c_{\epsilon}^2}$$

For any $0 < \delta < 4\pi$, we have by using the Hölder inequality and the classical Trudinger-Moser inequality (1),

$$\lambda_{\epsilon} = \int_{\Omega} u_{\epsilon}^2 e^{(4\pi - \epsilon)u_{\epsilon}^2} dx \le e^{\delta c_{\epsilon}^2} \int_{\Omega} u_{\epsilon}^2 e^{(4\pi - \epsilon - \delta)u_{\epsilon}^2} dx \le C e^{\delta c_{\epsilon}^2}$$

for some constant C depending only on δ . This leads to

$$r_{\epsilon}^{2} \leq C c_{\epsilon}^{-2} e^{-(4\pi - \epsilon - \delta)c_{\epsilon}^{2}} \to 0 \quad \text{as} \quad \epsilon \to 0.$$
⁽²⁸⁾

Let

$$\Omega_{\epsilon} = \{ x \in \mathbb{R}^2 : x_{\epsilon} + r_{\epsilon} x \in \Omega \}.$$

Define two blow-up sequences of functions on Ω_{ϵ} as

$$\psi_{\epsilon}(x) = c_{\epsilon}^{-1} u_{\epsilon}(x_{\epsilon} + r_{\epsilon}x), \quad \varphi_{\epsilon}(x) = c_{\epsilon}(u_{\epsilon}(x_{\epsilon} + r_{\epsilon}x) - c_{\epsilon}).$$

A direct computation shows

$$-\Delta\psi_{\epsilon} = \alpha r_{\epsilon}^{2}\psi_{\epsilon} + c_{\epsilon}^{-2}\psi_{\epsilon}e^{(4\pi-\epsilon)(u_{\epsilon}^{2}-c_{\epsilon}^{2})} \quad \text{in} \quad \Omega_{\epsilon},$$
⁽²⁹⁾

$$-\Delta\varphi_{\epsilon} = \alpha r_{\epsilon}^2 c_{\epsilon}^2 \psi_{\epsilon} + \psi_{\epsilon} e^{(4\pi - \epsilon)(1 + \psi_{\epsilon})\varphi_{\epsilon}} \quad \text{in} \quad \Omega_{\epsilon}.$$
(30)

We now investigate the convergence behavior of ψ_{ϵ} and φ_{ϵ} . Note that $\Omega_{\epsilon} \to \mathbb{R}^2$ as $\epsilon \to 0$. Since $|\psi_{\epsilon}| \le 1$ and $\Delta \psi_{\epsilon}(x) \to 0$ uniformly in $x \in \Omega_{\epsilon}$ as $\epsilon \to 0$, we have by elliptic estimates that $\psi_{\epsilon} \to \psi$

in $C^1_{\text{loc}}(\mathbb{R}^2)$, where ψ is a bounded harmonic function in \mathbb{R}^2 . Note that $\psi(0) = \lim_{\epsilon \to 0} \psi_{\epsilon}(0) = 1$. The Liouville theorem implies that $\psi \equiv 1$ on \mathbb{R}^2 . Thus we have

$$\psi_{\epsilon} \to 1 \quad \text{in} \quad C^1_{\text{loc}}(\mathbb{R}^2).$$
 (31)

By (28), we have $r_{\epsilon}^2 c_{\epsilon}^2 \to 0$ as $\epsilon \to 0$. Note also that

$$\varphi_{\epsilon}(x) \leq \varphi_{\epsilon}(0) = 0$$
 for all $x \in \Omega_{\epsilon}$

Thus $\Delta \varphi_{\epsilon}$ is uniformly bounded in Ω_{ϵ} . We then conclude by applying elliptic estimates to the equation (30) that

$$\varphi_{\epsilon} \to \varphi \quad \text{in} \quad C^1_{\text{loc}}(\mathbb{R}^2),$$
(32)

where φ satisfies

$$\Delta \varphi = -e^{8\pi\varphi} \quad \text{in} \quad \mathbb{R}^2$$
$$\varphi(0) = 0 = \sup_{\mathbb{R}^2} \varphi$$
$$\int_{\mathbb{R}^2} e^{8\pi\varphi} dx \le 1.$$

By a result of Chen-Li [11], we have

$$\varphi(x) = -\frac{1}{4\pi} \log(1 + \pi |x|^2)$$
(33)

and

$$\int_{\mathbb{R}^2} e^{8\pi\varphi} dx = 1$$

To understand the convergence behavior away from the blow-up point x_0 , we need to investigate how $c_{\epsilon}u_{\epsilon}$ converges. By a repetitive argument of ([26], Lemma 3.6), we have that

$$c_{\epsilon}u_{\epsilon} \rightharpoonup G$$
 weakly in $W_0^{1,q}(\Omega), \quad \forall 1 < q < 2,$ (34)

where $G \in C^1(\overline{\Omega} \setminus \{x_0\})$ is the Green function satisfying the equation

$$\begin{cases} -\Delta G - \alpha G = \delta_{x_0} & \text{in } \Omega \\ G = 0 & \text{on } \partial \Omega. \end{cases}$$
(35)

Moreover,

$$c_{\epsilon}u_{\epsilon} \to G \quad \text{in} \quad C^{1}_{\text{loc}}(\overline{\Omega} \setminus \{x_{0}\}).$$
 (36)

Step 4. Upper bound estimate

In view of (35) and (36), G can be represented by

$$G = -\frac{1}{2\pi} \log |x - x_0| + A_{x_0} + \psi_{\alpha}(x), \qquad (37)$$

where A_{x_0} is a constant depending on x_0 and $\alpha, \psi_{\alpha} \in C^1(\Omega)$ and $\psi_{\alpha}(x_0) = 0$. This leads to

$$\int_{\Omega \setminus B_{\delta}(x_0)} |\nabla G|^2 dx = \alpha \int_{\Omega \setminus B_{\delta}(x_0)} G^2 dx + \int_{\partial (\Omega \setminus B_{\delta}(x_0))} G \frac{\partial G}{\partial n} ds$$
$$= \frac{1}{2\pi} \log \frac{1}{\delta} + A_{x_0} + \alpha ||G||_2^2 + o_{\delta}(1).$$

Hence we obtain

$$\int_{\Omega \setminus B_{\delta}(x_0)} |\nabla u_{\epsilon}|^2 dx = \frac{1}{c_{\epsilon}^2} \left(\frac{1}{2\pi} \log \frac{1}{\delta} + A_{x_0} + \alpha ||G||_2^2 + o_{\delta}(1) + o_{\epsilon}(1) \right).$$
(38)

Let $s_{\epsilon} = \sup_{\partial B_{\delta}(x_0)} u_{\epsilon}$ and $\overline{u}_{\epsilon} = (u_{\epsilon} - s_{\epsilon})^+$. Then $\overline{u}_{\epsilon} \in W_0^{1,2}(B_{\delta}(x_0))$. By (38) and the fact that $\int_{B_{\delta}(x_0)} |\nabla u_{\epsilon}|^2 dx = 1 - \int_{\Omega \setminus B_{\delta}(x_0)} |\nabla u_{\epsilon}|^2 dx + \alpha \int_{\Omega} u_{\epsilon}^2$, we have

$$\int_{B_{\delta}(x_0)} |\nabla \overline{u}_{\epsilon}|^2 dx \le \tau_{\epsilon} = 1 - \frac{1}{c_{\epsilon}^2} \left(\frac{1}{2\pi} \log \frac{1}{\delta} + A_{x_0} + o_{\delta}(1) + o_{\epsilon}(1) \right). \tag{39}$$

This together with Lemma 5 (see the end of Section 2) leads to

$$\limsup_{\epsilon \to 0} \int_{B_{\delta}(x_0)} (e^{4\pi \overline{u}_{\epsilon}^2/\tau_{\epsilon}} - 1) dx \le \pi \delta^2 e.$$
⁽⁴⁰⁾

By (32), we have on $B_{Rr_{\epsilon}}(x_{\epsilon})$ that $u_{\epsilon}(x) = c_{\epsilon} + \frac{1}{c_{\epsilon}}\varphi(\frac{x-x_{\epsilon}}{r_{\epsilon}})$, which together with the fact that $c_{\epsilon}u_{\epsilon} \to G$ in $L^{2}(\Omega)$, gives on $B_{Rr_{\epsilon}}(x_{\epsilon})$,

$$\begin{aligned} (4\pi - \epsilon)u_{\epsilon}^{2} &\leq 4\pi(\overline{u}_{\epsilon} + s_{\epsilon})^{2} \\ &\leq 4\pi\overline{u}_{\epsilon}^{2} + 8\pi s_{\epsilon}\overline{u}_{\epsilon} + o_{\epsilon}(1) \\ &\leq 4\pi\overline{u}_{\epsilon}^{2} - 4\log\delta + 8\pi A_{x_{0}} + o_{\epsilon}(1) + o_{\delta}(1) \\ &\leq 4\pi\overline{u}_{\epsilon}^{2}/\tau_{\epsilon} - 2\log\delta + 4\pi A_{x_{0}} + o(1). \end{aligned}$$

Therefore

$$\begin{split} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi-\epsilon)u_{\epsilon}^{2}} dx &\leq \delta^{-2} e^{4\pi A_{x_{0}}+o(1)} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{4\pi \overline{u}_{\epsilon}^{2}/\tau_{\epsilon}} dx \\ &= \delta^{-2} e^{4\pi A_{x_{0}}+o(1)} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} (e^{4\pi \overline{u}_{\epsilon}^{2}/\tau_{\epsilon}}-1) dx + o(1) \\ &\leq \delta^{-2} e^{4\pi A_{x_{0}}+o(1)} \int_{B_{\delta}(x_{0})} (e^{4\pi \overline{u}_{\epsilon}^{2}/\tau_{\epsilon}}-1) dx. \end{split}$$

This together with (40) leads to

$$\limsup_{\epsilon \to 0} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi - \epsilon)u_{\epsilon}^2} dx \le \pi e^{1 + 4\pi A_{x_0}}.$$
(41)

By the same argument as in the proof of ([26], Lemma 3.3), we get

$$\lim_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dx \le |\Omega| + \lim_{R \to +\infty} \limsup_{\epsilon \to 0} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dx$$
(42)

Combining (41) and (42), we conclude

$$\sup_{u \in W_0^{1,2}(\Omega), \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{4\pi u^2} dx = \limsup_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)u_{\epsilon}^2} dx \le |\Omega| + \pi e^{1 + 4\pi A_{x_0}}.$$
 (43)

Step 5. Existence of extremal functions

We will construct a sequence of functions $\phi_{\epsilon} \in W_0^{1,2}(\Omega)$ such that $\|\phi_{\epsilon}\|_{1,\alpha} = 1$ and

$$\int_{\Omega} e^{4\pi\phi_{\epsilon}^2} dx > |\Omega| + \pi e^{1+4\pi A_{x_0}}$$

$$\tag{44}$$

for sufficiently small $\epsilon > 0$. The contradiction between (43) and (44) implies that c_{ϵ} must be bounded. Then applying elliptic estimates to (23), we conclude the existence of extremal function and finish the proof of Theorem 1.

To prove (44), we recall (37) and write $r(x) = |x - x_0|$. Set

$$\phi_{\epsilon} = \begin{cases} c + \frac{-\frac{1}{4\pi}\log(1+\pi\frac{r^{2}}{\epsilon^{2}})+B}{c} & \text{for } r \leq R\epsilon \\ \frac{G-\eta\psi_{\alpha}}{c} & \text{for } R\epsilon < r < 2R\epsilon \\ \frac{G}{c} & \text{for } r \geq 2R\epsilon, \end{cases}$$
(45)

where $R = -\log \epsilon$, $\eta \in C_0^{\infty}(B_{2R\epsilon}(x_0))$ verifying that $\eta = 1$ on $B_{R\epsilon}(x_0)$ and $\|\nabla \eta\|_{L^{\infty}} = O(\frac{1}{R\epsilon})$, *B* is a constant to be determined later, and *c* depending only on ϵ will also be chosen later such that $R\epsilon \to 0$ and $R \to +\infty$. In order to assure that $\phi_{\epsilon} \in W_0^{1,2}(\Omega)$, we set

$$c + \frac{1}{c} \left(-\frac{1}{4\pi} \log(1 + \pi R^2) + B \right) = \frac{1}{c} \left(-\frac{1}{2\pi} \log(R\epsilon) + A_{x_0} \right),$$

which gives

$$2\pi c^{2} = -\log \epsilon - 2\pi B + 2\pi A_{x_{0}} + \frac{1}{2}\log \pi + O(\frac{1}{R^{2}}).$$
(46)

A delicate but straightforward calculation shows

$$\begin{split} \int_{\Omega} |\nabla \phi_{\epsilon}|^2 dx &= \frac{1}{4\pi c^2} \left(2\log \frac{1}{\epsilon} + \log \pi - 1 + 4\pi A_{x_0} + 4\pi \alpha ||G||_2^2 \right. \\ &+ O(\frac{1}{R^2}) + O(R\epsilon \log(R\epsilon)) \right) \end{split}$$

and

$$\int_{\Omega} \phi_{\epsilon}^2 dx = \frac{1}{c^2} \left(\int_{\Omega} G^2 dx + O(R\epsilon \log(R\epsilon)) \right),$$

which yields

$$\begin{aligned} \|\phi_{\epsilon}\|_{1,\alpha}^{2} &= \int_{\Omega} (|\nabla\phi_{\epsilon}|^{2} - \alpha\phi_{\epsilon}^{2}) dx \\ &= \frac{1}{4\pi c^{2}} \left(2\log\frac{1}{\epsilon} + \log\pi - 1 + 4\pi A_{x_{0}} + O(\frac{1}{R^{2}}) + O(R\epsilon\log(R\epsilon)) \right) \end{aligned}$$

Set $\|\phi_{\epsilon}\|_{1,\alpha} = 1$, we have

$$c^{2} = -\frac{\log \epsilon}{2\pi} + \frac{\log \pi}{4\pi} - \frac{1}{4\pi} + A_{x_{0}} + O(\frac{1}{R^{2}}) + O(R\epsilon \log(R\epsilon)).$$
(47)

It follows from (46) and (47) that

$$B = \frac{1}{4\pi} + O(\frac{1}{R^2}) + O(R\epsilon \log(R\epsilon)).$$
(48)

Clearly we have on $B_{R\epsilon}(x_0)$

$$4\pi\phi_\epsilon^2 \geq 4\pi c^2 - 2\log(1+\pi\frac{r^2}{\epsilon^2}) + 8\pi B$$

This together with (47) and (48) yields

$$\int_{B_{R\epsilon}(x_0)} e^{4\pi\phi_{\epsilon}^2} dx \ge \pi e^{1+4\pi A_{x_0}} + O(\frac{1}{R^2}).$$
(49)

On the other hand,

$$\int_{\Omega \setminus B_{R\epsilon}(x_0)} e^{4\pi \phi_{\epsilon}^2} dx \geq \int_{\Omega \setminus B_{2R\epsilon}(x_0)} (1 + 4\pi \phi_{\epsilon}^2) dx$$

$$\geq |\Omega| + 4\pi \frac{||G||_2^2}{c^2} + o(\frac{1}{c^2}).$$
(50)

Recalling (47) and the choice of $R = -\log \epsilon$, we conclude (44) for sufficiently small $\epsilon > 0$ by combining (49) and (50).

Before proving Theorem 2, we state a special version of a regularity theorem due to S. Agmon ([6], page 444), which is essential for excluding boundary blow-up.

Lemma 6. Let Ω be a smooth bounded domain in \mathbb{R}^2 , $u \in L^r(\Omega)$ for some r > 1, and $f \in L^q(\Omega)$ for some q > 1. Suppose that for all functions $v \in C^2(\overline{\Omega}) \cap W_0^{1,q}(\Omega)$,

$$\int_{\Omega} u \Delta v dx = \int_{\Omega} f v dx.$$

Then $u \in W^{2,q}(\Omega) \cap W^{1,q}_0(\Omega)$.

Proof of Theorem 2. Firstly, we fix several notations concerning the function space E_{ℓ}^{\perp} defined as in (12). Let $\lambda_1(\Omega) < \lambda_2(\Omega) < \cdots$ be all distinct eigenvalues of the Laplacian with respect to Dirichlet boundary condition, and $E_{\lambda_i(\Omega)}$'s be associated eigenfunction spaces. It is known that $\lambda_i(\Omega) \to +\infty$ as $i \to +\infty$ and each space $E_{\lambda_i(\Omega)}$ has finite dimension (see [8], Theorem 9.31). We can assume

$$\dim E_{\lambda_i(\Omega)} = n_i, \quad i = 1, 2, \cdots$$

Moreover we can find a basis (e_{ij}) $(1 \le j \le n_i, 1 \le i \le \ell)$ of E_ℓ verifying

$$\begin{cases} E_{\lambda_{i}(\Omega)} = \operatorname{span}\{e_{i1}, \cdots, e_{in_{i}}\}, & i = 1, \cdots, \ell, \\ E_{\ell} = \operatorname{span}\{e_{11}, \cdots, e_{1n_{1}}, e_{21}, \cdots, e_{2n_{2}}, \cdots, e_{\ell 1}, \cdots, e_{\ell n_{\ell}}\}, \\ \int_{\Omega} |e_{ij}|^{2} dx = 1, \\ \int_{\Omega} e_{ij} e_{kl} dx = 0, & i \neq k \text{ or } j \neq l. \\ 13 \end{cases}$$
(51)

Note that

$$E_{\ell}^{\perp} = \left\{ u \in W_0^{1,2}(\Omega) : \int_{\Omega} u e_{ij} dx = 0, 1 \le j \le n_i, 1 \le i \le \ell \right\}.$$

Secondly, let $0 \le \alpha < \lambda_{\ell+1}(\Omega)$ be fixed, we shall find maximizers for subcritical Trudinger-Moser functionals. Analogous to Step 1 of the proof of Theorem 1, for any ϵ , $0 < \epsilon < 4\pi$, there exists some $u_{\epsilon} \in E_{\ell}^{\perp} \cap C^{1}(\overline{\Omega})$ with $||u_{\epsilon}||_{1,\alpha} = 1$ such that

$$\int_{\Omega} e^{(4\pi-\epsilon)u_{\epsilon}^2} dx = \sup_{u \in E_{\ell}^{\perp}, \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{(4\pi-\epsilon)u^2} dx,$$
(52)

where $\|\cdot\|_{1,\alpha}$ is defined as in (10). Moreover u_{ϵ} satisfies the Euler-Lagrange equation

$$-\Delta u_{\epsilon} - \alpha u_{\epsilon} = \frac{1}{\lambda_{\epsilon}} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_{i}} \frac{\gamma_{ij,\epsilon}}{\lambda_{\epsilon}} e_{ij} \text{ in } \Omega,$$

$$u_{\epsilon} \in E_{\ell}^{\perp} \cap C^{1}(\overline{\Omega}),$$

$$\lambda_{\epsilon} = \int_{\Omega} u_{\epsilon}^{2} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dx,$$

$$\gamma_{ij,\epsilon} = \int_{\Omega} e_{ij} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dx.$$
(53)

Without loss of generality we can assume

$$u_{\epsilon} \rightharpoonup u_0$$
 weakly in $W_0^{1,2}(\Omega)$, (54)

$$u_{\epsilon} \to u_0$$
 strongly in $L^p(\Omega), \ \forall p > 1,$ (55)

$$u_{\epsilon} \to u_0 \quad \text{a.e. in} \quad \Omega.$$
 (56)

Since $u_{\epsilon} \in E_{\ell}^{\perp}$, we have by (55)

$$\int_{\Omega} u_0 e_{ij} dx = \lim_{\epsilon \to 0} \int_{\Omega} u_\epsilon e_{ij} dx = 0, \quad 1 \le j \le n_i, \quad 1 \le i \le \ell,$$

which together with (54) implies that $u_0 \in E_{\ell}^{\perp}$ and $||u_0||_{1,\alpha} \leq 1$.

If u_{ϵ} is bounded in $C^{0}(\overline{\Omega})$, then for any $v \in E_{\ell}^{\perp}$ with $||v||_{1,\alpha} \leq 1$, (52), (56) and Lebesgue's dominated convergence theorem lead to

$$\int_{\Omega} e^{4\pi v^2} dx = \lim_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)v^2} dx \le \lim_{\epsilon \to 0} \int_{\Omega} e^{(4\pi - \epsilon)u_{\epsilon}^2} dx = \int_{\Omega} e^{4\pi u_0^2} dx.$$

Hence we have

$$\int_{\Omega} e^{4\pi u_0^2} dx = \sup_{u \in E_{\ell}^{\perp}, \|u\|_{1,a} \le 1} \int_{\Omega} e^{4\pi u^2} dx.$$
 (57)

It is easy to see that $||u_0||_{1,\alpha} = 1$. Applying elliptic estimates to the Euler-Lagrange equation of u_0 , we have $u_0 \in C^1(\overline{\Omega})$. Thus u_0 is the desired extremal function.

In the sequel we assume up to a subsequence

$$\|u_{\epsilon}\|_{C^{0}(\overline{\Omega})} = \max_{\overline{\Omega}} |u_{\epsilon}| \to +\infty \text{ as } \epsilon \to 0.$$

Thirdly, we perform blow-up analysis. Denote $c_{\epsilon} = |u_{\epsilon}(x_{\epsilon})| = ||u_{\epsilon}||_{C^{0}(\overline{\Omega})}$. Then $c_{\epsilon} \to +\infty$ as $\epsilon \to 0$. Without loss of generality we assume $c_{\epsilon} = u_{\epsilon}(x_{\epsilon})$. For otherwise u_{ϵ} can be replaced by

 $-u_{\epsilon}$ in the following blow-up analysis. Then up to a subsequence, $x_{\epsilon} \to x_0 \in \overline{\Omega}$. As in Step 2 of the proof of Theorem 1, we have $u_0 \equiv 0$ and $|\nabla u_{\epsilon}|^2 dx \to \delta_{x_0}$ weakly in sense of measure. The only difference is that $\phi u_{\epsilon} \in W_0^{1,2}(\mathbb{B}_r(x_0) \cap \Omega)$ in case $x_0 \in \partial\Omega$.

Set

$$\psi_{\epsilon}(x) = c_{\epsilon}^{-1} u_{\epsilon}(x_{\epsilon} + r_{\epsilon}x), \quad \varphi_{\epsilon}(x) = c_{\epsilon}(u_{\epsilon}(x_{\epsilon} + r_{\epsilon}x) - c_{\epsilon}), \quad x \in \Omega_{\epsilon},$$

where

$$r_{\epsilon} = \sqrt{\lambda_{\epsilon}} c_{\epsilon}^{-1} e^{-(2\pi - \epsilon/2)c_{\epsilon}^{2}}$$
(58)

and

$$\Omega_{\epsilon} = \{ x \in \mathbb{R}^2 : x_{\epsilon} + r_{\epsilon} x \in \Omega \}.$$

By (28), we have $r_{\epsilon} \to 0$ as $\epsilon \to 0$. Moreover we claim that up to a subsequence

$$r_{\epsilon}/\operatorname{dist}(x_{\epsilon},\partial\Omega) \to 0 \quad \text{as} \quad \epsilon \to 0.$$
 (59)

Let \mathbb{B} be a unit disc centered at $0 \in \mathbb{R}^2$. Since Ω is smooth, we have a a neighborhood $U \subset \mathbb{R}^2$ of x_0 and a bijective map $H : \mathbb{B} \to U$ such that $H \in C^2(\overline{\mathbb{B}})$, $J = H^{-1} \in C^2(\overline{U})$, $H(\mathbb{B}^+) = \Omega \cap U$, $H(\mathbb{B}_0) = \partial \Omega \cap U$. Here we denote $\mathbb{B}^+ = \mathbb{B} \cap \mathbb{R}^2_+$, $\mathbb{B}^0 = \mathbb{B} \cap \partial \mathbb{R}^2_+$, and $\mathbb{R}^2_+ = \{(x_1, x_2) \in \mathbb{R}^2 : x_2 > 0\}$. We write x = H(y) and $y = H^{-1}(x) = J(x)$. Furthermore we can assume (up to a linear transformation) the Jacobian matrix Jac H satisfies

$$\operatorname{Jac} H(0) = \left(\frac{\partial H_i}{\partial y_j}\right)_{y=0}, \ \frac{\partial H_i}{\partial y_j}(0) = \delta_{ij} = \begin{cases} 1, & i = j, \\ 0, & i \neq j. \end{cases}$$
(60)

In view of (53), we have

$$\int_{\Omega \cap U} \nabla u_{\epsilon} \nabla \varphi dx = \int_{\Omega \cap U} g_{\epsilon} \varphi dx, \quad \forall \varphi \in C_0^1(\Omega \cap U), \tag{61}$$

where

$$g_{\epsilon} = \frac{1}{\lambda_{\epsilon}} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} + \alpha u_{\epsilon} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_{i}} \frac{\gamma_{ij,\epsilon}}{\lambda_{\epsilon}} e_{ij}$$

Set

$$\widetilde{u}_{\epsilon}(y) = u_{\epsilon}(H(y)), \quad y \in \mathbb{B}^+.$$

Then (61) is transferred to

$$\sum_{k,\ell=1}^{2} \int_{\mathbb{B}^{+}} a_{k\ell} \frac{\partial \widetilde{u}_{\epsilon}}{\partial y_{k}} \frac{\partial \psi}{\partial y_{\ell}} dy = \int_{\mathbb{B}^{+}} \widetilde{g}_{\epsilon} \psi dy, \quad \forall \psi \in C_{0}^{1}(\mathbb{B}^{+}),$$
(62)

where

$$\widetilde{g}_{\epsilon} = (g_{\epsilon} \circ H) |\det \operatorname{Jac} H|,$$
$$a_{k,\ell} = \sum_{j=1}^{2} \frac{\partial J_k}{\partial x_j} \frac{\partial J_\ell}{\partial x_j} |\det \operatorname{Jac} H|,$$

and detJac *H* denotes the determinant of the Jacobian matrix Jac *H*. Note that $a_{k,\ell} \in C^1(\overline{\mathbb{B}^+})$ and that its ellipticity condition is satisfied.

Denote $\widetilde{x}_{\epsilon} = J(x_{\epsilon}) = (\widetilde{x}_{1,\epsilon}, \widetilde{x}_{2,\epsilon})$ and $\widetilde{x}'_{\epsilon} = (\widetilde{x}_{1,\epsilon}, 0)$. Set

$$v_{\epsilon}(y) = \frac{1}{c_{\epsilon}}\widetilde{u}_{\epsilon}(\vec{x}_{\epsilon} + r_{\epsilon}y), \quad y \in \overline{\mathbb{B}_{\epsilon}^{+}} = \left\{ y \in \mathbb{R}^{2} : \vec{x}_{\epsilon}' + r_{\epsilon}y \in \overline{\mathbb{B}} \right\}.$$

It follows from (62) that v_{ϵ} is a weak solution to the equation

$$-\frac{\partial}{\partial y_{\ell}}\left(a_{k\ell}(\widetilde{x}_{\epsilon}'+r_{\epsilon}y)\frac{\partial v_{\epsilon}}{\partial y_{k}}(y)\right) = \frac{r_{\epsilon}^{2}}{c_{\epsilon}}\widetilde{g}_{\epsilon}(\widetilde{x}_{\epsilon}'+r_{\epsilon}y), \quad y \in \mathbb{B}_{\epsilon}^{+}.$$
(63)

On one hand, by the definition of r_{ϵ} (see (58)), we have $r_{\epsilon}^2 c_{\epsilon}^{-1} \widetilde{g}_{\epsilon}(\widetilde{x}_{\epsilon} + r_{\epsilon} y)$ tends to zero uniformly in $y \in \mathbb{B}_{\epsilon}^+$ as $\epsilon \to 0$. On the other hand we have $|v_{\epsilon}(y)| \leq 1$ for all $y \in \mathbb{B}_{\epsilon}^+$. Note that $\mathbb{B}_{\epsilon}^+ \to \mathbb{R}_{+}^2$. Applying elliptic estimates to (63) and noticing (60), we obtain $v_{\epsilon} \to v$ in $C_{loc}^1(\mathbb{R}_{+}^2)$, where v satisfies

$$\begin{cases} -\Delta v = 0 & \text{in} & \mathbb{R}^2_+ \\ v = 0 & \text{on} & \partial \mathbb{R}^2_+ \\ |v| \le 1 & \text{in} & \mathbb{R}^2_+. \end{cases}$$

Obviously *v* can be extended to a bounded weak harmonic function in the whole \mathbb{R}^2 . Since v = 0 on $\partial \mathbb{R}^2_+$, Liouville theorem implies that $v \equiv 0$.

We now suppose that there exists some positive number v independent of ϵ such that

$$r_{\epsilon}/\operatorname{dist}(x_{\epsilon},\partial\Omega) \ge \nu > 0.$$
 (64)

We can find some constant C depending only on v and the bijective map H such that

$$\left|\widetilde{x}_{\epsilon} - \widetilde{x}_{\epsilon}'\right| \le Cr_{\epsilon}.$$

Note that

$$v_{\epsilon}\left(\frac{\widetilde{x}_{\epsilon}-\widetilde{x}_{\epsilon}}{r_{\epsilon}}\right) = \frac{1}{c_{\epsilon}}\widetilde{u}_{\epsilon}(\widetilde{x}_{\epsilon}) = \frac{1}{c_{\epsilon}}u_{\epsilon}(x_{\epsilon}) = 1.$$

We have $||v||_{L^{\infty}(\mathbb{B}^+_{2C})} = 1$, where $\mathbb{B}^+_{2C} = \{y \in \mathbb{R}^2_+ : |y| \le 2C\}$, since $v_{\epsilon} \to v$ in $C^1_{\text{loc}}(\overline{\mathbb{R}^2_+})$. This contradicts $v \equiv 0$. Therefore (64) is false and our claim (59) follows.

In view of (59), we conclude that

$$\Omega_{\epsilon} \to \mathbb{R}^2$$
 as $\epsilon \to 0$.

Using the argument in Step 2 of the proof of Theorem 1, we have

$$\begin{split} \psi_{\epsilon} &\to 1 \quad \text{in} \quad C^{1}_{\text{loc}}(\mathbb{R}^{2}), \\ \varphi_{\epsilon} &\to -\frac{1}{4\pi} \log(1+\pi|x|^{2}) \quad \text{in} \quad C^{1}_{\text{loc}}(\mathbb{R}^{2}), \\ c_{\epsilon}u_{\epsilon} &\to G \text{ weakly in } W^{1,q}_{0}(\Omega), \ \forall 1 < q < 2, \\ c_{\epsilon}u_{\epsilon} &\to G \text{ in } C^{1}_{\text{loc}}(\overline{\Omega} \setminus \{x_{0}\}), \\ 16 \end{split}$$

where *G* is a distributional solution to $-\Delta G - \alpha G = \delta_{x_0} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_i} e_{ij}(x_0) e_{ij}$, or equivalently

$$-\int_{\Omega} G\Delta\varphi dx + \alpha \int_{\Omega} G\varphi dx = \varphi(x_0) - \sum_{i=1}^{\ell} \sum_{j=1}^{n_i} e_{ij}(x_0) \int_{\Omega} \varphi e_{ij} dx, \quad \forall \varphi \in C^2(\overline{\Omega}).$$
(65)

Moreover,

$$\int_{\Omega} Ge_{ij} dx = \lim_{\epsilon \to 0} \int_{\Omega} c_{\epsilon} u_{\epsilon} e_{ij} dx = 0, \quad \forall 1 \le j \le n_i, \ 1 \le i \le \ell.$$

Hence we conclude

$$\int_{\Omega} Ghdv_g = 0, \quad \forall h \in E_{\ell}.$$
(66)

If $x_0 \in \partial \Omega$, testing the equation (65) by $\phi \in C^2(\overline{\Omega}) \cap W_0^{1,2}(\Omega)$, we have

$$-\int_{\Omega} G\Delta\phi dx + \alpha \int_{\Omega} G\phi dx = 0,$$

since $\phi = 0$ on $\partial\Omega$ (see [8], page 288). By the Sobolev embedding theorem, $G \in L^2(\Omega)$. By Lemma 6, we have $G \in W^{2,2}(\Omega) \cap W_0^{1,2}(\Omega)$. Hence G is an usual weak solution to the equation

$$\left\{ \begin{array}{ll} -\Delta G - \alpha G = 0 \quad \text{in} \quad \Omega, \\ G \in W_0^{1,2}(\Omega), \end{array} \right.$$

and thus $G \equiv 0$ in Ω , since $G \in E_{\ell}^{\perp}$ and $0 \le \alpha < \lambda_{\ell+1}(\Omega)$. Fourthly, we estimate the supremum (13) under the assumption that $c_{\epsilon} \to +\infty$ as $\epsilon \to 0$. If x_0 lies on the boundary $\partial \Omega$, we set

$$u_{\epsilon}^{*}(x) = \begin{cases} u_{\epsilon}(x), & x \in \Omega, \\ 0, & x \in \mathbb{R}^{2} \setminus \Omega. \end{cases}$$

Denote

$$s_{\epsilon} = \sup_{\partial B_{\delta}(x_0)} u_{\epsilon}^*, \quad \overline{u}_{\epsilon}^* = (u_{\epsilon}^* - s_{\epsilon})^+, \quad \tau_{\epsilon} = \int_{B_{\delta}(x_0)} |\nabla \overline{u}_{\epsilon}^*|^2 dx.$$

Since $c_{\epsilon}u_{\epsilon} \to 0$ in $C^1_{\text{loc}}(\overline{\Omega} \setminus \{x_0\}) \cap L^2(\Omega)$, we have

$$s_{\epsilon} = o_{\epsilon}(1)c_{\epsilon}^{-1} \tag{67}$$

and

$$\tau_{\epsilon} \leq \int_{B_{\delta}(x_{0})\cap\Omega} |\nabla u_{\epsilon}|^{2} dx$$

$$= 1 - \int_{\Omega \setminus B_{\delta}(x_{0})} |\nabla u_{\epsilon}|^{2} dx + \alpha \int_{\Omega} u_{\epsilon}^{2} dx$$

$$= 1 + \frac{o_{\epsilon}(1)}{c_{\epsilon}^{2}}.$$
 (68)

It follows from Lemma 5 that

$$\limsup_{\epsilon \to 0} \int_{B_{\delta}(x_0)} (e^{4\pi \overline{u}_{\epsilon}^{*2}/\tau_{\epsilon}} - 1) dx \le \pi \delta^2 e.$$
(69)

In view of (67) and (68), there holds on $B_{Rr_{\epsilon}}(x_{\epsilon})$,

$$\begin{aligned} (4\pi - \epsilon)u_{\epsilon}^{2} &\leq 4\pi u_{\epsilon}^{*2} \\ &\leq 4\pi (\overline{u}_{\epsilon}^{*} + s_{\epsilon})^{2} \\ &= 4\pi \overline{u}_{\epsilon}^{*2} + 8\pi s_{\epsilon} \overline{u}_{\epsilon}^{*} + 4\pi s_{\epsilon}^{2} \\ &= 4\pi \overline{u}_{\epsilon}^{*2} / \tau_{\epsilon} + o_{\epsilon}(1). \end{aligned}$$

This together with (69) leads to

$$\begin{split} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi-\epsilon)u_{\epsilon}^{2}} dx &\leq \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} (e^{4\pi \overline{u_{\epsilon}}^{2}} - 1) dx + o_{\epsilon}(1) \\ &\leq \int_{B_{\delta}(x_{0})} (e^{4\pi \overline{u_{\epsilon}}^{2}} - 1) dx + o_{\epsilon}(1) \\ &\leq \pi \delta^{2} e + o_{\epsilon}(1). \end{split}$$
(70)

By an analogue of (42), it follows from (70) that

$$\sup_{u\in E_{\ell}^{\perp}, ||u||_{1,\alpha}\leq 1}\int_{\Omega}e^{4\pi u^{2}}dx\leq |\Omega|+\pi\delta^{2}e.$$

Since $\delta > 0$ is arbitrary, we get

$$\sup_{u\in E_{\ell}^{\perp}, ||u||_{1,\alpha}\leq 1}\int_{\Omega}e^{4\pi u^{2}}dx\leq |\Omega|,$$

which is impossible. This excludes the possibility of $x_0 \in \partial \Omega$.

Now since $x_0 \in \Omega$, the Green function *G* given by (65) can be represented by

$$G(x) = -\frac{1}{2\pi} \log|x - x_0| + A_{x_0} + \psi_{\alpha}(x), \tag{71}$$

where A_{x_0} is a constant depending only on x_0 and $\alpha, \psi_{\alpha} \in C^1(\overline{\Omega})$ and $\psi_{\alpha}(x_0) = 0$. Repeating the argument of deriving (43), we get

$$\sup_{u \in E_{\ell}^{\perp}, \|u\|_{1,\alpha} \le 1} \int_{\Omega} e^{4\pi u^2} dx \le |\Omega| + \pi e^{1 + 4\pi A_{x_0}}.$$
(72)

Finally we prove the existence of extremal function. It suffices to construct a sequence of functions $\phi_{\epsilon}^* \in E_{\ell}^{\perp}$ with $\|\phi_{\epsilon}^*\|_{1,\alpha} = 1$ such that for sufficiently small $\epsilon > 0$,

$$\int_{\Omega} e^{4\pi\phi_{\epsilon}^{*2}} dx > |\Omega| + \pi e^{1+4\pi A_{x_0}}.$$
(73)

We shall adapt the test functions constructed in Step 5 of the proof of Theorem 1. Let ϕ_{ϵ} be defined by (45), *G* be as in (65), $R = -\log \epsilon$, c^2 be as in (47), and *B* be as in (48). In particular ϕ_{ϵ} satisfies the following three properties: (*i*) $\phi_{\epsilon} \in W_0^{1,2}(\Omega)$; (*ii*) $\|\phi_{\epsilon}\|_{1,\alpha} = 1$; (*iii*) there holds

$$\int_{\Omega} e^{4\pi\phi_{\epsilon}^2} dx \ge |\Omega| + \pi e^{1+4\pi A_{x_0}} + 4\pi \frac{\|G\|_2^2}{c^2} + o(\frac{1}{c^2}).$$

Recalling that (e_{ij}) is a basis of E_{ℓ} verifying (51), we set

•

$$\widetilde{\phi}_{\epsilon} = \phi_{\epsilon} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_i} (\phi_{\epsilon}, e_{ij}) e_{ij},$$

where

$$(\phi_{\epsilon}, e_{ij}) = \int_{\Omega} \phi_{\epsilon} e_{ij} dx.$$

Obviously $\widetilde{\phi}_{\epsilon} \in E_{\ell}^{\perp}$. Noting that $e_{ij} \in C^1(\overline{\Omega})$, $R = -\log \epsilon$, $c^2 = O(-\log \epsilon)$, B = O(1), and G can be represented by (71), we have

$$(\phi_{\epsilon}, e_{ij}) = \int_{B_{R\epsilon}(x_0)} \left(c + \frac{-\frac{1}{4\pi} \log(1 + \pi \frac{r^2}{\epsilon^2}) + B}{c} \right) e_{ij} dx$$
$$+ \int_{B_{2R\epsilon}(x_0) \setminus B_{R\epsilon}(x_0)} \frac{G - \eta \psi_{\alpha}}{c} e_{ij} dx + \int_{\Omega \setminus B_{R\epsilon}(x_0)} \frac{G}{c} e_{ij} dx$$
$$= o(\frac{1}{\log^2 \epsilon}).$$
(74)

Here we have used (66) to derive

$$\int_{\Omega\setminus B_{R\epsilon}(x_0)} \frac{G}{c} e_{ij} dx = -\int_{B_{R\epsilon}(x_0)} \frac{G}{c} e_{ij} dx = O(\epsilon^2 (-\log\epsilon)^{5/2}) = o(\frac{1}{\log^2\epsilon})$$

By (74) and property (*ii*) of ϕ_{ϵ} , we have

$$\widetilde{\phi}_{\epsilon} = \phi_{\epsilon} + o(\frac{1}{\log^2 \epsilon}),\tag{75}$$

$$\|\widetilde{\phi}_{\epsilon}\|_{1,\alpha}^2 = 1 + o(\frac{1}{\log^2 \epsilon}).$$
(76)

Combining (75), (76) and property (*iii*) of ϕ_{ϵ} , we obtain

$$\begin{split} \int_{\Omega} e^{4\pi \frac{\tilde{\phi}_{\epsilon}^{2}}{\|\phi_{\epsilon}\|_{1,\alpha}^{2}}} dx &= \int_{\Omega} e^{4\pi \phi_{\epsilon}^{2} + o(\frac{1}{\log \epsilon})} dx \\ &\geq (1 + o(\frac{1}{\log \epsilon})) \left(|\Omega| + \pi e^{1 + 4\pi A_{x_{0}}} + 4\pi \frac{\|G\|_{2}^{2}}{c^{2}} + o(\frac{1}{c^{2}}) \right) \\ &\geq |\Omega| + \pi e^{1 + 4\pi A_{x_{0}}} + 4\pi \frac{\|G\|_{2}^{2}}{c^{2}} + o(\frac{1}{c^{2}}). \end{split}$$

Set $\phi_{\epsilon}^* = \widetilde{\phi}_{\epsilon}/\|\widetilde{\phi}_{\epsilon}\|_{1,\alpha}$. Since $\widetilde{\phi}_{\epsilon} \in E_{\ell}^{\perp}$, we have $\phi_{\epsilon}^* \in E_{\ell}^{\perp}$. Moreover $\|\phi_{\epsilon}^*\|_{1,\alpha} = 1$ and (73) holds. The contradiction between (72) and (73) implies that c_{ϵ} must be bounded, and whence the existence of extremal function follows from (57) again. The proof of Theorem 2 is completely finished. \Box

4. The Riemannian surface case

In this section we shall combine Carleson-Chang's result (Lemma 5) and blow-up analysis to prove Theorems 3 and 4. We follow the lines of [21, 41, 23]. Throughout this section, we denote a geodesic ball centered at $q \in \Sigma$ with radius r by $B_r(q)$, while a Euclidean ball centered at $x \in \mathbb{R}^2$ with radius r is denoted by $\mathbb{B}_r(x)$.

Proof of Theorem 3. Let α , $0 \le \alpha < \lambda_1(\Sigma)$, be fixed. We divide the proof into several steps.

Step 1. Existence of maximizers for subcritical functionals

In this step, we shall prove for any $0 < \epsilon < 4\pi$, there exists some $u_{\epsilon} \in C^{1}(\Sigma)$ such that

$$||u_{\epsilon}||_{1,\alpha} = 1, \ \int_{\Sigma} u_{\epsilon} dv_g = 0,$$
 (77)

and that

$$\int_{\Sigma} e^{(4\pi-\epsilon)u_{\epsilon}^2} dv_g = \sup_{u \in W^{1,2}(\Sigma), \|u\|_{1,\alpha} \le 1, \int_{\Sigma} u dv_g = 0} \int_{\Sigma} e^{(4\pi-\epsilon)u^2} dv_g,$$
(78)

where $\|\cdot\|_{1,\alpha}$ is defined as in (14).

To do this, we choose a maximizing sequence u_j such that $||u_j||_{1,\alpha} \le 1$, $\int_{\Sigma} u_j dv_g = 0$ and

$$\int_{\Sigma} e^{(4\pi-\epsilon)u_j^2} dv_g \to \sup_{u \in W^{1,2}(\Sigma), \|u\|_{1,a} \le 1, \int_{\Sigma} u dv_g = 0} \int_{\Sigma} e^{(4\pi-\epsilon)u^2} dv_g.$$
(79)

It follows from $0 \le \alpha < \lambda_1(\Sigma)$ that u_j is bounded in $W^{1,2}(\Sigma)$. Then we can assume, up to a subsequence, $u_j \rightharpoonup u_{\epsilon}$ weakly in $W^{1,2}(\Sigma)$, $u_j \rightarrow u_{\epsilon}$ strongly in $L^2(\Sigma)$, and $u_j \rightarrow u_{\epsilon}$ a.e. in Σ . Similarly as in Step 1 of the proof of Theorem 1, we have $||u_{\epsilon}||_{1,\alpha} \le 1$ and

$$\int_{\Sigma} |\nabla_g u_j - \nabla_g u_{\epsilon}|^2 dv_g \le 1 - ||u_{\epsilon}||_{1,\alpha}^2 + o_j(1).$$

It follows from a manifold version of Lions' inequality ([41], Lemma 3.1) that $e^{(4\pi-\epsilon)u_j^2}$ is bounded in $L^q(\Sigma)$ for some q > 1. Hence $e^{(4\pi-\epsilon)u_j^2} \rightarrow e^{(4\pi-\epsilon)u_e^2}$ strongly in $L^1(\Sigma)$. This together with (79) leads to (78). Note that $\int_{\Sigma} u_e dv_g = 0$, since $\int_{\Sigma} u_j dv_g = 0$. We only need to confirm that $||u_e||_{1,\alpha} = 1$. Suppose not, we have $||u_e||_{1,\alpha} < 1$. Set $u^* = u_e/||u_e||_{1,\alpha}$. Then u^* satisfies (77) and

$$\int_{\Sigma} e^{(4\pi-\epsilon)u^{*2}} dv_g > \int_{\Sigma} e^{(4\pi-\epsilon)u_{\epsilon}^{2}} dv_g,$$

which contradicts (78). Therefore $||u_{\epsilon}||_{1,\alpha} = 1$.

It is not difficult to check that u_{ϵ} satisfies the Euler-Lagrange equation

$$\begin{cases} \Delta_g u_{\epsilon} - \alpha u_{\epsilon} = \frac{1}{\lambda_{\epsilon}} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^2} - \frac{\mu_{\epsilon}}{\lambda_{\epsilon}} \\ \lambda_{\epsilon} = \int_{\Sigma} u_{\epsilon}^2 e^{(4\pi - \epsilon)u_{\epsilon}^2} dv_g \\ \mu_{\epsilon} = \frac{1}{\operatorname{Vol}_g(\Sigma)} \int_{\Sigma} u_{\epsilon} e^{(4\pi - \epsilon)u_{\epsilon}^2} dv_g, \end{cases}$$

$$(80)$$

where Δ_g is the Laplace-Beltrami operator. Applying elliptic estimates to (80), we have that $u_{\epsilon} \in C^1(\Sigma)$.

Step 2. Blow-up analysis

Noting that

$$\int_{\Sigma} e^{(4\pi-\epsilon)u_{\epsilon}^2} dV_g \le \int_{\Sigma} \left(1 + (4\pi-\epsilon)u_{\epsilon}^2 e^{(4\pi-\epsilon)u_{\epsilon}^2} \right) dV_g = \operatorname{Vol}_g(\Sigma) + (4\pi-\epsilon)\lambda_{\epsilon}$$

and

$$\lim_{\epsilon \to 0} \int_{\Sigma} e^{(4\pi - \epsilon)u_{\epsilon}^2} dV_g = \sup_{u \in W^{1,2}(\Sigma), \|u\|_{1,\alpha} \le 1, \int_{\Sigma} u dv_g = 0} \int_{\Sigma} e^{(4\pi - \epsilon)u^2} dv_g$$

we have $\liminf_{\epsilon \to 0} \lambda_{\epsilon} > 0$. It then follows that $\mu_{\epsilon}/\lambda_{\epsilon}$ is a bounded sequence. Denote $c_{\epsilon} = |u_{\epsilon}(x_{\epsilon})| = \max_{\Sigma} |u_{\epsilon}|$. If c_{ϵ} is bounded, applying elliptic estimates to (80), we already conclude the existence of extremal function. Without loss of generality, we assume $x_{\epsilon} \to p \in \Sigma$ and $c_{\epsilon} = u_{\epsilon}(x_{\epsilon}) \to +\infty$ as $\epsilon \to 0$. Take an isothermal coordinate system (U, ϕ) near p such that the metric g can be represented by $g = e^{f}(dx_{1}^{2} + dx_{2}^{2})$, where $f \in C^{1}(\Omega, \mathbb{R}), \Omega = \phi(U) \subset \mathbb{R}^{2}$, and f(0) = 0. Denote $\widetilde{u_{\epsilon}} = u_{\epsilon} \circ \phi^{-1}, \widetilde{x} = \phi^{-1}(x)$ for $x \in \Omega$.

Let

$$\begin{aligned} r_{\epsilon} &= \sqrt{\lambda_{\epsilon}} c_{\epsilon}^{-1} e^{-(2\pi - \epsilon/2)c_{\epsilon}^{2}}, \\ \psi_{\epsilon}(x) &= c_{\epsilon}^{-1} \widetilde{u}_{\epsilon}(\widetilde{x_{\epsilon}} + r_{\epsilon}x), \end{aligned}$$

and

$$\varphi_{\epsilon}(x) = c_{\epsilon}(\widetilde{u}_{\epsilon}(\widetilde{x}_{\epsilon} + r_{\epsilon}x) - c_{\epsilon})$$

for $x \in \Omega_{\epsilon} = \{x \in \mathbb{R}^2 : \tilde{x}_{\epsilon} + r_{\epsilon}x \in \Omega\}$. By (80), we have

$$-\Delta_{\mathbb{R}^2}\psi_{\epsilon} = e^{f(\widetilde{x}_{\epsilon}+r_{\epsilon}x)} \left(\alpha r_{\epsilon}^2\psi_{\epsilon} + c_{\epsilon}^{-2}\psi_{\epsilon}e^{(4\pi-\epsilon)(\widetilde{u}_{\epsilon}^2-c_{\epsilon}^2)}\right),\tag{81}$$

$$-\Delta_{\mathbb{R}^2}\varphi_{\epsilon} = e^{f(\widetilde{x}_{\epsilon}+r_{\epsilon}x)} \left(\alpha r_{\epsilon}^2 c_{\epsilon}^2 \psi_{\epsilon} + \psi_{\epsilon} e^{(4\pi-\epsilon)(1+\psi_{\epsilon})\varphi_{\epsilon}}\right),\tag{82}$$

where $-\Delta_{\mathbb{R}^2}$ denotes the usual Laplacian operator. It is easy to see that $\Delta_{\mathbb{R}^2}\psi_{\epsilon} \to 0$ in $L^{\infty}_{\text{loc}}(\mathbb{R}^2)$, $|\psi_{\epsilon}| \leq 1$ and $\psi_{\epsilon}(0) = 1$. Applying elliptic estimates to (81) and using the Liouville theorem for harmonic function, we have

$$\psi_{\epsilon} \to 1$$
 in $C^1_{\text{loc}}(\mathbb{R}^2)$.

Since $\Delta_{\mathbb{R}^2} \varphi_{\epsilon}$ is bounded in $L^{\infty}_{\text{loc}}(\mathbb{R}^2)$ and $\varphi_{\epsilon}(x) \leq 0 = \varphi_{\epsilon}(0)$ for all $x \in \Omega_{\epsilon}$, we have by applying elliptic estimates to (82),

$$\varphi_{\epsilon} \to \varphi = -\frac{1}{4\pi} \log(1 + \pi |x|^2)$$
 in $C_{\text{loc}}^1(\mathbb{R}^2)$.
21

Moreover we have

$$\int_{\mathbb{R}^2} e^{8\pi\varphi} dx = 1.$$
(83)

Repeating the argument of proving ([41], Lemma 4.9), we obtain $c_{\epsilon}u_{\epsilon} \rightarrow G$ weakly in $W^{1,q}(\Sigma)$ for all 1 < q < 2, and $c_{\epsilon}u_{\epsilon} \rightarrow G$ in $C^{1}_{loc}(\Sigma \setminus \{p\}) \cap L^{2}(\Sigma)$, where G is a Green function defined by

$$\begin{cases} \Delta_g G - \alpha G = \delta_p - \frac{1}{\operatorname{Vol}_g(\Sigma)} & \text{in} \quad \Sigma \\ \int_{\Sigma} G dv_g = 0. \end{cases}$$
(84)

Clearly G can be represented by

$$G = -\frac{1}{2\pi}\log r + A_p + \psi, \tag{85}$$

where *r* denotes the geodesic distance from *p*, A_p is a constant real number, $\psi \in C^1(\Sigma)$ with $\psi(p) = 0$.

Step 3. Upper bound estimate

Similarly as we did in Step 4 of the proof of Theorem 1, we obtain by using Carleson-Chang's result (Lemma 5)

$$\lim_{R \to +\infty} \limsup_{\epsilon \to 0} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dv_{g} \le \pi e^{1 + 4\pi A_{p}},$$
(86)

where A_p is given by (85). Note that

$$\begin{split} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi-\epsilon)u_{\epsilon}} dv_{g} &= (1+o_{\epsilon}(1)) \int_{\mathbb{B}_{Rr_{\epsilon}}(\widetilde{x}_{\epsilon})} e^{(4\pi-\epsilon)\widetilde{u}_{\epsilon}} dx \\ &= (1+o_{\epsilon}(1)) \int_{\mathbb{B}_{R}(0)} e^{(4\pi-\epsilon)\widetilde{u}_{\epsilon}} r_{\epsilon}^{2} dx \\ &= (1+o_{\epsilon}(1)) \frac{\lambda_{\epsilon}}{c_{\epsilon}^{2}} \int_{\mathbb{B}_{R}(0)} e^{8\pi\varphi} dx. \end{split}$$

This together with (83) implies

$$\lim_{R \to +\infty} \limsup_{\epsilon \to 0} \int_{B_{Rr_{\epsilon}}(x_{\epsilon})} e^{(4\pi - \epsilon)u_{\epsilon}^{2}} dv_{g} = \limsup_{\epsilon \to 0} \frac{\lambda_{\epsilon}}{c_{\epsilon}^{2}},$$

which together with (86) and an analogue of ([41], Lemma 4.6) leads to

$$\sup_{u \in W^{1,2}(\Sigma), \|u\|_{1,\alpha} \le 1, \int_{\Sigma} u dv_g = 0} \int_{\Sigma} e^{4\pi u^2} dv_g = \limsup_{\epsilon \to 0} \int_{\Sigma} e^{(4\pi - \epsilon)u_\epsilon^2} dv_g \le \operatorname{Vol}_g(\Sigma) + \pi e^{1 + 4\pi A_p}.$$
(87)

Step 4. Existence of extremal function

In this step we will construct a blow-up sequence ϕ_{ϵ} such that

$$\int_{\Sigma} |\nabla_g \phi_{\epsilon}|^2 dv_g - \alpha \int_{\Sigma} (\phi - \overline{\phi}_{\epsilon})^2 dv_g = 1$$
(88)

and

$$\int_{\Sigma} e^{4\pi(\phi_{\epsilon} - \overline{\phi}_{\epsilon})^2} dv_g > \operatorname{Vol}(\Sigma) + \pi e^{1 + 4\pi A_p}$$
(89)

for sufficiently small $\epsilon > 0$, where

$$\overline{\phi}_{\epsilon} = \frac{1}{\operatorname{Vol}_g(\Sigma)} \int_{\Sigma} \phi_{\epsilon} dv_g.$$

The contradiction between (89) and (87) implies that c_{ϵ} must be bounded and elliptic estimates imply the existence of the desired extremal function. This completes the proof of Theorem 3.

Now we construct ϕ_{ϵ} verifying (88) and (89). Note that the Green function G defined as in (84) has the representation (85). Set

$$\phi_{\epsilon} = \begin{cases} c + \frac{-\frac{1}{4\pi} \log(1 + \pi \frac{r^2}{\epsilon^2}) + B}{c} & \text{for } r \leq R\epsilon \\ \frac{G - \eta \psi}{c} & \text{for } R\epsilon < r < 2R\epsilon \\ \frac{G}{c} & \text{for } r \geq 2R\epsilon, \end{cases}$$
(90)

where $R = -\log \epsilon$, $\eta \in C_0^{\infty}(B_{2R\epsilon}(p))$ verifying that $\eta = 1$ on $B_{R\epsilon}(p)$ and $\|\nabla_g \eta\|_{L^{\infty}} = O(\frac{1}{R\epsilon})$, *B* is a constant to be determined later, and *c* depending only on ϵ will also be chosen later such that $R\epsilon \to 0$ and $R \to +\infty$. In order to assure that $\phi_{\epsilon} \in W^{1,2}(\Sigma)$, we set

$$c + \frac{1}{c} \left(-\frac{1}{4\pi} \log(1 + \pi R^2) + B \right) = \frac{1}{c} \left(-\frac{1}{2\pi} \log(R\epsilon) + A_p \right),$$

which gives

$$2\pi c^2 = -\log \epsilon - 2\pi B + 2\pi A_p + \frac{1}{2}\log \pi + O(\frac{1}{R^2}).$$
(91)

We calculate

$$\begin{split} \int_{\Sigma} |\nabla_g \phi_{\epsilon}|^2 dv_g &= \frac{1}{4\pi c^2} \left(2\log \frac{1}{\epsilon} + \log \pi - 1 + 4\pi A_p + 4\pi \alpha ||G||_2^2 + O(\frac{1}{R^2}) + O(R\epsilon \log(R\epsilon)) \right), \end{split}$$

$$\begin{split} \int_{\Sigma} \phi_{\epsilon} dv_{g} &= \frac{1}{c} \left(\int_{r \geq 2R\epsilon} G dv_{g} + O(R\epsilon \log(R\epsilon)) \right) \\ &= \frac{1}{c} \left(- \int_{r < 2R\epsilon} G dv_{g} + O(R\epsilon \log(R\epsilon)) \right) \\ &= \frac{1}{c} O(R\epsilon \log(R\epsilon)), \end{split}$$

and

$$\begin{split} \int_{\Sigma} (\phi_{\epsilon} - \overline{\phi}_{\epsilon})^2 dv_g &= \int_{\Sigma} \phi_{\epsilon}^2 dv_g - \overline{\phi}_{\epsilon}^2 \mathrm{Vol}_g(\Sigma) \\ &= \frac{1}{c^2} \left(\int_{\Sigma} G^2 dv_g + O(R\epsilon \log(R\epsilon)) \right). \end{split}$$

This yields

$$\begin{split} \|\phi_{\epsilon} - \overline{\phi}_{\epsilon}\|_{1,\alpha}^{2} &= \int_{\Sigma} |\nabla_{g}\phi_{\epsilon}|^{2} dv_{g} - \alpha \int_{\Sigma} (\phi_{\epsilon} - \overline{\phi}_{\epsilon})^{2} dv_{g} \\ &= \frac{1}{4\pi c^{2}} \left(2\log\frac{1}{\epsilon} + \log\pi - 1 + 4\pi A_{p} + O(\frac{1}{R^{2}}) + O(R\epsilon\log(R\epsilon)) \right) \end{split}$$

Let ϕ_{ϵ} satisfy (88), i.e. $\|\phi_{\epsilon} - \overline{\phi}_{\epsilon}\|_{1,\alpha} = 1$. Then we have

$$c^{2} = -\frac{\log \epsilon}{2\pi} + \frac{\log \pi}{4\pi} - \frac{1}{4\pi} + A_{p} + O(\frac{1}{R^{2}}) + O(R\epsilon \log(R\epsilon)).$$
(92)

It follows from (91) and (92) that

$$B = \frac{1}{4\pi} + O(\frac{1}{R^2}) + O(R\epsilon \log(R\epsilon)).$$
(93)

Clearly we have on $B_{R\epsilon}(p)$

$$4\pi(\phi_{\epsilon}-\overline{\phi}_{\epsilon})^{2} \geq 4\pi c^{2} - 2\log(1+\pi\frac{r^{2}}{\epsilon^{2}}) + 8\pi B.$$

This together with (92) and (93) yields

$$\int_{B_{R\epsilon}(p)} e^{4\pi(\phi_{\epsilon} - \overline{\phi}_{\epsilon})^2} dv_g \ge \pi e^{1 + 4\pi A_p} + O(\frac{1}{(\log \epsilon)^2}).$$
(94)

On the other hand,

$$\int_{\Sigma \setminus B_{R\epsilon}(p)} e^{4\pi(\phi_{\epsilon} - \overline{\phi}_{\epsilon})^2} dv_g \geq \int_{\Sigma \setminus B_{2R\epsilon}(p)} (1 + 4\pi \phi_{\epsilon}^2) dv_g$$

$$\geq \operatorname{Vol}_g(\Sigma) + 4\pi \frac{\|G\|_2^2}{c^2} + o(\frac{1}{c^2}). \tag{95}$$

Recalling (92) and combining (94) and (95), we conclude (89) for sufficiently small $\epsilon > 0$. \Box

Proof of Theorem 4. Let $\lambda_1(\Sigma) < \lambda_2(\Sigma) < \cdots$ be all distinct eigenvalues of the Laplace-Beltrami operator Δ_g , and $E_{\lambda_i(\Sigma)}$'s be associated eigenfunction spaces. It is known that $\lambda_i(\Sigma) \rightarrow +\infty$ as $i \rightarrow +\infty$ and each space $E_{\lambda_i(\Sigma)}$ has finite dimension (see [10], Chapter I, Page 8). We can assume

$$\dim E_{\lambda_i(\Sigma)} = n_i, \quad i = 1, 2, \cdots.$$

Take a basis (e_{ij}) $(1 \le j \le n_i, 1 \le i \le \ell)$ of E_ℓ verifying

$$E_{\lambda_i(\Sigma)} = \operatorname{span}\{e_{i1}, \cdots, e_{in_i}\}, \quad i = 1, \cdots, \ell,$$

$$E_{\ell} = \operatorname{span}\{e_{11}, \cdots, e_{1n_1}, e_{21}, \cdots, e_{2,n_2}, \cdots, e_{\ell 1}, \cdots, e_{\ell n_{\ell}}\},$$

$$\int_{\Sigma} |e_{ij}|^2 dv_g = 1,$$

$$\int_{\Sigma} e_{ij} e_{kl} dv_g = 0, \quad i \neq k \text{ or } j \neq l.$$

$$24$$

Similar to Step 1 of the proof of Theorem 3, for any ϵ , $0 < \epsilon < 4\pi$, there exists some $u_{\epsilon} \in E_{\ell}^{\perp} \cap C^{1}(\Sigma)$ with $||u_{\epsilon}||_{1,\alpha} = 1$ such that

$$\int_{\Sigma} e^{(4\pi-\epsilon)u_{\epsilon}^{2}} dv_{g} = \sup_{u \in E_{\ell}^{\perp}, ||u||_{1,\alpha} \le 1} \int_{\Sigma} e^{(4\pi-\epsilon)u^{2}} dv_{g}.$$

Moreover u_{ϵ} satisfies the Euler-Lagrange equation

$$\begin{cases} \Delta_{g}u_{\epsilon} - \alpha u_{\epsilon} = \frac{1}{\lambda_{\epsilon}}u_{\epsilon}e^{(4\pi-\epsilon)u_{\epsilon}^{2}} - \frac{\mu_{\epsilon}}{\lambda_{\epsilon}} - \sum_{i=1}^{\ell}\sum_{j=1}^{n_{i}}\frac{\gamma_{ij,\epsilon}}{\lambda_{\epsilon}}e_{ij} \text{ in } \Sigma, \\ u_{\epsilon} \in E_{\ell}^{\perp} \cap C^{1}(\Sigma), \\ \lambda_{\epsilon} = \int_{\Sigma}u_{\epsilon}^{2}e^{(4\pi-\epsilon)u_{\epsilon}^{2}}dv_{g}, \\ \mu_{\epsilon} = \frac{1}{\operatorname{Vol}_{g}(\Sigma)}\int_{\Sigma}u_{\epsilon}e^{(4\pi-\epsilon)u_{\epsilon}^{2}}dv_{g}, \\ \gamma_{ij,\epsilon} = \int_{\Sigma}e_{ij}u_{\epsilon}e^{(4\pi-\epsilon)u_{\epsilon}^{2}}dv_{g}. \end{cases}$$

Let $c_{\epsilon} = \max_{\Sigma} |u_{\epsilon}|$. Without loss of generality, we assume $c_{\epsilon} = u_{\epsilon}(x_{\epsilon}) \to +\infty$ and $x_{\epsilon} \to p \in \Sigma$ as $\epsilon \to 0$. Take an isothermal coordinate system (U, ϕ) near p. Denote $\widetilde{u}_{\epsilon} = u_{\epsilon} \circ \phi^{-1}$, $\widetilde{x} = \phi^{-1}(x)$ for $x \in \phi(U) \subset \mathbb{R}^2$. Perform the same blow-up analysis as in the proof of Theorem 3. There holds

$$\begin{split} \varphi_{\epsilon} &\to \varphi = -\frac{1}{4\pi} \log(1 + \pi |x|^2) \quad \text{in} \quad C^1_{\text{loc}}(\mathbb{R}^2), \\ c_{\epsilon} u_{\epsilon} &\to G \text{ weakly in } W^{1,q}(\Sigma), \ \forall 1 < q < 2, \\ c_{\epsilon} u_{\epsilon} &\to G \text{ in } C^1_{\text{loc}}(\Sigma \setminus \{p\}) \cap L^2(\Sigma), \end{split}$$

where $\varphi_{\epsilon}(x) = c_{\epsilon}(\widetilde{u}_{\epsilon}(\widetilde{x}_{\epsilon} + r_{\epsilon}x) - c_{\epsilon}), G$ is a Green function defined by

$$\Delta_g G - \alpha G = \delta_p - \frac{1}{\operatorname{Vol}_g(\Sigma)} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_i} e_{ij}(p) e_{ij}.$$
(96)

Since $u_{\epsilon} \in E_{\ell}^{\perp}$, we have

$$\int_{\Sigma} Ge_{ij} dv_g = \lim_{\epsilon \to 0} \int_{\Sigma} c_{\epsilon} u_{\epsilon} e_{ij} dv_g = 0, \quad \forall 1 \le j \le n_i, \ 1 \le i \le \ell.$$

Clearly G can be written as

$$G = -\frac{1}{2\pi} \log r + A_p + \psi, \qquad (97)$$

where *r* denotes the geodesic distance from *p*, A_p is a constant real number, $\psi \in C^1(\Sigma)$ with $\psi(p) = 0$. Using Carleson-Chang's result (Lemma 5), we obtain

$$\sup_{u \in E_{\ell}^{\perp}, \|u\|_{1,\alpha} \le 1, \int_{\Sigma} u dv_g = 0} \int_{\Sigma} e^{4\pi u^2} dv_g = \limsup_{\epsilon \to 0} \int_{\Sigma} e^{(4\pi - \epsilon)u_{\epsilon}^2} dv_g \le \operatorname{Vol}_g(\Sigma) + \pi e^{1 + 4\pi A_p}.$$
(98)

Now we will construct a sequence of functions ϕ_{ϵ}^* such that $\phi_{\epsilon}^* \in E_{\ell}^{\perp}$, $\int_{\Sigma} \phi_{\epsilon}^* dv_g = 0$ and

$$\int_{\Sigma} e^{4\pi \phi_e^{*2}} d\nu_g > \operatorname{Vol}_g(\Sigma) + \pi e^{1+4\pi A_p}$$

$$25$$
(99)

for sufficiently small $\epsilon > 0$. The contradiction between (99) and (98) implies that c_{ϵ} must be bounded and elliptic estimates lead to the existence of the desired extremal function. This completes the proof of Theorem 4.

Let ϕ_{ϵ} be defined by (90), G be as in (96), $R = -\log \epsilon$, c^2 be as in (92), and B be as in (93). In particular ϕ_{ϵ} satisfies

$$\int_{\Sigma} |\nabla_g \phi_\epsilon|^2 dv_g - \alpha \int_{\Sigma} \phi_\epsilon^2 dv_g = 1$$
(100)

and

$$\int_{\Sigma} e^{4\pi(\phi_{\epsilon} - \overline{\phi}_{\epsilon})^2} d\nu_g \ge \operatorname{Vol}_g(\Sigma) + \pi e^{1 + 4\pi A_p} + 4\pi \frac{\|G\|_2^2}{c^2} + o(\frac{1}{c^2}), \tag{101}$$

where

$$\overline{\phi}_{\epsilon} = \frac{1}{\operatorname{Vol}_g(\Sigma)} \int_{\Sigma} \phi_{\epsilon} dv_g.$$

Set

$$\widetilde{\phi}_{\epsilon} = \phi_{\epsilon} - \overline{\phi}_{\epsilon} - \sum_{i=1}^{\ell} \sum_{j=1}^{n_i} (\phi_{\epsilon} - \overline{\phi}_{\epsilon}, e_{ij}) e_{ij},$$

where

$$(\phi_{\epsilon} - \overline{\phi}_{\epsilon}, e_{ij}) = \int_{\Sigma} (\phi_{\epsilon} - \overline{\phi}_{\epsilon}) e_{ij} dv_g.$$

Obviously $\tilde{\phi}_{\epsilon} \in E_{\ell}^{\perp}$. Note that $e_{ij} \in C^{1}(\Sigma)$, $R = -\log \epsilon$, $c^{2} = O(-\log \epsilon)$, B = O(1), and G can be represented by (97). We calculate

$$\begin{aligned} (\phi_{\epsilon} - \overline{\phi}_{\epsilon}, e_{ij}) &= \int_{B_{R\epsilon}(p)} \left(c + \frac{-\frac{1}{4\pi} \log(1 + \pi \frac{r^2}{\epsilon^2}) + B}{c} - \overline{\phi}_{\epsilon} \right) e_{ij} dv_g \\ &+ \int_{B_{2R\epsilon}(p) \setminus B_{R\epsilon}(p)} \left(\frac{G - \eta \psi_{\alpha}}{c} - \overline{\phi}_{\epsilon} \right) e_{ij} dv_g + \int_{\Sigma \setminus B_{R\epsilon}(p)} \left(\frac{G}{c} - \overline{\phi}_{\epsilon} \right) e_{ij} dv_g \\ &= I + II + III. \end{aligned}$$

Since $G \in E_{\ell}^{\perp}$, we have

$$\int_{\Sigma \setminus B_{R\epsilon}(p)} \frac{G}{c} dv_g = -\int_{B_{R\epsilon}(p)} \frac{G}{c} dv_g = \frac{1}{c} O(R^2 \epsilon^2 \log(R\epsilon)).$$

Note also that $\overline{\phi}_{\epsilon} = \frac{1}{c}O(R\epsilon \log(R\epsilon))$. Hence

$$III = \frac{1}{c}O(R\epsilon\log(R\epsilon)).$$

Clearly

$$I = O(cR^2\epsilon^2), \quad II = \frac{1}{c}O(R^2\epsilon^2\log(R\epsilon)).$$

Therefore

$$(\phi_{\epsilon} - \overline{\phi}_{\epsilon}, e_{ij}) = O(R^2 \epsilon^2 \sqrt{-\log \epsilon}) = o(\frac{1}{\log^2 \epsilon}).$$

This together with (100) leads to

$$\widetilde{\phi}_{\epsilon} = \phi_{\epsilon} - \overline{\phi}_{\epsilon} + o(\frac{1}{\log^2 \epsilon}), \tag{102}$$

$$\|\widetilde{\phi}_{\epsilon}\|_{1,\alpha}^2 = 1 + o(\frac{1}{\log^2 \epsilon}).$$
(103)

Combining (102), (103) and (101), we obtain

$$\begin{split} \int_{\Sigma} e^{4\pi \frac{\tilde{\phi}_{\varepsilon}^{2}}{\|\tilde{\phi}_{\varepsilon}\|_{1,\alpha}^{2}}} dv_{g} &= \int_{\Sigma} e^{4\pi (\phi_{\varepsilon} - \overline{\phi}_{\varepsilon})^{2} + o(\frac{1}{\log \varepsilon})} dv_{g} \\ &\geq (1 + o(\frac{1}{\log \varepsilon})) \left(\operatorname{Vol}_{g}(\Sigma) + \pi e^{1 + 4\pi A_{p}} + 4\pi \frac{\|G\|_{2}^{2}}{c^{2}} + o(\frac{1}{c^{2}}) \right) \\ &\geq \operatorname{Vol}_{g}(\Sigma) + \pi e^{1 + 4\pi A_{p}} + 4\pi \frac{\|G\|_{2}^{2}}{c^{2}} + o(\frac{1}{c^{2}}). \end{split}$$

Set $\phi_{\epsilon}^* = \widetilde{\phi}_{\epsilon}/\|\widetilde{\phi}_{\epsilon}\|_{1,\alpha}$. Since $\widetilde{\phi}_{\epsilon} \in E_{\ell}^{\perp}$, we have $\phi_{\epsilon}^* \in E_{\ell}^{\perp}$. Moreover $\|\phi_{\epsilon}^*\|_{1,\alpha} = 1$. Since $\Delta_g e_{ij} = \lambda_i e_{ij}$, we have $\int_{\Sigma} e_{ij} dv_g = 0$, and whence $\int_{\Sigma} \phi_{\epsilon}^* dv_g = 0$. Therefore ϕ_{ϵ}^* is the desired function sequence verifying (99).

Acknowledgements. This work is supported by the National Science Foundation of China (Grant No.11171347 and Grant No. 11471014). The author also gratefully acknowledges the helpful comments and suggestions of the referees, which have improved the presentation.

References

- Adimurthi, O. Druet, Blow-up analysis in dimension 2 and a sharp form of Trudinger-Moser inequality, Comm. Partial Differential Equations 29 (2004) 295-322.
- [2] Adimurthi, K. Sandeep, A singular Moser-Trudinger embedding and its applications, Nonlinear Differ. Equ. Appl. 13 (2007) 585-603.
- [3] Adimurthi, C. Tintarev, On compactness in the Trudinger-Moser inequality, Ann. Sc. Norm. Super. Pisa Cl. Sci. 13 (2014) 399-416.
- [4] Adimurthi, Y. Yang, An interpolation of Hardy inequality and Trudinger-Moser inequality in \mathbb{R}^N and its applications, Internat. Mathematics Research Notices 13 (2010) 2394-2426.
- [5] Adimurthi, M. Struwe, Global compactness properties of semilinear elliptic equation with critical exponential growth, J. Functional Analysis 175 (2000) 125-167.
- [6] S. Agmon, The L_p approach to the Dirichlet problem, Ann. Scuola Norm. Sup. Pisa 13 (1959) 405-448.
- [7] T. Aubin, Sur la function exponentielle, C. R. Acad. Sci. Paris, Series A 270 (1970) 1514-1514.
- [8] H. Brezis, Functional analysis, Sobolev spaces and PDEs, Springer, 2011.
- [9] L. Carleson, A. Chang, On the existence of an extremal function for an inequality of J. Moser, Bull. Sci. Math. 110 (1986) 113-127.
- [10] I. Chavel, Eigenvalues in Riemannian geometry, Academic Press, 1984.
- [11] W. Chen, C. Li, Classification of solutions of some nonlinear elliptic equations, Duke Math. J. 63 (1991) 615-622.
- [12] P. Cherrier, Une inégalité de Sobolev sur les variétés Riemanniennes, Bull. Sc. Math. 103 (1979) 353-374.
- [13] P. Cherrier, Cas d'exception du théorème d'inclusion de Sobolev sur les variétés Riemanniennes et applications, Bull. Sc. Math. 105 (1981) 235-288.
- [14] M. de Souza, J. M. do Ó, A sharp Trudinger-Moser type inequality in ℝ², to appear in Tran. Amer. Math. Soc., 2014.
- [15] W. Ding, J. Jost, J. Li, G. Wang, The differential equation $-\Delta u = 8\pi 8\pi he^u$ on a compact Riemann Surface, Asian J. Math. 1 (1997) 230-248.

- [16] M. Flucher, Extremal functions for Trudinger-Moser inequality in 2 dimensions, Comment. Math. Helv. 67 (1992) 471-497.
- [17] L. Fontana, Sharp borderline Sobolev inequalities on compact Riemannian manifolds, Comm. Math. Helv. 68 (1993) 415-454.
- [18] B. Gidas, W. Ni, L. Nirenberg, Symmetry and related properties via the maximum principle, Comm. Math. Phys. 68 (1979) 209-243.
- [19] K. Lin, Extremal functions for Moser's inequality, Trans. Amer. Math. Soc. 348 (1996) 2663-2671.
- [20] P. L. Lions, The concentration-compactness principle in the calculus of variation, the limit case, part I. Rev. Mat. Iber. 1 (1985) 145-201.
- [21] Y. Li, Moser-Trudinger inequality on compact Riemannian manifolds of dimension two, J. Part. Diff. Equations 14 (2001) 163-192.
- [22] Y. Li, P. Liu, Moser-Trudinger inequality on the boundary of compact Riemannian surface, Math. Z. 250 (2005) 363-386.
- [23] Y. Li, P. Liu, Y. Yang, Moser-Trudinger inequalities on vector bundles over a compact Riemannian manifold of dimension 2, Cal. Var. 28 (2007) 59-83.
- [24] Y. Li, B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in \mathbb{R}^N , Indiana Univ. Math. J. 57 (2008) 451-480.
- [25] P. Liu, A Moser-Trudinger type inequality and blow up analysis on compact Riemannian surface, Doctoral thesis, Max-Plank Institute, Germany, 2005.
- [26] G. Lu, Y. Yang, The sharp constant and extremal functions for Moser-Trudinger inequalities involving L^p norms, Discrete and continuous dynamical systems 25 (2009) 963-979.
- [27] G. Mancini, K. Sandeep, Moser-Trudinger inequality on conformal discs, Commun. Contemp. Math. 12 (2010) 1055-1068.
- [28] G. Mancini, K. Sandeep, Extremals for Sobolev and Moser inequalities in hyperbolic space, Milan J. Math. 79 (2011) 273-283.
- [29] G. Mancini, K. Sandeep, C. Tintarev, Trudinger-Moser inequality in the hyperbolic space \mathbb{H}^N , Adv. Nonlinear Anal. 2 (2013) 309-324.
- [30] J. Moser, A sharp form of an inequality by N.Trudinger, Ind. Univ. Math. J. 20 (1971) 1077-1091.
- [31] B. Osgood, R. Phillips, P. Sarnak, Extremals of determinants of Laplacians, J. Funct. Anal. 80 (1988) 148-211.
- [32] J. Peetre, Espaces d'interpolation et theoreme de Soboleff, Ann. Inst. Fourier (Grenoble) 16 (1966) 279-317.
- [33] S. Pohozaev, The Sobolev embedding in the special case pl = n, Proceedings of the technical scientific conference on advances of scientific reseach 1964-1965, Mathematics sections, 158-170, Moscov. Energet. Inst., Moscow, 1965.
- [34] B. Ruf, A sharp Trudinger-Moser type inequality for unbounded domains in \mathbb{R}^2 , J. Funct. Anal. 219 (2005) 340-367.
- [35] M. Struwe, Critical points of embeddings of $H_0^{1,n}$ into Orlicz spaces, Ann. Inst. H. Poincaré, Analyse Non Linéaire 5 (1988) 425-464.
- [36] C. Tintarev, Trudinger-Moser inequality with remainder terms, J. Funct. Anal. 266 (2014) 55-66.
- [37] N. Trudinger, On embeddings into Orlicz spaces and some applications, J. Math. Mech. 17 (1967) 473-484.
- [38] G. Wang, D. Ye, A Hardy-Moser-Trudinger inequality, Adv. Math. 230 (2012) 294-320.
- [39] Y. Yang, Extremal functions for Moser-Trudinger inequalities on 2-dimensional compact Riemannian manifolds with boundary, Inter. J. Math. 17 (2006) 313-330.
- [40] Y. Yang, A sharp form of Moser-Trudinger inequality in high dimension, J. Funct. Anal. 239 (2006) 100-126.
- [41] Y. Yang, A sharp form of the Moser-Trudinger inequality on a compact Riemannian surface, Trans. Amer. Math. Soc. 359 (2007) 5761-5776.
- [42] Y. Yang, A sharp form of trace Moser-Trudinger inequality on compact Riemannian surface with boundary, Math. Z. 255 (2007) 373-392.
- [43] Y. Yang, Trudinger-Moser inequalities on complete noncompact Riemannian manifolds, J. Funct. Anal. 263 (2012) 1894-1938.
- [44] V.I. Yudovich, Some estimates connected with integral operators and with solutions of elliptic equations, Sov. Math. Docl. 2 (1961) 746-749.