# ON FRACTIONAL SCHRÖDINGER EQUATIONS IN SOBOLEV SPACES

#### YOUNGHUN HONG AND YANNICK SIRE

ABSTRACT. Let  $\sigma \in (0, 1)$  with  $\sigma \neq \frac{1}{2}$ . We investigate the fractional nonlinear Schrödinger equation in  $\mathbb{R}^d$ :

$$
i\partial_t u + (-\Delta)^{\sigma} u + \mu |u|^{p-1} u = 0, \ u(0) = u_0 \in H^s,
$$

where  $(-\Delta)^{\sigma}$  is the Fourier multiplier of symbol  $|\xi|^{2\sigma}$ , and  $\mu = \pm 1$ . This model has been introduced by Laskin in quantum physics [\[Las02\]](#page-18-0). We establish local well-posedness and ill-posedness in Sobolev spaces for power-type nonlinearities.

#### CONTENTS



#### 1. INTRODUCTION

<span id="page-0-0"></span>Let  $\sigma \in (0, 1)$  with  $\sigma \neq \frac{1}{2}$  $\frac{1}{2}$ . We consider the Cauchy problem for the fractional nonlinear Schrödinger equation

$$
i\partial_t u + (-\Delta)^\sigma u + \mu |u|^{p-1} u = 0, \ u(0) = u_0 \in H^s,
$$
 (NLS<sub>\sigma</sub>)

where  $\mu = \pm 1$  depending on the focusing or defocusing case. The operator  $(-\Delta)^{\sigma}$  is the so-called fractional laplacian, a Fourier multiplier of  $|\xi|^{2\sigma}$ . The fractional laplacian

*Date*: January 8, 2015.

is the infinitesimal generator of some Levy processes [\[Ber96\]](#page-17-2). A rather extensive study of the potential theoretic aspects of this operator can be found in [\[Lan72\]](#page-18-1).

The previous equation is a fundamental equation of fractional quantum mechnics, a generalization of the standard quantum mechanics extending the Feynman path integral to Levy processes [\[Las02\]](#page-18-0).

The purpose of the present paper is to develop a general well-posedness and illposedness theory in Sobolev spaces. The one-dimensional case has been treated in [\[CHKL14\]](#page-17-3) for cubic nonlinearities, i.e.  $p = 3$ , and  $\sigma \in (\frac{1}{2}, 1)$ . Here, we consider a higher-dimensional version and other types of nonlinear terms. We also include all  $\sigma \in (0, 1)$  except  $\sigma = \frac{1}{2}$  $\frac{1}{2}$ ; furthermore, contrary to [\[CHKL14\]](#page-17-3) where the use of Bourgain spaces was crucial (since the main goal of their paper was to derive well-posedness theory on the flat torus), we rely only on standard Strichartz estimates and functional inequalities in  $\mathbb{R}^d$ . In the case of Hartree-type nonlinearities, the local well-posedness and blow-up have been investigated in [\[CHHO13\]](#page-17-4).

In the present paper, we will not consider global aspects with large data. For that, we refer the reader to [\[GSWZ13\]](#page-18-2) for a study of the energy-critical equation in the radial case, following the seminal work of Kenig and Merle [\[KM08,](#page-18-3) [KM06\]](#page-18-4). As a consequence, we do not consider blow-up phenomena, an aspect we will treat in a forthcoming work.

We introduce two important exponents for our purposes:

$$
s_c = \frac{d}{2} - \frac{2\sigma}{p - 1}
$$

and

$$
s_g = \frac{1 - \sigma}{2}.
$$

Here,  $s_c$  is the scaling-critical regularity exponent in the following sense: for  $\lambda > 0$ , the transformation

$$
u(t,x)\mapsto \frac{1}{\lambda^{2\sigma/(p-1)}}u\Big(\frac{t}{\lambda^{2\sigma}},\frac{x}{\lambda}\Big),\quad u_0(x)\mapsto \frac{1}{\lambda^{2\sigma/(p-1)}}u_0\Big(\frac{x}{\lambda}\Big)
$$

keeps the equation invariant and one can expect local-wellposedness for  $s \geq s_c$ , since the scaling leaves the  $\dot{H}^{s_c}$  norm invariant. On the other hand,  $s_g$  is the critical regularity in the "pseudo"-Galilean invariance (see the proof of ill-posedness below). Under the flow of the equation  $(NLS_{\sigma})$ , the following quantities are conserved:

$$
M[u] = \int_{\mathbb{R}^d} |u(t,x)|^2 dx
$$
 (mass),  
\n
$$
E[u] = \int_{\mathbb{R}^d} \frac{1}{2} |\nabla|^{\sigma} u(t,x)|^2 + \frac{\mu}{p+1} |u(t,x)|^{p+1} dx.
$$
 (energy).

An important feature of the equation under study is a loss of derivatives for the Strichartz estimates as proved in [\[COX11\]](#page-17-5). Unless additional assumptions are met such as radiality as in [\[GW14\]](#page-18-5), one has a loss of  $d(1 - \sigma)$  derivatives in the dispersion (see [\(2.1\)](#page-4-1)). This happens to be an issue in several arguments.

<span id="page-2-0"></span>**Main results.** The goal of this paper is to show that  $(NLS_{\sigma})$  is locally well-posed in  $H^s$  for  $s \ge \max(s_c, s_g, 0)$ , and it is ill-posed in  $H^s$  for  $s \in (s_c, 0)$ . We start with well-posedness results.

<span id="page-2-2"></span>**Theorem 1.1** (Local well-posedness in subcritical cases). Let

$$
\begin{cases}\ns \geqslant s_g & \text{when } d = 1 \text{ and } 2 \leqslant p < 5, \\
s > s_c & \text{when } d = 1 \text{ and } p \geqslant 5, \\
s > s_c & \text{when } d \geqslant 2 \text{ and } p \geqslant 3.\n\end{cases}
$$

Then,  $(NLS_{\sigma})$  is locally well-posed in  $H^s$ .

<span id="page-2-1"></span>**Theorem 1.2** (Local well-posedness in critical cases). Suppose that

$$
\begin{cases} p > 5 & when d = 1, \\ p > 3 & when d \geq 2. \end{cases}
$$

Then,  $(NLS_{\sigma})$  is locally well-posed in  $H^{s_c}$ .

The proof of Theorem [1.2](#page-2-1) is based on a new method, improving on estimates in  $[CKS<sup>+</sup>08]$  $[CKS<sup>+</sup>08]$ . This improvement, based on controlling the nonlinearity in a suitable space, is necessary due to the loss of derivatives in the Strichartz estimates.

As a by-product, we also prove small data scattering.

<span id="page-2-3"></span>**Theorem 1.3** (Small data scattering). Suppose that

$$
\begin{cases} p > 5 & when d = 1, \\ p > 3 & when d \geqslant 2. \end{cases}
$$

Then, there exists  $\delta > 0$  such that if  $||u_0||_{H^{s_c}} < \delta$ , then  $u(t)$  scatters in  $H^{s_c}$ . Precisely, there exist  $u_{\pm} \in H^{s_c}$  such that

$$
\lim_{t \to \pm \infty} \|u(t) - e^{it(-\Delta)^{\sigma}} u_{\pm}\|_{H^{s_c}} = 0.
$$

*Remark* 1.4. Contrary to the case  $\sigma \neq \frac{1}{2}$  $\frac{1}{2}$ , when  $\sigma = \frac{1}{2}$  $\frac{1}{2}$ , the fractional NLS does not have small data scattering. See [\[KLR13\]](#page-18-6).

Finally, our last theorem is the ill-posedness result. Note that our result is not optimal, since one should expect ill-posedness in  $H^s$  up to  $s_g = \frac{1-\sigma}{2}$  $\frac{-\sigma}{2}$ , which is nonnegative. We hope to come back to this issue in a forthcoming work.

<span id="page-3-1"></span>**Theorem 1.5** (Ill-posedness). Let  $d = 1, 2$  or 3 and  $\sigma \in (\frac{d}{4}, 1)$ . If p is not an odd integer, we further assume that  $p \ge k + 1$ , where k is an integer larger than  $\frac{d}{2}$ . Then, (NLS<sub> $\sigma$ </sub>) is ill-posed in H<sup>s</sup> for  $s \in (s_c, 0)$ .

An interesting feature of the previous ill-posedness result is the fact that, contrary to the standard NLS equation ( $\sigma = 1$ ) there is no exact Galilean invariance. However, one can introduce a new "pseudo-Galilean invariance" which is enough to our purposes. More precisely, for  $v \in \mathbb{R}^d$ , we define the transformation

$$
\mathcal{G}_{v}u(t,x) = e^{-iv\cdot x}e^{it|v|^{2\sigma}}u(t,x - 2t\sigma|v|^{2(\sigma-1)}v).
$$

Note that when  $\sigma = 1$ ,  $\mathcal{G}_v$  is simply a Galilean transformation, and that NLS is invariant under this transformation, that is, if  $u(t)$  solves NLS, so does  $\mathcal{G}_v u(t)$ . However, when  $\sigma \neq 1$ , (NLS<sub> $\sigma$ </sub>) is not exactly symmetric with respect to pseudo-Galilean transformations. This opens the construction of solitons for  $(NLS_{\sigma})$  which happen to be different from the ones constructed in the standard case  $\sigma = 1$ . Indeed, if we search for exact solutions of the type

$$
u(t,x) = e^{it(|v|^{2\sigma} - \omega^{2\sigma})} e^{-iv \cdot x} Q_{\omega}(x - 2t\sigma |v|^{2(\sigma - 1)}v), \qquad (1.1)
$$

then the profile  $Q_{\omega}$  solves the pseudo-differential equation

$$
\mathcal{P}_v Q_\omega + \omega^{2\sigma} Q_\omega - |Q_\omega|^{p-1} Q_\omega = 0,
$$
\n(1.2)

where

$$
\mathcal{P}_v = e^{iv \cdot x} (-\Delta)^\sigma e^{-iv \cdot x} - |v|^{2\sigma} - 2i\sigma |v|^{2\sigma - 2} v \cdot \nabla,
$$
\n(1.3)

i.e.,  $\mathcal{P}_v$  is a Fourier multiplier  $\widehat{\mathcal{P}_v f}(\xi) = p_v(\xi) \widehat{f}(\xi)$ , wiht symbol

$$
p_v(\xi) = |\xi - v|^{2\sigma} - |v|^{2\sigma} + 2\sigma |v|^{2\sigma - 2} v \cdot \xi.
$$
 (1.4)

<span id="page-3-0"></span>We plan to come back to this issue in future works.

## 2. Strichartz Estimates

In this section, we review Strichartz estimates for the linear fractional Schrödinger operators. We say that  $(q, r)$  is *admissible* if

$$
\frac{2}{q} + \frac{d}{r} = \frac{d}{2}, \quad 2 \leqslant q, r \leqslant \infty, \quad (q, r, d) \neq (2, \infty, 2).
$$

We define the Strichartz norm by

$$
\|u\|_{S^s_{q,r}(I)}:=\||\nabla|^{-d(1-\sigma)(\frac{1}{2}-\frac{1}{r})}u\|_{L^q_{t\in I}W_x^{s,r}},
$$

where  $I = [0, T)$ . Let  $\psi : \mathbb{R}^d \to [0, 1]$  be a compactly supported smooth function such that  $\sum_{N\in\mathbb{Z}}\psi_N = 1$ , where  $\psi_N(\xi) = \psi(\frac{\xi}{N})$  $\frac{\xi}{N}$ ). For dyadic  $N \in 2^{\mathbb{Z}}$ , let  $P_N$  be a Littlewood-Paley projection, that is,  $\widehat{P_N f}(\xi) = \psi(\frac{\xi}{N})$  $(\frac{\xi}{N})\hat{f}(\xi)$ . Then, we define a slightly stronger Strichartz norm by

$$
||u||_{\tilde{S}_{q,r}^s(I)} := \bigg( \sum_{N \in 2^{\mathbb{Z}}} ||P_N(|\nabla|^{-d(1-\sigma)(\frac{1}{2}-\frac{1}{r})} u) ||_{L_{t \in I}^q W_x^{s,r}}^2 \bigg)^{1/2}.
$$

<span id="page-4-2"></span>**Proposition 2.1** (Strichartz estimates [\[COX11\]](#page-17-5)). For an admissible pair  $(q, r)$ , we have

$$
||e^{it(-\Delta)^{\sigma}}u_0||_{S^s_{q,r}(I)}, ||e^{it(-\Delta)^{\sigma}}u_0||_{\tilde{S}^s_{q,r}(I)} \lesssim ||u_0||_{H^s},
$$
  

$$
\left\|\int_0^t e^{i(t-s)(-\Delta)^{\sigma}} F(s)ds\right\|_{S^s_{q,r}(I)} \lesssim ||F||_{L^1_{t\in I}H^s_x},
$$
  

$$
\left\|\int_0^t e^{i(t-s)(-\Delta)^{\sigma}} F(s)ds\right\|_{\tilde{S}^s_{q,r}(I)} \lesssim ||F||_{L^1_{t\in I}H^s_x}.
$$

Sketch of Proof. By the standard stationary phase estimate, one can show that

$$
\|e^{it(-\Delta)^\sigma} P_1\|_{L^1 \to L^\infty} \lesssim |t|^{-\frac{d}{2}},
$$

and by scaling,

<span id="page-4-1"></span>
$$
||e^{it(-\Delta)^{\sigma}} P_N||_{L^1 \to L^\infty} \lesssim N^{d(1-\sigma)} |t|^{-\frac{d}{2}}.
$$
\n(2.1)

Then, it follows from the argument of Keel-Tao [\[KT98\]](#page-18-7) that for any  $I \subset \mathbb{R}$ ,

$$
\|e^{it(-\Delta)^\sigma} P_N(|\nabla|^{-d(1-\sigma)(\frac{1}{2}-\frac{1}{r})} u_0)\|_{L^q_{t\in I}W^{s,r}_x} \lesssim \|P_N u_0\|_{H^s},
$$
  

$$
\Big\| \int_0^t e^{i(t-s)(-\Delta)^\sigma} P_N(|\nabla|^{-d(1-\sigma)(\frac{1}{2}-\frac{1}{r})} F)(s) ds \Big\|_{L^q_{t\in I}W^{s,r}_x} \lesssim \|P_N F\|_{L^1_{t\in I}H^s_x}.
$$

Squaring the above inequalities and summing them over all dyadic numbers in  $2^{\mathbb{Z}}$ , we prove Strichartz estimates.

The loss of derivatives is due to the Knapp phenomenon (see [\[GW14\]](#page-18-5)). However, in the radial case, one can overcome this loss as proved in [\[GW14\]](#page-18-5), restricting then the admissible powers of the fractional laplacian. Indeed, in [\[GW14\]](#page-18-5), this is proved that one has optimal Strichartz estimates if  $\sigma \in (d/(2d-1), 1)$ . In particular, the number  $d/(2d - 1)$  is larger than  $1/2$  and there is a gap between the Strichartz estimates for the wave operator  $\sigma = 1/2$  and the one occuring for higher powers. This issue suggests that a new phenomenon might occur for this range of powers.

## 3. Local Well-posedness

<span id="page-4-0"></span>We establish local well-posedness of the fractional NLS by the standard contraction mapping argument based on Strichartz estimates. Due to loss of regularity in Strichartz estimates, our proof relies on the  $L_x^{\infty}$  bounds (see Lemma 3.2 and 3.3).

<span id="page-5-0"></span>3.1. Subcritical cases. First, we consider the case that  $d = 1$  and  $2 \leq p < 5$ . In this case, the equation is scaling-subcritical in  $H^s$  for  $s > s_g$ , since  $s_g > s_c$ . We remark that in the proof, we control the  $L_{t \in I}^4 L_x^{\infty}$  norm simply by Strichartz estimates (see  $(3.1)$  and  $(3.2)$ ).

*Proof of Theorem [1.1](#page-2-2) when*  $d = 1$  and  $2 \leq p < 5$ . We define

$$
\Phi_{u_0}(u) := e^{it(-\Delta)^{\sigma}} u_0 + i\mu \int_0^t e^{i(t-s)(-\Delta)^{\sigma}} (|u|^{p-1}u)(s) ds.
$$

Let

<span id="page-5-2"></span><span id="page-5-1"></span>
$$
||u||_{X^s} := ||u||_{L^{\infty}_{t \in I} H^s_x \cap L^4_{t \in I} L^{\infty}_x},
$$

where  $I = [0, T)$ . Then, applying the 1d Strichartz estimates

$$
\|e^{it(-\Delta)^{\sigma}}u_{0}\|_{L_{t\in I}^{4}L_{x}^{\infty}} \lesssim \|u_{0}\|_{\dot{H}^{s_{g}}},
$$
\n
$$
\|e^{it(-\Delta)^{\sigma}}u_{0}\|_{L_{t\in I}^{\infty}H_{x}^{s}} \lesssim \|u_{0}\|_{H^{s}},
$$
\n
$$
\left\|\int_{0}^{t}e^{i(t-s)(-\Delta)^{\sigma}}F(s)ds\right\|_{L_{t\in I}^{4}L_{x}^{\infty}} \lesssim \|F\|_{L_{t\in I}^{1}H_{x}^{s_{g}}},
$$
\n
$$
\left\|\int_{0}^{t}e^{i(t-s)(-\Delta)^{\sigma}}F(s)ds\right\|_{L_{t\in I}^{\infty}H_{x}^{s}} \lesssim \|F\|_{L_{t\in I}^{1}H_{x}^{s}},
$$
\n
$$
(3.2)
$$

we get

$$
\|\Phi_{u_0}(u)\|_{X^s}\lesssim \|u_0\|_{H^s}+\||u|^{p-1}u\|_{L^1_{t\in I}H^s_x}.
$$

By the fractional chain rule

<span id="page-5-3"></span>
$$
\|\nabla|^s F(u)\|_{L^q} \lesssim \|F'(u)\|_{L^{p_1}} \|\nabla|^s u\|_{L^{p_2}},\tag{3.3}
$$

where  $s > 0$  and  $\frac{1}{q} = \frac{1}{p_1}$  $\frac{1}{p_1} + \frac{1}{p_2}$  $\frac{1}{p_2}$ , and Hölder inequality, we obtain

$$
\||u|^{p-1}u\|_{L^1_{t\in I}H^s_x}\lesssim \left\|\||u|^{p-1}\|_{L^\infty_x}\|u\|_{H^s_x}\right\|_{L^1_{t\in I}}\leqslant T^{\frac{5-p}{4}}\|u\|_{L^4_{t\in I}L^\infty_x}^{p-1}\|u\|_{L^\infty_{t\in I}H^s_x}.
$$

For the fractional chain rule [\(3.3\)](#page-5-3), we refer [\[CW91\]](#page-17-7), for example. We remark that one can choose  $p_1 = \infty$  in [\(3.3\)](#page-5-3). Indeed, this can be proved by a little modification of the last step in the proof of Proposition 3.1 in [\[CW91\]](#page-17-7). Thus, we have

$$
\|\Phi_{u_0}(u)\|_{X^s}\lesssim \|u_0\|_{H^s}+T^{\frac{5-p}{4}}\|u\|_{X^s}^p.
$$

Similarly, by Strichartz estimates,

$$
\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\|_{X^s} \lesssim \| |u|^{p-1}u - |v|^{p-1}v\|_{L^1_{t\in I}H^s_x}.
$$

Then, applying the fractional Leibniz rule and the fractional chain rule in [\[CW91\]](#page-17-7), we get

$$
|||u|^{p-1}u - |v|^{p-1}v||_{H_x^s} = \Big\|\int_0^1 p|v + t(u - v)|^{p-1}(u - v)dt\Big\|_{H_x^s}
$$
  
\n
$$
\leqslant p \int_0^1 ||v + t(u - v)|^{p-1}(u - v)||_{H_x^s}^{p-1}dt
$$
  
\n
$$
\leqslant \int_0^1 ||v + t(u - v)||_{L_x^{\infty}}^{p-1} ||u - v||_{H_x^s}
$$
  
\n
$$
+ |||v + t(u - v)|^{p-1} ||_{H_x^s} ||u - v||_{L_x^{\infty}} dt
$$
  
\n
$$
\leqslant \int_0^1 ||v + t(u - v)||_{L_x^{\infty}}^{p-1} ||u - v||_{H_x^s}
$$
  
\n
$$
+ ||v + t(u - v)||_{L_x^{\infty}}^{p-2} ||v + t(u - v)||_{H_x^s}^{p-1} ||u - v||_{L_x^{\infty}} dt
$$
  
\n
$$
\leqslant (||u||_{L_x^{\infty}}^{p-1} + ||v||_{L_x^{\infty}}^{p-1}) ||u - v||_{H_x^s}
$$
  
\n
$$
+ (||u||_{L_x^{\infty}}^{p-2} + ||v||_{L_x^{\infty}}^{p-2}) (||u||_{H_x^s} + ||v||_{H_x^s}) ||u - v||_{L_x^{\infty}}.
$$

Thus, it follows that

$$
\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\|_{X^s} \n\lesssim T^{\frac{5-p}{4}} \Big\{ (\|u\|_{L^4_{t\in I}L^\infty_x}^{p-1} + \|v\|_{L^4_{t\in I}L^\infty_x}^{p-1}) \|u - v\|_{L^\infty_{t\in I}H^s_x} \n+ (\|u\|_{L^4_{t\in I}L^\infty_x}^{p-2} + \|v\|_{L^4_{t\in I}L^\infty_x}^{p-2}) (\|u\|_{L^\infty_{t\in I}H^s_x} + \|v\|_{L^\infty_{t\in I}H^s_x}) \|u - v\|_{L^4_{t\in I}L^\infty_x} \Big\} \n\lesssim T^{\frac{5-p}{4}} (\|u\|_{X^s}^{p-1} + \|v\|_{X^s}^{p-1}) \|u - v\|_{X^s}.
$$

Choosing sufficiently small  $T > 0$ , we conclude that  $\Phi_{u_0}$  is a contraction on a ball

$$
B := \{ u : \|u\|_{X^s} \leqslant 2\|u_0\|_{H^s} \}
$$

equipped with the norm  $\|\cdot\|_{X^s}$ .

Next, we will prove Theorem [1.1](#page-2-2) when  $d = 1$  and  $p \ge 5$ , or  $d \ge 2$  and  $p \ge 3$ . In this case, we do not have a good control on the  $L_x^{\infty}$  norm from Strichartz estimates. Instead, we make use of Sobolev embedding.

**Lemma 3.1**  $(L_{t\in I}^{p-1}L_x^{\infty}$  bound). Suppose that  $d = 1$  and  $p \ge 5$ , or  $d \ge 2$  and  $p \ge 3$ . Let  $s > s_c$ . Then, we have

<span id="page-6-0"></span>
$$
||u||_{L_{t \in I}^{p-1} L_x^{\infty}} \lesssim T^{0+} ||u||_{S_{q_0, r_0}^s(I)},
$$
\n(3.4)

where  $(q_0, r_0) = ((p-1)^+, \left(\frac{2d(p-1)}{d(p-1)-q}\right)$  $d(p-1)-4$  $\int$  is an admissible pair. Here, we denote by  $c^+$ a number larger than  $c$  but arbitrarily close to  $c$ , and similarly for  $c^-$ .

Proof. We observe that

$$
\frac{1}{r_0} - \frac{s - d(1 - \sigma)(\frac{1}{2} - \frac{1}{r_0})}{d} < 0.
$$

Thus, by Sobolev inequality,

$$
||u||_{L_{t \in I}^{p-1} L_x^{\infty}} \leq T^{0+} ||u||_{L_{t \in I}^{q_0} L_x^{\infty}} \leq ||\nabla|^{-d(1-\sigma)(\frac{1}{2}-\frac{1}{r_0})} u||_{L_{t \in I}^{q_0} W_x^{s,r_0}} = ||u||_{S_{q_0,r_0}^s(I)}.
$$

We also employ a standard persistence of regularity argument.

<span id="page-7-0"></span>**Lemma 3.2** (Persistence of regularity). Let  $1 < q \le \infty$ ,  $1 < r < \infty$  and  $s_1 \ge s_2$ . Then,  $B = \{u : ||u||_{L^q_{t \in I} W^{s_1, r}_x} \leq R\}$ , equipped with the norm  $|| \cdot ||_{L^q_{t \in I} W^{s_2, r}_x}$ , is a complete metric space.

Proof. We recall:

**Theorem 3.3** (Theorem 1.2.5 in [\[Caz03\]](#page-17-8)). Consider two Banach spaces  $X \rightarrow Y$  and  $1 < p, q \le \infty$ . Let  $(f_n)_{n \ge 0}$  be a bouned sequence in  $L^q(I, Y)$  and let  $f : I \to Y$  be such that  $f_n(t) \rightharpoonup f(t)$  in Y as  $n \to \infty$ , for a.e.  $t \in I$ . If  $(f_n)_{n \geq 0}$  is bounded in  $L^p(I;X)$ and if X is reflexive, then  $f \in L^p(I;X)$  and  $||f||_{L^p(I;X)} \leq \liminf_{n \to \infty}||f_n||_{L^p(I;X)}$ .

Suppose that  $(f_n)_{n=1}^{\infty}$  be a Cauchy sequence in B. Then,  $f_n$  converges to f in  $L_{t\in I}^q W^{s_2,r}_{x}$ . Moreover, it follows from Theorem 1.2.5 in [\[Caz03\]](#page-17-8) that

$$
\|f\|_{L^q_{t\in I}W^{s_1,r}_x}\leqslant \liminf_{n\to\infty}\|f_n\|_{L^q_{t\in I}W^{s_1,r}_x}\leqslant R,
$$

and thus  $f \in B$ . Therefore, we conclude that B is complete.

*Proof of Theorem [1.1](#page-2-2) when*  $d = 1$  and  $p \ge 5$ , or  $d \ge 2$  and  $p \ge 3$ . Define the map  $\Phi_{u_0}$ as above, and let

$$
X^{\alpha} := L_{t \in I}^{\infty} H_x^{\alpha} \cap S_{q_0, r_0}^{\alpha}(I),
$$

where  $(q_0, r_0)$  is an admissible pair in Lemma 3.2. Then, by Strichartz estimates, the fractional chain rule and [\(3.4\)](#page-6-0), we get

$$
\|\Phi_{u_0}(u)\|_{X^s} \lesssim \|u_0\|_{H^s} + \| |u|^{p-1} u \|_{L^1_{t\in I} H^s_x}
$$
  
\n
$$
\lesssim \|u_0\|_{H^s} + \|u\|_{L^{p-1}_{t\in I} L^\infty_x}^{p-1} \|u\|_{L^\infty_{t\in I} H^s_x}
$$
  
\n
$$
\lesssim \|u_0\|_{H^s} + T^{0+} \|u\|_{S^s_{t_0, r_0}(I)}^{p-1} \|u\|_{L^\infty_{t\in I} H^s_x}
$$
  
\n
$$
\leq \|u_0\|_{H^s} + T^{0+} \|u\|_{X^s}^p,
$$

and similarly

<span id="page-8-1"></span>
$$
\begin{split} \|\Phi_{u_{0}}(u) - \Phi_{u_{0}}(v)\|_{X^{0}} &\lesssim \||u|^{p-1}u - |v|^{p-1}v\|_{L_{t\in I}^{1}L_{x}^{2}} \\ &\lesssim \|(|u|^{p-1} + |v|^{p-1})|u - v|\|_{L_{t\in I}^{1}L_{x}^{2}} \\ &\lesssim (\|u\|_{L_{t\in I}^{p-1}L_{x}^{\infty}}^{p-1} + \|v\|_{L_{t\in I}^{p-1}L_{x}^{\infty}}^{p-1})\|u - v\|_{L_{t\in I}^{\infty}L_{x}^{2}} \\ &\lesssim T^{0+}(\|u\|_{S_{q_{0},r_{0}}^{q_{0}}(I)}^{p-1} + \|v\|_{S_{q_{0},r_{0}}^{q_{0}}(I)}^{p-1})\|u\|_{L_{t\in I}^{\infty}L_{x}^{2}} \\ &\lesssim T^{0+}(\|u\|_{X^{s}}^{p-1} + \|v\|_{X^{s}}^{p-1})\|u - v\|_{X^{0}}. \end{split} \tag{3.5}
$$

Thus, for sufficiently small  $T > 0$ ,  $\Phi_{u_0}$  is contractive on a ball

$$
B := \{ u : \|u\|_{X^s} \leq 2 \|u_0\|_{H^s} \}
$$

equipped with the norm  $\|\cdot\|_{X^0}$ , which is complete by Lemma [3.2.](#page-7-0)

Remark 3.4. The standard persistence of regularity argument allows us to avoid derivatives in [\(3.5\)](#page-8-1). Indeed, for  $u \in B$ ,  $\|\langle \nabla \rangle^s u\|_{L^{p-1}_{t \in I} L^{\infty}_x}$  is not necessarily bounded.

<span id="page-8-0"></span>3.2. Scaling-critical cases. In the scaling-critical case, we use the following lemma, which plays the same role as [\(3.4\)](#page-6-0). We note that the norms in the lemma are defined via the Littlewood-Paley projection in order to overcome the failure of the Sobolev embedding  $W^{s,p} \hookrightarrow L^q$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{s}{d}$  $\frac{s}{d}$ , when  $q = \infty$ . Lemma 3.3 generalizes [\[CKS](#page-17-6)<sup>+</sup>08, Lemma 3.1].

**Lemma 3.5** (Scaling-critical  $L_{t \in I}^{p-1} L_x^{\infty}$  bound).

<span id="page-8-3"></span>
$$
||u||_{L_{\text{ter}}^{p-1}L_x^{\infty}}^{p-1} \leq \begin{cases} ||u||_{\tilde{S}_{2,\infty}^{sc}(I)}^{\frac{4}{3}} ||u||_{\tilde{S}_{\infty,2}^{sc}(I)}^{p-5} & when d = 1 \text{ and } p > 5, \\ ||u||_{\tilde{S}_{2+\infty}^{sc}(I)}^{\frac{2}{3}} ||u||_{\tilde{S}_{\infty,2}^{sc}(I)}^{p-3} & when d = 2 \text{ and } p > 3, \\ ||u||_{\tilde{S}_{2,\frac{2d}{d-2}}^{2}}^{p-3} (I)||u||_{\tilde{S}_{\infty,2}^{sc}(I)}^{p-3} & when d \geq 3 \text{ and } p > 3. \end{cases} (3.6)
$$

*Proof.* We will prove the lemma only when  $d \ge 3$ . By interpolation  $||f||_{L^{p_\theta}} \le$  $|| f ||_{L^{p_0}}^{\theta} ||f||_{L^{p_1}}^{1-\theta}, \frac{1}{p_\theta}$  $\frac{1}{p_\theta}$  =  $\frac{\theta}{p_0}$  $\frac{\theta}{p_0} + \frac{1-\theta}{p_1}$  $\frac{q-\theta}{p_1}$ ,  $0 < \theta < 1$ , it suffices to show the lemma for rational  $(p-1) = \frac{m}{n} > 2$  with  $gcd(m, n) = 1$ . First, we estimate

$$
A(t) = \left[\sum_{N} \|P_N u(t)\|_{L_x^{\infty}}\right]^m \sim \sum_{N_1 \geqslant \cdots \geqslant N_m} \prod_{i=1}^m \|P_{N_i} u(t)\|_{L_x^{\infty}}.
$$

Observe from Bernstein's inequality that

<span id="page-8-2"></span>
$$
||P_N u(t)||_{L_x^{\infty}} \lesssim N^{-\frac{\sigma(p-3)}{p-1}} d_N,
$$
\n(3.7)

$$
||P_N u(t)||_{L_x^{\infty}} \lesssim N^{\frac{2\sigma}{p-1}} d_N',\tag{3.8}
$$

where

$$
d_N = \|P_N u(t)\|_{\dot{W}^{s_c-(1-\sigma),\frac{2d}{d-2}}_x}, \ d'_N = \|P_N u(t)\|_{\dot{H}^{s_c}_x}.
$$

As a consequence, we have

<span id="page-9-0"></span>
$$
\|P_N u(t)\|_{L_x^{\infty}} \lesssim \left(N^{-\frac{\sigma(p-3)}{p-1}} d_N\right)^{\theta} \left(N^{\frac{2\sigma}{p-1}} d_N'\right)^{1-\theta} = N^{\frac{\sigma(p-3)}{(p-1)(p-2)}} (d_N)^{\theta} (d_N')^{1-\theta},\tag{3.9}
$$

where  $\theta = \frac{1}{n}$  $\frac{1}{p-2}$ . Hence, applying [\(3.7\)](#page-8-2) for  $i = 1, \dots, n$  and [\(3.9\)](#page-9-0) for  $i = n+1, \dots, m$ , we bound  $A(t)$  by

$$
\lesssim \sum_{N_1 \geqslant \cdots \geqslant N_m} \Big(\prod_{i=1}^n N_i^{-\frac{\sigma(p-3)}{p-1}} d_{N_i}\Big) \Big(\prod_{i=n+1}^m N_i^{\frac{\sigma(p-3)}{(p-1)(p-2)}} (d_{N_i})^{\theta} (d'_{N_i})^{1-\theta}\Big).
$$

For an arbitrarily small  $\epsilon > 0$ , we let

$$
\tilde{d}_N = \sum_{N' \in 2^{\mathbb{Z}}} \min \left( \frac{N}{N'}, \frac{N'}{N} \right)^{\epsilon} d_{N'}, \quad \tilde{d}'_N = \sum_{N' \in 2^{\mathbb{Z}}} \min \left( \frac{N}{N'}, \frac{N'}{N} \right)^{\epsilon} d'_{N'}.
$$

Then, since  $d_N \leq \tilde{d}_N$  and  $\tilde{d}_{N_i} \leq (\frac{N_1}{N_i})^{\epsilon} \tilde{d}_{N_1}$  and similarly for primes,  $A(t)$  is bounded by

$$
\lesssim \sum_{N_1 \geqslant \cdots \geqslant N_m} \bigg( \prod_{i=1}^n N_i^{- \frac{\sigma(p-3)}{p-1}} \Big(\frac{N_1}{N_i}\Big)^\epsilon \tilde{d}_{N_1} \bigg) \bigg( \prod_{i=n+1}^m N_i^{\frac{\sigma(p-3)}{(p-1)(p-2)}} \Big(\frac{N_1}{N_i}\Big)^\epsilon (\tilde{d}_{N_1})^{\theta} (\tilde{d}_{N_1}')^{1-\theta} \bigg).
$$

Summing in  $N_m, N_{m-1}, ..., N_{n+1}$  and using  $m - n = (p - 2)n$ ,

$$
A(t) \lesssim \sum_{N_1 \geq \dots \geq N_n} \Big( \prod_{i=1}^n N_i^{-\frac{\sigma(p-3)}{p-1}} \Big( \frac{N_1}{N_i} \Big)^{\epsilon} \tilde{d}_{N_1} \Big)
$$
  
\$\times N\_n^{-\frac{\sigma(p-3)(m-n)}{(p-1)(p-2)}} \Big( \frac{N\_1}{N\_n} \Big)^{(m-n)\epsilon} (\tilde{d}\_{N\_1})^{(m-n)\theta} (\tilde{d}'\_{N\_1})^{(m-n)(1-\theta)}\$  

$$
= \sum_{N_1 \geq \dots \geq N_n} \Big( \prod_{i=1}^n N_i^{-\frac{\sigma(p-3)}{p-1}} \Big( \frac{N_1}{N_i} \Big)^{\epsilon} \tilde{d}_{N_1} \Big)
$$
  
\$\times N\_n^{-\frac{\sigma(p-3)n}{p-1}} \Big( \frac{N\_1}{N\_n} \Big)^{(p-2)n\epsilon} (\tilde{d}\_{N\_1})^{(m-n)\theta} (\tilde{d}'\_{N\_1})^{(m-n)(1-\theta)},

and then summing in  $N_n, N_{n-1}, ..., N_1$ , we obtain that

$$
A(t) \lesssim \sum_{N_1} (\tilde{d}_{N_1})^{n+(m-n)\theta} (\tilde{d}'_{N_1})^{(m-n)(1-\theta)} = \sum_{N_1} (\tilde{d}_{N_1})^{2n} (\tilde{d}'_{N_1})^{m-2n},
$$

which is, by Hölder inequality and Young's inequality, bounded by

$$
\begin{array}{l} \lesssim \displaystyle \|\tilde{(d_N)}^{2n}\|_{\ell^2_N}\|(\tilde{d}'_N)^{m-2n}\|_{\ell^2_N}=\|\tilde{d}_N\|_{\ell^{4n}_N}^{2n}\|\tilde{d}'_N\|_{\ell^{2(m-2n)}_N}^{m-2n}\\ \leqslant \displaystyle \|\tilde{d}_N\|_{\ell^2_N}^{2n}\|\tilde{d}'_N\|_{\ell^2_N}^{m-2n}\lesssim \displaystyle \|d_N\|_{\ell^2_N}^{2n}\|d'_N\|_{\ell^2_N}^{m-2n}=\displaystyle \|d_N\|_{\ell^2_N}^{2n}\|d'_N\|_{\ell^2_N}^{(p-3)n}.\end{array}
$$

Finally, by the estimate for  $A(t)$ , we prove that

$$
\|u\|_{L^{p-1}_{t\in I}L^\infty_x}^{p-1}\leqslant \int_{I}A(t)^{p-1}dt=\int_{I}A(t)^{\frac{m}{n}}dt\\ \lesssim \int_{I}\|d_N\|_{\ell^2_N}^2\|d_N'\|_{\ell^2_N}^{p-3}dt\leqslant \|d_N\|_{L^2_{t\in I}\ell^2_N}^2\|d_N'\|_{L^{\infty}_{t\in I}\ell^2_N}^{p-3}\\ \lesssim \|u\|_{\tilde{S}^{s_c}_{\frac{2d}{d-2}}(I)}^2\|u\|_{\tilde{S}^{s_c}_{\infty,2}(I)}^{p-3}.
$$

*Proof of theorem [1.2.](#page-2-1)* For simplicity, we assume that  $d \geq 3$ . Indeed, with little modifications, we can prove the theorem when  $d = 1, 2$ . We define  $\Phi_{u_0}(u)$  as in the proof of Theorem [1.1.](#page-2-2) Then, by Strichartz estimates, the fractional chain rule and [\(3.6\)](#page-8-3), we have

$$
\begin{aligned} \|\Phi_{u_0}(u)\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)} &\leqslant \|e^{it(-\Delta)^{\sigma}}u_0\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)} + c_0 \| |u|^{p-1}u\|_{L^1_{t\in I}H^{sc}}\\ &\leqslant \|e^{it(-\Delta)^{\sigma}}u_0\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)} + c_1 \|u\|_{L^{{p-1}}_{t\in I}L^\infty_x}^{p-1} \|u\|_{L^\infty_{t\in I}H^{sc}}\\ &\leqslant \|e^{it(-\Delta)^{\sigma}}u_0\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)} + c \|u\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)}^{p-2} \|u\|_{\tilde{S}^{sc}_{\infty,2}(I)}^{p-2}. \end{aligned}
$$

Similarly, one can show that

$$
\|\Phi_{u_0}(u)\|_{\tilde{S}^{sc}_{\infty,2}(I)} \leq c\|u_0\|_{H^{s_c}} + c\|u\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)}^{2}\|u\|_{\tilde{S}^{sc}_{\infty,2}(I)}^{p-2}
$$

and

$$
\label{eq:20} \begin{split} &\|\Phi_{u_0}(u)-\Phi_{u_0}(v)\|_{\tilde{S}_{2,\frac{2d}{d-2}}^0(I)}+\|\Phi_{u_0}(u)-\Phi_{u_0}(v)\|_{\tilde{S}_{\infty,2}^0(I)}\\ &\leqslant c_0(\|u\|_{L^{p-1}_{t\in I}L^\infty_x}^{p-1}+\|v\|_{L^{p-1}_{t\in I}L^\infty_x}^{p-1})\|u-v\|_{L^\infty_{t\in I}L^2_x}\\ &\leqslant c(\|u\|_{\tilde{S}_{2,\frac{2d}{d-2}}^s(I)}^2\|u\|_{\tilde{S}_{\infty,2}^s(I)}^{p-2}+\|v\|_{\tilde{S}_{2,\frac{2d}{d-2}}^s(I)}^2\|v\|_{\tilde{S}_{\infty,2}^s(I)}^{p-2})\|u-v\|_{L^\infty_{t\in I}L^2_x}. \end{split}
$$

Now we let  $\delta = \delta(c, ||u_0||_{H^{s_c}}) > 0$  be a sufficiently small number to be chosen later, and then we pick  $T = T(u_0, \delta) > 0$  such that

$$
\|e^{it(-\Delta)^{\sigma}}u_0\|_{\tilde{S}^{s_c}_{2,\frac{2d}{d-2}}(I)} \leq \delta,
$$

Define

$$
B = \left\{ u : \|u\|_{\tilde{S}^{s_c}_{2,\frac{2d}{d-2}}(I)} \leq 2\delta \text{ and } \|u\|_{\tilde{S}^{s_c}_{\infty,2}(I)} \leq 2c\|u_0\|_{H^{s_c}} \right\}
$$

equipped with the norm

$$
\|u\|_X:=\|u\|_{\tilde{S}^0_{2,\frac{2d}{d-2}}(I)}+\|u\|_{\tilde{S}^0_{\infty,2}(I)}.
$$

 $\Box$ 

Then, for  $u \in B$ , we have

$$
\|\Phi_{u_0}(u)\|_{\tilde{S}^{sc}_{2,\frac{2d}{d-2}}(I)} \leq \delta + c(2\delta)^2 (2c\|u_0\|_{H^{sc}})^{p-2} \leq 2\delta,
$$
  

$$
\|\Phi_{u_0}(u)\|_{\tilde{S}^{sc}_{\infty,2}(I)} \leq c\|u_0\|_{H^{sc}} + c(2\delta)^2 (2c\|u_0\|_{H^{sc}})^{p-2} \leq 2c\|u_0\|_{H^{sc}}.
$$

Choosing sufficiently small  $\delta > 0$ , we prove that  $\Phi_{u_0}$  maps B to itself. Similarly, one can show

$$
\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\|_X \leq \frac{1}{2} \|u - v\|_X.
$$

Therefore, it follows that  $\Phi_{u_0}$  is a contraction mapping in B.

Remark 3.6. (i) In the proofs, the  $L_x^{\infty}$  norm bounds are crucial for the following reason. In Proposition [2.1,](#page-4-2) there is a loss of regularity except the trivial ones,

$$
\|e^{it(-\Delta)^\sigma}u_0\|_{L^\infty_{t\in I}L^2_x}=\|u_0\|_{L^2}
$$

and

$$
\Big\| \int_0^t e^{it(-\Delta)^\sigma} F(s) ds \Big\|_{L^\infty_{t \in I} L^2_x} \leqslant \|F\|_{L^1_{t \in I} L^2_x}.
$$

Hence, when we estimate the  $L_{t \in I}^{\infty} H_x^s$  norm of the integral term in  $\Phi_{u_0}(u)$ , we are forced to use the trivial one

$$
\left\| \int_0^t e^{it(-\Delta)^{\sigma}} |u|^{p-1} u(s) ds \right\|_{L^{\infty}_{t \in I} H^s_x} \leq \| |u|^{p-1} u \|_{L^1_{t \in I} H^s_x}.
$$
\n(3.10)

Indeed, otherwise, we have a higher regularity norm on the right hand side. Then, we cannot close the contraction mapping argument. Moreover, if  $u_0 \in H^s$ , there is no good bound for  $||e^{it(-\Delta)^{\sigma}}u_0||_{L^q_{teI}W_x^{s,r}}$  except the trivial one  $(q, r) = (\infty, 2)$ . Thus, we are forced to bound the right hand side of (3.10) by

$$
\|u\|_{L^{p-1}_{t\in I}L^\infty_x}^{p-1}\|u\|_{L^\infty_{t\in I}H^s_x}.
$$

Therefore, we should have a good control on  $||u||_{L^{p-1}_{t\in I}L^\infty_x}$ .

(*ii*) When  $p < 3$ , the  $L_{t \in I}^{p-1} L_x^{\infty}$  norm is scaling-supercritical. Thus, based on our method, the assumptions on p in Theorem [1.1](#page-2-2) and [1.2](#page-2-1) are optimal except  $p = 3$  in the critical case.

# 4. Small Data Scattering

<span id="page-11-0"></span>*Proof of Theorem [1.3.](#page-2-3)* For simplicity, we consider the case  $d \geq 3$  only. It follows from the estimates in the proof of Theorem [1.2](#page-2-1) that if  $||u_0||_{H^s}$  is small enough, then

$$
\|u(t)\|_{L^{p-1}_{t\in\mathbb R}L^\infty_x}+\|u(t)\|_{L^\infty_{t\in\mathbb R}H^{s_c}_x}\leqslant\|u(t)\|_{\tilde{S}^{s_c}_{2,\frac{2d}{d-2}}(\mathbb R)}+\|u(t)\|_{\tilde{S}^{s_c}_{\infty,2}(\mathbb R)}\lesssim\|u_0\|_{H^{s_c}}<\infty.
$$

By Strichartz estimates, the fractional chain rule and [\(3.6\)](#page-8-3), we prove that

$$
\|e^{-iT_1(-\Delta)^{\sigma}}u(T_1) - e^{-iT_2(-\Delta)^{\sigma}}u(T_2)\|_{H^{sc}}
$$
  
= 
$$
\Big\|\int_{T_1}^{T_2} e^{-is(-\Delta)^{\sigma}}(|u|^{p-1}u)(s)ds\Big\|_{H^{sc}}
$$
  

$$
\leq \|u(t)\|_{L^{p-1}_{t\in[T_1,T_2)}L^{\infty}_x}^{p-1} \|u(t)\|_{L^{\infty}_{t\in[T_1,T_2)}H^{sc}_x} \to 0
$$

as  $T_1, T_2 \to \pm \infty$ . Thus, the limits

$$
u_{\pm} = \lim_{t \to \pm \infty} e^{-it(-\Delta)^{\sigma}} u(t)
$$

exist in  $H^{s_c}$ . Repeating the above estimates, we show that

$$
||u(t) - e^{it(-\Delta)^{\sigma}} u_{\pm}||_{H^{s_c}} = ||e^{-it(-\Delta)^{\sigma}} u(t) - u_{\pm}||_{H^{s_c}} \to 0
$$

<span id="page-12-0"></span>as  $t \to \pm \infty$ .

#### 5. Ill-posedness

We will prove Theorem [1.5](#page-3-1) following the strategy in [\[CCT03a\]](#page-17-9). Throughout this section, we assume that  $d = 1, 2$  or 3 and  $\frac{d}{4} < \sigma < 1$ . If p is not an odd integer, we further assume that  $p \ge k + 1$ , where k is the smallest integer greater than  $\frac{d}{2}$ .

First, we construct an almost non-dispersive solution by small dispersion analysis.

<span id="page-12-1"></span>**Lemma 5.1** (Small dispersion analysis). Given a Schwartz function  $\phi_0$ , let  $\phi^{(\nu)}(t,x)$ be the solution to the fractional NLS

<span id="page-12-4"></span>
$$
i\partial_t u + \nu^{2\sigma} (-\Delta)^{\sigma} u + \mu |u|^{p-1} u = 0, \ u(0) = \phi_0,
$$
\n(5.1)

and  $\phi^{(0)}(t,x)$  be the solution to the ODE with no dispersion

$$
i\partial_t u + \mu |u|^{p-1}u = 0, \ u(0) = \phi_0,
$$

that is,

<span id="page-12-2"></span>
$$
\phi^{(0)}(t,x) = \phi_0(x)e^{it\omega|\phi_0(x)|^{p-1}}.\tag{5.2}
$$

Then there exist  $C, c > 0$  such that if  $0 < \nu \leq c$  is sufficiently small, then

<span id="page-12-3"></span>
$$
\|\phi^{(\nu)}(t) - \phi^{(0)}(t)\|_{H^k} \leq C\nu^{2\sigma} \tag{5.3}
$$

for all  $|t| \leq c |\log \nu|^c$ .

*Proof.* The proof closely follows the proof of Lemma 2.1 in [\[CCT03a\]](#page-17-9).  $\Box$ 

Obviously,  $\phi^{(\nu)}(t,\nu x)$  is a solution to (NLS<sub>σ</sub>). Moreover,  $\phi^{(\nu)}(t,\nu x)$  is bounded and almost flat in the following sense.

**Corollary 5.2.** Let  $\phi^{(\nu)}$ , v and c be in Lemma [5.1.](#page-12-1) Let  $s \geq 0$ . Then,

<span id="page-13-0"></span>
$$
\|\phi^{(\nu)}(t,\nu x)\|_{L_x^{\infty}} \sim 1\tag{5.4}
$$

and

<span id="page-13-2"></span>
$$
\|\phi^{(\nu)}(t,\nu x)\|_{\dot{H}^s_x} \sim \nu^{s-\frac{d}{2}}(c|\log \nu|^c)^s \tag{5.5}
$$

for all  $|t| \leq c |\log \nu|^c$ .

*Proof.* Since  $k > \frac{d}{2}$  $\frac{d}{2}$ , by the Sobolev inequality, we have

$$
\begin{aligned} \|\phi^{(\nu)}(t,\nu x) - \phi^{(0)}(t,\nu x)\|_{L_x^{\infty}} &= \|\phi^{(\nu)}(t,x) - \phi^{(0)}(t,x)\|_{L_x^{\infty}} \\ &\leq \|\phi^{(\nu)}(t) - \phi^{(0)}(t)\|_{H^k} \leq \nu^{2\sigma} .\end{aligned}
$$

Then, [\(5.4\)](#page-13-0) follows from the explicit formula [\(5.2\)](#page-12-2) for  $\phi^{(0)}(t, x)$ . It follows from [\(5.3\)](#page-12-3) and [\(5.2\)](#page-12-2) that

$$
\|\phi^{(\nu)}(t,\nu x)\|_{\dot{H}^s_x} \leqslant \nu^{s-\frac{d}{2}}(\|\phi^{(0)}(t)\|_{\dot{H}^s} + \|\phi^{(\nu)}(t) - \phi^{(0)}(t)\|_{\dot{H}^s}) \sim \nu^{s-\frac{d}{2}}(c|\log \nu|^c)^s.
$$

For  $v \in \mathbb{R}^d$ , we define the pseudo-Galilean transformation by

$$
\mathcal{G}_{v}u(t,x)=e^{-iv\cdot x}e^{it|v|^{2\sigma}}u(t,x-2t\sigma|v|^{2(\sigma-1)}v).
$$

Note that when  $\sigma = 1$ ,  $\mathcal{G}_v$  is simply a Galilean transformation, and that NLS is invariant under this transformation, that is, if  $u(t)$  solves NLS, so does  $\mathcal{G}_v u(t)$ . However, when  $\sigma \neq 1$ , (NLS<sub> $\sigma$ </sub>) is not exactly symmetric with respect to pseudo-Galilean transformations. Indeed, if  $u(t)$  solves  $(NLS_{\sigma})$ , then  $\tilde{u}(t) = \mathcal{G}_{v}u(t)$  obeys  $(NLS_{\sigma})$  with an error term

$$
i\partial_t \tilde{u} + (-\Delta)^\sigma \tilde{u} + \omega |\tilde{u}|^{p-1} \tilde{u} = e^{it|v|^{2\sigma}} e^{-iv \cdot x} (\mathcal{E}u)(t, x - 2\sigma t |v|^{2(\sigma-1)} v), \qquad (5.6)
$$

where

$$
\mathcal{E}\tilde{u}(\xi) = E(\xi)\hat{u}(\xi)
$$

with

$$
E(\xi) = |\xi - v|^{2\sigma} - |\xi|^{2\sigma} - |v|^{2\sigma} + 2\sigma |v|^{2(\sigma - 1)} v \cdot \xi.
$$

However, we note that

<span id="page-13-1"></span>
$$
|E(\xi)| \lesssim |\xi|^{2\sigma}.\tag{5.7}
$$

Indeed, if  $|\xi| \leq \frac{|v|}{100}$ , then

$$
E(\xi) = \left| |v|^{2\sigma} \left( |\frac{v}{|v|} - \frac{\xi}{|v|} |^{2\sigma} - 1 + 2\sigma \frac{v}{|v|} \cdot \frac{\xi}{|v|} \right) - |\xi|^{2\sigma} \right| \lesssim |v|^{2\sigma} \frac{|\xi|^2}{|v|^2} + |\xi|^{2\sigma} \lesssim |\xi|^{2\sigma}.
$$

Otherwise,

$$
E(\xi) \lesssim |\xi|^{2\sigma} + |v|^{2\sigma} + |\xi|^{2\sigma} + |v|^{2\sigma} + 2\sigma |v|^{2\sigma - 1} |\xi| \lesssim |\xi|^{2\sigma}.
$$

Therefore, one would expect an *almost* symmetry for an almost flat solution  $u(t)$ , such as  $\phi^{(\nu)}(t,\nu x)$  in Lemma [5.1.](#page-12-1) Precisely, we have the following lemma.

<span id="page-14-1"></span>**Lemma 5.3** (Pseudo-Galilean transformation). Let  $\phi^{(\nu)}$ , v and c be in Lemma [5.1.](#page-12-1) For  $v \in \mathbb{R}^d$ , we define

$$
\tilde{u}(t,x)=(\mathcal{G}_v\phi^{(\nu)}(\cdot,\nu\cdot))(t,x)=e^{-iv\cdot x}e^{it|v|^{2\sigma}}\phi^{(\nu)}\big(t,\nu(x-2t\sigma|v|^{2(\sigma-1)}v)\big),
$$

and let  $u(t, x)$  be the solution to  $(NLS_{\sigma})$  with the same initial data

$$
e^{-iv \cdot x} \phi^{(\nu)}(0, \nu x) = e^{-iv \cdot x} \phi_0(0, \nu x). \tag{5.8}
$$

Then, there exists  $\delta > 0$  such that

<span id="page-14-0"></span>
$$
\|e^{iv\cdot x}(u(t) - \tilde{u}(t))\|_{H_x^k} \lesssim \nu^{\delta} \tag{5.9}
$$

for all  $|t| \leq c |\log \nu|^c$ .

*Remark* 5.4. When  $p = 3$ , in [\[CHKL14\]](#page-17-3) the authors could use the counterexample in [\[CCT03b\]](#page-17-10). This counterexample is constructed by pseudo-conformal symmetry and Galilean transformation. A good thing is that this solution is very small in high Sobolev norms, too. Somehow, this smallness allows [\[CHKL14\]](#page-17-3) to show that the error in pseudo-Galilean transformation is also small. However, when  $p > 3$ , the counterexample in [\[CCT03b\]](#page-17-10) does not work. Later, Christ, Colliander and Tao [\[CCT03a\]](#page-17-9) constructed a different counterexample which works for more general  $p$ . Unfortunately, this counterexample is not small in high Sobolev norms. It is very large instead. In particular, for our purposes, it is hard to control the error from pseudo-Galilean transformation. But, our new counterexample still has small high Sobolev norm after translating it to its frequency center; this is the term  $e^{iv\cdot x}$  in equation [\(5.9\)](#page-14-0). Using this smallness, we can prove that pseudo-Galilean transformation is almost invariant. We also remark that the condition  $\sigma > \frac{d}{4}$  $\frac{d}{4}$  is to guarantee smallness of the error (see  $(5.10)$ ).

*Proof of Lemma [5.3.](#page-14-1)* Let  $R(t) = (u - \tilde{u})(t)$ . Then,  $R(t)$  satisfies

$$
i\partial_t R + (-\Delta)^{\sigma} R = \mu (|\tilde{u}|^{p-1}\tilde{u} - |u|^{p-1}u) - e^{it|v|^{2\sigma}} (\mathcal{E}\phi^{(\nu)}(t, \nu(x - 2\sigma t|v|^{2(\sigma-1)}v)),
$$

or equivalently

$$
R(t) = i \int_0^t e^{i(t-s)(-\Delta)\sigma} \left\{ \mu \left( |u|^{p-1}u - |\tilde{u}|^{p-1}\tilde{u} \right)(s) + e^{is|v|^{2\sigma}} \left( \mathcal{E}\phi^{(\nu)}\left(s, \nu(x - 2\sigma s|v|^{2(\sigma-1)}v)\right) \right) ds \right\} ds.
$$

Hence, by a trivial estimate, we get

$$
||e^{iv\cdot x}R(t)||_{H^k} \le \int_0^t ||e^{iv\cdot x} (|u|^{p-1}u - |\tilde{u}|^{p-1}\tilde{u})(s)||_{H^k} + ||\mathcal{E}\phi^{(\nu)}(s,\nu \cdot)||_{H^k} ds
$$
  
= 
$$
\int_0^t I(s) + II(s)ds.
$$

First, by  $(5.7)$  and  $(5.5)$ , we show that

<span id="page-15-0"></span>
$$
\int_0^t II(s)ds \lesssim \int_0^t \sum_{j=0}^k \|\phi^{(\nu)}(s,\nu\cdot)\|_{\dot{H}^{j+2\sigma}} ds \sim (c|\log \nu|^c)^{1+2\sigma-\frac{d}{2}}\nu^{2\sigma-\frac{d}{2}}.\tag{5.10}
$$

For  $I(s)$ , expanding  $u = \tilde{u} + R$  and then applying Hölder inequality and Sobolev inequalities, we bound  $I(s)$  by

$$
\lesssim \sum_{j=1}^{p} \|e^{iv \cdot x} R\|_{H^k}^j.
$$
\n(5.11)

For example, when  $p = 3$ ,

$$
I(s) \leq 2||e^{iv\cdot x}\tilde{u}|^2 e^{iv\cdot x}R||_{H^k} + ||(e^{iv\cdot x}\tilde{u})^2 e^{iv\cdot x}\overline{R}||_{H^k} + 2||e^{iv\cdot x}\tilde{u}|e^{iv\cdot x}R|^2||_{H^k}
$$
  
+  $||e^{iv\cdot x}\tilde{u}(e^{iv\cdot x}R)^2||_{H^k} + ||e^{iv\cdot x}R|^2 e^{iv\cdot x}R||_{H^k}$   
=:  $I_1(s) + I_2(s) + I_3(s) + I_4(s) + I_5(s)$ .

Consider

$$
I_1(s) = \sum_{|\alpha| \leq k} \|\nabla_{x_1}^{\alpha_1} \cdots \nabla_{x_d}^{\alpha_d} (|e^{iv \cdot x} \tilde{u}|^2 e^{iv \cdot x} R)(s)\|_{L^2} =: \sum_{|\alpha| \leq k} I_{1,\alpha}(s),
$$

where  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$  is a multi-index with  $|\alpha| = \sum_{i=1}^d \alpha_i$ . Observe that whenever a derivative hits

$$
e^{iv\cdot x}\tilde{u}(s) = e^{is|v|^{2\sigma}}\phi^{(\nu)}\big(s,\nu(x-2s\sigma|v|^{2(\sigma-1)}v)\big),\,
$$

we get a small factor  $\nu$ . Hence, after distributing derivatives by the Leibniz rule, the worst term we have in  $I_{1,\alpha}(s)$  is

$$
\| |e^{iv\cdot x}\tilde{u}(s)|^2 \nabla^\alpha e^{iv\cdot x} R(s)\|_{L^2},
$$

which is, by  $(5.4)$ , bounded by

$$
\|e^{iv\cdot x}\tilde{u}(s)\|_{L^{\infty}}^2 \|\nabla^{\alpha}e^{iv\cdot x}R(s)\|_{L^2} \sim \|\nabla^{\alpha}e^{iv\cdot x}R(s)\|_{L^2}.
$$

Likewise, we estimate other terms.

Collecting all,

$$
\|e^{iv\cdot x}R(t)\|_{H^k}\lesssim (c|\log \nu|^c)^{1+2\sigma-\frac{d}{2}}\nu^{2\sigma-\frac{d}{2}}+\int_0^t\sum_{j=1}^p\|e^{iv\cdot x}R(s)\|^j_{H^k}ds
$$

for  $|t| \leq c |\log \nu|^c$ . Then, by the standard nonlinear iteration argument, we prove the  $l$ emma.

Since we have solutions almost symmetric with respect to the pseudo-Galilean transformations, we can make use of the following decoherence lemma to construct counterexamples for local well-posedness.

<span id="page-16-0"></span>**Lemma 5.5** (Decoherence). Let  $s < 0$ . Fix a nonzero Schwartz function w. For  $a, a' \in \left[\frac{1}{2}, 1\right], 0 < \nu \leq \lambda \ll 1$  and  $v \in \mathbb{R}^d$  with  $|v| \geq 1$ , we define

$$
\tilde{u}^{(a,\nu,\lambda,v)}(t,x):=\mathcal{G}_{v}\Big(\lambda^{-\frac{2\sigma}{p-1}}\phi^{(a,\nu)}(\lambda^{-2\sigma}\cdot,\lambda^{-1}\nu\cdot)\Big)(t,x),
$$

where  $\phi^{(a,\nu)}$  is the solution to [\(5.1\)](#page-12-4) with initial data aw. Then, we have

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(0)\|_{H^s}, \|\tilde{u}^{(a',\nu,\lambda,v)}(0)\|_{H^s} \leq C|v|^s \lambda^{-\frac{2\sigma}{p-1}} \left(\frac{\lambda}{\nu}\right)^{d/2},
$$

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(0) - \tilde{u}^{(a',\nu,\lambda,v)}(0)\|_{H^s} \leq C|v|^s \lambda^{-\frac{2\sigma}{p-1}} \left(\frac{\lambda}{\nu}\right)^{d/2} |a - a'|
$$

and

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(t) - \tilde{u}^{(a',\nu,\lambda,v)}(t)\|_{H^s}
$$
\n
$$
\geq c|v|^s \lambda^{-\frac{2\sigma}{p-1}} \left(\frac{\lambda}{\nu}\right)^{d/2} \left\{ \|(\phi^{(a,\nu)}(\frac{t}{\lambda^{2\sigma}}) - \phi^{(a',\nu)}(\frac{t}{\lambda^{2\sigma}}) \|_{L^2} - C |\log \nu|^C (\frac{\lambda}{\nu})^{-k} |v|^{-s-k} \right\}
$$
\nfor all  $|t| \leq c |\log \nu|^c \lambda^{2\sigma}$ .

*Proof.* The proof closely follows the proof of Lemma 3.1 in [\[CCT03a\]](#page-17-9).  $\Box$ 

Proof of Theorem [1.5.](#page-3-1) The proof is very similar to that of Theorem 1 in [\[CCT03a\]](#page-17-9) except that in the last step, we need to use Lemma [5.3](#page-14-1) due to lack of exact symmetry. We give a proof for the readers' convenience.

Let  $\epsilon > 0$  be a given but arbitrarily small number. Let  $\nu = \lambda^{\alpha}$ , where  $\alpha > 0$  is a small number to be chosen later. Then, we pick  $v \in \mathbb{R}^d$  such that

$$
\lambda^{-\frac{2\sigma}{p-1}}|v|^s(\lambda/\nu)^{d/2}=\epsilon \Leftrightarrow |v|=\nu^{\frac{1}{s}\left(\frac{d(1-\alpha)}{2}+\frac{2\alpha\sigma}{p-1}\right)}\epsilon^{1/s}.
$$

Note that since  $s < 0$ ,  $\frac{1}{s}(\frac{d(1-\alpha)}{2} + \frac{2\alpha\sigma}{p-1})$  $\frac{2\alpha\sigma}{p-1}$ ) =  $\frac{1}{s}(\frac{d}{2} - \alpha s_c)$  < 0 for sufficiently small  $\alpha$ , and thus  $|v| \ge 1$ . Hence, it follows from Lemma [5.5](#page-16-0) that

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(0)\|_{H^s}, \|\tilde{u}^{(a',\nu,\lambda,v)}(0)\|_{H^s} \leqslant C\epsilon,
$$
\n
$$
(5.12)
$$

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(0) - \tilde{u}^{(a',\nu,\lambda,v)}(0)\|_{H^s} \leq C\epsilon |a - a'|,\tag{5.13}
$$

and

$$
\begin{aligned} &\|\tilde{u}^{(a,\nu,\lambda,v)}(t) - \tilde{u}^{(a',\nu,\lambda,v)}(t)\|_{H^s} \\ &\geqslant c\epsilon \Big\{\|(\phi^{(a,\nu)}(\tfrac{t}{\lambda^{2\sigma}}) - \phi^{(a',\nu)}(\tfrac{t}{\lambda^{2\sigma}})\|_{L^2} - C|\log \nu|^{C}(\tfrac{\lambda}{\nu})^{-k}|v|^{-s-k}\Big\} \end{aligned}
$$

for all  $|t| \leq c |\log \nu|^c \lambda^{2\sigma}$ . Now we observe from the explicit formula [\(5.2\)](#page-12-2) for  $\phi^{(a,0)}$ and [\(5.1\)](#page-12-4) that there exists  $T > 0$  such that  $\|\phi^{(a,\nu)}(T) - \phi^{(a',\nu)}(T)\|_{L^2} \geq c$ . Moverover, if  $\alpha > 0$  is sufficiently small,  $C|\log \nu|^C(\frac{\lambda}{\nu})$  $(\frac{\lambda}{\nu})^{-k}|v|^{-s-k} \to 0$  as  $\nu \to 0$ . Therefore, for  $\nu$ small enough, we have

$$
\|\tilde{u}^{(a,\nu,\lambda,v)}(\lambda^{2\sigma}T) - \tilde{u}^{(a',\nu,\lambda,v)}(\lambda^{2\sigma}T)\|_{H^s} \geqslant c\epsilon. \tag{5.14}
$$

Next, we replace  $\tilde{u}^{(a,\nu,\lambda,v)}$  and  $\tilde{u}^{(a',\nu,\lambda,v)}$  in (6.11), (6.12) and (6.13) by  $u^{(a,\nu,\lambda,v)}$  and  $u^{(a',\nu,\lambda,v)}$  by Lemma [5.3](#page-14-1) with  $O(\nu^{\delta})$  erorr. Then, making  $|a-a'|$  arbitrarily small and then sending  $\nu \to 0$  (so,  $\lambda^{2\sigma} T \to 0$ ), we complete the proof.

# <span id="page-17-0"></span>**ACKNOWLEDGEMENTS**

Y.H. would like to thank IHES for their hospitality and support while he visited in the summer of 2014. Y.S. would like to thank the hospitality of the Department of Mathematics at University of Texas at Austin where part of the work was initiated. Y.S. acknowledges the support of ANR grants "HAB" and "NONLOCAL".

#### <span id="page-17-1"></span>**REFERENCES**

- <span id="page-17-2"></span>[Ber96] Jean Bertoin. *L´evy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- <span id="page-17-8"></span>[Caz03] Thierry Cazenave. *Semilinear Schrödinger equations*, volume 10 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.
- <span id="page-17-9"></span>[CCT03a] M. Christ, J. Colliander, and T. Tao. Ill-posedness for nonlinear schrdingier and wave equations. *arXiv:math/0311048*, 2003.
- <span id="page-17-10"></span>[CCT03b] Michael Christ, James Colliander, and Terrence Tao. Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math.*, 125(6):1235–1293, 2003.
- <span id="page-17-4"></span>[CHHO13] Yonggeun Cho, Hichem Hajaiej, Gyeongha Hwang, and Tohru Ozawa. On the Cauchy problem of fractional Schrödinger equation with Hartree type nonlinearity. *Funkcial. Ekvac.*, 56(2):193–224, 2013.
- <span id="page-17-3"></span>[CHKL14] Y Cho, G Hwang, S. Kwon, and S. Lee. Well-posedness and ill-posedness for the cubic fractional schrödinger equations.  $http://arxiv.org/abs/1311.0082, 2014$ .
- <span id="page-17-6"></span>[CKS<sup>+</sup>08] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in  $\mathbb{R}^3$ . Ann. of Math. *(2)*, 167(3):767–865, 2008.
- <span id="page-17-5"></span>[COX11] Yonggeun Cho, Tohru Ozawa, and Suxia Xia. Remarks on some dispersive estimates. *Commun. Pure Appl. Anal.*, 10(4):1121–1128, 2011.
- <span id="page-17-7"></span>[CW91] F. M. Christ and M. I. Weinstein. Dispersion of small amplitude solutions of the generalized Korteweg-de Vries equation. *J. Funct. Anal.*, 100(1):87–109, 1991.
- <span id="page-18-2"></span>[GSWZ13] Z Guo, Y. Sire, Y Wang, and L. Zhao. On the energy-critical fractional schrödinger equation in the radial case. *Preprint*, 2013.
- <span id="page-18-5"></span>[GW14] Z Guo and Y. Wang. Improved strichartz estimates for a class of dispersive equations in the radial case and their applications to nonlinear schrodinger and wave equation. *to appear J. Anal. Math.*, 2014.
- <span id="page-18-6"></span>[KLR13] Joachim Krieger, Enno Lenzmann, and Pierre Raphaël. Nondispersive solutions to the L 2 -critical half-wave equation. *Arch. Ration. Mech. Anal.*, 209(1):61–129, 2013.
- <span id="page-18-4"></span>[KM06] Carlos E. Kenig and Frank Merle. Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case. *Invent. Math.*, 166(3):645–675, 2006.
- <span id="page-18-3"></span>[KM08] Carlos E. Kenig and Frank Merle. Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation. *Acta Math.*, 201(2):147–212, 2008.
- <span id="page-18-7"></span>[KT98] Markus Keel and Terence Tao. Endpoint Strichartz estimates. *Amer. J. Math.*, 120(5):955–980, 1998.
- <span id="page-18-1"></span>[Lan72] N. S. Landkof. *Foundations of modern potential theory*. Springer-Verlag, New York, 1972. Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- <span id="page-18-0"></span>[Las02] Nick Laskin. Fractional Schrödinger equation. *Phys. Rev. E* (3), 66(5):056108, 7, 2002.

University of Texas, Austin

Universit´e Aix-Marseille, I2M, UMR 7353,Marseille, France