GROUND STATES OF CRITICAL AND SUPERCRITICAL PROBLEMS OF BREZIS-NIRENBERG TYPE

MONICA CLAPP, ANGELA PISTOIA, AND ANDRZEJ SZULKIN ´

Abstract. We study the existence of symmetric ground states to the supercritical problem

$$
-\Delta v = \lambda v + |v|^{p-2} v \text{ in } \Omega, \qquad v = 0 \text{ on } \partial\Omega,
$$

in a domain of the form

 $\Omega = \{ (y, z) \in \mathbb{R}^{k+1} \times \mathbb{R}^{N-k-1} : (|y|, z) \in \Theta \},$

where Θ is a bounded smooth domain such that $\overline{\Theta} \subset (0,\infty) \times \mathbb{R}^{N-k-1}$, $1 \leq$ $k \leq N-3$, $\lambda \in \mathbb{R}$, and $p = \frac{2(N-k)}{N-k-2}$ is the $(k+1)$ -st critical exponent. We show that symmetric ground states exist for λ in some interval to the left of each symmetric eigenvalue, and that no symmetric ground states exist in some interval $(-\infty, \lambda_*)$ with $\lambda_* > 0$ if $k \geq 2$.

Related to this question is the existence of ground states to the anisotropic critical problem

 $-\text{div}(a(x)\nabla u) = \lambda b(x)u + c(x)|u|^{2^*-2}u$ in Θ , $u = 0$ on $\partial\Theta$,

where a, b, c are positive continuous functions on $\overline{\Theta}$. We give a minimax characterization for the ground states of this problem, study the ground state energy level as a function of λ , and obtain a bifurcation result for ground states.

KEY WORDS: Supercritical elliptic problem, anisotropic critical problem, ground states, bifurcation.

2010 MSC: 35J61 (35J20, 35J25).

1. INTRODUCTION

We consider the supercritical Brezis-Nirenberg type problem

$$
(\wp_{\lambda}) \qquad \qquad \left\{ \begin{array}{r} -\Delta v = \lambda v + |v|^{2_{N,k}^* - 2} v \quad \text{in } \Omega, \\ v = 0 \qquad \text{on } \partial \Omega, \end{array} \right.
$$

where Ω is given by

(1.1)
$$
\Omega := \{(y, z) \in \mathbb{R}^{k+1} \times \mathbb{R}^{N-k-1} : (|y|, z) \in \Theta\}
$$

for some bounded smooth domain Θ in \mathbb{R}^{N-k} such that $\overline{\Theta} \subset (0,\infty) \times \mathbb{R}^{N-k-1}$, $1 \leq k \leq N-3, \ \lambda \in \mathbb{R}, \text{ and } 2^*_{N,k} := \frac{2(N-k)}{N-k-2}$ $\frac{2(N-k)}{N-k-2}$ is the so-called $(k+1)$ -st critical exponent.

If $k = 0$ then $2_{N,0}^* = 2^*$ is the critical Sobolev exponent and problem (φ_λ) becomes

(1.2)
$$
\begin{cases}\n-\Delta v = \lambda v + |v|^{2^*-2} v & \text{in } \Theta, \\
v = 0 & \text{on } \partial \Theta.\n\end{cases}
$$

Date: October 15, 2018.

M. Clapp is partially supported by CONACYT grant 237661 and PAPIIT-DGAPA-UNAM grant IN104315 (Mexico), A. Pistoia is partially supported by PRIN 2009-WRJ3W7 grant (Italy).

A celebrated result by Brezis and Nirenberg [\[1\]](#page-13-0) states that [\(1.2\)](#page-0-1) has a ground state $v > 0$ if and only if $\lambda \in (0, \lambda_1)$ and $N \geq 4$, or if $\lambda \in (\lambda_*, \lambda_1)$ and $N = 3$, where λ_* is some number in $(0, \lambda_1)$. Moreover, they show that $\lambda_* = \frac{\lambda_1}{4} > 0$ if Θ is a ball. As usual, λ_m denotes the m-th Dirichlet eigenvalue of $-\Delta$ in Θ .

Problem [\(1.2\)](#page-0-1) has been widely investigated. Capozzi, Fortunato and Palmieri [\[2\]](#page-13-1) established the existence of solutions for all $\lambda > 0$ if $N \geq 5$ and for all $\lambda \neq \lambda_m$ if $N = 4$ (see also [\[11,](#page-14-0) [24\]](#page-14-1)). Several multiplicity results are also available, see e.g. [\[3,](#page-13-2) [7,](#page-14-2) [8,](#page-14-3) [9,](#page-14-4) [25\]](#page-14-5) and the references therein.

Recently, Szulkin, Weth and Willem [\[22\]](#page-14-6) gave a minimax characterization for the ground states of problem [\(1.2\)](#page-0-1) when $\lambda \geq \lambda_1$. They established the existence of ground states for $\lambda \neq \lambda_m$ if $N = 4$ and for all $\lambda \geq \lambda_1$ if $N \geq 5$.

Concerning the supercritical problem (φ_{λ}) with $k \geq 1$, Passaseo [\[16,](#page-14-7) [17\]](#page-14-8) showed that a nontrivial solution does not exist if $\lambda = 0$ and Θ is a ball. This statement was extended in [\[5\]](#page-14-9) to more general domains Θ, and to some unbounded domains in [\[6\]](#page-14-10). On the other hand, existence of multiple solutions has been established in [\[4,](#page-14-11) [14,](#page-14-12) [23\]](#page-14-13).

This work is concerned with the existence of symmetric ground states for the supercritical problem (\wp_{λ}) with $k \geq 1$. Note that the domain Ω is invariant under the action of the group $O(k+1)$ of linear isometries of \mathbb{R}^{k+1} on the first $k+1$ coordinates. A function $v : \Omega \to \mathbb{R}$ is called $O(k+1)$ -invariant if $v(gy, z) = v(y, z)$ for every $g \in O(k+1)$, $(y, z) \in \mathbb{R}^{k+1} \times \mathbb{R}^{N-k-1}$. The subspace

$$
H_0^1(\Omega)^{O(k+1)} := \{ v \in H_0^1(\Omega) : v \text{ is } O(k+1)\text{-invariant} \}
$$

of $H_0^1(\Omega)$ is continuously embedded in $L^{2_{N,k}^*}(\Omega)$, so the energy functional \mathcal{J}_λ : $H_0^1(\Omega)^{O(k+1)} \to \mathbb{R}$ given by

$$
\mathcal{J}_{\lambda}(v):=\frac{1}{2}\int_{\Omega}\left|\nabla v\right|^{2}-\frac{\lambda}{2}\int_{\Omega}v^{2}-\frac{1}{2^{*}}\int_{\Omega}\left|v\right|^{2^{*}_{N,k}}
$$

is well defined. Its critical points are the $O(k+1)$ -invariant solutions to problem (φ_{λ}) . An $O(k+1)$ -invariant $(PS)_{\tau}$ -sequence for \mathcal{J}_{λ} is a sequence (v_k) such that

$$
v_k \in H_0^1(\Omega)^{O(k+1)}, \qquad \mathcal{J}_\lambda(v_k) \to \tau \qquad \text{and} \qquad \mathcal{J}'_\lambda(v_k) \to 0 \text{ in } H^{-1}(\Omega).
$$

We set

 $\ell_{\lambda}^{O(k+1)}$ $Q_{\lambda}^{(k+1)} := \inf \{ \tau > 0 : \text{ there exists an } O(k+1) \text{-invariant } (PS)_{\tau} \text{-sequence for } \mathcal{J}_{\lambda} \}.$

This is the lowest possible energy level for a nontrivial $O(k+1)$ -invariant solution to problem (φ_{λ}) . An $O(k+1)$ *-invariant ground state of problem* (φ_{λ}) is a critical point $v \in H_0^1(\Omega)^{O(k+1)}$ of \mathcal{J}_λ such that $\mathcal{J}_\lambda(v) = \ell^{O(k+1)}_\lambda$ $\lambda^{\mathcal{O}(\kappa+1)}$. Since \mathcal{J}_{λ} does not satisfy the Palais-Smale condition, an $O(k+1)$ -invariant ground state does not necessarily exist.

Let $0 < \lambda_1^{[k]} < \lambda_2^{[k]} \leq \lambda_3^{[k]} \leq \cdots$ be the $O(k+1)$ -invariant eigenvalues of the problem

$$
-\Delta v = \lambda v \quad \text{in } \Omega, \qquad v \in H_0^1(\Omega)^{O(k+1)},
$$

counted with their multiplicity. Set $\lambda_0^{[k]} := 0$. We shall prove the following result for $O(k + 1)$ -invariant ground states.

Theorem 1.1. For every $1 \leq k \leq N-3$, the following statements hold true:

(a) *Problem* (\wp_{λ}) *does not have an* $O(k+1)$ *-invariant ground state if* $\lambda \leq 0$ *.*

- (b) *For each* $m \in \mathbb{N} \cup \{0\}$, there is a number $\lambda_{m,*}^{[k]} \in [\lambda_m^{[k]}, \lambda_{m+1}^{[k]})$ with the *property that problem* (\wp_{λ}) *has an* $O(k+1)$ *-invariant ground state for every* $\lambda \in (\lambda^{[k]}_{m,*},\lambda^{[k]}_{m+1})$ and does not have an $O(k+1)$ -invariant ground state for $any \lambda \in [\lambda_m^{[k]}, \lambda_{m,*}^{[k]}).$
- (c) Let $\beta := \max\{\text{dist}(x, \{0\} \times \mathbb{R}^{N-k-1}) : x \in \overline{\Theta}\}.$ Then,

$$
\lambda_{0,*}^{[k]} \geq \left\{ \begin{array}{ll} \frac{(k-1)^2}{4\beta^2} & \text{if } 3k \geq N, \\ \frac{k}{\left(2^*_{N,k}\beta\right)^2}\left((2^*_{N,k}-1)k-2^*_{N,k}\right) & \text{if } 3k \leq N. \end{array} \right.
$$

In particular, $\lambda_{0,*}^{[k]} > 0$ *if* $k \geq 2$.

This last statement stands in contrast with the case $k = 0$ where a ground state to problem [\(1.2\)](#page-0-1) exists for every $\lambda \in [0, \lambda_1)$ if $N \geq 4$. We also show that $\lambda_{0,*}^{[1]} > 0$ if Θ is thin enough, see Proposition [4.4.](#page-12-0)

As we shall see, the $O(k+1)$ -invariant ground states of problem (φ_{λ}) correspond to the ground states of the critical problem

(1.3)
$$
-\operatorname{div}(a(x)\nabla u) = \lambda b(x)u + c(x)|u|^{2^*-2}u \text{ in } \Theta, \quad u = 0 \text{ on } \partial \Theta,
$$

with $2^* = \frac{2n}{n-2}$, $n := \dim \Theta$, $a(x_1, ..., x_n) = x_1^k$ and $a = b = c$.

The critical problem [\(1.3\)](#page-2-0) with general coefficients $a \in C^1(\overline{\Theta}), b, c \in C^0(\overline{\Theta})$ has an interest in its own. We study it in section [2](#page-2-1) and give a minimax characterization for its ground states, similar to that in [\[22\]](#page-14-6). We study the properties of its ground state energy level as a function of λ , and obtain a bifurcation result for ground states, see Theorem [2.1.](#page-4-0)

Anisotropic critical problems of the form [\(1.3\)](#page-2-0) have been studied, for example, by Egnell [\[10\]](#page-14-14) and, more recently, by Hadiji et al. [\[12,](#page-14-15) [13\]](#page-14-16). They obtained existence and multiplicity results under some assumptions which involve flatness of the coefficient functions at some local maximum or minimum point in the interior of Θ. Note that the function $a(x_1,...,x_n) = x_1^k$ attains its minimum on the boundary of Θ . This produces a quite different behavior regarding the existence of ground states, as we shall see in the following sections.

Section [2](#page-2-1) is devoted to the study of the general anisotropic critical problem. In section [3](#page-8-0) we prove a nonexistence result for supercritical problems. It will be used in Section [4](#page-10-0) where we prove Theorem [1.1.](#page-1-0) In the last section we include some questions and remarks.

2. Ground states of the anisotropic critical problem

In this section we consider the anisotropic Brezis-Nirenberg type problem

(2.1)
$$
\begin{cases}\n-\operatorname{div}(a(x)\nabla u) = \lambda b(x)u + c(x)|u|^{2^*-2}u & \text{in } \Theta, \\
u = 0 & \text{on } \partial\Theta,\n\end{cases}
$$

where Θ is a bounded smooth domain in \mathbb{R}^n , $n \geq 3$, $\lambda \in \mathbb{R}$, $a \in \mathcal{C}^1(\overline{\Theta})$, $b, c \in$ $\mathcal{C}^0(\overline{\Theta})$ are strictly positive on $\overline{\Theta}$, and $2^* := \frac{2n}{n-2}$ is the critical Sobolev exponent in dimension n .

We take

(2.2)
$$
\langle u, v \rangle_a := \int_{\Theta} a(x) \nabla u \cdot \nabla v, \qquad ||u||_a := \left(\int_{\Theta} a(x) |\nabla u|^2 \right)^{1/2},
$$

to be the scalar product and the norm in $H_0^1(\Theta)$, and

$$
|u|_{b,2}:=\left(\int_{\Theta}b(x)u^{2}\right)^{1/2},\qquad |u|_{c,2^{*}}:=\left(\int_{\Theta}c(x)\left|u\right|^{2^{*}}\right)^{1/2^{*}},
$$

to be the norms in $L^2(\Theta)$ and $L^{2^*}(\Theta)$ respectively. They are, clearly, equivalent to the standard ones.

Let $0 < \lambda_1^{a,b} < \lambda_2^{a,b} \leq \lambda_3^{a,b} \leq \cdots$ be the eigenvalues of the problem

$$
-\text{div}(a(x)\nabla u) = \lambda b(x)u \quad \text{in } \Theta, \qquad u = 0 \quad \text{on } \partial \Theta,
$$

counted with their multiplicity, and e_1, e_2, e_3, \dots be the corresponding normalized eigenfunctions, i.e. $|e_j|_{b,2} = 1$. Set

$$
Z_0 := \{0\}, \qquad Z_m := \text{span}\{e_1, ..., e_m\},
$$

\n
$$
Y_m := \{w \in H_0^1(\Theta) : \langle w, z \rangle_a = 0 \text{ for all } z \in Z_m\},
$$

\n
$$
T_0 := (-\infty, \lambda_1^{a,b}), \quad \text{and} \quad T_m := [\lambda_m^{a,b}, \lambda_{m+1}^{a,b}) \text{ if } m \in \mathbb{N}.
$$

The solutions to problem [\(2.1\)](#page-2-2) are the critical points of the functional J_{λ} : $H_0^1(\Theta) \to \mathbb{R}$ given by

$$
J_{\lambda}(u) := \frac{1}{2} ||u||_a^2 - \frac{\lambda}{2} ||u||_{b,2}^2 - \frac{1}{2^*} ||u||_{c,2^*}^2.
$$

If $\lambda \in T_m$ we define

 $\mathcal{N}_{\lambda} \equiv \mathcal{N}_{\lambda}(\Theta) := \{u \in H_0^1(\Theta) \setminus Z_m : J'_{\lambda}(u)u = 0 \text{ and } J'_{\lambda}(u)z = 0 \text{ for all } z \in Z_m\}.$ This is a \mathcal{C}^1 -submanifold of codimension $m+1$ in $H_0^1(\Theta)$, cf. [\[22\]](#page-14-6). If $\lambda < \lambda_1^{a,b}$ it $_{0}$ (O), ci. [22]. ii $\lambda \leq \lambda_1$ is the usual Nehari manifold, and if $\lambda \geq \lambda_1^{a,b}$ it is the generalized Nehari manifold, introduced by Pankov in [\[15\]](#page-14-17) and studied by Szulkin and Weth in [\[20,](#page-14-18) [21\]](#page-14-19). Note that $J'_{\lambda}(z)z < 0$ for all $z \in Z_m \setminus \{0\}$. Clearly, the nontrivial critical points of J_{λ} belong to \mathcal{N}_{λ} . Moreover, they coincide with the critical points of its restriction $J_{\lambda}|_{\mathcal{N}_{\lambda}} : \mathcal{N}_{\lambda} \to \mathbb{R}$. The proof of these facts is completely analogous to the one given in [\[22\]](#page-14-6) for the autonomous case. Set

$$
\ell_{\lambda} \equiv \ell_{\lambda}^{a,b,c} := \inf_{\mathcal{N}_{\lambda}} J_{\lambda}.
$$

Following [\[20\]](#page-14-18) one shows that, for every $w \in Y_m \setminus \{0\}$, there exist unique $t_{\lambda,w} \in$ $(0, \infty)$ and $z_{\lambda,w} \in Z_m$ such that

$$
t_{\lambda,w}w+z_{\lambda,w}\in\mathcal{N}_{\lambda},
$$

and that

$$
J_\lambda(t_{\lambda,w}w+z_{\lambda,w})=\max_{t>0,\,z\in Z_m}J_\lambda(tw+z).
$$

Let $\Sigma_m := \{w \in Y_m : ||w||_a = 1\}$ be the unit sphere in Y_m . Then,

(2.3)
$$
\ell_{\lambda} = \inf_{w \in \Sigma_m} \max_{\substack{t > 0, \\ z \in Z_m}} J_{\lambda}(tw + z).
$$

As usual, we denote the best Sobolev constant for the embedding $H^1(\mathbb{R}^n) \hookrightarrow$ $L^{2^*}(\mathbb{R}^n)$ by S. We set

$$
\kappa^{a,c} := \left(\min_{x \in \overline{\Theta}} \frac{a(x)^{\frac{n}{2}}}{c(x)^{\frac{n-2}{2}}}\right) \frac{1}{n} S^{\frac{n}{2}},
$$

and define

$$
\lambda_{m,*}^{a,b,c} := \inf \{ \lambda \in T_m : \ell_\lambda < \kappa^{a,c} \}.
$$

Theorem 2.1. *For every* $m \in \mathbb{N} \cup \{0\}$ *the following statements hold true:*

(a) *The function* $\lambda \mapsto \ell_{\lambda}$ *is nonincreasing in* T_m *and*

$$
0 < \ell_{\lambda} \le \kappa^{a,c} \qquad \text{for all} \ \ \lambda \in T_m.
$$

- (b) ℓ_{λ} *is attained on* \mathcal{N}_{λ} *if* $\ell_{\lambda} < \kappa^{a,c}$.
- (c) *The function* $\lambda \mapsto \ell_{\lambda}$ *is continuous in* T_m *and*

$$
\lim_{\lambda \nearrow \lambda_{m+1}^{a,b}} \ell_{\lambda} = 0.
$$

Hence, $\lambda_{m,*}^{a,b,c} < \lambda_{m+1}^{a,b}$.

(d) ℓ_{λ} *is not attained if* $\lambda \in (-\infty, \lambda_{0,*}^{a,b,c})$ *or* $\lambda \in [\lambda_m^{a,b}, \lambda_{m,*}^{a,b,c}), m \ge 1$, *and is attained if* $\lambda \in (\lambda^{a,b,c}_{m,*}, \lambda^{a,b}_{m+1}).$

Remark 2.2. It follows from part (c) above that bifurcation (to the left) occurs at each $\lambda_m^{a,b}$. This fact is essentially known and can be obtained by other methods. However, we would like to emphasize that here we show that our bifurcating solutions are *ground states*.

Proof of Theorem 2.1. (a): Let $\lambda, \mu \in T_m$. If $\lambda \leq \mu$ then $J_{\lambda}(u) \geq J_{\mu}(u)$ for every $u \in H_0^1(\Theta)$. So $\ell_{\lambda} \geq \ell_{\mu}$ according to [\(2.3\)](#page-3-0). This proves that $\lambda \mapsto \ell_{\lambda}$ is nonincreasing in T_m .

If $\lambda \in T_m$ and $w \in \Sigma_m$ we have that

$$
(2.4) \qquad \max_{t>0, z \in Z_m} J_{\lambda}(tw+z) \ge \max_{t>0} J_{\lambda}(tw) = \frac{1}{n} \left(\frac{\|w\|_{a}^{2} - \lambda |w|_{b,2}^{2}}{|w|_{c,2^{*}}^{2}} \right)^{n/2}
$$

$$
\ge \frac{1}{n} \left(\frac{1 - \frac{\lambda}{\lambda_{m+1}}}{|w|_{c,2^{*}}^{2}} \right)^{n/2}.
$$

Using Sobolev's inequality we conclude that there is a positive constant C such that

$$
\max_{t>0, z \in Z_m} J_{\lambda}(tw + z) \ge C \quad \text{for all } w \in \Sigma_m.
$$

Therefore, $\ell_{\lambda} > 0$.

Let $\varphi_k \in C_c^{\infty}(\mathbb{R}^n)$ be a positive function such that $\text{supp}(\varphi_k) \subset B_{1/k}(0)$ and $\int |\nabla \varphi_k|^2 \to S^{n/2}, \int |\varphi_k|^{2^*} \to S^{n/2},$ where $B_r(\xi) = \{x \in \mathbb{R}^n : |x - \xi| < r\}.$ Let $\xi \in \overline{\Theta}$ be such that

$$
\frac{a(\xi)^{\frac{n}{2}}}{c(\xi)^{\frac{n-2}{2}}} = \min_{x \in \overline{\Theta}} \frac{a(x)^{\frac{n}{2}}}{c(x)^{\frac{n-2}{2}}}
$$

and choose $\nu \in \mathbb{R}^n$ with $|\nu|=1$ such that ν is the inward pointing unit normal at ξ if $\xi \in \partial \Theta$. Set $\xi_k := \xi + \frac{1}{k} \nu$ and $u_k(x) := \varphi_k(x - \xi_k)$. Then $u_k \in H_0^1(\Theta)$ for k large enough, and we have that

$$
(2.5) \qquad \max_{t>0} J_{\lambda}(tu_{k}) = \frac{1}{n} \left(\frac{\|u_{k}\|_{a}^{2} - \lambda |u_{k}|_{b,2}^{2}}{|u_{k}|_{c,2^{*}}^{2}} \right)^{\frac{n}{2}}
$$

$$
= \frac{1}{n} \left(\frac{\int_{B_{1/k}(\xi_{k})} a(x) |\nabla u_{k}|^{2} - \lambda \int_{B_{1/k}(\xi_{k})} b(x) u_{k}^{2}}{\left(\int_{B_{1/k}(\xi_{k})} c(x) |u_{k}|^{2^{*}}\right)^{2/2^{*}}} \right)^{\frac{n}{2}}
$$

$$
\longrightarrow \frac{1}{n} \left(\frac{a(\xi)^{\frac{n}{2}}}{c(\xi)^{\frac{n-2}{2}}} \right) S^{\frac{n}{2}} = \kappa^{a,c} \qquad \text{as } k \to \infty.
$$

Hence, $\ell_{\lambda} \leq \kappa^{a,c}$ for $\lambda < \lambda_1^{a,b}$.

Next, we assume that $\lambda \in T_m$ with $m \in \mathbb{N}$. We fix an open subset θ of Θ such that $\theta \cap B_{1/k}(\xi_k) = \emptyset$ for k large enough. If $z \in Z_m$ and $z = 0$ in θ then $z = 0$ in Θ , see [\[22,](#page-14-6) Lemma 3.3]. Hence, $(\int_{\theta} c(x)|z|^{2^*})^{1/2^*}$ is a norm in Z_m and, since Z_m is finite-dimensional, this norm is equivalent to $||z||_a$. In particular, there is a positive constant A such that $\int_{\theta} c(x) |z|^{2^*} \geq 2^* A \|z\|_{a}^{2^*}$ $\frac{2}{a}$ for all $z \in Z_m$. It follows by convexity that, for every $t > 0$ and every $z \in Z_m$, we have

$$
|tu_{k} + z|_{c,2^{*}}^{2^{*}} = \int_{\Theta \setminus \theta} c(x) |tu_{k} + z|^{2^{*}} + \int_{\theta} c(x) |z|^{2^{*}}
$$

$$
\geq t^{2^{*}} \int_{\Theta} c(x) u_{k}^{2^{*}} + 2^{*} t^{2^{*}-1} \int_{\Theta} c(x) u_{k}^{2^{*}-1} z + 2^{*} A ||z||_{a}^{2^{*}}.
$$

Therefore,

$$
(2.6) \qquad J_{\lambda}(tu_{k} + z) \leq J_{0}(tu_{k}) - \frac{\lambda}{2} |tu_{k}|_{b,2}^{2} + t \int_{\Theta} (a(x)\nabla u_{k}\nabla z - \lambda b(x)u_{k}z) + \frac{1}{2} (||z||_{a}^{2} - \lambda |z|_{b,2}^{2}) - t^{2^{*}-1} \int_{\Theta} c(x)u_{k}^{2^{*}-1}z - A ||z||_{a}^{2^{*}} \leq J_{0}(tu_{k}) + t \int_{\Theta} (a(x)\nabla u_{k}\nabla z - \lambda b(x)u_{k}z) - t^{2^{*}-1} \int_{\Theta} c(x)u_{k}^{2^{*}-1}z - A ||z||_{a}^{2^{*}}.
$$

Consequently,

$$
J_{\lambda}(tu_{k}+z) \leq B(t^{2}+t\left\|z\right\|_{a}+t^{2^{*}-1}\left\|z\right\|_{a})-C(t^{2^{*}}+\left\|z\right\|_{a}^{2^{*}})
$$

for some positive constants B and C. This implies that there exists $R > 0$ such that $J_{\lambda}(tu_k + z) \leq 0$ for all $t \geq R$, $z \in Z_m$ and k large enough. On the other hand, for $t \leq R$, $z \in Z_m$ and k large enough, since $\varphi_k \to 0$ weakly in $H_0^1(\Theta)$, inequalities (2.6) and (2.5) imply that

$$
J_{\lambda}(tu_{k}+z) \leq J_{0}(tu_{k}) + o(1) = \kappa^{a,c} + o(1).
$$

This proves that $\ell_{\lambda} \leq \kappa^{a,c}$ for $\lambda \geq \lambda_1^{a,b}$ and concludes the proof of statement (a).

(b): Let $I_{\lambda} : \Sigma_m \to \mathbb{R}$ be the function given by

$$
I_{\lambda}(w) := J_{\lambda}(t_{\lambda,w}w + z_{\lambda,w}).
$$

Then $\ell_{\lambda} := \inf_{w \in \Sigma_m} I_{\lambda}(w)$. It is shown in [\[20,](#page-14-18) [21\]](#page-14-19) that $I_{\lambda} \in C^1(\Sigma_m, \mathbb{R})$. Since Σ_m is a smooth submanifold of $H_0^1(\Theta)$, Ekeland's variational principle yields a Palais-Smale sequence (w_k) for I_λ such that $I_\lambda(w_k) \to \ell_\lambda$, cf. [\[24,](#page-14-1) Theorem 8.5]. Set $u_k := t_{\lambda,w_k} w_k + z_{\lambda,w_k}$. By Corollary 2.10 in [\[20\]](#page-14-18) or Corollary 33 in [\[21\]](#page-14-19), (u_k) is a Palais-Smale sequence for J_{λ} . Now, Corollary 3.2 in [\[4\]](#page-14-11) asserts that every Palais-Smale sequence (u_k) for J_λ such that $J_\lambda(u_k) \to \tau \lt \kappa^{a,c}$, contains a convergent subsequence. It follows that ℓ_{λ} is attained on \mathcal{N}_{λ} if $\ell_{\lambda} < \kappa^{a,c}$.

(c): Let $w \in \Sigma_m$. First, we will show that the function $\lambda \mapsto I_\lambda(w)$ is continuous in T_m . Let $\mu_j, \mu \in T_m$ be such that $\mu_j \to \mu$. A standard argument shows that $J_{\mu_j}(tw+z) \leq 0$ for every $j \in \mathbb{N}$ if $t^2 + ||z||_a^2$ is large enough. Therefore, the sequences $(t_{\mu_i,w})$ and $(z_{\mu_i,w})$ are bounded and, after passing to a subsequence, $t_{\mu_i,w} \to t_0$ in $[0, \infty)$ and $z_{\mu_i, w} \to z_0$ in Z_m . Hence,

$$
I_{\mu_j}(w) = J_{\mu_j}(t_{\mu_j,w}w + z_{\mu_j,w}) \to J_{\mu}(t_0w + z_0) \le I_{\mu}(w).
$$

If $J_{\mu}(t_0w+z_0) < I_{\mu}(w)$ then, since

$$
J_{\mu_j}(t_{\mu,w}w + z_{\mu,w}) \to J_{\mu}(t_{\mu,w}w + z_{\mu,w}) = I_{\mu}(w),
$$

we would have that, for j large enough,

$$
\max_{t>0, z \in Z_m} J_{\mu_j}(tw+z) = J_{\mu_j}(t_{\mu_j, w}w + z_{\mu_j, w}) < J_{\mu_j}(t_{\mu, w}w + z_{\mu, w}),
$$

which is a contradiction. Consequently, $I_{\mu_j}(w) \to I_{\mu}(w)$. This proves that $\lambda \mapsto$ $I_{\lambda}(w)$ is continuous in T_m for each $w \in \Sigma_m$.

Next, we prove that the function $\lambda \mapsto \ell_{\lambda}$ is continuous from the left in T_m . Let $\mu_i, \mu \in T_m$ be such that $\mu_i \leq \mu$ and $\mu_i \to \mu$. Since the infimum of any family of continuous functions is upper semicontinuous and $\lambda \mapsto \ell_{\lambda}$ is nonincreasing, we have that

$$
\limsup_{j \to \infty} \ell_{\mu_j} \le \ell_{\mu} \le \liminf_{j \to \infty} \ell_{\mu_j}.
$$

This proves that $\lambda \mapsto \ell_{\lambda}$ is continuous from the left in T_m .

To prove that $\lambda \mapsto \ell_{\lambda}$ is continuous from the right in T_m we argue by contradiction. Assume there are $\mu_j, \mu \in T_m$ such that $\mu_j \geq \mu$, $\mu_j \to \mu$ and $\sup_{j \in \mathbb{N}} \ell_{\mu_j} < \ell_{\mu}$. Then $\ell_{\mu_i} < \kappa^{a,c}$ and, by statement (b), there exists $w_j \in \Sigma_m$ such that $\ell_{\mu_i} =$ $J_{\mu_j}(t_{\mu_j,w}w_j+z_{\mu_j,w})$. Inequality [\(2.4\)](#page-4-1) asserts that

$$
\ell_{\mu} > \ell_{\mu_j} = J_{\mu_j}(t_{\mu_j, w_j}w_j + z_{\mu_j, w_j}) \geq \frac{1}{n} \left(\frac{1 - \frac{\mu_j}{\lambda_{m+1}}}{|w_j|_{c, 2^*}^2} \right)^{n/2}.
$$

This implies that $|w_j|_{c,i}^{2^*}$ $c_{c,2^*}^{2^*} \geq \varepsilon > 0$ for all $j \in \mathbb{N}$. Denote the closure of Y_m in $L^{2^*}(\Theta)$ by \widetilde{Y}_m . Since $\dim(Z_m) < \infty$, the projection $\widetilde{Y}_m \oplus Z_m \to \widetilde{Y}_m$ is continuous in $L^{2^*}(\Theta)$. Hence, there is a positive constant A_0 such that

$$
\ell_{\mu} \leq J_{\mu}(t_{\mu,w_j}w_j + z_{\mu,w_j})
$$
\n
$$
= \frac{t_{\mu,w_j}^2}{2}(1 - \mu |w_j|_{b,2}^2) + \frac{1}{2}(\left\|z_{\mu,w_j}\right\|_a^2 - \mu |z_{\mu,w_j}|_{b,2}^2) - \frac{1}{2^*}|t_{\mu,w_j}w_j + z_{\mu,w_j}|_{c,2^*}^{2^*}
$$
\n
$$
\leq \frac{t_{\mu,w_j}^2}{2} - A_0 \frac{t_{\mu,w_j}^2}{2^*}|w_j|_{c,2^*}^{2^*} \leq \frac{t_{\mu,w_j}^2}{2} - A_0 \varepsilon \frac{t_{\mu,w_j}^2}{2^*} \qquad \text{for all } j \in \mathbb{N}.
$$

It follows that (t_{μ,w_j}) is bounded. Hence, $(\|z_{\mu,w_j}\|_a)$ is bounded too. Consequently,

$$
\ell_{\mu} \leq J_{\mu}(t_{\mu,w_j}w_j + z_{\mu,w_j}) = J_{\mu_j}(t_{\mu,w_j}w_j + z_{\mu,w_j}) + (\mu - \mu_j) |t_{\mu,w_j}w_j + z_{\mu,w_j}|_{b,2}^2
$$

\n
$$
\leq J_{\mu_j}(t_{\mu_j,w_j}w_j + z_{\mu_j,w_j}) + o(1) = \ell_{\mu_j} + o(1) \leq \sup_{j \in \mathbb{N}} \ell_{\mu_j} + o(1) < \ell_{\mu} + o(1).
$$

This is a contradiction. It follows that the function $\lambda \mapsto \ell_{\lambda}$ is continuous in T_m . Finally, let $\mu_j \in T_m$ be such that $\mu_j \to \lambda_{m+1}$. We have that

$$
0 < \ell_{\mu_j} \leq J_{\mu_j}(t_{\mu_j, e_{m+1}} e_{m+1} + z_{\mu_j, e_{m+1}})
$$

= $\frac{t_{\mu_j, e_{m+1}}^2}{2} (\lambda_{m+1} - \mu_j) + \frac{1}{2} (\left\| z_{\mu_j, e_{m+1}} \right\|_a^2 - \mu_j \left| z_{\mu_j, e_{m+1}} \right\|_{b, 2}^2)$
 $- \frac{1}{2^*} \left| t_{\mu_j, e_{m+1}} e_{m+1} + z_{\mu_j, e_{m+1}} \right|_{c, 2^*}^{2^*}$
 $\leq \frac{t_{\mu_j, e_{m+1}}^2}{2} (\lambda_{m+1} - \mu_j) - A_0 \frac{t_{\mu_j, e_{m+1}}^{2^*}}{2^*} \left| e_{m+1} \right|_{c, 2^*}^{2^*}.$

It follows that $(t_{\mu_j, e_{m+1}})$ is bounded and, hence, that

$$
0 < \ell_{\mu_j} \le \frac{t_{\mu_j, e_{m+1}}^2}{2} (\lambda_{m+1} - \mu_j) = o(1).
$$

This proves that $\ell_{\mu_j} \to 0$ as $\mu_j \to \lambda_{m+1}$ from the left.

(d): If $\lambda \in T_m$, $\lambda \leq \lambda_{m,*}^{a,b,c}$, and $w \in \Sigma_m$ were such that $\ell_{\lambda} = I_{\lambda}(w)$ then for $\mu \in (\lambda, \lambda^{a,b,c}_{m,*}(\Theta))$ we would have that

$$
\kappa^{a,c} = \ell_{\mu} \le I_{\mu}(w) < I_{\lambda}(w) = \ell_{\lambda},
$$

contradicting (a). It follows that ℓ_{λ} is not attained if $\lambda \in [\lambda_m^{a,b}, \lambda_{m,*}^{a,b,c})$. Statement (b) implies that ℓ_{λ} is attained if $\lambda \in (\lambda^{a,b,c}_{m,*}, \lambda^{a,b}_{m+1})$.

Recall that a $(PS)_{\tau}$ -sequence for J_{λ} is a sequence (u_k) in $H_0^1(\Theta)$ such that $J_{\lambda}(u_k) \to \tau$ and $J'_{\lambda}(u_k) \to 0$ in $H^{-1}(\Theta)$. The value ℓ_{λ} is characterized as follows.

Corollary 2.3. $\ell_{\lambda} = \inf \{ \tau > 0 : \text{there exists a } (PS)_{\tau} \text{-sequence for } J_{\lambda} \}.$

Proof. The argument given in the proof of statement (b) of Theorem [2.1](#page-4-0) shows that there exists a $(PS)_{\ell_{\lambda}}$ -sequence for J_{λ} . To prove that ℓ_{λ} is the smallest positive number with this property, we argue by contradiction. Assume that $\tau < \ell_{\lambda}$ and that there exists a $(PS)_{\tau}$ -sequence for J_{λ} . Then $\tau < \kappa^{a,c}$ and Corollary 3.2 in [\[4\]](#page-14-11) asserts that (u_k) contains a subsequence which converges to a critical point u of J_λ with $J_{\lambda}(u) = \tau$. If $\tau \neq 0$ then $u \in \mathcal{N}_{\lambda}$ and, hence, $\ell_{\lambda} \leq \tau$. This is a contradiction. \square

For the classical Brezis-Nirenberg problem [\(1.2\)](#page-0-1) (where $a = b = c \equiv 1$) with $n \geq 4$, it is known that $\lambda_{0,*}^{a,b,c} = 0$ and $\lambda_{m,*}^{a,b,c} = \lambda_m$, the m-th Dirichlet eigenvalue of $-\Delta$ in Θ , for all $m \in \mathbb{N}$. Moreover, $\ell_{\lambda} = \frac{1}{n} S^{\frac{n}{2}} = \kappa^{a,c}$ for every $\lambda \leq 0$, but $\ell_{\lambda} < \frac{1}{n} S^{\frac{n}{2}}$ for every $\lambda > 0$ if $n \geq 5$, see [\[1,](#page-13-0) [11,](#page-14-0) [22\]](#page-14-6).

As we shall see below, this is not true in general: For the problem $(\wp_{\lambda}^{\#})$ in Section [4](#page-10-0) which arises from the supercritical one, one has that $\lambda_{0,*}^{a,b,c} > 0$ in most cases, see Propositions [4.3](#page-11-0) and [4.4.](#page-12-0) A special feature of that problem is that the value $\kappa^{a,c}$ is attained on the boundary of Θ. A different situation was considered by Egnell [\[10\]](#page-14-14) and Hadiji and Yazidi [\[13\]](#page-14-16). They showed for example that, if a attains its minimum at an interior point x_0 of Θ , $b = 1 = c$, and a is flat enough around x_0 , then $\lambda_{0,*}^{a,b,c} = 0$ for $n \geq 4$, as in the classical Brezis-Nirenberg case.

We do not know whether, in general, $\lambda_{0,*}^{a,b,c} \geq 0$. But this will be true in the special case we are interested in, see Proposition [4.1.](#page-10-2) The proof uses a nonexistence result for the supercritical problem, which we discuss in the following section.

3. Nonexistence of solutions to a supercritical problem

Let Θ be a bounded smooth domain in \mathbb{R}^{N-k} with $\overline{\Theta} \subset (0,\infty) \times \mathbb{R}^{N-k-1}$ and $0 \leq k \leq N-3$. Set

$$
\Omega := \{ (y, z) \in \mathbb{R}^{k+1} \times \mathbb{R}^{N-k-1} : (|y|, z) \in \Theta \}
$$

and consider the problem

(3.1)
$$
\begin{cases}\n-\Delta u = \lambda u + |u|^{p-2} u & \text{in } \Omega, \\
u = 0 & \text{on } \partial\Omega.\n\end{cases}
$$

Passaseo [\[16,](#page-14-7) [17\]](#page-14-8) showed that, if Θ is a ball, problem [\(3.1\)](#page-8-1) does not have a nontrivial solution for $\lambda = 0$ and $p \ge 2^*_{N,k} := \frac{2(N-k)}{N-k-2}$ $\frac{2(N-k)}{N-k-2}$. In [\[5\]](#page-14-9) it is shown that this is also true for doubly starshaped domains.

Definition 3.1. Θ *is doubly starshaped if there exist two numbers* $0 < t_0 < t_1$ *such that* $t \in (t_0, t_1)$ *for every* $(t, z) \in \Theta$ *and* Θ *is strictly starshaped with respect to* $\xi_0 := (t_0, 0)$ *and to* $\xi_1 := (t_1, 0)$ *, i.e.*

$$
\langle x - \xi_i, \nu_{\Theta}(x) \rangle > 0 \qquad \forall x \in \partial \Theta \setminus \{\xi_i\}, \quad i = 0, 1,
$$

where ν_{Θ} *is the outward pointing unit normal to* $\partial \Theta$ *.*

We denote the first Dirichlet eigenvalue of $-\Delta$ in Ω by $\lambda_1(\Omega)$.

Theorem 3.2. *If* Θ *is doubly starshaped,* $p \geq 2^*_{N,k}$ *and*

$$
\lambda \leq \frac{2(p-2^*_{N,k})}{2^*_{N,k}(p-2)}\lambda_1(\Omega),
$$

then problem [\(3.1\)](#page-8-1) *does not have a nontrivial solution.*

We point out that the geometric assumption on Θ cannot be dropped. Existence of multiple solutions to problem [\(3.1\)](#page-8-1) for $\lambda = 0$ and $p = 2^*_{N,k}$ in some domains where Θ is not doubly starshaped has been established in [\[4,](#page-14-11) [14,](#page-14-12) [23\]](#page-14-13).

The proof of Theorem [3.2](#page-8-2) follows the ideas introduced in [\[5,](#page-14-9) [16,](#page-14-7) [17\]](#page-14-8). Fix $\tau \in$ $(0, \infty)$ and let φ be the solution to the problem

$$
\left\{\begin{array}{ll} \varphi'(t)t+(k+1)\varphi(t)=1, & t\in (0,\infty),\\ \varphi(\tau)=0. & \end{array}\right.
$$

Explicitly, $\varphi(t) = \frac{1}{k+1} \left[1 - \left(\frac{\tau}{t}\right)^{k+1} \right]$. Note that φ is strictly increasing in $(0, \infty)$. For $y \neq 0$ we define

$$
\chi_{\tau}(y, z) := (\varphi(|y|)y, z).
$$

Lemma 3.3. *The vector field* χ_{τ} *has the following properties:*

- (a) div $\chi_{\tau} = N k$,
- (b) $\langle d\chi_{\tau}(y, z) [\xi], \xi \rangle \leq \max\{1 k\varphi(|y|), 1\} |\xi|^2$ *for every* $y \in \mathbb{R}^{k+1} \setminus \{0\},$ $z \in \mathbb{R}^{N-k-1}, \, \xi \in \mathbb{R}^N.$

Proof. See [\[17,](#page-14-8) Lemma 2.3] or [\[5,](#page-14-9) Lemma 4.2]. □

Proposition 3.4. *Assume there exists* $\tau \in (0,\infty)$ *such that* $|y| \in (\tau,\infty)$ *for every* $(y, z) \in \Omega$ *and* $\langle \chi_{\tau}, \nu_{\Omega} \rangle > 0$ *a.e. on* $\partial \Omega$ *. If* $p \geq 2^*_{N,k}$ *and*

$$
\lambda \leq \frac{2(p-2^*_{N,k})}{2^*_{N,k}(p-2)}\lambda_1(\Omega),
$$

then problem [\(3.1\)](#page-8-1) *does not have a nontrivial solution.*

Proof. The variational identity (4) in Pucci and Serrin's paper [\[19\]](#page-14-20) implies that, if $u \in C^2(\Omega) \cap C^1(\overline{\Omega})$ is a solution of (3.1) and $\chi \in C^1(\overline{\Omega}, \mathbb{R}^N)$, then

(3.3)
$$
\frac{1}{2} \int_{\partial \Omega} |\nabla u|^2 \langle \chi, \nu_{\Omega} \rangle d\sigma = \int_{\Omega} (div \chi) \left[\frac{1}{p} |u|^p + \frac{\lambda}{2} u^2 - \frac{1}{2} |\nabla u|^2 \right] dx + \int_{\Omega} \langle d\chi [\nabla u], \nabla u \rangle dx,
$$

where ν_{Ω} is the outward pointing unit normal to $\partial\Omega$ (in the notation of [\[19\]](#page-14-20) we have taken $\mathcal{F}(x, u, \nabla u) = \frac{1}{2} |\nabla u|^2 - \frac{1}{2} \lambda u^2 - \frac{1}{p} |u|^p$, $h = \chi$ and $a = 0$). Let $\chi := \chi_{\tau}$. Then, by Lemma [3.3,](#page-8-3)

$$
\mathrm{div}\chi_{\tau}=N-k.
$$

Moreover, since $1 - k\varphi(t) < 1$ for $t \in (\tau, \infty)$, and $|y| \in (\tau, \infty)$ for every $(y, z) \in \Omega$, Lemma [3.3](#page-8-3) yields

$$
\langle \mathrm{d}\chi_{\tau}(y,z) \left[\xi\right],\xi\rangle \leq \left|\xi\right|^2 \quad \forall (y,z) \in \Omega, \ \xi \in \mathbb{R}^N.
$$

By assumption, $\langle \chi_{\tau}, \nu_{\Omega} \rangle > 0$ a.e. on $\partial \Omega$. Therefore, if u is a nontrivial solution of (3.1) we have, using (3.3) , that

$$
0 < (N-k)\left(\frac{1}{p} - \frac{1}{2}\right) \int_{\Omega} \left[|\nabla u|^2 - \lambda u^2\right] dx + \int_{\Omega} |\nabla u|^2 dx
$$
\n
$$
= (N-k)\left(\frac{1}{p} - \frac{1}{2} + \frac{1}{N-k}\right) \int_{\Omega} |\nabla u|^2 dx - (N-k)\left(\frac{1}{p} - \frac{1}{2}\right) \lambda \int_{\Omega} u^2 dx,
$$

that is,

$$
\left(\frac{1}{2} - \frac{1}{p}\right) \lambda \int_{\Omega} u^2 dx > \left(\frac{1}{2_{N,k}^*} - \frac{1}{p}\right) \int_{\Omega} |\nabla u|^2 dx.
$$

Therefore, if $p \geq 2^*_{N,k}$ and

$$
\lambda \leq \frac{2(p-2^*_{N,k})}{2^*_{N,k}(p-2)}\inf_{\substack{u\in H^1_0(\Omega)\\ u\neq 0}}\frac{\int_{\Omega}|\nabla u|^2\,dx}{\int_{\Omega}u^2dx}=\frac{2(p-2^*_{N,k})}{2^*_{N,k}(p-2)}\lambda_1(\Omega),
$$

problem [\(3.1\)](#page-8-1) does not have a nontrivial solution in Ω , as claimed. \square

The following result was proved in [\[5\]](#page-14-9).

Proposition 3.5. *If* Θ *is doubly starshaped then* $\Theta \subset (t_0, \infty) \times \mathbb{R}^{N-k-1}$ *and* $\langle \chi_{t_0}, \nu_{\Theta} \rangle > 0$ *a.e.* on $\partial \Theta$, *with* t_0 *as in* Definition [3.1](#page-8-4)*.*

Proof. See the proof of (4.11) in [\[5\]](#page-14-9).

Proof of Theorem [3.2.](#page-8-2) The conclusion follows immediately from Propositions [3.4](#page-9-1) and [3.5.](#page-9-2) \Box

$$
\sqcup
$$

4. Existence and nonexistence of symmetric ground states to supercritical problems

Next, we come back to our original supercritical problem

$$
(\wp_{\lambda}) \qquad \qquad \left\{ \begin{array}{rcl} -\Delta v = \lambda v + |v|^{2_{N,k}^* - 2} \, v & \text{in } \Omega, \\ v = 0 & \text{on } \partial \Omega, \end{array} \right.
$$

where

$$
\Omega:=\{(y,z)\in\mathbb{R}^{k+1}\times\mathbb{R}^{N-k-1}:(\left|y\right|,z)\in\Theta\}
$$

for some bounded smooth domain Θ in \mathbb{R}^{N-k} with $\overline{\Theta} \subset (0,\infty) \times \mathbb{R}^{N-k-1}$, $1 \leq k \leq$ $N-3$, and $2^*_{N,k} := \frac{2(N-k)}{N-k-2}$ $\frac{2(N-\kappa)}{N-k-2}.$

An $O(k + 1)$ -invariant function $v : \Omega \to \mathbb{R}$ can be written as $v(y, z) = u(|y|, z)$ for some function $u : \Theta \to \mathbb{R}$. A straightforward computation shows that

(4.1)
$$
\Delta v = \frac{1}{a(x)} \text{div}(a(x)\nabla u),
$$

where $a(x_1, \ldots, x_{N-k}) := x_1^k$. Hence, v is an $O(k+1)$ -invariant solution of (φ_λ) if and only if u solves

$$
\left(\wp_{\lambda}^{\#}\right) \qquad \qquad \left\{ \begin{array}{rcl} -\text{div}(x_1^k \nabla u) = \lambda x_1^k u + x_1^k \left|u\right|^{2^*-2} u & \text{in } \Theta, \\ u = 0 & \text{on } \partial \Theta, \end{array} \right.
$$

where $2^* = 2^*_{N,k}$ is the critical exponent in dimension $n := N - k = \dim(\Theta)$. So this problem is a special case of the problem treated in section [2](#page-2-1) with $a(x_1, \ldots, x_n) := x_1^k$ and $a = b = c$.

For these functions a, b, c we simplify notation and write $\ell_{\lambda}^{[k]}$ $\frac{[k]}{\lambda}, \; \kappa^{[k]}, \; \lambda^{[k]}_m, \; \lambda^{[k]}_{m,*}$ instead of $\ell_{\lambda}^{a,b,c}, \kappa^{a,c}, \lambda_{m}^{a,b}, \lambda_{m,*}^{a,b,c}$. Note that

$$
\kappa^{[k]} = \left(\min_{x \in \overline{\Theta}} x_1^k\right) \frac{1}{n} S^{n/2}.
$$

Proposition 4.1. *If* $\alpha := \min_{x \in \overline{\Theta}} x_1$ *and* $\lambda \leq 0$, *then* $\ell_{\lambda}^{[k]} = \frac{\alpha^k}{n}$ $\frac{x^{\kappa}}{n}S^{n/2}$ and it is not *attained by* J_{λ} *on* $\mathcal{N}_{\lambda} \equiv \mathcal{N}_{\lambda}(\Theta)$.

Proof. By Theorem [2.1](#page-4-0) it is enough to show this for $\lambda = 0$. Arguing by contradiction, assume that $\ell_0^{[k]} < \frac{\alpha^k}{n}$ $\frac{\alpha^k}{n} S^{n/2}$. Then there exists $\varphi \in C_c^{\infty}(\Theta) \cap \mathcal{N}_0(\Theta)$ such that

$$
J_0(\varphi) < \frac{\alpha^k}{n} S^{n/2} = \kappa^{[k]}.
$$

Since supp (φ) is a compact subset of $(\alpha, \infty) \times \mathbb{R}^{n-1}$, there exists a $\varrho \in (\alpha, \infty)$ such that $\text{supp}(\varphi) \subset B := \{ x \in \mathbb{R}^n : (x_1 - \varrho)^2 + x_2^2 + \cdots + x_n^2 < (\alpha - \varrho)^2 \}$. Hence, $\varphi \in \mathcal{N}_0(B)$. Theorem [3.2](#page-8-2) and the discussion given at the beginning of this section imply that problem

$$
-\text{div}(x_1^k \nabla u) = x_1^k |u|^{2^*-2} u \quad \text{in } B, \qquad u = 0 \quad \text{on } \partial B
$$

does not have a nontrivial solution. So, by Theorem [2.1,](#page-4-0) $\inf_{\mathcal{N}_0(B)} J_0 = \kappa^{[k]} =$ α^k $\frac{\alpha^{\kappa}}{n}S^{n/2}$. But

$$
\inf_{\mathcal{N}_0(B)} J_0 \le J_0(\varphi) < \frac{\alpha^k}{n} S^{n/2}.
$$

This is a contradiction. We conclude that $\ell_0^{[k]} = \frac{\alpha^k}{n}$ $\frac{\alpha^{\kappa}}{n}S^{n/2}.$

Since this value is the same for every Θ such that $\alpha := \min_{x \in \overline{\Theta}} x_1$, a standard argument shows that $\ell_0^{[k]}$ is not attained by J_0 on $\mathcal{N}_0 \equiv \mathcal{N}_0(\Theta)$.

Set

$$
\alpha := \min_{x \in \overline{\Theta}} x_1, \qquad \beta := \max_{x \in \overline{\Theta}} x_1.
$$

Lemma 4.2. For every positive function $f \in C^2[\alpha, \beta]$ which satisfies (4.2) $kf^2(\alpha) \leq t^k f^2(t)$ and $t^k f^{2^*}(t) \leq \alpha^k f^{2^*}(\alpha)$ $\forall t \in [\alpha, \beta],$ *we have that*

$$
\lambda_{0,*}^{[k]}\geq\min_{t\in[\alpha,\beta]} \frac{-\left(t^k f'(t)\right)'}{t^k f(t)}
$$

.

Proof. Let $u \in H_0^1(\Theta)$, $u \neq 0$, and set $u(x) = f(x_1)w(x)$. Then

$$
\int_{\Theta} x_1^k |\nabla u|^2 = \int_{\Theta} \left(x_1^k f^2 |\nabla w|^2 + x_1^k f' f \frac{\partial w^2}{\partial x_1} + x_1^k (f')^2 w^2 \right)
$$

=
$$
\int_{\Theta} \left(x_1^k f^2 |\nabla w|^2 + x_1^k f' \frac{\partial (f w^2)}{\partial x_1} \right)
$$

=
$$
\int_{\Theta} \left(x_1^k f^2 |\nabla w|^2 - (x_1^k f')' f w^2 \right).
$$

So, if $\lambda \leq \frac{-\left(t^k f'(t)\right)'}{t^k f(t)}$ $\frac{t}{t^k f(t)}$ for all $t \in [\alpha, \beta]$, we have that

$$
\frac{\int_{\Theta} x_1^k |\nabla u|^2 - \lambda \int_{\Theta} x_1^k u^2}{\left(\int_{\Theta} x_1^k |u|^{2^*}\right)^{2/2^*}} = \frac{\int_{\Theta} x_1^k f^2 |\nabla w|^2 - \int_{\Theta} \left[\left(x_1^k f'\right)' + \lambda x_1^k f\right] f w^2}{\left(\int_{\Theta} x_1^k f^{2^*} |w|^{2^*}\right)^{2/2^*}} \times \frac{\left(\int_{\Theta} x_1^k f^{2^*} |w|^{2^*}\right)^{2/2^*}}{\left(\int_{\Theta} x_1^k f^{2^*} |w|^{2^*}\right)^{2/2^*}} \ge \frac{\alpha^k f^2(\alpha) \int_{\Theta} |\nabla w|^2}{\alpha^{2k/2^*} f^2(\alpha) \left(\int_{\Theta} |w|^{2^*}\right)^{2/2^*}} \times \frac{\alpha^2 k'^2}{\left(\int_{\Theta} |w|^2\right)^{2/2^*}} \ge \alpha^{2k/n} S > 0 \quad \text{for all } u \in H_0^1(\Theta), u \ne 0.
$$

This implies that $\lambda < \lambda_1^{[k]}$ and, hence, that

$$
\max_{t>0} J_{\lambda}(tu) = \frac{1}{n} \left(\frac{\int_{\Theta} x_1^k \left| \nabla u \right|^2 - \lambda \int_{\Theta} x_1^k u^2}{\left(\int_{\Theta} x_1^k |u|^{2^*} \right)^{2/2^*}} \right)^{n/2} \ge \frac{\alpha^k}{n} S^{n/2}
$$

for all $u \in H_0^1(\Theta), u \neq 0$. Therefore, $\ell_{\lambda}^{[k]} = \frac{\alpha^k}{n}$ $\sum_{n=0}^{\infty} S^{n/2}$ for every $\lambda \leq \min_{t \in [\alpha, \beta]} \frac{-(t^k f'(t))^{\lambda}}{t^k f(t)}$ $\frac{t^{k} f(t)}{t^{k} f(t)},$ and the conclusion follows.

We obtain the following estimates for $\lambda_{0,*}^{[k]}$.

Proposition 4.3. $\lambda_{0,*}^{[k]} \geq 0$ and

$$
\lambda_{0,*}^{[k]}\geq\left\{\begin{array}{ll}\frac{(k-1)^2}{4\beta^2} & \text{if }2k\geq n,\\ \frac{k}{(2^*\beta)^2}\left((2^*-1)k-2^*\right) & \text{if }2k\leq n.\end{array}\right.
$$

Therefore $\lambda_{0,*}^{[k]} > 0$ *if* $k \geq 2$.

Proof. Proposition [4.1](#page-10-2) implies that $\lambda_{0,*}^{[k]} \geq 0$.

Set $f(t) := t^{-\gamma}$ with $\frac{k}{2^*} \leq \gamma \leq \frac{k}{2}$. This function satisfies [\(4.2\)](#page-11-1) and, since

$$
\frac{-\left(t^{k}f'(t)\right)'}{t^{k}f(t)} = \frac{\gamma(k-\gamma-1)}{t^{2}},
$$

Lemma [4.2](#page-11-2) implies that

$$
\lambda_{0,*}^{[k]}\geq \frac{\gamma(k-\gamma-1)}{\beta^2}.
$$

Now observe that the the function $\phi(\gamma) := \gamma(k - \gamma - 1)$ attains its maximum on the interval $\left[\frac{k}{2^*}, \frac{k}{2}\right]$ at the point

$$
\gamma_*:=\max\left\{\frac{k-1}{2},\frac{k}{2^*}\right\}.
$$

Therefore $\lambda_{0,*}^{[k]} \geq \frac{1}{\beta^2} \phi(\gamma_*) = \frac{1}{\beta^2} \gamma_*(k - \gamma_* - 1)$, as claimed. Finally, note that $k > \frac{2^*}{2^*-1}$ $\frac{2^*}{2^*-1} = \frac{2n}{n+2}$ if $k \ge 2$. Hence, $\lambda_{0,*}^{[k]} > 0$ if $k \ge 2$. □

Proof of Theorem [1.1.](#page-1-0) Using Corollary [2.3](#page-7-0) it is easily seen that, if $v(y, z) = u(|y|, z)$, then v is an $O(k+1)$ -invariant ground state for problem (φ_{λ}) if and only if u is a

ground state for problem $(\wp_{\lambda}^{\#})$. So Theorem [1.1](#page-1-0) follows immediately from Proposition [4.1,](#page-10-2) Theorem [2.1](#page-4-0) and Proposition [4.3.](#page-11-0)

The following result shows that $\lambda_{0,*}^{[1]} > 0$ if the domain is thin enough in the x_1 -direction.

Proposition 4.4. *If* $\frac{\beta}{\alpha} \leq \frac{n}{n-2}$ *then* $\lambda_{0,*}^{[k]} \geq \frac{k^2}{4\beta^2} > 0$ *for all* $k \geq 1$. *Proof.* Set $f(t) := e^{-\gamma(t-\alpha)}$ with $\frac{k}{2^*\alpha} \leq \gamma \leq \frac{k}{2\beta}$, and write $g(t) := t^k f^2(t)$ and $h(t) :=$ $t^k f^{2^*}(t)$. Then $g'(t) = t^{k-1} e^{-2\gamma(t-\alpha)}(k-2\gamma t) \geq 0$ and $h'(t) = t^{k-1} e^{-2^*\gamma(t-\alpha)}(k-2\gamma t)$ $2^*\gamma t \leq 0$ for all $t \in [\alpha, \beta]$, so f satisfies [\(4.2\)](#page-11-1). Since

$$
\frac{-\left(t^k f'(t)\right)'}{t^k f(t)} = \frac{\gamma t^{k-1} e^{-\gamma(t-\alpha)} (k-\gamma t)}{t^k e^{-\gamma(t-\alpha)}} = \frac{\gamma(k-\gamma t)}{t},
$$

Lemma [4.2](#page-11-2) implies that

$$
\lambda_{0,*}^{[k]}\geq \frac{\gamma(k-\gamma\beta)}{\beta}.
$$

Now observe that the the function $\phi(\gamma) := \gamma (k - \gamma \beta)$ attains its maximum at the point $\gamma_* := \frac{k}{2\beta}$. Hence, $\lambda_{0,*}^{[k]} \ge \frac{k^2}{4\beta^2} > 0$, as claimed.

5. Some open questions and comments

Many questions remain open. Here are some of them.

Problem 1. Concerning problem $(\varphi_{\lambda}^{\#})$:

- (1) *Is it true that* $\lambda_{0,*}^{[1]} > 0$ *for any domain* Θ *, and not only for thin domains?*
- (2) For $m \geq 1$, is $\lambda_{m,*}^{[k]} > \lambda_m^{[k]}$, or is $\lambda_{m,*}^{[k]} = \lambda_m^{[k]}$ as in the classical Brezis-*Nirenberg case?*

(3) What happens in general at $\lambda_{m,*}^{[k]}$? *Is there, or not, a ground state of problem* $(\wp_{\lambda}^{\#})$ for $\lambda = \lambda_{m,*}^{[k]}$?

Problem 2. *Concerning the general anisotropic problem* [\(2.1\)](#page-2-2)*:*

- (1) *Is* $\lambda_{0,*}^{a,b,c}$ always nonnegative? Or are there examples where a ground state *exists for some* $\lambda < 0$? For all $\lambda < 0$?
- (2) *Can one give lower estimates for* $\lambda_{m,*}^{a,b,c}$ *in some cases?*
- (3) *Suppose that* $c \in C^1(\overline{\Theta})$ *in addition to our earlier assumptions.* If $\kappa^{a,c}$ *is attained only at points which are non-stationary for* $\frac{a(x)^{n/2}}{c(x)(n-2)}$ $\frac{a(x)-b}{c(x)(n-2)/2}$ and lie on *the boundary of* Θ *, is it then true that* $\lambda_{0,*}^{a,b,c} > 0$?

Two particular cases of (3) are: $c = 1$, and $a = b = c$. If the answer is positive in the first case, this would be in contrast to the results in [\[10\]](#page-14-14) and [\[13\]](#page-14-16). A positive answer in the second case would be a generalization of our results for $(\wp_{\lambda}^{\#})$. A partial answer can be given using Proposition [4.3.](#page-11-0) Consider, for example, the problem

(5.1)
$$
-\operatorname{div}(a(x)\nabla u) = \lambda b(x)u + |u|^{2^*-2}u \text{ in } \Theta, \quad u = 0 \text{ on } \partial \Theta,
$$

where Θ is a bounded smooth domain in \mathbb{R}^n , $n \geq 3$, $\lambda \in \mathbb{R}$, $a \in \mathcal{C}^1(\overline{\Theta})$, $b \in \mathcal{C}^0(\overline{\Theta})$ are strictly positive on $\overline{\Theta}$, and $2^* = \frac{2n}{n-2}$. Then, the following statement holds true.

Proposition 5.1. *If* $a(x) \ge x_1^k \ge b(x)$ *for all* $x \in \Theta$ *and* $\min_{x \in \overline{\Theta}} a(x) = (\min_{x \in \overline{\Theta}} x_1)^k >$ 0 for some $k \ge 2$, then $\lambda_{0,*}^{a,b,1} > 0$.

Proof. Let $\alpha := \min_{x \in \overline{\Theta}} x_1 > 0$. For every $u \in H_0^1(\Theta)$, $u \neq 0$, $\lambda \in [0, \lambda_{0,*}^{[k]}]$ we have that

$$
\frac{\int_{\Theta} a(x) |\nabla u|^2 - \lambda \int_{\Theta} b(x) u^2}{\alpha^{2k/2^*} \left(\int_{\Theta} |u|^{2^*} \right)^{2/2^*}} \ge \frac{\int_{\Theta} x_1^k |\nabla u|^2 - \lambda \int_{\Theta} x_1^k u^2}{\left(\int_{\Theta} x_1^k |u|^{2^*} \right)^{2/2^*}} > 0.
$$

Hence, $\lambda < \lambda_1^{a,b}$ and

$$
\frac{1}{n} \left(\frac{\int_{\Theta} a(x) \left| \nabla u \right|^2 - \lambda \int_{\Theta} b(x) u^2}{\left(\int_{\Theta} \left| u \right|^{2^*} \right)^{2/2^*}} \right)^{n/2} \ge \alpha^{\frac{k(n-2)}{2}} \frac{1}{n} \left(\frac{\int_{\Theta} x_1^k \left| \nabla u \right|^2 - \lambda \int_{\Theta} x_1^k u^2}{\left(\int_{\Theta} x_1^k \left| u \right|^{2^*} \right)^{2/2^*}} \right)^{n/2}
$$

$$
\ge \alpha^{\frac{k(n-2)}{2}} \frac{\alpha^k}{n} S^{n/2} = (\alpha^k)^{n/2} \frac{1}{n} S^{n/2} = \kappa^{a,1}.
$$

It follows from Theorem [2.1](#page-4-0) that

 $\ell_{\lambda}^{a,b,1} = \kappa^{a,1}$ for all $\lambda \in (-\infty, \lambda_{0,*}^{[k]}].$

Hence, by Proposition [4.3,](#page-11-0) $\lambda_{0,*}^{a,b,1} \geq \lambda_{0,*}^{[k]} > 0$, as claimed.

REFERENCES

- [1] H. Br´ezis, L. Nirenberg: Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. Comm. Pure Appl. Math. 36 (1983), 437–477.
- [2] A. Capozzi, D. Fortunato, G. Palmieri: An existence result for nonlinear elliptic problems involving critical Sobolev exponent. Ann. Inst. H. Poincaré, Anal. Non Linéaire 2 (1985), 463–440.
- [3] G. Cerami, S. Solimini, M. Struwe: Some existence results for superlinear elliptic boundary value problems involving critical exponents. J. Funct. Anal. 69 (1986), 289–306.

- [4] M. Clapp, J. Faya: Multiple solutions to anisotropic critical and supercritical problems in symmetric domains. Preprint.
- [5] M. Clapp, J. Faya, A. Pistoia: Nonexistence and multiplicity of solutions to elliptic problems with supercritical exponents. Calc. Var. Partial Differential Equations 48 (2013), 611–623.
- [6] M. Clapp, A. Szulkin: A supercritical elliptic problem in a cylindrical shell. Progress in Nonlinear Differential Equations and Their Applications 85 (2014), 233–242.
- [7] M. Clapp, T. Weth: Multiple solutions for the Brezis-Nirenberg problem. Adv. Differential Equations 10 (2005), 463–480.
- [8] G. Devillanova, S. Solimini, Concentration estimates and multiple solutions to elliptic problems at critical growth. Adv. Diff. Eq. 7 (2002), 1257–1280.
- [9] G. Devillanova, S. Solimini, A multiplicity result for elliptic equations at critical growth in low dimension. Comm. Contemp. Math. 5 (2003), 171–177.
- [10] H. Egnell: Semilinear elliptic equations involving critical Sobolev exponents. Arch. Rational Mech. Anal. 104 (1988), 27–56.
- [11] F. Gazzola, B. Ruf: Lower order perturbations of critical growth nonlinearities in semilinear elliptic equations. Adv. Diff. Eq. 2 (1997), 555–572.
- [12] R. Hadiji, R. Molle, D. Passaseo, H. Yazidi: Localization of solutions for nonlinear elliptic problems with critical growth. C. R. Math. Acad. Sci. Paris 343 (2006), 725–730.
- [13] R. Hadiji, H. Yazidi: Problem with critical Sobolev exponent and with weight. Chin. Ann. Math. Ser. B 28 (2007), 327–352.
- [14] S. Kim, A. Pistoia: Supercritical problems in domains with thin toroidal holes. Discrete Contin. Dyn. Syst. 34 (2014), 4671–4688.
- [15] A. Pankov: Periodic nonlinear Schrödinger equation with application to photonic crystals. Milan J. Math. 73 (2005), 259–287.
- [16] D. Passaseo: Nonexistence results for elliptic problems with supercritical nonlinearity in nontrivial domains. J. Funct. Anal. 114 (1993), 97–105.
- [17] D. Passaseo: New nonexistence results for elliptic equations with supercritical nonlinearity. Differential Integral Equations 8 (1995), 577–586.
- [18] S.I. Pohozhaev: Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$, Soviet Math. Dokl. 6 (1965). 1408–1411.
- [19] P. Pucci, J. Serrin: A general variational identity. Indiana Univ. Math. J. 35 (1986), 681–703.
- [20] A. Szulkin, T. Weth: Ground state solutions for some indefinite variational problems. J. Funct. Anal. 257 (2009), 3802–3822.
- [21] A. Szulkin, T. Weth: The method of Nehari manifold. Handbook of nonconvex analysis and applications, 597–632, Int. Press, Somerville, MA, 2010.
- [22] A. Szulkin, T. Weth, M. Willem: Ground state solutions for a semilinear problem with critical exponent. Differential Integral Equations 22 (2009), 913–926.
- [23] J. Wei, S. Yan: Infinitely many positive solutions for an elliptic problem with critical or supercritical growth. J. Math. Pures Appl. 96 (2011), 307–333.
- [24] M. Willem: Minimax theorems. Progress in Nonlinear Differential Equations and their Applications, 24. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [25] D. Zhang: On multiple solutions of $\Delta u + \lambda u + |u|^{\frac{4}{n-2}} = 0$. Nonlinear Anal. TMA 13 (1989), 353–372.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EX-TERIOR, C.U., 04510 MÉXICO D.F., MEXICO

E-mail address: monica.clapp@im.unam.mx

Dipartimento di Metodi e Modelli Matematici, Universita di Roma "La Sapienza", via ´ Antonio Scarpa 16, 00161 Roma, Italy

E-mail address: pistoia@dmmm.uniroma1.it

Department of Mathematics, Stockholm University, 106 91 Stockholm, Sweden E-mail address: andrzejs@math.su.se