

Isolated Singularities of Polyharmonic Operator in Even Dimension

Dhanya Rajendran^a, Abhishek Sarkar^{b,*}

^aDepartment of Mathematics
 Indian Institute of Science, Bangalore-560012, Karnataka, India.
^bTIFR Centre For Applicable Mathematics
 Post Bag No. 6503, Sharda Nagar, Bangalore-560065, Karnataka, India.

Abstract

We consider the equation $\Delta^2 u = g(x, u) \geq 0$ in the sense of distribution in $\Omega' = \Omega \setminus \{0\}$ where u and $-\Delta u \geq 0$. Then it is known that u solves $\Delta^2 u = g(x, u) + \alpha \delta_0 - \beta \Delta \delta_0$, for some non-negative constants α and β . In this paper we study the existence of singular solutions to $\Delta^2 u = a(x)f(u) + \alpha \delta_0 - \beta \Delta \delta_0$ in a domain $\Omega \subset \mathbb{R}^4$, a is a non-negative measurable function in some Lebesgue space. If $\Delta^2 u = a(x)f(u)$ in Ω' , then we find the growth of the nonlinearity f that determines α and β to be 0. In case when $\alpha = \beta = 0$, we will establish regularity results when $f(t) \leq Ce^{\gamma t}$, for some $C, \gamma > 0$. This paper extends the work of Soranzo (1997) where the author finds the barrier function in higher dimensions ($N \geq 5$) with a specific weight function $a(x) = |x|^\sigma$. Later we discuss its analogous generalization for the polyharmonic operator.

Keywords: Elliptic system; polyharmonic operator; existence of solutions; singularity
2010 MSC: 35J40, 35J61, 35J91

1. Introduction

Isolated singularities of elliptic operators are studied extensively, see for eg. [2],[10], [14], [15] and [16]. In this paper we wish to address the following problem and the questions related to it for the biharmonic(polyharmonic) operator in $\mathbb{R}^4(\mathbb{R}^{2m})$:-

Question: If a non negative measurable function u is known to solve a PDE in the sense of distribution in a punctured domain, then what can one say about the differential equation satisfied by u in the entire domain?

In [2], Brezis and Lions answered this question for the Laplace operator with the assumption that

$$0 \leq -\Delta u = f(u) \text{ in } \Omega \setminus \{0\}, u \geq 0, \liminf_{t \rightarrow \infty} \frac{f(t)}{t} > -\infty, \Omega \subset \mathbb{R}^N.$$

With the above hypotheses it was proved that both u and $f(u)$ belong to $L^1(\Omega)$, and satisfy $-\Delta u = f(u) + \alpha \delta_0$, for some $\alpha \geq 0$. For the dimension $N \geq 3$, P.L.Lions[10] found a sharp condition on f that determines whether α is zero or not in the previous expression. In [5], the authors further extended the result for dimension $N = 2$ by finding the minimal growth rate of the function f which guaranteed α to be 0.

Taliaferro, in his series of papers (see for e.g. [15], [16], [8]) studied the isolated singularities of non-linear elliptic inequalities. In [16] the author studied the asymptotic behaviour of the positive solution of the differential inequality

$$0 \leq -\Delta u \leq f(u) \tag{1.1}$$

*Corresponding author
 Email addresses: dhanya.tr@gmail.com (Dhanya Rajendran), abhishek@math.tifrbng.res.in (Abhishek Sarkar)

in a punctured domain under various assumptions on f . If $N \geq 3$ and the function f has a "super-critical" growth as in Lions[10], (i.e. $\lim_{t \rightarrow \infty} \frac{f(t)}{t^{\frac{N}{N-2}}} = \infty$,) then there exists arbitrarily 'large solutions' of (1.1). When $N = 2$, it was proved that there exists a punctured neighborhood of the origin such that (1.1) admits arbitrarily large solutions near the origin, provided that $\log f(t)$ has a superlinear growth at infinity. Moreover author characterizes the singularity at the origin of all solutions u of (1.1) when $\log f(t)$ has a sublinear growth. Later Taliaferro, Ghergu and Moradifam in [8] generalized these results to polyharmonic inequalities.

The study of the polyharmonic equations of the type $(-\Delta)^m u = h(x, u)$ is associated to splitting the equation into a non-linear coupled system involving Laplace operator alone. Orsina and Ponce[12] proved the existence of solutions to

$$(1) \quad \begin{cases} -\Delta u &= f(u, v) + \mu & \text{in } \Omega, \\ -\Delta v &= g(u, v) + \eta & \text{in } \Omega, \\ u &= v = 0 & \text{on } \partial\Omega. \end{cases}$$

with the assumption that the continuous functions f and g are non increasing in first and second variables respectively with $f(0, t) = g(s, 0) = 0$. But here the authors assumed that μ and η are diffusive measures and Dirac distribution is not a diffusive measure. Considerable amount of existence/non-existence results have been proved for the problem (1) when f is a function of v alone and g depends only on u and μ, η are Radon measures. For eg. see [1] where the authors assumed $f(u, v) = v^p$, $g(u, v) = u^q$ and with non-homogenous boundary condition. In [6] authors dealt with sign changing functions f and g , with a polynomial type growth at infinity and the measure μ and η were assumed to be multiples of δ_0 .

Our paper is closely related to the work of Soranzo [14] where author considers the equation:

$$\Delta^2 u = |x|^\sigma u^p \quad \text{with } u > 0, \quad -\Delta u > 0 \text{ in } \Omega \subset \mathbb{R}^N, N \geq 4 \text{ and } \sigma \in (-4, 0).$$

A complete description of the singularity was provided when $1 < p < \frac{N+\sigma}{N-4}$ for $N \geq 5$, or $1 < p < \infty$ when $N = 4$. In this work we prove that the results of Soranzo can be improved for the dimension $N = 4$ by replacing u^p by more general exponential type function.

2. Preliminaries

We assume that Ω is a bounded open set in \mathbb{R}^N , $N \geq 4$ with smooth boundary and $0 \in \Omega$. We denote Ω' to be $\Omega \setminus \{0\}$. In this section we discuss some of the well known results for biharmonic operator.

Theorem 2.1. (Brezis - Lions [2]) *Let $u \in L^1_{loc}(\Omega')$ be such that $\Delta u \in L^1_{loc}(\Omega')$ in the sense of distributions in Ω' , $u \geq 0$ in Ω such that*

$$-\Delta u + au \geq g \quad \text{a.e in } \Omega,$$

where a is a positive constant and $g \in L^1_{loc}(\Omega)$. Then there exist $\varphi \in L^1_{loc}(\Omega)$ and $\alpha \geq 0$ such that

$$-\Delta u = \varphi + \alpha \delta_0 \text{ in } \mathcal{D}'(B_R) \tag{2.1}$$

where δ_0 is the Dirac mass at origin. In particular, $u \in M^p_{loc}(B_R)$ ¹ where $p = N/N - 2$ when $N \geq 3$ and $1 \leq p < \infty$ is arbitrary when $N = 2$.

Theorem 2.2. (Weyl Lemma, Simader[13]) *Suppose $G \subset \mathbb{R}^N$ be open and let $u \in L^1_{loc}(G)$ satisfies*

$$\int_G u \Delta^2 \varphi dx = 0 \text{ for all } \varphi \in C_c^\infty(G), \text{ i.e. } \Delta^2 u = 0 \text{ in } \mathcal{D}'(G).$$

Then there exists $\tilde{u} \in C^\infty(G)$ with $\Delta^2 \tilde{u} = 0$ and $u = \tilde{u}$ a.e in G .

¹ $M^p_{loc}(B_R)$ denotes the Marcinkiewicz space

Theorem 2.3. (Weak maximum principle:) Let $u \in W^{4,r}(\Omega)$ be a solution of

$$\begin{cases} \Delta^2 u = f(x) \geq 0 & \text{in } \Omega \\ u \geq 0, -\Delta u \geq 0 & \text{on } \partial\Omega \end{cases}$$

Then we have $u \geq 0$ and $-\Delta u \geq 0$ in Ω .

Proof of maximum principle easily follows by splitting the equation into a (coupled) system of second order PDE's say: $w = -\Delta u$ and $-\Delta w = f$ with the corresponding boundary conditions. Using similar ideas we can in fact prove a maximum principle with weaker assumptions on the smoothness of u , which is stated below:

Theorem 2.4. Let $u, \Delta u \in L^1(\Omega)$ and $\Delta^2 u \geq 0$ in the sense of distributions. Also assume that $u, \Delta u$ are continuous near $\partial\Omega$ and $u > 0, -\Delta u > 0$ near $\partial\Omega$. Then $u(x) \geq 0$ in Ω .

Definition 2.1. Fundamental solution of Δ^2 is defined as a locally integrable function Φ in \mathbb{R}^N for which $\Delta^2 \Phi = \delta_0$ and precisely expressed as

$$\Phi(x) = a_N \begin{cases} |x|^{4-N} & \text{if } N \geq 5 \\ \log \frac{5}{|x|} & \text{if } N = 4 \\ |x| & \text{if } N = 3 \\ |x|^2 \log \frac{5}{|x|} & \text{if } N = 2 \end{cases}$$

for some constant $a_N > 0$.

Theorem 2.5. Suppose $g : \Omega' \times [0, \infty) \rightarrow \mathbb{R}^+$ be a measurable function and let $u, \Delta u$ and $\Delta^2 u \in L^1_{loc}(\Omega')$. Let $\Delta^2 u = g(x, u)$ in $\mathcal{D}'(\Omega')$ with $u \geq 0$ and $-\Delta u \geq 0$ a.e in Ω' . Then $u, g(x, u) \in L^1_{loc}(\Omega')$ and there exist a non-negative constants α, β such that $\Delta^2 u = g(x, u) + \alpha\delta_0 - \beta\Delta\delta_0$ in $\mathcal{D}'(\Omega')$.

Proof: Let us write $w = -\Delta u$. Then $-\Delta w = g(x, u) \geq 0$ in $\mathcal{D}'(\Omega')$ and also given that $w, g(x, u) \in L^1_{loc}(\Omega')$. Now as a direct application of Brezis-Lions Theorem 4.4, we obtain

$$-\Delta w = g(x, u) + \alpha\delta_0 \text{ for some } \alpha \geq 0 \quad (2.2)$$

and $w, g(x, u) \in L^1_{loc}(\Omega)$. Since $-\Delta u = w \geq 0$ in Ω' again by Theorem 4.4 $u \in L^1_{loc}(\Omega)$ and

$$-\Delta u = w + \beta\delta_0 \text{ for some } \beta \geq 0.$$

Now substituting $w = -\Delta u - \beta\delta_0$ in (2.2) we get

$$\Delta^2 u = g(x, u) + \alpha\delta_0 - \beta\Delta\delta_0. \quad (2.3)$$

Extending $g(x, u)$ to be zero outside Ω we get $\Delta^2(u - f(u) * \Phi - \alpha\Phi - \beta\Gamma) = 0$ in $\mathcal{D}'(\Omega)$. By Weyl's lemma for biharmonic operators, there exists a biharmonic function $h \in C^\infty(\Omega)$ and

$$u = g(x, u) * \Phi + \alpha\Phi + \beta\Gamma + h \text{ a.e in } \Omega.$$

Note that $\Gamma(x)$ belongs to Marcinkiewicz space $M^{\frac{N}{N-2}}(\Omega)$ when $N \geq 2$. By the property of the convolution of an L^1 function with the functions in $M^{\frac{N}{N-2}}(\mathbb{R}^N)$ we obtain $u \in M^{\frac{N}{N-2}}_{loc}(\Omega)$. \square . The above result has been proved in [14](see Theorem 2) as an application of their main result on the system of equations. Proof is essentially based on the idea of Brezis-Lions type estimates. We have instead given a direct alternative proof for the same result. Theorem 2.5 can be extended for polyharmonic operator in a standard way, for details see Theorem 4.1 .

3. Biharmonic operator in \mathbb{R}^4

In this section we will restrict ourselves to the dimension $N = 4$ and $g(x, u)$ to take a specific form $g(x, u) = a(x)f(u)$. Let Ω be a bounded open set in \mathbb{R}^4 , $0 \in \Omega$ and denote $\Omega' = \Omega \setminus \{0\}$. We assume

- (H1) $f : [0, \infty) \rightarrow [0, \infty)$ is a continuous function which is nondecreasing in \mathbb{R}^+ and $f(0) = 0$.
(H2) $a(x)$ is a non-negative measurable function in $L^k(\Omega)$ for some $k > \frac{4}{3}$.
(H3) There exists $r_0 > 0$ such that $\text{essinf}_{B_{r_0}} a(x) > 0$.

Let u be a measurable function which solves the following problem:

$$(P) \quad \begin{cases} \Delta^2 u = a(x)f(u) & \text{in } \Omega' \\ u \geq 0, \quad -\Delta u \geq 0 & \text{in } \Omega' \end{cases}$$

From Theorem 2.5 we know that u is a distributional solution of $(P_{\alpha, \beta})$

$$(P_{\alpha, \beta}) \quad \begin{cases} \Delta^2 u = a(x)f(u) + \alpha\delta_0 - \beta\Delta\delta_0 \\ u \geq 0, \quad -\Delta u \geq 0 \\ \alpha, \beta \geq 0, u \text{ and } a(x)f(u) \in L^1(\Omega). \end{cases} \text{ in } \Omega,$$

The assumption (H3) suggests that the presence of such a weight function does not reduce the singularity of $a(x)f(u)$ at origin. In particular, if $a(x) = |x|^\sigma$ for $\sigma \in (-3, 0)$, then $a(x)$ satisfies (H2) and (H3).

Now assume that

$$\lim_{t \rightarrow \infty} \frac{f(t)}{t^2} = c \in (0, \infty]. \quad (3.1)$$

i.e. $f(t)$ grows atleast at a rate of t^2 near infinity. Then for some t_0 large enough, we have $f(t) \geq \frac{c}{2}t^2$ for all $t \geq t_0$. Suppose u is a solution of $(P_{\alpha, \beta})$ and f satisfies 3.1. Then we know that for some biharmonic function h

$$u(x) = a(x)f(u) * \Phi + \alpha\Phi + \beta\Gamma + h \text{ a.e in } \Omega$$

where Φ is the fundamental solution of biharmonic operator in \mathbb{R}^4 and Γ is the fundamental solution of $-\Delta$ in \mathbb{R}^4 . Since α and $a(x)f(u)$ are non-negative, we have $u(x) \geq \beta\Gamma(x) + h(x)$. If $\beta \neq 0$, fix an $\tilde{r} \in (0, r_0)$ such that $u(x) \geq \frac{\beta}{2\pi^2|x|^2} \geq t_0$ whenever $|x| < \tilde{r}$. Now,

$$\int_{B_{\tilde{r}}} a(x)f(u) \geq C \int_{B_{\tilde{r}}} |x|^{-4} = \infty$$

which is a contradiction since $a(x)f(u) \in L^1(\Omega)$. Thus $\beta = 0$ if $f(t)$ grows at a rate faster than t^2 near infinity. We state this result in the next lemma.

Lemma 3.1. *Let f satisfies the condition (3.1) and u solves (P). Then for some α non-negative $\Delta^2 u = a(x)f(u) + \alpha\delta_0$ in $\mathcal{D}'(\Omega)$.*

Now onwards we assume that f satisfies (3.1). We would like to address following questions in this paper:

1. Can we find a sharp condition on f that determines whether $\alpha = 0$ or not in $(P_{\alpha, 0})$?
2. If $\alpha = 0$, is it true that u is regular in Ω ?

Definition 3.1. We call f a sub-exponential type function if

$$\lim_{t \rightarrow \infty} f(t)e^{-\gamma t} \leq C \quad \text{for some } \gamma, C > 0.$$

We call f to be of super-exponential type if it is not a sub-exponential type function.

We will show that the above two questions can be answered based on the non-linearity being a sub-exponential type function or not.

Theorem 3.1. (*Removable Singularity*) Let f be a super-exponential type function and u is a distributional solution of (P) . Then u extends as a distributional solution of $(P_{0,0})$.

Proof: Given u solves (P) , we know that $\Delta^2 u = a(x)f(u) + \alpha\delta_0 - \beta\Delta\delta_0$ for some $\alpha, \beta \geq 0$. To show the extendability of the distributional solution we need to prove $\alpha = \beta = 0$. Since f is of super exponential type function, from Lemma 3.1 it is clear that $\beta = 0$. Let us assume that $\alpha > 0$ and derive a contradiction. Note that we can find an r small enough such that $u(x) \geq -\frac{\alpha}{16\pi^2} \log|x|$ whenever $|x| < r$. Since f is not a sub-exponential type function, for a given $\gamma > 0$ there exists $t_0 > 0$ such that $f(t) \geq e^{\gamma t}$ for all $t \geq t_0$. Thus,

$$f(u(x)) \geq f\left(-\frac{\alpha}{16\pi^2} \log|x|\right) \geq e^{-\frac{\gamma\alpha}{16\pi^2} \log|x|}, \quad \text{for } |x| \ll 1.$$

Now if we choose $\gamma = \frac{64\pi^2}{\alpha}$ in the above inequality, it contradicts the fact $a(x)f(u) \in L^1(\Omega)$. Thus $\alpha = \beta = 0$ in $(P_{\alpha,\beta})$. \square

Theorem 3.2. If $f(t) = t^p$ where $1 \leq p < \frac{4+\sigma}{2}$ and $a(x) = |x|^\sigma$, for $\sigma \in (-2, 0)$ then $(P_{\alpha,\beta})$ is solvable for α, β small enough.

Proof follows from Theorem 4(ii) of Soranzo[14]. The idea was to split the equation into a coupled system and find a sub and super solution for the system. In the next theorem when f satisfies 3.1, we find a super solution for $(P_{\alpha,0})$ directly without splitting the equation into a coupled system and then use the idea of monotone iteration to show the existence of a non-negative solution for α small enough. When $\beta \neq 0$, such a direct monotone iteration technique is not applicable as $\Delta\delta_0$ is not a positive or a negative distribution, ie $\phi \geq 0$, does not imply $\langle \Delta\delta_0, \phi \rangle \geq 0$ or ≤ 0 .

Theorem 3.3. Let f and a satisfy the hypotheses (H1)–(H3). Additionally assume $\lim_{t \rightarrow \infty} \frac{f(t)}{t^2} = c \in (0, \infty]$. Then there exists an $\alpha_* > 0$ such that for all $\alpha \leq \alpha_*$ the problem $(P_{\alpha,0})$ admits a solution in $B_r(0)$.

Proof: We use the idea of sub and super solution to construct a distributional solution for $(P_{\alpha,0})$ for α small enough. Clearly $u_0 = 0$ is a subsolution for $(P_{\alpha,0})$. Given that f is a sub-exponential type nonlinearity, there exists a $\gamma > 0$ and a $C > 0$, such that $f(t) \leq Ce^{\gamma t}$ for all $t \in \mathbb{R}^+$.

Now define

$$\bar{u}(x) = \frac{-\log|x| + C\phi}{\gamma} \quad \text{in } B_1(0). \quad (3.2)$$

where ϕ is the unique solution of the following Navier boundary value problem,

$$\begin{cases} \Delta^2 \phi &= -\frac{a(x)}{|x|} \log|x| \quad \text{in } B_1(0) \\ \phi &= 0 = \Delta\phi \quad \text{on } \partial B_1(0). \end{cases} \quad (3.3)$$

We notice that since $a(x) \in L^k(\Omega)$, for some $k > \frac{4}{3}$, the term $a(x)|x|^{-1} \log|x| \in L^p(B_1)$ for some $p > 1$. Hence the existence of a unique weak solution $\phi \in W^{4,p}(B_1)$ is guaranteed by Gazzolla [7], Theorem 2.20. Now by maximum principle we have $\phi \geq 0, -\Delta\phi \geq 0$.

Therefore,

$$\bar{u} \geq 0 \quad \text{in } B_1(0), \quad (3.4)$$

$$-\Delta \bar{u} = \frac{2}{\gamma|x|^2} - \frac{C}{\gamma} \Delta \phi \geq 0. \quad (3.5)$$

and

$$\Delta^2 \bar{u} = \frac{\delta_0}{8\pi^2\gamma} + \frac{C}{\gamma|x|} a(x) |\log|x||. \quad (3.6)$$

Note that $a(x)f(\bar{u}) \leq \frac{C}{|x|} a(x)e^{C\phi}$. By Sobolev embedding, we know $W^{4,p}(\Omega) \hookrightarrow C(\bar{\Omega})$, and hence $e^{C\phi}$ is bounded in $B_1(0)$. Now we fix an $r > 0$ where $e^{C\phi} \leq \frac{|\log|x||}{\gamma}$ in $B_r(0)$. We let $\Omega = B_r(0)$ (where r depends only on γ and C) be a strict subdomain of $B_1(0)$ where $\frac{C}{\gamma|x|} a(x) |\log|x|| \geq a(x)f(\bar{u})$. Now from the choice of r and equations 3.4, (3.5) and (3.6) it is obvious that \bar{u} is a super solution of $(P_{\alpha,0})$ where $\alpha = \frac{1}{8\pi^2\gamma}$. Now let us define inductively with $u_0 = 0$

$$(P_{\alpha,0}^n) \begin{cases} \Delta^2 u_n = a(x)f(u_{n-1}) + \alpha\delta_0 & \text{in } \mathcal{D}'(\Omega) \\ u_n > 0, -\Delta u_n > 0 & \text{in } \Omega \\ u_n = \Delta u_n = 0 & \text{on } \partial\Omega \end{cases}$$

Existence of such a sequence $\{u_n\}$ can be obtained by writing $u_n = w_n + \alpha\Phi$ where

$$\begin{cases} \Delta^2 w_n = a(x)f(u_{n-1}) & \text{in } \Omega, \\ w_n = -\alpha\Phi, \Delta w_n = -\alpha\Delta\Phi & \text{on } \partial\Omega, \\ w_n \in W^{4,r}(\Omega) & \text{for some } r > 1. \end{cases}$$

Existence of w_1 is clear since $f(0) = 0$ and from Theorem 2.2 of [7]. First let us show the positivity of u_1 and $-\Delta u_1$ in Ω . Since w_1 is bounded, we can choose ϵ small enough so that $u_1 = w_1 + \alpha\Phi > 0$ and $-\Delta u_1 > 0$ in B_ϵ . In $\Omega \setminus B_\epsilon$ by weak comparison principle we can show that $u_1 > 0$ and $-\Delta u_1 > 0$. Next we need to show that $u_1 \leq \bar{u}$. Note that by construction, $\bar{u} > 0$ and $-\Delta \bar{u} > 0$ in $\bar{B}_r \setminus \{0\}$. Then, $\bar{u} - u_1$ satisfies the set of equations

$$\begin{cases} \Delta^2(\bar{u} - u_1) \geq 0 & \text{in } \mathcal{D}(\Omega), \\ \bar{u} - u_1 > 0, -\Delta(\bar{u} - u_1) > 0 & \text{near } \partial\Omega. \end{cases}$$

Now using the maximum principle for distributional solutions (Theorem 2.4) we find $u_1 \leq \bar{u}$.

Assume that there exists a function u_k solving $(P_{\alpha,0}^k)$ for $k = 1, 2 \dots n$ and

$$0 \leq u_1 \leq u_2 \dots \leq u_n \leq \bar{u} \text{ in } \Omega.$$

Since f is non-decreasing we have $a(x)f(u_n) \in L^p(\Omega)$, for some $p > 1$. Thus by Sobolev embedding there exists a $w_{n+1} \in C(\bar{\Omega}) \cap W^{4,p}(\Omega)$. Also,

$$\begin{cases} \Delta^2(u_{n+1} - u_n) = a(x)f(u_n) - a(x)f(u_{n-1}) \geq 0 & \text{in } \Omega \\ u_{n+1} = u_n, \Delta u_{n+1} = \Delta u_n & \text{on } \partial\Omega. \end{cases}$$

Again from weak comparison principle $0 < u_n \leq u_{n+1}$ and $0 \leq -\Delta u_n \leq -\Delta u_{n+1}$. As before one can show that $u_{n+1} \leq \bar{u}$. Now if we define $u(x) = \lim_{n \rightarrow \infty} u_n(x)$ one can easily verify that u is a solution of $(P_{\alpha,0})$ for $\alpha = \frac{1}{8\pi^2\gamma}$. For a given f sub-exponential type function we have found a ball of radius r such that $(P_{\alpha,0})$ posed on $B_r(0)$ has a solution u_α for $\alpha = \frac{1}{8\pi^2\gamma}$. This solution u_α is a supersolution for $(P_{\alpha',0})$ posed in $B_r(0)$ and for $\alpha' \in (0, \alpha)$. Thus one can repeat the previous iteration and show that for all $\alpha' \in (0, \alpha)$ there exists a weak solution for $(P_{\alpha',0})$ in $B_r(0)$. \square

Corollary 3.1. *Suppose for a given $\gamma > 0$ there exists a C_γ such that $f(t) \leq C_\gamma e^{\gamma t}$ for all $t \in \mathbb{R}^+$. Then $(P_{\alpha,0})$ has a solution in $B_{r_\alpha}(0)$ for all $\alpha \in (0, \infty)$. In particular if $f(t) = t^p$, $p > 2$ or e^{t^δ} , $\delta < 1$ then $(P_{\alpha,0})$ is solvable for all $\alpha > 0$.*

Next we recall a Brezis-Merle [3] type of estimate for Biharmonic operator in \mathbb{R}^4 . Let h be a distributional solution of

$$(2) \quad \begin{cases} \Delta^2 h = f & \text{in } \Omega \\ h = \Delta h = 0 & \text{on } \partial\Omega. \end{cases}$$

where Ω is a bounded domain in \mathbb{R}^4 .

Theorem 3.4. (C.S Lin [9]) *Let $f \in L^1(\Omega)$ and h is a distributional solution of (2). For a given $\delta \in (0, 32\pi^2)$ there exists a constant $C_\delta > 0$ such that the following inequality holds:*

$$\int_{\Omega} \exp\left(\frac{\delta h}{\|f\|_1}\right) dx \leq C_\delta (\text{diam}\Omega)^4$$

where $\text{diam } \Omega$ denote the diameter of Ω .

Theorem 3.5. *Let f be a sub-exponential type function. Let u be a solution of $(P_{0,0})$ with $u = \Delta u = 0$ on $\partial\Omega$. Then u is regular in Ω .*

Proof: Let u be a solution of $\Delta^2 u = a(x)f(u)$ in Ω with Navier boundary conditions. Write $g(x) = a(x)f(u)$, then $g \in L^1(\Omega)$. Fix a $l > 0$ and split $g = g_1 + g_2$ where $\|g_1\|_1 < \frac{1}{l}$ and $g_2 \in L^\infty(\Omega)$. Let u_2 be the unique solution of

$$\begin{cases} \Delta^2 u_2 = g_2 & \text{in } \Omega, \\ u_2 = 0, \Delta u_2 = 0 & \text{on } \partial\Omega. \end{cases}$$

Then

$$\begin{cases} \Delta^2 u_1 = g_1 & \text{in } \Omega, \\ u_1 = 0, \Delta u_1 = 0 & \text{on } \partial\Omega. \end{cases}$$

Choosing $\delta = 1$ in Theorem 3.4, we find $\int_{\Omega} \exp\left(\frac{|u_1|}{\|g_1\|_1}\right) < C_1 (\text{diam } \Omega)^4$. Thus $e^{l|u_1|} \in L^1(\Omega)$. Since $u_2 \in L^\infty(\Omega)$, we have $e^{l|u|} \in L^1(\Omega)$ for all $l > 0$. We use this higher integrability property of u in establishing its regularity.

We can show that $a(x)f(u) \in L^r(\Omega)$ for some $r > 1$. In fact,

$$\begin{aligned} \int_{\Omega} (a(x)f(u))^r &\leq \tilde{C} \int_{\Omega} a(x)^r e^{\gamma r u} \\ &\leq C_2 \left(\int_{\Omega} a(x)^{pr} \right)^{1/p} \left(\int_{\Omega} e^{p' \gamma r u} \right)^{1/p'} < \infty \end{aligned}$$

if we choose $p, r > 1$ close enough to 1 so that $1 < p.r \leq k$, where $a(x) \in L^k(\Omega)$. Now let v be the unique weak solution of

$$\begin{cases} \Delta^2 v = a(x)f(u) & \text{in } \Omega, \\ v = 0, \Delta v = 0 & \text{on } \partial\Omega. \end{cases}$$

We have $v \in C^{3,\gamma'}(\bar{\Omega})$ for all $\gamma' \in (0, 1)$. Now $u = v + h$ for some biharmonic function h . Therefore $u \in C^{3,\gamma'}(\Omega)$. \square

Remark 3.1. *The previous theorem is true even if $a(x) \in L^k(\Omega)$ for some $k > 1$.*

When f is super exponential in nature an arbitrary solution of $\Delta^2 u = a(x)f(u)$ in $\mathcal{D}'(\Omega)$ need not be bounded. We consider the following example.

Example 3.1. *Let $w(x) = (-4 \log|x|)^{\frac{1}{\mu}}$ for some $\mu > 1$. Then one can verify that whenever $x \neq 0$,*

$$\Delta^2 w = b_1 e^{w^\mu} w^{1-4\mu} [b_2 w^{2\mu} - b_3]$$

for some positive constants b_i . Since $f(w) = b_1 e^{w^\mu} w^{1-4\mu} [b_2 w^{2\mu} - b_3]$ is super exponential in nature, w extends as an unbounded distributional solution of $\Delta^2 w = f(w)$ in $B_r(0)$ for r small enough.

4. Polyharmonic Operator in \mathbb{R}^{2m}

We suppose Ω is a bounded domain in \mathbb{R}^N , $N \geq 2m$ with smooth boundary and $0 \in \Omega$. We denote Ω' as $\Omega \setminus \{0\}$.

Theorem 4.1. *Suppose $g : \Omega' \times [0, \infty) \rightarrow \mathbb{R}^+$ is a measurable function and $\Delta^k u \in L^1_{loc}(\Omega')$ for $k = 0, 1, \dots, m$. Let $(-\Delta)^m u = g(x, u)$ in $\mathcal{D}'(\Omega')$ with $(-\Delta)^k u \geq 0$ for $k = 0, 1, \dots, m-1$ a.e. in Ω' . Then $u, g(x, u) \in L^1_{loc}(\Omega)$ and there exist non-negative constants $\alpha_0, \dots, \alpha_{m-1}$ such that*

$$(-\Delta)^m u = g(x, u) + \sum_{i=0}^{m-1} \alpha_i (-\Delta)^i \delta_0 \text{ in } \mathcal{D}'(\Omega).$$

Now we restrict ourselves to dimension $N = 2m$ and $g(x, u)$ to take a specific form $g(x, u) = a(x)f(u)$. Throughout this section we make the following assumption:

(H1') $f : [0, \infty) \mapsto [0, \infty)$ is a continuous function which is non-decreasing in \mathbb{R}^+ and $f(0) = 0$.

(H2') $a(x)$ is non negative measurable function in $L^k(\Omega)$ for some $k > \frac{2m}{2m-1}$.

(H3') There exists $r_0 > 0$ such that $\text{essinf}_{B_{r_0}} a(x) > 0$.

Let u be a measurable function which satisfies the problem below,

$$(P^1) \quad \begin{cases} (-\Delta)^m u = a(x)f(u) & \text{in } \Omega' \\ (-\Delta)^k u \geq 0 & \text{in } \Omega', \quad k = 0, \dots, m-1 \\ u \in C^{2m}(\overline{\Omega} \setminus \{0\}). \end{cases}$$

Then by 4.1 we know that u is a distribution solution of $(P^1_{\alpha_0, \dots, \alpha_{m-1}})$

$$(P^1_{\alpha_0, \dots, \alpha_{m-1}}) \quad \begin{cases} (-\Delta)^m u = a(x)f(u) + \sum_{i=0}^{m-1} \alpha_i (-\Delta)^i \delta_0 & \text{in } \Omega \\ (-\Delta)^k u \geq 0, \quad k = 0, \dots, m-1 & \text{in } \Omega' \\ \alpha_i \geq 0, \text{ for } i = 0, \dots, m-1 \text{ and } u, a(x)f(u) \in L^1(\Omega). \end{cases}$$

In [4], Soranzo et.al considered a specific equation $(-\Delta)^m u = |x|^\sigma u^p$ in Ω' , with $\sigma \in (-2m, 0)$ and $(-\Delta)^k u \geq 0$, for $k = 0, 1, \dots, m$. By Corollary 1 of [4], if $N = 2m$ and $p > \max\{1, \frac{N+\sigma}{2}\}$ then $\alpha_1 = \alpha_2 = \dots = \alpha_{m-1} = 0$ in $(P^1_{\alpha_0, \dots, \alpha_{m-1}})$. This result can be sharpened for any weight function $a(x)$ satisfying (H3) in a standard way and we skip the details of the proof.

Remark 4.1. *Let u satisfy (P^1) and $\lim_{t \rightarrow \infty} \frac{f(t)}{t^m} = c \in (0, \infty]$. Then we have $\alpha_1 = \alpha_2 = \dots = \alpha_{m-1} = 0$ in $(P^1_{\alpha_0, \dots, \alpha_{m-1}})$ and hence u is a distributional solution of $(-\Delta)^m u = a(x)f(u) + \alpha_0 \delta_0$ in Ω .*

Now the following theorem gives us a sharp condition on f which determines $\alpha_0 = 0$ in $(P^1_{\alpha_0, 0, \dots, 0})$ and the proof is as similar to Theorem 3.1.

Theorem 4.2. *Let f be a super-exponential type function and u is distribution solution of (P^1) . Then u extends as a distributional solution of $(P^1_{0, 0, \dots, 0})$.*

Theorem 4.3. *Let f and a satisfy the hypotheses (H1')–(H3'). Additionally assume $\lim_{t \rightarrow \infty} \frac{f(t)}{t^m} = c \in (0, \infty]$. Then there exists an $\alpha_0 > 0$ such that for all $\alpha \leq \alpha_0$ the problem $(P^1_{\alpha, 0, \dots, 0})$ admits a solution in $B_r(0)$, where the radius of the ball depends on the nonlinearity f .*

Proof: We proceed as in Theorem 3.3, by constructing sub and super distributional solution for $(P_{\alpha,0,\dots,0}^1)$ for all α small enough. We note that $u_0 = 0$ is a sub-solution, and let

$$\bar{u}(x) = \frac{-\log|x| + C\phi}{\gamma} \text{ in } B_1(0) \quad (4.1)$$

where ϕ is the unique solution of the following Navier boundary value problem,

$$\begin{cases} (-\Delta)^m \phi &= -\frac{a(x)}{|x|} \log|x| \text{ in } B_1(0) \\ \phi = \Delta \phi &= \dots = (\Delta)^{m-1} \phi \text{ on } \partial B_1(0). \end{cases} \quad (4.2)$$

Then \bar{u} is a supersolution of $(P_{\alpha,0,\dots,0}^1)$ in a small ball $B_r(0)$. Rest of the proof follows exactly as in the case of biharmonic operator. \square

Next we state a Brezis-Merle type of estimates for poly-harmonic operator in \mathbb{R}^{2m} .

Theorem 4.4. (Martinazzi [11]) Let $f \in L^1(B_R(x_0))$, $B_R(x_0) \subset \mathbb{R}^{2m}$, and let v solve

$$\begin{cases} (-\Delta)^m v = f \text{ in } B_R(x_0), \\ v = \Delta^2 v = \dots = \Delta^{m-1} v = 0 \text{ on } \partial B_R(x_0) \end{cases}$$

Then, for any $p \in (0, \frac{\gamma_m}{\|f\|_{L^1(B_R(x_0))}})$, we have $e^{2mp|v|} \in L^1(B_R(x_0))$ and

$$\int_{B_R(x_0)} e^{2mp|v|} dx \leq C(p)R^{2m},$$

where $\gamma_m = \frac{(2m-1)!}{2} |S^{2m}|$.

Finally with the help of above theorem we prove a regularity result for the polyharmonic operator.

Theorem 4.5. Let $a(x)$ and f satisfies the properties as in $(H1')$ – $(H3')$ and also assume that f be a sub-exponential type function. Let u be a solution $(P_{0,0,\dots,0}^1)$ with $u = \Delta u = \dots = \Delta^{m-1} u = 0$ on $\partial\Omega$. Then $u \in C^{2m-1,\gamma'}(\Omega)$, for all $\gamma' \in (0, 1)$.

Acknowledgement : Dhanya.R was supported by UGC under Dr.D.S Kothari Postdoctoral fellowship scheme No.F.4-2/2006(BSR)/13-1045. Both the authors would like to thank Prof. S. Prashanth for various useful discussions.

References

- [1] Bidaut-Véron, Marie Françoise; Yarur, Cecilia; Semilinear elliptic equations and systems with measure data: existence and a priori estimates. Adv. Differential Equations 7 (2002), no. 3, 257-296.
- [2] H. Brezis and P.L Lions, A Note on Isolated Singularities for Linear Elliptic Equations, Mathematical Analysis and Applications, Part A Advances in Mathematics Supplementary Studies, Vol. 7A,263-266.
- [3] H. Brezis and Frank Merle, Uniform Estimates and blow-up behaviour for solutions of $-\Delta u = V(x)e^u$ in two dimensions, Communication in Partial Differential Equations, 16,(8 and 9), 1223-1253 (1991).
- [4] Caristi, Gabriella; Mitidieri, Enzo; Soranzo, Ramo;n Isolated singularities of polyharmonic equations. Dedicated to Prof. C. Vinti (Italian) (Perugia, 1996). Atti Sem. Mat. Fis. Univ. Modena 46 (1998), suppl., 257-294.

- [5] Dhanya, R.; Giacomoni, J.; Prashanth, S., Isolated singularities for the exponential type semi-linear elliptic equation in \mathbb{R}^2 . Proc. Amer. Math. Soc. 137 (2009), no. 12, 4099-4107.
- [6] García-Huidobro, Marta; Yarur, Cecilia; Existence of singular solutions for a Dirichlet problem containing a Dirac mass. Nonlinear Anal. 74 (2011), no. 8, 2831-2843.
- [7] F. Gazzola, H.-C. Grunau, G. Sweers, Polyharmonic Boundary Value Problems, 1st edition, Lecture Notes in Math., vol. 1991, Springer, 2010.
- [8] Ghergu, Marius; Moradifam, Amir; Taliaferro, Steven D., Isolated singularities of polyharmonic inequalities. J. Funct. Anal. 261 (2011), no. 3, 660-680.
- [9] C.S. Lin, A classification of solutions of a conformally invariant fourth order equation in \mathbb{R}^n . Comment. Math. Helv. 73 (1998), no. 2, 206-231.
- [10] P.L. Lions, Isolated Singularities in Semilinear Problems ,Journal of Differential Equations 38, 441-450(1980).
- [11] Luca Martinazzi, Concentration-compactness phenomena in the higher order Liouville's equation, Journal of Functional Analysis 256(2009), 3743-3741.
- [12] Orsina, Luigi; Ponce, Augusto C., Semilinear elliptic equations and systems with diffuse measures. J. Evol. Equ. 8 (2008), no. 4, 781-812.
- [13] Simader, Christian G., Mean value formulas, Weyl's lemma and Liouville theorems for Δ^2 and Stokes' system. Results Math. 22 (1992), no. 3-4, 761-780.
- [14] Soranzo, Ramon Isolated singularities of positive solutions of a superlinear biharmonic equation. Potential Anal. 6 (1997), no. 1, 57-85.
- [15] Taliaferro, Steven D., On the growth of superharmonic functions near an isolated singularity. I. J. Differential Equations 158 (1999), no. 1, 28-47.
- [16] Taliaferro, Steven D., On the growth of superharmonic functions near an isolated singularity. II. Comm. Partial Differential Equations 26 (2001), no. 5-6, 1003-1026.