

UNBOUNDED ROUGH DRIVERS

ISMAEL BAILLEUL AND MASSIMILIANO GUBINELLI

ABSTRACT. We propose a theory of linear differential equations driven by unbounded operator-valued rough signals. As an application we consider rough linear transport equations and more general linear hyperbolic symmetric systems of equations driven by time-dependent vector fields which are only distributions in the time direction.

1. INTRODUCTION

In this paper we would like to start a program of developing a general theory of rough PDEs aiming at extending classical PDE tools such as weak solutions, a priori estimates, compactness results, duality. This is a quite unexplored territory where few tools are available, so as a start, we will content ourselves in this work with the study of linear symmetric hyperbolic systems of the form

$$(1.1) \quad \partial_t f + a \nabla f = 0,$$

where f is an \mathbb{R}^N -valued space-time distribution on $\mathbb{R}_+ \times \mathbb{R}^d$, and $a : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow L(\mathbb{R}^d, \mathbb{R}^{N \times N})$ is a $N \times N$ matrix-valued family of time-dependent vector fields in \mathbb{R}^d . This setting includes as a particular case scalar transport equations. Moreover we restrict our attention to the case where the matrix-valued vector field a is only a distribution in the time variable, rather than a regular bounded function. We however retain some smoothness assumption in the space variable, as expected from the fact that general transport equations do not possess the regularisation properties needed to drive them with space-time irregular signals. Even in the classical setting it is known that non-regular coefficients can give rise to non-uniqueness of weak solutions [15].

When a is only a distribution in the time variable the above weak formulation is not available since in the classical setting solutions are considered in spaces like $C([0, T], \mathbb{L}^2(\mathbb{R}^d))$ for general symmetric systems or $L^\infty(\mathbb{R} \times \mathbb{R}^d)$ for scalar transport equation. In this case, the product $a \nabla f$ is not well-defined, not even in a distributional setting. Rough paths have their origin in the need to handle such difficulties in the case of ordinary differential equations driven by distribution valued signals [36, 38, 37, 20]. Controlled rough paths have been introduced in [22] as a setting for considering more general problems; they were used successfully in the study of some stochastic partial differential equations [3, 25, 11, 29, 23, 30, 31], including the remarkable solution by Hairer of the KPZ equation [27].

These developments have ultimately lead to the notion of paracontrolled distributions introduced by Gubinelli, Imkeller and Perkowski [24] and to Hairer's general theory of regularity structures [28], providing a framework for the analysis of non-linear operations on distributions. Despite their successes, these new tools and methods are somehow designed to deal with a prescribed class of singular PDEs which is far from exhausting the set of all interesting singular PDEs.

PDEs with irregular signals have been studied using directly rough path methods also by Friz and co-authors [4, 5, 18, 19, 16]. They have developed an approach to some fully non-linear PDEs and conservation laws driven by rough signals by interpreting these equations as transformations of classical PDEs, generalising the method of characteristics. Subsequently a combination of rough path stability and PDE stability allows to go from smooth signal to the wider class of signals described by rough paths. Entropy solutions to scalar conservation laws with rough drivers have also been analysed also by P.-L. Lions, Souganidis and coauthors [34, 35, 21]. A major drawback of this otherwise effective approach is that there is no intrinsic notion of solution to the PDE and that the study of the properties of the PDE has to be done on a global level.

In recent works intrinsic notions of weak solution to rough PDEs have been proposed by Tindel, Gubinelli and Torecilla [26] for viscosity solutions, Catellier [7] for weak solutions to linear transport equations (see also Hu and Le [32] for classical solutions to transport equations) and more recently by Diehl, Friz and Stannat [12] for a general class of parabolic equations. All these notions are based on a weak formulation of the equation where the irregularity of some data is taken into account via the framework of controlled paths introduced in [22]. However in all these papers explicit formulas involving the flow of rough characteristics play an important role, and this sets apart the study of rough PDEs with respect to the study of weak solutions to more regular PDEs. One of the main motivations of our investigations is an effort to understanding what kind of robust arguments can be put forward in the study a priori regularity of distributions satisfying certain rough PDEs, with the latter formulated, at least at this level, in the language of controlled path. Extensions to regularity structures or paracontrolled distributions will be considered in forthcoming work.

We study equation (1.1) by working in the technically easier setting of controlled paths. To motivate our formalism, note that a formal integration of the weak formulation (1.1) over any time interval $[s, t]$, gives an equation of the form

$$f_t = f_s + \int_s^t V_r f_r dr,$$

where $V_r = a_r \nabla$ is a matrix-valued vector-field and $f_r(x) = f(r, x)$, is a convenient notation of the distribution f evaluated at time r , assuming this make sense. An expansion for the time evolution of f is obtained by iterating the above equation, and reads

$$(1.2) \quad f_t = f_s + A_{ts}^1 f_s + A_{ts}^2 f_s + R_{ts},$$

where

$$A_{ts}^1 = \int_s^t V_r dr, \quad \text{and} \quad A_{ts}^2 = \int_s^t \int_s^r V_r V_{r'} dr' dr,$$

are respectively a first order differential operator (that is a vector field) and a second order differential operator, for each $s \leq t$. As a function of (s, t) , they satisfy formally *Chen's relation*

$$(1.3) \quad A_{ts}^2 = A_{tu}^2 + A_{us}^2 + A_{tu}^1 A_{us}^1$$

for all $0 \leq s \leq u \leq t$. It is a key observation of rough path theory that equation (1.2) can be used as a replacement for the differential or integral formulation of equation (1.1) if the remainder term can be shown to be sufficiently small as $t - s$ goes to 0, in a suitable sense.

We shall call a *rough driver* an operator-valued 2-index maps $\mathbf{A}_{ts} = (A_{ts}^1, A_{ts}^2)$ satisfying the operator Chen relation (1.3) and some regularity assumptions. Building on the above

picture, a path with values in some Banach space where the operators A_{ts}^1 and A_{ts}^2 act, will be said to solve the rough linear equation

$$df_s = \mathbf{A}(ds) f_s.$$

if the Taylor expansion (1.2) holds.

There is a complete theory of such equations in the case where the equation is set in a Banach algebra and the operators A_{ts}^1 and A_{ts}^2 are given by left multiplication by some elements of the algebra. It is however natural, in the present PDE setting, to consider also unbounded operators A^1, A^2 , which makes the use of rough paths ideas non-trivial, unless we work in the analytic category or in similar topologies.

We lay out in this work a theory of such rough linear equation driven by unbounded drivers \mathbf{A} , and obtain some a priori estimates that are used to study the well-posedness of some classes of linear symmetric systems in \mathbb{L}^2 and of the rough transport equation in \mathbb{L}^∞ . The major difficulty which has to be overcome is the lack of a Gronwall lemma for rough equations and the main contribution to this paper is to develop suitable a priori estimates on weak controlled solutions that replace the use of Gronwall lemma in classical proofs. Along the way we refine the standard theory of controlled path by introducing weighed norms compatible with the sewing map and by revisiting the theory of linear rough differential equations in the context of bounded drivers.

As a guide for the reader, here is how we have organised our work. Section 2 provides a refined version of the sewing lemma that allows to keep track of the growth in time of the additive function associated with an almost-additive 2-index map. This result is used in section 3 in the proof of the well-posed character of linear differential equations driven by the bounded rough drivers defined there. Unbounded rough drivers are introduced in section 4, where some fundamental a priori estimate is proved. An \mathbb{L}^2 theory of rough linear equations is developed for a class of unbounded drivers, that contains as a particular example the rough linear transport equation. Our main workhorse here is a novel renormalisation lemma obtained from a tensorization argument in the line of the "doubling of variables" method commonly used in the setting of transport equations or conservation laws. A complete \mathbb{L}^∞ theory of rough transport equations is given in section 6.

Notations – We gather here for reference a number of notations that will be used throughout the text.

- We shall denote by E a generic Banach space. Given two Banach spaces E and F , we denote by $L(E, F)$ the set of continuous linear maps from E to F .
- We shall denote by c a constant whose value may change from place to place.
- Given two positive real numbers α, β , we shall write $\alpha \lesssim \beta$ to say that $\alpha \leq c\beta$, for some positive constant c . To indicate that this constant c depends on a parameter λ , we write $\alpha \lesssim_\lambda \beta$.
- Denote by $\|\cdot\|_{\alpha; k}$ the α -Hölder norm of an E_k -valued path, for $k \in \mathbb{Z}$, and by $\|\cdot\|_{\alpha; E}$ the α -Hölder norm of an E -valued path.

2. WEIGHTED NORMS

We introduce some weighted norms that will be useful in getting a priori estimates on the growth of solutions to the linear differential equations studied in section 3. These

norms are modelled on Picard's well-known norms

$$\|f\| := \sup_{t \geq 0} e^{-\lambda^{-1}t} |f(t)|,$$

introduced in the study of ordinary differential equations in order to provide a functional setting where to get well-posedness results on the whole time interval $[0, \infty)$, as a direct consequence of the Banach fixed point theorem, and to get as a consequence a control on the growth of the size of solutions.

Let T be a possibly infinite positive time horizon. As is common in rough paths theory, we shall work with Banach space-valued multi-index maps, mainly 2 and 3-index maps, defined on the simplexes

$$\{(s, t) \in [0, T]^2; s \leq t\} \quad \text{and} \quad \{(s, u, t) \in [0, T]^3; s \leq u \leq t\}.$$

With Picard's norm in mind, we introduce a norm on the set of 2 and 3-index maps which captures both their Hölder size and their growth at infinity. For a fixed non-negative parameter λ , and $t \in \mathbb{R}$, set

$$|t|_\lambda := |t| \wedge \lambda.$$

Given an increasing non-negative function g defined on \mathbb{R}_+ , and a non-negative Hölder exponent γ , we define the (γ, φ) -norm of a 2-index map a , and a 3-index map b , by the formulae

$$\|a\|_{\gamma, g} := \sup_{0 \leq s < t < T} \frac{|a_{ts}|}{g(\lambda^{-1}t) |t - s|_\lambda^\gamma},$$

and

$$\|b\|_{\gamma, g} := \sup_{0 \leq s < u < t < T} \frac{|b_{tus}|}{g(\lambda^{-1}t) |t - s|_\lambda^\gamma};$$

note the following comparison: For $0 \leq \gamma' \leq \gamma$, we have

$$(2.1) \quad \|\cdot\|_{\gamma', g} \leq \lambda^{\gamma - \gamma'} \|\cdot\|_{\gamma, g}.$$

Recall that given an \mathbb{E} -valued 2-index map a , the *sewing lemma* [22, 14] asserts that if the inequality

$$(2.2) \quad |a_{ts} - (a_{tu} + a_{us})| \leq c|t - s|^\zeta$$

holds for all $0 \leq s \leq u \leq t < T$, with $t - s \leq 1$ say, for some exponent $\zeta > 1$ and some positive constant c , then there exists a unique map $A : [0, T) \rightarrow \mathbb{E}$ whose increments $\delta A_{ts} := A_t - A_s$ are well-approximated by a_{ts} , in the sense that

$$|\delta A_{ts} - a_{ts}| \lesssim |t - s|^\zeta,$$

for all $t - s \leq 1$ say. Moreover, if t_i denotes the times of a finite partition π_{ts} of an interval (s, t) , with mesh $|\pi_{ts}|$, we have

$$(2.3) \quad |\delta A_{ts} - \sum a_{t_{i+1}t_i}| \lesssim |t - s| |\pi_{ts}|^{\zeta-1}.$$

The **sewing map** associates to the above 2-index map a the 2-index map

$$\Lambda(a)_{ts} := \delta A_{ts} - a_{ts}.$$

For a given function $g : \mathbb{R} \rightarrow \mathbb{R}$ let

$$G(t) = g(t) + \int_0^t g(r) dr,$$

and write as usual δa_{tus} for $a_{ts} - (a_{tu} + a_{us})$, for any $0 \leq s \leq u \leq t < T$. The following lemma provides an estimate of the weighted norm of $\Lambda(a)$ in terms of the weighted norm of δa ; an estimate for the growth of A follows as a consequence.

LEMMA 1. *There exists a positive constant c_ζ , depending only on ζ , such that*

$$(\Lambda(a))_{\zeta, G} \leq c_\zeta (\delta a)_{\zeta, g}.$$

Remark that two different scales are used in the weighted norms; this is in accordance with the naive guess. If a grows at some rate g , one expects $A = \int da$ to grow at rate $\int g$. Note however that G is not equal to $\int g$.

PROOF – Given $0 \leq s < t$ and $n \geq 1$, set $t_i = s + i2^{-n}(t-s)$, and note that the telescopic sum

$$(\star) := \sum_{i=0}^{2^n-1} a_{t_{i+1}t_i} - a_{ts} = \sum_{k=0}^{n-1} \sum_{\ell=0}^{2^{n-(k+1)}} \left(a_{(\ell+2)2^k(\ell+1)2^k} + a_{(\ell+1)2^k\ell 2^k} - a_{(\ell+2)2^k\ell 2^k} \right)$$

provides a control of the quantity (\star) in terms of δa only. Sending n to infinity, we see by identity (2.3) that there exists a constant c_ζ depending only on ζ , such that

$$|\Lambda(a)_{ts}| \leq c_\zeta |t-s|^\zeta \sup \left\{ \frac{|\delta a_{t'u's'}|}{|t'-s'|^\zeta}; 0 \leq s < s' < u' < t' \leq t \right\}.$$

Now, if $|t-s| \leq \lambda$, we have $|t-s|_\lambda = |t-s|$, so the above inequality implies clearly that we have

$$(2.4) \quad \frac{|\Lambda(a)_{ts}|}{g(\lambda t) |t-s|_\lambda^\zeta} \leq c_\zeta |\delta a|_{\zeta, g}$$

in that case. If now $|t-s| > \lambda$, divide the interval $[s, t]$ into sufficiently many intervals $[t_k, t_{k+1}]$ of equal length no greater than λ , say N sub-intervals. As we have by definition

$$A_v - A_u = a_{vu} + \Lambda(a)_{vu}$$

for all $0 \leq u \leq v < T$, we have, as a result of a telescopic sum,

$$\Lambda(a)_{ts} + a_{ts} = \sum_{k=0}^{N-1} \left(a_{t_{k+1}t_k} + \Lambda(a)_{t_{k+1}t_k} \right).$$

As the choice of times t_k and inequality (2.4) guarantee that

$$\begin{aligned} \sum_{k=0}^{N-1} \left| \Lambda(a)_{t_{k+1}t_k} \right| &\leq c_\zeta \lambda^\zeta |\delta a|_{\zeta, g} \sum_{k=0}^{N-1} g(\lambda^{-1}t_{k+1}) \\ &\leq c_\zeta \lambda^\zeta |\delta a|_{\zeta, g} G(\lambda^{-1}t), \end{aligned}$$

the conclusion follows from the elementary inequality

$$\begin{aligned} \left| a_{ts} - \sum_{k=0}^{N-1} a_{t_{k+1}t_k} \right| &\leq \sum_{k=0}^{N-1} |\delta a_{t_{k+1}t_k s}| \leq \lambda^\zeta |\delta a|_{\zeta, g} \sum_{k=0}^{N-1} g(\lambda^{-1}t_{k+1}) \\ &\leq \lambda^\zeta |\delta a|_{\zeta, g} G(\lambda^{-1}t). \end{aligned}$$

▷

For the particular choice of function $g(t) = e^{\lambda^{-1}t}$, we have $G(t) = (1 + \lambda)e^{\lambda^{-1}t}$, and we shall set for a path $f : [0, T) \rightarrow E$

$$(2.5) \quad (\|f\bullet\|) := \sup_{t \geq 0} e^{-\lambda^{-1}t} |f(t)|,$$

and for a 2-index map a , and a 3-index map b ,

$$(2.6) \quad (\|a\|)_\gamma := \sup_{0 \leq s < t < T} e^{-\lambda^{-1}t} \frac{|a_{ts}|}{|t-s|^\gamma}, \quad \text{and} \quad (\|b\|)_\gamma := \sup_{0 \leq s < u < t < T} e^{-\lambda^{-1}t} \frac{|b_{tus}|}{|t-s|^\gamma}.$$

Note that these norms depend on a choice of parameter λ , which may be tuned on demand. This will be particularly useful in the statement and proof of theorem 3 below, giving the well-posed character of some linear rough differential equation. Note also that we can compare $(\|f\bullet\|)$ and $(\|\delta f\|)_\gamma$, as the inequality

$$(\|f\bullet\|) \leq |f_0| + e^{-1} \lambda^\gamma (\|\delta f\|)_\gamma$$

holds for all $\lambda > 0$. This will be our starting point in the proof of the a priori estimate (3.2) in theorem 3 below.

Last, a 2-index map a such that $\sup_{t-s \leq 1} \frac{|a_{ts}|}{|t-s|^\gamma}$ is finite will be called a γ -**Hölder map**.

3. LINEAR DIFFERENTIAL EQUATIONS WITH BOUNDED ROUGH DRIVERS

Let $(\mathcal{A}, |\cdot|)$ be a Banach algebra with unit $\mathbf{1}_\mathcal{A}$; one may think for instance to the space of continuous linear maps from some Hilbert space to itself, or to the truncated tensor algebra over some Banach space, equipped with a tensor norm and completed for that norm. We introduce in this section a notion of bounded rough driver in the Banach algebra \mathcal{A} , and show that they generate some flows on the algebra.

DEFINITION 2. Let $\frac{1}{3} < \gamma \leq \frac{1}{2}$. A **bounded γ -rough driver in \mathcal{A}** is a pair $\mathbf{A} = (A^1, A^2)$ of \mathcal{A} -valued 2-index maps satisfying Chen's relations

$$(3.1) \quad \delta A^1 = 0, \quad \text{and} \quad \delta A^2_{tus} = A^1_{tu} A^1_{us},$$

and such that A^1 is γ -Hölder and A^2 is 2γ -Hölder. The norm of \mathbf{A} is defined by the formula

$$\|\mathbf{A}\| := \sup_{0 \leq s < t < T; t-s \leq 1} \frac{|A^1_{ts}|}{|t-s|^\gamma} \vee \frac{|A^2_{ts}|}{|t-s|^{2\gamma}}.$$

As an elementary example, think of \mathcal{A} as the truncated tensor algebra $\bigoplus_{i=0}^N (\mathbb{R}^\ell)^{\otimes i}$ over \mathbb{R}^ℓ , for $N \geq 2$, and consider a weak geometric γ -Hölder rough path $\mathbf{X}_{ts} = 1 \oplus X_{ts} \oplus \mathbb{X}_{ts} \in \bigoplus_{i=0}^N (\mathbb{R}^\ell)^{\otimes i}$, with $2 \leq \gamma < 3$. Left multiplication by X_{ts} and \mathbb{X}_{ts} define operators A^1 and A^2 that are the components of a rough driver.

Recall the weighted norms $(\|\cdot\|)$ and $(\|\cdot\|)_\gamma$ defined by identities (2.5) and (2.6), respectively, depend on some parameter λ . The proof of the following well-posedness result for linear rough differential equations driven by bounded γ -rough drivers shows the interest of being flexible on the tuning of λ .

THEOREM 3 (Integration of bounded rough drivers). *Given any initial condition $J_0 \in \mathcal{A}$, there exists a unique γ -Hölder path f_\bullet starting from f_0 , and such that the formula*

$$\delta f_{t,s} - \left(A^1_{t,s} + A^2_{t,s} \right) f_s$$

defines a 3γ -Hölder 2-index map f^\sharp . Moreover, the following estimate holds

$$(3.2) \quad (\|f_\bullet\|) \leq 2 |f_0|$$

for all λ greater than some λ_0 depending only on $\|\mathbf{A}\|$ and γ . When $f_0 = \mathbf{1}_A$, we will use the notation $f_t = e_{t,0}^{\mathbf{A}}$; the flow property

$$e_{t,s}^{\mathbf{A}} = e_{t,u}^{\mathbf{A}} e_{u,s}^{\mathbf{A}},$$

holds for all $0 \leq s \leq u \leq t < T$.

Applied to the above example of rough driver in the truncated tensor product space, this well-posedness result provides a proof of Lyons' extension theorem [36].

PROOF – a) A priori estimate – Let us prove the a priori bound (3.2) first; the uniqueness claim in the theorem follows from this bound and the linear character of the problem. As mentioned above, we start from the inequality

$$\langle f_\bullet \rangle \leq |f_0| + c\lambda^\gamma \langle \delta f \rangle_\gamma,$$

and try and write $\langle \delta f \rangle_\gamma$ in terms of $\langle f_\bullet \rangle$; this can be done as follows.

By using Chen's relation (3.1) and the definition of the remainder f^\sharp , the identity

$$-\delta f_{t,u}^\sharp = A_{t,u}^1 \left(\delta f_{u,s} - A_{u,s}^1 f_s \right) + A_{t,u}^2 \delta f_{u,s} = A_{t,u}^1 \left(A_{t,s}^2 f_s + f_{s,t}^\sharp \right) + A_{t,u}^2 \delta f_{u,s},$$

gives us the estimate

$$\langle \delta f^\sharp \rangle_{3\gamma} \leq \|\mathbf{A}\|^2 \langle f_\bullet \rangle + \|\mathbf{A}\| \langle f^\sharp \rangle_{2\gamma} + \|\mathbf{A}\| \langle \delta f \rangle_\gamma.$$

But since $3\gamma > 1$, we have

$$f^\sharp = \Lambda \delta f^\sharp,$$

so the inequality

$$\langle f^\sharp \rangle_{3\gamma} \lesssim_\gamma \|\mathbf{A}\|^2 \langle f_\bullet \rangle + \|\mathbf{A}\| \langle f^\sharp \rangle_{2\gamma} + \|\mathbf{A}\| \langle \delta f \rangle_\gamma$$

follows from lemma 1. Using the inequality $\langle f^\sharp \rangle_{2\gamma} \leq \lambda^\gamma \langle f^\sharp \rangle_{3\gamma}$, emphasized in (2.1), the above equation gives

$$\langle f^\sharp \rangle_{3\gamma} \lesssim_\gamma \|\mathbf{A}\|^2 \langle f_\bullet \rangle + \lambda^\gamma \|\mathbf{A}\| \langle f^\sharp \rangle_{3\gamma} + \|\mathbf{A}\| \langle \delta f \rangle_\gamma.$$

For λ small enough so that $\lambda^\gamma \|\mathbf{A}\| \leq \frac{1}{2}$, we obtain

$$\langle f^\sharp \rangle_{3\gamma} \lesssim_\gamma \|\mathbf{A}\|^2 \langle f_\bullet \rangle + \|\mathbf{A}\| \langle \delta f \rangle_\gamma,$$

so, using again the definition of the remainder f^\sharp , and the observation that

$$\langle A^2 f \rangle_\gamma \lesssim_\gamma \lambda^\gamma \langle A^2 f \rangle_{2\gamma} \lesssim_\gamma \lambda^\gamma \|\mathbf{A}\| \langle f_\bullet \rangle,$$

we obtain the estimate

$$\begin{aligned} \langle \delta f \rangle_\gamma &\lesssim_\gamma \|\mathbf{A}\| \langle f_\bullet \rangle + \langle A^2 f \rangle_\gamma + \langle f^\sharp \rangle_\gamma \\ &\lesssim_\gamma \left\{ \|\mathbf{A}\| (1 + \lambda^{2\gamma} \|\mathbf{A}\|) + \lambda^\gamma \|\mathbf{A}\| \right\} \langle f_\bullet \rangle + \lambda^{2\gamma} \|\mathbf{A}\| \langle \delta f \rangle_\gamma. \end{aligned}$$

Taking λ small enough, depending only on $\|\mathbf{A}\|$, we eventually see that

$$\langle \delta f \rangle_\gamma \lesssim_\gamma \left\{ \|\mathbf{A}\| (1 + \lambda^{2\gamma} \|\mathbf{A}\|) + \lambda^\gamma \|\mathbf{A}\| \right\} \langle f_\bullet \rangle.$$

The a priori estimate $\langle f_\bullet \rangle \leq 2|f_0|$, follows now from a choice of sufficiently small parameter λ , since $\langle f_\bullet \rangle \leq |f_0| + c\lambda^\gamma \langle \delta f \rangle_\gamma$.

b) Existence – We can run a Picard iteration to prove the existence of a path satisfying the conditions of the theorem. Set first

$$f_t^0 = f_0, \quad \text{and} \quad f_t^1 = A_{t_0}^1 f_0$$

for all $t \in \mathbb{R}$. Given the paths $f_{\bullet}^{n-1}, f_{\bullet}^n$, the 2-index map

$$a_{ts}^n := A_{ts}^1 f_s^n + A_{ts}^2 f_s^{n-1}$$

satisfies the almost-additivity condition (2.2) with $\zeta = 3\gamma > 1$ here, so there is, by the sewing lemma, a unique γ -Hölder path f_{\bullet}^{n+1} for which the formula

$$\delta f_{ts}^{n+1} - \left(A_{ts}^1 f_s^n + A_{ts}^2 f_s^{n-1} \right)$$

defines 2-index 3γ -Hölder map $f^{n+1, \sharp}$. Setting $g_{\bullet}^0 := 0$ and

$$g_{\bullet}^{n+1} := f_{\bullet}^{n+1} - f_{\bullet}^n, \quad g^{n+1, \sharp} := f^{n+1, \sharp} - f^{n, \sharp},$$

for all $n \geq 0$, we have

$$\delta g_{ts}^{n+1} = A_{ts}^1 g_s^n + A_{ts}^2 g_s^{n-1} + g_{ts}^{n+1, \sharp}.$$

Note moreover that we have the identity

$$\begin{aligned} -\delta g_{t,u,s}^{n+1, \sharp} &= A_{t,u}^1 \left(\delta g_{u,s}^n - A_{u,s}^1 g_s^{n-1} \right) + A_{t,u}^2 \delta g_{u,s}^{n-1} \\ &= A_{t,u}^1 \left(A_{t,s}^2 g_s^{n-2} + g_{s,t}^{n, \sharp} \right) + A_{t,u}^2 \delta g_{u,s}^{n-1}, \end{aligned}$$

so, proceeding as in the proof of the a priori bound, we see that the inequality

$$\left(|g_{\bullet}^{n-1}| \right) + \left(|\delta g^n| \right)_{\gamma} + \left(|g^{n+1, \sharp}| \right)_{3\gamma} \lesssim_{\gamma, \|\mathbf{A}\|} \lambda^{\gamma} \left\{ \left(|g_{\bullet}^{n-2}| \right) + \left(|\delta g^{n-1}| \right)_{\gamma} + \left(|g^{n, \sharp}| \right)_{3\gamma} \right\}$$

holds, by choosing λ small enough. The estimate

$$\left(|g_{\bullet}^{n-1}| \right) + \left(|\delta g^n| \right)_{\gamma} + \left(|g^{n+1, \sharp}| \right)_{3\gamma} \lesssim_{\gamma, \|\mathbf{A}\|} \lambda^{\gamma n}$$

follows as a consequence, so the series $f^n = f^0 + \sum_{n \geq 1} g^n$ converges in the complete space of \mathcal{A} -valued γ -Hölder paths, and defines a path in \mathcal{A} satisfying the conditions of the theorem. \triangleright

REMARKS 4. (1) *Note that the proof given above gives back the known sharp growth rate $\exp((2\|\mathbf{A}\|)^{\gamma} t)$ for $|f_t|$; see [17]. Bounded rough drivers can also be integrated by defining recursively the $(n\gamma)$ -Hölder \mathcal{A} -valued 2-index map A^n using the formula*

$$\delta A_{t,u,s}^n = \sum_{k=1}^{n-1} A_{t,u}^{n-k} A_{u,s}^k,$$

and setting

$$e_{t,s}^{\mathbf{A}} = \sum_{n=0}^{\infty} A_{t,s}^n.$$

Standard estimates on the sewing map [22] show that δA^n has $n\gamma$ -Hölder norm no greater than $(n!)^{-\gamma}$, so the above series converges in \mathcal{A} for all $0 \leq s \leq t$. The flow property is obtained by a direct calculation, and setting $f_t := e_{t,0}^{\mathbf{A}} f_0$, we see that the path f_{\bullet} solves the problem.

(2) **Linear rough differential equations with a linear drift** – *The above theory extends easily to rough equations of the form*

$$(3.3) \quad \delta f_{t,s} = \int_s^t B_r f_r dr + A_{t,s}^1 f_s + A_{t,s}^2 f_s + f_{t,s}^{\natural}$$

where $B \in \mathbb{L}^\infty(\mathbb{R}; \mathcal{A})$ is a bounded measurable family of bounded operators. This equation is the rigorous meaning to give to solutions of the differential equation

$$\frac{d}{dt}f = (B_r + \dot{A}_r)f_r$$

where $\dot{A}_t = \partial_t A_{t0}^1$. In case $\dot{A}_t \in \mathbb{L}^\infty(\mathbb{R}; \mathcal{A})$, the two formulations are equivalent provided $\mathbf{A} = (A^1, A^2)$ is defined by the formula

$$A_{t,s}^1 = \int_s^t \dot{A}_r dr, \quad A_{t,s}^2 = \int_s^t \int_s^r \dot{A}_u \dot{A}_r dudr.$$

The proof of theorem 3 can be easily adapted in the present setting, and the final lower bound on λ gets an additional dependence on $\|B\|_\infty$. It yields moreover the following Duhamel formula

$$f_t = e_{t,0}^{\mathbf{A}} f_0 + \int_0^t e_{t,r}^{\mathbf{A}} B_r f_r dr.$$

Indeed, let f_\bullet be a function satisfying the above identity. If one computes the increment of the right hand side in the above equation, we get

$$\delta f_{t,s} = \int_s^t e_{t,r}^{\mathbf{A}} B_r f_r dr - (e_{t,s}^{\mathbf{A}} - \text{Id}) f_s = \int_s^t B_r f_r dr + A_{t,s}^1 f_s + A_{t,s}^2 f_s + f_{t,s}^\sharp$$

where

$$f_{s,t}^\sharp = \int_s^t (e_{t,r}^{\mathbf{A}} - \text{Id}) B_r f_r dr + (e_{t,s}^{\mathbf{A}} - \text{Id} - A_{t,s}^1 - A_{t,s}^2) f_s.$$

Using the bounds

$$\left| e_{t,r}^{\mathbf{A}} - \text{Id} \right| \lesssim_{\|\mathbf{A}\|} |t-r|^\gamma, \quad \text{and} \quad \left| e_{t,s}^{\mathbf{A}} - \text{Id} - A_{t,s}^1 - A_{t,s}^2 \right| \lesssim_{\|\mathbf{A}\|} |t-s|^{3\gamma},$$

this allows to conclude that $|f_{t,s}^\sharp| \lesssim |t-s|^{3\gamma}$, and that the path f_\bullet is indeed the unique solution to equation (3.3).

- (3) Bounded rough drivers have also been introduced previously in the work [8] of Coutin and Lejay, and studied in relation with the Magnus formula for what is called there the resolvent operator $e^{\mathbf{A}}$. The above short proof of theorem 3 can be considered an alternative proof of the main result of section 3 in [8]. They also consider perturbed linear equations, with an a priori given drift of the more general form C_{ts} , rather than $\int_s^t B_r f_r dr$, with C satisfying some regularity conditions.

4. UNBOUNDED ROUGH DRIVERS AND ROUGH LINEAR EQUATIONS

The above results apply in the particular case where \mathcal{A} is the Banach algebra of bounded operators on an Hilbert space H . We shall study in the remaining sections the integration problem

$$(4.1) \quad \delta f_{ts} = (A_{ts}^1 + A_{ts}^2) f_s + f_{ts}^\sharp,$$

for a particular class of drivers \mathbf{A} associated to a class of *unbounded* operators on H , or other Banach spaces, with in mind the model case of the **rough transport equation**

$$\delta f(\varphi)_{ts} = X_{ts} f_s(V^* \varphi) + \mathbb{X}_{ts} f_s(V^* V^* \varphi) + f_{ts}^\sharp(\varphi),$$

where $\mathbf{X} = (X, \mathbb{X})$ is an ℓ -dimensional γ -Hölder rough path and $V = (V_1, \dots, V_\ell)$ is a collection of ℓ vector fields on \mathbb{R}^d .

4.1. Rough drivers. To make sense of this equation we need to complete the functional setting by the datum of a scale of Banach spaces $(E_n, |\cdot|_n)_{n \geq 0}$, with E_{n+1} continuously embedded in E_n . For $n \geq 0$, we shall denote by $E_{-n} = E_n^*$ the dual space of E_n , equipped with its natural norm,

$$|e|_{-n} := \sup_{\varphi \in E_n, |\varphi|_n \leq 1} (\varphi, e), \quad e \in E_{-n}.$$

We require that the following continuous inclusions

$$E_n \subset \dots \subset E_2 \subset E_1 \subset E_0$$

hold for all $n \geq 2$. One can think of n as quantifying the 'regularity' of elements of some test functions, with the elements of E_n being more regular with n increasing. Denote by $\|\cdot\|_{(b,a)}$ for the norm of a linear operator from E_a to E_b . (Note that we use (b, a) and not (a, b) in the lower index for the norm.) We also assume the existence of a family $(J^\varepsilon)_{0 < \varepsilon \leq 1}$ of operators from E_0 to itself such that the estimates

$$(4.2) \quad \|J^\varepsilon - \text{Id}\|_{(n+k,n)} \leq c\varepsilon^k, \quad \|J^\varepsilon\|_{n+k} \leq c\varepsilon^{-k}$$

hold for all $n, k \geq 0$, for some positive constant c independent of ε . For $\varphi \in E_0$, the elements $\varphi_\varepsilon := J^\varepsilon \varphi$ are in particular 'smooth', that is in the intersection of all the spaces E_n , for $n \geq 0$. One can think for instance to scales of Sobolev spaces where smoothing operators can be used to construct explicitly the approximating family.

DEFINITION 5. Let $\frac{1}{3} < \gamma \leq \frac{1}{2}$ be given. An **unbounded γ -rough driver on the scales** $(E_n, |\cdot|_n)_{n \geq 0}$, is a pair $\mathbf{A} = (A^1, A^2)$ of 2-index maps, with

$$(4.3) \quad \begin{aligned} A_{ts}^1 &\in \text{L}(E_n, E_{n-1}), \text{ for } n \in \{-0, -2\}, \\ A_{ts}^2 &\in \text{L}(E_n, E_{n-2}), \text{ for } n \in \{-0, -1\}, \end{aligned}$$

for all $0 \leq s \leq t < T$, which satisfy Chen's relations (3.1), and such that A^1 is γ -Hölder and A^2 is 2γ -Hölder.

The forthcoming equation (4.5) will make it clear that unbounded rough drivers can be thought of a some kind of (dual objects to some) multi-scales velocity fields, with two time scales given by A^1 and A^2 . This is particularly clear on the following elementary example, where

$$A_{ts}^1 = \sum_{i=1}^{\ell} X_{ts}^i V_i, \quad \text{and} \quad A_{ts}^2 = \sum_{j,k=1}^{\ell} \mathbb{X}_{ts}^{jk} V_j V_k,$$

where $\mathbf{X} = (X, \mathbb{X})$ is a weak geometric γ -Hölder rough path over \mathbb{R}^ℓ , with $\frac{1}{3} < \gamma \leq \frac{1}{2}$, and V_1, \dots, V_ℓ are regular enough vector fields on \mathbb{R}^d with values in $N \times N$ matrices, understood here as first order differential operators. One can also take $V_i^* = -V_i - \text{div} V_i$ in the above definition of a rough driver.

In a probabilistic setting, the above rough path \mathbf{X} could be the (Stratonovich) Brownian rough path. More generally, one could take as first level A_{ts}^1 a semimartingale velocity field of Le Jan-Watanabe-Kunita type (or its dual), where noise (the above X_{ts}) and dynamics (the above V_i) cannot be separated from one another. It is shown in the forthcoming work [2] that these velocity fields can be lifted into a(n object very similar to a) rough driver

under some mild conditions on the semimartingale structure. In the same spirit, Catellier has shown in [6, 7] the interest for the study of rough transport equations with irregular drift, of considering velocity fields in \mathbb{R}^d given by the regularisation of some field V (that may even be a distribution) along some irregular path w , such as a typical trajectory of a fractional Brownian motion with any Hurst index. One deals in this particular setting with the 'Young analogue' of the above γ -rough drivers, corresponding to $\frac{1}{2} < \gamma \leq 1$, with only one level, and given in this example by the formula

$$A_{ts}^1(x) = \int_s^t V(x + w_u) du.$$

We denote by $\epsilon(e)$ the duality pairing between an element ϵ of E_{-n} and an element e of E_n , for any $n \in \mathbb{Z}$. We can now make sense of equation (4.1).

DEFINITION 6. *An E_{-0} -valued path f_\bullet is said to solve the linear rough differential equation*

$$(4.4) \quad df_s = \mathbf{A}(ds)f_s$$

if there exists an E_{-2} -valued 2-index map f^\sharp such that one has

$$(4.5) \quad \delta f_{ts}(\varphi) = f_s\left(\{(A_{ts}^1)^* + (A_{ts}^2)^*\}\varphi\right) + f_{ts}^\sharp(\varphi)$$

for all $0 \leq s \leq t < T$ and all $\varphi \in E_2$, and the map $f_{ts}^\sharp(\varphi)$ is 3γ -Hölder, for each $\varphi \in E_2$.

Let us insist on the fact that this notion of solution depends on the scale $(E_n)_{n \geq 0}$. We define, for all $0 \leq s \leq t < T$, an E_{-1} -valued 2-index map f^b setting, for $\varphi \in E_1$,

$$f_{ts}^b(\varphi) := \delta f_{ts}(\varphi) - f_s((A_{ts}^1)^*\varphi);$$

for $\varphi \in E_2$, one also has another expression for that quantity

$$f_{ts}^b(\varphi) = f_s((A_{ts}^2)^*\varphi) + f_{ts}^\sharp(\varphi).$$

4.2. A priori estimates. We show in this section that solutions f_\bullet of equation (4.4) satisfy some a priori bounds that depend only on certain norms on the rough driver A and on the uniform norm of f , when seen as a continuous path with values in some spaces of distributions.

The next lemma shows that the map f^b is actually 2γ -Hölder with values in a space of distributions "of order 2" in a certain sense while f^\sharp is only expected to be γ -Hölder from its definition. This *gain of time regularity, traded against a loss of 'space regularity'* may well be one of our main contribution, despite its elementary nature. It leads to a similar (and even better) result for f^\sharp , as expressed in theorem 8, that is the key technical result of this paper and which opens the road to a quite complete theory of linear rough equation.

Recall the notations $\|\cdot\|$ and $\|\cdot\|_\gamma$ for the norms introduced in equations (2.5) and (2.6). They depend implicitly on some positive parameter λ that we shall tune as needed along the way. Given $\varphi \in E_0$, set

$$K_1(\varphi) := \sup_{0 < \varepsilon \leq 1} \left\{ \|f((A^1)^*J^\varepsilon\varphi)\|_\gamma + \lambda^\gamma \varepsilon \|f((A^2)^*J^\varepsilon\varphi)\|_{2\gamma} \right. \\ \left. + \lambda^{2\gamma} \varepsilon^2 \|f^\sharp(J^\varepsilon\varphi)\|_{3\gamma} + \lambda^{-\gamma} \varepsilon^{-1} \|f((\text{Id} - J^\varepsilon)\varphi)\| \right\}$$

and

$$K_2(\varphi) := \sup_{0 < \varepsilon \leq 1} \left\{ \|f((A^2)^*\varphi)\|_{2\gamma} + \lambda^\gamma \varepsilon \|f^\sharp(J^\varepsilon\varphi)\|_{3\gamma} + \lambda^{-2\gamma} \varepsilon^{-2} \|f((\text{Id} - J^\varepsilon)\varphi)\| + \lambda^{-\gamma} \varepsilon^{-1} \|f((A^1)^*(\text{Id} - J^\varepsilon)\varphi)\|_\gamma \right\}$$

LEMMA 7. *We have $\|\delta f(\varphi)\|_\gamma \lesssim K_1(\varphi)$, and $\|f^\flat(\varphi)\|_{2\gamma} \lesssim K_2(\varphi)$.*

PROOF – The strategy is simple and consists, for $\varphi \in E_1$, in writing

$$|\delta f_{ts}(\varphi)| \leq \left| \delta f_{ts}(J^\varepsilon\varphi) \right| + \left| \delta f_{ts}((\text{Id} - J^\varepsilon)\varphi) \right|.$$

The second term above is bounded above by

$$\left| \delta f_{ts}((\text{Id} - J^\varepsilon)\varphi) \right| \leq \varepsilon |t - s|_\lambda^\gamma K_1(\varphi),$$

while the first term can be estimated using properties (4.2) and the defining identity (4.5); this will give us an upper bound for $|\delta f_{ts}(\varphi)|$ depending on ε , over which we shall optimise. Recall that the elements φ_ε are 'smooth', that is, they belong to all the spaces E_n , for $n \geq 0$.

- From the previous discussion, we have

$$\left| \delta f_{ts}(\varphi) \right| \leq \left(|t - s|_\lambda^\gamma + \lambda^{-\gamma} \varepsilon^{-1} |t - s|_\lambda^{2\gamma} + \lambda^{-2\gamma} \varepsilon^{-2} |t - s|_\lambda^{3\gamma} + \varepsilon \lambda^\gamma \right) K_1(\varphi).$$

By choosing $\varepsilon = \lambda^{-\gamma} |t - s|_\lambda^\gamma \in (0, 1]$, we obtain the upper bound on $\|\delta f(\varphi)\|_\gamma$ of the lemma.

- One can proceed in a similar way to estimate $|f_{ts}^\flat(\varphi)|$, starting from the inequality

$$|f_{ts}^\flat(\psi)| \leq |\delta f_{ts}(\psi)| + \left| f_s((A_{ts}^1)^*\psi) \right| + \left| f_s((A_{ts}^2)^*\psi) \right|$$

available for all $\psi \in E_2$. We get as a result the estimate

$$\begin{aligned} |f_{ts}^\flat(\varphi)| &\leq \left| f_s((A_{ts}^2)^*\varphi) \right| + \left| f_{ts}^\sharp(J^\varepsilon\varphi) \right| + \left| f_{ts}^\sharp((\text{Id} - J^\varepsilon)\varphi) \right| \\ &\leq K_2(\varphi) \left(|t - s|_\lambda^{2\gamma} + \varepsilon^{-1} |t - s|_\lambda^{3\gamma} + \varepsilon^2 + \varepsilon^{-1} |t - s|_\lambda^\gamma \right). \end{aligned}$$

Taking again $\varepsilon = \lambda^{-\gamma} |t - s|_\lambda^\gamma$, we get the upper bound on $|f_{ts}^\flat(\varphi)|$. \triangleright

The estimates proved in lemma 7 are all we need to give an upper bound on the 3γ -Hölder norm of f^\sharp in terms of f itself. As this result will be our workhorse for proving a number of results in different situations, we formulate here the result in some generality. Given a Banach space B and a B -valued path m_\bullet , we set

$$\|m\|_{\gamma; B} := \sup_{0 \leq s < t \leq T} e^{-\lambda^{-1}t} \frac{|m_{ts}|_B}{|t - s|_\lambda^\gamma},$$

for $\gamma \geq 0$. Set

$$N_1(\mathbf{A}) := \sup_{0 < \varepsilon \leq 1} \left\{ \varepsilon \|J^\varepsilon(A^1)^*\|_{\gamma; L(F, F)} + \varepsilon^2 \|J^\varepsilon(A^2)^*\|_{2\gamma; L(F, F)} \right\}$$

and

$$N_2(\mathbf{A}) := \sup_{0 < \varepsilon \leq 1} \left\{ \begin{aligned} & \left\| (A^1)^* J^\varepsilon (A^2)^* \right\|_{3\gamma; L(F, E)} + \varepsilon \left\| (A^2)^* J^\varepsilon (A^2)^* \right\|_{4\gamma; L(F, E)} \\ & + \varepsilon^{-1} \left\| (\text{Id} - J^\varepsilon) (A^2)^* \right\|_{2\gamma; L(F, E)} + \left\| (A^2)^* (A^1)^* \right\|_{3\gamma; L(F, E)} \\ & + \varepsilon^{-2} \left\| (\text{Id} - J^\varepsilon) (A^1)^* \right\|_{\gamma; L(F, E)} + \varepsilon^{-1} \left\| (A^1)^* (\text{Id} - J^\varepsilon) (A^1)^* \right\|_{2\gamma; L(F, E)} \end{aligned} \right\}$$

THEOREM 8. *Let $E \subset E_0$ and $F \subset E_3$ be two Banach spaces of distributions such that the above quantities $N_1(\mathbf{A})$ and $N_2(\mathbf{A})$ are finite. Let $\lambda \leq 1$ be sufficiently small so that $2\lambda^\gamma N_1(\mathbf{A}) \leq 1$. Then any solution of the rough linear equation (4.4) satisfies the a priori bound*

$$\|f^\sharp\|_{3\gamma; F^*} \lesssim \lambda^{-2\gamma} N_2(\mathbf{A}) \|f\|_{E^*}.$$

PROOF – Given $\varphi \in E_3$, note that

$$\delta f_{tus} = f_{us}^b((A_{tu}^1)^* \varphi) + \delta f_{us}((A_{tu}^2)^* \varphi),$$

is 3γ -Hölder by lemma 7, so the sewing lemma 1 can be used, giving

$$\begin{aligned} \|f^\sharp(\varphi)\|_{3\gamma} & \lesssim \|f^b((A^1)^* \varphi)\|_{3\gamma} + \|\delta f((A^2)^* \varphi)\|_{3\gamma} \\ & \lesssim \left\| K_2((A^1)^* \varphi) \right\|_{\gamma} + \left\| K_1((A^2)^* \varphi) \right\|_{2\gamma} \\ & \lesssim \lambda^{-2\gamma} K(\varphi) + \sup_{0 < \varepsilon \leq 1} \left\{ \lambda^\gamma \varepsilon \|f^\sharp(J^\varepsilon(A^1)^* \varphi)\|_{(3\gamma, \gamma)} + \lambda^{2\gamma} \varepsilon^2 \|f^\sharp(J^\varepsilon(A^2)^* \varphi)\|_{(3\gamma, 2\gamma)} \right\}, \end{aligned}$$

where $\|f^\sharp(m)\|_{(3\gamma, \gamma)}$ stands for the γ -Hölder norm of the function $\|f^\sharp(m_\bullet)\|_{3\gamma}$, with two different maps m above, and where we set

$$\begin{aligned} K(\varphi) & := \sup_{0 < \varepsilon \leq 1} \left\| f((A^1)^* J^\varepsilon (A^2)^* \varphi) \right\|_{3\gamma} + \varepsilon \left\| f((A^2)^* J_\varepsilon (A^2)^* \varphi) \right\|_{4\gamma} \\ & + \varepsilon^{-1} \left\| f((\text{Id} - J^\varepsilon) (A^2)^* \varphi) \right\|_{2\gamma} + \left\| f((A^2)^* (A^1)^* \varphi) \right\|_{3\gamma} \\ & + \varepsilon^{-2} \left\| f((\text{Id} - J^\varepsilon) (A^1)^* \varphi) \right\|_{\gamma} + \varepsilon^{-1} \left\| f((A^1)^* (\text{Id} - J^\varepsilon) (A^1)^* \varphi) \right\|_{2\gamma}; \end{aligned}$$

we assumed here $\lambda \leq 1$, with no loss of generality, to get a simpler expression for the formula. Subsequently, by maximizing over $\varphi \in F$, we get

$$\|f^\sharp\|_{3\gamma; F^*} \lesssim \lambda^{-2\gamma} \sup_{\varphi \in F} \frac{K(\varphi)}{\|\varphi\|_F} + \lambda^\gamma N_1(\mathbf{A}) \|f^\sharp\|_{3\gamma; F^*}.$$

Now, taking $\lambda \leq 1$ small enough so that $2\lambda^\gamma N_1(\mathbf{A}) \leq 1$, we obtain the awaited inequality

$$\|f^\sharp\|_{3\gamma; F^*} \lesssim 2\lambda^{-2\gamma} \sup_{\varphi \in F} \frac{K(\varphi)}{\|\varphi\|_F} \lesssim \lambda^{-2\gamma} N_2(\mathbf{A}) \|f\|_{E^*} \lesssim \lambda^{-2\gamma} N_2(\mathbf{A}) \|f\|_{E^*}.$$

▷

Now, if one takes

$$E = E_0, \quad \text{and} \quad F = E_3,$$

and one assumes that

$$(A_{ts}^1)^* \in L(E_1, E_0) \cap L(E_3, E_2)$$

and

$$(A_{ts}^2)^* \in L(E_3, E_1) \cap L(E_2, E_0)$$

for all $0 \leq s \leq t \leq T$, then

$$N_1(\mathbf{A}), N_2(\mathbf{A}) \lesssim C_0$$

where

$$(4.6) \quad C_0 := \|A^1\|_{\gamma; (-0, -1)} + \|A^2\|_{2\gamma; (-0, -2)} + \|A^1\|_{\gamma; (-2, -3)} + \|A^2\|_{2\gamma; (-1, -3)} < \infty$$

Note that it follows from Banach uniform boundedness principle that for a solution path (f, f^\sharp) , we have

$$\|f^\sharp\|_{3\gamma; -3} \leq \|f^\sharp\|_{3\gamma; -2} < \infty.$$

Theorem 8 shows that one has an explicit upper bound for $\|f_{ts}^\sharp\|_{3\gamma; -3}$ in terms of $(f)_{-0}$ and C_0 only.

THEOREM 9. *For any $\lambda \leq 1$ such that $\lambda^\gamma C_0 \lesssim 1$ we have the a priori bounds*

$$(4.7) \quad \max \left\{ (\delta f)_{\gamma; -1}, (f^\flat)_{2\gamma; -2}, (f^\sharp)_{3\gamma; -3} \right\} \lesssim_\gamma 2C_0 \lambda^{-2\gamma} (f)_{-0}.$$

5. INTEGRATION OF UNBOUNDED ROUGH DRIVERS IN HILBERT SPACES

We develop in this section the theory of integration of unbounded rough drivers in the Hilbert space $\mathbb{L}^2(\mathbb{R}^d) = E_0 = E_{-0}$. We are able to give a rather complete theory a class of drivers that enjoys the same algebraic properties as the rough drivers $\mathbf{A} = (XV, \mathbb{X}VV)$ involved in the rough transport equation, when the vector fields $V = (V_1, \dots, V_\ell)$ are divergence free. These drivers are called conservative. A general existence result for the rough linear equation $df_s = \mathbf{A}(ds)f_s$, driven by conservative drivers is given in section 5.1. To prove uniqueness of solutions to such equations, we develop in section (5.2) a robust tensorization argument for a larger class of unbounded rough drivers that is the key to obtain some a priori bounds. These bounds imply uniqueness for rough linear equations driven by conservative drivers under a mild additional assumption, but they also lead to a complete \mathbb{L}^2 -theory of rough transport equations, as illustrated in section (5.4).

Note that working in a Hilbert space setting, we have the continuous inclusions

$$(5.1) \quad E_n \subset \dots \subset E_1 \subset E_0 = E_{-0} \subset E_{-1} \subset \dots \subset E_{-n}.$$

5.1. Conservative drivers. We start with the simple situation where the driver is *conservative* according to the following definition.

DEFINITION 10. *A rough driver is said to be **conservative** if we have*

- (i) $(A_{ts}^1)^* + A_{ts}^1 = 0$, on E_1 ,
- (ii) $(A_{ts}^2)^* + A_{ts}^2 + (A_{ts}^1)^* A_{ts}^1 = 0$, on E_2 ,

for all $0 \leq s \leq t < T$.

Notice that the conservative conditions make sense because of the above continuous inclusions (5.1). As an elementary example of conservative unbounded driver, take as Banach spaces E_n the Sobolev spaces $W^{n,2}(\mathbb{R}^d)$, with norm $\|\varphi\|_n := \sum_{k=0}^n \|\nabla^k \varphi\|_0$, and consider the unbounded rough driver \mathbf{A} given by the formula

$$(5.2) \quad A_{ts}^1 = X_{ts}^i V_i, \quad A_{ts}^2 = \mathbb{X}_{ts}^{jk} V_j V_k,$$

for some ℓ -dimensional geometric γ -rough path, with $\frac{1}{3} < \gamma \leq 1$, and some *divergence-free* vector fields $(V_i)_{i=1..l}$ on \mathbb{R}^d , with the latter understood as first order differential operators.

Condition **(ii)** partly plays the role in our setting that the notion of weak geometric rough path plays in a rough paths setting. Also, the antisymmetry condition **(i)** holds due to the fact that V_i have null divergence, and condition **(ii)** holds as a consequence of the weak geometric character of \mathbf{X} . Indeed, in this setting we have

$$\begin{aligned} (A_{ts}^1)^* &= X_{ts}^i (V_i)^* = -X_{ts}^i V_i = -A_{ts}^1, \\ (A_{ts}^2)^* + A_{ts}^2 &= \mathbb{X}_{ts}^{jk} (V_k^* V_j^* + V_j V_k) = \frac{1}{2} X_{ts}^j X_{ts}^k (V_k V_j + V_j V_k) = -(A_{ts}^1)^* A_{ts}^1 \end{aligned}$$

on E_1 and E_2 respectively. The boundedness assumptions (4.3) that A_{ts}^1 and A_{ts}^2 need to satisfy hold if, for instance, the vector fields V_i are \mathcal{C}_b^2 . (Note that our setting is by no means restricted to working with vector fields. Working in the spaces $E_n = W^{kn,2}(\mathbb{R}^d)$, we may take V_i in the above formula for \mathbf{A} the operators $V_i = W_i^k$, for some divergence-free vector fields W_i . Less trivial operators appear in a number of examples.)

A general existence result holds for equations driven by conservative rough drivers under very mild conditions on the functional setting.

THEOREM 11. *Assume one can associate to the scale $(E_n)_{n \geq 0}$ a family of self-adjoint smoothing operators $(J^\epsilon)_{0 < \epsilon \leq 1}$, from E_0 to itself, satisfying the regularisation estimates (4.2), and let \mathbf{A} be a conservative unbounded γ -rough driver on the scales $(E_n)_{n \geq 0}$. Then for any $f_0 \in \mathbb{L}^2(\mathbb{R}^d)$, there exists an $\mathbb{L}^2(\mathbb{R}^d)$ -valued path f_\bullet , started from f_0 , such that we have*

$$\delta f_{ts}(\varphi) = f_s \left((A_{ts}^1)^* \varphi \right) + f_s \left((A_{ts}^2)^* \varphi \right) + f_{ts}^\#(\varphi)$$

for all $\varphi \in E_3$, with

$$|f_t|_0 \leq |f_0|_0,$$

for all $t \geq 0$, and we have, for each finite time horizon T ,

$$(5.3) \quad \left| f_{ts}^\#(\varphi) \right| \lesssim_{C_0, \mathbf{A}, T, |f_0|_0} |\varphi|_3 |t-s|^{3\gamma},$$

for $0 \leq s \leq t \leq T$.

(One can think for instance to the operators $J^\epsilon = (I - \epsilon \Delta)^{-j_0}$, for j_0 big enough, if one is working in the Sobolev spaces $E_n = W^{n,2}(\mathbb{R}^d)$.) The proof goes by approximating the unbounded rough driver \mathbf{A} by bounded rough drivers \mathbf{A}^ϵ , and by using the theory developed in section 3 to solve the equation

$$(5.4) \quad \delta f_{ts}^\epsilon(\varphi) = f_s^\epsilon \left((A_{ts}^{\epsilon,1})^* \varphi \right) + f_s^\epsilon \left((A_{ts}^{\epsilon,2})^* \varphi \right) + f_{ts}^{\epsilon,\#}(\varphi).$$

The a priori bound (4.7) is used to pass to the limit as ϵ goes to 0.

PROOF – More concretely, let

$$A_{ts}^{\epsilon,1} := J^\epsilon A_{ts}^1 J^\epsilon, \quad A_{ts}^{\epsilon,2} := J^\epsilon A_{ts}^{2,a} J^\epsilon - \frac{1}{2} A_{ts}^{\epsilon,1} (A_{ts}^{\epsilon,1})^*,$$

with J^ϵ understood as a continuous operator from E_0 to E_{k_0+2} , for k_0 big enough, and $A_{ts}^{2,a} = \frac{1}{2} (A_{ts}^2 - (A_{ts}^2)^*)$ stands for the antisymmetric part of A_{ts}^2 . Both $A^{\epsilon,1}$ and $A^{\epsilon,2}$ are bounded linear operators from E_0 to itself, and by construction $\mathbf{A}^\epsilon = (A^{\epsilon,1}, A^{\epsilon,2})$ is a *bounded* conservative γ -rough driver on the Banach algebra $L(E_0)$. So one can denote by M_\bullet^ϵ the solution path in $L(E_0)$ of the rough differential equation

$$\delta M_{ts}^\epsilon = \left(A_{ts}^{\epsilon,1} + A_{ts}^{\epsilon,2} \right) M_s^\epsilon + M_{ts}^{\epsilon,\#},$$

provided by the theory developed in section 3. Given $f_0 \in E_0$, the path $f_\bullet^\epsilon = M_\bullet^\epsilon f_0$ solves the rough differential equation (5.4). Whereas theorem 3 provides an exponential control of the growth of the E_0 -norm of f_t^ϵ , with an exponent λ^ϵ that may go to ∞ as ϵ goes to 0, the conservative character of \mathbf{A}^ϵ ensures the uniform bound

$$|f_t^\epsilon| = |f_0|_0.$$

Indeed, denoting by (\cdot, \cdot) the scalar product on E_0 , and setting $z_t^\epsilon = |f_t^\epsilon|_0^2$, one has

$$\begin{aligned} \delta z_{ts}^\epsilon &= 2(\delta f_{ts}^\epsilon, f_s) + (\delta f_{ts}^\epsilon, \delta f_{ts}^\epsilon) \\ &= 2\left((A_{ts}^{\epsilon,1} + A_{ts}^{\epsilon,2})f_s, f_s\right) + (f_{ts}^{\epsilon,\sharp}, f_s) \\ &\quad + \left((A_{ts}^{\epsilon,1} + A_{ts}^{\epsilon,2})f_s, (A_{ts}^{\epsilon,1} + A_{ts}^{\epsilon,2})f_s\right) + O(|t-s|^{3\gamma}) \\ &= 2\left(A_{ts}^{\epsilon,2}f_s, f_s\right) + \left(A_{ts}^{\epsilon,1}f_s, A_{ts}^{\epsilon,1}f_s\right) + O(|t-s|^{3\gamma}) \end{aligned}$$

by the conservative character of $A_{ts}^{\epsilon,1}$, which finally gives

$$\delta z_{ts}^\epsilon = O(|t-s|^{3\gamma}),$$

since the symmetric part of $A_{ts}^{\epsilon,2}$ is $-\frac{1}{2}(A_{ts}^{\epsilon,1})^* A_{ts}^{\epsilon,1}$. The above equality shows that z^ϵ is constant.

Remark at that point that the constant

$$C_0^\epsilon := \|A^{\epsilon,1}\|_{\gamma;(-0,-1)} + \|A^{\epsilon,2}\|_{2\gamma;(-0,-2)} + \|A^{\epsilon,1}\|_{\gamma;(-2,-3)} + \|A^{\epsilon,2}\|_{2\gamma;(-1,-3)} < \infty$$

is not only finite but also uniformly bounded above, independently of $\epsilon > 0$. So we have, by theorem 9 an ϵ -uniform upper bound on $\|f^{\epsilon,\sharp}\|_{3\gamma;-3}$, of the form

$$(5.5) \quad \|f^{\epsilon,\sharp}\|_{3\gamma;-3} \lesssim_{\gamma,T,|f_0|_0} 1.$$

These bounds ensure in particular that for each $\varphi \in E_3$, the functions $f_\bullet^\epsilon(\varphi)$ form a bounded family of γ -Hölder real-valued paths, so it has a subsequence converging uniformly to some γ -Hölder real-valued function. Moreover, by weak- \star compactness, the uniform bound (5.5) implies the existence of a sequence $(\epsilon_n)_{n \geq 0}$ converging to 0, such that the sequence f^{ϵ_n} converges weakly- \star in $\mathbb{L}^\infty([0, T], E_0)$ to some limit $f \in \mathbb{L}^\infty([0, T], E_0)$. In particular, for each $\varphi \in E_3$, the sequence $f^{\epsilon_n}(\varphi)$ converges weakly- \star in $\mathbb{L}^\infty([0, T], \mathbb{R})$ to $f(\varphi)$. As it has a uniformly converging subsequence, this shows the γ -Hölder character of each function $f(\varphi)$.

Assuming $\varphi \in E_3$, it follows that one can pass to the limit in equation (5.4) in the three terms involving f_s^ϵ . The limit $f_{ts}^\sharp(\varphi)$ is defined as a consequence, and the bound (5.3) follows as a direct consequence of (5.5).

It is elementary to extend the above solution defined on $[0, T]$ to a globally defined solution satisfying the statement of the theorem. \triangleright

Rather than working with a general scale of spaces satisfying some ad hoc conditions, we shall set

$$E_n = W^{n,2}(\mathbb{R}^d),$$

for the remainder of this section on linear rough differential equation on Hilbert spaces. So we shall essentially be working from now on with rough drivers given by (at most) first order rough (pseudo-)differential operators.

5.2. Tensorization. In order to study the problem of uniqueness and further properties of solutions to general linear rough equations associated to unbounded rough drivers, we develop in this section a tensorization argument which can be seen as a rough version of the (differential) second quantisation functor in Hilbert spaces [39], or the variables doubling method commonly used in the theory of transport equations and conservation laws after the pioneering work of Kruzhkov [33]. As far as applications are concerned, we shall not restrict much our range in assuming that the rough drivers we are working with enjoy the following property. Given a bounded function $\phi \in W^{n_0, \infty}$, denote by M_ϕ the multiplication operator by ϕ ; it is a bounded operator from $E_0 = \mathbb{L}^2(\mathbb{R}^d)$ to itself.

DEFINITION 12. *An unbounded rough driver \mathbf{A} is said to be **symmetric** if the symmetric operators*

$$\begin{aligned} \text{(i)} \quad & B_{ts}^1(\phi) = (A_{ts}^1)^* M_\phi + M_\phi A_{ts}^1, \\ \text{(ii)} \quad & B_{ts}^2(\phi) = M_\phi (A_{ts}^2)^* + A_{ts}^2 M_\phi + A_{ts}^1 M_\phi (A_{ts}^1)^*, \end{aligned}$$

define quadratic forms

$$g \mapsto \left(g, B_{ts}^i(\phi)g \right)$$

that are continuous on E_0 , for all $0 \leq s \leq t < T$, and for any test function $\phi \in W^{3, \infty}$.

For rough drivers $\mathbf{A} = (A^1, A^2)$ of the form

$$A_{ts}^1 = X_{ts}^i V_i, \quad A_{ts}^2 = \mathbb{X}_{ts}^{jk} V_j V_k,$$

for some vector fields V_i on \mathbb{R}^d , and some weak geometric rough path \mathbf{X} , the operator $B^1(\phi)$ is the multiplication operator by $(A_{ts}^1)^* \phi$, and the second quadratic form is of the form

$$\left(g, B_{ts}^2(\phi)g \right) = \left(g^2, h_{ts}^\phi \right),$$

for some explicit function h_{ts}^ϕ that turns \mathbf{A} into a symmetric unbounded rough driver if the vector fields V_i are \mathcal{C}_b^2 and $\phi \in W^{2, \infty}$.

Let now \mathbf{A} be a symmetric unbounded rough driver in the scale $(E_n)_{n \geq 0}$, and let f_\bullet be a solution to the linear rough differential equation

$$df_s = \mathbf{A}(ds)f_s.$$

Consider $f^{\otimes 2} = f \otimes f$, defined on $\mathbb{R}^d \times \mathbb{R}^d$ by the usual formula

$$f^{\otimes 2}(x, y) = f(x)f(y);$$

it satisfies the equation

$$\delta f^{\otimes 2} = \left(A^1 \otimes \mathbb{I} + \mathbb{I} \otimes A^1 \right) f^{\otimes 2} + \left(A^2 \otimes \mathbb{I} + \mathbb{I} \otimes A^2 + \tau(A^1 \otimes A^1) \right) f^{\otimes 2} + f^{\otimes 2, \sharp},$$

where \mathbb{I} stands for the identity map and

$$\tau(A^1 \otimes A^1)_{ts} = A_{ts}^1 \otimes A_{ts}^1.$$

Setting

$$\begin{aligned} \Gamma_{\mathbf{A}}^1 &:= A^1 \otimes \mathbb{I} + \mathbb{I} \otimes A^1, \\ \Gamma_{\mathbf{A}}^2 &:= A^2 \otimes \mathbb{I} + \mathbb{I} \otimes A^2 + \tau(A^1 \otimes A^1), \end{aligned}$$

and

$$\Gamma_{\mathbf{A}} := (\Gamma_{\mathbf{A}}^1, \Gamma_{\mathbf{A}}^2),$$

we have then

$$\delta\Gamma_{\mathbf{A}}^2 = (A^1 A^1) \otimes \mathbb{I} + \mathbb{I} \otimes (A^1 A^1) + A^1 \otimes A^1 + \sigma(A^1 \otimes A^1),$$

where $\sigma(v \otimes w) = w \otimes v$ is the exchange of the two factors in the tensor so that

$$\sigma(A^1 \otimes A^1)_{tus} = A_{us}^1 \otimes A_{tu}^1 + A_{tu}^1 \otimes A_{us}^1.$$

It follows that $\Gamma_{\mathbf{A}}$ satisfies Chen's relations (3.1). Endowing $E_n^{\otimes 2}$ with its natural Hilbert space structure, we turn $\Gamma_{\mathbf{A}}$ into a γ -Hölder unbounded rough driver in the scale $(E_n^{\otimes 2})_{n \geq 0}$, for which $f^{\otimes 2}$ happens to be a solution, in this scale of spaces, of the linear rough equation

$$df_s^{\otimes 2} = \Gamma_{\mathbf{A}}(ds)f_s^{\otimes 2}.$$

The idea of the doubling variable method consists in using the dynamics of $f^{\otimes 2}$ to gain some information of its behaviour near the diagonal of $\mathbb{R}^d \times \mathbb{R}^d$. Let us define for that purpose a space \mathcal{E}_n^{∇} of test functions on $\mathbb{R}^d \times \mathbb{R}^d$, by requiring that they have finite $\|\cdot\|_n^{\nabla}$ -norm, as defined by the formula

$$\|\Phi\|_n^{\nabla} := \sup_{z \in \mathbb{R}^d} \sup_{0 \leq k+\ell \leq n} \int_{\mathbb{R}^d} \left| ((\nabla^+)^k (\nabla^-)^{\ell} \Phi)(z+w, z-w) \right| (1 \wedge |w|)^{\ell} dw,$$

where

$$\nabla^{\pm} := \frac{1}{2}(\nabla_x \pm \nabla_y).$$

Smoothing operators can be introduced in a way compatible with these spaces. Recall the definition of the operators $B_{ts}^1(\phi)$ and $B_{ts}^2(\phi)$ used in the definition of a symmetric unbounded rough driver. We use below brackets $\langle \cdot, \cdot \rangle$ to denote the \mathbb{L}^2 scalar product. Note that f_s is an element of $E_{-0} = E_0 = \mathbb{L}^2(\mathbb{R}^d)$, so f_s^2 is in $\mathbb{L}^1(\mathbb{R}^d)$.

LEMMA 13 (renormalisation). *Let \mathbf{A} be a symmetric rough driver and f_{\bullet} be a solution of the equation*

$$df_s = \mathbf{A}(ds)f_s,$$

in the initial scale of spaces $(E_n)_{n \geq 0}$. Assume that $\Gamma_{\mathbf{A}}$ is an unbounded γ -rough driver in the scale of spaces $(\mathcal{E}_n^{\nabla})_{n \geq 0}$, so $f_{\bullet}^{\otimes 2}$ satisfies the equation

$$(5.6) \quad \delta f^{\otimes 2}(\Phi)_{ts} = f_s^{\otimes 2} \left((\Gamma_{\mathbf{A}}^1)_{ts}^* \Phi \right) + f_s^{\otimes 2} \left((\Gamma_{\mathbf{A}}^2)_{ts}^* \Phi \right) + f^{\otimes 2, \#}(\Phi)_{ts}$$

for all smooth functions Φ , with $\|f^{\otimes 2, \#}(\Phi)_{ts}\|_{3\gamma} \lesssim \|\Phi\|_3^{\nabla}$. Then the $\mathbb{L}^1(\mathbb{R}^d)$ -valued path f_{\bullet}^2 satisfies, for all $\phi \in W^{3,\infty}$, the equation

$$(5.7) \quad \delta f^2(\phi)_{ts} = \left\langle f_s, (B_{ts}^1(\phi) + B_{ts}^2(\phi))f_s \right\rangle + O\left(\|\phi\|_{W^{3,\infty}}|t-s|^{3\gamma}\right).$$

By polarisation the product fg satisfies an equation analogous to equation (5.7) if both f and g are solutions of the equation $df_s = \mathbf{A}(ds)f_s$, in the scale $(E_n)_{n \geq 0}$.

PROOF – Let us start with the equation (5.6) satisfied by $f^{\otimes 2}$, for any function $\Phi \in \mathcal{E}_3^{\nabla}$.

Equation (5.7) will come from taking in equation (5.6) some functions Φ of the form

$$\Phi^{\epsilon}(x, y) = \psi^{\epsilon}(x-y) \phi\left(\frac{x+y}{2}\right),$$

for some family $(\psi^{\epsilon})_{0 < \epsilon \leq 1}$ of smooth approximations of a Dirac mass at 0, and by letting ϵ tend to 0, after checking that some ϵ -uniform estimates hold for the different terms in (5.6).

With this aim in mind, note first that

$$\begin{aligned}
\|\Phi^\epsilon\|_3^\nabla &= \sup_{z \in \mathbb{R}^d} \sup_{0 \leq k+\ell \leq 3} \int_{\mathbb{R}^d} |(\nabla^\ell \psi^\epsilon)(2w)(\nabla^k \phi)(z)| (1 \wedge |w|)^\ell dw \\
(5.8) \quad &\lesssim \|\phi\|_{W^{3,\infty}(\mathbb{R}^d)} \sup_{0 \leq \ell \leq 3} \int_{\mathbb{R}^d} |(\nabla^\ell \psi^\epsilon)(2w)| (1 \wedge |w|)^\ell dw \\
&\leq C \|\phi\|_{W^{3,\infty}(\mathbb{R}^d)},
\end{aligned}$$

is bounded above uniformly in ϵ . So the 3γ -Hölder norm of the remainders $f^{\otimes 2, \#}(\Phi^\epsilon)$ are bounded by a constant multiple of $\|\phi\|_{W^{3,\infty}}$, independent of ϵ . Note also that Young's inequality provides the ϵ -uniform bound

$$\begin{aligned}
|f_s^{\otimes 2}(\Phi^\epsilon)| &= \left| \int_{\mathbb{R}^d \times \mathbb{R}^d} f_s(x) f_s(y) \psi^\epsilon(x-y) \phi\left(\frac{x+y}{2}\right) dx dy \right| \\
&\leq \|\phi\|_{\mathbb{L}^\infty} \int_{\mathbb{R}^d \times \mathbb{R}^d} |f_s(x)| |f_s(y)| |\psi^\epsilon(x-y)| dx dy \\
&\leq \|\phi\|_{\mathbb{L}^\infty} \|f_s\|_{\mathbb{L}^2}^2.
\end{aligned}$$

Now, given a positive constant δ , the fact that for any smooth function g which is δ -close in $\mathbb{L}^2(\mathbb{R}^d)$ of f_s , we have

$$|f_s^{\otimes 2}(\Phi^\epsilon) - g^{\otimes 2}(\Phi^\epsilon)| \lesssim 2\delta \|f\|_{\mathbb{L}^2} + \delta^2,$$

uniformly in ϵ , and

$$\lim_{\epsilon \rightarrow 0} g^{\otimes 2}(\Phi^\epsilon) = \int_{\mathbb{R}^d} |g(x)|^2 \phi(x) dx,$$

shows that

$$f_s^{\otimes 2}(\Phi^\epsilon) \xrightarrow{\epsilon \rightarrow 0} f_s^2(\phi).$$

We also have the convergence

$$g^{\otimes 2}((\Gamma_{\mathbf{A}}^1)_{ts}^* \Phi^\epsilon) = (A_{ts}^1 g \otimes g + g \otimes A_{ts}^1 g, \Phi^\epsilon) \xrightarrow{\epsilon \rightarrow 0} 2 \int_{\mathbb{R}^d} g(x) (A_{ts}^1 g)(x) \phi(x) dx,$$

which we can rewrite as

$$g^{\otimes 2}((\Gamma_{\mathbf{A}}^1)_{ts}^* \Phi^\epsilon) \xrightarrow{\epsilon \rightarrow 0} 2(g, (A_{ts}^1)^* M_\phi g) = (g, B_{ts}^1(\phi)g).$$

Now, since $\Gamma_{\mathbf{A}}$ is a rough driver in the scale \mathcal{E}_n^∇ , and we have equation (5.8), the family $((\Gamma_{\mathbf{A}}^1)_{ts}^* \Phi^\epsilon)_{0 < \epsilon \leq 1}$ is bounded in \mathcal{E}_0^∇ . Using in addition the boundedness on $\mathbb{L}^2(\mathbb{R}^d)$ of the quadratic form associated to $B_{ts}^1(\phi)$, one can then send g to f_s , in $\mathbb{L}^2(\mathbb{R}^d)$, in the above convergence result and conclude that

$$f_s^{\otimes 2}((\Gamma_{\mathbf{A}}^1)_{ts}^* \Phi^\epsilon) \rightarrow (f_s, B_{ts}^1(\phi) f_s).$$

Similarly, the boundedness in \mathcal{E}_0^∇ of the family $((\Gamma_{\mathbf{A}}^2)_{ts}^* \Phi^\epsilon)_{0 < \epsilon \leq 1}$, together with the boundedness on $\mathbb{L}^2(\mathbb{R}^d)$ of the quadratic form associated with $B_{ts}^2(\phi)$, show that

$$f_s^{\otimes 2}((\Gamma_{\mathbf{A}}^2)_{ts}^* \Phi^\epsilon) \rightarrow (f_s, B_{ts}^2(\phi) f_s);$$

equation (5.7) follows, as we have the ϵ -uniform bound $|f_{ts}^{\otimes 2, \#}(\phi)| \leq \|\phi\|_{W^{3,\infty}} |t-s|^{3\gamma}$.

▷

This result is sufficient to prove that rough linear equations driven by conservative drivers are unique if the driver is symmetric.

COROLLARY 14. *Let \mathbf{A} be a symmetric conservative unbounded γ -rough driver in the scale of spaces $(E_n)_{n \geq 0}$, such that $\Gamma_{\mathbf{A}}$ is an unbounded γ -rough driver in the scale of spaces $(\mathcal{E}_n^{\nabla})_{n \geq 0}$. Then the rough linear equation*

$$df_s = \mathbf{A}(ds)f_s$$

has a unique solution in $\mathbb{L}^2(\mathbb{R}^d)$, started from any initial condition $f_0 \in E_0$; it satisfies $\|f_t\|_0 = \|f_0\|_0$, for all times t .

PROOF – It suffices to notice that since the driver \mathbf{A} is conservative, we have $B_{ts}^1(\mathbf{1}) = B_{ts}^2(\mathbf{1}) = 0$, so it follows from equation (5.7) that any solution path f_{\bullet} has constant \mathbb{L}^2 -norm, which proves the uniqueness claim. Existence was proved in theorem 11. \triangleright

5.3. A priori bounds for closed symmetric drivers. One cannot use directly the renormalisation lemma to get some closed equation for f^2 when \mathbf{A} is non-conservative. We need for that purpose to assume that the symmetric unbounded rough driver \mathbf{A} enjoys the following property.

DEFINITION 15. *A symmetric unbounded rough driver \mathbf{A} in the scale of spaces $(W^{n,2})_{n \geq 0}$, is said to be **closed** if there exists some unbounded rough driver $\mathbf{B} = (B^1, B^2)$ in the scale of spaces $(W^{n,\infty})_{n \geq 0}$, such that we have*

$$\left(g, B_{ts}^1(\phi)g\right) = \left(g^2, (B_{ts}^1)^*\phi\right), \quad \text{and} \quad \left(g, B_{ts}^2(\phi)g\right) = \left(g^2, (B_{ts}^2)^*\phi\right),$$

for all $g \in E_0$.

As an example, it is elementary to check that the unbounded rough driver $\mathbf{A} = (XV, \mathbb{X}V)$ used in the rough transport equation

$$\delta f_{ts} = X V f_s + \mathbb{X} V V f_s + f_{ts}^{\sharp}$$

with some γ -Hölder weak geometric rough path $\mathbf{X} = (X, \mathbb{X})$, is closed and symmetric if the vector fields $V = (V_1, \dots, V_\ell)$ are C_b^3 , in which case $\mathbf{B} = \mathbf{A}$. Another interesting class of examples of closed symmetric unbounded rough driver in the scale of spaces $(E_n)_{n \geq 0}$, is provided by the lift to rough drivers of \mathcal{C}_b^3 -semimartingale velocity fields, as given in the theory of stochastic flows. This kind of stochastic velocity fields appear in the study of Navier–Stokes equation. See the work [2] for a thorough study of stochastic flows from this point of view.

Building on this notion of closed driver, the following statement provides amongst other things an a priori estimate on solutions of rough linear equations that plays in this setting the role played in the classical setting by a priori estimates obtained by any kind of Gronwall–type argument. The crucial point here is that no such Gronwall machinery was available so far in a rough path–like setting; despite its elementary nature, this result may well be one of our main contributions.

THEOREM 16. *Let \mathbf{A} be a closed symmetric unbounded γ -rough driver on the scales $(W^{n,2}(\mathbb{R}^d))_{n \in \mathbb{N}}$. Let \mathbf{B} be its associated driver, and assume that the inequality*

$$(5.9) \quad \left| (B_{t0}^1)^* \mathbf{1} \right| \vee \left| (B_{t0}^2)^* \mathbf{1} \right| \leq c_t$$

holds for all t , for some time-dependent positive constant c_t such that $e^{-\lambda t} c_t$ tends to 0 as t goes to infinity, for any positive parameter λ , so

$$(\mathbf{c}_{\bullet}) < \infty.$$

Assume that $\Gamma_{\mathbf{A}}$ is an unbounded γ -rough driver in the scale of spaces $(\mathcal{E}_n^\nabla)_{n \geq 0}$. Then, given any $f_0 \in \mathbb{L}^2(\mathbb{R}^d)$, there is at most one $\mathbb{L}^2(\mathbb{R}^d)$ -valued solution path f_\bullet to the equation

$$(5.10) \quad \delta f_{ts}(\varphi) = f_s \left((A_{ts}^1)^* \varphi \right) + f_s \left((A_{ts}^2)^* \varphi \right) + f_{ts}^\sharp(\varphi),$$

and we have, for each finite time horizon T ,

$$(5.11) \quad \left| f_{ts}^\sharp(\varphi) \right| \lesssim_{\mathbf{B}, T, |f_0|_0} \|\varphi\|_3 |t - s|^{3\gamma},$$

for all $\varphi \in W^{3,2}$, and all $0 \leq s \leq t \leq T$. It satisfies the upper bound

$$(5.12) \quad |f_t|_0 \lesssim_{\mathbf{B}, t} |f_0|_0$$

PROOF – Let f_\bullet be a solution to the rough linear equation (5.10) in the scale of spaces $(W^{n,2}(\mathbb{R}^d))_{n \in \mathbb{N}}$. Since \mathbf{A} is closed, the $\mathbb{L}^1(\mathbb{R}^d)$ -valued path f_\bullet^2 happens to be a solution to the rough linear equation

$$\delta f^2(\phi)_{ts} = f_s^2 \left((B_{ts}^1)^* \phi \right) + f_s^2 \left((B_{ts}^2)^* \phi \right) + (f^2)_{ts}^\sharp(\phi),$$

in the scale of spaces $(W^{n,\infty}(\mathbb{R}^d))_{n \in \mathbb{N}}$. Denote by $C_0^{\mathbf{B}}$ the finite constant associated to the unbounded rough driver \mathbf{B} , as defined by equation (4.6), with \mathbf{B} in the role of \mathbf{A} . It follows from the general a priori estimates on solutions of rough linear equations proved in theorem 8, and the fact that f^2 is in $\mathbb{L}^1(\mathbb{R}^d)$, that

$$(5.13) \quad \|(f^2)^\sharp\|_{3\gamma; (W^{3,\infty})^*} \lesssim_{\gamma, \lambda} C_0^{\mathbf{B}} \|(f^2)\|_{(\mathbb{L}^\infty)^*} \lesssim_{\gamma, \lambda} C_0^{\mathbf{B}} \|(f_\bullet^2)\|_{\mathbb{L}^1}.$$

But since we have the identity

$$f_t^2(\mathbf{1}) = f_0^2 \left(\mathbf{1} + (B_{ts}^1)^* \mathbf{1} + (B_{ts}^2)^* \mathbf{1} \right) + (f^2)_{t0}^\sharp(\mathbf{1})$$

and the bound (5.9), we also have the estimate

$$\begin{aligned} \|(f_\bullet^2)\|_{\mathbb{L}^1} &= \|(f_\bullet^2(\mathbf{1}))\| \lesssim_{(c_\bullet)} |f_0|_{\mathbb{L}^2} + \|(f^2)_{\bullet 0}^\sharp\|_{(W^{3,\infty})^*} \\ &\lesssim_{(c_\bullet)} \left(|f_0|_{\mathbb{L}^2} + \lambda^{3\gamma} \|(f^2)^\sharp\|_{3\gamma; (W^{3,\infty})^*} \right). \end{aligned}$$

(Note that $(f^2)^\sharp$, in the right hand side of the above inequality, is seen as a 2-index function.) Together with the bound (5.13), this gives the upper bound

$$\|(f_\bullet^2)\|_{\mathbb{L}^1} \lesssim_{(c_\bullet), C_0^{\mathbf{B}}} |f_0|_{\mathbb{L}^2}$$

for λ small enough, which implies uniqueness. \triangleright

5.4. Rough transport equation. Building on theorem 16, one can give a complete \mathbb{L}^2 -theory of rough transport equations

$$\delta f_{ts} = X V f_s + \mathbb{X} V V f_s + f_{ts}^\sharp$$

driven by non-divergence-free vector fields V_i of class $W^{3,\infty}$.

LEMMA 17. Let \mathbf{X} be a geometric γ -Hölder rough path on \mathbb{R}^ℓ , and V_1, \dots, V_ℓ be $W^{3,\infty}$ vector fields on \mathbb{R}^d . Then the operator $\Gamma_{\mathbf{A}}$ associated with $\mathbf{A} = (XV, \mathbb{X}VV)$, is an unbounded γ -rough driver in the scale of spaces $(\mathcal{E}_n^\nabla)_{n \geq 0}$.

PROOF – For a geometric rough path $\mathbf{X} = (X, \mathbb{X})$, the operator $\Gamma_{\mathbf{A}}$ takes the form $\Gamma_{\mathbf{A}} = (X\Gamma_V^1, \mathbb{X}\Gamma_V^2)$, with

$$\begin{aligned}\Gamma_V^1 &:= V \otimes \mathbb{I} + \mathbb{I} \otimes V, \\ \Gamma_V^2 &:= VV \otimes \mathbb{I} + \mathbb{I} \otimes VV + 2(V \otimes V) = \Gamma_V^1 \Gamma_V^1.\end{aligned}$$

So it is enough to show that the adjoints of these operators satisfy the inequalities

$$(5.14) \quad \|\Gamma_V^{1,*} \Phi\|_n^\nabla \lesssim \|V\|_{W^{n+1,\infty}} \|\Phi\|_{n+1}^\nabla, \quad \|\Gamma_V^{2,*} \Phi\|_m^\nabla \lesssim \|V\|_{W^{m+2,\infty}}^2 \|\Phi\|_{m+2}^\nabla$$

for $n = 0, 2$, $m = 0, 1$, for smooth test functions Φ , to prove the lemma.

Write $V = v_k \partial_k$ where $(v_k)_{k=1,\dots,d}$ are the coefficients of the vector fields in the canonical basis $(\partial_k)_{k=1,\dots,d}$ of derivations; with these notations, we have

$$V^* = -v_k \partial_k - d,$$

where $d := \operatorname{div} v$, is the divergence of the vector field V , and

$$\Gamma_V^{1,*} = v_k^+ \partial_k^+ + v_k^- \partial_k^- + d^+ + d^-,$$

where, for a real-valued function h on \mathbb{R}^d , we denote by

$$h^\pm(x, y) := h(x) \pm h(y)$$

its symmetric and antisymmetric lift to $\mathbb{R}^d \times \mathbb{R}^d$. The first estimate in (5.14) follows from the inequalities

$$\|a^\pm \nabla^\pm \Phi\|_n^\nabla \lesssim \|a\|_{W^{n+1,\infty}} \|\Phi\|_{n+1}^\nabla,$$

which we prove below. The second inequality in (5.14) is obtained by noting that we have

$$\begin{aligned}\|\Gamma_V^{2,*} \Phi\|_m^\nabla &= \|\Gamma_V^{1,*} \Gamma_V^{1,*} \Phi\|_m^\nabla \lesssim \|V\|_{W^{m+1,\infty}} \|\Gamma_V^{1,*} \Phi\|_{m+1}^\nabla \\ &\lesssim \|V\|_{W^{m+1,\infty}} \|V\|_{W^{m+2,\infty}} \|\Phi\|_{m+2}^\nabla.\end{aligned}$$

To prove the above inequality, note that the gradients involved in the definition of the norm of \mathcal{E}_n^∇ give rise to a number of terms of the form

$$((\nabla^+)^{\ell} (\nabla^-)^m a^\pm) ((\nabla^+)^{\ell'} (\nabla^-)^{m'} \nabla^\pm \Phi),$$

with $\ell + \ell' + m + m' \leq n$, and notice that $\nabla^+ a^- = (\nabla a)^-$, so we have

$$(\nabla^+)^{\ell} (\nabla^-)^m a^- = (\nabla^-)^m (\nabla^{\ell} a)^-.$$

Now, if $m = 0$, then

$$\begin{aligned}(1 \wedge |w|)^{m'} &\left| (\nabla^{\ell} a)^-(z+w, z-w) ((\nabla^+)^{\ell'} (\nabla^-)^{m'} \nabla^- \Phi)(z+w, z-w) \right| \\ &\lesssim (1 \wedge |w|)^{m'+1} \|\nabla^{\ell+1} a\|_{\mathbb{L}^\infty} \left| ((\nabla^+)^{\ell'} (\nabla^-)^{m'+1} \Phi)(z+w, z-w) \right|,\end{aligned}$$

since $|\varphi^-(z+w, z-w)| \lesssim \|\nabla \varphi\|_{\mathbb{L}^\infty}$. On the other hand, if $m > 0$

$$\begin{aligned}(1 \wedge |w|)^{m+m'} &\left| (\nabla^-)^m (\nabla^{\ell} a)^-(z+w, z-w) ((\nabla^+)^{\ell'} (\nabla^-)^{m'} \nabla^- \Phi)(z+w, z-w) \right| \\ &\lesssim (1 \wedge |w|)^{m+m'} \|\nabla^{\ell+m+1} a\|_{\mathbb{L}^\infty} \left| ((\nabla^+)^{\ell'} (\nabla^-)^{m'+1} \Phi)(z+w, z-w) \right| \\ &\lesssim (1 \wedge |w|)^{m'+1} \|\nabla^{\ell+m+1} a\|_{\mathbb{L}^\infty} \left| ((\nabla^+)^{\ell'} (\nabla^-)^{m'+1} \Phi)(z+w, z-w) \right|,\end{aligned}$$

since $\nabla^- \varphi^- = (\nabla \varphi)^+$ and $m' + 1 \leq m + m'$. This shows that the correct number of powers of $1 \wedge |w|$ remains available to balance the powers of ∇^- acting on the test function Φ ; estimates (5.14) follows easily from this observation. \triangleright

THEOREM 18. *Let \mathbf{X} be a geometric γ -Hölder rough path on \mathbb{R}^ℓ , and V_1, \dots, V_ℓ be $W^{3,\infty}$ vector fields on \mathbb{R}^d . Then the rough transport equation*

$$\delta f_s = (X_{ts} V + \mathbb{X}_{ts} V V) f_s + f_{ts}^\sharp$$

is well-posed.

PROOF – Notice first that the regularity assumption on the V_i puts us in a position to use the a priori bounds for symmetric closed drivers stated in theorem 11, with \mathbf{A} in the role of \mathbf{B} . So uniqueness is a direct consequence of the a priori bound (5.12) in theorem 16.

Let now $f_0 \in E_0$ be given. We prove the existence of a solution path to rough transport equation started from f_0 , by a classical approximation-compactness argument, relying in a crucial way on the a priori bound (5.12) on the \mathbb{L}^2 -norm of the solution to the approximate problem, and on the uniform estimate (5.3) for the remainder. Fix a finite time horizon T .

Given that \mathbf{X} is geometric, let $(\mathbf{X}^\epsilon)_{0 < \epsilon \leq 1}$ be a family of rough path lifts of smooth paths which converge to \mathbf{X} in a rough paths sense over the time interval $[0, T]$. Let also $(V^\epsilon)_{0 < \epsilon \leq 1}$ be a family of smooth vector fields that converge to V in $W^{3,\infty}$, and let $(\mathbf{A}^\epsilon)_{0 < \epsilon \leq 1}$ be their associated rough driver, defined by formula (5.2) with \mathbf{X}^ϵ and V^ϵ in place of \mathbf{X} and V respectively. One can choose $(\mathbf{X}^\epsilon)_{0 < \epsilon \leq 1}$ in such a way that the constant C_0^ϵ associated with \mathbf{A}^ϵ by formula (4.6) satisfies the inequality $C_0^\epsilon \lesssim C_0$, independently of $0 < \epsilon \leq 1$. Given the smooth character of the vector fields V^ϵ , one can solve uniquely the transport equation

$$\delta f_{ts}^\epsilon(\varphi) = f_s^\epsilon(V^{\epsilon,*} \varphi) X_{ts}^\epsilon + f_s^\epsilon(V^{\epsilon,*} V^{\epsilon,*} \varphi) \mathbb{X}_{ts}^\epsilon + f_{ts}^{\epsilon,\sharp}(\varphi), \quad \text{for } \varphi \in E_2,$$

by the method of characteristics, as the above equation is actually equivalent to the ordinary differential equation

$$\frac{df_t^\epsilon}{dt} = f_t^\epsilon(V^{\epsilon,*} \varphi) X_t^{\epsilon'}.$$

The solutions of this problem satisfy the uniform estimates

$$|f_t^\epsilon|_0 \lesssim_{C_0, T} |f_0|_0,$$

for all $0 \leq t \leq T$, as a consequence of (5.12), and we also have the uniform bound

$$\sup_{0 < \epsilon \leq 1} \|f^{\epsilon,\sharp}\|_{3\gamma; -3} \lesssim_{C_0, \gamma, T, |f_0|_0} 1,$$

by (5.3). These two a priori estimates are all we need to finish the proof of the theorem following word by word the end of the proof of theorem 16. \triangleright

It is perfectly possible to extend the present theory to deal with rough linear equations with a **drift**

$$df_s = W f_s ds + \mathbf{A}(ds) f_s,$$

where $W \in L(E_{-0}, E_{-2})$, such as the Laplacian operator in the $W^{n,2}(\mathbb{R}^d)$ scale of spaces. We refrain from giving the details here as this is not our main point and this does not

require the introduction of new tools or ideas. This provides an alternative road to some of the results of [12] in a slightly different setting.

6. THE \mathbb{L}^∞ THEORY OF ROUGH TRANSPORT EQUATIONS

We develop in this section an \mathbb{L}^∞ theory of the rough transport equation

$$(6.1) \quad \delta f_{ts} = X V f_s + \mathbb{X} V V f_s + f_{ts}^\sharp$$

and prove its well-posed character under the assumption that the vector fields be \mathcal{C}^3 , for some positive constant ν . We show for that purpose that all solutions are renormalised solutions, in the sense of di Perna–Lions, which classically leads to uniqueness and stability results in that setting.

6.1. A priori estimates and existence result. For developping that \mathbb{L}^∞ theory, we shall be working in the scale of Sobolev spaces

$$E_n = W^{n,1}(\mathbb{R}^d), \quad \text{for } n \geq 0,$$

with norm denoted by $|\cdot|_n$, and in which one has regularising operators $(J^\epsilon)_{0 < \epsilon \leq 1}$ for which estimates (4.2) hold. Our minimal regularity assumptions on the vector fields will be the existence of a positive constant C_1 such that the inequalities

$$(6.2) \quad |V_i^* \varphi|_0 \leq C_1 |\varphi|_1, \quad |V_i^* V_j^* \varphi|_0 \leq C_1 |\varphi|_2$$

hold for all $1 \leq i, j \leq \ell$. These conditions hold for instance if the vector fields V_i and $(V_i V_j)$ are all \mathcal{C}_b^1 ; we write here $(V_i V_j)$ for $(DV_j)(V_i)$. One proves the following existence result by proceeding exactly as in the proof of theorem 18, using the a priori \mathbb{L}^∞ -estimate

$$|f_t^\epsilon|_{\mathbb{L}^\infty} = |f_0^\epsilon|_{\mathbb{L}^\infty},$$

for the regularised equation, and using theorem 9 to get an ϵ -uniform control on $|f^{\epsilon, \sharp}|_{3\gamma; -3}$, in terms of \mathbf{X} and $|f_0^\epsilon|_{\mathbb{L}^\infty}$ only. It holds in particular if V is \mathcal{C}_b^2 .

THEOREM 19 (Existence for rough transport equations – \mathbb{L}^∞ setting). *Under the continuity assumptions (6.2) on the vector fields V_i , for any $f_0 \in \mathbb{L}^\infty(\mathbb{R}^d)$, there exists an $\mathbb{L}^\infty(\mathbb{R}^d)$ -valued path $(f_t)_{t \geq 0}$, started from f_0 , satisfying the equation*

$$\delta f_{ts}(\varphi) = f_s(V^* \varphi) X_{ts} + f_s(V^* V^* \varphi) \mathbb{X}_{ts} + f_{ts}^\sharp(\varphi)$$

for all $\varphi \in E_3$, and the bound

$$\sup_{t \geq 0} |f_t|_{\mathbb{L}^\infty(\mathbb{R}^d)} \leq |f_0|_{\mathbb{L}^\infty(\mathbb{R}^d)},$$

with a remainder $f^\sharp(\varphi)$ controlled by

$$(6.3) \quad |f_{ts}^\sharp(\varphi)| \lesssim_{C_1, \mathbf{X}, T, |f_0|_{\mathbb{L}^\infty}} |\varphi|_3 |t - s|^{3\gamma},$$

for $0 \leq s \leq t \leq T$.

6.2. renormalised solutions, uniqueness and stability. To proceed one step further, we show that a mild strengthening of the regularity conditions imposed on the vector fields V_i suffices to guarantee that all bounded solutions to the transport equation (6.1) are actually renormalised solution, in the sense of the following definition.

DEFINITION 20. *A solution f_\bullet to the transport equation (6.1) in the scales $(E_n)_{n \geq 0}$ is said to be a **renormalised solution** if for any function $H : \mathbb{R} \rightarrow \mathbb{R}$, of class \mathcal{C}_b^3 , the path $h_\bullet = H \circ f_\bullet$ is also a solution to equation (6.1) in the same scale $(E_n)_{n \geq 0}$.*

As expected, this property will lead below to uniqueness and stability results.

THEOREM 21. *Assume the vector fields V_i are \mathcal{C}_b^3 . Then every solution to the transport equation (6.1), bounded in $\mathbb{L}^\infty(\mathbb{R}^d)$, is a renormalised solution.*

PROOF – The renormalisation lemma 13 can be stated in the \mathbb{L}^∞ setting by choosing a slightly different scale \mathcal{F}_n^∇ of spaces of test functions, with norms modelled on \mathbb{L}^1

$$(6.4) \quad |\varphi|_n^\nabla := \sup_{0 \leq k+\ell \leq n} \int \int \left| (\nabla^+)^k (\nabla^-)^\ell \varphi(x, y) \right| |x - y|^\ell dx dy$$

rather than on a weighted \mathbb{L}^∞ space used in section 5.2. Identity (5.7) holds in that case with for functions $\phi \in W^{3,1}(\mathbb{R}^d)$, with an $O(\cdot)$ term involving the $W^{3,1}$ -norm of ϕ rather than its $W^{3,\infty}$ -norm, as the proof of lemma 13 works verbatim, provided we can prove that $\Gamma_{\mathbf{A}}$ is an unbounded rough driver in the scale of spaces $(\mathcal{F}_n^\nabla)_{n \geq 0}$ associated with the norm (6.4). (Note that we have in that case $|f_s^{\otimes 2}(\Phi^\epsilon)| \leq c_\phi |f_s|_{\mathbb{L}^\infty}^2 = c_\phi |f_s|_{-0}^2$.)

The proof that $\Gamma_{\mathbf{A}}$ is an unbounded rough driver follows the same pattern as the proof given in section 5.4. We invite the reader to complete the details.

So it follows from the renormalisation lemma that if f, g are two solutions the above argument also goes through and shows that fg is also a solution, so any power f^n of f is also a solution, with a size of the remainder that depends only on the \mathbb{L}^∞ norm of f^n . By linearity the result can be extended to any polynomial of f , and by density to any continuous function $H(f)$, with H defined on the interval $[-\|f\|_\infty, \|f\|_\infty]$. \triangleright

We can actually improve a bit this condition and require only a weak integrability for the third derivative of V ; it provides a significant strengthening of the previous statement when the vector fields V_i are divergence-free, giving some analogue of the traditional di Perna–Lions conditions in the classical setting.

THEOREM 22. *Assume that $V \in \mathcal{C}_b^2$, $\nabla^3 V \in L^1$ and $\operatorname{div} V \in \mathcal{C}_b^2$. Then every solution to the transport equation (6.1), bounded in $\mathbb{L}^\infty(\mathbb{R}^d)$, is a renormalised solution.*

PROOF – In the proof of the renormalisation lemma 13 we can use directly the general apriori estimate stated in Thm 8 applied to $\Gamma_{\mathbf{A}}$ with $F = \tilde{\mathcal{E}}_3^\nabla$ and $E = \mathcal{F}_0^\nabla$ (note the choice for F). Where $\tilde{\mathcal{E}}_n^\nabla$ are spaces of test functions, with norms modelled on \mathbb{L}^∞ like \mathcal{E}_n^∇ but with a small change given by an additional averaging over the auxiliary variable τ :

$$|\varphi|_n^\nabla := \sup_{0 \leq k+\ell \leq n} \sup_{x \in \mathbb{R}^d} \int_0^1 d\tau \int \left| (\nabla^+)^k (\nabla^-)^\ell \varphi(x - \tau w, x + (1 - \tau)w) \right| |w|^\ell dw$$

the reason of which will be clear below. In this case we can show that

$$N_1(\Gamma_{\mathbf{A}}) \lesssim (1 + \|V\|_{\mathcal{C}_b^2})^2$$

while

$$N_2(\Gamma_{\mathbf{A}}) \lesssim (1 + \|V\|_{\mathcal{C}_b^2} + \|\nabla^3 V\|_{L^1} + \|\operatorname{div} V\|_{\mathcal{C}_b^2})^3.$$

Indeed apart from many contributions which can be estimated as in the \mathbb{L}^2 or as in the previous theorem, a difficult term come from the estimation of norms like $\|\Gamma_V^* \Gamma_V^* \Gamma_V^*\|_{\mathbb{L}(\mathbb{F}, \mathbb{E})}$ of which the most singular contribution is given by $\|\Gamma_V \Gamma_V \Gamma_V\|_{\mathbb{L}(\mathbb{F}, \mathbb{E})}$. In this norm the contribution that requires more regularity to V is due to the first two vector fields Γ_V acting simultaneously on the third one giving terms of the form $\|v^+ v^+ (\nabla^2 v)^- \nabla^-\|_{\mathbb{L}(\mathbb{F}, \mathbb{E})}$ and easier ones. Now expanding $(\nabla^2 v)^-(x, y) = \int_0^1 d\tau (\nabla^3 v)(x + \tau(y - x))(y - x)$ we get

$$\begin{aligned} \|v^+ v^+ (\nabla^2 v)^- \nabla^-\Psi\|_E &= \int \int \left| (v^+ v^+ (\nabla^2 v)^- \Psi)(x, y) \right| dx dy \\ &\lesssim \|V\|_{L^\infty}^2 \int_0^1 d\tau \int \int |\nabla^3 v(x + \tau w)| |\Psi(x, x + w)| |w| dx dw \\ &\lesssim \|V\|_{L^\infty}^2 \int_0^1 d\tau \int dx |\nabla^3 v(x)| \int dw |\Psi(x - \tau w, x - \tau w + w)| |w| \\ &\lesssim \|V\|_{L^\infty}^2 \|\nabla^3 V\|_{L^1} \sup_{x \in \mathbb{R}^d} \int_0^1 d\tau \int dw |\Psi(x - \tau w, x - \tau w + w)| |w| \\ &\lesssim \|V\|_{L^\infty}^2 \|\nabla^3 V\|_{L^1} \|\Phi\|_F. \end{aligned}$$

Granted the bounds on $N_1(\Gamma_{\mathbf{A}})$ and $N_2(\Gamma_{\mathbf{A}})$ the proof continues as the proof of the previous theorem and gives the renormalisation result. \triangleright

As expected, theorem 21 on renormalised solutions to the transport equation (6.1) comes with a number of important consequences, amongst which is an equivalent of the missing Gronwall lemma, as given by the a priori estimate (6.5) below.

THEOREM 23. *Assume the vector fields V_i are \mathcal{C}_b^3 .*

- (1) **Uniqueness** – *Given an initial condition in $\mathbb{L}^\infty(\mathbb{R}^d)$, there exists a unique solution to the transport equation which remains bounded in $\mathbb{L}^\infty(\mathbb{R}^d)$.*
- (2) **Stability** – *Let the time horizon T be finite. Let $(V_i^{(n)})_{n \geq 0}$, $i = 1..l$ and $(f_0^{(n)})_{n \geq 0}$ be a sequence of approximating sequences with $V_i^{(n)}$ converging to V_i in \mathcal{C}_b^3 , and $f_0^{(n)}$ converging to f_0 in $\mathbb{L}(\mathbb{R}^d)$. Let also $(\mathbf{X}^{(n)})_{n \geq 0}$ be a sequence of weak geometric γ -rough paths above smooth paths, that converge in a rough paths sense to \mathbf{X} , over the time interval $[0, T]$. Then the solution paths $f_\bullet^{(n)}$ to the transport equation associated with $\mathbf{X}^{(n)}$, $V^{(n)}$ and $f_0^{(n)}$, converge weakly- \star in $\mathbb{L}^\infty([0, T], \mathbb{L}^\infty(\mathbb{R}^d))$, and in $\mathbb{L}_{\text{loc}}^1([0, T], \mathbb{L}^\infty(\mathbb{R}^d))$, to f_\bullet .*

PROOF – Uniqueness – We follow the same pattern of proof as that of theorem 16. Let f_\bullet and f'_\bullet be two solution paths to equation (6.1), bounded in $\mathbb{L}^\infty(\mathbb{R}^d)$, and started from the same initial condition. Let $H : \mathbb{R} \rightarrow \mathbb{R}$ be a non-negative function, of class \mathcal{C}_b^3 , null at 0 and positive elsewhere. Define the path

$$h_\bullet = H(f_\bullet - f'_\bullet);$$

it is also a positive solution to the transport equation (6.1) under the above regularity assumptions on the vector fields V_i , since all solutions are renormalised solution, by

theorem 21. Set $\psi(x) = (1 + |x|^2)^{-k_0}$, for $x \in \mathbb{R}^d$, and some exponent $k_0 > d$. That function satisfies

$$\begin{aligned} \operatorname{div}(\psi V) &= -V\psi - (\operatorname{div}V)\psi, \\ \operatorname{div}(\operatorname{div}(\psi V) V) &= V^2\psi + (\operatorname{div}V)V\psi + ((V\operatorname{div}V) + (\operatorname{div}V)^2)\psi \end{aligned}$$

with

$$\left| V\psi \right| \vee \left| V^2\psi + (\operatorname{div}V)V\psi \right| \lesssim \psi$$

as a consequence of the \mathcal{C}_b^1 character of the vector fields V_i .

Define the scale of spaces

$$E_n^\psi := \{ \varphi = \psi\phi; \phi \in L^\infty \}$$

with norm

$$\|\varphi\|_{E_n^\psi} := \|\phi\|_{W^{n,\infty}}$$

It is not difficult to check that $(VX, VV\mathbb{X})$ is a γ -rough driver also in this scale of spaces. In this case however we have

$$|h_t(\varphi)| = |h_t(\psi\phi)| \leq \|\phi\|_{L^\infty} |h_t(\psi)| = \|\varphi\|_{E_0^\psi} |z_t|$$

where

$$z_t := h_t(\psi),$$

and so

$$\langle h_\bullet \rangle_{(E_0^\psi)^*} \leq \langle z_\bullet \rangle$$

By the general a priori estimates we have that there exists λ and constants depending on A such that

$$\langle h^\sharp \rangle_{3\gamma; (E_3^\psi)^*} \lesssim \langle z_\bullet \rangle$$

But now

$$z_t = z_0 + h_0(V^*\psi)X_{0,t} + h_0(V^*V^*\psi)\mathbb{X}_{0,t} + h_{0,t}^\sharp(\psi)$$

so

$$\langle z_\bullet \rangle \leq |z_0| (1 + \langle X_{0,\bullet} \rangle + \langle \mathbb{X}_{0,\bullet} \rangle) + \langle h_{0,\bullet}^\sharp \rangle_{(E_3^\psi)^*}$$

and since $\langle h_{0,\bullet}^\sharp \rangle_{(E_3^\psi)^*} \leq \lambda^{3\gamma} \langle h^\sharp \rangle_{3\gamma; (E_3^\psi)^*} \leq \lambda^{3\gamma} \langle z_\bullet \rangle$, with h^\sharp considered as a 2-index map in its second occurrence, we have for λ small enough

$$(6.5) \quad \langle z_\bullet \rangle \leq 2|z_0| (1 + \langle X_{0,\bullet} \rangle + \langle \mathbb{X}_{0,\bullet} \rangle);$$

so $z_t = 0$, for all $t \geq 0$, if $z_0 = 0$.

Stability – Denote by $\mathbf{X}^{(n)}$ a smooth rough path converging to \mathbf{X} in the rough paths metric, and by $V_i^{(n)}$ a sequence of vector fields converging to V_i in \mathcal{C}_b^3 . Let $f_0^{(n)}$ be a smooth function converging to f_0 in $(\mathbb{L}^1)^*$. One solves the transport equation associated with $\mathbf{X}^{(n)}$ and $f_0^{(n)}$, using the elementary method of characteristics as the vector fields $V_i^{(n)}$ are sufficiently regular. It is elementary to use the uniform bound

$$\left\| f_t^{(n)} \right\|_{\mathbb{L}^\infty(\mathbb{R}^d)} \leq \left\| f_0^{(n)} \right\|_{\mathbb{L}^\infty(\mathbb{R}^d)} \leq C < \infty,$$

and the uniform a priori bound on $\|f^{(n),\sharp}\|_{3\gamma;-3}$ provided by theorem 9 and the convergence of $\mathbf{X}^{(n)}$ to \mathbf{X} , and $f_0^{(n)}$ to f_0 , to get the existence of a subsequence of $f_\bullet^{(n)}$ converging weakly- \star in $\mathbb{L}^\infty([0, T]; \mathbb{L}^\infty(\mathbb{R}^d))$ to some solution of the transport equation

(6.1), bounded in $\mathbb{L}^\infty(\mathbb{R}^d)$. Since this solution is unique, as proved above, the whole sequence $f_\bullet^{(n)}$ converges weakly- \star to f_\bullet in $\mathbb{L}^\infty([0, T]; \mathbb{L}^\infty(\mathbb{R}^d))$. As the same conclusion holds for $(f_\bullet^{(n)})^2$ and f^2 , by the renormalisation property, we classically get the convergence in $\mathbb{L}_{\text{loc}}^1([0, T]; \mathbb{L}^\infty(\mathbb{R}^d))$. \triangleright

REMARK 24. *It may be tempting, in the light of the results exposed in section 5.1, to try and develop an \mathbb{L}^∞ theory of differential equations driven by more general rough drivers $\mathbf{A}_{t,s}$ than those associated with the data of some vector fields V_1, \dots, V_ℓ and a weak geometric Hölder rough path over \mathbb{R}^ℓ , as in the transport equation (6.1). With a view towards the classical theory of stochastic flows, as developed by Le Jan-Watanabe, Kunita and many others, one may try, as a first step, to work with rough drivers whose first level are obtained as typical trajectories of semimartingale velocity fields. It is shown in [2] that such random fields can be lifted into some objects very similar to rough drivers, under some mild regularity conditions on the semimartingale, and that the use of the approximate flow machinery introduced in [1] leads to some well-posedness result for some dual evolution equation*

$$df_t(\varphi) = f_t(\mathbf{A}(dt)\varphi).$$

REFERENCES

- [1] Bailleul, I., Flows driven by rough paths. *Revista Mat. Iberoamericana*, 2015.
- [2] Bailleul, I. and Riedel, S., Rough flows. *Preprint*, 2014.
- [3] Bessaih, H. and Gubinelli, M. and Russo, F., The evolution of a random vortex filament. *Ann. Probab.*, 33(5):1825–1855, 2005.
- [4] Caruana, M. and Friz, P., Partial differential equations driven by rough paths. *J. Differential Equations*, 247(1):140–173, 2009.
- [5] Caruana, M. and Friz, P. and Oberhauser, H., A (rough) pathwise approach to a class of non-linear stochastic partial differential equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 28(1):27–46, 2011.
- [6] Catellier, R. and Gubinelli, M., Averaging along irregular curves and regularisation of ODEs. *arXiv:1205.1735*, 2014.
- [7] Catellier, R., Rough linear transport equation with an irregular drift. *Preprint*, 2014.
- [8] Coutin, L. and Lejay, A., Perturbed linear rough differential equations. *Annales Mathématiques Blaise Pascal*, 20(2):103–150, 2014.
- [9] Davie, A. M. Differential equations driven by rough paths: an approach via discrete approximation. *Appl. Math. Res. Express. AMRX*, (2), 2007.
- [10] de Lellis, C., ODEs with Sobolev coefficients: the Eulerian and the Lagrangian approach. *Discrete and cont. Dyn. Syst. Series S*, 1(3):405–426, 2008.
- [11] Deya, A. and Gubinelli, M., and Tindel, S., Non-linear rough heat equations. *Probab. Theory Related Fields*, 153(1-2):97–147, 2012.
- [12] Diehl, J. and Friz, P. and Stannat, W., Stochastic partial differential equations: a rough paths view. *arXiv:1412.6557v1*, 2014.
- [13] di Perna, R. and Lions, P.L., Ordinary differential equations, transport theory and Sobolev spaces. *Invent. mat.*, 98:511–547, 1989.
- [14] Feyel, D. and de La Pradelle, A., Curvilinear integrals along enriched paths. *Electron. J. Probab.*, 11:860–892, 2006.
- [15] Flandoli, F., The interaction between noise and transport mechanisms in PDEs, *Milan J. Math.* **79** (2011), no. 2, 543–560.
- [16] Friz, P. and Gess, B., Stochastic scalar conservation laws driven by rough paths. *arXiv:1403.6785v1*, 2014.
- [17] Friz, P. and Oberhauser, H., Rough path limits of a Wong-Zakai type with a modified drift term. *J. Funct. Anal.*, 256:3236–3256, 2009.

- [18] Friz, P. and Oberhauser, H., On the splitting-up method for rough (partial) differential equations. *J. Differential Equations*, 251:316–338, 2011.
- [19] Friz, P. and Oberhauser, H., Rough path stability of (semi-)linear SPDEs. *Probab. Theory Related Fields*, 158(1-2):401–434, 2014.
- [20] Friz, P., and Victoir, N., *Multidimensional stochastic processes as rough paths. Theory and applications*. Cambridge University Press, 2010.
- [21] Gess, B. and Souganidis, P.E., Scalar conservation laws with multiple rough fluxes. *arXiv:1406.2978v2*, 2014.
- [22] Gubinelli, M., Controlling rough paths. *J. Funct. Anal.*, 216(1):86–140, 2004.
- [23] Gubinelli, M., Rough solutions for the periodic Korteweg–de Vries equation. *Commun. Pure Appl. Anal.*, 11(2):709–733, 2012.
- [24] Gubinelli, M. and Imkeller, P. and Perkowski, N., Paracontrolled distributions and singular PDEs. *arXiv:1210.2684v3*, 2014.
- [25] Gubinelli, M. and Tindel, S., Rough evolution equations *Ann. Prob.*, 38:1–75, 2010.
- [26] Gubinelli, M. and Tindel, S. and Torrecilla, I., Controlled viscosity solutions of fully nonlinear rough PDEs *arXiv:1403.2832*, 2014.
- [27] Hairer, M., Solving the KPZ equation. *Annals Math.*, 178(2):559–664, 2013.
- [28] Hairer, M., A theory of regularity structures. *Invent. Math.*, 2014.
- [29] Hairer, M., Rough stochastic PDEs. *Comm. Pure Appl. Math.*, 64(11):1547–1585, 2011.
- [30] Hairer, M. and Maas, J. and Weber, H., Approximating rough stochastic PDEs. *arXiv preprint arXiv:1202.3094*, 2012.
- [31] Hairer, M. and Weber, H., Rough Burgers-like equations with multiplicative noise. *Probab. Theory Related Fields*, 155(1-2):71–126, 2013.
- [32] Hu, Y. and Le, K., Nonlinear Young integrals and differential systems in Hölder media *arXiv:1404.7582*, 2014.
- [33] Kruzhkov, S.N., First order quasilinear equations in several independent variables. *Math. USSR Sb.*, 10:217–243, 1970.
- [34] Lions, P.-L. and Perthame, B. and Souganidis, P.E., Scalar conservation laws with rough (stochastic) fluxes. *Stochastic Partial Differential Equations: Analysis and Computations*, 1(4):664–686, 2013.
- [35] Lions, P.-L. and Perthame, B. and Souganidis, P.E., Scalar conservation laws with rough (stochastic) fluxes: the spatially dependent case. *arXiv:1403.4424*, 2014.
- [36] Lyons, T., Differential equations driven by rough signals. *Rev. Matemática Iberoamericana*, 14(2):215–310, 1998.
- [37] Lyons, T.J. and Caruana, M. and Lévy, T., *Differential equations driven by rough paths*, volume 1908 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
- [38] Lyons, T. and Qian, Z., *System control and rough paths*. Oxford University Press, 2002.
- [39] Reed, L. and Simon, B., *Methods of modern mathematical physics*. Academic Press, 1980.

IRMAR, 263 AVENUE DU GENERAL LECLERC, 35042 RENNES, FRANCE
E-mail address: ismael.bailleul@univ-rennes1.fr

INSTITUT UNIVERSITAIRE DE FRANCE

CEREMADE & CNRS UMR 7534, UNIVERSITÉ PARIS DAUPHINE, PLACE DU MARÉCHAL DE
LATTRE DE TASSIGNY, 75775 PARIS CEDEX 16
E-mail address: massimiliano.gubinelli@ceremade.dauphine.fr