

# On a chain of fragmentation equations for duplication-mutation dynamics in DNA sequences

M.V. Koroteev<sup>1</sup>

<sup>1</sup>*School of Biochemistry and Cell Biology, University College Cork, Ireland\**

We study duplication-mutation models for processes in natural DNA sequences and length distributions of exact matches computed both from synthetic and natural sequences. For these models a hierarchy of equations for various number of exact matches is represented. It is demonstrated the reduction of these equations to one equation for pairs of exact repeats. Quantitative correspondence of solutions of the equation and simulations is demonstrated.

PACS numbers:

In recent years a series of duplication-mutation models related to processes occurring in natural DNA sequences received some attention in the literature[1–3]. The motivation for introducing these models were earlier empirical observations on length distribution[4] of identical repeats in natural DNA sequences[7, 8]. In part it was observed that when computing the length distributions within single chromosomes or whole genome sequences these distributions tended to exhibit power-law tails with the exponent close to  $-3$ [9]. These observations naturally drew attention to potential mechanisms accounting for them.

The first step in this direction was done in [1] where empirical computational models based on duplications were suggested. Duplications in that model for a synthetic chromosome as well as in the models that followed it were thought of as random events of copying and pasting a part of the chromosome. Each such event resulted in appearing a pair of identical sequences which then underwent further destruction by new duplication events and eventually disappeared but as the source of duplications generated new pairs at each time unit some balance of the number of duplicates might be expected. It was demonstrated that the model of random duplications results in forming length distributions of exact matches or maxmers[10] with power-law tails; it was also demonstrated that slope of these tails with the exponent  $-3$  can be obtained in the model by varying a parameter of the source of duplications. Here and in what follows by the source of duplications we imply a random mechanism of copying and pasting regions of a chromosome; it is characterized by several parameters, e.g., by the length of the region for copy-pasting which is chosen in accordance with some probability distribution.

The models independent of the source but incorporating additional mechanisms for generating exact repeats in the sequences were represented in [2, 3]. Two basic mechanisms utilized in the models, duplication as in [1] and point mutation represent those in natural chromosomes. It was demonstrated that length distributions[4]

of repetitive sequences simulated by the models correspond to those observed in natural chromosomes and that the form of those distributions also was close to algebraic with exponents, typically around  $-3$ . Thus the models in question were able to reproduce these exponents and even the amplitudes of the distributions were fitted[3] but unlike [1] the structure of the duplication source did not play an important role in those models.

The distinctive feature observed for the length distributions in [2] was the algebraic behavior for a broad range of parameters, while [3] demonstrated that when, simplistically speaking, mutations occurred as often as duplications the algebraic behavior disappeared; this point is discussed in more detail in [3]. In this paper we suggest dynamic equations reproducing both the exponent and the amplitude of the length distribution for pairs of exact matches and demonstrate its relation and the difference to the previous models. Our calculations demonstrate that the stationary equation reproducing the amplitude and the exponent for length distributions of pairs of exact repeats can be represented as a (infinite) sum of equations for different types of exact repeats which we discuss below.

The detailed explanation of the duplication-mutation models which we are interested in can be found, e.g., in [1, 3] but we resume them here for convenience; thus, this paper can be read independently of [3].

We look at a synthetic chromosome represented as a string of  $L$  bases chosen from a finite alphabet; in natural genomes the alphabet typically consists of four bases A, G, C, and T. The distance between bases is a length scale denoted by  $a$ ; for natural genomes it is close to  $1\text{\AA}$ . The fundamental sequence feature that we study is the set of repeated sequences within the chromosome, counted in an algorithm-independent way. The work [3] focused on ‘supermaximal repeats’(or ‘super maxmers’): sequence duplications neither copy of which is contained in any longer sequence duplication within the chromosome [1, 11]. However for the purposes of this paper we do not use the computation of supermaximal repeats; instead, we look at the output of the standard software for searching exact repeats in long sequences

---

\*Electronic address: maxim.koroteev@ucc.ie

called *mummer*[12]. *Mummer* searches maximal repeats or *maxmers*[10] in the sense similar to supermaxmers[13] but observations show that the output of these computations is different as it follows from comparison of the length distributions obtained in the models [2] and [3]. The question of accounting for this difference will be the subject of another paper: our aim here is the model capable to reproduce the simulated length distributions obtained with *mummer*. In the discussion below it is always implied that *mummer* is used with the option *-maxmatch* which according to *mummer* manual produces computations of exact matches ‘regardless their uniqueness’[14].

Within our models a subsequence of length  $m$  is chosen randomly within the chromosome according to a predetermined source distribution  $P(m)$  and is substituted for a sequence of length  $m$  at another randomly chosen position in the chromosome. As in [1], duplicates are realized not as fragments but as substrings within a single chromosomal polymer.

Let the number of pairs of duplicates of the length  $m$  at time moment  $t$  is  $g_2(t, m)$ . We assume that new duplication events occur with the rate  $\lambda$  per base, per time unit; in the same time the chromosome undergoes point mutation events occurring with the rate  $\mu$  per base, per time unit. Then the evolutionary (balance) equation for the average number of *pairs of duplicates*  $g_2$  can be written as [3]

$$\frac{\Delta g_2}{\Delta t} = -2 \left[ (m + D - a) \frac{a\lambda}{D} + \mu m \right] g_2(t, m) + 4 \left( \frac{a^2\lambda}{D} + a\mu \right) \sum_{k=m+1}^D g_2(t, k) + L \frac{a\lambda}{D} \delta_c(D - m). \quad (1)$$

The main difference of this equation from that of [3] is the alteration of notation (we use  $g_2$  here instead of  $f$ ). In addition, there is no prefactor 2 in the last term of the equation because in [3] we studied the number of duplicated *sequences* while here we look at the number of *pairs of duplicates*; thus, the source produces 1 pair of duplicates at each time step. We also we confine ourselves by the equation for the monoscale source using Kronecker delta function  $\delta_c(D - m)$ ; different source terms are also possible and will be presented elsewhere.

We will then focus on the stationary version of the equation implying that when  $t \rightarrow \infty$   $g_2(t, m) \rightarrow g_2(m)$  (this can be demonstrated by analytic calculation)

$$0 = -2 \left[ (m + D - a) \frac{a\lambda}{D} + \mu m \right] g_2(m) +$$

$$+ 4 \left( \frac{a^2\lambda}{D} + a\mu \right) \sum_{k=m+1}^D g_2(k) + L \frac{a\lambda}{D} \delta_c(D - m). \quad (2)$$

Now in the same way as we looked at *pairs of identical duplicates* we can look at triplets, quadruplets, etc. of identical sequences and write down the corresponding equations for them. For  $i$ -plets we will have the following stationary equation

$$0 = -i \left[ (m + D - a) \frac{a\lambda}{D} + \mu m \right] g_i(m) + 2i \left( \frac{a^2\lambda}{D} + a\mu \right) \sum_{k=m+1}^D g_i(k) + (i - 1) \left( \frac{a\lambda}{D} (D - m + a) \right) g_{i-1}(m) + 2(i - 1) \frac{a^2\lambda}{D} \sum_{k=m+1}^D g_{i-1}(k), \quad i > 2 \quad (3)$$

We see that unlike the equation for duplicates containing the source term with the delta function in it, other equations also have sources of new  $i$ -plets; these sources are  $i - 1$ -plets. Moreover, these sources have two terms: one produces  $i$ -plicates of  $i - 1$ -plicates of the same length  $m$  (the first term in the second line of (3)); the other generates  $i$ -plicates of longer  $i - 1$ -plicates by copying and pasting their parts of the length  $m$  (the second term in the second line of (3)), i.e., new duplicates  $g_2(m)$  gen-

erated by the source produce, in turn, triplicates  $g_3(m)$ , triplicates produce quadruplicates  $g_4$  etc. The first term in the first line of (3) is responsible for destruction of sequences by new duplications and point mutations; coefficients represent the corresponding rates. The second term in the first line of (3) shows that longer sequences are turned into shorter ones, again, by duplications and point mutations. The general mechanism has much in common with models studied in fragmentation




FIG. 1: Curves represent stationary length distributions obtained from simulations of duplication-mutation dynamics described in [3] with a monoscale source for various base substitution rates  $\mu$  and corresponding analytic solutions (orange) of (4). The chromosome length  $L = 10^6$ ; source length  $D = 10^3$ , duplication rate  $\lambda = 10^{-4}$ ; for simulations we always take  $a = 1$ . Length distributions for the same dynamics computed by mummer[12] were obtained using the following options `-maxmatch -n -b -l 20`. The results were then averaged over  $10^2$  realizations.

theory[15]. Thus for each  $m = 1 \dots D$  we have a set of equations for various sets of identical repeats (maxmers). We can sum up all the equations and find a new equation for the function  $G(m) = \sum i g_i(m)$ ; the equation has the form

$$-(\zeta + 2)mG(m) + 2aG(m) + 2(\zeta + 2)a \sum_{n>m} G(n) + L\delta_c(D - m) = 0, \quad (4)$$

where  $\zeta = D\mu/a\lambda$  is a dimensionless parameter[16].

Now we can compare the results of simulations with the solutions of (4); the comparison is represented in the fig. 1. Additional comparisons for different sets of parameters are given in supplemental figures (see Supplemental materials). Let us now compare solutions of the equation presented in [2] with the simulations of the same duplication-mutation dynamics. For that we used equation (5) of supplemental materials of [2]. Comparisons are represented in fig. 2. The solutions of [2] provide good agreement for sufficiently large mutation rates compared to duplication rate  $\lambda$  but fail to reproduce the amplitude of the length distributions for different regimes. In this regime it is observed saturation wrt. amplitude of the length distributions which is reproduced by solutions (4) as it is seen in fig. 1 and supplemental figures 1 and 2[17].

One then can easily understand the qualitative correspondence of length distributions observed in [2] and [3]: the growth of mutation rate  $\mu$  evidently affects  $g_i(m)$  for




FIG. 2: Curves represent stationary length distributions obtained from simulations of duplication-mutation dynamics with a monoscale source for various base substitution rates  $\mu$  and corresponding analytic solutions (magenta curves) of the eq. (5) of [2]. All parameters for simulations and the equation are the same as for fig. 1. The results of simulations were averaged over  $10^2$  realizations.

larger  $i$  as the growth of  $i$  means *more sequences* in the set. Thus the main contribution to  $G(m)$  comes from  $g_2(m)$ , i.e.,  $G(m) \sim g_2(m)$  as  $\zeta \rightarrow \infty$  and the dynamics is described by (2) in the main order. Also it is instructive to note that the situation  $\mu \gg \lambda$  generally implies  $\zeta \gg 1$  and one can neglect in (4) all terms compared to those containing  $\zeta$  and the source term with delta function to keep algebraic tail, hence  $L$  has to grow as  $\sim \zeta$ . However this is not applicable even for  $\zeta \sim 1$ . On the other hand, if  $\mu \ll \lambda$  then  $\zeta \rightarrow 0$  and we can write down the equation corresponding the limit of absent mutations

$$-2mG(m) + 2aG(m) + 4a \sum_{n>m} G(n) + L\delta_c(D - m) = 0. \quad (5)$$

If  $D$  is fixed as in figs. 1, 2, then the limit amplitude of the algebraic tail is controlled by the only parameter  $L$  and all distributions have the same saturation line (supplemental figure 3); this line establishes an upper boundary for fitting the model to the natural sequence. This also can be seen from the exact solution of (5) that has the form

$$G(m) = \begin{cases} \frac{aDL}{(m-a)m(m+a)}, & m < D \\ \frac{L}{2(D-a)}, & m = D \end{cases}$$

with obvious main order term  $\sim 1/m^3$  as  $a \ll m$ . The solution is applicable if  $a \ll D \ll L$ ; otherwise finite size effects turn out to be strong.

Finally we represent the correspondence of our dynamics for  $G(m)$  and natural DNA sequences. For that we plot fig. 3 in which there is a length distribution for *C. elegans chromosome 2* is portreyed. As all synthetic se-




FIG. 3: The length distribution for repeat-masked *C. elegans* chromosome 2 was computed using *mummer* with the options `-maxmatch -n -b -l 20`; self-hits were removed from the distribution. The length of the chromosome is  $\sim 10^7$ . The sequence of the same length was chosen for simulation with parameters  $D = 10^3$ ,  $\mu = 10^{-1}$ ,  $\lambda = 10^{-1}$ . The green line represents the solution of (4) for the same parameters.

quences when processed with *mummer* do not contain “self-hits”, i.e., identical sequences located exactly in the same positions for both copies of the chromosome, the self-hits were removed from the *mummer* output. A natural sequence does not provide information on the choice of  $D$ , though for monoscale source the algebraic regime is observed for  $m < D$ : thus we can have an upper bound

The equations (4) have several features deserving to be stressed. First of all, the equations we derived for  $G(m)$  allow to reproduce correctly the length distributions of exact matches computed by *mummer* in a broad range of parameters. That means, in part, that histograms computed by counting pairs of exact repeats with *mummer* can be understood as  $\sum ig_i(m)$  i.e., they consists of sums of all sequences of duplicates, triplicates, etc. It is worth noting that *mummer* output does not compute functions  $g_i(m)$  directly and thus the question of interpretation of  $g_i$  in terms of biologically meaningful sequences remains open: we observe only some cumulative effect of distributions for  $g_i(m)$ . On the other hand, the correspondence of functions  $g_2(m)$  to the length distribution of *supermaxmers* indicates a potential way to resolve this issue: if functions  $g_2(m)$  were interpreted as supermaxmers then the candidates for  $g_3(m)$ ,  $g_4(m)$  etc. could be so called ‘local maxmers’[18, 19]. In the same time the observed correspondence of *mummer* output and the function  $G(m)$  suggest we have an analytic interpretation for the length distributions computed by *mummer* for natural sequences: the length distributions for natural sequences exhibiting algebraic behaviour with the exponent  $-3$  can be understood in terms of the equations (3) and (4) and their solutions.

We proposed a hierarchy of equations for  $g_i$ , the first

of this equations, i.e. for  $g_2$  was derived in [3] and our observations here also demonstrate that the equations of [2] and [3] as well as those presented here treat different subjects. The work [3] deals with the collection of ‘supermaxmers’, specific pairs of exact repeats computed with additional conditions of maximality which are discussed in[18]; they are important as the equations for them not only account for the observed algebraic behaviour in length distributions of natural DNA sequences but demonstrate, in part, non-algebraic length distributions also observed both in simulations and natural DNA and also because their definition provides them with a natural biological interpretation[18]. Our equation (4) treats all pairs of exact matches neglecting their uniqueness and reproduces length distributions computed by *mummer*. Then  $G(m)$  in our interpretation may be represented as a sum of ‘supermaxmers’ for which biological interpretation was already discussed and other sets of sequences obtained by natural extension of the concept of supermaxmers; in this sense, we expect that such interpretation of  $g_i(m)$ ,  $m > 2$  is to appear soon.

The author is acknowledged to Kun Gao for helpful discussion.

- 
- [1] M.V. Koroteev, J. Miller. Phys. Rev. E 84, 061919 (2011)
  - [2] F. Massip, P.F. Arndt, Phys. Rev. Lett. 110, 148101 (2013).
  - [3] M.V. Koroteev, J. Miller. Fragmentation dynamics of DNA sequence duplication, preprint. arXiv:1304.1409v1
  - [4] The applications of length distributions in genomics have a long history (see [5, 6]). For the purposes of this paper a length distribution of matches or maxmers can be thought of as a histogram which has the length of maxmer on  $x$ -axis and the number of pairs (or triplets etc.) of identical matches on  $y$ -axis. We normally plot all length distributions in double log scale.
  - [5] S. Sawyer. Mol. Biol. Evol. 6(5), (1989).
  - [6] S.F. Altschul, W. Gish, W. Miller, E.W. Myers, D.J. Lipman. J. Mol. Biol. 215(3), (1990).
  - [7] W. Salerno, P. Havlak, J. Miller, Proc. Nat. Acad. Sci. USA, 103:13121 (2006).
  - [8] J. Miller, IPSJ SIG Technical Report No. 2009-BIO-17(7):1 (2009).
  - [9] K. Gao and J. Miller, PLoS One 6(7), (2011).
  - [10] There are multiple terms in the literature for exact copies of parts of DNA sequences; sometimes they are simply referred to as *exact matches*, in other cases it is important that exact matches are not contained in longer exact matches: in this case the term *maxmers* is used. Exact matches also can be referred to as *exact repeats*. For our purposes we do not distinguish these concepts and thus both terms are used interchangeably in the paper.
  - [11] E. Taillefer and J. Miller, in Proceedings of International Conference on Natural Computation, Shanghai, China, 2011, Vol. 3 (IEEE, New York, 2011), pp. 14801486; E. Taillefer, Miller, Int. Conf. on Computer Engineering and Bioinformatics, Cairo, Egypt, pp. 2229, 2011.

- [12] S. Kurtz, A. Phillippy, A.L. Delcher, M. Smoot, M. Shumway, C. Antonescu, and S.L. Salzberg, *Genome Biology* (2004), 5:R12.
- [13] both types of repeats include a condition of maximality of exact matches; this condition, however, turns out to be insufficient to obtain similar length distributions.
- [14] <http://mummer.sourceforge.net/manual/#maximal>
- [15] E. Ben-Naim, P.L. Krapivsky, *Phys. Lett. A*, 293(48), 2000. See also E. Ben-Naim, P.L. Krapivsky, *J. of Statistical Mechanics theory and experiment*, DOI: 10.1088/1742-5468/2005/10/L10002, 2005.
- [16] notice that the number of parameters was reduced from 5 to 3 compared to (3).
- [17] In the context of fig. 2 it is necessary to stress that com-

putations of [2] are different from those presented here as the authors of [2] use additional post-processing of *mummer* output as it is seen from the page 2 of supplemental materials of [2]. Therefore fig. 2 does not try to argue with the conclusions of [2] but only that these are different dynamics.

- [18] E. Taillefer, J. Miller. *J. Bioinformatics Comp. Biol.* 12(1), 2014.
- [19] Resolution of this issue will become possible when the software for computation of local maxmers is publicly available. The task of local maxmers computation and analysis is nontrivial and is out of the scope of the paper.

## Supplemental Materials: On a chain of fragmentation equations for duplication-mutation dynamics in DNA sequences

### SUPPLEMENTAL FIGURES




FIG. S1: Comparisons of simulations with solutions of the equation (4) of the main text. The parameters are:  $L = 10^6$ ,  $D = 10^3$ ,  $\lambda = 10^{-1}$ . Empirical length distributions were computed with the same switches of *mummer* as indicated in the caption for the figure 1 of the main text. The distributions were averaged over 100 realizations.




FIG. S2: Parameters of the model are:  $L = 10^6$ ,  $D = 10^4$ ,  $\lambda = 10^{-1}$ . All other parameters and options are the same as in the figure 1 of the main text and the supplemental figure 1.