

On Lawson's area-minimizing hypercones

Yongsheng ZHANG

School of Mathematics and Statistics, Northeast Normal University

ChangChun 130024, P.R. China

E-mail: yongsheng.chang@gmail.com

Abstract We show the area-minimality property of all homogeneous area-minimizing hypercones in Euclidean spaces (classified by Lawlor) following Lawson's original idea in his 72' Trans. A.M.S. paper "The equivariant Plateau problem and interior regularity". Moreover, each of them enjoys (coflat) calibrations singular only at the origin.

Keywords area-minimizing hypercone, coflat calibration, comass

MR(2010) Subject Classification 53C38, 28A75

1 Introduction

Let X be a Riemannian manifold and G a compact, connected group of isometries of X . In [6] Hsiang and Lawson showed that a G -invariant submanifold N of cohomogeneity k in X is minimal if and only if N^*/G is minimal in X^*/G with respect to a natural metric on the orbit space. Here $*$ means the union of principal orbits under action G . In particular they classified compact homogeneous minimal hypersurfaces in standard spheres.

In this paper $X = \mathbb{R}^{n+1}$ with Euclidean metric and M is a submanifold in the unit sphere S^n . We call

$$C(M) = \{tx : 0 \leq t < +\infty, x \in M\}$$

the cone over M and M the link of $C(M)$. It follows that M is minimal in S^n if and only if $C(M) \sim 0$ is minimal in \mathbb{R}^{n+1} .

In [8] Lawson first proved that there *always* exists a G -invariant solution to the Plateau problem for any G -invariant boundary M of codimension 2, and then he considered further which homogeneous minimal hypersurfaces in their classification give rise to area-minimizing cones, in the sense of Federer, namely the truncated cone by the unit ball centered at the origin is an area-minimizer among all integral currents (introduced in [3]) sharing the boundary M (these hypercones are closely related to the Bernstein problem, see [1, 13]). The question turns out to be equivalent to whether the line segment between the "0"-point and "M"-point in the orbit space is minimizing. He gained the following variety of minimizing hypercones by means of constructing calibration 1-forms on orbit spaces.

Received x x, 201x, accepted x x, 201x

Partially sponsored by the Fundamental Research Funds for the Central Universities, the SRF for ROCS, SEM, NSFC (Grant Nos. 11526048, 11601071) and the NSF (Grant No. 0932078 000) while the author was in residence at the MSRI during the 2013 Fall.

Theorem 1.1 (Lawson [8]) *There exist link manifolds M of types (I) $Sp(3)/Sp(1)^3$ in \mathbb{R}^{14} , $F_4/Spin(8)$ in \mathbb{R}^{26} , and moreover, (II) $S^{r-1} \times S^{s-1}$ in \mathbb{R}^{r+s} for $r, s \geq 2$ with $r + s \geq 10$ or with $r + s = 9$ and $|r - s| \leq 5$ or with $r = s = 4$, $SO(2) \times SO(k)/\mathbb{Z}_2 \times SO(k-2)$ in \mathbb{R}^{2k} for $k \geq 10$, $SU(2) \times SU(k)/T^1 \times SU(k-2)$ in \mathbb{R}^{4k} for $k \geq 5$, $U(5)/SU(2) \times SU(2) \times T^1$ in \mathbb{R}^{20} , $U(1) \cdot Spin(10)/T^1 \cdot SU(4)$ in \mathbb{R}^{32} , and $Sp(2) \times Sp(k)/Sp(1)^2 \times Sp(k-2)$ in \mathbb{R}^{8k} for $k \geq 2$, such that $C(M)$ is area-minimizing.*

Remark 1.2 Here we put overlapping types of (I) and (II) in [8] into class (II).

Remark 1.3 For example, cones $C_{m,n} \triangleq C(S^m(\sqrt{\frac{m}{m+n}}) \times S^n(\sqrt{\frac{n}{m+n}}))$ where $m = r - 1$ and $n = s - 1$ for r, s in the theorem are area-minimizing.

From the point of view of calibrated geometry, the calibration on the orbit space induces a coflat calibration in the Euclidean space (see §4) that is singular in a set of codimension at least two. Hence, by the coflat version of fundamental theorem of calibrated geometry (Theorem 4.9 in [5]), homogeneous hypercones with such calibrations are area-minimizing. In fact similar idea also can work for cones of higher codimension, for example, see [2].

By studying Lawson's calibrations, we give a proof of the *affirmative* part of the table on page 86 (classification of homogeneous area-minimizing hypercones) in [7] by verifying the rest cases.

Theorem 1.4 *M of class (II) in the above theorem can also be of type $S^2 \times S^4$ in \mathbb{R}^8 , $SO(2) \times SO(k)/\mathbb{Z}_2 \times SO(k-2)$ in \mathbb{R}^{2k} for $k = 9$, and $SU(2) \times SU(k)/T^1 \times SU(k-2)$ in \mathbb{R}^{4k} for $k = 4$, such that $C(M)$ is area-minimizing. Moreover, types of $S^1 \times S^5$ and $SO(2) \times SO(8)/\mathbb{Z}_2 \times SO(6)$ correspond to stable minimal hypercones.*

Remark 1.5 The area-minimality of $C_{2,4}$ in \mathbb{R}^8 was given by Simoes [11, 12].

According to Hardt and Simon [4], here comes a corollary.

Corollary 1.6 *All homogeneous area-minimizing hypercones are strictly area-minimizing.*

Remark 1.7 The strict area-minimality of $C_{2,4}$ was first proved by Lin [9].

We obtain the next theorem through a particular kind of modifications on Lawson's calibrations.

Theorem 1.8 *Each area-minimizing $C_{r-1,s-1} \subset \mathbb{R}^{r+s}$ has a coflat calibration singular only at the origin.*

An observation on the calibration O.D.E. (6.1) leads to a more general result.

Theorem 1.9 *Every homogeneous area-minimizing hypercone has a coflat calibration singular only at the origin.*

It may be interesting to consider whether every area-minimizing hypercone has such nice calibrations. See [17] for applications of these calibrations.

2 Preliminaries

Suppose a connected compact Lie group G has an orthogonal representation on X of cohomogeneity 2 and M_0 is the principal orbit of greatest volume in S^n . Then a celebrated classification due to [6] and [15] is the following table, based on which much interesting research has been

done, for example, recent papers [16] and [10]. Here we only list the needed information and refer readers to their original papers for details.

	G	type of M_0	A	V^2
1	$SO(r) \times SO(s)$	$S^{r-1} \times S^{s-1}$ $r, s \geq 2$	$\frac{\pi}{2}$	$c \cdot x^{2r-2} y^{2s-2}$
2	$SO(2) \times SO(k)$	$\frac{SO(2) \times SO(k)}{\mathbb{Z}_2 \times SO(k-2)}$ $k \geq 3$	$\frac{\pi}{4}$	$c \cdot (xy)^{2k-4} (x^2 - y^2)^2$
3	$SU(2) \times SU(k)$	$\frac{SU(2) \times SU(k)}{T^1 \times SU(k-2)}$ $k \geq 2$	$\frac{\pi}{4}$	$c \cdot (xy)^{4k-6} (x^2 - y^2)^4$
4	$Sp(2) \times Sp(k)$	$\frac{Sp(2) \times Sp(k)}{Sp(1)^2 \times Sp(k-2)}$ $k \geq 2$	$\frac{\pi}{4}$	$c \cdot (xy)^{8k-10} (x^2 - y^2)^8$
5	$U(5)$	$\frac{U(5)}{SU(2) \times SU(2) \times T^1}$	$\frac{\pi}{4}$	$c \cdot (xy)^2 \text{Im}\{(x + iy)^4\}^8$
6	$U(1) \cdot Spin(10)$	$\frac{U(1) \cdot Spin(10)}{T^1 \cdot SU(4)}$	$\frac{\pi}{4}$	$c (xy)^6 \text{Im}\{(x + iy)^4\}^{12}$
7	$SO(3)$	$\frac{SO(3)}{\mathbb{Z}_2 + \mathbb{Z}_2}$	$\frac{\pi}{3}$	$c \cdot \text{Im}\{(x + iy)^3\}^2$
8	$SU(3)$	$\frac{SU(3)}{T^2}$	$\frac{\pi}{3}$	$c \cdot \text{Im}\{(x + iy)^3\}^4$
9	$Sp(3)$	$\frac{Sp(3)}{Sp(1)^3}$	$\frac{\pi}{3}$	$c \cdot \text{Im}\{(x + iy)^3\}^8$
10	F_4	$\frac{F_4}{Spin(8)}$	$\frac{\pi}{3}$	$c \cdot \text{Im}\{(x + iy)^3\}^{16}$
11	$Sp(2)$	$\frac{Sp(2)}{T^2}$	$\frac{\pi}{4}$	$c \cdot \text{Im}\{(x + iy)^4\}^4$
12	G_2	$\frac{G_2}{T^2}$	$\frac{\pi}{6}$	$c \cdot \text{Im}\{(x + iy)^6\}^4$
13	$SO(4)$	$\frac{SO(4)}{\mathbb{Z}_2 + \mathbb{Z}_2}$	$\frac{\pi}{6}$	$c \cdot \text{Im}\{(x + iy)^6\}^2$

Table 1 Minimal homogeneous hypersurfaces in spheres

Each orbit space is a cone of angle A in \mathbb{R}^2 endowed with metric

$$ds^2 = V^2(dx^2 + dy^2),$$

where V is the volume function of orbits. Let π be the projection map to orbit space. Then it can be shown that (up to a constant factor) the volume of any G -invariant hypersurface N equals the length of $\pi(N)$.

In order to search for area-minimizing cone $C(M_0)$, Lawson made the following try for class **(II)**, i.e., Row 1 – 6 in the table.

First note that $A = \frac{\pi}{4}$ for Row 2–6. Consider $re^{i\theta} = z^2$ with $z = x + yi$. Now the orbit space is stretched to a cone Q of angle $\frac{\pi}{2}$ with metrics $r^{2k-3}s^{2k-4}c^2(r^2d\theta^2 + dr^2)$, $r^{4k-3}s^{4k-6}c^4(r^2d\theta^2 + dr^2)$, $r^{8k-3}s^{8k-10}c^8(r^2d\theta^2 + dr^2)$, $r^{17}s^{10}c^8(r^2d\theta^2 + dr^2)$ and $r^{29}s^{18}c^{12}(r^2d\theta^2 + dr^2)$ respectively. Here $c = \cos \theta$ and $s = \sin \theta$. For metric $ds^2 = r^l c^p s^q (r^2 d\theta + dr^2)$ on Q (including Row 1), the angle θ_0 corresponding to M_0 satisfies $\tan^2 \theta_0 = q/p$. Set $\tau = \cos^p \theta_0 \sin^q \theta_0$. Then, under $d\tilde{s}^2 = \frac{1}{\tau} ds^2$ on Q , Lawson considered the function of type

$$f = \alpha^{-1} r^\alpha \phi^\beta, \quad (2.1)$$

where $2\alpha = l + 2$, $\phi = \frac{c^p s^q}{\tau}$ and β is some real number to be determined. Since

$$\phi'(\theta) = \phi(\theta)(q \cot \theta - p \tan \theta),$$

it follows

$$df = r^{\alpha-1} \phi^\beta dr + \left(\frac{\beta}{\alpha}\right) r^\alpha \phi^\beta (q \cot \theta - p \tan \theta) d\theta.$$

Note that for a 1-form its comass (see [5]) and usual norm coincide. Therefore if for some β

$$r^{2\alpha-2}\phi^{2\beta}\left[1 + \left(\frac{\beta}{\alpha}\right)^2(q \cot \theta - p \tan \theta)^2\right] \leq \frac{r^l c^p s^q}{\tau}, \quad (2.2)$$

or simply

$$\phi^{2\beta-1}\left[1 + \left(\frac{\beta}{\alpha}\right)^2(q \cot \theta - p \tan \theta)^2\right] \leq 1, \quad (2.3)$$

then df is a calibration 1-form, smooth on Q° and continuous to ∂Q , calibrating the ray $\theta = \theta_0$, and hence $C(M_0)$ is area-minimizing. Denote the left-hand side of (2.3) by ψ . It is clear that ψ is pointwise the square of comass of df with respect to $d\bar{s}^2$.

3 Proof of Theorem 1.4

Consider the derivative of ψ .

$$\begin{aligned} \psi' = \phi^{2\beta-1}(q \cot \theta - p \tan \theta) & \left\{ (2\beta - 1) \left[1 + \left(\frac{\beta}{\alpha}\right)^2(q \cot \theta - p \tan \theta)^2 \right] \right. \\ & \left. + 2\left(\frac{\beta}{\alpha}\right)^2(q \cot \theta - p \tan \theta)' \right\} \end{aligned}$$

Let η_β stand for the part in braces. Then

$$\begin{aligned} \eta_\beta &= (2\beta - 1) \left[1 + \left(\frac{\beta}{\alpha}\right)^2(q \cot \theta - p \tan \theta)^2 \right] + 2\left(\frac{\beta}{\alpha}\right)^2\left(-\frac{q}{s^2} - \frac{p}{c^2}\right) \\ &= (2\beta - 1) \left[1 + \left(\frac{\beta}{\alpha}\right)^2(q^2 \cot^2 \theta + p^2 \tan^2 \theta - 2pq) \right] + 2\left(\frac{\beta}{\alpha}\right)^2\left(-\frac{q}{s^2} - \frac{p}{c^2}\right) \\ &= (2\beta - 1) - 2\left(\frac{\beta}{\alpha}\right)^2 [(2\beta - 1)pq + p + q] \\ & \quad + \left(\frac{\beta}{\alpha}\right)^2 \left\{ [(2\beta - 1)q^2 - 2q] \cot^2 \theta + [(2\beta - 1)p^2 - 2p] \tan^2 \theta \right\} \end{aligned}$$

Note that $\psi(\theta_0) = 1$. So for a triple (p, q, α) if there exists some β such that

$$\eta_\beta > 0 \text{ on } Q^\circ, \quad (3.1)$$

then (2.3) holds on Q .

A1. For simplicity, we try $\beta = 1$ for Row 1. Now

$$\begin{aligned} 2\alpha &= \Sigma + 2 \text{ where } \Sigma = p + q, \quad p = 2r - 2, \text{ and } q = 2s - 2. \\ 4\alpha^2\eta_1 &= 4\alpha^2 - 8(p + q + pq) + 4(q^2 - 2q) \cot^2 \theta + 4(p^2 - 2p) \tan^2 \theta \\ &\geq 4\alpha^2 - 8(p + q + pq) + 8\sqrt{(q^2 - 2q)(p^2 - 2p)} \quad (\text{“} > \text{” unless } p = q = 2) \\ &\geq 4\alpha^2 - 8(p + q + pq) + 8(q - 2)(p - 2) \\ &\geq (\Sigma + 2)^2 - 24\Sigma + 32 \\ &= (\Sigma - 2)(\Sigma - 18) \end{aligned} \quad (3.2)$$

Thus, when $2(r + s) - 4 = \Sigma \geq 18$, namely $r + s \geq 11$, we have $\eta_1 > 0$.

Actually from the first row of (3.2) one can get

$$\begin{aligned} 4\alpha^2\eta_1 \tan^2 \theta &= 4(p^2 - 2p) \tan^4 \theta + [4\alpha^2 - 8(p + q + pq)] \tan^2 \theta + 4(q^2 - 2q) \\ &= 4(p^2 - 2p)Y^2 + [4\alpha^2 - 8(p + q + pq)]Y + 4(q^2 - 2q) \end{aligned}$$

where $Y = \tan^2 \theta$. Easy to check that Δ of the quadratic is negative for $r, s \geq 3$ with $8 \leq r + s \leq 10$. So $\eta_1 > 0$. In particular $C_{2,4}$ for $(r, s) = (3, 5)$ is area-minimizing.

Due to Simons [14] and Simoes [11, 12] there are no area-minimizing cones for $r + s < 8$ or for $(r, s) = (2, 6)$. Therefore $(r, s) = (2, 7)$ and $(2, 8)$ are the only left cases to check. Although df cannot be a global calibration on Q for $(r, s) = (2, 6)$, it is indeed a local calibration in a very *thin* angular neighborhood of the ray $\theta = \theta_0$, e.g. $1.145 \leq \theta \leq 1.165$ shown in Figure 1 below. (At $\theta_0 = \arctan \sqrt{5} \approx 1.150262$, $\psi = 1$.) In fact the existence of such angular neighborhood can be seen from $\eta_1(\theta_0) > 0$. As a consequence, $C_{1,5}$ is a stable minimal hypercone in \mathbb{R}^8 .




Figure 1 Graph of ψ for $(r, s) = (2, 6)$ with $\beta = 1$.

A2. For the rest two cases as well as for our purpose later in C2., we consider $\beta = 1.2 > 1$. Similarly, it can be shown that $\eta_{1,2} > 0$ when $r, s \geq 2$ and $r + s \geq 9$.

B. For $SO(2) \times SO(k)/\mathbb{Z}_2 \times SO(k-2)$ in \mathbb{R}^{2k} for $k = 9$ (with $\alpha = 8.5$) one can use $\beta = 1.2$. There are two roots $\theta_{1,2}$ of $\eta_{1,2}$ satisfying $\theta_0 < \theta_1 < \theta_2$. Since $\psi(\theta_2) < 1$, df is a calibration on Q .




Figure 2 Graph of ψ with $\beta = 1.2$.

For $SO(2) \times SO(k)/\mathbb{Z}_2 \times SO(k-2)$ in \mathbb{R}^{2k} for $k = 8$ (with $\alpha = 7.5$), using $\beta = 1$, there is only one root θ_1 of η_1 and $\theta_1 > \theta_0$. So df is a local calibration in some angular neighborhood of $\theta = \theta_0$, for example $|\theta - \theta_0| < \theta_1 - \theta_0$.




Figure 3 Graph of ψ with $\beta = 1$.

Due to technique reason we save the the discussion on $SU(2) \times SU(k)/T^1 \times SU(k-2)$ in \mathbb{R}^{4k} for $k = 4$ to §5.

4 Proof of Theorem 1.8

By explanations on page 62 and Theorem 4.13 in [5], up to a constant,

$$\omega_f \triangleq \pi^*(df) \wedge \Omega, \quad (4.1)$$

where $\Omega = \Omega_0/V$ and Ω_0 is the oriented unit volume form of principal orbits, is a coflat calibration of the cone, smooth away from singular orbits in the Euclidean space.

In this section, we shall show that, for an area-minimizing hypercone in Row 1, either ω or its suitable modification is a calibration smooth away from the origin.

Observe that the principal orbit through $(\vec{X}, \vec{Y}) \in \mathbb{R}^r \times \mathbb{R}^s$, where neither \vec{X} nor \vec{Y} is a null vector, is $S^{r-1}(\|\vec{X}\|) \times S^{s-1}(\|\vec{Y}\|)$. Therefore

$$\Omega_0^{r,s} = \Omega_r(\vec{X}) \wedge \Omega_s(\vec{Y}), \text{ where}$$

$$\Omega_r(\vec{X}) = \frac{1}{\|\vec{X}\|} \sum_{i=1}^r (-1)^{i-1} x_i dx_1 \wedge \cdots \wedge \widehat{dx_i} \wedge \cdots \wedge x_r,$$

$$\Omega_s(\vec{Y}) = \frac{1}{\|\vec{Y}\|} \sum_{j=1}^s (-1)^{j-1} y_j dy_1 \wedge \cdots \wedge \widehat{dy_j} \wedge \cdots \wedge y_s,$$

and $\{x_i\}, \{y_j\}$ are standard basis of \mathbb{R}^r and \mathbb{R}^s respectively.

C1. When r and s are even integers with $r, s \geq 4$ in A1. ($\beta = 1$), there is a nonzero

constant C such that

$$\begin{aligned}
C \cdot \omega_f &= C \cdot d(\pi^* f) \wedge \frac{\Omega_0^{r,s}}{V} \\
&= d \left[R^{3-r-s} \cdot \|\vec{X}\|^{2r-2} \cdot \|\vec{Y}\|^{2s-2} \right] \wedge \frac{\Omega_0^{r,s}}{V} \quad \left(\text{Here } R = \sqrt{\|\vec{X}\|^2 + \|\vec{Y}\|^2} \right) \\
&= \left\{ R^{3-r-s} \left[(r-1) \|\vec{X}\|^{2r-4} \|\vec{Y}\|^{2s-2} d\|\vec{X}\|^2 + (s-1) \|\vec{X}\|^{2r-2} \|\vec{Y}\|^{2s-4} d\|\vec{Y}\|^2 \right] \right. \\
&\quad \left. + (3-r-s) R^{2-r-s} \|\vec{X}\|^{2r-2} \cdot \|\vec{Y}\|^{2s-2} dR \right\} \\
&\quad \wedge \frac{1}{\|\vec{X}\|^r} \left(\sum_{i=1}^r (-1)^{i-1} x_i dx_1 \wedge \cdots \wedge \widehat{dx}_i \wedge \cdots \wedge x_r \right) \\
&\quad \wedge \frac{1}{\|\vec{Y}\|^s} \left(\sum_{j=1}^s (-1)^{j-1} y_j dy_1 \wedge \cdots \wedge \widehat{dy}_j \wedge \cdots \wedge y_s \right) \\
&= \|\vec{X}\|^{r-4} \|\vec{Y}\|^{s-2} \cdot \Lambda_1 + \|\vec{X}\|^{r-2} \|\vec{Y}\|^{s-4} \cdot \Lambda_2 + \|\vec{X}\|^{r-2} \|\vec{Y}\|^{s-2} \cdot \Lambda_3
\end{aligned}$$

where Λ_i are smooth forms away from the origin. Hence so is ω_f .

C2. For remaining cases of Row 1 in A1. and A2., each reduced ray has a calibration df where $f = C\alpha^{-1}r^\alpha c^k s^t$ for some positive constant C and integers k, t with $2k - p > 2$ and $2t - q > 2$. (For such f one can use $\beta = 1$ when $p, q > 2$ and $\beta = 1.2$ when either p or q equals 2.) Inspired by C1., we wish to deform f to \tilde{f} so that $\omega_{\tilde{f}}$ behaves similarly as in C1. near singular orbits and $d\tilde{f}$ is a calibration of the cone.

Our strategy is to increase the exponential of f . We illustrate how to remove the singularity along $\theta = 0$ (with $r > 0$). Choose an integer N such that $N - \frac{q}{2} - 1$ (which will replace the exponential $s - 2$ of $\|\vec{Y}\|$ in C1.) is a positive even integer and $N > t$. Then search for a suitable smooth function λ of θ for the deformation:

$$\tilde{f} = C\alpha^{-1}r^\alpha c^k s^{t+\lambda}. \quad (4.2)$$

We wish $\lambda(\theta) \equiv N - t$ when θ is very small (for $\lambda = \lambda_{\epsilon_2}$ that we are about to construct, such an interval of θ can be $(0, \frac{\pi_0}{2} - \epsilon_2)$ introduced below) and $\lambda(\theta) \equiv 0$ when $\theta > \tilde{\theta}$ for some small positive $\tilde{\theta}$. By computation

$$d\tilde{f} = Cr^{\alpha-1}c^k s^{t+\lambda} dr + C\alpha^{-1}r^\alpha c^k s^{t+\lambda} (\lambda' \ln s + (t + \lambda) \cot \theta - k \tan \theta) d\theta$$

and similar to (2.3) the square of comass of $d\tilde{f}$ equals

$$\tau C^2 c^{2k-p} s^{2t+2\lambda-q} \left[1 + \left(\frac{1}{\alpha} \right)^2 (\lambda' \ln s + (t + \lambda) \cot \theta - k \tan \theta)^2 \right]. \quad (4.3)$$

In view of the square of comass of calibration df ,

$$\tau C^2 c^{2k-p} s^{2t-q} \left[1 + \left(\frac{1}{\alpha} \right)^2 (t \cot \theta - k \tan \theta)^2 \right] \leq 1 \text{ on } Q,$$

one only needs to show (4.3) is no large than one on the support of λ for $d\tilde{f}$ being a calibration.

We consider the following Lipschitzan function for a fixed $0 < x < \tilde{\theta}$:

$$\lambda_x(\theta) = \begin{cases} N - t & 0 \leq \theta \leq \frac{x}{2} \\ (N - t) \left[1 - \frac{1}{\int_{\frac{x}{2}}^x \frac{1}{\ln \sin \gamma} d\gamma} \cdot \int_{\frac{x}{2}}^{\theta} \frac{1}{\ln \sin \gamma} d\gamma \right] & \frac{x}{2} < \theta < x \\ 0 & x \leq \theta \end{cases}$$

By this choice (4.3) on $\frac{x}{2} < \theta < x$ becomes

$$\tau C^2 c^{2k-p} s^{2t+2\lambda_x-q} \left[1 + \left(\frac{1}{\alpha} \right)^2 \left(-\frac{N-t}{\int_{\frac{x}{2}}^x \frac{1}{\ln \sin \gamma} d\gamma} + (t + \lambda_x) \cot \theta - k \tan \theta \right)^2 \right]. \quad (4.4)$$

Focus on the behavior of (4.4) as x tends to zero. By the assumption $2t + 2\lambda_x - q \geq 2\rho \triangleq 2t - q > 2$, the expression limits to (for $\tilde{C}^2 = \frac{\tau C^2}{\alpha^2}$)

$$\lim_{x \rightarrow 0} \tilde{C}^2 s^{2t+2\lambda_x-q} \left(\frac{N-t}{\int_{\frac{x}{2}}^x \frac{1}{\ln \sin \gamma} d\gamma} \right)^2 \leq \tilde{C}^2 (N-t)^2 \left(\lim_{x \rightarrow 0} \frac{\sin^\rho x}{\int_{\frac{x}{2}}^x \frac{1}{\ln \sin \gamma} d\gamma} \right)^2. \quad (4.5)$$

Since $\int_0^a \frac{1}{\ln \sin \gamma} d\gamma$ is convergent for every $0 < a < \frac{\pi}{2}$, the denominator of the latter limit in parentheses goes to zero and one can apply L'Hôpital's rule. Due to

$$\lim_{x \rightarrow 0} \frac{\rho(2 \sin \frac{x}{2} \cos \frac{x}{2})^{\rho-1} \cos x}{\ln(2 \sin \frac{1}{2} \cos \frac{x}{2}) - \ln \sin \frac{x}{2}} = \lim_{x \rightarrow 0} \frac{\rho(2 \sin \frac{x}{2} \cos \frac{x}{2})^{\rho-1} \cos x}{-\ln 2 - \ln \cos \frac{x}{2}} = 0$$

the left hand side of (4.5) is zero (strictly less than one).

Not hard to see there exists a smooth approximation λ of some λ_x such that (4.3) does not exceed one. One construction can be given as follows. Choose x_0 small enough so that, for some small positive $\epsilon_1 < \frac{x_0}{5}$, the following analogous expression to (4.4)

$$\tau C^2 c^{2k-p} s^{2\rho} \left[1 + \left(\frac{1}{\alpha} \right)^2 \left(-\chi \cdot \frac{N-t}{\int_{\frac{x_0}{2}}^{x_0} \frac{1}{\ln \sin \gamma} d\gamma} + K \cot \theta - k \tan \theta \right)^2 \right] \quad (4.6)$$

is less than half of one pointwise for $\frac{x_0}{2} - \epsilon_1 \leq \theta \leq x_0 + \epsilon_1$, any $\chi \in [0, 2]$ and $K \in [0, N+1]$. Let λ_ϵ be an integral convolution of λ_{x_0} by a compactly supported, smooth, *even* mollifier function with averaging radius $\epsilon (< \epsilon_1)$. Define σ^ϵ by

$$\lambda'_\epsilon(y) = -\sigma^\epsilon(y) \frac{N-t}{\int_{\frac{x_0}{2}}^{x_0} \frac{1}{\ln \sin \gamma} d\gamma} \cdot \frac{1}{\ln \sin y} \quad \text{for } \frac{x_0}{2} - \epsilon_1 \leq y \leq x_0 + \epsilon_1.$$

Then σ^ϵ is a smooth function with

$$\lim_{\epsilon \rightarrow 0} \sigma^\epsilon(y) = \begin{cases} 1 & \text{for } \frac{x_0}{2} \leq y \leq x_0 \\ \frac{1}{2} & \text{at } x_0 \text{ or } \frac{x_0}{2} \\ 0 & \text{on } [\frac{x_0}{2} - \epsilon_1, \frac{x_0}{2}) \cup (x_0, x_0 + \epsilon_1] \end{cases}$$

Hence through a contradiction argument there exists some small $\epsilon_2 (< \epsilon_1)$ such that $\sigma^{\epsilon_2}(y) \leq 2$ on $[\frac{x_0}{2} - \epsilon_1, x_0 + \epsilon_1]$. Comparing (4.3) and (4.6) shows that $\lambda = \lambda_{\epsilon_2}$ meets our needs.

According to the discussion in C1. and (4.1), $N - \frac{q}{2} - 1$ being a positive even integer asserts the smoothness of $\omega_{\vec{f}}$ along $\{(\vec{X}, \vec{Y}) \in \mathbb{R}^r \times \mathbb{R}^s : \vec{X} = \vec{0} \text{ and } \vec{Y} \neq \vec{0}\}$. The same trick applies for removing the singularity along $\{(\vec{X}, \vec{Y}) \in \mathbb{R}^r \times \mathbb{R}^s : \vec{X} \neq \vec{0} \text{ and } \vec{Y} = \vec{0}\}$. Hence we finish the proof.

5 The Case of $SU(2) \times SU(k)/T^1 \times SU(k-2)$ in \mathbb{R}^{4k} for $k = 4$

In virtue of *Mathematica* it can be observed that the construction (2.1) cannot directly produce a calibration for any value β in this case. Instead we fix $\beta = 1$ and look at the graph of $\psi = \frac{1}{\tau} c^4 s^{10} \left[1 + \left(\frac{1}{7.5}\right)^2 (10 \cot \theta - 4 \tan \theta)^2 \right]$.




Figure 4

Since the second peak is relatively not quite high, we try to chop it off. To do so, we alter the order of term c by setting

$$f_1 = \frac{1}{7.5\tau} r^{7.5} c^{4+\lambda_1} s^{10},$$

where λ_1 is a function. Then the square of the comass of df_1 becomes

$$(\star) \quad \frac{1}{\tau} c^{4+2\lambda_1} s^{10} \left[1 + \left(\frac{1}{7.5}\right)^2 (\lambda_1' \ln c + 10 \cot \theta - (4 + \lambda_1) \tan \theta)^2 \right].$$

Notice that θ_0 (at where the first peak occurs) is less than 1.00686 and its local behavior is shown as follows:




Figure 5

Now we consider the ODE:

$$\lambda_1' = \frac{7.5 \sqrt{\frac{1}{c^{4+2\lambda_1} s^{10} (\frac{14}{4})^2 (\frac{14}{10})^5} - 1 + 10 \cot \theta - (4 + \lambda_1) \tan \theta}}{-\ln c},$$

with $\lambda_1(1.007) = 0$. Its solution tells us how to vary the exponential of c to keep the corresponding comass exactly one on $[1.007, 1.20]$. Running the following codes in *Mathematica*

```
sl =NDSolve[{y'[x] ==
((7.5)(1/(Cos[x]^ (4 + 2y[x])Sin[x]^ (10)(14/4)^2(14/10)^5) - 1)^(0.5)
+10Cot[x] - (4 + y[x])Tan[x])/(-Log[Cos[x]]),y[1.007] == 0},y,{x,1.007,1.2}]
```

```
Plot[Evaluate[y[x]/.sl], {x,1.007,1.2}]
```

we figure out the graph of λ_1 .




Figure 6

Denote by θ_1 the zero point of λ_1 near 1.2. Note that around 1.007 the integral curves Λ of the slope field and the horizontal lines form a coordinate chart. Let the integral curve through $(1.007, 0)$ be $\Lambda_{1.007}$. Then one can glue 0 and $\Lambda_{1.007}$ such that on the left side of $\Lambda_{1.007}$ it transversally intersects each of Λ at only one point. Do the same kind of gluing around θ_1 .




Figure 7

Now we get a smooth variation λ of λ_1 that vanishes near 1.00686 and 1.2. Since $10 \cot \theta - 4 \tan \theta < 0$ on $(\theta_0, 1.2]$, after replacing λ_1 by λ , (\star) is bounded from above by one. Hence we complete the proof of Theorem 1.4.

6 Proof of Theorem 1.9

Let us get back to the original calibration O.D.E. (or O.D.InE.) on the (stretched) orbit space Q of angle $\frac{\pi}{2}$ for area-minimizing hypercones of type **(II)**. Set

$$f = \alpha^{-1} r^\alpha y(\theta)$$

similar to (2.1), then (2.2) becomes (w.r.t. $d\tilde{s}^2$)

$$y^2 + \left(\frac{y'}{\alpha}\right)^2 \leq \frac{c^p s^q}{\tau}. \tag{6.1}$$

Remark. For Row 9-10 of type **(I)** with $V^2 = c \cdot \text{Im} \{(x + iy)^p\}^q$ and $A = \frac{\pi}{p}$. On the (stretched) orbit space Q of angle π with respect to (up to a constant factor)

$$ds^2 = r^{2\alpha-2} s^q (r^2 d\theta + dr^2) \quad \text{where} \quad 2\alpha - 2 = q - \frac{2p - 2}{p},$$

the corresponding calibration O.D.E. is

$$y^2 + \left(\frac{y'}{\alpha}\right)^2 \leq s^q. \tag{6.2}$$

In this case $\theta_0 = \frac{\pi}{2}$. See [8] for details.

Take type **(II)** for example (exactly the same for **(I)**). For each area-minimizing hypercone of **(II)**, (6.1) has a smooth solution Φ on $I = (0, \frac{\pi}{2})$ with value one at $\theta = \theta_0$ and value zero at the ending points of I . Consider the slope fields Γ_1, Γ_2 of

$$y'_1 = \alpha \sqrt{\frac{c^p s^q}{\tau} - y_1^2} \quad \text{on} \quad (0, \theta_0) \tag{6.3}$$

and

$$y'_2 = -\alpha \sqrt{\frac{c^p s^q}{\tau} - y_2^2} \quad \text{on} \quad I - (0, \theta_0]. \tag{6.4}$$

We observe (A) along the graph of Φ , Γ_1 is positively deeper than Φ' and Γ_2 is negatively deeper than Φ' and (B) for a fixed value $\theta < \theta_0$ (or $> \theta_0$), the absolute value of the slope of Γ_1 (or Γ_2) is decreasing in the variable y (in the region below $y = \sqrt{\frac{c^p s^q}{\tau}}$).




Figure 8

Therefore apparently there exists some nonnegative smooth solution Φ_0 of (6.1) which is zero in some neighborhoods around the ending points of I and has value one at θ_0 .




Figure 9

Then, according to (4.1), a coflat calibration singular only at the origin can be gained for each homogeneous area-minimizing hypercone. Now the proof is complete.

Acknowledgements The author would like to express deep gratitude to Professor H. Blaine Lawson, Jr. for his guidance and constant encouragement. He also wishes to thank Professor Hui Ma and Doctor Chao Qian for helpful comments, the referees for nice suggestions, and the MSRI for its warm hospitality.

References

- [1] Bombieri, E., De Giorgi, E., Giusti, E.: Minimal cones and the Bernstein problem, *Invent. Math.*, **7**, 243–268 (1969)
- [2] Cheng, B. N.: Area-minimizing cone-type surfaces and coflat calibrations, *Indiana Univ. Math. J.*, **37**, 505–535, (1988)
- [3] Federer, H., Fleming, W. H.: Normal and integral currents, *Ann. Math.*, **72**, 458–520 (1960)
- [4] Hardt, R., Simon, L.: Area minimizing hypersurfaces with isolated singularities, *J. Reine. Angew. Math.*, **362**, 102–129 (1985)
- [5] Harvey, F. R., Lawson, Jr., H. B.: Calibrated geometries, *Acta Math.*, **148**, 47–157 (1982)
- [6] Hsiang, W.-Y., Lawson, Jr., H. B.: Minimal submanifolds of low cohomogeneity, *J. Diff. Geom.*, **5**, 1–38 (1971)
- [7] Lawlor, G. R.: A Sufficient Criterion for a Cone to be Area-Minimizing, Mem. of the Amer. Math. Soc., **91**, No. 446, 111 pp., American Mathematical Society, Providence, Rhode Island, 1991
- [8] Lawson, Jr., H. B.: The equivariant Plateau problem and interior regularity, *Trans. Amer. Math. Soc.*, **173**, 231–249 (1972)

- [9] Lin, F.-H.: Minimality and stability of minimal hypersurfaces in \mathbb{R}^N , *Bull. Austral. Math. Soc.*, **36**, 209–214 (1987)
- [10] Ma, H., Ohnita, Y.: Hamiltonian stability of the Gauss images of homogeneous isoparametric hypersurfaces. I, *J. Diff. Geom.*, **97**, 275–348 (2014)
- [11] Simoes, P.: On a class of minimal cones in \mathbb{R}^n , *Bull. Amer. Math. Soc.*, **3**, 488–489 (1974)
- [12] Simoes, P.: A class of minimal cones in \mathbb{R}^n , $n \geq 8$, that minimize area, Ph.D. thesis, University of California, Berkeley, Calif., 1973
- [13] Simon, L.: Entire solutions of the minimal surface equation, *J. Diff. Geom.*, **30**, 643–688 (1989)
- [14] Simons, J.: Minimal varieties in riemannian manifolds, *Ann. of Math.*, **88**, 62–105 (1968)
- [15] Takagi, R., Takahashi, T.: On the principal curvatures of homogeneous hypersurfaces in a sphere, pp. 469–481, *Diff. Geom.*, in honor of K. Yano, Kinokuniya, Tokyo, 1972
- [16] Tang, Z.Z., Xie, Y.Q., Yan, W.J.: Shoen-Yau-Gromov-Lawson theory and isoparametric foliations, *Comm. Anal. Geom.*, **20**, 989–1018 (2012)
- [17] Zhang, Y.S.: On extending calibration pairs, accepted, preliminary version available at arXiv:1511.03953