

A new way to iterate Brzeziński crossed products

Leonard Dăuş *

Department of Mathematical Sciences, UAE University
PO-Box 15551, Al Ain, United Arab Emirates
e-mail: leonard.daus@uaeu.ac.ae

Florin Panaite†

Institute of Mathematics of the Romanian Academy
PO-Box 1-764, RO-014700 Bucharest, Romania
e-mail: Florin.Panaite@imar.ro

Abstract

If $A \otimes_{R,\sigma} V$ and $A \otimes_{P,\nu} W$ are two Brzeziński crossed products and $Q : W \otimes V \rightarrow V \otimes W$ is a linear map satisfying certain properties, we construct a Brzeziński crossed product $A \otimes_{S,\theta} (V \otimes W)$. This construction contains as a particular case the iterated twisted tensor product of algebras.

Introduction

The *twisted tensor product* of the associative unital algebras A and B is a new associative unital algebra structure built on the linear space $A \otimes B$ with the help of a linear map $R : B \otimes A \rightarrow A \otimes B$ called a *twisting map*. This construction, denoted by $A \otimes_R B$, appeared in several contexts and has various applications ([4], [10]). Concrete examples come especially from Hopf algebra theory, like for instance the smash product.

It was proved in [7] that twisted tensor products of algebras may be iterated. Namely, if $A \otimes_{R_1} B$, $B \otimes_{R_2} C$ and $A \otimes_{R_3} C$ are twisted tensor products and the twisting maps R_1, R_2, R_3 satisfy the braid relation $(id_A \otimes R_2) \circ (R_3 \otimes id_B) \circ (id_C \otimes R_1) = (R_1 \otimes id_C) \circ (id_B \otimes R_3) \circ (R_2 \otimes id_A)$, then one can define certain twisted tensor products $A \otimes_{T_2} (B \otimes_{R_2} C)$ and $(A \otimes_{R_1} B) \otimes_{T_1} C$ that are equal as algebras (and this algebra is called the iterated twisted tensor product).

The Brzeziński crossed product, introduced in [1], is a common generalization of twisted tensor products of algebras and the Hopf crossed product (containing also as a particular case the quasi-Hopf smash product introduced in [3]). If A is an associative unital algebra, V is a linear space endowed with a distinguished element 1_V and $\sigma : V \otimes V \rightarrow A \otimes V$ and $R : V \otimes A \rightarrow A \otimes V$ are linear maps satisfying certain conditions, then the Brzeziński crossed product is a certain associative unital algebra structure on $A \otimes V$, denoted by $A \otimes_{R,\sigma} V$.

In [9] was proved that Brzeziński crossed products may be iterated, in the following sense. One can define first a "mirror version" of the Brzeziński crossed product, denoted by $W \overline{\otimes}_{P,\nu} D$

*Research supported by the National Research Foundation - United Arab Emirates and by the United Arab Emirates University(NRF-UAEU) under Grant No. 31S076/2013.

†Work supported by a grant of the Romanian National Authority for Scientific Research, CNCS-UEFISCDI, project number PN-II-ID-PCE-2011-3-0635, contract nr. 253/5.10.2011.

(where D is an associative unital algebra, W is a linear space and P, ν are certain linear maps). Examples are twisted tensor products of algebras and the quasi-Hopf smash product introduced in [2]. Then it was proved that, if $W \overline{\otimes}_{P, \nu} D$ and $D \otimes_{R, \sigma} V$ are two Brzeziński crossed products and $Q : V \otimes W \rightarrow W \otimes D \otimes V$ is a linear map satisfying some conditions, then one can define certain Brzeziński crossed products $(W \overline{\otimes}_{P, \nu} D) \otimes_{\overline{R}, \overline{\sigma}} V$ and $W \overline{\otimes}_{\overline{P}, \overline{\nu}} (D \otimes_{R, \sigma} V)$ that are equal as algebras. Iterated twisted tensor products of algebras appear as a particular case of this construction, as well as the so-called quasi-Hopf two-sided smash product $A \# H \# B$ from [3].

The aim of this paper is to show that Brzeziński crossed products may be iterated in a different way, that will also contain as a particular case the iterated twisted tensor product of algebras. Namely, we prove that, if $A \otimes_{R, \sigma} V$ and $A \otimes_{P, \nu} W$ are two Brzeziński crossed products and $Q : W \otimes V \rightarrow V \otimes W$ is a linear map satisfying certain properties, then we can define two Brzeziński crossed products $A \otimes_{S, \theta} (V \otimes W)$ and $(A \otimes_{R, \sigma} V) \otimes_{T, \eta} W$ that are equal as algebras.

Our inspiration for looking at this new way of iterating Brzeziński crossed products came from the following result in graded ring theory: If G is a group, R is a G -graded ring, A and B are two finite left G -sets, then there exists a ring isomorphism between the smash products $R \# (A \times B)$ and $(R \# A) \# B$. This result was obtained in [5, Corollary 3.2], and it is useful in the study of the von Neumann regularity of rings of the type $R \# A$, cf. [5]. The smash product $R \# A$ of the G -graded ring R by a (finite) left G -set A was introduced in the paper [8] and it is a particular case of a more general construction. If H is a Hopf algebra, R an H -comodule algebra and C an H -module coalgebra, then we may consider the category ${}^C_R \mathcal{M}(H)$ of Doi-Koppinen Hopf modules (i.e. left R -modules and left C -comodules which satisfy certain compatibility relations). Then, the smash product $R \# A$ used in [5] is a particular smash product and it is the first example in the category ${}^C_R \mathcal{M}(H)$ (in the case when H is the groupring $k[G]$, R a G -graded ring and C the grouplike coalgebra $k[A]$ on a G -set A).

1 Preliminaries

We work over a commutative field k . All algebras, linear spaces etc. will be over k ; unadorned \otimes means \otimes_k . By "algebra" we always mean an associative unital algebra. The multiplication of an algebra A is denoted by μ_A or simply μ when there is no danger of confusion, and we usually denote $\mu_A(a \otimes a') = aa'$ for all $a, a' \in A$. The unit of an algebra A is denoted by 1_A or simply 1 when there is no danger of confusion.

We recall from [4], [10] that, given two algebras A, B and a k -linear map $R : B \otimes A \rightarrow A \otimes B$, with Sweedler-type notation $R(b \otimes a) = a_R \otimes b_R$, for $a \in A, b \in B$, satisfying the conditions $a_R \otimes 1_R = a \otimes 1, 1_R \otimes b_R = 1 \otimes b, (aa')_R \otimes b_R = a_R a'_r \otimes (b_R)_r, a_R \otimes (bb')_R = (a_R)_r \otimes b_r b'_R$, for all $a, a' \in A$ and $b, b' \in B$ (where r and R are two different indices), if we define on $A \otimes B$ a new multiplication, by $(a \otimes b)(a' \otimes b') = aa'_R \otimes b_R b'$, then this multiplication is associative with unit $1 \otimes 1$. In this case, the map R is called a *twisting map* between A and B and the new algebra structure on $A \otimes B$ is denoted by $A \otimes_R B$ and called the *twisted tensor product* of A and B afforded by the map R .

We recall from [1] the construction of Brzeziński's crossed product:

Proposition 1.1 ([1]) *Let $(A, \mu, 1_A)$ be an (associative unital) algebra and V a vector space equipped with a distinguished element $1_V \in V$. Then the vector space $A \otimes V$ is an associative algebra with unit $1_A \otimes 1_V$ and whose multiplication has the property that $(a \otimes 1_V)(b \otimes v) = ab \otimes v$, for all $a, b \in A$ and $v \in V$, if and only if there exist linear maps $\sigma : V \otimes V \rightarrow A \otimes V$ and*

$R : V \otimes A \rightarrow A \otimes V$ satisfying the following conditions:

$$R(1_V \otimes a) = a \otimes 1_V, \quad R(v \otimes 1_A) = 1_A \otimes v, \quad \forall a \in A, v \in V, \quad (1.1)$$

$$\sigma(1_V \otimes v) = \sigma(v \otimes 1_V) = 1_A \otimes v, \quad \forall v \in V, \quad (1.2)$$

$$R \circ (id_V \otimes \mu) = (\mu \otimes id_V) \circ (id_A \otimes R) \circ (R \otimes id_A), \quad (1.3)$$

$$\begin{aligned} & (\mu \otimes id_V) \circ (id_A \otimes \sigma) \circ (R \otimes id_V) \circ (id_V \otimes \sigma) \\ &= (\mu \otimes id_V) \circ (id_A \otimes \sigma) \circ (\sigma \otimes id_V), \end{aligned} \quad (1.4)$$

$$\begin{aligned} & (\mu \otimes id_V) \circ (id_A \otimes \sigma) \circ (R \otimes id_V) \circ (id_V \otimes R) \\ &= (\mu \otimes id_V) \circ (id_A \otimes R) \circ (\sigma \otimes id_A). \end{aligned} \quad (1.5)$$

If this is the case, the multiplication of $A \otimes V$ is given explicitly by

$$\mu_{A \otimes V} = (\mu_2 \otimes id_V) \circ (id_A \otimes id_A \otimes \sigma) \circ (id_A \otimes R \otimes id_V),$$

where $\mu_2 = \mu \circ (id_A \otimes \mu) = \mu \circ (\mu \otimes id_A)$. We denote by $A \otimes_{R,\sigma} V$ this algebra structure and call it the crossed product (or Brzeziński crossed product) afforded by the data (A, V, R, σ) .

If $A \otimes_{R,\sigma} V$ is a crossed product, we introduce the following Sweedler-type notation:

$$\begin{aligned} R : V \otimes A &\rightarrow A \otimes V, & R(v \otimes a) &= a_R \otimes v_R, \\ \sigma : V \otimes V &\rightarrow A \otimes V, & \sigma(v \otimes v') &= \sigma_1(v, v') \otimes \sigma_2(v, v'), \end{aligned}$$

for all $v, v' \in V$ and $a \in A$. With this notation, the multiplication of $A \otimes_{R,\sigma} V$ reads

$$(a \otimes v)(a' \otimes v') = aa'_R \sigma_1(v_R, v') \otimes \sigma_2(v_R, v'), \quad \forall a, a' \in A, v, v' \in V.$$

A twisted tensor product is a particular case of a crossed product (cf. [6]), namely, if $A \otimes_R B$ is a twisted tensor product of algebras then $A \otimes_R B = A \otimes_{R,\sigma} B$, where $\sigma : B \otimes B \rightarrow A \otimes B$ is given by $\sigma(b \otimes b') = 1_A \otimes bb'$, for all $b, b' \in B$.

Remark 1.2 The conditions (1.3), (1.4) and (1.5) for R, σ may be written in Sweedler-type notation respectively as

$$(aa')_R \otimes v_R = a_R a'_r \otimes (v_R)_r, \quad (1.6)$$

$$\begin{aligned} & \sigma_1(y, z)_R \sigma_1(x_R, \sigma_2(y, z)) \otimes \sigma_2(x_R, \sigma_2(y, z)) \\ &= \sigma_1(x, y) \sigma_1(\sigma_2(x, y), z) \otimes \sigma_2(\sigma_2(x, y), z), \end{aligned} \quad (1.7)$$

$$(a_R)_r \sigma_1(v_r, v'_R) \otimes \sigma_2(v_r, v'_R) = \sigma_1(v, v') a_R \otimes \sigma_2(v, v')_R, \quad (1.8)$$

for all $a, a' \in A$ and $x, y, z, v, v' \in V$, where r is another copy of R .

2 The main result and examples

Theorem 2.1 Let $A \otimes_{R,\sigma} V$ and $A \otimes_{P,\nu} W$ be two crossed products and $Q : W \otimes V \rightarrow V \otimes W$ a linear map, with notation $Q(w \otimes v) = v_Q \otimes w_Q$, for all $v \in V$ and $w \in W$. Assume that the following conditions are satisfied:

(i) Q is unital, in the sense that

$$Q(1_W \otimes v) = v \otimes 1_W, \quad Q(w \otimes 1_V) = 1_V \otimes w, \quad \forall v \in V, w \in W. \quad (2.1)$$

(ii) the braid relation for R, P, Q , i.e.

$$(id_A \otimes Q) \circ (P \otimes id_V) \circ (id_W \otimes R) = (R \otimes id_W) \circ (id_V \otimes P) \circ (Q \otimes id_A), \quad (2.2)$$

or, equivalently,

$$(a_R)_P \otimes (v_R)_Q \otimes (w_P)_Q = (a_P)_R \otimes (v_Q)_R \otimes (w_Q)_P, \quad \forall a \in A, v \in V, w \in W. \quad (2.3)$$

(iii) we have the following hexagonal relation between σ, P, Q :

$$(id_A \otimes Q) \circ (P \otimes id_V) \circ (id_W \otimes \sigma) = (\sigma \otimes id_W) \circ (id_V \otimes Q) \circ (Q \otimes id_V), \quad (2.4)$$

or, equivalently,

$$\sigma_1(v, v')_P \otimes \sigma_2(v, v')_Q \otimes (w_P)_Q = \sigma_1(v_Q, v'_q) \otimes \sigma_2(v_Q, v'_q) \otimes (w_Q)_q, \quad (2.5)$$

for all $v, v' \in V$ and $w \in W$, where q is another copy of Q .

(iv) we have the following hexagonal relation between ν, R, Q :

$$(id_A \otimes Q) \circ (\nu \otimes id_V) = (R \otimes id_W) \circ (id_V \otimes \nu) \circ (Q \otimes id_W) \circ (id_W \otimes Q), \quad (2.6)$$

or, equivalently,

$$\nu_1(w, w') \otimes v_Q \otimes \nu_2(w, w')_Q = \nu_1(w_q, w'_q)_R \otimes ((v_Q)_q)_R \otimes \nu_2(w_q, w'_q), \quad (2.7)$$

for all $v \in V$ and $w, w' \in W$, where q is another copy of Q .

Define the linear maps

$$\begin{aligned} S &: (V \otimes W) \otimes A \rightarrow A \otimes (V \otimes W), \quad S := (R \otimes id_W) \circ (id_V \otimes P), \\ \theta &: (V \otimes W) \otimes (V \otimes W) \rightarrow A \otimes (V \otimes W), \\ \theta &:= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes R \otimes id_W) \circ (\sigma \otimes \nu) \circ (id_V \otimes Q \otimes id_W), \\ T &: W \otimes (A \otimes V) \rightarrow (A \otimes V) \otimes W, \quad T := (id_A \otimes Q) \circ (P \otimes id_V), \\ \eta &: W \otimes W \rightarrow (A \otimes V) \otimes W, \\ \eta &(w \otimes w') = (\nu_1(w, w') \otimes 1_V) \otimes \nu_2(w, w'), \quad \forall w, w' \in W. \end{aligned}$$

Then we have a crossed product $A \otimes_{S, \theta} (V \otimes W)$ (with respect to $1_{V \otimes W} := 1_V \otimes 1_W$), we have a crossed product $(A \otimes_{R, \sigma} V) \otimes_{T, \eta} W$ and we have an algebra isomorphism $A \otimes_{S, \theta} (V \otimes W) \simeq (A \otimes_{R, \sigma} V) \otimes_{T, \eta} W$ given by the trivial identification.

Proof. We prove first that $A \otimes_{S, \theta} (V \otimes W)$ is a crossed product, i.e. we need to prove the relations (1.1)-(1.5) with R replaced by S , σ replaced by θ etc. The relations (1.1) and (1.2) follow immediately by (2.1) and the relations (1.1) and (1.2) for R, σ and P, ν . Note that the maps S and θ are defined explicitly by

$$\begin{aligned} S(v \otimes w \otimes a) &= (a_P)_R \otimes v_R \otimes w_P, \\ \theta(v \otimes w \otimes v' \otimes w') &= \sigma_1(v, v'_Q) \nu_1(w_Q, w')_R \otimes \sigma_2(v, v'_Q)_R \otimes \nu_2(w_Q, w'), \end{aligned}$$

for all $v, v' \in V$, $w, w' \in W$ and $a \in A$. We will denote by $R = r = \mathcal{R} = \overline{R}$, $Q = q = \overline{Q}$ and $P = p$ some more copies of R, Q and P .

Proof of (1.3):

$$\begin{aligned}
& S \circ (id_V \otimes id_W \otimes \mu_A)(v \otimes w \otimes a \otimes a') \\
&= S(v \otimes w \otimes aa') \\
&= ((aa')_P)_R \otimes v_R \otimes w_P \\
&\stackrel{(1.6)}{=} (a_P a'_P)_R \otimes v_R \otimes (w_P)_P \\
&\stackrel{(1.6)}{=} (a_P)_R (a'_P)_R \otimes (v_R)_R \otimes (w_P)_P \\
&= (\mu_A \otimes id_V \otimes id_W)((a_P)_R \otimes (a'_P)_R \otimes (v_R)_R \otimes (w_P)_P) \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes S)((a_P)_R \otimes v_R \otimes w_P \otimes a') \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes S) \circ (S \otimes id_A)(v \otimes w \otimes a \otimes a'), \quad q.e.d.
\end{aligned}$$

Proof of (1.4):

$$\begin{aligned}
& (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) \circ (S \otimes id_V \otimes id_W) \circ (id_V \otimes id_W \otimes \theta)(v \otimes w \otimes v' \otimes w' \otimes v'' \otimes w'') \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) \circ (S \otimes id_V \otimes id_W)(v \otimes w \otimes \sigma_1(v', v''_Q) \nu_1(w'_Q, w'')_R \\
&\quad \otimes \sigma_2(v', v''_Q)_R \otimes \nu_2(w'_Q, w'')) \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta)(([\sigma_1(v', v''_Q) \nu_1(w'_Q, w'')_R]_P)_R \otimes v_r \otimes w_P \\
&\quad \otimes \sigma_2(v', v''_Q)_R \otimes \nu_2(w'_Q, w'')) \\
&= ([\sigma_1(v', v''_Q) \nu_1(w'_Q, w'')_R]_P)_R \sigma_1(v_r, (\sigma_2(v', v''_Q)_R)_q) \nu_1((w_P)_q, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes \sigma_2(v_r, (\sigma_2(v', v''_Q)_R)_q)_{\mathcal{R}} \otimes \nu_2((w_P)_q, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.6)}{=} (\sigma_1(v', v''_Q)_P (\nu_1(w'_Q, w'')_R)_P)_R \sigma_1(v_r, (\sigma_2(v', v''_Q)_R)_q) \nu_1(((w_P)_p)_q, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes \sigma_2(v_r, (\sigma_2(v', v''_Q)_R)_q)_{\mathcal{R}} \otimes \nu_2(((w_P)_p)_q, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.6)}{=} (\sigma_1(v', v''_Q)_P)_{\overline{R}} ((\nu_1(w'_Q, w'')_R)_P)_R \sigma_1((v_{\overline{R}})_r, (\sigma_2(v', v''_Q)_R)_q) \nu_1(((w_P)_p)_q, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes \sigma_2((v_{\overline{R}})_r, (\sigma_2(v', v''_Q)_R)_q)_{\mathcal{R}} \otimes \nu_2(((w_P)_p)_q, \nu_2(w'_Q, w'')) \\
&\stackrel{(2.3)}{=} (\sigma_1(v', v''_Q)_P)_{\overline{R}} ((\nu_1(w'_Q, w'')_P)_R)_R \sigma_1((v_{\overline{R}})_r, (\sigma_2(v', v''_Q)_R)_q) \nu_1(((w_P)_q)_p, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes \sigma_2((v_{\overline{R}})_r, (\sigma_2(v', v''_Q)_R)_q)_{\mathcal{R}} \otimes \nu_2(((w_P)_q)_p, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.8)}{=} (\sigma_1(v', v''_Q)_P)_{\overline{R}} \sigma_1(v_{\overline{R}}, \sigma_2(v', v''_Q)_q) (\nu_1(w'_Q, w'')_P)_R \nu_1(((w_P)_q)_p, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes (\sigma_2(v_{\overline{R}}, \sigma_2(v', v''_Q)_q)_R)_{\mathcal{R}} \otimes \nu_2(((w_P)_q)_p, \nu_2(w'_Q, w'')) \\
&\stackrel{(2.5)}{=} \sigma_1(v'_{\overline{Q}}, (v''_Q)_q)_{\overline{R}} \sigma_1(v_{\overline{R}}, \sigma_2(v'_{\overline{Q}}, (v''_Q)_q)) (\nu_1(w'_Q, w'')_P)_R \nu_1(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes (\sigma_2(v_{\overline{R}}, \sigma_2(v'_{\overline{Q}}, (v''_Q)_q))_R)_{\mathcal{R}} \otimes \nu_2(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.7)}{=} \sigma_1(v, v'_{\overline{Q}}) \sigma_1(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q) (\nu_1(w'_Q, w'')_P)_R \nu_1(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w''))_{\mathcal{R}} \\
&\quad \otimes (\sigma_2(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q)_R)_{\mathcal{R}} \otimes \nu_2(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.6)}{=} \sigma_1(v, v'_{\overline{Q}}) \sigma_1(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q) [\nu_1(w'_Q, w'')_P \nu_1(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w''))]_R \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q)_R \otimes \nu_2(((w_{\overline{Q}})_q)_p, \nu_2(w'_Q, w'')) \\
&\stackrel{(1.7)}{=} \sigma_1(v, v'_{\overline{Q}}) \sigma_1(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q) [\nu_1((w_{\overline{Q}})_q, w'_Q) \nu_1(\nu_2((w_{\overline{Q}})_q, w'_Q), w'')]_R \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_{\overline{Q}}), (v''_Q)_q)_R \otimes \nu_2(\nu_2((w_{\overline{Q}})_q, w'_Q), w'')
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(1.8)}{=} \sigma_1(v, v'_Q) \{ [\nu_1((w_Q)_q, w'_Q) \nu_1(\nu_2((w_Q)_q, w'_Q), w'')]]_R \}_r \sigma_1(\sigma_2(v, v'_Q)_r, ((v''_Q)_q)_R) \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_Q)_r, ((v''_Q)_q)_R) \otimes \nu_2(\nu_2((w_Q)_q, w'_Q), w'') \\
&\stackrel{(1.6)}{=} \sigma_1(v, v'_Q) [\nu_1((w_Q)_q, w'_Q)_R \nu_1(\nu_2((w_Q)_q, w'_Q), w'')]_{\mathcal{R}}]_r \sigma_1(\sigma_2(v, v'_Q)_r, (((v''_Q)_q)_R)_{\mathcal{R}}) \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_Q)_r, (((v''_Q)_q)_R)_{\mathcal{R}}) \otimes \nu_2(\nu_2((w_Q)_q, w'_Q), w'') \\
&\stackrel{(2.7)}{=} \sigma_1(v, v'_Q) [\nu_1(w_Q, w') \nu_1(\nu_2(w_Q, w')_Q, w'')]_{\mathcal{R}}]_r \sigma_1(\sigma_2(v, v'_Q)_r, (v''_Q)_{\mathcal{R}}) \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_Q)_r, (v''_Q)_{\mathcal{R}}) \otimes \nu_2(\nu_2(w_Q, w')_Q, w'') \\
&\stackrel{(1.6)}{=} \sigma_1(v, v'_Q) \nu_1(w_Q, w')_R (\nu_1(\nu_2(w_Q, w')_Q, w'')]_{\mathcal{R}})_r \sigma_1((\sigma_2(v, v'_Q)_R)_r, (v''_Q)_{\mathcal{R}}) \\
&\quad \otimes \sigma_2((\sigma_2(v, v'_Q)_R)_r, (v''_Q)_{\mathcal{R}}) \otimes \nu_2(\nu_2(w_Q, w')_Q, w'') \\
&\stackrel{(1.8)}{=} \sigma_1(v, v'_Q) \nu_1(w_Q, w')_R \sigma_1(\sigma_2(v, v'_Q)_R, v''_Q) \nu_1(\nu_2(w_Q, w')_Q, w'')]_r \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_Q)_R, v''_Q)_r \otimes \nu_2(\nu_2(w_Q, w')_Q, w'') \\
&= (\mu_A \otimes id_V \otimes id_W) (\sigma_1(v, v'_Q) \nu_1(w_Q, w')_R \otimes \sigma_1(\sigma_2(v, v'_Q)_R, v''_Q) \nu_1(\nu_2(w_Q, w')_Q, w'')]_r \\
&\quad \otimes \sigma_2(\sigma_2(v, v'_Q)_R, v''_Q)_r \otimes \nu_2(\nu_2(w_Q, w')_Q, w'') \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) (\sigma_1(v, v'_Q) \nu_1(w_Q, w')_R \\
&\quad \otimes \sigma_2(v, v'_Q)_R \otimes \nu_2(w_Q, w') \otimes v'' \otimes w'') \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) \circ (\theta \otimes id_V \otimes id_W) (v \otimes w \otimes v' \otimes w' \otimes v'' \otimes w''), \quad q.e.d.
\end{aligned}$$

Proof of (1.5):

$$\begin{aligned}
&(\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) \circ (S \otimes id_V \otimes id_W) \circ (id_V \otimes id_W \otimes S) (v \otimes w \otimes v' \otimes w' \otimes a) \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) \circ (S \otimes id_V \otimes id_W) (v \otimes w \otimes (a_P)_R \otimes v'_R \otimes w'_P) \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes \theta) (((a_P)_R)_p)_r \otimes v_r \otimes w_p \otimes v'_R \otimes w'_P) \\
&= (((a_P)_R)_p)_r \sigma_1(v_r, (v'_R)_Q) \nu_1((w_p)_Q, w'_P)_{\mathcal{R}} \otimes \sigma_2(v_r, (v'_R)_Q)_{\mathcal{R}} \otimes \nu_2((w_p)_Q, w'_P) \\
&\stackrel{(2.3)}{=} (((a_P)_p)_R)_r \sigma_1(v_r, (v'_Q)_R) \nu_1((w_Q)_p, w'_P)_{\mathcal{R}} \otimes \sigma_2(v_r, (v'_Q)_R)_{\mathcal{R}} \otimes \nu_2((w_Q)_p, w'_P) \\
&\stackrel{(1.8)}{=} \sigma_1(v, v'_Q) ((a_P)_p)_R \nu_1((w_Q)_p, w'_P)_{\mathcal{R}} \otimes (\sigma_2(v, v'_Q)_R)_{\mathcal{R}} \otimes \nu_2((w_Q)_p, w'_P) \\
&\stackrel{(1.6)}{=} \sigma_1(v, v'_Q) [(a_P)_p \nu_1((w_Q)_p, w'_P)]_R \otimes \sigma_2(v, v'_Q)_R \otimes \nu_2((w_Q)_p, w'_P) \\
&\stackrel{(1.8)}{=} \sigma_1(v, v'_Q) [\nu_1(w_Q, w') a_P]_R \otimes \sigma_2(v, v'_Q)_R \otimes \nu_2(w_Q, w')_P \\
&\stackrel{(1.6)}{=} \sigma_1(v, v'_Q) \nu_1(w_Q, w')_R (a_P)_r \otimes (\sigma_2(v, v'_Q)_R)_r \otimes \nu_2(w_Q, w')_P \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes S) (\sigma_1(v, v'_Q) \nu_1(w_Q, w')_R \otimes \sigma_2(v, v'_Q)_R \otimes \nu_2(w_Q, w') \otimes a) \\
&= (\mu_A \otimes id_V \otimes id_W) \circ (id_A \otimes S) \circ (\theta \otimes id_A) (v \otimes w \otimes v' \otimes w' \otimes a),
\end{aligned}$$

so $A \otimes_{S, \theta} (V \otimes W)$ is indeed a crossed product. With a similar computation one can prove that $(A \otimes_{R, \sigma} V) \otimes_{T, \eta} W$ is a crossed product, so the only thing left to prove is that the multiplications of $A \otimes_{S, \theta} (V \otimes W)$ and $(A \otimes_{R, \sigma} V) \otimes_{T, \eta} W$ coincide. A straightforward computation shows that the multiplication of $A \otimes_{S, \theta} (V \otimes W)$ is given by the formula

$$(a \otimes v \otimes w) (a' \otimes v' \otimes w') = a (a'_P)_{\mathcal{R}} \sigma_1(v_{\mathcal{R}}, v'_Q) \nu_1((w_P)_Q, w')_r \otimes \sigma_2(v_{\mathcal{R}}, v'_Q)_r \otimes \nu_2((w_P)_Q, w').$$

We compute now the multiplication of $(A \otimes_{R,\sigma} V) \otimes_{T,\eta} W$:

$$\begin{aligned}
& (a \otimes v \otimes w)(a' \otimes v' \otimes w') \\
&= (a \otimes v)(a' \otimes v')_T \eta_1(w_T, w') \otimes \eta_2(w_T, w') \\
&= (a \otimes v)(a'_P \otimes v'_Q) \eta_1((w_P)_Q, w') \otimes \eta_2((w_P)_Q, w') \\
&= (a \otimes v)(a'_P \otimes v'_Q)(\nu_1((w_P)_Q, w') \otimes 1_V) \otimes \nu_2((w_P)_Q, w') \\
&= (a \otimes v)(a'_P \nu_1((w_P)_Q, w')_R \otimes (v'_Q)_R) \otimes \nu_2((w_P)_Q, w') \\
&= a[a'_P \nu_1((w_P)_Q, w')_R]_r \sigma_1(v_r, (v'_Q)_R) \otimes \sigma_2(v_r, (v'_Q)_R) \otimes \nu_2((w_P)_Q, w') \\
&\stackrel{(1.6)}{=} a(a'_P)_R (\nu_1((w_P)_Q, w')_R)_r \sigma_1((v_R)_r, (v'_Q)_R) \otimes \sigma_2((v_R)_r, (v'_Q)_R) \otimes \nu_2((w_P)_Q, w') \\
&\stackrel{(1.8)}{=} a(a'_P)_R \sigma_1(v_R, v'_Q) \nu_1((w_P)_Q, w')_r \otimes \sigma_2(v_R, v'_Q)_r \otimes \nu_2((w_P)_Q, w'),
\end{aligned}$$

and we can see that the two multiplications coincide. \square

Example 2.2 We recall from [7] what was called there an iterated twisted tensor product of algebras. Let A, B, C be associative unital algebras, $R_1 : B \otimes A \rightarrow A \otimes B$, $R_2 : C \otimes B \rightarrow B \otimes C$, $R_3 : C \otimes A \rightarrow A \otimes C$ twisting maps satisfying the braid equation

$$(id_A \otimes R_2) \circ (R_3 \otimes id_B) \circ (id_C \otimes R_1) = (R_1 \otimes id_C) \circ (id_B \otimes R_3) \circ (R_2 \otimes id_A).$$

Then we have an algebra structure on $A \otimes B \otimes C$ (called the iterated twisted tensor product) with unit $1_A \otimes 1_B \otimes 1_C$ and multiplication

$$(a \otimes b \otimes c)(a' \otimes b' \otimes c') = a(a'_{R_3})_{R_1} \otimes b_{R_1} b'_{R_2} \otimes (c_{R_3})_{R_2} c'.$$

We define $V = B$, $W = C$, $R = R_1$, $P = R_3$, $Q = R_2$ and the linear maps

$$\begin{aligned}
\sigma : V \otimes V &\rightarrow A \otimes V, & \sigma(b \otimes b') &= 1_A \otimes bb', \quad \forall b, b' \in V, \\
\nu : W \otimes W &\rightarrow A \otimes W, & \nu(c \otimes c') &= 1_A \otimes cc', \quad \forall c, c' \in W.
\end{aligned}$$

Then, for the crossed products $A \otimes_{R,\sigma} V = A \otimes_{R_1} B$, $A \otimes_{P,\nu} W = A \otimes_{R_3} C$ and the map Q , one can check that the hypotheses of Theorem 2.1 are satisfied and the crossed products $A \otimes_{S,\theta} (V \otimes W) \equiv (A \otimes_{R,\sigma} V) \otimes_{T,\eta} W$ (notation as in Theorem 2.1) coincide with the iterated twisted tensor product.

Example 2.3 Let $A \otimes_{R,\sigma} V$ be a crossed product and W an (associative unital) algebra. Define the linear maps

$$\begin{aligned}
P : W \otimes A &\rightarrow A \otimes W, & P(w \otimes a) &= a \otimes w, \quad \forall a \in A, w \in W, \\
\nu : W \otimes W &\rightarrow A \otimes W, & \nu(w \otimes w') &= 1_A \otimes ww', \quad \forall w, w' \in W,
\end{aligned}$$

so we have the crossed product $A \otimes_{P,\nu} W$ which is just the ordinary tensor product of algebras $A \otimes W$. Define the linear map $Q : W \otimes V \rightarrow V \otimes W$, $Q(w \otimes v) = v \otimes w$, for all $v \in V$, $w \in W$. Then one can easily check that the hypotheses of Theorem 2.1 are satisfied.

References

- [1] T. Brzeziński, Crossed products by a coalgebra, *Comm. Algebra* **25** (1997), 3551–3575.
- [2] D. Bulacu, F. Panaite, F. Van Oystaeyen, Quasi-Hopf algebra actions and smash products, *Comm. Algebra* **28** (2000), 631–651.
- [3] D. Bulacu, F. Panaite, F. Van Oystaeyen, Generalized diagonal crossed products and smash products for quasi-Hopf algebras. Applications, *Comm. Math. Phys.* **266** (2006), 355–399.
- [4] A. Cap, H. Schichl, J. Vanzura, On twisted tensor products of algebras, *Comm. Algebra* **23** (1995), 4701–4735.
- [5] L. Dăuş, C. Năstăsescu, M. Năstăsescu, Von Neumann regularity of smash products associated with G -set gradings, *J. Algebra* **331** (2011), 46–57.
- [6] C. Di Luigi, J. A. Guccione, J. J. Guccione, Brzeziński’s crossed products and braided Hopf crossed products, *Comm. Algebra* **32** (2004), 3563–3580.
- [7] P. Jara Martínez, J. López Peña, F. Panaite, F. Van Oystaeyen, On iterated twisted tensor products of algebras, *Internat. J. Math.* **19** (2008), 1053–1101.
- [8] C. Năstăsescu, Ş. Raianu, F. Van Oystaeyen, Modules graded by G -sets, *Math. Z.* **203** (1990), 605–627.
- [9] F. Panaite, Iterated crossed products, *J. Algebra Appl.* **13** (2014), 1450036 (14 pages).
- [10] A. Van Daele, S. Van Keer, The Yang–Baxter and Pentagon equation, *Compositio Math.* **91** (1994), 201–221.