# EXTENSION OF EUCLIDEAN OPERATOR RADIUS INEQUALITIES

MOHAMMAD SAL MOSLEHIAN<sup>1</sup>, MOSTAFA SATTARI<sup>1</sup> AND KHALID SHEBRAWI<sup>2</sup>

ABSTRACT. To extend the Euclidean operator radius, we define  $w_p$  for an *n*-tuples of operators  $(T_1, \ldots, T_n)$  in  $\mathbb{B}(\mathscr{H})$  by  $w_p(T_1, \ldots, T_n) := \sup_{\|x\|=1} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^p\right)^{\frac{1}{p}}$ for  $p \geq 1$ . We generalize some inequalities including Euclidean operator radius of two operators to those involving  $w_p$ . Further we obtain some lower and upper bounds for  $w_p$ . Our main result states that if f and g are nonnegative continuous functions on  $[0, \infty)$  satisfying f(t) g(t) = t for all  $t \in [0, \infty)$ , then

$$w_p^{rp}\left(A_1^*T_1B_1,\ldots,A_n^*T_nB_n\right) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( \left[ B_i^*f^2\left(|T_i|\right)B_i \right]^{rp} + \left[ A_i^*g^2\left(|T_i^*|\right)A_i \right]^{rp} \right) \right\|$$

for all  $p \ge 1$ ,  $r \ge 1$  and operators in  $\mathbb{B}(\mathscr{H})$ .

#### 1. INTRODUCTION

Let  $\mathbb{B}(\mathscr{H})$  be the  $C^*$ -algebra of all bounded linear operators on a Hilbert space  $(\mathscr{H}, \langle \cdot, \cdot \rangle)$ . The numerical radius of  $A \in \mathbb{B}(\mathscr{H})$  is defined by

$$w(A) = \sup\{|\langle Ax, x \rangle| : x \in \mathcal{H}, ||x|| = 1\}.$$

It is well known that  $w(\cdot)$  defines a norm on  $\mathbb{B}(\mathscr{H})$ , which is equivalent to the usual operator norm  $\|\cdot\|$ . Namely, we have

$$\frac{1}{2} \|A\| \le w(A) \le \|A\|.$$

for each  $A \in \mathbb{B}(\mathscr{H})$ . It is known that if  $A \in \mathbb{B}(\mathscr{H})$  is self-adjoint, then w(A) = ||A||. An important inequality for w(A) is the power inequality stating that  $w(A^n) \leq w^n(A)$  for  $n = 1, 2, \ldots$ . There are many inequalities involving numerical radius; see [2, 3, 4, 10, 11, 12] and references therein.

The Euclidean operator radius of an *n*-tuple  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)} := \mathbb{B}(\mathscr{H}) \times$ 

<sup>2010</sup> Mathematics Subject Classification. 47A12, 47B15, 47A30, 47A63.

*Key words and phrases.* Euclidean operator radius, numerical radius, Cartesian decomposition, self-adjoint operator.

 $\ldots \times \mathbb{B}(\mathscr{H})$  was defined in [9] by

$$w_e(T_1,\ldots,T_n) := \sup_{\|x\|=1} \left( \sum_{i=1}^n |\langle T_i x, x \rangle|^2 \right)^{\frac{1}{2}}.$$

The particular cases n = 1 and n = 2 are numerical radius and Euclidean operator radius. Some interesting properties of this radius were obtained in [9]. For example, it is established that

$$\frac{1}{2\sqrt{n}} \left\| \sum_{i=1}^{n} T_i T_i^* \right\|^{\frac{1}{2}} \le w_e(T_1, \cdots, T_n) \le \left\| \sum_{i=1}^{n} T_i T_i^* \right\|^{\frac{1}{2}}.$$
(1.1)

We also observe that if A = B + iC is the Cartesian decomposition of A, then

$$w_e^2(B,C) = \sup_{\|x\|=1} \{ |\langle Bx, x \rangle|^2 + |\langle Cx, x \rangle|^2 \} = \sup_{\|x\|=1} |\langle Ax, x \rangle|^2 = w^2(A).$$

By the above inequality and  $A^*A + AA^* = 2(B^2 + C^2)$ , we have

$$\frac{1}{16} \|A^*A + AA^*\| \le w^2(A) \le \frac{1}{2} \|A^*A + AA^*\|.$$

We define  $w_p$  for *n*-tuples of operators  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$  for  $p \ge 1$  by

$$w_p(T_1,\ldots,T_n) := \sup_{\|x\|=1} \left( \sum_{i=1}^n |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}}$$

It follows from Minkowski's inequality for two vectors  $a = (a_1, a_2)$  and  $b = (b_1, b_2)$ , namely,

$$(|a_1+b_1|^p+|a_2+b_2|^p)^{\frac{1}{p}} \le (|a_1|^p+|a_2|^p)^{\frac{1}{p}}+(|b_1|^p+|b_2|^p)^{\frac{1}{p}}$$
 for  $p>1$ 

that  $w_p$  is a norm.

Moreover  $w_p, p \ge 1$ , for *n*-tuple of operators  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$  satisfies the following properties:

(i) 
$$w_p(T_1, \ldots, T_n) = 0 \Leftrightarrow T_1 = \ldots = T_n = 0.$$
  
(ii)  $w_p(\lambda T_1, \ldots, \lambda T_n) = |\lambda| w_p(T_1, \ldots, T_n)$  for all  $\lambda \in \mathbb{C}$ .  
(iii)  $w_p(T_1 + T_1, \ldots, T_n + T_n) \leq w_p(T_1, \ldots, T_n) + w_p(T_1, \ldots, T_n)$  for  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$ .

(iv)  $w_p(X^*T_1X,\ldots,X^*T_nX) \le ||X||^2 w_p(T_1,\ldots,T_n)$  for  $X \in \mathbb{B}(\mathscr{H})$ .

Dragomir [1] obtained some inequalities for the Euclidean operator radius  $w_e(B, C) = \sup_{\|x\|=1} (|\langle Bx, x \rangle|^2 + |\langle Cx, x \rangle|^2)^{\frac{1}{2}}$  of two bounded linear operators in a Hilbert space. In section 2 of this paper we extend some his results including inequalities for the Euclidean operator radius of linear operators to  $w_p$  ( $p \ge 1$ ). In addition, we apply some known inequalities for getting new inequalities for  $w_p$  in two operators. In section 3 we prove inequalities for  $w_p$  for *n*-tuples of operators. Some of our result in this section, generalize some inequalities in section 2. Further, we find some lower and upper bounds for  $w_p$ .

### 2. Inequalities for $w_p$ for two operators

To prove our generalized numerical radius inequalities, we need several known lemmas. The first lemma is a simple result of the classical Jensen inequality and a generalized mixed Cauchy–Schwarz inequality [7, 8, 6].

**Lemma 2.1.** For  $a, b \ge 0$ ,  $0 \le \alpha \le 1$  and  $r \ne 0$ ,

- (a)  $a^{\alpha}b^{1-\alpha} \leq \alpha a + (1-\alpha)b \leq [\alpha a^r + (1-\alpha)b^r]^{\frac{1}{r}}$  for  $r \geq 1$ ,
- (b) If  $A \in \mathbb{B}(\mathscr{H})$ , then  $|\langle Ax, y \rangle|^2 \leq \langle |A|^{2\alpha}x, x \rangle \langle |A^*|^{2(1-\alpha)}y, y \rangle$  for all  $x, y \in \mathscr{H}$ , where  $|A| = (A^*A)^{\frac{1}{2}}$ .
- (c) Let  $A \in \mathbb{B}(\mathscr{H})$ , and f and g be nonnegative continuous functions on  $[0, \infty)$ satisfying f(t) g(t) = t for all  $t \in [0, \infty)$ . Then

$$|\langle Ax, y \rangle| \le ||f(|A|) x|| ||g(|A^*|) y||$$

for all  $x, y \in \mathcal{H}$ .

**Lemma 2.2** (McCarthy inequality [5]). Let  $A \in \mathbb{B}(\mathcal{H})$ ,  $A \ge 0$  and let  $x \in \mathcal{H}$  be any unit vector. Then

(a)  $\langle Ax, x \rangle^r \leq \langle A^r x, x \rangle$  for  $r \geq 1$ , (b)  $\langle A^r x, x \rangle \leq \langle Ax, x \rangle^r$  for  $0 < r \leq 1$ .

Inequalities of the following lemma were obtained for the first time by Clarkson[7].

**Lemma 2.3.** Let X be a normed space and  $x, y \in X$ . Then for all  $p \ge 2$  with  $\frac{1}{p} + \frac{1}{q} = 1$ ,

- (a)  $2(||x||^p + ||y||^p)^{q-1} \le ||x+y||^q + ||x-y||^q$ ,
- (b)  $2(||x||^p + ||y||^p) \le ||x+y||^p + ||x-y||^p \le 2^{p-1}(||x||^p + ||y||^p),$
- (c)  $||x + y||^p + ||x y||^p \le 2(||x||^q + ||y||^q)^{p-1}.$

If 1 the converse inequalities hold.

Making the transformations  $x \to \frac{x+y}{2}$  and  $y \to \frac{x-y}{2}$  we observe that inequalities (a) and (c) in Lemma 2.3 are equivalent and so are the first and the second inequalities of (b). First of all we obtain a relation between  $w_p$  and  $w_e$  for  $p \ge 1$ .

**Proposition 2.4.** Let  $B, C \in \mathbb{B}(\mathcal{H})$ . Then

$$w_p(B,C) \le w_q(B,C) \le 2^{\frac{1}{q} - \frac{1}{p}} w_p(B,C)$$

for  $p \ge q \ge 1$ . In particular

$$w_p(B,C) \le w_e(B,C) \le 2^{\frac{1}{2} - \frac{1}{p}} w_p(B,C)$$
 (2.1)

for  $p \geq 2$ , and

$$2^{\frac{1}{2} - \frac{1}{p}} w_p(B, C) \le w_e(B, C) \le w_p(B, C)$$

for  $1 \leq p \leq 2$ .

*Proof.* An application of Jensen's inequality says that for a, b > 0 and  $p \ge q > 0$ , we have

$$(a^p + b^p)^{\frac{1}{p}} \le (a^q + b^q)^{\frac{1}{q}}$$

Let  $x \in \mathscr{H}$  be a unit vector. Choosing  $a = |\langle Bx, x \rangle|$  and  $b = |\langle Cx, x \rangle|$ , we have

$$\left(\left|\langle Bx,x\rangle\right|^{p}+\left|\langle Cx,x\rangle\right|^{p}\right)^{\frac{1}{p}} \leq \left(\left|\langle Bx,x\rangle\right|^{q}+\left|\langle Cx,x\rangle\right|^{q}\right)^{\frac{1}{q}}.$$

Now the first inequality follows by taking the supremum over all unit vectors in  $\mathscr{H}$ . A simple consequence of the classical Jensen's inequality concerning the convexity or the concavity of certain power functions says that for  $a, b \ge 0, 0 \le \alpha \le 1$  and  $p \ge q$ , we have

$$(\alpha a^{q} + (1 - \alpha) b^{q})^{\frac{1}{q}} \le (\alpha a^{p} + (1 - \alpha) b^{p})^{\frac{1}{p}}.$$

For  $\alpha = \frac{1}{2}$ , we get

$$(a^{q} + b^{q})^{\frac{1}{q}} \le 2^{\frac{1}{q} - \frac{1}{p}} (a^{p} + b^{p})^{\frac{1}{p}}$$

Again let  $x \in \mathscr{H}$  be a unit vector. Choosing  $a = |\langle Bx, x \rangle|$  and  $b = |\langle Cx, x \rangle|$  we get

$$\left(\left|\langle Bx,x\rangle\right|^{q}+\left|\langle Cx,x\rangle\right|^{q}\right)^{\frac{1}{q}} \leq 2^{\frac{1}{q}-\frac{1}{p}} \left(\left|\langle Bx,x\rangle\right|^{p}+\left|\langle Cx,x\rangle\right|^{p}\right)^{\frac{1}{p}}.$$

Now the second inequality follows by taking the supremum over all unit vectors in  $\mathscr{H}$ .

On making use of inequality (2.1) we find a lower bound for  $w_p$   $(p \ge 2)$ .

**Corollary 2.5.** If  $B, C \in \mathbb{B}(\mathcal{H})$ , then for  $p \geq 2$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-2} \|B^*B + C^*C\|^{\frac{1}{2}}.$$

*Proof.* According to inequalities (1.1) and (2.1) we can write

$$w_e(B,C) \ge \frac{1}{2\sqrt{2}} \|B^*B + C^*C\|^{\frac{1}{2}}$$

and

$$w_p(B,C) \ge 2^{\frac{1}{p}-\frac{1}{2}}w_e(B,C),$$

respectively. We therefore get desired inequality.

The next result is concerned with some lower bounds for  $w_p$ . This consequence has several inequalities as special cases. Our result will be generalized to *n*-tuples of operators in the next section.

**Proposition 2.6.** Let  $B, C \in \mathbb{B}(\mathcal{H})$ . Then for  $p \geq 1$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} \max\left(w(B+C), w(B-C)\right).$$
 (2.2)

This inequality is sharp.

*Proof.* We use convexity of function  $f(t) = t^p$   $(p \ge 1)$  as follows:

$$(|\langle Bx, x \rangle|^{p} + |\langle Cx, x \rangle|^{p})^{\frac{1}{p}} \ge 2^{\frac{1}{p}-1}(|\langle Bx, x \rangle| + |\langle Cx, x \rangle|)$$
$$\ge 2^{\frac{1}{p}-1}|\langle Bx, x \rangle \pm \langle Cx, x \rangle|$$
$$= 2^{\frac{1}{p}-1}|\langle (B \pm C)x, x \rangle|.$$

Taking supremum over  $x \in \mathscr{H}$  with ||x|| = 1 yields that

$$w_p(B,C) \ge 2^{\frac{1}{p}-1}w(B\pm C).$$

For sharpness one can obtain the same quantity  $2^{\frac{1}{p}}w(B)$  on both sides of the inequality by putting B = C.

**Corollary 2.7.** If A = B + iC is the Cartesian decomposition of A, then for all  $p \ge 2$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} \max\left( \|B+C\|, \|B-C\| \right),$$

and

$$w(A) \ge 2^{\frac{1}{p}-2} \max\left( \|(1-i)A + (1+i)A^*\|, \|(1+i)A + (1-i)A^*\| \right)$$

*Proof.* Obviously by inequality (2.2) we have the first inequality. For the second we use inequality (2.1).

**Corollary 2.8.** If  $B, C \in \mathbb{B}(\mathcal{H})$ , then for  $p \geq 1$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} \max\{w(B), w(C)\}.$$
 (2.3)

In addition, if A = B + iC is the Cartesian decomposition of A, then for  $p \ge 2$ 

$$w(A) \ge 2^{\frac{1}{p}-2} \max\left( \|A + A^*\|, \|A - A^*\| \right).$$

*Proof.* By inequality (2.2) and properties of the numerical radius, we have

$$2w_p(B,C) \ge 2^{\frac{1}{p}-1}(w(B+C)+w(B-C)) \ge 2^{\frac{1}{p}-1}w(B+C+B-C).$$

 $\operatorname{So}$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-1}w(B)$$

By symmetry we conclude that

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} \max(w(B), w(C)).$$

While the second inequality follows easily from inequality (2.1).

Now we apply part (b) of Lemma 2.3 to find some lower and upper bounds for  $w_p \ (p > 1)$ .

Proposition 2.9. Let 
$$B, C \in \mathbb{B}(\mathscr{H})$$
. Then for all  $p \ge 2$ ,  
(i)  $2^{\frac{1}{p}-1}w_p(B+C, B-C) \le w_p(B, C) \le 2^{-\frac{1}{p}}w_p(B+C, B-C)$ ;  
(ii)  $2^{\frac{1}{p}-1}(w^p(B+C) + w^p(B-C))^{\frac{1}{p}} \le w_p(B, C) \le 2^{-\frac{1}{p}}(w^p(B+C) + w^p(B-C))^{\frac{1}{p}}$ .  
If  $1 these inequalities hold in the opposite direction.$ 

*Proof.* Let  $x \in \mathscr{H}$  be a unit vector. Part (b) of Lemma 2.3 implies that for any  $p \geq 2$ 

$$2^{1-p}(|a+b|^p + |a-b|^p) \le |a|^p + |b|^p \le \frac{1}{2}(|a+b|^p + |a-b|^p).$$

Replacing  $a = |\langle Bx, x \rangle|$  and  $b = |\langle Cx, x \rangle|$  in above inequalities we obtain the desired inequalities.

**Remark 2.10.** In inequality (2.3), if we take B + C and B - C instead of B and C, then for  $p \ge 1$ 

$$w_p(B+C, B-C) \ge 2^{\frac{1}{p}-1} \max\{w(B+C), w(B-C)\}$$

By employing the first inequality of part (i) of Proposition 2.9, we get

$$w_p(B,C) \ge 2^{\frac{2}{p}-2} \max\{w(B+C), w(B-C)\}$$

for  $p \geq 1$ .

Taking B + C and B - C instead of B and C in the second inequality of part (ii) of Proposition 2.9, we reach

$$w_p(B+C, B-C) \le 2^{1-\frac{1}{p}} \left( w^p(B) + w^p(C) \right)^{\frac{1}{p}}$$

for all  $p \ge 1$ .

Now by applying the second inequality of part (i) of Proposition 2.9, we infer for  $p \ge 1$  that

$$w_p(B,C) \le 2^{1-\frac{2}{p}} \left( w^p(B) + w^p(C) \right)^{\frac{1}{p}}.$$

 $\operatorname{So}$ 

$$2^{\frac{2}{p}-2}\max\{w(B+C), w(B-C)\} \le w_p(B,C) \le 2^{1-\frac{2}{p}} (w^p(B) + w^p(C))^{\frac{1}{p}}$$

Moreover if B and C are self-adjoint, then

$$2^{\frac{2}{p}-2}\max\{\|B+C\|, \|B-C\|\} \le w_p(B,C) \le 2^{1-\frac{2}{p}} \left(\|B\|^p + \|C\|^p\right)^{\frac{1}{p}}$$

for all  $p \ge 1$ .

In the following result we find another lower bound for  $w_p \ (p \ge 1)$ .

**Theorem 2.11.** Let  $B, C \in \mathbb{B}(\mathcal{H})$ . Then for  $p \geq 1$ 

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} w^{\frac{1}{2}} (B^2 + C^2).$$

*Proof.* It follows from (2.2) that

$$2^{\frac{2}{p}-2}w^2(B\pm C) \le w_p^2(B,C) \,.$$

Hence

$$2w_p^2(B,C) \ge 2^{\frac{2}{p}-2} \left[ w^2(B+C) + w^2(B-C) \right]$$
  
$$\ge 2^{\frac{2}{p}-2} \left[ w \left( (B+C)^2 \right) + w \left( (B-C)^2 \right) \right]$$
  
$$\ge 2^{\frac{2}{p}-2} \left[ w \left( (B+C)^2 + (B-C)^2 \right) \right] = 2^{\frac{2}{p}-1} w (B^2 + C^2) \,.$$

It follows that

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} w^{\frac{1}{2}} (B^2 + C^2).$$

**Corollary 2.12.** If A = B + iC is the Cartesian decomposition of A, then

$$w_p(B,C) \ge 2^{\frac{1}{p}-1} \|B^2 + C^2\|^{\frac{1}{2}}.$$

And

$$w(A) \ge 2^{\frac{1}{p} - \frac{3}{2}} \|A^*A + AA^*\|^{\frac{1}{2}}.$$

for any  $p \geq 2$ .

*Proof.* The first inequality is obvious. For the second we have  $A^*A + AA^* = 2(B^2 + C^2)$ . Now by using inequality (2.1) the proof is complete.

**Corollary 2.13.** If  $B, C \in \mathbb{B}(\mathcal{H})$ , then for  $p \geq 2$ 

$$w_p(B,C) \ge 2^{\frac{2}{p}-\frac{3}{2}} w^{\frac{1}{2}} (B^2 + C^2).$$

*Proof.* By choosing B + C and B - C instead of B and C in Theorem 2.11 and employing part (i) of Proposition 2.9 we conclude that the desired inequality.  $\Box$ 

The following result providing other bound for  $w_p$  (p > 1) may be stated as follows:

**Proposition 2.14.** Let  $B, C \in \mathbb{B}(\mathcal{H})$ . Then

$$w_p(B,C) \le w_q\left(\frac{B+C}{2},\frac{B-C}{2}\right)$$

for any  $p \ge 2, 1 < q \le 2$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . If 1 , the reverse inequality holds.

*Proof.* Let  $x \in \mathscr{H}$  be a unit vector. Part (a) of Lemma 2.3 implies that

$$|a|^{p} + |b|^{p} \le 2^{\frac{1}{1-q}} \left( |a+b|^{q} + |a-b|^{q} \right)^{\frac{1}{q-1}}$$
.

So

$$(|a|^p + |b|^p)^{\frac{1}{p}} \le 2^{\frac{1}{p(1-q)}} (|a+b|^q + |a-b|^q)^{\frac{1}{p(q-1)}}.$$

Now replacing  $a = \langle Bx, x \rangle$  and  $b = \langle Cx, x \rangle$  in the above inequality we conclude that

$$\left(\left|\langle Bx, x\rangle\right|^{p} + \left|\langle Cx, x\rangle\right|^{p}\right)^{\frac{1}{p}} \le \left(\left|\left\langle \left(\frac{B+C}{2}\right)x, x\right\rangle\right|^{q} + \left|\left\langle \left(\frac{B-C}{2}\right)x, x\right\rangle\right|^{q}\right)^{\frac{1}{q}}.$$
 (2.4)

By taking supremum over  $x \in \mathscr{H}$  with ||x|| = 1 we deduce that

$$w_p(B,C) \le w_q\left(\frac{B+C}{2},\frac{B-C}{2}\right)$$

for any  $p \ge 2, 1 < q \le 2$  with  $\frac{1}{p} + \frac{1}{q} = 1$ .

Corollary 2.15. Inequality (2.4) implies that

$$w_p(B,C) \le \left(w^q\left(\frac{B+C}{2}\right) + w^q\left(\frac{B-C}{2}\right)\right)^{\frac{1}{q}}$$

for any  $1 < q \le 2, p \ge 2$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . Further, if B and C are self-adjoint, then

$$w_p(B,C) \le \frac{1}{2} \left( \|B+C\|^q + \|B-C\|^q \right)^{\frac{1}{q}}.$$

If 1 , the converse inequalities hold.

**Corollary 2.16.** If  $B, C \in \mathbb{B}(\mathcal{H})$ , then

$$w_q\left(\frac{B+C}{2},\frac{B-C}{2}\right) \le 2^{\frac{1}{p}}w_p\left(\frac{B+C}{2},\frac{B-C}{2}\right).$$

for all  $1 with <math>\frac{1}{p} + \frac{1}{q} = 1$ . If  $p \ge 2$ , the above inequality is valid in the opposite direction.

*Proof.* By Proposition 2.14 we have

$$w_q\left(\frac{B+C}{2}, \frac{B-C}{2}\right) \le w_p(B, C).$$

for all  $1 with <math>\frac{1}{p} + \frac{1}{q} = 1$ . Proposition 2.9 follows that

$$w_p(B,C) \le 2^{\frac{1}{p}-1} w_p(B+C,B-C) = 2^{\frac{1}{p}} w_p\left(\frac{B+C}{2},\frac{B-C}{2}\right).$$

We therefore get the desired inequality.

# 3. Inequalities of $w_p$ for *n*-tuples of operators

In this section, we are going to obtain some numerical radius inequalities for *n*-tuples of operators. Some generalization of inequalities in the previous section are also established. According to the definition of numerical radius, we immediately get the following double inequality for  $p \geq 1$ 

$$w_p(T_1,\ldots,T_n) \le \left(\sum_{i=1}^n w^p(T_i)\right)^{\frac{1}{p}} \le \sum_{i=1}^n w(T_i).$$

An application of Holder's inequality gives the next result, which is a generalization of inequality (2.2).

**Theorem 3.1.** Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$  and  $0 \leq \alpha_i \leq 1, i = 1, \ldots n$ , with  $\sum_{i=1}^n \alpha_i = 1$ . Then

$$w_p(T_1,...,T_n) \ge w\left(\alpha_1^{1-\frac{1}{p}}T_1 \pm \alpha_2^{1-\frac{1}{p}}T_2 \pm ... \pm \alpha_n^{1-\frac{1}{p}}T_n\right)$$

for any p > 1.

*Proof.* In the Euclidean space  $\mathbb{R}^n$  with the standard inner product, Holder's inequality

$$\sum_{i=1}^{n} |x_i y_i| \le \left(\sum_{i=1}^{n} |x_i|^p\right)^{\frac{1}{p}} \left(\sum_{i=1}^{n} |y_i|^q\right)^{\frac{1}{q}}$$

holds, where p and q are in the open interval  $(1, \infty)$  with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $(x_1, \dots, x_n)$ ,  $(y_1, \dots, y_n) \in \mathbb{R}^n$ . For  $(y_1, \dots, y_n) = \left(\alpha_1^{1-\frac{1}{p}}, \dots, \alpha_n^{1-\frac{1}{p}}\right)$  we have  $\sum_{n=1}^{n} |z_1 - \frac{1}{2}| = \left(\sum_{n=1}^{n} |z_1 - \frac{1}{2}|^q\right)^{\frac{1}{q}}$ 

$$\sum_{i=1}^{n} \left| \alpha_{i}^{1-\frac{1}{p}} x_{i} \right| \leq \left( \sum_{i=1}^{n} \left| x_{i} \right|^{p} \right)^{\frac{1}{p}} \left( \sum_{i=1}^{n} \left| \alpha_{i}^{1-\frac{1}{p}} \right|^{q} \right)^{\frac{1}{q}}.$$

Thus

$$\left(\sum_{i=1}^{n} |x_i|^p\right)^{\frac{1}{p}} \ge \sum_{i=1}^{n} \left|\alpha_i^{1-\frac{1}{p}} x_i\right|.$$

Choosing  $x_i = |\langle T_i x, x \rangle|, i = 1, \dots n$ , we get

$$\begin{split} \left(\sum_{i=1}^{n} |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}} \\ &\geq \sum_{i=1}^{n} \left| \left\langle \alpha_i^{1-\frac{1}{p}} T_i x, x \right\rangle \right| \\ &\geq \left| \left\langle \alpha_1^{1-\frac{1}{p}} T_1 x, x \right\rangle \pm \left\langle \alpha_2^{1-\frac{1}{p}} T_2 x, x \right\rangle \pm \ldots \pm \left\langle \alpha_n^{1-\frac{1}{p}} T_n x, x \right\rangle \\ &= \left| \left\langle \left( \alpha_1^{1-\frac{1}{p}} T_1 \pm \alpha_2^{1-\frac{1}{p}} T_2 \pm \ldots \pm \alpha_n^{1-\frac{1}{p}} T_n \right) x, x \right\rangle \right|. \end{split}$$

Now the result follows by taking the supremum over all unit vectors in  $\mathscr{H}$ .  $\Box$ 

Now we give another upper bound for the powers of  $w_p$ . This result has several inequalities as special cases, which considerably generalize the second inequality of (1.1).

**Theorem 3.2.** Let  $(T_1, \ldots, T_n), (A_1, \ldots, A_n), (B_1, \ldots, B_n) \in \mathbb{B}(\mathscr{H})^{(n)}$  and let fand g be nonnegative continuous functions on  $[0, \infty)$  satisfying f(t) g(t) = t for all  $t \in [0, \infty)$ . Then

$$w_{p}^{rp}\left(A_{1}^{*}T_{1}B_{1},\ldots,A_{n}^{*}T_{n}B_{n}\right) \leq \frac{1}{2} \left\| \sum_{i=1}^{n} \left( \left[B_{i}^{*}f^{2}\left(|T_{i}|\right)B_{i}\right]^{rp} + \left[A_{i}^{*}g^{2}\left(|T_{i}^{*}|\right)A_{i}\right]^{rp} \right) \right\|$$

for  $p \ge 1$  and  $r \ge 1$ .

*Proof.* Let  $x \in \mathscr{H}$  be a unit vector.

$$\begin{split} \sum_{i=1}^{n} |\langle A_{i}^{*}T_{i}B_{i}x, x\rangle|^{p} \\ &= \sum_{i=1}^{n} |\langle T_{i}B_{i}x, A_{i}x\rangle|^{p} \\ &\leq \sum_{i=1}^{n} ||f\left(|T_{i}|\right)B_{i}x||^{p} ||g\left(|T_{i}^{*}|\right)A_{i}x||^{p} \quad (\text{by Lemma 2.1}(c)) \\ &= \sum_{i=1}^{n} \langle f\left(|T_{i}|\right)B_{i}x, f\left(|T_{i}|\right)B_{i}x\rangle^{\frac{p}{2}} \langle g\left(|T_{i}^{*}|\right)A_{i}x, g\left(|T_{i}^{*}|\right)A_{i}x\rangle^{\frac{p}{2}} \\ &= \sum_{i=1}^{n} \langle B_{i}^{*}f^{2}\left(|T_{i}|\right)B_{i}x, x\rangle^{\frac{p}{2}} \langle A_{i}^{*}g^{2}\left(|T_{i}^{*}|\right)A_{i}x, x\rangle^{\frac{p}{2}} \\ &\leq \sum_{i=1}^{n} \langle \left(B_{i}^{*}f^{2}\left(|T_{i}|\right)B_{i}\right)^{p}x, x\rangle^{\frac{1}{2}} \langle \left(A_{i}^{*}g^{2}\left(|T_{i}^{*}|\right)A_{i}\right)^{p}x, x\rangle^{\frac{1}{2}} \\ &\quad (\text{by Lemma 2.2}(a)) \\ &\leq \sum_{i=1}^{n} \left(\frac{1}{2}\left(\langle \left(B_{i}^{*}f^{2}\left(|T_{i}|\right)B_{i}\right)^{rp} + \left(A_{i}^{*}g^{2}\left(|T_{i}^{*}|\right)A_{i}\right)^{rp}\right)x, x\rangle^{\frac{1}{r}} \\ &\quad (\text{by Lemma 2.2}(a)) \\ &\leq \left(\frac{1}{2}\left\langle\sum_{i=1}^{n} \left(\left(\left(B_{i}^{*}f^{2}\left(|T_{i}|\right)B_{i}\right)^{rp} + \left(A_{i}^{*}g^{2}\left(|T_{i}^{*}|\right)A_{i}\right)^{rp}\right)x, x\rangle^{\frac{1}{r}} \right)^{\frac{1}{r}} \end{split}$$

Thus

$$\left(\sum_{i=1}^{n} |\langle A_i^* T_i B_i x, x \rangle|^p\right)^r$$
  
$$\leq \frac{1}{2} \left\langle \left(\sum_{i=1}^{n} \left( \left( B_i^* f^2\left(|T_i|\right) B_i \right)^{rp} + \left( A_i^* g^2\left(|T_i^*|\right) A_i \right)^{rp} \right) \right) x, x \right\rangle$$

Now the result follows by taking the supremum over all unit vectors in  $\mathscr{H}$ .

Choosing A = B = I, we get.

**Corollary 3.3.** Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$  and let f and g be nonnegative continuous functions on  $[0, \infty)$  satisfying f(t) g(t) = t for all  $t \in [0, \infty)$ . Then

$$w_p^{rp}(T_1,\ldots,T_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( f^{2rp}(|T_i|) + g^{2rp}(|T_i^*|) \right) \right\|$$

for  $p \ge 1$  and  $r \ge 1$ .

Letting  $f(t) = g(t) = t^{\frac{1}{2}}$ , we get.

**Corollary 3.4.** Let  $(T_1, \ldots, T_n), (A_1, \ldots, A_n), (B_1, \ldots, B_n)$  are in  $\mathbb{B}(\mathscr{H})^{(n)}$ . Then

$$w_p^{rp}\left(A_1^*T_1B_1,\ldots,A_n^*T_nB_n\right) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( \left(B_i^* \left|T_i\right| B_i\right)^{rp} + \left(A_i^* \left|T_i^*\right| A_i\right)^{rp} \right) \right\|$$

for  $p \ge 1$  and  $r \ge 1$ .

Corollary 3.5. Let  $(A_1, \ldots, A_n), (B_1, \ldots, B_n) \in \mathbb{B}(\mathscr{H})^{(n)}$ . Then

$$w_p^{rp}(A_1^*B_1,\ldots,A_n^*B_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( |B_i|^{2rp} + |A_i|^{2rp} \right) \right\|$$

for  $p \geq 1$  and  $r \geq 1$ .

Corollary 3.6. Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$ . Then

$$w_p^p(T_1,\ldots,T_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( |T_i|^{2\alpha p} + |T_i^*|^{2(1-\alpha)p} \right) \right\|$$

for  $0 \le \alpha \le 1$ , and  $p \ge 1$ . In particular.

$$w_p^p(T_1,\ldots,T_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( |T_i|^p + |T_i^*|^p \right) \right\|.$$

Corollary 3.7. Let  $B, C \in \mathbb{B}(\mathcal{H})$ . Then

$$w_p^p(B,C) \le \frac{1}{2} \left\| |B|^{2\alpha p} + |B^*|^{2(1-\alpha)p} + |C|^{2\alpha p} + |C^*|^{2(1-\alpha)p} \right\|$$

for  $0 \le \alpha \le 1$ , and  $p \ge 1$ . In particular.

$$w_p^p(B,C) \le \frac{1}{2} |||B|^p + |B^*|^p + |C|^p + |C^*|^p||.$$

The next results are related to some different upper bounds for  $w_p$  for *n*-tuples of operators, which have several inequalities as special cases.

**Proposition 3.8.** Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$ . Then

$$w_p(T_1,\ldots,T_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} \right)^p \right\|^{\frac{1}{p}}$$

for  $0 \le \alpha \le 1$ , and  $p \ge 1$ .

*Proof.* By using the arithmetic-geometric mean, for any unit vector  $x \in \mathscr{H}$  we have

Now the result follows by taking the supremum over all unit vectors in  $\mathcal{H}$ .  $\Box$ 

**Proposition 3.9.** Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}$ . Then

$$w_p(T_1, \dots, T_n) \le \left\| \sum_{i=1}^n \left( \alpha |T_i|^p + (1 - \alpha) |T_i^*|^p \right) \right\|^{\frac{1}{p}}$$

for  $0 \le \alpha \le 1$ , and  $p \ge 2$ .

*Proof.* For every unit vector  $x \in \mathscr{H}$ , we have

$$\begin{split} &\sum_{i=1}^{n} |\langle T_{i}x, x \rangle|^{p} \\ &= \sum_{i=1}^{n} \left( |\langle T_{i}x, x \rangle|^{2} \right)^{\frac{p}{2}} \\ &\leq \sum_{i=1}^{n} \left( \left\langle |T_{i}|^{2\alpha} x, x \right\rangle \left\langle |T_{i}^{*}|^{2(1-\alpha)} x, x \right\rangle \right)^{\frac{p}{2}} \quad \text{(by Lemma 2.1(b))} \\ &\leq \sum_{i=1}^{n} \left\langle |T_{i}|^{\alpha p} x, x \right\rangle \left\langle |T_{i}^{*}|^{(1-\alpha)p} x, x \right\rangle \qquad \text{(by Lemma 2.2(a))} \\ &\leq \sum_{i=1}^{n} \left\langle |T_{i}|^{p} x, x \right\rangle^{\alpha} \left\langle |T_{i}^{*}|^{p} x, x \right\rangle^{(1-\alpha)} \qquad \text{(by Lemma 2.2(b))} \\ &\leq \sum_{i=1}^{n} \left( \alpha \left\langle |T_{i}|^{p} x, x \right\rangle + (1-\alpha) \left\langle |T_{i}^{*}|^{p} x, x \right\rangle \right) \text{(by Lemma 2.1(a))} \\ &\leq \sum_{i=1}^{n} \left\langle \left( \alpha \left| T_{i} \right|^{p} + (1-\alpha) \left| T_{i}^{*} \right|^{p} \right) x, x \right\rangle \\ &= \left\langle \left( \sum_{i=1}^{n} \left( \alpha \left| T_{i} \right|^{p} + (1-\alpha) \left| T_{i}^{*} \right|^{p} \right) x, x \right\rangle. \end{split}$$

Now the result follows by taking the supremum over all unit vectors in  $\mathscr{H}$ .  $\Box$ 

## Remark 3.10. As special cases,

(1) For  $\alpha = \frac{1}{2}$ , we have

$$w_p^p(T_1,\ldots,T_n) \le \frac{1}{2} \left\| \sum_{i=1}^n \left( |T_i|^p + |T_i^*|^p \right) \right\|.$$

(2) For  $B, C \in \mathbb{B}(\mathcal{H}), 0 \leq \alpha \leq 1$ , and  $p \geq 1$ , we have

$$w_p^p(B,C) \le \|\alpha |B|^p + (1-\alpha) |B^*|^p + \alpha |C|^p + (1-\alpha) |C^*|^p\|.$$

In particular,

$$w_p^p(B,C) \le \frac{1}{2} |||B|^p + |B^*|^p + |C|^p + |C^*|^p||.$$

The next result reads as follows.

**Proposition 3.11.** Let  $(T_1, \ldots, T_n) \in \mathbb{B}(\mathscr{H})^{(n)}, 0 \leq \alpha \leq 1, r \geq 1$  and  $p \geq 1$ . Then

$$w_p(T_1,\ldots,T_n) \le \left(\sum_{i=1}^n \left\|\alpha \left|T_i\right|^{2r} + (1-\alpha) \left|T_i^*\right|^{2r}\right\|^{\frac{p}{2r}}\right)^{\frac{1}{p}}.$$

*Proof.* Let  $x \in \mathscr{H}$  be a unit vector.

$$\begin{split} &\sum_{i=1}^{n} |\langle T_{i}x, x \rangle|^{p} \\ &= \sum_{i=1}^{n} \left( |\langle T_{i}x, x \rangle|^{2} \right)^{\frac{p}{2}} \\ &\leq \sum_{i=1}^{n} \left( \left\langle |T_{i}|^{2\alpha} x, x \right\rangle \left\langle |T_{i}^{*}|^{2(1-\alpha)} x, x \right\rangle \right)^{\frac{p}{2}} \quad \text{(by Lemma 2.1(b))} \\ &\leq \sum_{i=1}^{n} \left( \left\langle |T_{i}|^{2} x, x \right\rangle^{\alpha} \left\langle |T_{i}^{*}|^{2} x, x \right\rangle^{(1-\alpha)} \right)^{\frac{p}{2}} \quad \text{(by Lemma 2.2(b))} \\ &\leq \sum_{i=1}^{n} \left( \alpha \left\langle |T_{i}|^{2} x, x \right\rangle^{r} + (1-\alpha) \left\langle |T_{i}^{*}|^{2} x, x \right\rangle^{r} \right)^{\frac{p}{2r}} \text{(by Lemma 2.1(a))} \\ &\leq \sum_{i=1}^{n} \left( \alpha \left\langle |T_{i}|^{2r} x, x \right\rangle + (1-\alpha) \left\langle |T_{i}^{*}|^{2r} x, x \right\rangle \right)^{\frac{p}{2r}} \text{(by Lemma 2.2(a))} \\ &\leq \sum_{i=1}^{n} \left\langle \left( \alpha |T_{i}|^{2r} + (1-\alpha) |T_{i}^{*}|^{2r} \right) x, x \right\rangle^{\frac{p}{2r}} . \end{split}$$

Now the result follows by taking the supremum over all unit vectors in  $\mathscr{H}$ .  $\Box$ 

Remark 3.12. Some special cases can be stated as follows:

(1) For  $\alpha = \frac{1}{2}$ , we have

$$w_p(T_1,\ldots,T_n) \le \left(\frac{1}{2^{\frac{p}{2r}}}\sum_{i=1}^n \left\||T_i|^{2r} + |T_i^*|^{2r}\right\|^{\frac{p}{2r}}\right)^{\frac{1}{p}}.$$

(2) For  $B, C \in \mathbb{B}(\mathscr{H}), 0 \leq \alpha \leq 1$ , and  $p \geq 1$ , we have

$$w_p(B,C) \le \left( \left\| \alpha \left| B \right|^{2r} + (1-\alpha) \left| B^* \right|^{2r} \right\|^{\frac{p}{2r}} + \left\| \alpha \left| C \right|^{2r} + (1-\alpha) \left| C^* \right|^{2r} \right\|^{\frac{p}{2r}} \right)^{\frac{1}{p}}.$$

In particular,

$$w_p(B,C) \le \frac{1}{2^{\frac{1}{2r}}} \left( \left\| |B|^{2r} + |B^*|^{2r} \right\|^{\frac{p}{2r}} + \left\| |C|^{2r} + |C^*|^{2r} \right\|^{\frac{p}{2r}} \right)^{\frac{1}{p}}.$$

#### References

- S. S. Dragomir, Some inequalities for the Euclidean operator radius of two operators in Hilbert Spaces, Linear Algebra and Its Appl, 419 (1) (2006), 256–264.
- 2. K. E. Gustufson and D. K. M. Rao, Numerical Range, Springer-Verlag, New york, 1997.
- O. Hirzallah, F. Kittaneh and K. Shebrawi, Numerical radius inequalities for certain 2 × 2 operator matrices, Integral Equations Operator Theory 71 (2011), no. 1, 129–147.
- O. Hirzallah and F. Kittaneh, Numerical radius inequalities for several operators, Math. Scand. 114 (2014), no. 1, 110-119.
- J. Pečarić, T. Furuta, J. Mićić Hot and Y. Seo, Mond-Pečarić Method in Operator Inequalities, Element, Zagreb, 2005.
- F. Kittaneh, Notes on some inequalities for Hilbert space operators Publ. Res. Inst. Math. Sci. 24 (1988), no. 2, 283–293.
- D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and new inequalities in analysis*, Kluwer Academic Publishers, 1993.
- J. I. Fujii, M. Fujii, M. S. Moslehian, J. E. Pečarić and Y. Seo, Reverses Cauchy–Schwarz type inequalities in pre-inner product C<sup>\*</sup>-modules, Hokkaido Math. J. 40 (2011), 1–17.
- G. Popescu, Unitary invariants in multivariable operator theory, Mem. Amer. Math. Soc., Vol. 200, no 941, 2009.
- A. Salemi and A. Sheikhhosseini, *Matrix Young numerical radius inequalities*, Math. Inequal. Appl. 16 (2013), no. 3, 783–791.
- T. Yamazaki, On upper and lower bounds for the numerical radius and an equality condition, Studia Math. 178 (2007), no. 1, 83–89.
- T. Yamazaki, On numerical range of the Aluthge transformation, Linear Algebra Appl. 341 (2002), 111–117.

<sup>1</sup> DEPARTMENT OF PURE MATHEMATICS, CENTER OF EXCELLENCE IN ANALYSIS ON AL-GEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN

E-mail address: moslehian@um.ac.ir E-mail address: msattari.b@gmail.com

<sup>2</sup> Department of Applied Sciences, Al-Balqa' Applied University Salt, Jordan; Department of Mathematics, College of Science, Qassim University, Qassim, Saudi Arabia

E-mail address: khalid@bau.edu.jo